

Construction of optimal locally recoverable codes and connection with hypergraph

Chaoping Xing* Chen Yuan†

Abstract

Recently, it was discovered by several authors that a q -ary optimal locally recoverable code, i.e., a locally recoverable code archiving the Singleton-type bound, can have length much bigger than $q + 1$. This is quite different from the classical q -ary MDS codes where it is conjectured that the code length is upper bounded by $q + 1$ (or $q + 2$ for some special case). This discovery inspired some recent studies on length of an optimal locally recoverable code. It was shown in [11] that a q -ary optimal locally recoverable code is unbounded for $d = 3, 4$. Soon after, it was proved in [5] that a q -ary optimal locally recoverable code with distance d and locality r can have length $\Omega_{d,r}(q^{\lceil 1+1/(d-3)/2 \rceil})$. Recently, an explicit construction of q -ary optimal locally recoverable codes for distance $d = 5, 6$ was given in [8].

In this paper, we further investigate construction of optimal locally recoverable codes along the line of using parity-check matrices. Inspired by classical Reed-Solomon codes and [8], we equip parity-check matrices with the Vandermonde structure. It turns out that a parity-check matrix with the Vandermonde structure produces an optimal locally recoverable code must obey certain disjoint property for subsets of \mathbb{F}_q . To our surprise, this disjoint condition is equivalent to a well-studied problem in extremal graph theory. With the help of extremal graph theory, we succeed to improve all of the best known results in [5] for $d \geq 7$. In addition, for $d = 6$, we are able to remove the constraint required in [8] that q is even.

1 Introduction

Motivated by applications in distributed and cloud storage systems, locally recoverable codes have been studied extensively in recent years. Informally speaking, a locally recoverable code (LRC for short) is a block code with an additional property called locality. For a locally recoverable code C of length n , dimension k and locality r , it was shown in [4] that the minimum distance $d(C)$ of C is upper bounded by

$$d(C) \leq n - k - \left\lceil \frac{k}{r} \right\rceil + 2. \quad (1)$$

*School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore. This research is supported by the National Research Foundation, Prime Minister's Office, Singapore under its Strategic Capability Research Centres Funding Initiative; and the Singapore MoE Tier 1 grants RG25/16 and RG21/18. Email: xingcp@ntu.edu.sg.

†Centrum Wiskunde & Informatica, Amsterdam, Netherlands. Most of this work was done when the author was visiting the School of Physical and Mathematical Science, Nanyang Technological University, Singapore. Research supported in part by ERC H2020 grant No.74079 (ALGSTRONGCRYPTO). Email: Chen.Yuan@cwi.nl

The bound (1) is called the Singleton-type bound for locally recoverable codes. A code achieving the above bound is usually called optimal.

1.1 Known results

Construction of optimal locally recoverable codes, i.e., block codes archiving the bound (1) is of both theoretical interest and practical importance. This is a challenging task and has attracted great attention in the last few years. In literature, there are a few constructions available and some classes of optimal locally recoverable codes are known. A class of codes constructed earlier and known as pyramid codes [7] are shown to be codes that are optimal. In [13], Silberstein *et al* proposed a two-level construction based on the Gabidulin codes combined with a single parity-check $(r + 1, r)$ code. Another construction [15] used two layers of MDS codes, a Reed-Solomon code and a special $(r + 1, r)$ MDS code. A common shortcoming of these constructions relates to the size of the code alphabet which in all the papers is an exponential function of the code length, complicating the implementation. There was an earlier construction of optimal locally recoverable codes given in [12] with alphabet size comparable to code length. However, the construction in [12] only produces a specific value of the length n , i.e., $n = \lceil \frac{k}{r} \rceil (r + 1)$. Thus, the rate of the code is very close to 1. There are also some existence results given in [12] and [14] with less restriction on locality r . But both results require large alphabet which is an exponential function of the code length.

A recent breakthrough construction was given in [14]. This construction naturally generalizes Reed-Solomon construction which relies on the alphabet of cardinality comparable to the code length n . The idea behind the construction is very nice. The only shortcoming of this construction is restriction on locality r . Namely, $r + 1$ must be a divisor of either $q - 1$ or q , or $r + 1$ is equal to a product of a divisor of $q - 1$ and a divisor of q for certain q , where q is the code alphabet. This construction was extended via automorphism group of rational function fields by Jin, Ma and Xing [9] and it turns out that there are more flexibility on locality and the code length can be $q + 1$. For particular locality such as $r = 2, 3, 5, 7, 11$ or 23 , it was shown that there exist q -ary optimal locally recoverable codes with length up to $q + 2\sqrt{q}$ via elliptic curves [10]. All these results are aimed at the optimal LRC with large distance.

Unlike classical MDS codes, it is surprising to discover that the optimal LRCs can have super-linear code length in alphabet size q . Barg et.al, [1] gave optimal LRCs by using algebraic surfaces of length $n \approx q^2$ when the distance $d = 3$ and $r \leq 4$. This inspired the construction of the optimal LRC with unbounded length and distance $d = 3, 4$ [11]. Furthermore, it was shown in [5] that an optimal LRC with $d \geq 5$ must have length upper bounded in terms of alphabet size q . More precisely, they showed that the length of an optimal q -ary linear LRC with distance $d \geq 5$ and locality r is upper bonded by $O\left(dq^{3+\frac{4}{d-4}}\right)$. As for the lower bound, they presented an explicit construction of optimal LRCs with code length $\Omega_r\left(q^{1+\frac{1}{\lceil (d-3)/2 \rceil}}\right)$ provided that $d \leq r + 2$, where Ω_r means that the implied constant depends on r . One can see that there is still huge gap between the lower bound and the upper bound. Following this discovery, there are several works dedicated to constructing the maximum length of optimal LRCs. The paper [8] aimed at the optimal LRC with small distance $d = 5$ or 6 . In particular, for $d = 6$, the results given in [8] are obtained subject to the constraint that q is even.

1.2 Our results, comparison and a conjecture

The main result of this paper can be summarized as follows.

Theorem 1.1. *Suppose that $r \geq d - 2$ and $(r + 1) | n$. Then*

- (i) *there exists an explicit construction of optimal locally recoverable codes with length $n = q^{2-o(1)}$, minimum distance d and locality r for $d = 7, 8$;*
- (ii) *there exists an explicit construction of optimal locally recoverable codes with length $n = q^{\frac{3}{2}-o(1)}$, minimum distance d and locality r for $d = 9, 10$;*
- (iii) *there exist optimal locally recoverable codes with length $n = \Omega_{r,d} \left(q(q \log q)^{\frac{1}{\lceil (d-3)/2 \rceil}} \right)$, minimum distance d and locality r for $d \geq 11$; and*
- (iv) *there exists an explicit construction of optimal locally recoverable code with length $n = \Omega_{r,d} \left(q^{1+\frac{1}{\lceil (d-3)/2 \rceil}} \right)$, minimum distance d and locality r for a constant $d \geq 11$.*

The first three results are derived from extremal graph theory (see Section 5). The last one is derived from the random arguments (see Section 4).

The first two results improve on the result in [5] which only achieves $n = \Omega(q^{3/2})$ for $d = 7, 8$ and $n = \Omega(q^{4/3})$ for $d = 9, 10$. The third one outperforms the result in [5] by a $(\log q)^{\frac{1}{\lceil (d-3)/2 \rceil}}$ multiplicative factor. In addition, for $d = 6$, we are able to remove the constraint required in [8] that q is even.

Although it was proved in [5] that the length of an optimal locally recoverable code is upper bounded by $q^{3+O(\frac{1}{d})}$, both the constructions in [5] and this paper show from different angles that the length of an optimal locally recoverable code only achieve $q^{1+O(\frac{1}{d})}$. Furthermore, via an upper bound from extremal graph theory, our construction in this paper can achieve at most $O \left(q^{1+\frac{2}{\lceil (d-3)/2 \rceil}} \right)$ (see Section 5). Thus, we make the following conjecture.

Conjecture 1.2. Every optimal locally recoverable code with minimum distance d and locality r has length upper bounded by $q^{1+O(\frac{1}{d})}$.

1.3 Our techniques

For minimum distance $d \geq 7$, the only optimal locally recoverable codes with super-linear code length was given in [5]. In this paper, we present another construction for optimal LRCs for $d \geq 5$. Our idea comes from generalized Reed-Solomon codes where parity-check matrices have the Vandermonde structure. This idea was already employed in [8] for $d = 5, 6$. Like in [8], we divide a parity-check matrix into disjoint blocks, each block with $r + 1$ columns. We require that each block of this matrix has a Vandermonde matrix structure. In order that the parity-check matrix with this structure produces an optimal locally recoverable code, elements in these blocks must satisfy certain disjoint property. It turns out that a necessary and sufficient condition for which a parity-check matrix with this structure produces an optimal locally recoverable code is obtained in terms of certain disjoint property for subsets of \mathbb{F}_q . This condition allows us to relate optimality of a locally recoverable graph code to a well-studied problem in extremal graph theory. With the help of extremal graph theory, we succeed to improve all of the best known results in [5] for $d \geq 7$.

Furthermore, by a random or probabilistic argument, we show an existence result. Moreover, for constant d the probabilistic method for the existence result can be converted into a deterministic algorithm via method of conditional probabilities. Thus, we obtain an algorithmic construction in polynomial time, i.e., Theorem 1.1(iv). The result of Theorem 1.1(iv) matches the result given in [5]. However, our parity-check matrix is more structured and this may lead to some other applications.

1.4 Organization

The paper is organized as follows. In Section 2, we briefly introduce locally recoverable codes and some basic notations on graph theory. Section 3 presents a necessary and sufficient condition for which a Vandermonde-type parity-check matrix produces an optimal locally recoverable code in terms of certain disjoint property for subsets of \mathbb{F}_q . In Section 4, we first show an existence result via a probabilistic method. Then this probabilistic method is converted into an algorithmic construction in polynomial time. Finally in Section 5, we show that the necessary and sufficient condition derived in Section 2 is equivalent to a central problem in extremal graph theory. By applying the known results from extremal graph theory, we obtain the desired results.

2 Preliminaries

2.1 Locally recoverable codes

Let q be a prime power and \mathbb{F}_q be the finite field with q elements and denote by $[n]$ the set $\{1, 2, \dots, n\}$. In this paper, we consider linear locally recoverable codes only. An $[n, k, d]$ linear code C is a k -dimensional subspace of \mathbb{F}_q^n with minimum (Hamming) distance d . The (Euclidean) dual code of C , denoted by C^\perp , is defined by $C^\perp = \{\mathbf{b} \in \mathbb{F}_q^n : \mathbf{c} \cdot \mathbf{b} = 0 \text{ for all } \mathbf{c} \in C\}$, where $\mathbf{c} \cdot \mathbf{b}$ denotes the standard inner product of the two vectors \mathbf{b} and \mathbf{c} .

Informally speaking, a block code is said with locality r if every coordinate of a given codeword can be recovered by accessing at most r other coordinates of this codeword. There are several equivalent definitions of locally recoverable codes. A formal definition of a locally recoverable code with locality r is given as follows.

Definition 1. A q -ary block code C of length n is called a locally recoverable code or locally repairable code (LRC for short) with locality r if for any $i \in [n]$, there exists a subset $R_i \subseteq [n] \setminus \{i\}$ of size r such that for any $\mathbf{c} = (c_1, \dots, c_n) \in C$, c_i can be recovered by $\{c_j\}_{j \in R_i}$, i.e., for any $i \in [n]$, there exists a subset $R_i \subseteq [n] \setminus \{i\}$ of size r such that for any $\mathbf{u}, \mathbf{v} \in C$, $\mathbf{u}_{R_i \cup \{i\}} = \mathbf{v}_{R_i \cup \{i\}}$ if and only if $\mathbf{u}_{R_i} = \mathbf{v}_{R_i}$. The set R_i is called a recovering set of i .

In literature, there are various definitions for locally recoverable code and all of them are equivalent. For example, we have the following two definitions that are equivalent to Definition 1. For the sake of completeness, we give a proof.

Lemma 2.1. *A q -ary code C of length n is a locally recoverable code if and only if one of the followings holds.*

- (i) *For any $i \in [n]$, there exists a subset $R_i \subseteq [n] \setminus \{i\}$ of size r such that position i of every codeword $\mathbf{c} \in C$ is determined by \mathbf{c}_{R_i} , i.e., there is a function $f_i(x_1, \dots, x_r)$ (independent*

of \mathbf{c} and only dependent on i) such that $c_i = f_i(\mathbf{c}_{R_i})$, where \mathbf{c}_{R_i} stands for the projection of \mathbf{c} at R_i .

(ii) For any $i \in [n]$, there exists a subset $R_i \subseteq [n] \setminus \{i\}$ of size r such that

$$C_{R_i}(i, \alpha) \cap C_{R_i}(i, \beta) = \emptyset$$

for any $\alpha \neq \beta \in \mathbb{F}_q$, where $C(i, \alpha) = \{\mathbf{c} \in C : c_i = \alpha\}$ and $C_{R_i}(i, \alpha)$ denotes the projection of $C(i, \alpha)$ on R_i .

Proof. Let C be a q -ary code of length n . Assume that the condition in Definition 1 is satisfied. For every $i \in [n]$, consider the subset $\{(c_i, \mathbf{c}_{R_i}) : \mathbf{c} \in C\}$ of \mathbb{F}_q^{r+1} . As \mathbf{c}_{R_i} determines c_i , we can find a function $f_i(x_1, \dots, x_r)$ from \mathbb{F}_q^r to \mathbb{F}_q (independent of \mathbf{c}) such that $c_i = f_i(\mathbf{c}_R)$ for every $\mathbf{c} \in C$. Conversely, if (i) holds, it is clear that C is a locally recoverable code with locality r .

Now assume that C is a locally recoverable code with locality r , i.e., (i) holds. Suppose that, for some $i \in [n]$ and all subsets $R_i \subseteq [n] \setminus \{i\}$ of size r , $C_{R_i}(i, \alpha) \cap C_{R_i}(i, \beta) \neq \emptyset$ for $\alpha \neq \beta \in \mathbb{F}_q$, i.e., there exist two codewords $\mathbf{u}, \mathbf{v} \in C$ such that $u_i = \alpha$, $v_i = \beta$ and $\mathbf{u}_{R_i} = \mathbf{v}_{R_i}$. This is a contradiction to the fact that $\alpha = u_i = f_i(\mathbf{u}_{R_i}) = f_i(\mathbf{v}_{R_i}) = v_i = \beta$. Conversely, assume that (ii) holds. we claim that $\mathbf{u}_{R_i \cup \{i\}} = \mathbf{v}_{R_i \cup \{i\}}$ if and only if $\mathbf{u}_{R_i} = \mathbf{v}_{R_i}$. Otherwise, one would have two codewords $\mathbf{u}, \mathbf{v} \in C$ such that $\mathbf{u}_{R_i} = \mathbf{v}_{R_i}$ and $u_i \neq v_i$. This implies that $C_{R_i}(i, u_i) \cap C_{R_i}(i, v_i)$ contains \mathbf{u}_{R_i} . This is a contradiction. \square

The Singleton (upper) bound in (1) is given in terms of minimum distance d . For convenience of this paper, we can rewrite this bound in terms of dimension k .

Lemma 2.2. *Let n, k, d, r be positive integers with $(r+1)|n$. If the Singleton-type bound (1) is achieved, then*

$$n - k = \frac{n}{r+1} + d - 2 - \left\lfloor \frac{d-2}{r+1} \right\rfloor. \quad (2)$$

Conversely, if $d-2 \not\equiv r \pmod{r+1}$ and the equality (2) is satisfied, then the Singleton-type bound (1) is achieved.

The proof is straightforward and can be found in [5].

Remark 1. If $d-2 \equiv r \pmod{r+1}$, one can verify that (2) implies that $r|k$. In this case, by [4, Corollary 10] one cannot achieve the Singleton-type bound (1) with equality and one must have $d \leq n - k - \left\lceil \frac{k}{r} \right\rceil + 1$. Therefore in this case we say an LRC attaining this latter bound as optimal.

Corollary 2.3. *If $r \geq d-2$, then an $[n, k, d]$ locally recoverable code with locality r is optimal if*

$$n - k - \frac{n}{r+1} = d - 2. \quad (3)$$

Proof. As $r \geq d-2$, $\left\lfloor \frac{d-2}{r+1} \right\rfloor = 0$. Hence, (2) and (3) are equivalent. \square

The locality of a locally recoverable code C can be determined by a parity-check matrix of C as follows. Assume that $(r+1)|n$. Let $m = \frac{n}{r+1}$ and let D_i be $(n-k-m) \times (r+1)$ matrices. Put

$$H = \left(\begin{array}{c|c|c|c} \mathbf{1} & \mathbf{0} & \cdots & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{1} & \cdots & \mathbf{0} \\ \hline \vdots & \vdots & \ddots & \vdots \\ \hline \mathbf{0} & \mathbf{0} & \cdots & \mathbf{1} \\ \hline D_1 & D_2 & \cdots & D_m \end{array} \right), \quad (4)$$

where $\mathbf{1}$ and $\mathbf{0}$ stand for the all-one row vector and the zero row vector of length $r+1$, respectively. Let C be the code with H as a parity-check matrix. Then it is clear that the dimension of C is at least k . Furthermore, we claim that the locality of C is r . Indeed, let $\mathbf{c} = (c_1, c_2, \dots, c_n)$ be a codeword of C , then $\sum_{j=1+(r+1)i}^{(r+1)(i+1)} c_j = 0$ for $0 \leq i \leq m-1$ as $H\mathbf{c}^T = \mathbf{0}$. Hence, a coordinate c_j with $j \in \{1+(r+1)i, \dots, (r+1)(i+1)\}$ for some $0 \leq i \leq m-1$ can be repaired by \mathbf{c}_{R_j} with $R_j = \{1+(r+1)i, \dots, (r+1)(i+1)\} \setminus \{j\}$.

In conclusion, to see if a linear code C with a parity-check matrix H of the form (4) is an optimal locally recoverable code, it is sufficient to check if the minimum distance of C satisfies (3) for $r \geq d-2$.

2.2 Graphs

A undirected graph G is a pair $G = (V, E)$, where V is a finite set and E is a set consisting of some subsets of size 2 of V . An element of V is called a vertex and an element of E is called an edge. A subgraph G' of a graph G is a graph whose vertex set and edge set are subsets of those of G . We say that G has a cycle (v_1, \dots, v_m) if $\{v_i, v_{i+1}\} \in E$ for $i = 1, \dots, m-1$ and $\{v_m, v_1\} \in E$. The following Lemma 2.4 provides a simple but useful way to determine if G contains a cycle. The proof can be found in any textbook about graph theory (see [3] for instance).

Lemma 2.4. *An undirected graph G contains a cycle if $|E| \geq |V|$.*

Apart from the above usual definition of graph, we also require some results on hypergraph in this paper. A hypergraph is a generalization of a graph in which an edge can join any number of vertices. Formally, a hypergraph H is a pair $H = (X, E)$ where X is a set of elements called vertices, and E is a set of non-empty subsets of X called hyperedges or edges. Therefore, E is a subset of $2^X \setminus \{\emptyset\}$, where 2^X stands for the power set of X .

Definition 2 (*r*-uniform Hypergraph (or *r*-hypergraph for short)). *A hypergraph $H = (X, E)$ is called *r*-uniform if every hyperedge in E has size *r*. In other words, every hyperedge of an *r*-uniform hypergraph connects exactly *r* vertices.*

There are several ways to define cycles in a hypergraph that coincide with the definition of cycles in the usual graph. In this paper, we use the Berge cycle as the generalization of cycles in the usual graph.

Definition 3 (Berge cycle). *A *r*-uniform hypergraph $H = (X, E)$ contains a Berge *k*-cycle (v_1, \dots, v_k) if there exist *k* hyperedges $e_1, \dots, e_k \in E$ such that $\{v_{i-1}, v_i\} \subseteq e_i$ for $i = 2, \dots, k$ and $\{v_1, v_k\} \subseteq e_1$.*

3 A criterion on minimum distance

It follows from Corollary 2.3 that for $d \leq r + 2$, a locally recoverable code with parity-check matrix H in (4) is optimal provided that any $d - 1$ columns of H are linearly independent and each D_i is a $(d - 2) \times (r + 1)$ matrix.

Let \mathbb{F}_q be a finite field and put $m = \frac{n}{r+1}$. Assume that A_1, \dots, A_m are subsets of \mathbb{F}_q , each of size $r + 1$. Let $A_i = \{a_{i,1}, \dots, a_{i,r+1}\}$ for $i = 1, \dots, m$. Let $\mathbf{a}_{i,j} = (a_{i,j}, a_{i,j}^2, \dots, a_{i,j}^{d-2})$ and put $D_i = (\mathbf{a}_{i,1}^T, \mathbf{a}_{i,2}^T, \dots, \mathbf{a}_{i,r+1}^T)$. Thus, D_i is a Vandermonde-type matrix. Let $\mathbf{e}_1, \dots, \mathbf{e}_m$ be the standard basis of vector space \mathbb{F}_q^m , i.e., all components of \mathbf{e}_i are 0 except that the i -th component is 1. Then, we can rewrite H as follow.

$$H = \begin{pmatrix} \mathbf{e}_1^T & \cdots & \mathbf{e}_1^T & \cdots & \mathbf{e}_m^T & \cdots & \mathbf{e}_m^T \\ \mathbf{a}_{1,1}^T & \cdots & \mathbf{a}_{1,r+1}^T & \cdots & \mathbf{a}_{m,1}^T & \cdots & \mathbf{a}_{m,r+1}^T \end{pmatrix}. \quad (5)$$

We now present a sufficient and necessary condition under which any $d - 1$ columns of the matrix H in (5) are linearly independent.

Theorem 3.1. *For $d \geq 5$, then any $d - 1$ columns of H defined in (5) are linearly independent if and only if $|\bigcup_{i \in S} A_i| \geq r|S| + 1$ for any $S \subseteq [m]$ of size no more than $t = \lfloor \frac{d-1}{2} \rfloor$.*

Proof. We first prove the “if” direction. Let $\mathbf{h}_{i,j}$ be the (i, j) th column of H , i.e., $\mathbf{h}_{i,j} = (\mathbf{e}_i, \mathbf{a}_{i,j})^T$ for $1 \leq i \leq m$ and $1 \leq j \leq r + 1$. Choose any $d - 1$ columns $\{\mathbf{h}_{i,j}\}_{1 \leq i \leq m; j \in S_i}$ of H , where S_i are subsets of $[r + 1]$ satisfying $\sum_{i=1}^m |S_i| = d - 1$. Let H' be the $(n - k - m) \times (d - 1)$ matrix consisting of these $d - 1$ columns. We are going to show that H' has rank $d - 1$. We assume that S_i is either empty or of size at least 2. Otherwise, the only column selected from D_i with $|S_i| = 1$ must be linearly independent from the rest $d - 2$ columns. We can consider the linear independence of the rest $d - 2$ columns instead. Now, we assume that there are at most t non-empty sets S_i . Let $A = \{a_{i,j}\}_{1 \leq i \leq m; j \in S_i}$. Assume that $A = \{a_1, \dots, a_s\}$ has s distinct elements. If $s = d - 1$, then by elementary row operations, one can find a $(d - 1) \times (d - 1)$ Vandermonde submatrix of the form

$$\begin{pmatrix} 1 & 1 & \cdots & 1 \\ \mathbf{a}_1^T & \mathbf{a}_2^T & \cdots & \mathbf{a}_{d-1}^T \end{pmatrix}$$

of H' , where $\mathbf{a}_i = (a_i, a_i^2, \dots, a_i^{d-2})$. Thus, the rank of H' is $d - 1$.

We proceed to the case where $s < d - 1$. By permuting the columns of H' , we obtain a matrix of the following form:

$$H_1 = \left(\begin{array}{ccc|ccc} \mathbf{e}_{i_1}^T & \mathbf{e}_{i_2}^T & \cdots & \mathbf{e}_{i_s}^T & \mathbf{e}_{i_{s+1}}^T & \cdots & \mathbf{e}_{i_{d-1}}^T \\ \mathbf{a}_1^T & \mathbf{a}_2^T & \cdots & \mathbf{a}_s^T & \mathbf{a}_{s+1}^T & \cdots & \mathbf{a}_{d-1}^T \end{array} \right),$$

where $1 \leq i_1 \leq i_2 \leq \dots \leq i_{d-1} \leq m$ and $\{a_{s+1}, \dots, a_{d-1}\}$ is a subset of A . Thus, a_j belongs to A_{i_j} for $1 \leq i \leq d - 1$. By elementary column operations, we can erase \mathbf{a}_{s+i}^T since it also appears in one of the first s columns. Hence, H_1 is equivalent to

$$H_2 = \left(\begin{array}{ccc|ccc} \mathbf{e}_{i_1}^T & \mathbf{e}_{i_2}^T & \cdots & \mathbf{e}_{i_s}^T & \mathbf{e}_{i_{s+1}}^T - \mathbf{e}_{k_{s+1}}^T & \cdots & \mathbf{e}_{i_{d-1}}^T - \mathbf{e}_{k_{d-1}}^T \\ \mathbf{a}_1^T & \mathbf{a}_2^T & \cdots & \mathbf{a}_s^T & \mathbf{0}^T & \cdots & \mathbf{0}^T \end{array} \right),$$

where $\{k_{s+1}, \dots, k_{d-1}\}$ is a subset of $\{i_1, \dots, i_s\}$. Since H_2 is an upper left triangular block matrix, showing that H_2 is a full-rank matrix is equivalent to showing both $(\mathbf{a}_1^T, \mathbf{a}_2^T, \dots, \mathbf{a}_s^T)$ and $(\mathbf{e}_{i_{s+1}}^T - \mathbf{e}_{k_{s+1}}^T, \dots, \mathbf{e}_{i_{d-1}}^T - \mathbf{e}_{k_{d-1}}^T)$ have full rank. Note that $(\mathbf{a}_1^T, \mathbf{a}_2^T, \dots, \mathbf{a}_s^T)$ is a $(d-2) \times s$ Vandermonde matrix and hence it has full rank s . It remains to show that $\mathbf{e}_{i_{s+1}} - \mathbf{e}_{k_{s+1}}, \dots, \mathbf{e}_{i_{d-1}} - \mathbf{e}_{k_{d-1}}$ are linearly independent. Suppose they were linearly dependent. Then there exist elements $\lambda_{s+1}, \dots, \lambda_{d-1} \in \mathbb{F}_q$ which are not all zero such that

$$\sum_{i=s+1}^{d-1} \lambda_i (\mathbf{e}_{j_i} - \mathbf{e}_{k_i}) = 0.$$

Let P be the subset of $\{s+1, \dots, d-1\}$ such that $\lambda_i \neq 0$ if and only if $i \in P$. It follows that

$$\sum_{i \in P} \lambda_i (\mathbf{e}_{j_i} - \mathbf{e}_{k_i}) = 0. \quad (6)$$

Let $U = \{j_i : i \in P\}$, $V = \{k_i : i \in P\}$ and $W = U \cup V$. As both U and V are subsets of $\{i \in [m] : |S_i| \geq 2\}$, we have $|W| \leq t = \lfloor \frac{d-1}{2} \rfloor$. Since λ_i is nonzero for all $i \in P$, every $\ell \in W$ must appear at least twice in the multiset consisting of elements of U and V . Otherwise, \mathbf{e}_ℓ could not be cancelled in (6). This implies $|W| \leq |P|$.

On the other hand, for each $a_i \in A$, there is exactly one subset A_{k_i} containing a_i since the first s columns have s distinct \mathbf{a}_i . Furthermore, let $t_i = |\{\ell \in U : a_i \in A_\ell\}|$. It follows that $\sum_{a_i \in A} t_i = |P|$ and a_i belongs to $t_i + 1$ subsets in $\{A_\ell : \ell \in W\}$. This implies

$$\left| \bigcup_{\ell \in W} A_\ell \right| \leq \sum_{\ell \in W} |A_\ell| - \sum_{i=1}^s t_i = (r+1)|W| - |P|.$$

Combining with the condition $|\bigcup_{\ell \in W} A_\ell| \geq r|W| + 1$ forces $|W| \geq |P| + 1$. A contradiction occurs and we complete the proof of the “if” direction.

We proceed to the “only if” direction. First, we claim that $|A_i \cap A_j| \leq 1$ for any $i \neq j$. Otherwise, we may assume that $A_i \cap A_j$ contains two distinct elements a_1 and a_2 . Thus, H contains the four linearly dependent columns $(\mathbf{e}_i, \mathbf{a}_1)^T, (\mathbf{e}_i, \mathbf{a}_2)^T, (\mathbf{e}_j, \mathbf{a}_1)^T$ and $(\mathbf{e}_j, \mathbf{a}_2)^T$.

We prove the “only if” part by contradiction. Without loss of generality, we assume that the first s subsets A_1, \dots, A_s do not satisfy the condition, i.e. $|\bigcup_{i=1}^s A_i| \leq sr$, where s satisfies $s \leq t$. Define an undirected graph $G = ([s], E)$ such that $\{i, j\} \in E$ if and only if $A_i \cap A_j \neq \emptyset$. By inclusion-exclusion principle, we have

$$rs \geq \left| \bigcup_{i=1}^s A_i \right| \geq \sum_{i=1}^s |A_i| - \sum_{(i,j) \in E} 1 = s(r+1) - |E|.$$

This implies $|E| \geq s$. By Lemma 2.4, there exists a cycle in this undirected graph. Without loss of generality, we may assume that $(1, \dots, \ell)$ is a cycle, i.e., $\{i, i+1\} \in E$ for $i = 1, \dots, \ell-1$ and $\{\ell, 1\} \in E$. By the definition of E , A_i and A_{i+1} contains a common element $\{a_{j_i}\}$. Then, we can pick two columns $(\mathbf{e}_i, \mathbf{a}_{j_{i-1}})^T$ ¹ and $(\mathbf{e}_i, \mathbf{a}_{j_i})^T$ from the i -th block D_i for $i = 1, \dots, \ell$. These 2ℓ columns are linearly dependent since

$$\sum_{i=1}^{\ell} \left((\mathbf{e}_i, \mathbf{a}_{j_{i-1}}) - (\mathbf{e}_i, \mathbf{a}_{j_i}) \right) = \sum_{i=1}^{\ell} (0, \mathbf{a}_{j_{i-1}} - \mathbf{a}_{j_i}) = \mathbf{0}.$$

¹Define $\mathbf{a}_{j_0} = \mathbf{a}_{j_\ell}$ for simplicity.

The proof is completed. □

By Theorem 3.1, we immediately obtain the following result.

Theorem 3.2. *If $t = \lfloor \frac{d-1}{2} \rfloor \geq 2$ and $(r+1)|n$, then there exists a q -ary optimal linear LRC with length n , minimum distance d and locality r provided that there are $m = \frac{n}{r+1}$ sets $A_1, \dots, A_m \subseteq \mathbb{F}_q$ such that*

$$\begin{aligned} |A_i| &= r+1 && \text{for } 1 \leq i \leq m, \\ |\bigcup_{i \in S} A_i| &\geq |S|r+1 && \text{for any } S \subseteq [m] \text{ of size at most } t. \end{aligned} \quad (7)$$

Remark 2. As we do not require that q is even, the constraint required in [8] that q is even for $d = 6$ can be removed.

4 Random and algorithmic constructions

In the previous section, we converted construction of optimal LRCs into a problem of finding subsets of \mathbb{F}_q satisfying (7). In this section, we first present a random construction of subsets satisfying (7). In addition, we can derandomize this random construction into a deterministic construction in polynomial time if d is constant.

The case $t = 2$, i.e., $d = 5$ and 6 , is equivalent to the design of constant weight codes [8]. In this section, we assume $t \geq 3$. Since the algebraic structure is not important for the union of set. We replace \mathbb{F}_q with $[q]$ from now on.

Theorem 4.1. *There exist $m = \left\lceil \frac{q^{1+\frac{1}{t-1}}}{2t^2(r+1)^{2+\frac{1}{t-1}}} \right\rceil$ sets A_1, \dots, A_m satisfying (7) provided q is large enough.*

Proof. Let $X_i = \{x_{i,1}, \dots, x_{i,r+1}\}$, $i = 1, \dots, 2m$ be the set picked uniformly at random over all $r+1$ -sized subsets of $[q]$. Define the binary random variable Y_S such that $Y_S = 1$ if $|\bigcup_{i \in S} X_i| \leq |S|r$ and 0 otherwise. Our goal is to bound the expectation $E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S \right]$. Without loss of generality, we may assume that $S = \{1, \dots, a\}$ for some $1 < a \leq t$. We order the random variables in X_i , $i = 1, \dots, a$, i.e., $x_{1,1}, \dots, x_{1,r+1}, \dots, x_{a,1}, \dots, x_{a,r+1}$. We want to bound the probability of the event $Y_S = 1$, i.e., at least a elements repeated in this sequence. Given an element $x_{i,j}$, the probability that $x_{i,j} \neq x_{i',j'}$ for some $x_{i',j'}$ prior to $x_{i,j}$ is at least $1 - \frac{(i-1)(r+1)+j}{q} \geq 1 - \frac{a(r+1)}{q}$. Taking over all sets of size at least a in this sequence, the probability of $Y_S = 1$ is at most

$$\sum_{i=a}^{a(r+1)} \binom{a(r+1)}{i} \left(\frac{a(r+1)}{q} \right)^i \leq \sum_{i=a}^{a(r+1)} \frac{(a(r+1))^i}{i!} \left(\frac{a(r+1)}{q} \right)^i \leq \frac{1.1}{a!} \left(\frac{a^2(r+1)^2}{q} \right)^a.$$

for $q \geq 10a^2(r+1)^2$. It follows that

$$\begin{aligned}
E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S \right] &= \sum_{i=2}^t \sum_{S \subseteq [2m], |S|=i} \Pr[Y_S = 1] \\
&\leq \sum_{i=2}^t \binom{2m}{i} \frac{1.1}{i!} \left(\frac{i^2(r+1)^2}{q} \right)^i \leq \sum_{i=2}^t 1.1 \left(\frac{1}{i!} \right)^2 \left(\frac{2mi^2(r+1)^2}{q} \right)^i \\
&\leq \sum_{i=2}^t 1.1 \left(\frac{1}{i!} \right)^2 \left(\frac{q}{(r+1)^2} \right)^{\frac{i}{t-1}} \leq 1.1 \times 1.5 \left(\frac{1}{t!} \right)^2 \left(\frac{q}{(r+1)^2} \right)^{\frac{t}{t-1}} \\
&\leq \frac{2}{4t^2} \left(\frac{q}{(r+1)^2} \right)^{\frac{t}{t-1}} \leq m.
\end{aligned}$$

for $q \geq t^{2t} 3^t (r+1)$ and $t \geq 3$. The second inequality is due to $\binom{2m}{i} \leq \frac{(2m)^i}{i!}$ and the third inequality is due to

$$\left(\frac{1}{i!} \right)^2 \left(\frac{q}{(r+1)^2} \right)^{\frac{i}{t-1}} \geq 3 \left(\frac{1}{(i-1)!} \right)^2 \left(\frac{q}{(r+1)^2} \right)^{\frac{i-1}{t-1}}.$$

That means there exists $2m(r+1)$ -sized sets A_1, \dots, A_{2m} such that there are at most m subsets $S \subseteq [2m]$ with $|\bigcup_{i \in S} A_i| \leq |S|r$. For each of these m subsets S , remove one set from $A_i, i \in S$. The desired result follows as we remove at most m sets. \square

Theorem 4.1 is an existence proof. However, if t is a constant, it is possible to turn this argument into an algorithm via the method of conditional probabilities.

Theorem 4.2. *There exists a polynomial-time deterministic algorithm to find m sets in Theorem 4.1 provided that t is a constant.*

Proof. We follow the same notation in Theorem 4.1. Let $X_i = \{x_{i,1}, \dots, x_{i,r+1}\}$ be a random set of size $r+1$. Our goal is to minimize $E[\sum_{S \subseteq [2m], |S| \leq t} Y_S]$ by fixing the set X_i one by one. Since

$$\begin{aligned}
E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S \right] &= \sum_{A \subseteq [q], |A|=r+1} E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S | X_1 = A \right] \Pr[X_1 = A] \\
&= \frac{1}{\binom{q}{r+1}} \sum_{A \subseteq [q], |A|=r+1} E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S | X_1 = A \right],
\end{aligned}$$

there exists a set A such that $E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S | X_1 = A \right] \leq E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S \right]$. If $r+1$ is a constant, we only need to enumerate all subsets of size $r+1$ in polynomial time. However, if $r+1$ is not a constant, we enumerate $x_{1,1} \in X_1$ instead of the whole set, i.e., we minimize $E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S | x_{1,1} = a_{1,1} \right]$ for $a_{1,1} \in [q]$. It remains to show how to compute this expectation. Given a subset $S \subseteq [2m]$ of size t , let us show how to compute $E[Y_S | x_{1,1} = a_{1,1}]$. Without loss of generality, we assume $S = \{1, \dots, t\}$. We list $t(r+1)$ random elements

$x_{1,1} = a_{1,1}, x_{1,2}, \dots, x_{1,r+1}, \dots, x_{t,1}, \dots, x_{t,r+1}$. For large enough q , it suffices to compute $E[Y_S | x_{1,1} = a_{1,1}]$ by counting the number of sequences where there are exact t repetitions. There are $\binom{(r+1)t}{t}$ combinations of these t positions. Let $R \subseteq [t] \times [r+1]$ be any set of t positions. We first remove these t positions from the sequence. The remaining tr positions in the sequence must have distinct elements and there are $\prod_{i=0}^{rt-1} (q-i)$ ways to pick these tr elements. Now we assign $1, \dots, rt$ to these tr positions and then determine the rest of sequence. To obtain our final result, we multiply it by $\prod_{i=0}^{rt-1} (q-i)$. For each $(i, j) \in R$, we enumerate all possible choices of $x_{i,j}$, $(i, j) \in R$ and find out the number of combinations that there are exact t repetitions in the resulting sequence. There are at most q^t ways to do the enumeration. Then, we obtain the exact value of $E[Y_S | x_{1,1} = a_{1,1}]$. Observe that there are at most $\sum_{i=2}^t \binom{n}{i}$ subsets S . Thus, this expectation can be computed in polynomial time as t is a constant. We do it $r+1$ times so as to fix all elements in X_1 . Given A_1, \dots, A_k , our goal is to find $X_{k+1} = A_{k+1}$ to minimize the expectation

$$E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S | X_1 = A_1, \dots, X_k = A_k \right] \leq E \left[\sum_{S \subseteq [2m], |S| \leq t} Y_S \right].$$

It can be done in the same way as X_1 is already fixed. After we fix all these $2m$ sets, we will obtain A_1, \dots, A_{2m} with the same property as Theorem 4.1 claims. Then, we enumerate all t -sized subsets $S \subseteq [q]$ and do the same as Theorem 4.1 does. The resulting subsets are the output of our algorithm. The number of these subsets is at least m . Since t is constant, all this operation is done in polynomial time. The proof is completed. \square

The following is a direct consequence of Theorem 3.1 and Theorem 4.1.

Theorem 4.3. *For $d \geq 5$, put $t = \lfloor \frac{d-1}{2} \rfloor$. If $r \geq d-2$, $(r+1)|n$ and q is sufficiently large, then there exists a q -ary $[n, k, d]$ optimal locally recoverable code with locality r and $n \geq \frac{q^{1+\frac{1}{t-1}}}{2t^2(r+1)^{1+\frac{1}{t-1}}}$. The parity matrix of this code has the form of (5). Moreover, if d is a constant, there exists a deterministic algorithm running in polynomial time to construct this code.*

5 The connection with extremal graph theory

To our surprise, it turns out that finding a collection of sets satisfying (7) is equivalent to constructing an $(r+1)$ -uniform hypergraph avoiding the small cycle. The latter is one of the central problems in extremal graph theory and this problem is extremely difficult.

Lemma 5.1. *There exist m sets satisfying (7) if and only if there exists an $(r+1)$ -hypergraph $H = ([q], E)$ with $|E| = m$ that does not have any Berge ℓ -cycles for all $\ell \leq t$.*

Proof. To see the equivalence of these two problems, we define an $(r+1)$ -hypergraph as follows: Let $H = (V, E)$ with $V = [q]$ and $E = \{A_1, \dots, A_m\}$. It is clear that H is an $(r+1)$ -hypergraph. Assume that there exists $k \leq t$ subsets A_{i_1}, \dots, A_{i_k} does not satisfy the condition that $|\bigcup_{j=1}^k A_{i_j}| \geq rk+1$. The same argument in Theorem 3.1 implies that there exists a cycle $(1, 2, \dots, \ell)$ such that $j \in A_{i_j} \cap A_{i_{j+1}}$. That means $\{j-1, j\} \subseteq A_{i_j}$

for $j = 2, \dots, \ell$ and $\{1, \ell\} \subseteq A_{i_1}$. By the definition of Berge cycle, the $(r+1)$ -hypergraph H contains this Berge ℓ -cycle $(1, 2, \dots, \ell)$. On the other hand, assume that there exists a Berge ℓ -cycle in H . Denote the ℓ edges of this cycle $A_{i_1}, \dots, A_{i_\ell}$. The results follows since $|A_{i_j} \cap A_{i_{j+1}}| \geq 1$ for $i = 1, \dots, \ell - 1$ and $|A_{i_1} \cap A_{i_\ell}| \geq 1$. \square

The equivalence of both the problems allow us to make use of known results in this area. Let \mathcal{F} be a family of $r+1$ -hypergraph. Denote by $ex_{r+1}(n, \mathcal{F})$ the maximum number of edges in an $(r+1)$ -hypergraph that does not contain any subgraphs in \mathcal{F} . Denote by BC_k the set of k -cycles. Let $\mathcal{B}_k = \{BC_2, \dots, BC_k\}$. One upper bound on $ex_{r+1}(n, \mathcal{B}_t)$ is obtained by reducing this problem to an $m \times n$ bipartite graph with girth more than $2t$ and apply the result in [6].

Proposition 5.2 ([17]). *$ex_{r+1}(n, \mathcal{B}_t)$ is upper bounded by*

$$(i) \frac{n}{r} \left(\frac{n}{r+1} \right)^{\frac{2}{t-1}} + \frac{n}{r+1} \text{ if } t \text{ is odd,}$$

$$(ii) \frac{n}{r(r+1)} n^{\frac{2}{t}} + \frac{n}{r+1} \text{ if } t \text{ is even.}$$

Since these two problems are equivalent, Proposition 5.2 gives an upper bound on the number m of sets A_i . For $t = 3, 4$, the following two propositions show that this upper bound is asymptotically tight. However, constructing such hypergraph requires sophisticated knowledge in this area which is beyond the scope of this paper. We summarize the results as follows.

Proposition 5.3 ([16]). *There exists explicit construction of $(r+1)$ -hypergraph $H = ([q], E)$ with $|E| = q^{2-o(1)}$ that contains no subgraph in \mathcal{B}_3 .*

Proposition 5.4 (Theorem 23 [17]). *There exists explicit construction of $(r+1)$ -hypergraph $H = ([q], E)$ with $|E| = q^{\frac{3}{2}-o(1)}$ that contains no subgraph in \mathcal{B}_4 .*

Determining the exact value of $ex_{r+1}(n, \mathcal{B}_t)$ for $r \geq 2$ and $t \geq 3$ is extremely difficult. A major open problem in this area is whether $ex_{r+1}(n, \mathcal{B}_t) = \Omega(n^{1+\frac{2}{t}})$. A tighter lower bound for general t can be obtained from H -free random process [2]. The method in [2] can also be applied to hypergraph and add a log factor above the probabilistic method in Theorem 4.1. Again this technique is beyond our scope.

Proposition 5.5 ([16]). *$ex_{r+1}(n, \mathcal{B}_t) = \Omega_{r,t}(n(n \log n)^{\frac{1}{t-1}})$.*

Theorem 1.1 summarizes all above results in the language of codes.

Acknowledgement

We sincerely thank Prof. J. Verstraëte for his linking our condition (7) with the problem in extremal graph theory. He also provided us some references for latest results on extremal graph theory. We would also like to express our great gratitude to Profs. V. Guruswami, Q. Xiang and M. Lu for discussions and help.

References

- [1] Alexander Barg, Kathryn Haymaker, Everett W. Howe, Gretchen L. Matthews, and Anthony Várilly-Alvarado. Locally recoverable codes from algebraic curves and surfaces. *CoRR*, abs/1701.05212, 2017.
- [2] Tom Bohman and Peter Keevash. The early evolution of the h-free process. *Inventiones mathematicae*, 181(2):291–336, Aug 2010.
- [3] Belá Bollobás. *Modern Graph Theory*. Springer, New York, 1998.
- [4] Parikshit Gopalan, Cheng Huang, Huseyin Simitci, and Sergey Yekhanin. On the locality of codeword symbols. *IEEE Trans. Information Theory*, 58(11):6925–6934, 2012.
- [5] Venkatesan Guruswami, Chaoping Xing, and Chen Yuan. How long can optimal locally repairable codes be? In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, APPROX/RANDOM 2018, August 20-22, 2018 - Princeton, NJ, USA*, pages 41:1–41:11, 2018.
- [6] Shlomo Hoory. The size of bipartite graphs with a given girth. *J. Comb. Theory, Ser. B*, 86(2):215–220, 2002.
- [7] Cheng Huang, Minghua Chen, and Jin Li. Pyramid codes: Flexible schemes to trade space for access efficiency in reliable data storage systems. In *Sixth IEEE International Symposium on Network Computing and Applications (NCA 2007), 12 - 14 July 2007, Cambridge, MA, USA*, pages 79–86, 2007.
- [8] Lingfei Jin. Explicit construction of optimal locally recoverable codes of distance 5 and 6 via binary constant weight codes. *CoRR*, abs/1808.04558, 2018.
- [9] Lingfei Jin, Liming Ma, and Chaoping Xing. Construction of optimal locally repairable codes via automorphism groups of rational function fields. *CoRR*, abs/1710.09638, 2017.
- [10] Xudong Li, Liming Ma, and Chaoping Xing. Optimal locally repairable codes via elliptic curves. *CoRR*, abs/1712.03744, 2017.
- [11] Yuan Luo, Chaoping Xing, and Chen Yuan. Optimal locally repairable codes of distance 3 and 4 via cyclic codes. *CoRR*, abs/1801.03623, 2018.
- [12] N. Prakash, Govinda M. Kamath, V. Lalitha, and P. Vijay Kumar. Optimal linear codes with a local-error-correction property. In *Proceedings of the 2012 IEEE International Symposium on Information Theory, ISIT 2012, Cambridge, MA, USA, July 1-6, 2012*, pages 2776–2780, 2012.
- [13] Natalia Silberstein, Ankit Singh Rawat, Onur Ozan Koyluoglu, and Sriram Vishwanath. Optimal locally repairable codes via rank-metric codes. In *Proceedings of the 2013 IEEE International Symposium on Information Theory, Istanbul, Turkey, July 7-12, 2013*, pages 1819–1823, 2013.
- [14] Itzhak Tamo and Alexander Barg. A family of optimal locally recoverable codes. *IEEE Trans. Information Theory*, 60(8):4661–4676, 2014.

- [15] Itzhak Tamo, Dimitris S. Papailiopoulos, and Alexandros G. Dimakis. Optimal locally repairable codes and connections to matroid theory. *IEEE Trans. Information Theory*, 62(12):6661–6671, 2016.
- [16] Jacques Verstraëte. Personal communication.
- [17] Jacques Verstraëte. *Extremal problems for cycles in graphs*, pages 83–116. Springer International Publishing, Cham, 2016.