Finite time blow up for a model of incompressible Navier-Stokes Equations

Zhentao Jin^a, Yi Zhou^{a,*}

^aSchool of Mathematical Sciences, Fudan University, Shanghai 200433, China

Abstract

In [15], T. Tao constructed an averaged Navier-Stokes equations which obey an energy identity. Nevertheless, he proved that smooth solutions can blow up in finite time. This demonstrates that any proposed positive solution to the famous regularity problem for the three dimensional incompressible Navier-Stokes equations which does not use the finer structure of the nonlinearity cannot possibly be successful. In this paper, we propose a very simple model of the Navier Stokes equation which also obey the energy identity and establish a similar result of finite-time blow up . The drawback of our result is that we only work in space dimensions $n \geq 5$.

Keywords: incompressible Navier Stokes equations, energy identity, finite time blow up.

2010 MSC: 35K40;35K55.

1. Introduction

Consider the Cauchy problem of 3D incompressible Navier-Stokes equations

$$\partial_t u + (u \cdot \nabla)u + \nabla p = \Delta u,$$

$$\nabla \cdot u = 0,$$

$$u(0, \cdot) = u_0.$$

(1.1)

^{*}Corresponding Author

Email addresses: ztjin180fudan.edu.cn (Zhentao Jin), yizhou@fudan.edu.cn (Yi Zhou)

It is well-known that if $u_0 \in (L^2(\mathbf{R}^3))^3$, then the smooth solution obeys the following energy identity

$$\frac{1}{2} \|u(t,\cdot)\|_{L^2(\mathbf{R}^3)}^2 + \int_0^t \|\nabla u(\tau,\cdot)\|_{L^2(\mathbf{R}^3)}^2 d\tau = \frac{1}{2} \|u_0\|_{L^2(\mathbf{R}^3)}^2.$$
(1.2)

It's a clay institute's million dollar problem [3] [16] that if the solution of (1.1) is smooth for all time for smooth initial data with finite energy. In [15], T. Tao constructed an averaged Navier-Stokes equations which obey an energy identity. Nevertheless, he proved that smooth solutions can blow up in finite time. This demonstrates that any proposed positive solution to the regularity problem for the three dimensional incompressible Navier-Stokes equations which does not use the finer structure of the nonlinearity cannot possibly be successful. In this paper, we propose a very simple model of the Navier Stokes equation which also obey the energy identity and establish a similar result of finite-time blow up. The drawback of our result is that we only work in space dimensions $n \geq 5$.

We consider the following systems of nonlinear heat equation in space dimensions $n \ge 5$:

$$\begin{cases} \partial_t u + \nabla \cdot (u \otimes u) + \frac{1}{2} \nabla |u|^2 = \Delta u, \\ u(0, \cdot) = u_0. \end{cases}$$
(1.3)

This system of nonlinear heat equations obeys the energy identity. We shall find some initial data u_0 such that the smooth solutions of (1.3) blow up in finite time.

We pause to review previous work in this direction. If we do not require energy identity, then the blow up solutions for various Navier-Stokes type equations have been constructed in the literature. For instance, finite time blow up for a "cheep Navier-Stokes equation" $\partial_t u = \Delta u + \sqrt{-\Delta}(u^2)$ (in which u now a scalar field) was demonstrated in [4] [12]. In a slightly different direction, finite-time blow up was established in [11] for a complexified version of the Navier-Stokes equations, in which the energy identity was again unavailable. Further models of Navier Stokes type, which obey an energy identity, were introduced by Plechac and Sverak in [13] [14], by Katz and Pavlovic [10] and by Hou and Lei [6], [7], [8]. Our main result can be stated as follows

Theorem 1.1. Consider the Cauchy problem of (1.3) in $n \ (n \ge 5)$ space dimensions with initial data

$$u_0(x) = -\frac{x}{|x|}\phi_0(|x|)$$
(1.4)

then the solution will blow up before time t = 1 provided that $\int_0^{+\infty} r^{2.1} e^{-r} \phi_0(r) dr$ is nonegative and sufficiently large.

Noting (1.4), by rotational invariance of the problem, we conclude that the solution has the form

$$u(t,x) = -\frac{x}{|x|}\psi(t,|x|)$$
(1.5)

Then, by a simple computation, we see that ψ satisfies the equations

$$\psi_t - \psi_{rr} - \frac{n-1}{r}\psi_r + \frac{n-1}{r^2}\psi = \frac{3}{2}(\psi^2)_r + \frac{n-1}{r}\psi^2$$
(1.6)

with $\psi = 0$, if r = 0. The philosophy for this kind of model is that many important theorems for the incompressible Navier Stoke equations at present not just hold for the Navier Stokes equations but also hold for any system of nonlinear heat equations which has a quadratic nonlinearality with a derivative or pseudoderivative before the nonlinearity and obeys an energy identity. Our result shows that for such system of heat equations smooth solutions may still blow up in finite time. But then by our philosophy the Caffarelli-Kohn-Nirenberg partial regularity theorem [1] also hold for our systems in three dimensions. This would rule out any finite time blow up at (t_0, r_0) for $r_0 \neq 0$, otherwise u(t, x)would blow up at a two dimensional sphere $|x| = r_0$, which contradicts the fact that singularity set is at most one dimensional. Moreover, at $r = 0, \psi = 0$ so, solutions will never blow up. By a theorem in [5], similar result as Caffarelli-Kohn-Nirenberg partial regularity theorem also hold in four space dimensions which rule out finite time blow up for ψ in four space dimensions. Thus, our blow up theorem in $n \ge 5$ dimensions is sharp. Compare to the averaged Navier Stokes equation of Tao, Our model is too simple and does not has incompressible conditions and only blow up at high dimensions. However, at least compare to "cheap" Navier Stokes equations it has the advantage of energy identity and its proof of the finite time blow up result is even simpler than that of the "cheap" Navier Stokes equation as we will see in the next section.

2. Proof of Theorem 1.1.

We argue by contradiction, assume that we have a smooth solution on the time interval [0,1). Let F(r) be a test function such that F(0) = 0, we multiply (1.6) by F(r) and integration by parts to get

$$\frac{d}{dt} \left(\int_{0}^{+\infty} F(r)\psi(t,r)dr \right)$$

$$= \int_{0}^{+\infty} \left[-\frac{3}{2}F_{r}(r) + \frac{n-1}{r}F(r) \right]\psi^{2}(t,r)dr$$

$$+ \int_{0}^{+\infty} \left(F_{rr}(r) - \frac{n-1}{r}F_{r}(r) \right)\psi(t,r)dr$$
(2.1)

Take $F = r^{p}e^{-r}$, then $F_{r} = pr^{p-1}e^{-r} - r^{p}e^{-r} = p\frac{F}{r} - F$, $F_{rr} = r^{p}e^{-r} - 2pr^{p-1}e^{-r} + p(p-1)r^{p-2}e^{-r} = F - 2p\frac{F}{r} + p(p-1)\frac{F}{r^{2}}$.

We have:

$$\frac{d}{dt} \left(\int_{0}^{+\infty} F(r)\psi(t,r)dr \right) \tag{2.2}$$

$$= \int_{0}^{+\infty} \left[-\frac{3}{2}F_{r} + \frac{n-1}{r}F \right] \psi^{2}dr + \int_{0}^{+\infty} (F_{rr} - \frac{n-1}{r}F_{r})\psi(t,r)dr + \int_{0}^{+\infty} \left[\frac{3}{2}F + (n-1-\frac{3}{2}p)\frac{F}{r} \right] \psi^{2}(t,r)dr + \int_{0}^{+\infty} \left[F + (n-1-2p)\frac{F}{r} + (p^{2}-np)\frac{F}{r^{2}} \right] \psi dr$$

Since:

$$-[F + (n - 1 - 2p)\frac{F}{r} + (p^2 - np)\frac{F}{r^2}]\psi$$

$$\leq \frac{r[F + (p^2 - np)\frac{F}{r^2} + (n - 1 - 2p)\frac{F}{r}]^2}{2F} + \frac{1}{2}\frac{F}{r}\psi^2$$
(2.3)

For p > 2, we have:

$$M_0 = \int_0^\infty \frac{r[F + (p^2 - np)\frac{F}{r^2} + (n - 1 - 2p)\frac{F}{r}]^2}{F} dr < +\infty$$

Thus:

$$\frac{d}{dt} \left(\int_{0}^{+\infty} F(r)\psi(t,r)dr \right)$$

$$\geq \int_{0}^{+\infty} \left[\frac{3}{2}F + \left(n - \frac{3}{2} - \frac{3}{2}p\right) \frac{F}{r} \right] \psi^{2}dr - M_{0}$$
(2.4)

Since $n \ge 5$, we can take $p = 2.1 < \frac{2}{3}n - 1$. Then

$$\frac{d}{dt} \left(\int_0^{+\infty} F(r)\psi(t,r)dr \right)$$

$$\geq \frac{3}{2} \int_0^{+\infty} F\psi^2 dr - M_0$$
(2.5)

Take $X(t) = \int_0^{+\infty} F(r)\psi(t,r)dr$, then

$$X^{2}(t) \leq \int_{0}^{+\infty} F(r)dr \int_{0}^{+\infty} F(r)\psi^{2}(t,r)dr$$
 (2.6)

Since $F = r^p e^{-r}$, $c_0 = \int_0^{+\infty} F(r) dr < +\infty$. We have:

$$\frac{d}{dt}X(t) \ge \frac{3}{2c_0}X^2(t) - M_0;$$
(2.7)

If $X(0) \ge 2M_0$, since we consider $0 \le t < 1$, integrate this inequality from 0

to t, we get:

$$X(t) \ge X(0) - M_0 t + \frac{3}{2c_0} \int_0^t X^2(\tau) d\tau$$

$$\ge \frac{X(0)}{2} + \frac{3}{2c_0} \int_0^t X^2(\tau) d\tau$$
(2.8)

Let $Y(t) = \frac{X(0)}{2} + \frac{3}{2c_0} \int_0^t X^2(\tau) d\tau$, then:

$$\frac{dY}{dt} = \frac{3}{2c_0} X^2(t) \ge \frac{3}{2c_0} Y^2(t)$$

$$Y(0) = \frac{X(0)}{2}$$
(2.9)

At last, we get the inequality:

$$X(t) \ge Y(t) \ge \frac{1}{\frac{2}{X(0)} - \frac{3}{2c_0}t}$$
(2.10)

This means X(t) will be infinite before t=1 if X(0) is sufficiently large, which is a contradiction.

Finally, we remark that if we add one more dimension, it is also possible to build models which have an incompressible conditions. Let $y = (x, z) \in$ $R^n \times [0, \pi/2]$, We consider the initial boundary value problem

$$\begin{cases} \partial_t u + \nabla_x \cdot (u \otimes u) + \nabla_x p = \Delta_y u, \\ \partial_t v - \frac{1}{2} \partial_z |u|^2 + \partial_z p = \Delta_y v, \\ \nabla_x u + \partial_z v = 0, \\ u = 0, v_z = p = 0, z = 0; u = 0, v = 0, z = \pi/2, \\ u(0, \cdot) = u_0, v(0, \cdot) = v_0, (\nabla_x u_0 + \partial_z v_0 = 0). \end{cases}$$

$$(2.11)$$

Let
$$u(t, x, z) = \frac{x}{|x|} \psi(t, |x|, z), v(t, x, z) = v(t, |x|, z), p = \frac{1}{2} |u|^2 + q(t, |x|, z)$$
, then

$$\begin{cases}
\partial_t \psi + \frac{3}{2} (\psi^2)_r + \frac{n-1}{r} \psi^2 + \partial_r q = \psi_{rr} - \frac{n-1}{r} \psi_r + \frac{n-1}{r^2} \psi + \psi_{zz}, \\
\partial_t v + \partial_z q = \Delta_y v, \\
\partial_r \psi + \frac{n-1}{r} \psi + \partial_z v = 0, \\
\psi = 0, v_z = p = 0; z = 0; \psi = 0, v = 0, z = \pi/2 \\
\psi(0, \cdot) = \psi_0, v(0, \cdot) = v_0,
\end{cases}$$
(2.12)

Noting the incompressible condition, $v_z = 0, z = 0, \pi/2$. We may also assume that u, p are even functions with respect to $\pi/2$, v is an odd function with respect to $\pi/2$. Thus, $\psi_z = 0$ at $z = \pi/2$. By the second equation,

$$\left(\partial_t - \Delta_y\right) \left(\partial_r \psi + \frac{n-1}{r}\psi\right) = q_{zz} \tag{2.13}$$

Noting that q is determined only up to an arbitrarily function of r, that is why we assume q = 0 at z = 0. Test the first equation of (2.12) by $F(r) \sin z$, second equation of (2.12) by $F'(r) \cos z$, sum over, we still can prove a blow up result similar to Theorem 1.1, we omit the details.

References

References

- Caffarelli L, Kohn R, Nirenberg L, Partial regularity of suitable weak solutions of the Navier-Stokes equations, *Comm. Pure Appl. Math.*, 35, (1982), no. 6, 771-831.
- [2] A. Cheskidov, Blow-up in finite time for the dyadic model of the Navier-Stokes equations, *Trans. Am. Math. Soc.*, **360**, (2010), 5101-5120.
- [3] Charles L. Fefferman, Existence and smoothness of the Navier-Stokes equation, The millennium prize problems, *Clay Math. Inst.*, Cambridge, MA, 2006, 57-67.

- [4] I. Gallagher, M. Paicu, Remarks on the blow-up of solutions to a toy model for the Navier-Stokes equations, *Proc. Amer. Math. Soc.*, 137 (2009), no. 6, 2075-2083.
- [5] Dong Hongjie, Gu Xumin, Boundary partial regularity for the high dimensional Navier-Stokes equations, J. Funct. Anal., 267 (2014), no. 8, 2606-2637.
- [6] Thomas Y. Hou, Zhen Lei, On the stabilizing effect of convection in threedimensional incompressible flows, *Comm. Pure Appl. Math.*, 62 (2009), no. 4, 501-564.
- [7] Thomas Y. Hou, Zhen Lei, Guo Luo, Shu Wang, and Chen Zou, On finite time singularity and global regularity of an axisymmetric model for the 3D Euler equations, Arch. Ration. Mech. Anal., 212 (2014), no. 2, 683-706.
- [8] T. Y. Hou, R. Li, Blowup or no blowup? The interplay between theory and numerics, *Physica D*, 237 (2008), 1937-1944.
- [9] Thomas Y. Hou, Zuoqiang Shi, and Shu Wang, On singularity formation of a 3D model for incompressible Navier-Stokes equations, Adv. Math., 230 (2012), no. 2, 607-641.
- [10] N. H. Katz, N. Pavlovic, Finite time blow-up for a dyadic model of the Euler equations, *Trans. Am. Math. Soc.*, **357**, (2005), 695-708.
- [11] D. Li, Ya. Sinai , Blowups of complex solutions of the 3d-Navier-Stokes system and renormalization group method, J. Eur. Math. Soc., 10 (2008), no.2, 267-313.
- [12] Stephen Montgomery-Smith, Finite time blow up for a Navier-Stokes like equation, Proc. Amer. Math. Soc., 129 (2001), no. 10, 3025-3029.
- [13] Petr Plechac and Vladimr Sverak, On self-similar singular solutions of the complex Ginzburg-Landau equation, *Comm. Pure Appl. Math.*, 54(2001), no. 10, 1215-1242.

- [14] Petr Plechac and Vladimr Sverak, Singular and regular solutions of a nonlinear parabolic system, *Nonlinearity*, 16(2003), no. 6, 2083-2097.
- [15] T. Tao, Finite time blowup for an averaged three-dimensional Navier-Stokes equation, J. Amer. Math. Soc., 29(2016), no. 3, 601-674.
- [16] T. Tao, Localisation and compactness properties of the Navier-Stokes global regularity problem, Anal. PDE, 6 (2013), no. 1, 25-107.