

Relatively bounded operators and the operator E-norms (addition to arXiv:1806.05668)

M.E. Shirokov*

Abstract

In this brief note we describe relations between the well known notion of a relatively bounded operator and the operator E-norms considered in [arXiv:1806.05668].

We show that the set of all \sqrt{G} -bounded operators equipped with the E-norm induced by a positive operator G is the Banach space of all operators with finite E-norm and that the \sqrt{G} -bound is a continuous seminorm on this space.

We also show that the set of all \sqrt{G} -infinitesimal operators (operators with zero \sqrt{G} -bound) equipped with the E-norm induced by a positive operator G is the completion of the algebra $\mathfrak{B}(\mathcal{H})$ of bounded operators w.r.t. this norm.

Some properties of \sqrt{G} -infinitesimal operators are considered.

1 Introduction

A linear operator A on a Hilbert space \mathcal{H} is called relatively bounded w.r.t. a linear operator B (briefly, B -bounded) if $\mathcal{D}(B) \subseteq \mathcal{D}(A)$ and

$$\|A\varphi\|^2 \leq a^2\|\varphi\|^2 + b^2\|B\varphi\|^2 \quad \forall \varphi \in \mathcal{D}(B) \quad (1)$$

for some nonnegative numbers a and b . The infimum over b for which (1) holds (with some a) is called the B -bound for A . If this B -bound is equal to zero then A is called B -infinitesimal operator (infinitesimally bounded w.r.t. B). These notions are widely used in the modern operator theory, in particular, in analysis of perturbations of unbounded operators in a Hilbert space [3, 8].

The operator E-norm $\|A\|_E^G$ of a bounded operator A on a Hilbert space \mathcal{H} induced by a positive unbounded operator G is introduced¹ in [5] as the maximum of $\sqrt{\text{Tr}A\rho A^*}$ over all states (positive operators with unit trace) ρ such that $\text{Tr}G\rho \leq E$.² This norm was used in [5] to obtain the modification of the Kretschmann-Schlingemann-Werner

*Steklov Mathematical Institute, RAS, Moscow, email:msh@mi.ras.ru

¹As far as I know. I would be grateful for any references.

²The value of $\text{Tr}G\rho$ (finite or infinite) is defined as $\sup_n \text{Tr}P_n G\rho$, where P_n is the spectral projector of G corresponding to the interval $[0, n]$.

theorem³ which quantifies continuity of the Stinespring dilation of a quantum channel w.r.t. the strong convergence topology on the set of channels and the strong operator topology on the set of Stinespring isometries.⁴ These norms are studied in detail in [6], where they are extended to unbounded operators. The extended operator E-norms and the corresponding Banach spaces of unbounded operators have different applications described in [6, Section 5],[7].

In this note we describe relations between the notion of a relatively bounded operator and the operator E-norms extended to unbounded operators.

We show that the set of all \sqrt{G} -bounded operators equipped with the E-norm induced by a positive operator G coincides with the Banach space of all operators with finite E-norm denoted by $\mathfrak{B}_G(\mathcal{H})$ in [6] and that the \sqrt{G} -bound of an operator is a continuous seminorm on $\mathfrak{B}_G(\mathcal{H})$. We obtain an explicit formula for the E-norm $\|A\|_E^G$ in terms of the set of coefficients (a, b) for which (1) holds with $B = \sqrt{G}$ and the expression for the \sqrt{G} -bound of an operator A via $\|A\|_E^G$.

We also show that the set of all \sqrt{G} -infinitesimal operators (operators with zero \sqrt{G} -bound) equipped with the E-norm $\|\cdot\|_E^G$ coincides with the completion of the algebra $\mathfrak{B}(\mathcal{H})$ of bounded operators w.r.t. this norm denoted by $\mathfrak{B}_G^0(\mathcal{H})$ in [6].

2 Definitions and the main result

Let \mathcal{H} be a separable infinite-dimensional Hilbert space, $\mathfrak{B}(\mathcal{H})$ – the algebra of all bounded operators on \mathcal{H} with the operator norm $\|\cdot\|$ and $\mathfrak{T}(\mathcal{H})$ – the Banach space of all trace-class operators on \mathcal{H} with the trace norm $\|\cdot\|_1$ (the Schatten class of order 1) [3, 8]. Let $\mathfrak{S}(\mathcal{H})$ be the set of quantum states – positive operators in $\mathfrak{T}(\mathcal{H})$ with unit trace [2].

Let G be a positive (semidefinite) operator on \mathcal{H} with a dense domain $\mathcal{D}(G)$ such that

$$\inf \{ \|G\varphi\| \mid \varphi \in \mathcal{D}(G), \|\varphi\| = 1 \} = 0. \quad (2)$$

For a given linear (bounded or unbounded) operator A such that $\mathcal{D}(\sqrt{G}) \subseteq \mathcal{D}(A)$ the operator E-norm induced by G is defined in [6] as

$$\|A\|_E^G \doteq \sup \left\{ \sqrt{\text{Tr} A \rho A^*} \mid \rho \in \mathfrak{S}(\mathcal{H}), \text{Tr} G \rho \leq E, \text{rank} \rho < +\infty \right\} \quad (3)$$

where we assume that⁵

$$A \rho A^* \doteq \sum_i |\alpha_i\rangle \langle \alpha_i|, \quad |\alpha_i\rangle = A |\varphi_i\rangle, \quad (4)$$

³The original Kretschmann-Schlingemann-Werner theorem obtained in [4] quantifies continuity of the Stinespring dilation of CP linear maps w.r.t. the diamond norm (*cb*-norm) topology on the set of CP linear maps and the operator norm topology on the set of Stinespring operators.

⁴If G is a unbounded operator with discrete spectrum of finite multiplicity then any of the E-norms $\|\cdot\|_E^G$, $E > 0$, generates the strong operator topology on the unit ball of $\mathfrak{B}(\mathcal{H})$ [5, Proposition 2].

⁵This assumption is made to avoid the notion of adjoint operator.

provided that $\rho = \sum_i |\varphi_i\rangle\langle\varphi_i|$ (by using Schrodinger's mixture theorem (see [1, Ch.8]) it is easy to show that the r.h.s. of (4) does not depend on this decomposition of ρ).⁶ By using purification of a state it is easy to see that

$$\|A\|_E^G = \sup_n \sup \left\{ \|A \otimes I_{\mathcal{H}_n} \varphi\| \mid \varphi \in \mathcal{H} \otimes \mathcal{H}_n, \|\varphi\| = 1, \|\sqrt{G} \otimes I_{\mathcal{H}_n} \varphi\| \leq \sqrt{E} \right\}, \quad (5)$$

where \mathcal{H}_n and $I_{\mathcal{H}_n}$ denote, respectively, a n -dimensional Hilbert space and the unit operator in this space.

For any given operator A the nonnegative nondecreasing function $E \mapsto [\|A\|_E^G]^2$ is concave on \mathbb{R}_+ and tends to $\|A\| \leq +\infty$ as $E \rightarrow +\infty$ [6]. This implies that

$$\|A\|_{E_1}^G \leq \|A\|_{E_2}^G \leq \sqrt{E_2/E_1} \|A\|_{E_1}^G \quad \text{for any } E_2 > E_1 > 0. \quad (6)$$

So, for given G all the norms $\|A\|_E^G$ are equivalent. In particular, if $\|A\|_E^G$ is finite for some $E > 0$ then $\|A\|_E^G$ is finite for all $E > 0$.

It is shown in [6] that the set of all operators A with finite $\|A\|_E^G$ equipped with the norm $\|\cdot\|_E^G$ and naturally defined linear operations is a nonseparable Banach space denoted therein by $\mathfrak{B}_G(\mathcal{H})$.⁷ The completion $\mathfrak{B}_G^0(\mathcal{H})$ of the algebra $\mathfrak{B}(\mathcal{H})$ w.t.r. to the norm $\|\cdot\|_E^G$ is a proper subspace of $\mathfrak{B}_G(\mathcal{H})$ determined by the condition

$$\|A\|_E^G = o(\sqrt{E}) \quad \text{as } E \rightarrow +\infty. \quad (7)$$

If G is a unbounded operator with discrete spectrum of finite multiplicity then the Banach space $\mathfrak{B}_G^0(\mathcal{H})$ is separable and for any $A \in \mathfrak{B}_G^0(\mathcal{H})$ its E-norm can be defined by the simple formula

$$\|A\|_E^G = \sup \left\{ \|A\varphi\| \mid \varphi \in \mathcal{H}, \|\varphi\| = 1, \|\sqrt{G}\varphi\| \leq \sqrt{E} \right\}, \quad (8)$$

which means that the first supremum in (5) is achieved at $n = 1$ and the supremum in (3) can be taken over pure states [6, Theorem 3F]. Due to the assumption (2) the condition $\|\varphi\| = 1$ in (5) and (8) can be replaced by $\|\varphi\| \leq 1$ [6, Proposition 3A].⁸

According to the general definition mentioned in the Introduction an operator A is called relatively bounded w.r.t. the operator \sqrt{G} (briefly, \sqrt{G} -bounded) if $\mathcal{D}(\sqrt{G}) \subseteq \mathcal{D}(A)$ and

$$\|A\varphi\|^2 \leq a^2 \|\varphi\|^2 + b^2 \|\sqrt{G}\varphi\|^2, \quad \forall \varphi \in \mathcal{D}(\sqrt{G}) \quad (9)$$

for some nonnegative numbers a and b . Denote by $\Gamma_{\sqrt{G}}(A)$ the set of all pairs (a, b) for which (9) holds. It is easy to see that $\Gamma_{\sqrt{G}}(A)$ is a closed subset of \mathbb{R}_+^2 . The \sqrt{G} -bound of A (denoted by $b_{\sqrt{G}}(A)$ in what follows) is defined as

$$b_{\sqrt{G}}(A) = \inf \{ b \mid (a, b) \in \Gamma_{\sqrt{G}}(A) \}.$$

⁶For any vector $|\alpha\rangle$ the symbol $|\alpha\rangle\langle\alpha|$ denotes the 1-rank operator mapping a vector $|\beta\rangle$ to $\langle\alpha|\beta\rangle|\alpha\rangle$.

⁷We identify operators coinciding on the set $\mathcal{D}(\sqrt{G})$.

⁸The question about coincidence of (5) and (8) for any positive operator G is open. It is easy to show that this coincidence is equivalent to concavity of the r.h.s. of (8) as a function of E .

Lemma 1. *A pair (a, b) belongs to the set $\Gamma_{\sqrt{G}}(A)$ if and only if $\|A\|_E^G \leq \sqrt{a^2 + b^2E}$ for all $E > 0$.*

Proof. If $\|A\|_E^G \leq \sqrt{a^2 + b^2E}$ then definition (5) implies that

$$\|A\varphi\| \leq \|A\|_{\|\sqrt{G}\varphi\|^2}^G \leq \sqrt{a^2 + b^2\|\sqrt{G}\varphi\|^2}$$

for any unit vector φ in $\mathcal{D}(\sqrt{G})$. Hence $(a, b) \in \Gamma_{\sqrt{G}}(A)$.

If $(a, b) \in \Gamma_{\sqrt{G}}(A)$ then it is easy to show that $(a, b) \in \Gamma_{\sqrt{G} \otimes I_{\mathcal{H}_n}}(A \otimes I_{\mathcal{H}_n})$, where \mathcal{H}_n is a n -dimensional Hilbert space, for any n [8, Theorem 7.1.20]. Hence

$$\sup \left\{ \|A \otimes I_{\mathcal{H}_n} \varphi\| \mid \varphi \in \mathcal{H} \otimes \mathcal{H}_n, \|\varphi\| = 1, \|\sqrt{G} \otimes I_{\mathcal{H}_n} \varphi\| \leq \sqrt{E} \right\} \leq \sqrt{a^2 + b^2E}$$

for any n and $E > 0$. So, definition (5) implies that $\|A\|_E^G \leq \sqrt{a^2 + b^2E}$. \square

Proposition 1. A) *The Banach space $\mathfrak{B}_G(\mathcal{H})$ coincides (as a set) with the set of all \sqrt{G} -bounded operators. If $A \in \mathfrak{B}_G(\mathcal{H})$ and $E > 0$ then*

$$\|A\|_E^G = \inf \left\{ \sqrt{a^2 + b^2E} \mid (a, b) \in \Gamma_{\sqrt{G}}(A) \right\} \quad \text{and} \quad b_{\sqrt{G}}(A) = \lim_{E \rightarrow +\infty} \|A\|_E^G / \sqrt{E}.$$

The limit in the last formula can be replaced by the infimum over all $E > 0$.

B) *The completion $\mathfrak{B}_G^0(\mathcal{H})$ of $\mathfrak{B}_G(\mathcal{H})$ w.r.t. the norm $\|\cdot\|_E^G$ coincides (as a set) with the set of all \sqrt{G} -infinitesimal operators, i.e. operators with the \sqrt{G} -bound equal to 0.*

C) *The function $b_{\sqrt{G}}(\cdot)$ is a continuous seminorm on $\mathfrak{B}_G(\mathcal{H})$ s.t. $b_{\sqrt{G}}^{-1}(0) = \mathfrak{B}_G^0(\mathcal{H})$. Quantitatively,*

$$|b_{\sqrt{G}}(A) - b_{\sqrt{G}}(B)| \leq b_{\sqrt{G}}(A - B) \leq \|A - B\|_E^G / \sqrt{E} \quad (10)$$

for arbitrary A, B in $\mathfrak{B}_G(\mathcal{H})$ and any $E > 0$.

Proof. Since $E \mapsto [\|A\|_E^G]^2$ is a concave nonnegative function on \mathbb{R}_+ , it coincides with the infimum of all linear functions $E \mapsto a^2 + b^2E$ such that $[\|A\|_E^G]^2 \leq a^2 + b^2E$ for all $E > 0$ and the function $E \mapsto [\|A\|_E^G]^2 / E$ is non-increasing. So, the assertions A and B can be easily derived from Lemma 1.

To prove C note first that the seminorm properties of $b_{\sqrt{G}}(\cdot)$ follow from the second formula in A, while B implies $b_{\sqrt{G}}^{-1}(0) = \mathfrak{B}_G^0(\mathcal{H})$. So, since the function $E \mapsto [\|A\|_E^G]^2 / E$ is non-increasing for any given A , the inequality (10) follows from the triangle inequality for $b_{\sqrt{G}}(\cdot)$. \square

Due to Proposition 1 one can reformulate the results in Section 4 in [6] using the notions of \sqrt{G} -bounded and \sqrt{G} -infinitesimal operators. In particular, Theorem 3 in [6] implies the following characterization of \sqrt{G} -infinitesimal operators.

Corollary 1. *An operator A defined on $\mathcal{D}(\sqrt{G})$ is \sqrt{G} -infinitesimal if and only if for any separable Hilbert space \mathcal{K} the operator $A \otimes I_{\mathcal{K}}$ (naturally defined on the set $\mathcal{D}(\sqrt{G}) \otimes \mathcal{K}$) has a continuous linear extension to the set*

$$\mathcal{V}_{\sqrt{G} \otimes I_{\mathcal{K}}, E} = \left\{ \varphi \in \mathcal{H} \otimes \mathcal{K} \mid \|\sqrt{G} \otimes I_{\mathcal{K}} \varphi\|^2 \leq E \right\}$$

for any $E > 0$. If A is a \sqrt{G} -infinitesimal operator then

$$\|A \otimes I_{\mathcal{K}}(\varphi - \psi)\| \leq \varepsilon \|A\|_{4E/\varepsilon^2}^G \quad (11)$$

for any φ and ψ in $\mathcal{V}_{G \otimes I_{\mathcal{K}}, E}$ such that $\|\varphi - \psi\| \leq \varepsilon$. The r.h.s. of (11) tends to zero as $\varepsilon \rightarrow 0^+$ by condition (7).

Proposition 6 in [6] and Proposition 1B imply that any 2-positive linear map $\Phi : \mathfrak{B}(\mathcal{H}) \rightarrow \mathfrak{B}(\mathcal{H})$ such that $\Phi(I_{\mathcal{H}}) \leq I_{\mathcal{H}}$ having the predual map $\Phi_* : \mathfrak{T}(\mathcal{H}) \rightarrow \mathfrak{T}(\mathcal{H})$ with finite⁹

$$Y_{\Phi}(E) \doteq \sup \{ \text{Tr} G \Phi_*(\rho) \mid \rho \in \mathfrak{S}(\mathcal{H}), \text{Tr} G \rho \leq E \}$$

is uniquely extended to a linear transformation of the set of all \sqrt{G} -infinitesimal operators bounded w.r.t. the norm $\|\cdot\|_E^G$.¹⁰

Finally, consider application of the formula for the \sqrt{G} -bound in Proposition 1A.

Example. Let $\mathcal{H} = L_2(\mathbb{R})$ and $S(\mathbb{R})$ be the set of infinitely differentiable rapidly decreasing functions with all the derivatives tending to zero quicker than any degree of $|x|$ when $|x| \rightarrow +\infty$. Consider the operators q and p defined on the set $S(\mathbb{R})$ by setting

$$(q\varphi)(x) = x\varphi(x) \quad \text{and} \quad (p\varphi)(x) = \frac{1}{i} \frac{d}{dx} \varphi(x).$$

These operators are essentially self-adjoint. They represent (sharp) real observables of position and momentum of a quantum particle in the system of units where Planck's constant \hbar is equal to 1 [2, Ch.12]. For given $\omega > 0$ consider the operators

$$a = (\omega q + ip)/\sqrt{2\omega} \quad \text{and} \quad a^\dagger = (\omega q - ip)/\sqrt{2\omega} \quad (12)$$

defined on $S(\mathbb{R})$. The operator $N = a^\dagger a = aa^\dagger - I_{\mathcal{H}}$ is positive and essentially self-adjoint. It represents (sharp) real observable of the number of quanta of the harmonic oscillator with frequency ω . In [6, Section 5] the following estimates are obtained

$$\sqrt{\frac{2E + 1/2}{\omega}} < \|q\|_E^N \leq \sqrt{\frac{2E + 1}{\omega}}, \quad \sqrt{(2E + 1/2)\omega} < \|p\|_E^N \leq \sqrt{(2E + 1)\omega} \quad (13)$$

(the E -norms of q and p depend on ω , since the operator N depends on ω). Thus, the second formula in Proposition 1A implies that $b_{\sqrt{N}}(p) = \sqrt{2/\omega}$ and $b_{\sqrt{N}}(q) = \sqrt{2\omega}$. \square

I am grateful to T.V.Shulman for the help and useful discussion.

⁹It is easy to see that $E \mapsto Y_{\Phi}(E)$ is a concave function. So, finiteness of $Y_{\Phi}(E)$ for some $E > 0$ implies finiteness of $Y_{\Phi}(E)$ for all $E > 0$ and boundedness of the function $E \mapsto Y_{\Phi}(E)/E$.

¹⁰If G is a Hamiltonian of a quantum system then the quantity $Y_{\Phi}(E)/E$ can be treated as an energy amplification factor of Φ . Quantum channels Φ with finite $Y_{\Phi}(E)$ called energy-limited in [9] naturally appear as realistic quantum dynamical maps.

References

- [1] I.Bengtsson, K.Zyczkowski, "Geometry of Quantum States: An Introduction to Quantum Entanglement", 2nd edition. Cambridge: Cambridge University Press, 2017.
- [2] A.S.Holevo, "Quantum systems, channels, information. A mathematical introduction", Berlin, DeGruyter, 2012.
- [3] T.Kato, "Perturbation Theory for Linear Operators", Springer-Verlag, New York-Heidelberg-Berlin, 1980.
- [4] D.Kretschmann, D.Schlingemann, R.F.Werner, "A Continuity Theorem for Stinespring's Dilation", J. Funct. Anal. V.255, N.8, 1889-1904 (2008); arXiv:0710.2495.
- [5] M.E.Shirokov, "Strong convergence of quantum channels: continuity of the Stinespring dilation and discontinuity of the unitary dilation", arXiv:1712.03219.
- [6] M.E.Shirokov, "Operator E-norms and their use", arXiv:1806.05668.
- [7] M.E.Shirokov, "On completion of the cone of CP linear maps with respect to the energy-constrained diamond norm", arXiv:1810.10922.
- [8] B.Simon, "Operator Theory: A Comprehensive Course in Analysis", Part IV American Mathematical Society, 2015.
- [9] A.Winter, "Energy-constrained diamond norm with applications to the uniform continuity of continuous variable channel capacities", arXiv:1712.10267.