Electroweak Baryogenesis From Dark CP Violation

Marcela Carena,^{1,2,3} Mariano Quirós,^{4,5} and Yue Zhang^{1,6}

¹Theoretical Physics Department, Fermilab, P.O. Box 500, Batavia, IL 60510, USA

²Enrico Fermi Institute, University of Chicago, Chicago, Illinois, 60637, USA

³Kavli Institute for Cosmological Physics, University of Chicago, Chicago, Illinois, 60637, USA

⁴Department of Physics, University of Notre Dame, 225 Nieuwland Hall, Notre Dame, IN 46556, USA

⁵Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science

and Technology (BIST), Campus UAB, 08193 Bellaterra (Barcelona) Spain

⁶Department of Physics and Astronomy, Northwestern University, Evanston, IL 60208, USA

(Dated: November 27, 2018)

We present a novel mechanism of electroweak baryogenesis where CP violation occurs in a dark sector, comprised of standard model gauge singlets, thereby evading the strong electric dipole moment constraints. In this framework, the background of time-like component of a new gauge boson Z'_{μ} , generated at electroweak temperatures, drives the electroweak sphaleron processes to create the required baryon asymmetry. We first discuss the crucial ingredients for this mechanism to work, and then show that all of them can be elegantly embedded in ultraviolet completions with spontaneously broken gauged lepton number. The models under consideration have a rich phenomenology and can be experimentally probed in leptophilic Z' searches, dark matter searches, heavy Majorana neutrino searches, as well as through hunting for new Higgs portal scalars in multi-lepton channels at colliders.

The observed matter-antimatter asymmetry in the universe is widely believed to yield strong evidence for new phenomena beyond the standard model (SM) of particle physics. Electroweak baryogenesis (EWBG) is an elegant mechanism [1–15] that generates the observed baryon asymmetry at the electroweak phase transition (EWPT). This demands new physics close to the electroweak scale, to account for CP violating effects larger than those present in the SM. Moreover, the requirement of a sufficiently strong EWPT, along with the precision measurements of the Higgs boson properties, demands an extended scalar sector affecting the out-of-equilibrium processes.

The impressive recent progress in electric dipole moment (EDM) experiments [16–19] imposes strong constraints on the required new sources of CP violation in SM extensions, such as two-Higgs-doublet models and supersymmetry [20–35]. This provides a strong motivation to consider CP violation triggered in dark sectors through SM gauge singlets [36], which may naturally suppress contributions to EDMs. Such an enticing idea, however, leaves the challenging task of finding the suitable mechanism to transfer the CP violation from the dark sector to the visible sector, to successfully create the baryon asymmetry at electroweak temperatures.

In this *Letter*, we propose a new mechanism of EWBG where the transfer of CP violation to the visible sector is achieved by means of a vector boson Z'_{μ} which couples to the SM leptons, and to dark fermions with CPviolating Yukawa interactions involving additional SM singlet scalars. Such scalars may provide, through the Higgs portal, a sufficiently strong first-order EWPT. The time-like component of the new gauge boson, Z'_0 , is CPodd and can transfer CP violation to the visible sector. During EWPT, the CP violating source yields a nonzero background $\langle Z'_0 \rangle$, which acts as a chemical potential for the SM leptons, providing a thermal equilibrium lepton number asymmetry. In the absence of any primordial asymmetries, such source term, through the electroweak sphaleron processes, generates an equal amount of baryon and lepton number asymmetries, which freeze in after the EWPT is completed and the sphalerons become inactive. We point out that the current which couples to Z'_{μ} must be anomalous with respect to the SM weak interaction during the EWBG epoch.

In the following, we will present the main ingredients of the proposed EWBG mechanism, and briefly explore its phenomenological consequences. We refer to the companion paper [37] for a more detailed, quantitative presentation.

The Need of An Anomalous Current Coupled to Z'. The effective Lagrangian relevant to our discussion contains the couplings of a vector boson Z'_{μ} to a current involving SM leptons and quarks, J^{μ} ,

$$\mathcal{L}_{eff} = \mathcal{L}_{\rm SM} - \frac{1}{4} Z'_{\mu\nu} Z'^{\mu\nu} + \frac{1}{2} M_{Z'}^2 Z'_{\mu} Z'^{\mu} + g' Z'_{\mu} J^{\mu} , \quad (1)$$

$$J^{\mu} = \sum_{i=1}^{3} \left[q_{L_{L_{i}}} \bar{L}_{L_{i}} \gamma^{\mu} L_{L_{i}} + q_{e_{R_{i}}} \bar{e}_{R_{i}} \gamma^{\mu} e_{R_{i}} + q_{\nu_{R_{i}}} \bar{\nu}_{R_{i}} \gamma^{\mu} \nu_{R_{i}} \right. \\ \left. + q_{Q_{L_{i}}} \bar{Q}_{L_{i}} \gamma^{\mu} Q_{L_{i}} + q_{u_{R_{i}}} \bar{u}_{R_{i}} \gamma^{\mu} u_{R_{i}} + q_{d_{R_{i}}} \bar{d}_{R_{i}} \gamma^{\mu} d_{R_{i}} \right], (2)$$

where L_{L_i}, e_{R_i} and ν_{R_i} are the SM leptons and righthanded neutrinos, Q_{L_i}, u_{R_i} and d_{R_i} are SM quarks, and q_F is the charge of the corresponding fermion F.

We assume that the above effective Lagrangian describes a period of the early universe when EWBG occurs. Moreover, we further assume that the vector field Z'_{μ} develops a time-like background, $\langle Z'_0 \rangle \neq 0$, sourced by a U(1) charge density, whose origin will be addressed later on. Through Eq. (1) the Z'_0 background acts as a chemical potential for the fermions. Of particular interest to us are the SM quark and lepton doublets. In the presence of a chemical potential, if the lepton or quark number were allowed to change independently, the particle-antiparticle number density asymmetry, defined as $\Delta n_F \equiv n_F - n_{\bar{F}}$ with $F = L_{L_i}, Q_{L_i}$, will be generated and evolve toward its thermal equilibrium value

$$\Delta n_F^{\rm EQ} = \frac{2N_c}{3} T_c^2 g' q_F \left\langle Z_0' \right\rangle \ , \tag{3}$$

where $N_c = 3$ (1) for quarks (leptons) is the color factor, whereas T_c is the EWPT critical temperature. This expression is exact in the electroweak symmetric phase where all SM fermions are massless.

Within the context of EWBG, there is only one process where the lepton (L) and baryon (B) numbers are simultaneously violated – the electroweak sphalerons. Each sphaleron process violates B + L but conserves B - L among the left-handed $SU(2)_L$ doublets, where the baryon and lepton asymmetries are defined as

$$\Delta n_{B_L} \equiv \frac{1}{3} \sum_{i=1}^{3} \Delta n_{Q_{L_i}}, \quad \Delta n_{L_L} \equiv \sum_{i=1}^{3} \Delta n_{L_{L_i}} .$$
 (4)

The $(B \pm L)_L$ asymmetries satisfy the Boltzmann equations

$$\frac{\partial \Delta n_{(B+L)_L}}{\partial t} = \Gamma_{\rm sph} \left(\mathcal{S} - \Delta n_{(B+L)_L} \right) ,$$

$$\frac{\partial \Delta n_{(B-L)_L}}{\partial t} = 0, \qquad \mathcal{S} = \sum_{i=1}^3 \left(\Delta n_{L_{L_i}}^{\rm EQ} + \Delta n_{Q_{L_i}}^{\rm EQ} \right) ,$$
⁽⁵⁾

where $\Gamma_{\rm sph} \simeq 120 \alpha_w^5 T_c$ is the sphaleron rate in the electroweak unbroken phase [38] and it is exponentially suppressed in the broken phase. S serves as the source for creating a net B + L asymmetry, with $\Delta n_{L_{L_i}}^{\rm EQ}$ and $\Delta n_{Q_{L_i}}^{\rm EQ}$ contributing to it democratically, the same way as sphalerons act on every $SU(2)_L$ doublet.

Starting from a primordially symmetric universe implies, $\Delta n_{B_L} = \Delta n_{L_L} = (1/2)\Delta n_{(B+L)_L}$. With this, the first equation in (5) simplifies to

$$\frac{\partial \Delta n_{B_L}}{\partial t} = \Gamma_{\rm sph} \left(\frac{1}{2} \mathcal{S} - \Delta n_{B_L} \right) , \qquad (6)$$

and from Eq. (3) it is straightforward to derive

$$S = \frac{2}{3}T^2 g' \sum_{i=1}^{3} \left(q_{L_{L_i}} + 3q_{Q_{L_i}} \right) \langle Z'_0 \rangle .$$
 (7)

Remarkably Eq. (7) is proportional to the nonconservation of the current J^{μ} , *i.e.*, the coefficient appearing its chiral anomaly with respect to $SU(2)_L^2$,

$$\partial_{\mu}J^{\mu} \propto \sum_{i=1}^{3} \left(q_{L_{L_{i}}} + 3q_{Q_{L_{i}}} \right) \operatorname{tr}(W\widetilde{W}) , \qquad (8)$$

where W(W) is the $SU(2)_L$ field (dual) strength. Hence we have found a *necessary* condition for the proposed EWBG mechanism to work, namely, the current to which the Z'_{μ} couples must be anomalous with respect to $SU(2)_L^2$. Had the charges q_F in Eq. (2) been arranged such that the current J^{μ} were conserved, the source term in Eq. (6) would have vanished and, in turn, no net baryon asymmetry would have been created.

The effective Lagrangian of Eq. (1) can be obtained from a UV complete U(1) gauge theory whose gauge boson is Z'_{μ} and q_F are the corresponding SM fermion U(1) charges. In the case of a non-conserved J^{μ} , additional fermions (anomalons) are required to render the U(1) anomaly free. The total current of the U(1) gauge symmetry is the sum of J^{μ} in Eq. (2) and that of the anomalons, J^{μ}_{a} , such that the anomaly cancellation condition imposes, $\partial_{\mu}(J^{\mu} + J^{\mu}_{a}) = 0$. The anomalon fields, once introduced, will also contribute to the source term \mathcal{S} , Eq. (7). Here, however, we assume that the U(1)gauge symmetry is spontaneously broken above the electroweak scale (e.g., at TeV scales), and that the anomalons get symmetry breaking masses. If the anomalons have masses much larger than the EWPT temperature, their population, as well as their impact on the electroweak sphalerons, will become Boltzmann suppressed. In such case, the anomalons, although canceling the gauge anomalies, have negligible contribution to \mathcal{S} .

Good candidates for such a U(1) symmetry include gauged lepton number, baryon number, or any flavor dependent combination of the two that remains anomalous, within the SM, with respect to $SU(2)_L^2$. In contrast, the proposed EWBG mechanism cannot work if the U(1) is already anomaly free given the SM fermion content (plus right-handed neutrinos), for example, B-L, $L_{\mu}-L_{\tau}$, etc. In the following we discuss the realization of our EWBG mechanism in a UV complete model with gauged lepton number, $U(1)_{\ell}$. As we shall see, a dark matter candidate also naturally emerges in the theory.

As another useful remark, interestly the above discussion remains valid even if the $\langle Z'_0 \rangle$ background is spacetime inhomogeneous. Indeed, we will consider a first order EWPT which temporarily creates $\langle Z'_0 \rangle$ in front of the expanding bubble walls.

CP Violation and the Electroweak Phase Transition. We will now address the origin of the Z'_0 field background, as well as the dynamics of the EWPT. In analogy to a static electric potential, the $\langle Z'_0 \rangle$ background is *C*, *CP* and *CPT* odd, and can be generated by a net $U(1)_\ell$ charge distribution near the bubble wall. To this end we introduce a fermionic particle χ with *CP* violating microscopic interactions with the bubble wall. Since χ is a SM gauge singlet that cannot couple to the Higgs field through renormalizable interactions, we will introduce a SM scalar singlet *S* to interact with it,

$$\bar{\chi}_L(m_0 + \lambda e^{i\theta_\lambda}S)\chi_R + \text{h.c.}$$
 (9)

Within the bubble wall, the Higgs VEV turns on, while the S VEV simultaneously turns off. Such a transition to the electroweak broken phase has been studied and involves a two-step process from the original vacuum with $\langle S \rangle = \langle H \rangle = 0$ [36, 39–42]. The necessary ingredient to allow for a strongly first order EWPT is a sizable scalar quartic term, $|S|^2|H|^2$, by which the Higgs becomes a portal to the dark sector.

We will consider the following ansatz for the S profile across the bubble wall, $|S(z)| = s_0 \left[1 + \tanh(z/L_{\omega})\right]/2$. The coordinate z is defined in the rest frame of the bubble wall which is located at z = 0, whereas L_{ω} is the wall width. Observe that to accommodate a physical CP violating effect through Eq. (9), we need a scalar potential that fixes the phase of S. During the EWPT, the VEV of S contributes to the χ mass through Eq. (9), whereas the bare mass term m_0 has its origin in the spontaneous breaking of $U(1)_{\ell}$. If the two mass terms carry different, space-time dependent phases, the dispersion relations of χ_L, χ_R and their antiparticles will be modified in a *CP* violating way. This affects the phase space distributions of such particles and yields a non-trivial solution to the corresponding diffusion equations, leading to net number density asymmetries in χ_L, χ_R ,

$$\Delta n_{\chi}(z) \equiv n_{\chi_L} - n_{\chi_L^c} = -(n_{\chi_R} - n_{\chi_R^c}) \neq 0 . \quad (10)$$

The spatial distribution of $\Delta n_{\chi}(z)$ will peak around the bubble wall. For details on solving the diffusion equations and the numerical computation of $\Delta n_{\chi}(z)$, we refer the reader to the companion paper [37]. If χ_L and χ_R carry different $U(1)_{\ell}$ quantum numbers, the above chiral asymmetries will give a net $U(1)_{\ell}$ charge density distribution around the bubble wall,

$$\rho_{\ell}(z) = (q_{\chi_L} - q_{\chi_R}) \Delta n_{\chi}(z) .$$
 (11)

Neglecting the curvature of the bubble wall, the $\langle Z'_0 \rangle$ background sourced by ρ_ℓ can be calculated in cylindrical coordinates to be

$$\langle Z'_0(z) \rangle = \frac{g'}{2M_{Z'}} \int_{-\infty}^{\infty} dy \ \rho_\ell(y) \ e^{-M_{Z'}|z-y|} \ .$$
 (12)

Given this $\langle Z'_0 \rangle$ background, the final baryon asymmetry generated can be obtained by solving Eq. (6),

$$\Delta n_B = \frac{\Gamma_{\rm sph}}{v_\omega} \int_0^\infty dz \ \mathcal{S}(z) \ e^{-\Gamma_{\rm sph} z/v_\omega} \ , \qquad (13)$$

where v_{ω} is the bubble wall expansion velocity. The parametric dependence in today's baryon to entropy ratio is, $\eta_B = \Delta n_B/s \sim g'^2 N_g^2 T_c^3 L_{\omega} \alpha_W^5 / (M_{Z'}^2 v_{\omega}).$

UV Complete Models. Next, we discuss unifying all the above ingredients for EWBG into a UV complete framework with gauged (anomaly free) lepton number symmetry, $U(1)_{\ell}$. There are several choices to define the lepton number ℓ . The most obvious one is to gauge all

the three SM families universally by taking $\ell = L_e + L_{\mu} + L_{\tau}$. Alternatively, one could also gauge only two lepton flavors such as $\ell = L_{\mu} + L_{\tau}$. We will consider these two cases as benchmark models. The minimal set of new fermion content is given in Table I [43, 44], where q is an arbitrary real number. The index *i* runs through $e, \mu, \tau \ (\mu, \tau)$ in the first (second) model, and $N_g = 3$ (2) defines the number of families charged under the $U(1)_{\ell}$, correspondingly.

Particle	$SU(3)_c$	$SU(2)_L$	$U(1)_Y$	$U(1)_{\ell}$
L_{L_i}	1	2	-1/2	1
e_{R_i}	1	1	-1	1
ν_{R_i}	1	1	0	1
$L' = (\nu'_L, e'_L)^T$	1	2	-1/2	q
e'_R	1	1	-1	q
χ_R	1	1	0	q
$R' = (\nu_R'', e_R'')^T$	1	2	-1/2	$\mathbf{q} + N_g$
$e_L^{\prime\prime}$	1	1	-1	$\mathbf{q} + N_g$
χ_L	1	1	0	$\mathbf{q} + N_g$
Φ, S	1	1	0	N_g

TABLE I. UV completion of the effective theory.

To spontaneously break the $U(1)_{\ell}$, and at the same time give masses to the new fermions, we introduce the complex scalar Φ carrying $U(1)_{\ell}$ number N_g . We assume that Φ picks up a VEV, $v_{\Phi}/\sqrt{2}$, above the electroweak scale. This VEV gives mass to the gauge boson $Z', M_{Z'} = N_g g' v_{\Phi}/2$, which can still be light if the gauge coupling g' is sufficiently small. We can also write down Yukawa couplings of the form,

$$\left(c_L \bar{R}' L' + c_e \bar{e}'_L e_R'' + c_\chi \bar{\chi}_L \chi_R\right) \Phi + \text{h.c.} , \qquad (14)$$

which will give vectorlike masses (with respect to the SM) to the new fermions. We assume $c_L \sim c_e$ to be large enough so that L', R', e''_L, e'_R are sufficiently heavy in comparison with the critical temperature of the EWPT. As noted earlier, the fermions L' and R' are needed to cancel the $U(1)_{\ell} \otimes SU(2)_L^2$ gauge anomaly, whereas decoupling them from the thermal bath provides the necessary condition for our EWBG mechanism to work. On the other hand, we assume the parameter c_{χ} to be sufficiently small so that χ is light and remains populated in the thermal bath during the EWPT. Their $U(1)_{\ell}$ charges are fixed in the UV theory: $q_{\chi_L} = \mathbf{q} + N_g, q_{\chi_R} = \mathbf{q}$, and their difference does not depend on \mathbf{q} . This helps to eliminate a free parameter from Eq. (11), so that $\rho_{\ell}(z) = N_q \Delta n_{\chi}$.

The fermion χ will source *CP* violation when it interacts with the expanding bubble wall. The χ - Φ interaction in Eq. (14) is responsible for generating the $m_0 = c_{\chi} v_{\Phi} / \sqrt{2}$ mass term in Eq. (9). In addition, as discussed before, another complex scalar *S* with the same




FIG. 1. We scan broadly over the model parameters to find points that allow for successful EWBG as proposed in this work (blue points). We show various experimental constraints (LEP, BaBar, beam dump, electron and muon g - 2, CCFR) by the correspondingly labeled shaded regions, in the gauged $L_e + L_\mu + L_\tau$ (left) and $L_\mu + L_\tau$ (right) models.

quantum numbers as Φ and a scalar potential that fixes its phase is required to yield a physical *CP* phase, barring the redefinitions of fermion fields. Moreover, such a complex scalar will also be responsible for the strong EWPT through a two-step phase transition.

To summarize, we have argued that all the key ingredients for our EWBG mechanism can be elegantly accommodated in a UV theory of gauged lepton number. The UV completion is useful in shedding light into the parameters of the low energy effective theory.

Neutrino Cosmology. It is worth commenting on the neutrino sector and implications of cosmological measurements on additional neutrino degrees of freedom, ΔN_{eff} [45–48], for the two benchmark models. The $U(1)_{\ell}$ gauge interaction could thermalize, in the very early universe, all the new fermions charged under it, and in particular the right-handed neutrinos ν_{R_i} . To avoid an excessive contribution to ΔN_{eff} , one option is to make the $U(1)_{\ell}$ interaction decouple early enough, preferably above the QCD phase transition temperature, $T_{\rm QCD} \sim 100 \,{\rm MeV}$. This implies $v_{\Phi} \gtrsim 10 \,{\rm TeV}$ if $M_{Z'} \gg T_{\rm QCD}$, or $g' \lesssim 10^{-5}$ if $M_{Z'} \ll T_{\rm QCD}$. The other option is to implement the seesaw mechanism by giving Majorana masses to ν_{R_i} . If all the ν_{R_i} are heavier than $\sim 500 \,\mathrm{MeV}$, they will decay before the big-bang nucleosynthesis and have no effect in ΔN_{eff} [49]. Interestingly, this option could easily be achieved in the gauged $L_{\mu} + L_{\tau}$ model where both Φ and S have charge $N_q = 2$. The experimental search for heavy Majorana neutrinos is of great phenomenological interests [50], especially as the new $U(1)_{\ell}$ gauge interaction allows them to be more copiously produced. We will investigate this exciting opportunity in a future work.

Experimental Probes. Here we summarize the phenomenological predictions unique to our EWBG mechanism and discuss the present experimental bounds in the two benchmark models presented above.

– The main motivation for this work is to provide the necessary amount of CP violation for baryogenesis, without being in tension with EDM measurements. In the gauged $U(1)_{\ell}$ UV complete models presented here, the fermion field χ responsible for CP violation is a SM singlet, thereby eliminating its Barr-Zee type [51] contribution to EDMs. In Ref. [37], we will show that the leading contribution arises at the four-loop level.

– In our EWBG mechanism, the Z'_{μ} gauge boson transmits the dark CP violation to the SM sector, and thereby the generation of the observed baryon asymmetry, $\eta_B \simeq 0.9 \times 10^{-10}$, restricts the values of $M_{Z'}$ as a function of its gauge coupling g'. In Fig. 1, the blue points are obtained from a scan over the parameter space that can account for the correct η_B . The allowed Z'masses are in the MeV to TeV range, with decreasing g'values for lighter Z', in agreement with the parametric dependence estimated below Eq. (13). In the left (right) panel of Fig. 1, we show the gauged $L_e + L_{\mu} + L_{\tau}$ ($L_{\mu} + L_{\tau}$) benchmark models.

– The search for Z' provides a handle on these EWBG scenarios. In the gauged $L_e + L_\mu + L_\tau$ model, the Z' has a coupling to the electron, which is subject to constraints from electron g - 2, e^+e^- colliders (LEP, BaBar) and electron beam dump experiments [52], as shown by the correspondingly labeled shaded regions in the left panel of Fig. 1. On the other hand, the gauged $L_\mu + L_\tau$ model is free from the above constraints. There is, however, a relevant constraint from neutrino trident production [53] which excludes the magenta region (labeled by CCFR) in the right panel of Fig. 1. Interestingly, Fig. 1 shows that there is a region of parameter space that can explain the muon g-2 anomaly (yellow band), and in the case of the $L_\mu + L_\tau$ model, such a region is allowed and favored by EWBG, with $M_{Z'} \lesssim 200$ MeV.

– The models considered here provide a dark matter

candidate, χ . The $U(1)_{\ell}$ gauge invariance implies that the SM singlet fermion χ can only interact with the SM particles via the Z' exchange, making χ a leptophilic dark matter candidate [54]. Its thermal relic density and detection prospects will be discussed in detail in Ref. [37].

– It is possible to search for the dark scalar S at high energy colliders, where it can be pair produced through the Higgs portal interaction. If S is lighter than twice the χ mass, it must decay via a χ loop into a pair of Z'bosons, yielding four leptons in the final state. The S- χ interaction is inherently CP violating and such decay can provide a test of dark CP violation via interference effects in the golden 4ℓ -channel.

Summary. We have proposed a novel mechanism for EWBG in which the CP violation occurs in a dark sector and is transmitted to the observable sector via the timelike background of a Z'_μ vector boson during a strong first-order EWPT. The Z'_μ is the gauge boson of a $U(1)_\ell$ gauge symmetry, and couples to an anomalous SM lepton number current. After the spontaneous $U(1)_{\ell}$ symmetry breaking, the new $SU(2)_L$ doublet fermions required to render the theory anomaly free become massive and decouple from the thermal bath before the EWPT. Because the *CP* violating interactions are active in the dark sector, its effects on EDMs are highly suppressed and evade present bounds. We show two benchmark scenarios with gauged $U(1)_{L_e+L_\mu+L_\tau}$ and $U(1)_{L_\mu+L_\tau}$ symmetries which provide concrete examples of UV completions. The models under consideration provide a rich phenomenology that can be probed in searches for leptophilic Z', dark matter, heavy Majorana neutrinos, and new scalars in multi-lepton channels at the LHC or prospective high energy colliders.

We thank Zackaria Chacko, James Cline, Bogdan Dobrescu, Paddy Fox, Ian Low, David Morrissey, Tim Tait and Carlos Wagner for useful discussions and comments. This manuscript has been authored by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics. Work at University of Chicago is supported in part by U.S. Department of Energy grant number DE-FG02-13ER41958. The work of M.Q. is partly supported by Spanish MINEICO under Grant CICYT-FEDER-FPA2014-55613-P and FPA2017-88915-P, and by the Severo Ochoa Excellence Program of MINEICO under Grant SEV-2016-0588. The work of Y.Z. is also supported by the DoE under contract number DE-SC0007859. Y.Z. would like to thank the Aspen Center for Physics, which is supported by National Science Foundation grant PHY-1607611, where part of this work was performed, and Colegio De Fisica Fundamental E Interdisciplinaria De Las Americas (COFI) for a travel support during the completion of this work.

- V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, Phys. Lett. **155B**, 36 (1985).
- [2] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, Phys. Lett. B245, 561 (1990).
- [3] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, Phys. Lett. B263, 86 (1991).
- [4] G. R. Farrar and M. E. Shaposhnikov, Phys. Rev. Lett. **70**, 2833 (1993), [Erratum: Phys. Rev. Lett.71,210(1993)], arXiv:hep-ph/9305274 [hep-ph].
- [5] M. Quiros, Helv. Phys. Acta 67, 451 (1994).
- [6] P. Huet and A. E. Nelson, Phys. Lett. B355, 229 (1995), arXiv:hep-ph/9504427 [hep-ph].
- [7] P. Huet and A. E. Nelson, Phys. Rev. D53, 4578 (1996), arXiv:hep-ph/9506477 [hep-ph].
- [8] A. Riotto, Phys. Rev. D53, 5834 (1996), arXiv:hep-ph/9510271 [hep-ph].
- [9] M. Carena, M. Quiros, A. Riotto, I. Vilja, and C. E. M. Wagner, Nucl. Phys. B503, 387 (1997), arXiv:hepph/9702409 [hep-ph].
- [10] M. Quiros, in Proceedings, Summer School in High-energy physics and cosmology: ICTP, Trieste, Italy, June 29-July 17, 1998 (1999) pp. 187–259, arXiv:hep-ph/9901312 [hep-ph].
- [11] M. Carena, J. M. Moreno, M. Quiros, M. Seco, and C. E. M. Wagner, Nucl. Phys. **B599**, 158 (2001), arXiv:hep-ph/0011055 [hep-ph].
- [12] J. M. Cline, M. Joyce, and K. Kainulainen, JHEP 07, 018 (2000), arXiv:hep-ph/0006119 [hep-ph].
- [13] M. Carena, M. Quiros, M. Seco, and C. E. M. Wagner, Nucl. Phys. B650, 24 (2003), arXiv:hep-ph/0208043 [hep-ph].
- [14] C. Lee, V. Cirigliano, and M. J. Ramsey-Musolf, Phys. Rev. **D71**, 075010 (2005), arXiv:hep-ph/0412354 [hep-ph].
- [15] J. M. Cline, in Les Houches Summer School Session 86: Particle Physics and Cosmology: The Fabric of Spacetime Les Houches, France, July 31-August 25, 2006 (2006) arXiv:hep-ph/0609145 [hep-ph].
- [16] V. Andreev et al. (ACME), Nature 562, 355 (2018).
- [17] J. Baron *et al.* (ACME), Science **343**, 269 (2014), arXiv:1310.7534 [physics.atom-ph].
- [18] W. C. Griffith, M. D. Swallows, T. H. Loftus, M. V. Romalis, B. R. Heckel, and E. N. Fortson, Phys. Rev. Lett. **102**, 101601 (2009).
- [19] C. A. Baker *et al.*, Phys. Rev. Lett. **97**, 131801 (2006), arXiv:hep-ex/0602020 [hep-ex].
- [20] J. Shu and Y. Zhang, Phys. Rev. Lett. 111, 091801 (2013), arXiv:1304.0773 [hep-ph].
- [21] S. Ipek, Phys. Rev. D89, 073012 (2014), arXiv:1310.6790 [hep-ph].
- [22] M. Jung and A. Pich, JHEP 04, 076 (2014), arXiv:1308.6283 [hep-ph].
- [23] T. Abe, J. Hisano, T. Kitahara, and K. Tobioka, JHEP 01, 106 (2014), [Erratum: JHEP04,161(2016)], arXiv:1311.4704 [hep-ph].
- [24] S. Inoue, M. J. Ramsey-Musolf, and Y. Zhang, Phys. Rev. D89, 115023 (2014), arXiv:1403.4257 [hep-ph].
- [25] K. Cheung, J. S. Lee, E. Senaha, and P.-Y. Tseng, JHEP 06, 149 (2014), arXiv:1403.4775 [hep-ph].
- [26] L. Bian, T. Liu, and J. Shu, Phys. Rev. Lett. 115, 021801 (2015), arXiv:1411.6695 [hep-ph].

- [27] C.-Y. Chen, S. Dawson, and Y. Zhang, JHEP 06, 056 (2015), arXiv:1503.01114 [hep-ph].
- [28] K. Fuyuto, J. Hisano, and E. Senaha, Phys. Lett. B755, 491 (2016), arXiv:1510.04485 [hep-ph].
- [29] M. Jiang, L. Bian, W. Huang, and J. Shu, Phys. Rev. D93, 065032 (2016), arXiv:1502.07574 [hep-ph].
- [30] N. Blinov, J. Kozaczuk, D. E. Morrissey, and C. Tamarit, Phys. Rev. **D92**, 035012 (2015), arXiv:1504.05195 [hepph].
- [31] C. Balazs, G. White, and J. Yue, JHEP 03, 030 (2017), arXiv:1612.01270 [hep-ph].
- [32] L. Bian and N. Chen, Phys. Rev. D95, 115029 (2017), arXiv:1608.07975 [hep-ph].
- [33] C.-Y. Chen, H.-L. Li, and M. Ramsey-Musolf, Phys. Rev. **D97**, 015020 (2018), arXiv:1708.00435 [hep-ph].
- [34] C. Cesarotti, Q. Lu, Y. Nakai, A. Parikh, and M. Reece, (2018), arXiv:1810.07736 [hep-ph].
- [35] D. Egana-Ugrinovic and S. Thomas, (2018), arXiv:1810.08631 [hep-ph].
- [36] J. M. Cline, K. Kainulainen, and D. Tucker-Smith, Phys. Rev. **D95**, 115006 (2017), arXiv:1702.08909 [hep-ph].
- [37] M. Carena, M. Quiros, and Y. Zhang, to appear.
- [38] D. Bodeker, G. D. Moore, and K. Rummukainen, Phys. Rev. **D61**, 056003 (2000), arXiv:hep-ph/9907545 [hep-ph].
- [39] J. R. Espinosa, T. Konstandin, and F. Riva, Nucl. Phys. B854, 592 (2012), arXiv:1107.5441 [hep-ph].
- [40] H. H. Patel, M. J. Ramsey-Musolf, and M. B. Wise, Phys. Rev. D88, 015003 (2013), arXiv:1303.1140 [hepph].
- [41] C. Cheung and Y. Zhang, JHEP 09, 002 (2013),

arXiv:1306.4321 [hep-ph].

- [42] D. Curtin, P. Meade, and C.-T. Yu, JHEP 11, 127 (2014), arXiv:1409.0005 [hep-ph].
- [43] P. Fileviez Perez and M. B. Wise, Phys. Rev. D82, 011901 (2010), [Erratum: Phys. Rev.D82,079901(2010)], arXiv:1002.1754 [hep-ph].
- [44] P. Schwaller, T. M. P. Tait, and R. Vega-Morales, Phys. Rev. D88, 035001 (2013), arXiv:1305.1108 [hep-ph].
- [45] V. Barger, P. Langacker, and H.-S. Lee, Phys. Rev. D67, 075009 (2003), arXiv:hep-ph/0302066 [hep-ph].
- [46] D. K. Ghosh, G. Senjanovic, and Y. Zhang, Phys. Lett. B698, 420 (2011), arXiv:1010.3968 [hep-ph].
- [47] V. Barger, P. Fileviez Perez, and S. Spinner, Phys. Lett. B696, 509 (2011), arXiv:1010.4023 [hep-ph].
- [48] J. Hamann, S. Hannestad, G. G. Raffelt, and Y. Y. Y. Wong, JCAP **1109**, 034 (2011), arXiv:1108.4136 [astroph.CO].
- [49] J. M. Berryman, A. de GouvĀła, K. J. Kelly, and Y. Zhang, Phys. Rev. **D96**, 075010 (2017), arXiv:1706.02722 [hep-ph].
- [50] A. Atre, T. Han, S. Pascoli, and B. Zhang, JHEP 05, 030 (2009), arXiv:0901.3589 [hep-ph].
- [51] S. M. Barr and A. Zee, Phys. Rev. Lett. 65, 21 (1990),
 [Erratum: Phys. Rev. Lett.65,2920(1990)].
- [52] J. Alexander et al. (2016) arXiv:1608.08632 [hep-ph].
- [53] W. Altmannshofer, S. Gori, M. Pospelov, and I. Yavin, Phys. Rev. Lett. **113**, 091801 (2014), arXiv:1406.2332 [hep-ph].
- [54] P. J. Fox, R. Harnik, J. Kopp, and Y. Tsai, Phys. Rev. D84, 014028 (2011), arXiv:1103.0240 [hep-ph].