

# On diagonal solutions of the reflection equation

Zengo Tsuboi

Osaka city university advanced mathematical institute  
3-3-138 Sugimoto, Sumiyoshi-ku Osaka 558-8585, Japan

November 27, 2018

## Abstract

We study solutions of the reflection equation associated with the quantum affine algebra  $U_q(\hat{gl}(N))$  and obtain diagonal K-operators in terms of the Cartan elements of a quotient of  $U_q(gl(N))$ . We also consider intertwining relations for these K-operators and find an augmented q-Onsager algebra like symmetry behind them.

Key words: augmented q-Onsager algebra; intertwining relation; K-matrix; L-operator; quantum algebra; reflection equation

## 1 Introduction

The reflection equation [1] is a fundamental object in quantum integrable systems with open boundary conditions [2]. It has the following form

$$R_{12} \left( \frac{y}{x} \right) K_1(x) R_{21}(xy) K_2(y) = K_2(y) R_{12} \left( \frac{1}{xy} \right) K_1(x) R_{21} \left( \frac{x}{y} \right), \quad x, y \in \mathbb{C}. \quad (1.1)$$

Here  $R(x)$  is a solution (R-matrix) of the Yang-Baxter equation and  $K(x)$  is a K-matrix. The indices 1,2 denote the space where the operators act non-trivially. In particular, there is a  $N \times N$  diagonal matrix solution [3] of the reflection equation associated with the R-matrices [4] for the  $N$ -dimensional fundamental representation of  $U_q(\hat{gl}(N))$ . These R-matrices are specialization of more general operators called  $L$ -operators:  $\mathbf{L}_{12}(x), \bar{\mathbf{L}}_{12}(x) \in U_q(gl(N)) \otimes \text{End}(\mathbb{C}^N)$ . Namely, they are given by  $R_{12}(x) = (\pi \otimes 1) \mathbf{L}_{12}(x)$ ,  $R_{21}(x) = (\pi \otimes 1) \bar{\mathbf{L}}_{12}(x)$ , where  $\pi$  is the fundamental representation of  $U_q(gl(N))$ . In this context, a natural problem is to investigate the solutions of the reflection equation associated with the L-operators:

$$\mathbf{L}_{12} \left( \frac{y}{x} \right) \mathbf{K}_1(x) \bar{\mathbf{L}}_{12}(xy) K_2(y) = K_2(y) \mathbf{L}_{12} \left( \frac{1}{xy} \right) \mathbf{K}_1(x) \bar{\mathbf{L}}_{12} \left( \frac{x}{y} \right), \quad (1.2)$$

where the K-operator  $\mathbf{K}(x)$  is an operator in  $U_q(\mathfrak{gl}(N))$ . In this paper, we propose diagonal solutions of (1.2), namely solutions written in terms of only the Cartan elements of (a quotient of)  $U_q(\mathfrak{gl}(N))$ . Evaluation of the K-operator for the fundamental representation of  $U_q(\mathfrak{gl}(N))$  reproduces the  $N \times N$  diagonal K-matrix:  $K(x) = \pi(\mathbf{K}(x))$ . In the context of Baxter Q-operators for integrable systems with open boundaries, the essentially same K-operator for  $N = 2$  case was previously proposed in [5]. This paper extends this to  $N \geq 3$  case in part.

We also rewrite (1.2) as intertwining relations for the K-operator and speculate an explicit form of the underlying symmetry algebra. For  $N = 2$  case, it is known that the augmented q-Onsager algebra [6, 7], which is a co-ideal subalgebra of  $U_q(\widehat{\mathfrak{sl}}(2))$ , serves as this. We find an augmented q-Onsager algebra like symmetry still exists for an oscillator representation even for  $N \geq 3$  case. At a little more abstract level, the relevant algebras will be the reflection equation algebras [2] and some coideal subalgebras of quantum affine algebras [8]. In addition, symmetries of the transfer matrix of open spin chains with diagonal K-matrices are discussed in [9]. Intertwining relations for K-matrices are studied, for example, in [10]. However, an explicit expression for the higher rank analogue of the augmented q-Onsager algebra seems to be missing in the literatures. We expect that our results can be a cue for further study on this.

There are several versions of the reflection equation ('left', 'right', 'twisted', 'untwisted'). However, their solutions, K-matrices, are considered to be related each other by some transformations. Then we concentrate on one of them. Throughout this paper, we use the general gradation<sup>1</sup> of  $U_q(\widehat{\mathfrak{gl}}(N))$ . This does not produce particularly new results since the difference of the gradation reduces to a simple similarity transformation and a rescaling of the spectral parameter on the level of the R-matrices. However, there is a merit to use it since the parameters in the general gradation play a role as 'markers' and make it easier to trace the actions of the generators of  $U_q(\widehat{\mathfrak{gl}}(N))$ .

## 2 The quantum affine and finite algebras

In this section, we review the quantum algebras [12, 13, 14] of type A and associated L-operators. We also refer the books [15, 16] for review of this subject.

### 2.1 The quantum affine algebra $U_q(\widehat{\mathfrak{gl}}(N))$

We introduce a q-commutator  $[X, Y]_q = XY - qYX$ , and set  $[X, Y]_1 = [X, Y]$ . The quantum affine algebra  $U_q(\widehat{\mathfrak{gl}}(N))$  is a Hopf algebra generated by the generators<sup>2</sup>  $e_i, f_i, k_i$ , where  $i \in \{1, \dots, N\}$ . For  $i, j \in \{1, 2, \dots, N\}$ , the defining relations of the

---

<sup>1</sup>We emulate [11] and their subsequent papers for this.

<sup>2</sup>In this paper, we do not use the degree operator  $d$ . We will only consider level zero representations. The notation  $e_0, f_0$  conventionally used in literatures corresponds to  $e_N, f_N$ . We assume that the deformation parameter  $q = e^{\hbar}$  ( $\hbar \in \mathbb{C}$ ) is not a root of unity.

algebra are given by

$$\begin{aligned}
[k_i, k_j] &= 0, & [k_i, e_j] &= (\delta_{ij} - \delta_{i,j+1})e_j, & [k_i, f_j] &= -(\delta_{ij} - \delta_{i,j+1})f_j, \\
[e_i, f_j] &= \delta_{ij} \frac{q^{h_i} - q^{-h_i}}{q - q^{-1}}, \\
[e_i, e_j] &= [f_i, f_j] = 0 \quad \text{for } |i - j| \geq 2, \\
[e_i, [e_i, [e_i, e_j]_{q^2}]_{q^{-2}}] &= [f_i, [f_i, [f_i, f_j]_{q^{-2}}]_{q^2}] = 0 \quad \text{for } N = 2, \quad i \neq j, \\
[e_i, [e_i, e_j]_q]_{q^{-1}} &= [f_i, [f_i, f_j]_{q^{-1}}]_q = 0 \quad \text{for } N \geq 3, \quad |i - j| = 1,
\end{aligned} \tag{2.1}$$

where  $h_i = k_i - k_{i+1}$ , and  $i, j$  should be interpreted modulo  $N$ : ( $N+1 \equiv 1$ ).  $\hat{c} = \sum_{i=1}^N k_i$  is a central element of the algebra. The algebra has the co-product  $\Delta : U_q(\hat{gl}(N)) \rightarrow U_q(\hat{gl}(N)) \otimes U_q(\hat{gl}(N))$  defined by

$$\begin{aligned}
\Delta(e_i) &= e_i \otimes 1 + q^{-h_i} \otimes e_i, \\
\Delta(f_i) &= f_i \otimes q^{h_i} + 1 \otimes f_i, \\
\Delta(k_i) &= k_i \otimes 1 + 1 \otimes k_i.
\end{aligned} \tag{2.2}$$

We will also use the opposite co-product defined by

$$\Delta' = \sigma \circ \Delta, \quad \sigma \circ (X \otimes Y) = Y \otimes X, \quad X, Y \in U_q(\hat{gl}(N)). \tag{2.3}$$

In addition to these, there are anti-poiside and co-unit, which will not be used here. The Borel subalgebras  $\mathcal{B}_+$  (resp.  $\mathcal{B}_-$ ) is generated by the elements  $e_i, k_i$  (resp.  $f_i, k_i$ ), where  $i \in \{1, \dots, N\}$ .

There exists a unique element [12, 17]  $\mathcal{R} \in \mathcal{B}_+ \otimes \mathcal{B}_-$  called the universal R-matrix which satisfies the following relations

$$\begin{aligned}
\Delta'(a) \mathcal{R} &= \mathcal{R} \Delta(a) \quad \text{for } \forall a \in U_q(\hat{gl}(N)), \\
(\Delta \otimes 1) \mathcal{R} &= \mathcal{R}_{13} \mathcal{R}_{23}, \\
(1 \otimes \Delta) \mathcal{R} &= \mathcal{R}_{13} \mathcal{R}_{12}.
\end{aligned} \tag{2.4}$$

where <sup>3</sup>  $\mathcal{R}_{12} = \mathcal{R} \otimes 1$ ,  $\mathcal{R}_{23} = 1 \otimes \mathcal{R}$ ,  $\mathcal{R}_{13} = (\sigma \otimes 1) \mathcal{R}_{23}$ . The Yang-Baxter equation

$$\mathcal{R}_{12} \mathcal{R}_{13} \mathcal{R}_{23} = \mathcal{R}_{23} \mathcal{R}_{13} \mathcal{R}_{12} \tag{2.5}$$

is a corollary of these relations (2.4). The universal R-matrix can be written in the form

$$\mathcal{R} = \overline{\mathcal{R}} q^{\sum_{i=1}^N k_i \otimes k_i}. \tag{2.6}$$

Here  $\overline{\mathcal{R}}$  is the reduced universal R-matrix, which is a series in  $e_j \otimes 1$  and  $1 \otimes f_j$  and does not contain Cartan elements.

---

<sup>3</sup>We will use similar notation for the L-operators to indicate the space on which they non-trivially act.

## 2.2 The quantum finite algebra $U_q(\mathfrak{gl}(N))$

The finite quantum algebra  $U_q(\mathfrak{gl}(N))$  is generated by the generators  $\{e_{i,i+1}, e_{i+1,i}\}_{i=1}^{N-1}$  and  $\{e_{ii}\}_{i=1}^N$  obeying the following defining relations:

$$\begin{aligned} [e_{ii}, e_{jj}] &= 0, & [e_{ii}, e_{j,j+1}] &= (\delta_{i,j} - \delta_{i,j+1})e_{j,j+1}, & [e_{ii}, e_{j+1,j}] &= -(\delta_{i,j} - \delta_{i,j+1})e_{j+1,j}, \\ [e_{i,i+1}, e_{j+1,j}] &= \delta_{ij} \frac{q^{e_{ii}-e_{i+1,i+1}} - q^{-e_{ii}+e_{i+1,i+1}}}{q - q^{-1}}, \\ [e_{i,i+1}, e_{j,j+1}] &= [e_{i+1,i}, e_{j+1,j}] = 0 & \text{for } |i-j| \geq 2, \\ [e_{i,i+1}, [e_{i,i+1}, e_{j,j+1}]_q]_{q^{-1}} &= [e_{i+1,i}, [e_{i+1,i}, e_{j+1,j}]_{q^{-1}}]_q = 0 & \text{for } |i-j| = 1. \end{aligned} \quad (2.7)$$

Note that  $c = e_{11} + e_{22} + \cdots + e_{NN}$  is a central element of the algebra. We will also use elements defined recursively by

$$\begin{aligned} e_{ij} &= [e_{ik}, e_{kj}]_q & \text{for } i > k > j, \\ e_{ij} &= [e_{ik}, e_{kj}]_{q^{-1}} & \text{for } i < k < j, \end{aligned} \quad (2.8)$$

$$\begin{aligned} \bar{e}_{ij} &= [\bar{e}_{ik}, \bar{e}_{kj}]_{q^{-1}} & \text{for } i > k > j, \\ \bar{e}_{ij} &= [\bar{e}_{ik}, \bar{e}_{kj}]_q & \text{for } i < k < j, \end{aligned} \quad (2.9)$$

where  $\bar{e}_{i,i+1} = e_{i,i+1}$ ,  $\bar{e}_{i+1,i} = e_{i+1,i}$ . We summarize relations among these elements in the appendix.

There is an evaluation map  $\mathbf{ev}_x: U_q(\hat{\mathfrak{gl}}(N)) \mapsto U_q(\mathfrak{gl}(N))$ :

$$\begin{aligned} e_N &\mapsto x^{s_N} q^{-e_{11}} e_{N1} q^{-e_{NN}}, \\ f_N &\mapsto x^{-s_N} q^{e_{NN}} e_{1N} q^{e_{11}}, \\ e_i &\mapsto x^{s_i} e_{i,i+1}, & f_i &\mapsto x^{-s_i} e_{i+1,i} & \text{for } 1 \leq i \leq N-1, \\ k_i &\mapsto e_{ii} & \text{for } 1 \leq i \leq N, \end{aligned} \quad (2.10)$$

where  $x \in \mathbb{C}$  is a spectral parameter; and the parameters  $s_i \in \mathbb{Z}$  fix the gradation of the algebra. We will also use another evaluation map  $\bar{\mathbf{ev}}_x: U_q(\hat{\mathfrak{gl}}(N)) \mapsto U_q(\mathfrak{gl}(N))$ , which is defined by

$$\begin{aligned} e_N &\mapsto x^{s_N} q^{e_{11}} \bar{e}_{N1} q^{e_{NN}}, \\ f_N &\mapsto x^{-s_N} q^{-e_{NN}} \bar{e}_{1N} q^{-e_{11}}, \\ e_i &\mapsto x^{s_i} e_{i,i+1}, & f_i &\mapsto x^{-s_i} e_{i+1,i} & \text{for } 1 \leq i \leq N-1, \\ k_i &\mapsto e_{ii} & \text{for } 1 \leq i \leq N. \end{aligned} \quad (2.11)$$

## 2.3 Representations of quantum algebras

Let  $\pi$  be the fundamental representation of  $U_q(\mathfrak{gl}(N))$ , and  $E_{ij}$  be a  $N \times N$  matrix unit whose  $(k, l)$ -element is  $\delta_{i,k} \delta_{j,l}$ . We chose the basis of the representation space so that  $\pi(e_{ij}) = \pi(\bar{e}_{ij}) = E_{ij}$  holds. The composition  $\pi_x = \pi \circ \mathbf{ev}_x = \pi \circ \bar{\mathbf{ev}}_x$  gives an evaluation representation <sup>4</sup> of  $U_q(\hat{\mathfrak{gl}}(N))$ .

<sup>4</sup>Evaluation representations based the map  $\mathbf{ev}_x$  do not necessary coincide with the ones based on the map  $\bar{\mathbf{ev}}_x$  for more general representations.

The  $q$ -oscillator algebra is generated by the generators  $\mathbf{c}_i, \mathbf{c}_i^\dagger, \mathbf{n}_i$  ( $i \in \{1, 2, \dots, N\}$ ) obeying the defining relations:

$$\begin{aligned} [\mathbf{c}_i, \mathbf{c}_j^\dagger]_{q^{\delta_{ij}}} &= \delta_{ij} q^{-\mathbf{n}_i}, & [\mathbf{c}_i, \mathbf{c}_j^\dagger]_{q^{-\delta_{ij}}} &= \delta_{ij} q^{\mathbf{n}_i}, \\ [\mathbf{n}_i, \mathbf{c}_j] &= -\delta_{ij} \mathbf{c}_j, & [\mathbf{n}_i, \mathbf{c}_j^\dagger] &= \delta_{ij} \mathbf{c}_j^\dagger, & [\mathbf{n}_i, \mathbf{n}_j] &= [\mathbf{c}_i, \mathbf{c}_j] = [\mathbf{c}_i^\dagger, \mathbf{c}_j^\dagger] = 0. \end{aligned} \quad (2.12)$$

A Fock space is given by the actions of the generators on the vacuum vector defined by  $\mathbf{c}_i|0\rangle = \mathbf{n}_i|0\rangle = 0$ . A highest weight representation of  $U_q(\mathfrak{gl}(N))$  with the highest weight  $(m, 0, \dots, 0)$  and the highest weight vector  $|0\rangle$  ( $e_{ii}|0\rangle = m\delta_{i,1}|0\rangle$ ,  $e_{j,j+1}|0\rangle = 0$ ,  $1 \leq i \leq N$ ,  $1 \leq j \leq N-1$ ) can be realized in terms of the  $q$ -oscillator algebra (cf.  $q$ -analogue of the Holstein-Primakoff realization [18, 19]) :

$$\begin{aligned} e_{11} &= m - \mathbf{n}_2 - \dots - \mathbf{n}_N, \\ e_{kk} &= \mathbf{n}_k, & \text{for } 2 \leq k \leq N \\ e_{1j} &= \mathbf{c}_j q^{\sum_{k=2}^{j-1} \mathbf{n}_k}, & \bar{e}_{1j} &= \mathbf{c}_j q^{-\sum_{k=2}^{j-1} \mathbf{n}_k} & \text{for } 2 \leq j \leq N, \\ e_{i1} &= \mathbf{c}_i^\dagger \left[ m - \sum_{k=2}^N \mathbf{n}_k \right]_q q^{-\sum_{k=2}^{i-1} \mathbf{n}_k}, & \bar{e}_{i1} &= \mathbf{c}_i^\dagger \left[ m - \sum_{k=2}^N \mathbf{n}_k \right]_q q^{\sum_{k=2}^{i-1} \mathbf{n}_k} & \text{for } 2 \leq i \leq N, \\ e_{ij} &= \mathbf{c}_i^\dagger \mathbf{c}_j q^{\sum_{k=i+1}^{j-1} \mathbf{n}_k}, & \bar{e}_{ij} &= \mathbf{c}_i^\dagger \mathbf{c}_j q^{-\sum_{k=i+1}^{j-1} \mathbf{n}_k} & \text{for } 2 \leq i < j \leq N, \\ e_{ij} &= \mathbf{c}_i^\dagger \mathbf{c}_j q^{-\sum_{k=j+1}^{i-1} \mathbf{n}_k}, & \bar{e}_{ij} &= \mathbf{c}_i^\dagger \mathbf{c}_j q^{\sum_{k=j+1}^{i-1} \mathbf{n}_k} & \text{for } 2 \leq j < i \leq N, \end{aligned} \quad (2.13)$$

where  $[x]_q = (q^x - q^{-x})/(q - q^{-1})$ .

## 2.4 L-operators

The so-called L-operators are images of the universal R-matrix, which are given by  $\mathbf{L}(xy^{-1}) = \phi(xy^{-1})(\mathbf{ev}_x \otimes \pi_y)\mathcal{R}$ ,  $\bar{\mathbf{L}}(xy^{-1}) = \bar{\phi}(xy^{-1})(\bar{\mathbf{ev}}_x \otimes \pi_y)\mathcal{R}_{21}$ , where  $\phi(xy^{-1})$  and  $\bar{\phi}(xy^{-1})$  are overall factors. They are solutions of the intertwining relations following from (2.4):

$$((\mathbf{ev}_x \otimes \pi_y)\Delta'(a)) \mathbf{L}(xy^{-1}) = \mathbf{L}(xy^{-1}) ((\mathbf{ev}_x \otimes \pi_y)\Delta(a)), \quad (2.14)$$

$$((\bar{\mathbf{ev}}_x \otimes \pi_y)\Delta(a)) \bar{\mathbf{L}}(xy^{-1}) = \bar{\mathbf{L}}(xy^{-1}) ((\bar{\mathbf{ev}}_x \otimes \pi_y)\Delta'(a)) \quad \forall a \in U_q(\hat{\mathfrak{gl}}(N)). \quad (2.15)$$

One can solve <sup>5</sup> these to get (cf. [14, 20])

$$\mathbf{L}(xy^{-1}) = \sum_{j,k=1}^N \left\{ (xy^{-1})^{-\xi_k + \xi_j} L_{kj}^+ - (xy^{-1})^{s - \xi_k + \xi_j} L_{kj}^- \right\} \otimes E_{kj}, \quad (2.16)$$

$$\bar{\mathbf{L}}(xy^{-1}) = \sum_{j,k=1}^N \left\{ - (xy^{-1})^{-s - \xi_k + \xi_j} \bar{L}_{kj}^+ + (xy^{-1})^{-\xi_k + \xi_j} \bar{L}_{kj}^- \right\} \otimes E_{kj}, \quad (2.17)$$

---

<sup>5</sup>Taking note on the expression (2.6), we adopt the solutions which satisfy  $\mathbf{L}(0) = \bar{\mathbf{L}}(\infty) = \sum_{i=1}^N q^{e_{ii}} \otimes E_{ii}$  for the case  $s_k > 0$  (for all  $k \in \{1, 2, \dots, N\}$ ).

where  $\xi_k = s_k + s_{k+1} + \cdots + s_N$ ,  $s = \xi_1$ , and the coefficients are related to the generators of  $U_q(\mathfrak{gl}(N))$  as

$$\begin{aligned} L_{ii}^+ &= q^{e_{ii}}, & L_{ii}^- &= q^{-e_{ii}}, \\ L_{ij}^+ &= (q - q^{-1})e_{ji}q^{e_{jj}}, & L_{ij}^- &= 0 \quad \text{for } i > j, \end{aligned} \quad (2.18)$$

$$L_{ij}^- = -(q - q^{-1})q^{-e_{ii}}e_{ji}, \quad L_{ij}^+ = 0 \quad \text{for } i < j,$$

$$\begin{aligned} \bar{L}_{ii}^+ &= q^{-e_{ii}}, & \bar{L}_{ii}^- &= q^{e_{ii}}, \\ \bar{L}_{ij}^+ &= -(q - q^{-1})q^{-e_{ii}}\bar{e}_{ji}, & \bar{L}_{ij}^- &= 0 \quad \text{for } i > j, \end{aligned} \quad (2.19)$$

$$\bar{L}_{ij}^- = (q - q^{-1})\bar{e}_{ji}q^{e_{jj}}, \quad \bar{L}_{ij}^+ = 0 \quad \text{for } i < j.$$

We remark that the second L-operator (2.17) follows from the first one (2.16) by the transformation

$$\begin{aligned} e_{ij} &\mapsto \bar{e}_{N+1-i, N+1-j}, & E_{ij} &\mapsto \bar{E}_{N+1-i, N+1-j} \quad \text{for } 1 \leq i, j \leq N, \\ s_l &\mapsto -s_{N-l} \quad \text{for } 1 \leq l \leq N-1, \\ s_N &\mapsto -s_N, \end{aligned} \quad (2.20)$$

which is composition of automorphisms of  $U_q(\hat{\mathfrak{gl}}(N))$  and  $U_q(\mathfrak{gl}(N))$ . In fact, the intertwining relation (2.15) follows from (2.14) by the same transformation (2.20).

Evaluating the L-operators for the fundamental representation <sup>6</sup>, we obtain R-matrices [4],

$$\begin{aligned} R(x) &= (\pi \otimes 1)\mathbf{L}(x) = \sum_{i=1}^N (q - x^s q^{-1}) E_{ii} \otimes E_{ii} + \sum_{i \neq j} (1 - x^s) E_{ii} \otimes E_{jj} \\ &\quad + (q - q^{-1}) \sum_{i < j} x^{\xi_i - \xi_j} E_{ij} \otimes E_{ji}, + (q - q^{-1}) \sum_{i > j} x^{s + \xi_i - \xi_j} E_{ij} \otimes E_{ji}, \end{aligned} \quad (2.21)$$

$$\begin{aligned} \bar{R}(x) &= (\pi \otimes 1)\bar{\mathbf{L}}(x) = \sum_{i=1}^N (q - x^{-s} q^{-1}) E_{ii} \otimes E_{ii} + \sum_{i \neq j} (1 - x^{-s}) E_{ii} \otimes E_{jj} \\ &\quad + (q - q^{-1}) \sum_{i > j} x^{\xi_i - \xi_j} E_{ij} \otimes E_{ji}, + (q - q^{-1}) \sum_{i < j} x^{-s + \xi_i - \xi_j} E_{ij} \otimes E_{ji}. \end{aligned} \quad (2.22)$$

### 3 The reflection equation and its solutions

In this section, we will derive intertwining relations from the reflection equation associated with the L-operators, and obtain K-operators in terms of the Cartan elements of (a quotient of)  $U_q(\mathfrak{gl}(N))$ .

---

<sup>6</sup>Up to an overall factor, (2.22) coincides with  $(\pi_x \otimes \pi_1)\mathcal{R}_{21}$  since (3.27) and  $c = 1$  hold true for the fundamental representation.

We start from the following form of the reflection equation for the R-matrices (2.21) and (2.22):

$$R_{12} \left( \frac{y}{x} \right) K_1(x) \bar{R}_{12}(xy) K_2(y) = K_2(y) R_{12} \left( \frac{1}{xy} \right) K_1(x) \bar{R}_{12} \left( \frac{x}{y} \right). \quad (3.1)$$

It is known that this equation allows a  $N \times N$  diagonal matrix solution (K-matrix) [3]. In our convention, it reads

$$K(x) = \sum_{k=1}^a x^{2(s-\xi_k)} (\epsilon_- + \epsilon_+ x^s) E_{kk} + \sum_{k=a+1}^N x^{2(s-\xi_k)} (\epsilon_- + \epsilon_+ x^{-s}) E_{kk}, \quad (3.2)$$

where  $a \in \{0, 1, \dots, N\}$ ,  $\epsilon_{\pm} \in \mathbb{C}$ . Now we would like to consider the reflection equation for the L-operators (2.16) and (2.17):

$$\mathbf{L}_{12} \left( \frac{y}{x} \right) \mathbf{K}_1(x) \bar{\mathbf{L}}_{12}(xy) K_2(y) = K_2(y) \mathbf{L}_{12} \left( \frac{1}{xy} \right) \mathbf{K}_1(x) \bar{\mathbf{L}}_{12} \left( \frac{x}{y} \right). \quad (3.3)$$

The reflection equation (3.1) is the image of (3.3) for  $\pi \otimes 1$ . Expanding (3.3) with respect to  $y$ , we obtain

$$\sum_{k=1}^N x^{-2\xi_k} \left( L_{ik}^+ \mathbf{K}(x) \bar{L}_{kj}^+ - L_{ik}^- \mathbf{K}(x) \bar{L}_{kj}^- \right) = 0 \quad \text{for } i, j \leq a \quad \text{or} \quad i, j > a, \quad (3.4)$$

$$\sum_{k=1}^N x^{-2\xi_k} \left\{ \left( x^s L_{ik}^+ \mathbf{K}(x) \bar{L}_{kj}^- + x^{-s} L_{ik}^- \mathbf{K}(x) \bar{L}_{kj}^+ \right) \epsilon_+ + L_{ik}^- \mathbf{K}(x) \bar{L}_{kj}^- \epsilon_- \right\} = 0$$

for  $i \leq a < j$ , (3.5)

$$\sum_{k=1}^N x^{-2\xi_k} \left\{ \left( x^s L_{ik}^+ \mathbf{K}(x) \bar{L}_{kj}^- + x^{-s} L_{ik}^- \mathbf{K}(x) \bar{L}_{kj}^+ \right) \epsilon_+ + L_{ik}^+ \mathbf{K}(x) \bar{L}_{kj}^+ \epsilon_- \right\} = 0$$

for  $j \leq a < i$ . (3.6)

One can rewrite <sup>7</sup> these in terms of  $e_{ij}$  and  $\bar{e}_{ij}$ :

$$q^{2e_{ii}} \mathbf{K}(x) = \mathbf{K}(x) q^{2e_{ii}}, \quad (3.7)$$

$$x^{-2\xi_j} e_{ji} \mathbf{K}(x) - (q - q^{-1}) \sum_{k=j+1}^{i-1} x^{-2\xi_k} e_{ki} \mathbf{K}(x) \bar{e}_{jk} - x^{-2\xi_i} \mathbf{K}(x) \bar{e}_{ji} = 0$$

for  $j < i \leq a$  or  $a < j < i$ , (3.8)

---

<sup>7</sup>The commutation relations for the Cartan elements from (3.4) are of the form  $q^{e_{ii}} \mathbf{K}(x) q^{-e_{ii}} = q^{-e_{ii}} \mathbf{K}(x) q^{e_{ii}}$ , from which  $e_{ii} \mathbf{K}(x) = \mathbf{K}(x) e_{ii}$  are deduced (under expansion of  $e_{ii} = \log q^{e_{ii}} / \log q$ ). We have used these to simplify (3.8)-(3.11).

$$x^{-2\xi_j} e_{ji} \mathbf{K}(x) + (q - q^{-1}) \sum_{k=i+1}^{j-1} x^{-2\xi_k} e_{ki} \mathbf{K}(x) \bar{e}_{jk} - x^{-2\xi_i} \mathbf{K}(x) \bar{e}_{ji} = 0$$

for  $i < j \leq a$  or  $a < i < j$ ,

(3.9)

$$x^{-2\xi_j} e_{ji} (\epsilon_+ x^{-s} q^{-2e_{jj}} + \epsilon_-) \mathbf{K}(x) - (q - q^{-1}) \left( \epsilon_+ x^s \sum_{k=1}^{i-1} x^{-2\xi_k} q^{e_{ii}} e_{ki} q^{e_{kk}} \mathbf{K}(x) \bar{e}_{jk} \right. \\ \left. + \epsilon_+ x^{-s} \sum_{k=j+1}^N x^{-2\xi_k} e_{ki} \mathbf{K}(x) q^{-e_{kk}} \bar{e}_{jk} q^{-e_{jj}} - \epsilon_- \sum_{k=i+1}^{j-1} x^{-2\xi_k} e_{ki} \mathbf{K}(x) \bar{e}_{jk} \right) \\ - x^{-2\xi_i} \mathbf{K}(x) \bar{e}_{ji} (\epsilon_+ x^s q^{2e_{ii}-2} + \epsilon_-) = 0 \quad \text{for } i \leq a < j,$$
(3.10)

$$x^{-2\xi_j} e_{ji} (\epsilon_+ x^s q^{2e_{jj}} + \epsilon_-) \mathbf{K}(x) + (q - q^{-1}) \left( \epsilon_+ x^s \sum_{k=1}^{j-1} x^{-2\xi_k} e_{ki} q^{e_{kk}} \mathbf{K}(x) \bar{e}_{jk} q^{e_{jj}} \right. \\ \left. + \epsilon_+ x^{-s} \sum_{k=i+1}^N x^{-2\xi_k} q^{-e_{ii}} e_{ki} \mathbf{K}(x) q^{-e_{kk}} \bar{e}_{jk} - \epsilon_- \sum_{k=j+1}^{i-1} x^{-2\xi_k} e_{ki} \mathbf{K}(x) \bar{e}_{jk} \right) \\ - x^{-2\xi_i} \mathbf{K}(x) \bar{e}_{ji} (\epsilon_+ x^{-s} q^{-2e_{ii}+2} + \epsilon_-) = 0 \quad \text{for } j \leq a < i.$$
(3.11)

For  $i$  and  $j$  satisfying  $|i - j| = 1$ , the relations (3.8) and (3.9) reduce to

$$x^{-2\xi_j} e_{j,j+1} \mathbf{K}(x) = x^{-2\xi_{j+1}} \mathbf{K}(x) e_{j,j+1}, \quad x^{-2\xi_{j+1}} e_{j+1,j} \mathbf{K}(x) = x^{-2\xi_j} \mathbf{K}(x) e_{j+1,j},$$

for  $j \neq a$ .

(3.12)

Taking note on (2.8) and (2.9) (and (3.7)), one can derive

$$x^{-2\xi_j} \bar{e}_{ji} \mathbf{K}(x) = x^{-2\xi_i} \mathbf{K}(x) \bar{e}_{ji} \quad \text{for } i, j \leq a \text{ or } a < i, j,$$
(3.13)

$$x^{-2\xi_j} e_{ji} \mathbf{K}(x) = x^{-2\xi_i} \mathbf{K}(x) e_{ji} \quad \text{for } i, j \leq a \text{ or } a < i, j$$
(3.14)

from (3.12). Then the relations (3.8) and (3.9) boil down to the relations

$$e_{ji} - (q - q^{-1}) \sum_{k=j+1}^{i-1} e_{ki} \bar{e}_{jk} - \bar{e}_{ji} = 0 \quad \text{for } j < i,$$
(3.15)

$$e_{ji} + (q - q^{-1}) \sum_{k=i+1}^{j-1} e_{ki} \bar{e}_{jk} - \bar{e}_{ji} = 0 \quad \text{for } i < j,$$
(3.16)

which are special cases of (A2) and (A3), under (3.13) and (3.14). Thus it suffices to consider (3.12) instead of (3.8) and (3.9). We further rewrite (3.10) and (3.11) under



the relations (3.13) and (3.14), in the form of intertwining relations:

$$\begin{aligned}
x^{-2\xi_j}(\epsilon_+x^{-s}A_{ji} + \epsilon_-B_{ji})\mathbf{K}(x) &= x^{-2\xi_i}\mathbf{K}(x)(\epsilon_+x^sC_{ji} + \epsilon_-D_{ji}), \\
A_{ji} &= e_{ji}q^{-2e_{jj}} - (q - q^{-1}) \sum_{k=j+1}^N e_{ki}\bar{e}_{jk}q^{-e_{kk}-e_{jj}+1}, \\
B_{ji} &= e_{ji} + (q - q^{-1}) \sum_{k=a+1}^{j-1} e_{ki}\bar{e}_{jk}, \\
C_{ji} &= \bar{e}_{ji}q^{2e_{ii}-2} + (q - q^{-1}) \sum_{k=1}^{i-1} e_{ki}\bar{e}_{jk}q^{e_{kk}+e_{ii}-2}, \\
D_{ji} &= \bar{e}_{ji} - (q - q^{-1}) \sum_{k=i+1}^a e_{ki}\bar{e}_{jk} \quad \text{for } i \leq a < j, \tag{3.17}
\end{aligned}$$

$$\begin{aligned}
x^{-2\xi_j}(\epsilon_+x^sA_{ji} + \epsilon_-B_{ji})\mathbf{K}(x) &= x^{-2\xi_i}\mathbf{K}(x)(\epsilon_+x^{-s}C_{ji} + \epsilon_-D_{ji}), \\
A_{ji} &= e_{ji}q^{2e_{jj}} + (q - q^{-1}) \sum_{k=1}^{j-1} e_{ki}\bar{e}_{jk}q^{e_{kk}+e_{jj}-1}, \\
B_{ji} &= e_{ji} - (q - q^{-1}) \sum_{k=j+1}^a e_{ki}\bar{e}_{jk}, \\
C_{ji} &= \bar{e}_{ji}q^{-2e_{ii}+2} - (q - q^{-1}) \sum_{k=i+1}^N e_{ki}\bar{e}_{jk}q^{-e_{kk}-e_{ii}+2}, \\
D_{ji} &= \bar{e}_{ji} + (q - q^{-1}) \sum_{k=a+1}^{i-1} e_{ki}\bar{e}_{jk} \quad \text{for } j \leq a < i. \tag{3.18}
\end{aligned}$$

We remark that the relation  $B_{ji} = D_{ji}$  holds due to (A2) and (A3). We find that the following operators <sup>8</sup> satisfy (3.7), (3.12), (3.17) and (3.18) for the  $q$ -oscillator representation (2.13), and thus solve the reflection equation (3.3).

$$\begin{aligned}
\mathbf{K}(x) &= q^{2(c-1)\sum_{k=1}^a e_{kk}} x^{2\sum_{k=1}^N (s\theta(1 \leq k \leq a) - \xi_k) e_{kk}} \frac{\left(-\frac{\epsilon_-}{\epsilon_+} x^s q^{2\sum_{k=a+1}^N e_{kk}-2}; q^{-2}\right)_{\infty}}{\left(-\frac{\epsilon_-}{\epsilon_+} x^{-s} q^{-2\sum_{k=1}^a e_{kk}}; q^{-2}\right)_{\infty}} \\
&\quad \text{for } |q| > 1, \text{ or} \tag{3.19} \\
&= q^{2(c-1)\sum_{k=1}^a e_{kk}} x^{2\sum_{k=1}^N (s\theta(1 \leq k \leq a) - \xi_k) e_{kk}} \frac{\left(-\frac{\epsilon_-}{\epsilon_+} x^{-s} q^{-2\sum_{k=1}^a e_{kk}+2}; q^2\right)_{\infty}}{\left(-\frac{\epsilon_-}{\epsilon_+} x^s q^{2\sum_{k=a+1}^N e_{kk}}; q^2\right)_{\infty}}
\end{aligned}$$

---

<sup>8</sup>In 2016, we were informed by S. Belliard that he found a (non-diagonal) solution for the rational case  $Y(sl(2))$ .

$$\text{for } |q| < 1, \text{ or} \quad (3.20)$$

$$= x^{-2\sum_{k=1}^N \xi_k e_{kk}} \frac{\left(-\frac{\epsilon_+}{\epsilon_-} x^s q^{2\sum_{k=1}^a e_{kk}-2}; q^{-2}\right)_\infty}{\left(-\frac{\epsilon_+}{\epsilon_-} x^{-s} q^{-2\sum_{k=a+1}^N e_{kk}}; q^{-2}\right)_\infty} \text{ for } |q| > 1, \text{ or} \quad (3.21)$$

$$= x^{-2\sum_{k=1}^N \xi_k e_{kk}} \frac{\left(-\frac{\epsilon_+}{\epsilon_-} x^{-s} q^{-2\sum_{k=a+1}^N e_{kk}+2}; q^2\right)_\infty}{\left(-\frac{\epsilon_+}{\epsilon_-} x^s q^{2\sum_{k=1}^a e_{kk}}; q^2\right)_\infty} \text{ for } |q| < 1, \quad (3.22)$$

where  $c = \sum_{k=1}^N e_{kk}$ ,  $\theta(\text{True}) = 1$ ,  $\theta(\text{False}) = 0$ , and  $(x; q)_\infty = \prod_{j=0}^{\infty} (1 - xq^j)$ . One can directly check that (3.19)-(3.22) satisfy (3.7) and (3.12) for the generic generators of  $U_q(\mathfrak{gl}(N))$ . One can also show that the operators (3.19)-(3.22) satisfy the following relations

$$\mathbf{K}(x)\mathbf{E}_{ji} = x^{2(\xi_i - \xi_j)} \mathbf{E}_{ji}\mathbf{K}(x) \left( \frac{\epsilon_+ x^s q^{2\sum_{k=1}^a e_{kk}} + \epsilon_-}{\epsilon_+ x^{-s} q^{-2\sum_{k=a+1}^N e_{kk}+2} + \epsilon_-} \right) \text{ for } j \leq a < i, \quad (3.23)$$

$$\mathbf{K}(x)\mathbf{E}_{ji} = x^{2(\xi_i - \xi_j)} \mathbf{E}_{ji}\mathbf{K}(x) \left( \frac{\epsilon_+ x^{-s} q^{-2\sum_{k=a+1}^N e_{kk}} + \epsilon_-}{\epsilon_+ x^s q^{2\sum_{k=1}^a e_{kk}-2} + \epsilon_-} \right) \text{ for } i \leq a < j, \quad (3.24)$$

where  $\mathbf{E}_{ji}$  is any operator obeying  $q^{e_{kk}} \mathbf{E}_{ji} q^{-e_{kk}} = q^{\delta_{kj} - \delta_{ki}} \mathbf{E}_{ji}$ . Thus (3.17) and (3.18) reduce to

$$B_{ji} = A_{ji} q^{2\sum_{k=a+1}^N e_{kk}} = C_{ji} q^{-2\sum_{k=1}^a e_{kk}+2} \text{ for } i \leq a < j, \quad (3.25)$$

$$B_{ji} = A_{ji} q^{-2\sum_{k=1}^a e_{kk}} = C_{ji} q^{2\sum_{k=a+1}^N e_{kk}-2} \text{ for } j \leq a < i. \quad (3.26)$$

These relations (3.25) and (3.26) give a constraint on the class of representations of  $U_q(\mathfrak{gl}(N))$ . In other words, we have to consider the quotient of  $U_q(\mathfrak{gl}(N))$  by the relations (3.25) and (3.26). Moreover, we find that the first evaluation map (2.10) is almost<sup>9</sup> equivalent to the second one (2.11) under these:

$$\text{ev}_x(e_N) = q^{-2(c-1)} \overline{\text{ev}}_x(e_N), \quad \text{ev}_x(f_N) = q^{2(c-1)} \overline{\text{ev}}_x(f_N) \quad [\text{under (3.25) and (3.26)}]. \quad (3.27)$$

One can check that (2.13) satisfies (3.25) and (3.26). In addition, the fundamental representation of  $U_q(\mathfrak{gl}(N))$ , which is a quotient of (2.13) at  $m = 1$ , also fulfills (3.25) and (3.26). Then we recover (3.2) from (3.19):

$$\pi(\mathbf{K}(x)) = \frac{x^{-s} \left(-\frac{\epsilon_-}{\epsilon_+} x^s q^{-2}; q^{-2}\right)_\infty}{\epsilon_+ \left(-\frac{\epsilon_-}{\epsilon_+} x^{-s}; q^{-2}\right)_\infty} K(x). \quad (3.28)$$

---

<sup>9</sup>The factor  $q^{\mp 2(c-1)}$  in the right hand side can be eliminated if one considers composition of the automorphism  $e_i \mapsto q^{-2(\hat{c}-1)\delta_i} e_i$ ,  $f_i \mapsto q^{2(\hat{c}-1)\delta_i} f_i$ ,  $k_i \mapsto k_i$  of  $U_q(\mathfrak{gl}(N))$  and the evaluation map  $\overline{\text{ev}}_x$ .

It is important to note that the relations (3.25) and (3.26) become trivial for  $N = 2$  case. This means that the K-operators (3.19)-(3.22) written in terms of the generic Cartan elements solve the reflection equation (3.3) without any constraint on the representations for  $U_q(\mathfrak{gl}(2))$  case. In fact, (3.19) for  $N = 2$  reproduces <sup>10</sup> the K-operator proposed in [5].

Let us consider a product of the L-operators

$$\begin{aligned} \mathbf{L}(x)q^{\frac{2(c-1)}{s}\sum_{j=1}^N \xi_j e_{jj}} \bar{\mathbf{L}}(xq^{-\frac{2(c-1)}{s}})q^{-\frac{2(c-1)}{s}\sum_{j=1}^N \xi_j e_{jj}} = \\ = \sum_{i=1}^N (-x^s - x^{-s} + G_i) \otimes E_{ii} + (q - q^{-1}) \sum_{i < j} (C_{ji} - A_{ji}q^{2(c-1)})q^{e_{jj} - e_{ii} + 1} \otimes E_{ij} \\ + (q - q^{-1}) \sum_{i > j} (A_{ji} - C_{ji}q^{2(c-1)}) \otimes E_{ij}, \end{aligned} \quad (3.29)$$

where  $G_i = G_i^{(+)} + G_i^{(-)}$  is defined by

$$G_i^{(+)} = q^{2e_{ii}} + (q - q^{-1})^2 \sum_{k=1}^{i-1} e_{ki} \bar{e}_{ik} q^{e_{kk} + e_{ii} - 1}, \quad (3.30)$$

$$G_i^{(-)} = q^{2(c-1)} \left( q^{-2e_{ii}} + (q - q^{-1})^2 \sum_{k=i+1}^N e_{ki} \bar{e}_{ik} q^{-e_{kk} - e_{ii} + 1} \right). \quad (3.31)$$

Here  $A_{ji}$  and  $C_{ji}$  for  $i < j$  (resp.  $i > j$ ) are the ones in (3.17) (resp. (3.18)) [dependence on the parameter  $a$  is irrelevant]. In deriving (3.29), we have used (A2) and (A3). Note that (3.29) becomes diagonal under the relations

$$A_{ji}q^{2(c-1)} = C_{ji} \quad \text{for } i < j, \quad \text{and} \quad A_{ji} = C_{ji}q^{2(c-1)} \quad \text{for } j < i. \quad (3.32)$$

We remark that a part of the relations (3.25) and (3.26) automatically follows from (3.32). For  $N = 2$ , the relation (3.32) becomes trivial, and the diagonal elements  $G_1$  and  $G_2$  reduce to a Casimir element of  $U_q(\mathfrak{gl}(2))$  [cf. eq. (3.5) in [5]]. On the other hand, for  $N \geq 3$  case, each diagonal element is not a Casimir element, but rather twisted sums <sup>11</sup>  $\sum_{k=1}^N G_k^{(+)} q^{2k}$ ,  $\sum_{k=1}^N G_k^{(-)} q^{2k}$  and  $\sum_{k=1}^N G_k q^{2k}$  are (cf. eq. (41) in [24]). However, there are cases where simplifications occur for particular representations. In fact, one can check that the q-oscillator representation (2.13) satisfies the condition (3.32), and  $G_i = q^{2m} + q^{-2}$  holds for any  $i$ . This is one of the desirable properties for construction of mutually commuting transfer matrices.

## 4 Augmented q-Onsager algebra like symmetry

In this section, we will reconsider the intertwining relations in the previous section and point out their connection to the generators of  $U_q(\hat{\mathfrak{gl}}(N))$ . We will also mention our

<sup>10</sup>  $x^{s_2 c} q^{-\frac{(c-1)((s+s_2)c+s_1 H)}{s}} \mathbf{K}(xq^{-\frac{c-1}{s}})$  for  $N = 2$  coincides with eq. (4.9) in [5].

<sup>11</sup>  $\sum_{k=1}^N G_k q^{2k}$  is a part of the twisted trace of (3.29) (by  $1 \otimes \pi(q^{2\sum_{k=1}^N k e_{kk}})$ ).

observation on an underlying symmetry behind them.

We find that the intertwining relations (3.13), (3.14), (3.17) and (3.18) can be written in terms of the generators of  $U_q(\hat{gl}(N))$ :

$$\text{ev}_{x^{-1}}(Z_{ji})\mathbf{K}(x) = \mathbf{K}(x)\overline{\text{ev}}_x(Z_{ji}), \quad (4.1)$$

where  $Z_{ji}$  is defined as follows:

For  $1 \leq i = j \leq N$ ,

$$Z_{ii} = k_i. \quad (4.2)$$

For  $1 \leq j < i \leq a$  or  $a+1 \leq j < i \leq N$ ,

$$Z_{ji} = e_{[j,i-1]}. \quad (4.3)$$

For  $1 \leq i < j \leq a$  or  $a+1 \leq i < j \leq N$ ,

$$Z_{ji} = f_{[i,j-1]}. \quad (4.4)$$

For  $j \leq a < i$ ,

$$Z_{ji} = \epsilon_+ Z_{ji}^+ + \epsilon_- Z_{ji}^-,$$

$$Z_{ji}^+ = \begin{cases} [[\bar{f}_{[1,j-1]}, f_N]_{q^{-1}}, f_{[i,N-1]}]_q q^{k_j - k_i + 1} & \text{for } 2 \leq j \leq a < i \leq N-1 \\ [f_N, f_{[i,N-1]}]_q q^{k_1 - k_i + 1} & \text{for } 1 = j \leq a < i \leq N-1 \\ [\bar{f}_{[1,j-1]}, f_N]_{q^{-1}} q^{k_j - k_N + 1} & \text{for } 2 \leq j \leq a < i = N \\ f_N q^{k_1 - k_N + 1} & \text{for } 1 = j \leq a < i = N, \end{cases} \quad (4.5)$$

$$Z_{ji}^- = \begin{cases} [\bar{e}_{[j,a]}, e_{[a+1,i-1]}]_{q^{-1}} & \text{for } 1 \leq j \leq a < a+1 < i \leq N \\ \bar{e}_{[j,a]} & \text{for } 1 \leq j \leq a < a+1 = i \leq N \\ e_{[a,i-1]} & \text{for } 1 \leq j = a < i \leq N. \end{cases} \quad (4.6)$$

For  $i \leq a < j$ :

$$Z_{ji} = \epsilon_+ Z_{ji}^+ + \epsilon_- Z_{ji}^-,$$

$$Z_{ji}^+ = \begin{cases} [[\bar{e}_{[j,N-1]}, [e_N, e_{[1,i-1]}]_{q^{-1}}]_q & \text{for } 2 \leq i \leq a < j \leq N-1 \\ [\bar{e}_{[j,N-1]}, e_N]_q & \text{for } 1 = i \leq a < j \leq N-1 \\ [e_N, e_{[1,i-1]}]_{q^{-1}} & \text{for } 2 \leq i \leq a < j = N \\ e_N & \text{for } 1 = i \leq a < j = N, \end{cases} \quad (4.7)$$

$$Z_{ji}^- = \begin{cases} [\bar{f}_{[a+1,j-1]}, f_{[i,a]}]_{q^{-1}} q^{k_j - k_i + 1} & \text{for } 1 \leq i \leq a < a+1 < j \leq N \\ f_{[i,a]} q^{k_{a+1} - k_i + 1} & \text{for } 1 \leq i \leq a < a+1 = j \leq N \\ \bar{f}_{[a,j-1]} q^{k_j - k_a + 1} & \text{for } 1 \leq i = a < j \leq N. \end{cases} \quad (4.8)$$

Here we use the following notation for root vectors. For  $i < j$ , we define

$$\begin{aligned}
e_{[i,j]} &= [e_i, [e_{i+1}, \dots, [e_{j-2}, [e_{j-1}, e_j]_{q^{-1}}]_{q^{-1}} \dots]_{q^{-1}}]_{q^{-1}}, \\
\bar{e}_{[i,j]} &= [e_i, [e_{i+1}, \dots, [e_{j-2}, [e_{j-1}, e_j]_q]_q \dots]_q]_q, \\
f_{[i,j]} &= [f_j, [f_{j-1}, \dots, [f_{i+2}, [f_{i+1}, f_i]_q]_q \dots]_q]_q, \\
\bar{f}_{[i,j]} &= [f_j, [f_{j-1}, \dots, [f_{i+2}, [f_{i+1}, f_i]_{q^{-1}}]_{q^{-1}} \dots]_{q^{-1}}]_{q^{-1}},
\end{aligned} \tag{4.9}$$

and  $e_{[i,i]} = \bar{e}_{[i,i]} = e_i$ ,  $f_{[i,i]} = \bar{f}_{[i,i]} = f_i$ . We remark that the third case in (4.6) (resp. (4.8)) is a special case of the first or the second ones. In deriving these, we have used (A2)-(A10).

We introduce generators composed of Cartan elements

$$\mathfrak{k}_{ji}^+ = \epsilon_+ q^{\sum_{l=1}^j k_l - \sum_{l=i}^N k_l}, \quad \mathfrak{k}_{ji}^- = \epsilon_- q^{-\sum_{l=j}^a k_l + \sum_{l=a+1}^i k_l} \quad \text{for } 1 \leq j \leq a < i \leq N, \tag{4.10}$$

and use notation  $Z_{a,a+1} = \mathfrak{e}_a$ ,  $Z_{N,1} = \mathfrak{e}_N$ ,  $Z_{a+1,a} = \mathfrak{f}_a$ ,  $Z_{1,N} = \mathfrak{f}_N$ ,  $\mathfrak{k}_a^+ = \mathfrak{k}_{a,a+1}^+$ ,  $\mathfrak{k}_a^- = \mathfrak{k}_{a,a+1}^-$ ,  $\mathfrak{k}_N^+ = \mathfrak{k}_{1,N}^+$ ,  $\mathfrak{k}_N^- = \mathfrak{k}_{1,N}^-$ . We have observed<sup>12</sup> the following commutation relations among generators<sup>13</sup> for the  $q$ -oscillator representation of  $U_q(\hat{\mathfrak{gl}}(N))$  under the evaluation map (2.10) and (2.13), in addition to the relations (2.1) restricted to the generators  $\{e_i, f_i\}_{i \neq a, N}$  and  $\{k_i\}_{i=1}^N$ .

$$[\mathfrak{k}_{ji}^+, \mathfrak{k}_{rs}^-] = [\mathfrak{k}_{ji}^\pm, \mathfrak{k}_{rs}^\pm] = [\mathfrak{k}_{ji}^\pm, k_l] = 0 \quad \text{for } 1 \leq j, r \leq a < i, s \leq N, \quad 1 \leq l \leq N; \tag{4.11}$$

$$[k_l, Z_{ji}] = (\delta_{lj} - \delta_{il})Z_{ji} \quad \text{for } 1 \leq l, i, j \leq N; \tag{4.12}$$

$$\begin{aligned}
\mathfrak{k}_{ji}^+ Z_{rs} &= q^{\theta(1 \leq r \leq j) - \theta(i \leq r \leq N) - \theta(1 \leq s \leq j) + \theta(i \leq s \leq N)} Z_{rs} \mathfrak{k}_{ji}^+, \\
\mathfrak{k}_{ji}^- Z_{rs} &= q^{-\theta(j \leq r \leq a) + \theta(a+1 \leq r \leq i) + \theta(j \leq s \leq a) - \theta(a+1 \leq s \leq i)} Z_{rs} \mathfrak{k}_{ji}^-
\end{aligned} \tag{4.13}$$

for  $1 \leq j \leq a < i \leq N, \quad 1 \leq r, s \leq N;$

$$\begin{aligned}
[\mathfrak{e}_a, [\mathfrak{e}_a, [\mathfrak{e}_a, \mathfrak{f}_a]_{q^2}]]_{q^{-2}} &= \tilde{\rho} \mathfrak{e}_a ((\mathfrak{k}_a^+)^2 - (\mathfrak{k}_a^-)^2) \mathfrak{e}_a, \\
[\mathfrak{e}_N, [\mathfrak{e}_N, [\mathfrak{e}_N, \mathfrak{f}_N]_{q^2}]]_{q^{-2}} &= \tilde{\rho} \mathfrak{e}_N ((\mathfrak{k}_N^-)^2 - (\mathfrak{k}_N^+)^2) \mathfrak{e}_N, \\
[\mathfrak{f}_a, [\mathfrak{f}_a, [\mathfrak{f}_a, \mathfrak{e}_a]_{q^2}]]_{q^{-2}} &= \tilde{\rho} \mathfrak{f}_a ((\mathfrak{k}_a^-)^2 - (\mathfrak{k}_a^+)^2) \mathfrak{f}_a, \\
[\mathfrak{f}_N, [\mathfrak{f}_N, [\mathfrak{f}_N, \mathfrak{e}_N]_{q^2}]]_{q^{-2}} &= \tilde{\rho} \mathfrak{f}_N ((\mathfrak{k}_N^+)^2 - (\mathfrak{k}_N^-)^2) \mathfrak{f}_N \quad \text{for } 1 \leq a \leq N-1;
\end{aligned} \tag{4.14}$$

$$\begin{aligned}
[\mathfrak{e}_a, [\mathfrak{e}_a, [\mathfrak{e}_a, \mathfrak{e}_N]_{q^2}]]_{q^{-2}} &= [\mathfrak{f}_a, [\mathfrak{f}_a, [\mathfrak{f}_a, \mathfrak{f}_N]_{q^2}]]_{q^{-2}} = 0, \\
[\mathfrak{e}_N, [\mathfrak{e}_N, [\mathfrak{e}_N, \mathfrak{e}_a]_{q^2}]]_{q^{-2}} &= [\mathfrak{f}_N, [\mathfrak{f}_N, [\mathfrak{f}_N, \mathfrak{f}_a]_{q^2}]]_{q^{-2}} = 0, \\
[\mathfrak{f}_N, \mathfrak{e}_a]_{q^{-2}} &= [\mathfrak{e}_N, \mathfrak{f}_a]_{q^2} = 0 \quad \text{for } 2 \leq a \leq N-2;
\end{aligned} \tag{4.15}$$

<sup>12</sup>We have checked these by Mathematica.

<sup>13</sup>Note that the  $q$ -commutators  $[A, [A, B]_{q_1}]_{q_2}$  and  $[A, [A, [A, B]_{q_1}]_{q_2}]_{q_3}$  are invariant under the permutation on  $\{q_1, q_2\}$  and  $\{q_1, q_2, q_3\}$ , respectively.

$$\begin{aligned}
[e_{a-1}, [e_{a-1}, \mathbf{e}_a]_q]_{q^{-1}} &= [\mathbf{e}_a, [e_a, e_{a-1}]_q]_{q^{-1}} = [f_{a-1}, [f_{a-1}, \mathbf{f}_a]]_{q^2} = [\mathbf{f}_a, [f_a, f_{a-1}]]_{q^{-2}} = 0, \\
[e_1, [e_1, \mathbf{e}_N]_q]_{q^{-1}} &= [\mathbf{e}_N, [e_N, e_1]_q]_{q^{-1}} = [f_1, [f_1, \mathbf{f}_N]]_{q^2} = [\mathbf{f}_N, [f_N, f_1]]_{q^{-2}} = 0, \\
[e_{a-1}, \mathbf{f}_a]_{q^{-1}} &= [e_1, \mathbf{f}_N]_{q^{-1}} = 0 \quad \text{for } 2 \leq a \leq N-1;
\end{aligned} \tag{4.16}$$

$$\begin{aligned}
[e_{a+1}, [e_{a+1}, \mathbf{e}_a]_q]_{q^{-1}} &= [\mathbf{e}_a, [e_a, e_{a+1}]_q]_{q^{-1}} = [f_{a+1}, [f_{a+1}, \mathbf{f}_a]]_{q^2} = [\mathbf{f}_a, [f_a, f_{a+1}]]_{q^{-2}} = 0, \\
[e_{N-1}, [e_{N-1}, \mathbf{e}_N]_q]_{q^{-1}} &= [\mathbf{e}_N, [e_N, e_{N-1}]_q]_{q^{-1}} = 0, \\
[f_{N-1}, [f_{N-1}, \mathbf{f}_N]]_{q^2} &= [\mathbf{f}_N, [f_N, f_{N-1}]]_{q^{-2}} = 0, \\
[e_{a+1}, \mathbf{f}_a]_{q^{-1}} &= [e_{N-1}, \mathbf{f}_N]_{q^{-1}} = 0 \quad \text{for } 1 \leq a \leq N-2;
\end{aligned} \tag{4.17}$$

$$[e_i, \mathbf{e}_a] = [e_i, \mathbf{f}_a] = [f_i, \mathbf{f}_a] = 0 \quad \text{for } 1 \leq i \leq a-2 \quad \text{or } a+2 \leq i \leq N-1; \tag{4.18}$$

$$[e_i, \mathbf{e}_N] = [e_i, \mathbf{f}_N] = [f_i, \mathbf{f}_N] = 0 \quad \text{for } 2 \leq i \leq a-1 \quad \text{or } a+1 \leq i \leq N-2; \tag{4.19}$$

$$[f_i, \mathbf{e}_a] = [f_i, \mathbf{e}_N] = 0 \quad \text{for } 1 \leq i \leq a-1 \quad \text{or } a+1 \leq i \leq N-1; \tag{4.20}$$

$$[e_{i-1}, Z_{ij}]_q = Z_{i-1,j}, \quad [Z_{ji}, f_{i-1}] = Z_{j,i-1} q^{-1+k_{i-1}-k_i} \quad \text{for } 2 \leq i \leq a < j \leq N; \tag{4.21}$$

$$[Z_{ij}, e_j]_{q^{-1}} = Z_{i,j+1}, \quad [f_j, Z_{ji}] = Z_{j+1,i} q^{k_j-k_{j+1}} \quad \text{for } 1 \leq i \leq a < j \leq N-1; \tag{4.22}$$

$$\begin{aligned}
[f_1, [f_1, [f_1, \mathbf{f}_N]_{q^3}]_q]_{q^{-1}} &= q^{-1} \rho \mathbf{f}_1 (\mathbf{e}_1^-)^2 \mathbf{f}_1 Z_{2N}, \\
[f_N, [f_N, [f_N, \mathbf{f}_1]_{q^{-3}}]_{q^{-1}}]_q &= -q^{-4} \rho \mathbf{f}_N (\mathbf{e}_1^-)^2 \mathbf{f}_N Z_{2N},
\end{aligned} \tag{4.23}$$

$$[\mathbf{e}_1, [\mathbf{e}_1, [\mathbf{e}_1, \mathbf{e}_N]_{q^3}]_q]_{q^{-1}} = -q^5 \rho \mathbf{e}_1 (\mathbf{e}_N^+)^2 \mathbf{e}_1 Z_{N2} q^{k_N-k_2},$$

$$[\mathbf{e}_N, [\mathbf{e}_N, [\mathbf{e}_N, \mathbf{e}_1]_{q^{-3}}]_{q^{-1}}]_q = q^2 \rho \mathbf{e}_N (\mathbf{e}_N^+)^2 \mathbf{e}_N Z_{N2} q^{k_N-k_2} \quad \text{for } a=1, \quad N > 2;$$

$$[f_{N-1}, [f_{N-1}, [f_{N-1}, \mathbf{f}_N]_{q^3}]_q]_{q^{-1}} = -q^2 \rho \mathbf{f}_{N-1} (\mathbf{e}_N^-)^2 \mathbf{f}_{N-1} Z_{1,N-1} q^{2k_1},$$

$$[f_N, [f_N, [f_N, \mathbf{f}_{N-1}]_{q^{-3}}]_{q^{-1}}]_q = q \rho \mathbf{f}_N (\mathbf{e}_N^-)^2 \mathbf{f}_N Z_{1,N-1} q^{2k_1},$$

$$[\mathbf{e}_{N-1}, [\mathbf{e}_{N-1}, [\mathbf{e}_{N-1}, \mathbf{e}_N]_{q^3}]_q]_{q^{-1}} = \rho \mathbf{e}_{N-1} (\mathbf{e}_{N-1}^+)^2 \mathbf{e}_{N-1} Z_{N-1,1} q^{-k_1-k_{N-1}}, \tag{4.24}$$

$$\begin{aligned}
[\mathbf{e}_N, [\mathbf{e}_N, [\mathbf{e}_N, \mathbf{e}_{N-1}]_{q^{-3}}]_{q^{-1}}]_q &= -q^{-1} \rho \mathbf{e}_N (\mathbf{e}_{N-1}^+)^2 \mathbf{e}_N Z_{N-1,1} q^{-k_1-k_{N-1}} \\
&\quad \text{for } a=N-1, \quad N > 2,
\end{aligned}$$

where  $\rho = (q^3 - q^{-3})(q^2 - q^{-2})$ ,  $\tilde{\rho} = \rho/(q - q^{-1})$ . We remark that the parameter  $a$  is a fixed parameter in the diagonal K-matrix (3.2). These commutation relations are valid without any restriction of the class of representations for  $N = 2$  case, and define the so-called augmented q-Onsager algebra [6, 7], a realization of which in terms of the generators of  $U_q(\hat{sl}(2))$  is known in [7]. In contrast, not all of them hold true for

generic representations for  $N \geq 3$  case. Whether the elements  $Z_{ji}$  for  $N \geq 3$  satisfy closed commutation relations on the level of the algebra remains to be clarified.

## 5 Rational limit

In this section, we consider the rational limit of some of the formulas in previous sections.

Let  $\epsilon_{\pm} = \mp q^{-2p_{\pm}}$ ,  $p = p_+ - p_-$ , and  $u \in \mathbb{C}$ . The rational limit of the L-operators (2.16), (2.17), the K-matrix (3.2), the K-operators (3.19)-(3.22), the intertwining relations (3.13), (3.14), (3.17), and (3.18) are given, respectively, by

$$\mathbf{L}(u) = \lim_{q \rightarrow 1} (q - q^{-1})^{-1} \mathbf{L}(q^{-2u}) = \sum_{i,j} (su\delta_{ij} + e_{ji}) \otimes E_{ij}, \quad (5.1)$$

$$\bar{\mathbf{L}}(u) = \lim_{q \rightarrow 1} (q - q^{-1})^{-1} \bar{\mathbf{L}}(q^{-2u}) = \sum_{i,j} (-su\delta_{ij} + e_{ji}) \otimes E_{ij}, \quad (5.2)$$

$$K(u) = \lim_{q \rightarrow 1} (q - q^{-1})^{-1} K(q^{-2u}) = \sum_{k=1}^a (su + p) E_{kk} + \sum_{k=a+1}^N (-su + p) E_{kk}, \quad (5.3)$$

$$\begin{aligned} \mathbf{K}(u) &= \lim_{|q| \rightarrow 1 \pm 0} \mathbf{K}(q^{-2u}) (1 - q^{\mp 2})^{1-c+2su} \\ &= \frac{\Gamma(-su - p + \sum_{k=1}^a e_{kk})}{\Gamma(su - p + 1 - \sum_{k=a+1}^N e_{kk})} \quad \text{from (3.19) or (3.22), or} \end{aligned} \quad (5.4)$$

$$= \frac{\Gamma(-su + p + \sum_{k=a+1}^N e_{kk})}{\Gamma(su + p + 1 - \sum_{k=1}^a e_{kk})} \quad \text{from (3.20) or (3.21),} \quad (5.5)$$

$$e_{ji} \mathbf{K}(u) = \mathbf{K}(u) e_{ji} \quad \text{for } i, j \leq a \quad \text{or } i, j \geq a + 1, \quad (5.6)$$

$$\begin{aligned} \left( (su - p) e_{ji} - \sum_{k=a+1}^N e_{ki} e_{jk} \right) \mathbf{K}(u) &= \mathbf{K}(u) \left( (-su - p) e_{ji} + \sum_{k=1}^a e_{ki} e_{jk} \right) \\ &\quad \text{for } i \leq a < j, \end{aligned} \quad (5.7)$$

$$\begin{aligned} \left( (-su - p) e_{ji} + \sum_{k=1}^a e_{ki} e_{jk} \right) \mathbf{K}(u) &= \mathbf{K}(u) \left( (su - p) e_{ji} - \sum_{k=a+1}^N e_{ki} e_{jk} \right) \\ &\quad \text{for } j \leq a < i, \end{aligned} \quad (5.8)$$

where  $e_{ij}$  are generators of  $gl(N)$ :

$$[e_{ij}, e_{kl}] = \delta_{jk} e_{il} - \delta_{li} e_{kj}. \quad (5.9)$$

We find that (5.4) and (5.5) solve the relations (5.6)-(5.8), and thus the reflection equation

$$L_{12}(v-u)K_1(u)\bar{L}_{12}(u+v)K_2(v) = K_2(v)L_{12}(-u-v)K_1(u)\bar{L}_{12}(u-v) \quad \text{for } u, v \in \mathbb{C}, \quad (5.10)$$

which is the rational limit of (3.3), if the following conditions<sup>14</sup> are satisfied.

$$e_{ji} \sum_{k=a+1}^N e_{kk} = \sum_{k=a+1}^N e_{ki}e_{jk}, \quad e_{ji} \left( \sum_{k=1}^a e_{kk} - 1 \right) = \sum_{k=1}^a e_{ki}e_{jk} \quad \text{for } i \leq a < j, \quad (5.11)$$

$$e_{ji} \sum_{k=1}^a e_{kk} = \sum_{k=1}^a e_{ki}e_{jk}, \quad e_{ji} \left( \sum_{k=a+1}^N e_{kk} - 1 \right) = \sum_{k=a+1}^N e_{ki}e_{jk} \quad \text{for } j \leq a < i. \quad (5.12)$$

The relations (5.11) and (5.12) correspond to the rational limit of (3.25) and (3.26), respectively. Then the rational limit of (2.13) satisfies (5.11) and (5.12). Note that the above relations reproduce (a part of) a kind of generalized rectangular condition<sup>15</sup> [21].

$$\sum_{k=1}^N e_{ki}e_{jk} = \alpha e_{ji} + \beta \delta_{ij} \quad \alpha, \beta \in \mathbb{C}. \quad (5.13)$$

In fact, sum of the first and the second relations in (5.11) or (5.12) gives

$$e_{ji} \left( \sum_{k=1}^N e_{kk} - 1 \right) = \sum_{k=1}^N e_{ki}e_{jk} \quad \text{for } i \leq a < j \quad \text{or} \quad j \leq a < i. \quad (5.14)$$

This corresponds to (5.13) for  $\beta = 0$  since  $\alpha := \sum_{k=1}^N e_{kk} - 1$  is a central element of  $gl(N)$ .

## 6 Concluding remarks

We have derived intertwining relations (3.13), (3.14), (3.17) and (3.18) from the reflection equation (3.3) for L-operators associated with  $U_q(\hat{gl}(N))$  and have obtained the diagonal K-operators (3.19)-(3.22) in terms of the Cartan elements of a quotient of  $U_q(\hat{gl}(N))$  by the relations (3.25) and (3.26). The intertwining relations can be expressed in terms of generators of  $U_q(\hat{gl}(N))$  as in (4.1). In particular, the elements  $\{Z_{ji}\}$  (4.2)-(4.8) in the intertwining relations satisfy augmented q-Onsager algebra like

<sup>14</sup>These are sufficient conditions. If the parameter  $p$  (in addition to  $u$ ) is interpreted as a free one, these are necessary conditions as well. On the other hand, (3.25) and (3.26) are necessary and sufficient conditions independent of  $\epsilon_{\pm}$ .

<sup>15</sup>In case the representations are finite dimensional, this condition is satisfied by the representations labeled by rectangular Young diagrams.



commutation relations for the q-oscillator representation (2.13) under the evaluation map (2.10). It will be an interesting problem to investigate the algebra generated by  $\{Z_{ji}\}$  (with some constraint on generators of  $U_q(\hat{gl}(N))$ , if any), which could be a version of higher rank generalization of the augmented q-Onsager algebra. The reflection equation (3.1) defines [2] the reflection equation algebra if one adds the third component in the K-matrix:

$$R_{12} \left( \frac{y}{x} \right) K_{13}(x) \bar{R}_{12}(xy) K_{23}(y) = K_{23}(y) R_{12} \left( \frac{1}{xy} \right) K_{13}(x) \bar{R}_{12} \left( \frac{x}{y} \right), \quad (6.1)$$

$$K_{13}(x) = \sum_{i,j=1}^N E_{ij} \otimes 1 \otimes k_{ij}(x), \quad K_{23}(x) = \sum_{i,j=1}^N 1 \otimes E_{ij} \otimes k_{ij}(x),$$

where the elements  $\{k_{ij}(x)\}$  generate the algebra. In this context, it will be desirable to clarify the direct connection between  $\{Z_{ji}\}$  and a certain specialization of  $\{k_{ij}(x)\}$  for the diagonal K-operators.

In this paper, we have defined the L-operator  $\bar{\mathbf{L}}(x)$  based on the second evaluation map (2.11). It will be worthwhile to define it based on the first one (2.10) as  $\bar{\mathbf{L}}'(xy^{-1}) = \bar{\phi}'(xy^{-1})(\mathbf{ev}_x \otimes \pi_y) \mathcal{R}_{21}$  and investigate solutions of the reflection equation. The reason why we avoided  $\bar{\mathbf{L}}'(x)$  was that its expression based on the generators  $e_{ij}$  or  $\bar{e}_{ij}$  might be involved for  $N \geq 3$  case. The conditions (3.25) and (3.26) for the solutions of the reflection equation are the ones that the L-operator  $\bar{\mathbf{L}}(x)$  coincides with  $\bar{\mathbf{L}}'(x)$  up to a fine tune by the central element (see the relation (3.27)). The technical difficulty on  $\bar{\mathbf{L}}'(x)$  may be resolved if we start from the FRT formulation of the algebra (based Yang-Baxter relations) [22] and use the matrix elements of the L-operators as the generators of the algebra instead of  $e_{ij}$  or  $\bar{e}_{ij}$ .

Another important problem is construction of Baxter Q-operators for open boundary conditions in the light of [23, 5]. The K-operators obtained in this paper could be building blocks of Q-operators after taking limits on the parameter  $m$  of the q-oscillator representation, as was already discussed in [5] for  $N = 2$  case.

## Acknowledgments

The author would like to thank Pascal Baseilhac, Sergey Khoroshkin and Masato Okado for correspondence.

## Appendix A: Relations for $U_q(gl(N))$

Here we review relations among the generators of  $U_q(gl(N))$ , some of which are used in the main text.

$$\begin{aligned}
e_{ab} &= [e_{ac}, e_{cb}]_q \quad \text{for } a > c > b, \\
e_{ab} &= [e_{ac}, e_{cb}]_{q^{-1}} \quad \text{for } a < c < b, \\
[e_{ab}, e_{ba}] &= \frac{q^{e_{aa}-e_{bb}} - q^{-e_{aa}+e_{bb}}}{q - q^{-1}} \quad \text{for } a < b, \\
[e_{dc}, e_{ba}] &= -(q - q^{-1})e_{da}e_{bc} \quad \text{for } b < d < a < c \\
&\quad \text{or } a < c < b < d, \\
[e_{dc}, e_{ba}] &= 0 \quad \text{for } d < c < b < a \text{ or } d > c > b > a \text{ or } d < b < a < c \text{ or} \\
&\quad d > b > a > c \text{ or } d < c \leq a < b \text{ or } c < d \leq b < a \text{ or } d < a < b < c \text{ or} \\
&\quad c < b < a < d, \\
[e_{dc}, e_{ba}] &= -(q - q^{-1})q^{e_{aa}-e_{cc}}e_{da}e_{bc} \\
&\quad \text{for } d < a < c < b, \\
[e_{dc}, e_{ba}] &= (q - q^{-1})e_{da}e_{bc}q^{e_{bb}-e_{dd}} \\
&\quad \text{for } a < d < b < c, \\
[e_{ba}, e_{ac}] &= e_{bc}q^{e_{bb}-e_{aa}} \quad \text{for } a < b < c, \\
[e_{ba}, e_{ac}] &= q^{e_{aa}-e_{cc}}e_{bc} \quad \text{for } a < c < b, \\
[e_{db}, e_{ba}] &= e_{da}q^{e_{bb}-e_{dd}} \quad \text{for } a < d < b, \\
[e_{db}, e_{ba}] &= q^{e_{aa}-e_{bb}}e_{da} \quad \text{for } d < a < b, \\
[e_{da}, e_{ba}]_{q^{-1}} &= 0 \quad \text{for } a < b < d \text{ or } b < d < a, \\
[e_{bc}, e_{ba}]_q &= 0 \quad \text{for } c < a < b \text{ or } b < c < a.
\end{aligned} \tag{A1}$$

The relations for  $\bar{e}_{ij}$  can be obtained from the above relations through  $q \rightarrow q^{-1}$ . In addition, one can prove the following relations based on the above relations and induction. We remark that (A3) for  $l = j - 1$  corresponds to eq. (10) in [24].

$$\begin{aligned}
e_{ji} - (q - q^{-1}) \sum_{k=j+1}^l e_{ki} \bar{e}_{jk} &= \bar{e}_{ji} + (q - q^{-1}) \sum_{k=l+1}^{i-1} e_{ki} \bar{e}_{jk} = \\
&= \begin{cases} \bar{e}_{ji} & \text{for } j \leq l = i - 1, \\ [\bar{e}_{j,l+1}, e_{l+1,i}]_{q^{-1}} & \text{for } j \leq l < i - 1, \\ e_{ji} & \text{for } j = l < i, \end{cases} \tag{A2}
\end{aligned}$$

$$\bar{e}_{ji} - (q - q^{-1}) \sum_{k=i+1}^l e_{ki} \bar{e}_{jk} = e_{ji} + (q - q^{-1}) \sum_{k=l+1}^{j-1} e_{ki} \bar{e}_{jk} =$$

$$= \begin{cases} e_{ji} & \text{for } i \leq l = j - 1, \\ [\bar{e}_{j,l+1}, e_{l+1,i}]_{q^{-1}} & \text{for } i \leq l < j - 1, \\ \bar{e}_{ji} & \text{for } i = l < j, \end{cases} \quad (\text{A3})$$

$$e_{ji} + (q - q^{-1}) \sum_{k=l}^{j-1} e_{ki} \bar{e}_{jk} q^{e_{kk} - e_{jj} - 1} = [\bar{e}_{jl}, e_{li}]_{q^{-2}} q^{e_{ll} - e_{jj}} \quad \text{for } l < j < i, \quad (\text{A4})$$

$$\bar{e}_{ji} + (q - q^{-1}) \sum_{k=l}^{i-1} e_{ki} \bar{e}_{jk} q^{e_{kk} - e_{ii}} = [\bar{e}_{jl}, e_{li}]_{q^{-2}} q^{e_{ll} - e_{ii} + 1} \quad \text{for } l < i < j, \quad (\text{A5})$$

$$\bar{e}_{ji} - (q - q^{-1}) \sum_{k=i+1}^l e_{ki} \bar{e}_{jk} q^{e_{ii} - e_{kk}} = [\bar{e}_{jl}, e_{li}]_{q^2} q^{e_{ii} - e_{ll} - 1} \quad \text{for } j < i < l, \quad (\text{A6})$$

$$e_{ji} - (q - q^{-1}) \sum_{k=j+1}^l e_{ki} \bar{e}_{jk} q^{e_{jj} - e_{kk} + 1} = [\bar{e}_{jl}, e_{li}]_{q^2} q^{e_{jj} - e_{ll}} \quad \text{for } i < j < l, \quad (\text{A7})$$

$$[\bar{e}_{jl}, e_{li}]_{q^2} = [\bar{e}_{j,l+1}, e_{l+1,i}] q^{e_{ll} - e_{l+1,l+1}} \quad \text{for } i < j < l \quad \text{or } j < i < l, \quad (\text{A8})$$

$$[\bar{e}_{jl}, e_{li}]_{q^{-2}} = [\bar{e}_{j,l-1}, e_{l-1,i}] q^{e_{l-1,l-1} - e_{ll}} \quad \text{for } l < i < j \quad \text{or } l < j < i, \quad (\text{A9})$$

$$[\bar{e}_{j,l-1}, e_{l-1,i}]_q = [\bar{e}_{jl}, e_{li}]_{q^{-1}} \quad \text{for } i < l - 1 < l < j \quad \text{or } j < l - 1 < l < i. \quad (\text{A10})$$

We also remark that the third relation in the right hand side of (A2) (and (A3)) is a special case of the second one.

## References

- [1] I. V. Cherednik, Factorizing particles on a half-line and root systems, Theor. Math. Phys. 61 (1984) 977-983.
- [2] E.K. Sklyanin, Boundary conditions for integrable quantum systems, J. Phys. A 21 (1988) 2375-2389.
- [3] H. J. de Vega, A. González Ruiz, Boundary K-Matrices for the Six Vertex and the  $n(2n - 1) A_{n-1}$  Vertex Models, J.Phys. A26 (1993) L519-L524 [arXiv:hep-th/9211114].
- [4] I. V. Cherednik, On a method of constructing factorized  $S$  matrices in elementary functions, Theor. Math. Phys. 43 (1980) 356-358.
- [5] P. Baseilhac, Z. Tsuboi, Asymptotic representations of augmented q-Onsager algebra and boundary K-operators related to Baxter Q-operators, Nucl. Phys. B 929 (2018) 397-437 [arXiv:1707.04574].

- [6] T. Ito, P. Terwilliger, The augmented tridiagonal algebra, *Kyushu J. Math.* 64 (2010) 81-144 [arXiv:0904.2889 [math.QA]].
- [7] P. Baseilhac and S. Belliard, The half-infinite XXZ chain in Onsager's approach, *Nucl. Phys. B* 873 (2013) 550-583 [arXiv:1211.6304 [math-ph]].
- [8] A. Molev, E. Ragoucy, P. Sorba, Coideal subalgebras in quantum affine algebras, *Rev. Math. Phys* 15 (2003) 789-822 [arXiv:math/0208140].
- [9] A. Doikou, R. I. Nepomechie, Duality and quantum-algebra symmetry of the  $A_{N-1}^{(1)}$  open spin chain with diagonal boundary fields, *Nucl. Phys. B* 530 (1998) 641-664 [arXiv:hep-th/9807065].
- [10] G. W. Delius, N. J. MacKay, Quantum group symmetry in sine-Gordon and affine Toda field theories on the half-line, *Commun.Math.Phys.* 233 (2003) 173-190 [arXiv:hep-th/0112023].
- [11] H. Boos, F. Göhmann, A. Klümper, K.S. Nirov and A.V. Razumov, Exercises with the universal R-matrix, *J. Phys. A: Math. Theor.* 43 (2010) 415208 [arXiv:1004.5342 [math-ph]].
- [12] V. Drinfeld, Hopf algebras and the quantum Yang-Baxter equations, *Sov. Math. Dokl.* 32 (1985) 254-258.
- [13] M. Jimbo, A q-difference analogue of  $U(g)$  and the Yang-Baxter equation, *Lett. Math. Phys.* 10 (1985) 63-69.
- [14] M. Jimbo, A q-analogue of  $U(\mathfrak{gl}(N+1))$ , Hecke algebra, and the Yang-Baxter equation, *Lett. Math. Phys.* 11 (1986) 247-252.
- [15] V. Chari, A. Pressley, A guide to quantum groups, Cambridge University Press (1994).
- [16] A. Klimyk, K. Schmüdgen, Quantum groups and their representations, Springer-Verlag Berlin Heidelberg (1997).
- [17] S. M. Khoroshkin, V. N. Tolstoy, The uniqueness theorem for the universal R-matrix, *Lett. Math. Phys.* 24 (1992) 231-244.
- [18] T. D. Palev, A Holstein-Primakoff and a Dyson realization for the quantum algebra  $U_q[\mathfrak{sl}(n+1)]$ , *J. Phys. A* 31 (1998) 5145-5148 [arXiv:math/9804017 [math.QA]].
- [19] T. Hayashi, Q-analogues of Clifford and Weyl algebras-spinor and oscillator representations of quantum enveloping algebras, *Commun. Math. Phys.* 127 (1990) 129-144.
- [20] S. Khoroshkin, S. Pakuliak, A computation of an universal weight function for the quantum affine algebra  $U_q(\hat{\mathfrak{gl}}_N)$ , *J. Math. Kyoto Univ.* 48 (2008), 277-321 [arXiv:0711.2819 [math.QA]].

- [21] R. Frassek, T. Lukowski, C. Meneghelli, M. Staudacher, Baxter operators and Hamiltonians for gnearly allh integrable closed  $gl(n)$  spin chains, Nucl. Phys. B 874 (2013) 620-646 [arXiv:1112.3600 [math-ph]].
- [22] L. D. Faddeev, N. Y. Reshetikhin and L. A. Takhtajan, “Quantization of Lie Groups and Lie Algebras,” Leningrad Math. J. **1**, 193-225 (1990) [Alg. Anal. **1**, 178-206 (1989)].
- [23] R. Frassek, I. M. Szecsenyi, Q-operators for the open Heisenberg spin chain, Nucl. Phys. B 901 (2015) 229-248 [arXiv:1509.04867 [math-ph]].
- [24] R. B. Zhang, Universal  $L$  operator and invariants of the quantum supergroup  $U_q(gl(m/n))$ , J. Math. Phys. 33 (1992) 1970-1979.