

GENERAL DECAY RATE OF SOLUTION FOR NONLINEAR LOVE-EQUATION WITH INFINITE MEMORY

KHALED ZENNIR

ABSTRACT. In this paper, we consider a nonlinear Love-equation with infinite memory. By certain properties of convex functions, we use an appropriate Lyapunov functional to find a very general rate of decay for energy (2.3).

1. INTRODUCTION

Denote $y = y(x, t)$, $y' = y_t = \frac{\partial y}{\partial t}(x, t)$, $y'' = y_{tt} = \frac{\partial^2 y}{\partial t^2}(x, t)$, $y_x = \frac{\partial y}{\partial x}(x, t)$, $y_{xx} = \frac{\partial^2 y}{\partial x^2}(x, t)$, $x \in \Omega = (0, L)$, $L > 0$, $t > 0$. In this article, we consider a nonlinear Love-equation in the form

$$\begin{aligned} y'' - (y_x + y'_x + y''_x)_x + \int_{-\infty}^t \mu(t-s)y_{xx}(s)ds \\ = F[y] - (F[y])_x + f(x, t), \quad x \in \Omega, \quad 0 < t < T, \end{aligned} \quad (1.1)$$

where

$$F[y] = F(x, t, y, y_x, y', y'_x) \in C^1([0, 1] \times \mathbb{R}^+ \times \mathbb{R}^4), \quad (1.2)$$

The given functions μ, f are specified later. With $F = F(x, t, y_1, \dots, y_4)$, we put $D_1 F = \frac{\partial F}{\partial x}$, $D_2 F = \frac{\partial F}{\partial t}$, $D_{i+2} F = \frac{\partial F}{\partial y_i}$, with $i = 1, \dots, 4$.

Equation (1.1) satisfies the homogeneous Dirichlet boundary conditions:

$$y(0, t) = y(L, t) = 0, \quad t > 0, \quad (1.3)$$

and the following initial conditions

$$y(x, -t) = y_0(x, t), \quad y'(x, 0) = y_1(x). \quad (1.4)$$

To deal with a wave equation with infinite history, we assume that the kernel function μ satisfies the following hypothesis:

(Hyp1:) $\mu : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a non-increasing C^1 function such that

$$1 - \int_0^\infty \mu(s)ds = l > 0, \quad \mu(0) > 0. \quad (1.5)$$

and there exists an increasing strictly convex function $H : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ of $C^1(\mathbb{R}^+)$ satisfying

$$H(0) = H'(0) = 0, \quad \text{and} \quad \lim_{t \rightarrow \infty} H'(t) = \infty. \quad (1.6)$$

2010 *Mathematics Subject Classification.* 35L20, 35L70, 37B25, 93D15.

Key words and phrases. Nonlinear Love-equation; Existence; General Decay rate; Infinite Memory.

such that

$$\int_0^\infty \frac{\mu(s)}{H^{-1}(-\mu'(s))} ds + \sup_{s \in \mathbb{R}^+} \frac{\mu(s)}{H^{-1}(-\mu'(s))} ds < \infty \quad (1.7)$$

(Hyp2:) $y_0(0), y_1 \in H_0^1 \cap H^2$;

We need the following assumptions on source forces:

(Hyp3:) $f \in H^1((0, 1) \times (0, T))$;

(Hyp4:) $F \in C^1([0, 1] \times [0, T] \times \mathbb{R}^4)$, such that $F(0, t, 0, y_2, 0, y_4) = F(1, t, 0, y_2, 0, y_4) = 0$ for all $t \in [0, T]$, $y_2, y_4 \in \mathbb{R}$.

This kind of systems appears in the models of nonlinear Love waves or Love type waves. It is a generalization of a model introduced by [6], [22] and [23]. The original equation is

$$u'' - \frac{E}{\rho} u_{xx} - 2\mu^2 w^2 u''_{xx} = 0, \quad (1.8)$$

This type of problem describes the vertical oscillations of a rod, was established from Euler's variational equation of an energy functional associated with (1.8). A classical solution of problem (1.8) with null boundary conditions and asymptotic behavior are obtained by using the Fourier method and method of small parameter. The results are very interesting in the application point of view and, ad for as I know, new that is there is no results for equations of Love waves or Love type waves with the presence of finite/infinite memory term ([4], [6], [18], [19], [20], [21], [22], [23], [26], [27], ...).

Without infinite memory term, when $\lambda = 0$ in (1.1), Triet and his collaborator in [21] considered an IBVP for a nonlinear Kirchhoff-Love equation

$$u_{tt} - \frac{\partial}{\partial x} (u_x + \lambda_1 u_{xt} + u_{xtt}) + \lambda u_t = F(x, t, u, u_x, u_t, u_{xt}) - \frac{\partial}{\partial x} [G(x, t, u, u_x, u_t, u_{xt})] + f(x, t), \quad x \in \Omega = (0, 1), 0 < t < T, \quad (1.9)$$

$$u(0, t) = u(1, t) = 0, \quad (1.10)$$

$$u(x, 0) = \tilde{u}_0(x), \quad u_t(x, 0) = \tilde{u}_1(x), \quad (1.11)$$

where $\lambda > 0$, $\lambda_1 > 0$ are constants and $\tilde{u}_0, \tilde{u}_1 \in H_0^1 \cap H^2$; f , F and G are given functions. First, under suitable conditions, the existence of a unique local weak solution has been proved and a blow up result for solutions with negative initial energy is also established. A sufficient condition guaranteeing the global existence and exponential decay of weak solutions is given in the last section. This results will be improved in [26], [27] to the Kirchhoff typ.

The existence/ nonexistence, exponential decay of solutions and blow up results for viscoelastic wave equations with finite history, have been extensively studied and many results have been obtained by many authors (see [2], [5], [8], [9], [12], [14], [15], [31], [33] ...).

Concerning problems with infinite history, we mention the work [8] in which considered the following semi-linear hyperbolic equation, in a bounded domain of \mathbb{R}^3 ,

$$u'' - K(0)\Delta u - \int_0^\infty K'(s)\Delta u(t-s)ds + g(u) = f,$$

with $K(0), K(\infty) > 0, K' \leq 0$ and gave the existence of global attractors for the problem. Next in [28], the authors considered a fourth-order suspension bridge equation with nonlinear damping term and source term. The authors found necessary and sufficient condition for global existence and energy decay results without considering the relation between m and p . Moreover, when $p > m$, they gave sufficient condition for finite time blow-up of solutions. The lower bound of the blow-up time is also established.

Recently, in [24], the authors studied a three-dimensional (3D) visco-elastic wave equation with nonlinear weak damping, supercritical sources and prescribed past history, $t \leq 0$ in

$$u'' - k(0)\Delta u - \int_0^\infty k'(s)\Delta u(t-s)ds + |u'|^{m-1}u' = |u|^{p-1}u,$$

where the relaxation function k is monotone decreasing with $k(+\infty) = 1, m \geq 1, 1 \leq p < 6$. When the source is stronger than dissipations, i.e. $p > \max\{m, \sqrt{k(0)}\}$, they obtained some finite time blow-up results with positive initial energy. In particular, they obtained the existence of certain solutions which blow up in finite time for initial data at arbitrary energy level. (see [13], [16], [17], [24], [25], ...).

In addition to the existence results in Theorem 2.3 and Theorem 2.8 obtained by a new combined methods, our decay rate, which is very general, obtained in the fourth section, Theorem 3.1 extend that obtained in [31] and [13], where they established a general decay rate result for relaxation functions satisfying

$$g'(t) \leq -H(g(t)), t \geq 0, \quad H(0) = 0 \quad (1.12)$$

for a positive function $H \in C^1(\mathbb{R}^+)$ and H is linear or strictly increasing and strictly convex C^2 function on $(0, r], 1 > r$. This improves the conditions introduced by [2] on the relaxation functions

$$g'(t) \leq -\chi(g(t)), \quad \chi(0) = \chi'(0) = 0 \quad (1.13)$$

where χ is a non-negative function, strictly increasing and strictly convex on $(0, k_0], k_0 > 0$. They required that

$$\int_0^{k_0} \frac{dx}{\chi(x)} = +\infty, \quad \int_0^{k_0} \frac{xdx}{\chi(x)} < 1, \quad \liminf_{s \rightarrow 0^+} \frac{\chi(s)/s}{\chi'(s)} > \frac{1}{2} \quad (1.14)$$

and proved a decay result for the energy in a bounded domain. In addition to these assumptions, if

$$\limsup_{s \rightarrow 0^+} \frac{\chi(s)/s}{\chi'(s)} < 1, \quad (1.15)$$

then, in this case, an explicit rate of decay is given.

2. THE EXISTENCE OF SOLUTION

We define the weak solution to of (1.1)–(1.4) as follows.

Definition 2.1. *A function y is said to be a weak solution of (1.1)–(1.4) on $[0, T]$ if*

$$y, y', y'' \in L^\infty(0, T; H_0^1 \cap H^2),$$

such that y satisfies the variational equation

$$\begin{aligned} & \int_{\Omega} y'' w \, dx + \int_{\Omega} (y_x + y'_x + y''_x) w_x \, dx \\ & - \int_{\Omega} \int_0^{\infty} \mu(s) y_x(t-s) \, ds w_x \, dx \\ & = \int_{\Omega} f w \, dx + \int_{\Omega} F[y] w \, dx + \int_{\Omega} F'[y] w_x \, dx, \end{aligned} \quad (2.1)$$

for all test function $w \in H_0^1$, for almost all $t \in (0, T)$.

The following famous and widely used technical lemma will play an important role in the sequel.

Lemma 2.2. [32] For any $v \in C^1(0, T, H_0^1)$ we have

$$\begin{aligned} & \int_{\Omega} \int_0^{\infty} \mu(s) v_{xx}(t-s) v'(t) \, ds \, dx \\ & = \frac{1}{2} \frac{d}{dt} \int_{\Omega} \mu(s) \int_{\Omega} |v(t-s) - v(t)|^2 \, dx \, ds - \frac{1}{2} \frac{d}{dt} \int_{\Omega} \mu(s) \, ds \int_{\Omega} |v_x(t)|^2 \, dx \\ & \quad - \frac{1}{2} \int_{\Omega} \mu'(s) \int_{\Omega} |v(t-s) - v(t)|^2 \, dx \, ds. \end{aligned}$$

Now, we state the existence of a local solution for (1.1)–(1.4).

Theorem 2.3. [32], Theorem 2.3] Let $y_0(0), y_1 \in H_0^1 \cap H^2$ be given. Assume that (Hyp1)–(Hyp4) hold. Then Problem (1.1)–(1.4) has a unique local solution y and

$$y, y', y'' \in L^{\infty}(0, T_*; H_0^1 \cap H^2), \quad (2.2)$$

for some $T_* > 0$ small enough.

Here, we consider problem (1.1) with $p \geq 2$ and $f \equiv 0$ with the boundary conditions (1.3) and the initial conditions (1.4). We introduce the energy functional $E(t)$ associated with our problem

$$E(t) = \frac{1}{2} \int_{\Omega} |y'|^2 \, dx + \frac{1}{2} \int_{\Omega} |y'_x|^2 \, dx + J(t). \quad (2.3)$$

where

$$\begin{aligned} J(t) &= \frac{1}{2} \int_{\Omega} \left(1 - \int_0^{\infty} \mu(s) \, ds\right) |y_x|^2 \, dx \\ &+ \frac{1}{2} \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 \, ds \, dx \\ &+ \frac{1}{p} \int_{\Omega} |y_x|^p \, dx - \frac{1}{p} \int_{\Omega} |y|^p \, dx \end{aligned} \quad (2.4)$$

Now, we introduce the stable set as follows (see [29], [30])

$$W = \{y \in H_0^1 \cap H^2 : I(t) > 0, J(t) < d\} \cup \{0\} \quad (2.5)$$

where

$$\begin{aligned} I(t) &= \int_{\Omega} \left(1 - \int_0^{\infty} \mu(s) \, ds\right) |y_x|^2 \, dx \\ &+ \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 \, ds \, dx \end{aligned}$$

$$+ \int_{\Omega} |y_x|^p dx - \int_0^L |y|^p dx \quad (2.6)$$

We notice that the mountain pass level d given in (2.5) defined by

$$d = \inf \left\{ \sup_{y \in H_0^1 \cap H^2 \setminus \{0\}, \nu \geq 0} J(\nu y) \right\}, \quad (2.7)$$

Also, by introducing the so called "Nehari manifold"

$$\mathcal{N} = \{y \in H_0^1 \cap H^2 \setminus \{0\} : I(t) = 0\}$$

It is readily seen that the potential depth d is also characterized by

$$d = \inf_{y \in \mathcal{N}} J(t). \quad (2.8)$$

This characterization of d shows that

$$\text{dist}(0, \mathcal{N}) = \min_{y \in \mathcal{N}} \|y\|_{H_0^1 \cap H^2} \quad (2.9)$$

It is note hard to see this Lemma.

Lemma 2.4. *Suppose that (Hyp1) holds. Let y be solution of our equation. Then the energy functional (2.3) is a non-increasing function, i.e., for all $t \geq 0, \nu > 0$,*

$$\frac{d}{dt} E(t) \leq - \int_{\Omega} |y'_x|^2 dx + \frac{1}{2} \int_{\Omega} \int_0^{\infty} \mu'(s) |y_x(t) - y_x(t-s)|^2 ds dx \quad (2.10)$$

Proof. Multiplying (1.1) with $p \geq 2$ and $f \equiv 0$ by $y'(x, t)$ and integrating over Ω , using Lemma 2.2 we obtain th result. \square

As in [32], we will prove the invariance of the set W . That is if for some $t_0 > 0$ if $y(t_0) \in W$, then $y(t) \in W, \forall t \geq t_0$, beginning by the existence of the potential depth in the next Lemma.

Lemma 2.5 ([32], Lemma 3.2). *d is positive constant.*

Lemma 2.6 ([32], Lemma 3.3). *W is a bounded neighborhood of 0 in $H_0^1 \cap H^2$.*

Now, we will show that our local solution y is global in time, for this purpose it suffices to prove that the norm of the solution is bounded, independently of t , this is equivalent to prove the following theorem.

Theorem 2.7 ([32], Theorem 3.4). *Suppose that (Hyp1) and*

$$C^p l^{(1-p)} \left(\frac{2p}{p-2} E(0) \right)^{(p-2)} < l. \quad (2.11)$$

hold, where C is the best Poincare's constant. If $y_0(0) \in W, y_1 \in H_0^1$, then the solution $y \in W, \forall t \geq 0$.

The next Theorem shows that our local solution is global in time.

Theorem 2.8. *Suppose that (Hyp1), $p \geq 2$ and (2.11) hold. If $y_0(0) \in W, y_1 \in H_0^1$. Then the local solution y is global in time such that $y \in G_T$ where*

$$G_T = \left\{ y : \begin{array}{l} y \in L^\infty(\mathbb{R}^+; H_0^1 \cap H^2), \\ y' \in L^\infty(\mathbb{R}^+; H_0^1) \end{array} \right\}. \quad (2.12)$$

Proof. Now, it suffices to show that the following norm

$$\int_{\Omega} |y'|^2 dx + \int_{\Omega} |y_x|^2 dx, \quad (2.13)$$

is bounded independently of t .

To achieve this, we use (2.3), (2.4) and (2.10) to get

$$\begin{aligned} E(0) &\geq E(t) = J(t) + \frac{1}{2} \int_{\Omega} |y'|^2 dx \\ &\geq \left(\frac{p-2}{2p}\right) \left[\int_{\Omega} \left(1 - \int_0^{\infty} \mu(s) ds\right) |y_x|^2 dx + \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \right] \\ &\quad + \frac{1}{2} \int_{\Omega} |y'|^2 dx + \frac{1}{p} I(t) \\ &\geq \left(\frac{p-2}{2p}\right) \left[l \int_{\Omega} |y_x|^2 dx + \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \right] + \frac{1}{2} \int_{\Omega} |y'|^2 dx + \frac{1}{p} I(t) \\ &\geq \left(\frac{l(p-2)}{2p}\right) \int_{\Omega} |y_x|^2 dx + \frac{1}{2} \int_{\Omega} |y'|^2 dx \end{aligned}$$

since $I(t)$ and $\int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx$ are positive, hence

$$\left(\frac{l(p-2)}{2p}\right) \int_{\Omega} |y_x|^2 dx + \frac{1}{2} \int_{\Omega} |y'|^2 dx \leq CE(0),$$

where C is a positive constant depending only on p and l . This completes the proof. \square

3. GENERAL DECAY RATE

Theorem 3.1. *Suppose that (Hyp1), $p \geq 2$ and (2.11) hold. If $y_0(0) \in H_0^1 \cap H^2$, $y_1 \in H_0^1$. Then the energy function (2.3) satisfies*

$$E(t) \leq \kappa_1 H_1^{-1}(\kappa t + \kappa_0), \quad \forall t \geq 0, \quad (3.1)$$

where

$$H_1(\tau) = \int_{\tau}^1 (sH'(\kappa s))^{-1} ds, \quad \kappa_0, \kappa_1, \kappa > 0. \quad (3.2)$$

In order to prove the main Theorem 3.1, we need to introduce a several Lemmas. To this end, let us introduce the functionals

$$\varphi(t) = \int_{\Omega} yy' dx + \frac{1}{2} \int_{\Omega} |y_x|^2 dx + \int_{\Omega} y_x y'_x dx \quad (3.3)$$

and

$$\begin{aligned} \xi(t) &= - \int_{\Omega} y' \int_0^{\infty} \mu(s) [y(t) - y(t-s)] ds dx \\ &\quad - \int_0^t \int_{\Omega} y'_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds dx d\tau \\ &\quad - \int_0^t \int_{\Omega} y''_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds dx d\tau, \end{aligned} \quad (3.4)$$

Lemma 3.2. *Assume that (Hyp1), $p \geq 2$ and (2.11) hold. Then the functional $\varphi(t)$ introduced in (3.3) satisfies, along the solution, the estimate*

$$\begin{aligned} \varphi'(t) &\leq \int_{\Omega} y'^2 dx + \int_{\Omega} y_x'^2 dx \\ &\quad - \left[\frac{l}{2} - \left(\frac{2p}{(p-2)l} E(0) \right)^{(p-2)/2} \right] \int_{\Omega} |y_x|^2 dx + \frac{(1-l)}{2l} \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t-s) - y_x|^2 ds dx \end{aligned}$$

Proof.

$$\begin{aligned} \varphi'(t) &= \int_{\Omega} y'^2 dx + \int_{\Omega} y_x'^2 dx - \int_{\Omega} y_x^2 dx + \int_{\Omega} |y|^p dx - \int_{\Omega} |y_x|^p dx \\ &\quad + \int_{\Omega} y_x \int_0^{\infty} \mu(s) y_x(t-s) ds dx \end{aligned}$$

The next term can be treated as follows

$$\begin{aligned} \int_{\Omega} y_x \int_0^{\infty} \mu(s) y_x(t-s) ds dx &\leq \frac{1}{2} \int_{\Omega} y_x^2 dx + \frac{1}{2} \int_{\Omega} \left(\int_0^{\infty} \mu(s) y_x(t-s) ds \right)^2 dx \\ &\leq \frac{1}{2} \int_{\Omega} y_x^2 dx + \frac{1}{2} \int_{\Omega} \left(\int_0^{\infty} \mu(s) |y_x(t-s) - y_x| + |y_x| ds \right)^2 dx \end{aligned}$$

Thanks to Cauchy-Schwarz inequality, Young's inequality to obtain, for some $\nu > 0$,

$$\begin{aligned} &\int_{\Omega} \left(\int_0^{\infty} \mu(s) |y_x(t-s) - y_x| + |y_x| ds \right)^2 dx \\ &\leq \int_{\Omega} \left(\int_0^{\infty} \mu(s) |y_x(t-s) - y_x| ds \right)^2 dx + \int_{\Omega} \left(\int_0^{\infty} \mu(s) |y_x| ds \right)^2 dx \\ &\quad + 2 \int_{\Omega} \left(\int_0^{\infty} \mu(s) |y_x(t-s) - y_x| ds \right) \left(\int_0^{\infty} \mu(s) |y_x| ds \right) dx \\ &\leq \left(1 + \frac{1}{\nu} \right) \int_{\Omega} \left(\int_0^{\infty} \mu(s) |y_x(t-s) - y_x| ds \right)^2 dx \\ &\quad + (1 + \nu) \int_{\Omega} \left(\int_0^{\infty} \mu(s) ds |y_x| \right)^2 dx \\ &\leq \left(1 + \frac{1}{\nu} \right) (1-l) \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t-s) - y_x|^2 ds dx + (1 + \nu) (1-l)^2 \int_{\Omega} |y_x|^2 dx. \end{aligned}$$

Then

$$\begin{aligned} \varphi'(t) &\leq \int_{\Omega} y'^2 dx + \int_{\Omega} y_x'^2 dx \\ &\quad + \left[\frac{1}{2} + \frac{1}{2} (1 + \nu) (1-l)^2 - 1 \right] \int_{\Omega} |y_x|^2 dx \\ &\quad + \frac{1}{2} \left(1 + \frac{1}{\nu} \right) (1-l) \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t-s) - y_x|^2 ds dx \\ &\quad + \int_{\Omega} |y|^p dx. \end{aligned} \tag{3.5}$$

By the continuous embedding for $p \geq 2$, we have

$$\begin{aligned} \int_{\Omega} |y|^p dx &\leq c^p \left(\int_{\Omega} |y_x|^2 dx \right)^{p/2} \\ &\leq c^p \left(\int_{\Omega} |y_x|^2 dx \right)^{(p-2)/2} \int_{\Omega} |y_x|^2 dx \end{aligned}$$

$$\leq c^p \left(\frac{2p}{(p-2)l} E(0) \right)^{(p-2)/2} \int_{\Omega} |y_x|^2 dx \quad (3.6)$$

Since (2.11), we can choose $\nu = \frac{l}{1-l}$ to get

$$\begin{aligned} \varphi'(t) &\leq \int_{\Omega} y^2 dx + \int_{\Omega} y_x'^2 dx \\ &\quad - \left[\frac{l}{2} - c^p l^{1-p} \left(\frac{2p}{(p-2)} E(0) \right)^{(p-2)} \right] \int_{\Omega} |y_x|^2 dx \\ &\quad + \frac{(1-l)}{2l} \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t-s) - y_x|^2 ds dx \end{aligned}$$

□

Lemma 3.3. *Assume that (Hyp1), $p \geq 2$ hold. Then the functional introduced in (3.4) satisfies, along the solution, the estimate Then, for $\nu < (1-l)$*

$$\begin{aligned} \xi'(t) &\leq -a \int_{\Omega} |y_x|^2 dx - ((1-l) - \nu) \int_{\Omega} y'^2 dx \\ &\quad + b \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \\ &\quad + \frac{c\mu(0)}{4\nu} \int_{\Omega} \int_0^{\infty} (-\mu'(s)) |y(t) - y(t-s)|^2 ds dx. \end{aligned}$$

where

$$a = c(\nu) \left(1 + 2(1-l)^2 - \left(\frac{2p}{(p-2)l} E(0) \right)^{(p-2)/2} \right) > 0$$

and for all $\nu > 0$

$$b = \frac{1-l}{4\nu} + (2\nu + \frac{1}{4\nu})(1-l) + 2\nu(1-l)^{p-1} c \left(\frac{8}{1-l} E(0) + 2m_0^2 \right)^{(p-2)/2} > 0.$$

Proof. We have

$$\begin{aligned} \xi'(t) &= \int_{\Omega} y_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds dx \\ &\quad - \int_{\Omega} \int_0^{\infty} \mu(s) y_x(t-s) ds \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds dx \\ &\quad - \int_{\Omega} |y|^{p-2} y \int_0^{\infty} \mu(s) [u(t) - u(t-s)] ds dx \\ &\quad + \int_{\Omega} |y_x|^{p-2} y_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds dx \\ &\quad - \int_{\Omega} y' \int_0^{\infty} \mu'(s) [y(t) - y(t-s)] ds dx - (1-l) \int_{\Omega} y'^2 dx. \end{aligned}$$

For $\nu > 0$, we have

$$\begin{aligned} &\int_{\Omega} y_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds dx \\ &\leq \nu \int_{\Omega} |y_x|^2 dx + \frac{1-l}{4\nu} \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx, \end{aligned}$$

and

$$\begin{aligned} & \int_{\Omega} \int_0^{\infty} \mu(s) y_x(t-s) ds \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds dx \\ & \leq 2\nu(1-l)^2 \int_{\Omega} |y_x|^2 dx + (2\nu + \frac{1}{4\nu})(1-l) \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx, \end{aligned}$$

we have the estimate

$$\begin{aligned} & \int_{\Omega} |y(t) - y(t-s)|^2 dx \\ & \leq 2 \int_{\Omega} |y_x(t)|^2 dx + 2 \int_{\Omega} |y_x(t-s)|^2 dx \\ & \leq 4 \sup_{s>0} \int_{\Omega} |y_x(t)|^2 dx + 2 \sup_{\tau<0} \int_{\Omega} |y_x(\tau)|^2 dx \\ & \leq \sup_{s>0} \int_{\Omega} |y_x(t)|^2 dx + 2 \sup_{\tau<0} \int_{\Omega} |y_{0x}(\tau)|^2 dx \\ & \leq \frac{8}{1-l} E(0) + 2m_0^2, \end{aligned} \tag{3.7}$$

where $\int_{\Omega} |y_{0x}(\tau)|^2 dx \leq m_0, m_0 \geq 0, \quad \forall \tau > 0$. Since $p \geq 2$, we have by using (3.6) and the previous estimate

$$\begin{aligned} & \int_{\Omega} |y|^{p-2} y \int_0^{\infty} \mu(s) [y(t) - y(t-s)] ds dx \\ & \leq \nu \int_{\Omega} \left| \int_0^{\infty} \mu(s) |y(t) - y(t-s)| ds \right|^p dx + c(\nu) \int_{\Omega} |y|^p dx \\ & \leq \nu(1-l)^{p-1} \int_{\Omega} \int_0^{\infty} \mu(s) |y(t) - y(t-s)|^p ds dx + c(\nu) \int_{\Omega} |y|^p dx \\ & \leq \nu(1-l)^{p-1} c \int_0^{\infty} \mu(s) \left(\int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right)^{p/2} ds + c(\nu) \int_{\Omega} |y|^p dx \\ & \leq \nu(1-l)^{p-1} c \left(\frac{8}{1-l} E(0) + 2m_0^2 \right)^{(p-2)/2} \int_0^{\infty} \mu(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \\ & \quad + c(\nu) \int_{\Omega} |y|^p dx \\ & \leq \nu(1-l)^{p-1} c \left(\frac{8}{1-l} E(0) + 2m_0^2 \right)^{(p-2)/2} \int_0^{\infty} \mu(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \\ & \quad + c(\nu) \left(\frac{2p}{(p-2)l} E(0) \right)^{(p-2)/2} \int_{\Omega} |y_x|^2 dx. \end{aligned}$$

Similarly to estimate

$$\begin{aligned} & \int_{\Omega} |y_x|^{p-2} y_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds dx \\ & \leq \nu \int_{\Omega} \left| \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)| ds \right|^p dx + c(\nu) \int_{\Omega} |y_x|^p dx \\ & \leq \nu(1-l)^{p-1} \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^p ds dx + c(\nu) \int_{\Omega} |y_x|^p dx \\ & \leq \nu(1-l)^{p-1} c \int_0^{\infty} \mu(s) \left(\int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right)^{p/2} ds + c(\nu) \int_{\Omega} |y_x|^p dx \end{aligned}$$

$$\begin{aligned} &\leq \nu(1-l)^{p-1}c\left(\frac{8}{1-l}E(0) + 2m_0^2\right)^{(p-2/2)} \int_0^\infty \mu(s) \int_\Omega |y_x(t) - y_x(t-s)|^2 dx \\ &+ c(\nu) \int_\Omega |y_x|^2 dx. \end{aligned}$$

The next term estimated as

$$\begin{aligned} & - \int_\Omega y' \int_0^\infty \mu'(s)[y(t) - y(t-s)] ds dx \\ & \leq \nu \int_\Omega |y'|^2 dx + \frac{c\mu(0)}{4\nu} \int_\Omega \int_0^\infty (-\mu'(s))|y(t) - y(t-s)|^2 ds dx. \end{aligned}$$

A combination of all estimates gives

$$\begin{aligned} \xi'(t) &\leq \nu \int_\Omega |y_x|^2 dx + \frac{1-l}{4\nu} \int_\Omega \int_0^\infty \mu(s)|y_x(t) - y_x(t-s)|^2 ds dx \\ &+ 2\nu(1-l)^2 \int_\Omega |y_x|^2 dx + (2\nu + \frac{1}{4\nu})(1-l) \int_\Omega \int_0^\infty \mu(s)|y_x(t) - y_x(t-s)|^2 ds dx \\ &+ \nu(1-l)^{p-1}c\left(\frac{8}{1-l}E(0) + 2m_0^2\right)^{(p-2/2)} \int_0^\infty \mu(s) \int_\Omega |y_x(t) - y_x(t-s)|^2 dx \\ &+ c(\nu)\left(\frac{2p}{(p-2)l}E(0)\right)^{(p-2/2)} \int_\Omega |y_x|^2 dx \\ &+ \nu(1-l)^{p-1}c\left(\frac{8}{1-l}E(0) + 2m_0^2\right)^{(p-2/2)} \int_0^\infty \mu(s) \int_\Omega |y_x(t) - y_x(t-s)|^2 dx \\ &+ c(\nu) \int_\Omega |y_x|^2 dx \\ &+ \nu \int_\Omega |y'|^2 dx + \frac{c\mu(0)}{4\nu} \int_\Omega \int_0^\infty (-\mu'(s))|y(t) - y(t-s)|^2 ds dx - (1-l) \int_\Omega y'^2 dx. \end{aligned}$$

Then, for $\nu < (1-l)$

$$\begin{aligned} \xi'(t) &\leq -a \int_\Omega |y_x|^2 dx - ((1-l) - \nu) \int_\Omega y'^2 dx \\ &+ b \int_\Omega \int_0^\infty \mu(s)|y_x(t) - y_x(t-s)|^2 ds dx \\ &+ \frac{c\mu(0)}{4\nu} \int_\Omega \int_0^\infty (-\mu'(s))|y(t) - y(t-s)|^2 ds dx. \end{aligned}$$

where by (2.11), we have

$$a = c(\nu)\left(1 + 2(1-l)^2 - \left(\frac{2p}{(p-2)l}E(0)\right)^{(p-2/2)}\right) > 0$$

and for all $\nu > 0$, we have

$$b = \frac{1-l}{4\nu} + (2\nu + \frac{1}{4\nu})(1-l) + 2\nu(1-l)^{p-1}c\left(\frac{8}{1-l}E(0) + 2m_0^2\right)^{(p-2/2)} > 0.$$

□

Let define the functional

$$L(t) = \varepsilon_1 E(t) + \varphi(t) + \varepsilon_2 \xi(t), \quad \varepsilon_1, \varepsilon_2 > 0 \quad (3.8)$$

We need the next lemma, which means that there is equivalent between the Lyapunov and energy functions

Lemma 3.4. For $\varepsilon_1, \varepsilon_2 > 1$, we have

$$L \sim E, \quad (3.9)$$

Proof. By (3.8) we have

$$\begin{aligned} |L(t) - \xi_1 E(t)| &\leq |\varphi(t)| + \varepsilon_2 |\xi_2(t)| \\ &\leq \int_{\Omega} |yy'| dx + \frac{1}{2} \int_{\Omega} |y_x|^2 dx + \int_{\Omega} |y_x y'_x| dx \\ &\quad + \int_{\Omega} \left| y' \int_0^{\infty} \mu(s) [y(t) - y(t-s)] ds \right| dx \\ &\quad + \int_0^t \int_{\Omega} \left| y'_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds d\tau \right| dx \\ &\quad + \int_0^t \int_{\Omega} \left| y''_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds d\tau \right| dx, \end{aligned}$$

Thanks to Holder and Young's inequalities, we have

$$\begin{aligned} \int_{\Omega} |yy'| dx &\leq \left(\int_{\Omega} |y|^2 dx \right)^{1/2} \left(\int_{\Omega} |y'|^2 dx \right)^{1/2} \\ &\leq \frac{1}{2} \left(\int_{\Omega} |y|^2 dx \right) + \frac{1}{2} \left(\int_{\Omega} |y'|^2 dx \right) \\ &\leq \frac{c}{2} \left(\int_{\Omega} |y_x|^2 dx \right) + \frac{1}{2} \left(\int_{\Omega} |y'|^2 dx \right) \end{aligned}$$

Similarly

$$\int_{\Omega} |y_x y'_x| dx \leq \frac{1}{2} \left(\int_{\Omega} |y_x|^2 dx \right) + \frac{1}{2} \left(\int_{\Omega} |y'_x|^2 dx \right)$$

and

$$\begin{aligned} &\int_{\Omega} \left| y' \int_0^{\infty} \mu(s) [y(t) - y(t-s)] ds \right| dx \\ &\leq \frac{1}{2} \left(\int_{\Omega} |y'|^2 dx \right) + \frac{c}{2} \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \end{aligned}$$

and

$$\begin{aligned} &\int_0^t \int_{\Omega} \left| y'_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds d\tau \right| dx \\ &\leq \frac{1}{2} \int_0^t \left(\int_{\Omega} |y'_x|^2 dx \right) d\tau + \frac{1}{2} \int_0^t \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx d\tau \end{aligned}$$

and

$$\begin{aligned} &\int_0^t \int_{\Omega} \left| y''_x \int_0^{\infty} \mu(s) [y_x(t) - y_x(t-s)] ds d\tau \right| dx \\ &\leq \frac{1}{2} \int_0^t \left(\int_{\Omega} |y''_x|^2 dx \right) d\tau + \frac{1}{2} \int_0^t \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx d\tau \end{aligned}$$

Then, for some $c > 0$, we have

$$|L(t) - \varepsilon_1 E(t)| \leq c(E(t))$$

Therefore, we can choose ε_1 so that

$$L(t) \sim E(t). \quad (3.10)$$

□

Lemma 3.5. *Assume (Hyp1) hold. Then there exist strictly positive constants c such that*

$$L'(t) \leq -\varepsilon_1 E(t) + c \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \quad (3.11)$$

Proof. By Lemma 2.4, Lemma 3.2 and Lemma 3.3, we have

$$\begin{aligned} L'(t) &= \varepsilon_1 E'(t) + \varphi'(t) + \varepsilon_2 \xi'(t) \\ &\leq -[\varepsilon_2[(1-l) - \nu] - 1] \int_0^L |y'|^2 dx - \left(1 - \frac{\nu}{2}\right) \varepsilon_1 - 1 \int_{\Omega} |y'_x|^2 dx \\ &\quad - \left[\frac{l}{2} - \left(\frac{2p}{(p-2)l} E(0)\right)^{(p-2)/2} - \varepsilon_2 a\right] \int_{\Omega} |y_x|^2 dx \\ &\quad + \left[\frac{(1-l)}{2l} + b\varepsilon_2\right] \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t-s) - y_x|^2 ds dx \\ &\quad + \left[\frac{\varepsilon_1}{2} - \varepsilon_2 \frac{c\mu(0)}{4\nu}\right] \int_{\Omega} \int_0^{\infty} \mu'(s) |y_x(t) - y_x(t-s)|^2 ds dx, \end{aligned}$$

where, by (2.11), we have

$$a = c(\nu) \left(1 + 2(1-l)^2 - \left(\frac{2p}{(p-2)l} E(0)\right)^{(p-2)/2}\right) > 0$$

and for all $\nu > 0$

$$b = \frac{1-l}{4\nu} + (2\nu + \frac{1}{4\nu})(1-l) + 2\nu(1-l)^{p-1} c \left(\frac{8}{1-l} E(0) + 2m_0^2\right)^{(p-2)/2} > 0.$$

Now, we choose ν , and whence this constant is fixed. we can choice $\varepsilon_1, \varepsilon_2$ small enough such that for $p \geq 2$ there exist $c > 0$ and (3.11) yields. □

Lemma 3.6. *Assume that (Hyp1) hold. Then, there exist $\gamma, \gamma_0 > 0$ such that for all $t > 0$*

$$\int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \leq \frac{-\gamma E'(t)}{H'(\gamma_0 E(t))} + \gamma \gamma_0 E(t) \quad (3.12)$$

Proof. Let H^* be the convex conjugate of g in the sense of Young (see [1], pages 61-64), then

$$\begin{aligned} H^*(s) &= s(H')^{-1}(s) - H[(H')^{-1}(s)] \\ &\leq s(H')^{-1}(s), \quad s \in (0, H'(r)) \end{aligned} \quad (3.13)$$

and satisfies the following Young's inequality

$$AB \leq H^*(A) + H(B), \quad A \in (0, H'(r)), B \in (0, r]. \quad (3.14)$$

for

$$\begin{aligned} A &= H^{-1} \left(-r_2 \mu'(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right) \\ B &= \frac{r_1 H'(\gamma_0 E(t)) \mu(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx}{H^{-1} \left(-r_2 \mu'(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right)}. \end{aligned}$$

Then, for $r_1, r_2 > 0$, we have (see [17])

$$\begin{aligned}
& \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \\
&= \frac{1}{r_1 H'(\gamma_0 E(t))} \int_0^{\infty} \left\{ H^{-1} \left(-r_2 \mu'(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right) \right. \\
&\quad \times \left. \frac{r_1 H'(\gamma_0 E(t)) \mu(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx}{H^{-1} \left(-r_2 \mu'(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right)} \right\} ds \\
&\leq -\frac{r_2}{r_1 H'(\gamma_0 E(t))} \int_{\Omega} \int_0^{\infty} \mu'(s) |y_x(t) - y_x(t-s)|^2 ds dx \\
&\quad + \frac{1}{r_1 H'(\gamma_0 E(t))} \int_0^{\infty} H^* \left(\frac{r_1 H'(\gamma_0 E(t)) \mu(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx}{H^{-1} \left(-r_2 \mu'(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right)} \right) ds
\end{aligned}$$

By (2.10), (3.13), we have

$$\begin{aligned}
& \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \leq \frac{2r_2}{r_1 H'(\gamma_0 E(t))} E'(t) \\
&\quad + \int_0^{\infty} \left\{ \frac{\mu(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx}{H^{-1} \left(-r_2 \mu'(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right)} \right. \\
&\quad \times \left. H'^{-1} \left(\frac{r_1 H'(\gamma_0 E(t)) \mu(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx}{H^{-1} \left(-r_2 \mu'(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right)} \right) \right\} ds
\end{aligned}$$

By the fact that H^{-1} is concave and $H^{-1}(0) = 0$, the function $h(s) = \frac{s}{H^{-1}(s)}$, such that for $0 \leq s_1 < s_2$, we have

$$h(s_1) \leq h(s_2).$$

Therefore, using (3.7) to get

$$\begin{aligned}
& H'^{-1} \left(\frac{r_1 H'(\gamma_0 E(t)) \mu(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx}{H^{-1} \left(-r_2 \mu'(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right)} \right) \\
&= H'^{-1} \left[\frac{r_1 H'(\gamma_0 E(t)) \mu(s)}{-r_2 \mu'(s)} h \left(-r_2 \mu'(s) \int_{\Omega} |y_x(t) - y_x(t-s)|^2 dx \right) \right] \\
&\leq H'^{-1} \left[\frac{r_1 H'(\gamma_0 E(t)) \mu(s)}{-r_2 \mu'(s)} h \left(-r_2 \mu'(s) \left(\frac{8}{1-l} E(0) + 2m_0^2 \right) \right) \right] \\
&\leq H'^{-1} \left[\frac{r_1 \left(\frac{8}{1-l} E(0) + 2m_0^2 \right) H'(\gamma_0 E(t)) \mu(s)}{H^{-1} \left(-r_2 \mu'(s) \left(\frac{8}{1-l} E(0) + 2m_0^2 \right) \right)} \right].
\end{aligned}$$

Then,

$$\begin{aligned}
& \int_{\Omega} \int_0^{\infty} \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \\
&\leq -\frac{2r_2}{r_1 H'(\gamma_0 E(t))} E'(t) \\
&\quad + \left(\frac{8}{1-l} E(0) + 2m_0^2 \right) \int_0^{\infty} \left\{ \frac{\mu(s)}{H^{-1} \left(-r_2 \left(\frac{8}{1-l} E(0) + 2m_0^2 \right) \mu'(s) \right)} \right.
\end{aligned}$$

$$\times H'^{-1} \left[\frac{r_1 \left(\frac{8}{1-l} E(0) + 2m_0^2 \right) H'(\gamma_0 E(t)) \mu(s)}{H^{-1} \left(-r_2 \mu'(s) \left(\frac{8}{1-l} E(0) + 2m_0^2 \right) \right)} \right] ds$$

By (Hyp1), we have

$$\sup_{s \in \mathbb{R}^+} \frac{\mu(s)}{H^{-1} \left(-\mu'(s) \right)} = \kappa_1 < \infty$$

$$\int_0^\infty \frac{\mu(s)}{H^{-1} \left(-\mu'(s) \right)} = \kappa_2 < \infty.$$

Since H'^{-1} is nondecreasing, we choose r_1, r_2 such that

$$\begin{aligned} & \int_{\Omega} \int_0^\infty \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \\ & \leq -\frac{2\kappa_2}{H'(\gamma_0 E(t))} E'(t) + \left(\frac{8}{1-l} E(0) + 2m_0^2 \right) H'^{-1} H'(\gamma_0 E(t)) \int_0^\infty \frac{\mu(s)}{H^{-1} \left(-\mu'(s) \right)} \\ & \leq -\frac{2\kappa_2}{H'(\gamma_0 E(t))} E'(t) + \left(\frac{8}{1-l} E(0) + 2m_0^2 \right) \gamma_0 E(t). \end{aligned}$$

This completes the proof. \square

Proof. (of Theorem 3.1) Multiplying (3.11) by $H'(\gamma_0 E(t))$ and using results in (3.12)

$$\begin{aligned} H'(\gamma_0 E(t)) L'(t) & \leq -\varepsilon_1 H'(\gamma_0 E(t)) E(t) + c H'(\gamma_0 E(t)) \int_{\Omega} \int_0^\infty \mu(s) |y_x(t) - y_x(t-s)|^2 ds dx \\ & \leq -[\varepsilon_1 - c\gamma\gamma_0] H'(\gamma_0 E(t)) E(t) - c\gamma E'(t). \end{aligned} \tag{3.15}$$

We choose γ_0 small enough so that $\varepsilon_1 - c\gamma\gamma_0 > 0$.

Put

$$g(t) = H'(\gamma_0 E(t)) L(t) + c\gamma E(t) \sim E(t),$$

then,

$$g'(t) \leq -\kappa g(t) H'(\gamma_0 g(t)),$$

which implies that $(H_1(g))' \geq \kappa$, where

$$H_1(\tau) = \int_\tau^1 \frac{1}{s H'(\gamma_0 s)} ds, \quad 0 < \tau < 1.$$

Integrating $(H_1(g))' \geq \kappa$ over $[0, t]$ we get

$$g(t) \leq H_1^{-1}(\kappa t + \kappa_0)$$

the equivalence between $E(t)$ and $g(t)$ gives the result. \square

REFERENCES

- [1] Arnold, V. I., *Mathematical Methods of Classical Mechanics*, Springer-Verlag, New York, 1989.
- [2] F. Alabau-Boussouira and P. Cannarsa, *A general method for proving sharp energy decay rates for memory-dissipative evolution equations*, C. R. Math. Acad. Sci. Paris, Ser. I **347**, (2009), 867-872.
- [3] E. L. A. Coddington, N. Levinson; *Theory of ordinary differential equations*, McGraw-Hill, 1955, p. **43**.
- [4] A. Chattopadhyay, S.Gupta, A. K. Singh, S. A. Sahu; *Propagation of shear waves in an irregular magneto-elastic monoclinic layer sandwiched between two isotropic half-spaces* , International Journal of Engineering, Science and Technology, **1** (2009), 228-244.
- [5] C. M. Dafermos, *On the existence and the asymptotic stability of solution to the equations of linear thermo-elasticity*, Arch. Ration. Mech. Anal., **29**, (1968) 241-271.
- [6] S. Dutta, *On the propagation of Love type waves in an infinite cylinder with rigidity and density varying linearly with the radial distance*, Pure Appl. Geophys., **98**, (1972) 35-39.
- [7] Erhan, Piskin and Necat, Polat, *Global existence, exponential and polynomial decay solutions for a system class of nonlinear higher-order wave equations with damping and source terms*, Inter. j. Pure and Appl. Math., 76(2012), no. 4, 559-570.
- [8] C. Giorgi, J. Muenoz Rivera and V. Pata, *Global Attractors for a Semilinear Hyperbolic Equation in visco-elasticity*, J. Math. Anal. Appl., **260**(1), (2001) 83-99.
- [9] X. Han and M. Wang, *Global existence and blow-up of solutions for a system of nonlinear visco-elastic wave equations with damping and source*, Nonli. Anal., **71**(2009), 5427-5490.
- [10] W. Hrusa, *Global existence and asymptotic stability for a semi-linear Volterra equation with large initial data*, SIAMJ. Math. Anal. **16**(1)(1985), 110-134.
- [11] H. A. Levine, S. R. Park, and J. Serrin, *Global existence and global nonexistence of solutions of the Cauchy problem for a nonlinearly damped wave equation*, J. Math. Anal. Appl., 228(1998), no. 1, 181-205.
- [12] M. Mustafa, *Well posedness and asymptotic behavior of a coupled system of nonlinear visco-elastic equations*, Nonlin. Anal. (**13**), (2012), 452-463.
- [13] Muhammad I. Mustafa and S. A. Messaoudi, *General stability result for viscoelastic wave equations*, J. Math. Phys. **53**, 053702 (2012).
- [14] S. A. Messaoudi; *Blow up and global existence in nonlinear visco-elastic wave equations*, Math. Nachrich. **260** (2003), 58-66.
- [15] S. A. Messaoudi; *General decay of solutions of a viscoelastic equation*, J. Math. Anal. Appl. 341(2) (2008), 1457-1467.
- [16] S. A. Messaoudi, Muhammad M. Al-Gharabli; *A general stability result for a nonlinear wave equation with infinite memory*, Applied Mathematics Letters, **26** (2013) 1082-1086
- [17] M. M. Al-Gharabli; *A general decay result of a viscoelastic equation with infinite history and nonlinear damping*, Applicable Analysis, **97** (2018) 382-399
- [18] L. T. P. Ngoc, N. T. Duy, N. T. Long; *A linear recursive scheme associated with the Love's equation*, Acta Mathematica Vietnamica, **38** (4) (2013), 551-562.
- [19] L. T. P. Ngoc, N. T. Duy, N. T. Long; *Existence and properties of solutions of a boundary problem for a Love's equation*, Bull. Malays. Math. Sci. Soc., **2** (37)(2014), 997-1016.
- [20] L. T. P. Ngoc, N. T. Duy, N. T. Long; *On a high-order iterative scheme for a nonlinear Love equation*, Applications of Mathematics, **60** (3)(2015), 285-298.
- [21] L. T. P. Ngoc, N. T. Long; *Existence, blow-up and exponential decay for a nonlinear Love equation associated with Dirichlet conditions*, Applications of Mathematics, **61**(2) (2016), 165-196.
- [22] M. K. Paul; *On propagation of Love-type waves on a spherical model with rigidity and density both varying exponentially with the radial distance*, Pure Appl. Geophys, **59** (1964), 33-37.
- [23] V. Radochova; *Remark to the comparison of solution properties of Loves equation with those of wave equation*, Apl. Mat., **23** (1978), 199-207.
- [24] Fenglong Sun, Lishan Liu and Yonghong Wu; *Blow-up of solutions for a nonlinear visco-elastic wave equation with initial data at arbitrary energy level*, Applicable Analysis, DOI: 10.1080/00036811.2018.1460812
- [25] Sun-Hye PARK; *Stability of a von karman equation with infinite memory*, Acta Mathematica Scientia, 2017, 37B(4):965-973

- [26] N. A. Triet, V. T. T. Mai, L. T. P. Ngoc, N. T. Long, *A Dirichlet problem for a nonlinear wave equation of Kirchhoff - Love type*, Nonlinear Functional Analysis and Applications, **22** (3) (2017), 595-626.
- [27] N. A. Triet, V. T. T. Mai, L. T. P. Ngoc, N. T. Long, *Existence, blow-up and exponential decay for Kirchhoff-love equations with Dirichlet conditions*, Electronic Journal of Differential Equations, Vol. 2018 (2018), No. **167**, pp. 1-26.
- [28] Wenjun Liu and H. Zhuang; *Global existence, asymptotic behavior and blow-up of solutions for a suspension bridge equation with nonlinear damping and source terms*, Nonlinear Differ. Equ. Appl., (2017) **24**: 67. <https://doi.org/10.1007/s00030-017-0491-5>
- [29] Yaojun Ye, *Global existence and asymptotic stability for coupled nonlinear Klein-Gordon equations with nonlinear damping terms*, Dyn. Syst., **28**(2013), no. 2, 287-298.
- [30] Kh. zennir and Amar Guesmia, *Existence of solutions to nonlinear κ -th-order coupled Klein-Gordon equations with nonlinear sources and memory terms*. Applied Mathematics E-Notes, **15**(2015), 121-136.
- [31] Kh. zennir, *General decay of solutions for damped wave equation of Kirchhoff type with density in \mathbb{R}^n* . Ann. Univ. Ferrara **61**(2015) 381-394.
- [32] Kh. zennir and Toufic El Arwadi, *Results on local and Global existence of solution for Love-equation with infinite memory*. Preprint
- [33] S. Zitouni, and Kh. Zennir, *On the existence and decay of solution for viscoelastic wave equation with nonlinear source in weighted spaces*. Rend. Circ. Mat. Palermo, II. Ser, Vol. **66**, No 3 (2017), pp 337-353.

KHALED ZENNIR

FIRST ADDRESS: LABORATORY LAMAHIS, UNIVERSITY 20 AOÛT 1955- SKIKDA, 21000, ALGERIA

SECOND ADDRESS: DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCES AND ARTS, AL-RAS,

QASSIM UNIVERSITY, KINGDOM OF SAUDI ARABIA.

E-mail address: `khaledzennir4@yahoo.vom`