Ze Li*

Abstract

In this paper and the companion work [28], we prove that the Schrödinger map flows from \mathbb{R}^d with $d \ge 2$ to compact Kähler manifolds with small initial data in critical Sobolev spaces are global. The main difficulty compared with the constant sectional curvature case is that the gauged equation now is not self-contained due to the curvature part. Our main idea is to use a novel bootstrap-iteration scheme to reduce the gauged equation to an approximate constant curvature system in finite times of iteration. This paper with the companion work [28] solves the open problem raised by Tataru.

Contents

1	Introduction	1
2	Preliminaries	14
3	Iteration for Heat Flows	15
4	Evolution of SMF solutions along the heat direction	31
5	Evolution along the Schrödinger map flow direction	49
6	Iteration scheme	54
7	Proof of Theorem 1.1 and Theorem 1.2	60
8	Appendix A. Bilinear estimates	66
9	Appendix B. Proof of Remained Claims	68
Re	teferences	

1 Introduction

Let (N, J, h) be a Kähler manifold, the Schrödinger map flow (SMF) on Euclidean spaces is a map $u : (t, x) \in \mathbb{R} \times \mathbb{R}^d \mapsto N$ which satisfies

$$\begin{cases} u_t = J(\sum_{i=1}^d \nabla_i \partial_i u) \\ u \upharpoonright_{t=0} = u_0, \end{cases}$$
(1.1)

where ∇ denotes the induced covariant derivative in the pullback bundle u^*TN .

^{*}School of Mathematics, School of Mathematics and Statistics, Ningbo University, Ningbo, 315211, Zhejiang, P.R. China. Email: rikudosennin@163.com

Assume that N is isometrically embedded into \mathbb{R}^N , then (1.1) can be formulated as

$$\begin{cases} u_t = JP_u^N(\Delta_{\mathbb{R}^d} u) \\ u \upharpoonright_{t=0} = u_0, \end{cases}$$
(1.2)

where P_u^N denotes the orthogonal projection from \mathbb{R}^N onto $T_u \mathcal{N}$.

(1.1) plays a fundamental role in solid-state physics and is usually referred as the Landau-Lifshitz flow in physics literature. The various forms of SMF are commonly used in micromagnetics to model the effects of a magnetic filed on ferromagnetic materials (e.g. [27]). In the d = 1 and d = 2 case with $\mathcal{N} = \mathbb{S}^2$, SMF is referred as the ferromagnetic chain equation and the continuous isotropic Heisenberg spin model respectively (e.g. [42]).

The Schrödinger map flow can be viewed as a Hamiltonian flow on infinite dimensional symplectic manifolds, see Ding [8]. One of the conservation law of SMF is its energy defined by

$$E(u) = \frac{1}{2} \int_{\mathbb{R}^d} |\partial_x u|^2 dx.$$

And SMF has the scaling invariance property: $u(t, x) \mapsto u(\lambda^2 t, \lambda x)$. Thus d = 2 is the energy critical case. In the case $\mathcal{N} = \mathbb{S}^2$, SMF has masses as another conservation law:

$$M(u) = \frac{1}{2} \int_{\mathbb{R}^d} |u - P|^2 dx, \text{ if } ||u_0 - P||_{L^2_x} < \infty, \text{ for some } P \in \mathbb{S}^2.$$

However, the mass does not conserve for general target N.

We recall the following non-exhaustive list of works on Cauchy problems. The local well-posedness theory of Schrödinger map flows was developed by Sulem-Sulem-Bardos [32], Ding-Wang [9], McGahagan [30]. The global well-posedness theory was started by Chang-Shatah-Uhlenbeck [7] and Nahmod-Stefanov-Uhlenbeck [22]. And the d = 1 case with general targets was studied by Rodnianski-Rubinstein-Staffilani [31]. The global existence for small data in critical Besov spaces was proved by Ionescu-Kenig [14] and Bejenaru [1] independently. The small data global well-posedness theory in critical Sobolev spaces was done by Bejenaru-Ionescu-Kenig [3] for high dimensions $d \ge 4$. The two dimension case, which is energy critical, was studied by Bejenaru-Ionescu-Kenig-Tataru [4] where the global well-posedness theory for small data in critical Sobolev spaces was established for $\mathcal{N} = S^2$ with $d \ge 2$. And Dodson-Smith [10] studied the conditional global regularity problem for d = 2.

The stationary solutions of SMF are harmonic maps. So far, the dynamical behavior of SMF near harmonic maps is partly known in the equivariant case with d = 2, $\mathcal{N} = \mathbb{S}^2$. The works of Gustafson, Kang, Tsai, Nakanish [11, 12] proved asymptotic stability v.s. wind oscillating near harmonic maps in high equivariant classes. Bejenaru-Tataru [2] proved global stability v.s. instability of harmonic maps for 1-equivariant 2D SMF. The type II blowup solutions were constructed by Merle-Raphael-Rodnianski [21] and Perelman [26] for 1-equivariant 2D SMF. And the below threshold conjecture was verified for equivariant SMF from \mathbb{R}^2 into \mathbb{S}^2 or \mathbb{H}^2 by Bejenaru-Ionescu-Kenig-Tataru [5, 6].

All the mentioned global well-posedness results of SMF with $d \ge 2$ are for targets of \mathbb{S}^2 or \mathbb{H}^2 . Tataru raised the proof of small data global well-posedness in critical Sobolev spaces for general compact Kähler targets as an open problem in the survey report [19]. This work, which deals with the energy critical case d = 2, together with our companion work [28] solves this problem.

1.1 Main results

Before stating our main results, we introduce some notations on working spaces. For geometric PDEs, it is convenient to work in both intrinsic Sobolev spaces and extrinsic Sobolev spaces. For smooth maps from $\mathbb{R}^2 \to \mathcal{N}$ the intrinsic norms are defined by

$$||u||_{W^{k,p}}^{p} := \sum_{j=1}^{k} ||\nabla^{j}u||_{L^{p}_{x}(\mathbb{R}^{d})}^{p},$$

where ∇ denotes the induced covariant derivative in u^*TN .

Given a point $Q \in \mathcal{N}$, we define the extrinsic Sobolev space H_Q^k by

$$H_O^k := \{ u : \mathbb{R}^d \to \mathbb{R}^N \mid u(x) \in \mathcal{N}a.e. \text{ in } \mathbb{R}^d, \|u - Q\|_{H^k(\mathbb{R}^d)} < \infty \},\$$

which is equipped with the metric $d_Q(f, g) = ||f - g||_{H^k}$. Define \mathcal{H}_Q to be

$$\mathcal{H}_Q := \bigcap_{k=1}^{\infty} H_Q^k.$$

Our main results are the following two.

Theorem 1.1. Let d = 2, N be a 2n-dimensional compact Kähler manifold which is isometrically embedded into \mathbb{R}^N , and let $Q \in N$ be a given point. There exists a sufficiently small constant $\epsilon_* > 0$ such that if $u_0 \in \mathcal{H}_O$ satisfies

$$\|\partial_x u_0\|_{L^2_r} \le \epsilon_*,\tag{1.3}$$

then (1.1) with initial data u_0 evolves into a global unique solution $u \in C(\mathbb{R}; \mathcal{H}_Q)$. Moreover, as $|t| \to \infty$ the solution u converges to the constant map Q in the sense that

$$\lim_{|t| \to \infty} \|u(t) - Q\|_{L^{\infty}_{x}} = 0.$$
(1.4)

Furthermore, in the energy space, we also have

$$\lim_{t \to \infty} \|u(t) - \sum_{j=1}^{n} \Re(e^{it\Delta}h_{+}^{j}) - \sum_{j=1}^{n} \Im(e^{it\Delta}g_{+}^{j})\|_{\dot{H}_{x}^{1}} = 0,$$
(1.5)

for some functions $h^j_+, g^j_+ : \mathbb{R}^2 \to \mathbb{C}^N$ belonging to \dot{H}^1 with j = 1, ..., n.

Remark 1.1. The asymptotic behaviors (1.4) and (1.5) are new for SMF. The analogous result of (1.4) for wave maps was obtained in Part VII of Tao [36]. Similar result like (1.5) was recently obtained by the author [29] in the setting of SMF on hyperbolic planes. One can see (1.5) is natural by checking the trivial target $N = \mathbb{R}^{2n}$, see Remark 1.1 of [29] for instance.

We also prove the uniform bounds and well-posedness results analogous to that of [4].

Theorem 1.2. Let d = 2, $\sigma_1 \ge 0$. Let N be a compact Kähler manifold which is isometrically embedded into \mathbb{R}^N , and let $Q \in N$ be a given point. There exists a sufficiently small constant $\epsilon_{\sigma_1} > 0$ depending only on σ_1 such that the global solution $u = S_O(t)u_0 \in C(\mathbb{R}; \mathcal{H}_O)$ constructed in Theorem 1.1 satisfies the uniform bounds

$$\sup_{t \in \mathbb{R}} \|u(t) - Q\|_{H_x^{\sigma+1}} \le C_{\sigma}(\|u_0 - Q\|_{H_x^{\sigma+1}}), \ \forall \sigma \in [0, \sigma_1].$$
(1.6)

In addition, for any $\sigma \in [0, \sigma_1]$, the operator S_O admits a continuous extension

$$S_Q: \mathfrak{B}^{\sigma}_{\epsilon_{\sigma_1}} \to C(\mathbb{R}; H^{\sigma+1}),$$

where we denote

$$\mathfrak{B}^{\sigma}_{\epsilon} := \{ f \in H^{\sigma+1}_{O} : \| f - Q \|_{\dot{H}^1} \le \epsilon \}.$$

Remark 1.2. Theorem 1.1 holds as well for $d \ge 3$. This will be proved in our companion work [28]. The main proof of higher dimensions uses ideas of this work and some additional ingredients in heat flows. We will explain this issue at the end of introduction.

1.2 Caloric gauge and heat flows

For dispersive geometric PDEs, especially for critical problems, it is important to choose suitable gauges and function spaces adapted to the structure of nonlinearities (e.g. null structure). Most of these tools were developed in the study of wave map equations, see for instance [16, 17, 23, 33, 34, 18, 39, 40]. In this work, we will use Tao's caloric gauge and function spaces developed by [4, 13]. As observed by [4, 35], the caloric gauge is essential for eliminating bad frequency interactions in dimension two compared with Coulomb gauge. For the convenience of statement, we briefly recall the definition of caloric gauge.

First, let's recall the moving frame dependent quantities and some identities related to them, see [24] and [31] for more expositions. Let Greek indices run in $\{1, ..., n\}$. Let Roman indices run in $\{1, ..., 2n\}$ or $\{1, ..., d\}$ according to the context. Denote $\overline{\beta} = \beta + n$ for $\beta \in \{1, ..., n\}$.

Let N be a 2n-dimensional compact Kähler manifold. Since $\mathbb{R}^2 \times [-T, T]$ is contractible, there must exist global orthonormal frames for $u^*(TN)$. Using the complex structure one can assume the orthonormal frames are of the form

$$\mathbf{E} := \{ e_1(t, x), Je_1(t, x), \dots, e_n(t, x), Je_n(t, x) \}.$$
(1.7)

Let $\psi_i = (\psi_i^1, \psi_i^{\bar{1}}, ..., \psi_i^n, \psi_i^{\bar{n}})$ for i = 0, 1, 2 be the components of $\partial_{t,x} u$ in the frame **E**:

$$\psi_i^{\alpha} = \langle \partial_i u, e_{\alpha} \rangle, \\ \psi_i^{\overline{\alpha}} = \langle \partial_i u, J e_{\alpha} \rangle.$$
(1.8)

We always use 0 to represent *t* in lower index. The isomorphism of \mathbb{R}^{2n} to \mathbb{C}^n induces a \mathbb{C}^n -valued function defined by $\phi_i^{\beta} = \psi_i^{\beta} + \sqrt{-1}\psi_i^{\overline{\beta}}$ with $\beta = 1, ..., n$. Conversely, given function $\phi : [-T, T] \times \mathbb{R}^2 \to \mathbb{C}^n$, we associate it with a section $\phi \mathbf{E}$ of the bundle $u^*(T\mathcal{N})$ via

$$\phi \Longleftrightarrow \phi \mathbf{E} := \mathfrak{R}(\phi^{\beta})e_{\beta} + \mathfrak{I}(\phi^{\beta})Je_{\beta}, \tag{1.9}$$

where $(\phi^1, ..., \phi^n)$ denotes the components of ϕ . Then *u* induces a covariant derivative on the trivial complex vector bundle over base manifold $[-T, T] \times \mathbb{R}^d$ with fiber \mathbb{C}^n defined by

$$D_i\varphi^{\beta} = \partial_i\varphi^{\beta} + \sum_{\alpha=1}^n \left([A_i]_{\alpha}^{\beta} + \sqrt{-1} [A_i]_{\alpha}^{\overline{\beta}} \right) \varphi^{\alpha},$$

where the induced connection coefficient matrices are defined by

$$[A_i]_q^p = \left\langle \nabla_i e_p, e_q \right\rangle$$

Schematically we write $D_i = \partial_i + A_i$. Recall the torsion free identity and the commutator identity

$$D_i \phi_i = D_j \phi_i \tag{1.10}$$

$$[D_i, D_j]\varphi = \left(\partial_i A_j - \partial_j A_i + [A_i, A_j]\right)\varphi \Longleftrightarrow \mathbf{R}(\partial_i u, \partial_j u)(\varphi \mathbf{E}).$$
(1.11)

Schematically, we write $[D_i, D_j] = \mathcal{R}(\phi_i, \phi_j)$. With the notations given above, (1.1) can be written as

$$\phi_t = \sqrt{-1} \sum_{i=1}^2 D_i \phi_i.$$
(1.12)

[25] proved the heat flow with initial data u(t, x) below threshold energy would converge to Q as $s \to \infty$ in the topology of $C([-T, T]; C_x^{\infty})$. Tao's caloric gauge is defined as follows:

Definition 1.1. Let $u(t, x) : [-T, T] \times \mathbb{R}^2 \to N$ be a solution of (1.1) in $C([-T, T]; \mathcal{H}_Q)$. For a given orthonormal frame $E^{\infty} := \{e_1^{\infty}, Je_1^{\infty}, ..., e_n^{\infty}, Je_n^{\infty}\}$ for T_QN , a caloric gauge is a tuple consisting of a map $v : \mathbb{R}^+ \times [-T, T] \times \mathbb{R}^2 \to N$ and orthonormal frames $\mathbf{E}(v(s, t, x)) := \{e_1, Je_1, ..., e_n, Je_n\}$ such that

$$\begin{cases} \partial_s v = \sum_{i=1}^2 \nabla_i \partial_i v \\ v(0,t,x) = u(t,x) \end{cases}$$
(1.13)

and

$$\begin{cases}
\nabla_s e_k = 0, \ k = 1, ..., n. \\
\lim_{s \to \infty} e_k = e_k^{\infty}.
\end{cases}$$
(1.14)

Denote

$$\mathcal{H}_Q(T) := C([-T,T];\mathcal{H}_Q).$$

Proposition 1.1. Let $u \in \mathcal{H}_Q(T)$ solve SMF with $u_0 \in \mathcal{H}_Q$. For any fixed frame $E^{\infty} := \{e_k^{\infty}, Je_k^{\infty}\}_{k=1}^n$ for T_QN , there exists a unique corresponding caloric gauge defined in Definition 1.1. Moreover, we have for i = 1, 2 and p, q = 1, ..., 2n

$$\lim_{s \to \infty} [A_i]_p^q(s, t, x) = 0$$
$$\lim_{s \to \infty} [A_t]_p^q(s, t, x) = 0$$

Particularly, we have for i = 1, 2, s > 0,

$$\begin{split} & [A_i]_q^p(s,t,x) = -\int_s^\infty \langle \mathbf{R} \left(\partial_s v(\widetilde{s}) \right), \partial_i v(\widetilde{s}) e_p, e_q \rangle d\widetilde{s} \\ & [A_t]_q^p(s,t,x) = -\int_s^\infty \langle \mathbf{R} \left(\partial_s v(\widetilde{s}), \partial_t v(\widetilde{s}) \right) e_p, e_q \rangle d\widetilde{s}. \end{split}$$

Proof. The proof is standard (see e.g. [25]). The only new issue here is the complex structure J. But this will not cause any trouble since J commutes with ∇_s .

Given $u \in \mathcal{H}_Q(T)$ which solves (1.1), let $v : \mathbb{R}^+ \times [-T, T] \times \mathbb{R}^2 \to \mathcal{N}$ be the solution to (1.13). Let $\{e_\alpha, Je_\alpha\}_{\alpha=1}^n$ be the corresponding caloric gauge. Define the heat tension field ϕ_s to be

$$\phi_s^{\alpha} = \langle \partial_s v, e_{\alpha} \rangle + \sqrt{-1} \langle \partial_s v, J e_{\alpha} \rangle, \ \alpha = 1, ..., n$$

And define the differential fields to be

$$\phi_i^{\alpha} = \langle \partial_i v, e_{\alpha} \rangle + \sqrt{-1} \langle \partial_i v, J e_{\alpha} \rangle, \ \alpha = 1, ..., n,$$

where i = 1, 2 refers to the variable x_i , i = 1, 2, and i = 0 refers to the variable *t*.

Lemma 1.1. The heat tension filed ϕ_s satisfies

$$\phi_s = \sum_{j=1}^2 D_j \phi_j.$$
(1.15)

The differential fields $\{\phi_i\}_{i=1}^2$ *along the heat flow satisfy*

$$\partial_{s}\phi_{i} = \sum_{j=1}^{2} D_{j}D_{j}\phi_{i} + \sum_{j=1}^{2} \mathcal{R}(\phi_{i}, \phi_{j})\phi_{j}.$$
(1.16)

And when s = 0, along the Schrödinger flow direction, $\{\phi_i\}_{i=1}^2$ satisfy

$$-\sqrt{-1}D_t\phi_i = \sum_{j=1}^2 D_j D_j\phi_i + \sum_{j=1}^2 \mathcal{R}(\phi_i, \phi_j)\phi_j.$$
(1.17)

1.3 Function Spaces built by [4]

We recall the spaces developed by Bejenaru-Ionescu-Kenig-Tataru [4]. Given a unite vector $\mathbf{e} \in \mathbb{S}^1$ we denote its orthogonal complement of \mathbb{R}^2 by \mathbf{e}^{\perp} . The lateral space $L_{\mathbf{e}}^{p,q}$ is defined by the norm

$$\|f\|_{L^{p,q}_{\mathbf{e}}} = \left(\int_{\mathbb{R}} \left(\int_{\mathbf{e}^{\perp} \times \mathbb{R}} \left|f(t, x_1 \mathbf{e} + x')\right|^q dx' dt\right)^{\frac{p}{q}} dx_1\right)^{\frac{1}{p}},$$

with standard modifications when either $p = \infty$ or $q = \infty$. And for any given $\lambda \in \mathbb{R}$, $W \subset \mathbb{R}$, we define the spaces $L_{\mathbf{e},\lambda}^{p,q}$, $L_{\mathbf{e},W}^{p,q}$ with norms

$$\begin{split} \|f\|_{L^{p,q}_{\mathbf{e},\lambda}} &= \|G_{\lambda\mathbf{e}}(f)\|_{L^{p,q}_{\mathbf{w}}} \\ \|f\|_{L^{p,q}_{\mathbf{e},W}} &= \inf_{f = \sum_{\lambda \in W} f_{\lambda}} \sum_{\lambda \in W} \|f_{\lambda}\|_{L^{p,q}_{\mathbf{e},\lambda}}, \end{split}$$

where G_a denotes the Galilean transform:

$$G_{\mathbf{a}}(f)(t,x) = e^{-\frac{1}{2}ix\cdot\mathbf{a}}e^{-\frac{i}{4}t|\mathbf{a}|^2}f(x+t\mathbf{a},t).$$

We remark that the lateral space was developed by Linares-Ponce [20], Kenig-Ponce-Vega [15] and Ionescu-Kenig [14].

The main dyadic function spaces $N_k(T)$, $F_k(T)$, $G_k(T)$ are recalled as follows: Given $T \in \mathbb{R}^+$, $k \in \mathbb{Z}$, let $I_k := \{\eta \in \mathbb{R}^2 : 2^{k-1} \le |\eta| \le 2^{k+1}\}$ and

$$L_k^2(T) := \{ g \in L^2([-T, T] \times \mathbb{R}^2) : \mathcal{F}g(t, \eta) \text{ is supoorted in } \mathbb{R} \times I_k \}.$$
(1.18)

Given $\mathcal{L} \in \mathbb{Z}_+$, $T \in (0, 2^{2\mathcal{L}}]$, $k \in \mathbb{Z}$, define

$$W_k = \{\lambda \in [-2^{2k}, 2^{2k}] : 2^{k+2\mathcal{L}} \lambda \in \mathbb{Z}\}.$$
(1.19)

The $N_k(T)$, $F_k(T)$, $G_k(T)$ spaces are Banach spaces of functions in $L_k^2(T)$ for which the associated norms are finite:

$$\begin{split} \|g\|_{F_{k}^{0}(T)} &:= \|g\|_{L_{t}^{\infty}L_{x}^{2}} + 2^{-\frac{k}{2}} \|g\|_{L_{k}^{4}L_{t}^{\infty}} + \|g\|_{L^{4}} + 2^{-\frac{k}{2}} \sup_{\mathbf{e} \in \mathbb{S}^{1}} \|g\|_{L_{\mathbf{e},W_{k+40}}^{2,\infty}} \\ \|g\|_{F_{k}(T)} &:= \inf_{j \in \mathbb{Z}_{+}, n_{1}, \dots, n_{j} \in \mathbb{N}} \inf_{g = g_{n_{1}} + \dots + g_{n_{j}}} \sum_{l=1}^{j} 2^{n_{l}} \|g_{n_{l}}\|_{F_{k+n_{l}}^{0}} \\ \|g\|_{G_{k}(T)} &:= \|g\|_{F_{k}^{0}} + 2^{-\frac{k}{6}} \sup_{\mathbf{e} \in \mathbb{S}^{1}} \|g\|_{L_{\mathbf{e}}^{3,6}} + 2^{\frac{k}{6}} \sup_{|k-j| \leq 20} \sup_{\mathbf{e} \in \mathbb{S}^{1}} \|P_{j,\mathbf{e}}g\|_{L_{\mathbf{e},l}^{6,3}} \\ &+ 2^{\frac{k}{2}} \sup_{|k-j| \leq 20} \sup_{\mathbf{e} \in \mathbb{S}^{1}} \sup_{|\lambda| < 2^{k-40}} \|P_{j,\mathbf{e}}g\|_{L_{\mathbf{e},l}^{\infty,2}} \\ \|g\|_{N_{k}(T)} &:= \inf_{g = g_{1} + g_{2} + g_{3} + g_{4}} \|g_{1}\|_{L^{\frac{4}{3}}} + 2^{\frac{k}{6}} \|g_{2}\|_{L^{\frac{2}{3},\frac{5}{5}}} + 2^{\frac{k}{6}} \|g_{3}\|_{L^{\frac{2}{3},\frac{5}{5}}} \\ &+ 2^{-\frac{k}{2}} \sup_{\mathbf{e} \in \mathbb{S}^{1}} \|g_{4}\|_{L^{\frac{1,2}{6},W_{k-40}}}, \end{split}$$

where $\{\mathbf{e}_1, \mathbf{e}_2\} \subset \mathbb{S}^1$ consist of the standard basis of \mathbb{R}^2 . The G_k, F_k spaces were built by Bejenaru-Ionescu-Kenig-Tataru [4].

1.4 Overview of the proof

Main difficulty for general targets. The new difficulty arising in the case of general targets is to control the curvature terms in frequency localized spaces. Since the curvature term relates with the map itself, it cannot be written in a self-closed form of differential fields and heat tension fields { ϕ_x , ϕ_s }. Thus directly working with the moving frame dependent quantities may lose control of curvature terms, which is much serious when frequency interactions are considered. In the wave map setting, the general targets case was solved by Tataru [41] using Tao's micro-local gauge and Tataru's function spaces. It is important that the wave map equation is semilinear in the extrinsic form, and the micro-local gauge adapts to the extrinsic formulation well. However, for SMF, on one side, since the extrinsic form equation is not a self-contained system where the curvature term is not determined by differential fields. The two conflict sides make solving SMF for general targets challenging.

Outline of proof for d = 2. Let us sketch the outline of proof in the d = 2 case. The whole proof is divided into ten steps. Given $\delta > 0$, let $\{a_k\}_{k \in \mathbb{Z}}$ be a positive sequence, we call it a frequency envelope of order δ if

$$\sum_{k \in \mathbb{Z}} a_k^2 < \infty, \text{ and } a_j \le 2^{\delta |l-j|} a_l, \ \forall j, l \in \mathbb{Z}.$$

We call the frequency envelope $\{a_k\}$ an ϵ -envelope if it additionally satisfies

$$\sum_{k\in\mathbb{Z}}a_k^2\leq\epsilon^2.$$

Step 1. Tracking $L_t^{\infty} L_x^2$ bounds along the heat flow direction. Recall the extrinsic formulations of heat flows: Assume that the target manifold N is isometrically embedded into \mathbb{R}^N , then the heat flow equation can be formulated as

$$\partial_{s}v^{l} - \Delta v^{l} = \sum_{a=1}^{2} \sum_{i,j=1}^{N} S^{l}_{ij} \partial_{a}v^{i} \partial_{a}v^{j}, \ l = 1, ..., N,$$
(1.20)

where $S = \{S_{ii}^l\}$ denotes the second fundamental form of the embedding $\mathcal{N} \hookrightarrow \mathbb{R}^N$. For $u \in \mathcal{H}_Q(T)$, define

$$\gamma_k(\sigma) = \sup_{k' \in \mathbb{Z}} 2^{-\delta|k-k'|} 2^{\sigma k'+k'} ||P_{k'}u||_{L^{\infty}_t L^2_x}, \ \sigma \ge 0, \ \delta = \frac{1}{800}$$

Denote $\{\gamma_k\}$ the frequency envelope for the energy norm, i.e. $\gamma_k = \gamma_k(0)$. The first result of Step 1 is stated for $\sigma \in [0, \frac{5}{4}]$:

Proposition 1.2. Assume that $u \in \mathcal{H}_Q(T)$ satisfies

$$\|\partial_x u\|_{L^\infty_x L^2_x} = \epsilon_1 \ll 1. \tag{1.21}$$

And let v(s, t, x) be the solution of heat flow (1.20) with initial data u(t, x). Then v satisfies

$$\sup_{s \ge 0} (1 + s2^{2k})^{31} 2^k ||P_k v||_{L_t^{\infty} L_x^2} \lesssim 2^{-\sigma k} \gamma_k(\sigma)$$

for all $\sigma \in [0, \frac{99}{100}]$, $k \in \mathbb{Z}$. Moreover, for any $\sigma \in [\frac{99}{100}, \frac{5}{4}]$, $k \in \mathbb{Z}$, we have

$$\sup_{s\geq 0} (1+s2^{2k})^{30} 2^{\sigma k+k} ||P_k v||_{L^{\infty}_t L^2_x} \leq_M \gamma_k(\sigma) + \gamma_k(\sigma-\frac{3}{8}) \gamma_k(\frac{3}{8}).$$

Remark 1.3. The power of $(1 + s2^{2k})$ in Proposition 1.2 can be chosen to be any $M \in \mathbb{Z}_+$ with additionally assuming that ϵ_1 is sufficiently small depending on M. See Proposition 1.3 below.

The second result of this step is bounds of $2^{(\sigma+1)k} \|P_k v\|_{L^{\infty}_t L^2_x}$ for $\sigma \in [0, \frac{j+1}{4}]$:

Proposition 1.3. (j-th iteration) Let $j \in \mathbb{N}$, $M \in \mathbb{Z}_+$. Assume that $u \in \mathcal{H}_Q(T)$ satisfies (1.21) with ϵ_1 sufficiently small depending on j + M. Let v(s, t, x) be the solution of heat flow (1.20) with initial data u(t, x). Then for $\sigma \in [0, 1 + \frac{j}{4}]$ and any $k \in \mathbb{Z}$, v satisfies

$$\sup_{s \in [0,\infty)} (1 + s2^{2k})^M 2^{k+\sigma k} \|P_k v\|_{L_t^{\infty} L_x^2} \lesssim \gamma_k^{(j)}(\sigma).$$
(1.22)

where $\{\gamma_{k}^{(j)}(\sigma)\}$ are defined in (3.21)-(3.23).

Step 2. Pretreat curvature terms. The curvature part in master equation (1.17) can be schematically written as

$$\mathfrak{R}[\mathcal{R}(\phi_i,\phi_j)\phi_j]^{\alpha} = \sum_{1 \le j_0, j_1, j_2 \le 2n} \langle \mathbf{R}(e_{j_0},e_{j_1})e_{j_2},e_{\alpha} \rangle \psi_i^{i_0}\psi_j^{i_1}\psi_j^{i_2}$$
$$\mathfrak{I}[\mathcal{R}(\phi_i,\phi_j)\phi_j]^{\alpha} = \sum_{1 \le j_0, j_1, j_2 \le 2n} \langle \mathbf{R}(e_{j_0},e_{j_1})e_{j_2},e_{\overline{\alpha}} \rangle \psi_i^{i_0}\psi_j^{i_1}\psi_j^{i_2}.$$

With abuse of notations, we denote

$$\mathcal{G} = \langle \mathbf{R}(e_{j_0}, e_{j_1})e_{j_2}, e_{j_3} \rangle,$$

for any given index $j_0, ..., j_3 \in \{1, ..., 2n\}$. And let $\phi_i \diamond \phi_j$ denote the linear combinations of multiplications of real and imaginary parts of ϕ_i, ϕ_j , i.e. $\sum_{ij} c_{ij} \phi_i^{\pm} \phi_j^{\pm}$, where we denote $\phi_j^{+} = \Re \phi_j, \phi_j^{-} = \Im \phi_j$. Then the master equation (1.17) is now schematically written as

$$-\sqrt{-1}D_t\phi_i = \sum_{j=1}^2 D_j D_j\phi_i + \sum_{j=1}^2 \mathcal{G}\phi_i \diamond \phi_j \diamond \phi_j.$$
(1.23)

Meanwhile, the connection coefficients in D_i also depend on curvatures and can be schematically written as

$$[A_j]_q^p(s) = -\int_s^\infty \phi_s \diamond \phi_j \mathcal{G} ds'.$$

We shall perform dynamical separation to G. In fact, by caloric condition and twice dynamic separation, G can be decomposed into

$$\begin{aligned} \mathcal{G}(s) &= \langle \mathbf{R}(e_{j_0}, e_{j_1})e_{j_2}, e_{j_3} \rangle(s) \\ &= \Gamma^{\infty} - \Gamma_l^{\infty,(1)} \int_s^{\infty} \psi_s^l(\widetilde{s})d\widetilde{s} - \int_s^{\infty} \psi_s^l(\widetilde{s}) \left(\int_{\widetilde{s}}^{\infty} \psi_s^p(s')(\widetilde{\nabla}^2 \mathbf{R})(e_l, e_p; e_{j_0}, ..., e_{j_3})ds' \right) d\widetilde{s}. \\ &:= \Gamma^{\infty} + \mathcal{U}_{00} + \mathcal{U}_{01} + \mathcal{U}_I + \mathcal{U}_{II}, \end{aligned}$$

where we denote

constant limit part
$$\Gamma^{\infty} := \lim_{s \to \infty} \mathcal{G}(s); \quad \Gamma_l^{\infty,(1)} := \lim_{s \to \infty} (\widetilde{\nabla} \mathbf{R})(e_l; e_{j_0}, ..., e_{j_3})$$

one order terms $\mathcal{U}_{00} := -\Gamma_l^{\infty} \int_s^{\infty} \sum_{i=1}^2 (\partial_i \psi_i)^l ds',$

and quadratic terms by

$$\begin{aligned} \mathcal{U}_{01} &:= -\int_{s}^{\infty} \sum_{i=1}^{2} (\partial_{i}\psi_{i}) \left((\widetilde{\nabla}\mathbf{R})(e_{l}; e_{j_{0}}, ..., e_{j_{3}}) - \Gamma_{l}^{\infty,(1)} \right) ds' \\ \mathcal{U}_{I} &:= -\Gamma_{l}^{\infty,(1)} \int_{s}^{\infty} \sum_{i=1}^{2} (A_{i}\psi_{i})^{l} ds' \\ \mathcal{U}_{II} &:= -\int_{s}^{\infty} \sum_{i=1}^{2} (A_{i}\psi_{i})^{l} (\widetilde{s}) \left(\int_{\widetilde{s}}^{\infty} \psi_{s}^{p}(s') (\widetilde{\nabla}^{2}\mathbf{R})(e_{l}, e_{p}; e_{j_{0}}, ..., e_{j_{3}}) ds' \right) d\widetilde{s} \\ &= -\int_{s}^{\infty} \sum_{i=1}^{2} (A_{i}\psi_{i})^{l} (\widetilde{s}) \left((\widetilde{\nabla}\mathbf{R})(e_{l}; e_{j_{0}}, ..., e_{j_{3}}) - \Gamma_{l}^{\infty,(1)} \right) d\widetilde{s}. \end{aligned}$$

Here, with abuse of notations, A_j denotes the $2n \times 2n$ real-valued matrix with elements $\{[A_j]_q^p\}_{p,q=1}^{2n}$. The constant limit part and the one order terms will be dominated by frequency envelopes of $\{\phi_i\}$, the bounds of quadratic terms essentially rely on a delicate bootstrap on the term:

$$\widetilde{\mathcal{G}}_l^{(1)} := (\widetilde{\nabla} \mathbf{R})(e_l; e_{j_0}, ..., e_{j_3}) - \Gamma_l^{\infty, (1)}$$

We also need higher order derivatives of \mathcal{G} . Given $k \in \mathbb{N}$, let

$$\begin{aligned} \mathcal{G}_{l_1,...,l_k}^{(k)} &:= (\widetilde{\nabla}^{(k)} \mathbf{R})(e_{l_1},...,e_{l_k};e_{j_0},...,e_{j_3}) \\ \widetilde{G}_{l_1,...,l_k}^{(k)} &:= \mathcal{G}_{l_1,...,l_k}^{(k)} - \Gamma_{l_1,...,l_k}^{\infty,(k)}, \end{aligned}$$

where we denote

$$\Gamma_{l_1,\ldots,l_k}^{\infty,(k)} := \lim_{s \to \infty} \mathcal{G}_{l_1,\ldots,l_k}^{(k)}(s).$$

Similarly, we perform dynamical separation to frames. In fact, let \mathcal{P} be the isometric embedding of \mathcal{N} into \mathbb{R}^N , and let $\{e_l\}_{l=1}^{2n}$ be the caloric gauge built in Prop. 1.1. With abuse of notations, we denote

$$[d\mathcal{P}]^{(k)} := (\mathbf{D}^k d\mathcal{P})(\underbrace{e, \dots, e}_k; e)$$
$$[\widetilde{d\mathcal{P}}]^{(k)} := [d\mathcal{P}]^{(k)} - \lim_{s \to \infty} [d\mathcal{P}]^{(k)}$$

Step 3. Tracking $L^4 \cap L^{\infty}_t L^2_x$ bounds for curvature terms and frames along heat direction.

Proposition 1.4. Let d = 2. And let $u \in \mathcal{H}_Q(T)$ be a solution of SMF. Denote v(s, t, x) the solution to heat flow with initial data u(t, x), and denote $\{\phi_i\}_{i=0}^2$ the corresponding differential fields under the caloric gauge. Assume that $\{\beta_k(\sigma)\}$ is a frequency envelope of order δ such that for all $i = 1, 2, k \in \mathbb{Z}$,

$$2^{\sigma k} \|\phi_i(\uparrow_{s=0})\|_{L^{\infty}_t L^2_x \cap L^4_{t_x}} \le \beta_k(\sigma).$$
(1.24)

• There exists a sufficiently small constant $\epsilon > 0$ such that if

$$\sum_{k \in \mathbb{Z}} |\beta_k(0)|^2 < \epsilon, \tag{1.25}$$

then we have for any $m \in \mathbb{N}$, any $\sigma \in [0, \frac{99}{100}]$, $s \in [2^{2j-1}, 2^{2j+1})$, $j, k \in \mathbb{Z}$,

$$\begin{split} \|P_k \widetilde{\mathcal{G}}^{(m)}\|_{L^4 \cap L^{\infty}_t L^2_x} &\lesssim_m 2^{-\sigma k-k} \beta_k(\sigma) (2+s2^{2k})^{-30} \\ \|P_k \phi_s\|_{L^4 \cap L^{\infty}_t L^2_x} &\lesssim 2^{-\sigma k+k} \left[1_{k+j \ge 0} (1+s2^{2k})^{-30} \beta_k(\sigma) + 1_{k+j \le 0} \sum_{k \le l \le -j} \beta_l(\sigma) \beta_l \right] \\ \|P_k([\widetilde{d\mathcal{P}}]^{(m)})\|_{L^{\infty}_t L^2_x \cap L^4} &\lesssim_m \beta_k(\sigma) (1+s2^{2k})^{-29} 2^{-\sigma k-k} \\ \|P_k A_i\|_{L^{\infty}_t L^2_x} &\lesssim \beta_{k,s}(\sigma) (1+s2^{2k})^{-27} 2^{-\sigma k}. \end{split}$$

• Furthermore, given $j, M \in \mathbb{Z}_+$, if $\{\beta_k(\sigma)\}$ is a frequency envelope of order $\frac{1}{2^j}\delta$, then similar results hold for $\sigma \in [0, \frac{1}{4}j+1]$ and ϵ sufficiently small depending only on $j, M \in \mathbb{Z}_+$. Particularly, for any $m \in \mathbb{N}$, $k \in \mathbb{Z}$ and any $\sigma \in [0, \frac{1}{4}j+1]$, one has

$$\begin{aligned} &(1+s2^{2k})^{M+2}2^{\sigma k+k} \|P_k\widetilde{\mathcal{G}}^{(m)}\|_{L^4 \cap L^\infty_t L^2_x} \lesssim_{m,M} \beta_k^{(j)}(\sigma) \\ &(1+s2^{2k})^{M+1}2^{\sigma k+k} \|P_k([\widetilde{d\mathcal{P}}]^{(m)})\|_{L^\infty_t L^2_x \cap L^4_{t,x}} \lesssim_{m,M} \beta_k^{(j)}(\sigma) \\ &(1+s2^{2k})^M 2^{\sigma k} \|P_k A_i\|_{L^\infty_t L^2_x} \lesssim_M \beta_{k,s}^{(j)}(\sigma). \end{aligned}$$

Remark 1.4. The $\{\beta_k^{(j)}\}$ and $\{\beta_{k,s}^{(j)}(\sigma)\}$ are defined as (3.21)-(3.23).

Step 4.1. $F_k \cap S_k^{\frac{1}{2}}$ **bounds for connections along the heat direction.** In this step, we prove **Lemma 1.2.** Given $\sigma \in [0, \frac{99}{100}]$, let $\{h_k(\sigma)\}$ be frequency envelopes defined by

$$h_k(\sigma) = \sup_{k' \in \mathbb{Z}, j=1,2} 2^{-\delta|k-k'|} 2^{\sigma k'} (1 + s 2^{2k'})^4 ||P_{k'}\phi_j||_{F_{k'}(T)}.$$
(1.26)

Let $\{b_k\}$ be a ε -frequency envelope. Assume that for any $k, j \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1})$, there holds

$$2^{\frac{1}{2}k} \| P_k \widetilde{\mathcal{G}}^{(1)} \|_{L^4_x L^\infty_t(T)} \le \varepsilon^{-\frac{1}{4}} h_k [(1 + s2^{2k})^{-9} \mathbf{1}_{j+k \ge 0} + \mathbf{1}_{j+k \le 0} 2^{\delta |k+j|}].$$
(1.27)

Then, if $\varepsilon > 0$ is sufficiently small, for $\sigma \in [0, \frac{99}{100}]$ one has

$$\|P_k A_i(s)\|_{F_k(T) \cap S_k^{\frac{1}{2}}} \lesssim h_{k,s}(\sigma) 2^{-\sigma k} (1+s 2^{2k})^{-4}.$$

Step 4.2. F_k bounds along the heat direction without assuming (1.27).

Lemma 1.3. Let $\{b_k\}$ be a ε -frequency envelope. Given $\sigma \in [0, \frac{99}{100}]$, suppose that $\{b_k(\sigma)\}$ are also frequency envelopes, and $\{h_k(\sigma)\}$ are the frequency envelopes defined by (1.26). Assume that for i = 1, 2,

$$\begin{aligned} \|P_k \phi_i \uparrow_{s=0} \|_{F_k(T)} &\leq b_k(\sigma) 2^{-\sigma k}, \ \sigma \in [0, \frac{99}{100}] \\ \|P_k \phi_i \uparrow_{s=0} \|_{L^4_{t,x}} &\leq b_k(\sigma) 2^{-(\sigma-1)k}, \ \sigma \in [0, \frac{99}{100}] \\ \|P_k \phi_i(s)\|_{F_k(T)} &\leq \varepsilon^{-\frac{1}{2}} b_k(1 + s 2^{2k})^{-4}. \end{aligned}$$

Then, if $\varepsilon > 0$ is sufficiently small, for $\sigma \in [0, \frac{99}{100}]$ one has

$$\begin{split} \|P_{k}A_{i}(s)\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}} &\leq h_{k,s}(\sigma)2^{-\sigma k}(1+s2^{2k})^{-4} \\ \|P_{k}\phi_{i}(s)\|_{F_{k}(T)} &\leq b_{k}(\sigma)2^{-\sigma k}(1+s2^{2k})^{-4} \\ \|P_{k}A_{i}\uparrow_{s=0}\|_{L_{t,x}^{4}} &\leq b_{k}(\sigma)2^{-\sigma k}, \ i=1,2. \\ \|P_{k}\phi_{t}(s)\|_{L_{t,x}^{4}} &\leq b_{k}(\sigma)2^{-(\sigma-1)k}(1+2^{2k}s)^{-2} \\ \|P_{k}A_{t}\uparrow_{s=0}\|_{L_{t,x}^{2}} &\leq \varepsilon b_{k}(\sigma)2^{-\sigma k}, \ \sigma \in [\frac{1}{100}, \frac{99}{100}] \\ \|P_{k}A_{t}\uparrow_{s=0}\|_{L_{t,x}^{2}} &\leq \varepsilon^{2}, \ \sigma \in [0, \frac{99}{100}]. \end{split}$$

Step 5. G_k bounds along the SMF direction for $\sigma \in [0, \frac{99}{100}]$.

Proposition 1.5. Assume that $\sigma \in [0, \frac{99}{100}]$. Given any $\mathcal{L} \in \mathbb{Z}_+$, assume that $T \in (0, 2^{2\mathcal{L}}]$. Let ϵ_0 be a sufficiently small constant. Assume that $\{c_k\}$ is an ϵ_0 -frequency envelope of order δ . And let $\{c_k(\sigma)\}$ be another frequency envelope of order δ . Let $u \in \mathcal{H}_O(T)$ be the solution to SMF with initial data u_0 which satisfies

$$\begin{aligned} \|P_k \nabla u_0\|_{L^2_x} &\leq c_k \\ \|P_k \nabla u_0\|_{L^2_x} &\leq c_k(\sigma) 2^{-\sigma k}. \end{aligned}$$

Denote $\{\phi_i\}$ the corresponding differential fields of the heat flow initiated from u. Suppose also that at the heat initial time s = 0,

$$||P_k\phi_i||_{G_k(T)} \le \epsilon_0^{-\frac{1}{2}}c_k.$$

Then, when s = 0, we have for all $i = 1, 2, k \in \mathbb{Z}$,

$$\begin{aligned} \|P_k\phi_i\|_{G_k(T)} &\leq c_k \\ \|P_k\phi_i\|_{G_k(T)} &\leq c_k(\sigma)2^{-\sigma k}. \end{aligned}$$

Step 6. Improve F_k bounds of $P_k \widetilde{\mathcal{G}}^{(1)}$ once.

Lemma 1.4. Let $u \in \mathcal{H}_Q(T)$ solve SMF with data u_0 . Let $\{c_k\}$ be ϵ_0 -frequency envelopes of order $\frac{1}{2}\delta$. Given any $\sigma \in [0, \frac{99}{100}]$, let $\{c_k(\sigma)\}$ be another frequency envelopes of order δ such that

$$\|P_k \nabla u_0\|_{L^2_x} \le c_k$$
$$\|P_k \nabla u_0\|_{L^2_x} \le c_k(\sigma) 2^{-\sigma k}.$$

Then for ϵ_0 sufficiently small there holds

$$2^{k} \| P_{k} \widetilde{\mathcal{G}}^{(1)} \|_{F_{k}(T)} \le c_{k}(\sigma) 2^{-\sigma k} [(1 + 2^{2k+2k_{0}})^{-20} \mathbf{1}_{k+k_{0} \ge 0} + \mathbf{1}_{k+k_{0} \le 0} 2^{\delta |k+k_{0}|}]$$

for any $\sigma \in [0, \frac{99}{100}]$, $k, k_0 \in \mathbb{Z}$, $s \in [2^{2k_0-1}, 2^{2k_0+1})$.

Step 7. Improve $F_k \cap S_k^{\frac{1}{2}}$ bounds of $\{P_k A_j\}_{j=0}^2$ and derive parabolic estimates for $\{\phi_j\}_{j=0}^2$ with $\sigma \in [0, \frac{5}{4}]$.

Lemma 1.5. Let $u \in \mathcal{H}_Q(T)$ be solution to SMF with initial data $u_0 \in \mathcal{H}_Q$. Define frequency envelopes $\{c^{(1)}(\sigma)\}$ as Def. 6.1. Given any $\sigma \in [0, \frac{5}{4}]$, let $\{b_k(\sigma)\}$ be frequency envelopes of order δ . And assume that

$$b_k(\sigma) \leq c_k^{(1)}(\sigma), \ \forall \sigma \in [0, \frac{99}{100}].$$

Assume also that for i = 1, 2,

$$\begin{aligned} \|P_k\phi_i \upharpoonright_{s=0}\|_{F_k(T)} &\leq b_k(\sigma) 2^{-\sigma k}, \ \sigma \in [0, \frac{5}{4}] \\ \|P_k\phi_t \upharpoonright_{s=0}\|_{L^4_{t,x}} &\leq b_k(\sigma) 2^{-(\sigma-1)k}, \ \sigma \in [0, \frac{5}{4}]. \end{aligned}$$

Then, if $\varepsilon > 0$ is sufficiently small, for $\sigma \in [0, \frac{5}{4}]$ one has for any $\sigma \in [0, \frac{5}{4}]$

$$\begin{split} \|P_{k}A_{i}(s)\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}} &\leq h_{k,s}^{(1)}(\sigma)2^{-\sigma k}(1+s2^{2k})^{-4} \\ \|P_{k}\phi_{i}(s)\|_{F_{k}(T)} &\leq b_{k}^{(1)}(\sigma)2^{-\sigma k}(1+s2^{2k})^{-4} \\ \|P_{k}A_{i}\uparrow_{s=0}\|_{L_{t,x}^{4}} &\leq b_{k}^{(1)}(\sigma)2^{-\sigma k}, \ i=1,2. \\ \|P_{k}\phi_{t}(s)\|_{L_{t,x}^{4}} &\leq b_{k}^{(1)}(\sigma)2^{-(\sigma-1)k}(1+2^{2k}s)^{-2} \\ \|P_{k}A_{t}\uparrow_{s=0}\|_{L_{t,x}^{2}} &\leq \varepsilon b_{k}^{(1)}(\sigma)2^{-\sigma k}, \ \sigma \in [\frac{1}{100},\frac{5}{4}] \\ \|P_{k}A_{t}\uparrow_{s=0}\|_{L_{t,x}^{2}} &\leq \varepsilon^{2}, \ \sigma \in [0,\frac{5}{4}]. \end{split}$$

Remark 1.5. In Lemma 1.5, the $\{b_k^{(j)}\}, \{h_k^{(j)}\}, \{b_{k,s}^{(j)}\}\$ are defined as Def. 6.2.

Step 8. G_k bounds along the SMF direction for $\sigma \in [0, \frac{5}{4}]$.

Lemma 1.6. Assume that $\sigma \in [\frac{99}{100}, \frac{5}{4}]$. Let $\epsilon_0 > 0$ be a sufficiently small constant. Given any $\mathcal{L} \in \mathbb{Z}_+$, assume that $T \in (0, 2^{2\mathcal{L}}]$. Let $\{c_k(\sigma)\}$ be frequency envelopes of order $\frac{1}{2}\delta$. And let $\{c_k\}$ be an ϵ_0 -frequency envelope of order $\frac{1}{2}\delta$. Let $u \in \mathcal{H}_Q(T)$ be the solution to SMF with initial data u_0 which satisfies

$$\begin{aligned} \|P_k \nabla u_0\|_{L^2_x} &\leq c_k \\ \|P_k \nabla u_0\|_{L^2_x} &\leq c_k(\sigma) 2^{-\sigma k}. \end{aligned}$$

Denote $\{\phi_i\}$ the corresponding differential fields of the heat flow initiated from u. Then, when s = 0, given $\sigma \in [\frac{99}{100}, \frac{5}{4}]$, one has

$$\|P_k\phi_i\|_{s=0} \|_{G_k(T)} \lesssim \left(c_k(\sigma) + c_k(\sigma - \frac{3}{8})c_k(\frac{3}{8})\right) 2^{-\sigma k}$$

Step 9. F_k bounds of $P_k \widetilde{\mathcal{G}}^{(1)}$ and G_k bounds of ϕ_x along the SMF direction for $\sigma \in [0, 1 + \frac{j}{4}]$.

Lemma 1.7. Given $j \ge 2$, assume that $\sigma \in [0, 1 + \frac{j}{4}]$. Let $Q \in N$ be a fixed point and ϵ_0 be a sufficiently small constant depending on j. Given any $\mathcal{L} \in \mathbb{Z}_+$, assume that $T \in (0, 2^{2\mathcal{L}}]$. Let $u \in \mathcal{H}_Q(T)$ be the solution to SMF with initial data u_0 . Let $\{c_k^{(j)}(\sigma)\}$ be frequency envelopes defined in Def. 6.1. And assume that $\{c_k^{(j)}(0)\}$ is an ϵ_0 -frequency envelope with $0 < \epsilon_0 \ll 1$.

• Then, for $\sigma \in [0, 1 + \frac{j-1}{4}]$, we have

$$2^{k} \| P_{k} \widetilde{\mathcal{G}}^{(1)} \|_{F_{k}(T)} \le c_{k}^{(j)}(\sigma) 2^{-\sigma k} [(1 + 2^{2k + 2k_{0}})^{-20} \mathbf{1}_{k+k_{0} \ge 0} + \mathbf{1}_{k+k_{0} \le 0} 2^{\delta |k+k_{0}|}],$$

for any $s \in [2^{2k_0-1}, 2^{2k_0+1}), k_0, k \in \mathbb{Z}$.

• Denote $\{\phi_i\}$ the corresponding differential fields of the heat flow initiated from u. Then, for $\sigma \in [0, \frac{j}{4} + 1]$ there holds

$$\|P_k\phi_i\|_{s=0}\|_{G_k(T)} \lesssim c_k^{(j)}(\sigma) 2^{-\sigma k}$$

Step 10. Global regularity, global well-posedness and asymptotic behaviors.

As Step 9, proceeding the bootstrap-iteration scheme for *K* times gives bounds of $2^{\sigma k} ||P_k \phi_j||_{G_k}$ for $\sigma \in [0, \frac{K}{4} + 1]$. Then, transforming the bounds of $\{\phi_j\} \upharpoonright_{s=0}$ back to the solution *u* gives

$$||u||_{L_t^{\infty}\dot{H}_x^{\sigma+1}(T)} \leq ||u_0||_{\dot{H}_x^1 \cap \dot{H}_x^{\sigma+1}}$$

Noticing that an $\dot{H}^1 \cap \dot{H}^{2+}$ uniform bound will rule out blow-up for SMF in \mathbb{R}^2 , one step iteration suffices to show *u* is global. And the ϵ_0 depends only on the dimension and the target manifold \mathcal{N} . Moreover, we proceed the bootstrap-iteration scheme for *K* times and obtain uniform bounds for higher Sobolev norms.

The asymptotic behaviors stated in (1.5) will be proved following our recent work [29] on SMF on hyperbolic planes.

1.5 Main ideas

Let us explain the main ideas. To control the curvature terms, which is the non-selfclosed part, we use dynamical separation and a bootstrap-iteration scheme to obtain an approximate constant sectional curvature nonlinearity with controllable remainder in finite steps of iteration. The essential advantage of this scheme is that it reduces estimates in frequency localized spaces such as F_k , G_k to decay estimates in Lebesgue spaces along the heat direction.

Iteration scheme for Step 1. We describe the iteration scheme for heat flows. The starting point of heat flow iteration is bounds for $\partial_s v$. <u>1. First time iteration</u>. Suppose that we have obtained parabolic decay estimates of $||P_k \partial_s v||_{L^{\infty}L^2_v}$ such as

$$\|P_k \partial_s v\|_{L^{\infty}_t L^2_x} \lesssim (1 + 2^{2k} s)^{-M_1} \gamma_{k,s}(\sigma) 2^{-\sigma k}, \ \sigma \in [0, \frac{99}{100}].$$

By applying dynamical separation

$$S_{ij}^{l}(v(s)) = S_{ij}^{l}(Q) - \int_{s}^{\infty} (DS_{ij}^{l})(v(s')) \cdot \partial_{s}v ds'$$

bounds for $||P_k \partial_s v||_{L_t^{\infty} L_x^2}$ yield improved frequency localized bounds for the second fundamental form term, i.e. $||P_k S_{ij}^l(v(s))||_{L_t^{\infty} L_x^2}$. The only potential trouble to do this is the High × Low interaction of $(DS_{ij}^l)(v(s')) \cdot \partial_s v$. But we will see this interaction can be handled with additionally proving the decay estimates

$$\|\partial_x^{L+1} DS_{ij}^l(v(s))\|_{L^{\infty}_t L^2} \lesssim_L \epsilon_1 s^{-\frac{L}{2}}, \ \forall \ L \in \mathbb{N}.$$

Then back to the extrinsic map *v*, using the heat flow equation will give an improved bound for $||P_k \partial_s v||_{L_t^{\infty} L_x^2}$ for $\sigma \in [1, \frac{5}{4}]$. <u>2</u>. *m*-th time iteration. Using dynamical separation of the schematic form

$$D^{m-1}S(v(s)) = D^{m-1}S(Q) - \int_{s}^{\infty} D^{m}S(v(s'))\partial_{s}vds$$

and the decay estimates

$$\|\partial_x^{L+1} D^{m-1} S_{ii}^l(v(s))\|_{L^{\infty}_t L^2} \lesssim_{L,m} \epsilon_1 s^{-\frac{L}{2}}, \ \forall \ L \in \mathbb{N},$$

one gets frequency localized bounds of the extrinsic map v for $\sigma \in [1, 1 + \frac{m}{4}]$:

$$2^{\sigma k} \|P_k v(s)\|_{L^{\infty}_{t, k}L^2_{x}} \lesssim \epsilon_1 (1 + 2^{2k} s)^{-M_1 + m} \gamma_k^{(m)}(\sigma)$$

The motivation of decomposition in Step 2. To bound the curvature terms, by a kind of dynamical separation we have the decomposition of curvature terms denoted by G:

G = constant + one order terms + quadratic terms,

(see Step 2). We observe that to control \mathcal{G} in the F_k space, it suffices to prove parabolic decay estimates of $\mathcal{G}^{(j)}$. The same idea will be applied to bound the frames in frequency localized spaces. We remark that dynamical separation was previously used by [18, 36] to expose the implicit null structures. Here, we apply dynamical separation to do iterations. Besides using dynamical separation, in order to give a bound for connection coefficients which is the heart for bootstrap, we further decompose the curvature term into differential fields ϕ_i dominated terms and relatively smaller quadratic terms. By an appropriate bootstrap argument, bounding connection coefficients $\{A_j\}_{j=1}^2$ in the $F_k \cap S_k^{1/2}$ space reduces to derive parabolic decay estimates of covariant derivatives of curvatures $\mathcal{G}^{(j)}$ in the simpler $L_i^{\infty} C_k^2 \cap L^4$ spaces.

The motivation of adding (1.27) in Step 4.1. The key difficulty in bounding curvature involved terms is the High × Low \rightarrow High interaction of curvatures and differential fields or heat tension fields, i.e. the frequency of curvatures occupies the dominate position compared with differential fields or heat tension fields. First of all, we observe that it suffices to control the F_k norms of curvatures \mathcal{G} . Then we further clarify that among the four blocks of F_k space only the three blocks $L_t^{\infty} L^2$, L^4 , $L_x^4 L_t^{\infty}$ need to be estimated for curvatures. Second, we find that using dynamic separation in the heat direction

$$\mathcal{G} := \mathbf{R}(e_{i_0}, e_{i_1}, e_{i_2}, e_{i_3}) = \Gamma^{\infty} - \int_s^{\infty} \psi_s^l(\widetilde{\nabla}\mathbf{R})(e_l; e_{i_0}, e_{i_1}, e_{i_2}, e_{i_3})ds'$$
$$:= \Gamma^{\infty} - \Gamma_l^{\infty,(1)} \int_s^{\infty} \psi_s^l ds' - \int_s^{\infty} \psi_s^l \widetilde{\mathcal{G}}_l^{(1)} ds'$$

and the heat flow iteration scheme, the $L_t^{\infty}L^2$, L^4 norms of curvatures can be controlled by corresponding norms of differential fields.

The troublesome block is the $L_x^4 L_t^\infty$ norm. This norm of curvatures can not be obtained by dynamic separation in the heat direction and the heat flow iteration scheme as before. The problem is the High × Low \rightarrow High interaction of $\tilde{\mathcal{G}}_l^{(1)} \psi_s^l$ in $L_x^4 L_t^\infty$ fails if one only previously has bounds for $\tilde{\mathcal{G}}^{(1)}$ in $L_t^\infty L_x^2 \cap L^4$. (Estimates of $\tilde{\mathcal{G}}^{(1)}$ in $L_t^\infty L_x^2 \cap L^4$ are obtained in Step 3.) So we add a bootstrap assumption (1.27) to bound $L_x^4 L_t^\infty$ of $\tilde{\mathcal{G}}^{(1)}$. The key is that one can indeed improve the assumption (1.27) and then close bootstrap.

How to drop (1.27) in Step 4.2. Let us explain how to improve (1.27) and thus close bootstrap of Step 4. (I) With assumption (1.27) we can prove bounds of A_t and ϕ_t in L^4 by envelopes of differential fields ϕ_x , loosely speaking say

$$\begin{aligned} \|P_k(A_t)\|_{L^4} &\leq 2^{-\sigma k+k} h_k(\sigma) [(1+2^{2k+2k_0})^{-1} \mathbf{1}_{k+k_0 \ge 0} + \mathbf{1}_{k+k_0 \le 0} 2^{\delta |k+k_0|}] \\ \|P_k(\phi_t)\|_{L^4} &\leq 2^k b_k (1+2^{2k}s)^{-2} \end{aligned}$$

where $k, k_0 \in \mathbb{Z}, s \in [2^{2k_0-1}, 2^{2k_0+1}), \{h_k(\sigma)\}$ is the frequency envelope associated with differential fields $\{\phi_i\}_{i=1}^2$ defined in (1.26), and $\{b_k\}$ is some frequency envelope with $\|b_k\|_{\ell^2}$ sufficiently small. This will give bounds for $\|P_k\partial_t \widetilde{\mathcal{G}}^{(1)}\|_{L^4}$. (II) We improve the assumption (1.27) by interpolation

$$\|P_{k}\widetilde{\mathcal{G}}^{(1)}\|_{L_{x}^{4}L_{t}^{\infty}} \leq \|P_{k}\widetilde{\mathcal{G}}^{(1)}\|_{L^{4}}^{\frac{4}{3}} \|P_{k}\partial_{t}\widetilde{\mathcal{G}}^{(1)}\|_{L^{4}}^{\frac{1}{4}},$$
(1.28)

and the L^4 estimates obtained from Step 3 (Prop. 1.4 for heat flow iteration)

$$2^{k} \| P_{k} \widetilde{\mathcal{G}}^{(1)} \|_{L^{4}} \le h_{k} (1 + 2^{2k} s)^{-M}.$$

In fact, we can prove

$$2^{\frac{1}{2}k} \| P_k \widetilde{\mathcal{G}}^{(1)} \|_{L^4_x L^\infty_t} \le h_k 2^{-\sigma k} [(1+2^{2k+2k_0})^{-\frac{3}{4}M} \mathbf{1}_{k+k_0 \ge 0} + \mathbf{1}_{k+k_0 \le 0} 2^{\delta [k+k_0]}].$$

Then choosing sufficiently large M, one obtains better bounds of $\tilde{\mathcal{G}}^{(1)}$ than the assumption (1.27). This presents the way to bound curvatures in the block space $L_x^4 L_t^{\infty}$ of F_k .

In (I), the key step is to obtain bounds of connection coefficients, see Lemma 4.1. To prove Lemma 4.1, as mentioned before, we decompose the curvature term into differential fields ϕ_i dominated terms and the remained quadratic terms, see Step 2 of Lemma 4.1. The additional smallness gain yielded by the remained quadratic terms gives us the chance to use bootstrap argument to control connection coefficients.

Iteration in Step 6 to Step 9. With these new ideas and [4]'s framework, the range of $\sigma \in [0, \frac{99}{100}]$ can be reached before performing iteration for the SMF evolution. In order to reach larger σ , one combine the heat flow iteration with an SMF iteration. For the SMF iteration scheme, the key is to improve the estimate $||P_k \widetilde{\mathcal{G}}^{(1)}||_{L^4_* L^\infty}$ step by step to reach larger σ .

1.6 Idea for higher dimensions

Let us give a prevue of the higher dimensions. As d = 2, for higher dimensions in order to track the curvature terms \mathcal{G} in the F_k space along the heat flow direction, it suffices to control the one order covariant derivative of curvature term $\widetilde{\mathcal{G}}^{(1)}$ in the simpler Lebesgue spaces in the heat direction. Thus the parabolic decay estimates of moving frame dependent quantities for $d \ge 3$ should be established. The difficulty is to bound all geometric quantities in the fractional Sobolev spaces when d is odd. We solve this problem by using the geodesic parallel transpose and difference characterization of Besov spaces. The idea is that the difference characterization reduces bounding fractional Sobolev norms to bound difference of all these geometric quantities and their covariant derivatives in Lebesgue spaces. And the geodesic parallel transpose gives us the difference of geometric quantities at different points of the base manifold.

We divide the whole Theorem for $d \ge 2$ into two papers to make the main idea clear and avoid the paper being too long.

Notations Let $\mathbb{Z}_+ = \{1, 2, ...\}, \mathbb{N} = \{0, 1, 2, ...\}$. We apply the notation $X \leq Y$ whenever there exists some constant C > 0 so that $X \leq CY$. Similarly, we will use $X \sim Y$ if $X \leq Y \leq X$. We sometimes drop the integral variable in the integration if no confuse occurs. And we closely follow the notations of [4] for reader's convenience.

Let \mathcal{F} denote the Fourier transformation in \mathbb{R}^2 .

Let $\chi : \mathbb{R} \to [0, 1]$ be a given smooth even function which is supported in $\{z \in \mathbb{R} : |z| \le \frac{8}{5}\}$ and equals to 1 for $\{z \in \mathbb{R} : |z| \le \frac{5}{4}\}$. Define $\chi_k(z) = \chi(\frac{z}{2^k}) - \chi(\frac{z}{2^{k-1}}), k \in \mathbb{Z}$. The Littlewood-Paley projection operators with Fourier multiplier $\eta \mapsto \chi_k(|\eta|)$ are denoted by $P_k, k \in \mathbb{Z}$. For $I \subset R$, let $\chi_I = \sum_{i \in I} \chi_i(|\xi|)$. The low frequency cutoff operator with Fourier multiplier $\eta \mapsto \chi_{(-\infty,k]}(|\xi|)$ is denoted by $P_{\leq k}$. And the high frequency cutoff is defined by $P_{\geq k} = I - P_{\leq k}$. Given $\mathbf{e} \in \mathbb{S}^1$, $k \in \mathbb{Z}$, denote $P_{k,\mathbf{e}}$ the operator with Fourier multiplier $\xi \mapsto \chi_k(\xi \cdot \mathbf{e})$.

The Riemannian curvature tensor on N is denoted by **R**. The covariant derivative on N is denoted by $\overline{\nabla}$. And we denote ∇ the induced covariant derivative on u^*TN . The metric tensor of N is denoted by $\langle \cdot, \cdot \rangle$. Let E be a Riemannian manifold with connection ∇ , and \mathbb{T} be a (0, r) type tensor. For $k, r \in \mathbb{Z}_+$, we define the (0, r + k) type tensor $\nabla^k \mathbb{T}$ by

$$\nabla^{k}\mathbb{T}(X_{1},...,X_{k};Y_{1},...,Y_{r}):=(\nabla_{X_{k}}(\nabla^{k-1}\mathbb{T}))(X_{1},...,X_{k-1};Y_{1},...,Y_{r})$$

for arbitrary tangent vector fields $X_1, ..., X_k, Y_1, ..., Y_r$ on E.

Preliminaries 2

2.1 Linear estimates

The following is the main linear estimates established by [4].

Proposition 2.1 ([4]). Given $\mathcal{L} \in \mathbb{Z}_+$, assume that $T \in (0, 2^{2\mathcal{L}}]$. Then for every $u_0 \in L^2_x$ with frequency localized in I_k and every $F \in N_k(T)$, we have the inhomogeneous estimate: If u solves

$$\begin{cases} i\partial_t u + \Delta u = F, \\ u(0, x) = u_0(x), \end{cases}$$
(2.1)

then

$$\|u\|_{G_k(T)} \lesssim \|u_0\|_{L^2_x} + \|F\|_{N_k(T)}.$$
(2.2)

Recall the refined space for $F_k(T)$: Let $S_k^{\omega}(T)$ denote the normed space of functions in $L_k^2(T)$ for which

$$\|g\|_{S_k^{\omega}(T)} = 2^{k\omega} \left(\|g\|_{L_t^{\infty} L_x^{2\omega}} + \|g\|_{L_t^4 L_x^{p_{\omega}^*}} + 2^{-\frac{k}{2}} \|g\|_{L_x^{p_{\omega}^*} L_x^{\infty}} \right)$$
(2.3)

is finite, where the exponents 2_{ω} and p_{ω}^* are defined via

$$\frac{1}{2_{\omega}} - \frac{1}{2} = \frac{1}{p_{\omega}^*} - \frac{1}{4} = \frac{\omega}{2}.$$
(2.4)

The following lemma will be used widely.

Lemma 2.1 ([4]). For $f \in L^2_k(T)$, there hold

$$\|P_k f\|_{L^4} \le \|f\|_{F_k(T)} \tag{2.5}$$

- -

$$\|P_k f\|_{F_k(T)} \lesssim \|f\|_{L^2_x L^\infty_t} + \|f\|_{L^4}$$
(2.6)

$$\|P_k f\|_{L^2_x L^\infty_x} \le \|f\|_{S^{\frac{1}{2}}_k},\tag{2.7}$$

and

$$\|e^{s\Delta}g\|_{F_k(T)} \leq (1+s2^{2k})^{-20} \|g\|_{F_k(T)},$$
(2.8)

provided that the RHS is finite.

Frequency Envelopes 2.2

We recall the definition of envelopes introduced by Tao.

Definition 2.1. Let $\{a_k\}_{k \in \mathbb{Z}}$ be a positive sequence, we call it a frequency envelope if

$$\sum_{k \in \mathbb{Z}} a_k^2 < \infty, \text{ and } a_j \le 2^{\delta |l-j|} a_l, \ \forall j, l \in \mathbb{Z}.$$
(2.9)

We call the frequency envelope $\{a_k\}$ an ϵ -envelope if it additionally satisfies

$$\sum_{k\in\mathbb{Z}}a_k^2\leq\epsilon^2.$$

For any nonnegative sequence $\{a_i\} \in \ell^2$, we define its frequency envelope by

$$\tilde{a}_j := \sup_{j' \in \mathbb{Z}} a_{j'} 2^{-\delta|j-j'|}$$

And $\{\tilde{a}_i\}$ satisfies

$$|a_j| \leq \tilde{a}_j, \forall j \in \mathbb{Z}; \sum_{j \in \mathbb{Z}} \tilde{a}_j^2 \lesssim \sum_{j \in \mathbb{Z}} a_j^2.$$

Generally the δ in Definition 2.2 is not important if it has been fixed throughout the paper. But due to our iteration argument we shall introduce different δ in different steps of iterations. So we call $\{a_k\}$ satisfying (2.9) frequency envelope of order δ .

In this paper, each time we mention frequency envelopes, we will clearly state its order.

We recall the following two facts on envelopes: (a) If $d_k \le b_k$ for all $k \in \mathbb{Z}$ and $\{b_k\}$ is a frequency envelope of order $\delta > 0$ then $\widetilde{d}_k \le b_k$ for all $k \in \mathbb{Z}$ as well, where $\{\widetilde{d}_k\}$ denotes the envelope of $\{d_k\}$ of the same order $\delta > 0$:

$$\tilde{d}_k := \sup_{j \in \mathbb{Z}} d_j 2^{-\delta|k-j|}$$

(b) If $\{d_k\}$ is already an envelope of order $\delta > 0$ then $d_k = \tilde{d}_k$ for all $k \in \mathbb{Z}$. We recall the classic result obtained by [25, 38].

Lemma 2.2 ([25, 38]). Assume that $u \in \mathcal{H}_O(T)$ satisfies

$$\|\partial_x u\|_{L^\infty_x L^2_x} = \epsilon_1 \ll 1. \tag{2.10}$$

And let v(s, t, x) be the solution of heat flow (1.13) with initial data u(t, x). Then

$$\|\partial_x^{j+1}v\|_{L^\infty_t L^2_x} \lesssim s^{-\frac{1}{2}}\epsilon_1,\tag{2.11}$$

and the corresponding differential fields and connection coefficients satisfy

$$s^{\frac{j}{2}} \|\partial_x^j \phi_i\|_{L^{\infty}_t L^2_x} \lesssim \epsilon_1 \tag{2.12}$$

$$s^{\frac{1}{2}} \|\partial_x^J A_i\|_{L^{\infty}_t L^2_x} \lesssim \epsilon_1 \tag{2.13}$$

$$s^{\frac{j}{2}} \|\partial_x^j \phi_i\|_{L^{\infty}_t L^{\infty}_x} \lesssim \epsilon_1 \tag{2.14}$$

$$s^{\frac{j+1}{2}} \|\partial_x^j A_i\|_{L^\infty_t L^\infty_x} \lesssim \epsilon_1, \tag{2.15}$$

for all $s \in [0, \infty)$, i = 1, 2 and any nonnegative integer j.

3 Iteration for Heat Flows

3.1 Main results on the extrinsic map v solving HF

For $u \in \mathcal{H}_Q(T)$, define

$$\gamma_k(\sigma) = \sup_{k' \in \mathbb{Z}} 2^{-\delta|k-k'|} 2^{\sigma k'+k'} \|P_{k'}u\|_{L^\infty_t L^2_x}, \ \sigma \ge 0, \ \delta = \frac{1}{800}.$$
(3.1)

Denote $\{\gamma_k\}$ the frequency envelope for the energy norm, i.e.,

 $\gamma_k = \gamma_k(0).$

Thus

$$2^{k} \|P_{k}u\|_{L^{\infty}_{x}L^{2}_{x}} \leq 2^{-\sigma k} \gamma_{k}(\sigma), \ \forall \sigma \geq 0.$$

Before going ahead, we recall the extrinsic formulations of heat flows. Assume that the target manifold N is isometrically embedded into \mathbb{R}^N , then the heat flow equation can be formulated as

$$\partial_{s}v^{l} - \Delta v^{l} = \sum_{a=1}^{2} \sum_{i,j=1}^{N} S^{l}_{ij} \partial_{a}v^{i} \partial_{a}v^{j}, \ l = 1, ..., N,$$
(3.2)

where $S = \{S_{ij}^l\}$ denotes the second fundamental form of the embedding $\mathcal{N} \hookrightarrow \mathbb{R}^N$.

Recall that $\{\gamma_k(\sigma)\}\$ are defined to be the frequency envelopes of $u \in \mathcal{H}_Q(T)$. (See (3.1) and *u* **neednot** solve SMF.) The frequency localized estimates of the extrinsic map *v* solving (3.2) with initial data *u* are given below.

Proposition 3.1. Assume that $u \in \mathcal{H}_O(T)$ satisfies

$$\|\partial_x u\|_{L^{\infty}_x L^2_x} = \epsilon_1 \ll 1. \tag{3.3}$$

And let v(s, t, x) be the solution of heat flow (3.2) with initial data u(t, x). Then v satisfies

$$\sup_{s \ge 0} (1 + s2^{2k})^{31} 2^k \|P_k v\|_{L^{\infty}_t L^2_x} \le 2^{-\sigma k} \gamma_k(\sigma)$$

for all $\sigma \in [0, \frac{99}{100}]$, $k \in \mathbb{Z}$. Moreover, for any $\sigma \in [\frac{99}{100}, \frac{5}{4}]$, $k \in \mathbb{Z}$, we have

$$\sup_{s\geq 0} (1+s2^{2k})^{30} 2^{\sigma k+k} \|P_k v\|_{L^{\infty}_t L^2_x} \leq \gamma_k(\sigma) + \gamma_k(\sigma-\frac{3}{8})\gamma_k(\frac{3}{8}).$$

Remark 3.1. The power of $(1 + s2^{2k})$ in Proposition 3.1 can be chosen to be any $M \in \mathbb{Z}_+$ with additionally assuming that ϵ_1 is sufficiently small depending on M. See Proposition 3.4 below.

3.2 Before Iteration

Given an initial data $v_0 \in \mathcal{H}_Q$ with energy sufficiently small, by Lemma 2.2 the corresponding heat flow is global with (2.11) holding. Then combining this with local Cauchy theory in Sobolev spaces for heat flows yields the following lemma.

Lemma 3.1. Assume that $v_0 \in \mathcal{H}_Q$ has sufficiently small energy. And let v(s, x) be the solution of heat flow (3.2) with initial data v_0 . Given arbitrary $L \in \mathbb{Z}_+$, there exist constants $C_L > 0$, $C_s > 0$, such that for any $s \ge 0, 0 \le j \le L$,

$$\|\partial_x^{j+1}v(s)\|_{H^L_x} \le C_L(1+s)^{-\frac{1}{2}}; \|v(s)-Q\|_{L^2_x} \le C_s.$$

Proposition 3.2. Assume that $u \in \mathcal{H}_O(T)$ satisfies

$$\sum_{k \in \mathbb{Z}} 2^{2k} \|P_k u\|_{L_t^\infty L_x^2}^2 = \epsilon_1^2 \ll 1.$$
(3.4)

And let v(s, t, x) be the solution of heat flow (3.2) with initial data u(t, x). Then for $\sigma \in [0, \frac{99}{100}]$ and all $k \in \mathbb{Z}$, v satisfies

$$\sup_{s \in [0,\infty)} (1 + s2^{2k})^{31} 2^{k+\sigma k} \|P_k v\|_{L^{\infty}_t L^2_x} \leq \gamma_k(\sigma).$$
(3.5)

Proof. Since v converges to a fixed point $Q \in \mathcal{N}$ as $s \to \infty$, we put

$$S_{ij}^{l}(v) = S_{ij}^{l}(Q) + (S_{ij}^{l}(v) - S_{ij}^{l}(Q)).$$

The $\{S_{ij}^{l}(Q)\}$ part is constant and makes acceptable contribution to the final estimates by [Lemma 8.3,[4]]. Moreover, by Lemma 2.2, for all nonnegative integers *L* the remained part satisfies

$$\|\partial_x^{L+1}(S_{ij}^l(v) - S_{ij}^l(Q))\|_{L_t^{\infty}L_x^2} \lesssim_L s^{-\frac{L}{2}} \|\nabla u\|_{L_t^{\infty}L_x^2}.$$

Thus for all $j \in \mathbb{Z}_+$ we have the bound

$$(1+2^{2k}s)^{j} \left\| \partial_{x} \left[P_{k}(S_{ij}^{l}(v) - S_{ij}^{l}(Q)) \right] \right\|_{L_{t}^{\infty}L_{x}^{2}} \lesssim_{j} \epsilon_{1}.$$
(3.6)

Now, let's use [Lemma 8.3,[4]]'s arguments. Given $\sigma \in [0, \frac{99}{100}]$, define $B_{1,\sigma}(S)$ to be

$$B_{1,\sigma}(S) = \sup_{k \in \mathbb{Z}, s \in [0,S)} \gamma_k^{-1}(\sigma) (1 + s2^{2k})^{31} 2^{\sigma k} 2^k ||P_k v||_{L_t^{\infty} L_x^2}.$$

By Lemma 3.1 and the fact $\{\gamma_k(\sigma)\}\$ is a frequency envelope, $B_{1,\sigma}(S)$ is well-defined for $S \ge 0$ and continuous in S with $\lim_{S \to 0} B_{1,\sigma}(S) = 1.$ Then by trilinear Littlewood-Paley decomposition (see (8.2) in Lemma 8.1), we have

$$\begin{split} 2^{k} \| P_{k} S_{ij}^{l}(f) \partial_{a} f^{i} \partial_{a} f^{j} \|_{L_{t}^{\infty} L_{x}^{2}} \lesssim 2^{k} \sum_{k_{1} \leq k} \mu_{k_{1}} 2^{k_{1}} \mu_{k} + \sum_{k_{2} \geq k} 2^{2k} \mu_{k_{2}}^{2} + a_{k} (\sum_{k_{1} \leq k} 2^{k_{1}} \mu_{k_{1}})^{2} \\ &+ \sum_{k_{2} \geq k} 2^{2k} 2^{-k_{2}} a_{k_{2}} \mu_{k_{2}} \sum_{k_{1} \leq k_{2}} 2^{k_{1}} \mu_{k_{1}}, \end{split}$$

where $\{a_k\}, \{\mu_k\}$ denote

$$a_{k} := \sum_{|k-k'| \le 20} \sum_{l,i,j=1}^{N} \|\partial_{x} P_{k'}(S_{ij}^{l}(v))\|_{L_{t}^{\infty} L_{x}^{2}}; \ \mu_{k} := \sum_{l=1}^{N} \sum_{|k'-k| \le 20} 2^{k'} \|P_{k'} v^{l}\|_{L_{t}^{\infty} L_{x}^{2}}.$$
(3.7)

Then by definition of $B_1(S)$ and slow variation of envelopes, for $s \in [0, S], \sigma \in [0, \frac{99}{100}]$, we get

$$2^{k} \|P_{k}S_{ij}^{l}(f)\partial_{a}f^{i}\partial_{a}f^{j}\|_{L_{t}^{\infty}L_{x}^{2}}$$

$$\leq B_{1,\sigma}B_{1,0}(1+s2^{2k})^{-31}\gamma_{k}\sum_{k_{1}\leq k}2^{k_{1}-\sigma k_{1}+k}\gamma_{k_{1}}(\sigma) + B_{1,\sigma}B_{1,0}\sum_{k_{2}\geq k}2^{2k-\sigma k_{2}}(1+s2^{2k_{2}})^{-62}\gamma_{k_{2}}\gamma_{k_{2}}(\sigma)$$

$$+ B_{1,\sigma}B_{1,0}2^{k}a_{k}(\sum_{k_{1}\leq k}2^{k_{1}}(1+s2^{2k_{1}})^{-31}\gamma_{k_{1}})(\sum_{k_{1}\leq k}2^{k_{1}-\sigma k_{1}}(1+s2^{2k_{1}})^{-31}\gamma_{k_{1}}(\sigma))$$

$$+ B_{1,\sigma}B_{1,0}\sum_{k_{2}\geq k}2^{2k}(1+s2^{2k_{2}})^{-31}2^{-\sigma k_{2}}a_{k_{2}}\gamma_{k_{2}}(\sigma)\gamma_{k_{2}}$$

$$\leq B_{1,\sigma}B_{1,0}(1+s2^{2k})^{-62}2^{2k-\sigma k}\gamma_{k}\gamma_{k}(\sigma) + B_{1,\sigma}B_{1,0}\sum_{k_{2}\geq k}2^{2k-\sigma k_{2}}(1+s2^{2k_{2}})^{-31}\gamma_{k_{2}}\gamma_{k_{2}}(\sigma)$$

$$+ B_{1,\sigma}B_{1,0}\sum_{k_{2}\geq k}2^{-\sigma k_{2}}2^{2k}(1+s2^{2k_{2}})^{-31}a_{k_{2}}\gamma_{k_{2}}\gamma_{k_{2}}(\sigma) + a_{k}2^{-\sigma k}B_{1,\sigma}B_{1,0}2^{2k}\gamma_{k}\gamma_{k}(\sigma)$$

$$(3.9)$$

Applying (3.6) to $\{a_k\}$, (3.9) is further bounded by

$$(3.9) \leq 2^{-\sigma k} B_{1,\sigma} B_{1,0} 2^{2k} \sum_{k_2 \geq k} (1 + s 2^{2k_2})^{-31} \gamma_k \gamma_{k_2}(\sigma).$$

Therefore, for $s \ge 0$, we conclude that (3.8) is dominated by

$$2^{k} \| P_{k} S_{ij}^{l}(v) \partial_{a} v^{i} \partial_{a} v^{j} \|_{L_{t}^{\infty} L_{x}^{2}} \lesssim 2^{-\sigma k} 2^{2k} B_{1,\sigma} B_{1,0} \sum_{k_{2} \ge k} (1 + s 2^{2k_{2}})^{-31} \gamma_{k_{2}} \gamma_{k_{2}}(\sigma) .$$

Hence by Duhamel principle:

$$\begin{aligned} (1+s2^{2k})^{31}2^{k+\sigma k} \|P_k v\|_{L^{\infty}_t L^2_x} &\lesssim (1+s2^{2k})^{31} e^{-s2^{2k}} 2^{k+\sigma k} \|P_k u\|_{L^{\infty}_t L^2_x} \\ &+ B_{1,\sigma} B_{1,0} (1+s2^{2k})^M \int_0^s e^{-(s-\tau)2^{2k}} 2^{k+\sigma k} \|P_k S^l_{ij}(v) \partial_a v^i \partial_a v^j\|_{L^{\infty}_t L^2_x} d\tau, \end{aligned}$$

and the inequality

$$\int_0^s e^{-(s-\tau)\lambda} (1+\tau\lambda_1)^{-31} d\tau \leq s(1+\lambda s)^{-31} (1+\lambda_1 s)^{-1},$$
(3.10)

we get

$$\begin{aligned} (1+s2^{2k})^{31}2^{k+\sigma k} \|P_k v\|_{L^{\infty}_{t}L^{2}_{x}} &\lesssim \gamma_k(\sigma) + B_{1,\sigma} B_{1,0}(S) 2^{2k} s \sum_{k_2 \ge k} \gamma_{k_2} \gamma_{k_2}(\sigma) (1+2^{2k_2} s)^{-1} \\ &\lesssim \gamma_k(\sigma) + B_{1,\sigma}(S) B_{1,0}(S) \epsilon_1 \gamma_k(\sigma). \end{aligned}$$

Then $B_{1,0} \leq 1 + \epsilon_1 B_{1,0}^2$. By $B_{1,0}(0) \leq 1$ and ϵ_1 is sufficiently small, we have $B_{1,0}(S) \leq 1$ for all $S \geq 0$. Then using $B_{1,\sigma} \leq 1 + \epsilon_1 B_{1,0} B_{1,\sigma}$ and $B_{1,\sigma}(0) \leq 1$, we get $B_{1,\sigma}(S) \leq 1$ for any $\sigma \in [0, \frac{99}{100}]$ and any $S \geq 0$ provided that ϵ_1 is sufficiently small. Thus (3.5) has been proved.

Remark 3.2. The power of $(1 + s2^{2k})$ in Proposition 3.2 can be chosen to be any $M \in \mathbb{Z}_+$ with additionally assuming that ϵ_1 is sufficiently small depending on M. See Proposition 3.4 below.

3.3 First time iteration

We state the first time iteration in the the following Proposition.

Proposition 3.3. Assume that $u \in \mathcal{H}_Q(T)$ satisfies (3.4). And let v(s, t, x) be the solution of heat flow (3.2) with initial data u(t, x). Then for $\sigma \in [\frac{99}{100}, \frac{5}{4}]$ and any $k \in \mathbb{Z}$, v satisfies

$$\sup_{s \in [0,\infty)} (1 + s2^{2k})^{30} 2^{k+\sigma k} ||P_k v||_{L_t^\infty L_x^2} \lesssim \gamma_k(\sigma) + \gamma_k(\sigma - \frac{3}{8})\gamma_k(\frac{3}{8}).$$
(3.11)

Proof. The key point is to improve the bounds of $\{a_k\}$ defined by (3.7). For this, we use dynamic separation again. One has

$$S_{ij}^{l}(v)(s) = S_{ij}^{l}(Q) - \int_{s}^{\infty} (DS_{ij}^{l})(v) \cdot \partial_{s} v ds'.$$
(3.12)

By Proposition 3.2, for $\sigma \in [0, \frac{99}{100}]$ and any $k \in \mathbb{Z}$, we get

$$2^{k+\sigma k} \|P_k \Delta v\|_{L^{\infty}_t L^2_x} \lesssim (2^{2k}s+1)^{-31} 2^{2k} \gamma_k(\sigma)$$

And repeating the proof of Proposition 3.2 gives

$$\sum_{a=1,2} 2^{k+\sigma k} \|S_{ij}^{l}(\partial_a v^i, \partial_a v^j)\|_{L^{\infty}_{t}L^2_{x}} \lesssim 2^{2k} \sum_{k_1 \ge k} (2^{2k_1}s + 1)^{-31} \gamma_{k_1} \gamma_{k_1}(\sigma).$$

Thus, given $s \in [2^{2k_0-1}, 2^{2k_0+1})$, by the heat flow equation we get for all $k \in \mathbb{Z}$ and $\sigma \in [0, \frac{99}{100}]$

$$2^{k+\sigma k} \|P_k \partial_s v\|_{L^{\infty}_{t} L^2_{x}} \leq (2^{2k}s+1)^{-31} 2^{2k} \gamma_k(\sigma) + \sum_{k_1 \geq k} (2^{2k}s+1)^{-31} 2^{2k} \gamma_{k_1} \gamma_{k_1}(\sigma)$$

$$\leq (2^{2k}s+1)^{-31} 2^{2k} \gamma_k(\sigma) + 1_{k+k_0 \geq 0} (2^{2k}s+1)^{-31} 2^{2k} \gamma_k \gamma_k(\sigma)$$

$$+ 2^{2k} 1_{k+k_0 \leq 0} \sum_{k \leq l \leq -k_0} \gamma_l \gamma_l(\sigma).$$
(3.13)

Recall the bound

$$2^{k} \|P_{k}[(DS)(v)]\|_{L_{t}^{\infty}L_{x}^{2}} \lesssim \epsilon_{1} (2^{2k}s + 1)^{-j}$$
(3.14)

for all $j \in \mathbb{Z}_+$ and $k \in \mathbb{Z}$. Then for $s \in [2^{2k_0-1}, 2^{2k_0+1})$, repeating bilinear arguments, (3.12) shows that if $k + k_0 \ge 0$ then

$$\begin{split} \left\| P_{k}[S_{ij}^{l}(v)(s)] \right\|_{L_{t}^{\infty}L_{x}^{2}} &\lesssim \int_{s}^{\infty} \sum_{|k_{1}-k| \leq 4} \| P_{\leq k-4}(DS(v)) \|_{L_{t,x}^{\infty}} \| P_{k_{1}}\partial_{s}v \|_{L_{t}^{\infty}L_{x}^{2}} ds' \\ &+ \int_{s}^{\infty} 2^{k} \sum_{|k_{1}-k_{2}| \leq 8, k_{1}, k_{2} \geq k-4} \| P_{k_{2}}(DS(v)) \|_{L_{t}^{\infty}L_{x}^{2}} \| P_{k_{1}}\partial_{s}v \|_{L_{t}^{\infty}L_{x}^{2}} ds' \\ &+ \int_{s}^{\infty} \sum_{|k_{2}-k| \leq 4, k_{1} \leq k-4} 2^{k_{1}} \| P_{k_{2}}(DS(v)) \|_{L_{t}^{\infty}L_{x}^{2}} \| P_{k_{1}}\partial_{s}v \|_{L_{t}^{\infty}L_{x}^{2}} ds' \\ &\lesssim 2^{-\sigma k-k} (2^{2k+2k_{0}}+1)^{-31} 2^{2k+2k_{0}} \gamma_{k}(\sigma) \gamma_{k}, \end{split}$$
(3.15)

provided $\sigma \in [0, \frac{99}{100}]$, where we applied (3.14), (3.13) in the last line. Moreover, for any $\sigma \in [0, \frac{99}{100}]$, $k_0 \in \mathbb{Z}$, and $s \in [2^{2k_0-1}, 2^{2k_0+1}]$, in the case $k + k_0 \leq 0$ one has

$$\left\|P_k[S_{ij}^l(\nu)(s)]\right\|_{L_t^{\infty}L_x^2} \lesssim \sum_{k_0 \le j \le -k} 2^{-\sigma k} 2^{j+2\delta|k+j|} \gamma_k(\sigma) \gamma_k \lesssim 2^{-\sigma k-k} \gamma_k(\sigma) \gamma_k.$$
(3.16)

Thus (3.16), (3.15) yield the following bounds for $\{a_k\}$:

$$2^{\sigma k}a_k \leq (1+2^{2k}s)^{-30}\gamma_k(\sigma)\gamma_k, \tag{3.17}$$

provided that $\sigma \in [0, \frac{99}{100}]$. Now define the function $B_{2,\sigma}(S)$ for a given $\sigma \in [\frac{99}{100}, \frac{5}{4}]$ by

$$B_{2,\sigma}(S) = \sup_{k \in \mathbb{Z}, s \in [0,S)} \left(\gamma_k^{(1)}(\sigma) \right)^{-1} 2^{\sigma k} (1 + s 2^{2k})^{30} 2^k ||P_k v||_{L^{\infty}_t L^2_x},$$

where we denote

$$\gamma_k^{(1)}(\sigma) := \begin{cases} \gamma_k(\sigma), & \sigma \in [0, \frac{99}{100}] \\ \gamma_k(\sigma) + \gamma_k(\sigma - \frac{3}{8})\gamma_k(\frac{3}{8}), & \sigma \in (\frac{99}{100}, \frac{5}{4}] \end{cases}$$

Moreover, by Lemma 3.1 and the fact that $\{\gamma_k^{(1)}(\sigma)\}$ is a frequency envelope of order 2δ , it is clear that $B_{2,\sigma} : [0, \infty) \to \mathbb{R}^+$ is well-defined and continuous in $S \ge 0$ with $\lim_{S \to 0} B_{2,\sigma}(S) = 1$. Then by trilinear Littlewood-Paley decomposition (see (8.2) in Lemma 8.1), the definition of $B_{2,\sigma}$ and slow variation of envelopes, we get for $s \in [0, S], \sigma \in [\frac{99}{100}, \frac{5}{4}]$ that

$$2^{k} \left\| P_{k} [S_{ij}^{l}(v)\partial_{a}v^{i}\partial_{a}v^{j}] \right\|_{L_{t}^{\infty}L_{x}^{2}}$$

$$(3.18)$$

$$\lesssim B_{2,\sigma} B_{1,0} (1 + s2^{2k})^{-30} 2^{-\sigma k} \gamma_k^{(1)}(\sigma) \sum_{k_1 \le k} 2^{k_1 + k} \gamma_{k_1}$$
(3.19)

$$+ B_{2,\sigma}B_{1,0}\sum_{k_2 \ge k} 2^{2k-\sigma k_2} (1+s2^{2k_2})^{-60} \gamma_{k_2} \gamma_{k_2}^{(1)}(\sigma) + B_{1,0}B_{1,\frac{3}{8}}a_k (\sum_{k_1 \le k} 2^{k_1} (1+s2^{2k_1})^{-30} \gamma_{k_1}) (\sum_{k_1 \le k} 2^{k_1-\frac{3}{8}\sigma k_1} (1+s2^{2k_1})^{-30} \gamma_{k_1}(\frac{3}{8})) + B_{2,\sigma}B_{1,0}\sum_{k_2 \ge k} 2^{2k} (1+s2^{2k_2})^{-30} 2^{-\sigma k_2} a_{k_2} \gamma_{k_2}^{(1)}(\sigma) \gamma_{k_2} \lesssim B_{2,\sigma}B_{1,0}\sum_{k_2 \ge k} 2^{2k-\sigma k_2} (1+s2^{2k_2})^{-30} \gamma_{k_2} \gamma_{k_2}^{(1)}(\sigma) + B_{2,\sigma}B_{1,0}\sum_{k_2 \ge k} 2^{-\sigma k_2} 2^{2k} (1+s2^{2k_2})^{-30} a_{k_2} \gamma_{k_2} \gamma_{k_2}^{(1)}(\sigma) + a_k 2^{-\frac{3}{8}\sigma k} B_{1,0} B_{1,\frac{3}{8}} 2^{2k} \gamma_k \gamma_k(\frac{3}{8}).$$
(3.20)

Then applying the trivial bound (3.6) to the RHS of (3.20) except the last term and applying (3.17) to $\{a_k\}$ in the last term, we get for all $\sigma \in (\frac{99}{100}, \frac{5}{4}]$ that

$$\begin{split} 2^{k+\sigma k} \left\| P_k [S_{ij}^l(v) \partial_a v^i \partial_a v^j] \right\|_{L_t^\infty L_x^2} &\lesssim B_{1,0} B_{2,\sigma} 2^{2k} \sum_{k_2 \ge k} (1 + s 2^{2k_2})^{-30} \gamma_{k_2} \gamma_{k_2}^{(1)}(\sigma) \\ &+ B_{1,0} B_{1,\frac{3}{8}} 2^{2k} 2^{-\sigma k} (1 + s 2^{2k})^{-30} \gamma_k(\sigma - \frac{3}{8}) \gamma_k(\frac{3}{8}) \gamma_k 1_{k+k_0 \ge 0} \\ &+ B_{1,0} B_{1,\frac{3}{8}} 2^{2k} 2^{-\sigma k} 2^{2\delta |k+k_0|} \gamma_k(\sigma - \frac{3}{8}) \gamma_k(\frac{3}{8}) \gamma_k 1_{k+k_0 \le 0} \end{split}$$

if $s \in [2^{2k_0-1}, 2^{2k_0+1})$. Then using Duhamel principle, (3.10) and the following inequality

$$(1+2^{2k}s)^{30}e^{-2^{2k}s}\int_0^s e^{s'2^{2k}}(s'2^{2k})^{-\delta}\mathbf{1}_{s'\leq 2^{-2k}}ds' \leq 2^{-2k},$$

we obtain

$$2^{k+\sigma k} (1+2^{2k}s)^{30} \|P_kv\|_{L^{\infty}_t L^2_x} \lesssim (1+\epsilon_1 B_{1,0} B_{1,\frac{3}{8}} + \epsilon_1 B_{2,\sigma} B_{1,0}) \gamma_k^{(1)}(\sigma).$$

Since $B_{1,\tilde{\sigma}} \leq 1$ for $\tilde{\sigma} \in [0, \frac{99}{100}]$ has been proved in Proposition 3.2, we arrive at

$$B_{2,\sigma} \lesssim 1 + \epsilon_1 B_{2,\sigma}, \ \forall \sigma \in (\frac{99}{100}, \frac{5}{4}],$$

which shows $B_{2,\sigma} \leq 1$, thus finishing our proof.

We define the frequency envelope $\gamma_k^{(j)}(\sigma),\,j=0,1$ by :

$$\gamma_k^{(0)}(\sigma) = \gamma_k(\sigma), 0 \le \sigma < \frac{99}{100}$$
(3.21)

$$\gamma_k^{(1)}(\sigma) = \begin{cases} \gamma_k^{(0)}(\sigma), 0 \le \sigma \le \frac{99}{100} \\ \gamma_k(\sigma) + \gamma_k^{(0)}(\sigma - \frac{3}{8})\gamma_k(\frac{3}{8}), \frac{99}{100} < \sigma \le \frac{5}{4} \end{cases}$$
(3.22)

And the frequency envelopes $\gamma_k^{(j)}(\sigma), \ j \ge 2$ are defined by induction:

$$\gamma_{k}^{(j)}(\sigma) = \begin{cases} \gamma_{k}^{(j-1)}(\sigma), & 0 \le \sigma \le \frac{j+3}{4} \\ \gamma_{k}(\sigma) + \gamma_{k}^{(j-1)}(\sigma - \frac{3}{8})\gamma_{k}(\frac{3}{8}), & \frac{j+3}{4} < \sigma \le \frac{j+4}{4} \end{cases}$$
(3.23)

Define the sequence $\{\gamma_{k,s}^{(j)}(\sigma)\}_{k\in\mathbb{Z}}$ with $j\in\mathbb{N}$ by

$$\gamma_{k,s}^{(j)}(\sigma) = \begin{cases} 2^{k+k_0} \gamma_k^{(j)}(\sigma) \gamma_{-k_0}^{(j)}(0), & k+k_0 \ge 0\\ \sum_{l=k}^{-k_0} \gamma_l^{(j)}(\sigma) \gamma_l^{(j)}(0), & k+k_0 \le 0 \end{cases}$$
(3.24)

for $s \in [2^{2k_0-1}, 2^{2k_0+1}), k, k_0 \in \mathbb{Z}$.

We state the *j*-th time iteration in the the following proposition.

Proposition 3.4. (*j-th iteration*) Let $j \in \mathbb{N}$, $M \in \mathbb{Z}_+$. Assume that $u \in \mathcal{H}_Q(T)$ satisfies (3.4) with ϵ_1 sufficiently small depending on j + M. Let v(s, t, x) be the solution of heat flow (3.2) with initial data u(t, x). Then for $\sigma \in [0, 1 + \frac{j}{4}]$ and all $s \ge 0$, v satisfies

$$\sup_{s \in [0,\infty)} (1 + s2^{2k})^M 2^{k+\sigma k} \|P_k v\|_{L^{\infty}_t L^2_x} \leq \gamma_k^{(j)}(\sigma).$$
(3.25)

Proof. Define intervals $\{\mathbb{I}_l\}_{l=0}^{\infty}$ by

$$\mathbb{I}_0 := [0, \frac{99}{100}]; \ \mathbb{I}_1 = (\frac{99}{100}, \frac{5}{4}]; \ \mathbb{I}_l = (\frac{3+l}{4}, \frac{4+l}{4}], l \ge 2.$$

Given $K \in \mathbb{Z}_+$, $\sigma \in \mathbb{I}_l$, $l \in \mathbb{N}$, we denote

$$B_{l+1,\sigma,K}(S) := \sup_{s \in [0,S), k \in \mathbb{Z}} \frac{1}{\gamma_k^{(l)}(\sigma)} (1 + s2^{2k})^K 2^{k+\sigma k} \|P_k v\|_{L^{\infty}_t L^2_x}$$

And let

$$\mathbb{B}_{l+1,K}(S) := \sup_{\substack{\sigma \in \bigcup_{\ell \leq l} \mathbb{I}_{\ell}}} B_{l+1,\sigma,K}(S).$$

(In this notation, Proposition 3.2 and Proposition 3.3 have proved B_{2,30}(S) ≤ 1.) Moreover, the argument of Proposition 3.3 indeed shows:
(i) For all K₀ ≥ 2, j ∈ N, 0 ≤ a ≤ j + 1,

$$2^{k} \|P_{k}[D^{a}S(v)]\|_{L^{\infty}_{t}L^{2}_{x}} \leq C_{K_{0},j} \epsilon (1+2^{2k}s)^{-K_{0}-(j+1)}$$

(ii) For all $K_0 \ge 2, j \in \mathbb{N}$, if

$$\begin{cases} 2^{k} \left\| P_{k}[D^{j+1}S(v)] \right\|_{L_{v}^{\infty}L_{x}^{2}} \lesssim \epsilon (1+2^{2k}s)^{-K_{0}-j-1} \\ \| P_{k}v \|_{L_{v}^{\infty}L_{x}^{2}} \lesssim 2^{-\sigma k} \gamma_{k}^{(0)}(\sigma) (1+2^{2k}s)^{-K_{0}-j-1} \end{cases},$$

then

$$2^k \left\| P_k[D^j S(v)] \right\|_{L^\infty_t L^2_x} \lesssim 2^{-\sigma k} \gamma^{(0)}_k(\sigma) (1+2^{2k}s)^{-K_0-j}.$$

where the implicit constant in the conclusion is of the form $C(1 + C_1^2 + C_2^2)$ if we denote C_1, C_2 to be the implicit constants in the conditions of (ii). Here, *C* is universal and C_1, C_2 may depend on *j*, K_0 . (iii) For all $K_0 \ge 2, j \in \mathbb{N}, 0 \le a \le j + 1$, if

$$\left(\begin{array}{c} 2^{k} \left\| P_{k}[D^{a+1}S(v)] \right\|_{L_{t}^{c}L_{x}^{2}} \lesssim 2^{-\sigma k} \gamma_{k}^{(j-(a+1))}(\sigma)(1+2^{2k}s)^{-K_{0}-(a+1)}, \\ \left\| P_{k}v \right\|_{L_{t}^{c}L_{x}^{2}} \lesssim 2^{-\sigma k} \gamma_{k}^{(j-a)}(\sigma)(1+2^{2k}s)^{-K_{0}-(a+1)} \end{array} \right)$$

then

$$2^{k} \|P_{k}[D^{a}S(v)]\|_{L^{\infty}_{t}L^{2}_{x}} \lesssim 2^{-\sigma k} \gamma_{k}^{(j-a)}(\sigma)(1+2^{2k}s)^{-K_{0}-a},$$

where the implicit constant in the conclusion is of the form $C(1 + C_1^2 + C_2^2)$ if we denote C_1, C_2 to be the implicit constants in the conditions of (iii). Here, *C* is universal and C_1, C_2 may depend on *j*, K_0 . (iv) For any $K \ge 2$, $j \ge 1$, $1 \le a \le j + 1$, $\sigma \in \mathbb{I}_a$, if

$$2^{k} \|P_{k}[S(v)]\|_{L_{t}^{\infty}L_{x}^{2}} \leq C_{K} 2^{-\sigma k} \gamma_{k}^{(a-1)}(\sigma)(1+2^{2k}s)^{-K} \\ \|P_{k}v\|_{L_{x}^{\infty}L_{x}^{2}} \leq B_{a+1,\sigma,K} 2^{-\sigma k} \gamma_{k}^{(a)}(\sigma)(1+2^{2k}s)^{-K}$$

then for all $S \in [0, \infty)$ there holds

$$B_{a+1,\sigma,K}(S) \le C_*(1+\epsilon_1 \mathbb{B}_{a,K} B_{a+1,\sigma,K}(S) + C_K \sum_{l=1}^a \sup_{S \in [0,\infty)} \mathbb{B}_{l,K+l+1}^2(S)).$$

where C_* depends only on *d* and emerges from the Littlewood-Paley trilinear decomposition. Then our proposition follows by iteration. To be concrete, we make several remarks. First, in order to get the *M*-power decay in (3.25), it suffices to set $K_0 = M + 4$ and the top involved derivative order is $D^{j+1}S$. Second, let us describe the iteration in a clearer way: In the first step, one verifies

$$\sup_{S \in [0,\infty)} B_{1,K_0+j+1}(S) \le C_{K_0,j},\tag{3.26}$$

i.e. the second conditions in (ii). This was presented in Proposition 3.2. (We emphasize that in this step ϵ_1 shall be sufficiently small depending on $K_0 + j$.) In the second step, one verifies $\sup_{S \in [0,\infty)} B_{2,K_0+j}(S) \leq C_{K_0,j}$, and in the *a*-th step one verifies $\sup_{S \in [0,\infty)} B_{a,K_0+j+2-a}(S) \leq C_{K_0,j}$. This is presented as (iii) and (iv). Thus in the *j*-th step, we get (3.25).

3.4 Rough dynamical separation

Recall the notations $\psi_i^{\alpha} = \langle \partial_i v, e_{\alpha} \rangle$, $\psi_i^{\overline{\alpha}} = \langle \partial_i v, Je_{\alpha} \rangle$, $\alpha = 1, ..., n, i = 0, ..., 2, 3$, and $\phi_i^{\alpha} = \psi_i^{\alpha} + \sqrt{-1}\psi_i^{\overline{\alpha}}$. Here, i = 0 refers to the *t* variable and i = 3 refers to the *s* variable.

We aim to bound connection coefficients in the localized frequency spaces. As a preparation, we first derive a suitable form of connection coefficients. By definitions, we see

$$\begin{aligned} \mathbf{R}(\mathbf{E}\phi_i,\mathbf{E}\phi_s) &= \mathbf{R}\left((\mathfrak{R}\phi_i^{\alpha})e_{\alpha} + (\mathfrak{I}\phi_i^{\alpha})e_{\overline{\alpha}},(\mathfrak{R}\phi_s^{\alpha})e_{\beta} + (\mathfrak{I}\phi_s^{\alpha})e_{\overline{\beta}}\right) \\ &= (\phi_i^{\alpha} \wedge \phi_s^{\beta})\mathbf{R}(e_{\alpha},e_{\overline{\beta}}) + (\phi_i^{\alpha} \cdot \phi_s^{\beta})\mathbf{R}(e_{\alpha},e_{\beta}). \end{aligned}$$

where we denote $z_1 \wedge z_2 = -\Im(z_1\overline{z_2}), z_1 \cdot z_2 = \Re z_1 \Re z_2 + \Im z_1 \Im z_2$ for complex numbers z_1, z_2 . Thus schematically under the frame $\mathbf{E} = \{e_\alpha, e_{\overline{\alpha}}\}_{\alpha=1}^n$ we can write

$$\begin{cases} (\Re[A_i])^{\gamma}_{\theta} = \sum \int_{s}^{\infty} (\phi_i^{\alpha} \diamond \phi_s^{\beta}) \langle \mathbf{R}(e_{\alpha}, e_{\beta,\overline{\beta}})(e_{\gamma}), e_{\theta} \rangle ds', \\ (\Im[A_i])^{\gamma}_{\overline{\theta}} = \sum \int_{s}^{\infty} (\phi_i^{\alpha} \diamond \phi_s^{\beta}) \langle \mathbf{R}(e_{\alpha}, e_{\beta,\overline{\beta}})(e_{\gamma}), e_{\overline{\theta}} \rangle ds', \end{cases}$$
(3.27)

where $\diamond = `` \land "$ when $e_{\beta,\overline{\beta}} = e_{\overline{\beta}}$, and $\diamond = `` \cdot "$ when $e_{\beta,\overline{\beta}} = e_{\beta}$. For simplicity, we schematically write

$$A_i(s) = \int_s^\infty (\phi_i \diamond \phi_s) \langle \mathbf{R}(e_{j_0}, e_{j_1})(e_{j_2}), e_{j_3} \rangle ds',$$

where $\{j_c\}_{c=0}^3$ run in $\{1, ..., 2n\}$, and *i* runs in $\{0, 1, 2\}$. Recall also that $\phi_s = \sum_{l=1}^2 D_l \phi_l$.

With abuse of notations, denote

$$\mathcal{G}(s) = \langle \mathbf{R}(e_{j_0}, e_{j_1})e_{j_2}, e_{j_3} \rangle(s).$$
(3.28)

for any given $j_0, ..., j_3 \in \{1, 2, ..., 2n\}$. We expand \mathcal{G} as

$$\langle \mathbf{R}(e_{j_0}, e_{j_1})e_{j_2}, e_{j_3} \rangle(s) = \lim_{s \to \infty} \langle \mathbf{R}(e_{j_0}, e_{j_1})e_{j_2}, e_{j_3} \rangle - \int_s^\infty \partial_s \langle \mathbf{R}(e_{j_0}, e_{j_1})e_{j_2}, e_{j_3} \rangle ds'$$
$$= \Gamma^\infty - \int_s^\infty \psi_s^l(\widetilde{\nabla}\mathbf{R})(e_l; e_{j_0}, ..., e_{j_3})ds',$$

where Γ^{∞} denotes the limit part which is constant, and we used the identity $\nabla_s e_p = 0$ for all p = 1, ..., 2n in the last line. Here, we view **R** as a type (0, 4) tensor.

With the above notations, we write

$$A_i(s) = \sum_{j_0, j_1, j_2, j_3} \int_s^\infty (\phi_i \diamond \phi_s) \mathcal{G} ds',$$
(3.29)

and \mathcal{G} is decomposed as

$$\mathcal{G} = \Gamma^{\infty} - \int_{s}^{\infty} \psi_{s}^{l}(\widetilde{\nabla}\mathbf{R})(e_{l}; e_{j_{0}}, ..., e_{j_{3}})ds'.$$

Of course, one can perform this separation for any time as desired. Denote

$$\mathcal{G}^{(j)} = (\overline{\nabla}^{j} \mathbf{R})(\underbrace{e, ..., e}_{j}; e_{j_0}, ..., e_{j_3})$$

and

$$\Gamma^{\infty,(j)} = \lim_{s \to \infty} \mathcal{G}^{(j)}(s).$$

Then we can schematically write

$$\mathcal{G} = \Gamma^{\infty} - \int_{s}^{\infty} \psi_{s}(s_{1}) ds_{1} \left(\Gamma^{\infty,(1)} - \int_{s_{1}}^{\infty} \psi_{s}(s_{2}) ds_{2}(\Gamma^{\infty,(2)} + \dots) \right)$$

For simplicity we also denote

$$\widetilde{\mathcal{G}} = \mathcal{G} - \Gamma^{\infty}, \ \widetilde{\mathcal{G}}^{(j)} = \mathcal{G}^{(j)} - \Gamma^{\infty,(j)}.$$

3.5 Intrinsic v.s. Extrinsic formulations in localized frequency pieces

Proposition 3.5. Let d = 2. And let $u \in \mathcal{H}_Q(T)$ satisfy

$$\|\partial_x u\|_{L^\infty_t L^2_x} = \epsilon_1 \ll 1. \tag{3.30}$$

Here, we do not require u to solve SMF. Denote v(s, t, x) *the solution to heat flow with data* u(t, x)*, and denote* $\{\phi_i\}$ *the corresponding differential fields under the caloric gauge. Assume that* $\{\eta_k(\sigma)\}$ *is a frequency envelope of order* δ *such that for all* $i = 1, 2, k \in \mathbb{Z}$ *,*

$$2^{\sigma k} \|\phi_i(\uparrow_{s=0})\|_{L^{\infty}_t L^2_x} \le \eta_k(\sigma).$$
(3.31)

Then we have

$$\gamma_k(\sigma) \leq \eta_k(\sigma) \tag{3.32}$$

$$(1+s2^{2k})^{30}2^{\sigma k}\|P_kA_i\|_{L^{\infty}_tL^2_x} \lesssim \eta^{(0)}_{k,s}(\sigma).$$
(3.33)

for any $\sigma \in [0, \frac{99}{100}]$, $k \in \mathbb{Z}$. Furthermore, assume that for $\sigma \in [0, \frac{5}{4}]$, $\{\eta_k(\sigma)\}$ is a frequency envelope of order $\frac{1}{2}\delta$ such that for all $i = 1, 2, k \in \mathbb{Z}$, (3.31) holds. Then for any $\sigma \in [0, \frac{5}{4}]$, $k \in \mathbb{Z}$, one has

$$\gamma_k^{(1)}(\sigma) \leq \eta_k^{(1)}(\sigma) \tag{3.34}$$

$$(1+s2^{2k})^{29}2^{\sigma k}||P_kA_i||_{L^{\infty}_tL^2_x} \leq \eta^{(1)}_{k,s}(\sigma).$$
(3.35)

Proof. Step 1.1. $\sigma \in [0, \frac{99}{100}]$. Let $\mathcal{P} : \mathcal{N} \to \mathbb{R}^N$ be the isometric embedding. By definition, we see

$$\partial_{i}v = \sum_{l=1}^{2n} \psi^{l} d\mathcal{P}(e_{l}) = \sum_{l=1}^{2n} \psi^{l}_{i} \chi^{\infty}_{l} + \sum_{l=1}^{2n} \psi^{l}_{i} \left(d\mathcal{P}(e_{l}) - \chi^{\infty}_{l} \right),$$
(3.36)

where $\{\chi_l^{\infty}\}$ are the corresponding limit of $(d\mathcal{P})(e_l)$ as $s \to \infty$ which are constant vectors belonging to \mathbb{R}^N . Denote

$$\omega_k(s) = \sum_{|k-k'| \le 20} \|P_{k'}\psi_i(s)\|_{L_t^\infty L_x^2}, \ \nu_k(s) = \sum_{|k-k'| \le 20} 2^{k'} \|P_{k'}\left(d\mathcal{P}(e_l) - \chi_l^\infty\right)\|_{L_t^\infty L_x^2}.$$
(3.37)

Then we see by Lemma 2.2 that

$$\|\{\nu_k\}\|_{\ell^2} \lesssim \|\partial_i \left(d\mathcal{P}(e_l) - \chi_l^{\infty} \right)\|_{L^2_x} \lesssim \|\partial_i \nu\|_{L^2_x} + \|A_i\|_{L^2_x} \lesssim \epsilon_1.$$

Moreover, direct calculations give the inequality

$$\|\partial_x^L \left((d\mathcal{P})(e_l) - \chi_l^{\infty} \right)\|_{L_t^{\infty} L_x^2} \lesssim \sum_{0 \le p,q \le L} \sum_{\mathcal{A}} |\partial_x^{\alpha_1} \phi_x|^{l_1} \dots |\partial_x^{\alpha_p} \phi_x|^{l_p} |\partial_x^{\beta_1} A|^{n_1} \dots |\partial_x^{\beta_q} A|^{n_q},$$
(3.38)

where \mathcal{A} is the set of nonnegative indexes $l_1, ..., n_q \in \mathbb{Z}$ and $(\alpha_1, ..., \beta_q) \in \mathbb{Z}^2 \times ... \times \mathbb{Z}^2$ which satisfy

$$l_1(|\alpha_1|+1) + \dots + l_p(|\alpha_p|+1) + n_1(|\beta_1|+1) + \dots + n_q(|\beta_q|+1) = L.$$

Suppose $l_1 \ge 1$, by Hölder and Lemma 2.2, we get

$$\begin{split} \|\partial_{x}^{L} \left((d\mathcal{P})(e_{l}) - \chi_{l}^{\infty} \right)\|_{L_{x}^{2}} \\ & \lesssim \epsilon_{1} \sum_{s} s^{-\frac{\alpha_{1}}{2}} s^{-(l_{1}-1)\frac{\alpha_{1}+1}{2}} s^{-\frac{l_{2}(\alpha_{2}+1)}{2} - \dots - \frac{l_{p}(\alpha_{p}+1)}{2}} s^{-\frac{(\beta_{1}|+1)n_{1}}{2} - \dots - \frac{(\beta_{q}|+1)n_{q}}{2}} \\ & \lesssim \epsilon_{1} s^{-\frac{L-1}{2}}. \end{split}$$

Suppose that $n_1 \ge 1$, then we also obtain the same bound as above. Thus we arrive at

$$\|\{(1+s2^{2k})^M \nu_k(s)\}\|_{\ell^2} \leq_M \epsilon_1, \tag{3.39}$$

for $M \in \mathbb{Z}_+$. Meanwhile, we see $\|(d\mathcal{P})(e_l) - \chi_l^{\infty}\|_{L^{\infty}} \leq 1$. Thus by [[4],(8.4)], we obtain

$$2^{k} \|P_{k}(\partial_{i}\nu)\|_{L_{t}^{\infty}L_{x}^{2}} \lesssim 2^{k}\omega_{k} + \nu_{k}\sum_{k_{1} \le k} \omega_{k_{1}}2^{k_{1}} + \sum_{k_{1} \ge k} 2^{-2|k_{1}-k|}\omega_{k_{1}}2^{k_{1}}\nu_{k_{1}}.$$
(3.40)

Since when s = 0, $\omega_k(0) \le 2^{-\sigma k} \eta_k(\sigma)$, by slow variation of envelopes one deduces

$$\begin{split} & 2^{k} \| P_{k}(\partial_{i} v) \|_{L_{t}^{\infty} L_{x}^{2}} \\ & \leq 2^{k} 2^{-\sigma k} \eta_{k}(\sigma) + \nu_{k} \sum_{k_{1} \leq k} 2^{k_{1} + \delta |k - k_{1}|} 2^{-\sigma k_{1}} \eta_{k_{1}}(\sigma) + \sum_{k_{1} \geq k} 2^{-2(k_{1} - k)} 2^{\delta |k_{1} - k|} 2^{k_{1}} \nu_{k_{1}} 2^{-\sigma k_{1}} \eta_{k_{1}}(\sigma) \\ & \leq 2^{k} 2^{-\sigma k} \eta_{k}(\sigma) (1 + \epsilon_{1}), \end{split}$$

for $\sigma \in [0, \frac{99}{100}]$, s = 0. Thus since $\{\eta_k(\sigma)\}$ is an envelope, by the definition of $\{\gamma_k(\sigma)\}$ we obtain

$$\gamma_k(\sigma) \leq \eta_k(\sigma).$$
 (3.41)

Hence, (3.32) has been done for $\sigma \in [0, \frac{99}{100}]$. **Step 1.2.** $\sigma \in (\frac{99}{100}, \frac{5}{4}]$. Recall $\mathcal{P} : \mathcal{N} \hookrightarrow \mathbb{R}^N$ is the given isometric embedding. Viewing $d\mathcal{P}$ as a section of $T^*\mathcal{N} \otimes T\mathbb{R}^N$, then the connection on \mathcal{N} induces a covariant derivative **D** on the bundle $T^*\mathcal{N} \otimes T\mathbb{R}^N$. We have the identity

$$d\mathcal{P}(e_l) - \chi_l^{\infty} = \int_s^{\infty} \psi_s^j \mathbf{D} d\mathcal{P}(e_j; e_l) ds'.$$
(3.42)

where we used the caloric condition $\nabla_s e_l = 0$ for all l = 1, ..., 2n. Similar to (3.39), direct calculations give

$$\|P_k(\mathbf{D}d\mathcal{P}(e_j;e_l))\|_{L^{\infty}_x L^2_x} \lesssim_M \epsilon_1 2^{-k} (1+s2^{2k})^{-M}$$
(3.43)

for any $M \in \mathbb{Z}_+, k \in \mathbb{Z}$.

By (3.13), we have the bound for $\partial_s v$:

$$2^{\sigma k+k} \|P_k(\partial_s v)\|_{L^{\infty}_t L^2_x} \lesssim 2^{2k} \left[(1+2^{2k}s)^{-31} \mathbf{1}_{k+k_0 \ge 0} \gamma_k(\sigma) + \sum_{k \le l \le -k_0} \gamma_l(\sigma) \gamma_l \right],$$
(3.44)

if $s \in [2^{2k_0-1}, 2^{2k_0+1}), k_0 \in \mathbb{Z}$. And using the identity $\psi_s^l = (d\mathcal{P}e_l) \cdot \partial_s v$, (3.39) and (3.44) instead yield

$$\|P_k\psi_s\|_{L^{\infty}_t L^2_x} \lesssim 2^{k-\sigma k} \left(\mathbf{1}_{k+k_0 \ge 0} (1+s2^{2k})^{-31} \gamma_k(\sigma) + \mathbf{1}_{k+k_0 \le 0} \sum_{k \le l \le -k_0} \gamma_l(\sigma) \gamma_l \right)$$
(3.45)

for all $k \in \mathbb{Z}$ and $\sigma \in [0, \frac{99}{100}]$, $s \in [2^{2k_0-1}, 2^{2k_0+1})$, $k_0 \in \mathbb{Z}$.

Then applying bilinear Littlewood-Paley decomposition to (3.42), (3.45) and (3.43) yield for any $\sigma \in [0, \frac{99}{100}]$

$$(1 + s2^{2k})^{30} 2^k \|P_k \left((d\mathcal{P})(e_l) - \chi_l^{\infty} \right)\|_{L_t^{\infty} L_x^2} \lesssim 2^{-\sigma k} \gamma_k(\sigma) \lesssim 2^{-\sigma k} \eta_k(\sigma),$$
(3.46)

where we applied (3.41) in the last inequality. Then by the identity (3.36), (3.46) and bilinear Littlewood-Paley decomposition, one has when s = 0

$$\begin{split} \|P_{k}\partial_{i}v\|_{L_{t}^{\infty}L_{x}^{2}} &\leq \|P_{k}\psi_{i}\|_{L_{t}^{\infty}L_{x}^{2}}\|P_{\leq k-4}[(d\mathcal{P})(e_{l}) - \chi_{l}^{\infty}]\|_{L^{\infty}} \\ &+ \|P_{k}[(d\mathcal{P})(e_{l}) - \chi_{l}^{\infty}]\|_{L_{t}^{\infty}L_{x}^{2}} \sum_{k_{1}\leq k-4} 2^{k_{1}}\|P_{k_{1}}\psi_{i}\|_{L_{t}^{\infty}L_{x}^{2}} \\ &+ 2^{k} \sum_{|k_{1}-k_{2}|\leq 8,k_{1},k_{2}\geq k-4} \|P_{k_{1}}\psi_{i}\|_{L_{t}^{\infty}L_{x}^{2}}\|P_{k}[(d\mathcal{P})(e_{l}) - \chi_{l}^{\infty}]\|_{L_{t}^{\infty}L_{x}^{2}} \\ &\leq 2^{-\sigma k}\eta_{k}(\sigma) + 2^{-\sigma k}\eta_{k}(\frac{3}{8})\eta_{k}(\sigma - \frac{3}{8}). \end{split}$$
(3.47)

Thus (3.47) gives $\gamma_k^{(1)}(\sigma) \leq \eta_k^{(1)}(\sigma)$ for $\sigma \in (\frac{99}{100}, \frac{5}{4}]$. Combining Step 1.1 and Step 1.2, we have proved (3.34) and (3.32). **Step 2.1. Bounds of connections for** $\sigma \in [0, \frac{99}{100}]$. Applying Proposition 3.1 gives

$$(1+2^{2k}s)^{31}2^{\sigma k+k} \|P_k v\|_{L^{\infty}_{t}L^2_{x}} \leq \gamma_k(\sigma)$$
(3.48)

for all $\sigma \in [0, \frac{99}{100}]$ and $s \ge 0$. Then using the identity $\psi_i^l = (d\mathcal{P}e_l) \cdot \partial_i v$ and the bound (3.39), by (3.48) we infer from the bilinear Littlewood-Paley decomposition that

$$2^{\sigma k} \|P_k \phi_i\|_{L^{\infty}_t L^2_x} \lesssim (1 + s 2^{2k})^{-31} \eta_k(\sigma)$$

for all $k \in \mathbb{Z}$ and $\sigma \in [0, \frac{99}{100}]$. Then, by (3.44), applying bilinear Littlewood-Paley decomposition again gives

$$\|P_k(\phi_s \diamond \phi_i)\|_{L^{\infty}_t L^2_x} \lesssim 2^{-\sigma k} (1 + 2^{2j+2k})^{-31} \left(2^{-j+k} \eta_{-j} \eta_k(\sigma) + 2^{-2j} \eta_{-j} \eta_{-j}(\sigma) \right)$$
(3.49)

for $j + k \ge 0$, $s \in [2^{2j-1}, 2^{2j+1})$, $\sigma \in [0, \frac{99}{100}]$. Recall that Section 3.4 shows A_i can be written in the form of

$$A_i(s) = \int_s^\infty (\phi_s \diamond \phi_i) \mathcal{G} ds'.$$
(3.50)

Direct calculations and Lemma 2.2 imply \mathcal{G} modulate a constant part Γ^{∞} satisfies

$$2^{k} \| P_{k} \widetilde{\mathcal{G}} \|_{L_{t}^{\infty} L_{x}^{2}} \lesssim_{M_{1}} (1 + s 2^{2k})^{-M_{1}} \epsilon_{1},$$
(3.51)

for all $M_1 \in \mathbb{Z}_+$, $k \in \mathbb{Z}$ and $s \ge 0$. Then applying bilinear Littlewood-Paley decomposition and (3.49) leads to

$$\int_{s}^{\infty} \|P_{k}((\phi_{i} \diamond \phi_{s})\mathcal{G})\|_{L_{t}^{\infty}L_{x}^{2}} ds' \lesssim 2^{-\sigma k} (1 + s2^{2k})^{-30} \eta_{k}(\sigma) \eta_{-j}$$

$$(3.52)$$

for all $j \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1})$ and $k + j \ge 0$. Moreover, one has

$$\int_{s}^{\infty} \|P_{k}((\phi_{i} \diamond \phi_{s})\mathcal{G})\|_{L_{t}^{\infty}L_{x}^{2}} ds' \leq 2^{-\sigma k} \sum_{k \leq l \leq -j} \eta_{l}(\sigma)\eta_{l}$$

for any $j \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1})$ and $k + j \leq 0$.

Hence, (3.33) is done. **Step 2.2.** $\sigma \in [\frac{99}{100}, \frac{5}{4}]$. Recall that (3.32), (3.34) has given

$$\gamma_k(\sigma) \lesssim \eta_k(\sigma), \ \gamma_k^{(1)}(\sigma) \lesssim \eta_k^{(1)}(\sigma)$$

Now, we are ready to estimate A_x for $\sigma \in [\frac{99}{100}, \frac{5}{4}]$. By the identity $\psi_i^l = (d\mathcal{P}e_l) \cdot \partial_i v$, the bound (3.46) and

$$|P_k \partial_i v||_{L^{\infty}_t L^2_x} \lesssim 2^{-\sigma k} (1 + s 2^{2k})^{-30} \eta_k^{(1)}(\sigma),$$

one obtains by bilinear Littlewood-Paley decomposition that

$$\|P_k\phi_i\|_{L^{\infty}_t L^2_x} \lesssim 2^{-\sigma k} (1+s2^{2k})^{-30} \eta_k^{(1)}(\sigma), \tag{3.53}$$

for any $\sigma \in [\frac{99}{100}, \frac{5}{4}]$, $k \in \mathbb{Z}$, $s \ge 0$. For any $\sigma \in [\frac{99}{100}, \frac{5}{4}]$, the proof of Proposition 3.3 yields the bound

$$\|P_k \partial_s v\|_{L^{\infty}_t L^2_x} \leq 2^{-\sigma k+k} \left[\mathbf{1}_{k+j \ge 0} (1+s2^{2k})^{-30} \eta_k^{(1)}(\sigma) + \mathbf{1}_{k+j \le 0} \sum_{k \le l \le -j} \eta_l^{(1)}(\sigma) \eta_l \right],$$

which combined with (3.46) instead gives

$$\|P_k\phi_s\|_{L_t^{\infty}L_x^2} \lesssim 2^{-\sigma k+k} \left[\mathbf{1}_{k+j\geq 0} (1+s2^{2k})^{-30} \eta_k^{(1)}(\sigma) + \mathbf{1}_{k+j\leq 0} \sum_{k\leq l\leq -j} \eta_l^{(1)}(\sigma) \eta_l \right],$$
(3.54)

for any $k \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1}), j \in \mathbb{Z}$, $\sigma \in [\frac{99}{100}, \frac{5}{4}]$.

In order to apply (3.50), we also need to improve the bound of $\widetilde{\mathcal{G}}$ stated in (3.51). Recall the formula

$$\mathcal{G} := \langle \mathbf{R}(e_{j_0}, e_{j_1})(e_{j_2}), e_{j_3} \rangle = \Gamma^{\infty} - \int_s^{\infty} \psi_s^p(\widetilde{\nabla} \mathbf{R})(e_p; e_{j_0}, ..., e_{j_3}) ds'.$$

By Lemma 2.2 and the direct calculations (see Step 1.1 for instance) we have the bounds:

$$2^{k} \| P_{k} \left((\widetilde{\nabla} \mathbf{R})(e_{l}; e_{j_{0}}, ..., e_{j_{3}}) - \Gamma_{l}^{\infty, (1)} \right) \|_{L_{t}^{\infty} L_{x}^{2}} \lesssim_{M} (1 + s2^{2k})^{-M}$$
(3.55)

for all $M \in \mathbb{Z}_+$, $k \in \mathbb{Z}$. Hence one obtains by (3.45) and bilinear Littlewood-Paley decomposition that

$$2^{k} \| P_{k} (\mathcal{G} - \Gamma^{\infty}) \|_{L_{t}^{\infty} L_{x}^{2}} \lesssim 2^{-\sigma k} (1 + s 2^{2k})^{-30} \gamma_{k}(\sigma)$$
(3.56)

for any $k \in \mathbb{Z}$, $\sigma \in [0, \frac{99}{100}]$.

Then by (3.56), (3.54), (3.53) using trilinear Littlewood-Paley decomposition as Step 2.1 with additional modifications in the Low × High interaction of $P_k((\phi_s \circ \phi_i)\widetilde{\mathcal{G}})$ (see Proposition 3.2 for instance) we conclude that

$$\int_{s}^{\infty} \|P_k\left((\phi_s \diamond \phi_i)\mathcal{G}\right)\|_{L^{\infty}_t L^2_x} ds' \lesssim 2^{-\sigma k} (1+s2^{2k})^{-29} \eta^{(1)}_{k,s}(\sigma)$$

for any $k \in \mathbb{Z}$, $\sigma \in (\frac{99}{100}, \frac{5}{4}]$. Thus, (3.35) is done.

Lemma 3.2. Let d = 2, $j, M \in \mathbb{Z}_+$. And let $u \in \mathcal{H}_Q(T)$, and v(s, t, x) be the solution of heat flow with data u(t, x). Denote $\{\phi_i\}$ the differential fields of v under the caloric gauge. Assume that $\sigma \in [0, 1 + \frac{j}{4}]$, and $\{\eta_k(\sigma)\}$ are frequency envelopes of order $\frac{1}{2^j}\delta$ such that for all $i = 1, 2, k \in \mathbb{Z}$,

$$2^{\sigma k} \|\phi_i(\uparrow_{s=0})\|_{L^{\infty}_{x}L^2_{x}} \le \eta_k(\sigma).$$

$$(3.57)$$

Then, given $j, M \in \mathbb{Z}_+$, there exists a sufficiently small constant ϵ_j depending only on M, j such that if $\|\nabla u\|_{L^{\infty}_t L^2_x} \leq \epsilon_j$, then we have

$$\gamma_k^{(j)}(\sigma) \leq \eta_k^{(j)}(\sigma) \tag{3.58}$$

for $\sigma \in [0, \frac{j}{4} + 1]$, $k \in \mathbb{Z}$. Moreover, for l = 0, ..., j, we have

$$(1+s2^{2k})^{M} \|P_{k}\phi_{i}\|_{L_{t}^{\infty}L_{x}^{2}} \lesssim 2^{-\sigma k} \eta_{k}^{(j)}(\sigma), \ i = 1, 2, \ \sigma \in [0, 1+j/4)$$
(3.59)

$$(1+s2^{2k})^{M} \|P_{k}[\widetilde{d\mathcal{P}}]^{(l)}\|_{L_{t}^{\infty}L_{x}^{2}} \lesssim 2^{-\sigma k} \eta_{k}^{(j-l)}(\sigma), \ \sigma \in [0, 1+(j-l)/4)$$
(3.60)

$$(1+s2^{2k})^{M} \| P_k \widetilde{\mathcal{G}}^{(l)} \|_{L^{\infty}_t L^2_x} \lesssim 2^{-\sigma k} \eta_k^{(j-l)}(\sigma), \ \sigma \in [0, 1+(j-l)/4)$$
(3.61)

$$(1+s2^{2k})^{M} \|P_{k}A_{i}\|_{L_{t}^{\infty}L_{x}^{2}} \lesssim 2^{-\sigma k} \eta_{k,s}^{(j)}(\sigma), \ \sigma \in [0,1+j/4)$$
(3.62)

where we denote $[d\mathcal{P}]^{(l)} = (\mathbf{D}^{l} d\mathcal{P})(\underbrace{e, ..., e}_{l}; e)$, and $[\widetilde{d\mathcal{P}}]^{(l)} = [d\mathcal{P}]^{(l)} - \lim_{s \to \infty} [d\mathcal{P}]^{(l)}$.

Proof. The case $\sigma \in [0, 5/4]$ has been done in Proposition 3.5. Let $\sigma \in [1 + j/4, 1 + (j + 1)/4]$. The general case of (3.58) follows by iteration. The highest covariant derivative order of \mathcal{G} and $d\mathcal{P}(e)$ one needs for the *j*-th iteration is *j*+1, and it suffices to take the decay power M + 2 + j, i.e.,

$$\begin{aligned} \left\| \partial_{x}^{L+1} \mathcal{G}^{(j+1)} \right\|_{L_{t}^{\infty} L_{x}^{2}} \lesssim_{L,j} \epsilon s^{-\frac{L}{2}}, \forall L \in [0, M+2+j] \\ \left\| \partial_{x}^{L+1} [d\mathcal{P}]^{(j+1)} \right\|_{L_{t}^{\infty} L_{x}^{2}} \lesssim_{L,j} \epsilon s^{-\frac{L}{2}}, \forall L \in [0, M+2+j], \end{aligned}$$

where we denote $\mathcal{G}^{(k)} = (\widetilde{\nabla}^k \mathbf{R})(\underbrace{e, ..., e}_{k}; \underbrace{e, ..., e}_{4})$, and $[d\mathcal{P}]^{(k)} = (\mathbf{D}^k d\mathcal{P})(\underbrace{e, ..., e}_{k}; e)$. These decay estimates are easy to check by

using Lemma 2.2. If these decay estimates along heat direction are verified, then (3.58) follows by repeating the arguments of Step 2 in Proposition 3.5 for *j* times. Moreover, the left (3.59), (3.60), (3.61), (3.62) follow along the same path by applying dynamical separation and iterations for *j* times. \Box

Similar to Proposition 3.5, one also have

Corollary 3.1. Let d = 2. And let $v_0 \in \mathcal{H}_O$ satisfy

$$\|\partial_x v_0\|_{L^2_x} = \epsilon_1 \ll 1.$$

• Let $\{d_k(\sigma)\}$ with $k \in \mathbb{Z}$, $\sigma \in [0, \frac{5}{4}]$ be frequency envelopes of order $\frac{1}{2}\delta$ satisfying

$$2^{\sigma k+k} \|P_k v_0\|_{L^2_x} \le d_k(\sigma). \tag{3.63}$$

Denote v(s, x) the solution to heat flow with data v_0 , and denote $\{\phi_i\}$ the corresponding differential fields under the caloric gauge. Then we have

$$\|\phi_i(\uparrow_{s=0})\|_{L^2_x} \le 2^{-\sigma k} d_k^{(1)}(\sigma).$$
(3.64)

• Let $j \in \mathbb{Z}_+$. Assume that $\{d_k(\sigma)\}$ with $k \in \mathbb{Z}$, $\sigma \in [0, 1 + \frac{1}{4}j]$ are frequency envelopes of order $\frac{1}{2^j}\delta$. Then for ϵ_1 sufficiently small depending only on j, similar results hold with $d_k^{(1)}(\sigma)$ replaced by $d_k^{(j)}(\sigma)$.

Proof. Using (3.63) Proposition 3.2 and Proposition 3.3 show for $\sigma \in [0, \frac{5}{4})$

$$(1 + s2^{2k})^{30} 2^{k + \sigma k} \|P_k v\|_{L^2_x} \leq d_k^{(1)}(\sigma).$$
(3.65)

Let's first consider $\sigma \in [0, \frac{99}{100}]$. Recall

$$2^{k} \|P_{k}(d\mathcal{P}(e_{l}) - \chi_{l}^{\infty})\|_{L^{2}_{x}} \leq \epsilon_{1}(1 + s2^{2k})^{-29}.$$
(3.66)

Then by the identity $\psi_i^l = d\mathcal{P}(e_l) \cdot \partial_i v$, (3.65), (3.66), we obtain from the bilinear Littlewood-Paley decomposition

$$\begin{split} \|P_{k}(d\mathcal{P}(e_{l}) \cdot \partial_{i}v)\|_{L^{2}_{x}} \\ &\lesssim 2^{-\sigma k}d_{k}(\sigma)\|P_{\leq k-4}d\mathcal{P}(e_{l})\|_{L^{\infty}} + 2^{k}\sum_{k_{1}\geq k-4, |k_{1}-k_{2}|\leq 8} 2^{-\sigma k_{1}}d_{k_{1}}(\sigma)\|P_{k_{2}}d\mathcal{P}(e_{l})\|_{L^{\infty}_{t}L^{2}_{x}} \\ &+ \sum_{|k-k_{2}|\leq 4}\|P_{k_{2}}(d\mathcal{P}(e_{l}))\|_{L^{2}_{x}}\sum_{k_{1}\leq k-4} 2^{k_{1}-\sigma k_{1}}d_{k_{1}}(\sigma) \end{split}$$

that for any $\sigma \in [0, \frac{99}{100}], k \in \mathbb{Z}$,

$$\|P_k\psi_i(\upharpoonright_{s=0})\|_{L^2_x} \leq 2^{-\sigma k} d_k(\sigma).$$

Using this $\sigma \in [0, \frac{99}{100}]$ bound and similar arguments as before one can improve (3.66) to

$$2^{k} \|P_{k}(d\mathcal{P}(e_{l}) - \chi_{l}^{\infty})\|_{L^{2}_{x}} \leq 2^{-\sigma k} d_{k}(\sigma)(1 + s2^{2k})^{-29}, \ \sigma \in [0, \frac{99}{100}],$$

and thus giving (3.64). The second item claimed in our corollary follows by similar arguments and Lemma 3.2.

Lemma 3.3. Let $u \in \mathcal{H}_Q(T)$ solve SMF. And let v(s, t, x) be the solution of heat flow (3.2) with initial data u(t, x). Then given $L \in \mathbb{Z}_+$, $L \ge 200$, for any $0 \le \sigma \le 2L$, there exist constants $\epsilon_L > 0$, $C_L > 0$, $C_{L,T}$, such that if $\|\partial_x u\|_{L^{\infty}_t L^2_x} \le \epsilon_L \ll 1$, then for any $s \ge 0$, $i = 1, 2, \rho = 0, 1, m = 0, 1, ..., L$,

$$\|\partial_t^{\rho}\partial_x^m(v-Q)\|_{L_t^{\infty}H_x^L} \le C_{L,T}(s+1)^{-\frac{m}{2}}$$
(3.67)

$$(2^{-\frac{1}{2}k}1_{k\le 0} + 2^{\sigma k}) \|P_k\phi_i\|_{L^\infty_v L^2_x} \le C_L (2^{2k}s + 1)^{-30}$$
(3.68)

$$(2^{-\frac{1}{2}k}1_{k\le 0} + 2^{\sigma k}) \|P_k A_i\|_{L^{\infty}_{t}L^2_x} \le C_L (2^{2k}s + 1)^{-28}$$
(3.69)

$$2^{mk} \|P_k \partial_t \phi_i\|_{L^{\infty}_t L^2_x} \le C_{L,T} (2^{2k} s + 1)^{-25}$$
(3.70)

$$2^{mk} \|P_k \partial_t A_i\|_{L^\infty_v L^2_v} \le C_{L.T} (2^{2k} s + 1)^{-25}.$$
(3.71)

Proof. Fix arbitrary $L \in \mathbb{N}$, $L \ge 200$. Let $\lambda_k(\sigma)$ be the frequency envelope

$$\lambda_k(\sigma) := \sup_{k' \in \mathbb{Z}} 2^{-\frac{1}{2^j} \delta |k-k'|} 2^{\sigma k'} \|P_{k'}(u-Q)\|_{L^{\infty}_t L^2_x}.$$

for $\sigma \in \mathbb{I}_i \cap [0, 2L]$. And define

$$\widetilde{B}_{j,\sigma,K}(S) = \sup_{k \in \mathbb{Z}, s \in [0,S)} [\lambda_k^{(j)}(\sigma)]^{-1} (1 + s2^{2k})^K 2^{\sigma k} \|P_k v\|_{L^\infty_t L^2_x},$$

for $\sigma \in \mathbb{I}_j \cap [0, 2L]$, $j \in \mathbb{N}$, $K \in \mathbb{Z}_+$. By Lemma 3.1 and the fact $\{\lambda_k^{(j)}(\sigma)\}$ are frequency envelopes, $\widetilde{B}_{j,\sigma,K}(S)$ is well-defined for $S \ge 0$ and continuous in S with $\lim_{S \to 0} \widetilde{B}_{j,\sigma,K}(S) = 1$. Then applying Proposition 3.2, Proposition 3.3, Proposition 3.4 and their arguments, we get $\widetilde{B}_{j,\sigma,30}(S) \le 1$. Then applying similar arguments of Corollary 3.1 one obtains

$$2^{\sigma k} \|P_k \phi_i(s=0)\|_{L^{\infty}_t L^2_x} \lesssim \lambda_k^{(j)}(\sigma)$$

for $\sigma \in \mathbb{I}_j \cap [0, 2L]$. Then similar arguments of Proposition 3.5 yield for any $\sigma \in [0, L]$

$$2^{\sigma k} \|P_k \phi_i\|_{L^{\infty}_t L^2_x} \lesssim C_L (2^{2k} s + 1)^{-30}$$

$$2^{\sigma k} \|P_k A_i\|_{L^{\infty}_x L^2_x} \lesssim C_L (2^{2k} s + 1)^{-28}$$

which verifies half of (3.68)-(3.69). Moreover, one has

$$2^{\sigma k} \|P_k(d\mathcal{P}e)\|_{L^{\infty}_t L^2_x} \leq 2^{-k} (1 + s2^{2k})^{-30}$$
(3.72)

for $\sigma \in [0, 2L]$. Let's check the left half of (3.68)-(3.69). Recall the bounds

$$\|P_k v\|_{L^2_x} \lesssim 2^{-\sigma k} \lambda_k^{(0)}(\sigma) (1 + s 2^{2k})^{-30}$$

for $\sigma \in [0, \frac{99}{100}]$. Then (3.66) and bilinear Littlewood-Paley decomposition show that

$$\begin{split} \|P_{k}(\phi_{i})\|_{L_{t}^{\infty}L_{x}^{2}} &= \sum_{l}^{2n} \|P_{k}(d\mathcal{P}(e_{l}) \cdot \partial_{i}\nu)\|_{L_{t}^{\infty}L_{x}^{2}} \\ &\lesssim 2^{k}\lambda_{k}^{(0)}(0)(1+s^{2k})^{-30}\|P_{\leq k-4}d\mathcal{P}(e_{l})\|_{L^{\infty}} + 2^{k}\sum_{k_{1}\geq k-4, |k_{1}-k_{2}|\leq 8} 2^{\frac{1}{2}k_{1}}\lambda_{k_{1}}^{(0)}(\frac{1}{2})\|P_{k_{2}}d\mathcal{P}(e_{l})\|_{L_{t}^{\infty}L_{x}^{2}} \\ &+ \sum_{|k-k_{2}|\leq 4} \|P_{k_{2}}(d\mathcal{P}(e_{l}))\|_{L_{x}^{2}}\sum_{k_{1}\leq k-4} 2^{2k_{1}}\lambda_{k_{1}}^{(0)}(0) \\ &\lesssim 2^{\frac{k}{2}}(1+s^{2k})^{-30} \end{split}$$

for any $i = 1, 2, k \le 0$. Similar arguments give

$$\|P_k(A_i)\|_{L^{\infty}_{u,L^2}} \lesssim 2^{\frac{k}{2}} (1+s2^{2k})^{-28}$$
(3.73)

for any $i = 1, 2, k \le 0$. Hence, (3.68) and (3.68) have been done.

Since $\partial_t v = J(v)(\sum_{i=1,2} D_i \phi)$ at s = 0, we observe from $u \in \mathcal{H}_Q(T)$ that

$$\|\partial_t v(\uparrow_{s=0})\|_{L^\infty_* H^l_x} \le C_{l,T}, \ \forall \ l \in \mathbb{N}.$$

Thus using the smoothing estimates of heat semigroups and applying ∂_t to (3.2), one obtains (3.67). From (3.67), (3.66), (3.72) and the identity

$$\partial_x^l \phi_t^a = \sum_{l_1=0}^l \partial_x^{l_1} (d\mathcal{P}(e_a)) \partial_x^{l-l_1} (\partial_t v)$$

we get

$$2^{mk} \|P_k \phi_t\|_{L^{\infty}_{t}L^2_{x}} \lesssim (1 + s2^{2k})^{-28} \ \forall \ 0 \le m \le L,$$

which further gives bounds of $||P_k A_t||_{L_t^{\infty} L_x^2}$. Then applying similar bounds of $A_i \phi_i$, $A_t \phi_i$ and the identity $\partial_t \phi_i = -A_t \phi_i + D_i \phi_t$, we obtain (3.70). For (3.71), we use $\phi_s = D_i \phi_i$ and

$$|\partial_x^l \partial_t A_i| \le \sum_{l_1=0}^l \int_s^\infty |D_x^{l-l_1} D_t \phi_i| |D_x^{l_1} \phi_s| ds' + \int_s^\infty |D_x^{l-l_1} \phi_i| |D_x^{l_1} D_t \phi_s| ds'.$$

3.6 Additional decay estimates for dynamical caloric gauge

Proposition 3.6. Let d = 2. And let $u \in \mathcal{H}_Q(T)$ be solution of SMF. Denote v(s, t, x) the solution to heat flow with data u(t, x), and denote $\{\phi_i\}_{i=0}^2$ the corresponding differential fields under the caloric gauge. Assume that $\{\beta_k(\sigma)\}$ is a frequency envelope of order δ such that for all $i = 1, 2, k \in \mathbb{Z}$,

$$2^{\sigma k} \|\phi_i(\uparrow_{s=0})\|_{L^{\infty}_t L^2_x \cap L^4_{t_x}} \le \beta_k(\sigma).$$
(3.74)

• There exists a sufficiently small constant $\epsilon > 0$ such that if

$$\sum_{k\in\mathbb{Z}} |\beta_k(0)|^2 < \epsilon, \tag{3.75}$$

then we have for any $l \in \mathbb{N}$

$$\|P_k \widetilde{\mathcal{G}}^{(l)}\|_{L^4 \cap L^{\infty}_t L^2_x} \lesssim_l 2^{-\sigma k - k} \beta_k(\sigma) (2 + s 2^{2k})^{-30}$$
(3.76)

$$\|P_k\phi_s\|_{L^4\cap L^\infty_t L^2_x} \lesssim 2^{-\sigma k+k} \left[\mathbf{1}_{k+j\ge 0} (1+s2^{2k})^{-30} \beta_k(\sigma) + \mathbf{1}_{k+j\le 0} \sum_{k\le l\le -j} \beta_l(\sigma) \beta_l \right]$$
(3.77)

$$(1+s2^{2k})^{29}2^{\sigma k+k} \|P_k(d\mathcal{P}(e))\|_{L^4} \leq \beta_k(\sigma)$$
(3.78)

for any $\sigma \in [0, \frac{99}{100}]$, $s \in [2^{2j-1}, 2^{2j+1})$, $j, k \in \mathbb{Z}$. Furthermore, assume that for $\sigma \in [0, \frac{5}{4}]$, $\{\beta_k(\sigma)\}$ is a frequency envelope of order $\frac{1}{2}\delta$ such that for all $i = 1, 2, k \in \mathbb{Z}$, (3.74), (3.75) hold. Then for any $\sigma \in [0, \frac{5}{4}]$, $k \in \mathbb{Z}$, one has

$$(1+s2^{2k})^{27}2^{\sigma k} \|P_k A_i\|_{L^{\infty}_t L^2_x} \leq \beta^{(1)}_{k,s}(\sigma).$$
(3.79)

• If $\{\beta_k(\sigma)\}$ is a frequency envelope of order $\frac{1}{2^j}\delta$, then similar results hold for $\sigma \in [0, \frac{1}{4}j + 1)$ and ϵ sufficiently small depending only on $j \in \mathbb{Z}_+$. (See Prop. 1.4 for instance)

Proof. By Proposition 3.5 and its proof, we have

$$(1 + s2^{2k})^{31} 2^{\sigma k} 2^{k} \|P_k v\|_{L^{\infty}_{x}L^2_{x}} \leq \beta_k(\sigma).$$
(3.80)

$$(1+s2^{2k})^{30}2^{\sigma k}2^{k}||P_{k}S(v)||_{L_{t}^{\omega}L_{x}^{2}} \leq \beta_{k}(\sigma).$$
(3.81)

$$\|P_k\phi_s\|_{L^{\infty}_t L^2_x} \lesssim 2^{-\sigma k+k} \left[\mathbf{1}_{k+j \ge 0} (1+s2^{2k})^{-30} \beta_k(\sigma) + \mathbf{1}_{k+j \le 0} \sum_{k \le l \le -j} \beta_l(\sigma) \beta_l \right]$$
(3.82)

$$(1 + s2^{2k})^{30}2^k \|P_k((d\mathcal{P})(e_l) - \chi_l^{\infty})\|_{L^2_x} \leq 2^{-\sigma k}\beta_k(\sigma)$$
(3.83)

$$\sum_{l=1,2} (1 + s2^{2k})^{29} 2^{\sigma k} \|P_k A_l\|_{L^{\infty}_t L^2_x} \leq \beta_{k,s}^{(0)}(\sigma).$$
(3.84)

for any $\sigma \in [0, \frac{99}{100}]$, $j, k \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1})$, if $\{\beta_k(\sigma)\}$ is a frequency envelope of order δ . And Proposition 3.5 and its proof also give similar results as above for $\sigma \in [0, \frac{5}{4}]$, if $\{\beta_k(\sigma)\}$ is a frequency envelope of order $\frac{1}{2}\delta$. **Step 1.** When s = 0, using $\partial_i v = \psi_i^l \mathcal{P}(e_l)$, we get from the bilinear Littlewood-Paley decomposition

$$\begin{aligned} \|P_{k}(fg)\|_{L^{4}} &\lesssim \sum_{|k-k_{2}| \leq 4} \|P_{\leq k-4}f\|_{L^{\infty}_{t,x}} \|P_{k_{2}}g\|_{L^{4}_{xL^{\infty}_{t}}} + 2^{k} \sum_{k_{1},k_{2} \geq k-4, |k_{1}-k_{2}| \leq 8} \|P_{k_{1}}f\|_{L^{\infty}_{t}L^{2}_{x}} \|P_{k_{2}}g\|_{L^{4}} \\ &+ \sum_{k_{2} \leq k-4, |k_{1}-k| \leq 4} 2^{\frac{1}{2}k_{1}} \|P_{k_{1}}f\|_{L^{\infty}_{t}L^{2}_{x}} 2^{\frac{1}{2}k_{2}} \|P_{k_{2}}g\|_{L^{4}} \end{aligned}$$
(3.85)

and (3.74), (3.83) that

$$\|P_k(\partial_i v)\|_{L^4} \lesssim 2^{-\sigma k} \beta_k(\sigma) \tag{3.86}$$

for s = 0 and any $\sigma \in [0, \frac{99}{100}]$, $k \in \mathbb{Z}$. Then we turn to study the heat flow equation (1.13) to obtain

$$(1 + s2^{2k})^{30} \|P_k(\partial_i v)\|_{L^4} \lesssim 2^{-\sigma k} \beta_k(\sigma)$$
(3.87)

for any $s \ge 0, \sigma \in [0, \frac{99}{100}], k \in \mathbb{Z}$. In fact, define

$$Z_1(S) := \sup_{s \in [0,S), k \in \mathbb{Z}} \frac{1}{\beta_k(\sigma)} 2^{\sigma k} (1 + s 2^{2k})^{30} ||P_k(\partial_i v)||_{L^4}$$

 $Z_1(S)$ is well-defined, continuous and tends to 1 as $S \rightarrow 0$ by (3.86). Using the following trilinear Littlewood-Paley decomposition

$$2^{k} \|P_{k}S(v)(\partial_{x}v,\partial_{x}v)\|_{L^{4}_{t,x}} \leq 2^{k}\widetilde{\beta}_{k}\sum_{k_{1}\leq k}\widetilde{\beta}_{k_{1}}2^{k_{1}} + \sum_{k_{2}\geq k}2^{-2|k-k_{2}|}2^{2k_{2}}\widetilde{\beta}_{k_{2}}^{2}$$
$$+ 2^{\frac{1}{2}k}\widetilde{\alpha}_{k}(\sum_{k_{1}\leq k}2^{k_{1}}\widetilde{\beta}_{k_{1}})^{2} + \sum_{k_{2}\geq k}2^{2k-k_{2}}\widetilde{\alpha}_{k_{2}}\widetilde{\beta}_{k_{2}}\sum_{k_{1}\leq k_{2}}2^{k_{1}}\widetilde{\beta}_{k_{1}}.$$

$$(3.88)$$

where we denote

$$\widetilde{\beta}_{k} = \sum_{|k'-k| \le 30} 2^{k'} ||P_{k'}v||_{L^{\infty}_{t}L^{2}_{x} \cap L^{4}_{t,x}}; \ \widetilde{\alpha}_{k} = \sum_{|k'-k| \le 30} 2^{k'} ||P_{k'}(S(v))||_{L^{\infty}_{t}L^{2}_{x}}$$

using similar arguments as Proposition 3.2 we obtain by (3.81), (3.80) that

$$Z_1(S) \leq 1 + \epsilon Z_1^2(S)$$

for any $S \ge 0$. Then $Z_1(S) \le 1$ since $\lim_{S \to 0} Z_1(S) = 1$. Thus (3.87) follows.

Step 2. With (3.87) in hand, using the heat flow equation one obtains bounds for $||P_k\partial_s v||_{L^4}$. Then the bound of $||P_k\phi_s||_{L^4}$ follows by the bilinear Littlewood-Paley decomposition (3.85) and (3.83). By performing dynamical separation for $d\mathcal{P}(e)$, we get bounds of $||P_k(d\mathcal{P}(e))||_{L^4}$ from $||P_k\phi_s||_{L^4}$ and $||P_k(\mathbf{D}d\mathcal{P}(e;e))||_{L^\infty_t L^2_x}$. And then one obtains bounds of $||P_k(\mathcal{G})||_{L^4}$, $||P_k\phi_s||_{L^4}$ for all $s \ge 0$, which further yields $||A_i||_{L^4}$ for any $s \ge 0$. See Proposition 3.5 for the details.

Outline of Proof before Iteration

One of the ingredients of proof before iteration is the framework due to [4]. The other main ingredient is the decomposition of curvatures mentioned in Step 2 of Section 1.4. And in the technic level we need bootstrap assumption of $||P_k \widetilde{\mathcal{G}}^{(1)}||_{L^4_x L^\infty_t(T)}$ and ideas to improve this bound, see Step 4.1 and Step 4.2 of Section 1.4 for instance.

We outline the framework of [4] for reader's convenience. Since the gauged equation is now not self-closed due to the curvature terms, several key new ideas as mentioned above shall be used. But to leave the reader a whole picture, we just sketch the framework of [4] rather than presenting all technically complex issues.

The proof is a bootstrap argument. Let $\sigma \in [0, \frac{99}{100})$ be given. Given $\mathcal{L} \in \mathbb{Z}_+$, $Q \in \mathcal{N}$, let $T \in (0, 2^{2\mathcal{L}}]$. Assume that $\{c_k\}$ is an ϵ_0 -frequency envelope of order δ and $\{c_k(\sigma)\}$ is another frequency envelope of order δ . Let u_0 be the initial data of SMF which satisfies

$$\|P_k \nabla u_0\|_{L^2_x} \le c_k(\widetilde{\sigma}) 2^{-\widetilde{\sigma}k}, \ \widetilde{\sigma} \in [0, \frac{99}{100}].$$
(3.89)

Denote u the solution to SMF with initial data u_0 . Assume that u satisfies

Bootstrap I.
$$||P_k \nabla u||_{L^{\infty}_t L^2_x} \leq \epsilon_0^{-\frac{1}{2}} c_k$$

Denote v(s, t, x) the solution of heat flow with initial data u(t, x), and A_i, A_t, A_s the corresponding connection coefficients. And denote the heat tension field by ϕ_s and the differential fields by $\{\phi_i\}$, ϕ_t respectively. Suppose that $\{\phi_i\}_{i=1}^2$ satisfy the bootstrap condition at s = 0:

Bootstrap II.
$$||P_k\phi_i||_{s=0} ||_{G_k(T)} \leq \epsilon_0^{-\frac{1}{2}} c_k$$

In Step 1, by studying the heat equations (1.15), (1.16), we prove **Bootstrap I,II** in fact give parabolic estimates for A_i, A_t and $\phi_{i,t}$ along the heat flow direction:

$$\begin{split} \|P_k\phi_i(s)\|_{F_k(T)} &\leq c_k(\sigma)2^{-\sigma k}(1+s2^{2k})^{-4}, \sigma \in [0, \frac{99}{100}]\\ \|P_k\phi_t(s)\|_{L^4_{t,x}} &\leq c_k(\sigma)2^{-\sigma k}(1+s2^{2k})^{-2}, \sigma \in [0, \frac{99}{100}].\\ \|P_kA_i \upharpoonright_{s=0}\|_{L^4_{t,x}} &\leq c_k(\sigma)2^{-\sigma k}, \sigma \in [0, \frac{99}{100}]\\ \|P_kA_t \upharpoonright_{s=0}\|_{L^2_{t,x}} &\leq \epsilon_0. \end{split}$$

In Step 2, by studying the Schrödinger equations (1.17), we prove **Bootstrap I**, **II** indeed yield improved estimates for ϕ_i along the Schrödinger flow direction:

$$\|P_k\phi_i\|_{s=0} \|_{G_k(T)} \leq c_k(\sigma) 2^{-\sigma k}, \ \sigma \in [0, \frac{99}{100}].$$

In Step 3, we prove

$$\|P_k\phi_i\|_{s=0} \|_{G_k(T)} \lesssim c_k \tag{3.90}$$

with Bootstrap I, II dropped.

4 Evolution of SMF solutions along the heat direction

4.1 Parabolic Estimates for differential fields

The main result of this section is the following Proposition.

Proposition 4.1. Let $\{b_k\}$ be a ε -frequency envelope. Assume that for i = 1, 2, there hold

$$\|P_k\phi_i\|_{s=0} \|_{F_k(T)} \le b_k(\sigma) 2^{-\sigma k}, \ \sigma \in [0, \frac{99}{100}]$$
(4.1)

$$\|P_k\phi_t\|_{s=0} \|_{L^4_{t,x}} \lesssim b_k(\sigma) 2^{-(\sigma-1)k}, \ \sigma \in [0, \frac{99}{100}].$$
(4.2)

and

$$\|P_k\phi_i(s)\|_{F_k(T)} \le \varepsilon^{-\frac{1}{2}} b_k (1 + s2^{2k})^{-4}.$$
(4.3)

Then if $\varepsilon > 0$ is sufficiently small, for $\sigma \in [0, \frac{99}{100}]$ one has

$$\|P_k\phi_i(s)\|_{F_k(T)} \leq b_k(\sigma)2^{-\sigma k}(1+s2^{2k})^{-4}$$
(4.4)

$$\|P_k A_i\|_{s=0} \|_{L^4_{t,x}} \lesssim b_k(\sigma) 2^{-\sigma k}, \ i = 1, 2.$$
(4.5)

$$\|P_k \phi_t(s)\|_{L^4_{t,x}} \lesssim b_k(\sigma) 2^{-(\sigma-1)k} (1+2^{2k}s)^{-2}$$

$$\|P_k A_t \upharpoonright_{s=0}\|_{L^2_{t,x}} \lesssim \varepsilon b_k(\sigma) 2^{-\sigma k}, \ \sigma \in [\frac{1}{100}, \frac{99}{100}]$$
(4.6)

$$\|P_k A_t \upharpoonright_{s=0}\|_{L^2_{t,x}} \lesssim \varepsilon^2, \ \sigma \in [0, \frac{99}{100}].$$

Remark. Assumption (4.3) can be dropped. It suffices to apply Sobolev embeddings, Lemma 3.3 and [[4], Page 1463]'s argument.

4.2 **Proof of Proposition 4.1**

Now we turn to prove the parabolic estimates in Proposition 4.1.

Denote

$$h(k) = \sup_{s \ge 0} (1 + s2^{2k})^4 \sum_{i=1}^2 \|P_k \phi_i(s)\|_{F_k(T)}.$$
(4.7)

Define the corresponding envelope by

$$h_k(\sigma) = \sup_{k' \in \mathbb{Z}} 2^{\sigma k'} 2^{-\delta |k'-k|} h(k').$$
(4.8)

Assume that

$$2^{\frac{1}{2}k} \| P_k(\widetilde{\mathcal{G}}^{(1)}) \|_{L^4_x L^\infty_t(T)} \le \varepsilon^{-\frac{1}{4}} h_k [(1+2^{2k}s)^{-9} \mathbf{1}_{j+k \ge 0} + \mathbf{1}_{j+k \le 0} 2^{\delta|k+j|}]$$
(4.9)

for any $s \in [2^{2j-1}, 2^{2j+1}), k, j \in \mathbb{Z}$.

Lemma 4.1. Under the assumptions of Proposition 4.1 and (4.9), for any $k \in \mathbb{Z}$, $s \ge 0$, i = 1, 2, we have

$$\|P_k(A_i(s))\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \lesssim 2^{-\sigma k} (1 + s 2^{2k})^{-4} h_{k,s}(\sigma),$$
(4.10)

where the sequences $\{h_{k,s}\}$ when $2^{2k_0-1} \leq s < 2^{2k_0+1}, k_0 \in \mathbb{Z}$ are defined by

$$h_{k,s}(\sigma) = \begin{cases} 2^{k+k_0} h_{-k_0} h_k(\sigma) \text{ if } k + k_0 \ge 0\\ \sum_{l=k}^{-k_0} h_l h_l(\sigma) \text{ if } k + k_0 \le 0. \end{cases}$$
(4.11)

Proof. By assumption (4.3) of Proposition 4.1 and noticing $\{b_k\}$ is a ε -envelope, we have

$$\|\{h_k\}\|_{\ell^2}^2 \le \varepsilon. \tag{4.12}$$

In order to prove (4.10), let B_1 denote the smallest number in $[1, \infty)$ for which it holds for all $\sigma \in [0, \frac{99}{100}]$, $s \ge 0, k \in \mathbb{Z}$, i = 1, 2,

$$\|P_k(A_i(s))\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \le B_1 2^{-\sigma k} (1 + s 2^{2k})^{-4} h_{k,s}(\sigma).$$
(4.13)

Recall that Section 3.4 shows A_i is schematically written as

$$A_{i}(s) = \sum_{j_{0}, j_{1}, j_{2}, j_{3}} \int_{s}^{\infty} (\phi_{i} \diamond \phi_{s}) \langle \mathbf{R}(e_{j_{0}}, e_{j_{1}})(e_{j_{2}}), e_{j_{3}} \rangle ds',$$
(4.14)

where $\{j_c\}_{c=0}^3$ run in $\{1, ..., 2n\}$, and *i* runs in $\{1, 2\}$. Recall also that $\phi_s = \sum_{l=1}^2 D_l \phi_l$. Applying P_k to (4.14) we have

$$\begin{split} \|P_{k}(A_{i}(s))\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}(T)} \\ &\leq \sum_{|k_{1}-k_{2}|\leq 8,k_{1},k_{2}\geq k-4} \int_{s}^{\infty} \left\|P_{k}[P_{k_{1}}(\phi_{i} \diamond \phi_{s}) P_{k_{2}}\langle \mathbf{R}(e_{j_{0}},e_{j_{1}})(e_{j_{2}}),e_{j_{3}}\rangle]\right\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}(T)} ds' \\ &+ \sum_{|k_{1}-k|\leq 4} \int_{s}^{\infty} \left\|P_{k}[P_{k_{1}}(\phi_{i} \diamond \phi_{s}) P_{\leq k-4}\langle \mathbf{R}(e_{j_{0}},e_{j_{1}})(e_{j_{2}}),e_{j_{3}}\rangle]\right\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}(T)} ds' \\ &+ \sum_{|k_{2}-k|\leq 4,k_{1}\leq k-4} \int_{s}^{\infty} \left\|P_{k}[P_{k_{1}}(\phi_{i} \diamond \phi_{s}) P_{k_{2}}\langle \mathbf{R}(e_{j_{0}},e_{j_{1}})(e_{j_{2}}),e_{j_{3}}\rangle]\right\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}(T)} ds'. \end{split}$$
(4.15)

The above three subcases according to their order are usually called (a) High \times High \rightarrow Low, (b) High \times Low \rightarrow High, (c) Low \times High \rightarrow High.

Case b. High× Low \rightarrow High. In [[4], Lemma 5.2, Page 1470], the authors have proved

$$\sum_{i=1}^{2} \int_{s}^{\infty} \left\| P_{k} \left(\phi_{i} \diamond \phi_{s} \right) \right\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} ds' \lesssim \varepsilon B_{1} 2^{-\sigma k} (1 + s 2^{2k})^{-4} h_{k,s}(\sigma).$$
(4.16)

with slightly different notations. Thus in the case (b), by (8.5) and applying the trivial bound

$$\|\langle \mathbf{R}(e_{j_0}, e_{j_1})(e_{j_2}), e_{j_3} \rangle\|_{L^{\infty}_{t,s,s}} \lesssim K(\mathcal{N}), \tag{4.17}$$

to the $P_{\leq k}$ part and (4.16) to the P_{k_1} part, we obtain that

$$\sum_{|k_1-k| \le 4} \int_{s}^{\infty} \left\| P_k \left(P_{k_1}(\phi_i \diamond \phi_s) P_{\le k-4} \langle \mathbf{R}(e_{j_0}, e_{j_1}) e_{j_2}, e_{j_3} \rangle \right) \right\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \\ \lesssim \varepsilon B_1 2^{-\sigma k} (1 + s 2^{2k})^{-4} h_{k,s}(\sigma).$$
(4.18)

Step 2. Refined Dynamic Separation. For the Low \times High and High \times High part, we need to further decompose the curvature term. The dynamic separation performed in Section 3.4 also needs to be refined. Recall the notation

$$\mathcal{G}(s) = \langle \mathbf{R}(e_{j_0}, e_{j_1})e_{j_2}, e_{j_3} \rangle(s),$$

for any given $j_0, ..., j_3 \in \{1, ..., 2n\}$, and the decomposition of \mathcal{G} in Section 3.4. Thus using $\psi_s = \sum_{i=1,2} (\partial_i + A_i) \psi_i$, after the second time dynamic separation \mathcal{G} can be decomposed into

$$\mathcal{G}(s) = \langle \mathbf{R}(e_{j_0}, e_{j_1})e_{j_2}, e_{j_3} \rangle (s)$$
(4.19)

$$=\Gamma^{\infty}-\Gamma_{l}^{\infty,(1)}\int_{s}^{\infty}\psi_{s}^{l}(\widetilde{s})d\widetilde{s}-\int_{s}^{\infty}\psi_{s}^{l}(\widetilde{s})\left(\int_{\widetilde{s}}^{\infty}\psi_{s}^{p}(s')(\widetilde{\nabla}^{2}\mathbf{R})(e_{l},e_{p};e_{j_{0}},...,e_{j_{3}})ds'\right)d\widetilde{s}.$$

$$(4.20)$$

$$:=\Gamma^{\infty}+\mathcal{U}_{00}+\mathcal{U}_{01}+\mathcal{U}_{I}+\mathcal{U}_{II}.$$

where we denote

$$\begin{aligned} \mathcal{U}_{00} &:= -\Gamma_l^{\infty,(1)} \int_s^{\infty} \sum_{i=1}^2 (\partial_i \psi_i) ds' \\ \mathcal{U}_{01} &:= -\int_s^{\infty} \sum_{i=1}^2 (\partial_i \psi_i)^l \left((\widetilde{\nabla} \mathbf{R}) (e_l; e_{j_0}, ..., e_{j_3}) - \Gamma_l^{\infty,(1)} \right) ds' \\ \mathcal{U}_I &:= -\Gamma_l^{\infty,(1)} \int_s^{\infty} \sum_{i=1}^2 (A_i \psi_i)^l ds' \\ \mathcal{U}_{II} &:= -\int_s^{\infty} \sum_{i=1}^2 (A_i \psi_i)^l (\widetilde{s}) \left(\int_{\widetilde{s}}^{\infty} \psi_s^p (s') (\widetilde{\nabla}^2 \mathbf{R}) (e_l, e_p; e_{j_0}, ..., e_{j_3}) ds' \right) d\widetilde{s} \\ &= -\int_s^{\infty} \sum_{i=1}^2 (A_i \psi_i)^l (\widetilde{s}) \left((\widetilde{\nabla} \mathbf{R}) (e_l; e_{j_0}, ..., e_{j_3}) - \Gamma_l^{\infty,(1)} \right) d\widetilde{s}. \end{aligned}$$

It is easy to prove

$$\left\| \int_{2^{2k_0-1}}^{\infty} \sum_{i=1}^{2} (\partial_i \psi_i) ds' \right\|_{F_k(T)} \le 2^{-\sigma k} h_k(\sigma) (1_{k_0+k \le 0} 2^{-k} + 1_{k_0+k \ge 0} 2^{2k_0+k}) (1 + 2^{2k_0+2k})^{-4}.$$
(4.21)

And recall that [[4], Lemma 5.2] has shown for $s \in [2^{2j-1}, 2^{2j+2})$ there holds

$$\|P_k(\phi_i \diamond \phi_s)(s)\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)}$$
(4.22)

$$\lesssim \begin{cases} 2^{-\sigma k} (1+2^{2k}s)^{-4} \left(\widetilde{h}_{k,s}(\sigma) + B_1 \varepsilon 2^{-2j} h_{k,s}(\sigma) \right), \text{ if } k+j \ge 0\\ 2^{-\sigma k} \left(\widetilde{h}_{k,s}(\sigma) + B_1 \varepsilon 2^{-2j} h_{-j} h_{-j}(\sigma) \right), \text{ if } k+j \le 0 \end{cases}$$

$$(4.23)$$

where $\tilde{h}_{k,s}(\sigma)$ is defined by

 $\widetilde{h}_{k,s}(\sigma) = 2^{-j} h_{-j} \left(2^k h_k(\sigma) + 2^{-j} h_{-j}(\sigma) \right).$

Then repeating the bilinear estimates [Lemma 5.1, [4]], we have

$$\int_{s}^{\infty} \|P_{k}(\mathcal{U}_{00}(\phi_{s} \diamond \phi_{i}))\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} ds' \lesssim (1 + \varepsilon B_{1})2^{-\sigma k}h_{k,s}(\sigma)(1 + 2^{2k_{0}+2k})^{-4}.$$
(4.24)

The Γ^{∞} part

$$\int_{s}^{\infty} \|P_{k}(\Gamma^{\infty}(\phi_{s} \diamond \phi_{i}))\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} ds' \leq (1 + \varepsilon B_{1})2^{-\sigma k}h_{k,s}(\sigma)(1 + 2^{2k_{0}+2k})^{-4}$$

follows by directly applying Lemma 5.2 of [[4]], since Γ^{∞} is just a constant.

Recall the notation $\widetilde{\mathcal{G}}^{(1)} = (\widetilde{\nabla} \mathbf{R})(e; e_{j_0}, e_{j_1}, e_{j_2}, e_{j_3}) - \Gamma^{\infty,(1)}$. For \mathcal{U}_{01} , applying (3.76) which says

$$2^{k} \| P_{k} \widetilde{\mathcal{G}}^{(1)} \|_{L^{\infty}_{t} L^{2}_{x}} \lesssim 2^{-\sigma k} h_{k}(\sigma) (1 + 2^{2k} s)^{-30}, \tag{4.25}$$

and (4.9) which says for $s \in [2^{2j-1}, 2^{2j+1})$

$$2^{\frac{k}{2}} \| P_k \widetilde{\mathcal{G}}^{(1)} \|_{L^4_x L^\infty_t} \lesssim 2^{-\sigma k} h_k(\sigma) [(1 + 2^{2k} s)^{-9} \mathbf{1}_{j+k \ge 0} + 2^{\delta|j+k|} \mathbf{1}_{k+j \le 0}],$$
(4.26)

we obtain by Lemma 4.2 that

$$\begin{split} \|P_{k}\left((\partial_{i}\psi_{i})\widehat{\mathcal{G}}^{(1)}\right)\|_{F_{k}(T)} \\ &\lesssim \sum_{|k_{1}-k|\leq 4} \|P_{k_{1}}\partial_{i}\phi_{i}\|_{F_{k_{1}}(T)}\|P_{\leq k-4}\widetilde{\mathcal{G}}^{(1)}\|_{L^{\infty}} + \sum_{|k_{2}-k|\leq 4,k_{1}\leq k-4} 2^{\frac{k_{1}}{2}}\|P_{k_{1}}\partial_{i}\phi_{i}\|_{F_{k_{1}}(T)}\|P_{k_{2}}\widetilde{\mathcal{G}}^{(1)}\|_{L^{4}_{x}L^{6}_{t}} \\ &+ \sum_{|k_{2}-k_{1}|\leq 8,k_{1},k_{2}\geq k-4} \|P_{k_{1}}\partial_{i}\phi_{i}\|_{F_{k_{1}}(T)}\|P_{k_{2}}\widetilde{\mathcal{G}}^{(1)}\|_{L^{\infty}} + 2^{\frac{1}{2}k_{1}}\|P_{k_{2}}\widetilde{\mathcal{G}}^{(1)}\|_{L^{4}_{x}L^{\infty}_{t}}). \end{split}$$

Thus by the slow variation of envelopes we further have

$$\|P_k\left((\partial_i\psi_i)\widetilde{\mathcal{G}}^{(1)}\right)\|_{F_k(T)} \lesssim 2^{-\sigma k} h_k(\sigma) \left(1_{k+j\geq 0} 2^k (1+2^{2k+2j})^{-4} + 1_{k+j\leq 0} 2^{-j} 2^{\delta|j+k|}\right)$$

for $s \in [2^{2j-1}, 2^{2j+1})$, $k, j \in \mathbb{Z}$. Notice that the large constant $\varepsilon^{-\frac{1}{4}}$ is absorbed by $||\{h_k\}||_{\ell^{\infty}} \leq \varepsilon^{\frac{1}{2}}$. And notice that in the Low × High interaction of $(\partial_i \psi_i) \widetilde{\mathcal{G}}^{(1)}$ it is possible to deduce $2^{-\sigma k}$ for $\sigma \in [0, \frac{99}{100}]$ from $\partial_i \psi_i$ due to fact that the series $\sum_{k_1 \leq k-4} 2^{2k-\sigma k} h_k(\sigma)$ is summable for $\sigma < 2$. Thus for $s \in [2^{2k_0-1}, 2^{2k_0+1})$ summing the above formula in $j \geq k_0$ yields

$$\int_{s}^{\infty} \|P_{k}\left((\partial_{i}\psi_{i})\widetilde{\mathcal{G}}^{(1)}\right)\|_{F_{k}(T)} ds' \leq 2^{-\sigma k} h_{k}(\sigma) \left(1_{k+k_{0} \geq 0} 2^{k+2k_{0}} (1+2^{2k+2k_{0}})^{-4} + 1_{k+k_{0} \leq 0} 2^{-k}\right)$$

which is the same as (4.21). Thus (4.22) and bilinear estimates give

$$\int_{s}^{\infty} \|P_{k}(\mathcal{U}_{01}(\phi_{s} \diamond \phi_{i}))\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} ds' \leq (1 + \varepsilon B_{1})2^{-\sigma k}h_{k,s}(\sigma)(1 + 2^{2k_{0}+2k})^{-4}.$$

Therefore, it remains to estimate $\mathcal{U}_{I}, \mathcal{U}_{II}$.

Step 3. Prove our lemma with a bootstrap condition. We first prove our lemma with additional bootstrap condition. And in the final step we will drop the bootstrap condition and finish the whole proof.

Bootstrap Assumption A. Assume that for all $k, j \in \mathbb{Z}, s \in [2^{2j-1}, 2^{2j+1})$, there holds

$$\|P_k \mathcal{U}_I\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \le \varepsilon^{-\frac{1}{2}} (1 + 2^{k+j})^{-7} T_{k,j} h_k c_0^*$$
(4.27)

$$\|P_k \mathcal{U}_{II}\|_{F_k(T)} \le \varepsilon^{-\frac{1}{2}} (1 + 2^{k+j})^{-7} T_{k,j} h_k c_0^*, \tag{4.28}$$

where $c_0^* := ||\{h_k\}||_{\ell^2}$ and we denote

$$T_{k,j} = \mathbf{1}_{k+j \le 0} 2^{-k} + \mathbf{1}_{k+j \ge 0} 2^j.$$
(4.29)

Our aim for this step is to prove B_1 defined by (4.15) satisfies

$$B_1 \lesssim 1 + \varepsilon B_1$$

assuming Bootstrap Assumption A.

For $s \in [2^{2j-1}, 2^{2j+1}), j \in \mathbb{Z}, (4.27)$ and (4.28) show $\mathcal{U} := \mathcal{U}_I + \mathcal{U}_{II}$ satisfies

$$2^{k}(1+2^{k+j})^{6} \|P_{k}\mathcal{U}\|_{F_{k}(T)} \leq 1.$$
(4.30)

Case a. High × High → Low. The bound (4.30) suffices to control the High × High interaction. For $k + k_0 \ge 0$, applying the bounds (4.30), (4.22) and (8.4) of Lemma 8.2 with $\omega = \frac{1}{2}$, one obtains that in High × High case

$$\sum_{j\geq k_{0}} \int_{2^{2j+1}}^{2^{2j+1}} \sum_{|k_{1}-k_{2}|\leq 8,k_{1},k_{2}\geq k-4} \left\| P_{k} \left(P_{k_{1}}(\phi_{i} \diamond \phi_{s}) P_{k_{2}}\mathcal{U} \right) \right\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}(T)} ds'$$

$$\lesssim 2^{-\sigma k} \sum_{j\geq k_{0}} \sum_{k_{1}\geq k-4} 2^{\frac{k-k_{1}}{2}} (1+2^{2k_{1}+2j})^{-6} h_{-j} \left(2^{k_{1}+j} h_{k_{1}}(\sigma) + h_{-j}(\sigma) \right)$$

$$+ 2^{-\sigma k} \sum_{j\geq k_{0}} \sum_{k_{1}\geq k-4} 2^{\frac{k-k_{1}}{2}} B_{1} \varepsilon (1+2^{2k_{1}+2j})^{-6} h_{k_{1},2^{2j}}(\sigma)$$

$$(4.31)$$

which by slow variation of envelopes is further bounded by

$$\begin{split} &2^{-\sigma k} \sum_{j \geq k_0} \sum_{k_1 \geq k-4} 2^{\frac{k-k_1}{2}} (1+2^{k_1+j})^{-10} 2^{\delta |j-k_0|} h_{-k_0} h_k(\sigma) \left(2^{k_1+j+\delta |k_1-k|} + 2^{\delta |k+j|} \right) \\ &+ 2^{-\sigma k} \sum_{j \geq k_0} \sum_{k_1 \geq k-4} 2^{\frac{k-k_1}{2}} B_1 \varepsilon (1+2^{k_1+j})^{-10} 2^{\delta |j-k_0|} 2^{\delta |k_1-k|} h_{-k_0} h_k(\sigma). \end{split}$$

Since $k_1 + j \ge k + j \ge k + k_0 \ge 0$, it is easy to see the above formula is acceptable by

$$\begin{split} &\sum_{j\geq k_0} 2^{-\sigma k} (1+\varepsilon B_1) h_{-k_0} h_k(\sigma) 2^{\delta |j-k_0|} 2^{\delta |k+j|} (1+2^{k+j})^{-8} \\ &\lesssim (1+\varepsilon B_1) 2^{-\sigma k} h_{-k_0} h_k(\sigma) 2^{k_0+k} (1+2^{k+k_0})^{-8}. \end{split}$$

Therefore, the $k + k_0 \ge 0$ case has been done for the High × High interaction.

Assume $k + k_0 \le 0$. Applying the bounds (4.22), (4.30) with $\sigma = 0$ and (8.4) of Lemma 8.2 with $\omega = \frac{1}{2}$, for $j + k \le 0$, by slow variation of envelopes we have

$$\begin{split} &\sum_{k_0 \leq j \leq -k} \int_{2^{2j+1}}^{2^{2j+1}} \sum_{|k_1-k_2| \leq 8, k_1, k_2 \geq k-4} \left\| P_k \left(P_{k_1}(\phi_i \diamond \phi_s) P_{k_2} \mathcal{U} \right) \right\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} ds' \\ &\lesssim \sum_{k_0 \leq j \leq -k} \sum_{k_1 \geq k-4} 2^{\frac{k-k_1}{2}} 2^{-\sigma k_1} (1 + 2^{2k_1 + 2j})^{-6} h_{-j} \left(2^{k_1 + j} h_{k_1}(\sigma) + h_{-j}(\sigma) \right) \\ &+ B_1 \varepsilon 2^{-\sigma k} \sum_{k_0 \leq j \leq -k} \left(\sum_{k-4 \leq k_1 \leq -j} 2^{\frac{k-k_1}{2}} h_{-j} h_{-j}(\sigma) + \sum_{k_1 \geq -j} 2^{\frac{k-k_1}{2}} (1 + 2^{2j+2k_1})^{-4} h_{k_1, 2^{2j}}(\sigma) \right) \\ &\lesssim \sum_{k_0 \leq j \leq -k} 2^{-\sigma k} (1 + \varepsilon B_1) h_{-j} h_{-j}(\sigma). \end{split}$$

Therefore, for $k_0 + k \le 0$, the High × High part is bounded by

$$\sum_{j\geq k_{0}} \int_{2j-1}^{2j+1} \sum_{|k_{1}-k_{2}|\leq 8,k_{1},k_{2}\geq k-4} \|P_{k}(P_{k_{1}}(\phi_{i} \diamond \phi_{s})P_{k_{2}}\mathcal{U})\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}(T)} d\tau$$

$$\lesssim \sum_{k_{0}\leq j\leq -k} C(1+2^{k+j})^{-12}2^{-\sigma k}h_{-j}h_{-j}(\sigma)2^{(1\pm\delta)(k+j)} + \sum_{j\geq -k} (...)$$

$$\lesssim (1+B_{1}\varepsilon)\sum_{k_{0}\leq j\leq -k} 2^{-\sigma k}h_{-j}h_{-j}(\sigma) + (1+B_{1}\varepsilon)2^{-\sigma k}h_{k}h_{k}(\sigma)$$

$$\lesssim \sum_{k_{0}\leq j\leq -k} (1+B_{1}\varepsilon)2^{-\sigma k}h_{-j}h_{-j}(\sigma).$$
(4.32)

where the $\sum_{j \ge -k} (...)$ part in (4.32) is bounded by $(1+B_1\varepsilon)2^{-\sigma k}h_kh_k(\sigma)$ via directly using results of the $k+k_0 \ge 0$ case. Therefore, we conclude that

$$\begin{split} &\sum_{j \ge k_0} \int_{2j-1}^{2j+1} \sum_{|k_1 - k_2| \le 8, k_1, k_2 \ge k-4} \|P_k(P_{k_1}(\phi_i \diamond \phi_s) P_{k_2} \mathcal{U})\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} d\tau \\ &\lesssim (1 + \varepsilon B_1) 2^{-\sigma k} (1 + 2^{j+k})^{-8} h_{k,s}(\sigma). \end{split}$$

Case c. Low × High \rightarrow High. In Case c, we assume $|k - k_2| \le 4$ and $k \ge k_1 + 4$, i.e. Low × High \rightarrow High. We applied the bound

$$\|P_{k}(f_{k_{1}}g_{k_{2}})\|_{F_{k}\cap S_{k}^{\frac{1}{2}}} \leq 2^{k_{1}}\|P_{k_{1}}f\|_{F_{k_{1}}\cap S_{k_{1}}^{\frac{1}{2}}}\|P_{k_{2}}g\|_{F_{k_{2}}(T)},$$

provided that $|k - k_2| \le 20$. To avoid too long formula, we recall the notation

$$T_{k,j} = (1_{k+j \ge 0} 2^j + 1_{k+j \le 0} 2^{-k}).$$

Thus by (4.22), (4.27) and (4.28), for $j \ge k_0$, $\sigma \in [0, \frac{99}{100}]$, we obtain that the Low × High \rightarrow High part is bounded by

$$\begin{split} &\sum_{j\geq k_0} \int_{2^{2j+1}}^{2^{2j+1}} \sum_{|k_2-k|\leq 4,k_1\leq k-4} \|P_k \left(P_{k_1}(\phi_s \diamond \phi_i) P_{k_2} \mathcal{U} \right) \|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \\ &\lesssim \sum_{j\geq k_0} \sum_{|k-k_2|\leq 4} \sum_{k_1\leq k-4} \int_{2^{2j+1}}^{2^{2j+1}} 2^{k_1} \|P_{k_1}(\phi_s \diamond \phi_i)\|_{F_{k_1} \cap S_{k_1}^{\frac{1}{2}}(T)} \|P_{k_2}(\mathcal{U})\|_{F_{k_2}(T)} \\ &\lesssim \sum_{j\geq k_0} h_k T_{k,j} (1+2^{j+k})^{-7} \sum_{k_1\leq k} 2^{k_1-\sigma k_1} h_{-j} \left(2^{k_1+j} h_{k_1}(\sigma) + h_{-j}(\sigma) \right) (1+2^{k_1+j})^{-7} \\ &+ \sum_{j\geq k_0} h_k T_{k,j} (1+2^{j+k})^{-7} B_1 \varepsilon 1_{k+j\geq 0} \sum_{k_1\leq -j} 2^{k_1-\sigma k_1} h_{-j} h_{-j}(\sigma) \\ &+ \sum_{j\geq k_0} h_k T_{k,j} (1+2^{j+k})^{-7} B_1 \varepsilon 1_{k+j\geq 0} \sum_{-j\leq k_1\leq k} 2^{k_1-\sigma k_1} 2^{k_1+j} h_{-j} h_{k_1}(\sigma) (1+2^{k_1+j})^{-7} \\ &+ \sum_{j\geq k_0} h_k T_{k,j} (1+2^{j+k})^{-7} B_1 \varepsilon 1_{k+j\geq 0} \sum_{k_1\leq k} 2^{k_1-\sigma k_1} h_{-j} h_{-j}(\sigma). \end{split}$$

Therefore, for $k + k_0 \ge 0$ we conclude

$$\begin{split} &\sum_{j\geq k_0} \int_{2^{2j+1}}^{2^{2j+1}} \sum_{|k_2-k|\leq 4, k_1\leq k-4} \|P_k \left(P_{k_1}(\phi_s \diamond \phi_i) P_{k_2} \mathcal{U} \right)\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \\ &\lesssim 2^{-\sigma k} \sum_{j\geq k_0} (1+B_1 \varepsilon) (1+2^{k+j})^{-7} 2^{\delta|j-k_0|} h_{-k_0} h_k(\sigma) \\ &\lesssim 2^{-\sigma k} \left(1+B_1 \varepsilon \right) (1+2^{2k+2j})^{-4} h_{k,2^{2k_0}}(\sigma). \end{split}$$

And for $k + k_0 \le 0$, we also have

$$\begin{split} &\sum_{j\geq k_0}\sum_{|k_2-k|\leq 8,k_1\leq k}\int_{2^{2j-1}}^{2^{2j+1}}\|P_k\left(P_{k_1}(\phi_s\diamond\phi_i)P_{k_2}\mathcal{U}\right)\|_{F_k(T)\cap S_k^{\frac{1}{2}}(T)} \\ &\lesssim 2^{-\sigma k}\left(1+B_1\varepsilon^{\frac{1}{2}}\right)(1+2^{2k+2j})^{-4}h_{k,2^{2k_0}}(\sigma). \end{split}$$

Thus the Low \times High part has been done for \mathcal{U} as well.

Therefore, combining the three cases, we summarize

$$\|P_k(A_i(s))\|_{F_k \cap S_k^{\frac{1}{2}}} \lesssim (\varepsilon^{\frac{1}{2}}B_1 + 1)2^{-\sigma k}(1 + 2^{k+k_0})^{-8}h_{k,2^{2k_0}}(\sigma).$$
(4.33)

Then (4.33) shows

$$B_1 \lesssim \varepsilon^{\frac{1}{2}} B_1 + 1. \tag{4.34}$$

Hence $B_1 \leq 1$. So, we have obtained our lemma for $\sigma \in [0, \frac{99}{100}]$ with assuming Bootstrap Assumption A. **Step 4.** In this step, we prove our lemma remains valid as if we drop the Bootstrap condition A in (4.28), (4.27). First, we prove a claim.

Claim A. If (4.27)-(4.28) hold, then for all $k, j \in \mathbb{Z}, \sigma \in [0, \frac{99}{100}], s \in [2^{2j-1}, 2^{2j+1}),$

$$\|P_k \mathcal{U}_l\|_{F_{k_2}(T) \cap S_k^{\frac{1}{2}}(T)} \le c_0^* 2^{-\sigma k} (1 + 2^{k+j})^{-7} T_{k,j} h_k(\sigma)$$
(4.35)

$$\|P_k \mathcal{U}_{II}\|_{F_{k_2}(T)} \le c_0^* 2^{-\sigma k} (1+2^{k+j})^{-7} T_{k,j} h_k(\sigma).$$
(4.36)

Recall the definition of \mathcal{U}_I

$$\mathcal{U}_I = -\Gamma_l^{\infty,(1)} \int_s^\infty \sum_{i=1,2} (A_i \psi_i)^l (s') ds',$$

For U_{II} , it is better to use

$$\mathcal{U}_{II} = -\int_{s}^{\infty} \sum_{i=1,2} (A_{i}\psi_{i})^{p}(s') \left[(\widetilde{\nabla}\mathbf{R})(e_{p}; e_{j_{0}}, e_{j_{1}}, e_{j_{2}}, e_{j_{3}}) - \Gamma_{p}^{\infty,(1)} \right] ds'.$$

Recall the notation

$$\widetilde{\mathcal{G}}^{(1)} = (\widetilde{\nabla} \mathbf{R})(e; e_{j_0}, ..., e_{j_3}) - \Gamma^{\infty, (1)}.$$

Moreover, we have by (3.76) and $c_0^* := \|\{h_k\}\|_{\ell^2}$ that

$$\|P_k(\widetilde{\mathcal{G}}^{(1)})(s)\|_{L^{\infty}_{t,x}} \lesssim 2^{-\sigma k} h_k(\sigma)(1+s2^{2k})^{-20}, \ \forall \sigma \in [0, \frac{99}{100}].$$
(4.37)

Thus in order to prove Claim A for \mathcal{U}_{II} , it suffices to prove

$$\int_{s}^{\infty} \|(A_{i}\psi_{i})\widetilde{\mathcal{G}}^{(1)}\|_{F_{k}(T)} ds' \lesssim (1+2^{k}s^{\frac{1}{2}})^{-7}c_{0}^{*}T_{k,j}.$$
(4.38)

The bound claimed for \mathcal{U}_{I} is easier to verify. Since now $B_{1} \leq 1$, applying bilinear Lemma 8.2 to $A_{i}\psi_{i}$, one has for $j + k \geq 0$, $s \in [2^{2j-1}, 2^{2j+1})$

$$\begin{aligned} \|(A_{i}\psi_{i})\|_{F_{k}(T)\cap S_{k_{1}}^{\frac{1}{2}}(T)} \\ \lesssim \left(1_{k+j\geq 0}2^{-j}+1_{k+j\leq 0}2^{\frac{k-j}{2}}\right)(1+2^{2k+2j})^{-4}2^{-\sigma k}c_{0}^{*}h_{k,s}(\sigma). \end{aligned}$$

$$\tag{4.39}$$

Summing the above formula w.r.t. $j \ge k_0$, we get

$$\int_{s}^{\infty} \|A_{i}\psi_{i}\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}(T)} \, ds' \lesssim c_{0}^{*} 2^{-\sigma k} T_{k,k_{0}}(1+2^{k_{0}+k})^{-7} h_{k}(\sigma).$$

$$(4.40)$$

Thus Claim A has been verified for \mathcal{U}_I .

For \mathcal{U}_{II} , we will use the following inequality (see (4.47))

$$\|P_{k}(P_{k_{1}}fP_{k_{2}}g)\|_{F_{k}(T)} \leq \|P_{k_{2}}g\|_{L_{x}^{\infty}}\|P_{k_{1}}f\|_{F_{k_{1}}(T)\cap S_{k_{1}}^{\frac{1}{2}}(T)}.$$

Then by Littlewood-Paley bilinear decomposition,

$$\begin{split} \|P_{k}((A_{i}\psi_{i})\widetilde{\mathcal{G}}^{(1)})\|_{F_{k}(T)} &\lesssim \sum_{|k_{1}-k| \leq 4} \|P_{k_{1}}(A_{i}\psi_{i})\|_{F_{k_{1}}(T) \cap S^{\frac{1}{2}}_{k_{1}}(T)} \|P_{\leq k-4}\widetilde{\mathcal{G}}^{(1)}\|_{L^{\infty}} \\ &+ \sum_{|k_{1}-k_{2}| \leq 8, k_{1}, k_{2} \geq k-4} \|P_{k_{1}}(A_{i}\psi_{i})\|_{F_{k_{1}}(T) \cap S^{\frac{1}{2}}_{k_{1}}(T)} \|P_{k_{2}}\widetilde{\mathcal{G}}^{(1)}\|_{L^{\infty}} \\ &+ \sum_{|k_{2}-k| \leq 4, k_{1} \leq k-4} \|P_{k_{1}}(A_{i}\psi_{i})\|_{F_{k_{1}}(T) \cap S^{\frac{1}{2}}_{k_{1}}(T)} \|P_{k_{2}}\widetilde{\mathcal{G}}^{(1)}\|_{L^{\infty}}. \end{split}$$

Thus by (4.39), the High × Low part of $(A_i\psi_i)\widetilde{\mathcal{G}}^{(1)}$ is dominated by

$$\begin{split} &\sum_{|k-k_1|\leq 4} \|P_k(P_{k_1}(A_i\psi_i)P_{\leq k-4}\widetilde{\mathcal{G}}^{(1)})\|_{F_k(T)} \\ &\lesssim c_0^* 2^{-\sigma k} \left(\mathbf{1}_{k+j\leq 0} 2^{\frac{k-j}{2}+\delta|k+j|} h_k(\sigma) + \mathbf{1}_{k+j\geq 0} 2^{-j} (1+2^{k+j})^{-7} h_k(\sigma) h_{-j} \right). \end{split}$$

Summing in $j \ge k_0$ yields

$$\begin{split} &\sum_{j\geq k_0} 2^{2j} \sum_{|k-k_1|\leq 4} \|P_k(P_{k_1}(A_i\psi_i)P_{\leq k}\widetilde{\mathcal{G}}^{(1)})\|_{F_k(T)} \\ &\lesssim c_0^* 2^{-\sigma k} h_k(\sigma) \Big(\mathbf{1}_{k+k_0\geq 0} 2^{k_0} (1+2^{k+k_0})^{-7} + \mathbf{1}_{k+k_0\leq 0} 2^{-k} \Big). \end{split}$$

Using (4.37) and (4.39), the High × High part of $(A_i\psi_i)\widetilde{\mathcal{G}}^{(1)}$ is dominated by

$$\begin{split} &\sum_{|k_2-k_1|\leq 8,k_1,k_2\geq k-4} ||P_k(P_{k_1}(A_i\psi_i)P_{k_2}\widetilde{\mathcal{G}}^{(1)})||_{F_k(T)} \\ &\lesssim c_0^* \mathbf{1}_{k+j\geq 0} \sum_{k_1\geq k-4} 2^{-\sigma k_1} (1+2^{2j+2k})^{-7} 2^{k_1} h_{k_1}(\sigma) \\ &+ c_0^* \mathbf{1}_{k+j\leq 0} \left[\sum_{k-4\leq k_1\leq -j} 2^{\frac{k_1-j}{2}} 2^{\delta|k_1+j|} 2^{-\sigma k_1} h_{k_1}(\sigma) + \sum_{k_1\geq -j} 2^{-\sigma k_1} (1+2^{2j+2k})^{-7} 2^{k_1} h_{k_1}(\sigma) \right] \\ &\lesssim c_0^* \mathbf{1}_{k+j\geq 0} 2^{-j} (1+2^{j+k})^{-10} 2^{-\sigma k} h_k(\sigma) + c_0^* \mathbf{1}_{k+j\leq 0} 2^{-j} 2^{\delta|k+j|} 2^{-\sigma k} h_k(\sigma). \end{split}$$

Summing in $j \ge k_0$ also gives

$$\sum_{j\geq k_0} 2^{2j} \sum_{\substack{|k_2-k_1|\leq 8,k_1,k_2\geq k-4 \\ \leq c_0^* 2^{-\sigma k} \mathbf{1}_{k+k_0\geq 0} 2^{k_0} (1+2^{k+k_0})^{-7} h_k(\sigma) + c_0^* \mathbf{1}_{k+k_0\leq 0} 2^{-k} 2^{-\sigma k} h_k(\sigma).$$

Then using (4.37), the Low × High part of $(A_i\psi_i)\widetilde{\mathcal{G}}^{(1)}$ is bounded as

$$\begin{split} &\sum_{|k-k_2| \le 4, k_1 \le k-4} \|P_k(P_{k_1}(A_i\psi_i)P_{k_2}\widetilde{\mathcal{G}}^{(1)})\|_{F_k(T)} \\ &\lesssim h_k(\sigma) 2^{-\sigma k} \mathbf{1}_{k+j \le 0} \sum_{k_1 \le k} h_{k_1} 2^{\frac{1}{2}(k_1-j)} 2^{\delta|k_1+j|} \\ &+ h_k(\sigma) 2^{-\sigma k} (1+2^{2j+2k})^{-20} \mathbf{1}_{k+j \ge 0} \left[\sum_{-j \le k_1 \le k} h_{k_1} 2^{k_1} (1+2^{2j+2k_1})^{-4} \right] \\ &+ h_k(\sigma) 2^{-\sigma k} (1+2^{2j+2k})^{-20} \mathbf{1}_{k+j \ge 0} \left[\sum_{k_1 \le -j} h_{k_1} 2^{\frac{k_1-j}{2}} 2^{\delta|k_1+j|} \right] \\ &\lesssim c_0^* 2^{-\sigma k} \mathbf{1}_{k+j \ge 0} 2^{-j} (1+2^{j+k})^{-7} 2^{-\sigma k} h_k(\sigma) + c_0^* 2^{-\sigma k} \mathbf{1}_{k+j \le 0} 2^{\frac{k-j}{2}} 2^{\delta|k+j|} 2^{-\sigma k} h_k(\sigma) . \end{split}$$

Summing in $j \ge k_0$ as well yields

$$\begin{split} &\sum_{j\geq k_0} 2^{2j} \sum_{\substack{|k-k_2|\leq 4,k_1\leq k+4 \\ \leqslant c_0^* 2^{-\sigma k} \mathbf{1}_{k+k_0\geq 0} 2^{k_0} (1+2^{k+k_0})^{-7} h_k(\sigma) + c_0^* \mathbf{1}_{k+k_0\leq 0} 2^{-k} 2^{-\sigma k} h_k(\sigma). \end{split}$$

Thus back to the LHS of (4.38), we conclude if (4.28) holds, then

$$\|P_k\mathcal{U}_{II}\|_{F_k(T)} \lesssim c_0^* 2^{-\sigma k} \mathbf{1}_{k+k_0 \ge 0} 2^{k_0} (1+2^{k+k_0})^{-7} h_k(\sigma) + c_0^* \mathbf{1}_{k+k_0 \le 0} 2^{-k_0} 2^{-\sigma k} h_k(\sigma).$$

Particularly (4.28) holds, thus proving Claim A.

Now we are ready to prove our lemma with (4.27) and (4.28) being dropped. Define the function on $T' \in [0, T]$

$$\Phi(T') = \sum_{\{j_c\} \subset \{1,...,2n\}} \sup_{\substack{k,j \in \mathbb{Z}}} \sup_{s \in [2^{2j-1}, 2^{2j+1}]} (c_0^*)^{-1} (1 + 2^k 2^{\frac{s}{2}})^7 T_{k,j}^{-1} h_k^j$$

 $\times (||P_k \mathcal{U}_I||_{F_k(T')} + ||P_k \mathcal{U}_{II}||_{F_k(T')}).$

Using Lemma 3.3 and Sobolev embeddings, we find that $\Phi(T')$ is a continuous function in $T' \in [0, T]$. Then in order to prove our lemma, it suffices to prove $\Phi \leq 1$. It is easy to see Φ is also an increasing continuous function on $T' \in [0, T]$. And Claim A shows

$$\Phi(T') \le \varepsilon^{-\frac{1}{2}} \Longrightarrow \Phi(T') \le 1.$$

Hence it suffices to verify

$$\lim_{T'\to 0} \Phi(T') \lesssim 1.$$

This reduces to prove for all $j, k \in \mathbb{Z}, s \in [2^{2j-1}, 2^{2j+1})$

$$\sum_{\substack{\{j_c\} \subset \{1,...,2n\}}} \left\| P_k \int_s^\infty (A_i \psi_i)^q \left[(\widetilde{\nabla} \mathbf{R}) (e_q; e_{j_0}, ..., e_{j_3}) - \Gamma_q^{\infty,(1)} \right] \right\|_{L^2_x} + \int_s^\infty \| P_k (A_i \psi_i) \|_{L^2_x} ds \leq c_0^* (1 + 2^k 2^{\frac{s}{2}})^{-7} T_{k,j} h_k,$$

where all these fields ψ_i , matrices A_i are associated with the heat flow with initial data u_0 . This can be proved by applying results of Section 3. In fact, by the definition of $h_k(\sigma)$ one has

$$2^{\sigma k} \|\phi_i\|_{s=0} \|_{L^{\infty}_t L^2_x} \le h_k(\sigma),$$

if $\sigma \in [0, \frac{99}{100}]$. Then by Proposition 3.5 with $\eta_k(\sigma) = h_k(\sigma)$ we get

$$(1+2^{2k}s)^{29} \|P_k A_i(s)\|_{L_t^{\infty} L_x^2} \le 2^{-\sigma k} h_{k,s}(\sigma).$$
(4.41)

And proof of Proposition 3.5 shows

$$(1+2^{2k}s)^{30} \|P_k\psi_i(s)\|_{L^\infty_t L^2_x} \le 2^{-\sigma k} h_k(\sigma).$$
(4.42)

if $\sigma \in [0, \frac{99}{100}]$. Then by (4.42), (4.41) and bilinear Littlewood-Paley decomposition, one obtains

$$(1+2^{2k}s)^{28}\int_s^\infty \|P_k(A_i\psi_i)\|_{L^\infty_t L^2_x} ds' \lesssim \|\{h_k\}\|_{\ell^2} T_{k,j}h_k(\sigma) 2^{-\sigma k}$$

for $s \in [2^{2j-1}, 2^{2j+1}), k, j \in \mathbb{Z}$. The left is to prove

$$(1+2^{2k}s)^{28}\int_{s}^{\infty} \|P_{k}[A_{i}\psi_{i}\widetilde{\mathcal{G}}^{(1)}]\|_{L_{i}^{\infty}L_{x}^{2}}ds' \lesssim \|\{h_{k}\}\|_{\ell^{2}}T_{k,j}h_{k}(\sigma)2^{-\sigma k}, \ \sigma \in [0,\frac{99}{100}].$$
(4.43)

This follows by (4.41), (4.42), (3.55) and bilinear Littlewood-Paley decomposition as well.

Remark 4.1. Checking the proof of Lemma 4.1, we see the range of $\sigma \in [0, \frac{99}{100}]$ was only used in the Low × High interaction of $(\partial_i \psi_i) \widetilde{\mathcal{G}}^{(1)}$, $(A_i \psi_i) (\mathcal{U}_I + \mathcal{U}_{II})$ of Step 2 and Step 3 respectively.

Lemma 4.2. *If* $|k_1 - k| \le 4$, *then*

$$\|P_k(P_{k_1}fP_{\leq k-4}g)\|_{F_k(T)} \lesssim \|P_{k_1}g\|_{F_{k_1}(T)}\|P_{\leq k-4}g\|_{L^{\infty}}.$$
(4.44)

If $|k_2 - k_1| \le 8$, $k_1, k_2 \ge k - 4$, then

$$\|P_{k}(P_{k_{1}}fP_{k_{2}}g)\|_{F_{k}(T)} \leq \|P_{k_{1}}f\|_{F_{k_{1}}(T)}(\|P_{k_{2}}g\|_{L^{\infty}} + 2^{\frac{1}{2}k_{2}}\|P_{k_{2}}g\|_{L^{4}_{x}L^{\infty}_{t}}).$$

$$(4.45)$$

If $|k_2 - k| \le 4$, $k_1 \le k - 4$, then

$$\|P_{k}(P_{k_{1}}fP_{k_{2}}g)\|_{F_{k}(T)} \leq 2^{\frac{k_{1}}{2}} \|P_{k_{1}}f\|_{F_{k_{1}}(T)} \|P_{k_{2}}g\|_{L^{4}_{x}L^{\infty}_{t}} + 2^{k_{1}} \|P_{k_{1}}f\|_{F_{k_{1}}(T)} \|P_{k_{2}}g\|_{L^{4}}.$$
(4.46)

For any $k_1, k_2, k \in \mathbb{Z}$ *, one has*

$$\|P_{k}(P_{k_{1}}fP_{k_{2}}g)\|_{F_{k}(T)} \lesssim \|P_{k_{1}}f\|_{F_{k_{1}}(T) \cap S^{\frac{1}{2}}_{k_{1}}(T)}\|P_{k_{2}}g\|_{L^{\infty}}.$$
(4.47)

Proof. (4.44) has been given in [4]. And (4.45) follows by Hölder and [(3.17), [4]] which says

$$\|P_k f\|_{F_k(T)} \lesssim \|P_k f\|_{L^2_x L^\infty_t} + \|P_k f\|_{L^4_{t,x}}.$$

(4.46) follows by the same reason with additionally using Bernstein inequality. Moreover, by definition

$$||P_k f||_{L^4_{t,x}} \le ||f||_{F_k(T)}, ||P_k f||_{L^2_x L^\infty_t} \le ||f||_{S^{\frac{1}{2}}_k(T)}.$$

Thus one obtains

$$\begin{split} \|P_{k}(P_{k_{1}}fP_{k_{2}}g)\|_{F_{k}(T)} &\leq \|P_{k_{1}}fP_{k_{2}}g\|_{L_{x}^{2}L_{t}^{\infty}} + \|P_{k_{1}}fP_{k_{2}}g\|_{L_{t,x}^{4}} \\ &\leq \left(\|P_{k_{1}}f\|_{L_{x}^{2}L_{t}^{\infty}} + \|P_{k_{1}}f\|_{L_{t,x}^{4}}\right)\|P_{k_{2}}g\|_{L^{\infty}} \\ &\leq \|P_{k_{2}}g\|_{L^{\infty}}\|P_{k_{1}}f\|_{F_{k_{1}}(T)\cap S^{\frac{1}{2}}_{k_{1}}(T)}. \end{split}$$

The proof of Lemma 4.1 yields

Corollary 4.1. Under the assumptions of Proposition 4.1 and (4.9), for $s \in [2^{2j-1}, 2^{2j+1}]$, $\sigma \in [0, \frac{99}{100}]$, $j, k \in \mathbb{Z}$, there holds

$$||P_k(\mathcal{G})||_{F_k(T)} \leq 2^{-\sigma k} h_k(\sigma) T_{k,j} (1+2^{j+k})^{-7}.$$

where $T_{k,j}$ is defined by (4.29). When s = 0, we have

$$\|P_k(\widetilde{\mathcal{G}})\|_{s=0}\|_{F_k(T)} \leq 2^{-\sigma k} h_k(\sigma) 2^{-k}.$$

Proof. Lemma 4.1 gives

$$\begin{split} \|P_{k}(\mathcal{U}_{00})\|_{F_{k}(T)} + \|P_{k}(\mathcal{U}_{01})\|_{F_{k}(T)} &\leq 2^{-\sigma k} h_{k}(\sigma) T_{k,j}(1+2^{j+k})^{-7} \\ \|P_{k}\mathcal{U}_{I}\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} &\leq 2^{-\sigma k} T_{k,j} h_{k}(\sigma)(1+2^{j+k})^{-7} \\ \|P_{k}\mathcal{U}_{II}\|_{F_{k}(T)} &\leq 2^{-\sigma k} T_{k,j} h_{k}(\sigma)(1+2^{j+k})^{-7}. \end{split}$$

Then our corollary follows by the decomposition

$$\overline{\mathcal{G}} = \mathcal{G} - \Gamma^{\infty} = \mathcal{U}_{00} + \mathcal{U}_{01} + \mathcal{U}_{I} + \mathcal{U}_{II}$$

and the inequality $(1 + 2^{j+k})^{-1}T_{k,j} \le 2^{-k}$ for all $j, k \in \mathbb{Z}$.

4.3 Evolution of differential fields along the heat flow

Recall the evolution equation for ϕ_i along the heat flow:

$$(\partial_s - \Delta)\phi_i = K_i$$

$$K_i := 2\sum_{j=1}^2 \partial_j (A_j\phi_i) + \sum_{j=1}^2 (A_j^2 - \partial_j A_j)\phi_i + \sum_{j=1}^2 \phi_j \diamond \phi_i \diamond \phi_j \mathcal{G}.$$
(4.48)

Now we control the nonlinearities in the above equations.

Lemma 4.3. Under the assumptions of Proposition 4.1 and (4.9), for all $s \in [0, \infty)$, i = 1, 2 and $\sigma \in [0, \frac{99}{100}]$, we have

$$\|\int_{0}^{s} e^{(s-\tau)\Delta} P_{k} K_{i}(\tau) d\tau\|_{F_{k}(T)} \lesssim \varepsilon (1+s2^{2k})^{-4} 2^{-\sigma k} h_{k}(\sigma).$$
(4.49)

Proof. First, we consider the quartic term $\mathcal{G}(\phi_i \diamond \phi_j) \diamond \phi_j$ in K_i . [[4], (5.25)] have proved that for $\tau \in [2^{2j-1}, 2^{2j+1})$

$$\|P_k\left(\phi_i \diamond \phi_p \diamond \phi_l\right)(\tau)\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \lesssim \varepsilon 2^{-\sigma k} 2^{2k} (1 + 2^{2k+2j})^{-4} \left[h_k(\sigma) + 2^{-\frac{3}{2}(k+j)} h_{-j}(\sigma)\right].$$
(4.50)

Recall that $\mathcal{G} = \widetilde{\mathcal{G}} + \Gamma^{\infty}$. The constant part follows by (4.50). By bilinear Littlewood-Paley decomposition we have

$$\begin{split} \|P_{k}\left(\phi_{i} \diamond \phi_{p} \diamond \phi_{l}\mathcal{G}\right)\|_{F_{k}(T)} \\ &\lesssim \sum_{k_{1} \geq k-4}^{|k_{1}-k_{2}| \leq 8} \|P_{k_{1}}\left(\phi_{i} \diamond \phi_{p} \diamond \phi_{l}\right)P_{k_{2}}\widetilde{\mathcal{G}}\|_{F_{k}(T)} + \sum_{k_{1} \leq k-4}^{|k_{2}-k| \leq 4} \|P_{k_{1}}\left(\phi_{i} \diamond \phi_{p} \diamond \phi_{l}\right)P_{k_{2}}\widetilde{\mathcal{G}}\|_{F_{k}(T)} \\ &+ \sum_{|k_{1}-k| \leq 4} \|P_{k_{1}}\left(\phi_{i} \diamond \phi_{p} \diamond \phi_{l}\right)P_{\leq k-4}\widetilde{\mathcal{G}}\|_{F_{k}(T)} \end{split}$$

For the High × Low term, directly applying $\|\widetilde{\mathcal{G}}\|_{L^{\infty}_{t,x}} \leq K(\mathcal{N})$ gives

$$\begin{split} \|P_{k_1}\left(\phi_i \diamond \phi_p \diamond \phi_l\right) P_{\leq k-4} \widetilde{\mathcal{G}}\|_{F_k(T)} &\lesssim \|P_k\left(\phi_i \diamond \phi_p \diamond \phi_l\right)\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \\ &\lesssim \varepsilon 2^{-\sigma k} 2^{2k} (1+2^{2k+2j})^{-4} \left[h_k(\sigma) + 2^{-\frac{3}{2}(k+j)} h_{-j}(\sigma)\right]. \end{split}$$

For the High × High term, denoting $\mathcal{V} := \phi_i \diamond \phi_p \diamond \phi_l$, Corollary 4.1 and (8.4) show

$$\sum_{|k_{1}-k_{2}|\leq 8,k_{1},k_{2}\geq k-4} \|P_{k_{1}}\left(\phi_{i} \diamond \phi_{p} \diamond \phi_{l}\right)P_{k_{2}}\mathcal{G}\|_{F_{k}(T)}$$

$$\lesssim \sum_{|k_{1}-k_{2}|\leq 8,k_{1},k_{2}\geq k-4} 2^{\frac{k_{1}+k}{2}} \|P_{k_{1}}\mathcal{V}\|_{F_{k_{1}}(T)} \cap S_{k_{1}}^{\omega}(T)\|P_{k_{2}}\mathcal{G}\|_{F_{k_{2}}(T)}$$

$$\lesssim \sum_{k_{1}\geq k,|k_{1}-k_{2}|\leq 2} 2^{\frac{k_{1}+k}{2}} 2^{-\sigma k_{1}+2k_{1}} (1+2^{k_{1}+j})^{-15} \left[h_{k_{1}}(\sigma)+2^{-\frac{3}{2}(k_{1}+j)}h_{-j}(\sigma)\right] T_{k_{1},j}h_{k_{1}}.$$
(4.51)

If $k + j \ge 0$, then by slow variation of envelopes, (4.51) is bounded by

$$2^{3k+j}2^{-\sigma k}(1+2^{k+j})^{-14}h_k(\sigma)h_k.$$

If $k + j \le 0$, using $(1 + 2^{k_1 + j})^{-1} T_{k_1, j} \le 2^{-k_1}$ for all $k_1 \in \mathbb{Z}$, (4.51) is dominated by

$$2^{\frac{k}{2}-\frac{3j}{2}}h_k(\sigma)h_k2^{\delta|j+k|}.$$

Therefore, for the High × High interaction, we have if $s \in [2^{2j-1}, 2^{2j+1})$ then

$$\sum_{|k_1-k_2|\le 8, k_1, k_2\ge k-4} \|P_{k_1}\left(\phi_i \diamond \phi_p \diamond \phi_l\right) P_{k_2} \widetilde{\mathcal{G}}\|_{F_k(T)} \lesssim 2^{-\sigma k-\frac{3}{2}j} 2^{\frac{1}{2}k} 2^{\delta|j+k|} h_k h_k(\sigma) (1+2^{2k+2j})^{-5}.$$
(4.52)

To finish estimates for the High × High interaction, we turn to verify the corresponding part in (4.49). We use (4.52) to verify (4.49). Let $s \in [2^{2k_0-1}, 2^{2k_0+1})$ with $k_0 \in \mathbb{Z}$ fixed. For $k + k_0 \le 0$, by (2.8) one has

$$\begin{split} \| \int_{0}^{s} e^{(s-\tau)\Delta} \sum_{k_{1} \ge k-4, |k_{1}-k_{2}| \le 8} P_{k}(P_{k_{1}} \mathcal{V} P_{k_{2}} \widetilde{\mathcal{G}}) d\tau \|_{F_{k}(T)} \\ \lesssim \sum_{j \le k_{0}} \int_{2^{2j+1}}^{2^{2j+1}} \sum_{k_{1} \ge k-4, |k_{1}-k_{2}| \le 8} \| P_{k}(P_{k_{1}} \mathcal{V} p_{k_{2}} \widetilde{\mathcal{G}}) \|_{F_{k}(T)} d\tau \lesssim \varepsilon \sum_{j \le k_{0}} 2^{-\sigma k} h_{k}(\sigma) \left(2^{\frac{1}{2}(j+k)} + 2^{(\frac{1}{2} \pm \delta)(k+j)} \right) \\ \lesssim \varepsilon 2^{-\sigma k} h_{k}(\sigma). \end{split}$$

For $k + k_0 \ge 0$, by (2.8), (4.52) one has

$$\begin{split} &\int_{0}^{s} \|e^{(s-\tau)\Delta} \sum_{k_{1} \geq k-4, |k_{1}-k_{2}| \leq 8} P_{k}(P_{k_{1}} \mathcal{V} P_{k_{2}} \widetilde{\mathcal{G}})\|_{F_{k}(T)} d\tau \lesssim \int_{0}^{\frac{s}{2}} ... d\tau + \int_{\frac{s}{2}}^{s} ... d\tau \\ &\lesssim \sum_{j \leq -k_{0}-1} \int_{2^{2j+1}}^{2^{2j+1}} 2^{-20(k+k_{0})} \sum_{k_{1} \geq k-4, |k_{1}-k_{2}| \leq 8} \|P_{k}(P_{k_{1}} \mathcal{V} P_{k_{2}} \widetilde{\mathcal{G}})\|_{F_{k}(T)} d\tau \\ &+ 2^{2k_{0}} \sum_{k_{1} \geq k-4, |k_{1}-k_{2}| \leq 8} \|P_{k}(P_{k_{1}} \mathcal{V} P_{k_{2}} \widetilde{\mathcal{G}})\|_{F_{k}(T)} \upharpoonright_{\tau \in [2^{2k_{0}-2}, 2^{2k_{0}+1}]} \\ &\lesssim \varepsilon 2^{-20(k+k_{0})} \sum_{j \leq -k_{0}-1} 2^{-\sigma k} h_{k}(\sigma) \left(2^{\frac{1}{2}(j+k)} + 2^{(\frac{1}{2}\pm\delta)(j+k)}\right) \\ &+ \varepsilon 2^{-\sigma k} h_{k}(\sigma) (1 + 2^{k+k_{0}})^{-10} \left[2^{\frac{1}{2}(k_{0}+k)} + 2^{(\frac{1}{2}\pm\delta)(k_{0}+k)}\right] \\ &\lesssim \varepsilon (1 + 2^{k+k_{0}})^{-8} 2^{-\sigma k} h_{k}(\sigma). \end{split}$$

Thus we conclude

$$\int_0^s \|e^{(s-\tau)\Delta} \sum_{k_1 \ge k-4, |k_1-k_2| \le 8} P_k(P_{k_1} \mathcal{W} P_{k_2} \widetilde{\mathcal{G}})\|_{F_k(T)} d\tau \lesssim \varepsilon^2 2^{-\sigma k} h_k(\sigma) (1+2^{2k}s)^{-4}.$$

For the Low \times High term, using

$$\|P_{k_1}(\mathcal{V})\|_{L^{\infty}_{t,x}} \lesssim 2^{k_1} \varepsilon 2^{-\sigma k_1 + 2k_1} (1 + 2^{k_1 + j})^{-8} \left[h_{k_1}(\sigma) + 2^{-\frac{3}{2}(k_1 + j)} h_{-j}(\sigma)\right]$$

we obtain by Corollary 4.1 and (8.4) that

$$\begin{split} &\sum_{k_1 \leq k-4, |k-k_2| \leq 4} \left\| P_k \left(P_{k_1}(\phi_i \diamond \phi_p \diamond \phi_l) P_{k_2} \widetilde{\mathcal{G}} \right) \right\|_{F_k(T)} \\ &\lesssim 2^{-\sigma k} T_{k,j} (1+2^{j+k})^{-7} h_k(\sigma) \varepsilon \sum_{k_1 \leq k-4} 2^{3k_1} (1+2^{k_1+j})^{-8} \left(1+2^{-\frac{3}{2}(k_1+j)} \right). \end{split}$$

For $k + j \ge 0$, we have

$$\sum_{k_1 \le k-4, |k-k_2| \le 4} \left\| P_k(P_{k_1}\left((\phi_i \diamond \phi_p \diamond \phi_l) P_{k_2} \widetilde{\mathcal{G}} \right) \right\|_{F_k(T)} \le \varepsilon 2^{-\sigma} 2^{-2j} (1 + 2^{k+j})^{-7} h_k(\sigma).$$
(4.53)

For $k + j \le 0$, we have

$$\sum_{k_1 \le k-4, |k-k_2| \le 4} \left\| P_k \left(P_{k_1}(\phi_i \diamond \phi_p \diamond \phi_l) P_{k_2} \widetilde{\mathcal{G}} \right) \right\|_{F_k(T)} \le \varepsilon 2^{-\sigma k} 2^{\frac{1}{2}k - \frac{3}{2}j} (1 + 2^{k+j})^{-7} h_k(\sigma).$$
(4.54)

As a summary, we use (4.53), (4.54) to verify (4.49). Let $s \in [2^{2k_0-1}, 2^{2k_0+1})$ with k_0 fixed. For $k + k_0 \le 0$, by (2.8) one has for the Low × High interaction that

$$\begin{split} \int_0^s \|e^{(s-\tau)\Delta} \sum_{k_1 \le k-4, |k-k_2| \le 4} P_k(P_{k_1} \mathcal{V} P_{k_2} \widetilde{\mathcal{G}})\|_{F_k(T)} d\tau &\lesssim \sum_{j \le k_0} \int_{2^{2j-1}}^{2^{2j+1}} \dots d\tau \\ &\lesssim \varepsilon \sum_{j \le k_0} 2^{\frac{j}{2} + \frac{k}{2}} 2^{-\sigma k} h_k(\sigma) \lesssim 2^{-\sigma k} \varepsilon h_k(\sigma). \end{split}$$

For $k + k_0 \ge 0$, similarly we have

$$\begin{split} &\int_{0}^{s} \|e^{(s-\tau)\Delta} \sum_{k_{1} \leq k-4, |k-k_{2}| \leq 4} P_{k}(P_{k_{1}}\mathcal{W}P_{k_{2}}\widetilde{\mathcal{G}})\|_{F_{k}(T)} d\tau \\ &\lesssim \varepsilon 2^{-20(k+k_{0})} 2^{-\sigma k} h_{k}(\sigma) \sum_{j \leq -k_{0}-1} 2^{\frac{1}{2}(j+k)} + \varepsilon 2^{-\sigma k} h_{k}(\sigma)(1+2^{k+k_{0}})^{-7} 2^{-2k_{0}-2k} \\ &\lesssim \varepsilon (1+2^{k+k_{0}})^{-8} 2^{-\sigma k} h_{k}(\sigma). \end{split}$$

Thus we conclude

$$\int_0^s \|e^{(s-\tau)\Delta} \sum_{k_1 \leq k-4, |k-k_2| \leq 4} P_k(P_{k_1} \mathcal{W} P_{k_2} \widetilde{\mathcal{G}})\|_{F_k(T)} d\tau \lesssim \varepsilon (1+2^{k+k_0})^{-8} 2^{-\sigma k} h_k(\sigma).$$

And the High \times Low case is easy by repeating the same argument or directly applying the result of [[4], Lemma 5.3]. Therefore, the curvature term has been done:

$$\left\|\int_0^s e^{(s-\tau)\Delta} P_k\left(\phi_i \diamond \phi_p \diamond \phi_l \widetilde{\mathcal{G}}\right) d\tau\right\|_{F_k(T)} \lesssim \varepsilon (1+s2^{2k})^{-4} 2^{-\sigma k} h_k(\sigma).$$

Step 2. Connection coefficient term. In this step, we turn to estimate the terms $\partial_l(A_l\psi_i)$, $\partial_lA_l\phi_i$ and $A_l^2\phi_i$. With Lemma 4.1 in hand, all these terms follow directly by repeating arguments of [[4],Lemma 5.3].

Lemma 4.4. Under the assumptions of Proposition 4.1 and (4.9), for all $k \in \mathbb{Z}$, $s \ge 0$, i = 1, 2, we have

$$\|P_k\phi_i(s)\|_{F_k(T)} \leq b_k(\sigma)2^{-\sigma k}(1+s2^{2k})^{-4} \ \sigma \in [0, \frac{99}{100}].$$
(4.55)

Proof. By Duhamel principle and (4.49), we get

$$\sup_{s \ge 0} (1 + s2^{2k})^4 2^{\sigma k} \|P_k \phi_i(s)\|_{F_k(T)} \le b_k(\sigma) + \varepsilon h_k(\sigma).$$
(4.56)

Since the RHS of (4.56) is a frequency envelope of δ order, by the definition of $\{h_k(\sigma)\}$ we get

$$h_k(\sigma) \leq b_k(\sigma) + \varepsilon h_k(\sigma),$$
(4.57)

which by letting ε be sufficiently small yields

$$h_k(\sigma) \leq b_k(\sigma).$$
 (4.58)

Lemma 4.5. Under the assumptions of Proposition 4.1 and (4.9), for all $k \in \mathbb{Z}$, $s \ge 0$, i = 1, 2, we have

$$\|P_k A_i\|_{s=0} \|_{L^4_{u_s}} \lesssim b_k(\sigma) 2^{-\sigma k}.$$
(4.59)

Proof. Lemma 4.4 has shown (4.58). Then the previous bounds in Lemma 4.1 now hold with $h_k(\sigma)$ replaced by $b_k(\sigma)$. Recall that [[4], Page 1473-1474] proved

$$\|P_k\phi_s\|_{L^4_{t,x}} \lesssim 2^k 2^{-\sigma k} b_k(\sigma) (2^{2k} s)^{-\frac{3}{8}} (1 + s 2^{2k})^{-3}.$$
(4.60)

We also recall the bilinear estimate of [[4], Lemma 5.4] in our Appendix A, Lemma 8.4. Then (4.60) and (4.55) show

$$\begin{split} \|P_k(\phi_i \diamond \phi_s)\|_{L^4_{t,x}} &\lesssim 2^{-\sigma k} \sum_{l \leq k} b_k(\sigma) b_l 2^{l+k} (s 2^{2k})^{-\frac{3}{8}} (1+s 2^{2k})^{-3} \\ &+ 2^{-\sigma k} \sum_{l \leq k} b_k(\sigma) b_l 2^{2l} 2^{\frac{1}{2}(k-l)} (2^{2k} s)^{-\frac{3}{8}} (1+s 2^{2k})^{-4} \\ &+ \sum_{l \geq k} 2^{-\sigma l} b_l(\sigma) b_l 2^{k+l} (2^{2l} s)^{-\frac{3}{8}} (1+s 2^{2l})^{-7}. \end{split}$$

Thus given $s \in [2^{2j-1}, 2^{2j+2})$ with $j \in \mathbb{Z}$, we conclude for $k + j \ge 0$,

$$\|P_k(\phi_i \diamond \phi_s)\|_{L^4_{t,s}} \lesssim b_k(\sigma)b_k 2^{2k-\sigma k} (s2^{2k})^{-\frac{3}{8}} (1+s2^{2k})^{-3}, \tag{4.61}$$

and for $k + j \le 0$,

$$\|P_k(\phi_i \diamond \phi_s)\|_{L^4_{t_x}} \le b_k(\sigma)b_k 2^{2\delta|k+j|} 2^{-\sigma k} 2^k 2^{-j}.$$
(4.62)

Recalling that for i = 1, 2,

$$A_i(0) = \int_0^\infty ((\phi_i \diamond \phi_s)\mathcal{G})ds.$$
(4.63)

we see the left is to deal with the interaction of $\phi_s \diamond \phi_i$ with \mathcal{G} . Recall that $\mathcal{G} = \Gamma^{\infty} + \widetilde{\mathcal{G}}$ with

$$\|P_k(\widetilde{\mathcal{G}})\|_{F_k(T)} \lesssim 2^{-\sigma k} T_{k,j} (1+2^{k+j})^{-7} h_k(\sigma), \tag{4.64}$$

for $s \in [2^{2j-1}, 2^{2j+1})$ with $j \in \mathbb{Z}$. (4.62), (4.61) show the constant part Γ^{∞} contributes to $||A_i(0)||_{L^4_{t,x}}$ by $b_k(\sigma)2^{-\sigma k}$. Thus it suffices to control $(\phi_i \diamond \phi_s)\widetilde{\mathcal{G}}$.

As before, we consider three cases according to Littlewood-Paley decomposition. Using the trivial bound $\|\widetilde{\mathcal{G}}\|_{L^{\infty}_{t,x}} \leq K(\mathcal{N})$ in the High × Low part gives

$$\begin{split} &\sum_{j\in\mathbb{Z}} \int_{2^{2j+1}}^{2^{2j+1}} \sum_{|k_1-k|\leq 4} \|P_k \left[P_{k_1}(\phi_s \diamond \phi_i) P_{\leq k-4} \widetilde{\mathcal{G}} \right] \|_{L^4_{t,x}} ds \\ &\lesssim \sum_{j\geq -k} \mathbf{1}_{k+j\geq 0} b_k(\sigma) b_k 2^{2k+2j-\sigma k} (2^{2k+2j})^{-\frac{3}{8}} (1+2^{2k+2j})^{-3} + \sum_{j\leq -k} \mathbf{1}_{k+j\leq 0} b_k(\sigma) b_k 2^{2\delta|k+j|} 2^{-\sigma k} 2^k 2^j \\ &\lesssim b_k(\sigma) b_k 2^{-\sigma k}. \end{split}$$

Notice that Lemma 8.4 shows

$$\sum_{|k_1-k_2|\leq 8, k_1, k_2\geq k-4} \left\| P_k(P_{k_1}fP_{k_2}g) \right\|_{L^4_{t,x}} \lesssim \sum_{k_1\geq k} 2^{k(1+\omega)} 2^{-\omega k_1} \mu_{k_1} \nu_{k_1}$$

where $\mu_k = \sum_{|k-k'| \le 20} \|P_{k'}f\|_{S_{k'}^{\omega}}, v_k = \sum_{|k-k'| \le 20} \|P_{k'}g\|_{L_{t_x}^4}$. Thus using (4.64), we have by choosing $\omega = 0$ that

$$\begin{split} &\int_{0}^{\infty} \sum_{|k_{1}-k_{2}| \leq 8, k_{1}, k_{2} \geq k-4} \left\| P_{k} \left[P_{k_{1}}(\phi_{s} \diamond \phi_{i}) P_{k_{2}} \widetilde{\mathcal{G}} \right] \right\|_{L^{4}_{t,x}} ds \\ &\lesssim \sum_{j \geq -k} 2^{2j} \varepsilon 2^{k} \sum_{k_{1} \geq k-4} b_{k_{1}} b_{k_{1}}(\sigma) 2^{k_{1}} 2^{-\sigma k_{1}} (1 + 2^{2j+2k_{1}})^{-5} \\ &+ \sum_{j \leq -k} 2^{-\sigma k} 2^{k} \sum_{k-4 \leq k_{1} \leq -j} 2^{j} b_{k_{1}}(\sigma) b_{k_{1}} 2^{2\delta|k_{1}+j|} \\ &+ 2^{-\sigma k} \sum_{j \leq -k} 2^{2j} 2^{k} \sum_{k_{1} \geq -j} b_{k_{1}}(\sigma) b_{k_{1}} 2^{2\delta|k_{1}+j|} 2^{k_{1}} (2^{2j+2k_{1}})^{-\frac{3}{8}} (1 + 2^{2j+2k_{1}})^{-5} \\ &\lesssim \sum_{j \geq -k} 2^{-\sigma k} 2^{2j+2k} (1 + 2^{2k+2j})^{-5} b_{k} b_{k}(\sigma) + \sum_{j \leq -k} 2^{-\sigma k} 2^{(k+j)+\delta|k+j|} b_{k} b_{k}(\sigma) \\ &\lesssim b_{k} b_{k}(\sigma) 2^{-\sigma k}, \end{split}$$

where we used $(1 + 2^{k+j})^{-1}T_{k,j} \le 2^{-k}$ for all $k \in \mathbb{Z}$. In the Low × High part, Lemma 8.4 shows

$$\sum_{k_2-k|\leq 4, k_1\leq k-4} \left\| P_k(P_{k_1}fP_{k_2}g) \right\|_{L^4_{t,x}} \lesssim \sum_{l\leq k} 2^l \mu_l \nu_k,$$

where $v_k = \sum_{|k'-k| \le 20} \|P_{k'}f\|_{S_k^{\omega}}$, $v_k = \sum_{|k'-k| \le 20} \|P_{k'}g\|_{L^4_{t,x}}$. Then by (4.64) we have

$$\begin{split} &\sum_{\substack{|k_2-k|\leq 4,k_1\leq k-4}} \left\| P_k \left[P_{k_1}(\phi_s \diamond \phi_i) P_{k_2} \widetilde{\mathcal{G}} \right] \right\|_{L^4_{t,x}} \\ &\lesssim 2^{-\sigma k} b_k(\sigma) \varepsilon T_{k,j} (1+2^{j+k})^{-7} \sum_{l\leq k} 2^l b_l b_l \left(1_{l+j\leq 0} 2^{2\delta|l+j|} 2^{l-j} + 1_{j+l\geq 0} 2^{2l} 2^{-\frac{3}{4}(j+l)} (1+2^{2j+2l})^{-3} \right) \\ &\lesssim 2^{-\sigma k} b_k(\sigma) b_k \left(1_{k+j\geq 0} 2^{-2j} (1+2^{k+j})^{-4} + 1_{k+j\leq 0} 2^{k-j} 2^{\delta|k+j|} \right). \end{split}$$

Hence we conclude for the Low × High part

$$\begin{split} &\int_0^\infty \sum_{\substack{|k_2-k| \le 4, k_1 \le k-4 \\ j \in \mathbb{Z}}} \left\| P_k \left[P_{k_1}(\phi_s \diamond \phi_i) P_{k_2} \widetilde{\mathcal{G}} \right] \right\|_{L^4_{t,x}} ds \\ &\lesssim 2^{-\sigma k} \sum_{j \in \mathbb{Z}} b_k b_k(\sigma) \mathbf{1}_{k+j \ge 0} (1+2^{k+j})^{-4} + 2^{-\sigma k} \sum_{j \in \mathbb{Z}} b_k b_k(\sigma) \mathbf{1}_{k+j \le 0} 2^{k+j} 2^{\delta|k+j} \\ &\lesssim b_k b_k(\sigma) 2^{-\sigma k}. \end{split}$$

Therefore, we get

$$\|P_k(A_i(0))\|_{L^4_{t,x}} \lesssim 2^{-\sigma k} b_k(\sigma).$$

Now we turn to the bounds for ϕ_t stated in Proposition 4.1.

Lemma 4.6. Assume that the assumptions of Proposition 4.1 and (4.9) hold. Then for $\sigma \in [0, \frac{99}{100}]$, one has

$$\|P_k\phi_t(s)\|_{L^4_{t,x}} \lesssim b_k(\sigma)2^{-(\sigma-1)k}(1+2^{2k}s)^{-2}.$$
(4.65)

Proof. Recall that ϕ_t satisfies

$$\begin{aligned} \partial_s \phi_t - \Delta \phi_t &= L(\phi_t) \\ L(\phi_t) &= L_1(\phi_t) + L_2(\phi_t) \\ L_1(\phi_t) &= \sum_{i=1}^2 2\partial_i (A_i \phi_t) + (\sum_{l=1}^2 A_l^2 - \partial_l A_l) \phi_t \\ L_2(\phi_t) &= \sum_{i=1}^2 (\phi_t \diamond \phi_i) \diamond \phi_i \mathcal{G}. \end{aligned}$$

By Duhamel principle, ϕ_t can be written as

$$\phi_t = e^{s\Delta}\phi_t(\uparrow_{s=0}) + \int_0^s e^{(s-\tau)\Delta} L(\phi_t(\tau))d\tau.$$
(4.66)

By a uniqueness argument as [[4], Lemma 5.6], in order to prove (4.65), it suffices to show

$$\|P_k\phi_t(s)\|_{L^4_{t,x}} \lesssim b_k(\sigma)2^{-(\sigma-1)k}(1+2^{2k}s)^{-2}$$
(4.67)

$$\int_{0}^{s} \|e^{(s-\tau)\Delta} L(\phi_{t}(\tau))\|_{L^{4}_{t,x}} d\tau \leq \varepsilon^{2} b_{k}(\sigma) 2^{-(\sigma-1)k} (1+2^{2k}s)^{-2}.$$
(4.68)

The $L_1(\phi_t)$ part of (4.68) has been done in [[4], Lemma 5.6]. It suffices to prove (4.68) for $L_2(\phi_t)$ under the assumption of (4.67). Recall also that $\mathcal{G} = \Gamma^{\infty} + \widetilde{\mathcal{G}}$ satisfies

$$\|P_k(\mathcal{G} - \Gamma^{\infty})\|_{F_k(T)} \lesssim 2^{-\sigma k} (1 + 2^{j+k})^{-7} T_{k,j}.$$
(4.69)

By the proof of [[4], Lemma 5.6],

$$\|P_k(\phi_l(s) \diamond \phi_i \diamond \phi_l)\|_{L^4_{l,x}} \lesssim b_k^2 2^{-(\sigma-3)k} (1+2^{2k}s)^{-2} (s2^{2k})^{-\frac{7}{8}} b_k(\sigma).$$
(4.70)

 \implies

Then the Γ^{∞} part of $L_2(\phi_l)$ follows directly by proof of [[4], Lemma 5.6]. Denote $\mathbf{P} = \phi_l(s) \diamond \phi_i \diamond \phi_l$. In order to control $\mathbf{P}(\mathcal{G} - \Gamma^{\infty})$, we first control $\|(\phi_i \diamond \phi_l)\widetilde{\mathcal{G}}\|_{S_k^{\frac{1}{2}}(T)}$. We have seen

$$\|P_{k}(\phi_{i} \diamond \phi_{l})\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} \lesssim 2^{-\sigma k} (1 + 2^{2k+2j})^{-4} 2^{-j} b_{-j} b_{\max(k,-j)}(\sigma).$$
(4.71)

Thus applying bilinear Littlewood-Paley decomposition, we obtain by (4.69) that

$$\begin{split} \|P_{k}(\phi_{i} \diamond \phi_{l}\widetilde{\mathcal{G}})\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} &\lesssim \sum_{|k_{1}-k| \leq 4} \|P_{k_{1}}(\phi_{i} \diamond \phi_{l})\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} \|P_{\leq k-4}\widetilde{\mathcal{G}}\|_{L^{\infty}} \\ &+ \sum_{|k_{1}-k_{2}| \leq 8, k_{1}, k_{2} \geq k-4} 2^{\frac{k+k_{1}}{2}} \|P_{k_{1}}(\phi_{i} \diamond \phi_{l})\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} \|P_{k_{2}}\widetilde{\mathcal{G}}\|_{F_{k_{2}}(T)} \\ &+ \sum_{|k_{2}-k| \leq 4, k_{1} \leq k-4} 2^{k_{1}} \|P_{k_{1}}(\phi_{i} \diamond \phi_{l})\|_{F_{k}(T) \cap S_{k}^{\frac{1}{2}}(T)} \|P_{k_{2}}\widetilde{\mathcal{G}}\|_{F_{k_{2}}(T)} \\ &\lesssim 2^{-\sigma k} b_{k}(\sigma) b_{k} 2^{\delta |k+j|} \Big(1_{k+j \leq 0} 2^{-j} + 2^{k} 1_{k+j \geq 0} (1+2^{k+j})^{-7} \Big). \end{split}$$

Then using Lemma 8.4 with $\omega = \frac{1}{2}$ and (4.69), $\mathbf{P}\widetilde{\mathcal{G}}$ is dominated by

$$\begin{split} \|P_k \left(P_{k_1} \phi_l P_{k_2}(\phi_i \diamond \phi_l \widetilde{\mathcal{G}}) \right)\|_{L^4_{l,x}} &\lesssim \sum_{|k_1 - k| \le 4, k_2 \le k - 4} 2^{k_2} \|P_{k_1} \phi_l\|_{L^4_{l,x}} \|P_{k_2}(\phi_i \diamond \phi_l \widetilde{\mathcal{G}})\|_{S^{\frac{1}{2}}_{k}(T)} \\ &+ 2^k \sum_{|k_1 - k_2| \le 8, k_2, k_1 \ge k - 4} 2^{-\frac{1}{2}(k_1 - k)} \|P_{k_1} \phi_l\|_{L^4_{l,x}} \|P_{k_2}(\phi_i \diamond \phi_l \widetilde{\mathcal{G}})\|_{S^{\frac{1}{2}}_{k}(T)} \\ &+ \sum_{|k_2 - k| \le 4, k_1 \le k - 4} 2^{\frac{k + k_1}{2}} \|P_{k_1} \phi_l\|_{L^4_{l,x}} \|P_{k_2}(\phi_i \diamond \phi_l \widetilde{\mathcal{G}})\|_{S^{\frac{1}{2}}_{k}(T)} \\ &\lesssim 2^{-\sigma k} b_k(\sigma) b_k 2^{\delta |k+j|} \mathbf{1}_{k+j \le 0} \left(2^{\frac{3}{2}k - \frac{3}{2}j} + 2^{2k - j} \right) \\ &+ 2^{-\sigma k} b_k(\sigma) b_k \mathbf{1}_{k+j \ge 0} \left(2^{\delta |k+j|} 2^{3k} (1 + 2^{k+j})^{-10} + 2^{\frac{3}{2}(k-j)} (1 + 2^{k+j})^{-7} + 2^{k-2j} (1 + 2^{k+j})^{-4} \right) \end{split}$$

for $s \in [2^{2j-1}, 2^{2j+1})$, $j, k \in \mathbb{Z}$. As a summary, inserting this bound to the following heat estimates

$$\int_0^{\widetilde{s}} \left\| e^{(\widetilde{s}-s)\Delta} P_k[\phi_t \diamond \phi_i \diamond \phi_l \widetilde{\mathcal{G}}] \right\|_{L^4_{t,x}} ds \lesssim \int_0^{\widetilde{s}} (1+|\widetilde{s}-s|)^{-N}(...) ds,$$

we conclude that

$$\left\|P_k\int_0^s e^{(s-\tau)\Delta}L_2(\phi_t(\tau))d\tau\right\|_{L^4_{t,x}} \lesssim \varepsilon(1+2^{2k}s)^{-2}2^{-\sigma k+k}b_k(\sigma).$$

Thus since L_1 has been done before, we finish our proof.

Lemma 4.7. With the assumptions of Proposition 4.1, the bootstrap assumption (4.9) can be improved to be

$$2^{\frac{1}{2}k} \| P_k \widetilde{\mathcal{G}}^{(1)} \|_{L^4_x L^\infty_t} \lesssim h_k (\mathbf{1}_{k+j \ge 0} (1+s2^{2k})^{-20} + 2^{\delta |k+j|} \mathbf{1}_{k+j \le 0}),$$

for any $k, j \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1})$.

Proof. The proof of Lemma 4.1 has shown for any $\sigma \in [0, \frac{99}{100}]$, $k, j \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1})$,

$$\|P_k(\mathcal{G}D_i\phi_i)\|_{L^4\cap L^\infty_i L^2_x} \lesssim h_k(\sigma) 2^{-\sigma k+k} (1+s2^{2k})^{-3} 1_{j+k\geq 0} + 2^{-j} 2^{\delta|k+j|} 1_{k+j\leq 0}.$$
(4.72)

Meanwhile, Lemma 4.6 yields

$$\|P_k\phi_t\|_{L^4} \lesssim b_k 2^k (1+s2^{2k})^{-2}. \tag{4.73}$$

Recall that $b_k \leq \varepsilon^{\frac{1}{2}}$, for any $k \in \mathbb{Z}$. Then bilinear Littlewood-Paley decomposition shows

$$\|P_k(\phi_t(D_i\phi_i\mathcal{G}))\|_{L^4} \lesssim h_k(\sigma)2^{-\sigma k+3k}(1+s2^{2k})^{-2}1_{k+j\geq 0} + 2^{-2j}2^{k-\sigma k}2^{2\delta|k+j|}h_k(\sigma)h_k1_{k+j\leq 0}$$

for i = 1, 2, any $\sigma \in [0, \frac{99}{100}]$, $k, j \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1})$. Here, in the High × Low interaction of $\phi_t(D_i\phi_i\mathcal{G})$ we use

$$\sum_{\substack{|k_1-k| \le 4, k_2 \le k-4 \\ \le b_k 2^k \sum_{k_2 \le k-4} h_{k_2}(\sigma) 2^{-\sigma k_2 + 2k_2} \left((1 + s 2^{2k_2})^{-3} \mathbf{1}_{j+k_2 \ge 0} + 2^{-j} 2^{\delta |k_2+j|} \mathbf{1}_{k_2+j \le 0} \right)}$$

The other two frequency interactions are standard. Thus

$$\int_{s}^{\infty} \|P_{k}(\phi_{t}D_{i}\phi_{i})\mathcal{G}\|_{L^{4}}ds' \leq h_{k}(\sigma)2^{-\sigma k+k}(1+2^{2k+2k_{0}})^{-1}\mathbf{1}_{k+k_{0}\geq0} + 2^{k-\sigma k}h_{k}(\sigma)(1+2^{2\delta|k+k_{0}|}h_{k})\mathbf{1}_{k+k_{0}\leq0}$$

$$(4.74)$$

for any $\sigma \in [0, \frac{99}{100}], k, k_0 \in \mathbb{Z}, s \in [2^{2k_0-1}, 2^{2k_0+1})$. Recall that

$$A_t = \int_s^\infty (\phi_t \diamond \phi_s) \mathcal{G} ds'; \ \phi_s = \sum_{i=1,2} D_i \phi_i.$$

we see $||P_kA_t||_{L^4}$ is bounded by the RHS of (4.74). By the schematic formula

$$\partial_t(\widetilde{\mathcal{G}}^{(1)}) = \phi_t \mathcal{G}^{(2)} + A_t \mathcal{G}^{(1)}$$

and the bounds

$$\|P_k(\widetilde{\mathcal{G}}^{(l)})\|_{L^4 \cap L^\infty_t L^2_x} \lesssim (1 + s2^{2k})^{-30} 2^{-\sigma k - k} h_k(\sigma), \; \forall \; l = 1, 2,$$

we deduce from bilinear Littlewood-Paley decomposition that

$$\|P_k \partial_t (\widetilde{\mathcal{G}}^{(1)})\|_{L^4} \lesssim h_k 2^k (1 + 2^{2\delta |k+k_0|} h_k 1_{k+k_0 \le 0}).$$

Then by Gagliardo-Nirenberg inequality we get

$$2^{\frac{1}{2}k} \|P_k(\widetilde{\mathcal{G}}^{(1)})\|_{L^4_x L^\infty_t} \lesssim \|P_k(\widetilde{\mathcal{G}}^{(1)})\|_{L^4}^{\frac{1}{4}} \|\partial_t P_k(\widetilde{\mathcal{G}}^{(1)})\|_{L^4}^{\frac{1}{4}} \\ \lesssim h_k (1+s2^{2k})^{-20} \mathbf{1}_{k+k_0 \ge 0} + h_k 2^{\delta|k+k_0|} \mathbf{1}_{k+k_0 \le 0}.$$

for $k_0, k \in \mathbb{Z}$, $s \in [2^{2k_0-1}, 2^{2k_0+1})$.

4.4 End of proof of Proposition 4.1

By Lemma 4.7, the assumption (4.9) in Lemma 4.4, Lemma 4.5 and Lemma 4.6 can be dropped. In fact, let

$$\widetilde{\Phi}(T') := \sup_{k,j \in \mathbb{Z}} \sup_{s \in [2^{2j-1}, 2^{2j+1})} h_k^{-1} \left(\mathbf{1}_{k+j \ge 0} (1 + s2^{2k})^{-20} + 2^{\delta|k+j|} \mathbf{1}_{k+j \le 0} \right)^{-1} 2^{\frac{1}{2}k} \| P_k \widetilde{\mathcal{G}}^{(1)} \|_{L_x^4 L_t^\infty(T')}$$

Lemma 3.3 and Sobolev embeddings imply $\widetilde{\Phi}$ is an increasing continuous function on $T' \in [0, T]$. Lemma 4.7 shows $\widetilde{\Phi}(T') \le \varepsilon^{-\frac{1}{4}} \Longrightarrow \widetilde{\Phi}(T') \le 1$. Then by Bernstein inequality and letting $T' \to 0$, it remains to prove

$$\|P_k\widetilde{\mathcal{G}}^{(1)}\|_{L^2_x} \lesssim h_k \left(\mathbf{1}_{k+j \ge 0} (1+s2^{2k})^{-20} + 2^{\delta|k+j|} \mathbf{1}_{k+j \le 0} \right)$$

along the heat flow initiated from u_0 for any $k, j \in \mathbb{Z}, s \in [2^{2j-1}, 2^{2j+1})$. This follows by (3.76).

By Lemma 3.3 and similar arguments, assumption (4.3) can be also dropped. Thus Lemma 4.4, Lemma 4.5 and Lemma 4.6 all hold with only assuming (4.1), (4.2) of Proposition 4.1. The left for Proposition 4.1 is to prove the $L_{t,x}^2$ bound for A_t .

Lemma 4.8. With the assumptions (4.1), (4.2) of Proposition 4.1, for all $k \in \mathbb{Z}$, one has

$$\|P_k A_t\|_{s=0} \|_{L^2_{t,x}} \lesssim \varepsilon b_k(\sigma) 2^{-\sigma k} \text{ if } \sigma \in [\frac{1}{100}, \frac{99}{100}]$$
(4.75)

$$\|A_t \upharpoonright_{s=0}\|_{L^2_{t,x}} \lesssim \varepsilon^2 \text{ if } \sigma \in [0, \frac{99}{100}].$$

$$(4.76)$$

Proof. Recall that [[4], Lemma 5.7] has proved

$$\|P_{k}(\phi_{l} \diamond \phi_{s})\|_{L^{2}_{l,x}} \lesssim \sum_{l \le k} 2^{-\sigma l} 2^{l+k} b_{k}(\sigma) b_{l}(s 2^{2l})^{-\frac{3}{8}} (1 + s 2^{2k})^{-2} + \sum_{l \ge k} 2^{-\sigma l} 2^{2l} b_{l}(\sigma) b_{l}(s 2^{2l})^{-\frac{3}{8}} (1 + s 2^{2l})^{-4}.$$
(4.77)

Denote the RHS of (4.77) by $\mathbf{a}_k(\sigma)$ for simplicity. Since $A_t(0) = \int_0^{\infty} (\phi_t \diamond \phi_s) \mathcal{G} ds$, (4.76) follows by directly applying [[4], Lemma 5.7] since $\|\mathcal{G}\|_{L^{\infty}} \leq K(\mathcal{N})$. For (4.75), we need to clarify the frequency interaction between $(\phi_t \diamond \phi_s)$ and \mathcal{G} as before. The constant part of \mathcal{G} follows by (4.77). It remains to deal with the $\widetilde{\mathcal{G}}$ part. In the High × Low part of $P_k[(\phi_t \diamond \phi_s)\widetilde{\mathcal{G}}]$, we have

$$\sum_{|k_1-k|\leq 4} \|P_{k_1}(\phi_t \diamond \phi_s)P_{\leq k-4}\widetilde{\mathcal{G}}\|_{L^2_{t,x}} \lesssim \sum_{|k_1-k|\leq 4} \|P_{k_1}(\phi_t \diamond \phi_s)\|_{L^2_{t,x}} \|\widetilde{\mathcal{G}}\|_{L^\infty} \lesssim \mathbf{a}_k(\sigma).$$

Then the High × Low part is acceptable by directly repeating [[4], Lemma 5.7]. From now on to the end of this lemma, we assume $\sigma \in [\frac{1}{100}, \frac{99}{100}]$. In the High×High part of $P_k[(\phi_t \diamond \phi_s)\widetilde{\mathcal{G}}]$, by $||P_k f||_{L^{\infty}} \leq 2^k ||f||_{F_k}$ and (4.64), we have

$$\sum_{\substack{|k_1-k_2| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_2| \le 8, k_1, k_2 \ge k-4 \\ \le \sum_{\substack{|k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ = \sum_{\substack{|k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ \le \sum_{\substack{|k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ = \sum_{\substack{|k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ \le \sum_{\substack{|k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ = \sum_{\substack{|k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ = \sum_{\substack{|k_1-k_1| \le 8, k_1, k_2 \ge k-4 \\ |k_1-k_1| \le 8, k_1 \le k-4 \\ |k_1-$$

$$+\sum_{k_1 \ge k-4}^{k_1 \ge k-4} 2^{-\sigma k_1} b_{k_1}(\sigma) (1+s2^{2k_1})^{-3} \sum_{l \ge k_1}^{l_2 \le l} 2^{2l} b_l b_l (s2^{2l})^{-\frac{3}{8}} (1+s2^{2l})^{-4}.$$
(4.79)

Thus for $j \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1})$, when $k + j \ge 0$, the above formula is bounded by

$$(1+2^{2j+2k})^{-2}2^{2k-\sigma k}b_kb_k(\sigma)(2^{2k+2j})^{-\frac{3}{8}}.$$

When $k + j \le 0$, by (4.78), (4.79), the High × High part is dominated by

$$\begin{split} &(\sum_{k_{1}\geq -j}+\sum_{k-4\leq k_{1}\leq -j})(1+2^{2k_{1}+2j})^{-3}2^{2k_{1}-\sigma k_{1}}b_{k_{1}}^{2}b_{k_{1}}(\sigma)2^{-\frac{3}{4}(k_{1}+j)}\\ &+\sum_{k_{1}\geq k-4}1_{k_{1}+j\geq 0}(1+2^{2k_{1}+2j})^{-3}2^{-\sigma k_{1}}b_{k_{1}}(\sigma)[\sum_{l\geq k_{1}}2^{2l}b_{l}b_{l}2^{-\frac{3}{4}(j+l)}(1+2^{2l+2j})^{-4}]\\ &+\sum_{k_{1}\geq k-4}1_{k_{1}+j\leq 0}(1+2^{2k_{1}+2j})^{-3}2^{-\sigma k_{1}}b_{k_{1}}(\sigma)[(\sum_{l\geq -j}+\sum_{k_{1}\leq l\leq -j})2^{2l}b_{l}b_{l}2^{-\frac{3}{4}(j+l)}(1+2^{2l+2j})^{-4}], \end{split}$$

which is further bounded by

$$\begin{split} &\sum_{k_1 \ge -j} b_{k_1}(\sigma) b_{-j}^2 2^{2\delta|k_1+j|} 2^{2k_1 - \sigma k_1} 2^{-\frac{3}{4}(k_1+j)} 2^{-6(k_1+j)} + \sum_{k \le k_1 \le -j} b_{k_1}(\sigma) b_{-j}^2 2^{-\sigma k_1} 2^{2\delta|k_1+j|} 2^{2k_1} 2^{-\frac{3}{4}(k_1+j)} \\ &+ b_{-j}^2 b_{-j}(\sigma) \sum_{k_1 \ge k-4} \mathbf{1}_{k_1+j \ge 0} (1 + 2^{2k_1+2j})^{-7} 2^{2k_1 - \sigma k_1} 2^{\delta|k_1+j|} 2^{-\frac{3}{4}(j+k_1)} \\ &+ \sum_{k_1 \ge k-4} \mathbf{1}_{k_1+j \le 0} b_{-j}^2 b_{k_1}(\sigma) [2^{-\sigma k_1} 2^{-2j} + 2^{-\sigma k_1} \sum_{k_1 \le l \le -j} 2^{2l} 2^{\delta|l+j|} 2^{-\frac{3}{4}(j+l)}] \\ &\lesssim b_{-j}^2 b_{-j}(\sigma) 2^{\sigma j} 2^{-2j} + \sum_{k_1 \ge k-4} \mathbf{1}_{k_1+j \le 0} b_{-j}^2 b_{k_1}(\sigma) 2^{-\sigma k_1} 2^{-2j} \\ &\lesssim b_{-j}^2 b_{-j}(\sigma) 2^{\sigma j} 2^{-2j} + 2^{-\sigma k} b_k(\sigma) b_{-j}^2 2^{-2j}, \end{split}$$

where in the last line we used $\sigma \ge \frac{1}{100}$. Summing over $j \ge k_0$ we see the High × High part satisfies

$$\begin{split} &\int_{0}^{\infty} \sum_{\substack{|k_{1}-k_{2}| \leq 8, k_{1}, k_{2} \geq k-4 \\ \leq \sum_{j \leq -k} 2^{\sigma j} b_{-j}(\sigma) b_{-j}^{2} + \sum_{j \leq -k} b_{-j}^{2} 2^{-\sigma k} b_{k}(\sigma) + \sum_{j \geq -k} (1 + 2^{2j+2k})^{-2} 2^{2k+2j-\sigma k} b_{k} b_{k}(\sigma) (2^{2k+2j})^{-\frac{3}{8}} \\ &\lesssim \varepsilon^{2} 2^{-\sigma k} b_{k}(\sigma), \end{split}$$

where we applied $\sigma \ge \frac{1}{100}$ in the last line again. Now let us consider the Low × High part of $P_k[(\phi_t \diamond \phi_s)\widetilde{\mathcal{G}}]$. By the same reason as High × High, the Low × High part is dominated by

$$\begin{split} \int_0^\infty \sum_{|k_2-k| \leq 4} \|P_{\leq k-4}(\phi_t \diamond \phi_s) P_{k_2} \widetilde{\mathcal{G}}\|_{L^2_{t,x}} ds' \lesssim \int_0^\infty \sum_{|k_2-k| \leq 4} \|P_{\leq k-4}(\phi_t \diamond \phi_s)\|_{L^2_{t,x}} 2^{k_2} \|P_{k_2} \widetilde{\mathcal{G}}\|_{L^\infty_{t} L^2_x} ds' \\ \lesssim b_k(\sigma) 2^{-\sigma k} \int_0^\infty \|(\phi_t \diamond \phi_s)\|_{L^2_{t,x}} ds' \lesssim b_k(\sigma) 2^{-\sigma k} \varepsilon^2, \end{split}$$

where we applied (4.64) and $2^k T_{k,j}(1+2^{j+k})^{-1} \leq 1$ in the third inequality.

5 Evolution along the Schrödinger map flow direction

In this section, we prove the following proposition, which is the key to close the bootstrap for solutions in the Schrödinger evolution direction.

Proposition 5.1. Assume that $\sigma \in [0, \frac{99}{100}]$. Let $Q \in N$ be a fixed point and ϵ_0 be a sufficiently small constant. Given any $\mathcal{L} \in \mathbb{Z}_+$, assume that $T \in (0, 2^{2\mathcal{L}}]$. Let $\{c_k\}$ be an ϵ_0 -frequency envelope of order δ . And let $\{c_k(\sigma)\}$ be another frequency envelope of order δ . Let $u \in \mathcal{H}_Q(T)$ be the solution to SMF with initial data u_0 which satisfies

$$\|P_k \nabla u_0\|_{L^2_{\infty}} \le c_k \tag{5.1}$$

$$\|P_k \nabla u_0\|_{L^2} \le c_k(\sigma) 2^{-\sigma k} \tag{5.2}$$

Denote $\{\phi_i\}$ the corresponding differential fields of the heat flow initiated from u. Suppose also that at the heat initial time s = 0,

$$\|P_k\phi_i\|_{G_k(T)} \le \epsilon_0^{-\frac{1}{2}} c_k.$$
(5.3)

Then when s = 0, we have for all $i = 1, 2, k \in \mathbb{Z}$,

$$\|P_k\phi_i\|_{G_k(T)} \lesssim c_k \tag{5.4}$$

$$\|P_k\phi_i\|_{G_k(T)} \leq c_k(\sigma)2^{-\sigma k}.$$
(5.5)

The proof of Proposition 5.1 will be divided into several lemmas. First of all, Corollary 3.1 shows

$$\sum_{i=1}^{2} \|P_k \phi_i(\uparrow_{s=0,t=0})\|_{L^2_x} \leq 2^{-\sigma k} c_k(\sigma)$$
(5.6)

for any $k \in \mathbb{Z}$, $\sigma \in [0, \frac{99}{100}]$.

Second, we reduce the proof to frequency envelope bounds. Let

$$b(k) = \sum_{i=1}^{2} \|P_k \phi_i\|_{s=0} \|_{G_k(T)}.$$
(5.7)

For $\sigma \in [0, \frac{99}{100}]$, define the frequency envelopes:

$$b_{k}(\sigma) = \sup_{k' \in \mathbb{Z}} 2^{\sigma k'} 2^{-\delta |k-k'|} b(k').$$
(5.8)

By Proposition 3.1 and Sobolev embeddings, they are finite and ℓ^2 summable. And

$$\|P_k\phi_i\|_{s=0} \|_{G_k(T)} \lesssim 2^{-\sigma k} b_k(\sigma).$$

$$(5.9)$$

To prove (5.4) and (5.5), it suffices to show

$$b_k(\sigma) \leq c_k(\sigma).$$
 (5.10)

By (4.3), we have $b_k \leq \varepsilon_0^{-\frac{1}{2}} c_k$, and particularly,

$$\sum_{k\in\mathbb{Z}}b_k^2 \le \epsilon_0.$$
(5.11)

The assumption (4.1) of Proposition 4.1 follows from the inclusion $G_k \subset F_k$. The following lemma will show the assumption (4.2) holds as a corollary of (5.9) if *u* solves SMF.

Lemma 5.1. If $\{b_k(\sigma)\}$ are defined above. Then the field ϕ_t at heat initial time s = 0 satisfies

$$\|P_k\phi_t\|_{s=0} \|_{L^4_{t,s}} \lesssim b_k(\sigma) 2^{-(\sigma-1)k}.$$
(5.12)

Proof. When s = 0, $\phi_t(0) = \sqrt{-1} \sum_{i=1}^d \partial_i \phi_i(0) + A_i(0) \phi_i(0)$. The terms $\psi_i(0), A_i(0)$ are estimated before in Section 4. Thus copying the proof of [[4],Lemma 6.1] gives (5.12).

Thus both the assumption (4.1) and the assumption (4.2) of Proposition 4.1 are verified. Now one can apply Proposition 4.1, since (4.3) can be dropped. We summarize the results in the following:

$$\begin{cases} \|P_k(\phi_i(s))\|_{F_k(T)} \leq 2^{-\sigma k} b_k(\sigma)(1+2^{2k}s)^{-4}, \\ \|P_k(D_i\phi_i(s))\|_{F_k(T)} \leq 2^k 2^{-\sigma k} b_k(\sigma)(s2^{2k})^{-\frac{3}{8}}(1+2^{2k}s)^{-2}, \end{cases}$$
(5.13)

and for $F \in \{\psi_i \diamond \psi_j, A_l^2\}_{l,i,j=1}^2 \upharpoonright_{s=0}$

$$\|P_k F\|_{L^2_{tx}} \lesssim 2^{-\sigma k} b^2_{>k}(\sigma), \ \|F\|_{L^2_{tx}} \lesssim \epsilon_0.$$
(5.14)

The at $s = 0, A_t$ satisfies

$$\begin{aligned} \|A_t(0)\|_{L^2_{t,x}} &\leq \epsilon_0, \text{ if } \sigma \in [0, \frac{99}{100}] \\ \|P_k A_t(0)\|_{L^2_{t,x}} &\leq 2^{-\sigma k} b_k(\sigma), \text{ if } \sigma \in [\frac{1}{100}, \frac{99}{100}] \end{aligned}$$

Recall that when s = 0 the evolution equation of ϕ_i along the Schrödinger map flow direction (see Lemma 1.1) is :

$$-\sqrt{-1}D_t\phi_i = \sum_{j=1}^2 D_j D_j\phi_i + \sum_{j=1}^2 \mathcal{R}(\phi_i, \phi_j)\phi_j.$$
(5.15)

5.1 Control of nonlinearities

Now let us deal with the nonlinearities in (5.15). In this section we always assume s = 0. Denote

$$L'_{j} = A_{t}\phi_{j} + \sum_{i=1}^{2} A_{i}^{2}\phi_{j} + 2\sum_{i=1}^{2} \partial_{i}(A_{i}\phi_{j}) - \sum_{i=1}^{2} (\partial_{i}A_{i})\phi_{j}.$$
(5.16)

Proposition 5.2 ([4]). *For all* $j \in \{1, 2\}$ *and* $\sigma \in [0, \frac{99}{100}]$ *we have*

$$\|P_k(L'_j)\|_{s=0} \|_{N_k(T)} \leq \epsilon_0 2^{-\sigma k} b_k(\sigma)$$
(5.17)

$$\sum_{j_0,j_1,j_3=1}^2 \|P_k\left(\phi_{j_0} \diamond \phi_{j_1} \diamond \phi_{j_3}\right)|_{s=0}\|_{N_k(T)} \lesssim \epsilon_0 2^{-\sigma k} b_k(\sigma).$$
(5.18)

Proof. (5.17) and (5.18) have been proved in [[4], Proposition 6.2]. We emphasize that to bound $||A_t\phi_i||_{N_k}$, [[4], Proposition 6.2] used $||A_t||_{L^2_{t,x}} \leq \varepsilon^2$ when $\sigma \in [0, \frac{1}{12}]$ and $||P_kA_t||_{L^2_{t,x}} \leq 2^{-\sigma k}b_k(\sigma)$ when $\sigma \geq \frac{1}{12}$. Thus our bounds (4.75), (4.76) suffice to bound $||A_t\phi_i||_{N_k}$ as well although (4.75)-(4.76) itself differs from the bounds stated by [[4], Lemma 5.7].

Now we turn to the remained curvature term in (5.15).

Proposition 5.3. *For all* $k \in \mathbb{Z}$ *and* $\sigma \in [0, \frac{99}{100}]$ *we have*

$$\sum_{j_0,j_1,j_3=1}^2 \|P_k\left((\phi_{j_0} \diamond \phi_{j_1} \diamond \phi_{j_3})\mathcal{G}\right)\|_{N_k(T)} \lesssim 2^{-\sigma k} \epsilon_0 b_k(\sigma).$$
(5.19)

Proof. Recall $\mathcal{G} = \Gamma^{\infty} + \widetilde{\mathcal{G}}$. The constant part Γ^{∞} satisfies (5.19) by directly applying (5.18). It suffices to control $\widetilde{\mathcal{G}}$ part. As a preparation, we first prove the following estimate

$$\sum_{i=1}^{2} \|P_{k}(\widetilde{\mathcal{G}}\phi_{i})\|_{F_{k}(T)} \lesssim \begin{cases} 2^{-\sigma k}b_{k}(\sigma), & \frac{1}{100} < \sigma \le \frac{99}{100} \\ 2^{-\sigma k}\sum_{j \ge k}b_{j}b_{j}(\sigma), & 0 \le \sigma \le \frac{1}{100} \end{cases}$$
(5.20)

This follows directly by applying Corollary 4.1 and [Lemma 5.1, [4]]: If $\sigma > \frac{1}{100}$, then

$$\begin{split} \|P_k(\widetilde{\mathcal{G}}\phi_l)\|_{F_k(T)} &\lesssim 2^{-\sigma k} b_k(\sigma) + 2^{-k-\sigma k} b_k(\sigma) \sum_{l \leq k} 2^{\delta |k-l|} 2^l b_l + b_k(\sigma) \sum_{j \geq k} 2^{-\sigma j} 2^{2\delta |k-j|} \\ &\lesssim 2^{-\sigma k} b_k(\sigma). \end{split}$$

If $\sigma \in [0, \frac{1}{100}]$, for the High × High interaction we directly use

$$\begin{split} \sum_{|k_1-k_2| \le 8, k_1, k_2 \ge k-4} \|P_k(P_{k_1} \widetilde{\mathcal{G}} P_{k_2} \phi_i)\|_{F_k(T)} &\lesssim \sum_{j \ge k-4} 2^j (\sum_{|k_1-j| \le 28} \|P_{k_1} \widetilde{\mathcal{G}}\|_{F_{k_1}(T)}) (\sum_{|k_2-j| \le 28} \|P_{k_2} \phi_i\|_{F_{k_2}(T)}) \\ &\lesssim 2^{-\sigma k} \sum_{j \ge k} b_j b_j(\sigma). \end{split}$$

The other two interactions are all the same as $\sigma \ge \frac{1}{100}$. Thus (5.20) follows. As before, denoting $\mathbf{F} = \phi_{j_0} \diamond \phi_{j_1}$, by bilinear Littlewood-Paley decomposition, we have

$$\begin{split} \|P_{k}\left(\mathbf{F} \diamond (\phi_{j_{3}}\mathcal{G})\right)\|_{N_{k}(T)} \\ &= \sum_{|l-k| \leq 4} \|P_{k}(P_{< k-100}\mathbf{F}P_{l}(\widetilde{\mathcal{G}}\phi_{j_{3}}))\|_{N_{k}(T)} + \sum_{|k_{1}-k| \leq 4} \|P_{k}(P_{k_{1}}\mathbf{F}P_{< k-100}(\widetilde{\mathcal{G}}\phi_{j_{3}}))\|_{N_{k}(T)} \\ &+ \sum_{k_{1},k_{2} \geq k-100}^{|k_{1}-k_{2}| \leq 120} \|P_{k}(P_{k_{1}}\mathbf{F}P_{k_{2}}(\widetilde{\mathcal{G}}\phi_{j_{3}}))\|_{N_{k}(T)}. \end{split}$$
(5.21)

For the first RHS term of (5.21), applying (8.12) and the trivial bound

$$\|\widetilde{\mathcal{G}}\|_{L^{\infty}_{t,r}} \lesssim 1 \tag{5.22}$$

$$\|\phi_x\|_{L^4_{t,x}} \lesssim \epsilon_0, \tag{5.23}$$

and (5.20), for $\sigma \in [\frac{1}{100}, \frac{99}{100}]$ we get

$$\sum_{|k_0-k|\leq 4} \|P_k(P_{< k-100} \mathbf{F} P_{k_0}(\widetilde{\mathcal{G}}\phi_{j_3}))\|_{N_k(T)} \lesssim \|\phi_{j_0}\phi_{j_1}\|_{L^2_{t,x}} \|P_k(\widetilde{\mathcal{G}}\phi_{j_3})\|_{F_k(T)} \lesssim \epsilon_0 2^{-\sigma k} b_k(\sigma) b_k.$$

For the first RHS term of (5.21), when $\sigma \in [0, \frac{1}{100}]$, one further decomposes $P_{[k-4,k+4]}(\widetilde{\mathcal{G}}\phi_{j_3})$ into High × High, Low × High, High \times Low. We schematically write

$$\sum_{|k_0-k|\leq 4} \|P_k\left(P_{

$$\lesssim \sum_{|l-k|\leq 8} \|P_k\left((P_{
(5.24)$$$$

+
$$\sum_{|l-k| \le 8} ||P_k \left((P_{ (5.25)$$

+
$$\sum_{|k_1-k_2| \le 16, k_1, k_2 \ge k-8} ||P_k((P_{ (5.26)$$

Since for all $\sigma \in [0, \frac{99}{100}]$, the Low × High (denoted by P_k^{lh} for short) and High × Low (denoted by P_k^{hl} for short) interactions lead to $\|(P_k^{lh} + P_k^{hl})(\widetilde{\mathcal{G}}\phi_{j_3})\|_{F_k} \leq 2^{-\sigma k}b_k(\sigma)$, we conclude that

$$(5.24) + (5.25) \leq \|\phi_{j_0}\phi_{j_1}\|_{L^2_{t,x}} \left(\|P_k^{lh}(\mathcal{G}\phi_{j_3})\|_{F_k(T)} + \|P_k^{hl}(\mathcal{G}\phi_{j_3})\|_{F_k(T)} \right) \\ \leq \epsilon_0 2^{-\sigma k} b_k(\sigma).$$

For the (5.26) term, applying (8.14) yields

$$(5.26) \lesssim \sum_{k_{2} \ge k-8} \sum_{|k_{1}-k_{2}| \le 16} \left\| P_{k} \left[\left((P_{< k-100} \mathbf{F}) P_{k_{2}} \widetilde{\mathcal{G}} \right) P_{k_{1}} \phi_{j_{3}} \right] \right\|_{N_{k}} \\ \lesssim \sum_{k_{2} \ge k-8, |k_{1}-k_{2}| \le 16} \left\| (P_{< k-100} \mathbf{F}) P_{k_{2}} \widetilde{\mathcal{G}} \right\|_{L^{2}_{t,x}} 2^{\frac{k-k_{1}}{6}} \| P_{k_{1}} \phi_{j_{3}} \|_{G_{k_{1}}} \\ \lesssim \sum_{k_{1} \ge k-12} \left\| \mathbf{F} \right\|_{L^{2}_{t,x}} 2^{\frac{k-k_{1}}{6}} 2^{-\sigma k_{1}} b_{k_{1}}(\sigma) \\ \lesssim \epsilon_{0} 2^{-\sigma k} b_{k}(\sigma).$$

Thus the first RHS term of (5.21) has been done.

For the second RHS term of (5.21), we further divide **F** into

$$\sum_{\substack{|k_1-k|\leq 4\\ \leq}} \|P_k(P_{k_1} \mathbf{F} P_{< k-100}(\mathcal{G}\phi_{j_3}))\|_{N_k(T)}$$

$$\lesssim \sum_{\substack{|l-k|\leq 8\\ k-100}} \|P_k(P_l \phi_{j_0})(P_{\leq k-8} \phi_{j_1})P_{\leq k-100}(\mathcal{G}\phi_{j_3})\|_{N_k(T)}$$
(5.27)

$$+\sum_{|l-k|\leq 8} ||P_k(P_l\phi_{j_1})(P_{\leq k-8}\phi_{j_0})P_{\leq k-100}(\mathcal{G}\phi_{j_3})||_{N_k(T)}$$
(5.28)

+
$$\sum_{|l_1-l_2| \le 16, l_1, l_2 \ge k-8} \|P_k[(P_{l_1}\phi_{j_0})(P_{l_2}\phi_{j_1})P_{\le k-100}(\mathcal{G}\phi_{j_3})]\|_{N_k(T)}.$$
 (5.29)

For the first two terms on the RHS, using again (8.12) and the bounds (5.22), (5.23), we obtain

 $(5.28) + (5.27) \leq \|P_k(\phi_x)\|_{F_k(T)} \|\phi_x\|_{L^4_{t,x}}^2 \leq \epsilon_0 2^{-\sigma k} b_k(\sigma).$

And similarly, for $\sigma > \frac{1}{100}$,

$$(5.29) \lesssim \|\widetilde{\mathcal{G}}\phi_{j_3}\|_{L^4_{t,x}} \|\phi_{j_1}\|_{L^4_{t,x}} \sum_{l \ge k-8} \|P_l\phi_{j_0}\|_{F_l(T)} \lesssim \epsilon_0 2^{-\sigma k} b_k(\sigma).$$

For $0 \le \sigma \le \frac{1}{100}$, using again (8.14) and the bounds (5.22), (5.23), we have

$$(5.29) \lesssim \sum_{l \ge k} 2^{\frac{k-l}{6}} \left\| \left(P_{\le k-100}(\widetilde{\mathcal{G}}\phi_{j_3}) \right) P_l \phi_{j_1} \right\|_{L^2_{t,x}} \|P_l \phi_{j_0}\|_{G_l(T)}$$

$$\lesssim \|\phi_x\|^2_{L^4_{t,x}} \sum_{l \ge k} 2^{\delta |l-k|} 2^{\frac{k-l}{6}} 2^{-\sigma l} b_l b_l(\sigma) \lesssim 2^{-\sigma k} b_k(\sigma) \epsilon_0.$$

Thus the first two RHS terms of (5.21) are done.

For the third term of (5.21), applying Littlewood-Paley decomposition to F shows

$$\sum_{k_1,k_2 \ge k-100}^{|k_1-k_2| \le 120} \|P_k(P_{k_1} \mathbf{F} P_{k_2}(\widetilde{\mathcal{G}}\phi_{j_3}))\|_{N_k}$$

=
$$\sum_{k_1,k_2 \ge k-100}^{|k_1-k_2| \le 120} \sum_{|l-k_1| \le 4} \|P_k\left[P_l\phi_{j_0} P_{\le k_1-8}\phi_{j_1} P_{k_2}(\widetilde{\mathcal{G}}\phi_{j_3})\right]\|_{N_k}$$
(5.30)

$$+\sum_{k_1,k_2 \ge k-100}^{|k_1-k_2| \le 120} \sum_{|l-k_1| \le 4} \|P_k \left[P_l \phi_{j_1} P_{\le k_1-8} \phi_{j_0} P_{k_2}(\widetilde{\mathcal{G}}\phi_{j_3}) \right]\|_{N_k}$$
(5.31)

$$+\sum_{k_1,k_2 \ge k-100}^{|k_1-k_2| \le 120} \sum_{l_1,l_2 \ge k_1-8, |l_1-l_2| \le 16} \|P_k \left[P_{l_1} \phi_{j_1} P_{l_2} \phi_{j_0} P_{k_2}(\widetilde{\mathcal{G}}\phi_{j_3}) \right]\|_{N_k}.$$
(5.32)

By (8.14) and (5.20), (5.23), $\|P_k f\|_{L^4_{t,x}} \le \|P_k f\|_{F_k}$, the first two terms are bounded as

$$(5.30) + (5.31) \lesssim \sum_{k_1 \ge k-100} \sum_{|k_1 - k_2| \le 120} \sum_{|l-k_1| \le 4} 2^{\frac{k-l}{6}} \|P_l \phi_x\|_{G_l(T)} \|\phi_x\|_{L^4_{l,x}} \|P_{k_2}(\widetilde{\mathcal{G}}\phi_{j_3})\|_{F_{k_2}(T)}$$

$$\lesssim \sum_{k_1 \ge k-100} \varepsilon_0 b_{k_1} \left[2^{\frac{k-k_1}{6}} 2^{-\sigma k_1} b_{k_1}(\sigma) \right]$$

$$\lesssim \epsilon_0 2^{-\sigma k} b_k(\sigma).$$

And using (8.14) and (8.12), we see

$$(5.32) \lesssim \sum_{k_{1},k_{2} \ge k-100}^{|k_{1}-k_{2}| \le 120} \sum_{l_{1},l_{2} \ge k_{1}-8, |l_{1}-l_{2}| \le 16} 2^{\frac{k-l_{1}}{6}} \|P_{l_{1}}\phi_{j_{1}}\|_{G_{l_{1}}(T)} \|(P_{l_{2}}\phi_{j_{0}})P_{k_{2}}(\widetilde{\mathcal{G}}\phi_{j_{3}})\|_{L_{t,x}^{2}}$$

$$\lesssim \sum_{k_{1},k_{2} \ge k-100}^{|k_{1}-k_{2}| \le 120} \sum_{l_{1},l_{2} \ge k_{1}-8, |l_{1}-l_{2}| \le 16} 2^{\frac{k-l_{1}}{6}} \|P_{l_{1}}\phi_{j_{1}}\|_{G_{l_{1}}(T)} \|P_{l_{2}}\phi_{j_{0}}\|_{L^{4}} \|P_{k_{2}}(\widetilde{\mathcal{G}}\phi_{j_{3}})\|_{L^{4}}$$

$$\lesssim \epsilon_{0} \sum_{k_{1} \ge k-100}^{|k_{1}-k_{2}| \le 16} \sum_{l_{1}-k_{1}|l_{1}-l_{2}| \le 16}^{|k_{1}-l_{1}|} 2^{\frac{k-l_{1}}{6}} b_{l_{1}} 2^{-\sigma l_{2}} b_{l_{2}}(\sigma)$$

$$\lesssim 2^{-\sigma k} b_{k} b_{k}(\sigma),$$

where we used the embedding $L_k^4(T) \hookrightarrow F_k(T) \hookrightarrow G_k(T)$ and the fact $\|P_{k_2}(\widetilde{\mathcal{G}}\phi_{j_3})\|_{L^4} \lesssim \|\phi_x\|_{L^4}$ in the third inequality. Thus the third RHS term of (5.21) has been done. Hence, we have finished the proof.

Corollary 5.1. (*Proof of Proposition 5.1*) Under the assumptions of Proposition 5.1, for all $i \in \{1, 2\}$ and $\sigma \in [0, \frac{99}{100}]$ we have

$$\|P_k\phi_i\|_{G_k(T)} \leq 2^{-\sigma k} c_k(\sigma). \tag{5.33}$$

Proof. (5.6) shows for any $k \in \mathbb{Z}$, $\sigma \in [0, \frac{99}{100}]$,

$$2^{\sigma k} \|P_k \phi_i(0,0,\cdot)\|_{L^2_x} \le c_k(\sigma).$$
(5.34)

Then by Proposition 5.3, Proposition 5.2 and the linear estimates of Proposition 2.1, one has

$$b_k(\sigma) \leq c_k(\sigma) + \epsilon_0 b_k(\sigma),$$
 (5.35)

for all $\sigma \in [0, \frac{99}{100}]$. Thus $b_k(\sigma) \leq c_k(\sigma)$, and our result follows by the definition of $\{b_k(\sigma)\}$ in Section 5.

5.2 Unform bounds for $\sigma \in [0, \frac{99}{100}]$

We end the arguments for $\sigma \in [0, \frac{99}{100}]$ with the following proposition.

Proposition 5.4. Assume that $\sigma \in [0, \frac{99}{100}]$. Let $Q \in N$ be a fixed point and ϵ_0 be a sufficiently small constant. Given any $\mathcal{L} \in \mathbb{Z}_+$, assume that $T \in (0, 2^{2\mathcal{L}}]$. Let $\{c_k\}$ be an ϵ_0 -frequency envelope of order δ . And let $\{c_k(\sigma)\}$ be another frequency envelope of order δ . Let $u \in \mathcal{H}_O(T)$ be the solution to SMF with initial data u_0 which satisfies

$$\|P_k \nabla u_0\|_{L^2_x} \le c_k \tag{5.36}$$

$$\|P_k \nabla u_0\|_{L^2_x} \le c_k(\sigma) 2^{-\sigma \kappa}.$$
(5.37)

Denote $\{\phi_i\}$ the corresponding differential fields of the heat flow initiated from u. Then we have for all $i = 1, 2, k \in \mathbb{Z}$, $\sigma \in [0, \frac{99}{100}]$

$$\|P_k\phi_i\|_{s=0} \|_{G_k(T)} \lesssim c_k \tag{5.38}$$

$$\|P_k\phi_i\|_{s=0} \|_{G_k(T)} \lesssim c_k(\sigma) 2^{-\sigma k}$$
(5.39)

$$\sup_{s>0} (1+s2^{2k})^4 \|P_k\phi_i(s)\|_{F_k(T)} \lesssim c_k(\sigma)2^{-\sigma k}.$$
(5.40)

Proof. Define the function $\Theta : [-T.T] \to \mathbb{R}^+$ by

$$\Theta(T') := \sup_{k \in \mathbb{Z}} c_k^{-1} \left(\|P_k \phi_i \upharpoonright_{s=0} \|_{G_k(T')} + \|P_k \nabla u\|_{L^\infty L^2_x(T')} \right).$$

By Lemma 3.3, the function $\Theta(T')$ is continuous in $T' \in [0, T]$. Then Proposition 5.1 implies

$$\Theta(T') \leq \epsilon_0^{-\frac{1}{2}} \Longrightarrow \sup_{k \in \mathbb{Z}} c_k^{-1} \left(\| P_k \phi_i \|_{S=0} \|_{G_k(T')} \right) \leq 1.$$

And by Proposition 3.5,

$$\sup_{k\in\mathbb{Z}}c_k^{-1}\left(\|P_k\phi_i\uparrow_{s=0}\|_{G_k(T')}\right)\lesssim 1\Longrightarrow \sup_{k\in\mathbb{Z}}c_k^{-1}\left(\|P_k\nabla u\|_{L^\infty_t L^2_x(T')}\right)\lesssim 1.$$

Hence, we conclude

$$\Theta(T') \leq \epsilon_0^{-\frac{1}{2}} \Longrightarrow \Theta(T') \lesssim 1.$$

And it is easy to see $\Theta(T')$ is continuous and increasing. Moreover, we have the limit

$$\lim_{T'\to 0} \Theta(T') \lesssim 1,$$

by the definition of $\Theta(T')$, $G_k(T')$ and Corollary 3.1. Therefore, from the continuity of Θ we conclude that (5.36), (5.37) suffice to yield

$$\Theta(T) \leq 1$$
,

thus giving (5.38). Then Proposition 5.1 yields (5.39), and (5.40) follows by the inclusion $G_k \subset F_k$ and Proposition 4.1.

6 Iteration scheme

From now on, the notations $a_k^{(j)}(\sigma)$ and $a_{k,s}^{(j)}(\sigma)$ differ from the ones defined in Section 3. They are defined as follows. Definition 6.1. Assume that $u_0 \in \mathcal{H}_Q$. Given $j \in \mathbb{N}$, let

$$c_{k,(j)}(\sigma) = \sup_{k' \in \mathbb{Z}} 2^{-\frac{1}{2^j}\delta|k-k'|} \|P_{k'} \nabla u_0\|_{L^2_x}, \ \forall \sigma \in \mathbb{I}_j, \ k \in \mathbb{Z},$$

where $\mathbb{I}_0 = [0, \frac{99}{100}]$ and $\mathbb{I}_j = [0, \frac{j}{4} + 1]$ for $j \in \mathbb{Z}_+$.

• For $\sigma \in [0, \frac{99}{100}]$, define

$$c_k^{(0)}(\sigma) = c_{k,(0)}(\sigma), \ \forall \sigma \in [0, \frac{99}{100}].$$

• For $\sigma \in [0, \frac{5}{4}]$, define

$$c_k^{(1)}(\sigma) = \begin{cases} c_{k,(1)}(\sigma), & \sigma \in [0, \frac{99}{100}] \\ c_{k,(1)}(\sigma) + c_{k,(1)}(\frac{3}{8})c_{k,(1)}(\sigma - \frac{3}{8}), & \sigma \in (\frac{99}{100}, \frac{5}{4}]. \end{cases}$$

• Given an integer $j \ge 2$, for $\sigma \in [0, \frac{j}{4} + 1]$, define $\{c_k^{(j)}(\sigma)\}$ by induction:

$$c_{k}^{(j)}(\sigma) = \begin{cases} c_{k,(j)}(\sigma), & \sigma \in [0, \frac{99}{100}] \\ c_{k,(j)}(\sigma) + c_{k,(j)}(\frac{3}{8})c_{k,(j)}(\sigma - \frac{3}{8}), & \sigma \in [0, \frac{5}{4}] \\ \dots \\ c_{k,(j)}(\sigma) + c_{k,(j)}(\frac{3}{8})c_{k}^{(j)}(\sigma - \frac{3}{8}), & \sigma \in [\frac{m+3}{4}, \frac{m}{4} + 1] \\ \dots \\ c_{k,(j)}(\sigma) + c_{k,(j)}(\frac{3}{8})c_{k}^{(j)}(\sigma - \frac{3}{8}), & \sigma \in [\frac{j+3}{4}, \frac{j}{4} + 1]. \end{cases}$$

Definition 6.2. • Assume that $\{a_k(\sigma)\}$ are frequency envelopes of order δ with $\sigma \in [0, \frac{99}{100}]$. Define

$$a_k^{(0)}(\sigma) = c_k^{(0)}(\sigma), \ \forall \sigma \in [0, \frac{99}{100}].$$

• Assume that $\{a_k(\sigma)\}\ are\ frequency\ envelopes\ of\ order\ \delta\ with\ \sigma\in[0,\frac{99}{100}].$ Define

$$a_k^{(1)}(\sigma) = \begin{cases} c_k^{(1)}(\sigma), & \sigma \in [0, \frac{99}{100}] \\ a_k(\sigma) + c_k^{(1)}(\frac{3}{8})c_k^{(1)}(\sigma - \frac{3}{8}), & \sigma \in (\frac{99}{100}, \frac{5}{4}]. \end{cases}$$

• Given an integer $j \ge 2$, assume that $\{a_k(\sigma)\}$ are frequency envelopes of order δ with $\sigma \in [0, \frac{j}{4} + 1]$. Define

$$a_k^{(j)}(\sigma) = \begin{cases} c_k^{(j)}(\sigma), & \sigma \in [0, \frac{j+3}{4}] \\ a_k(\sigma) + c_k^{(j)}(\frac{3}{8})c_k^{(j)}(\sigma - \frac{3}{8}), & \sigma \in (\frac{j+3}{4}, \frac{j}{4} + 1] \end{cases}$$

Given an integer $j \in \mathbb{N}$, assume that $\{a_k(\sigma)\}$ are frequency envelopes of order δ with $\sigma \in [0, \frac{j}{4} + 1]$, we also define

$$a_{k,s}^{(j)}(\sigma) = \begin{cases} 2^{k+k_0} a_{-k_0}(0) a_k^{(j)}(\sigma) \text{ if } k+k_0 \ge 0\\ \sum_{l=k}^{-k_0} a_l(0) a_l^{(j)}(\sigma) \text{ if } k+k_0 \le 0, \end{cases}$$

for $s \in [2^{2k_0-1}, 2^{2k_0+1}), k, k_0 \in \mathbb{Z}$.

Remark 6.1. Given $j \ge 2$, we infer from Def. 6.1 that $\{c^{(j)}(\sigma)\}$ is of order $\frac{1}{2^m}\delta$ if $\sigma \in [\frac{m+3}{4}, \frac{m}{4} + 1]$, $2 \le m \le j$. Particularly, $\{c^{(j)}(\sigma)\}$ is of order δ for all $\sigma \in [0, \frac{j}{4} + 1]$. One can also see from Def. 6.2 that $\{b^{(j)}(\sigma)\}$ are of order δ for all $\sigma \in [0, \frac{j}{4} + 1]$.

Now we iterate the argument of previous sections to obtain uniform bounds for all $\sigma \in [0, \frac{5}{4}]$. We aim to prove the following proposition:

Proposition 6.1. Assume that $\sigma \in [0, \frac{5}{4}]$. Let $Q \in N$ be a fixed point and ϵ_0 be a sufficiently small constant. Given any $\mathcal{L} \in \mathbb{Z}_+$, assume that $T \in (0, 2^{2\mathcal{L}}]$. Let $u \in \mathcal{H}_Q(T)$ be the solution to SMF with initial data u_0 . Let $\{c_k^{(1)}(\sigma)\}$ be frequency envelopes defined by Definition 6.1, and assume that $\{c_k^{(1)}(0)\}$ is an ϵ_0 -frequency envelope. Denote $\{\phi_i\}$ the corresponding differential fields of the heat flow initiated from u. Then for all $i = 1, 2, k \in \mathbb{Z}$, $\sigma \in [0, \frac{5}{4}]$, we have

$$2^{\sigma k} \| P_k \phi_i |_{s=0} \|_{G_k(T)} \leq c_k^{(1)}(\sigma).$$

As before, this proposition will be divided into two propositions for proof, one is for heat flow evolution and the other is for the Schrödinger map flow evolution. In the statement of following propositions or lemmas, the notation \checkmark means the line it lies in can be dropped.

Proposition 6.2. Let $\sigma \in [0, \frac{5}{4}]$. Let $\{b_k(\sigma)\}$ be frequency envelopes of order δ such that $b_k(\sigma) \leq c_k^{(1)}(\sigma)$ for $\sigma \in [0, \frac{99}{100}]$. Assume that $\{c_k^{(1)}(0)\}$ is an ϵ_0 -frequency envelope.

• Assume that for i = 1, 2,

$$\|P_k\phi_i\|_{s=0} \|_{F_k(T)} \le b_k(\sigma') 2^{-\sigma' k} \sigma' \in [0, \frac{5}{4}],$$
(6.1)

$$\sqrt{\|P_k\phi_i(s)\|_{F_k(T)}} \le \varepsilon^{-1} b_k^{(1)}(0)(1+s2^{2k})^{-4}.$$
(6.2)

Then for $\sigma \in [0, \frac{5}{4}]$ *, i* = 1, 2,

$$\|P_k\phi_i(s)\|_{F_k(T)} \lesssim 2^{-\sigma k} (1+s2^{2k})^{-4} b_k^{(1)}(\sigma)$$
(6.3)

$$\|P_k A_i\|_{s=0} \|_{L^4_{t,r}} \lesssim b_k^{(1)}(\sigma) 2^{-\sigma k}.$$

$$(6.4)$$

• Assume further that

$$\|P_k\phi_t\|_{s=0} \|_{L^4_{t,x}} \lesssim b_k(\sigma') 2^{-(\sigma'-1)k} \sigma' \in [0, \frac{5}{4}].$$
(6.5)

Then for $\sigma \in [0, \frac{5}{4}]$, one has

$$\|A_t\|_{s=0} \|_{L^2_{r_x}} \lesssim \varepsilon^2 \tag{6.6}$$

$$\|P_k\phi_t(s)\|_{L^4_{r,r}} \lesssim b_k^{(1)}(\sigma) 2^{-(\sigma-1)k} (1+2^{2k}s)^{-2}$$
(6.7)

$$\|P_k A_t\|_{s=0} \|_{L^2_{tx}} \lesssim \varepsilon b_k^{(1)}(\sigma) 2^{-\sigma k}.$$
(6.8)

Proof. Recalling definitions of $c_k^{(1)}(\sigma)$, $b_k^{(1)}(\sigma)$ in Def. 6.1 and Def. 6.2, by Proposition 5.4, we see (6.3), (6.4), (6.7), (6.8) are already done for $\sigma \in [0, \frac{99}{100}]$. Moreover, (6.6) and the assumption (6.2) hold naturally. The left is to prove (6.3), (6.4), (6.7), (6.8) for $\sigma \in [\frac{99}{100}, \frac{5}{4}]$.

The key and starting point for SMF iteration scheme is to improve $||P_k \widetilde{\mathcal{G}}^{(1)}||_{L^4_t L^\infty_{\omega}}$ step by step.

Lemma 6.1. Let $u \in \mathcal{H}_Q(T)$ solve SMF with data u_0 . Given any $\sigma \in [0, \frac{99}{100}]$, let $\{c_k^{(1)}(\sigma)\}$ be frequency envelopes defined in Def. 6.1. Assume also that $\{c_k^{(1)}(0)\}$ is an ϵ_0 -frequency envelope. Then for ϵ_0 sufficiently small there holds

$$2^{\frac{1}{2}k} \|P_k \widetilde{\mathcal{G}}^{(1)}\|_{L^4_x L^\infty_t} \le c_k^{(1)}(\sigma) 2^{-\sigma k} [(1+2^{2k+2k_0})^{-20} \mathbf{1}_{k+k_0 \ge 0} + \mathbf{1}_{k+k_0 \le 0} 2^{\delta |k+k_0|}]_{L^\infty_x L^\infty_t}$$

for any $\sigma \in [0, \frac{99}{100}]$, $k, k_0 \in \mathbb{Z}$, $s \in [2^{2k_0-1}, 2^{2k_0+1})$.

Proof. We obtain by combining Proposition 5.1 and Proposition 4.1 that

$$\begin{split} \|P_k\phi_t\|_{L^4} &\lesssim (1+s2^{2k})^{-2}2^{-\sigma k+k}c_k^{(1)}(\sigma), \ \sigma \in [0,\frac{99}{100}]\\ \|P_k\phi_i\|_{L^4} &\lesssim (1+s2^{2k})^{-4}2^{-\sigma k}c_k^{(1)}(\sigma), \ \sigma \in [0,\frac{99}{100}]. \end{split}$$

Then Proposition 3.6 yields

$$\begin{split} \left\| P_k \widetilde{\mathcal{G}} \right\|_{L^4_{t,x} \cap L^\infty_t L^2_x} &\lesssim (1 + s 2^{2k})^{-30} 2^{-\sigma k - k} c^{(1)}_k(\sigma), \ \sigma \in [0, \frac{99}{100}] \\ \left\| P_k \widetilde{\mathcal{G}}^{(m)} \right\|_{L^4_{t,x} \cap L^\infty_t L^2_x} &\lesssim (1 + s 2^{2k})^{-30} 2^{-\sigma k - k} c^{(1)}_k(\sigma), \ m = 1, 2, \ \sigma \in [0, \frac{99}{100}]. \end{split}$$

Then using the schematic formula

$$A_t = \int_s^\infty \phi_t \diamond (D_i \phi_i) \mathcal{G} ds'$$

and bilinear Littlewood-Paley decomposition (see Lemma 4.7), we get

$$\|P_k A_t\|_{L^4} \le c_k^{(1)}(\sigma) 2^{-\sigma k+k} [(1+2^{2k+2j})^{-1} \mathbf{1}_{k+j \ge 0} + \mathbf{1}_{k+j \le 0} c_k^{(1)} 2^{\delta [k+j]}]$$

for any $\sigma \in [0, \frac{99}{100}]$, $k, j \in \mathbb{Z}$, $s \in [2^{2j-1}, 2^{2j+1})$. And then using $\partial_t \widetilde{\mathcal{G}}^{(1)} = A_t \mathcal{G}^{(1)} + \mathcal{G}^{(2)} \phi_t$ and interpolation (see Lemma 4.7), one deduces that

$$2^{\frac{1}{2}k} \| P_k \widetilde{\mathcal{G}}^{(1)} \|_{L^4_x L^\infty_t} \le c_k^{(1)}(\sigma) 2^{-\sigma k} [(1 + 2^{2k+2k_0})^{-20} \mathbf{1}_{k+k_0 \ge 0} + \mathbf{1}_{k+k_0 \le 0} 2^{\delta |k+k_0|}]$$

for any $\sigma \in [0, \frac{99}{100}], k, k_0 \in \mathbb{Z}, s \in [2^{2k_0-1}, 2^{2k_0+1}).$

As before, we start with the bound for connection forms. Lemma 6.2. Let $\sigma \in [\frac{99}{100}, \frac{5}{4}]$. Denote

$$h(k) = \sup_{s \ge 0} (1 + s2^{2k})^4 \sum_{i=1}^2 ||P_k \phi_i(s)||_{F_k(T)}.$$
(6.9)

Define the corresponding envelope by

$$h_{k}(\sigma) = \sup_{k' \in \mathbb{Z}} 2^{\sigma k'} 2^{-\delta |k'-k|} h(k').$$
(6.10)

Then under the assumptions of Proposition 6.2, for all $k \in \mathbb{Z}$, $s \ge 0$, i = 1, 2, we have

$$\|P_k(A_i(s))\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \leq 2^{-\sigma k} (1 + s 2^{2k})^{-4} h_{k,s}^{(1)}(\sigma),$$
(6.11)

where the sequence $\{h_{k,s}^{(1)}\}$ when $2^{2k_0-1} \leq s < 2^{2k_0+1}, k_0 \in \mathbb{Z}$, is defined by

$$h_{k,s}^{(1)}(\sigma) = \begin{cases} 2^{k+k_0} h_{-k_0} h_k^{(1)}(\sigma) \text{ if } k + k_0 \ge 0\\ \sum_{l=k}^{-k_0} h_l h_l^{(1)}(\sigma) \text{ if } k + k_0 \le 0 \end{cases}$$
(6.12)

with

$$h_k^{(1)}(\sigma') = \begin{cases} c_k^{(1)}(\sigma'), & \sigma' \in [0, \frac{99}{100}] \\ h_k(\sigma') + c_k^{(1)}(\frac{3}{8})c_k^{(1)}(\sigma' - \frac{3}{8}), & \sigma' \in (\frac{99}{100}, \frac{5}{4}]. \end{cases}$$
(6.13)

Proof. The proof is almost the same as Lemma 4.1. The difference is that more concerns are needed for the High × Low interaction of $P_k[\widetilde{\mathcal{G}}^{(1)}\psi_s]$ in Step 4 of Lemma 4.1. First of all we point out (5.40) of Proposition 5.4 shows for all $\sigma' \in [0, \frac{99}{100}]$

$$h_k(\sigma') \leq c_k^{(1)}(\sigma'). \tag{6.14}$$

Let $B_1^{(1)}$ be the smallest constant such that for all $\sigma \in [\frac{99}{100}, \frac{5}{4}]$, $s \ge 0, k \in \mathbb{Z}$, there holds

$$\|P_k(A_i(s))\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \lesssim B_1^{(1)} 2^{-\sigma k} (1 + s 2^{2k})^{-4} h_{k,s}^{(1)}(\sigma).$$
(6.15)

Recall the following decomposition of G:

$$\mathcal{G} = \Gamma^{\infty} - \Gamma_p^{\infty,(1)} \int_s^\infty \psi_s^p ds' - \int_s^\infty \psi_s^p \widetilde{\mathcal{G}}^{(1)} ds'.$$

By $\psi_s = \sum_{i=1}^2 \partial_i \psi_i + A_i \psi_i$, we separate the ψ_i part away. And thus schematically one has

$$\mathcal{G} = \Gamma^{\infty} - \Gamma_l^{\infty,(1)} \int_s^{\infty} (\partial_i \psi_i)^l ds' - \int_s^{\infty} (\partial_i \psi_i)^l \widetilde{\mathcal{G}}_l^{(1)} ds$$
$$- \Gamma_l^{\infty,(1)} \int_s^{\infty} (A_i \psi_i)^l ds' - \int_s^{\infty} (A_i \psi_i)^l \widetilde{\mathcal{G}}_l^{(1)} ds'$$

In order to prove our lemma, as before we first prove $B_1^{(1)} \leq 1$ under the **Bootstrap Assumption B**: For the fixed given $\sigma \in (\frac{99}{100}, \frac{5}{4}]$, there hold

$$\int_{s}^{\infty} \|P_{k}(A_{i}\psi_{i})\|_{F_{k}(T)} ds' \leq \varepsilon^{-\frac{1}{2}} 2^{-\sigma k} T_{k,j} (1+s^{\frac{1}{2}}2^{k})^{-7} h_{k}^{(1)}(\sigma) c_{0}^{*}$$
$$\int_{s}^{\infty} \|P_{k}[(A_{i}\psi_{i})\widetilde{\mathcal{G}}^{(1)}]\|_{F_{k}(T)} ds' \leq \varepsilon^{-\frac{1}{2}} 2^{-\sigma k} T_{k,j} (1+s^{\frac{1}{2}}2^{k})^{-7} h_{k}^{(1)}(\sigma) c_{0}^{*}.$$

where $c_0^* := ||\{h_k\}||_{\ell^2}$, $s \in [2^{2j-1}, 2^{2j+1})$, and $T_{k,j}$ is defined in (4.29). This part is the same as Step 2 of Lemma 4.1 except controlling

$$\left\|P_k\left(\int_s^\infty (\partial_i\psi_i)(\widetilde{\mathcal{G}}^{(1)})ds'\right)\right\|_{F_k(T)},\tag{6.16}$$

which was labeled as \mathcal{U}_{01} in Lemma 4.1. To estimate (6.16), recall the bounds in Lemma 6.1 and Proposition 3.6 for $\widetilde{\mathcal{G}}^{(1)}$:

$$2^{k} \| P_{k}(\widetilde{\mathcal{G}}^{(1)}) \|_{L_{t}^{\infty} L_{x}^{2} \cap L^{4}} + 2^{\frac{1}{2}k} \| P_{k}(\widetilde{\mathcal{G}}^{(1)}) \|_{L_{x}^{4} L_{t}^{\infty}}$$

$$\lesssim 2^{-\widetilde{\sigma}k} c_{k}(\widetilde{\sigma}) [1_{k+j \ge 0} (1+2^{2k}s)^{-20} + 1_{k+j \le 0} 2^{\delta[k+j]}]$$
(6.17)

for any $k, j \in \mathbb{Z}, s \in [2^{2j-1}, 2^{2j+1}), \widetilde{\sigma} \in [0, \frac{99}{100}]$. By bilinear Littlewood-Paley decomposition and Lemma 4.2, we have

$$\begin{split} \|P_{k}((\partial_{i}\psi_{i})\mathcal{G}^{(1)})\|_{F_{k}(T)} \\ &\lesssim \sum_{|k_{1}-k|\leq4} \|P_{k_{1}}(\partial_{i}\psi_{i})\|_{F_{k_{1}}(T)} \|P_{\leq k-4}\widetilde{\mathcal{G}}^{(1)}\|_{L^{\infty}} \\ &+ \sum_{|k_{1}-k_{2}|\leq8,k_{1},k_{2}\geq k-4} \|P_{k_{1}}(\partial_{i}\psi_{i})\|_{F_{k_{1}}(T)} (\|P_{k_{2}}(\widetilde{\mathcal{G}}^{(1)})\|_{L^{\infty}} + 2^{\frac{k_{1}}{2}} \|P_{k_{2}}(\widetilde{\mathcal{G}}^{(1)})\|_{L^{4}_{x}L^{\infty}_{t}}) \\ &+ \sum_{|k_{2}-k|\leq4,k_{1}\leq k-4} 2^{\frac{k_{1}}{2}} \|P_{k_{1}}(\partial_{i}\psi_{i})\|_{F_{k_{1}}(T)} \|P_{k_{2}}(\widetilde{\mathcal{G}}^{(1)})\|_{L^{4}_{x}L^{\infty}_{t}} + 2^{k_{1}} \|P_{k_{1}}(\partial_{i}\psi_{i})\|_{F_{k_{1}}(T)} \|P_{k_{2}}(\widetilde{\mathcal{G}}^{(1)})\|_{L^{4}_{x}L^{\infty}_{t}} + 2^{k_{1}} \|P_{k_{1}}(\partial_{i}\psi_{i})\|_{F_{k_{1}}(T)} \|P_{k_{2}}(\widetilde{\mathcal{G}}^{(1)})\|_{L^{4}_{x}L^{\infty}_{t}} \\ &\lesssim 2^{-\sigma k} h_{k}^{(1)}(\sigma) + 2^{-\sigma k} h_{k} h_{k}^{(1)}(\sigma) \left[1_{k+j\geq0} 2^{k}(1+2^{2k}s)^{-4} + 1_{k+j\leq0} 2^{\delta|k+j|}2^{-j}\right] \\ &+ R_{j,k} 2^{-(\sigma-\frac{3}{8})k} c_{k}^{(1)}(\sigma-\frac{3}{8}) [2^{-\frac{1}{2}k} \sum_{k_{1}\leq k-4} 2^{\frac{3}{2}k_{1}-\frac{3}{8}k_{1}} c_{k_{1}}^{(1)}(\frac{3}{8}) + 2^{-k} \sum_{k_{1}\leq k-4} 2^{2k_{1}-\frac{3}{8}k_{1}} c_{k_{1}}^{(1)}(\frac{3}{8})] \end{split}$$

where we denote $R_{j,k} := 1_{k+j \ge 0} (1 + 2^{2k}s)^{-20} + 1_{k+j \le 0} 2^{\delta|k+j|} h_k$ and have used (6.14). Thus by slow variation of envelopes we get

$$\|P_k((\partial_i \psi_i)\widetilde{\mathcal{G}}^{(1)})\|_{F_k(T)} \lesssim 2^{-\sigma k} h_k^{(1)}(\sigma) \left(\mathbf{1}_{k+j \ge 0} 2^k (1+2^{2k_2}s)^{-4} + \mathbf{1}_{k+j \le 0} 2^{-j} 2^{\delta|k+j|} \right).$$

for $s \in [2^{2j-1}, 2^{2j+1})$, $j, k \in \mathbb{Z}$. This bound is the same as \mathcal{U}_{01} in Lemma 4.1 and acceptable.

In the third step, we prove the claim: If **Bootstrap Assumption B** holds, then

$$\int_{s}^{\infty} \|P_{k}(A_{i}\psi_{i})\|_{F_{k}(T)} ds' \leq 2^{-\sigma k} T_{k,j} (1 + s^{\frac{1}{2}} 2^{k})^{-7} h_{k}^{(1)}(\sigma) c_{0}^{*}$$
(6.18)

$$\int_{s}^{\infty} \|P_{k}(A_{i}\psi_{i})\widetilde{\mathcal{G}}^{(1)}\|_{F_{k}(T)} ds' \leq 2^{-\sigma k} T_{k,j}(1+s^{\frac{1}{2}}2^{k})^{-7} h_{k}^{(1)}(\sigma) c_{0}^{*}.$$
(6.19)

The proof of (6.18) is the same as Step 4 of Lemma 4.1. For (6.19), the Low × High interaction of $P_k[(A_i\psi_i)\widetilde{\mathcal{G}}^{(1)}]$ is different due to the larger σ . The other two interactions are all the same. We present the modifications as follows. Since under Bootstrap Assumption B one has $B^{(1)} \leq 1$, $P_k(A_i\psi_i)$ enjoys the same $F_k \cap S_k^{\frac{1}{2}}$ bound as Lemma 4.1 with $h_k(\sigma)$ replaced by $h_k^{(1)}(\sigma)$:

$$\begin{aligned} \|P_{k}(A_{i}\psi_{i})\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}(T)} &\lesssim c_{0}^{*}2^{-\sigma k}\mathbf{1}_{k+j\leq 0}h_{k}^{(1)}(\sigma)2^{\frac{1}{2}(k-j)}2^{\delta|k+j|} \\ &+ c_{0}^{*}2^{-\sigma k}\mathbf{1}_{k+j\geq 0}h_{k}^{(1)}(\sigma)2^{k}(1+2^{j+k})^{-8}. \end{aligned}$$

for all $\sigma \in [0, \frac{5}{4}]$, $s \in [2^{2j-1}, 2^{2j+1})$, $j, k \in \mathbb{Z}$.

Then by (6.17), (6.14) and (4.47), the Low × High part of $(A_i\psi_i)\widetilde{\mathcal{G}}^{(1)}$ is dominated by

$$\begin{split} &\sum_{|k-k_2| \le 4, k_1 \le k+4} \|P_k(P_{k_1}(A_i\psi_i)P_{k_2}\widetilde{\mathcal{G}}^{(1)})\|_{F_k(T)} \\ &\lesssim c_0^* 2^{-(\sigma-\frac{3}{8})k} c_k^{(1)}(\sigma-\frac{3}{8}) \mathbf{1}_{k+j \le 0} \sum_{k_1 \le k-4} c_{k_1}^{(1)}(\frac{3}{8}) 2^{\frac{1}{2}(k_1-j)} 2^{\delta|k_1+j|} 2^{-\frac{3}{8}k_1} \\ &+ c_0^* 2^{-(\sigma-\frac{3}{8})k} (1+2^{2j+2k})^{-20} c_k^{(1)}(\sigma-\frac{3}{8}) \mathbf{1}_{k+j \ge 0} \left[\sum_{-j \le k_1 \le k} c_{k_1}^{(1)}(\frac{3}{8}) 2^{k_1-\frac{3}{8}k_1} (1+2^{2j+2k_1})^{-4} \right] \\ &+ c_0^* 2^{-(\sigma-\frac{3}{8})k} (1+2^{2j+2k})^{-20} c_k^{(1)}(\sigma-\frac{3}{8}) \mathbf{1}_{k+j \ge 0} \left[\sum_{k_1 \le -j} c_{k_1}^{(1)}(\frac{3}{8}) 2^{\frac{k_1-j}{2}} 2^{\delta|k_1+j|} 2^{-\frac{3}{8}k_1} \right] \\ &\lesssim c_0^* 2^{-\sigma k} c_k^{(1)}(\sigma-\frac{3}{8}) c_k^{(1)}(\frac{3}{8}) \left(\mathbf{1}_{k+j \ge 0} 2^{-j} (1+2^{j+k})^{-7} + \mathbf{1}_{k+j \le 0} 2^{\frac{k_j}{2}} 2^{\delta|k+j|} \right). \end{split}$$

Summing in $j \ge k_0$ as well yields

$$\sum_{j \ge k_0} 2^{2j} \sum_{|k-k_2| \le 4, k_1 \le k+4} \|P_k(P_{k_1}\psi_s P_{k_2}\widetilde{\mathcal{G}}^{(1)})\|_{F_k(T)}$$

$$\lesssim c_0^* 2^{-\sigma k} h_k^{(1)}(\sigma) \Big(\mathbf{1}_{k+k_0 \ge 0} 2^{k_0} (1+2^{k+k_0})^{-7} + 2^{-k} \mathbf{1}_{k+k_0 \le 0} \Big)$$

for $s \in [2^{2k_0-1}, 2^{2k_0+1}), k_0, k \in \mathbb{Z}$. This bound is again the same as \mathcal{U}_{II} in Lemma 4.1 and acceptable.

Finally, we need to prove (6.18), (6.19) of Bootstrap Assumption B hold when $T \to 0$. Let's verify it. Using (3.35), (3.55) and putting $\frac{3}{8}$ order derivatives on $(A_i\psi_i)$ while estimating the Low × High interaction of $(A_i\psi_i)\widetilde{\mathcal{G}}^{(1)}$, we also have

$$\int_{s}^{\infty} \|P_{k}[A_{i}\psi_{i}\widetilde{\mathcal{G}}^{(1)}]\|_{L_{t}^{\infty}L_{x}^{2}} ds' \lesssim \|\{h_{k}\}\|_{\ell^{2}} T_{k,j}h_{k}^{(1)}(\sigma) 2^{-\sigma k}$$

if $\sigma \in [\frac{99}{100}, \frac{5}{4}]$.

Therefore, combining the above four steps gives Lemma 6.2.

The proof of Lemma 6.2 gives an F_k bound for \mathcal{G} .

Lemma 6.3. *For all* $\sigma \in [\frac{99}{100}, \frac{5}{4}], k \in \mathbb{Z}$,

$$\left\| P_k(\widetilde{\mathcal{G}}) \right\|_{F_k(T)} \lesssim \begin{cases} 2^{-\sigma k} (1 + s2^{2k})^{-4} 2^j h_k^{(1)}(\sigma), \text{ if } j + k \ge 0\\ 2^{-\sigma k} 2^{-k} h_k^{(1)}(\sigma), \text{ if } j + k \le 0 \end{cases}$$
(6.20)

when $2^{2j-1} \leq s \leq 2^{2j+1}$, $j \in \mathbb{Z}$. Moreover, for s = 0

$$\left\| P_k \widetilde{\mathcal{G}} \upharpoonright_{s=0} \right\|_{F_k(T)} \lesssim 2^{-k-\sigma k} h_k^{(1)}(\sigma).$$
(6.21)

Proof of Proposition 6.2. With this improved bound of $\widetilde{\mathcal{G}}$, running the program of Section 4 again gives

$$\sup_{s\geq 0} 2^{\sigma k} (1+s2^{2k})^4 \sum_{i=1}^{2} \|P_k \phi_i(s)\|_{F_k(T)} \leq b_k(\sigma) + \varepsilon h^{(1)}(\sigma).$$

Since the right side is a frequency envelope of order δ , there holds

$$h_k(\sigma) \leq b_k(\sigma) + \varepsilon h_k^{(1)}(\sigma)$$

By the definition of $h_k^{(1)}(\sigma)$, we conclude for $\sigma \in [\frac{99}{100}, \frac{5}{4}]$

$$h_k(\sigma) \leq b_k(\sigma) + c_k^{(1)}(\frac{3}{8})c_k^{(1)}(\sigma - \frac{3}{8}),$$

thus proving (6.3). The left (6.4), (6.7), (6.8) are the same.

In the following proposition, we finish iteration of σ in the Schrödinger direction.

Proposition 6.3. Given $\mathcal{L} \in \mathbb{Z}_+$, suppose that $T \in (0, 2^{2\mathcal{L}}]$ and $Q \in \mathcal{N}$. Assume that $\sigma \in [0, \frac{5}{4}]$. Let $u \in \mathcal{H}_Q(T)$ be a solution to SMF with initial data u_0 , and set $\{c_k^{(1)}(\sigma)\}_{k\in\mathbb{Z}}$ to be frequency envelopes defined in Def. 6.1. And assume that $\{c_k^{(1)}(0)\}$ is an ϵ_0 -frequency envelope with $0 < \epsilon_0 \ll 1$. Then for any $\sigma \in [0, \frac{5}{4}]$, $k \in \mathbb{Z}$, we have

$$\|P_k\phi_i\|_{s=0} \|_{G_k(T)} \leq c_k^{(1)}(\sigma).$$
 (6.22)

Proof. (6.22) has been proved for $\sigma \in [0, \frac{99}{100}]$ in Section 5. Thus, it suffices to consider $\sigma \in (\frac{99}{100}, \frac{5}{4}]$ only. Let

$$b(k) = \sum_{i=1}^{2} ||P_k \phi_i||_{s=0} ||_{G_k(T)}.$$

For $\sigma \in [0, \frac{5}{4}]$, define the frequency envelopes:

$$b_k(\sigma) = \sup_{k' \in \mathbb{Z}} 2^{\sigma k'} 2^{-\delta|k-k'|} b(k').$$

By Proposition 3.1, they are finite and ℓ^2 summable. And

$$\|P_k\phi_i\|_{s=0}\|_{G_k(T)} \leq 2^{-\sigma k}b_k(\sigma).$$

The assumption (6.5) holds by repeating the same argument of Lemma 4.1. Thus using Proposition 6.2, we see (6.3)-(6.8) hold. With Lemma 6.3, repeating the argument in Section 5, one obtains when s = 0,

$$\|P_k\phi_i\|_{s=0} \|_{G_k(T)} \lesssim c_k^{(1)}(\sigma) + \epsilon_0 \left(b_k(\sigma) + c_k^{(1)}(\frac{3}{8}) c_k^{(1)}(\sigma - \frac{3}{8}) \right), \ \forall \sigma \in (\frac{99}{100}, \frac{5}{4}].$$

Since the RHS is frequency envelope of order δ , we conclude

$$b_k(\sigma) \leq c_k^{(1)}(\sigma).$$

This gives (6.22) and finishes our proof.

7 **Proof of Theorem 1.1 and Theorem 1.2**

7.1 Global Regularity

In order to prove *u* is global, it suffices to verify (see Appendix B)

$$\|\nabla u\|_{L^{\infty}_{t,x}} \lesssim 1. \tag{7.1}$$

To prove (7.1), it suffices to give uniform bound for $||u(t)||_{\dot{H}^1 \cap \dot{H}^{2+}}$. Since energy preserves, it reduces to bound $||u(t)||_{\dot{H}^{2+}}$, which is related to frequency envelopes with $\sigma = 1+$. Thus we need to transform the intrinsic bound (6.22) to bounds for u.

The following lemma follows directly by Corollary 3.1.

Lemma 7.1. Let $u \in \mathcal{H}_Q(T)$ solve SMF with data u_0 of small energy. For $\sigma \in [0, \frac{5}{4}]$, suppose that $\{c_k^{(1)}(\sigma)\}$ are frequency envelopes defined in Def. 6.1. And assume that the differential fields $\{\phi_i\}$ associated with u under the caloric gauge satisfy

$$\|P_k\psi_i\|_{s=0} \|_{L^{\infty}_t L^2_x} \le 2^{-\sigma k} c^{(1)}_k(\sigma), \ \forall \ k \in \mathbb{Z}.$$

$$(7.2)$$

Then we have

$$2^{k} \|P_{k}u\|_{L^{\infty}_{t}L^{2}_{x}} \leq 2^{-\sigma k} c_{k}^{(1)}(\sigma), \ \forall \ k \in \mathbb{Z}.$$

$$(7.3)$$

Proposition 6.3 shows the assumption (7.2) of Lemma 7.1 holds. And thus by applying Lemma 7.1, we conclude

$$\|u\|_{\dot{H}^{\rho}\cap\dot{H}^{1}} \leq C(\|u_{0}\|_{\dot{H}^{\rho}\cap\dot{H}^{1}}), \tag{7.4}$$

for all $\rho \in [0, \frac{9}{4}]$. Particularly $\|\nabla u\|_{L^{\infty}_{t,x}} \leq 1$ by Sobolev embedding. Therefore, *u* is global by Appendix B and the global regularity follows by local theory of [30].

The remained part for Theorem 1.1 is (1.4) and (1.5). They will be proved in Section 7.4 and Section 7.5 respectively.

7.2 Uniform Sobolev norm bounds of solutions to SMF

To get uniform Sobolev norm bounds for SMF up to $\sigma = 1 + \frac{K}{4}$, $K \in \mathbb{Z}_+$, in the heat flow iteration scheme it suffices to begin with the parabolic decay estimates

$$\begin{aligned} \left\| \partial_x^{L+1} \mathcal{G}^{(K+1)} \right\|_{L^{\infty}_t L^2_x} &\lesssim \epsilon s^{-\frac{L}{2}}, \forall L \in [0, 100 + K] \\ \left| \partial_x^{L+1} [d\mathcal{P}]^{(K+1)} \right\|_{L^{\infty}_t L^2_x} &\lesssim \epsilon s^{-\frac{L}{2}}, \forall L \in [0, 100 + K]. \end{aligned}$$

And in the SMF iteration scheme, for the *j*-th iteration we always begin with proving

$$2^{\frac{1}{2}k} \| P_k \widetilde{\mathcal{G}}^{(1)} \|_{L^4_x L^\infty_t} \le c_k^{(j)}(\sigma) 2^{-\sigma k} [(1+2^{2k+2k_0})^{-20} \mathbf{1}_{k+k_0 \ge 0} + \mathbf{1}_{k+k_0 \le 0} 2^{\delta |k+k_0|}], \ \sigma \in [0, 1+\frac{j-1}{4}],$$

for any $s \in [2^{2k_0-1}, 2^{2k_0+1}), k_0, k \in \mathbb{Z}$. Then repeating the argument of first time iteration for K times we obtain

$$\begin{aligned} \|P_k d\mathcal{P}(e)(\uparrow_{s=0})\|_{L^{\infty}_t L^2_x} &\leq 2^{-\sigma k} c^{(K)}_k(\sigma) \\ \|P_k \phi_x(\uparrow_{s=0})\|_{L^{\infty}_t L^2_x} &\leq 2^{-\sigma k+k} c^{(K)}_k(\sigma). \end{aligned}$$

By bilinear estimates we then arrive at

$$\|P_k \partial_x u\|_{L^\infty_* L^2_x} \lesssim 2^{-\sigma k+k} c_k^{(K)}(\sigma), \tag{7.5}$$

by which the uniform Sobolev bounds follow. Each time iteration requires ϵ_* to be smaller in our arguments. We emphasize that the key for the succeeding SMF iterations is to improve $\|P_k \mathcal{G}^{(1)}\|_{L^4_r L^\infty_r}$ step by step. (see e.g. Lemma 6.1)

Therefore, we have the following result:

Proposition 7.1. For any $j \ge 1$, there exists a constant $\epsilon_j > 0$ such that if $u_0 \in \mathcal{H}_Q$ with $||u_0||_{\dot{H}^1} \le \epsilon_j$, then $||u(t)||_{\dot{H}^j_x} \le C(||u_0||_{\dot{H}^1 \cap \dot{H}^j})$ for all $t \in \mathbb{R}$.

Since the mass of SMF solutions doesnot conserve, the $||u - Q||_{L^2_x}$ norm should be handled separately. This will be proved as a corollary of the well-posedness, see the next section.

7.3 Well-posedness

In fact, the well-posedness stated in Theorem 1.2 follows closely by [41] and [4]'s original arguments. We sketch it for reader's convenience.

[Tataru [41], Prop. 3.13] proves that: given two maps $u_0^0, u_0^1 \in \mathcal{H}_Q$ with $||u_0^h||_{\dot{H}^1} \ll 1, h = 0, 1$, there exists a smooth one parameter family of initial data $\{u_0^h\}_{h\in[0,1]} \in C^{\infty}([0,1];\mathcal{H}_Q)$ which satisfies

$$\|u_0^h\|_{\dot{H}^1} \ll 1, \ h \in [0, 1] \tag{7.6}$$

$$\int_0^1 \left\| P_k \partial_x u^h \right\|_{L^2_x} dh \approx \| u_0^0 - u_0^1 \|_{L^2_x}.$$
(7.7)

Given $h \in [0, 1]$, Theorem 1.1 yields a solution $u^h(t, x) \in C(\mathbb{R}; \mathcal{H}_Q)$ with initial data u_0^h . Then under the caloric gauge $\{e_\alpha, Je_\alpha\}$ for $u^h(t, x)$, define the differential field ϕ_h by

$$\phi_h^{\alpha} = \langle \partial_h u^h, e_{\alpha} \rangle + \sqrt{-1} \partial_h u^h, J e_{\alpha} \rangle, \ \alpha = 1, ..., n,$$
(7.8)

and define $\{\phi_i\}_{i=0}^2$ as before. Since $-\sqrt{-1}\phi_t = \sum_{i=1,2} D_i\phi_i$ at s = 0 (because for all $h \in [0, 1]$ $u^h(t, x)$ solves SMF), applying $D_h = \partial_h + A_h$ to the both sides gives

$$-\sqrt{-1}D_t\phi_h = \sum_{i=1}^2 D_i D_i \phi_h + \sum_{i=1}^2 \mathcal{R}(u^h(t,x))(\phi_i,\phi_h)\phi_i, \text{ when } s = 0.$$

which as before can be further schematically written as

$$-\sqrt{-1}D_t\phi_h = \sum_{i=1}^2 D_i D_i \phi_h + \sum (\phi_i \diamond \phi_h)\phi_i \mathcal{G}, \text{ when } s = 0.$$
(7.9)

Given $\sigma \in [0, 1 + \frac{j}{4})$ with $j \in \mathbb{Z}_+$, let $\{c_{k,(j)}(\sigma)\}$ be

$$c_{k,(j),h}(\sigma) = \sup_{k' \in \mathbb{Z}} 2^{-\frac{1}{2^{j}}\delta|k'-k|} 2^{\sigma k'+k'} \|P_{k'}u_{0}^{h}\|_{L^{2}_{x}}.$$

And define $\{c_{k,h}^{(j)}(\sigma)\}$ as Def. 6.1. Then Section 7.2 gives

$$\sum_{i=1}^{2} 2^{\sigma k} \| P_k \phi_i(s=0,h,\cdot,\cdot) \|_{G_k(T)} \lesssim c_{k,h}^{(j)}(\sigma),$$
(7.10)

and thus

$$2^{\sigma k} \| P_k \widetilde{\mathcal{G}}(s=0,h,\cdot,\cdot) \|_{F_k(T)} \leq c_{k,h}^{(j)}(\sigma).$$

$$(7.11)$$

Using (7.10), (7.11) we obtain by (7.9) that

$$\sum_{k \in \mathbb{Z}} \|P_k \phi_h(s=0)\|_{G_k(T)}^2 \lesssim \|\phi_h(s=0,t=0)\|_{L^2_x}^2$$

Transforming this bound to $\partial_h u^h$ yields

 $\|\partial_h u^h\|_{L^\infty_t L^2_x} \lesssim \|\partial_h u^h_0\|_{L^2_x}.$

Then (7.7) leads to

$$\|u^{1} - u^{0}\|_{L_{t}^{\infty}L_{x}^{2}} \leq \|u_{0}^{1} - u_{0}^{0}\|_{L_{x}^{2}}.$$
(7.12)

With (7.13) in hand, the continuity of S_Q from $\mathfrak{B}^{\sigma}_{\epsilon}$ to $C(\mathbb{R}; H^{\sigma+1}_Q)$ follows by the same arguments of [[4],1467-1468], if $\epsilon > 0$ is sufficiently small depending only on *j* thus σ . Moreover, letting $u_0^1 = Q$, $u_0^0 = u_0$ in (7.13) one obtains

$$\|u - Q\|_{L_t^{\infty} L_x^2} \lesssim \|u_0 - Q\|_{L_x^2}. \tag{7.13}$$

which combined with Proposition 7.1 gives (1.6).

Asymptotic behavior 7.4

Let us prove (1.4). First, we notice

$$|u(t,x) - Q| = \int_0^\infty |\partial_s v(s,t,x)| ds' \lesssim \int_0^\infty |\phi_s| ds'.$$
(7.14)

Step 1.1. Recall the definition of $\{c_k^{(j)}(\sigma)\}$ in Def. 6.1. Applying (3.74) with $\beta_k(\sigma) = c_k^{(0)}(\sigma)$, and its analogies in succeeding iterations, we get by Bernstein inequality that

$$\|\phi_s\|_{L^4_t L^{\infty}_x} \lesssim s^{-\frac{1}{4}} \sum_{k \in \mathbb{Z}} c_k^{(1)}(1)$$
(7.15)

$$\|\phi_s\|_{L^4_t L^\infty_x} \lesssim s^{-\frac{3}{4}} \sum_{k \in \mathbb{Z}} c_k^{(0)}(0).$$
(7.16)

We find by Young's inequality and triangle inequality that

$$2^{\frac{1}{2^{j+4}}\delta|k|}c_k^{(j)} \lesssim \sup_{k' \in \mathbb{Z}} 2^{\frac{1}{2^{j+4}}\delta|k'|} \|P_{k'}\nabla u_0\|_{L^2_x}$$

thus there holds

$$\sum_{k \in \mathbb{Z}} c_k^{(j)} \lesssim \sup_{k' \in \mathbb{Z}} 2^{\frac{1}{2^{j+4}} \delta |k'|} \|P_{k'} \nabla u_0\|_{L^2_x} \lesssim 1,$$
(7.17)

since $u_0 \in \mathcal{H}_Q$. Then (7.16), (7.15) show

$$\|\phi_s\|_{L^4_t L^\infty_x} \lesssim \min(s^{-\frac{1}{4}}, s^{-\frac{3}{4}}). \tag{7.18}$$

We see (7.18) is not enough to put $\|\phi_s\|_{L_x^{\infty}}$ in L_s^1 , but useful for Step 2 below. **Step 1.2.** Applying (3.74) with $\beta_k(\sigma) = c_k^{(0)}(\sigma)$, $\sigma = 0$, and by interpolation, we see for any $p \in (4, \infty)$, $\tilde{p} \in (2, 4)$ satisfying $\frac{1}{p} + \frac{1}{\tilde{p}} = \frac{1}{2}$, there holds

$$\|\phi_s\|_{L^p_t L^{\bar{p}}_x} \lesssim 2^k \mathbf{1}_{k+j \geq 0} (1+2^{2j+2k})^{-4} c^{(0)}_k(0) + 2^k \mathbf{1}_{k+j \leq 0} 2^{\delta |k+j|} c^{(0)}_k(0),$$

for $s \in [2^{2j-1}, 2^{2j+1}), k, j \in \mathbb{Z}$. Then we get by Bernstein inequality that

$$\int_{0}^{\infty} \|\phi_{s}\|_{L_{t}^{p}L_{x}^{\infty}} ds' \lesssim \sum_{k \in \mathbb{Z}} \sum_{j \leq -k} 2^{2j+k} 2^{\frac{2k}{p}} c_{k}^{(0)}(0) 2^{\delta|k+j|} + \sum_{k \in \mathbb{Z}} \sum_{j \geq -k} 2^{2j+k} 2^{\frac{2k}{p}} (1+2^{k+j})^{-8} c_{k}^{(0)}(0)$$
$$\lesssim \sum_{k \in \mathbb{Z}} 2^{(\frac{2}{p}-1)k} c_{k}^{(0)}(0).$$
(7.19)

Taking $\tilde{p} \in (2, 4)$ such that $|\frac{2}{\tilde{p}} - 1| \le \frac{1}{8}\delta$, one finds (7.19) is finite by (7.17). Hence, there exists a $p \in (4, \infty)$ such that

$$\int_0^\infty \|\phi_s\|_{L^p_t L^\infty_x} ds' \lesssim 1.$$
(7.20)

Step 1.3. We aim to prove

$$\lim_{t \to \infty} \int_0^\infty \|\phi_s(t)\|_{L^\infty_x} ds' = 0.$$
(7.21)

If (7.21) fails, then for some $\rho > 0$, there exists a time sequence $\{t_{\nu}^{1}\}$, such that $\lim_{\nu \to \infty} t_{\nu}^{1} = \infty$,

$$\int_0^\infty \|\phi_s(t_\nu^1)\|_{L^\infty_x} ds' > \varrho, \ \forall \nu \in \mathbb{Z}_+.$$

$$(7.22)$$

We can also assume $t_{\nu}^1 \leq t_{\nu+1}^1 - 4$ for any $\nu \in \mathbb{Z}_+$. Thus by (7.20) there must exist a sufficiently large constant N and a time sequence $\{t_{\nu}^2\}$ such that

$$t_{\nu}^{1} - 1 \le t_{\nu}^{2} \le t_{\nu}^{1} + 1 \tag{7.23}$$

$$\int_0^\infty \|\phi_s(t_\nu^2)\|_{L^\infty_x} ds' \le \frac{1}{8}\varrho, \ \forall \nu \ge N.$$
(7.24)

Step 2. On the other hand, we have

$$\partial_t \phi_s = D_t \phi_s - A_t \phi_s = D_s \phi_t - A_t \phi_s$$

= $\Delta \phi_t + \sum_{i=1,2} 2A_i \partial_i \phi_t + A_i A_i \phi_t + \phi_t \partial_i A_i + \mathcal{R}(\phi_i, \phi_t) \phi_i - A_t \phi_s.$

Using Proposition 6.2 ((6.12), (6.13)) with $b_k(\sigma)$ replaced by $c_k^{(1)}(\sigma)$ and similar results for succeeding iterations, we see

$$\begin{split} \|\phi_t\|_{L^4_t L^{\infty}_x} &\lesssim \sum_{k \in \mathbb{Z}} c_k^{(2)}(\frac{3}{2}) \lesssim 1 \\ \|\partial_x \phi_t\|_{L^4_t L^{\infty}_x} &\lesssim \sum_{k \in \mathbb{Z}} c_k^{(6)}(\frac{5}{2}) \lesssim 1 \\ \|\partial_x^2 \phi_t\|_{L^4_t L^{\infty}_x} &\lesssim \sum_{k \in \mathbb{Z}} c_k^{(10)}(\frac{7}{2}) \lesssim 1, \end{split}$$

since as before one has

$$\sum_{k\in\mathbb{Z}} 2^{\frac{1}{2^{j+4}}\delta|k|} c_k^{(j)}(\sigma) \lesssim 1.$$

And by the same reason there hold

$$\begin{aligned} \|\partial_x^2 \phi_l\|_{L^4_t L^\infty_x} &\lesssim s^{-\frac{5}{4}} \sum_{k \in \mathbb{Z}} c^{(1)}_k(1) \lesssim s^{-\frac{5}{4}} \\ \|\partial_x \phi_l\|_{L^4_t L^\infty_x} &\lesssim s^{-\frac{3}{4}} \sum_{k \in \mathbb{Z}} c^{(1)}_k(1) \lesssim s^{-\frac{3}{4}}. \end{aligned}$$

Meanwhile, Lemma 3.3 shows

$$\begin{split} \|\phi_i\|_{L^{\infty}} &\lesssim (1+s)^{-\frac{3}{4}}, \ i=1,2\\ \|\partial_x^j A_i\|_{L^{\infty}} &\lesssim (1+s)^{-\frac{3}{4}-\frac{1}{2}j}, \ j=0,1. \end{split}$$

Thus we arrive at

$$\|\Delta\phi_t\|_{L^4_tL^\infty_x} + \sum_{i=1,2} \|2A_i\partial_i\phi_t + A_iA_i\phi_t + \phi_t\partial_iA_i + \mathcal{R}(\phi_i,\phi_t)\phi_i\|_{L^4_tL^\infty_x} \leq 1.$$

For the rest $A_t \phi_s$, by the proof of Lemma 6.1 and its analogies in succeeding iterations, we see

$$\begin{aligned} \|A_t\|_{L^4_t L^\infty_x} &\lesssim s^{-\frac{1}{4}} \sum_{k \in \mathbb{Z}} c_k^{(1)}(1) \lesssim s^{-\frac{1}{4}} \\ \|A_t\|_{L^4_t L^\infty_x} &\lesssim s^{-\frac{3}{4}} \sum_{k \in \mathbb{Z}} c_k^{(0)}(0) \lesssim s^{-\frac{3}{4}}. \end{aligned}$$

Hence, (7.18) implies

$$\int_0^\infty \|A_t\phi_s\|_{L^2_t L^\infty_x} ds' \lesssim 1.$$

Therefore, we conclude in this step that there exists a decomposition of $\partial_t \phi_s = I_1 + I_2$ such that

$$\int_{0}^{\infty} \|I_{1}\|_{L^{4}_{t}L^{\infty}_{x}} ds' \leq 1, \quad \int_{0}^{\infty} \|I_{2}\|_{L^{2}_{t}L^{\infty}_{x}} ds' \leq 1.$$
(7.25)

Step 3. (7.25) and (7.23) show

$$\int_{0}^{\infty} \|\phi_{s}(t_{\nu}^{1}) - \phi_{s}(t_{\nu}^{2})\|_{L_{x}^{\infty}} ds' \lesssim \int_{0}^{\infty} \left(\|I_{1}\|_{L_{t}^{2}L_{x}^{\infty}([t_{\nu}^{2}-1,t_{\nu}^{2}+1]\times\mathbb{R}^{d})} + \|I_{2}\|_{L_{t}^{2}L_{x}^{\infty}([t_{\nu}^{2}-1,t_{\nu}^{2}+1]\times\mathbb{R}^{d})}\right) ds'.$$

$$(7.26)$$

Then as $\nu \to \infty$, (7.25) further implies the RHS of (7.26) goes to zero. Thus (7.24) yields

$$\int_0^\infty \|\phi_s(t_v^1)\|_{L^\infty_x} ds' \leq \frac{1}{4}\varrho,$$

for ν sufficiently large, which contradicts with (7.22). So we have verified (7.21). Similar to (7.21) we also have

$$\lim_{t\to-\infty}\int_0^\infty \|\phi_s(t)\|_{L^\infty_x}ds'=0.$$

Then (1.4) follows by (7.14).

7.5 **Proof of (1.5)**

The proof of (1.5) can be reduced to the following lemma.

Lemma 7.2. Given s > 0, there exists a function $f_s : \mathbb{R}^2 \to \mathbb{C}^n$ belonging to \dot{H}^1 such that

$$\lim_{t\to\infty} \|\phi_s(t) - e^{it\Delta} f_s\|_{\dot{H}^1_x} = 0$$

Moreover, f_s satisfies

$$||f_s||_{\dot{H}^1_x} \lesssim \mathbf{1}_{s \in [0,1]} + \mathbf{1}_{s \ge 1} s^{-\frac{3}{2}}.$$

Now, let's prove (1.5) by assuming Lemma 7.2. Recall that

$$\phi_i = -\int_s^\infty (\partial_i \phi_s + A_i \phi_s) ds'.$$

Since $||A||_{L^{\infty}_{s,t}L^{2}_{x}} \leq 1$, (7.21) shows

$$\lim_{t\to\infty} \|\int_s^\infty |A_i\phi_s| ds'\|_{L^2_x} = 0.$$

Then Lemma 7.2 yields

$$\lim_{t \to \infty} \|\phi(0, t, x) - \nabla e^{it\Delta} f_+\|_{L^2_x} = 0, \tag{7.27}$$

where $f_+ \in \dot{H}^1$ is defined by

$$f_+ = -\int_0^\infty f_s ds'.$$

Let \mathcal{P} denote the isometric embedding of \mathcal{N} into \mathbb{R}^N . Recall that $\{e_\alpha, Je_\alpha\}_{\alpha=1}^n$ denotes the caloric gauge. Then the caloric gauge condition shows

$$\sum_{l=1}^{2n} |d\mathcal{P}(e_l) - d\mathcal{P}(e_l^{\infty})| \lesssim \int_0^\infty |\phi_s| ds'.$$

which combined with (7.21) implies for s = 0

$$\lim_{t \to \infty} \|d\mathcal{P}(e_l) - d\mathcal{P}(e_l^{\infty})\|_{L_x^{\infty}} = 0, \ \forall l = 1, ..., 2n.$$
(7.28)

Thus, we deduce from

$$\partial_{j}u = \sum_{\alpha=1}^{n} \Re(\phi_{j}^{\alpha})e_{\alpha} + \Im(\phi_{j}^{\alpha})Je_{\alpha},$$

that for s = 0

$$\begin{split} \|d\mathcal{P}(\nabla u) &- \sum_{\alpha=1}^{n} \mathfrak{R}(\nabla e^{it\Delta} f_{+})^{\alpha} d\mathcal{P}(e_{\alpha}^{\infty}) - \sum_{\alpha=1}^{n} \mathfrak{I}(\nabla e^{it\Delta} f_{+})^{\alpha} d\mathcal{P}(Je_{\alpha}^{\infty})\|_{L^{2}_{x}} \\ &\lesssim \|\phi - \nabla e^{it\Delta} f_{+}\|_{L^{2}_{x}} + \||e^{it\Delta} \nabla f_{+}\| d\mathcal{P}(e) - d\mathcal{P}(e^{\infty})\|\|_{L^{2}_{x}} \\ &+ \||\phi - \nabla e^{it\Delta} f_{+}\| d\mathcal{P}(e) - d\mathcal{P}(e^{\infty})\|\|_{L^{2}_{x}}. \end{split}$$

Therefore, (7.28) and (7.27) give

$$\lim_{t\to\infty} \|d\mathcal{P}(\nabla u) - \sum_{\alpha=1}^n \Re(\nabla e^{it\Delta}f_+)^\alpha d\mathcal{P}(e_\alpha^\infty) - \Im(\nabla e^{it\Delta}f_+)^\alpha d\mathcal{P}(Je_\alpha^\infty)\|_{L^2_x} = 0$$

Then, letting $\vec{v}_{\alpha} = d\mathcal{P}(e_{\alpha}^{\infty}), \vec{v}_{\alpha+n} = d\mathcal{P}(Je_{\alpha}^{\infty})$, we get

$$\lim_{t \to \infty} \|u(t) - \sum_{j=1}^{n} \Re(e^{it\Delta}f_{+})^{j} \vec{v}_{j} - \sum_{j=1}^{n} \Im(e^{it\Delta}f_{+})^{j} \vec{v}_{j+n}\|_{\dot{H}^{1}_{x}} = 0.$$
(7.29)

Thus, (1.5) follows form (7.29) by setting

$$h^{j}_{+} := f^{j}_{+} \vec{v}_{j}, \ g^{j}_{+} := f^{j}_{+} \vec{v}_{j+n}, \ j = 1, ..., n.$$

Now, let's prove Lemma 7.2. The convenient way to verify Lemma 7.2 is to introduce the so-called Schrödinger map tension field $Z := \phi_s - i\phi_t$. Then the heat tension field ϕ_s satisfies for any $s \ge 0$

$$(i\partial_t + \Delta)\phi_s = \mathbf{N} \tag{7.30}$$

$$\mathbf{N} := -\left(\sum_{k=1}^{2} \partial_k A_k\right)\phi_s - \sum_{j=1}^{2} 2A_j \partial_j \phi_s - A_j A_j \phi_s + i \partial_s Z + \sum_{j=1}^{2} \mathcal{R}(\phi_j, \phi_s)\phi_j.$$
(7.31)

And the Schrödinger map tension field Z satisfies the heat equation

$$\begin{aligned} (\partial_s - \Delta)Z &= (\sum_{k=1}^2 \partial_k A_k)Z + \sum_{j=1}^2 [2A_j \partial_j Z + A_j A_j Z] \\ &+ \sum_{j=1}^2 [\mathcal{R}(Z, \phi_j)\phi_j + i\mathcal{R}(\phi_j, \phi_s)\phi_j - \mathcal{R}(\phi_j, i\phi_s)\phi_j] \\ Z(0, t, x) &= 0. \end{aligned}$$
(7.32)

To prove Lemma 7.2, it suffices to verify

$$||\{2^k || P_k \mathbf{N} ||_{N_k}\}||_{\ell^2} \leq (1+s)^{-\frac{3}{2}},$$

where **N** is given by (7.31). Except for the $\partial_s Z$ term in **N**, the other terms have been handled with before. It remains to dominate $||P_k \partial_s Z||_{N_k}$. In fact, one can prove a stronger result for *Z*:

$$\|\{(1+2^{2k})2^k \|P_k Z\|_{L^{\frac{4}{3}}}\}\|_{\ell^2} \lesssim (1+s)^{-\frac{3}{2}}.$$
(7.33)

We see (7.33) follows by bootstrap and (7.32). Therefore, Lemma 7.2 follows.

7.6 Conclusion

Hence, we have finished the proof of Theorem 1.1 and Theorem 1.2.

8 Appendix A. Bilinear estimates

Lemma 8.1. Let $S : \mathbb{R}^N \to \mathbb{R}$ be a smooth function in $y \in \mathbb{R}^N$ and $f : (-T, T) \times \mathbb{R}^2 \to \mathbb{R}^N$ be smooth w.r.t. $(t, x) \in (-T, T) \times \mathbb{R}^2$. Let

$$\mu_k = \sum_{|k_1 - k| \le 20} 2^{k_1} ||P_{k_1} f||_{L^{\infty}_{L^2_x}}.$$
(8.1)

Assume that $||f||_{L_x^{\infty}} \leq 1$ and $\sup_{k \in \mathbb{Z}} \mu_k \leq 1$. Then

$$2^{k} \|P_{k}S(f)(\partial_{a}f\partial_{b}f)\|_{L_{t}^{\infty}L_{x}^{2}} \leq 2^{k} \sum_{k_{1} \leq k} \mu_{k_{1}} 2^{k_{1}} \mu_{k} + \sum_{k_{2} \geq k} 2^{2k} \mu_{k_{2}}^{2} + a_{k} \left(\sum_{k_{1} \leq k} 2^{k_{1}} \mu_{k_{1}}\right)^{2} + \sum_{k_{2} \geq k} 2^{2k} 2^{-k_{2}} a_{k_{2}} \mu_{k_{2}} \sum_{k_{1} \leq k_{2}} 2^{k_{1}} \mu_{k_{1}}.$$

$$(8.2)$$

where $\{a_k\}$ denotes

$$a_k := \|\nabla P_k(S(f))\|_{L^\infty_t L^2_x}.$$
(8.3)

Proof. The same proof of [[4], Lemma 8.2] shows

$$\begin{aligned} &2^{k} \| P_{k} S(f) (\partial_{a} f \partial_{b} f) \|_{L_{t}^{\infty} L_{x}^{2}} \\ &\lesssim 2^{2k} \sum_{k_{1} \leq k} \mu_{k_{1}} 2^{k} \mu_{k} + \sum_{k_{2} \geq k} 2^{-2(k_{2}-k)} 2^{2k_{2}} \mu_{k_{2}}^{2} \\ &+ a_{k} (\sum_{k_{1} \leq k} 2^{k_{1}} \mu_{k_{1}})^{2} + \sum_{k_{2} \geq k} 2^{2k} 2^{-2k_{2}} 2^{k_{2}} a_{k_{2}} \mu_{k_{2}} \sum_{k_{1} \leq k_{2}} 2^{k_{1}} \mu_{k_{1}}. \end{aligned}$$

The only difference is that we use

$$\|P_k(S(f))\|_{L^2_x} \le 2^{-k} \|\nabla P_k(S(f))\|_{L^2_x}$$

when S(f) lies in the high frequency w.r.t. $\partial_a f \partial_b f$, and the trivial bound

$$\|P_k(S(f))\|_{L^\infty_x} \leq 1$$

when S(f) lies in the relatively low frequency.

Denote $H^{\infty,\infty}(T)$ the set of functions f defined in $(t, x) \in [-T, T] \times \mathbb{R}^2$ satisfying $\partial_t^{b_1} \partial_x^{b_2} f \in L^2([-T, T] \times \mathbb{R}^2)$ for any $b_1, b_2 \in \mathbb{N}$.

Lemma 8.2 ([4],Lemma 5.1). Given $\mathcal{L} \in \mathbb{Z}_+$, $\omega \in [0, \frac{1}{2}]$, $T \in (0, 2^{2\mathcal{L}}]$. Suppose that $f, g \in H^{\infty,\infty}(T)$, let

$$\alpha_k := \sum_{|k-k'| \leq 20} \|f_{k'}\|_{S_{k'}^{\omega}(T) \cap F_{k'}(T)}, \ \beta_k := \sum_{|k-k'| \leq 20} \|g_{k'}\|_{S_{k'}^0(T)},$$

If $|k_1 - k_2| \le 8$, *then*

$$\|P_{k}(P_{k_{1}}fP_{k_{2}}g)\|_{F_{k}(T)\cap S_{k}^{\frac{1}{2}}(T)} \leq 2^{\frac{kd}{2}}2^{(k_{2}-k)(\frac{2d}{d+2}-\omega)}\alpha_{k_{1}}\beta_{k_{2}}.$$
(8.4)

If $|k - k_1| \le 4$, *then*

$$\|P_k(gP_{k_1}f)\|_{F_k(T) \cap S_k^{\frac{1}{2}}(T)} \lesssim \|g\|_{L^{\infty}} \alpha_{k_1}.$$
(8.5)

Lemma 8.3 ([4],Lemma 5.4). *Given* $\mathcal{L} \in \mathbb{Z}_+, \omega \in [0, \frac{1}{2}], T \in (0, 2^{2\mathcal{L}}]$. *Then for* $f, g \in H^{\infty,\infty}(T)$

$$\|P_{k}(fg)\|_{L^{4}_{l,x}} \lesssim \sum_{l \le k} 2^{l} (\mathbf{a}_{l} \mathbf{b}_{k} + 2^{\frac{1}{2}(k-l)} \mathbf{a}_{k} \mathbf{b}_{l}) + 2^{k} \sum_{l \ge k} 2^{-\omega(l-k)} \mathbf{a}_{l} \mathbf{b}_{l}.$$
(8.6)

where we denote

$$\mathbf{a}_{k} := \sum_{|l-k| \le 20} ||P_{k}f||_{S_{l}^{\omega}(T)}, \ \mathbf{b}_{k} := \sum_{|l-k| \le 20} ||P_{k}g||_{L_{t,x}^{4}(T)}.$$
(8.7)

Lemma 8.4 ([4],Lemma 5.4). Given $\mathcal{L} \in \mathbb{Z}_+$, $\omega \in [0, \frac{1}{2}]$, $T \in (0, 2^{2\mathcal{L}}]$. Suppose that $f, g \in H^{\infty,\infty}(T)$, $P_k f \in S_k^{\omega}(T)$, $P_k g \in L^4_{t,x}$ for all $k \in \mathbb{Z}$. Let

$$\mu_k := \sum_{|l-k| \le 20} \|P_k f\|_{S_l^{\omega}}(T), \ \nu_k := \sum_{|l-k| \le 20} \|P_k g\|_{L^4_{t,x}(T)}.$$
(8.8)

If $|k_2 - k| \le 4$, $k_1 \le k - 4$, *then*

$$\|P_k(f_{k_1}g_{k_2})\|_{L^4_{tx}} \lesssim 2^{k_1}\mu_{k_2}\nu_k.$$
(8.9)

If $|k_1 - k| \le 4, k_2 \le k - 4$, then

$$\|P_k(f_{k_1}g_{k_2})\|_{L^4_{t,x}} \lesssim 2^{k_2} 2^{\frac{1}{2}(k-k_2)} \mu_k \nu_{k_2}.$$
(8.10)

If $|k_1 - k_2| \le 8$, $k_1, k_2 \ge k - 4$, then

$$\|P_k(f_{k_1}g_{k_2})\|_{L^4_{t,x}} \lesssim 2^{k(1+\omega)} 2^{-\omega k_2} \mu_{k_2} \nu_{k_2}.$$
(8.11)

Lemma 8.5 ([4], Lemma 6.3). • If $|l - k| \le 80$ and $f \in F_l(T)$, then

$$\|P_k(gf)\|_{N_k(T)} \leq \|g\|_{L^2_t L^2_x} \|f\|_{F_l(T)}.$$
(8.12)

- $\|P_k(gf)\|_{N_k(T)} \lesssim 2^{\frac{l-k}{2}} \|g\|_{L^2_t L^2_v} \|f\|_{F_l(T)}.$ (8.13)
- If $k \leq l 80$ and $f \in G_l(T)$, then

• If $l \le k - 80$ and $f \in F_l(T)$, then

$$\|P_k(gf)\|_{N_k(T)} \leq 2^{\frac{k-1}{6}} \|g\|_{L^2_t L^2_x} \|f\|_{G_l(T)}.$$
(8.14)

Lemma 8.6 ([4], Lemma 6.5).
• If
$$k \le l$$
 and $f \in F_k(T), g \in F_l(T)$ then
 $\|fg\|_{L^2_{t_x}} \le \|f\|_{F_k(T)} \|g\|_{F_l(T)}.$
(8.15)

1. 1

• If
$$k \leq l$$
 and $f \in F_k(T), g \in G_l(T)$ then

$$\|fg\|_{L^{2}_{tx}} \leq 2^{\frac{k-l}{2}} \|f\|_{F_{k}(T)} \|g\|_{G_{l}(T)}.$$
(8.16)

9 Appendix B. Proof of Remained Claims

It seems that the following blow-up criterion was not explicitly written down in literature of SMF. This result is well-known in energy critical heat flows. For completeness, we give a proof.

Proposition 9.1. Suppose that $u_0 \in H_Q^L$ with $L \ge 4$ is the initial data to SMF. If in the time interval [-T, T], the SMF solution *u* satisfies

$$\|u(t)\|_{L^{\infty}_{tx}(T)} \le B < \infty, \tag{9.1}$$

then u has the bound

$$\|u(t)\|_{L^{\infty}_{x}H^{L}_{x}} \le C(B, T, \|u_{0}\|_{H^{L}_{x}}) < \infty.$$
(9.2)

As a corollary, if (9.1) holds then u can be extended beyond [-T, T] to $C([-T - \rho, T + \rho]; H_Q^L)$ for some $\rho > 0$.

Proof. Recall the tension field $\tau(u) = \sum_{j=1}^{2} \nabla_j \partial_j u$. By integration by parts,

$$\int_{\mathbb{R}^2} \langle \tau(u), \tau(u) \rangle dx = \int_{\mathbb{R}^2} \sum_{j,k=1}^2 \langle \nabla_j \partial_j u, \nabla_k \partial_k u \rangle dx$$
$$= \int_{\mathbb{R}^2} \langle \nabla_k \nabla_j \partial_k u, \nabla_k \nabla_j \partial_k u \rangle + \int_{\mathbb{R}^2} O(|du|^4) dx.$$
(9.3)

Since *u* solves *SMF*, by integration by parts, we get

$$\begin{split} &\frac{d}{dt} \int_{\mathbb{R}^2} \langle \tau(u), \tau(u) \rangle dx = 2 \sum_{j=1}^2 \int_{\mathbb{R}^2} \langle \nabla_j \partial_j \partial_t u, \tau(u) \rangle dx + \int_{\mathbb{R}^2} O(|du|^2 |\partial_t u| |\tau(u)|) dx \\ &= 2 \sum_{j=1}^2 \int_{\mathbb{R}^2} \langle \nabla_j J \tau(u), \nabla_j \tau(u) \rangle + \int_{\mathbb{R}^2} O(|du|^2 |\partial_t u| |\tau(u)|) dx. \end{split}$$

Since J commutes with ∇_i , $\langle JX, X \rangle = 0$, we then arrive at

$$\frac{d}{dt} \|\tau(u)\|_{L^2_x}^2 \lesssim \|du\|_{L^\infty_{t,x}}^2 \|\tau(u)\|_{L^2_x}^2$$

Gronwall inequality and (9.1) show

$$\|\tau(u)\|_{L^2_x} \lesssim e^{Bt} \|\tau(u_0)\|_{L^2_x}$$

Using the energy bound

$$\|\nabla u\|_{L^{\infty}_{t}L^{2}_{x}} \lesssim \|\nabla u_{0}\|_{L^{2}_{x}}$$

and (9.3) give

$$\|u(t)\|_{W^{2,2}} \leq B \|\nabla u_0\|_{L^2_{r}} + e^{Bt} \|\tau(u_0)\|_{L^2_{r}}.$$
(9.4)

By integration by parts,

$$\begin{split} &\int_{\mathbb{R}^2} \langle \nabla_i \tau(u), \nabla_i \tau(u) \rangle dx = \int_{\mathbb{R}^2} \sum_{j,k=1}^2 \langle \nabla_i \nabla_j \partial_j u, \nabla_i \nabla_k \partial_k u \rangle dx = \int_{\mathbb{R}^2} \langle \nabla_i \nabla_j \partial_k u, \nabla_i \nabla_j \partial_k u \rangle \\ &+ \int_{\mathbb{R}^2} O(|du|^3 |\nabla^2 du| + |\nabla u|^2 |\nabla du|^2 + |\nabla du| |du|^2) dx. \end{split}$$

Thus we have

$$\begin{aligned} \|\nabla^{2} du(t)\|_{L^{2}_{x}}^{2} &\lesssim \|\nabla\tau(u)\|_{L^{2}_{x}}^{2} + \|du\|_{L^{6}_{x}}^{6} + \|du\|_{L^{\infty}_{x}}^{2} \|\nabla du\|_{L^{2}_{x}}^{2} + \|du\|_{L^{4}_{x}}^{2} \|\nabla du\|_{L^{2}_{x}}^{2} \\ &\lesssim \|\nabla\tau(u)\|_{L^{2}_{x}}^{2} + C(B, t, \|u_{0}\|_{W^{2,2}}) \end{aligned}$$

$$\tag{9.5}$$

And applying integration by parts furthermore gives

$$\begin{split} &\frac{1}{2} \frac{d}{dt} ||\nabla \tau(u)||_{L_x^2}^2 = \sum_{i,j} \langle \nabla_i \nabla_j \partial_j u, \nabla_t \nabla_i \nabla_j \partial_j u \rangle \\ &= \sum_{i,j} \langle \nabla_i \tau(u), \nabla_i \nabla_j \nabla_j \partial_t u \rangle + \int_{\mathbb{R}^2} |\nabla \tau(u)| |\nabla \partial_t u| |du|^2 dx \\ &+ \int_{\mathbb{R}^2} |\nabla^2 du| \nabla du| |\partial_t u| |du| dx + \int_{\mathbb{R}^2} |du|^3 |\partial_t u| |\nabla^2 du| dx \\ &\lesssim - \langle \sum_i \nabla_i \nabla_i \tau(u), J \sum_j \nabla_j \nabla_j \tau(u) \rangle + B^2 ||\nabla \tau(u)||_{L_x^2}^2 + B ||\nabla \tau(u)||_{L_x^2} ||\nabla du||_{L_x^4}^2 \\ &+ B^3 ||\nabla \tau(u)||_{L_x^2} ||\tau(u)||_{L_x^2} \\ &\lesssim B^2 ||\nabla \tau(u)||_{L_x^2}^2 + B ||\nabla \tau(u)||_{L_x^2} ||\nabla^2 du||_{L_x^2} ||\nabla du||_{L_x^2} + B^3 ||\nabla \tau(u)||_{L_x^2} ||\tau(u)||_{L_x^2}. \end{split}$$

Hence, denoting $F(t) = \|\nabla \tau(u)\|_{L^2_{Y}}$, (9.4) and (9.5) show

$$\frac{1}{2}\frac{d}{dt}F^{2}(t) \leq C_{1}(B,T)F(t)[F(t) + C_{2}(B,T)],$$

where $C_1(B, T)$ and $C_2(B, T)$ are smooth functions of B, T. So the Sobolev norm of u has a uniform bound in [-T, T] up to order three. This with the classical local existence theory (see [9] or [30]) implies u can be extended to $[-\rho - T, T + \rho]$ for some $\rho > 0$. And the bounds for the higher order Sobolev norms follow by Theorem 3.3 of [30] or induction. Then by Sobolev embedding u is smooth in $[-\rho - T, T + \rho]$ if $u_0 \in \mathcal{H}_Q$.

References

- [1] I. Bejenaru, On Schrodinger maps, Amer. J. Math. 130, 1033-1065, 2008.
- [2] I. Bejenaru, T. Tataru. Near soliotn evolution for equivariant Schrödigner maps in two spatial dimensions. Memories of American Mathematical Society, 2010.
- [3] I. Bejenaru, A. Ionescu, and C. Kenig. Global existence and uniqueness of Schrödinger maps in dimensions $d \ge 4$. Adv. Math., 215, 263-291, 2007.
- [4] I. Bejenaru, A. Ionescu, C. Kenig, D. Tataru. Global Schrödinger maps in dimensions $d \ge 2$: Small data in the critical Sobolev spaces. Ann. Math., **173**, 1443-1506, 2011.
- [5] I. Bejenaru, A. Ionescu, C. Kenig, D. Tataru. Equivariant Schrödinger maps in two spatial dimensions. Duke Math. J., 162, 1967-2025, 2013.
- [6] I. Bejenaru, A. Ionescu, C. Kenig, D. Tataru. Equivariant Schrödinger Maps in two spatial dimensions: the H² target. Kyoto J. Math. 56, 283-323, 2016.
- [7] N.H. Chang, J. Shatah, and K. Uhlenbeck. Schrödinger maps. Comm. Pure Appl. Math., 53, 590-602, 2000.
- [8] W.Y. Ding, On the Schrödinger flows, pp. 283-291 in Proceedings of the International Congress of Mathematicians (Beijing, 2002), vol.2, Higher Education Press, 2002.
- [9] W.Y. Ding, and Y.D. Wang. Local Schrödinger flow into Kähler manifolds. Sci. China Math., Ser. A 44, 1446-1464, 2001.
- [10] B. Dodson, P. Smith. A controlling norm for energy-critical Schrödinger maps. Transactions of AMS, 367(10), 7193-7220, 2015.
- [11] S. Gustafson, K. Kang, T.P. Tsai. Asymptotic stability of harmonic maps under the Schrödinger flow. Duke Math. J., **145**, 537-583, 2008.
- [12] S. Gustafson, K. Nakanishi, T.P. Tsai, Asymptotic stability, concentration, and oscillation in harmonic map heat-flow, Landau-ifshitz, and Schröinger maps on *R*². Comm. Math. Phys. **300**, 205-242, 2010.
- [13] A. D. Ionescu and C. E. Kenig, Low-regularity Schrodinger maps, Differential Integral Equations, 19, 1271-1300, 2006.
- [14] A. D. Ionescu and C. E. Kenig, Low-regularity Schrodinger maps. II. Global well-posedness in dimen sions $d \ge 3$, Comm. Math. Phys. **271**, 523-559, 2007.
- [15] C. E. Kenig, G. Ponce, and L. Vega, Smoothing effects and local existence theory for the generalized nonlinear Schrödinger equations, Invent. Math. 134, 489-545, 1998.
- [16] S. Klainerman and M. Machedon, Space-time estimates for null forms and the local existence theorem, Comm. Pure Appl. Math. 46, 1221-1268, 1993.
- [17] S. Klainerman and S. Selberg, Remark on the optimal regularity for equations of wave maps type, Comm. Partial Differential Equations, 22, 901-918, 1997.
- [18] J. Krieger, Global regularity of wave maps from R^{2+1} to H^2 . Small energy, Comm. Math. Phys. 250, 507-580, 2004.
- [19] H. Koch, D. Tataru, and M. Visan. Dispersive Equations and Nonlinear Waves. Oberwolfach Seminars Springer Basel, 2014. Part II, Chapter 5.
- [20] F. Linares and G. Ponce, On the Davey-Stewartson systems, Ann. Inst. H. Poincare Anal. Non Lineaire 10, 523-548, 1993.
- [21] F. Merle, P. Raphael, I. Rodnianski. Blowup dynamics for smooth data equivariant solutions to the critical Schrödinger map problem. Invent. Math., **193**(2), 249-365, 2013.
- [22] A. Nahmod, A. Stefanov, K. Uhlenbeck, On Schrödinger maps, Comm. Pure Appl. Math. 56, 114-151, 2003. Erratum: On Schrödinger map. Comm. Pure Appl. Math. 57, 833-839.
- [23] A. Nahmod, A. Stefanov, K. Uhlenbeck, On the well-posedness of the wave map problem in high dimensions, Comm. Anal. Geom. 11, 49-83, 2003.
- [24] A. Nahmod, J. Shatah, L. Vega, and C. Zeng, Schrodinger maps and their associated frame systems. Int. Math. Res. Notices, 2007, No. 21, article ID rnm088, 2007.
- [25] P. Smith, Geometric renormalization below the ground state. Int. Math. Res. Notices, **2012**, 3800-3844, 2011.
- [26] G. Perelman. Blow up dynamics for equivariant critical Schrödinger maps. Comm. Math. Phys., 330, 69-105, 2014.
- [27] L. Landau, and E. Lifshitz. On the theory of the dispersion of magnetic permeability in ferromagnetic bodies. Phys. Z. Sovietunion, 8, 153-169, 1935.
- [28] Z. Li. Global Schrödinger map flows to Kähler manifolds with small data in critical Sobolev spaces: High dimensions. arxiv preprint 2019.
- [29] Z. Li. On global dynamics of Schrödinger map flows on hyperbolic planes near harmonic maps. arxiv preprint 2020.
- [30] H. McGahagan. An approximation scheme for Schrödinger maps. Comm. Partial Differential Equations, 32, 375-400, 2007.
- [31] I. Rodnianski, Y. Rubinstein, G. Staffilani. On the global well-posedness of the one-dimensional Schrödinger map flow. Anal. PDE, **2**, 187-209, 2009.
- [32] P.L. Sulem, C. Sulem, C. Bardos. On the continuous limit for a system of classical spins. Comm. Math. Phys., 107, 431-454, 1986.
- [33] T. Tao. Global regularity of wave maps, I: small critical Sobolev norm in high dimension. Int. Math. Res. Not. IMRN, 2001, 299-328, 2001.
- [34] T. Tao. Global Regularity of Wave Maps II. Small Energy in Two Dimensions. Comm. Math. Phys., 224, 443-544, 2001.
- [35] T. Tao, Gauges for the Schrödinger map, unpublished. http://www.math.ucla.edu/ tao/preprints/Expository.
- [36] T. Tao. Global regularity of wave maps. III-VII. arXiv preprint 2008-2009.

- [37] T. Tao. Nonlinear Dispersive Equations, Local and global analysis. CBMS Regional Conf. Series Math. no. 106, Published for the Conference Board of the Mathematical Sciences, Washington DC, 2006.
- [38] T. Tao. Geometric renormalization of large energy wave maps. Journees equations aux derivees partielles, 1-32, 2004.
- [39] D. Tataru. On global existence and scattering for the wave maps equation. Amer. J. Math., **123**, 37-77, 2001.
- [40] D. Tataru. Local and global results for the wave maps I. Comm. Partial Differential Equations, 23, 1781-1793, 1998.
- [41] D. Tataru. Rough solutions for the wave maps equation. Amer. J. Math., 127, 293-377, 2005.
- [42] Y.L. Zhou, B.L. Guo, S.B. Tan. Existence and uniqueness of smooth solution for system of ferromagnetic chain. Sci. China Math., Ser. A, 34, 257-266, 1991.