

MMMMMM YYYY, Volume VV, Issue II. [doi: 10.18637/jss.v000.i00](http://dx.doi.org/10.18637/jss.v000.i00)

CVEK: Robust Estimation and Testing for Nonlinear Effects using Kernel Machine Ensemble

Wenying Deng Harvard University **Jeremiah Zhe Liu** Harvard University **Erin Lake**

Harvard University

Brent A. Coull Harvard University

Abstract

The R package CVEK introduces a suite of flexible machine learning models and robust hypothesis tests for learning the joint nonlinear effects of multiple covariates in limited samples. It implements the *Cross-validated Ensemble of Kernels* (CVEK)[\(Liu and Coull](#page-29-0) [2017\)](#page-29-0), an ensemblebased kernel machine learning method that adaptively learns the joint nonlinear effect of multiple covariates from data, and provides powerful hypothesis tests for both main effects of features and interactions among features. The R Package CVEK provides a flexible, easy-to-use implementation of CVEK, and offers a wide range of choices for the kernel family (for instance, polynomial, radial basis functions, Matérn, neural network, and others), model selection criteria, ensembling method (averaging, exponential weighting, cross-validated stacking), and the type of hypothesis test (asymptotic or parametric bootstrap). Through extensive simulations we demonstrate the validity and robustness of this approach, and provide practical guidelines on how to design an estimation strategy for optimal performance in different data scenarios.

Keywords: robust hypothesis test, nonlinear effect, Gaussian process, CVEK, R, kernel ensemble.

1. Introduction

In recent years, kernel machine methods have seen widespread application in biomedical studies for learning the complex, nonlinear effects of multivariate genetic or environmental exposures. Given data and feature $\{y, \mathbf{x}\}\$, practitioners are often interested in constructing a function $\hat{h}(\mathbf{x})$ that best describes the data generation mechanism $y = \mu + h(\mathbf{x}) + \epsilon$. Further, given groups of features $\{x_1, x_2\} \subset x$, interest may focus on conducting a hypothesis test for either the overall effect of x_1 on y , or on the interaction effect between two feature groups \mathbf{x}_1 and \mathbf{x}_2 based on *h*.

Traditionally, kernel machine regression (KMR) handles this task by specifying a kernel

function $k(\mathbf{x}, \mathbf{x}')$ that gives rise to a large function space \mathcal{H}_k , and estimates the data generation function h through regularized regression. Using the estimated h , hypothesis tests for feature effects can become simply variance component tests by interpreting the KMR as a linear mixed model [\(Liu, Lin, and Ghosh](#page-29-1) [2007\)](#page-29-1). Given unlimited data and proper choice of kernel family, kernel machine regression enjoys a theoretical guarantee of learning arbitrary continuous target functions defined over a compact input space [\(Micchelli, Xu,](#page-29-2) [and Zhang](#page-29-2) [2006\)](#page-29-2), thereby inducing a valid hypothesis test with correct Type I error.

In practice, however, the performance of KMR in limited samples is known to be extremely sensitive to the choices of the kernel function. To guarantee reasonable performance, *k* must be selected carefully so that its mathematical properties reflect those of the data generating mechanism. Selecting an overly smooth kernel will result in an h[†] that underfits the data, inducing an invalid hypothesis test with inflated Type I error. Selecting an overly flexible kernel function will lead to \hat{h} that overfits the data, leading to an underpowered test. For most applications in epidemiology and the natural sciences, it is often difficult to specify a kernel family *a priori* for a complex, nonlinear data generation mechanism *h*. This leads to challenges in both estimation and testing.

CVEK is an R package that provides a suite of robust estimation and testing procedures that adaptively learn the proper kernel function from the data through the use of kernel ensembling, thereby achieving unbiased effect estimation and valid hypothesis testing in limited samples. Specifically, CVEK implements the *Cross-validated Ensemble of Kernels* (CVEK) [\(Liu and Coull](#page-29-0) [2017;](#page-29-0) [Liu, Lee, Lin, Valeri, Christiani, Bellinger, Wright,](#page-29-3) [Mazumdar, and Coull](#page-29-3) [2019\)](#page-29-3), an ensemble-based kernel machine learning procedure that automatically discerns the most appropriate kernel for the data using a cross-validated approach. CVEK offers a range of choices in base kernel families, model selection criteria, and ensemble methods, so the practitioner can flexibly design a modeling strategy for the data at hand. Section [2](#page-1-0) introduces CVEK and such choices in detail. Section [3](#page-7-0) presents CVEK's implementation of asymptotic and bootstrap-based hypothesis tests for the overall effect of a single feature/group, as well as the interaction effect between features/groups. Section [4](#page-11-0) provides a hands-on tutorial in learning and testing gene-environment interactions with **CVEK**. Section [5](#page-16-0) discusses the impact of choice of estimation strategy (i.e., choices of model selection criteria and ensemble strategy) on the performance of the resulting hypothesis test. This is demonstrated through a comprehensive simulation study that evaluates validity (i.e., Type I error) and power of the implemented tests using diverse modeling strategies under a wide range of data generation mechanisms.

2. Robust estimation using kernel ensemble via CVEK

2.1. Gaussian process regression

Assume we observe data from *n* independent subjects. For the i^{th} subject, let y_i be a continuous response, \mathbf{x}_i be the set of p continuous features that have a potentially nonlinear effect on y_i . We assume that the outcome y_i depends on features \mathbf{x}_i through the data generating model

$$
y_i = \mu + h(\mathbf{x}_i) + \epsilon_i
$$
, where $\epsilon_i \stackrel{iid}{\sim} N(0, \lambda)$.

We assume $h : \mathbb{R}^p \to \mathbb{R}$ follows the Gaussian process (GP) prior $\mathcal{GP}(0, k)$ governed by the positive definite kernel function *k*, such that the function evaluated at the observed covariates follows the multivariate normal (MVN) distribution

$$
\mathbf{h} = [h(\mathbf{x}_1), \dots, h(\mathbf{x}_n)] \sim MVN(\mathbf{0}, \mathbf{K})
$$

with covariance matrix having elements $\mathbf{K}_{ij} = k(\mathbf{x}_i, \mathbf{x}_j)$. Under this construction, the predictive distribution of *h* evaluated at the samples is also multivariate normal,

$$
\mathbf{h} | \{y_i, \mathbf{x}_i\}_{i=1}^n \sim MVN(\mathbf{h}^*, \mathbf{K}^*),
$$

$$
\mathbf{h}^* = \mathbf{K}(\mathbf{K} + \lambda \mathbf{I})^{-1}(\mathbf{y} - \boldsymbol{\mu}),
$$

$$
\mathbf{K}^* = \mathbf{K} - \mathbf{K}(\mathbf{K} + \lambda \mathbf{I})^{-1}\mathbf{K}.
$$

To understand the impact of λ and k on h^* , recall that operationally, a Gaussian process can be understood as the Bayesian version of kernel machine regression, where **h** ∗ equivalently arises from the optimization problem

$$
\mathbf{h}^* = \underset{h \in \mathcal{H}_k}{\operatorname{argmin}} \ \|\mathbf{y} - \boldsymbol{\mu} - h(\mathbf{x})\|^2 + \lambda \|h\|_{\mathcal{H}}^2,
$$

where \mathcal{H}_k is the reproducing kernel Hilbert space (RKHS) generated by kernel function k . From this perspective, \mathbf{h}^* is the element in a spherical ball in \mathcal{H}_k that best approximates the observed data **y**. The mathematical properties (e.g., smoothness, spectral density) of **h** ∗ are governed by the kernel function *k*. The norm of \mathbf{h}^* , $\|h\|_{\mathcal{H}}^2$, is constrained by the tuning parameter *λ*.

Consequently, choice of kernel function k and tuning parameter λ critically impact the quality of the final estimate **h**. To this end, **CVEK** offers a wide range of choices for the kernel family and tuning parameter selection strategies, which we review below.

Kernel Family

In this section we review some commonly-used kernel functions that are implemented in CVEK, including three stationary covariance functions (Gaussian radial basis function, Matérn and rational quadratic), as well as non-stationary covariance functions (polynomial and neural network).

• **Intercept Kernel** (intercept)

The intercept kernel implements the simplest of all kernel functions

$$
k(\mathbf{x}, \mathbf{x}') = 1,
$$

which corresponds to the intercept under the (generalized) linear model.

• **Linear Kernel** (linear)

The linear kernel is

$$
k(\mathbf{x}, \mathbf{x}') = \langle \mathbf{x}, \mathbf{x}' \rangle,
$$

where $\langle \mathbf{x}, \mathbf{x}' \rangle := \mathbf{x}^\top \mathbf{x}'$, the inner product of **x** and **x'**. It is useful when dealing with large, sparse data vectors **x**.

• **Polynomial Kernel** (polynomial)

The polynomial kernel is

$$
k(\mathbf{x}, \mathbf{x}') = (1 + \langle \mathbf{x}, \mathbf{x}' \rangle)^p,
$$

which is commonly used with support vector machines (SVMs). The polynomial kernel becomes the intercept kernel when $p = 0$, and the linear kernel when $p = 1$.

• **Gaussian Radial Basis Function (RBF) Kernel** (rbf)

The Gaussian radial basis function kernel is

$$
k(\mathbf{x}, \mathbf{x}') = \exp\Big(-\frac{|\mathbf{x} - \mathbf{x}'|^2}{2l^2}\Big),\,
$$

where *l* is the *characteristic length-scale*. It is typically used when knowledge about the form of the exposure-response relationship exists.

• **Matérn Kernel** (matern)

The Matérn kernel is

$$
k(\mathbf{x}, \mathbf{x}') = \frac{2^{1-\nu}}{\Gamma(\nu)} \Big(\frac{\sqrt{2\nu|\mathbf{x} - \mathbf{x}'|}}{l}\Big)^{\nu} K_{\nu} \Big(\frac{\sqrt{2\nu|\mathbf{x} - \mathbf{x}'|}}{l}\Big)
$$

with positive parameters ν and *l*, where K_{ν} is a modified Bessel function [\(Abramowitz](#page-28-0) [1974\)](#page-28-0). The Matérn kernel is commonly used to define the statistical covariance between measurements made at two points that are $|\mathbf{x} - \mathbf{x}'|$ units distant from each other. The most interesting cases for machine learning are $\nu = 3/2$ and $\nu = 5/2$, for which

$$
k_{\nu=3/2}(\mathbf{x}, \mathbf{x}') = \left(1 + \frac{\sqrt{3}|\mathbf{x} - \mathbf{x}'|}{l}\right) \exp\left(-\frac{\sqrt{3}|\mathbf{x} - \mathbf{x}'|}{l}\right),
$$

$$
k_{\nu=5/2}(\mathbf{x}, \mathbf{x}') = \left(1 + \frac{\sqrt{5}|\mathbf{x} - \mathbf{x}'|}{l} + \frac{5|\mathbf{x} - \mathbf{x}'|^2}{3l^2}\right) \exp\left(-\frac{\sqrt{5}|\mathbf{x} - \mathbf{x}'|}{l}\right),
$$

since for $\nu = 1/2$ the process becomes very rough, and for $\nu \geq 7/2$, in the absence of explicit prior knowledge about the existence of higher order derivatives, it is probably very hard from finite noisy training examples to distinguish between values of $\nu \geq 7/2$.

• **Rational Quadratic Kernel** (rational)

The rational quadratic kernel is

$$
k(\mathbf{x}, \mathbf{x}') = \left(1 + \frac{|\mathbf{x} - \mathbf{x}'|^2}{2\alpha l^2}\right)^{-\alpha}
$$

with α , $l > 0$ can be seen as a *scale mixture* (an infinite sum) of squared exponential (SE) covariance functions with different characteristic length-scales (sum of covariance functions is a valid covariance). The limit of the rational quadratic covariance as $\alpha \to \infty$ is the SE covariance function with characteristic length-scale *l*.

• **Neural Network Kernel** (nn)

The neural network kernel is

$$
k(\mathbf{x}, \mathbf{x}') = \frac{2}{\pi} \sin^{-1} \left(\frac{2\sigma \tilde{\mathbf{x}}^\top \tilde{\mathbf{x}}'}{\sqrt{(1 + 2\sigma \tilde{\mathbf{x}}^\top \tilde{\mathbf{x}})(1 + 2\sigma \tilde{\mathbf{x}}'^\top \tilde{\mathbf{x}}')}} \right),
$$

where $\tilde{\mathbf{x}} = (1, x_1, ..., x_d)^\top$ is an augmented input vector and σ is the covariance coefficient.

Model Selection Criteria

In practice, the tuning parameter λ is selected by minimizing certain objective functions in a process known as *Model Selection* that measures the model's degree of "appropriateness" given certain values of *λ*. Depending on the specific criteria, such "appropriateness" can be the distance between the current model and the true model, the model's out-of-sample prediction error, or the model likelihood. Here we review some commonly used model selection criteria.

In kernel machine regression, most of the model selection criteria can be expressed as a function of λ through the model's predictive "hat" matrix \mathbf{A}_{λ} :

$$
\mathbf{A}_{\lambda} = \mathbf{K}(\mathbf{X}, \mathbf{X})[\mathbf{K}(\mathbf{X}, \mathbf{X}) + \lambda \mathbf{I}]^{-1}.
$$

In this way, $tr(\mathbf{A}_{\lambda})$ is the effective number of model parameters. It decreases monotonically with $\lambda > 0$. For notational simplicity we assume **y** is centered: $\mathbf{y} = \mathbf{y} - \hat{\boldsymbol{\mu}}$, where $\hat{\mu} = \frac{1}{n}$ $\frac{1}{n}\sum_{i=1}^n y_i$.

• **Akaike Information Criterion** (AIC) and its small-sample variant (AICc)

AIC handles the trade-off between the goodness of fit of the model and the simplicity of the model:

$$
\lambda_{AIC} = \underset{\lambda \in \Lambda}{\text{argmin}} \Big\{ \log \ \mathbf{y}^\top (\mathbf{I} - \mathbf{A}_{\lambda})^2 \mathbf{y} + \frac{2[\text{tr}(\mathbf{A}_{\lambda}) + 2]}{n} \Big\},\,
$$

where Λ is the set that contains all possible values of λ . When *n* is small (e.g., $n/p < 40$) [\(Burnham and Anderson](#page-28-1) [2002\)](#page-28-1), extreme overfitting is possible, giving small bias/ large variance estimates. The AIC small-sample correction [\(Hurvich and Tsai](#page-29-4) [1989;](#page-29-4) [Hurvich](#page-29-5) [Clifford M., Simonoff Jeffrey S., and Tsai Chih Ling](#page-29-5) [2002\)](#page-29-5) is derived by modifying the penalty as the product of the original penalty, $2[\text{tr}(\mathbf{A}_{\lambda}) + 2]/n$ and $n/[n - \text{tr}(\mathbf{A}_{\lambda}) - 3]$, where we plug in \mathbf{A}_{λ} and $\hat{\sigma}^2$. In this case, we obtain our small-sample objective function AICc,

$$
\lambda_{AICc} = \underset{\lambda \in \Lambda}{\text{argmin}} \Big\{ \log \mathbf{y}^\top (\mathbf{I} - \mathbf{A}_{\lambda})^2 \mathbf{y} + \frac{2[\text{tr}(\mathbf{A}_{\lambda}) + 2]}{n - \text{tr}(\mathbf{A}_{\lambda}) - 3} \Big\}.
$$

• **Bayesian Information Criterion** (BIC)

The Bayesian information criterion is

$$
\lambda_{BIC} = \underset{\lambda \in \Lambda}{\text{argmin}} \Big\{ \log \mathbf{y}^\top (\mathbf{I} - \mathbf{A}_{\lambda})^2 \mathbf{y} + \frac{\log(n)[\text{tr}(\mathbf{A}_{\lambda}) + 2]}{n} \Big\}.
$$

The formula for BIC is similar to the one for AIC. It is more conservative in the selection process with penalty for number of parameters to be $log(n)$, instead of 2 for AIC.

• **Leave-one-out Cross Validation** (loocv)

Suppose we perform *K*-fold cross-validation, which partitions observations into *K* groups, $\kappa(1), \ldots, \kappa(K)$, and calculates $\mathbf{A}_{\lambda} K$ times, each time leaving out group $\kappa(i)$, to get

$$
\mathbf{A}_{\lambda}^{-\kappa(1)}, \mathbf{A}_{\lambda}^{-\kappa(2)}, \ldots, \mathbf{A}_{\lambda}^{-\kappa(K)}
$$

.

A value of $K = 10$ is very common in the field of applied machine learning. For $\mathbf{A}_{\lambda}^{-\kappa(i)}$ λ , λ , λ cross-validated residuals are calculated on the observations in $\kappa(i)$, which did not contribute to the estimation of **A**. The objective function estimates prediction error and is the sum of the squared cross-validated residuals,

$$
\lambda_{K-CV} = \underset{\lambda \in \Lambda}{\text{argmin}} \left\{ \log \sum_{i=1}^{K} [\mathbf{y}_{\kappa(i)} - \mathbf{A}_{\lambda}^{-\kappa(i)} \mathbf{y}_{\kappa(i)}]^{\top} [\mathbf{y}_{\kappa(i)} - \mathbf{A}_{\lambda}^{-\kappa(i)} \mathbf{y}_{\kappa(i)}] \right\}.
$$

Note that loocv corresponds to $K = n$. In this case, we can write our objective function as [\(Golub, Heath, and Wahba](#page-28-2) [1979\)](#page-28-2),

$$
\lambda_{loocv} = \underset{\lambda \in \Lambda}{\text{argmin}} \left\{ \log \mathbf{y}^\top [\mathbf{I} - \text{diag}(\mathbf{A}_\lambda) - \frac{1}{n} \mathbf{I}]^{-1} (\mathbf{I} - \mathbf{A}_\lambda)^2 [\mathbf{I} - \text{diag}(\mathbf{A}_\lambda) - \frac{1}{n} \mathbf{I}]^{-1} \mathbf{y} \right\}.
$$
 (1)

• **Generalized Cross Validation** (GCV) and its small-sample variant (GCVc) In [\(1\)](#page-5-0), if we approximate each diagonal element of the matrix \mathbf{A}_{λ} , $A_{\lambda[ii]}$ with its mean $tr(A_\lambda)$ $\frac{A_{\lambda}}{n}$, in a sense we give equal weight to all observations. We then get the generalized cross validation objective function,

$$
\lambda_{GCV} = \underset{\lambda \in \Lambda}{\operatorname{argmin}} \Big\{ \log \ \mathbf{y}^\top (\mathbf{I} - \mathbf{A}_{\lambda})^2 \mathbf{y} - 2 \log[1 - \frac{\text{tr}(\mathbf{A}_{\lambda})}{n} - \frac{1}{n}] \Big\},\tag{2}
$$

where $\cdot - \frac{1}{n}$ $\frac{1}{n}$ ['] is due to GCV counting μ as part of the model complexity, but not σ^2 . This motivates the proposed small-sample correction to GCV [\(Boonstra, Mukherjee, and Taylor](#page-28-3) [2015\)](#page-28-3), which does count σ^2 as a parameter,

$$
\lambda_{GCVc} = \underset{\lambda \in \Lambda}{\operatorname{argmin}} \Big\{ \log \ \mathbf{y}^\top (\mathbf{I} - \mathbf{A}_{\lambda})^2 \mathbf{y} - 2 \log[1 - \frac{\text{tr}(\mathbf{A}_{\lambda})}{n} - \frac{2}{n}]_+ \Big\}.
$$

• **Generalized Maximum Profile Marginal Likelihood** (gmpml)

The generalized maximum profile marginal likelihood is defined as

$$
\lambda_{gmpml} = \underset{\lambda \in \Lambda}{\text{argmin}} \Big\{ \log \ \mathbf{y}^\top (\mathbf{I} - \mathbf{A}_{\lambda}) \mathbf{y} - \frac{1}{n-1} \log |\mathbf{I} - \mathbf{A}_{\lambda}| \Big\}.
$$

This is a likelihood-based method, where λ is interpreted as the variance component of a mixed-effects model.

2.2. Cross-Validated ensemble of kernels

Traditional applications of Gaussian process estimate *h* using a single kernel function *k* for $h \in \mathcal{H}_k$, therefore imposing *a priori* assumption on the mathematical properties of *h*

through *k*. In such case, choosing a kernel function that is too restrictive or too flexible will lead to either model underfit or overfit, rendering the subsequent hypothesis tests invalid. Recently, [Liu and Coull](#page-29-0) [\(2017\)](#page-29-0); Liu *[et al.](#page-29-3)* [\(2019\)](#page-29-3) addressed the challenge by proposing Cross-Validated Ensemble of Kernels (CVEK), an ensemble-based estimator that adaptively learns the form of the kernel function from data. CVEK estimates *h* using the ensemble of GP predictions generated from a library of (fixed) base kernel functions ${k_d}_{d=1}^D$,

$$
\hat{h}(\mathbf{x}) = \sum_{d=1}^{D} u_d \hat{h}_d(\mathbf{x}), \quad \mathbf{u} \in \Delta := \{ \mathbf{u} \mid \mathbf{u} \ge 0, \| \mathbf{u} \|_1 = 1 \},
$$
\n(3)

where \hat{h}_d is the kernel predictor generated by the d^{th} base kernel k_d . The exact algorithm proceeds in three stages as follows (see Algorithm 1, which can be found in Section [6\)](#page-20-0).

Stage 1: Estimate Base Model Cross-Validation Error

For each base kernel in the library ${k_d}_{d=1}^D$, we first standardize the kernel matrix by its trace $\mathbf{K}_d = \mathbf{K}_d/\text{tr}(\mathbf{K}_d)$, and then estimate the prediction based on each kernel as $\hat{\mathbf{h}}_{d,\hat{\lambda}_d} = \mathbf{K}_d(\mathbf{K}_d + \hat{\lambda}_d \mathbf{I})^{-1} \mathbf{y}, d \in \{1, \dots, D\}$, where the tuning parameter $\hat{\lambda}_d$ is selected by minimizing one of the model selection criterion introduced in Section [2.1.2.](#page-4-0) In the case of leave-one-out cross validation (loocv), the cross-validation error can be expressed in closed-form:

$$
CV(\lambda|k_d) = [\mathbf{I} - \text{diag}(\mathbf{A}_{d,\lambda})]^{-1} (\mathbf{y} - \mathbf{\hat{h}}_{d,\lambda}), \text{ where } \mathbf{A}_{d,\lambda} = \mathbf{K}_d(\mathbf{K}_d + \lambda \mathbf{I})^{-1}.
$$

We denote the final estimated loocv error for d^{th} kernel as $\hat{\xi}_d = \text{CV}(\hat{\lambda}_d | k_d)$

Stage 2: Estimate Ensemble

Using the estimated individual model cross-validation errors $\{\hat{\xi}_d\}_{d=1}^D$, we estimate the ensemble weights $\mathbf{u} = \{u_d\}_{d=1}^D$ according to one of the ensemble strategies that will be introduced in Section [2.2.1.](#page-7-1) After estimating \hat{u} , the final ensemble prediction is estimated as:

$$
\widehat{\mathbf{h}} = \sum_{d=1}^{D} \widehat{u}_d \widehat{\mathbf{h}}_d = \sum_{d=1}^{D} \widehat{u}_d \mathbf{A}_{d,\widehat{\lambda}_d} \mathbf{y} = \widehat{\mathbf{A}} \mathbf{y},
$$

where $\hat{\mathbf{A}} = \sum_{d=1}^{D} \hat{u}_d \mathbf{A}_{d,\hat{\lambda}_d}$ is the ensemble hat matrix.

Stage 3: Estimate Ensemble Kernel Matrix

Using the ensemble hat matrix $\hat{\mathbf{A}}$, estimate the ensemble kernel matrix $\hat{\mathbf{K}}$ by solving:

$$
\widehat{\mathbf{K}}(\widehat{\mathbf{K}} + \lambda \mathbf{I})^{-1} = \widehat{\mathbf{A}}.
$$

Specifically, if we denote \mathbf{U}_A and $\{\delta_{A,k}\}_{k=1}^n$ as the eigenvectors and eigenvalues of $\widehat{\mathbf{A}}$, respectively, then the ensemble kernel matrix $\hat{\mathbf{K}}$ adopts the form:

$$
\widehat{\mathbf{K}} = \lambda_{\mathbf{K}} * \left[\mathbf{U}_{A} \text{diag}\left(\frac{\delta_{A,k}}{1 - \delta_{A,k}}\right) \mathbf{U}_{A}^{\top} \right],\tag{4}
$$

where we recommend setting $\lambda_{\mathbf{K}} = \min \left[1, \left(\sum_{k=1}^{n} \frac{\delta_{A,k}}{1 - \delta_{A,k}} \right)^{-1} \right]$.

Ensemble Strategies

We now introduce the choices for an ensemble strategy to be used in Stage 2 of CVEK. Briefly, an ensemble strategy estimates the ensemble weights $\mathbf{u} = \{u_d\}_{d=1}^D$ from individual model cross-validation errors $\{\hat{\xi}_d\}_{d=1}^D$. Choices available are:

- **Averaging Ensemble** (avg) Motivated by existing literature on the omnibus kernel [\(Zhan, Plantinga, Zhao, and Wu](#page-29-6) [2017\)](#page-29-6), a simple way to choose the weights is $u_d = 1/D$ for $d = 1, 2, ...D$.
- **Exponential Weighting** (exp) [Dalalyan and Tsybakov](#page-28-4) [\(2007\)](#page-28-4) proposed estimating **u** using the estimated errors $\{\hat{\xi}_d\}_{d=1}^D$ as:

$$
u_d(\beta) = \frac{\exp(-\|\hat{\xi}_d\|_2^2/\beta)}{\sum_{d=1}^D \exp(-\|\hat{\xi}_d\|_2^2/\beta)}.
$$

From the perspective of optimal model aggregation, the authors showed that under squared loss, the error bound in excessive risk of exponential weighting converges at the fast rate of $O(\frac{1}{n})$ $\frac{1}{n}$). Here, β serves as a tuning parameter to determine how different the ensemble weights are. Increasing *β* results in more similar weights, based on their estimated errors. An infinite *β* results in an averaging ensemble. Exponential weighting can also be viewed as the frequentist version for Bayesian model averaging under Gaussian noise [\(Yang and](#page-29-7) [Dunson](#page-29-7) [2014\)](#page-29-7).

• **Cross-Validated Stacking** (stack)

Alternatively, we can estimate **u** such that it minimizes the overall cross-validation error. After obtaining the estimated errors $\{\hat{\xi}_d\}_{d=1}^D$, we estimate the ensemble weights $\mathbf{u} = \{u_d\}_{d=1}^D$ such that it minimizes the overall error:

$$
\hat{\mathbf{u}} = \underset{\mathbf{u} \in \Delta}{\operatorname{argmin}} \|\sum_{d=1}^{D} u_d \hat{\xi}_d\|^2 \quad \text{where } \Delta = \{\mathbf{u} \mid \mathbf{u} \ge 0, \|\mathbf{u}\|_1 = 1\}.
$$

3. Hypothesis testing for nonlinear effects

3.1. Testing for general nonlinear effect

We use the classical variance component test [\(Lin](#page-29-8) [1997\)](#page-29-8) to construct a testing procedure for the hypothesis about a Gaussian process. Specifically, recall the assumed model:

$$
y_i = \mu + h(\mathbf{x}_i) + \epsilon_i
$$
, where $\epsilon_i \stackrel{iid}{\sim} N(0, \lambda)$.

We are interested in testing the null hypothesis:

$$
H_0: h \in \mathcal{H}_0.
$$

We first express this hypothesis in terms of model parameters. The key to our approach is to assume that *h* lies in a RKHS generated by a *garrote kernel function* $k_{\delta}(\mathbf{z}, \mathbf{z}')$ [\(Maity and](#page-29-9) [Lin](#page-29-9) [2011\)](#page-29-9), which is constructed by including an extra *garrote parameter* δ in a given kernel function. When $\delta = 0$, the garrote kernel function $k_0(\mathbf{x}, \mathbf{x}') = k_\delta(\mathbf{x}, \mathbf{x}')|_{\delta=0}$ generates \mathcal{H}_0 , the space of functions under the null hypothesis. In order to focus on a particular hypothesis of interest, practitioners need only to specify the form of the garrote kernel such that \mathcal{H}_0 corresponds to the null hypothesis. As a result, the general null hypothesis is equivalent to,

$$
H_0: \delta = 0. \tag{5}
$$

We now construct a test statistic \hat{T}_0 for [\(5\)](#page-8-0) by noticing that the garrote parameter δ can be treated as a variance component parameter in a linear mixed model (LMM). This is because the Gaussian process under a garrote kernel can be formulated into the LMM

$$
\mathbf{y} = \boldsymbol{\mu} + \mathbf{h} + \boldsymbol{\epsilon}, \quad \text{where} \quad \mathbf{h} \sim N(\mathbf{0}, \tau \mathbf{K}_{\delta}), \quad \boldsymbol{\epsilon} \sim N(\mathbf{0}, \sigma^2 \mathbf{I}),
$$

and \mathbf{K}_{δ} is the kernel matrix generated by $k_{\delta}(\mathbf{z}, \mathbf{z}')$. Consequently, we can derive a variance component test for H_0 by calculating the squared derivative of the Restricted Maximum Likelihood (REML) with respect to δ under H_0 [\(Lin](#page-29-8) [1997\)](#page-29-8),

$$
\hat{T}_0 = \hat{\tau} * (\mathbf{y} - \hat{\boldsymbol{\mu}})^{\top} \mathbf{V}_0^{-1} [\partial \mathbf{K}_0] \mathbf{V}_0^{-1} (\mathbf{y} - \hat{\boldsymbol{\mu}}),
$$
\n(6)

where $\tau = \frac{\sigma^2}{\lambda}$ $\frac{\partial^2}{\partial \lambda^2}$ and $\mathbf{V}_0 = \hat{\sigma}^2 \mathbf{I} + \hat{\tau} \mathbf{K}_0$. In this expression, $\mathbf{K}_0 = \mathbf{K}_\delta |_{\delta=0}$, and $\partial \mathbf{K}_0$ is the null derivative kernel matrix whose (i, j) th entry is $\frac{\partial}{\partial \delta} k_{\delta}(\mathbf{x}, \mathbf{x}') |_{\delta=0}$. Appendix [E](#page-40-0) also provides the derivation of the REML based test statistic.

Extension for Interaction Testing

In the previous section, we assume that we are able to obtain a k_{δ} that generates \mathcal{H}_0 exactly. However, depending on the exact hypothesis of interest, identifying such a k_0 is not always straightforward. In this section, we revisit the case of interaction testing and consider how to build a k_0 for the hypothesis of interest.

$$
\mathcal{H}_0: h(\mathbf{x}) = h_1(\mathbf{x}_1) + h_2(\mathbf{x}_2),
$$

$$
\mathcal{H}_a: h(\mathbf{x}) = h_1(\mathbf{x}_1) + h_2(\mathbf{x}_2) + h_{12}(\mathbf{x}_1, \mathbf{x}_2),
$$

where h_{12} is the "pure interaction" function that is orthogonal to main effect function h_1 and *h*2. This hypothesis is difficult to formulate with Gaussian process models, since the kernel functions $k(\mathbf{x}, \mathbf{x}')$ in general do not explicitly separate the main and the interaction effect. Therefore rather than directly defining k_0 , we need to first construct \mathcal{H}_0 and \mathcal{H}_a that correspond to the null and alternative hypotheses, respectively, and then identify the garrote kernel function k_{δ} such that corresponds to \mathcal{H}_0 when $\delta = 0$ and \mathcal{H}_a when $\delta > 0$. We build \mathcal{H}_0 using the tensor-product constructions of RKHS on the product domain $(\mathbf{x}_{1,i}, \mathbf{x}_{2,i}) \in \mathbb{R}^{p_1} \times \mathbb{R}^{p_2}$ [\(Gu](#page-28-5) [2013\)](#page-28-5), due to this approach's unique ability to explicitly characterize the space of "pure interaction" functions. Let $\mathbf{1} = \{f \mid f \propto 1\}$ be the RKHS of constant functions, and \mathcal{H}_1 , \mathcal{H}_2 be the RKHS of centered functions for \mathbf{x}_1 , \mathbf{x}_2 respectively. We can then define the full space as $\mathcal{H} = \otimes_{m=1}^2 (\mathbf{1} \oplus \mathcal{H}_m)$. H describes the space of functions that depends jointly on $\{x_1, x_2\}$. It adopts the orthogonal decomposition,

$$
\begin{aligned} \mathcal{H} &= (\mathbf{1} \oplus \mathcal{H}_1) \otimes (\mathbf{1} \oplus \mathcal{H}_2) \\ &= \mathbf{1} \otimes \{\mathcal{H}_1 \oplus \mathcal{H}_2\} \oplus \{\mathcal{H}_1 \otimes \mathcal{H}_2\} = \mathbf{1} \oplus \mathcal{H}_{12}^\perp \oplus \mathcal{H}_{12}, \end{aligned}
$$

where we have denoted $\mathcal{H}_{12}^{\perp} = \mathcal{H}_1 \oplus \mathcal{H}_2$ and $\mathcal{H}_{12} = \mathcal{H}_1 \otimes \mathcal{H}_2$ respectively. We see that \mathcal{H}_{12} is indeed the space of "pure interaction" functions, since \mathcal{H}_{12} contains functions on the product domain $\mathbb{R}^{p_1} \times \mathbb{R}^{p_2}$, but is orthogonal to the space of additive main effect functions \mathcal{H}_{12}^{\perp} . To summarize, we have identified two function spaces \mathcal{H}_0 and \mathcal{H}_a that have the desired interpretation,

$$
\mathcal{H}_0=\mathcal{H}_{12}^{\perp},\quad \mathcal{H}_a=\mathcal{H}_{12}^{\perp}\oplus\mathcal{H}_{12}.
$$

We are now ready to identify the garrote kernel $k_{\delta}(\mathbf{x}, \mathbf{x}')$. To this end, we notice that both \mathcal{H}_{12}^\perp and \mathcal{H}_{12} are composite spaces built from basis RKHSs using direct sum and tensor products. If we denote $k_m(\mathbf{x}_m, \mathbf{x}'_m)$ as the reproducing kernel associated with \mathcal{H}_m , we can construct kernel functions for composite spaces \mathcal{H}_{12}^{\perp} and \mathcal{H}_{12} as

$$
k_0(\mathbf{x}, \mathbf{x}') = k_1(\mathbf{x}_1, \mathbf{x}_1) + k_2(\mathbf{x}_2, \mathbf{x}_2),
$$

$$
k_{12}(\mathbf{x}, \mathbf{x}') = k_1(\mathbf{x}_1, \mathbf{x}_1) k_2(\mathbf{x}_2, \mathbf{x}_2).
$$

Hence, the garrote kernel function for \mathcal{H}_a is

$$
k_{\delta}(\mathbf{x}, \mathbf{x}') = k_0(\mathbf{x}, \mathbf{x}') + \delta k_{12}(\mathbf{x}, \mathbf{x}').
$$

Finally, using the chosen form of the garrote kernel function, the (i, j) th element of the null derivative kernel matrix is $\frac{\partial}{\partial \delta} k_{\delta}(\mathbf{x}, \mathbf{x}')\Big|_{\delta=0} = k_{12}(\mathbf{x}, \mathbf{x}')$, i.e., the null derivative kernel matrix *∂***K**⁰ is simply the kernel matrix **K**¹² that corresponds to the interaction space. Thus, we also call it the alternative kernel. Therefore the score test statistic \hat{T}_0 in [\(6\)](#page-8-1) simplifies to

$$
\hat{T}_0 = \hat{\tau} * (\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}})^{\top} \mathbf{V}_0^{-1} \mathbf{K}_{12} \mathbf{V}_0^{-1} (\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}),
$$

where $\mathbf{V}_0 = \hat{\sigma}^2 \mathbf{I} + \hat{\tau} \mathbf{K}_0$.

There exist multiple approaches for estimating the null distribution of \hat{T}_0 . Specifically, assuming ∂ **K**₀ is a fixed matrix (e.g., in the case of interaction testing, fixing the k_{12} to be a linear kernel), one can derive the closed form expression of the asymptotic distribution of \hat{T}_0 (i.e., a mixture of χ^2 distributions). Alternatively, one can approximate the null distribution in a data-driven manner using bootstrap sampling. Compared to the bootstrap, the asymptotic method is advantageous in that it is more powerful if the null model is correctly specified, but is restrictive in that it requires the alternative kernel *∂***K**⁰ to be fixed *a priori*. This requirement further prevents practitioners from improving test power due to the need to learn the optimal *∂***K**⁰ from data. The bootstrap test, on the other hand, does not require *∂***K**⁰ to be fixed and hence does not suffer from this limitation. We introduce these two types of procedures in detail in Section [3.2,](#page-9-0) and discuss a data-adaptive strategy for estimating *∂***K**⁰ in Section [3.3.](#page-10-0)

3.2. Null distribution estimation

Asymptotic Approximation

Assuming fixed *∂***K**, the null distribution of *T*ˆ can be approximated with a scaled chi-square distribution $\kappa \chi^2_{\nu}$ using the Satterthwaite method that matches the first two moments of *T*,

$$
\kappa * \nu = \mathcal{E}(T) = \hat{\tau} * \text{tr}(\mathbf{V}_0 \partial \mathbf{K}_0), \quad 2 * \kappa^2 * \nu = Var(T) = \hat{\mathbf{I}}_{\delta \delta}.
$$

This procedure yields the solution

$$
\hat{\kappa} = \hat{\mathbf{I}}_{\delta\delta}/[\hat{\tau} * \text{tr}(\mathbf{V}_0^{-1}\partial \mathbf{K}_0)], \quad \hat{\nu} = [\hat{\tau} * \text{tr}(\mathbf{V}_0^{-1}\partial \mathbf{K}_0)]^2/(2 * \hat{\mathbf{I}}_{\delta\theta}),
$$

where $\hat{\mathbf{I}}_{\delta\delta} = \mathbf{I}_{n,\delta\delta} - \mathbf{I}_{\delta\theta}^{\top} \mathbf{I}_{\delta\theta}^{-1} \mathbf{I}_{\delta\theta}$ is the efficient information of δ REML. $\mathbf{I}_{\delta\delta}$, $\mathbf{I}_{\theta\theta}$ and $\mathbf{I}_{\delta\theta}$ are sub-matrices of the REML information matrix. Numerically more accurate, but computationally less efficient, approximation methods are also available [\(Bodenham and](#page-28-6) [Adams](#page-28-6) [2016\)](#page-28-6).

Finally, the p-value of this test is calculated using the tail probability of $\hat{\kappa}\chi^2_{\hat{\nu}}$,

$$
p = P(\hat{\kappa}\chi_{\hat{\nu}}^2 > \hat{T}) = P(\chi_{\hat{\nu}}^2 > \hat{T}/\hat{\kappa}).
$$

A complete summary of the proposed testing procedure is available in Algorithm 2, which can be found in Section [6.](#page-20-0)

Parametric Bootstrap

When the sample size is small, we make valid inferences about a population using resampling. A commonly used resampling method is the bootstrap, which gives valid tests in small to moderate sample sizes.

Testing in a regression model framework requires computing the distribution of the test statistic under the null hypothesis. We approximate this null distribution using a bootstrap sample of the test statistic resampled from the fit of the null model. For instance, when testing [\(5\)](#page-8-0), we first fit the model under the null,

$$
E(\mathbf{y}^{\star}) = \mathbf{K}_0(\mathbf{K}_0 + \lambda \mathbf{I})^{-1} \mathbf{y} = \mathbf{A}_0 \mathbf{y},
$$

and generate \mathbf{Y}^* with a random noise, whose variance is also estimated. We then compute the test statistic for this simulated sample, and repeat this process *B* times. The empirical distribution of the test statistic provides an estimate of the test statistic's distribution under the null. Correspondingly, p-values are calculated as the proportion of simulated test statistics that are as or more extreme than the observed value.

Like the classical bootstrap, this approach samples from a distribution based on the observed data, but the simulations are from a fitted parametric model rather than the empirical distribution. To obtain a valid test, the fitted parametric model is chosen so that the null hypothesis is satisfied. A complete summary of the proposed testing procedure is available in Algorithm 3, which can be found in Section [6.](#page-20-0)

3.3. Strategy for estimating alternative kernel

As mentioned previously, the asymptotic test requires the alternative kernel *∂***K**⁰ in the test statistic [\(6\)](#page-8-1) to be fixed *a priori*, due to the need to approximate the null distribution analytically. Traditionally, *∂***K**⁰ is fixed to be either linear or a specific kernel family with fixed hyperparameters (e.g., Gaussian RBF family with fixed length-scale). Consequently, the form of the alternative kernel needs to be correctly specified in order to sufficiently describe the interaction effect, since otherwise a misspecified alternative kernel may lead to a loss of power. On the other hand, the bootstrap test allows $\partial \mathbf{K}_0$ to be estimated adaptively from the data in order to better represent the alternative hypothesis space.

To this end, we propose a strategy for data-adaptive estimation of the alternative kernel in the bootstrap test. Specifically, we estimate the alternative kernel using the ensemble weights $\{\hat{\mu}_d\}_{d=1}^D$ obtained from the ensemble procedure as described in [\(3\)](#page-6-0), i.e.,

$$
\partial \mathbf{K}_0 = \sum_{d=1}^D \hat{u}_d \ast \partial \mathbf{K}_{0,d}.\tag{7}
$$

Consequently, if the true interaction effect is not linear, a bootstrap test with an adaptively estimated alternative kernel can better describe the interaction effect from the data, and therefore will have better power when compared to an asymptotic or bootstrap test with the *∂***K**⁰ fixed to be a linear kernel. In Section [5.2,](#page-19-0) we empirically investigate the effectiveness of this approach through extensive simulation.

4. The CVEK package

Using a library of base kernels, **CVEK** learns the generating function from data by directly minimizing the ensemble model's error, and tests whether the data is generated by the RKHS under the null hypothesis. Section [4.1](#page-11-1) presents a simple example to conduct Gaussian process regression and hypothesis testing using the cvek function on simulated data. Section [4.2](#page-14-0) shows a real-world application where we use CVEK to understand whether the per capita crime rate impacts the relationship between local socioeconomic status and the housing price in Boston, MA, U.S.A.

4.1. Tutorial using simulated dataset

Generate Data and Define Model

We generate a simulated dataset using the **linear** kernel, and set the relative interaction strength to be 0.2. The outcome y_i is generated as,

$$
y_i = h_1(\mathbf{x}_{i,1}) + h_2(\mathbf{x}_{i,2}) + 0.2 * h_{12}(\mathbf{x}_{i,1}, \mathbf{x}_{i,2}) + \epsilon_i,
$$

where h_1 , h_2 , h_{12} are sampled from RKHSs \mathcal{H}_1 , \mathcal{H}_2 , \mathcal{H}_{12} , generated using the corresponding linear kernel. We standardize all sampled functions to have unit form, so that 0*.*2 represents the strength of interaction relative to the main effect.

```
> set.seed(0726)
> n <- 60 # including training and test observations
> d < -4> int_effect <- 0.2
> data \leq matrix(rnorm(n * d), ncol = d)
> Z1 <- data[, 1:2]
> Z2 <- data[, 3:4]
>
> kern <- generate_kernel(method = "linear")
> w <- rnorm(n)> w12 < -rnorm(n)
```

```
> K1 < - \text{kern}(Z1, Z1)> K2 <- kern(Z2, Z2)
> K1 <- K1 / sum(diag(K1)) # standardize kernel
> K2 < - K2 / sum(diag(K2))> h0 <- K1 \frac{9}{8} * W + K2 \frac{9}{8} * W
> h0 <- h0 / sqrt(sum(h0 \hat{ } 2)) # standardize main effect
>
> h1_prime <- (K1 * K2) %*% w12 # interaction effect
>
> # standardize sampled functions to have unit norm, so that 0.2
> # represents the interaction strength relative to main effect
> Ks \leq svd(K1 + K2)
> len <- length(Ks$d[Ks$d / sum(Ks$d) > .001])
> U0 <- Ks$u[, 1:len]
> h1_prime_hat <- fitted(lm(h1_prime ~ U0))
> h1 <- h1_prime - h1_prime_hat
>
> h1 <- h1 / sqrt(sum(h1 \hat{ } 2)) # standardize interaction effect
> Y <- h0 + int_effect * h1 + rnorm(1) + rnorm(n, 0, 0.01)
> data <- as.data.frame(cbind(Y, Z1, Z2))
> colnames(data) <- c("y", paste0("z", 1:d))
>
> data train \leq data[1:40, ]
> data_test <- data[41:60, ]
```
The resulting data look as follows.

Now we can apply the cvek function to conduct Gaussian process regression. Table 1 is a detailed list of all the arguments of the function cvek.

Suppose we want our model library to contain three kernels: linear, polynomial with p=2, and rbf with l=1 (the effective parameter for polynomial is p and the effective parameter for rbf is l, so we can set anything to l for polynomial kernel and p for rbf kernel). We then first apply define_library.

```
> kern par \leq data.frame(method = c("linear", "polynomial", "rbf"),
                         1 = rep(1, 3), p = 1:3, stringsAsFactors = FALSE)> # define model library
> kern_func_list <- define_library(kern_par)
```


Table 1: Arguments of the function cvek().

The null model is then $y \sim z1 + z2 + k(z3, z4)$.

> formula <- $y \sim z1 + z2 + k(z3, z4)$

Estimation and Testing

With all these parameters specified, we can conduct Gaussian process regression.

> est_res <- cvek(formula, kern_func_list = kern_func_list, data = data_train) > est_res\$lambda

[1] 4.539993e-05

> est_res\$u_hat

[1] 0.994864707 0.000000000 0.005135293

We can see that the ensemble weight assigns 0.99 to the linear kernel, which is the true kernel. This illustrates the accuracy and efficiency of the CVEK method.

We next specify the testing procedure. Note that we can use the same function cvek to perform hypothesis testing, as we did for estimation, but we need to provide formula_test, which is the user-supplied formula indicating the additional alternative effect (e.g., interactions) to test for. Specifically, we will first show how to conduct the classic score test by specifying test="asymp", followed by a bootstrap test where we specify test="boot", and the number of bootstrap samples B=200.

```
> formula_test <- y - k(z1, z2):k(z3, z4)>
> cvek(formula, kern_func_list = kern_func_list,
+ data = data_train, formula_test = formula_test,
+ mode = "loocv", strategy = "stack",
+ beta exp = 1, lambda = exp(seq(-10, 5)),
      test = "asymp", alt\_kernel\_type = "ensemble",
+ verbose = FALSE)$pvalue
            [,1]
[1,] 1.493613e-08
> cvek(formula, kern_func_list = kern_func_list,
+ data = data_train, formula_test = formula_test,
+ mode = "loocv", strategy = "stack",
      beta\_exp = 1, lambda = exp(seq(-10, 5)),
+ test = "boot", alt_kernel_type = "ensemble",
+ B = 200, verbose = FALSE)$pvalue
```
 $\lceil 1 \rceil$ 0

Both tests come to the same conclusion. At the significance level 0*.*05, we reject the null hypothesis that there's no interaction effect, which matches our data generation mechanism. Additionally, we can predict new outcomes based on estimation results est_res.

```
> y_pred <- predict(est_res, data_test[, 2:5])
> data_test_pred <- cbind(y_pred, data_test)
```


4.2. Detecting nonlinear interactions in Boston Housing Prices

In this section, we show an example of using the cvek test to detect nonlinear interactions between socioeconomic factors that contribute to housing price in the city of Boston, Massachusetts, USA. We consider the Boston dataset (available in the MASS package), which is collected by the U.S Census Service about the median housing price (medv) in Boston, along with additional variables describing local socioeconomic information such as per capita crime rate, proportion of non-retail business, number of rooms per household, etc. Table 2 lists the 14 variables.

Variables	Description
crim	Per capita crime rate by town.
zn	Proportion of residential land zoned for lots over 25,000 sq.ft.
indus	Proportion of non-retail business acres per town.
chas	Charles River dummy variable $(= 1$ if tract bounds river; 0 otherwise).
nox	Nitric oxides concentration (parts per 10 million).
rm	Average number of rooms per dwelling.
age	Proportion of owner-occupied units built prior to 1940.
dis	Weighted distances to five Boston employment centres.
rad	Index of accessibility to radial highways.
tax	Full-value property-tax rate per USD 10,000.
ptratio	Pupil-teacher ratio by town.
black	$1000(B-0.63)^2$ where B is the proportion of blacks by town.
lstat	Percentage of lower socioeconomic status in the population.
medv	Median value of owner-occupied homes in USD 1000's.

Table 2: Variables of the Boston dataset.

Here we use cvek to study whether the per capita crime rate (crim) impacts the relationship between the local socioeconomic status (lstat) and the housing price. The null model is,

$$
medv \sim \mathbf{x}^\top \boldsymbol{\beta} + k(crim) + k(lstat),
$$

where $\mathbf{x}^\top =$ (1, zn, indus, chas, nox, rm, age, dis, rad, tax, ptratio, black), and $k()$ is specified as a semi-parametric model with a model library that includes linear and rbf kernels with $l = 1$. This inclusion of nonlinearity (i.e., the rbf kernel) is important, since per classic results in the macroeconomics literature, the crime rates and socioeconomic status of local community are known to have a nonlinear association with the local housing price [\(Harrison and Rubinfeld](#page-28-7) [1978\)](#page-28-7).

```
> kern par \leq data.frame(method = c("linear", "rbf"),
l = rep(1, 2), p = 1:2, strings As Factors = FALSE)> # define kernel library
> kern_func_list <- define_library(kern_par)
```
To this end, the hypothesis regarding whether the crime rate (crim) impacts the association between local socioeconomic status (lstat) and the housing price (medv) is equivalent to testing whether there exists a nonlinear interaction between crim and lstat in predicting

medv, i.e.,

```
\mathcal{H}_0: medv \sim \mathbf{x}^\top \boldsymbol{\beta} + k(crim) + k(lstat)\mathcal{H}_a: \textit{medv} \sim \mathbf{x}^\top \boldsymbol{\beta} + k(\textit{crim}) + k(\textit{lstat}) + k(\textit{crim}): k(\textit{lstat}).
```
To test this hypothesis using cvek, we specify the null model using formula, and specify the additional interaction term (*k*(*crim*) : *k*(*lstat*)) in the alternative model using formula_test, as shown below:

```
> formula <- medv \sim zn + indus + chas + nox + rm + age + dis +
   rad + tax + ptratio + black + k(crim) + k(lstat)> formula_test <- medv ~ k(crim):k(lstat)
> fit bos<- cvek(formula, kern func list = kern func list, data = Boston,
                formula test = formula test,
+ lambda = exp(seq(-3, 5)), test = "asymp")
```
Given the fitted object (fit bos), the p-value of the cvek test can be extracted as below:

> fit bos\$pvalue

[1] 4.614106e-06

Since $p < 0.05$, we reject the null hypothesis that there's no crim: latat interaction, and conclude that the data does suggest an impact of the crime rate on the relationship between the local socioeconomic status and the housing price. In Appendix [F,](#page-42-0) we provide additional code showing how to visualize the interaction effect from a cvek model.

5. Simulation and practical recommendations

In this section, we conduct a simulation study to evaluate the finite-sample performance of the CVEK hypothesis tests in a setting that is analogous to a typical nutrition-environment interaction study (Liu *[et al.](#page-29-3)* [2019\)](#page-29-3). We generate two groups of input features $(\mathbf{x}_{i,1}, \mathbf{x}_{i,2}) \in \mathbb{R}^{p_1} \times \mathbb{R}^{p_2}$ independently, both within and between group from a standard Gaussian distribution, representing a subject's level of exposure to p_1 environmental pollutants and the levels of a subject's intake of *p*² nutrients. Across all simulation scenarios, we keep $n = 100$, and $p_1 = p_2 = 2$. We generate the outcome y_i as,

$$
y_i = h_1(\mathbf{x}_{i,1}) + h_2(\mathbf{x}_{i,2}) + \delta * h_{12}(\mathbf{x}_{i,1}, \mathbf{x}_{i,2}) + \epsilon_i,
$$
\n(8)

where h_1 , h_2 , h_{12} are sampled from RKHSs H_1 , H_2 , H_{12} , generated using a ground-truth main effect kernel k_{main} (related to h_1 , h_2) and an interaction kernel k_{int} (related to h_{12}), and $\epsilon_i \sim N(0, \sigma^2 = 0.01^2)$. We standardize all sampled functions to have unit form, so that δ represents the strength of interaction relative to the main effect. Additional simulation results for correlated exposures are presented in Appendix [B.](#page-35-0)

For each simulation scenario, we first generate data using δ , k_{main} and k_{int} as in [\(8\)](#page-16-1), then select a *kmodel* to estimate the null model and obtain a p-value using either an asymptotic approximation (Algorithm 2) or parametric bootstrap (Algorithm 3). We repeat each scenario 200 times, and evaluate the test performance using the empirical probability $\hat{P}(p \leq 0.05)$. Under the null hypothesis, a correct test should produce \hat{P} that is smaller or equal to the significance level 0.05. Under the alternative hypothesis $H_a: \delta > 0$, $\hat{P}(p \le 0.05)$ estimates the test's power, and should ideally approach 1.0 quickly as the strength of interaction δ increases.

In this study, we vary the combination of *kmain* (the true main effect) and *kint* (the true interaction effect) to produce data generating functions $h_\delta(\mathbf{x}_{i,1}, \mathbf{x}_{i,2})$ with different smoothness and complexity properties, and vary k_{model} to reflect different common modeling strategies for the null model in addition to using CVEK. We then evaluate how these two aspects impact the Type I error and power of the resulting hypothesis tests.

Specifically, we consider the following seven types of data generation mechanism where $k_{main} = k_{int}$ (i.e., the main effect function and the interaction function belong to the same family). We also consider two additional scenarios where $k_{main} \neq k_{int}$.

The data generation mechanisms are:

- Three simple (non)linear kernels that can be sufficiently modeled using finite-dimensional, parametric functions.
	- **–** linear: *kmain* is a polynomial kernel with degree 1.
	- **–** quadratic: *kmain* is a polynomial kernel with degree 2.
	- **–** cubic: *kmain* is a polynomial kernel with degree 3.
- Four flexible nonlinear kernels that each represent the space of all continuous functions with a prespecified set of mathematical properties (e.g., differentiability or complexity). Data generated by these kernels are usually more difficult to model.
	- **–** rbf_1: A Gaussian RBF kernel with length-scale 1. This kernel represents the space of functions that are *smooth* (i.e., infinitely differentiable) and have reasonable *complexity* (i.e., does not have fast-varying fluctuations that are difficult to model).
	- **–** rbf_0.5: A Gaussian RBF kernel with length-scale 0*.*5. Compared to rbf_1, rbf_0.5 has the same degree of smoothness but is more complex, i.e., has fast-varying local fluctuations.
	- matern_2.5_1: A Matérn $\frac{5}{2}$ kernel with length-scale 1. Compared to rbf_1, matern_2.5_1 has the same degree of complexity but is less smooth, in the sense that it represents the space of twice-differentiable functions, but is not necessarily infinitely differentiable.
	- **–** matern_1.5_0.5: A Matérn ³ 2 kernel with length-scale 0*.*5. Compared to matern_2.5_1, matern_1.5_0.5 is more complex but less smooth. It represents the space of once-differentiable functions.
- Two data generation mechanisms where the true main effect $h_1 + h_2$ and the true interaction effect h_{12} are generated from two separate kernel families (i.e., $k_{main} \neq k_{int}$). The data generating mechanisms are

– quadratic_rbf: The main effects are generated from polynomial with degree 2, while the interaction effect is generated from rbf with length-scale 1. This combination generates data under quadratic main effects but a flexible nonlinear interaction:

$$
y = x_1^2 \beta_1 + x_2^2 \beta_2 + \delta * h_{12}(x_1, x_2) + \epsilon.
$$

– rbf_lnr_0.5: The main effects are generated from rbf with length-scale 0.5, while the interaction effect is generated from polynomial with degree 1. This combination generates data under flexible nonlinear main effects but with a linear interaction:

$$
y = h_1(x_1) + h_2(x_2) + \delta * x_1 x_2 + \epsilon.
$$

The types of model libraries *kmodel* **considered are**:

- **Polynomial**: A library of three parametric, polynomial kernels with degree $p = 1, 2, 3$. This library represents a parametric model with polynomial nonlinearity.
- **RBF** A library of three nonparametric, rbf kernels with length-scale $l = 0.5, 1, 2$. This library represents a nonparametric model with a high degree of smoothness (i.e., infinitely differentiable) that can incorporate more general types of nonlinearity.
- **Polynomial+RBF** A library of three polynomial kernels with $p = 1, 2, 3$ and three rbf kernels with $l = 0.5, 1, 2$. This library represents a semi-parametric model with a mixture of parametric and nonparametric kernels.

To understand how the choices of model-selection criteria and ensemble strategy impact the model performance, for each combination of data generation mechanism and model library, we estimate the null model $y = h_1(x_1) + h_2(x_2)$ under all possible choices of model selection criteria (loocv, AIC, AICc, BIC, GCV, GCVc, gmpml) and ensemble strategy (avg, exp, stack). In Sections [5.1](#page-18-0)[-5.3,](#page-20-1) unless otherwise specified, we report results using loocv for tuning parameter selection and stack for the ensemble strategy, which corresponds to the default setting in CVEK. In general, loocv guarantees correct Type I error, except for Cubic data, which is hard to fit. Also, while correct Type I error is guaranteed, stack leads to better power. We provide the results for the other settings in Appendix [A.](#page-30-0)

Figures 1-4 present the results. They show the estimated $\hat{P}(p < 0.05)$ (y-axis) as a function of interaction strength $\delta \in [0,1]$ (x-axis). Each panel in the figure represents the result for a specific data generating mechanism (Linear, Quadratic, Cubic, RBF $l = 1$, RBF $l = 0.5$, Matérn $\nu = 5/2$ with $l = 1$ and Matérn $\nu = 3/2$ with $l = 0.5$), while the different lines represent results from different modeling choices (e.g., choice of null model model library, type of hypothesis test, choice of model selection criteria, etc).

Summary of Recommendations Our key recommendations are (1) for kernel library design, it is beneficial to include flexible nonlinear kernels such as the **RBF** (Section [5.1\)](#page-18-0), (2) for hypothesis testing, use the parametric bootstrap test and the linear alternative kernel when the sample size is small (Section [5.2\)](#page-19-0), and (3) choose loocv as the model selection criterion and stack as the ensemble strategy (Section [5.3\)](#page-20-1).

5.1. Impact of model library for null model

Figure [1](#page-22-0) compares the performance of the hypothesis test constructed when the null model is fixed to match the true model (**Oracle**), or estimated using one of the three different types of model libraries (**Polynomial**, **RBF**, **Polynomial + RBF**). Here, the test is based on the bootstrap test with linear alternative kernel. Results for other test types are reported in the Appendix [A.](#page-30-0)

Generally speaking, when all kernels in the model library are parametric (i.e., **Polynomial** library), the resulting test is more powerful for polynomial data (e.g., linear, quadratic, cubic in Figure [1\)](#page-22-0), but loses power for data with nonlinearities (e.g., quadratic rbf, rbf_0.5, rbf_lnr_0.5, matern_1.5_0.5 in Figure [1\)](#page-22-0). On the other hand, model libraries that involves nonparametric kernels (e.g., **RBF** library and **Polynomial + RBF** library) lead to tests that yield slightly less power for polynomial data, but more more powerful for data with more complex nonlinearities (e.g., quadratic rbf, rbf 0.5 , rbf lnr 0.5, matern 1.5 0.5 in Figure [1\)](#page-22-0). Comparing the test power between a purely non-parametric model library (**RBF**) versus a semi-parametric model library (**Polynomial + RBF**), **RBF** performs slightly better than **Polynomial + RBF** library, especially when data comes from quadratic main effect with RBF interaction. This is likely due to the fact that adding less flexible polynomial kernels doesn't result in much of an increase in bias, but introduces more variance due to the need to estimate additional weights. Consequently, we recommend designing the model library to include the nonparametric kernels (e.g., **RBF** library or **Polynomial + RBF** library).

5.2. Impact of test type and choice for the alternative kernel

Figure [2](#page-23-0) compares the performance of three types of hypothesis test (lnr asymp refers to asymptotic test with linear alternative kernel, lnr_boot which is the bootstrap test with linear alternative kernel and ens_boot which is the bootstrap test with ensemble alternative kernel) under different data generation mechanisms. We fix the the model library for null model to **RBF**.

We first compare the asymptotic to the bootstrap test under the linear alternative kernel. Here, the asymptotic test is observed to be more powerful when data is generated from parametric models with polynomial nonlinearity (e.g., linear, quadratic, cubic in Figure [2\)](#page-23-0), but has difficulty in guaranteeing correct Type I error under more complex data (e.g., the cubic). Meanwhile, when data are generated from more flexible kernels (e.g., rbf_0.5, matern_1.5_0.5 in Figure [2\)](#page-23-0), the bootstrap test is observed to be slightly more powerful than the asymptotic test. So we recommend choosing the test type to be bootstrap.

Now, fixing the test type to be bootstrap test, we consider the impact of the choice of the alternative kernel on test performance. Specifically, we are interested in whether the test with an adaptively estimated alternative kernel (using the strategy outlined in Section [3.3\)](#page-10-0) leads to better power compared to the one with linear kernel as the alternative kernel. In general, the test based on the ensemble alternative kernel is more powerful than that generated by the linear alternative kernel, especially when data are generated having high complexity with quickly-varying local fluctuations (e.g., rbf_0.5, matern_1.5_0.5 in Figure [2\)](#page-23-0). Interestingly, when the data are simulated under nonlinearities generated from the RBF or the Matérn kernel, the test power sometimes decreases as the data generation mechanism moves away from the null (e.g., rbf_1 , matern 2.5 1 in Figure [2\)](#page-23-0). The simulation in Appendix [A](#page-30-0) also shows this phenomenon. This result suggests the test

statistic [\(6\)](#page-8-1) calculated with the ensemble alternative kernel can be unstable and yields an underestimate of the true alternative kernel under the alternative. Since the linear alternative kernel is more stable, we recommend choosing a linear alternative kernel in combination with the parametric bootstrap for testing.

5.3. Impact of model selection criterion and ensemble strategy

Figure [3](#page-24-0) compares the performance of the proposed hypothesis test when null models are selected from different tuning parameter criteria. This figure presents results when fixing the model library for null model to **RBF** and test type to bootstrap test with linear alternative kernel.

Among the seven tuning parameter selection methods (loocv, AIC, AICc, BIC, GCV, $GCVc$, gmpml), we notice that loocv is generally better at guaranteeing correct Type I error, except for Cubic data, which is hard to fit for all selection methods. Furthermore, it is important to note that some conventional hyper-parameter selection criteria (e.g., AIC in red and BIC in purple) produces suboptimal tests with inflated Type I error and weak power (e.g., cubic, quadratic_rbf, rbf_0.5 in Figure [3\)](#page-24-0). The issue with test power is especially severe when data comes from polynomial main effect, as the test power is weak and sometimes even decreases as the data generation mechanism moves away from the null (e.g., cubic, quadratic_rbf in Figure [3\)](#page-24-0). This result suggests poor fits of the null model as estimated using AIC/BIC. Consequently, we recommend using loocv and avoid using AIC/BIC-type criteria under these scenarios.

Figure [4](#page-25-0) compares the performance of hypothesis test when null models are selected from different ensemble strategies, fixing the model library for null model to **RBF** and test type to bootstrap test with linear alternative kernel. Interestingly, while Cubic data is hard to fit and stack is more powerful, avg and exp can guarantee correct Type I error. This is likely due to the fact that avg and exp have closed-form solutions for weights while the weights of stack need to be estimated, thus introducing more variance. On the other hand, among the three ensemble strategies, we notice that stack is generally more powerful than avg and exp. Therefore, we recommend choosing stack as the ensemble strategy.

6. Conclusion

In this paper we describe **CVEK**, a powerful and flexible toolkit in R for learning nonlinear, multivariate feature effects in limited samples. Given data, CVEK efficiently learns the complex, nonlinear data generation mechanism using an ensemble of kernel machine regressions. The package offers flexible choices for both kernel specification and hyper-parameter tuning for constructing the base models in a given library, as well as different ensemble strategies for constructing the final ensemble. CVEK also offers a suite of hypothesis tests for both the main effects and the interaction effects for the kernel features, and provides an asymptotic approximation and the parametric bootstrap as options for estimating the null distribution of the resulting test statistic. Further, one can use either the linear alternative kernel or an ensemble alternative kernel to test interaction effect. Through comprehensive simulation, we show that the hypothesis tests offered by CVEK are valid and powerful even under complex, non-smooth data generating mechanisms where the classical approaches fail. In practice, for a robust option for general purpose use, we recommend

users employ an RBF model library for the ensemble, select the hyper-parameter using loocv, and construct the ensemble using cross-validated stacking. For testing, we recommend one chooses the linear alternative kernel in combination with the parametric bootstrap. We encourage practitioners to conduct sensitivity analysis in order to examine the robustness of their conclusions, utilizing the wide range of options provided by CVEK.

Tables and Figures

Library \rightarrow Oracle \rightarrow Polynomial \rightarrow RBF \rightarrow Polynomial+RBF

Figure 1: Power of hypothesis test using the true model (Oracle) and the three *kmodel*'s, fixing the tuning parameter selection method to loocv, ensemble strategy to stack, and the test type to bootstrap test with linear alternative kernel.

type \rightarrow lnr_asymp \rightarrow lnr_boot \rightarrow ens_boot

Figure 2: Power of hypothesis test using different types of tests, fixing the tuning parameter selection method to loocv, ensemble strategy to stack, and model library for null model to

RBF.

Figure 3: Performance of hypothesis test when null models are selected from different tuning parameter selections, fixing the ensemble strategy to stack, test type to bootstrap test with linear alternative kernel, and model library for null model to **RBF**.

Figure 4: Performance of hypothesis test when null models are selected from different ensemble strategies, fixing the tuning parameter selection method to loocv, test type to bootstrap test with linear alternative kernel, and model library for null model to **RBF**.

Algorithm 1 Cross validated ensemble of kernels (CVEK)

1: **procedure** CVEK **Input:** A library of kernels $\{k_j\}_{j=1}^K$, Data (\mathbf{y}, \mathbf{x}) **Output:** Ensemble Kernel Matrix **K**^b $#$ Stage 1: Estimate λ and CV error for each kernel 2: **for** $j = 1$ to K do 3: $\mathbf{K}_j = \mathbf{K}_j/\text{tr}(\mathbf{K}_j)$ 4: $\widehat{\lambda}_j = \text{argmin }$ LOOCV $\left(\lambda \mid \mathbf{K}_j\right)$ 5: $\hat{\xi}_j = \text{CV}(\hat{\lambda}_j \mid \mathbf{K}_j)$ 6: **end for** # Stage 2: Estimate ensemble weights $\mathbf{u}_{K \times 1} = \{u_1, \ldots, u_K\}$ 7: $\hat{\mathbf{u}} = \underset{\mathbf{u} \in \Delta}{argmin} \parallel \sum_{j=1}^{K} u_j \hat{\xi}_j \parallel^2$ where $\Delta = {\mathbf{u} \mid \mathbf{u} \geq 0, \parallel \mathbf{u} \parallel_1=1}$ **u**∈∆ $\#$ Stage 3: Assemble the ensemble kernel matrix \mathbf{K}_{ens} 8: $\hat{\mathbf{A}} = \sum_{j=1}^K \hat{\mu}_j \mathbf{A}_{\widehat{\lambda}_j, k_j}$ 9: $U_A, \delta_A = \text{spectral_decomp}(\hat{A})$ 10: $\lambda_{\mathbf{K}} = min \Big(1, \left(\sum_{k=1}^{n} \frac{\delta_{A,k}}{1 - \delta_A} \right)$ $\left(\frac{\delta_{A,k}}{1-\delta_{A,k}}\right)^{-1}, min\big(\{\widehat{\lambda}_j\}_{j=1}^K\big)\right)$ 11: $\hat{\mathbf{K}} = \lambda_{\mathbf{K}} * \hat{\mathbf{U}}_A \ diag\left(\frac{\delta_{A,k}}{1-\delta_{A,k}}\right) \hat{\mathbf{U}}_A^{\top}$ 12: **end procedure**

Algorithm 2 Variance component test for $h \in \mathcal{H}_0$

1: **procedure** VCT FOR INTERACTION **Input:** Null Kernel Matrix **K**₀, Derivative Kernel Matrix ∂ **K**₀, Data (*y*, **x**) **Output:** Hypothesis Test p-value *p* # Stage 1: Estimate Null Model using REML 2: $(\hat{\boldsymbol{\mu}}, \hat{\tau}, \hat{\sigma}^2) = argmax L_{REML}(\boldsymbol{\mu}, \tau, \sigma^2 \mid \mathbf{K}_0)$ # Stage 2: Compute Test Statistic and Null Distribution Parameters $\hat{T}_0 = \hat{\tau} * (\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}})^{\top} \mathbf{V}_0^{-1} \partial \mathbf{K}_0 \mathbf{V}_0^{-1} (\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}})^{\top}$ $\hat{\kappa} = \hat{\mathbf{I}}_{\delta\delta}/[\hat{\tau} * tr(\mathbf{V}_0^{-1}\partial \mathbf{K}_0)], \quad \hat{\nu} = [\hat{\tau} * tr(\mathbf{V}_0^{-1}\partial \mathbf{K}_0)]^2/(2*\hat{\mathbf{I}}_{\delta\theta})$ # Stage 3: Compute p-value and reach conclusion 5: $p = P(\hat{\kappa}\chi^2_{\hat{\nu}} > \hat{T}) = P(\chi^2_{\hat{\nu}} > \hat{T}/\hat{\kappa})$ 6: **end procedure**

Algorithm 3 Parametric bootstrap test

1: **procedure** PARAMETRIC BOOTSTRAP TEST **Input:** Null Kernel Matrix \mathbf{K}_0 , Derivative Kernel Matrix $\partial \mathbf{K}_0$, Data (**y***,* **x**) **Output:** Hypothesis Test p-value *p* # Stage 1: Estimate Null Model using Gaussian Process Regression 2: $\hat{\boldsymbol{\mu}} = \mathbf{A}_0 \mathbf{y}, \quad \hat{\sigma}^2 = \frac{\mathbf{y}^\top (\mathbf{I} - \mathbf{A}_0) \mathbf{y}}{n - tr(\mathbf{A}_0)}$ *n*−*tr*(**A**0) *, τ*ˆ $#$ Stage 2: Sample response from the fitted model obtain in Step 1 $#$ and compute the test statistic based on fitting the alternative $#$ model, repeat for B times 3: **for** $b = 1$ to B **do** 4: $\mathbf{y}^* = \hat{\boldsymbol{\mu}} + \boldsymbol{\epsilon}, \quad \boldsymbol{\epsilon} \sim \mathrm{N}(0, \hat{\sigma}^2)$ 5: $\hat{T}_{0b} = \hat{\tau} * (\mathbf{y}^* - \hat{\boldsymbol{\mu}})^{\top} \mathbf{V}_0^{-1} \partial \mathbf{K}_0 \mathbf{V}_0^{-1} (\mathbf{y}^* - \hat{\boldsymbol{\mu}})$ 6: **end for** $#$ Stage 3: Compute the test statistic for the original data, based $#$ on fitting the alternative hypothesis model 7: $\hat{T}_0 = \hat{\tau} * (\mathbf{y} - \hat{\boldsymbol{\mu}})^{\top} \mathbf{V}_0^{-1} \partial \mathbf{K}_0 \mathbf{V}_0^{-1} (\mathbf{y} - \hat{\boldsymbol{\mu}})$ $\#$ Stage 4: Compute p-value and reach conclusion 8: $p = \frac{1}{E}$ $\frac{1}{B}\sum_{b=1}^{B} I(\hat{T}_{0b} > \hat{T}_{0})$ 9: **end procedure**

Computational details

The results in this paper were obtained using R 3.6.1 with the CVEK 0.1-2 package.

References

- Abramowitz M (1974). *Handbook of Mathematical Functions, With Formulas, Graphs, and Mathematical Tables,*. Dover Publications, Inc., New York, NY, USA. ISBN 978-0-486-61272-0.
- Bodenham DA, Adams NM (2016). "A comparison of efficient approximations for a weighted sum of chi-squared random variables." *Statistics and Computing*, **26**(4), 917–928. ISSN 1573-1375. [doi:10.1007/s11222-015-9583-4](http://dx.doi.org/10.1007/s11222-015-9583-4). URL <https://doi.org/10.1007/s11222-015-9583-4>.
- Boonstra PS, Mukherjee B, Taylor JMG (2015). "A Small-Sample Choice of the Tuning Parameter in Ridge Regression." *Statistica Sinica*, **25**(3), 1185–1206. ISSN 1017-0405. [doi:10.5705/ss.2013.284](http://dx.doi.org/10.5705/ss.2013.284).
- Buja A, Hastie T, Tibshirani R (1989). "Linear Smoothers and Additive Models." *The Annals of Statistics*, **17**(2), 453–510. ISSN 0090-5364. [doi:10.1214/aos/1176347115](http://dx.doi.org/10.1214/aos/1176347115). URL <http://projecteuclid.org/euclid.aos/1176347115>.
- Burnham KP, Anderson DR (2002). *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. 2 edition. Springer-Verlag, New York. ISBN 978-0-387-95364-9. [doi:10.1007/b97636](http://dx.doi.org/10.1007/b97636). URL <https://www.springer.com/gp/book/9780387953649>.
- Dalalyan AS, Tsybakov AB (2007). "Aggregation by Exponential Weighting and Sharp Oracle Inequalities." In *Learning Theory*, Lecture Notes in Computer Science, pp. 97–111. Springer, Berlin, Heidelberg. ISBN 978-3-540-72925-9 978-3-540-72927-3. [doi:10.1007/978-3-540-72927-3_9](http://dx.doi.org/10.1007/978-3-540-72927-3_9). URL https://link.springer.com/chapter/10.1007/978-3-540-72927-3_9.
- Golub GH, Heath M, Wahba G (1979). "Generalized Cross-Validation as a Method for Choosing a Good Ridge Parameter." *Technometrics*, **21**(2), 215–223. ISSN 0040-1706. [doi:10.1080/00401706.1979.10489751](http://dx.doi.org/10.1080/00401706.1979.10489751). URL <https://www.tandfonline.com/doi/abs/10.1080/00401706.1979.10489751>.
- Gu C (2013). *Smoothing Spline ANOVA Models*. Springer Series in Statistics, 2 edition. Springer-Verlag, New York. ISBN 978-1-4614-5368-0. URL <//www.springer.com/us/book/9781461453680>.
- Harrison D, Rubinfeld DL (1978). "Hedonic housing prices and the demand for clean air." *Journal of Environmental Economics and Management*, **5**(1), 81–102. ISSN 0095-0696. [doi:10.1016/0095-0696\(78\)90006-2](http://dx.doi.org/10.1016/0095-0696(78)90006-2). URL <http://www.sciencedirect.com/science/article/pii/0095069678900062>.
- Hurvich CM, Tsai CL (1989). "Regression and time series model selection in small samples." *Biometrika*, **76**(2), 297–307. ISSN 0006-3444. [doi:10.1093/biomet/76.2.297](http://dx.doi.org/10.1093/biomet/76.2.297). URL <https://academic.oup.com/biomet/article/76/2/297/265326>.
- Hurvich Clifford M, Simonoff Jeffrey S, Tsai Chih Ling (2002). "Smoothing parameter selection in nonparametric regression using an improved Akaike information criterion." *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **60**(2), 271–293. ISSN 1369-7412. [doi:10.1111/1467-9868.00125](http://dx.doi.org/10.1111/1467-9868.00125). URL <https://rss.onlinelibrary.wiley.com/doi/abs/10.1111/1467-9868.00125>.
- Lin X (1997). "Variance component testing in generalised linear models with random effects." *Biometrika*, **84**(2), 309–326. ISSN 0006-3444. [doi:10.1093/biomet/84.2.309](http://dx.doi.org/10.1093/biomet/84.2.309). URL <https://academic.oup.com/biomet/article/84/2/309/233889>.
- Lin X, Zhang D (1999). "Inference in generalized additive mixed modelsby using smoothing splines." *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, **61**(2), 381–400. ISSN 1467-9868. [doi:10.1111/1467-9868.00183](http://dx.doi.org/10.1111/1467-9868.00183). URL <https://rss.onlinelibrary.wiley.com/doi/abs/10.1111/1467-9868.00183>.
- Liu D, Lin X, Ghosh D (2007). "Semiparametric Regression of Multidimensional Genetic Pathway Data: Least-Squares Kernel Machines and Linear Mixed Models." *Biometrics*, **63**(4), 1079–1088. ISSN 0006-341X. [doi:10.1111/j.1541-0420.2007.00799.x](http://dx.doi.org/10.1111/j.1541-0420.2007.00799.x). URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2665800/>.
- Liu J, Coull B (2017). "Robust Hypothesis Test for Nonlinear Effect with Gaussian Processes." In I Guyon, UV Luxburg, S Bengio, H Wallach, R Fergus, S Vishwanathan, R Garnett (eds.), *Advances in Neural Information Processing Systems 30*, pp. 795–803. Curran Associates, Inc. URL [http://papers.nips.cc/paper/](http://papers.nips.cc/paper/6681-robust-hypothesis-test-for-nonlinear-effect-with-gaussian-processes.pdf) [6681-robust-hypothesis-test-for-nonlinear-effect-with-gaussian-processes.](http://papers.nips.cc/paper/6681-robust-hypothesis-test-for-nonlinear-effect-with-gaussian-processes.pdf) [pdf](http://papers.nips.cc/paper/6681-robust-hypothesis-test-for-nonlinear-effect-with-gaussian-processes.pdf).
- Liu JZ, Lee J, Lin PiD, Valeri L, Christiani DC, Bellinger DC, Wright RO, Mazumdar MM, Coull BA (2019). "A Cross-validated Ensemble Approach to Robust Hypothesis Testing of Continuous Nonlinear Interactions: Application to Nutrition-Environment Studies." *arXiv:1904.10918 [stat]*. ArXiv: 1904.10918, URL <http://arxiv.org/abs/1904.10918>.
- Maity A, Lin X (2011). "Powerful tests for detecting a gene effect in the presence of possible gene-gene interactions using garrote kernel machines." *Biometrics*, **67**(4), 1271–1284. ISSN 1541-0420. [doi:10.1111/j.1541-0420.2011.01598.x](http://dx.doi.org/10.1111/j.1541-0420.2011.01598.x).
- Micchelli CA, Xu Y, Zhang H (2006). "Universal Kernels." *J. Mach. Learn. Res.*, **7**, 2651–2667. ISSN 1532-4435. URL <http://dl.acm.org/citation.cfm?id=1248547.1248642>.
- Yang Y, Dunson DB (2014). "Minimax Optimal Bayesian Aggregation." *arXiv:1403.1345 [math, stat]*. URL <http://arxiv.org/abs/1403.1345>.
- Zhan X, Plantinga A, Zhao N, Wu MC (2017). "A fast small-sample kernel independence test for microbiome community-level association analysis." *Biometrics*, **73**(4), 1453–1463. ISSN 0006-341X. [doi:10.1111/biom.12684](http://dx.doi.org/10.1111/biom.12684). URL <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5592124/>.

A. Further simulation results under the settings in Section 5

Figure A.1: Power of hypothesis test using different libraries combined with different types of tests, fixing the tuning parameter selection method to loocv and ensemble strategy to stack.

A.2. Performances of different libraries combined with different ensemble strategies

Figure A.2: Power of hypothesis test using different libraries combined with different ensemble strategies, fixing the test type to bootstrap test with linear alternative kernel and tuning parameter selection method to loocv.

A.3. Performances of different testing types combined with different ensemble strategies

Figure A.3: Power of hypothesis test using different testing types combined with different ensemble strategies, fixing the model library for null model to **RBF** and tuning parameter selection method to loocv.

Figure A.4: Power of hypothesis test using different tuning parameter selections combined with different types of tests, fixing the model library for null model to **RBF** and ensemble strategy to stack.

A.5. Performances of different beta's of exponential weighting

Below shows the performances under three different beta's of exponential weighting: $\min\{RSS\}_{d=1}^D/10$, $\text{median}\{RSS\}_{d=1}^D$ and $\max\{RSS\}_{d=1}^D * 2$. Here $\{RSS\}_{d=1}^D$ are the set of residual sum of squares of *D* base kernels. We can see that their performances are quite similar.

Figure A.5: Power of hypothesis test using different beta's of exponential weighting, fixing the model library for null model to **RBF**, test type to bootstrap test with linear alternative kernel, tuning parameter selection method to loocv and ensemble strategy to stack.

B. Correlated settings

B.1. Performances of different libraries under correlated setting (without between group correlation)

Figure B.1: Power of hypothesis test using the true model (Oracle) and the three *kmodel*'s when within group correlation coefficients are 0.3 and 0.7 respectively, and between group correlation coefficient is 0, fixing the tuning parameter selection method to loocv, ensemble strategy to stack, and the test type to bootstrap test with linear alternative kernel.

B.2. Performances of different testing types under correlated setting (without between group correlation)

Figure B.2: Power of hypothesis test using different types of tests when within group correlation coefficients are 0.3 and 0.7 respectively, and between group correlation coefficient is 0, fixing the tuning parameter selection method to loocv, ensemble strategy to stack and model library for null model to **RBF**.

B.3. Performances of different libraries under correlated setting (with between group correlation)

●● ● ●● ●● ●● ● ● ● ● ● ● ●●●● ●● ●● ●● ●● ●● ●● ● ●● ●● ●● ● ● ● ●●●●● ●● ●●● ●●● ●● $\overline{\mathbb{Z}}$ ●● ●● ● ● ● ● ● ● ● ●● ●● ● ●● ● ●● ● ●● ● ● ● ●● ● $\Bigg)$ ● ● ●● ●● ●● ●● ●● ●● ●● ● ●● ● ● ● ●● ● ● ● matern_2.5_1 ●● ●● ● ● ●● ● ● ● ● ● ● ● ● ● rbf_lnr_0.5 matern_1.5_0.5 matern_2.5_1 quadratic_rbf rbf_0.5 rbf_1 linear and the cubic 0.00 0.25 0.50 0.75 1.00 0.00 0.25 0.50 0.75 1.00 0.00 0.25 0.50 0.75 1.00 0.00 0.25 0.50 0.75 1.00 0.00 0.25 ፱
≷ 0.50
Ⴍ 0.75 1.00 $0.00 \frac{1}{0.00}$ 0.25 0.50 0.75 1.00 interaction strength, δ

Library \rightarrow Oracle \rightarrow Polynomial \rightarrow RBF \rightarrow Polynomial+RBF

Figure B.3: Power of hypothesis test using the true model (Oracle) and the three *kmodel*'s when within group correlation coefficients are 0.3 and 0.7 respectively, and between group correlation coefficient is 0.2, fixing the tuning parameter selection method to loocv, ensemble strategy to stack and the test type to bootstrap test with linear alternative kernel.

B.4. Performances of different testing types under correlated setting (with between group correlation)

Figure B.4: Power of hypothesis test using different types of tests when within group correlation coefficients are 0.3 and 0.7 respectively, and between group correlation coefficient is 0.2, fixing the tuning parameter selection method to loocv, ensemble strategy to stack and model library for null model to **RBF**.

C. Backfitting algorithm for multiple kernels

```
Algorithm 4 Backfitting algorithm for multiple kernels
```

```
1: procedure Fitting Multiple Kernels
      Input: A set of kernel matrices \{K_d\}_{d=1}^D, Data (\mathbf{y}, \mathbf{X}), \lambdaOutput: Estimators \boldsymbol{\beta}, \{\boldsymbol{\alpha}_d\}_{d=1}^D\# Stage 1: Initialize parameters \beta and \{\boldsymbol{\alpha}_d\}_{d=1}^D2: \qquad \boldsymbol{\beta} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}3: for d = 1 to D do
 4: \boldsymbol{\alpha}_d = (K_d + \lambda \mathbf{I})^{-1} (\mathbf{y} - \mathbf{X}\boldsymbol{\beta})5: end for
     # Stage 2: Iterative update
 6: for step = 1 to max\_step do
 7: \beta = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top (\mathbf{y} - \sum_{d=1}^D K_d \alpha_d)8: for d = 1 to D do
 9: \boldsymbol{\alpha}_d = (K_d + \lambda \mathbf{I})^{-1} (\mathbf{y} - \mathbf{X}\boldsymbol{\beta} - \sum_{d' \neq d} K_{d'} \boldsymbol{\alpha}_{d'})10: end for
11: end for
12: until convergence
13: end procedure
```
D. Projection matrices for multiple kernels

Basically, for the problem [\(Buja, Hastie, and Tibshirani](#page-28-8) [1989\)](#page-28-8),

$$
\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \sum_{d=1}^{D} K_d \boldsymbol{\alpha}_d,
$$

define below projection matrices to the linear and the *d th* kernel as,

$$
H = \mathbf{X}(\mathbf{X}^{\top}\mathbf{X})^{-1}\mathbf{X}^{\top},
$$

\n
$$
S_d = K_d(K_d + \lambda \mathbf{I})^{-1}.
$$

Then we can compute the "projection matrices" to the overall kernel space *B* as,

$$
A_d = (\mathbf{I} - S_d)^{-1} S_d,
$$

\n
$$
A = \sum_{d=1}^{D} A_d,
$$

\n
$$
B = (\mathbf{I} + A)^{-1} A.
$$

Consequently, the final projections to the kernel effect space and the fixed effect space are,

$$
P_K = (\mathbf{I} - BH)^{-1}B(\mathbf{I} - H),
$$

\n
$$
P_X = H(\mathbf{I} - P_K),
$$

such that $\mathbf{X}\hat{\boldsymbol{\beta}} = P_X\mathbf{y}$ and $\sum_{d=1}^{D} K_d\hat{\boldsymbol{\alpha}}_d = P_K\mathbf{y}$.

E. Derivation of the REML based test statistic

E.1. Derivation of the score test statistic

In this section, we derive the score test statistic based on REML [\(Maity and Lin](#page-29-9) [2011\)](#page-29-9). Denote $\mathbf{V}(\boldsymbol{\theta}) = \sigma^2 \mathbf{V}_{\lambda} = \sigma^2 \mathbf{I} + \tau \mathbf{K}_{\delta}$, where $\boldsymbol{\theta} = (\delta, \tau, \sigma^2)$. The REML

$$
l_R(\mu, \lambda, \sigma^2 | \mathbf{y}) = -\frac{1}{2} \Big[\log |\sigma^2 \mathbf{V}_{\lambda}| + (\mathbf{y} - \boldsymbol{\mu})^T (\sigma^2 \mathbf{V}_{\lambda})^{-1} (\mathbf{y} - \boldsymbol{\mu}) + \log |\sigma^{-2} \mathbf{1}^T \mathbf{V}_{\lambda}^{-1} \mathbf{1}| \Big], \quad (9)
$$

can be rewritten as

$$
l_R = -\frac{1}{2} \Big[\log |\mathbf{V}(\boldsymbol{\theta})| + \log |\mathbf{1}^T \mathbf{V}(\boldsymbol{\theta})^{-1} \mathbf{1}| + (\mathbf{y} - \boldsymbol{\mu})^T \mathbf{V}(\boldsymbol{\theta})^{-1} (\mathbf{y} - \boldsymbol{\mu}) \Big]. \tag{10}
$$

Under $H_0: \delta = 0$ (2.2.2), we set $\theta_0 = (0, \tau, \sigma^2)$ and

$$
\mathbf{P}_0(\boldsymbol{\theta}_0) = \mathbf{V}(\boldsymbol{\theta}_0)^{-1} - \mathbf{V}(\boldsymbol{\theta}_0)^{-1} \mathbf{1} [\mathbf{1}^T \mathbf{V}(\boldsymbol{\theta}_0)^{-1} \mathbf{1}]^{-1} \mathbf{1}^T \mathbf{V}(\boldsymbol{\theta}_0)^{-1}.
$$

Take the derivative of [\(10\)](#page-40-1) with respect to δ ,

$$
\frac{\partial l_R}{\partial \delta} = -\frac{1}{2} \Big[\frac{\partial log |\mathbf{V}(\boldsymbol{\theta})|}{\partial \delta} + \frac{\partial log |\mathbf{1}^T \mathbf{V}(\boldsymbol{\theta})^{-1} \mathbf{1}|}{\partial \delta} + \frac{\partial (\mathbf{y} - \boldsymbol{\mu})^T \mathbf{V}(\boldsymbol{\theta})^{-1} (\mathbf{y} - \boldsymbol{\mu})}{\partial \delta} \Big]
$$
\n
$$
= -\frac{1}{2} \Big[tr(\mathbf{V}(\boldsymbol{\theta})^{-1} \frac{\partial \mathbf{V}(\boldsymbol{\theta})}{\partial \delta}) + tr(|\mathbf{1}^T \mathbf{V}(\boldsymbol{\theta})^{-1} \mathbf{1}|^{-1} \mathbf{1}^T \frac{\partial \mathbf{V}(\boldsymbol{\theta})^{-1}}{\partial \delta} \mathbf{1} \Big]
$$
\n
$$
+ (\mathbf{y} - \boldsymbol{\mu})^T \frac{\partial \mathbf{V}(\boldsymbol{\theta})^{-1}}{\partial \delta} (\mathbf{y} - \boldsymbol{\mu}) \Big]
$$
\n
$$
= -\frac{1}{2} \Big[tr(\mathbf{V}(\boldsymbol{\theta})^{-1} \tau(\partial \mathbf{K}_{\delta})) - tr(\tau(\partial \mathbf{K}_{\delta}) \mathbf{V}(\boldsymbol{\theta})^{-1} \mathbf{1} [\mathbf{1}^T \mathbf{V}(\boldsymbol{\theta})^{-1} \mathbf{1}]^{-1} \mathbf{1}^T \mathbf{V}(\boldsymbol{\theta})^{-1} \Big]
$$
\n
$$
- (\mathbf{y} - \boldsymbol{\mu})^T \mathbf{V}(\boldsymbol{\theta})^{-1} \tau(\partial \mathbf{K}_{\delta}) \mathbf{V}(\boldsymbol{\theta})^{-1} (\mathbf{y} - \boldsymbol{\mu}) \Big]
$$
\n
$$
= \frac{1}{2} (\mathbf{y} - \boldsymbol{\mu})^T \mathbf{V}(\boldsymbol{\theta})^{-1} \tau(\partial \mathbf{K}_{\delta}) \mathbf{V}(\boldsymbol{\theta})^{-1} (\mathbf{y} - \boldsymbol{\mu})
$$
\n
$$
- \frac{1}{2} tr \Big[\tau(\partial \mathbf{K}_{\delta}) [\mathbf{V}(\boldsymbol{\theta})^{-1} - \mathbf{V}(\boldsymbol{\theta})^{-1} \mathbf{1} [\mathbf{1}^T \mathbf{V}(\boldsymbol{\theta})^{-1} \mathbf
$$

where $\partial \mathbf{K}_{\delta}$ is the derivative kernel matrix whose (i, j) th entry is $\frac{\partial k_{\delta}(\mathbf{x}, \mathbf{x}')}{\partial \delta}$. If we further denote $\mathbf{K}_0 = \mathbf{K}_\delta |_{\delta=0}$ and $\partial \mathbf{K}_0 = (\partial \mathbf{K}_\delta) |_{\delta=0}$, we get the REML based score function of δ evaluated at H_0

$$
S_{\delta=0} = \frac{1}{2}(\mathbf{y} - \boldsymbol{\mu})^T \mathbf{V}(\boldsymbol{\theta}_0)^{-1} \tau (\partial \mathbf{K}_0) \mathbf{V}(\boldsymbol{\theta}_0)^{-1}(\mathbf{y} - \boldsymbol{\mu}) - \frac{1}{2} tr[\tau (\partial \mathbf{K}_0) \mathbf{P}_0].
$$

To test for $H_0: \delta = 0$, we propose to use the score-based test statistic

$$
\hat{T}_0 = \hat{\tau}(\mathbf{y} - \hat{\boldsymbol{\mu}})^T \mathbf{V}_0^{-1} (\partial \mathbf{K}_0) \mathbf{V}_0^{-1} (\mathbf{y} - \hat{\boldsymbol{\mu}}), \tag{12}
$$

where $\mathbf{V}_0 = \hat{\sigma}^2 \mathbf{I} + \hat{\tau} \mathbf{K}_0$.

E.2. The null distribution of the test statistic

For simplicity, we denote

$$
\mathbf{V} = \mathbf{V}(\boldsymbol{\theta}),
$$

$$
\mathbf{P} = \mathbf{P}(\boldsymbol{\theta}) = \mathbf{V}^{-1} - \mathbf{V}^{-1} \mathbf{1} [\mathbf{1}^T \mathbf{V}^{-1} \mathbf{1}]^{-1} \mathbf{1}^T \mathbf{V}^{-1}.
$$

With similar derivation as [\(11\)](#page-40-2), for each $\theta_i \in \boldsymbol{\theta} = (\delta, \tau, \sigma^2)$, we have

$$
\frac{\partial l_R}{\partial \theta_i} = -\frac{1}{2} \Big[tr\left(\mathbf{P} \frac{\partial \mathbf{V}}{\partial \theta_i} \right) - (\mathbf{y} - \boldsymbol{\mu})^T \mathbf{V}^{-1} \left(\frac{\partial \mathbf{V}}{\partial \theta_i} \right) \mathbf{V}^{-1} (\mathbf{y} - \boldsymbol{\mu}) \Big]. \tag{13}
$$

From (Liu *[et al.](#page-29-1)* [2007\)](#page-29-1) we know $\hat{\boldsymbol{\mu}} = [\mathbf{1}^T \mathbf{V}^{-1} \mathbf{1}]^{-1} \mathbf{1}^T \mathbf{V}^{-1} \mathbf{y}$. Plug it in [\(Lin and Zhang](#page-29-10) [1999\)](#page-29-10), and we obtain

$$
(\mathbf{y} - \boldsymbol{\mu})^T \mathbf{V}^{-1} = \mathbf{y}^T (\mathbf{I} - \mathbf{1} [\mathbf{1}^T \mathbf{V}^{-1} \mathbf{1}]^{-1} \mathbf{1}^T \mathbf{V}^{-1})^T \mathbf{V}^{-1} = \mathbf{y}^T \mathbf{P}.
$$

Then [\(13\)](#page-41-0) becomes

$$
\frac{\partial l_R}{\partial \theta_i} = -\frac{1}{2} \Big[tr(\mathbf{P} \frac{\partial \mathbf{V}}{\partial \theta_i}) - \mathbf{y}^T \mathbf{P} \big(\frac{\partial \mathbf{V}}{\partial \theta_i} \big) \mathbf{P} \mathbf{y} \Big].
$$

The second-order partial derivatives with respect to θ_i and θ_j are

$$
\frac{\partial^2 l_R}{\partial \theta_i \partial \theta_j} = -\frac{1}{2} \Big[tr \Big(\frac{\partial \mathbf{P}}{\partial \theta_j} \frac{\partial \mathbf{V}}{\partial \theta_i} \Big) + tr \Big(\mathbf{P} \frac{\partial^2 \mathbf{V}}{\partial \theta_i \partial \theta_j} \Big) + \mathbf{y}^T \mathbf{P} \Big(\frac{\partial \mathbf{V}}{\partial \theta_i} \Big) \mathbf{P} \Big(\frac{\partial \mathbf{V}}{\partial \theta_j} \Big) \mathbf{P} \mathbf{y} + \mathbf{y}^T \mathbf{P} \Big(\frac{\partial \mathbf{V}}{\partial \theta_j} \Big) \mathbf{P} \Big(\frac{\partial \mathbf{V}}{\partial \theta_i} \Big) \mathbf{P} \mathbf{y} - \mathbf{y}^T \mathbf{P} \frac{\partial^2 \mathbf{V}}{\partial \theta_i \partial \theta_j} \mathbf{P} \mathbf{y} \Big],
$$
\n(14)

where we have used the fact that

$$
\frac{\partial \mathbf{P}}{\partial \theta_j} = -\mathbf{V}^{-1} \frac{\partial \mathbf{V}}{\partial \theta_j} \mathbf{V}^{-1} + \mathbf{V}^{-1} \frac{\partial \mathbf{V}}{\partial \theta_j} \mathbf{V}^{-1} \mathbf{1} [\mathbf{1}^T \mathbf{V}^{-1} \mathbf{1}]^{-1} \mathbf{1}^T \mathbf{V}^{-1} \n+ \mathbf{V}^{-1} \mathbf{1} [\mathbf{1}^T \mathbf{V}^{-1} \mathbf{1}]^{-1} \mathbf{1}^T \mathbf{V}^{-1} \frac{\partial \mathbf{V}}{\partial \theta_j} \mathbf{V}^{-1} \n- \mathbf{V}^{-1} \mathbf{1} ([\mathbf{1}^T \mathbf{V}^{-1} \mathbf{1}]^{-1} \mathbf{1}^T \mathbf{V}^{-1} \frac{\partial \mathbf{V}}{\partial \theta_j} \mathbf{V}^{-1} \mathbf{1} [\mathbf{1}^T \mathbf{V}^{-1} \mathbf{1}]^{-1}) \mathbf{1}^T \mathbf{V}^{-1} \n= -\mathbf{P} \frac{\partial \mathbf{V}}{\partial \theta_j} \mathbf{P}.
$$

Then [\(14\)](#page-41-1) becomes

$$
\frac{\partial^2 l_R}{\partial \theta_i \partial \theta_j} = -\frac{1}{2} \Big[-tr(\mathbf{P} \frac{\partial \mathbf{V}}{\partial \theta_j} \mathbf{P} \frac{\partial \mathbf{V}}{\partial \theta_i}) + tr(\mathbf{P} \frac{\partial^2 \mathbf{V}}{\partial \theta_i \partial \theta_j}) + \mathbf{y}^T \mathbf{P} (\frac{\partial \mathbf{V}}{\partial \theta_i}) \mathbf{P} (\frac{\partial \mathbf{V}}{\partial \theta_j}) \mathbf{P} \mathbf{y} + \mathbf{y}^T \mathbf{P} (\frac{\partial \mathbf{V}}{\partial \theta_j}) \mathbf{P} (\frac{\partial \mathbf{V}}{\partial \theta_i}) \mathbf{P} \mathbf{y} - \mathbf{y}^T \mathbf{P} \frac{\partial^2 \mathbf{V}}{\partial \theta_i \partial \theta_j} \mathbf{P} \mathbf{y} \Big].
$$
\n(15)

Since

$$
E(\mathbf{P} \mathbf{y} \mathbf{y}^T) = \mathbf{P}[Var(\mathbf{y}) + (E\mathbf{y})(E\mathbf{y})^T] = \mathbf{P}[\mathbf{V} + \boldsymbol{\mu}\boldsymbol{\mu}^T] = \mathbf{P}\mathbf{V},
$$

$$
\mathbf{P}\mathbf{V}\mathbf{P} = \mathbf{P}[\mathbf{I} - \mathbf{1}[\mathbf{1}^T\mathbf{V}^{-1}\mathbf{1}]^{-1}\mathbf{1}^T\mathbf{V}^{-1}] = \mathbf{P},
$$

we get

$$
E\Big[\mathbf{y}^T \mathbf{P} \big(\frac{\partial \mathbf{V}}{\partial \theta_j} \big) \mathbf{P} \big(\frac{\partial \mathbf{V}}{\partial \theta_i} \big) \mathbf{P} \mathbf{y} \Big] = tr\Big(E\Big[\mathbf{P} \big(\frac{\partial \mathbf{V}}{\partial \theta_j} \big) \mathbf{P} \big(\frac{\partial \mathbf{V}}{\partial \theta_i} \big) \mathbf{P} \mathbf{y} \mathbf{y}^T\Big]\Big) = tr\Big(\mathbf{P} \big(\frac{\partial \mathbf{V}}{\partial \theta_j} \big) \mathbf{P} \big(\frac{\partial \mathbf{V}}{\partial \theta_i} \big) \mathbf{P} \mathbf{V} \Big) = tr\Big(\mathbf{P} \big(\frac{\partial \mathbf{V}}{\partial \theta_j} \big) \mathbf{P} \big(\frac{\partial \mathbf{V}}{\partial \theta_i} \big) \Big),
$$

$$
E\Big[\mathbf{y}^T \mathbf{P} \frac{\partial^2 \mathbf{V}}{\partial \theta_i \partial \theta_j} \mathbf{P} \mathbf{y} \Big] = tr\Big(\mathbf{P} \frac{\partial^2 \mathbf{V}}{\partial \theta_i \partial \theta_j} \Big).
$$

Therefore,

$$
\mathbf{I}_{\theta_i,\theta_j} = -E\Big[\frac{\partial^2 l_R}{\partial \theta_i \partial \theta_j}\Big] = \frac{1}{2}tr\Big(\mathbf{P}\big(\frac{\partial \mathbf{V}}{\partial \theta_j})\mathbf{P}\big(\frac{\partial \mathbf{V}}{\partial \theta_i}\big)\Big).
$$

F. Visualizing nonlinear interaction in Boston Housing Price

In this section we show an example of how to visualize the fitted interaction from a cvek model. Here we consider the Boston example in Section [4.2.](#page-14-0) We visualize the interaction effects by creating five datasets: Fix all confounding variables to their means, vary lstat in a reasonable range (i.e., from 12*.*5 to 17*.*5, since the original range of lstat in Boston dataset is (1*.*73*,* 37*.*97)), and respectively set crim value to its 5%*,* 25%*,* 50%*,* 75% and 95% quantiles.

```
> # first fit the alternative model
> formula_alt <- medv ~ zn + indus + chas + nox + rm + age + dis +
   rad + tax + ptratio + black + k(crin):k(lstat)> fit bos_alt <- cvek(formula = formula_alt, kern_func_list = kern_func_list,
+ data = Boston, lambda = exp(seq(-3, 5)))
>
> # mean-center all confounding variables not involved in the interaction
> # so that the predicted values are more easily interpreted
> pred_name <- c("zn", "indus", "chas", "nox", "rm", "age",
+ "dis", "rad", "tax", "ptratio", "black")
> covar_mean <- apply(Boston, 2, mean)
> pred_cov <- covar_mean[pred_name]
> pred_cov_df <- t(as.data.frame(pred_cov))
> lstat_list \leq seq(12.5, 17.5, length.out = 100)
> crim_quantiles \leq quantile(Boston$crim, probs = c(.05, .25, .5, .75, .95))
>
> # crim is set to its 5% quantile
> data_test1 <- data.frame(pred_cov_df, lstat = lstat_list,
                              crim = crim quantiles [1])
row names were found from a short variable and have been discarded
> data_test1_pred <- predict(fit_bos_alt, data_test1)
```

```
> # crim is set to its 25% quantile
> data_test2 <- data.frame(pred_cov_df, lstat = lstat_list,
                           crim = crim quantiles [2])
row names were found from a short variable and have been discarded
> data_test2_pred <- predict(fit_bos_alt, data_test2)
>
> # crim is set to its 50% quantile
> data test3 <- data.frame(pred cov df, lstat = lstat list,
+ crim = crim_quantiles[3])
row names were found from a short variable and have been discarded
> data_test3_pred <- predict(fit_bos_alt, data_test3)
>
> # crim is set to its 75% quantile
> data_test4 <- data.frame(pred_cov_df, lstat = lstat_list,
                           crim = crim quantiles [4])
row names were found from a short variable and have been discarded
> data_test4_pred <- predict(fit_bos_alt, data_test4)
>
> # crim is set to its 95% quantile
> data_test5 <- data.frame(pred_cov_df, lstat = lstat_list,
                           crim = crim quantiles [5])
row names were found from a short variable and have been discarded
> data_test5_pred <- predict(fit_bos_alt, data_test5)
>
> # combine five sets of prediction data together
> medv <- rbind(data_test1_pred, data_test2_pred, data_test3_pred,
              data_test4_pred, data_test5_pred)
> data_pred \le - data.frame(lstat = rep(lstat_list, 5), medv = medv,
+ crim = rep(c("5% quantile", "25% quantile",
+ "50% quantile", "75% quantile",
                                  "95% quantile"), each = 100))
> data_pred$crim <- factor(data_pred$crim,
+ levels = c("5% quantile", "25% quantile",
+ "50% quantile", "75% quantile",
+ "95% quantile"))
>
> data_label <- data_pred[which(data_pred$lstat == 17.5), ]
> data_label$value <- c("0.028%", "0.082%", "0.257%", "3.677%", "15.789%")
> data_label$value <- factor(data_label$value, levels =
+ c("0.028%", "0.082%", "0.257%",
+ "3.677%", "15.789%"))
>
> ggplot(data = data_pred, aes(x = lstat, y = medv, color = crim)) +
     geom\_point(size = 0.1) ++ geom_text_repel(aes(label = value), data = data_label,
+ color = "black", size = 3.6) +
```
>

```
+ scale_colour_manual(values = c("firebrick1", "chocolate2",
+ "darkolivegreen3", "skyblue2",
+ "purple2")) +
+ geom_line() + theme_set(theme_bw()) +
+ theme(panel.grid = element blank(),
+ axis.title.x = element_text(size = 12),
+ axis.title.y = element text(size = 12),
+ legend.title = element_text(size = 12, face = "bold"),
+ legend.text = element text(size = 12)) +
+ labs(x = "percentage of lower status",
+ y = "median value of owner-occupied homes ($1000)",
+ col = "per capita crime rate")
```
Figure [F.1](#page-22-0) shows the medv - lstat relationship under different levels of crim. Numbers at the end of each curve indicate the actual values of crim rate (per capita crime rate by town) at the corresponding quantiles. From the figure we see that crime rate does impact the relationship between the local socioeconomic status v.s. housing price. Building on this code, user can continue to refine the visualization (e.g., by adding in confidence levels) and use it to improve the the model fit based on domain knowledge (e.g., by experimenting with different kernels / hyper-parameters).

Figure F.1: medv - lstat relationship under different levels of crim

Affiliation:

Wenying Deng, Jeremiah Zhe Liu, Erin Lake, Brent A. Coull

Department of Biostatistics Harvard University Cambridge, MA 02138 USA E-mail: [{wdeng@g, zhl112@mail, eklake@hsph, bcoull@hsph}.harvard.edu](mailto:\{wdeng@g, zhl112@mail, eklake@hsph, bcoull@hsph\}.harvard.edu)

