# SHARP BOUNDS FOR FINITELY MANY EMBEDDED EIGENVALUES OF PERTURBED STARK TYPE OPERATORS

WENCAI LIU

ABSTRACT. For perturbed Stark operators  $Hu = -u'' - xu + qu$ , the author has proved that lim sup<sub>x→∞</sub>  $x^{\frac{1}{2}}|q(x)|$  must be larger than  $\frac{1}{\sqrt{2}}N^{\frac{1}{2}}$  in order to create N linearly independent eigensolutions in  $L^2(\mathbb{R}^+)$  [\[25](#page-11-0)]. In this paper, we apply *generalized Wigner-von Neumann*  $type$  functions to construct embedded eigenvalues for a class of Schrödinger operators, including a proof that the bound  $\frac{1}{\sqrt{2}}N^{\frac{1}{2}}$  is sharp.

### 1. INTRODUCTION

The Stark operator  $Hu = -u'' - xu + qu$  describes a charged quantum particle in a constant electric field with an additional electric potential  $q$ . It has attracted a lot of attentions from both mathematics and physics [\[1](#page-11-1)[–3,](#page-11-2) [5,](#page-11-3) [7,](#page-11-4) [8,](#page-11-5) [10,](#page-11-6) [12,](#page-11-7) [18,](#page-11-8) [19,](#page-11-9) [21](#page-11-10), [22](#page-11-11), [37](#page-12-0), [39](#page-12-1)].

In this paper, we consider a class of more general operators, Stark type operators on  $L^2(\mathbb{R}^+)$ :

(1) Hu = −u ′′ − x <sup>α</sup>u + qu,

where  $0 < \alpha < 2$ . Denote by  $H_0 u = -u'' - x^{\alpha} u$  and regard q as a perturbation.

It is well known that for any  $0 < \alpha < 2$ ,  $\sigma_{\text{ess}}(H_0) = \sigma_{\text{ac}}(H_0) = \mathbb{R}$  and  $H_0$  does not have any eigenvalue. The criteria for the perturbation such that the associated perturbed Stark type operator has single eigenvalue, finitely many eigenvalues or countably many eigenvalues have been obtained in [\[25\]](#page-11-0).

Define  $P \subset \mathbb{R}$  as

 $P = \{ E \in \mathbb{R} : -u'' - x^{\alpha}u + qu = Eu \text{ has an } L^2(\mathbb{R}^+) \text{ solution} \}.$ 

In [\[25](#page-11-0)], the author proved that

<span id="page-0-0"></span>**Theorem 1.1.** [\[25](#page-11-0), Theorem 1.5] Let a be given by

(2) 
$$
a = \limsup_{x \to \infty} x^{1 - \frac{\alpha}{2}} |q(x)|.
$$

Then we have

(3) 
$$
a \ge \frac{2-\alpha}{\sqrt{2}} (\#P)^{\frac{1}{2}}.
$$

<span id="page-0-2"></span>**Theorem 1.2.** [\[25](#page-11-0), Theorem 1.6] For any  $\{E_j\}_{j=1}^N \subset \mathbb{R}$  and any  $\{\theta_j\}_{j=1}^N \subset [0, \pi]$ , there exist potentials  $q \in C^{\infty}[0, +\infty)$  such that

<span id="page-0-1"></span>
$$
\limsup_{x \to \infty} x^{1 - \frac{\alpha}{2}} |q(x)| \le (2 - \alpha) e^{2\sqrt{\ln N}} N,
$$

and for any  $j = 1, 2, \cdots, N$ ,  $-u'' - x^{\alpha}u + qu = E_ju$  has an  $L^2(\mathbb{R}^+)$  solution u with the boundary condition

$$
\frac{u'(0)}{u(0)} = \tan \theta_j.
$$

#### 2 WENCAI LIU

Theorem [1.1](#page-0-0) implies that in order to create N linearly independent eigensolutions in  $L^2(\mathbb{R}^+),$ the quantity a given by [\(2\)](#page-0-1) must be equal or larger than  $\frac{2-\alpha}{\sqrt{2}}N^{\frac{1}{2}}$ . However, Theorem [1.2](#page-0-2) shows that if we allow  $a \ge (2-\alpha)e^{2\sqrt{\ln N}} N$ , one can create N eigensolutions in  $L^2(\mathbb{R}^+)$  for arbitrary N. There is a gap between  $N^{\frac{1}{2}}$  and  $e^{2\sqrt{\ln N}}N$ . It is natural to ask what is the sharp bound of a to create N linearly independent eigensolutions in  $L^2(\mathbb{R}^+)$ .

**Question 1:** What is the minimum of  $\gamma$  such that for any N, there is a potential q on  $\mathbb{R}^+$ such that  $\#P \geq N$  and

$$
\limsup_{x \to \infty} x^{1 - \frac{\alpha}{2}} |q(x)| \le C(\gamma) N^{\gamma}.
$$

Theorems [1.1](#page-0-0) and [1.2](#page-0-2) imply  $\gamma \in [\frac{1}{2}, 1]$ .

Our first result in this paper is to show that for any  $\alpha$  satisfying  $\frac{2}{3} < \alpha < 2$ ,  $\gamma = \frac{1}{2}$  is the solution to Question 1.

<span id="page-1-0"></span>**Theorem 1.3.** Suppose  $\frac{2}{3} < \alpha < 2$ . Then for any  $N > 0$ , there exists a potential q on  $\mathbb{R}^+$ such that

(4) 
$$
\limsup_{x \to \infty} x^{1-\frac{\alpha}{2}} |q(x)| \leq 48(2-\alpha)\sqrt{N \ln N}
$$

and  $\#P = N$ .

For some technical reasons, currently we can only give the proof for  $\frac{2}{3} < \alpha < 2$ . We believe it that  $\gamma = \frac{1}{2}$  is the solution to Question 1 for all  $0 < \alpha < 2$ .

Question 1 and Theorem [1.3](#page-1-0) do not care about the locations of the corresponding energies. If we take the distribution of energies into consideration, what is the sharp upper bound? We formulate it as the following question.

**Question 2:** What is the minimum of  $\gamma$  such that for any  $\{E_j\}_{j=1}^N$ , there exists a potential q on  $\mathbb{R}^+$  such that  $-u'' - x^{\alpha}u + qu = E_ju$  has an  $L^2(\mathbb{R}^+)$  solution for each  $j = 1, 2, \dots, N$  and

$$
\limsup_{x \to \infty} x^{1 - \frac{\alpha}{2}} |q(x)| \le C(\gamma) N^{\gamma}
$$

.

Theorems [1.1](#page-0-0) and [1.2](#page-0-2) imply  $\gamma \in [\frac{1}{2}, 1]$ . We conjecture that  $\gamma = 1$  is the solution to Question 2.

During the proof Theorem [1.3,](#page-1-0) we are able to improve the bound in Theorem [1.2.](#page-0-2)

<span id="page-1-1"></span>**Theorem 1.4.** For any  $\varepsilon > 0$ ,  $\{E_j\}_{j=1}^N \subset \mathbb{R}$  and  $\{\theta_j\}_{j=1}^N \subset [0, \pi]$ , there exist local  $L^1(\mathbb{R}^+)$ potentials q such that

(5) 
$$
\limsup_{x \to \infty} x^{1 - \frac{\alpha}{2}} |q(x)| \le (2 - \alpha + \varepsilon) N,
$$

and for each  $j = 1, 2, \cdots, N$ ,  $-u'' - x^{\alpha}u + qu = E_ju$  has an  $L^2(\mathbb{R}^+)$  solution u with the boundary condition

<span id="page-1-2"></span>
$$
\frac{u'(0)}{u(0)} = \tan \theta_j.
$$

Remark 1.5. • Theorem [1.4](#page-1-1) gives better bounds than those in Theorem [1.1,](#page-0-0) but less regularity in potentials.

• Applying additional piecewise constructions in the proof of Theorem [1.4,](#page-1-1) it is possible to show that the upper bound in [\(5\)](#page-1-2) can be improved to  $(2-\alpha)N$ . We refer the readers to the critical case of [\[25](#page-11-0), Theorem 1.2 ] for details.

The proof of both Theorems [1.3](#page-1-0) and [1.4](#page-1-1) are inspired by the methods tackling perturbed free Schrödinger operators. Let us turn to perturbed free Schrödinger operators  $-D^2+V$  first. Naboko [\[33](#page-12-2)] and Simon [\[36](#page-12-3)] constructed power-decaying potentials V such that  $-D^2 + V$  has

dense eigenvalues. Before that, Wigner-von Neumann type functions can only create one  $L^2$ solution [\[38](#page-12-4)]. Recently, there have been several important developments on the problem of embedded eigenvalues for Schrödinger operators, Laplacians on manifolds or other models [\[11,](#page-11-12) [13](#page-11-13)[–16,](#page-11-14) [23](#page-11-15), [25,](#page-11-0) [27](#page-11-16)[–30,](#page-12-5) [32\]](#page-12-6). For perturbed Stark type operators, under the rational independence assumption of set  $\{E_i\}$ , Naboko and Pushnitskii [\[34\]](#page-12-7) constructed operators with given a set  ${E_i}$  as embedded eigenvalues. The author [\[25\]](#page-11-0) constructed perturbed Stark type operators with any given  ${E_i}$  as a set of eigenvalues with the quantitative bound (see Theorem [1.2\)](#page-0-2). However, the potential can not be given explicitly. One of the motivations of this paper is to approach the problem in an explicit way.

In [\[36\]](#page-12-3), Simon used Wigner-von Neumann type functions  $V(x) = \frac{a}{1+x} \sum_j \sin(2\lambda_j x + 2\phi_j) \chi_{[a_j,\infty)},$ to complete his constructions. It turns out that Wigner-von Neumann type function is a good way to create embedded eigenvalues [\[14,](#page-11-17) [16,](#page-11-14) [24](#page-11-18), [29](#page-12-8)[–31\]](#page-12-9). Moreover, Wigner-von Neumann type functions can also be used to achieve the optimal bounds. Denote by

<span id="page-2-0"></span>
$$
S = \{ E > 0 : -u''(x) + V(x)u(x) = Eu(x) \text{ has an } L^{2}(\mathbb{R}^{+}) \text{ solution } \}.
$$

Kiselev-Last-Simon [\[20](#page-11-19)] proved if  $\limsup_{x\to\infty} x|V(x)| < \infty$ , then the set S is countable and

(6) 
$$
\sum_{E_i \in S} E_i < \infty.
$$

This result has been extended to perturbed periodic operators by the author [\[26](#page-11-20)]. By Wignervon Neumann type functions and additional probability arguments from [\[17](#page-11-21)], Remling [\[35\]](#page-12-10) proved that there are potentials  $V(x) = O(x^{-1})$  with  $\sum_{E_i \in S} E_i^p = \infty$  for every  $p < 1$ . Remling's result implies that [\(6\)](#page-2-0) can not be improved in some sense, which answers a question in [\[20](#page-11-19)].

Another motivation of the present paper is to find the substitution of Wigner-von Neumann type functions to deal with perturbed Stark type operators, so that we can use the ideas of Simon, Remling, and among others to address our problems. We found a good type of substituted functions  $\frac{\sin(\int_0^{\xi} \sqrt{1-\beta E_j(x)} dx + t_j)}{\xi}$  $\frac{\sum_{i=1}^{n} (x_i - x_i - x_j)}{\xi}$ , which is called the *generalized Wigner-von Neumann* type function. See the definition of  $\beta_{E_j}(x)$  below.

Although the arguments in this paper are inspired by those in dealing with perturbed free Schrödinger operators, the details are much more delicate and difficult, in particular, oscillated integrals and resonant phenomena. The spectra of perturbed free Schrödinger operators and Stark operators are quite different. Under the assumption  $V(x) = O(x^{-1})$ , perturbed free Schrödinger operators can have infinitely many eigenvalues. However, under the corresponding assumption, perturbed Stark operators can only have finitely many eigenvalues by Theorem [1.1.](#page-0-0) Moreover, for free cases, there is only one leading entry dominating each Prüfer angle and the leading entries are distinguished by energies. For Stark type cases, there are finitely many entries (the number of entries depends on  $\alpha$ ) dominating each Prüfer angle, and the first/leading entry is 1 for any Prüfer angle, which leads to resonance. Similar resonance has been studied in [\[25\]](#page-11-0). In the proof of Theorem [1.4,](#page-1-1) we are able to deal with the resonance coming from the first dominating entry and all the other dominating entries at the same time. However, for the technical reason, we can only deal with the first two dominating entry for the topic in Theorem [1.3,](#page-1-0) and the assumption  $\alpha > \frac{2}{3}$  will guarantee that there are exact two entries to dominate Prüfer angles.

## <span id="page-2-1"></span>2. Preparations

Let  $v_\alpha(x) = x^\alpha$  for  $x \in \mathbb{R}^+$  and consider the Schrödinger equation on  $\mathbb{R}^+$ ,

(7) 
$$
-u''(x) - x^{\alpha}u(x) + q(x)u(x) = Eu(x).
$$

The Liouville transformation (see [\[3,](#page-11-2) [34\]](#page-12-7)) is given by

(8) 
$$
\xi(x) = \int_0^x \sqrt{v_\alpha(t)} dt, \phi(\xi) = v_\alpha(x(\xi))^{\frac{1}{4}} u(x(\xi)).
$$

We define a weight function  $p_{\alpha}(\xi)$  by

(9) 
$$
p_{\alpha}(\xi) = \frac{1}{v_{\alpha}(x(\xi))}.
$$

We also define a potential by

(10) 
$$
Q_{\alpha}(\xi, E) = -\frac{5}{16} \frac{|v_{\alpha}'(x(\xi))|^2}{v_{\alpha}(x(\xi))^3} + \frac{1}{4} \frac{v_{\alpha}''(x(\xi))}{v_{\alpha}(x(\xi))^2} + \frac{q(x(\xi)) - E}{v_{\alpha}(x(\xi))}.
$$

Let  $c = (1 + \frac{\alpha}{2})^{\frac{2}{2+\alpha}}$ . Direct computations imply that

(11) 
$$
x = c\xi^{\frac{2}{2+\alpha}}, \phi(\xi, E) = c^{\frac{\alpha}{4}}\xi^{\frac{\alpha}{2(2+\alpha)}}u(c\xi^{\frac{2}{2+\alpha}}),
$$

<span id="page-3-4"></span>(12) 
$$
p(\xi) = \frac{1}{c^{\alpha} \xi^{\frac{2\alpha}{2+\alpha}}},
$$

and

(13) 
$$
Q_{\alpha}(\xi, E) = -\frac{5}{4} \frac{\alpha^2}{(2+\alpha)^2} \frac{1}{\xi^2} + \frac{\alpha(\alpha-1)}{(2+\alpha)^2} \frac{1}{\xi^2} + \frac{q(c\xi^{\frac{2}{2+\alpha}}) - E}{c^{\alpha}\xi^{\frac{2\alpha}{2+\alpha}}}.
$$

Notice that the potential  $Q_{\alpha}(\xi, E)$  depends on q,  $\alpha$  and E. In the following, we always fix  $\alpha \in (0, 2)$ . For simplicity, we drop off its dependence. Let

<span id="page-3-0"></span>(14) 
$$
V(\xi) = \frac{q(c\xi^{\frac{2}{2+\alpha}})}{c^{\alpha}\xi^{\frac{2\alpha}{2+\alpha}}}.
$$

Then

<span id="page-3-1"></span>(15) 
$$
Q(\xi, E) = -\frac{5}{4} \frac{\alpha^2}{(2+\alpha)^2} \frac{1}{\xi^2} + \frac{\alpha(\alpha-1)}{(2+\alpha)^2} \frac{1}{\xi^2} - \frac{E}{c^{\alpha} \xi^{\frac{2\alpha}{2+\alpha}}} + V(\xi)
$$

$$
= -\frac{E}{c^{\alpha} \xi^{\frac{2\alpha}{2+\alpha}}} + V(\xi) + \frac{O(1)}{\xi^2}.
$$

Suppose  $u \in L^2(\mathbb{R}^+)$  is a solution of [\(7\)](#page-2-1). It follows that  $\phi$  satisfies

(16) 
$$
-\frac{d^2\phi}{d\xi^2} + Q(\xi, E)\phi = \phi,
$$

and  $\phi \in L^2(\mathbb{R}^+, p(\xi)d\xi)$ .

Below,  $\epsilon > 0$  always depends on  $\alpha$  in an explicit way. Denote by

<span id="page-3-2"></span>
$$
\beta_E(\xi) = -\frac{E}{c^{\alpha} \xi^{\frac{2\alpha}{2+\alpha}}}.
$$

When  $\xi$  is large, one has  $|\beta_E(\xi)| < 1$ . By shifting the equation, we always assume  $\beta_E(\xi)$  is sufficiently small. So  $\sqrt{1 - \beta_E(\xi)}$  is well defined.

<span id="page-3-3"></span>**Proposition 2.1.** Suppose  $a \neq 0$ . Then following estimates hold for  $\xi > \xi_0 > 1$  and  $\gamma \in \mathbb{R}$ : (1)

(17) 
$$
\int_{\xi_0}^{\xi} \frac{\sin(a \int_0^s \sqrt{1 - \beta_E(x)} dx + \gamma)}{s} ds = \frac{O(1)}{\xi_0^{\xi}}.
$$

<span id="page-4-0"></span>(2) for any  $E_1 \neq E_2 \in \mathbb{R}$ ,

(18) 
$$
\int_{\xi_0}^{\xi} \frac{\sin(a \int_0^s \sqrt{1 - \beta_{E_1}(x)} dx \pm a \int_0^s \sqrt{1 - \beta_{E_2}(x)} dx + \gamma)}{s} ds = \frac{O(1)}{\xi_0^{\epsilon}}
$$

Proof. We only give the proof of case "−" in [\(18\)](#page-4-0). The rest can be proceeded in a similar way. Denote by  $\beta(\xi) = a \int_0^{\xi} \left[ \sqrt{1 - \beta_{E_1}(x)} dx - a \int_0^{\xi} \sqrt{1 - \beta_{E_2}(x)} dx + \gamma$ . Then

$$
\beta'(\xi) = a\sqrt{1 - \beta_{E_1}(\xi)} - a\sqrt{1 - \beta_{E_2}(\xi)} = a\frac{E_1 - E_2}{2c^{\alpha}\xi^{\frac{2\alpha}{2+\alpha}}} + \frac{o(1)}{\xi^{\frac{2\alpha}{2+\alpha}}},
$$

and

$$
\beta''(\xi) = \frac{O(1)}{\xi^{1 + \frac{2\alpha}{2 + \alpha}}}.
$$

Integration by part, we have

$$
\int_{\xi_0}^{\xi} \frac{\sin(a \int_0^s \sqrt{1 - \beta_{E_1}(x)} dx - a \int_0^s \sqrt{1 - \beta_{E_2}(x)} dx + \gamma)}{s}
$$
\n
$$
= \int_{\xi_0}^{\xi} \frac{\sin \beta(s)}{s} ds = \int_{\xi_0}^{\xi} \frac{\beta'(s) \sin \beta(s)}{\beta'(s)s} ds
$$
\n
$$
= O(\xi_0^{-\epsilon}) + O(1) \int_{\xi_0}^{\xi} \frac{\cos \beta(s) \beta''}{s \beta'^2} ds
$$
\n
$$
= O(\xi_0^{-\epsilon}).
$$

### 3. Asymptotical behavior of solutions for a class of linear systems

The proof of this section is inspired by the WKB method. We refer the readers to papers [\[4,](#page-11-22) [6,](#page-11-23) [9\]](#page-11-24) for arguments.

<span id="page-4-2"></span>**Theorem 3.1.** Suppose  $a > 0$  is a constant. Suppose  $\{E_j\} \in \mathbb{R}$  are distinct. Define  $V(\xi) = 0$ for  $\xi \in [0,1]$  and

<span id="page-4-1"></span>(19) 
$$
V(\xi) = \frac{4a}{\xi} \sum_{j=1}^{N} \sin \left( \int_0^{\xi} 2\sqrt{1 - \beta_{E_j}(x)} dx + 2t_j \right),
$$

for  $\xi > 1$ .

Define  $q(x)$  on  $[0, \infty)$  such that [\(14\)](#page-3-0) holds for [\(19\)](#page-4-1). Let  $Q(\xi, E)$  be given by [\(15\)](#page-3-1). Then the following asymptotics hold as  $\xi$  goes to infinity,

(1) if  $E \neq E_j$  for any  $j = 1, 2, \cdots N$ , then there exists a fundamental system of solutions  ${y_1(\xi), y_2(\xi)}$  of [\(16\)](#page-3-2) such that

$$
\begin{bmatrix} y_1(\xi) \\ y'_1(\xi) \end{bmatrix} = \begin{bmatrix} \cos(\int_0^{\xi} \sqrt{1 - \beta_E(x)} dx + t_j) \\ -\sin(\int_0^{\xi} \sqrt{1 - \beta_E(x)} dx + t_j) \end{bmatrix} + O(\xi^{-\epsilon})
$$

and

$$
\begin{bmatrix} y_2(\xi) \\ y'_2(\xi) \end{bmatrix} = \begin{bmatrix} \sin(\int_0^{\xi} \sqrt{1 - \beta_E(x)} dx + t_j) \\ \cos(\int_0^{\xi} \sqrt{1 - \beta_E(x)} dx + t_j) \end{bmatrix} + O(\xi^{-\epsilon}).
$$

 $\Box$ 

.

### 6 WENCAI LIU

(2) if  $E = E_j$  for some j, then there exists a fundamental system of solutions  $\{y_1(\xi), y_2(\xi)\}$ of [\(16\)](#page-3-2) such that

$$
\begin{bmatrix} y_1(\xi) \\ y'_1(\xi) \end{bmatrix} = \xi^a \begin{bmatrix} \cos(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) \\ -\sin(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) \end{bmatrix} + O(\xi^{a-\epsilon})
$$

and

$$
\begin{bmatrix} y_2(\xi) \\ y_2'(\xi) \end{bmatrix} = \xi^{-a} \begin{bmatrix} \sin(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) \\ \cos(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) \end{bmatrix} + O(\xi^{-a-\epsilon}).
$$

*Proof.* In order to avoid repetition, we only give the proof of the case the case  $E = E_j$  for some  $j = 1, 2, \cdots, N$ . Denote by

(20) 
$$
\tilde{V}(\xi) = \frac{4a}{\xi} \sum_{i=1, i \neq j}^{N} \sin(2 \int_{0}^{\xi} \sqrt{1 - \beta_{E_i}(x)} dx + 2t_i).
$$

Rewrite the second order differential equation of [\(16\)](#page-3-2) as the linearly differential equations,

$$
\begin{bmatrix} y \\ y' \end{bmatrix}' = \begin{bmatrix} 0 & 1 \\ \beta_{E_j}(\xi) + V(\xi) + \frac{O(1)}{\xi^2} - 1 & 0 \end{bmatrix} \begin{bmatrix} y \\ y' \end{bmatrix}.
$$

$$
\begin{bmatrix} u_1 \end{bmatrix} = \begin{bmatrix} \sqrt{1 - \beta_{E_j}(\xi)} & 0 \end{bmatrix} \begin{bmatrix} y \\ y' \end{bmatrix}.
$$

Let

$$
\left[\begin{array}{c} u_1 \\ u_2 \end{array}\right] = \left[\begin{array}{cc} \sqrt{1-\beta_{E_j}(\xi)} & 0 \\ 0 & 1 \end{array}\right] \left[\begin{array}{c} y \\ y' \end{array}\right].
$$

We obtain a new equation

$$
\begin{bmatrix} u_1 \\ u_2 \end{bmatrix}' = \left( \begin{bmatrix} 0 \\ -\sqrt{1-\beta_{E_j}(\xi)} + \frac{V}{\sqrt{1-\beta_{E_j}(\xi)}} & 1 \end{bmatrix} + \frac{O(1)}{\xi^{1+\epsilon}} \right) \begin{bmatrix} y \\ y' \end{bmatrix}.
$$

Let

$$
\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} \cos(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) & -\sin(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) \\ \sin(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) & \cos(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix}.
$$

Obviously, one has

$$
\begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} \cos(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) & \sin(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) \\ -\sin(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) & \cos(\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + t_j) \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}.
$$

After some calculations, we have

$$
\begin{bmatrix} y_1 \\ y_2 \end{bmatrix}' = (\Lambda(\xi) + H(\xi) + O(\xi^{-1-\epsilon})) \begin{bmatrix} y_1 \\ y_2 \end{bmatrix},
$$

where

(21) 
$$
\Lambda(\xi) = \begin{bmatrix} -2a \frac{\sin^2(2\int_0^{\xi} \sqrt{1-\beta_{E_j}(x)} dx + 2t_j)}{\xi} & 0\\ 0 & 2a \frac{\sin^2(2\int_0^{\xi} \sqrt{1-\beta_{E_j}(x)} dx + 2t_j)}{\xi} \end{bmatrix}
$$

and

$$
H(\xi) = \left[ \begin{array}{cc} H_{11}(\xi) & H_{12}(\xi) \\ H_{21}(\xi) & H_{22}(\xi) \end{array} \right].
$$

The explicit formulas for  $H_{ij}$ ,  $i, j = 1, 2$  are

$$
H_{11} = -\frac{1}{2}\tilde{V}(\xi)\sin\left(2\int_0^{\xi}\sqrt{1-\beta_{E_j}(x)}dx + 2t_j\right),
$$

$$
H_{22}(\xi) = \frac{1}{2}\tilde{V}(\xi)\sin\left(2\int_0^{\xi}\sqrt{1-\beta_{E_j}(x)}dx + 2t_j\right),
$$
  

$$
H_{12}(\xi) = -\frac{1}{2}V(\xi)\left(1-\cos\left(2\int_0^{\xi}\sqrt{1-\beta_{E_j}(x)}dx + 2t_j\right)\right),
$$

and

$$
H_{21}(\xi) = \frac{1}{2}V(\xi)\left(1 + \cos\left(2\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)}dx + 2t_j\right)\right).
$$
  
1, one has

By Proposition [2.1,](#page-3-3) one has

<span id="page-6-1"></span>
$$
Q(\xi) \equiv -\int_{\xi}^{\infty} H(s)ds = O(\xi^{-\epsilon}).
$$

Assume  $\xi$  is large, then  $||Q|| \leq \frac{1}{2}$ . Let  $\begin{bmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{bmatrix}$  $\tilde y_2$  $= (I + Q)^{-1} \left[ \begin{array}{c} y_1 \\ y_2 \end{array} \right]$  $y_2$ . We obtain

(22) 
$$
\begin{bmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{bmatrix}' = (\Lambda(\xi) + R(\xi)) \begin{bmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{bmatrix},
$$

where  $R(\xi) = (R_{ij}) = O(\xi^{-1-\epsilon})$ . Let  $\varphi(\xi) = \begin{bmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{bmatrix}$  $\tilde{y}_2$  and  $\lambda(\xi) = 2a \frac{\sin^2(2\int_0^{\xi} \sqrt{1-\beta_{E_j}(x)}dx + 2t_j)}{\xi}$  $\frac{1}{\xi}$ .

Let us consider the integral equation,

<span id="page-6-0"></span>(23) 
$$
\varphi(\xi) = \begin{bmatrix} e^{-\int_1^{\xi} \lambda(s)ds} \\ 0 \end{bmatrix} - \int_{\xi}^{\infty} \begin{bmatrix} e^{-\int_{\xi}^{y} \lambda(s)ds} & 0 \\ 0 & e^{\int_{\xi}^{y} \lambda(s)ds} \end{bmatrix} R(y)\varphi(y)dy.
$$

If [\(23\)](#page-6-0) has a solution  $\|\varphi(\xi)\| \leq 2e^{-\int_1^{\xi} \lambda(s)ds}$ , then (by direct computation)  $\varphi(\xi)$  is a solution of equation [\(22\)](#page-6-1). Moreover

<span id="page-6-2"></span>
$$
\|\int_{\xi}^{\infty} \begin{bmatrix} e^{-\int_{\xi}^{y} \lambda(s)ds} & 0\\ 0 & e^{\int_{\xi}^{y} \lambda(s)ds} \end{bmatrix} R(y)\varphi(y)dy\| = O(1) \int_{\xi}^{\infty} e^{\int_{\xi}^{y} \lambda(s)} ||R(y)||e^{-\int_{1}^{y} \lambda(s)ds}
$$
  

$$
= O(1)e^{-\int_{1}^{\xi} \lambda(s)ds} \int_{\xi}^{\infty} ||R(y)||
$$
  

$$
= e^{-\int_{1}^{\xi} \lambda(s)ds} O(\xi^{-\epsilon}).
$$

Define iteration equations:

(25) 
$$
\varphi_k(\xi) = \begin{bmatrix} e^{-\int_1^{\xi} \lambda(s)ds} \\ 0 \end{bmatrix} - \int_{\xi}^{\infty} \begin{bmatrix} e^{-\int_{\xi}^{y} \lambda(s)ds} & 0 \\ 0 & e^{\int_{\xi}^{y} \lambda(s)ds} \end{bmatrix} R(y) \varphi_{k-1}(y) dy,
$$

with  $\varphi = 0$ . By induction that  $||\varphi_k(\xi) - \varphi_{k-1}(\xi)|| \leq \frac{1}{2^{k+1}} e^{-\int_1^{\xi} \lambda(s) ds}$ , one can show [\(23\)](#page-6-0) has a solution  $||\varphi(\xi)|| \leq 2e^{-\int_1^{\xi} \lambda(s)ds}$  (see p.94 in [\[4](#page-11-22)] for all the details).

By  $(22)$ ,  $(23)$  and  $(24)$ , we get a solution

$$
\begin{bmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{bmatrix} = e^{-\int_1^{\xi} \lambda(s)ds} \left( \begin{bmatrix} 1 \\ 0 \end{bmatrix} + O(\xi^{-\epsilon}) \right).
$$

By Proposition [2.1](#page-3-3) again, we have

$$
\int_{1}^{\xi} \lambda(s)ds = 2a \int_{1}^{\xi} \frac{\sin^{2}(2 \int_{0}^{s} \sqrt{1 - \beta_{E_{j}}(x)} dx + 2t_{j})}{s} ds
$$
  
\n
$$
= a \int_{1}^{\xi} \frac{1}{s} ds - a \int_{1}^{\xi} \frac{\cos(4 \int_{0}^{s} \sqrt{1 - \beta_{E_{j}}(x)} dx + 4t_{j})}{s} ds
$$
  
\n
$$
= a \int_{1}^{\xi} \frac{1}{s} ds - a \int_{1}^{\infty} \frac{\cos(4 \int_{0}^{s} \sqrt{1 - \beta_{E_{j}}(x)} dx + 4t_{j})}{s} ds
$$
  
\n
$$
+ a \int_{\xi}^{\infty} \frac{\cos(4 \int_{0}^{s} \sqrt{1 - \beta_{E_{j}}(x)} dx + 4t_{j})}{s} ds
$$
  
\n
$$
= \ln \xi^{a} - c + O(\xi^{-\epsilon}),
$$

where the constant  $c$  equals

<span id="page-7-0"></span>
$$
a\int_1^\infty \frac{\cos(4\int_0^s \sqrt{1-\beta_{E_j}(x)}dx+4t_j)}{s}ds.
$$

Thus [\(22\)](#page-6-1) has a solution

(26) 
$$
\begin{bmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{bmatrix} = \xi^{-a} \begin{bmatrix} 1 \\ 0 \end{bmatrix} + O(\xi^{-a-\epsilon}).
$$

By the similar argument (see [\[4\]](#page-11-22) again), we obtain that [\(22\)](#page-6-1) has a solution

(27) 
$$
\begin{bmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{bmatrix} = \xi^a \begin{bmatrix} 1 \\ 0 \end{bmatrix} + O(\xi^{a-\epsilon}).
$$

Now the Theorem follows from  $(26)$  and  $(27)$ .

## <span id="page-7-1"></span>4. Proof of Theorems [1.3](#page-1-0) and [1.4](#page-1-1)

**Proof of Theorem [1.4.](#page-1-1)** Let  $a > \frac{2-\alpha}{2(2+\alpha)}$ . Define potentials

$$
V(\xi) = \frac{4a}{\xi} \sum_{j=1}^{N} \sin(2 \int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + 2t_j) \chi_{[a_j,\infty)},
$$

with large  $a_i$ .

Obviously,

<span id="page-7-3"></span><span id="page-7-2"></span>
$$
|\xi V(\xi)| \le 4aN,
$$

so that

(28) 
$$
|\xi^{1-\frac{\alpha}{2}}q(\xi)| \leq 2(2+\alpha)aN.
$$

By [\(12\)](#page-3-4),

(29) 
$$
p(\xi)|\phi(\xi, E_j)|^2 \leq O(1)\xi^{-2a - \frac{2\alpha}{2+\alpha}} \leq O(1)\xi^{-1-\epsilon}.
$$

By [\(29\)](#page-7-2), [\(16\)](#page-3-2) has a solution  $\phi(\xi, E_j) \in L^2(\mathbb{R}^+, p(\xi)d\xi)$  for each  $j = 1, 2, \cdots, N$ . However,  $\phi(\xi, E_j)$  may not satisfy the given boundary condition. This can be done by adjusting  $a_j$  and additional functions W with support in  $(1, 2)$ . We refer readers to [\[36\]](#page-12-3) for rigorous arguments. Now the Theorem follows from  $(28)$ .

Let

$$
\tau=\frac{2+\alpha}{2(2-\alpha)}\frac{1}{(1+\frac{\alpha}{2})^{\frac{2\alpha}{2+\alpha}}},
$$

so that

$$
\tau \frac{d\xi^{\frac{2-\alpha}{2+\alpha}}}{d\xi} = \frac{1}{2} \frac{1}{c^{\alpha} \xi^{\frac{2\alpha}{2+\alpha}}}.
$$

For any given  $N$ , let

$$
E_j = \frac{j}{N\tau}
$$
, for  $j = 1, 2, \dots, N$ .

Before giving the proof of Theorem [1.3,](#page-1-0) one key Lemma is needed, which is motivated by [\[35\]](#page-12-10).

<span id="page-8-0"></span>**Lemma 4.1.** There exist  $\theta_j \in [0, 1)$  for  $j = 1, 2, \dots, N$  such that for any  $\xi > 0$ 

(30) 
$$
|\sum_{j=1}^{N} \sin(2\tau E_j \xi + 2\pi \theta_j)| + |\sum_{j=1}^{N} \cos(2\tau E_j \xi + 2\pi \theta_j)| \le 4\sqrt{2N \ln(8(N+1)N)}.
$$

Proof. Let

(31) 
$$
\tilde{E}_j = \frac{j}{N}, \text{ for } j = 1, 2, \cdots, N.
$$

It suffices to show there exist  $\theta = (\theta_1, \theta_2, \cdots, \theta_N) \in [0, 1)^N$  such that for any  $\xi > 0$ ,

(32) 
$$
f(\xi, \theta) \equiv |\sum_{j=1}^{N} \sin(2\tilde{E}_j \xi + 2\pi \theta_j)| + |\sum_{j=1}^{N} \cos(2\tilde{E}_j \xi + 2\pi \theta_j)| \le 4\sqrt{2N \ln(8(N+1)N)}.
$$

Let

$$
f_1(\xi, \theta) = \sum_{j=1}^N \sin(2\tilde{E}_j \xi + 2\pi\theta_j),
$$

and

$$
M_1(\theta) = \sup_{\xi \in \mathbb{R}^+} |f_1(\xi, \theta)| = \max_{0 \le \xi \le \pi N} |f_1(\xi, \theta)|.
$$

Pick  $\xi_0$  so that  $|f_1(\xi_0, \theta)| = M_1(\theta)$ . Notice that

$$
1 = \frac{2}{\pi N} \int_0^{\pi N} f_1(\xi, \theta) \sin(2\tilde{E}_j \xi + 2\pi \theta_j) d\xi \le 2M_1(\theta),
$$

and

$$
\left| \frac{df_1(\xi, \theta)}{d\xi} \right| = 2|\sum_{j=1}^N \tilde{E}_j \cos(2\tilde{E}_j \xi + \theta_j)|
$$
  

$$
\leq 2|\sum_{j=1}^N \tilde{E}_j|
$$
  

$$
= N + 1.
$$

It follows that there exits a interval  $I(\theta)$  of  $\xi$  centered at  $\xi_0$  with size  $\frac{1}{2(N+1)}$   $(I(\theta)$  means the interval depends on  $\theta$ ) such that

$$
|f_1(\xi,\theta)| \ge M_1(\theta) - \frac{1}{4(N+1)}(N+1) \ge \frac{M_1(\theta)}{2},
$$

for all  $\xi \in I(\theta)$ .

Integration yields (p.493, [\[35](#page-12-10)])

$$
\int_0^{2\pi} e^{a\sin(b+2\pi y)} dy \le e^{\frac{a^2}{4}}.
$$

Thus

$$
\frac{1}{2(N+1)} \int_{(\mathbb{R}/\mathbb{Z})^N} e^{\frac{1}{2}\lambda M_1(\theta)} d\theta \le \int_{(\mathbb{R}/\mathbb{Z})^N} d\theta \int_{I(\theta)} (e^{\lambda f_1(\xi,\theta)} + e^{-\lambda f_1(\xi,\theta)}) d\xi
$$
  
\n
$$
\le \int_{(\mathbb{R}/\mathbb{Z})^N} d\theta \int_0^{\pi N} (e^{\lambda f_1(\xi,\theta)} + e^{-\lambda f_1(\xi,\theta)}) d\xi
$$
  
\n
$$
= \int_0^{\pi N} d\xi \int_{(\mathbb{R}/\mathbb{Z})^N} (e^{\lambda f_1(\xi,\theta)} + e^{-\lambda f_1(\xi,\theta)}) d\theta
$$
  
\n
$$
\le \int_0^{\pi N} 2e^{\frac{N}{4}\lambda^2} d\xi
$$
  
\n
$$
= 2\pi N e^{\frac{N}{4}\lambda^2}.
$$

It implies

(33) 
$$
\int_{(\mathbb{R}/\mathbb{Z})^N} e^{\frac{1}{2}\lambda(M_1(\theta)-\frac{N}{2}\lambda-\frac{2}{\lambda}\ln 8N(N+1)-\frac{2}{\lambda}\ln 2\pi)} d\theta \leq \frac{1}{4}.
$$

By [\(33\)](#page-9-0), we have for any  $\lambda > 0$ , there exists a subset  $S_1(\lambda) \subset [0,1)^N$  such that  $\text{Leb}(S_1) \geq \frac{3}{4}$ and

<span id="page-9-0"></span>
$$
M_1(\theta) \le \frac{N}{2}\lambda + \frac{2}{\lambda}\ln 8N(N+1) + \frac{2}{\lambda}\ln 2\pi,
$$

for all  $\theta \in S_1.$ 

Let

$$
f_2(\xi, \theta) = \sum_{j=1}^{N} \cos(2\tilde{E}_j \xi + 2\pi\theta_j),
$$

and

$$
M_2(\theta) = \sup_{\xi \in \mathbb{R}^+} |f_1(\xi, \theta)| = \max_{0 \le \xi \le \pi N} |f_2(\xi, \theta)|.
$$

Similarly, for any  $\lambda > 0$ , there exists a subset  $S_2(\lambda) \subset [0,1)^N$  such that  $\text{Leb}(S_2) \geq \frac{3}{4}$  and

$$
M_2(\theta) \leq \frac{N}{2}\lambda + \frac{2}{\lambda}\ln 8N(N+1) + \frac{2}{\lambda}\ln 2\pi
$$
  
 
$$
\leq M_2(\theta) \leq \frac{N}{2}\lambda + \frac{4}{\lambda}\ln 8N(N+1)).
$$

for all  $\theta \in S_2$ .

Let

$$
\lambda = \sqrt{\frac{8 \ln 8N(N+1)}{N}},
$$

and  $S = S_1(\lambda) \cap S_2(\lambda)$ . Then we have  $\text{Leb}(S) \geq \frac{1}{4}$ , and

$$
M_1(\theta) + M_2(\theta) \le 4\sqrt{2N\ln 8N(N+1)},
$$

for all  $\theta \in S$ . We finish the proof.

$$
E_j = \frac{j}{N\tau}, \text{ for } j = 1, 2, \cdots, N.
$$

Let  $a > \frac{2-\alpha}{2(2+\alpha)}$ . By Lemma [4.1,](#page-8-0) there exist  $t_j \in [0, \pi)$  such that for any  $\xi > 0$ ,

<span id="page-10-0"></span>(34) 
$$
|\sum_{j=1}^{N} \sin(2\tau E_j \xi + 2t_j)| + |\sum_{j=1}^{N} \cos(2\tau E_j \xi + 2t_j)| \le 4\sqrt{2N \ln(8(N+1)N)}.
$$

Let

<span id="page-10-1"></span>(35) 
$$
V(\xi) = \frac{4a}{\xi} \sum_{j=1}^{N} \sin \left( 2 \int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + 2t_j \right).
$$

By Tayor series, one has

<span id="page-10-2"></span>(36)  
\n
$$
\int_0^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx = \int_0^{\xi} \left( 1 - \frac{1}{2} \beta_{E_j}(x) + O(1) \beta_{E_j}(x)^2 \right) dx
$$
\n
$$
= \xi + \tau E_j \xi^{\frac{2-\alpha}{2+\alpha}} + \frac{O(1)}{\xi^{\frac{4\alpha}{2+\alpha} - 1}} + \tilde{t}_j
$$
\n
$$
= \xi + \tau E_j \xi^{\frac{2-\alpha}{2+\alpha}} + O(\xi^{-\epsilon}) + \tilde{t}_j,
$$

since  $\alpha > \frac{2}{3}$ .

By [\(34\)](#page-10-0), [\(35\)](#page-10-1) and [\(36\)](#page-10-2), one has

<span id="page-10-3"></span>
$$
|\xi V(\xi)| = 4a| \sum_{j=1}^{N} \sin \left( 2 \int_{0}^{\xi} \sqrt{1 - \beta_{E_j}(x)} dx + 2t_j \right) |
$$
  
\n
$$
= 4a| \sum_{j=1}^{N} \sin \left( 2\xi + 2\tau E_j \xi^{\frac{\alpha}{2 + \alpha}} + 2t_j + 2\tilde{t}_j \right) | + O(\xi^{-\epsilon})
$$
  
\n
$$
\leq 4a| \sin(2\tau E_j \xi^{\frac{\alpha}{2 + \alpha}} + 2t_j + 2\tilde{t}_j)| + 4a| \cos(2\tau E_j \xi^{\frac{\alpha}{2 + \alpha}} + 2t_j + 2\tilde{t}_j)| + O(\xi^{-\epsilon})
$$
  
\n
$$
\leq 16a\sqrt{2N \ln(8(N+1)N)} + O(\xi^{-\epsilon})
$$
  
\n(37) 
$$
\leq 96a\sqrt{N \ln N},
$$

for large ξ. Define  $q(x)$  on  $[0, \infty)$  such that [\(14\)](#page-3-0) holds for [\(35\)](#page-10-1). Then by [\(37\)](#page-10-3), we have

(38) 
$$
\xi^{1-\frac{\alpha}{2}}|q(\xi)| \le (2+\alpha)48a\sqrt{N\ln N}.
$$

By Theorem [3.1,](#page-4-2) for any  $E_j$ ,  $j = 1, 2, \dots, N$ , [\(16\)](#page-3-2) has a solution  $\phi(\xi, E_j)$  satisfying

<span id="page-10-4"></span>
$$
|\phi(\xi, E_j)| \le 2\xi^{-a}
$$

for large  $\xi$ . By [\(12\)](#page-3-4),

(39) 
$$
p(\xi)|\phi(\xi, E_j)|^2 \le O(1)\xi^{-2a - \frac{2\alpha}{2+\alpha}} \le O(1)\xi^{-1-\epsilon}.
$$

It implies  $\phi(\xi, E_j) \in L^2(\mathbb{R}^+, p(\xi)d\xi)$  and then  $u \in L^2(\mathbb{R}^+)$ . The Theorem follows immediately by applying  $a = \frac{2-\alpha}{2+\alpha}$  to [\(38\)](#page-10-4).  $\Box$ 

## Acknowledgments

The research was supported by NSF DMS-1700314 and NSF DMS-1401204.

#### 12 WENCAI LIU

#### **REFERENCES**

- <span id="page-11-1"></span>[1] S. Agmon, I. Herbst, and S. Maad Sasane. Persistence of embedded eigenvalues. J. Funct. Anal., 261(2):451–477, 2011.
- [2] J. E. Avron and I. W. Herbst. Spectral and scattering theory of Schrödinger operators related to the Stark effect. Comm. Math. Phys., 52(3):239–254, 1977.
- <span id="page-11-2"></span>[3] M. Christ and A. Kiselev. Absolutely continuous spectrum of Stark operators. Ark. Mat., 41(1):1–33, 2003.
- <span id="page-11-22"></span>[4] E. A. Coddington and N. Levinson. Theory of ordinary differential equations. McGraw-Hill Book Company, Inc., New York-Toronto-London, 1955.
- <span id="page-11-3"></span>[5] M. Courtney, N. Spellmeyer, H. Jiao, and D. Kleppner. Classical, semiclassical, and quantum dynamics in the lithium Stark system. Physical Review A, 51(5):3604, 1995.
- <span id="page-11-23"></span>[6] J. D. Dollard and C. N. Friedman. Product integrals and the Schrödinger equation. J. Mathematical Phys., 18(8):1598–1607, 1977.
- <span id="page-11-4"></span>[7] P. S. Epstein. The stark effect from the point of view of Schrödinger's quantum theory. Physical Review, 28(4):695, 1926.
- <span id="page-11-5"></span>[8] S. Graffi and V. Grecchi. Resonances in the Stark effect of atomic systems. Comm. Math. Phys., 79(1):91– 109, 1981.
- <span id="page-11-24"></span>[9] W. A. Harris, Jr. and D. A. Lutz. Asymptotic integration of adiabatic oscillators. J. Math. Anal. Appl., 51:76–93, 1975.
- <span id="page-11-6"></span>[10] I. Herbst and J. Rama. Instability of pre-existing resonances under a small constant electric field. Ann. Henri Poincaré, 16(12):2783-2835, 2015.
- <span id="page-11-12"></span>[11] J. Janas and S. Naboko. On the point spectrum of periodic Jacobi matrices with matrix entries: elementary approach. J. Difference Equ. Appl., 21(11):1103–1118, 2015.
- <span id="page-11-7"></span>[12] A. Jensen and T. Ozawa. Classical and quantum scattering for Stark Hamiltonians with slowly decaying potentials. Ann. Inst. H. Poincaré Phys. Théor., 54(3):229-243, 1991.
- <span id="page-11-13"></span>[13] S. Jitomirskaya and W. Liu. Noncompact complete riemannian manifolds with dense eigenvalues embedded in the essential spectrum of the Laplacian.  $arXiv$  preprint  $arXiv:1805.01072$ , 2018.
- <span id="page-11-17"></span>[14] E. Judge, S. Naboko, and I. Wood. Eigenvalues for perturbed periodic Jacobi matrices by the Wigner–von Neumann approach. Integral Equations Operator Theory, 85(3):427–450, 2016.
- [15] E. Judge, S. Naboko, and I. Wood. Embedded eigenvalues for perturbed periodic Jacobi operators using a geometric approach. J. Difference Equ. Appl., 24(8):1247–1272, 2018.
- <span id="page-11-14"></span>[16] E. Judge, S. Naboko, and I. Wood. Spectral results for perturbed periodic Jacobi matrices using the discrete Levinson technique. Studia Math., 242(2):179–215, 2018.
- <span id="page-11-21"></span>[17] J.-P. Kahane. Some random series of functions, volume 5 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, second edition, 1985.
- <span id="page-11-8"></span>[18] B. Khosropour. The generalized uncertainty principle and the Stark effect. Acta Phys. Polon. B, 48(2):217– 228, 2017.
- <span id="page-11-9"></span>[19] A. Kiselev. Absolutely continuous spectrum of perturbed Stark operators. Trans. Amer. Math. Soc., 352(1):243–256, 2000.
- <span id="page-11-19"></span>[20] A. Kiselev, Y. Last, and B. Simon. Modified Prüfer and EFGP transforms and the spectral analysis of one-dimensional Schrödinger operators. Comm. Math. Phys., 194(1):1–45, 1998.
- <span id="page-11-10"></span>[21] E. L. Korotyaev. Resonances for 1d Stark operators. J. Spectr. Theory, 7(3):699–732, 2017.
- <span id="page-11-11"></span>[22] E. L. Korotyaev. Asymptotics of resonances for 1D Stark operators. Lett. Math. Phys., 108(5):1307–1322, 2018.
- <span id="page-11-15"></span>[23] H. Kumura. The radial curvature of an end that makes eigenvalues vanish in the essential spectrum. I. Math. Ann., 346(4):795–828, 2010.
- <span id="page-11-18"></span>[24] P. Kurasov and S. Naboko. Wigner-von Neumann perturbations of a periodic potential: spectral singularities in bands. Math. Proc. Cambridge Philos. Soc., 142(1):161–183, 2007.
- <span id="page-11-0"></span>[25] W. Liu. Criteria for eigenvalues embedded into the absolutely continuous spectrum of perturbed Stark type operators. J. Funct. Anal. to appear.
- <span id="page-11-20"></span>[26] W. Liu. Asymptotical behaviour of embedded eigenvalues for perturbed periodic operators. arXiv preprint [arXiv:1809.04699](http://arxiv.org/abs/1809.04699), 2018.
- <span id="page-11-16"></span>[27] W. Liu. Criteria for embedded eigenvalues for discrete Schrödinger operators. arXiv preprint [arXiv:1805.02817](http://arxiv.org/abs/1805.02817), 2018.
- [28] W. Liu and D. C. Ong. Sharp spectral transition for eigenvalues embedded into the spectral bands of perturbed periodic operators. J. Anal. Math. to appear.
- <span id="page-12-8"></span>[29] M. Lukic. Schrödinger operators with slowly decaying Wigner-von Neumann type potentials. J. Spectr. Theory, 3(2):147–169, 2013.
- <span id="page-12-5"></span>[30] M. Lukic. A class of Schrödinger operators with decaying oscillatory potentials. Comm. Math. Phys., 326(2):441–458, 2014.
- <span id="page-12-9"></span>[31] M. Lukic and D. C. Ong. Wigner-von Neumann type perturbations of periodic Schrödinger operators. Trans. Amer. Math. Soc., 367(1):707–724, 2015.
- <span id="page-12-6"></span>[32] M. Lukic and D. C. Ong. Generalized Prüfer variables for perturbations of Jacobi and CMV matrices. J. Math. Anal. Appl., 444(2):1490–1514, 2016.
- <span id="page-12-2"></span>[33] S. N. Naboko. On the dense point spectrum of Schrödinger and Dirac operators. Teoret. Mat. Fiz., 68(1):18–28, 1986.
- <span id="page-12-7"></span>[34] S. N. Naboko and A. B. Pushnitskii. Point spectrum on a continuous spectrum for weakly perturbed Stark type operators. Functional Analysis and its Applications, 29(4):248–257, 1995.
- <span id="page-12-10"></span>[35] C. Remling. Schrödinger operators with decaying potentials: some counterexamples. Duke Math. J., 105(3):463–496, 2000.
- <span id="page-12-3"></span>[36] B. Simon. Some Schrödinger operators with dense point spectrum. Proc. Amer. Math. Soc., 125(1):203– 208, 1997.
- <span id="page-12-0"></span>[37] J. C. Solem. Variations on the Kepler problem. Foundations of Physics, 27(9):1291–1306, 1997.
- <span id="page-12-4"></span>[38] J. von Neuman and E. Wigner. Uber merkwürdige diskrete Eigenwerte. Uber das Verhalten von Eigenwerten bei adiabatischen Prozessen. Zhurnal Physik, 30:467–470, 1929.
- <span id="page-12-1"></span>[39] K. Yajima. Spectral and scattering theory for Schrödinger operators with Stark effect. J. Fac. Sci. Univ. Tokyo Sect. IA Math., 26(3):377–390, 1979.

(Wencai Liu) Department of Mathematics, University of California, Irvine, California 92697-3875, USA

E-mail address: liuwencai1226@gmail.com