

# The Number of Fractional Powers in the Fibonacci Word

Huang Yuke<sup>1</sup> Wen Zhiying<sup>2</sup>

## ABSTRACT

The Fibonacci word is the fixed point beginning with the letter  $a$  of morphism  $\sigma(a) = ab, \sigma(b) = a$  defined over the alphabet  $\{a, b\}$ . In this paper, we get explicit expression of the number of distinct fractional powers in each factor of the Fibonacci word.

**Key words** the Fibonacci word; repetition; factor spectrum; the return word sequence.

**2010 MR Subject Classification** 68R15

## 1 Introduction

Let  $\mathcal{A}$  be a finite set called alphabet. Let  $\omega = x_1x_2 \cdots x_N$  be a finite word over  $\mathcal{A}$ . We denote by  $|\omega|$  (resp.  $|\omega|_\alpha, \#\omega$ ) the number of letters (resp. letter  $\alpha$ , distinct letters) in  $\omega$ . For  $1 \leq i \leq j \leq N$ , we define  $\omega[i, j] = x_i x_{i+1} \cdots x_{j-1} x_j$  or  $[x_i, x_{i+1}, \dots, x_{j-1}, x_j]$  by convention,  $\omega[i] = \omega[i, i] = x_i$  and  $\omega[i, i-1] = \varepsilon$  (empty word). We call  $\omega[i, j]$  (resp.  $\omega[1, j], \omega[i, N]$ ) a factor (resp. prefix, suffix) of  $\omega$ , denoted by  $\omega[i, j] \prec \omega$  (resp.  $\omega[1, j] \triangleleft \omega, \omega[i, N] \triangleright \omega$ ). For any  $i \geq 1$  and  $i+n-1 \leq N$ , we define  $\omega[i; n] = \omega[i, i+n-1]$ . The position of factor  $\omega[i, j]$  in  $\omega$  is defined by  $i$ . An infinite word is said to be recurrent if every factor occurs infinitely often [1]. Let  $\omega$  be a factor of a recurrent infinite word. We denote the position of the  $p$ -th occurrence of  $\omega$  by  $(\omega)_p$ . When  $|\omega| = 1$ , we denote  $(\omega)_p$  by  $\omega_p$  for short. Let  $\omega$  and  $\nu$  be two words.  $\omega \cdot \nu$  is the concatenation of  $\omega$  and  $\nu$ . Let  $S$  be a set. We denote  $\#S$  the number of elements in  $S$ . For a real number  $r$ , the floor function  $\lfloor r \rfloor$  is the greatest integer less than or equal to  $r$ , and the ceiling function  $\lceil r \rceil$  is the least integer greater than or equal to  $r$ .

The Fibonacci morphism  $\sigma$  over alphabet  $\{a, b\}$  is a substitution defined by  $\sigma(a) = ab$  and  $\sigma(b) = a$ . The Fibonacci word  $\mathbb{F}$  is defined to be the fixed point beginning with the letter  $a$  of the Fibonacci morphism. It is a recurrent infinite word [1]. Define  $F_m = \sigma^m(a)$  for  $m \geq 0$ , by convention,  $F_{-1} = b$  and  $F_{-2} = \varepsilon$ . The  $m$ -th Fibonacci number  $f_m$  is equal to  $|F_m|$ . As a classical example of sequences over the binary alphabet, the Fibonacci word  $\mathbb{F}$  has many remarkable properties. We refer to Lothaire [16, 17], Allouche-Shallit [1] and Berstel [2].

Let  $R_a$  and  $R_b$  be two words over alphabet  $\mathcal{A}$ . The Fibonacci word over alphabet  $\{R_a, R_b\}$  is denoted by  $\mathbb{F}(R_a, R_b)$ . In this paper, we always assume that there exists an integer  $m \geq 0$  such that  $|R_a| = f_m$  and  $|R_b| = f_{m-1}$ . In particular, when  $\mathcal{A} = \{a, b\}$ ,  $R_a = a$  and  $R_b = b$ , we have  $\mathbb{F}(a, b) = \mathbb{F}$ . We also regard  $\mathbb{F}(R_a, R_b)$  still a sequence over alphabet  $\mathcal{A}$  if no confusion happens. In this case we use  $\mathbb{F}(R_a, R_b)[i]$  as the  $i$ -th letter in the sequence. For instance, let  $\mathcal{A} = \{A, B, C, D, E\}$ ,  $R_a = ABC$  and  $R_b = DE$ , then  $\mathbb{F}(R_a, R_b) = R_a R_b R_a R_a R_b \cdots = ABC \cdot DE \cdot ABC \cdot ABC \cdot DE \cdots$ . And  $\mathbb{F}(R_a, R_b)[5] = E$ .

By Zeckendorf numeration system [22], every positive integer  $n$  can be written uniquely as  $n = \sum_{j \geq 0} a_j f_j$  with  $a_j \in \{0, 1\}$  and  $a_{j+1} \times a_j = 0$  for  $j \geq 0$ . If  $a_k = 1$  and  $a_j = 0$  for all  $j > k$ , we call  $\text{Fib}(n) = a_k a_{k-1} \cdots a_1 a_0$  the canonical Fibonacci representation of  $n$ . Let  $a_m a_{m-1} \cdots a_1 a_0$  be a word over alphabet  $\{0, 1\}$ , by [5], the word can be written as a number  $[a_m a_{m-1} \cdots a_1 a_0]_F = \sum_{j \geq 0} a_j f_j$ , even if  $a_m = 0$  or  $a_{j+1} \times a_j \neq 0$  for some  $j \geq 0$ .

Let  $u = x_1 x_2 \cdots x_N$  be a finite word over alphabet  $\mathbb{Z}$  (the set of integers). We denote  $\Sigma u = x_1 + x_2 + \cdots + x_N$ . If an infinite word  $\rho$  satisfies that  $\rho[i] = 0$  for all  $i \geq 1$ , we call it a zero sequence. Moreover,  $\underbrace{[n \rightarrow]}_N := \underbrace{[n, n, \dots, n]}_N$ ,  $\underbrace{[n \nearrow]}_N := \underbrace{[n, n+1, \dots, n+N-1]}_N$  and  $\underbrace{[n \searrow]}_N := \underbrace{[n, n-1, \dots, n-N+1]}_N$ .

<sup>1</sup>School of Science, Beijing University of Posts and Telecommunications, Beijing, 100876, P. R. China. E-mail address: hyk@bupt.edu.cn, hyg03ster@163.com (Corresponding author).

<sup>2</sup>Department of Mathematical Sciences, Tsinghua University, Beijing, 100084, P. R. China. E-mail address: wenyzy@tsinghua.edu.cn.

The *fractional power* is a topic dealing with repetitions in words. We say a (finite or infinite) word  $\omega$  contains a  $r$ -power (real  $r > 1$ ) if  $\omega$  has a factor of the form  $x^{\lfloor r \rfloor} x'$  where  $x'$  is a prefix of  $x$  and  $|x^{\lfloor r \rfloor} x'| \geq r|x|$  see [1]. In this case, we call  $x^{\lfloor r \rfloor} x'$  a  $r$ -power with *size*  $|x|$ . For instance, taking  $x = ab$ , then  $\mathbb{F}[4, 8] = ababa$  is a  $\frac{5}{2}$ -power of size  $|ab| = 2$  in  $\mathbb{F}$ . Obviously the notion  $r$ -power is a generalization of square (2-power) and cube (3-power).

The study of power of a word has a long history. There are many significant contributions, for example [4, 6, 8, 9, 10, 15, 18, 19, 20, 21]. In particular, Iliopoulos-Moore-Smyth [15] computed the positions of all squares in  $\mathbb{F}$ . Fraenkel-Simpson [8, 9] obtained the number of squares in  $F_m = \sigma^m(a)$ . Du-Mousavi-Schaeffer-Shallit [6] obtained the numbers of repeated squares and cubes in  $\mathbb{F}[1, n]$  for all  $n \geq 1$ . All their numerations start from the first letter of the sequence.

In this paper, we count the number of distinct  $r$ -powers in  $\mathbb{F}[i; n]$  for all  $i, n \geq 1$  and  $r \geq 2$ , denoted by  $D(r, i, n)$ . Our numeration can start from any letter of the sequence, comparing starting from the first letter, there are some difficulties. To overcome these difficulties, we develop some new techniques by using the structure of return word sequences introduced in [12, 13, 14]. To get the general numeration of distinct  $r$ -powers, using again the structure of return word sequences, we establish a simple correspondent relation between the positions of squares and  $r$ -powers.

Our method is based on *factor spectrum*, *return word sequences* and *kernel words*, established in Huang-Wen [12, 13, 14]. This method can be used to study some other topics in combinatorics on words. For instance, using similar method, we can count the number of palindromes in  $\rho[i; n]$ . Here  $\rho$  is a substitution sequence and  $i, n \geq 1$ .

We give our main idea and some simple examples in Section 2. Then we list main results in Section 3. Sections 4, 5 and 7 will be devoted to the details of the proofs. We give some properties of function  $D(2, i, n)$  in Section 6.

## 2 Main idea and some simple examples

We first recall *factor spectrum* introduced in Huang-Wen [12, 13].

Let  $\rho$  be an infinite word and  $\Omega$  be the set of all factors in  $\rho$ . A factor  $\omega \in \Omega$  may occur many times (even infinity many times) at different positions in  $\rho$ . We note  $(\omega, j)$  the factor  $\omega$  which occurs at the position  $j$ . As we have mentioned before, any factor of the Fibonacci word will appear infinity many times. Let  $\mathcal{P}$  be a property. Our aim is to find out all factors which are located at different positions in  $\rho$  satisfying the property.

**Definition 2.1** (Factor Spectrum). *Let  $\mathcal{P}$  be a property, the factor spectrum with respect to  $\mathcal{P}$  is defined by*

$$\mathcal{S}_{\mathcal{P}} = \{(\omega, j) \mid \rho[j; |w|] = \omega \text{ and } (\omega, j) \text{ satisfies the property } \mathcal{P}\}.$$

*Notice that  $\mathcal{S}_{\mathcal{P}}$  is a subset of  $\Omega \times \mathbb{Z}^+$ . And it maybe a finite or infinite set according to different property  $\mathcal{P}$ .*

We notice that factor spectrum with respect to  $\mathcal{P}$  can be regarded a function of two variables, factor variable  $\omega$  and position variable  $j$ . Factor spectrum plays an important role in the paper. By factor spectrum, we get not only all positions of some fixed squares, but also all positions starting a square. Let us start from a simple question.

**Q1. How many repeated factor  $aa$  in  $\mathbb{F}[i; n]$  for  $i, n \geq 1$ ?**

Let us consider factor spectrum  $\mathcal{S}_{\mathcal{P}_1} = \{(aa, j) \mid \mathbb{F}[j; 2] = aa\}$ . It shows all positions of factor  $aa$  in  $\mathbb{F}$ . A powerful tool to determine  $\mathcal{S}_{\mathcal{P}_1}$  is the structure of the *return word sequence* which is introduced and studied in Huang-Wen [12, 13], see Property 2.2 and Figure 1 below. The definition of *return word* is from Durand [3]. Let  $\omega$  be a factor of  $\mathbb{F}$ . The  $p$ -th *return word* of  $\omega$  is  $R_p(\omega) := \mathbb{F}[(\omega)_p, (\omega)_{p+1} - 1]$  for  $p \geq 1$ . The sequence  $\{R_p(\omega)\}_{p \geq 1}$  is called the return word sequence of the factor  $\omega$ .

**Property 2.2** (Theorem 2.11 in [12]). *For each factor  $\omega \prec \mathbb{F}$ , the return word sequence  $\{R_p(\omega)\}_{p \geq 1}$  is itself still a Fibonacci word over the alphabet  $\{R_1(\omega), R_2(\omega)\}$ .*

$$\mathbb{F}[1, 31] = \text{a b } \underbrace{\text{a a b a b}}_{R_0 \text{ A}} \underbrace{\text{a a b}}_{\text{B}} \underbrace{\text{a a b a b}}_{\text{A}} \underbrace{\text{a a b a b}}_{\text{A}} \underbrace{\text{a a b a b}}_{\text{B}} \underbrace{\text{a a b a b a a b}}_{\text{A}}$$

Figure 1: The return word sequence  $\{R_p(aa)\}_{p \geq 1}$  is a Fibonacci word over the alphabet  $\{A, B\} := \{R_1(aa), R_2(aa)\} = \{aabab, aab\}$  with prefix  $R_0 = ab$ .

Obviously, the return word sequence gives a decomposition of  $\mathbb{F}$ . By Property 2.2, we have

$$(aa)_p = |R_0| + |\mathbb{F}[1, p-1]_a| \times |R_1(aa)| + |\mathbb{F}[1, p-1]_b| \times |R_2(aa)| + 1.$$

Thus  $\{(aa)_p \mid p \geq 1\} = \{3, 8, 11, 16, 21, 24, \dots\}$ . Define a sequence  $P_{aa} = \{P_{aa}[j]\}_{j \geq 1}$  as below, called the *position sequence* of factor  $aa$ .

$$P_{aa}[j] = \begin{cases} 1, & \text{if } j \in \{(aa)_p \mid p \geq 1\}, \\ 0, & \text{otherwise.} \end{cases}$$

Then  $P_{aa} = 001000010\dots$ , see the second line in Figure 2. Moreover the number of repeated  $aa$  in  $\mathbb{F}[i; n]$  is  $\Sigma P_{aa}[i; n - |aa| + 1]$ .

**Remark 1.** For any factor  $\omega \prec \mathbb{F}$ ,  $\omega \prec \mathbb{F}[i; n]$  is equivalent to  $\Sigma P_\omega[i; n - |\omega| + 1] \geq 1$ . Now we consider that how many distinct squares in  $\mathbb{F}[i; n]$  for  $i, n \geq 1$ ? A direct method is that: we find out all squares in  $\mathbb{F}$  denoted by set  $\mathcal{S}$ , and count  $\sum_{\omega \prec \mathcal{S}} \min\{1, \Sigma P_\omega[i; n - |\omega| + 1]\}$ . But unfortunately, the set  $\mathcal{S}$  is infinity. The counting will be complicated and difficult to find an explicit expression.

**Remark 2.** A known result is that all squares in  $\mathbb{F}$  are of length  $2f_m$  for  $m \geq 0$ , see [21] for instance. Thus in order to count the number of squares in  $\mathbb{F}[i; n]$ , we only need to count the number of squares of length  $2f_m$  in  $\mathbb{F}[i; n]$  for each  $m \geq 0$  first.

## Q2. How many squares of length $2f_m$ in $\mathbb{F}[i; n]$ for $i, n \geq 1$ ?

We consider factor spectrum  $\mathcal{S}_{P_2} = \{(\omega, j) \mid \mathbb{F}[j; 2f_m] = \omega\omega\}$ , i.e., the set of all squares of length  $2f_m$  with their positions in  $\mathbb{F}$ . By Wen-Wen [21], the number of distinct squares of length  $2f_m$  in  $\mathbb{F}$  is exact  $f_m$  for  $m \geq 0$ . We will use a tool called *kernel word* introduced and studied in [12, 13, 14] (we call them singular word in [21]). By this way, we can divided all squares of length  $2f_m$  into two types (for details see Subsection 4.1), and then get the counting expressions in Section 5.

The  $m$ -th kernel word  $K_m$  is defined by  $K_m = F_{m+1}[f_{m+1}]F_m[1, f_m - 1]$  for  $m \geq -1$ , by convention,  $K_{-2} = \varepsilon$ . Let  $\text{Ker}(\omega)$  be the maximal kernel word occurring in the factor  $\omega$ , called the *kernel* of  $\omega$ . By Theorem 1.9 in [12],  $\text{Ker}(\omega)$  occurs in  $\omega$  uniquely and only once. In Subsection 4.1, we prove that  $\{\text{Ker}(\omega^2) \mid |\omega| = f_m, \omega^2 \prec \mathbb{F}\} = \{K_{m+1}, K_{m-1}\}$  for  $m \geq 2$ , that is, only two kernel words. Thus all squares of length  $2f_m$  can be divided into two types according to their kernels. By Properties 2.3 and 2.4 below, we can determine the positions of squares in each type by the positions of their kernels.

**Property 2.3** (Property 4.1 in [14]). *For  $m \geq -1$ ,  $p \geq 1$  and  $\phi = \frac{\sqrt{5}-1}{2}$ , the position of the  $p$ -th occurrence of  $K_m$ , denoted by  $(K_m)_p$ , is equal to  $(K_m)_p = pf_{m+1} + \lfloor p\phi \rfloor f_m$ . In particular,  $a_p = p + \lfloor p\phi \rfloor$ ,  $b_p = 2p + \lfloor p\phi \rfloor$  for  $p \geq 1$ , and  $(K_m)_1 = f_{m+1}$  for  $m \geq -1$ .*

**Property 2.4** (Theorem 2.8 in [12]). *Let  $\omega \prec \mathbb{F}$  be a factor. For all  $p \geq 1$ , the difference*

$$(\text{Ker}(\omega))_p - (\omega)_p = (\text{Ker}(\omega))_1 - (\omega)_1$$

*is independent of  $p$ .*

Take  $m = 2$  for instance. There are three different squares of length  $2f_2 = 6$  in  $\mathbb{F}$ :  $abaaba = (aba)^2$ ,  $baabaa = (baa)^2$  and  $aabaab = (aab)^2$ . We define  $\text{code}(aba) = 1$ ,  $\text{code}(baa) = 2$  and  $\text{code}(aab) = 3$ . Define the position sequence  $\mathcal{D} = \{\mathcal{D}[j]\}_{j \geq 1}$  as below.

$$\mathcal{D}[j] = \begin{cases} \text{code}(\mathbb{F}[j; 3]), & \text{if } \mathbb{F}[j; 3] = \mathbb{F}[j+3; 3], \\ 0, & \text{if } \mathbb{F}[j; 3] \neq \mathbb{F}[j+3; 3]. \end{cases}$$

Obviously, it shows the positions of all squares of length 6, see the 4-th line in Figure 2. Thus the number of distinct squares of length 6 in  $\mathbb{F}[i; n]$  is equal to the number of distinct letters in  $\mathcal{D}[i; n - 5]$ . It is equal to  $\#\mathcal{D}[i; n - 5]$ .

Now we have two position sequences:  $P_\omega$  and  $\mathcal{D}$ .  $P_\omega$  can be determined by the property of return word sequence. Using the method below, we can determine  $\mathcal{D}$  by  $P_\omega$ . By the definition of kernel,  $\text{Ker}(abaaba) = aa$  and  $\text{Ker}(baabaa) = \text{Ker}(aabaab) = aabaa$ . According to this, we split  $\mathcal{D}$  into  ${}^1\mathcal{D} + {}^2\mathcal{D}$ , see lines 4 to 6 in Figure 2. Here  ${}^1\mathcal{D}$  and  ${}^2\mathcal{D}$  are the position sequences of  $abaaba$  and  $\{baabaa, aabaab\}$  respectively. By Property 2.4, the difference  $(aa)_p - (abaaba)_p = 2$ . So the relation between  ${}^1\mathcal{D}$  and  $P_{aa}$  is  ${}^1\mathcal{D}[j] = P_{aa}[j + 2] \times \text{code}(aba)$ , see lines 2 and 5 in Figure 2. Similarly,  ${}^2\mathcal{D}[j] = P_{aabaa}[j + 1] \times \text{code}(baa) + P_{aabaa}[j] \times \text{code}(aab)$ , see lines 3 and 6 in Figure 2.

$$\begin{array}{l}
\mathbb{F}[1, 31] = a \ b \ \overline{a \ a} \ b \ a \ b \ \overline{a \ a} \ b \ \overline{a \ a} \ b \ a \ b \ \overline{a \ a} \ b \ a \ b \ \overline{a \ a} \ b \ \overline{a \ a} \ b \ a \ b \ a \ a \ b \\
P_{aa}[1, 26] = \left| \begin{array}{c|c|c|c|c|c} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \end{array} \right| \begin{array}{l} \leftarrow \text{The position sequence of } aa \\ \leftarrow \text{The position sequence of } aabaa \end{array} \\
P_{aabaa}[1, 26] = \left| \begin{array}{c|c|c|c|c|c} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right| \begin{array}{l} \leftarrow \text{The position sequence of } aabaa \\ \leftarrow \text{The position sequence of } aabaa \end{array} \\
\mathcal{D}[1, 26] = \left| \begin{array}{c|c|c|c|c|c} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 3 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{array} \right| \begin{array}{l} \leftarrow \text{Ker}(abaaba) = aa \\ \leftarrow \left\{ \begin{array}{l} \text{Ker}(baabaa) = aabaa \\ \text{Ker}(aabaab) = aabaa \end{array} \right. \end{array} \\
\begin{array}{l} \parallel \\ {}^1\mathcal{D}[1, 26] = \\ + \\ {}^2\mathcal{D}[1, 26] = \end{array} \left| \begin{array}{c|c|c|c|c|c} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{array} \right| \begin{array}{l} \leftarrow \text{Ker}(abaaba) = aa \\ \leftarrow \left\{ \begin{array}{l} \text{Ker}(baabaa) = aabaa \\ \text{Ker}(aabaab) = aabaa \end{array} \right. \end{array}
\end{array}$$

Figure 2: The relations among  $\mathbb{F}$  and the position sequences  $P_{aa}$ ,  $P_{aabaa}$ ,  $\mathcal{D}$ .

**Q3. How many  $r$ -powers of size  $f_m$  in  $\mathbb{F}[i; n]$  for  $i, n \geq 1$  and  $r \geq 2$ ?**

Taking  $m = 2$  for example, the key step is to determine the position sequence of  $r$ -powers of size  $f_m = 3$ , denoted by  $\mathcal{D}^r$ , where  $r = 2 + \frac{h}{3}$ ,  $h = 1, 2, \dots$ . Then the number of distinct  $r$ -powers of size  $f_2$  in  $\mathbb{F}[i; n]$  is  $\#\mathcal{D}^r[i; n - rf_2 + 1]$ .

Using the method below, we can determine  $\mathcal{D}^r$  by  $\mathcal{D}$  (the position sequence of squares of size 3). Notice that, if  $\mathcal{D}[j] \times \mathcal{D}[j + 1] \neq 0$ , both  $\mathbb{F}[j; 6]$  and  $\mathbb{F}[j + 1; 6]$  are squares. So  $\mathbb{F}[j; 7]$  is a  $(2 + \frac{1}{3})$ -power. This means  $\mathcal{D}^{2+\frac{1}{3}}[j] \neq 0$ . Similarly, if  $\mathcal{D}[j] \times \mathcal{D}[j + 1] = 0$ , then  $\mathcal{D}^{2+\frac{1}{3}}[j] = 0$ , see the first two lines in Figure 3. By an analogous argument, we can determine sequence  $\mathcal{D}^{2+\frac{h+1}{3}}$  by  $\mathcal{D}^{2+\frac{h}{3}}$  for  $h = 1, 2, \dots$ , see lines 2 to 4 in Figure 3.

$$\begin{array}{l}
\mathcal{D}[1, 26] = 1 \ 0 \ 0 \ 0 \ 0 \ 1 \ 2 \ 3 \ 1 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 1 \ 2 \ 3 \ 1 \ 0 \ 0 \ 0 \ 0 \\
\mathcal{D}^r[1, 26] = 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 2 \ 3 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 2 \ 3 \ 0 \ 0 \ 0 \ 0 \ 0 \ \text{for } r = 2 + \frac{1}{3} \\
\mathcal{D}^r[1, 26] = 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 2 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 2 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ \text{for } r = 2 + \frac{2}{3} \\
\mathcal{D}^r[1, 26] = 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 1 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ \text{for } r = 3
\end{array}$$

Figure 3: The relations among sequences  $\mathcal{D}$  and  $\mathcal{D}^r$  for  $r = 2 + \frac{h}{3}$ ,  $h = 1, 2, 3$ .

**Q4. The number of positions starting a square.**

Harju-Kärki-Nowotka [11] considered the number of positions that do not start a square in a binary word. As an application of factor spectrum, we prove that all positions in the Fibonacci word start a square of length  $2f_m$  for some  $0 \leq m \leq 3$ , see Property 5.4. Moreover, all positions in the Fibonacci word start infinite distinct squares, see Property 5.5.

### 3 Main results

In this paper, we count the number of distinct  $r$ -powers in  $\mathbb{F}[i; n]$  for all  $i, n \geq 1$  and  $r \geq 2$ , denoted by  $D(r, i, n)$ . We give precise results for  $r \in \{2, 2 + \epsilon, 3\}$  in this paper, where  $\epsilon$  is a small positive number. In this section, we show the results and some examples for  $r = 2$ . The results for  $r \in \{2 + \epsilon, 3\}$  are given in Section 7.

**Theorem 3.1** ( $D(2, 1, n)$ ). For  $n \geq 3$ , let  $h \geq 0$  such that  $2f_h \leq n < 2f_{h+1}$ . Then the number of distinct squares in  $\mathbb{F}[1, n]$  is equal to  $D(2, 1, n) = \min\{n - f_{h-1} - 2, f_{h+1} + f_{h-1} - 3\}$ .

**Remark.** Let  $n = f_{h+2}$  for  $h \geq 0$ . Since  $2f_h \leq n < 2f_{h+1}$  and  $n - f_{h-1} - 2 \leq f_{h+1} + f_{h-1} - 3$ ,  $D(2, 1, n) = n - f_{h-1} - 2 = 2f_h - 2$ . We obtain thus the result of Fraenkel-Simpson [8].

**Theorem 3.2** ( $D(2, i, n)$ ). For  $i \geq 1$  and  $n \geq 2$ , let  $h \geq 0$  such that  $2f_h \leq n < 2f_{h+1}$ . Then the number of distinct squares in  $\mathbb{F}[i; n]$  is equal to

$$D(2, i, n) = \begin{cases} [0, n - 2, 1, 0, 0][\hat{i}], & n = 2, 3, \\ [1, 1, n - 3, n - 3, n - 3, 1, 1, 1][\hat{i}], & n = 4, 5. \\ [R_a, R_b][\hat{i}] + f_{h-1} - 2, & n \geq 6 \text{ and } n \in \text{I} \cup \text{II} \cup \text{III} \\ [R_a, R_b][\hat{j}] + f_{h-1} - 2, & \text{otherwise.} \end{cases} \quad (1)$$

Here  $\text{Fib}(i - 1) = a_k a_{k-1} \dots a_1 a_0$ ,  $\hat{i} = [a_{h+1} a_h \dots a_1 a_0]_F + 1$  and  $\hat{j} = [a_{h+2} a_{h+1} \dots a_1 a_0]_F + 1$ . The definitions of Ranges I to V see Figure 4(A). The expressions of  $R_a$  and  $R_b$  see Figure 4(B).

(A) Divide  $2f_h \leq n < 2f_{h+1}$  into five ranges.

Range I:  $2f_h \leq n < 2f_h + f_{h-3} - 1$ ;

Range II:  $2f_h + f_{h-3} - 1 \leq n < 2f_h + f_{h-2} - 1$ ;

Range III:  $2f_h + f_{h-2} - 1 \leq n < f_{h+1} + 2f_{h-1}$ ;

Range IV:  $f_{h+1} + 2f_{h-1} \leq n < 3f_h$ ;

Range V:  $3f_h \leq n < 2f_{h+1}$ .

(B) The expressions of  $R_a$  and  $R_b$  in Theorem 3.2.

$$\begin{aligned} n \in \text{I}: \quad R_a &= [ \underbrace{n - 2f_{h-1} \rightarrow}_{f_{h+1}+f_{h-1}-n-1}, \underbrace{n - 2f_{h-1} \searrow}_{f_h \rightarrow}, \underbrace{2n - f_{h+1} - 3f_{h-1} + 1 \rightarrow}_{f_h - 1 \searrow}, \underbrace{2n - f_{h+1} - 3f_{h-1} + 2 \nearrow}_{f_h - f_{h-4} - 1 \rightarrow}, \underbrace{n - 2f_{h-1} \rightarrow}_{f_h - f_{h-4} \nearrow}, \underbrace{n - 2f_{h-1} + 1 \nearrow}_{f_h \rightarrow}, \underbrace{n - 2f_{h-1} \rightarrow}_{f_h - 1 \searrow}, \underbrace{n - 2f_{h-1} + 1 \nearrow}_{f_{h+1}+f_{h-1}-n-1} ]; \\ R_b &= [ \underbrace{n - 2f_{h-1} \rightarrow}_{f_h - 1 \searrow}, \underbrace{2n - f_{h+1} - 3f_{h-1} + 1 \rightarrow}_{f_{h+1}+f_{h-1}-n-1}, \underbrace{2n - f_{h+1} - 3f_{h-1} + 2 \nearrow}_{f_h - 4}, \underbrace{n - 2f_{h-1} \rightarrow}_{n - 2f_{h+1}}, \underbrace{n - 2f_{h-1} + 1 \nearrow}_{f_{h+1}+f_{h-1}-n-1} ]; \\ n \in \text{II}: \quad R_a &= [ \underbrace{n - 2f_{h-1} \rightarrow}_{f_{h+1}+f_{h-1}-n-1}, \underbrace{n - 2f_{h-1} \searrow}_{n - 3f_{h-1} + 1}, \underbrace{f_h \rightarrow}_{f_h - 1 \searrow}, \underbrace{f_h - 1 \searrow}_{f_{h+1}+f_{h-1}-n-2}, \underbrace{n - f_h - f_{h-2} + 1 \rightarrow}_{n - 2f_{h+1} + 2} ]; \\ R_b &= [ \underbrace{n - 2f_{h-1} \rightarrow}_{f_{h+2}+f_{h-3}-n}, \underbrace{n - f_{h+1} - f_{h-1} + 1 \rightarrow}_{n - f_h - f_{h-2} + 2}, \underbrace{f_h \rightarrow}_{f_h \nearrow}, \underbrace{f_h - 3 \nearrow}_{f_{h+1}+2f_{h-2}-n-2}, \underbrace{n - f_{h+1} - f_{h-1} + 3 \rightarrow}_{n - f_{h+1} - f_{h-1} + 3} ]; \\ n \in \text{III}: \quad R_a &= [ \underbrace{n - 2f_{h-1} \rightarrow}_{f_{h+1}+2f_{h-2}-n-2}, \underbrace{n - 2f_{h-1} \searrow}_{f_{h-3}}, \underbrace{f_h \rightarrow}_{n - f_{h+1} - f_{h-1} + 1}, \underbrace{f_h \nearrow}_{f_{h-3}}, \underbrace{n - 2f_{h-1} \rightarrow}_{f_h} ]; \\ R_b &= [ \underbrace{n - 2f_{h-1} \rightarrow}_{f_h} ]; \\ n \in \text{IV}: \quad R_a &= [ \underbrace{f_{h+1} - 1 \searrow}_{f_{h+2}+f_{h-3}-n-1}, \underbrace{f_h \rightarrow}_{n - f_{h+1} - f_{h-1} + 1}, \underbrace{f_h + 1 \nearrow}_{3f_h - n - 1}, \underbrace{n - 2f_{h-1} \rightarrow}_{n - f_{h+1} - f_{h-1} + 1}, \underbrace{n - 2f_{h-1} - 1 \searrow}_{f_h - 3} ]; \\ R_b &= [ \underbrace{f_{h+1} - 1 \searrow}_{f_{h-1}}, \underbrace{f_h \rightarrow}_{f_{h+2}+f_{h-2}-n}, \underbrace{f_h + 1 \nearrow}_{n - f_{h+2} + f_{h-2}}, \underbrace{f_{h+1} - 1 \rightarrow}_{f_{h+1}+4f_{h-1}-n}, \underbrace{n - 2f_{h+1} + f_{h-2} \rightarrow}_{n - 2f_{h+1} + f_{h-2}} ]; \\ n \in \text{V}: \quad R_a &= [ \underbrace{f_{h+1} - 1 \searrow}_{f_{h-1}}, \underbrace{n - 2f_h \rightarrow}_{f_{h+2}+f_{h-2}-n}, \underbrace{n - 2f_h + 1 \nearrow}_{f_{h-1}-2}, \underbrace{n - 2f_{h-1} \rightarrow}_{n - f_{h+2} + 2}, \underbrace{n - 2f_{h-1} - 1 \searrow}_{n - 2f_{h-1} - 1}, \underbrace{f_h + 2f_{h-2} - 1 \rightarrow}_{f_h + 2f_{h-2}}, \underbrace{f_h + 2f_{h-2} \nearrow}_{n - 3f_{h+1}} ]; \\ R_b &= [ \underbrace{f_{h+1} - 1 \searrow}_{f_{h+2}+f_h-n}, \underbrace{n - 2f_h \rightarrow}_{n - 3f_h}, \underbrace{n - 2f_h + 1 \nearrow}_{2f_{h-2}}, \underbrace{f_{h+1} - 1 \rightarrow}_{f_{h+2}+f_{h-1}-n}, \underbrace{n - 2f_h \rightarrow}_{n - 3f_h}, \underbrace{n - 2f_h + 1 \nearrow}_{f_{h+2}+f_h-n-1}, \underbrace{f_{h+1} - 1 \rightarrow}_{f_{h+1} - 1} ]; \end{aligned}$$

Figure 4: The expressions of  $R_a$  and  $R_b$  in Theorem 3.2.

**Example.** We consider the number of distinct squares in  $\mathbb{F}[333; 20]$ . Since  $n = 20$ , we have  $h = 4$ . By the definitions of Ranges I to V in Figure 4(A), we have  $n = 20 \in \text{Range III} = \{18, 19, \dots, 22\}$ . By the expressions of  $R_a$  and  $R_b$  in Figure 4(B), for  $n = 20$  and  $h = 4$

$$\begin{aligned} R_a &= [ \underbrace{n - 2f_{h-1} \rightarrow}_{f_{h+2}+f_{h-3}-n-1}, \underbrace{n - 2f_{h-1} \searrow}_{n - f_{h+1} - f_{h-1} + 1}, \underbrace{f_h \rightarrow}_{3f_h - n - 1}, \underbrace{f_h \nearrow}_{n - f_{h+1} - f_{h-1} + 1}, \underbrace{n - 2f_{h-1} \rightarrow}_{f_{h-3}} ]; \\ &= [ \underbrace{10 \rightarrow}_2, \underbrace{10 \searrow}_3, \underbrace{8 \rightarrow}_3, \underbrace{8 \nearrow}_3, \underbrace{10 \rightarrow}_2 ] = [10, 10, 10, 9, 8, 8, 8, 8, 8, 9, 10, 10, 10]; \\ R_b &= [ \underbrace{n - 2f_{h-1} \rightarrow}_{f_h} ] = [ \underbrace{10 \rightarrow}_8 ] = [10, 10, 10, 10, 10, 10, 10, 10]. \end{aligned}$$

Since  $i = 333$  and  $h = 4$ ,  $\text{Fib}[i - 1] = \text{Fib}[332] = 101000010010$ ,  $\hat{i} = [a_{h+1} a_h \dots a_1 a_0]_F + 1 = [10010]_F + 1 = 11$ . By Equation (1),  $D(2, 333, 20) = [R_a, R_b][\hat{i}] + f_{h-1} - 2 = 10 + 5 - 2 = 13$ .

**Remark.** A famous conjecture of Fraenkel-Simpson [7] states that a word of length  $n$  contains fewer than  $n$  distinct squares. By Figure 4(B), For fixed  $n \geq 2$  and  $2f_h \leq n < 2f_{h+1}$ , the maximal element in  $R_a$  and  $R_b$  is  $f_h$  for  $n \in \text{Range I}$ , and  $n - 2f_{h-1}$  otherwise. So

$$\max_{1 \leq i \leq \infty} D(2, i, n) = \begin{cases} f_h + f_{h-1} - 2 = f_{h+1} - 2, & n \in \text{Range I} \\ n - 2f_{h-1} + f_{h-1} - 2 = n - f_{h-1} - 2, & \text{otherwise} \end{cases}$$

Thus  $D(2, i, n) < n$  for all  $i, n \geq 1$ , and the conjecture holds for the Fibonacci word.

**Property 3.3.** *Any word of length  $n$  in  $\mathbb{F}$  contains fewer than  $n$  distinct square factors.*

In Section 6, we give some properties of function  $D(2, i, n)$ , such as: (1) for  $n$  large enough,

$$\begin{cases} 0.7236n \leq \max_{1 \leq i \leq \infty} D(2, i, n) \leq 0.8090n, & 0.5393n \leq \min_{1 \leq i \leq \infty} D(2, i, n) \leq 0.6806n, \\ 0.0528n \leq \max_{1 \leq i, j \leq \infty} |D(2, i, n) - D(2, j, n)| \leq 0.2547n; \end{cases}$$

and (2) for fixed  $n$  and any  $M$  such that  $\max_{1 \leq i \leq \infty} D(2, i, n) \leq M \leq \min_{1 \leq i \leq \infty} D(2, i, n)$ , there exists  $i_0$  such that  $D(2, i_0, n) = M$ .

## 4 The position sequences of squares

In Section 2 (Q2), we claim that we can divided all squares of length  $2f_m$  into two types by their kernels. We give the details in subsection below.

### 4.1 Two cases of squares

By Theorem 1.9 in [12] and Property 2.2,

$$\{R_p(K_m), p \geq 1\} = \{R_1(K_m), R_2(K_m)\} = \{K_m K_{m+1}, K_m K_{m-1}\}.$$

Moreover, the return word sequence  $\{R_p(K_m)\}_{p \geq 1}$  is still a Fibonacci word over  $\{R_1(K_m), R_2(K_m)\}$  and  $bb \not\prec \mathbb{F}$ , so  $R_2(K_m)R_2(K_m) \not\prec \mathbb{F}$ . Thus

$$\begin{aligned} & \{|R_p(K_m)R_{p+1}(K_m)|, p \geq 1\} \\ & = \{|R_1(K_m)R_1(K_m)|, |R_1(K_m)R_2(K_m)|, |R_2(K_m)R_1(K_m)|\} = \{2f_{m+2}, f_{m+3}\}. \end{aligned}$$

On the other hand, by Definition 2.9 and Corollary 2.10 in [12], any factor  $\omega$  with kernel  $K_m$  can be expressed uniquely as  $\omega = K_{m+1}[i, f_{m+1}]K_m K_{m+1}[1, j]$ , where  $2 \leq i \leq f_{m+1} + 1$  and  $0 \leq j \leq f_{m+1} - 1$ . So  $\omega\omega \prec \mathbb{F}$  means

$$K_{m+1}[i, f_{m+1}] \underbrace{K_m K_{m+1}[1, j] K_{m+1}[i, f_{m+1}]}_{\text{denoted by } R(K_m)} K_m K_{m+1}[1, j] \prec \mathbb{F}.$$

By the ranges of  $i$  and  $j$ ,  $|\omega| = |R(K_m)| = f_{m+2} + j - i + 1 \leq f_{m+3} - 2$ .

So  $|R(K_m)| \leq \min\{2f_{m+2}, f_{m+3}\}$ . This means  $K_m$  occurs in  $R(K_m)$  only once. Thus  $R(K_m) \in \{R_p(K_m), p \geq 1\} = \{K_m K_{m+1}, K_m K_{m-1}\}$ . Two cases have to be considered.

**Case 1.**  $R(K_m) = R_1(K_m) = K_m K_{m+1}$ . In this case  $|R(K_m)| = |R_1(K_m)|$  implies  $j = i - 1$ . So  $0 \leq j \leq f_{m+1} - 1$  gives a range of  $i$ . Comparing this range with  $2 \leq i \leq f_{m+1} + 1$ , we have  $2 \leq i \leq f_{m+1}$  and  $m \geq 0$ . Moreover  $|\omega| = |R(K_m)| = f_{m+2}$  and

$$\begin{aligned} \omega^2 &= K_{m+1}[i, f_{m+1}]K_m K_{m+1} K_m K_{m+1}[1, i - 1] \\ &= K_{m+2}[i, f_{m+2}] \underline{K_{m+1}} K_{m+2}[1, f_m + i - 1] = K_{m+4}[i, 2f_{m+2} + i - 1]. \end{aligned}$$

The second and third equalities hold by Lemma 2.2 in [12], i.e.,

$$K_{m+2} = F_{m+1}[f_{m+1}]K_{m+1}[2, f_{m+1}]K_m = K_m K_{m+1}[1, f_{m+1} - 1]F_{m+1}[f_{m+1}]$$

and  $K_{m+4} = K_{m+2}K_{m+1}K_{m+2}$ . Thus  $K_{m+1} \prec \omega\omega \prec K_{m+4}[2, f_{m+4} - 1]$ . By the cylinder structure of palindromes [14], both  $K_{m+2}$  and  $K_{m+3}$  are not the factors of  $\omega\omega$ . Thus  $K_{m+1}$  is the maximal kernel word in  $\omega\omega$ , i.e.  $\text{Ker}(\omega\omega) = K_{m+1}$  for  $m \geq 0$ .

**Case 2.**  $R(K_m) = R_2(K_m) = K_m K_{m-1}$ .  $|R(K_m)| = |R_2(K_m)|$  implies  $j = i - f_m - 1$ . Consequently,  $f_m + 1 \leq i \leq f_{m+1} + 1$  and  $m \geq -1$ . Moreover  $|\omega| = |R(K_m)| = f_{m+1}$  and

$$\begin{aligned} \omega^2 &= K_{m+1}[i, f_{m+1}]K_m K_{m-1} K_m K_{m+1}[1, i - f_m - 1] \\ &= K_{m+3}[f_{m+2} + i, f_{m+3}] \underline{K_{m+2}} K_{m+3}[1, i - f_m - 1] = K_{m+5}[f_{m+2} + i, f_{m+3} + f_{m+1} + i - 1]. \end{aligned}$$

Thus  $K_{m+2} \prec \omega\omega \prec K_{m+5}[2, f_{m+5} - 1]$ . By an analogous argument  $\text{Ker}(\omega\omega) = K_{m+2}$ .

Thus we get the expressions of all squares in the Fibonacci word.

**Definition 4.1.** For all  $m, p \geq 1$ ,

$$\begin{cases} K_{m,p}^1 = \{(\omega^2)_p \mid \text{Ker}(\omega^2) = K_m, |\omega| = f_{m+1}, \omega^2 \prec \mathbb{F}\}, \\ K_{m,p}^2 = \{(\omega^2)_p \mid \text{Ker}(\omega^2) = K_m, |\omega| = f_{m-1}, \omega^2 \prec \mathbb{F}\}. \end{cases} \quad (2)$$

Each set contains several consecutive positive integers.

By Case 1 above,  $K_{m,p}^1 = \{(\omega^2)_p \mid \omega^2 = K_{m+1}[i, f_{m+1}]K_m K_{m+1}[1, f_{m-1} + i - 1], 2 \leq i \leq f_m\}$  for  $m \geq 1$ . For each fixed  $i$ ,  $\text{Ker}(\omega^2) = K_m$ . By Property 2.4,

$$(K_m)_p - (\omega^2)_p = (K_m)_1 - (\omega^2)_1 = |K_{m+1}[i, f_{m+1}]| = f_{m+1} - i + 1.$$

So  $K_{m,p}^1 = \{(K_m)_p - f_{m+1} + i - 1 \mid 2 \leq i \leq f_m\}$  for  $m \geq 1$ . Similarly, for  $m \geq 1$

$$\begin{aligned} K_{m,p}^2 &= \{(\omega^2)_p \mid \omega^2 = K_{m+1}[f_m + i, f_{m+1}]K_m K_{m+1}[1, i - f_{m-2} - 1], f_{m-2} + 1 \leq i \leq f_{m-1} + 1\} \\ &= \{(K_m)_p - f_{m-1} + i - 1 \mid f_{m-2} + 1 \leq i \leq f_{m-1} + 1\}. \end{aligned}$$

Thus for  $m, p \geq 1$

$$\begin{cases} K_{m,p}^1 = \{(K_m)_p - f_{m+1} + 1, \dots, (K_m)_p - f_{m-1} - 1\}, & \#K_{m,p}^1 = f_m - 1, \\ K_{m,p}^2 = \{(K_m)_p - f_{m-3}, \dots, (K_m)_p\}, & \#K_{m,p}^2 = f_{m-3} + 1. \end{cases} \quad (3)$$

By Property 2.3,  $(K_m)_1 = f_{m+1}$ . Thus for  $m \geq 1$

$$K_{m,1}^1 = \{1, \dots, f_m - 1\} \text{ and } K_{m,1}^2 = \{f_m + f_{m-2}, \dots, f_{m+1}\}.$$

For instance,  $(K_1)_1 = 3$ ,  $(K_1)_2 = 8$ ,  $(K_2)_1 = 5$  and  $(K_2)_2 = 13$ . Thus the first few values of  $K_{m,p}^1$  and  $K_{m,p}^2$  are:  $K_{1,1}^1 = \{1\}$ ,  $K_{1,2}^1 = \{6\}$ ,  $K_{2,1}^1 = \{1, 2\}$ ,  $K_{2,2}^1 = \{9, 10\}$ ,  $K_{1,1}^2 = \{3\}$ ,  $K_{1,2}^2 = \{8\}$ ,  $K_{2,1}^2 = \{4, 5\}$ ,  $K_{2,2}^2 = \{12, 13\}$ .

## 4.2 The position sequences of squares

Now we consider all squares of length  $2f_m$ . For  $m \geq 2$ , there are two types of distinct squares of length  $2f_m$ , whose positions are in sets  $K_{m-1,p}^1$  and  $K_{m+1,p}^2$ , respectively.

1. Recall that  $K_{m-1,1}^1 = \{1, 2, \dots, f_{m-1} - 1\}$  for  $m \geq 2$ . We denote square  $\mathbb{F}[j; 2f_m]$  by code  $j$  for  $1 \leq j \leq f_{m-1} - 1$ . We define sequence  ${}^1\mathcal{D}^{2,m}$  for  $i \geq 1$  and  $1 \leq j \leq f_{m-1} - 1$  that

$${}^1\mathcal{D}^{2,m}[i] = \begin{cases} j, & \text{if } \mathbb{F}[i; 2f_m] = \mathbb{F}[j; 2f_m], \\ 0, & \text{otherwise.} \end{cases}$$

We call it the position sequences of squares of length  $2f_m$  with kernel  $K_{m-1}$  for  $m \geq 2$ .

By Equation (3) and Property 2.3,  ${}^1\mathcal{D}^{2,m}$  is a Fibonacci word over the alphabet  $\{{}^1R_a^{2,m}, {}^1R_b^{2,m}\}$  with prefix  ${}^1R_0^{2,m}$  for  $m \geq 2$ , where

$$\left\{ \begin{array}{l} {}^1R_a^{2,m} = {}^1\mathcal{D}^{2,m}[\min K_{m-1,1}^1, \min K_{m-1,2}^1 - 1] = {}^1\mathcal{D}^{2,m}[1, f_{m+1}] \\ \quad = [1, 2, \dots, f_{m-1} - 1, \underbrace{0, \dots, 0}_{f_{m+1}}], \\ {}^1R_b^{2,m} = {}^1\mathcal{D}^{2,m}[\min K_{m-1,2}^1, \min K_{m-1,3}^1 - 1] = {}^1\mathcal{D}^{2,m}[f_{m+1} + 1, f_{m+2}] \\ \quad = {}^1\mathcal{D}^{2,m}[f_{m+1} + 1; f_m] = [1, 2, \dots, f_{m-1} - 1, \underbrace{0, \dots, 0}_{f_{m-2}+1}], \\ {}^1R_0^{2,m} = {}^1\mathcal{D}^{2,m}[1, \min K_{m-1,1}^1 - 1] = \varepsilon. \end{array} \right. \quad (4)$$

2. Recall that  $K_{m+1,1}^2 = \{f_{m+1} + f_{m-1}, \dots, f_{m+2}\}$  for  $m \geq 0$ . We denote square  $\mathbb{F}[f_{m+1} + j; 2f_m]$  by code  $j$  for  $f_{m-1} \leq j \leq f_m$ . We define sequence  ${}^2\mathcal{D}^{2,m}$  for  $i \geq 1$  and  $f_{m-1} \leq j \leq f_m$  that

$${}^2\mathcal{D}^{2,m}[i] = \begin{cases} j, & \text{if } \mathbb{F}[i; 2f_m] = \mathbb{F}[f_{m+1} + j; 2f_m], \\ 0, & \text{otherwise.} \end{cases}$$

We call it the position sequences of squares of length  $2f_m$  with kernel  $K_{m+1}$  for  $m \geq 0$ .

By Equation (3) and Property 2.3,  ${}^2\mathcal{D}^{2,m}$  is a Fibonacci word over the alphabet  $\{\tilde{R}_a^{2,m}, \tilde{R}_b^{2,m}\}$  with prefix  $\tilde{R}_0^{2,m}$  for  $m \geq 2$ , where

$$\left\{ \begin{array}{l} \tilde{R}_a^{2,m} = {}^2\mathcal{D}^{2,m}[\min K_{m+1,1}^2, \min K_{m+1,2}^2 - 1] = {}^2\mathcal{D}^{2,m}[f_{m+1} + f_{m-1}, 4f_{m+1} - 1] \\ \quad = {}^2\mathcal{D}^{2,m}[f_{m+1} + f_{m-1}; f_{m+3}] = [f_{m-1}, f_{m-1} + 1, \dots, f_m, \underbrace{0, \dots, 0}_{2f_{m+1}+f_{m-1}-1}], \\ \tilde{R}_b^{2,m} = {}^2\mathcal{D}^{2,m}[\min K_{m+1,2}^2, \min K_{m+1,3}^2 - 1] = {}^2\mathcal{D}^{2,m}[4f_{m+1}, f_{m+2} + 4f_{m+1} - 1] \\ \quad = {}^2\mathcal{D}^{2,m}[4f_{m+1}; f_{m+2}] = [f_{m-1}, f_{m-1} + 1, \dots, f_m, \underbrace{0, \dots, 0}_{f_{m+1}+f_{m-1}-1}], \\ \tilde{R}_0^{2,m} = {}^2\mathcal{D}^{2,m}[1, \min K_{m+1,1}^2 - 1] = {}^2\mathcal{D}^{2,m}[1, f_{m+1} + f_{m-1} - 1] = [ \underbrace{0, \dots, 0}_{f_{m+1}+f_{m-1}-1} ]. \end{array} \right.$$

Notice that both  $\tilde{R}_a^{2,m}$  and  $\tilde{R}_b^{2,m}$  have suffix  $\tilde{R}_0^{2,m}$ , we can shift to left each return word  $f_{m+1} + f_{m-1} - 1$  letters. Thus we get the equivalent expressions that  $\hat{R}_0^{2,m} = \varepsilon$  and

$$\left\{ \begin{array}{l} \hat{R}_a^{2,m} = [ \underbrace{0, \dots, 0}_{f_{m+1}+f_{m-1}-1}, f_{m-1}, f_{m-1} + 1, \dots, f_m, \underbrace{0, \dots, 0}_{f_{m+1}} ], \\ \hat{R}_b^{2,m} = [ \underbrace{0, \dots, 0}_{f_{m+1}+f_{m-1}-1}, f_{m-1}, f_{m-1} + 1, \dots, f_m ]. \end{array} \right.$$

Furthermore, the Fibonacci morphism  $\sigma$  over the alphabet  $\{{}^2R_a^{2,m}, {}^2R_b^{2,m}\}$  where

$${}^2R_a^{2,m} = \underbrace{[0, \dots, 0]}_{f_{m+1}} \text{ and } {}^2R_b^{2,m} = \underbrace{[0, \dots, 0]}_{f_{m-1}-1}, f_{m-1}, f_{m-1} + 1, \dots, f_m \quad (5)$$

satisfies that  $\sigma^2({}^2R_a^{2,m}) = \hat{R}_a^{2,m}$  and  $\sigma^2({}^2R_b^{2,m}) = \hat{R}_b^{2,m}$ . This means  ${}^2\mathcal{D}^{2,m} = \mathbb{F}({}^2R_a^{2,m}, {}^2R_b^{2,m})$ .

By the argument above, we give the definition of position sequences of squares.

**Definition 4.2.** We define sequence  $\mathcal{D}^{2,m}$  ( $m \geq 0$ ) that for  $i \geq 1$  and  $1 \leq j \leq f_m$

$$\mathcal{D}^{2,m}[i] = \begin{cases} j, & \text{if } \mathbb{F}[i; 2f_m] = \mathbb{F}[j; 2f_m] \text{ for } 1 \leq j \leq f_{m-1} - 1, \\ j, & \text{if } \mathbb{F}[i; 2f_m] = \mathbb{F}[f_{m+1} + j; 2f_m] \text{ for } f_{m-1} \leq j \leq f_m, \\ 0, & \text{otherwise.} \end{cases}$$

We call  $\mathcal{D}^{2,m}$  the position sequences of squares of length  $2f_m$  for  $m \geq 0$ .

When  $m \geq 2$ , if there exist an integer  $p$  such that  $i \in K_{m-1,p}^1$  (resp.  $i \in K_{m+1,p}^2$ ),  $\mathbb{F}[i; 2f_m]$  is a square with kernel  $K_{m-1}$  (resp.  $K_{m+1}$ ). Since the kernel of a factor is unique [12], sets  $K_{m-1,p}^1$  and  $K_{m+1,q}^2$  are disjoint for all  $p, q \geq 1$ . This means  ${}^1\mathcal{D}^{2,m}[i] \times {}^2\mathcal{D}^{2,m}[i] = 0$  for  $i \geq 1$ . Thus  $\mathcal{D}^{2,m} = {}^1\mathcal{D}^{2,m} + {}^2\mathcal{D}^{2,m}$ . By Equations (4) and (5),  $\mathcal{D}^{2,m} = \mathbb{F}(R_a^{2,m}, R_b^{2,m})$  where

$$\begin{cases} R_a^{2,m} = {}^1R_a^{2,m} + {}^2R_a^{2,m} = [1, 2, \dots, f_{m-1} - 1, \underbrace{0, \dots, 0}_{f_m+1}], \\ R_b^{2,m} = {}^1R_b^{2,m} + {}^2R_b^{2,m} = [1, 2, \dots, f_m]. \end{cases} \quad (6)$$

When  $m = 0, 1$ , there are  $\#K_{m+1,p}^2 = f_{m-2} + 1$  distinct squares of length  $2f_m$ . Thus sequence  $\mathcal{D}^{2,m}$  is exact sequence  ${}^2\mathcal{D}^{2,m}$ . But since all digits in  ${}^1R_a^{2,m}$  and  ${}^1R_b^{2,m}$  are zero for  $m = 0, 1$ , the expressions in Equation (6) still hold.

**Theorem 4.3.**  $\mathcal{D}^{2,m}$  is a Fibonacci word over  $\{R_a^{2,m}, R_b^{2,m}\}$  given in Equation (6) for  $m \geq 0$ .

## 5 The number of squares in $\mathbb{F}[i; n]$

By the definition of sequence  $\mathcal{D}^{2,m}$ ,  $\mathcal{D}^{2,m}[i] = j$  ( $1 \leq j \leq f_m$ ) means  $\mathbb{F}[i; 2f_m]$  is a square of length  $2f_m$  encoded by  $j$ ; and  $\mathcal{D}^{2,m}[i] = 0$  means  $\mathbb{F}[i; 2f_m]$  is not a square. Thus in order to count the number of distinct squares of length  $2f_m$  in  $\mathbb{F}[i; n]$ , we only need to consider the number of distinct letters in  $\mathcal{D}^{2,m}[i; n - 2f_m + 1]$ , i.e.,  $\#\mathcal{D}^{2,m}[i; n - 2f_m + 1]$ .

When  $n < 2f_{h+1}$ , there is no square of length  $2f_m$  ( $m \geq h + 1$ ) in  $\mathbb{F}[i; n]$ . Thus we only need to consider sequences  $\mathcal{D}^{2,m}$  for  $0 \leq m \leq h$  when  $2f_h \leq n < 2f_{h+1}$ .

### 5.1 Three useful lemmas

Let  $R_a$  and  $R_b$  be two words of lengths  $f_m$  and  $f_{m-1}$  for  $m \geq 0$ . We give three useful lemmas about the sequence  $\mathbb{F}(R_a, R_b)$  in this subsection.

**Lemma 5.1.** For  $i > f_m$ , let  $\text{Fib}(i - 1) = a_k a_{k-1} \dots a_1 a_0$  then  $k \geq m$  and

$$\mathbb{F}(R_a, R_b)[i] = (R_a R_b)[[a_m a_{m-1} \dots a_0]_F + 1].$$

**Example.** Take  $i = 20$ ,  $R_a = ABC$  and  $R_b = DE$ . Then  $\text{Fib}(i - 1) = 101001$ . Moreover,  $\mathbb{F}(R_a, R_b) = ABCDEABCABCDEABCDEABC\dots$ , so  $\mathbb{F}(R_a, R_b)[i] = B$ . On the other hand,  $m = 2$  and

$$(R_a R_b)[[a_m a_{m-1} \dots a_0]_F + 1] = (ABCDE)[[001]_F + 1] = ABCDE[2] = B.$$

**Lemma 5.2.** Let  $h \geq 0$  be an integer, and

$$M = \begin{cases} h, & h \geq m + 2, \\ m + 2, & h = m + 1, \\ m + 1, & h = 1, 2, \dots, m, \\ m, & h = 0. \end{cases}$$

Then the sequence  $\Theta = \{\mathbb{F}(R_a, R_b)[i; f_h]\}_{i \geq 1}$  is a Fibonacci word over  $\{\Theta[1, f_M], \Theta[f_M + 1, f_{M+1}]\}$  without prefix.

$$\begin{aligned}
\mathbb{F}(ab, c) &= a \ b \ c \ a \ b \ a \ b \ c \ a \ b \ c \ a \ b \ a \ \dots \\
\{\mathbb{F}(ab, c)[i; 2]\}_{i \geq 1} &= ab \ bc \ ca \ ab \ ba \ ab \ bc \ ca \ ab \ bc \ ca \ ab \ ba \ \dots \\
\mathbb{F}(ABC, AD) &= \underbrace{A \ B \ C}_{ABC} \ \underbrace{A \ D}_{AD} \ \underbrace{A \ B \ C}_{ABC} \ \underbrace{A \ B \ C}_{ABC} \ \underbrace{A \ D}_{AD} \ \dots
\end{aligned}$$

Figure 5: For instance, the sequence  $\{\mathbb{F}(ab, c)[i; 2]\}_{i \geq 1} = \mathbb{F}(ABC, AD)$ .

**Example.** Let  $m = h = 1$ . By Lemma 5.2,  $M = m + 1 = 2$  and the sequence  $\{\mathbb{F}(ab, c)[i; 2]\}_{i \geq 1}$  is a Fibonacci word over  $\{\Theta[1, f_M], \Theta[f_M + 1, f_{M+1}]\} = \{ABC, AD\}$  where  $A = ab$ ,  $B = bc$ ,  $C = ca$  and  $D = ba$ . We check this conclusion in Figure 5.

*Proof.* (1) We first let  $h = m + 2$ . Since  $\mathbb{F} = \mathbb{F}(F_2, F_1)$ , a decomposition of  $\mathbb{F}$  is

$$\mathbb{F} = F_2 \cdot F_1 \cdot F_2 \cdot F_2 \cdot F_1 \cdots \quad (7)$$

Here  $\omega \cdot \nu$  is the concatenation of two words  $\omega$  and  $\nu$ . In this decomposition,  $F_2$  and  $F_1$  are used as building blocks of  $\mathbb{F}$ . Since  $bb \not\prec \mathbb{F}$ , each occurrence of letter  $a$  in  $\mathbb{F}$  is followed by word  $ba$  or  $a$ . Thus each block  $F_2$  in Equation (7) is followed by the blocks  $F_1 \cdot F_2 = ababa$  or  $F_2 = aba$ . This means, each block  $F_2 = aba$  is followed by word  $aba$ . Notice that, the last word  $aba$  may not be a block. Similarly,  $\mathbb{F}(R_a, R_b)$  has a decomposition with blocks  $R_a R_b R_a$  and  $R_a R_b$  that

$$\mathbb{F}(R_a, R_b) = R_a R_b R_a \cdot R_a R_b \cdot R_a R_b R_a \cdot R_a R_b R_a \cdot R_a R_b \cdots \quad (8)$$

And each block  $R_a R_b R_a$  in Equation (8) is followed by word  $R_a R_b R_a$  (may not be a block). On the other hand, since  $|R_a R_b R_a| = f_{m+2}$ , each word starting in block  $R_a R_b R_a$  of length  $f_{m+2}$  belongs in  $R_a R_b R_a \cdot R_a R_b R_a$ . This means, for fixed  $1 \leq i \leq f_{m+2}$ , the factor of length  $f_{m+2}$  starting in the  $i$ -th letter in any block  $R_a R_b R_a$  are  $(R_a R_b R_a R_a R_b R_a)[i; f_{m+2}]$ .

Similarly, each block  $R_a R_b$  is followed by the word  $R_a R_b R_a$  too. And for  $1 \leq i \leq f_{m+1}$ , the factor of length  $f_{m+2}$  starting in the  $i$ -th letter in any block  $R_a R_b$  are  $(R_a R_b R_a R_b R_a)[i; f_{m+2}]$ . Thus the conclusion holds for  $h = m + 2$ .

(2) Since  $\mathbb{F}(R_a, R_b)$  is also a Fibonacci word over  $\{R_a R_b, R_a\}$  and  $|R_a R_b| = f_{m+1}$ , the conclusion holds for  $h = m + 3$  by (1) above. And the conclusion holds for all  $h \geq m + 3$  by induction.

(3) When  $h = m + 1$ ,  $\mathbb{F}(R_a, R_b)[i; f_h]$  is prefix of  $\mathbb{F}(R_a, R_b)[i; f_{m+2}]$ . By (1), the conclusion holds.

(4) When  $h = m$ ,  $\mathbb{F}(R_a, R_b)$  has a decomposition  $\mathbb{F}(R_a, R_b) = R_a R_b \cdot R_a \cdot R_a R_b \cdot R_a R_b \cdot R_a \cdots$ . Each block  $R_a R_b$  (resp.  $R_a$ ) is followed by the word  $R_a$ . On the other hand,  $|R_a| = f_m$ . This means, for  $1 \leq i \leq f_{m+1}$  (resp.  $1 \leq i \leq f_m$ ), the factor of length  $f_m$  starting in the  $i$ -th letter in any block  $R_a R_b$  (resp.  $R_a$ ) are  $(R_a R_b R_a)[i; f_m]$  (resp.  $(R_a R_a)[i; f_m]$ ). Thus the conclusion holds for  $h = m$ .

(5) When  $h = 1, \dots, m - 1$ , by (4) and an analogous argument in (2), the conclusion holds too.  $\square$

**Lemma 5.3.** Let  $2f_h \leq n < 2f_{h+1}$  and  $0 \leq m \leq h$ .  $\Theta = \{\mathcal{D}^{2,m}[i; n - 2f_m + 1]\}_{i \geq 1}$  is a Fibonacci word over  $\{\Theta[1, f_{h+3}], \Theta[f_{h+3} + 1, f_{h+4}]\}$  without prefix.

*Proof.* By Theorem 4.3,  $\mathcal{D}^{2,m}$  is a Fibonacci word over  $\{\mathcal{D}^{2,m}[1, f_{m+1}], \mathcal{D}^{2,m}[f_{m+1} + 1, f_{m+2}]\}$ . On the other hand,  $n - 2f_m + 1 < n < 2f_{h+1} < f_{h+3}$ . Thus by Lemma 5.2 the conclusion holds.  $\square$

## 5.2 Distinct squares in $\mathbb{F}[i; n]$

Since the return word sequence of letter  $b$  in  $\mathbb{F}$  is  $\mathbb{F}(baa, ba)$ , two successional occurrences of letter  $b$  have two cases:  $baab$  and  $bab$ . On the other hand,  $\mathcal{D}^{2,m}$  is a Fibonacci word over

$$\{R_a^{2,m}, R_b^{2,m}\} = \{[1, 2, \dots, f_{m-1} - 1, \underbrace{0, \dots, 0}_{f_{m+1}}, [1, 2, \dots, f_m]\}.$$

Two successional occurrences of  $R_b^{2,m}$  have two cases:  $[R_b^{2,m}, R_a^{2,m}, R_a^{2,m}, R_b^{2,m}]$  and  $[R_b^{2,m}, R_a^{2,m}, R_b^{2,m}]$ . This means when  $n = |[R_b^{2,m}, R_a^{2,m}, R_a^{2,m}]| = f_{m+3}$ , one of the two cases below holds

$$\begin{cases} R_b^{2,m}[1, j] \triangleright \mathcal{D}^{2,m}[i; n] \text{ and } R_b^{2,m}[j+1, f_m] \triangleleft \mathcal{D}^{2,m}[i; n] \\ R_b^{2,m} \prec \mathcal{D}^{2,m}[i; n]. \end{cases} \quad \text{for some integer } j \text{ such that } 0 \leq j \leq f_m,$$

Thus when  $n - 2f_m + 1 \geq f_{m+3}$ ,  $\mathcal{D}^{2,m}[i; n - 2f_m + 1]$  contains all elements in  $R_b^{2,m}$ . By Equation (6),  $R_b^{2,m}$  gives positions of all distinct squares of length  $2f_m$ . In this situation  $\#\mathcal{D}^{2,m}[i; n - 2f_m + 1] = f_m$ , and  $\mathbb{F}[i; n]$  contains all distinct squares of length  $2f_m$ .

This means  $\#\mathcal{D}^{2,m}[i; n - 2f_m + 1] = f_m$  for  $0 \leq m \leq h - 3$ . So

$$D(2, i, n) = \sum_{m=0}^{h-3} f_m + \sum_{m=h-2}^h \#\mathcal{D}^{2,m}[i; n - 2f_m + 1].$$

Since  $\sum_{j=0}^m f_j = f_{m+2} - 2$ , the number of distinct squares in  $\mathbb{F}[i; n]$  is equal to

$$D(2, i, n) = \begin{cases} \#\mathcal{D}^{2,0}[i; n - 1], & n = 2, 3, \\ \#\mathcal{D}^{2,0}[i; n - 1] + \#\mathcal{D}^{2,1}[i; n - 3], & n = 4, 5, \\ \sum_{m=h-2}^h \#\mathcal{D}^{2,m}[i; n - 2f_m + 1] + f_{h-1} - 2, & n \geq 6. \end{cases} \quad (9)$$

Now we turn to simplify  $\sum_{m=h-2}^h \#\mathcal{D}^{2,m}[i; n - 2f_m + 1]$  for  $2f_h \leq n < 2f_{h+1}$ . First we divide  $2f_h \leq n < 2f_{h+1}$  into five ranges, see Figure 4(A).

**Conclusion 1.** By Lemma 5.3,  $\{\#\mathcal{D}^{2,m}[i; n - 2f_m + 1]\}_{i \geq 1}$  is a Fibonacci word over  $\{\tilde{R}_a, \tilde{R}_b\}$  for fixed  $m \in \{h, h-1, h-2\}$  and fixed  $n$ , where  $|\tilde{R}_a| = f_{h+3}$  and  $|\tilde{R}_b| = f_{h+2}$ . We must determine the expressions of  $\tilde{R}_a$  and  $\tilde{R}_b$  for each  $m \in \{h, h-1, h-2\}$ .

Since the proof is tediously long, we only list the expressions of  $R_a$  and  $R_b$  in Figure 6.

$$\begin{cases} \tilde{R}_a = R_a R_b R_a, \tilde{R}_b = R_a R_b, |R_a| = f_{h+1}, |R_b| = f_h, & \text{if } n \in \text{I} \cup \text{II} \cup \text{III}(1), \\ \tilde{R}_a = R_a R_b, \tilde{R}_b = R_a, |R_a| = f_{h+2}, |R_b| = f_{h+1}, & \text{otherwise.} \end{cases}$$

Thus in all cases above,  $\{\#\mathcal{D}^{2,m}[i; n - 2f_m + 1]\}_{i \geq 1}$  is a Fibonacci word over  $\{R_a, R_b\}$  without prefix.

**Conclusion 2.** By Conclusion 1,  $\{\sum_{m=h-2}^h \#\mathcal{D}^{2,m}[i; n - 2f_m + 1]\}_{i \geq 1}$  is also a Fibonacci word for fixed  $n$ . We give the expressions of  $R_a$  and  $R_b$  in Figure 4(B). When  $n \in \text{I} \cup \text{II} \cup \text{III}$ ,  $|R_a| = f_{h+1}$  and  $|R_b| = f_h$ ; otherwise  $|R_a| = f_{h+2}$  and  $|R_b| = f_{h+1}$ .

**Conclusion 3.** By Conclusion 2 and Lemma 5.1,

$$\sum_{m=h-2}^h \#\mathcal{D}^{2,m}[i; n - 2f_m + 1] = \begin{cases} [R_a, R_b][\hat{i}], & \text{if } n \in \text{I} \cup \text{II} \cup \text{III}, \\ [R_a, R_b][\hat{j}], & \text{otherwise.} \end{cases}$$

Here  $\text{Fib}(i-1) = a_k a_{k-1} \dots a_1 a_0$ ,  $\hat{i} = [a_{h+2} a_{h+1} \dots a_1 a_0]_F + 1$  and  $\hat{j} = [a_{h+3} a_{h+2} \dots a_1 a_0]_F + 1$ .

By an analogous argument, we simplify Equation (9) for  $2 \leq n \leq 5$ . Thus we get Theorem 3.2.

### 5.3 Distinct squares in $\mathbb{F}[1, n]$

By analyzing the first number for each  $n$  in Figure 4(B), we have

$$D(2, 1, n) - f_{h-1} + 2 = \begin{cases} n - 2f_{h-1}, & 2f_h \leq n < f_{h+1} + 2f_{h-1} \text{ (Range I} \cup \text{II} \cup \text{III)}, \\ f_{h+1} - 1, & f_{h+1} + 2f_{h-1} \leq n < 2f_{h+1} \text{ (Range V} \cup \text{VI)}. \end{cases}$$

Let  $Z[n] = D(2, 1, n+1) - D(2, 1, n)$ ,  $Z_1 = Z[1, 4]$  and  $Z_h = Z[2f_h - 1, 2f_{h+1} - 2]$  for  $h \geq 2$ . Then  $D(2, 1, n) = \sum_{j=1}^{n-1} Z[j]$ . We can prove that  $Z_1 = [0, 0, 1, 0]$  for  $h \geq 2$

$$Z_h = \underbrace{[1, \dots, 1]}_{f_{h-1} + f_{h-3}} \underbrace{[0, \dots, 0]}_{f_{h-2}} \text{ and } \sum_{j=1}^{2f_h-2} Z[j] = f_h + f_{h-2} - 3.$$

First we divide Range III into Range III(1) ( $2f_h + f_{h-2} - 1 \leq n < f_{h+2} - 1$ ) and Range III(2) ( $f_{h+2} - 1 \leq n < f_{h+1} + 2f_{h-1}$ ).

**When  $m = h$ .**  $n \in I \cup II \cup III(1)$ :  $R_a = [n - 2f_h + 1 \rightarrow, n - 2f_h \searrow, 0 \rightarrow, 1 \nearrow]$ ;  $R_b = [n - 2f_h + 1 \rightarrow]$ .

$n \in III(2)$ :  $R_a = [f_{h-1} - 1 \searrow, 0 \rightarrow, 1 \nearrow, n - 2f_h + 1 \rightarrow, n - 2f_h \searrow]$ ;  $R_b = [f_{h-1} - 1 \searrow, 0 \rightarrow, 1 \nearrow, f_{h-1} - 1 \rightarrow]$ .

$n \in IV$ :  $R_a = [f_{h-1} - 1 \searrow, 0 \rightarrow, 1 \nearrow, n - 2f_h + 1 \rightarrow, n - 2f_h \searrow]$ ;  $R_b = [f_{h-1} - 1 \searrow, 0 \rightarrow, 1 \nearrow, f_{h-1} - 1 \rightarrow]$ .

$n \in V$ :  $R_a = [f_{h-1} - 1 \searrow, n - 3f_h \rightarrow, n - 3f_h + 1 \nearrow, f_h \rightarrow, f_h - 1 \searrow]$ ;  $R_b = [f_{h-1} - 1 \searrow, n - 3f_h \rightarrow, n - 3f_h + 1 \nearrow, f_{h-1} - 1 \rightarrow]$ .

**When  $m = h - 1$ .**  $n \in I$ :  $R_a = [f_{h-2} - 1 \searrow, n - 3f_{h-1} \rightarrow, n - 3f_{h-1} + 1 \nearrow, f_{h-1} \rightarrow, f_{h-1} - 1 \searrow]$ ;  $R_b = [f_{h-2} - 1 \searrow, n - 3f_{h-1} \rightarrow, n - 3f_{h-1} + 1 \nearrow, f_{h-2} - 1 \rightarrow]$ .

$n \in II \cup III(1)$ :  $R_a = [f_{h-2} - 1 \rightarrow, f_{h-2} \nearrow, f_{h-1} \rightarrow, f_{h-1} - 1 \searrow]$ ;  $R_b = [f_{h-2} - 1 \rightarrow]$ .

$n \in III(2)$ :  $R_a = [n - f_{h+1} - f_{h-1} + 1 \nearrow, f_{h-1} \rightarrow, f_{h-1} - 1 \searrow, f_{h-2} - 1 \rightarrow, f_{h-2} \nearrow]$ ;  $R_b = [n - f_{h+1} - f_{h-1} + 1 \nearrow, f_{h-1} \rightarrow, f_{h-1} - 1 \searrow, n - f_{h+1} - f_{h-1} + 1 \rightarrow]$ .

$n \in IV \cup V$ :  $R_a = [f_{h-1} \rightarrow, f_{h-1} - 1 \searrow, f_{h-2} - 1 \rightarrow, f_{h-2} \nearrow, f_{h-1} \rightarrow]$ ;  $R_b = [f_{h-1} \rightarrow]$ .

**When  $m = h - 2$ .**  $n \in I$ :  $R_a = [f_{h-2} \rightarrow, f_{h-2} - 1 \searrow, f_{h-3} - 1 \rightarrow, f_{h-3} \nearrow, f_{h-2} \rightarrow]$ ;  $R_b = [f_{h-2} \rightarrow]$ .

$n \in II$ :  $R_a = [f_{h-2} \rightarrow, f_{h-2} - 1 \searrow, n - 2f_h + 1 \rightarrow, n - 2f_h + 1 \nearrow, f_{h-2} \rightarrow]$ ;  $R_b = [f_{h-2} \rightarrow]$ .

$n \in III(1)$ :  $R_a = [f_{h-2} \rightarrow]$ ;  $R_b = [f_{h-2} \rightarrow]$ .

$n \in III(2) \cup IV \cup V$ :  $R_a = [f_{h-2} \rightarrow]$ ;  $R_b = [f_{h-2} \rightarrow]$ .

Figure 6: The expressions of  $R_a$  and  $R_b$  in Conclusion 1 for  $r = 2$ .

Thus when  $2f_h - 1 \leq n < 2f_{h+1} - 1$ ,  $D(2, 1, n) = \sum_{j=1}^{n-1} Z[j] = \sum_{j=1}^{2f_h-2} Z[j] + \sum_{j=2f_h-1}^{n-1} Z[j]$ . So

$$D(2, 1, n) = f_h + f_{h-2} - 3 + \min\{n - 2f_h + 1, f_{h-1} + f_{h-3}\}.$$

Thus  $D(2, 1, n) = \min\{n - f_{h-1} - 2, f_{h+1} + f_{h-1} - 3\}$ . Now we obtain Theorem 3.1.

## 5.4 All positions starting a square

Harju-Kärki-Nowotka [11] considered the number  $N(\omega)$  of positions that do not start a square in a binary word  $\omega$ . Letting  $N(n)$  be the maximum of  $N(\omega)$  for length  $|\omega| = n$ , they showed that  $\lim_{n \rightarrow \infty} \frac{N(n)}{n} = \frac{15}{31}$ . As an application of factor spectrum, we have

**Property 5.4.** *All positions in  $\mathbb{F}$  start a square of length  $2f_m$  for some  $0 \leq m \leq 3$ .*

Define sequence  $\mathbb{D}$  by that  $\mathbb{D}[i] = \sum_{m=0}^3 \mathcal{D}^{2,m}[i]$ . The property is equivalent to prove  $\mathbb{D}[i] > 0$  for all  $i$ . Since all of  $\mathcal{D}^{2,m}$  ( $0 \leq m \leq 3$ ) are Fibonacci words over  $\{\mathcal{D}^{2,m}[1, f_4], \mathcal{D}^{2,m}[f_4 + 1, f_5]\}$ , we only need to prove that  $\mathbb{D}[i] > 0$  for all  $1 \leq i \leq f_5 = 13$ .

By the expressions of  $R_a^{2,m}$  and  $R_b^{2,m}$  given in Theorem 4.3 and Equation (6),

$$\begin{cases} \mathcal{D}^{2,0}[1, 13] = 0010000100100, & \mathcal{D}^{2,1}[1, 13] = 0001200000012, \\ \mathcal{D}^{2,2}[1, 13] = 1000012310000, & \mathcal{D}^{2,3}[1, 13] = 1200000012345. \end{cases}$$

This obtain the conclusion in Property 5.4.

**Remark.** Let us consider factor spectrum  $\mathcal{S}_{\mathcal{P}_3} = \{(\omega, j) \mid \mathbb{F}[j; 2|\omega|] = \omega\omega\}$ . This property means that for all  $j \geq 1$  there exists a factor  $\omega$  of length  $f_m$  ( $0 \leq m \leq 3$ ) such that  $(\omega, j) \in \mathcal{S}_{\mathcal{P}_3}$ .



By Equation (10),

When  $n \in \text{Range I}$ ,  $\overline{Max} = \overline{\lim}_{n \rightarrow \infty} \frac{f_{h+1}-2}{n} = \lim_{h \rightarrow \infty} \frac{f_{h+1}-2}{2f_h} = \frac{\alpha}{2} \doteq 0.8090$

$$\underline{Max} = \underline{\lim}_{n \rightarrow \infty} \frac{f_{h+1}-2}{n} = \lim_{h \rightarrow \infty} \frac{f_{h+1}-2}{2f_h+f_{h-3}-1} = \frac{\alpha^4}{2\alpha^3+1} \doteq 0.7236$$

Otherwise,  $\frac{\max_{1 \leq i \leq \infty} D(2, i, n)}{n} = \frac{n-f_{h-1}}{n} = 1 - \frac{f_{h-1}}{n}$ . Thus

$$\overline{Max} = 1 - \lim_{h \rightarrow \infty} \frac{f_{h-1}}{2f_{h+1}-1} = 1 - \frac{1}{2\alpha^2} \doteq 0.8090;$$

$$\underline{Max} = 1 - \lim_{h \rightarrow \infty} \frac{f_{h-1}}{2f_h+f_{h-3}-1} = 1 - \frac{\alpha^2}{2\alpha^3+1} \doteq 0.7236.$$

This means  $0.7236n \leq \max_{1 \leq i \leq \infty} D(2, i, n) \leq 0.8090n$  for  $n$  large enough.

By Equation (11) and an analogous argument,

$$\left\{ \begin{array}{ll} \underline{Min} = 2 - \frac{\alpha^2+2}{2\alpha} \doteq 0.5729, & \overline{Min} = 2 - \frac{2}{7}(\alpha^2 + 2) \doteq 0.6806, \quad n \in \text{Range I(1)}, \\ \underline{Min} = \frac{\alpha^5-1}{2\alpha^4+\alpha} \doteq 0.6583, & \overline{Min} = \frac{2(\alpha^5-1)}{7\alpha^3} \doteq 0.6806, \quad n \in \text{Range I(2)}, \\ \underline{Min} = 1 - \frac{2\alpha}{2\alpha^3+1} \doteq 0.6583, & \overline{Min} = 1 - \frac{2}{2\alpha^2+1} \doteq 0.6793, \quad n \in \text{Range II}, \\ \underline{Min} = \frac{\alpha}{3} \doteq 0.5393, & \overline{Min} = \frac{\alpha^3}{2\alpha^2+1} \doteq 0.6793, \quad n \in \text{Range III} \cup \text{IV}, \\ \underline{Min} = 1 - \frac{\alpha^2+1}{3\alpha^2} \doteq 0.5393, & \overline{Min} = 1 - \frac{\alpha^2+1}{2\alpha^3} \doteq 0.5729, \quad n \in \text{Range V}. \end{array} \right.$$

Since  $\min\{0.5729, 0.6583, 0.5393\} = 0.5393$  and  $\max\{0.6806, 0.6793, 0.5729\} = 0.6806$ , we have  $0.5393n \leq \min_{1 \leq i \leq \infty} D(2, i, n) \leq 0.6806n$  for  $n$  large enough.

Similarly,

$$\left\{ \begin{array}{ll} \underline{\Delta} = \frac{2 \times (\alpha^3+3)}{7} - 2 \doteq 0.0674, & \overline{\Delta} = \frac{\alpha^3+3}{2\alpha} - 2 \doteq 0.2361, \quad n \in \text{Range I(1)}, \\ \underline{\Delta} = \frac{1}{2\alpha^4+\alpha} \doteq 0.0653, & \overline{\Delta} = \frac{2}{7\alpha^3} \doteq 0.0675, \quad n \in \text{Range I(2)}, \\ \underline{\Delta} = \frac{1}{2\alpha^4+2\alpha^2} \doteq 0.0528, & \overline{\Delta} = \frac{1}{2\alpha^4+\alpha} \doteq 0.0653, \quad n \in \text{Range II}, \\ \underline{\Delta} = 1 - \frac{\alpha^3+\alpha}{2\alpha^2+1} \doteq 0.0613, & \overline{\Delta} = 1 - \frac{\alpha^2+1}{3\alpha} \doteq 0.2547, \quad n \in \text{Range III} \cup \text{IV}, \\ \underline{\Delta} = \frac{1}{\alpha^3} \doteq 0.2361, & \overline{\Delta} = \frac{2}{3\alpha^2} \doteq 0.2547, \quad n \in \text{Range V}. \end{array} \right.$$

Since  $\min\{0.0674, 0.0653, 0.0528, 0.0613, 0.2361\} = 0.0528$  and  $\max\{0.2361, 0.0675, 0.0653, 0.2547\} = 0.2547$ , we have  $0.0528n \leq \max_{1 \leq i, j \leq \infty} |D(2, i, n) - D(2, j, n)| \leq 0.2547n$  for  $n$  large enough.

### 6.3 Property of intermediate values

By Figure 4(B) and Theorem 3.2,  $|D(2, i+1, n) - D(2, i, n)| \leq 1$ . Thus for fixed  $n$  and any  $M$  such that  $\max_{1 \leq i \leq \infty} D(2, i, n) \leq M \leq \min_{1 \leq i \leq \infty} D(2, i, n)$ , there exists  $i_0$  such that  $D(2, i_0, n) = M$ .

## 7 The number of r-powers in $\mathbb{F}[i; n]$

First of all, we prove some facts, which are fit for all sequence including  $\mathbb{F}$ .

Fact 1. If both  $P$  and  $P+1$  are the positions of some squares of size  $n$ , then  $P$  is the position of some  $(2 + \frac{1}{n})$ -powers of size  $n$ . In fact, since  $P$  is the position of some squares of size  $n$ ,  $\mathbb{F}[P, P+2n-1]$  has expression  $x_1x_2 \cdots x_nx_1x_2 \cdots x_n$  for  $x_i \in \{a, b\}$ . Since  $P+1$  is the position of some squares of size  $n$  too,  $\mathbb{F}[P+2n] = x_1$ . This means  $\mathbb{F}[P, P+2n] = x_1x_2 \cdots x_nx_1x_2 \cdots x_nx_1$ . Thus  $P$  is the position of the  $(2 + \frac{1}{n})$ -power factor  $\mathbb{F}[P, P+2n]$ .

Fact 2. If both  $P$  and  $P+1$  are the positions of some  $(2 + \frac{i}{n})$ -powers of size  $n$ , then  $P$  is the position of some  $(2 + \frac{i+1}{n})$ -powers of size  $n$  for  $i \geq 0$ .

Fact 3. If each integer in set  $\{P, P+1, \dots, P+h\}$  is the position of some  $(2 + \frac{i}{n})$ -powers of size  $n$ , then each integer in  $\{P, P+1, \dots, P+h-j\}$  is the position of some  $(2 + \frac{i+j}{n})$ -powers of size  $n$  for

$h \geq j$ . Now we consider the sequences  $\mathcal{D}^{r,m}$ . Let  $i \geq 0$  and  $n, h \geq 1$  such that

$$\begin{aligned} \mathcal{D}^{2+\frac{i}{f_m},m}[n; h+2] &= [0, 1, 2, \dots, h, 0] \\ \text{then } \mathcal{D}^{2+\frac{i+j}{f_m},m}[n; h+2] &= [0, 1, 2, \dots, h-j, \underbrace{0, \dots, 0}_{j+1}] \end{aligned} \quad (12)$$

for  $0 < j \leq h$ . For real number  $r \geq 2$ ,  $r$ -power  $\omega^r$  of size  $|\omega|$  means  $h$ -power  $\omega^h$ , where  $h = \frac{\lceil r|\omega| \rceil}{|\omega|}$ . So we can generalize the conclusion above to all real number  $r \geq 2$ .

We first generalize the definition of  $\mathcal{D}^{2,m}$  (see Definition 4.2) to  $\mathcal{D}^{r,m}$  as below.

**Definition 7.1.** We define sequence  $\mathcal{D}^{r,m}$  ( $r \geq 2$  and  $m \geq 0$ ) that for  $i \geq 1$  and  $1 \leq j \leq f_m$

$$\mathcal{D}^{r,m}[i] = \begin{cases} j, & \text{if } \mathcal{D}^{2,m}[i] = j \text{ and } \mathbb{F}[i; \lceil rf_m \rceil] \text{ is an } r\text{-power;} \\ 0, & \text{otherwise.} \end{cases}$$

We call  $\mathcal{D}^{r,m}$  the position sequence of  $r$ -powers of size  $f_m$  for  $m \geq 0$ .

By Fact 3 above and Theorem 4.3, we have

**Property 7.2.**  $\mathcal{D}^{r,m}$  is a Fibonacci word over  $\{R_a^{r,m}, R_b^{r,m}\}$  for  $r \geq 2$  and  $m \geq 0$ . Specifically,

- (1)  $\mathcal{D}^{r,0}$  is a zero sequence for  $r > 2$ ;
- (2)  $\mathcal{D}^{r,1}$  is a zero sequence for  $r > 2.5$ ;  $R_a^{r,1} = [0, 0, 0]$  and  $R_b^{r,1} = [1, 0]$  for  $2 < r \leq 2.5$ ;
- (3)  $\mathcal{D}^{r,m}$  ( $m \geq 2$ ) is a zero sequence for  $r > 3 + \frac{f_{m-1}-1}{f_m}$ ; and

$$\begin{cases} R_a^{r,m} = [1, 2, \dots, f_{m-1} - x, \underbrace{0, \dots, 0}_{f_m+x}], & R_b^{r,m} = [1, 2, \dots, f_m], & 2 \leq r \leq 2 + \frac{f_{m-1}-1}{f_m} \\ R_a^{r,m} = [\underbrace{0, \dots, 0}_{f_{m+1}}], & R_b^{r,m} = [1, 2, \dots, f_m - y, \underbrace{0, \dots, 0}_y], & 2 < r - \frac{f_{m-1}-1}{f_m} \leq 3. \end{cases}$$

Here  $x = \lceil rf_m \rceil - 2f_m + 1$  and  $y = \lceil rf_m \rceil - f_{m+2} + 1$ .

**Remark.** Given a sequence  $\rho$ , it is an interesting and challenging task to determine its critical exponent  $e$ , such that  $\rho$  contains  $r$ -powers for all  $r < e$ , but has no  $r$ -powers for all  $r > e$  [1]. The critical exponent of  $\mathbb{F}$  is  $\lim_{m \rightarrow \infty} 3 + \frac{f_{m-1}-1}{f_m} = 3 + \frac{1}{\alpha} \approx 3.618$ , where  $\alpha = \frac{1+\sqrt{5}}{2}$ . This is an immediate corollary of Property 7.2.

Now we turn to consider two special cases:  $r = 2 + \epsilon$  and  $r = 3$ .

When  $r = 2 + \epsilon$ ,  $\mathcal{D}^{r,0}$  is a zero sequence;  $R_a^{r,1} = [0, 0, 0]$  and  $R_b^{r,1} = [1, 0]$ ; for  $m \geq 2$

$$R_a^{r,m} = [1, 2, \dots, f_{m-1} - 2, \underbrace{0, \dots, 0}_{f_m+2}] \text{ and } R_b^{r,m} = [1, 2, \dots, f_m]. \quad (13)$$

When  $r = 3$ , both  $\mathcal{D}^{r,0}$  and  $\mathcal{D}^{r,1}$  are zero sequences; for  $m \geq 2$

$$R_a^{r,m} = [\underbrace{0, \dots, 0}_{f_{m+1}}] \text{ and } R_b^{r,m} = [1, 2, \dots, f_{m-1} - 1, \underbrace{0, \dots, 0}_{f_{m-2}+1}]. \quad (14)$$

## 7.1 Distinct $r$ -powers for $r = 2 + \epsilon$

For  $n \geq 9$ , let  $h \geq 2$  such that  $2f_h \leq n < 2f_{h+1}$ . Then

$$D(2 + \epsilon, 1, n) = \min\{n - f_{h-1} - 6, f_{h+1} + f_{h-1} - 6\}.$$

By Equation (13),  $\#\mathcal{D}^{2+\epsilon,1}[i; n - 2f_1] = 1$  and  $\#\mathcal{D}^{2+\epsilon,m}[i; n - 2f_m] = f_m$  for  $2 \leq m \leq h - 3$ . Thus  $\sum_{m=1}^{h-3} \#\mathcal{D}^{2,m}[i; n - 2f_m] = \sum_{m=2}^{h-3} f_m + 1 = f_{h-1} - 4$ . For  $i \geq 1$  and  $n \geq 5$ , let  $h \geq 1$  such that  $2f_h + 1 \leq n < 2f_{h+1} + 1$ . Then

$$D(2 + \epsilon, i, n) = \begin{cases} [0, 0, n - 5, 1, 0, 0, 0, 0], & n = 5, 6, \\ [n - 7, 1, 1, 1, n - 7, n - 6, n - 6, 1, n - 7, 1, 1, 1, 0][\hat{j}], & n = 7, 8, \\ [1, 1, n - 8, n - 7, n - 7, 3, n - 7, 2, 1, 1, 1, 1, 1][\hat{j}], & n = 9, 10, \\ [R_a, R_b][\hat{i}] + f_{h-1} - 4, & n \geq 11 \text{ and } n \in \text{I} \cup \text{II} \cup \text{III} \cup \text{IV}, \\ [R_a, R_b][\hat{j}] + f_{h-1} - 4, & \text{otherwise.} \end{cases}$$

Here  $\text{Fib}(i - 1) = a_k a_{k-1} \dots a_1 a_0$ ,  $\hat{i} = [a_{h+1} a_h \dots a_1 a_0]_F + 1$  and  $\hat{j} = [a_{h+2} a_{h+1} \dots a_1 a_0]_F + 1$ . The definitions of Ranges I to VI, and the expressions of  $R_a$  and  $R_b$  see Figure 7.

**Property 7.3.** *All positions in  $\mathbb{F}$  starting a  $(2 + \epsilon)$ -power of size  $f_m$  for some  $2 \leq m \leq 6$ .*

The proof could be obtained by an analogous argument in Property 5.4. Let us consider factor spectrum  $\mathcal{S}_{\mathcal{P}_4} = \{(\omega, j) \mid \mathbb{F}[j; 2|\omega| + 1] = \omega\omega \cdot \omega[1]\}$ . This property means that for all  $j \geq 1$  there exists a factor  $\omega$  such that  $(\omega, j) \in \mathcal{S}_{\mathcal{P}_4}$ .

## 7.2 Distinct r-powers for $r = 3$

For  $n \geq 14$ , let  $h \geq 3$  such that  $2f_h - 1 \leq n < 2f_{h+1} - 1$ . Then

$$D(3, 1, n) = \max\{n - f_{h+1} - f_{h-1} - h + 1, f_{h-1} - h\}.$$

By Equation (14),  $\#\mathcal{D}^{2+\epsilon,m}[i; n - 3f_m] = f_{m-1} - 1$  for  $2 \leq m \leq h - 3$ . Thus

$$\sum_{m=2}^{h-3} \#\mathcal{D}^{2,m}[i; n - 3f_m] = \sum_{m=2}^{h-3} (f_{m-1} + 1) = f_{h-1} + h - 9.$$

For  $i \geq 1$  and  $n \geq 9$ , let  $h \geq 2$  such that  $3f_h \leq n < 3f_{h+1}$ . Then

$$D(3, i, n) = \begin{cases} [0, \dots, 0, \underbrace{1, \dots, 1}_{n-8}, 0, \dots, 0, ][\hat{i}], & 9 \leq n \leq 14, \\ [R_a, R_b][\hat{i}] + f_{h-1} + h - 9, & n \geq 15 \text{ and } n \notin \text{Range X}, \\ [R_a, R_b][\hat{j}] + f_{h-1} + h - 9, & \text{otherwise.} \end{cases}$$

Here  $\text{Fib}(i - 1) = a_k a_{k-1} \dots a_1 a_0$ ,  $\hat{i} = [a_{h+2} a_h \dots a_1 a_0]_F + 1$  and  $\hat{j} = [a_{h+3} a_{h+1} \dots a_1 a_0]_F + 1$ . The definitions of Ranges I to X, and the expressions of  $R_a$  and  $R_b$  see Figure 8.

There exists some positions that do not start a cube. In fact,  $\mathbb{F}[1, n]$  is not a cube for all  $n \geq 1$ .

**Property 7.4.** *For  $m \geq 0$  the number of positions starting a cube in  $\mathbb{F}[1, f_m]$  is*

$$C(m) = \begin{cases} f_{2k} - k^2 - k - 1, & m = 2k, \\ f_{2k+1} - k^2 - 2k - 2, & m = 2k + 1. \end{cases}$$

*Proof.* We only need to prove the relation below by induction

$$C(m) = C(m - 1) + C(m - 2) - C(m - 3) + \#R_b^{3,m-2}.$$

Here  $\#R_b^{3,m-2} = \#[1, 2, \dots, f_{m-3} - 1, \underbrace{0, \dots, 0}_{f_{m-4} + 1}] = f_{m-3} - 1$ . We give this relation in Figure 9 for  $m = 7$ . □

**Remark.** Since  $\lim_{m \rightarrow \infty} \frac{C(m)}{f_m} = 1$ , almost all positions in  $\mathbb{F}$  start some cubes. Let us consider factor spectrum  $\mathcal{S}_{\mathcal{P}_5} = \{(\omega, j) \mid \mathbb{F}[j; 3|\omega|] = \omega\omega\omega\}$ . The asymptotic solution is equivalent to that for almost all  $j \geq 1$  there exists a factor  $\omega$  such that  $(\omega, j) \in \mathcal{S}_{\mathcal{P}_5}$ .

*Acknowledgements.* The research is supported by the Grant NSFC No.11701024 and No.11431007.

Divide  $2f_h + 1 \leq n < 2f_{h+1} + 1$  into six ranges:

Range I:  $2f_h + 1 \leq n < 2f_h + f_{h-3}$ ;

Range II:  $n = 2f_h + f_{h-3}$ ;

Range III:  $2f_h + f_{h-3} + 1 \leq n < 2f_h + f_{h-2}$ ;

Range IV:  $2f_h + f_{h-2} \leq n < f_{h+1} + 2f_{h-1} + 1$ ; Range V:  $f_{h+1} + 2f_{h-1} + 1 \leq n < 3f_h + 1$ ; Range VI:  $3f_h + 1 \leq n \leq 2f_{h+1}$ .

$$\begin{aligned}
n \in \text{I}: \quad R_a &= \underbrace{[n - f_h - f_{h-3} - 2 \searrow, 2n - f_{h+1} - 3f_{h-1} - 2 \rightarrow, 2n - f_{h+1} - 3f_{h-1} - 1 \nearrow, n - f_h - f_{h-3} - 2 \rightarrow]}_{\substack{f_{h+1}+f_{h-1}-n \\ n - f_h - f_{h-3} - 1 \nearrow, f_h \rightarrow, f_{h-1} \searrow, f_{h-1} + f_{h-3} - 2 \rightarrow, f_{h-1} + f_{h-3} - 1 \nearrow, \\ f_{h+1}+f_{h-1}-n+1 \quad n-2f_h-1 \quad f_{h-4}+1 \quad f_{h+1}+f_{h-1}-n-1 \quad f_{h-4}+1 \\ f_h \rightarrow, f_{h-1} \searrow, n - f_h - f_{h-3} - 2 \rightarrow]}; \\
R_b &= \underbrace{[n - f_h - f_{h-3} - 2 \searrow, 2n - f_{h+1} - 3f_{h-1} - 2 \rightarrow, 2n - f_{h+1} - 3f_{h-1} - 1 \nearrow, n - f_h - f_{h-3} - 2 \rightarrow]}_{n-2f_h-1 \quad f_{h+1}+f_{h-1}-n+2 \quad n-2f_h-1}. \\
n \in \text{II}: \quad R_a &= \underbrace{[f_h - 2 \rightarrow, f_h - 1, f_h \rightarrow, f_h - 1 \searrow, f_h - f_{h-4} \rightarrow, f_h - f_{h-4} + 1 \nearrow, f_h \rightarrow, f_h - 1, f_h - 2 \rightarrow]}_{\substack{f_{h+1}+f_{h-1}-n \\ f_{h-3}-1 \quad f_{h-4}-1 \quad 3 \quad f_{h-4}-1 \quad f_{h-3}-1 \quad f_{h-3} \\ n - f_h - f_{h-3} - 2 \rightarrow, n - f_h - f_{h-3} - 2 \searrow, f_h \rightarrow, f_h - 1 \searrow, n - f_h - f_{h-2} \rightarrow, n - f_h - f_{h-2} + 1 \nearrow, \\ f_{h+1}+2f_{h-1}-n \quad n-f_{h+1}-f_{h-1}-1 \quad f_{h-3} \quad 2f_h+f_{h-2}-n-1 \quad n-f_{h+1}-f_{h-1}+3 \quad 2f_h+f_{h-2}-n-1 \\ f_h \rightarrow, f_h \nearrow, n - f_h - f_{h-3} - 2 \rightarrow];} \\
n \in \text{III}: \quad R_a &= \underbrace{[n - f_h - f_{h-3} - 2 \rightarrow, n - f_h - f_{h-3} - 2 \searrow, f_h \rightarrow, f_h - 1 \searrow, n - f_h - f_{h-2} \rightarrow, n - f_h - f_{h-2} + 1 \nearrow]}_{\substack{f_{h-3} \quad n-f_{h+1}-f_{h-1}-1 \quad f_{h-3}+1 \quad f_h \rightarrow, f_h \nearrow, n - f_h - f_{h-3} - 2 \rightarrow];} \\
n \in \text{IV}: \quad R_a &= \underbrace{[n - f_h - f_{h-3} - 2 \rightarrow, n - f_h - f_{h-3} - 2 \searrow, f_h \rightarrow, f_h \nearrow, n - f_h - f_{h-3} - 2 \rightarrow]}_{\substack{f_{h+1}+2f_{h-1}-n \quad n-f_{h+1}-f_{h-1}-1 \quad 3f_h-n+1 \quad n-f_{h+1}-f_{h-1}-1 \quad f_{h-3}+1 \\ f_{h+1}-2 \searrow, f_h \rightarrow, f_h \nearrow, n - f_h - f_{h-3} - 2 \rightarrow, n - f_h - f_{h-3} - 3 \searrow];} \\
n \in \text{V}: \quad R_a &= \underbrace{[f_{h+1} - 2 \searrow, f_h \rightarrow, f_h \nearrow, n - f_h - f_{h-3} - 2 \rightarrow, n - f_h - f_{h-3} - 3 \searrow]}_{\substack{f_{h-1}-1 \quad f_{h+2}+f_{h-2}-n+1 \quad n-f_{h+1}-f_{h-1}-2 \quad n-f_{h+3}+f_{h-4}-3 \quad n-f_{h+2}-f_{h-3}-1 \\ f_{h+1}-2 \searrow, f_h \rightarrow, f_h \nearrow, f_{h+1}-2 \rightarrow.} \\
n \in \text{VI}: \quad R_a &= \underbrace{[f_{h+1} - 2 \searrow, n - 2f_h - 2 \rightarrow, n - 2f_h - 1 \nearrow, n - f_h - f_{h-3} - 2 \rightarrow, n - f_h - f_{h-3} - 3 \searrow, f_{h+1} + f_{h-4} - 2 \rightarrow]}_{\substack{f_{h+2}+f_h-n \quad n-f_{h+2}-f_{h-2}-1 \quad 2f_h-2-1 \quad f_{h+2}+f_{h-1}-n+3 \quad n-f_{h+2}-f_{h-2}-1 \quad f_{h+2}+f_h-n-1 \\ f_{h+1} + f_{h-4} - 1 \nearrow, n - f_h - f_{h-3} - 2 \rightarrow, n - f_h - f_{h-3} - 3 \searrow, \\ n-f_{h+2}-f_{h-2}-1 \quad f_{h+2}+f_{h-1}-n+3 \quad n-f_{h+2}-f_{h-3}-1 \\ f_{h+1}-2 \searrow, n-2f_h-2 \rightarrow, n-2f_h-1 \nearrow, f_{h+1}-2 \rightarrow.} \\
R_b &= \underbrace{[f_{h+1} - 2 \searrow, n - 2f_h - 2 \rightarrow, n - 2f_h - 1 \nearrow, f_{h+1} - 2 \rightarrow]}_{\substack{f_{h+2}+f_h-n \quad n-f_{h+2}-f_{h-2}-1 \quad f_{h+2}+f_h-n-1 \quad n-f_{h+2}+2}}.
\end{aligned}$$

Figure 7: The expressions of  $R_a$  and  $R_b$  for  $r = 2 + \epsilon$ .  $|R_a| = f_{h+1}$  and  $|R_b| = f_h$  for  $n \in \text{I} \cup \text{II} \cup \text{III} \cup \text{IV}$ ;  $|R_a| = f_{h+2}$  and  $|R_b| = f_{h+1}$  otherwise.

## References

- [1] J. P. Allouche, J. Shallit. *Automatic sequences: Theory, applications, generalizations*. Cambridge University Press, Cambridge, 2003.
- [2] J. Berstel. Recent results on Sturmian words. *Salomaa Development in Language Theory World Scientific*. (1995) 13-24.
- [3] F. Durand. A characterization of substitutive sequences using return words, *Discrete Math*. 179 (1998) 89-101.
- [4] D. Damanik, D. Lenz, Powers in Sturmian sequences, *European Journal of Combinatorics*. 24.4 (2003) 377-390.
- [5] C.-F. Du, H. Mousavi, L. Schaeffer, J. Shallit. Decision Algorithms for Fibonacci-Automatic Words, with Applications to Pattern Avoidance. Presented at 15<sup>e</sup> Journées Montoises d'Informatique Théorique, 23-26 September 2014, Nancy, France.
- [6] C.-F. Du, H. Mousavi, L. Schaeffer, J. Shallit. Decision Algorithms for Fibonacci-Automatic Words, III: Enumeration and Abelian Properties. *Internat. J. Found. Comput. Sci*. 27.8 (2016) 943-963.
- [7] A. S. Fraenkel, J. Simpson, How Many Squares Can a String Contain? *Journal of Combinatorial Theory*. 82.1 (1998) 112-120.



- [8] A. S. Fraenkel, J. Simpson, The exact number of squares in Fibonacci words, *Theoretical Computer Science*. 218 (1999) 95-106.
- [9] A. S. Fraenkel, J. Simpson, Corrigendum to “The exact number of squares in Fibonacci words”, *Theoretical Computer Science*. 547 (2014) 122.
- [10] A. Glen, *On Sturmian and Episturmian Words, and Related Topics*, Ph.D. thesis. The University of Adelaide, Australia, 2006.
- [11] T. Harju, T. Kärki, D. Nowotka. The number of positions starting a square in binary words. *Electronic Journal of Combinatorics*. 18.1 (2011), 106-108.
- [12] Y.-K. Huang, Z.-Y. Wen. The sequence of return words of the Fibonacci sequence, *Theoretical Computer Science*. 593 (2015) 106-116.
- [13] Y.-K. Huang, Z.-Y. Wen, Kernel words and gap sequence of the Tribonacci sequence, *Acta Mathematica Scientia (Series B)*. 36.1 (2016) 173-194.
- [14] Y.-K. Huang, Z.-Y. Wen, The numbers of repeated palindromes in the Fibonacci and Tribonacci sequences. *Discrete Applied Mathematics*. 230 (2017), 78-90.
- [15] C. S. Iliopoulos, D. Moore, and W. F. Smyth. A characterization of the squares in a Fibonacci string. *Theoretical Computer Science*. 172 (1997), 281-291.
- [16] M. Lothaire. *Combinatorics on words, in: Encyclopedia of Mathematics and its applications*. Vol.17, Addison-Wesley, Reading, MA, 1983.
- [17] M. Lothaire. *Algebraic combinatorics on words*, Cambridge Univ. Press, Cambridge, 2002.
- [18] H. Mousavi, J. Shallit, Mechanical proofs of properties of the Tribonacci word, in *Combinatorics on Words*. Springer International Publishing. (2014) 170-190.
- [19] K. Saari, Everywhere  $\alpha$ -repetitive sequences and Sturmian words, *European Journal of Combinatorics*. 31.1 (2010) 177-192.
- [20] B. Tan, Z.-Y. Wen, Some properties of the Tribonacci sequence. *European Journal of Combinatorics*. 28 (2007) 1703-1719.
- [21] Z.-X. Wen, Z.-Y. Wen. Some properties of the singular words of the Fibonacci word, *European Journal of Combinatorics*. 15 (1994) 587-598.
- [22] E. Zeckendorf. Représentation des nombres naturels par une somme de nombres de Fibonacci ou de nombres de Lucas. *Bulletin De La Societe Royale Des Sciences De Liège*, 41(1972) 179-182.