

Hamiltonian cycles and paths in hypercubes with disjoint faulty edges

Janusz Dybizbański, Andrzej Szepietowski

Institute of Informatics,

Faculty of Mathematics, Physics and Informatics,

University of Gdańsk, 80-308 Gdańsk, Poland

jdybiz@inf.ug.edu.pl, matszp@inf.ug.edu.pl

Abstract

We consider hypercubes with pairwise disjoint faulty edges. An n -dimensional hypercube Q_n is an undirected graph with 2^n nodes, each labeled with a distinct binary strings of length n . The parity of the vertex is 0 if the number of ones in its labels is even, and is 1 if the number of ones is odd. Two vertices a and b are connected by the edge iff a and b differ in one position. If a and b differ in position i , then we say that the edge (a, b) goes in direction i and we define the parity of the edge as the parity of the end with 0 on the position i . It was already known that Q_n is not Hamiltonian if all edges going in one direction and of the same parity are faulty.

In this paper we show that if $n \geq 4$ then all other hypercubes are Hamiltonian. In other words, every cube Q_n , with $n \geq 4$ and disjoint faulty edges is Hamiltonian if and only if for each direction there are two healthy crossing edges of different parity.

Keywords: Hamiltonian cycle, hypercube, fault tolerance, disjoint faulty edges.

1 Introduction

For definitions and notations we follow [3, 4]. An n -dimensional hypercube (cube), denoted by Q_n , is an undirected graph with 2^n nodes, each labeled with a distinct binary string $b_{n-1} \dots b_1 b_0$, where $b_i \in \{0, 1\}$. A vertex $x = b_{n-1} \dots b_i \dots b_0$ and the vertex $x^{(i)} = b_{n-1} \dots \bar{b}_i \dots b_0$ are connected by an edge along dimension i , where \bar{b}_i is the negation of b_i . The hypercube Q_n is bipartite, the set of nodes is the union of two sets: nodes of parity 0 (the number of ones in their labels is even), and nodes of parity 1 (the number of ones is odd), and each edge connects nodes of different parity. We shall denote by $Par(x)$ the parity of the vertex x .

For any dimension i , there is a partition of Q_n into two subcubes $Q_0^i = \{x \in Q_n : x_i = 0\}$ and $Q_1^i = \{x \in Q_n : x_i = 1\}$. We shall often use simpler notations Q_0 or Q_1 , if the dimension is obvious or irrelevant. For any edge $e = (u, v)$, if $u \in Q_0^i$ and $v \in Q_1^i$, then we say that e is crossing and goes in i dimension. We say that e is of parity 0 (or 1) if u is of parity 0 (or 1).

A path or a cycle is *Hamiltonian* if it visits each node in the cube exactly once. The hypercube is Hamiltonian, i.e., it contains a Hamiltonian cycle. Moreover, it is *laceable*, which means that any two vertices of different parity can be connected by a Hamiltonian path. Note that vertices of the same parity cannot be connected by a Hamiltonian path. An important property of the hypercube is its fault tolerance. It contains Hamiltonian cycles and paths even if some edges or vertices are faulty.

In this paper we consider Hamiltonian cycles and paths in hypercube Q_n with faulty edges (we shall denote the set of faulty edges by F). Tsai et al. [11] showed that if $|F| \leq n - 2$, then Q_n is Hamiltonian. The hypercube Q_n is n -regular, so it is not Hamiltonian if it contains a vertex with incident $n - 1$ faulty edges. We shall call such a set of faulty edges a local trap. Xu et al. [12] proved that if each vertex is incident with at least two healthy edges and $|F| \leq n - 1$, then the cube is still Hamiltonian. Chan and Lee [1] showed that also with $|F| \leq 2n - 5$ faulty edges Q_n is Hamiltonian. Shih et al. [8] observed that the bound $2n - 5$ is optimal, because with $2n - 4$ faulty edges it is possible to build another kind of local trap, namely a (healthy) cycle (u, v, w, x) , where all edges going out of the cycle from u and w are faulty. This kind of trap is called f_4 -cycle in [6]. Szepietowski [9] observed that these two kinds of traps are examples of a more general scheme, namely subgraphs disconnected halfway.

Definition 1. *A proper subgraph $T \subset Q_n$ is disconnected halfway if half of the nodes of T have parity 0 and either (1) all edges joining the nodes of parity 0 in T with nodes outside T , are faulty, or (2) all edges joining the nodes of parity 1 in T with nodes outside T , are faulty.*

Lemma 2. *([9]) The cube Q_n is not Hamiltonian if it has a trap disconnected halfway.*

If Q_n has a node u with $n - 1$ faulty incident edges, then u with its neighbor form a subcube of dimension 1 disconnected halfway. We shall denote it by Q_1 -DHW. If T is a cycle $C_4 = (u, v, w, x)$ (subcube of dimension 2), then we obtain the trap described in [8] and mentioned above. Similarly, in Q_n , with $n \geq 4$, using $4(n - 3)$ faulty edges we can disconnect halfway the 3-dimensional subcube Q_3 . Szepietowski [9] proved that the cube Q_n with $n \geq 5$ and with $2n - 4$ faulty edges is not Hamiltonian only if it contains a Q_1 -DHW or Q_2 -DHW trap. Liu and Wang [7] showed that the same is true for hypercubes with $3n - 8$ faulty edges. The bound $3n - 8$ is sharp, because with $3n - 7$ faulty edges it is possible to build another kind of local trap, namely a (healthy) cycle (u, v, w, x, y, z) , where all edges going out of the cycle from u , w , and y are faulty, see [6, 7, 9]. Li et al. [6] proved that with $n \geq 6$ and with $3n - 7$ faulty edges Q_n is not Hamiltonian only if it contains a Q_1 -DHW, Q_2 -DHW, or C_6 -DHW trap.

In [9], another type of trap was described which is not a set disconnected halfway. Consider a node u that has three neighbors v , w and x , each of degree 2, see Fig. 1. Then, the cube Q_n is not Hamiltonian. We call them claw traps.




Figure 1: A claw trap.

In Q_3 claw consists of three disjoint faulty edges. If $n \geq 4$, the faulty edges in Claw are not disjoint. Each of the trap mentioned above consists of small collection of faulty edges where the number of faulty edges is bounded by a linear function.

In this paper we consider cubes Q_n with pairwise disjoint faulty edges, i.e. for any two faulty edges $e, f \in F$, $e \cap f = \emptyset$. In [3] authors described all nonhamiltonian cubes with $n = 3$ and $n = 4$. From their results it follows that if F consists of pairwise disjoint edges, then the cube Q_3 is not Hamiltonian if and only if it contains a Q_2 -DHW or a Claw trap, and that Q_4 is not Hamiltonian if and only if it contains a Q_3 -DHW trap. It is easy to see that Q_3 is laceable if and only if $|F| \leq 1$. Using computer we checked that Q_4 with disjoint faulty edges is not laceable only if it contains either Q_3 -DHW or six parallel faulty edges.

If the set F consists of disjoint edges, then $|F| \leq 2^{n-1}$ and only one trap of the kind DHW is possible. Namely Q_{n-1} -DHW.

Definition 3. We say that Q_n with $n \geq 3$, has a SCDHW (subcube disconnected halfway) along dimension i , if all crossing edges of parity 0 (or 1) going in dimension i are faulty.

Suppose that F consists of disjoint edges and Q_n with $n \geq 3$, has a SCDHW along dimension i . We may assume that all vertices of parity 0 in Q_0^i are cut off from the vertices in Q_1^i . Then there are faulty edges neither in Q_0^i nor in Q_1^i . However some other edges going in dimension i can be faulty. By Lemma 2, Q_n is not Hamiltonian, hence, it is not laceable. Moreover, if Q_n contains a Hamiltonian path, then one of its ends is of parity 0 and belongs to Q_0^i and the other end is of parity 1 and belongs to Q_1^i , see Lemma 10 in [4].

Definition 4. We say that Q_n has DTBCE (dimension with two balanced crossing edges), if there is a dimension i and two vertices $u, w \in Q_0^i$ of different parity such that: all crossing edges in dimension i , except two $(u, u^{(i)})$ and $(w, w^{(i)})$, are faulty.

Note that it is possible that the vertices u and w (or $u^{(i)}$ and $w^{(i)}$) form a faulty edge. Note also that if F consists of disjoint edges then no other faulty

edges are possible, hence, there is at most one faulty edge in Q_0^i and at most one in Q_1^i . The following lemma is easy to prove,

Lemma 5. *The cube Q_n with DTBCE:*

- (i) *has a Hamiltonian cycle, even if $(u, w) \in F$ and $(u^{(i)}, w^{(i)}) \in F$*
- (ii) *is not laceable. There is neither a HP going from u to w nor a HP going from $u^{(i)}$ to $w^{(i)}$.*

In this paper we show that if $n \geq 4$ and F consists of pairwise disjoint edges, then only the cubes which has SCDHW are not Hamiltonian. In other words Q_n with $n \geq 4$, is Hamiltonian if and only if for each dimension i , there is a healthy crossing edge of parity 0 and a healthy crossing edge of parity 1. Moreover we show that Q_n is not laceable if and only if it contains either SCDHW or DTBCE. In other words Q_n is laceable if and only if for each dimension i , there are at least three healthy edges not of the same parity.

2 Preliminaries

Let f_0 denote the number of faulty edges in Q_0 and f_1 denote the set of faulty edges in Q_1 . By $dist(A, B)$, we denote the distance between the vertices A and B . We shall use abbreviations:

- HP $x \rightarrow y$ to denote a Hamiltonian path from x to y ,
- HP $x \rightarrow_F y$ to denote a Hamiltonian path from x to y passing through one faulty edge,
- $x \not\rightarrow y$ to denote that there are no HP going from x to y ,
- $x \not\rightarrow_F y$ to denote that there are no HP going from x to y and passing through one faulty edge.

Lemma 6. ([11]) *If $n \geq 3$ and $|F| \leq n - 2$, then Q_n is laceable.*

Lemma 7. ([5], see also [11]) *Suppose that x, y , and v are three vertices in Q_n with $n \geq 2$, and that $Par(x) = Par(y) \neq Par(v)$. Then there is a HP $x \rightarrow y$ in $Q_n - v$.*

Lemma 8. ([2]) *Suppose that p, q, r , and s are four different vertices in Q_n and $Par(p) = Par(q) = 0, Par(r) = Par(s) = 1$. Then there is a partition of Q_n into two disjoint paths $p \rightarrow r$ and $q \rightarrow s$.*

Lemma 9. *For any two vertices x, y , with $Par(x) \neq Par(y)$ and an edge $e \neq (x, y)$, there is HP $x \rightarrow y$ passing through e .*

Proof. If neither x nor y is incident to e , then the lemma follows from Lemma 8. If x or y is incident to e , then the lemma follows from Lemma 7. \square

We shall use the following notations. The vertex x is free if it is not incident to any faulty edges. For two dimensions $i \neq j$, by $Q_{00}^{i,j}$ we denote the set $\{x \in Q_n : x_i = 0 \text{ and } x_j = 0\}$. Similarly we define $Q_{01}^{i,j}$, $Q_{10}^{i,j}$, and $Q_{11}^{i,j}$.

3 Partition

Lemma 10. *In Q_n , with $n \geq 5$ and disjoint faulty edges, there is a dimension m such that both Q_0^m and Q_1^m have neither SCDHW nor DTBCE.*

In order to prove the lemma we shall need two lemmas.

Lemma 11. *Suppose that in Q_n with $n \geq 4$, there are two different dimensions, i and k .*

(i) *If there are two vertices $u, v \in Q_{00}^{i,k}$ or $u, v \in Q_{01}^{i,k}$ such that: $\text{Par}(u) \neq \text{Par}(v)$ and the edges $(u, u^{(k)})$ and $(v, v^{(k)})$ are healthy. Then there is no SCDHW along dimension k in Q_0^i .*

(ii) *If there are two vertices $u, v \in Q_{10}^{i,k}$ or $u, v \in Q_{11}^{i,k}$ such that: $\text{Par}(u) \neq \text{Par}(v)$ and the edges $(u, u^{(k)})$ and $(v, v^{(k)})$ are healthy. Then there is no SCDHW along dimension k in Q_1^i .*

Proof. (i) If the subcube Q_0^i has SCDHW along dimension k , then both in $Q_{00}^{i,k}$ and in $Q_{01}^{i,k}$ either all vertices of parity 0 or all vertices of parity 1 are incident to faulty edges going in dimension k . \square

Lemma 12. *Suppose that $n \geq 4$ and the subcube Q_0^i (or Q_1^i) has SCDHW along dimension k , with $k \neq i$. We may assume that $i = 1$ and $k = 2$. Then:*

(i) *for each $j > 2$, neither Q_0^j nor Q_1^j has SCDHW along dimension $m = 1$.*

(ii) *for each $j > 2$, neither Q_0^j nor Q_1^j has SCDHW along dimension m , with $m > 2$ and $m \neq j$.*

(iii) *neither Q_0^2 nor Q_1^2 has SCDHW along dimension m , with $m > 2$.*

Proof. We may assume that every vertex of parity 0 in $Q_{00}^{1,2}$ is joined by a faulty edge with its neighbour in $Q_{01}^{1,2}$ (and every vertex of parity 1 in $Q_{01}^{1,2}$ is joined by a faulty edge with its neighbour in $Q_{00}^{1,2}$). Consider partition of Q_n into Q_0^j and Q_1^j , for any dimension $j \geq 3$, let say $j = 3$, see Fig. 2.

(i) First consider two vertices $u = 000^{n-2}$ and $v = 010^{n-2}$ in $Q_{00}^{1,3}$. They are of different parity and are incident to faulty edges going in dimension 2, so the edges $(u, u^{(1)})$ and $(v, v^{(1)})$ are healthy. Hence, by Lemma 11, Q_0^3 has no SCDHW along dimension $m = 1$. Next consider two vertices $x = 00110^{n-4}$ and $y = 01110^{n-4}$ in $Q_{01}^{1,3}$. They are of different parity and the edges $(x, x^{(1)})$ and $(y, y^{(1)})$ are healthy. Hence, by Lemma 11, Q_1^3 has no SCDHW along dimension $m = 1$.

(ii) First suppose that Q_0^3 has SCDHW along dimension $m > 3$, let say $m = 4$. There are two vertices $u = 00000^{n-4}$ and $v = 01000^{n-4}$ in $Q_{00}^{3,4}$. They are of different parity and the edges $(u, u^{(4)})$ and $(v, v^{(4)})$ are healthy. Hence, by Lemma 11, Q_0^3 has no SCDHW along dimension $m = 4$. In order to prove that Q_1^3 does not have SCDHW along dimension 4, consider two vertices $u = 00110^{n-4}$ and $v = 01110^{n-4}$ in $Q_{11}^{3,4}$. They are of different parity and the edges $(u, u^{(4)})$ and $(v, v^{(4)})$ are healthy.

(iii) Suppose that Q_0^2 has SCDHW along dimension $m > 2$, let say $m = 3$. Then either each vertex of the form $000\dots$ and of parity 0 is incident to faulty edge in dimension 3, or each vertex of the form $001\dots$ and of parity 0 is incident to faulty edges going in dimension 3. This is not possible because all vertices of the form $00\dots$ and of parity 0 are incident to faulty edges going in dimension 2. Similarly, we can show that in Q_1^2 there are no SCDHW along dimension m .

□




Figure 2: Q_4 with SCDHW along dimension 2 in Q_0^1 .

Proof of Lemma 10. First suppose that for some dimension i , the subcube Q_0^i or Q_1^i has DTBCE along some dimension $j \neq i$. We may assume that the subcube Q_0^1 has DTBCE along dimension 2. Consider partition of Q_n into Q_0^2 and Q_1^2 . Then neither Q_0^2 nor Q_1^2 has SCDHW or DTBCE. In order to prove this, observe that in the subcube $Q_{00}^{1,2}$ all vertices of parity 0 except one (there are $2^{n-3} - 1 \geq 3$ such vertices) and all vertices of parity 1 except one are incident to faulty edges in dimension 2, hence they are free inside Q_0^2 . Similarly, in the subcube $Q_{01}^{1,2}$ at least three vertices of parity 0 and at least three vertices of parity 1 are free inside Q_1^2 . Observe that in Q_0^2 :

- DTBCE is not possible, because a subcube with DTBCE has at most two free vertices of parity 0.

- SCDHW along dimension 1 is not possible because in such a case all vertices of parity 0 (or 1) in Q_{00}^{12} (there are $2^{n-3} \geq 4$ such vertices) are incident to faulty edges inside Q_0^2 .
- SCDHW along dimension $k > 2$, let say $k = 3$, is not possible, because in such a case all vertices of the form 000... and parity 0 (or 1) (there are $2^{n-4} \geq 2$ such vertices) are incident to faulty edges inside Q_0^2 .

Similarly we can show that Q_1^2 has neither DTBCE nor SCDHW.

Next suppose that for each dimension i , neither Q_0^i nor Q_1^i has DTBCE, and that for some i , Q_0^i has SCDHW along dimension k with $k \neq i$. We may assume that Q_0^1 has SCDHW along direction 2. Now, consider the partition along direction 2. By Lemma 12, the subcubes Q_0^2 or Q_1^2 may have SCDHW only along direction 1. If they have, then consider partition along direction 3. By Lemma 12, neither Q_0^3 nor Q_1^3 has any SCDHW. \square

4 Main results

Theorem 13. *The cube Q_n with $n \geq 4$ and pairwise disjoint faulty edges, is not Hamiltonian if and only if it has SCDHW.*

Theorem 14. *The Q_n with $n \geq 4$ and disjoint faulty edges is laceable if for every dimension i , there are at least three healthy crossing edges, and they are not of the same parity.*

These two theorems will follow from the lemma:

Lemma 15. *Suppose that in Q_n with $n \geq 4$, the set of faulty edges F consists of pairwise disjoint edges and there is neither SCDHW nor DTBCE. Then:*

(La1) *For any two vertices $A, B \in Q_n$ with $Par(A) \neq Par(B)$, there is a HP $A \rightarrow B$,*

(La2) *For any two vertices $A, B \in Q_n$ with $Par(A) \neq Par(B)$, if $|F| \geq 2$ or $|F| = 1$ and A, B do not form the edge $(A, B) \in F$, then there is a faulty edge e and a HP $A \rightarrow B$ passing through e and not through any other faulty edges.*

Proof. If $|F| \leq 1$, then the theorem follows from Lemmas 9 and 6. So we assume that $|F| \geq 2$. Moreover, without loss of generality we may assume that if A and B form an edge, then this edge is healthy. Indeed, any HP $A \rightarrow B$ (or $A \rightarrow_F B$) does not use the edge (A, B) . We may also assume that $Par(A) = 0$ and $Par(B) = 1$.

We prove the theorem by induction on n . The case $n = 4$ we have checked by computer. We have run the program which checks the all 4-dimensional hypercubes with disjoint faulty edges.

Suppose that $n \geq 5$ and that the theorem is valid for smaller cubes. By Lemma 10, there is a dimension m , let say $m = 1$, such that after partition of Q_n into Q_0^1 and Q_1^1 , neither Q_0^1 nor Q_1^1 has SCDHW or DTBCE. Hence, by the induction hypothesis, both Q_0^1 and Q_1^1 satisfy (La1) and (La2).

(La1).

Case 1. $A \in Q_0$ and $B \in Q_1$. Because Q_n has no SCDHW, there is a healthy crossing edge (u, v) of parity 1. By the induction hypothesis, there is a HP $A \rightarrow u$ in Q_0 and a HP $v \rightarrow B$ in Q_1 . We can build a HP $A \rightarrow B$ in Q_n using these two paths and the edge (u, v) , see Fig. 3.




Figure 3: Case 1.




Figure 4: Subcase 2.1

Case 2. $A, B \in Q_0$.

Subcase 2.1. There is a faulty edge $f \in F$ with $f \in Q_0$. By the induction hypothesis, there is HP $A \rightarrow_F B$ in Q_0 passing through a faulty edge $f' = (u, v)$ (it is possible that $f' \neq f$), and there is HP $u^{(1)} \rightarrow v^{(1)}$ in Q_1 . The crossing edges $(u, u^{(1)})$, $(v, v^{(1)})$ are healthy, because they are incident to the faulty edge (u, v) , see Fig. 4.

Subcase 2.2. There are no faulty edges in Q_0 . Because Q_n has neither SCDHW nor DTBCE, there are two healthy crossing edges $(x, x^{(1)})$, $(y, y^{(1)})$ such that: $x, y \in Q_0$, $Par(x) \neq Par(y)$, and $\{x, y\} \neq \{A, B\}$. We may assume that $Par(A) \neq Par(x)$. By the induction hypothesis, there is HP $x^{(1)} \rightarrow y^{(1)}$ in Q_1 . If $\{A, B\} \cap \{x, y\} = \emptyset$, then by Lemma 8, there is partition of Q_0 into two paths $A \rightarrow x$ and $B \rightarrow y$, see Fig. 5.

If $\{A, B\} \cap \{x, y\} \neq \emptyset$ then we may assume that $A = y$ and $B \neq x$, and by Lemma 7, there is a HP $x \rightarrow B$ in $Q_0 - A$, see Fig. 6.

(La2).

Case 3. $A \in Q_0$ and $B \in Q_1$.

Subcase 3.1. There is a faulty crossing edge (u, v) of parity 1. The Hamiltonian path in Q_n goes from A to u , next crosses to v , and then goes from v to B , see Fig. 7.

Subcase 3.2. All crossing edges of parity 1 are healthy and there is a faulty edge $f = (y, z)$ in Q_0 . In this case we choose a healthy crossing edge (u, v) with $u \in Q_0$, $Par(u) = 1$ and $u \notin f$. By the induction hypothesis, there is a HP $A \rightarrow_F u$ in Q_0 , and there is a HP $v \rightarrow B$ in Q_1 , see Fig. 8.

Subcase 3.3. All crossing edges of parity 1 are healthy and there are faulty edges neither in Q_0 nor in Q_1 . Because $|F| \geq 2$, there are at least two faulty




Figure 5: Subcase 2.2, $\{A, B\} \cap \{x, y\} = \emptyset$




Figure 6: Subcase 2.2, $\{A, B\} \cap \{x, y\} \neq \emptyset$




Figure 7: Subcase 3.1




Figure 8: Subcase 3.2

crossing edges and we can choose an edge $(x, x^{(i)})$ such that $Par(x) = 0$ and $x \neq A$. Take two crossing edges $(y, y^{(i)})$, and $(z, z^{(i)})$ with $y, z \in Q_0$ and $Par(y) = Par(z) = 1$. There is a partition of Q_0 into two disjoint paths $A \rightarrow y$ and $x \rightarrow z$. If $B \neq x^{(i)}$, then by Lemma 8, we use a partition of Q_1 into two disjoint paths $z^{(i)} \rightarrow B$ and $y^{(i)} \rightarrow x^{(i)}$, see Fig. 9. If $B = x^{(i)}$, then by Lemma 7, we use a HP $y^{(i)} \rightarrow z^{(i)}$ in $Q_1 - B$, see Fig. 10.

Case 4. $A, B \in Q_0$

Subcase 4.1. There are faulty crossing edges. By the induction hypothesis, there is a HP $A \rightarrow B$, in Q_0 . There are two neighbours y and z on the path such that the crossing edge $(y, y^{(1)}) \in F$ and the edge $(z, z^{(1)}) \notin F$. There is a HP $y^{(i)} \rightarrow z^{(i)}$ in Q_1 , see Fig. 11.

Subcase 4.2. There are no faulty crossing edges and there is a faulty edge $f \in Q_0$ with $f \neq (A, B)$. By the induction hypothesis, there is a HP $A \rightarrow_F B$ in Q_0 . Choose a healthy edge (x, y) on the path. There is a HP $x^{(i)} \rightarrow y^{(i)}$ in Q_1 , see Fig. 12.




Figure 9: Subcase 3.3. $B \neq x^{(i)}$




Figure 10: Subcase 3.3. $B = x^{(i)}$




Figure 11: Subcase 4.1




Figure 12: Subcase 4.2

Subcase 4.3. There are no faulty crossing edges and no faulty edges in Q_0 . Then there are at least two faulty edges in Q_1 . By the induction hypotheses, there is a HP $A \rightarrow B$ in Q_0 . Choose a (healthy) edge (x, y) on the path. There is a HP $x^{(i)} \rightarrow_F y^{(i)}$ in Q_1 , see Fig. 13. □

Proof of Theorem 13. By Lemma 2, Q_n is not Hamiltonian if it has SCDHW. Suppose now that Q_n is not Hamiltonian, so it is not laceable and by Lemma 15, it has SCDHW or DTBCE. However, by Lemma 5, Q_n is Hamiltonian if it has DTBCE. □

Corollary 16. *If $n \geq 4$, faulty edges F are pairwise disjoint, and $|F| < 2^{n-2}$, then Q_n is Hamiltonian and laceable.*

Corollary 17. *If $n \geq 4$, faulty edges F are pairwise disjoint, and there are two nonparallel faulty edges, then Q_n is Hamiltonian.*

Corollary 18. *If $n \geq 4$, faulty edges F are pairwise disjoint, and there are three faulty edges of three different dimensions, then Q_n laceable.*




Figure 13: Subcase 4.3

References

- [1] M.Y. Chan, S.J. Lee, *On the existence of Hamiltonian circuits in faulty hypercubes*, SIAM Journal on Discrete Mathematics, 4 (1991) 511–527.
- [2] T. Dvorak, *Hamiltonian cycles with prescribed edges in hypercubes*, SIAM Journal on Discrete Mathematics, 19 (2005) 1, 135–144.
- [3] J. Dybizbański, A. Szepietowski, *Hamiltonian cycles in hypercubes with faulty edges*, International Journal of Computer Mathematics: Computer Systems Theory 3 (2018) 244–261.
- [4] J. Dybizbański, A. Szepietowski, *Hamiltonian paths in hypercubes with local traps*, Information Sciences, 375 (2017) 258–270
- [5] M. Lewinter, W. Widulski, *Hyper-Hamilton Laceable and Caterpillar-Spannable Product Graphs*, Computers & Mathematics with Applications, 34 (1997) 99–104.
- [6] J. Li, D. Liu, X. Gao, *Hamiltonian cycles in hypercubes with more faulty edges*, International Journal of Computer Mathematics, 94 (2017) 1155–1171.
- [7] J.-J. Liu, Y.-L. Wang, *Hamiltonian cycles in hypercubes with faulty edges*, Information Sciences, 256 (2014) 225–233.
- [8] L.-M. Shih, J.J.M. Tan, L.H. Hsu, *Edge-bipancyclicity of conditional faulty hypercubes*, Information Processing Letters 105 (2007) 20–25.
- [9] A. Szepietowski, *Hamiltonian cycles in hypercubes with $2n - 4$ faulty edges*, Information Sciences, 215 (2012) 75–82.
- [10] C.-H. Tsai, *Linear array and ring embeddings in conditional faulty hypercubes*, Theoretical Computer Science, 314 (2004) 431–443.

- [11] C.-H. Tsai, J.J.M. Tan, T. Liang, L.-H. Hsu, *Fault-tolerant Hamiltonian laceability of hypercubes*, Information Processing Letters 83 (2002) 301–306.
- [12] J.-M. Xu, Z.-Z. Du, M. Xu, *Edge-fault-tolerant edge-bipancyclicity of hypercubes*, Information Processing Letters 96 (2005) 146–150.