

A REMARK ON INFINITE INITIAL VALUES FOR QUASILINEAR PARABOLIC EQUATIONS

PETER LINDQVIST AND MIKKO PARVIAINEN

ABSTRACT. We study the possibility of prescribing infinite initial values for solutions of the Evolutionary p -Laplace Equation in the fast diffusion case $p > 2$. This expository note has been extracted from our previous work. When infinite values are prescribed on the whole initial surface, such solutions can exist only if the domain is a space-time cylinder.

We are interested in a violent initial value problem for the Evolutionary p -Laplace Equation

$$\frac{\partial u}{\partial t} = \operatorname{div}(|\nabla u|^{p-2} \nabla u), \quad p > 2,$$

in a cylindrical domain $\Omega_T = \Omega \times (0, T)$, where Ω is an open connected domain in the Euclidean n -dimensional space \mathbb{R}^n . We study the slow diffusion case $p > 2$. For $p = 2$, the equation is the heat equation

$$\frac{\partial u}{\partial t} = \Delta u,$$

but the phenomenon we describe is impossible for the heat equation.

The objective of our note is the possibility of prescribing *infinite* initial values in some subset E of Ω , i.e.

$$\lim_{t \rightarrow 0^+} u(x, t) = \infty \quad \text{at every } x \in E. \quad (1)$$

We assume that $u \geq 0$ and do not mind the boundary values at the lateral boundary $\partial\Omega \times [0, T]$, since they now play a minor role in comparison to the violent effects of the ∞ initial values. The solutions are defined as weak solutions belonging to the Sobolev space $L_{\text{loc}}^p(0, T; W_{\text{loc}}^{1,p}(\Omega))$ or even to $L_{\text{loc}}^p(0, T; W^{1,p}(\Omega))$, and they are continuous in Ω_T . See for example [DiB93] or [KLP16, Definition 5].

We have detected a strange phenomenon, which can be described as follows.

Date: November 29, 2018.

2010 Mathematics Subject Classification. 35J92, 35J62.

Key words and phrases. Friendly giant, parabolic p -Laplace equation.

- If the n -dimensional Lebesgue measure $|E| > 0$ of E is strictly positive, then $E = \Omega$.
- There exist solutions, if $E = \Omega$ and Ω is bounded.
- There is no solution, if $\Omega = \mathbb{R}^n$ and $|E| > 0$.

To spell it out, the condition that

$$\liminf_{t \rightarrow 0^+} u(x_0, t) < \infty$$

at a point x_0 in Ω is incompatible with the condition that (1) holds in a set E of positive n -dimensional measure. In the case when Ω is the whole space \mathbb{R}^n , the *global* restriction that $u \geq 0$ is decisive.¹

We remark that if Ω is bounded and regular, then there exists a separable solution of the form

$$V(x, t) = t^{-\frac{1}{p-2}} U(x) \quad (2)$$

where $U \in C(\overline{\Omega})$, $U > 0$ in Ω and $U|_{\partial\Omega} = 0$. The function U solves the auxiliary equation

$$\operatorname{div}(|\nabla U|^{p-2} \nabla U) + \frac{1}{p-2} U = 0.$$

The function V can serve as a minorant: every other solution $u \geq 0$ with $E = \Omega$, the lateral boundary values of which are not necessarily zero, has to grow faster than V . That is

$$u(x, t) \geq t^{-\frac{1}{p-2}} U(x) \quad \text{in } \Omega_T.$$

Proposition 1. *If a weak solution $u \geq 0$, satisfies*

$$\lim_{t \rightarrow 0^+} u(x, t) = \infty$$

at every point x in a set of positive n -dimensional measure, then the limit is infinite at each point $x \in \Omega$. Moreover,

$$\liminf_{(x,t) \rightarrow (x_0,0)} \left[t^{\frac{1}{p-2}} u(x, t) \right] > 0 \quad (3)$$

at every point x_0 in Ω , and of course $(x, t) \in \Omega_T$.

The case $|E| = 0$ remains. For example, the celebrated Barenblatt solution for $t > 0$

$$\mathcal{B}(x, t) = t^{-n/\lambda} \left\{ C - \frac{p-2}{p} \lambda^{\frac{1}{1-p}} \left(\frac{|x|}{t^{1/\lambda}} \right)^{\frac{p}{p-1}} \right\}_+^{\frac{p-1}{p-2}}, \quad (4)$$

¹Otherwise one can construct an odd periodic extension of U , first defined in the box $\Omega = [0, 1] \times \cdots \times [0, 1]$, to the whole \mathbb{R}^n . The corresponding V will do as an example.

where $\lambda = n(p - 2) + p$, the constant $C > 0$ can be chosen freely, and $\{\cdot\}_+$ means $\max\{\cdot, 0\}$, exhibits that it is possible that E consists of a single point: $E = \{(0, 0)\}$. We have that $\mathcal{B}(x, 0) = c\delta_0$ (Dirac's delta) in the sense that

$$\lim_{t \rightarrow 0^+} \int_{\mathbb{R}^n} \mathcal{B}(x, t) \phi(x) dx = c\phi(0)$$

for all test functions $\phi \in C_0^\infty(\mathbb{R}^n)$. In general, the question about whether there exists a Radon measure μ such that $'u = \mu'$ as $t = 0$ has drawn much attention, see for example [AC83, BCP84, DK84] in the case of the porous medium equation. It is clear that no σ -finite measure will do in the case $E = \Omega$, since (3) holds. However, if $|E| = 0$ this is possible. The initial values can even be prescribed as a Radon measure, see [DH89] and [KP09].

Finally, we mention a surprising result, when the domain is of the form

$$\Omega^\Psi = \{(x, t) : x \in \Omega, \Psi(x) < t < T\}$$

where Ψ is a given smooth function. Assume now that

$$\lim_{(y, \tau) \rightarrow (x, \Psi(x))} u(x, t) = +\infty$$

at every point $x \in \Omega$, where of course (y, τ) is in Ω^Ψ . Indeed, such a solution does not exist, except when $\Psi(x) \equiv \text{constant}$. Then we are back to the case $\Omega \times (0, T)$. *Infinite initial values cannot be prescribed on a whole surface in the (x, t) -space which is not a timeslice $t = \text{constant}$.* It is arresting, that not even a domain restricted by a hyperplane

$$t_0 + a_1 x_1 + \dots + a_n x_n < t < T, \quad |x| < 1,$$

is possible, except when $a_1 = \dots = a_n = 0$.

For the proof we let $x_0 \in \Omega$ be a point such that a small ball $B = B(x_0, r)$ is divided by the smooth level surface $C_0 = \{x \in \Omega : \Psi(x) = \Psi(x_0)\}$ into two pieces, say $B^+ = \{x \in B : \Psi(x) > \Psi(x_0)\}$ and $B^- = \{x \in B : \Psi(x) < \Psi(x_0)\}$. Then let $\omega = \omega(x)$ be the solution of the elliptic problem

$$\Delta_p \omega = 0 \quad \text{in } B^- \quad \text{and} \quad \omega = \begin{cases} 0 & \text{in } B^- \cap \partial B \\ 1 & \text{in } C_0 \cap B. \end{cases}$$

(If desired, we may round off the boundary values so that the value 1 is prescribed in a smaller set, yet one of positive area). On the parabolic boundary of the domain

$$\Psi(x) < t < \Psi(x_0), \quad x \in B^-$$

we have $u(x, t) \geq k\omega(x)$ for any constant k . By the comparison principle, see [KL96] and [BBGP15, Theorem 2.4], the inequality $u(x, t) \geq k\omega(x)$ holds in this domain. Sending k to ∞ we obtain that $u \equiv \infty$ in an open set of \mathbb{R}^{n+1} . This contradicts the continuity of u .

For the Porous medium equation

$$\frac{\partial u}{\partial t} = \Delta(u^m), \quad m > 1,$$

a similar theory is valid. The separable solution corresponding to (2) is called 'the friendly giant' by Dahlberg and Kenig [DK88], see also Vazquez [Váz07, p. 111].

As mentioned above, most of this note has been extracted from our work [KLP16]. For more recent related works, see [KL16, KLLP, PV].

REFERENCES

- [AC83] D. G. Aronson and L. A. Caffarelli. The initial trace of a solution of the porous medium equation. *Trans. Amer. Math. Soc.*, 280(1):351–366, 1983.
- [BBGP15] A. Björn, J. Björn, U. Gianazza, and M. Parviainen. Boundary regularity for degenerate and singular parabolic equations. *Calc. Var. Partial Differential Equations*, 52(3-4):797–827, 2015.
- [BCP84] P. Bénilan, M. G. Crandall, and M. Pierre. Solutions of the porous medium equation in \mathbf{R}^N under optimal conditions on initial values. *Indiana Univ. Math. J.*, 33(1):51–87, 1984.
- [DH89] E. DiBenedetto and M. A. Herrero. On the Cauchy problem and initial traces for a degenerate parabolic equation. *Trans. Amer. Math. Soc.*, 314(1):187–224, 1989.
- [DiB93] E. DiBenedetto. *Degenerate Parabolic Equations*. Universitext. Springer-Verlag, New York, 1993.
- [DK84] B. E. J. Dahlberg and C. E. Kenig. Nonnegative solutions of the porous medium equation. *Comm. Partial Differential Equations*, 9(5):409–437, 1984.
- [DK88] B. E. J. Dahlberg and C. E. Kenig. Nonnegative solutions of the initial-Dirichlet problem for generalized porous medium equations in cylinders. *J. Amer. Math. Soc.*, 1(2):401–412, 1988.
- [KL96] T. Kilpeläinen and P. Lindqvist. On the Dirichlet boundary value problem for a degenerate parabolic equation. *SIAM J. Math. Anal.*, 27(3):661–683, 1996.
- [KL16] J. Kinnunen and P. Lindqvist. Unbounded supersolutions of some quasi-linear parabolic equations: a dichotomy. *Nonlinear Anal.*, 131:229–242, 2016.
- [KLLP] J. Kinnunen, P. Lehtelä, P. Lindqvist, and M. Parviainen. Supercaloric functions for the porous medium equation. *arXiv preprint arXiv:1801.04121*. To appear in *J. Evol. Equ.*

- [KLP16] T. Kuusi, P. Lindqvist, and M. Parviainen. Shadows of infinities. *Ann. Mat. Pura Appl. (4)*, 195(4):1185–1206, 2016.
- [KP09] T. Kuusi and M. Parviainen. Existence for a degenerate Cauchy problem. *Manuscripta Math.*, 128(2):213–249, 2009.
- [PV] M. Parviainen and J. L. Vázquez. Equivalence between radial solutions of different parabolic gradient-diffusion equations and applications. *arXiv preprint arXiv:1801.00613*.
- [Váz07] J. L. Vázquez. *The Porous Medium Equation*. Oxford Math. Monogr. The Clarendon Press Oxford University Press, Oxford, 2007. Mathematical theory.

DEPARTMENT OF MATHEMATICS AND STATISTICS, NORWEGIAN UNIVERSITY
OF SCIENCE AND TECHNOLOGY, NO-7491 TRONDHEIM, NORWAY

E-mail address: lqvist@math.ntnu.no

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ,
PO BOX 35, FI-40014 JYVÄSKYLÄ, FINLAND

E-mail address: mikko.j.parviainen@jyu.fi