Simple rules of functional integration in the Schwarzian theory: SYK correlators

Vladimir V. Belokurov^{*a,b*} and Evgeniy T. Shavgulidze^{*a*}

^bInstitute for Nuclear Research of the Russian Academy of Sciences, 60th October Anniversary Prospect 7a, Moscow, 117312, Russia

E-mail: vvbelokurov@yandex.ru, shavgulidze@bk.ru

ABSTRACT: We derive the general rules of functional integration in the theories of the Schwarzian type, and evaluate explicitly the functional integrals assigning correlation functions in the SYK model.

KEYWORDS: SYK model, Schwarzian theory, correlation functions, quasi-invariant measures

ARXIV EPRINT: 1811.11863v2

^aLomonosov Moscow State University,

Leninskie gory 1, Moscow, 119991, Russia

Contents

1	Introduction	1
2	Quasi-invariance of the measure as a key to functional integration	3
3	Splitting the interval multiplies functional integrals	7
4	SYK correlation functions	10
5	Concluding remarks	11
\mathbf{A}	Convolution of the \mathcal{E} functions	12
в	Correlation functions in terms of ordinary integrals	13
\mathbf{C}	Functional integrals over the space $Diff_+^1(\mathbf{R})$	15

1 Introduction

The appearance of a dynamics common to different physical models in different fields may indicate the existence of a hidden symmetry behind it. It has recently become clear that a quantum mechanical model of Majorana fermions with a random interaction (Sachdev-Ye-Kitaev model), the holographic description of the Jackiw-Teitelboim dilaton gravity, open string theory and some other models lead to the same effective theory with the Schwarzian action

$$I_{Schw} = -\frac{1}{\sigma^2} \int_{S^1} \left[\mathcal{S}_{\varphi}(t) + 2\pi^2 \left(\varphi'(t) \right)^2 \right] dt , \qquad (1.1)$$

where S^1 is the unit circle, and

$$S_{\varphi}(t) = \left(\frac{\varphi''(t)}{\varphi'(t)}\right)' - \frac{1}{2} \left(\frac{\varphi''(t)}{\varphi'(t)}\right)^2$$
(1.2)

is the Schwarzian derivative.

The list of the papers includes (but is not confined to) [1] - [46], and it will not rest here.

In some approximations, these physical models appear to be reparametrization invariant. And $SL(2, \mathbf{R})$ -invariant action (1.1) inherits the emergent reparametrization symmetry.

Let us remind briefly how the Schwarzian action originates in the SYK model and in 2D JT gravity. Functional integrals in the SYK model can be rewritten in terms of the bilocal bosonic functions \tilde{G} and $\tilde{\Sigma}$ with the effective bosonic action $A_{eff}(\tilde{G}, \tilde{\Sigma})$

$$\int F(\tilde{G},\,\tilde{\Sigma})\,\exp\{-A_{eff}(\tilde{G},\,\tilde{\Sigma})\}\,d\tilde{G}\,d\tilde{\Sigma}\,.$$

The saddle points of the infrared essential part of the effective bosonic action A_{eff}

$$G_f(t_1, t_2) = \frac{(f'(t_1)f'(t_2))^{\frac{1}{4}}}{|f(t_2) - f(t_1)|^{\frac{1}{2}}}$$
(1.3)

form the space \mathcal{F} of one-time differentiable functions f on the unit circle.

The rest (infrared inessential) part of the effective bosonic action A_{eff} results (for three-time differentiable f) in the factor

$$\exp\left\{\frac{1}{\sigma^2} \int\limits_{S^1} \left[\left(\frac{f''(t)}{f'(t)}\right)' - \frac{1}{2} \left(\frac{f''(t)}{f'(t)}\right)^2 \right] dt \right\}$$

in the integrand.

Thus the averaged saddle point solution (or two-point SYK correlation function) has the form

$$\langle G_f(t_1, t_2) \rangle = \int_{\mathcal{F}} \frac{(f'(t_1)f'(t_2))^{\frac{1}{4}}}{|f(t_2) - f(t_1)|^{\frac{1}{2}}} \exp\left\{\frac{1}{\sigma^2} \int_{S^1} \mathcal{S}_f(t) dt\right\} df.$$
 (1.4)

The action of Jackiw-Teitelboim gravity is

$$A_{JT} = Topological \ term - \alpha \left(\int d^2 x \sqrt{g} \,\phi \left(R+2\right) + 2 \int_{bdy} \sqrt{\gamma} \,\phi K \right) \,, \tag{1.5}$$

where ϕ is the dilaton field and K is the extrinsic curvature. While the equation of motion for dilaton demands the space to be AdS_2 , variations of the closed boundary curve with the proper boundary condition for ϕ transform the last term in the action into

$$\frac{1}{\sigma^2} \int\limits_{S^1} \mathcal{S}_f(t) \, dt \,,$$

where t is a parameter of the closed boundary curve.

Functional integrals in these theories are the integrals with the measure

$$\mu_{\sigma}(d\varphi) = \exp\left\{\frac{1}{\sigma^2} \int_{S^1} S_{\varphi}(t) dt\right\} d\varphi$$
(1.6)

over the group of diffeomorphisms $Diff_+^1(S^1)$.

In [48], [49], we explicitly evaluated some nontrivial functional integrals of this type using the quasi-invariance of the measure μ .

Although one can use the invariance of the Schwarzian theory to link it to another theory where the corresponding calculations are much simpler than in the original one [23] - [27], [50], [51], we consider it essential to elaborate a special technique of functional integration in the Schwarzian theory.

Here, we would like to emphasize that there is a significant difference between functional integration over the group $Diff_+^1(S^1)$ and that over the group $Diff_+^1(\mathbf{R})$.

The group $SL(2, \mathbf{R})$ is turned into the subgroup of $Diff^1_+(S^1)$ by linear-fractional transformation

$$\exp\{i2\pi\psi\} = \frac{\alpha \exp\{i2\pi\phi\} - \beta}{\bar{\beta}\exp\{i2\pi\phi\} - \bar{\alpha}}, \qquad |\alpha|^2 - |\beta|^2 = 1.$$

The measure (1.6) is $SL(2, \mathbf{R})$ invariant.

Therefore to obtain finite results in case of $SL(2, \mathbf{R})$ invariant integrands, we should integrate over the quotient space $Diff_{+}^{1}(S^{1})/SL(2, \mathbf{R})$.

However $SL(2, \mathbf{R})$ is not a subgroup of the group $Diff_+^1(\mathbf{R})$. For example, the transformation $g_a \circ f = g_a(f) = (f+a)^{-1}$ $(g_a \in SL(2, \mathbf{R}), f \in Diff_+^1(\mathbf{R}))$ removes $g_a \circ f$ from $Diff_+^1(\mathbf{R})$.

At the same time, the subgroup P = AN of $SL(2, \mathbf{R})$ consisting of transformations $f \to af + b$ (see [52] Ch. III, par. 1) is a subgroup of $Diff_+^1(\mathbf{R})$. The group P is noncompact, and for invariant integrands, one should factorize the integration space, that is one should consider the functional integrals over the quotient space $Diff_+^1(\mathbf{R})/P$.

Thus it is quite natural that the functional integrals over different integration spaces give different results for the same integrand and determine different quantum theories.

In this paper, we complete the elaboration of the functional integrals calculus with the measure (1.6).

In section 2, we review and refine the approach to functional integration based on the quasi-invariance of the measure, and define functional integration over the quotient space $Diff_{+}^{1}(S^{1})/SL(2, \mathbf{R})$.

In section 3, we derive the simple universal rules of functional integration in the theories of the Schwarzian type and demonstrate how the functional integrals are reduced to ordinary multiple integrals. Although the method is developed for functional integrals over the space $Diff_{+}^{1}(S^{1})/SL(2, \mathbf{R})$, it can also be modified to evaluate functional integrals over the space $Diff_{+}^{1}(\mathbf{R})/P$.

In section 4 and in appendices B, C, we evaluate the functional integrals assigning correlation functions in the corresponding models.

In section 5, we discuss the difference between the results in these two cases and give the concluding remarks.

2 Quasi-invariance of the measure as a key to functional integration

Invariant measures analogous to the Haar measure on finite-dimensional groups do not exist for the noncompact groups H [57]. However, there can exist measures that are

quasi-invariant with respect to the action of a more smooth subgroup $G \subset H$. The quasiinvariance means that under the action of the subgroup G the measure transforms to itself multiplied by a function $\mathcal{R}_g(h)$ (the Radon-Nikodim derivative) parametrized by the elements of the subgroup $g \in G$.

The measure (1.6) on the group of diffeomorphisms $Diff_+^1(S^1)$ is quasi-invariant with respect to the action of the subgroup $Diff_+^3(S^1)$ [53]-[55].

To evaluate functional integrals over the group $Diff_+^1(S^1)$, it is convenient to represent them as integrals over the group $Diff_+^1([0,1])$ with the ends of the interval [0, 1] glued $(\varphi'(0) = \varphi'(1))$. After fixing a point t = 0 on the circle of unit length S^1 , the integral $\int_{S^1} S^1$ is

written as \int_{0}^{1} .

The measure on the group of diffeomorphisms of the interval $Diff_+^1([0,1])$ ($\varphi(0) = 0, \varphi(1) = 1, \varphi'(t) > 0$) has the form [53]-[55]

$$\mu_{\sigma}(d\varphi) = \frac{1}{\sqrt{\varphi'(0)\varphi'(1)}} \exp\left\{\frac{1}{\sigma^2} \left[\frac{\varphi''(0)}{\varphi'(0)} - \frac{\varphi''(1)}{\varphi'(1)}\right]\right\} \exp\left\{\frac{1}{\sigma^2} \int_0^1 \mathcal{S}_{\varphi}(t) \, dt\right\} d\varphi \,. \tag{2.1}$$

Now, the integral over $Diff_{+}^{1}(S^{1})$ turns into the the integral over $Diff_{+}^{1}([0,1])$ as follows [49]:

$$\frac{1}{\sqrt{2\pi\sigma}} \int_{Diff^1_+(S^1)} F(\varphi)\mu_\sigma(d\varphi) = \int_{Diff^1_+([0,1])} \delta\left(\frac{\varphi'(1)}{\varphi'(0)} - 1\right) F(\varphi)\mu_\sigma(d\varphi) \,. \tag{2.2}$$

The measure $\mu_{\sigma}(d\varphi)$ is generated by the Wiener measure under some special substitution of variables [53]-[55] (see also [56]). Namely, if we consider a continuous function on the interval [0, 1] $\xi(t)$ satisfying the boundary condition $\xi(0) = 0$ ($\xi \in C_0([0, 1])$), then under the substitution

$$\varphi(t) = \frac{\int_{0}^{t} e^{\xi(\tau)} d\tau}{\int_{0}^{1} e^{\xi(\eta)} d\eta}, \qquad \xi(t) = \log \varphi'(t) - \log \varphi'(0), \qquad (2.3)$$

the measure $\mu_{\sigma}(d\varphi)$ on the group $Diff^1_+([0, 1])$ turns into the Wiener measure $w_{\sigma}(d\xi)$ on $C_0([0, 1])$. In this way, we get the following equality of functional integrals

$$\int_{Diff_{+}^{1}([0,1])} F(\varphi) \, \mu_{\sigma}(d\varphi) = \int_{C_{0}([0,1])} F(\varphi(\xi)) \, w_{\sigma}(d\xi) \,.$$
(2.4)

After the substitution $f(t) = -\cot \pi \varphi(t)$ in (1.4), $\langle G_f(t_1, t_2) \rangle$ is rewritten in the form

$$< G(t_1, t_2) >= \sqrt{2\pi}\sigma \int_{Diff^1_+([0, 1])} \delta\left(\frac{\varphi'(1)}{\varphi'(0)} - 1\right)$$

$$\times \frac{\left(\varphi'(t_1)\varphi'(t_2)\right)^{\frac{1}{4}}}{\left|\sin\left[\pi\varphi(t_2) - \pi\varphi(t_1)\right]\right|^{\frac{1}{2}}} \exp\left\{\frac{2\pi^2}{\sigma^2} \int_0^1 \left(\varphi'(t)\right)^2 dt\right\} \mu_{\sigma}(d\varphi) \,. \tag{2.5}$$

The quasi-invariance of the measure (2.1) with respect to the action of the subgroup $Diff^3_+([0,\,1])$ gives

$$\int_{Diff_{+}^{1}([0,1])} F(\varphi)\mu_{\sigma}(d\varphi) = \frac{1}{\sqrt{g'(0)g'(1)}} \int_{Diff_{+}^{1}([0,1])} F(g(\varphi))$$

$$\times \exp\left\{\frac{1}{\sigma^{2}} \left[\frac{g''(0)}{g'(0)}\varphi'(0) - \frac{g''(1)}{g'(1)}\varphi'(1)\right] + \frac{1}{\sigma^{2}} \int_{0}^{1} \mathcal{S}_{g}\left(\varphi(t)\right) \left(\varphi'(t)\right)^{2} dt\right\} \mu_{\sigma}(d\varphi). \quad (2.6)$$

In what follows, we assume the function g to be

$$g(t) = g_{\alpha}(t) = \frac{1}{2} \left[\frac{1}{\tan \frac{\alpha}{2}} \tan \left(\alpha(t - \frac{1}{2}) \right) + 1 \right].$$
(2.7)

In this case,

$$g'_{\alpha}(0) = g'_{\alpha}(1) = \frac{\alpha}{\sin \alpha}, \quad -\frac{g''_{\alpha}(0)}{g'_{\alpha}(0)} = \frac{g''_{\alpha}(1)}{g'_{\alpha}(1)} = 2\alpha \tan \frac{\alpha}{2}, \quad \mathcal{S}_{g_{\alpha}}(t) = 2\alpha^2, \quad (2.8)$$

and the equation (2.6) looks like:

$$\int_{Diff^{1}([0,1])} F(g_{\alpha}(\varphi)) \exp\left\{-\frac{2\alpha}{\sigma^{2}} \tan\frac{\alpha}{2} \left(\varphi'(0) + \varphi'(1)\right)\right\} \exp\left\{\frac{2\alpha^{2}}{\sigma^{2}} \int_{0}^{1} \left(\varphi'(t)\right)^{2} dt\right\} \mu_{\sigma}(d\varphi)$$

$$= \frac{\alpha}{\sin \alpha} \int_{Diff^{1}([0,1])} F(\varphi) \,\mu_{\sigma}(d\varphi) \,.$$
(2.9)

Denote

$$\Psi(\varphi) = \exp\left\{-\frac{2\alpha}{\sigma^2}\tan\frac{\alpha}{2}\left(\varphi'(0) + \varphi'(1)\right)\right\} F(g_\alpha(\varphi)).$$

Then

$$F(\varphi) = \exp\left\{\frac{2\alpha}{\sigma^2}\tan\frac{\alpha}{2}\left(\left(g_{\alpha}^{-1}(\varphi)\right)'(0) + \left(g_{\alpha}^{-1}(\varphi)\right)'(1)\right)\right\}\Psi\left(\left(g_{\alpha}^{-1}(\varphi)\right)\right).$$
(2.10)

Thus the integral

$$J^{\alpha} = \int_{Diff^{1}([0,1])} \Psi(\varphi) \exp\left\{\frac{2\alpha^{2}}{\sigma^{2}} \int_{0}^{1} \left(\varphi'(t)\right)^{2} dt\right\} \mu_{\sigma}(d\varphi)$$
(2.11)

is transformed into

$$J^{\alpha} = \frac{\alpha}{\sin \alpha}$$

$$\times \int_{Diff^{1}([0,1])} \exp\left\{\frac{2\alpha}{\sigma^{2}} \tan\frac{\alpha}{2} \left(\left(g_{\alpha}^{-1}(\varphi)\right)'(0) + \left(g_{\alpha}^{-1}(\varphi)\right)'(1)\right)\right\} \Psi\left(\left(g_{\alpha}^{-1}(\varphi)\right)\right) \mu_{\sigma}(d\varphi) \,.$$

$$(2.12)$$

For the function g_{α} given by (2.7), the inverse function is

$$\left(g_{\alpha}^{-1}(\varphi)\right)(t) = \frac{1}{\alpha} \arctan\left[\tan\frac{\alpha}{2}\left(2\varphi(t)-1\right)\right] + \frac{1}{2},$$

and

$$\left(g_{\alpha}^{-1}(\varphi)\right)'(0) = \frac{\sin\alpha}{\alpha}\varphi'(0), \qquad \left(g_{\alpha}^{-1}(\varphi)\right)'(1) = \frac{\sin\alpha}{\alpha}\varphi'(1),$$
$$\exp\left\{\frac{2\alpha}{\sigma^2}\tan\frac{\alpha}{2}\left(\left(g_{\alpha}^{-1}(\varphi)\right)'(0) + \left(g_{\alpha}^{-1}(\varphi)\right)'(1)\right)\right\} = \exp\left\{\frac{4\sin^2\frac{\alpha}{2}}{\sigma^2}\left(\varphi'(0) + \varphi'(1)\right)\right\}$$

For the factor

$$\frac{(\varphi'(t_1)\varphi'(t_2))^{\frac{1}{4}}}{\left|\sin\left[\pi\varphi(t_2)-\pi\varphi(t_1)\right]\right|^{\frac{1}{2}}}$$

in the integrand of (2.5), we have

$$\lim_{\alpha \to \pi \to 0} \frac{\left(\left(g_{\alpha}^{-1}(\varphi) \right)'(t_1) \left(g_{\alpha}^{-1}(\varphi) \right)'(t_2) \right)^{\frac{1}{4}}}{\left| \sin \left[\pi \left(g_{\alpha}^{-1}(\varphi) \right)(t_2) - \pi \left(g_{\alpha}^{-1}(\varphi) \right)(t_1) \right] \right|^{\frac{1}{2}}} = \pi^{-\frac{1}{2}} \frac{\left(\varphi'(t_1) \varphi'(t_2) \right)^{\frac{1}{4}}}{\left| \varphi(t_2) - \varphi(t_1) \right|^{\frac{1}{2}}}$$
(2.13)

Now the functional integral for the two-point SYK correlation function (2.5) is transformed into

$$< G_{2}(t_{1}, t_{2}) >_{SYK}^{\alpha} = \sqrt{2}\sigma \frac{\alpha}{\sin \alpha}$$

$$\times \int_{Diff_{+}^{1}([0, 1])} \delta \left(\frac{\varphi'(1)}{\varphi'(0)} - 1 \right) \frac{(\varphi'(t_{1})\varphi'(t_{2}))^{\frac{1}{4}}}{|\varphi(t_{2}) - \varphi(t_{1})|^{\frac{1}{2}}} \exp \left\{ \frac{4}{\sigma^{2}} \left(\varphi'(0) + \varphi'(1) \right) \right\} \mu_{\sigma}(d\varphi) + O(1).$$

$$(2.14)$$

Note that the integral (2.14) is invariant under the shifts of the variable t (that is, under the choice of the point on the S^1 corresponding to the left end of the interval [0, 1]):

$$< G_2(t_1, t_2) >^{\alpha}_{SYK} = < G_2(0, t_2 - t_1) >^{\alpha}_{SYK}$$
.

Generally speaking, the functional integrals (2.12) converge for $0 < \alpha < \pi$, and diverge for $\alpha = \pi$. The integral (2.14) is a typical example. The point is that the Schwarzian action is invariant under the noncompact group of linear-fractional transformations. To get the finite results for functional integrals in the Schwarzian theory, one should factor the infinite input of this noncompact group out.

Technically, it is convenient to exclude the input of the $SL(2, \mathbf{R})$ group that is a subgroup of $Diff_{+}^{1}(S^{1})$, or to integrate over the quotient space $Diff_{+}^{1}(S^{1})/SL(2, \mathbf{R})$.

To this end, we propose first to evaluate regularized ($\alpha < \pi$) functional integrals over the group $Diff^{1}([0,1])$ (with the ends glued) and then normalize them to the corresponding integrals over the group $SL(2, \mathbf{R})$. Thus, we define the renormalized functional integral J^R as the limit

$$J^{R} = \lim_{\alpha \to \pi - 0} \frac{J^{\alpha}}{\int_{SL(2,\mathbf{R})} \Psi(\varphi_{z}) \exp\left\{\frac{2\alpha^{2}}{\sigma^{2}} \int_{0}^{1} (\varphi_{z}'(t))^{2} dt\right\} d\nu_{H}}.$$
 (2.15)

Here, J^{α} is given by (2.11), $\varphi_z \in SL(2, \mathbf{R})$ and $d\nu_H$ is the invariant Haar measure on $SL(2, \mathbf{R})$.

The factor

$$\frac{\left(\varphi'(t_1)\varphi'(t_2)\right)^{\frac{1}{4}}}{\left|\varphi(t_2)-\varphi(t_1)\right|^{\frac{1}{2}}}$$

in the integrand of (2.14) is $SL(2, \mathbf{R})$ invariant.

Therefore, for renormalized SYK correlation functions we have

$$\langle G \rangle_{SYK}^{R} = \lim_{\alpha \to \pi - 0} \frac{\langle G \rangle_{SYK}^{\alpha}}{V_{SL(2,\mathbf{R})}^{\alpha}},$$

$$(2.16)$$

where $V_{SL(2,\mathbf{R})}^{\alpha}$ is the regularized volume of the group $SL(2,\mathbf{R})$ [49]:

$$V_{SL(2,\mathbf{R})}^{\alpha} = \int_{SL(2,\mathbf{R})} \exp\left\{-\frac{2\left[\pi^{2} - \alpha^{2}\right]}{\sigma^{2}} \int_{0}^{1} (\varphi'(t))^{2} dt\right\} d\nu_{H} = \frac{\pi\sigma^{2}}{\pi^{2} - \alpha^{2}} \exp\left\{-\frac{2\left(\pi^{2} - \alpha^{2}\right)}{\sigma^{2}}\right\}$$

Thus, the renormalization excludes the singularity $(\pi - \alpha)^{-1}$ in the functional integrals (2.12), and we get the finite results for SYK correlation functions.

3 Splitting the interval multiplies functional integrals

In the case when the functional F in (2.4) has the special form

$$F(\varphi) = \Phi(\varphi(t_1); \, \varphi'(0), \, \varphi'(t_1), \, \varphi'(1)) \,,$$

the equation (2.4) looks like

$$\int_{Diff^{1}_{+}([0,1])} \Phi(\varphi(t_{1}); \varphi'(0), \varphi'(t_{1}), \varphi'(1)) \mu_{\sigma}(d\varphi)$$

=
$$\int_{C_{0}([0,1])} \Phi(\varphi(\xi(t_{1})); \varphi'(0), \varphi'(\xi(t_{1})), \varphi'(\xi(1))) w_{\sigma}(d\xi).$$
(3.1)

It is convenient to split the interval [0, 1] into the two intervals $[0, t_1]$ and $[t_1, 1]$, and substitute

$$\xi(t) = \eta_0 \left(\frac{t}{t_1}\right), \quad 0 \le t \le t_1; \qquad \qquad \xi(t) = \eta_0(1) + \eta_1 \left(\frac{t - t_1}{1 - t_1}\right), \quad t_1 \le t \le 1.$$
(3.2)

Now the exponent of the Wiener measure is written as

$$\frac{1}{\sigma^2} \left(\int_0^{t_1} \left(\xi'(t) \right)^2 dt + \int_{t_1}^1 \left(\xi'(t) \right)^2 dt \right) = \frac{1}{\sigma^2 t_1} \int_0^1 \left(\eta'_0(t) \right)^2 dt + \frac{1}{\sigma^2 (1 - t_1)} \int_0^1 \left(\eta'_1(t) \right)^2 dt \,.$$

Due to the Markov property of the Wiener process $\xi(t)$, the measure $w_{\sigma}(d\xi)$ turns into the product of the two measures

$$w_{\sigma\sqrt{t_1}}(d\eta_0) w_{\sigma\sqrt{1-t_1}}(d\eta_1)$$
.

To return to the integrals over the group of diffeomorphisms, define the functions ψ_0 , $\psi_1 \in Diff_+^1([0, 1])$

$$\psi_0(t) = \frac{\int\limits_0^t e^{\eta_0(\tau)} d\tau}{\int\limits_0^1 e^{\eta_0(\tau)} d\tau}, \qquad \psi_1(t) = \frac{\int\limits_0^t e^{\eta_1(\tau)} d\tau}{\int\limits_0^1 e^{\eta_1(\tau)} d\tau}.$$
(3.3)

In this way, we get

$$\int_{Diff_{+}^{1}([0,1])} \Phi(\varphi(t_{1}); \varphi'(0), \varphi'(t_{1}), \varphi'(1)) \mu_{\sigma}(d\varphi)$$

=
$$\int_{Diff_{+}^{1}([0,1])} \int_{Diff_{+}^{1}([0,1])} \Phi(\varphi(t_{1}); \varphi'(0), \varphi'(t_{1}), \varphi'(1)) \mu_{\sigma\sqrt{t_{1}}}(d\psi_{0}) \mu_{\sigma\sqrt{1-t_{1}}}(d\psi_{1}). \quad (3.4)$$

The functions φ and ψ (and their derivatives) are related by the following equations:

$$\varphi(t) = \frac{t_1 \,\psi_1'(0) \,\psi_0\left(\frac{t}{t_1}\right)}{t_1 \psi_1'(0) + (1-t_1)\psi_0'(1)} \,, \qquad 0 \le t \le t_1 \,; \tag{3.5}$$

$$\varphi(t) = \frac{t_1 \,\psi_1'(0) + (1 - t_1) \,\psi_0'(1) \,\psi_1\left(\frac{t - t_1}{1 - t_1}\right)}{t_1 \psi_1'(0) + (1 - t_1) \psi_0'(1)} \,, \qquad t_1 \le t \le 1 \,, \tag{3.6}$$

and

=

$$\varphi'(t) = \frac{\psi_1'(0)\,\psi_0'\left(\frac{t}{t_1}\right)}{t_1\psi_1'(0) + (1-t_1)\psi_0'(1)}, \qquad 0 \le t \le t_1\,; \tag{3.7}$$

$$\varphi'(t) = \frac{\psi_0'(1)\,\psi_1'\left(\frac{t-t_1}{1-t_1}\right)}{t_1\psi_1'(0) + (1-t_1)\psi_0'(1)}\,,\qquad t_1 \le t \le 1\,. \tag{3.8}$$

In particular, the arguments in the integrand are

$$\varphi(t_1) = \frac{t_1 \,\psi_1'(0)}{t_1 \psi_1'(0) + (1 - t_1)\psi_0'(1)}; \qquad \varphi'(t_1) = \frac{\psi_0'(1)\psi_1'(0)}{t_1 \psi_1'(0) + (1 - t_1)\psi_0'(1)}, \tag{3.9}$$

$$\varphi'(0) = \frac{\psi_0'(0)\psi_1'(0)}{t_1\psi_1'(0) + (1-t_1)\psi_0'(1)}, \qquad \varphi'(1) = \frac{\psi_0'(1)\psi_1'(1)}{t_1\psi_1'(0) + (1-t_1)\psi_0'(1)}.$$
(3.10)

Note that the substitution (2.3) is nonlocal, and for diffeomorphisms $\varphi(t)$ the Markov property is not valid.

Now we can represent all functional integrals with integrands depending on $\varphi(t_1)$, $\varphi'(0)$, $\varphi'(t_1)$, $\varphi'(1)$ in a similar way. To this end, we define the basic functional integral:

$$\mathcal{E}_{\sigma}(u, v) = \int_{Diff^{1}([0,1])} \delta\left(\varphi'(0) - u\right) \delta\left(\varphi'(1) - v\right) \mu_{\sigma}(d\varphi), \qquad (3.11)$$

and rewrite the functional integral (3.4) as

$$\int_{Diff_{+}^{1}([0,1])} \int_{Diff_{+}^{1}([0,1])} \Phi\left(\varphi(t_{1}); \varphi'(0), \varphi'(t_{1}), \varphi'(1)\right) \mu_{\sigma\sqrt{t_{1}}}(d\psi_{0}) \mu_{\sigma\sqrt{1-t_{1}}}(d\psi_{1})$$

$$= \int_{0}^{+\infty} \cdots \int_{0}^{+\infty} du_{0} dv_{0} du_{1} dv_{1} \Phi\left(\varphi_{uv}(t_{1}); \varphi'_{uv}(0), \varphi'_{uv}(t_{1}), \varphi'_{uv}(1)\right) \mathcal{E}_{\sigma\sqrt{t_{1}}}(u_{0}, v_{0}) \mathcal{E}_{\sigma\sqrt{1-t_{1}}}(u_{1}, v_{1})$$

$$(3.12)$$

Instead of the variables (u_0, v_0, u_1, v_1) , it is convenient to use the variables

$$z_1 = \varphi(t_1), \quad x_0 = \varphi'(0), \quad y_1 = \varphi'(t_1), \quad x_1 = \varphi'(1).$$

Eqs. (3.9) and (3.10) lead to

$$u_0 = \frac{t_1}{z_1} x_0, \quad v_0 = \frac{t_1}{z_1} y_1, \quad u_1 = \frac{1 - t_1}{1 - z_1} y_1, \quad v_1 = \frac{1 - t_1}{1 - z_1} x_1,$$

and

$$\left|\det \frac{\partial(u_1, v_1, u_2, v_2)}{\partial(z, x_0, x_1, y_1)}\right| = \frac{[t_1(1-t_1)]^2}{[z_1(1-z_1)]^3} y_1.$$

Thus, in this case, we get the transparent rule of the functional integration

$$\int_{Diff_{+}^{1}([0,1])} \Phi\left(\varphi(t_{1});\varphi'(0),\varphi'(t_{1}),\varphi'(1)\right) \ \mu_{\sigma}(d\varphi)$$

$$= [t_{1}(1-t_{1})]^{2} \int_{0}^{1} [z_{1}(1-z_{1})]^{-3} \ dz_{1} \int_{0}^{+\infty} dx_{0} \int_{0}^{+\infty} y_{1} \ dy_{1} \int_{0}^{+\infty} dx_{1}$$

$$\times \Phi\left(z_{1};x_{0},y_{1},x_{1}\right) \mathcal{E}_{\sigma\sqrt{t_{1}}}\left(\frac{t_{1}}{z_{1}}x_{0},\frac{t_{1}}{z_{1}}y_{1}\right) \mathcal{E}_{\sigma\sqrt{1-t_{1}}}\left(\frac{1-t_{1}}{1-z_{1}}y_{1},\frac{1-t_{1}}{1-z_{1}}x_{1}\right).$$
(3.13)

If the integrand Φ depends on values of the function φ and its derivative φ' at other points $t_1, t_2, \dots t_k$ of the interval, we can continue the described procedure

$$\varphi \to (\psi_0, \psi_1), \quad \psi_1 \to (\psi_2, \psi_3), \quad \psi_3 \to (\psi_4, \psi_5), \quad \dots \quad \psi_{2k-3} \to (\psi_{2(k-1)}, \psi_{2k-1}),$$

and obtain the generalization of (3.13)

$$\int_{Diff_+^1([0,1])} \Phi\left(\varphi(t_1), \dots \varphi(t_k); \varphi'(0), \varphi'(t_1), \dots \varphi'(t_k), \varphi'(1)\right) \ \mu_{\sigma}(d\varphi)$$

$$= [t_{1}(t_{2}-t_{1})\cdots(t_{k}-t_{k-1})(1-t_{k})]^{2} \int_{0}^{1} dz_{1}\dots\int_{z_{k-1}}^{1} dz_{k} [z_{1}(z_{2}-z_{1})\cdots(z_{k}-z_{k-1})(1-z_{k})]^{-3}$$

$$\times \int_{0}^{+\infty} dx_{0} \int_{0}^{+\infty} y_{1} dy_{1}\dots\int_{0}^{+\infty} y_{k} dy_{k} \int_{0}^{+\infty} dx_{1} \Phi (z_{1},\dots,z_{k};x_{0},y_{1},\dots,y_{k},x_{1}) \mathcal{E}_{\sigma\sqrt{t_{1}}} \left(\frac{t_{1}}{z_{1}}x_{0},\frac{t_{1}}{z_{1}}y_{1}\right)$$

$$\times \mathcal{E}_{\sigma\sqrt{t_{2}-t_{1}}} \left(\frac{t_{2}-t_{1}}{z_{2}-z_{1}}y_{1},\frac{t_{2}-t_{1}}{z_{2}-z_{1}}y_{2}\right)\cdots\mathcal{E}_{\sigma\sqrt{t_{k}-t_{k-1}}} \left(\frac{t_{k}-t_{k-1}}{z_{k}-z_{k-1}}y_{k-1},\frac{t_{k}-t_{k-1}}{z_{k}-z_{k-1}}y_{k}\right)$$

$$\times \mathcal{E}_{\sigma\sqrt{1-t_{k}}} \left(\frac{1-t_{k}}{1-z_{k}}y_{k},\frac{1-t_{k}}{1-z_{k}}x_{1}\right). \tag{3.14}$$

In this case, the Jacobian of the substitution is

$$\left|\det \frac{\partial(u_1, v_1, \dots, u_{k+1}, v_{k+1})}{\partial(z_1, \dots, z_k, x_0, y_1, \dots, y_k, x_1)}\right| = [t_1 \cdots (1 - t_k)]^2 [z_1 \cdots (1 - z_k)]^{-3} y_1 \cdots y_k.$$

Note that (3.11) is the only one functional integral we need to evaluate. In [49], we performed the functional integration explicitly and represent the functional integral (3.11) in the form of the ordinary integral:

$$\mathcal{E}_{\sigma}(u, v) = \left(\frac{2}{\pi\sigma^2}\right)^{\frac{3}{2}} \frac{1}{\sqrt{uv}} \exp\left\{\frac{2}{\sigma^2}\left(\pi^2 - u - v\right)\right\}$$
$$\times \int_{0}^{+\infty} \exp\left\{-\frac{2}{\sigma^2}\left(2\sqrt{uv}\cosh\theta + \theta^2\right)\right\} \sin\left(\frac{4\pi\theta}{\sigma^2}\right)\sinh(\theta)\,d\theta\,.$$
(3.15)

Now the functional integrals (3.13) and (3.14) are reduced to ordinary multiple integrals.

4 SYK correlation functions

The renormalized two-point SYK correlation function looks like

$$< G_{2}^{n}(0, t_{1}) >_{SYK}^{R}$$

$$= \frac{2\pi}{\sigma^{2}} \int_{Diff^{1}([0,1])} \delta\left(\frac{\varphi'(1)}{\varphi'(0)} - 1\right) \left(\frac{(\varphi'(t_{1})\varphi'(0))^{\frac{1}{4}}}{|\varphi(t_{1})|^{\frac{1}{2}}}\right)^{n} \exp\left\{\frac{8}{\sigma^{2}}\left(\varphi'(0)\right)\right\} \mu_{\sigma}(d\varphi)$$

$$= \frac{2\pi}{\sigma^{2}} \pi^{-\frac{n}{2}} \left[t_{1}\left(1 - t_{1}\right)\right]^{2} \int_{0}^{1} dz_{1} z_{1}^{-3 - \frac{n}{2}} (1 - z_{1})^{-3} \int_{0}^{+\infty} \int_{0}^{+\infty} dx_{0} dy_{1} (x_{0} y_{1})^{1 + \frac{n}{4}}$$

$$\times \exp\left\{\frac{8}{\sigma^{2}} x_{0}\right\} \mathcal{E}_{\sigma\sqrt{t_{1}}}\left(\frac{t_{1}}{z_{1}} x_{0}, \frac{t_{1}}{z_{1}} y_{1}\right) \mathcal{E}_{\sigma\sqrt{1 - t_{1}}}\left(\frac{1 - t_{1}}{1 - z_{1}} y_{1}, \frac{1 - t_{1}}{1 - z_{1}} x_{0}\right). \tag{4.1}$$

In the similar way, we can write the time-ordered (TO) and the out-of-time-ordered (OTO) renormalized four point correlation functions:

$$< G_4^{TO}(0, t_1; t_2, t_3) >^R_{SYK}$$

$$= \frac{2}{\sigma^2} \int_{Diff^1([0,1])} \delta\left(\frac{\varphi'(1)}{\varphi'(0)} - 1\right) \frac{(\varphi'(t_1)\varphi'(0))^{\frac{1}{4}}}{|\varphi(t_1)|^{\frac{1}{2}}} \frac{(\varphi'(t_3)\varphi'(t_2))^{\frac{1}{4}}}{|\varphi(t_3) - \varphi(t_2)|^{\frac{1}{2}}} \exp\left\{\frac{8}{\sigma^2} \left(\varphi'(0)\right)\right\} \mu_{\sigma}(d\varphi),$$
(4.2)

and

$$< G_4^{OTO}(0, t_2; t_1, t_3) >_{SYK}^R$$

$$= \frac{2}{\sigma^2} \int_{Diff^1([0,1])} \delta\left(\frac{\varphi'(1)}{\varphi'(0)} - 1\right) \frac{(\varphi'(t_2)\varphi'(0))^{\frac{1}{4}}}{|\varphi(t_2)|^{\frac{1}{2}}} \frac{(\varphi'(t_3)\varphi'(t_1))^{\frac{1}{4}}}{|\varphi(t_3) - \varphi(t_1)|^{\frac{1}{2}}} \exp\left\{\frac{8}{\sigma^2} \left(\varphi'(0)\right)\right\} \mu_{\sigma}(d\varphi).$$

$$(4.3)$$

(In the both equations, we assume that $0 < t_1 < t_2 < t_3 < 1$.)

In terms of the functions ${\mathcal E}\,,$ the four-point correlation functions look like

0.000

$$< G_4^{TO(OTO)} \left(0, t_{1(2)}; t_{2(1)}, t_3 \right) >_{SYK}^R = \frac{2}{\sigma^2} \left[t_1 (t_2 - t_1) (t_3 - t_2) (1 - t_3) \right]^2$$

$$\times \int_0^1 dz_1 \int_{z_1}^1 dz_2 \int_{z_2}^1 dz_3 \ \chi^{TO(OTO)} (z_1, z_2, z_3) \int_0^{+\infty} dx_0 \int_0^{+\infty} dy_1 \int_0^{+\infty} dy_2 \int_0^{+\infty} dy_3 \left[x_0 y_1 y_2 y_3 \right]^{\frac{5}{4}}$$

$$\times \exp \left\{ \frac{8}{\sigma^2} x_0 \right\} \ \mathcal{E}_{\sigma\sqrt{t_1}} \left(\frac{t_1}{z_1} x_0, \frac{t_1}{z_1} y_1 \right) \ \mathcal{E}_{\sigma\sqrt{t_2 - t_1}} \left(\frac{t_2 - t_1}{z_2 - z_1} y_1, \frac{t_2 - t_1}{z_2 - z_1} y_2 \right)$$

$$\times \ \mathcal{E}_{\sigma\sqrt{t_3 - t_2}} \left(\frac{t_3 - t_2}{z_3 - z_2} y_2, \frac{t_3 - t_2}{z_3 - z_2} y_3 \right) \ \mathcal{E}_{\sigma\sqrt{1 - t_3}} \left(\frac{1 - t_3}{1 - z_3} y_3, \frac{1 - t_3}{1 - z_3} x_0 \right),$$

$$(4.4)$$

where

$$\chi^{TO}(z_1, z_2, z_3) = \left[z_1(z_2 - z_1)(z_3 - z_2)(1 - z_3)\right]^{-3} \left[z_1(z_3 - z_2)\right]^{-\frac{1}{2}}, \quad (4.5)$$

and

$$\chi^{OTO}(z_1, z_2, z_3) = \left[z_1(z_2 - z_1)(z_3 - z_2)(1 - z_3)\right]^{-3} \left[z_2(z_3 - z_1)\right]^{-\frac{1}{2}}.$$
 (4.6)

Note that the only difference between (4.5) and (4.6) is in the dependence of the integrands on the variables z_i .

Neither the OTO four-point correlation function nor the TO four-point correlation function has the form of the product of two two-point correlation functions.

In appendix B, we obtain the representations for the correlation functions in terms of the ordinary integrals that can be analyzed numerically.

Using the general rule (3.14) we can write any n-point correlator in the same way.

5 Concluding remarks

The paper completes the elaboration of the functional integrals calculus in the theories invariant under the groups of diffeomorphisms started in [48] and [49].

The general rules derived here give a straightforward scheme to evaluate functional integrals in the theories of the Schwarzian type. The great merit of the method is that it reduces various problems to the evaluation of the functional integral (3.11) only.

The functional integral (3.11)

$$\mathcal{E}_{\sigma}(u, v) = \int_{Diff^{1}([0,1])} \delta\left(\varphi'(0) - u\right) \delta\left(\varphi'(1) - v\right) \mu_{\sigma}(d\varphi)$$

is evaluated explicitly. And the result is written in the form of the ordinary integral (3.15). Therefore the evaluations of other functional integrals of the form (3.14) lead also to ordinary (multiple) integrals.

In the introduction, we explain the difference between functional integration over the group $Diff_{+}^{1}(S^{1})$ and that over the group $Diff_{+}^{1}(\mathbf{R})$. Depending on the problem (e.g., closed or open boundary in AdS_{2} is considered), one should decide what kind of functional integrals is adequate.

The functional integrals over $Diff_+^1(\mathbf{R})/P$ corresponding to SYK correlation functions were evaluated in many papers (see,e.g., [23], [24], [25], [30]).

As far as we know, the results for correlation functions defined as functional integrals over $Diff_+^1(S^1)/SL(2, \mathbf{R})$ has been obtained here for the first time.

The fact that in this case, the four-point correlation functions cannot be represented in the form of the product of two two-point correlation functions is connected with the apparent non-Markov behaviour of the function $\varphi(t)$. Although ξ is the Wiener process, the Markov property is violated by the nonlocal substitution (2.3). The non-Markov property of the measure (1.6) also manifests itself in (4.1) and (4.4). The regions with $t > t_1$ (for $\langle G_2(0, t_1) \rangle_{SYK}^R$), and with $t > t_3$ (for $\langle G_4(0, t_{1(2)}; t_{2(1)}, t_3) \rangle_{SYK}^R$) give the nonzero inputs into the integrals for correlation functions.

If one considers t as the time variable, then one can say that the present is influenced by the future. It is a direct consequence of the fact that the ends of the interval are glued together and the theory is $SL(2, \mathbf{R})$ invariant.

In this paper, we obtain the correlation functions in the form that can be analyzed numerically. The analysis is planned to be the subject of another paper.

A Convolution of the \mathcal{E} functions

Let us take the integrand of (3.11) for Φ . In this case, we get the convolution rule for two \mathcal{E} functions:

$$\mathcal{E}_{\sigma}(u, v) = [t_1(1-t_1)]^2 \int_0^1 [z_1(1-z_1)]^{-3} dz_1 \int_0^{+\infty} y_1 dy_1$$

 $\times \mathcal{E}_{\sigma\sqrt{t_1}} \left(\frac{t_1}{z_1} u, \frac{t_1}{z_1} y_1\right) \mathcal{E}_{\sigma\sqrt{1-t_1}} \left(\frac{1-t_1}{1-z_1} y_1, \frac{1-t_1}{1-z_1} v\right).$ (A.1)

We would like to stress that this convolution rule has been proved by direct explicit evaluation of functional integrals.

However, to demonstrate once more that the dependence of the right-hand side of (A.1) on t_1 is fictitious, we consider the integral

$$\int_{0}^{+\infty} \mathcal{E}_{\sigma}(u, u) u \, du = \frac{1}{\sqrt{2\pi^{3}\sigma^{2}}} \int_{0}^{+\infty} d\theta \, \exp\left\{-\frac{2}{\sigma^{2}} \left[\theta^{2} - \pi^{2}\right]\right\} \, \sin\left(\frac{4\pi\,\theta}{\sigma^{2}}\right) \tanh(\frac{\theta}{2}) \,. \tag{A.2}$$

At the same time, we have from the right-hand side of (A.1)

$$\frac{2}{\pi^{3}\sigma^{2}}[t_{1}(1-t_{1})]^{-\frac{1}{2}}\int_{0}^{+\infty}\int_{0}^{+\infty}d\theta_{1} d\theta_{2} \exp\left\{-\frac{2}{\sigma^{2}}\left[\frac{\theta_{1}^{2}-\pi^{2}}{t_{1}}+\frac{\theta_{2}^{2}-\pi^{2}}{(1-t_{1})}\right]\right\}$$

$$\times \sin\left(\frac{4\pi\theta_{1}}{\sigma^{2}t_{1}}\right)\sin\left(\frac{4\pi\theta_{2}}{\sigma^{2}(1-t_{1})}\right)\sinh\theta_{1}\sinh\theta_{2}\int_{0}^{1}dz_{1}\left[z_{1}(1-z_{1})\right]^{-2}$$

$$\times\int_{0}^{+\infty}\int_{0}^{+\infty}du\,dy_{1}\exp\left\{-\frac{1}{z_{1}(1-z_{1})}\left(u+y_{1}+2\left[(1-z_{1})\cosh\theta_{1}+z_{1}\cosh\theta_{2}\right]\sqrt{uy_{1}}\right)\right\}$$

$$=\frac{2}{\pi^{3}\sigma^{2}}\left[t_{1}\left(1-t_{1}\right)\right]^{-\frac{1}{2}}\int_{0}^{+\infty}\int_{0}^{+\infty}d\theta_{1}\,d\theta_{2}\exp\left\{-\frac{2}{\sigma^{2}}\left[\frac{\theta_{1}^{2}-\pi^{2}}{t_{1}}+\frac{\theta_{2}^{2}-\pi^{2}}{(1-t_{1})}\right]\right\}\sinh\theta_{1}\sinh\theta_{2}$$

$$\times\sin\left(\frac{4\pi\theta_{1}}{\sigma^{2}t_{1}}\right)\sin\left(\frac{4\pi\theta_{2}}{\sigma^{2}(1-t_{1})}\right)\int_{0}^{1}dz_{1}\left[\frac{b}{(b^{2}-1)^{\frac{3}{2}}}arc\cosh b-\frac{1}{b^{2}-1}\right],\qquad(A.3)$$

where

$$b = \left[(1 - z_1) \cosh \theta_1 + z_1 \cosh \theta_2 \right] \,.$$

To obtain (A.3), we have re-scaled the variables

$$\bar{u} = \frac{u}{z_1(1-z_1)}, \quad \bar{y_1} = \frac{y_1}{z_1(1-z_1)}.$$

And then after the substitutions

$$\bar{u} = \rho \cos^2 \omega, \quad \bar{y}_1 = \rho \sin^2 \omega, \quad \xi = \tan \omega,$$
 (A.4)

we have used the table of integrals [58] (eq. 2.2.9.15).

The results of computations of (A.3) at ten different points of the interval [0, 1] coincide with the result of computation of (A.2) with very high accuracy (e.g., the results are 0.07978 for $\sigma = 5$, and 0.03989 for $\sigma = 10$).

B Correlation functions in terms of ordinary integrals

In terms of the ordinary integrals, the two-point correlation function (4.1) has the form

$$< G_{2}^{n}(0, t_{1}) >_{SYK}^{R} = \frac{1}{\pi^{2}} \left(\frac{\sigma^{2}}{2\pi}\right)^{\frac{n}{2}} [t_{1}(1-t_{1})]^{-\frac{1}{2}}$$

$$\times \int_{0}^{+\infty} \int_{0}^{+\infty} d\theta_{1} d\theta_{2} \exp\left\{-\frac{2}{\sigma^{2}} \left[\frac{\theta_{1}^{2} - \pi^{2}}{t_{1}} + \frac{\theta_{2}^{2} - \pi^{2}}{(1-t_{1})}\right]\right\} \sin\left(\frac{4\pi \theta_{1}}{\sigma^{2} t_{1}}\right) \sin\left(\frac{4\pi \theta_{2}}{\sigma^{2} (1-t_{1})}\right)$$

$$\times \sinh(\theta_1) \sinh(\theta_2) \int_{0}^{1} dz_1 (1-z_1)^{\frac{n}{2}} \int_{0}^{+\infty} \int_{0}^{+\infty} dx_0 dy_1 (x_0 y_1)^{\frac{n}{4}}$$

 $\times \exp\left\{-\left(x_0 + y_1 + 2\left[(1 - z_1)\cosh\theta_1 + z_1\cosh\theta_2\right]\sqrt{x_0y_1} - 4z_1(1 - z_1)x_0\right)\right\}.$ (B.1)

After the substitutions (A.4), the integrals over x_0 and y_1 are transformed into the integral

$$\int_{0}^{+\infty} \frac{2\xi^{1+\frac{n}{2}}}{\left[\xi^{2}+2b\xi+1-4z_{1}(1-z_{1})\right]^{2+\frac{n}{2}}} d\xi \,. \tag{B.2}$$

It is convenient to use the table of integrals [58] (eq. 2.2.9.8).

$$< G_{2}^{n}(0,t_{1}) >_{SYK}^{R} = \frac{2}{\pi^{2}} \left(\frac{\sigma^{2}}{2\pi} \right)^{\frac{n}{2}} \Gamma^{3} \left(\frac{n+4}{2} \right) \Gamma^{-1}(n+4) \left[t_{1}(1-t_{1}) \right]^{-\frac{1}{2}}$$

$$\times \int_{0}^{+\infty} \int_{0}^{+\infty} d\theta_{1} d\theta_{2} \exp \left\{ -\frac{2}{\sigma^{2}} \left[\frac{\theta_{1}^{2} - \pi^{2}}{t_{1}} + \frac{\theta_{2}^{2} - \pi^{2}}{(1-t_{1})} \right] \right\} \sin \left(\frac{4\pi \theta_{1}}{\sigma^{2} t_{1}} \right) \sin \left(\frac{4\pi \theta_{2}}{\sigma^{2} (1-t_{1})} \right)$$

$$\times \sinh(\theta_{1}) \sinh(\theta_{2}) \int_{0}^{1} dz_{1} \left(1 - z_{1} \right)^{\frac{n}{2}} \left[1 - 4z_{1}(1-z_{1}) \right]^{-\frac{n+4}{4}}$$

$$\times F \left(\frac{n+4}{2}, \frac{n+4}{2}; \frac{n+5}{2}; \frac{1}{2} \left(1 - \frac{b}{1-4z_{1}(1-z_{1})} \right) \right), \qquad (B.3)$$

where F is the Gaussian hypergeometric function, and

$$b = (1 - z_1)\cosh\theta_1 + z_1\cosh\theta_2.$$
(B.4)

For n = 2k, $k = 0, 1 \dots$, we can also use eq. 2.2.9.12 of [58], and rewrite the result in the form

$$< G_{2}^{n}(0, t_{1}) >_{SYK}^{R} = (-1)^{k+1} \frac{1}{(k+1)!} \frac{1}{2\pi^{2}} \left(\frac{\sigma^{2}}{4\pi}\right)^{k} \Gamma(k+2) \left[t_{1}(1-t_{1})\right]^{-\frac{1}{2}}$$

$$\times \int_{0}^{+\infty} \int_{0}^{+\infty} d\theta_{1} d\theta_{2} \exp\left\{-\frac{2}{\sigma^{2}} \left[\frac{\theta_{1}^{2} - \pi^{2}}{t_{1}} + \frac{\theta_{2}^{2} - \pi^{2}}{(1-t_{1})}\right]\right\} \sin\left(\frac{4\pi \theta_{1}}{\sigma^{2} t_{1}}\right) \sin\left(\frac{4\pi \theta_{2}}{\sigma^{2} (1-t_{1})}\right)$$

$$\times \sinh(\theta_{1}) \sinh(\theta_{2}) \int_{0}^{1} dz_{1} (1-z_{1})^{k}$$

$$\times \frac{\partial^{k+1}}{\partial b^{k+1}} \left(\frac{1}{\sqrt{b^{2} - 1 + 4z_{1}(1-z_{1})}} \log \frac{b + \sqrt{b^{2} - 1 + 4z_{1}(1-z_{1})}}{b - \sqrt{b^{2} - 1 + 4z_{1}(1-z_{1})}}\right), \quad (B.5)$$

where b is given by (B.4).

Here, we also present the four-point correlation functions in the form that can be analyzed numerically. It is

$$< G_4^{TO(OTO)} \left(0, t_{1(2)}; t_{2(1)}, t_3 \right) >_{SYK}^R = \left(\frac{2}{\pi^3 \sigma^2} \right)^2 \left[t_1(t_2 - t_1)(t_3 - t_2)(1 - t_3) \right]^{-\frac{1}{2}}$$

$$\times \int_0^1 dz_1 \int_{z_1}^1 dz_2 \int_{z_2}^1 dz_3 \ \tilde{\chi}^{TO(OTO)} \left(z_1, z_2, z_3 \right)$$

$$\times \int_0^{+\infty} d\theta_1 \dots \int_0^{+\infty} d\theta_4 \exp \left\{ -\frac{2}{\sigma^2} \left[\frac{\theta_1^2 - \pi^2}{\tau_1} + \frac{\theta_2^2 - \pi^2}{(\tau_2 - \tau_1)} + \frac{\theta_3^2 - \pi^2}{(\tau_3 - \tau_2)} + \frac{\theta_4^2 - \pi^2}{(1 - \tau_3)} \right] \right\}$$

$$\times \sin \left(\frac{4\pi \theta_1}{\sigma^2 \tau_1} \right) \sin \left(\frac{4\pi \theta_2}{\sigma^2 (\tau_2 - \tau_1)} \right) \sin \left(\frac{4\pi \theta_3}{\sigma^2 (\tau_3 - \tau_2)} \right)$$

$$\times \sinh \theta_1 \sinh \theta_2 \sinh \theta_3 \sinh \theta_4 \int_0^{+\infty} dx_0 \int_0^{+\infty} dy_1 \int_0^{+\infty} dy_2 \int_0^{+\infty} dy_3 \left[x_0 y_1 y_2 y_3 \right]^{\frac{1}{4}}$$

$$\times \exp \left\{ 4x_0 - \frac{1}{z_1} \left[x_0 + y_1 + 2\sqrt{x_0 y_1} \cosh \theta_1 \right] - \frac{1}{z_2 - z_1} \left[y_1 + y_2 + 2\sqrt{y_1 y_2} \cosh \theta_2 \right] \right\}$$

$$\exp \left\{ -\frac{1}{z_3 - z_2} \left[y_2 + y_3 + 2\sqrt{y_2 y_3} \cosh \theta_3 \right] - \frac{1}{1 - z_3} \left[y_3 + x_0 + 2\sqrt{y_3 x_0} \cosh \theta_4 \right] \right\},$$

$$(B.6)$$

$$\exp \left\{ -\frac{1}{z_1} \left[x_0 + y_1 + 2\sqrt{y_2 y_3} \cosh \theta_3 \right] - \frac{1}{1 - z_3} \left[y_3 + x_0 + 2\sqrt{y_3 x_0} \cosh \theta_4 \right] \right\},$$

where

 \times

$$\tilde{\chi}^{TO}(z_1, z_2, z_3) = [z_1(z_2 - z_1)(z_3 - z_2)(1 - z_3)]^{-2} [z_1(z_3 - z_2)]^{-\frac{1}{2}},$$

and

$$\tilde{\chi}^{OTO}(z_1, z_2, z_3) = [z_1(z_2 - z_1)(z_3 - z_2)(1 - z_3)]^{-2} [z_2(z_3 - z_1)]^{-\frac{1}{2}}.$$

C Functional integrals over the space $Diff_+^1(\mathbf{R})$

In this appendix, we evaluate functional integrals over $Diff_+^1({\bf R})$ of the form

$$\int_{Diff_{+}^{1}(\mathbf{R})} \Psi(f(\tau_{1}), f(\tau_{2}), ...) \exp\left\{\frac{1}{\sigma^{2}} \int_{-\infty}^{+\infty} \mathcal{S}_{f}(\tau) d\tau\right\} df.$$
(C.1)

Consider the two-point correlation function in this theory

$$\mathcal{G}_{2}^{n}(0,\,\tau) = \int_{Diff_{+}^{1}(\mathbf{R})/P} \frac{(f'(0)f'(\tau))^{\frac{n}{4}}}{|f(\tau) - f(0)|^{\frac{n}{2}}} \exp\left\{\frac{1}{\sigma^{2}} \int_{-\infty}^{+\infty} \mathcal{S}_{f}(t) \, dt\right\} \, df \, .$$

As it is explained in the introduction, we should factorize the integration space. To exclude the non-relevant degrees of freedom in another way, we can fix the values

$$f(0) = 0$$
, $f'(0) = 1$.

After the substitution $f(\tau) = \int_{0}^{\tau} \exp\{\xi(t)\}dt$, $\xi(0) = 0$ ([23], [24]), it looks like

$$\mathcal{G}_{2}^{n}(0,\,\tau) = \int_{C((-\infty,\,+\infty))} \frac{e^{\frac{n}{4}(\xi(\tau))}}{\left(\int\limits_{0}^{\tau} e^{\xi(t)}dt\right)^{\frac{n}{2}}} \exp\left\{-\frac{1}{2\sigma^{2}}\int\limits_{-\infty}^{+\infty} \left(\xi'(t)\right)^{2}\,dt\right\}\,d\xi\,.$$

Due to the Markov property of the Wiener measure, it can be rewritten as

$$\mathcal{G}_{2}^{n}(0,\tau) = \int_{C([0,\tau])} \frac{e^{\frac{n}{4}(\xi(\tau))}}{\left(\int\limits_{0}^{\tau} e^{\xi(t)}dt\right)^{\frac{n}{2}}} \exp\left\{-\frac{1}{2\sigma^{2}}\int\limits_{0}^{\tau} \left(\xi'(t)\right)^{2} dt\right\} d\xi.$$
(C.2)

Instead of transforming the Wiener integral (C.2) to the functional integral in Liouville theory as it was proposed in [23], [24], we use the method described above. First, we re-scale the time variable $t = \tau \bar{t}$ and substitute $\eta(\bar{t}) = \xi(t)$. Then (C.2) has the form

$$\mathcal{G}_{2}^{n}(0,\tau) = \tau^{-\frac{n}{2}} \int_{C([0,1])} \frac{e^{\frac{n}{4}\eta(1)}}{\left(\int_{0}^{1} e^{\eta(\bar{t})} d\bar{t}\right)^{\frac{n}{2}}} \exp\left\{-\frac{1}{2\sigma^{2}\tau} \int_{0}^{1} \left(\eta'(\bar{t})\right)^{2} d\bar{t}\right\} d\eta.$$
(C.3)

In terms of the function

$$\varphi(t) = \frac{\int\limits_{0}^{t} e^{\eta(\bar{t})} d\bar{t}}{\int\limits_{0}^{1} e^{\eta(\bar{t})} d\bar{t}},$$

(C.3) is written as

$$\tau^{-\frac{n}{2}} \int_{Diff^{1}_{+}([0,1])} \left(\varphi'(0)\varphi'(1)\right)^{\frac{n}{4}} \mu_{\sigma\sqrt{\tau}}(d\varphi) = \tau^{-\frac{n}{2}} \int_{0}^{+\infty} \int_{0}^{+\infty} \int_{0}^{+\infty} \mathcal{E}_{\sigma\sqrt{\tau}}(u,v) \, du \, dv \,. \quad (C.4)$$

Integrating over u and v as in the appendix A, we obtain

$$\mathcal{G}_{2}^{n}(0,\tau) = 2\pi^{-\frac{3}{2}} \left(\frac{\sigma^{2}}{2}\right)^{\frac{n-1}{2}} B\left(\frac{n+1}{2},\frac{n+1}{2}\right) \tau^{-\frac{1}{2}} \int_{0}^{+\infty} d\theta \sin\left(\frac{4\pi\theta}{\sigma^{2}\tau}\right) \sinh\theta$$

$$\times \exp\left\{-\frac{2}{\sigma^{2}\tau} \left[\theta^{2} - \pi^{2}\right]\right\} F\left(\frac{n+1}{2},\frac{n+1}{2};\frac{n+1}{2} + 1;\frac{1-\cosh\theta}{2}\right). \quad (C.5)$$

In the case n = 2, it looks like

$$\mathcal{G}_2^2(0,\,\tau) = \left(\frac{\sigma^2}{2\pi^3}\right)^{\frac{1}{2}} \tau^{-\frac{1}{2}} \int_0^{+\infty} d\theta \exp\left\{-\frac{2}{\sigma^2 \tau} \left[\theta^2 - \pi^2\right]\right\} \sin\left(\frac{4\pi\theta}{\sigma^2 \tau}\right) \left[\frac{\theta\cosh\theta}{\sinh^2\theta} - \frac{1}{\sinh\theta}\right].$$
(C.6)

The asymptotic form of $\mathcal{G}_2^n(0,\,\tau)$ at $\tau\to\infty\;$,

$$\left(\mathcal{G}_{2}^{n}(0,\tau)\right)_{As} = 4\pi^{-\frac{1}{2}} \left(\frac{\sigma^{2}}{2}\right)^{\frac{n-3}{2}} B\left(\frac{n+1}{2},\frac{n+1}{2}\right) \tau^{-\frac{3}{2}} \\ \times \int_{0}^{+\infty} \theta \sinh\theta F\left(\frac{n+1}{2},\frac{n+1}{2};\frac{n+1}{2}+1;\frac{1-\cosh\theta}{2}\right) d\theta, \qquad (C.7)$$

immediately follows from (C.5).

In the same way, we can explicitly evaluate functional integrals for other correlation functions in this theory.

In particular, for $\mathcal{G}_4^{TO}(0, \tau_1; \tau_2, \tau_3)$ we have

$$\mathcal{G}_{4}^{TO}(0,\,\tau_{1};\,\tau_{2},\,\tau_{3}) = \int_{Diff_{+}^{1}(\mathbf{R})/P} \frac{(f'(0)f'(\tau_{1}))^{\frac{1}{4}}}{|f(\tau_{1}) - f(0)|^{\frac{1}{2}}} \frac{(f'(\tau_{2})f'(\tau_{3}))^{\frac{1}{4}}}{|f(\tau_{3}) - f(\tau_{2})|^{\frac{1}{2}}} \exp\left\{\frac{1}{\sigma^{2}} \int_{-\infty}^{+\infty} \mathcal{S}_{f}(t) \, dt\right\} \, df$$
$$= \mathcal{G}_{2}(0,\,\tau_{1}) \, \mathcal{G}_{2}(\tau_{2},\,\tau_{3}) \, . \tag{C.8}$$

The OTO four-point correlation function is given by $(0 < \tau_1 < \tau_2 < \tau_3)$

$$\begin{aligned} \mathcal{G}_{4}^{OTO}(0,\,\tau_{2};\,\tau_{1},\,\tau_{3}) &= \int_{Diff_{+}^{4}(\mathbf{R})/P} \frac{(f'(0)f'(\tau_{2}))^{\frac{1}{4}}}{|f(\tau_{2}) - f(0)|^{\frac{1}{2}}} \frac{(f'(\tau_{1})f'(\tau_{3}))^{\frac{1}{4}}}{|f(\tau_{3}) - f(\tau_{1})|^{\frac{1}{2}}} \exp\left\{\frac{1}{\sigma^{2}} \int_{-\infty}^{+\infty} \mathcal{S}_{f}(t) \, dt\right\} \, df \\ &= \pi^{\frac{9}{2}} \left(\frac{\sigma^{2}}{2}\right)^{\frac{1}{2}} [\tau_{1}(\tau_{2} - \tau_{1})(\tau_{3} - \tau_{2})]^{-\frac{1}{2}} \int_{0}^{1} dz_{1} \int_{z_{1}}^{1} dz_{2} \left[z_{1}(z_{2} - z_{1})(1 - z_{2})\right]^{-2} \left[z_{2}(1 - z_{1})\right]^{-\frac{1}{2}} \\ &\times \int_{0}^{+\infty} \int_{0}^{+\infty} \int_{0}^{+\infty} d\theta_{1} \, d\theta_{2} \, d\theta_{3} \exp\left\{-\frac{2}{\sigma^{2}} \left[\frac{\theta_{1}^{2} - \pi^{2}}{\tau_{1}} + \frac{\theta_{2}^{2} - \pi^{2}}{(\tau_{2} - \tau_{1})} + \frac{\theta_{3}^{2} - \pi^{2}}{(\tau_{3} - \tau_{2})}\right]\right\} \\ &\times \sin\left(\frac{4\pi \, \theta_{1}}{\sigma^{2} \, \tau_{1}}\right) \sin\left(\frac{4\pi \, \theta_{2}}{\sigma^{2} \, (\tau_{2} - \tau_{1})}\right) \sin\left(\frac{4\pi \, \theta_{3}}{\sigma^{2} \, (\tau_{3} - \tau_{2})}\right) \sinh\theta_{1} \sinh\theta_{2} \sinh\theta_{3} \\ &\times \int_{0}^{+\infty} dx_{0} \int_{0}^{+\infty} dy_{1} \int_{0}^{+\infty} dy_{2} \int_{0}^{+\infty} dx_{1} \left[x_{0}x_{1}\right]^{-\frac{1}{4}} \left[y_{1}y_{2}\right]^{\frac{1}{4}} \exp\left\{-\frac{1}{z_{1}} \left[x_{0} + y_{1} + 2\sqrt{x_{0}y_{1}} \cosh\theta_{1}\right]\right\} \\ &\times \exp\left\{-\frac{1}{z_{2} - z_{1}} \left[y_{1} + y_{2} + 2\sqrt{y_{1}y_{2}} \cosh\theta_{2}\right] - \frac{1}{1 - z_{2}} \left[y_{2} + x_{1} + 2\sqrt{y_{2}x_{1}} \cosh\theta_{3}\right]\right\}. (C.9) \end{aligned}$$

In contrast to the correlation functions in the theory considered in the main part of the paper, the correlation functions (C.2), (C.8), (C.9) have no input from the regions $t > \tau$, or $t > \tau_3$.

References

- [1] S. Sachdev and J. Ye, Gapless spin fluid ground state in a random quantum Heisenberg magnet, Phys. Rev. Lett. **70** (1993) 3339, arXiv:cond-mat/9212030 [cond-mat].
- [2] A. Kitaev, Hidden correlations in the Hawking radiation and thermal noise, Talk at KITP, http://online.kitp.ucsb.edu/online/joint98/kitaev/, Febrary, 2015.
- [3] A. Kitaev, A simple model of quantum holography, Talks at KITP, http://online.kitp.ucsb.edu/online/entangled15/kitaev/ and http://online.kitp.ucsb.edu/online/entangled15/kitaev2/, April and May, 2015.
- [4] J. Maldacena and D. Stanford, Remarks on the Sachdev-Ye-Kitaev model, Phys. Rev. D 94(2016) 106002, [arXiv:1604.07818].
- [5] J. Polchinski and V. Rosenhaus, The spectrum in the Sachdev-Ye-Kitaev model, JHEP 04 (2017) 001, [arXiv:1601.06768].
- [6] A. Jevicki, K. Suzuki and J. Yoon, Bi-local holography in the SYK model, JHEP 07 (2016) 007, [arXiv:1603.06246].
- [7] A. Jevicki and K. Suzuki, Bi-local holography in the SYK model: perturbations, JHEP 11 (2016) 046, [arXiv:1608.07567].
- [8] D. J. Gross and V. Rosenhaus, A Generalization of Sachdev-Ye-Kitaev, JHEP 02 (2017) 093, [arXiv:1610.01569].
- [9] D. J. Gross and V. Rosenhaus, The bulk dual of SYK: cubic couplings, JHEP 05 (2017) 092, [arXiv:1702.08016].
- S.R. Das, A. Jevicki and K. Suzuki, Three dimensional view of the SYK/AdS duality, JHEP 09 (2017) 017, [arXiv:1704.07208].
- [11] S.R. Das, A.Ghosh, A. Jevicki and K. Suzuki, Space-time the SYK model, JHEP 07 (2018) 184, [arXiv:1712.02725].
- [12] A. Almheiri and J. Polchinski, Models of AdS₂ backreaction and holography, JHEP 11 (2015) 014, [arXiv:1402.6334].
- [13] K. Jensen, Chaos in AdS₂ holography, Phys. Rev. Lett. **117** (2016) 111601, [arXiv:1605.06098].
- [14] J. Maldacena, D. Stanford and Z. Yang, Conformal symmetry and its breaking in two dimensional Nearly Anti-de-Sitter space, PTEP 2016, no. 12, 12C104 (2016), [arXiv:1606.01857].
- [15] J. Engelsoy, T. G. Mertens, and H. L. Verlinde, An investigation of AdS₂ backreaction and holography, JHEP 07 (2016) 139, [arXiv:1606.03438].
- [16] J.S. Cotler, G. Gur-Ari, M. Hanada, J. Polchinski, P. Saad, S.H. Shenker, D. Stanford, A. Streicher and M. Tezuka, *Black Holes and Random Matrices*, *JHEP* 05 (2017) 118, [arXiv:1611.04650].
- [17] A. M. Garcia-Garcia and J. J. M. Verbaarschot, Analytical Spectral Density of the Sachdev-Ye-Kitaev Model at finite N, Phys. Rev. D 96(2017) 066012, [arXiv:1701.06593].
- [18] G. J. Turiaci and H. L. Verlinde, Towards a 2d QFT Analog of the SYK Model, JHEP 10 (2017) 167, [arXiv:1701.00528].

- [19] G. Mandal, P. Nayak and S. R. Wadia, Coadjoint orbit action of Virasoro group and two-dimentional quantum gravity dual SYK/tensor models, JHEP 11 (2018) 046, [arXiv:1702.04266].
- [20] A. Kitaev and S. J. Suh, The soft mode in the Sachdev-Ye-Kitaev model and its gravity dual, JHEP 05 (2018) 183, [arXiv:1711.08467].
- [21] G. Sarosi, AdS₂ holography and the SYK model, PoS (Modave2017)001 (2018), [arXiv:1711.08482].
- [22] J. Maldacena, S.H. Shenker and D. Stanford, A bound on chaos, JHEP 08(2016) 106, [arXiv:1503.01409].
- [23] D. Bagrets, A. Altland and A. Kamenev, Sachdev-Ye-Kitaev Model as Liouville Quantum Mechanics, Nucl. Phys. B 911 (2016) 191, [arXiv:1607.00694].
- [24] D. Bagrets, A. Altland and A. Kamenev, Power-law out of time order correlation functions in the SYK model, Nucl. Phys. B 921 (2017) 727, [arXiv:1702.08902].
- [25] T. G. Mertens, G. J. Turiaci and H. L. Verlinde, Solving the Schwarzian via the Conformal Bootstrap, JHEP 08 (2017) 136, [arXiv:1705.08408].
- [26] D. J. Gross and V. Rosenhaus, All point correlation functions in SYK, JHEP 12 (2017) 148, [arXiv:1710.08113].
- [27] D. Stanford and E. Witten, Fermionic Localization of the Schwazian Theory, JHEP 10 (2017) 008, [arXiv:1703.04612].
- [28] T. G. Mertens, The Schwarzian Theory Origins, JHEP 05 (2018) 036, [arXiv:1801.09605].
- [29] H.T Lam, T. G. Mertens, G. J. Turiaci and H. L. Verlinde, Shockwave S-matrix from Schwarzian Quantum Mechanics, arXiv:1804.09834.
- [30] A. Blommaert, T. G. Mertens and H. Verschelde, The Schwarzian Theory A Wilson Line Perspective, arXiv:1806.07765.
- [31] A. Gaikwad, L. K. Joshi, G. Mandal and S. R. Wadia Holographic dual to charged SYK from 3D Gravity and Chern-Simons, arXiv:1802.07746.
- [32] I. Aref'eva and I. Volovich, Notes on the SYK model in real time, Theor. Math. Phys. 197(2018) 1650, [arXiv:1801.08118].
- [33] P. Nayak, A. Shukla, R. M. Soni, S. P. Trivedi and V. Vishal, On the Dynamics of Near-Extremal Black Holes, JHEP 09 (2018) 048, [arXiv:1802.09547].
- [34] D. A. Roberts, D. Stanford and A. Streicher, Operator grouth in the SYK model, arXiv:1802.02633.
- [35] P. Saad, S.H. Shenker and D. Stanford, A semiclassical ramp in SYK and in gravity, arXiv:1806.06840.
- [36] G. Gur-Ari, R. Mahajan and A. Vaezi, Does the SYK model have a spin glass phase?, JHEP 11 (2018) 070, [arXiv:1806.10145].
- [37] V. Rosenhaus, An introduction to the SYK model, arXiv:1807.03334.
- [38] J. Cotler and K. Jensen, A theory of reparametrization for AdS₂ gravity, arXiv:1808.03263.
- [39] A. Kitaev and S. J. Suh, *Statistical mechanics of a two-dimensional black hole* arXiv:1808.07032.

- [40] A. Banerjee, A. Kundu and R. P. Poojary, String, Brains, Schwarzian Action and Maximal Chaos, arXiv:1809.02090.
- [41] M. Berkooz, M. Isachenkov, V. Narovlansky and G. Torrents, Towards a full solution of the large N double-scaled SYK model, arXiv:1811.02584.
- [42] I. Aref'eva, M. Khramtsov, M. Tikhanovskaya and I. Volovich, *Replica-nondiagonal solutions in the SYK model*, arXiv:1811.04831.
- [43] H. Wang, D. Bagrets, A. L. Chudnovskiy and A. Kamenev On the replica structure of Sachdev-Ye-Kitaev model, arXiv:1812.02666.
- [44] A. Blommaert, T. G. Mertens and H. Verschelde, Fine Structure of Jackiw-Teitelboim Quantum Gravity, arXiv:1812.00918.
- [45] A. M. Garcia-Garcia, T. Nosaka, D. Rosa and J. J. M. Verbaarschot Quantum chaos transition in a two-site SYK model dual to an eternal traversable wormhole, arXiv:1901.06031.
- [46] S. Sachdev, Universal low temperature theory of charged black holes with AdS₂ horisons, arXiv:1902.04078.
- [47] A. Blommaert, T. G. Mertens and H. Verschelde, Clocks and Rods in Jackiw-Teitelboim Quantum Gravity, arXiv:1902.11194.
- [48] V. V. Belokurov and E. T. Shavgulidze, Exact solution of the Schwarzian theory, Phys. Rev. D 96 (2017) 101701(R), [arXiv:1705.02405].
- [49] V. V. Belokurov and E. T. Shavgulidze, Correlation functions in the Schwarzian theory, JHEP 11 (2018) 036, [arXiv:1804.00424].
- [50] V. V. Belokurov and E. T. Shavgulidze, Unusual view of the Schwarzian theory, Mod. Phys. Lett. A 33 (2018) 1850221, [arXiv:1806.05605].
- [51] V. V. Belokurov and E. T. Shavgulidze, Polar decomposition of the Wiener measure: Schwarzian theory versus conformal quantum mechanics, arXiv:1812.04039, (to appear in Theor. Math. Phys.).
- [52] S. Lang, " $SL_2(\mathbf{R})$ ". Addison-Wesley Publishing. 1975.
- [53] E. T. Shavgulidze, An example of a measure quasi-invariant with respect to the action of a group of diffeomorphisms of the circle, Funct. Anal. Appl. 12 (1978) 203.
- [54] E. T. Shavgulidze, A measure quasi-invariant with respect to the action of a group of diffeomorphisms of a finite-dimensional manifold, Sov. Math. Dokl. 38 (1988) 622.
- [55] E. T. Shavgulidze, Some Properties of Quasi-Invariant Measures on Groups of Diffeomorphisms of the Circle, Russ. J. Math. Phys. 7 (2000) 464.
- [56] V. V. Belokurov and E. T. Shavgulidze, Extraordinary Properties of Functional Integrals and Groups of Diffeomorphisms, Phys. Part. Nucl. 48 (2017) 267.
- [57] A. Weil, L'integration dans les groupes topologiques et ses applications, Actual. Scient. et Ind., 869, Paris: Herman, 1940.
- [58] A. P. Prudnikov, Yu. A. Brychkov, O. I. Marichev, Integrals and Series. V. 1. Elementary Functions, Gordon Breach: New York-London, 1986.