

A Note on the Well-posedness of Prandtl Equations in Dimension Two

Jincheng Gao[†] Daiwen Huang[‡] Zheng-an Yao[†]

[†]School of Mathematics, Sun Yat-sen University,
Guangzhou, 510275, P.R.China

[‡]Institute of Applied Physics and Computational Mathematics,
Beijing, 100088, P.R.China

Abstract

In this paper, we investigate the local-in-time well-posedness for the two-dimensional Prandtl equations in weighted Sobolev spaces under the Oleinik's monotonicity condition. Due to the loss of tangential derivative caused by vertical velocity appearing in convective term, we add with artificial horizontal viscosity term to construct an approximate system that can obtain local-in-time well-posedness results easily. For this approximate system, we construct a new weighted norm for the vorticity to derive a positive life time (independent of artificial viscosity coefficient) and obtain uniform bound for vorticity in this weighted norm. Then, based on compactness argument, we prove the solution of approximate system converging to the solution of original Prandtl equations, and hence, obtain the local-in-time well-posedness result for the Prandtl equations with any large initial data, which improves the recent work [10].

1 Introduction

Throughout this paper, we are concerned with the two-dimensional Prandtl equations, derived by Ludwig Prandtl [1], in a periodic domain $\mathbb{T} \times \mathbb{R}^+ := \{(x, y) : x \in \mathbb{R}/\mathbb{Z}, 0 \leq y < +\infty\}$:

$$\begin{cases} \partial_t u + u \partial_x u + v \partial_y u - \partial_y^2 u + \partial_x p = 0, \\ \partial_x u + \partial_y v = 0, \\ u|_{y=0} = v|_{y=0} = 0, \quad \lim_{y \rightarrow +\infty} u = U(t, x), \\ u|_{t=0} = u_0, \end{cases} \quad (1.1)$$

where the velocity field $(u, v) := (u(t, x, y), v(t, x, y))$ is unknown, and the initial data $u_0 := u_0(x, y)$ and the outer flow $U := U(t, x)$ are given and satisfy the compatibility conditions:

$$u_0|_{y=0} = 0, \quad \lim_{y \rightarrow +\infty} u_0 = U|_{t=0}. \quad (1.2)$$

Furthermore, the given scalar pressure $p := p(t, x)$ and the outer flow U satisfy the Bernoulli's law:

$$\partial_t U + U \partial_x U = -\partial_x p. \quad (1.3)$$

Note that the Prandtl equations mentioned above arise from the vanishing viscosity limit of Navier-Stokes equations in a domain with Dirichlet boundary condition. This is due to the formation of a

Email: gaojch5@mail.sysu.edu.cn(J.C.Gao), hdw55@tom.com(D.W.Huang) mcsyao@mail.sysu.edu.cn(Z.A.Yao).

boundary layer, where the solution undergoes a sharp transition from a solution of the Euler system to the zero non-slip boundary condition on boundary of the Navier-Stokes system. This boundary layer satisfies the Prandtl boundary layer equations formally. The first systematic work in rigorous mathematics was obtained by Oleinik [2, 3] in which she established the local in time well-posedness of Prandtl equations in dimension two by applying the Crocco transformation under the monotonicity condition on the tangential velocity field in the normal direction to the boundary. For more extensional mathematical results, the interested readers can refer to the classical book finished by Oleinik and Samokhin [4]. In addition to Oleinik’s monotonicity assumption on the velocity field, by imposing a so-called favorable condition on the pressure, Xin and Zhang [5] obtained the existence of global weak solutions to the Prandtl equation. As mentioned in [6–8], the local in time well-posedness of Prandtl equation for the initial data in Sobolev space is an open problem. Then, the researchers in [9] and [10] independently used the nonlinear cancelation method to establish well-posedness theory for the two-dimensional Prandtl equations in the framework of Sobolev spaces. *Note that the local-in-time well-posedness result in $H^{s,\gamma}$ Sobolev framework, obtained by Masmoudi and Wong [10], required there exists a small constant δ_0 such that*

$$\sum_{|\alpha|\leq 2} |(1+y)^{\sigma+\alpha_2} D^\alpha w_0|^2 \leq (2\delta_0)^{-2}, \quad (1.4)$$

and for $s = 4$, they required additionally

$$\|w_0\|_{H^{s,\gamma}} \leq C\delta_0^{-1}. \quad (1.5)$$

Thus, the main target in this paper is to remove the conditions (1.4)-(1.5). In other words, the local-in-time well-posedness results for the Prandtl equations (1.1) will be established for any large initial data.

Finally, we point out that it is an outstanding open problem to rigorously justify the validity of expansion in the inviscid limit. On one hand, within space of functions that are analytic, Sammartino and Caffisch [11, 12] obtained the well-posedness in the framework of analytic functions without the monotonicity condition on the velocity field and justified the boundary layer expansion for the unsteady incompressible Navier-Stokes equations. On the other hand, for more “realistic” functional settings, Guo and Nguyen[13] justified the boundary layer expansion for the steady incompressible flow with a non-slip boundary condition on a moving plate. This result was generalized to the case of non-moving or with external forcing(cf.[14, 15]). For more results in this direction, the interested readers can refer to [16–18] and references therein.

In this work, we will consider the Prandtl equations (1.1) under Oleinik’s monotonicity assumption:

$$w := \partial_y u > 0.$$

Under this hypothesis, one must further assume $U > 0$. Let us first introduce some weighted Sobolev spaces for later use. Denoting the vorticity $w := \partial_y u$, we define the weighted Sobolev space $H^{s,\gamma}$ for w by

$$H^{s,\gamma} := \{w : \mathbb{T} \times \mathbb{R}^+ \rightarrow \mathbb{R} : \|w\|_{H^{s,\gamma}} < \infty\},$$

where the weighted $H^{s,\gamma}$ norm is defined by

$$\|w\|_{H^{s,\gamma}}^2 := \sum_{|\alpha|\leq s} \|(1+y)^{\gamma+\alpha_2} D^\alpha w\|_{L^2(\mathbb{T}\times\mathbb{R}^+)}^2,$$

where $D^\alpha = \partial_x^{\alpha_1} \partial_y^{\alpha_2}$. Here, the main idea is adding an extra weight $(1+y)$ for each y -derivative. This corresponds to the weight $\frac{1}{y}$ in the Hardy-type inequality. Let us define

$$\|w(t)\|_{\mathcal{B}^{s,\gamma,\sigma}}^2 := \|w(t)\|_{H^{s,\gamma}}^2 + \sum_{1\leq|\alpha|\leq 2} \|(1+y)^{\sigma+\alpha_2} D^\alpha w(t)\|_{L^\infty}^2, \quad (1.6)$$

and introduce the space

$$\tilde{H}_{\sigma, \delta_0}^{s, \gamma} := \{w : \mathbb{T} \times \mathbb{R}^+ \rightarrow \mathbb{R} \mid \|w(t)\|_{\mathcal{B}^{s, \gamma, \sigma}}^2 < +\infty, \quad (1+y)^\sigma w \geq \delta_0, \}$$

where $s \geq 4, \gamma \geq 1, \sigma > \gamma + \frac{1}{2}, \delta_0 \in (0, \frac{1}{2})$. Now, we can state our main result:

Theorem 1.1. *Let $s \geq 4$ be an even integer, $\gamma \geq 1, \sigma > \gamma + \frac{1}{2}$ and $\delta_0 \in (0, \frac{1}{2})$. Suppose the outer flow U satisfies*

$$M_U := \sum_{k=0}^{s/2+1} \sup_{0 \leq t \leq T} \|\partial_t^k U\|_{H^{s-2k+2}(\mathbb{T})}^2 < +\infty. \quad (1.7)$$

Assume that the initial tangential velocity $u_0 - U|_{t=0} \in H^{s, \gamma-1}$ and the initial vorticity $w_0 := \partial_y u_0 \in \tilde{H}_{\sigma, 2\delta_0}^{s, \gamma}$. Then there exist a times $T = T(s, \gamma, \sigma, \delta_0, \|w_0\|_{\mathcal{B}^{s, \gamma, \sigma}}, M_U) > 0$ and a unique classical solution (u, v) to the Prandtl equations (1.1)-(1.3) such that

$$\sup_{0 \leq t \leq T} \|w(t)\|_{\mathcal{B}^{s, \gamma, \sigma}}^2 \leq C_{s, \gamma, \sigma} \{1 + \|w_0\|_{\mathcal{B}^{s, \gamma, \sigma}}^8 + M_U^4\} < +\infty,$$

and

$$\min_{\mathbb{T} \times \mathbb{R}^+} (1+y)^\sigma w(t) \geq \delta_0,$$

for all $t \in [0, T]$.

Remark 1.1. *Defined the space*

$$H_{\sigma, \delta_0}^{s, \gamma} := \{w : \mathbb{T} \times \mathbb{R}^+ \rightarrow \mathbb{R} : \|w\|_{H^{s, \gamma}}^2 < +\infty, \quad \sum_{|\alpha| \leq 2} |(1+y)^{\sigma+\alpha_2} D^\alpha w|^2 \leq \frac{1}{\delta_0^2}, \quad (1+y)^\sigma w \geq \delta_0\},$$

Masmoudi and Wong [10] established the local-in-time well-posedness theory for the Prandtl equations (1.1) if the initial vorticity w_0 belongs to $H_{\sigma, 2\delta_0}^{s, \gamma}$ instead of $\tilde{H}_{\sigma, 2\delta_0}^{s, \gamma}$ (required in Theorem 1.1). In other words, they required the initial vorticity itself, first and second order derivatives with weight in L^∞ -norm should be controlled by δ_0^{-1} rather than being sufficiently large.

Remark 1.2. *When the well-posedness for the Prandtl equations (1.1) in the $H^{4, \gamma}$ -framework, Masmoudi and Wong [10] required the condition (1.5) additionally, which we do not need in Theorem 1.1.*

We now explain main difficulties of proving Theorem 1.1 as well as our strategies for overcoming them. In order to solve the Prandtl equations (1.1) in certain H^s Sobolev space, the main difficulty comes from the vertical velocity $v = -\partial_y^{-1} \partial_x u$ creates a loss of tangential derivative, so the standard energy methods can not apply directly. The main idea of establishing the well-posedness of Prandtl equations (1.1) is to apply the so-called vanishing viscosity and nonlinear cancellation methods. To this end, we consider the following approximate system (or regularized Prandtl equations cf.[10]):

$$\begin{cases} \partial_t u^\varepsilon + u^\varepsilon \partial_x u^\varepsilon + v^\varepsilon \partial_y u^\varepsilon - \varepsilon^2 \partial_x^2 u^\varepsilon - \partial_y^2 u^\varepsilon + \partial_x p^\varepsilon = 0, \\ \partial_x u^\varepsilon + \partial_y v^\varepsilon = 0, \\ u^\varepsilon|_{t=0} = u_0, \\ u^\varepsilon|_{y=0} = v^\varepsilon|_{y=0} = 0, \quad \lim_{y \rightarrow +\infty} u^\varepsilon(t, x, y) = U(t, x), \end{cases} \quad (1.8)$$

for any $\varepsilon > 0$. Here the quantities p^ε and U satisfy a regularized Bernoulli's law:

$$\partial_t U + U \partial_x U = \varepsilon^2 \partial_x^2 U - \partial_x p^\varepsilon. \quad (1.9)$$

We point out that the approximate system (1.8) will turn into the original Prandtl equations (1.1) as the parameter ε tends to zero. For any $\varepsilon > 0$, the local-in-time well-posedness of approximate system (1.8) is obtained easily in life time $[0, T^\varepsilon]$ (T^ε may depends on parameter ε), and hence, we hope to prove that the solution of (1.8) in life time $[0, T^\varepsilon]$ will converge to the solution of (1.1) when artificial viscosity ε tends to zero. For this purpose, we need to prove the time of existence T^ε stays bounded away from zero. Since the domain considered in this article is periodic, the main part of the boundary layer will vanish or being stable. Thus, the main difficulty to prove T^ε staying bounded away from zero arises from the vertical velocity v^ε . This can be overcome by the nonlinear cancellation methods developed in [10]. Here we also mention that the readers interested in the vanishing viscosity limit for incompressible Navier-Stokes equations can refer to [19–21] and references therein.

Now, let us explain the main idea to prove T^ε staying bounded away from zero and the main novelty to relax to additional conditions (1.4) and (1.5) required in [10]. First of all, we derive the weighted L^2 estimate for $D^\alpha w^\varepsilon$ for $|\alpha| \leq s$ and $\alpha_1 \leq s - 1$. This works since we are allowed to loss at least one x -regularity in these cases. Secondly, we introduce the quantity g_s^ε (cf.[10]):

$$g_s^\varepsilon := \partial_x^s w^\varepsilon - \frac{\partial_y w^\varepsilon}{w^\varepsilon} \partial_x^s (u^\varepsilon - U) \quad (1.10)$$

and the weighted norm

$$\|w^\varepsilon\|_{H_g^{s,\gamma}(\mathbb{T} \times \mathbb{R}^+)}^2 := \|(1+y)^\gamma g_s^\varepsilon\|_{L^2(\mathbb{T} \times \mathbb{R}^+)}^2 + \sum_{\substack{|\alpha| \leq s \\ \alpha_1 \leq s-1}} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2(\mathbb{T} \times \mathbb{R}^+)}^2, \quad (1.11)$$

provided $w^\varepsilon := \partial_y u^\varepsilon > 0$. As mentioned in [10], this quantity g_s^ε can avoid the loss of x -derivative by the nonlinear cancellation; in other words, the quantity g_s^ε in L^2 -norm will have uniform bound independent of ε . Without the additional condition (1.4) (required in [10]), we need to control the quantity $(1+y)^{\sigma+\alpha_2} D^\alpha w^\varepsilon$ ($1 \leq |\alpha| \leq 2$) in L^∞ -norm to close the estimate. Due to the weight index $\sigma > \gamma + \frac{1}{2}$, it is not easy to apply the quantity $\|w^\varepsilon\|_{H_g^{s,\gamma}}$ to control these quantities by the Sobolev inequality. Define

$$\|w^\varepsilon(t)\|_{\mathcal{B}_g^{s,\gamma,\sigma}}^2 := \|w^\varepsilon(t)\|_{H_g^{s,\gamma}}^2 + \sum_{1 \leq |\alpha| \leq 2} \|(1+y)^{\sigma+\alpha_2} D^\alpha w^\varepsilon(t)\|_{L^\infty}^2,$$

we may apply the maximum principle of heat equation to control quantities $(1+y)^{\sigma+\alpha_2} D^\alpha w^\varepsilon$ ($1 \leq |\alpha| \leq 2$) in L^∞ -norm by $\|w^\varepsilon(t)\|_{\mathcal{B}_g^{s,\gamma,\sigma}}^2$, initial and boundary data, which can be controlled by the quantity $\|w^\varepsilon\|_{H_g^{s,\gamma}}$ after using the Sobolev inequality. An important remark is that $\|w^\varepsilon(t)\|_{\mathcal{B}_g^{s,\gamma,\sigma}}$ is equivalent to $\|w^\varepsilon(t)\|_{\mathcal{B}^{s,\gamma,\sigma}}$ (see (B.5) and (B.6)), which has uniform bound on the life span time $[0, T^\varepsilon]$ (see Lemma C.1 in Appendix C). Thus, based on the estimates obtained above, we can choose the life span time T_a independent of ε such the quantity $\|w^\varepsilon(t)\|_{\mathcal{B}^{s,\gamma,\sigma}}$ has uniform bound on $[0, T_a]$. Then we can pass the limit $\varepsilon \rightarrow 0^+$ and obtain the existence and uniqueness of solution to the original Prandtl equations (1.1) by the approximate system (1.8). The weighed norm $\|\cdot\|_{\mathcal{B}_g^{s,\gamma,\sigma}}$ is not only equivalent to norm $\|\cdot\|_{\mathcal{B}^{s,\gamma,\sigma}}$, but also avoids the loss of tangential derivatives without the condition (1.4). This is the main novelty in our paper and help us improve the recent result [10]. But we should point out that the idea, which overcomes the loss of tangential derivative arising by vertical velocity $v = -\partial_y^{-1} \partial_x u$, comes from the nonlinear cancellation method developed in [10].

The rest of this paper is organized as follows. In Section 2, one establishes the a priori estimates for the approximate system (1.8). Some useful inequalities and important equivalent relations will be stated in Appendix A and B. Before we proceed, let us comment on our notation. Through this paper, all constants C may be different from line to line. Subscript(s) of a constant illustrates the dependence of the constant, for example, C_s is a constant depending on s only. Denote by ∂_y^{-1} the inverse of the derivative ∂_y , i.e., $(\partial_y^{-1} f)(y) := \int_0^y f(z) dz$.

2 A priori estimates

In this section, we will derive a priori estimates (independent of ε), which are crucial to prove the local-in-time well-posedness theory of solutions to original Prandtl equations (1.1). Denote vorticity $w^\varepsilon := \partial_y u^\varepsilon$, using the regularized Prandtl equations (1.8), we find that this vorticity satisfies the following evolution equations:

$$\begin{cases} \partial_t w^\varepsilon + u^\varepsilon \partial_x w^\varepsilon + v^\varepsilon \partial_y w^\varepsilon = \varepsilon^2 \partial_x^2 w^\varepsilon + \partial_y^2 w^\varepsilon, \\ w^\varepsilon|_{t=0} = w_0 := \partial_y u_0, \\ \partial_y w^\varepsilon|_{y=0} = \partial_x p^\varepsilon, \end{cases} \quad (2.1)$$

where the velocity field $(u^\varepsilon, v^\varepsilon)$ is given by

$$u^\varepsilon(t, x, y) := U(t, x) - \int_y^{+\infty} w^\varepsilon(t, x, \eta) d\eta, \quad (2.2)$$

and

$$v^\varepsilon(t, x, y) := - \int_0^y \partial_x u^\varepsilon(t, x, \eta) d\eta. \quad (2.3)$$

Next, we derive a life existence time T_a (independent of ε) such the quantity $\|w^\varepsilon(t)\|_{\mathcal{B}^{s,\gamma,\sigma}}$ owning a uniform bound. More precisely, we have the following results.

Theorem 2.1 (a priori estimates). *Let $s \geq 4$ be an even integer, $\gamma \geq 1, \sigma > \gamma + \frac{1}{2}, \delta_0 \in (0, \frac{1}{2})$, and $\varepsilon \in (0, 1]$, the smooth solution $(u^\varepsilon, v^\varepsilon, w^\varepsilon)$, defined on $[0, T^\varepsilon]$, to the regularized Prandtl equations (2.1)-(2.3). Under the assumptions of Theorem 1.1, there exists a time $T_a := T_a(s, \gamma, \sigma, \delta_0, \|w_0\|_{\mathcal{B}^{s,\gamma,\sigma}}, M_U) > 0$ independent of ε such the following estimates hold on*

$$\Omega(t) := \sup_{0 \leq \tau \leq t} \|w^\varepsilon(\tau)\|_{\mathcal{B}^{s,\gamma,\sigma}}^2 \leq C_{s,\gamma,\sigma} \{1 + \|w_0\|_{\mathcal{B}^{s,\gamma,\sigma}}^8 + M_U^4\}, \quad (2.4)$$

and

$$\min_{\mathbb{T} \times \mathbb{R}^+} (1+y)^\sigma w^\varepsilon(t, x, y) \geq \delta_0, \quad (2.5)$$

for all $t \in [0, \min(T_a, T^\varepsilon)]$.

Remark 2.1. *After having the results in Theorem 2.1 at hand, we can pass to the limit $\varepsilon \rightarrow 0^+$ in the regularized Prandtl equations (1.8)₁-(1.8)₄ and the regularized Bernoulli's law (1.9). Thus, it is easy to check that the limit functions (u, v) will solve the original Prandtl equations (1.1) with the Bernoulli's law (1.3) in the classical sense (cf. [10]). On the other hand, the uniqueness of Prandtl equations (1.1) has already been derived in [10] without the condition (1.4). In other words, the local-in-time well-posedness theory of solutions to the Prandtl equations (1.1) in Theorem 1.1 is a direct consequence of Theorem 2.1.*

Throughout this section, for any small constant $\delta \in (0, \frac{1}{2})$, we assume a priori assumption

$$(1+y)^\sigma w^\varepsilon(t, x, y) \geq \delta, \quad \forall (t, x, y) \in [0, T^\varepsilon] \times \mathbb{T} \times \mathbb{R}^+, \quad (2.6)$$

holds on. Let us define

$$\|w^\varepsilon(t)\|_{\mathcal{B}_g^{s,\gamma,\sigma}}^2 := \|w^\varepsilon(t)\|_{H_g^{s,\gamma}}^2 + Q(t), \quad \text{and} \quad \Omega_g(t) := \sup_{0 \leq \tau \leq t} \|w^\varepsilon(\tau)\|_{\mathcal{B}_g^{s,\gamma,\sigma}}^2, \quad (2.7)$$

where $Q(t)$ is defined by

$$Q(t) := \sum_{1 \leq |\alpha| \leq 2} \|(1+y)^{\sigma+\alpha_2} D^\alpha w^\varepsilon(t)\|_{L^\infty}^2. \quad (2.8)$$

2.1. Weighted Energy Estimates

In this subsection, we will derive the uniform weighted estimates for the vorticity w^ε , which plays an important role for us to find the uniform existence life time. Since one order tangential derivative loss is allowed, we may apply the energy method to establish the weighted estimates for the vorticity $D^\alpha w^\varepsilon$ ($|\alpha| \leq s, \alpha_1 \leq s - 1$).

Lemma 2.2. *Under the hypotheses of Theorem 2.1, we have the following estimate:*

$$\begin{aligned} & \frac{d}{dt} \sum_{\substack{|\alpha| \leq s \\ \alpha_1 \leq s-1}} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2}^2 + \sum_{\substack{|\alpha| \leq s \\ \alpha_1 \leq s-1}} \|(1+y)^{\gamma+\alpha_2} (\varepsilon \partial_x D^\alpha w^\varepsilon, \partial_y D^\alpha w^\varepsilon)\|_{L^2}^2 \\ & \leq C_{s,\gamma,\sigma,\delta} \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})}^8 + C_{s,\gamma,\sigma,\delta} (1 + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}^8 + \|w^\varepsilon\|_{H_g^{s,\gamma}}^8) \\ & \quad + C_{s,\gamma} (1 + \|w^\varepsilon\|_{H_g^{s,\gamma}})^{s-2} \|w^\varepsilon\|_{H_g^{s,\gamma}}^2 + C_s \sum_{k=0}^{s/2} \|\partial_t^k \partial_x p^\varepsilon\|_{H^{s-2k}(\mathbb{T})}^2, \end{aligned}$$

where the positive constants $C_s, C_{s,\gamma}$ and $C_{s,\gamma,\sigma,\delta}$ are independent of ε .

Proof. Differentiating the vorticity equation (2.1) with respect to x α_1 times and y α_2 times, and multiplying the resulting equality by $(1+y)^{2\gamma+2\alpha_2} D^\alpha w^\varepsilon$, we get after integrating over $\mathbb{T} \times \mathbb{R}^+$,

$$\frac{1}{2} \frac{d}{dt} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2}^2 + \varepsilon^2 \|(1+y)^{\gamma+\alpha_2} \partial_x D^\alpha w^\varepsilon\|_{L^2}^2 = J_1 + J_2 + J_3 + J_4, \quad (2.9)$$

where J_1, J_2, J_3 and J_4 are defined by

$$\begin{aligned} J_1 &= \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2} D^\alpha w^\varepsilon \partial_y^2 D^\alpha w^\varepsilon dx dy, \\ J_2 &= (\gamma + \alpha_2) \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2-1} v^\varepsilon |D^\alpha w^\varepsilon|^2 dx dy, \\ J_3 &= - \sum_{0 < \beta \leq \alpha} \binom{\alpha}{\beta} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2} D^\alpha w^\varepsilon \cdot D^\beta u^\varepsilon \partial_x D^{\alpha-\beta} w^\varepsilon dx dy, \\ J_4 &= - \sum_{0 < \beta \leq \alpha} \binom{\alpha}{\beta} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2} D^\alpha w^\varepsilon \cdot D^\beta v^\varepsilon \partial_y D^{\alpha-\beta} w^\varepsilon dx dy. \end{aligned}$$

First of all, integrating by part and applying the Cauchy inequality, we get

$$J_1 \leq -\frac{3}{4} \|(1+y)^{\gamma+\alpha_2} \partial_y D^\alpha w^\varepsilon\|_{L^2}^2 - \int_{\mathbb{T}} D^\alpha w^\varepsilon \partial_y D^\alpha w^\varepsilon|_{y=0} dx + C_{s,\gamma} \|w^\varepsilon\|_{H^{s,\gamma}}^2.$$

Next, integrating by part and applying the divergence-free condition, it follows

$$|J_2| \leq C_{s,\gamma} \left\| \frac{v^\varepsilon}{1+y} \right\|_{L^\infty} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2}^2.$$

Using the divergence-free condition, Sobolev and Hardy inequalities, we get for $\gamma \geq 1$

$$\begin{aligned} \left\| \frac{v^\varepsilon}{1+y} \right\|_{L^\infty} &\leq \left\| \frac{v^\varepsilon + y \partial_x U}{1+y} \right\|_{L^\infty} + \left\| \frac{y \partial_x U}{1+y} \right\|_{L^\infty} \\ &\leq C \left\{ \left\| \frac{v^\varepsilon + y \partial_x U}{1+y} \right\|_{L^2} + \left\| \frac{\partial_x v^\varepsilon + y \partial_x^2 U}{1+y} \right\|_{L^2} + \|\partial_y^2 \left\{ \frac{v^\varepsilon + y \partial_x U}{1+y} \right\}\|_{L^2} \right\} + \left\| \frac{y \partial_x U}{1+y} \right\|_{L^\infty} \\ &\leq C (\|\partial_x w^\varepsilon\|_{L^2} + \|\partial_x (u^\varepsilon - U)\|_{L^2} + \|\partial_x^2 (u^\varepsilon - U)\|_{L^2}) + \|\partial_x U\|_{L^\infty(\mathbb{T})} \\ &\leq C (\|w^\varepsilon\|_{H^{s,\gamma}} + \|\partial_x U\|_{L^\infty(\mathbb{T})}), \end{aligned} \quad (2.10)$$

and hence, J_2 can be estimated as

$$|J_2| \leq C_{s,\gamma} (\|\partial_x U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}) \|w^\varepsilon\|_{H^{s,\gamma}}^2.$$

Deal with term J_3 . Using the Hölder inequality, we get for $0 < \beta \leq \alpha$

$$\begin{aligned} & \left| \int (1+y)^{2\gamma+2\alpha_2} D^\alpha w^\varepsilon \cdot D^\beta u^\varepsilon \partial_x D^{\alpha-\beta} w^\varepsilon dx dy \right| \\ & \leq \|(1+y)^{\gamma+\alpha_2} D^\beta (u^\varepsilon - U) D^{\alpha+e_1-\beta} w^\varepsilon\|_{L^2} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2} \\ & \quad + \|(1+y)^{\gamma+\alpha_2} \partial_x^{\beta_1} U \partial_x^{\alpha_1+1-\beta_1} \partial_y^{\alpha_2} w^\varepsilon\|_{L^2} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2}. \end{aligned} \quad (2.11)$$

Using the Moser and Hardy inequalities, it follows

$$\|(1+y)^{\gamma+\alpha_2} D^\beta (u^\varepsilon - U) D^{\alpha+e_1-\beta} w^\varepsilon\|_{L^2} \leq C_{s,\gamma} \|w^\varepsilon\|_{H^{s,\gamma}}^2. \quad (2.12)$$

Applying the Sobolev and Wirtinger inequalities, we get

$$\|(1+y)^{\gamma+\alpha_2} \partial_x^{\beta_1} U \partial_x^{\alpha_1+1-\beta_1} \partial_y^{\alpha_2} w^\varepsilon\|_{L^2} \leq C_s \|\partial_x^s U\|_{L^\infty(\mathbb{T})} \|w^\varepsilon\|_{H^{s,\gamma}}. \quad (2.13)$$

Substituting estimates (2.12) and (2.13) into (2.11), we obtain

$$|J_3| \leq C_{s,\gamma} (\|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}) \|w^\varepsilon\|_{H^{s,\gamma}}^2.$$

Deal with the term J_4 . Using the Hölder inequality, we get for $0 < \beta \leq \alpha$

$$\begin{aligned} & \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2} D^\alpha w^\varepsilon \cdot D^\beta v^\varepsilon \partial_y D^{\alpha-\beta} w^\varepsilon dx dy \right| \\ & \leq \|(1+y)^{\gamma+\alpha_2} D^\beta (v^\varepsilon + y \partial_x U) D^{\alpha+e_2-\beta} w^\varepsilon\|_{L^2} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2} \\ & \quad + \|(1+y)^{\gamma+\alpha_2} D^\beta (y \partial_x U) D^{\alpha+e_2-\beta} w^\varepsilon\|_{L^2} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2}. \end{aligned} \quad (2.14)$$

Using the Sobolev and Wirtinger inequalities, it follows

$$\|(1+y)^{\gamma+\alpha_2} D^\beta (y \partial_x U) D^{\alpha+e_2-\beta} w^\varepsilon\|_{L^2} \leq C \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})} \|w^\varepsilon\|_{H^{s,\gamma}}. \quad (2.15)$$

For $|\alpha| \leq s-1$, we apply the Moser and Hardy inequalities to get for $i = 1, 2$,

$$\begin{aligned} & \|(1+y)^{\gamma+\alpha_2} D^\beta (v^\varepsilon + y \partial_x U) D^{\alpha+e_2-\beta} w^\varepsilon\|_{L^2} \\ & \leq C \|D^{e_i} (v^\varepsilon + y \partial_x U)\|_{H^{s-2, i-2}} \|w^\varepsilon\|_{H^{s-1, \gamma}} \leq C \|w^\varepsilon\|_{H^{s,\gamma}}^2. \end{aligned} \quad (2.16)$$

Similar, for $|\alpha| = s$ and $\alpha_1 \leq s-1$, we get for $\beta_2 \geq 1$

$$\begin{aligned} & \|(1+y)^{\gamma+\alpha_2} D^\beta (v^\varepsilon + y \partial_x U) D^{\alpha+e_2-\beta} w^\varepsilon\|_{L^2} \\ & = \|(1+y)^{\gamma+\alpha_2} \partial_x^{\beta_1} \partial_y^{\beta_2-1} \partial_x (u^\varepsilon - U) \partial_x^{\alpha_1-\beta_1} \partial_y^{\alpha_2-\beta_2} \partial_y w^\varepsilon\|_{L^2} \leq C \|w^\varepsilon\|_{H^{s,\gamma}}^2, \end{aligned} \quad (2.17)$$

and for $\beta_2 = 0$

$$\begin{aligned} & \|(1+y)^{\gamma+\alpha_2} D^\beta (v^\varepsilon + y \partial_x U) D^{\alpha+e_2-\beta} w^\varepsilon\|_{L^2} \\ & = \|(1+y)^{\gamma+\alpha_2} \partial_x^{\beta_1} (v^\varepsilon + y \partial_x U) \partial_x^{\alpha_1-\beta_1} \partial_y^{\alpha_2-1} \partial_y w^\varepsilon\|_{L^2} \\ & \leq \|v^\varepsilon + y \partial_x U\|_{H^{s-1, -1}} \|w^\varepsilon\|_{H^{s,\gamma}} \leq C \|w^\varepsilon\|_{H^{s,\gamma}}^2. \end{aligned} \quad (2.18)$$

Substituting the estimates (2.15)-(2.18) into (2.14), it follows

$$|J_4| \leq C_s (\|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}) \|w^\varepsilon\|_{H^{s,\gamma}}^2.$$

Plugging the estimates of J_1 through J_4 into the equality (2.9), we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2}^2 + \varepsilon^2 \|(1+y)^{\gamma+\alpha_2} \partial_x D^\alpha w^\varepsilon\|_{L^2}^2 + \frac{3}{4} \|(1+y)^{\gamma+\alpha_2} \partial_y D^\alpha w^\varepsilon\|_{L^2}^2 \\ & \leq - \int_{\mathbb{T}} D^\alpha w^\varepsilon \partial_y D^\alpha w^\varepsilon|_{y=0} dx + C_{s,\gamma} (1 + \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})}) + \|w^\varepsilon\|_{H^{s,\gamma}} \|w^\varepsilon\|_{H^{s,\gamma}}^2, \end{aligned} \quad (2.19)$$

which, together with the relation (B.7), yields directly

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2}^2 + \varepsilon^2 \|(1+y)^{\gamma+\alpha_2} \partial_x D^\alpha w^\varepsilon\|_{L^2}^2 + \frac{3}{4} \|(1+y)^{\gamma+\alpha_2} \partial_y D^\alpha w^\varepsilon\|_{L^2}^2 \\ & \leq - \int_{\mathbb{T}} D^\alpha w^\varepsilon \partial_y D^\alpha w^\varepsilon|_{y=0} dx + C_{s,\gamma,\sigma,\delta} (1 + \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})}^8 + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}^8 + \|w^\varepsilon\|_{H_g^{s,\gamma}}^8). \end{aligned} \quad (2.20)$$

Note the boundary term above can be estimated as follows(cf.[10]) for $|\alpha| \leq s-1$,

$$\left| \int_{\mathbb{T}} D^\alpha w^\varepsilon \partial_y D^\alpha w^\varepsilon|_{y=0} dx \right| \leq \frac{1}{12} \|(1+y)^{\gamma+\alpha_2+1} \partial_y^2 D^\alpha w^\varepsilon\|_{L^2}^2 + C \|w^\varepsilon\|_{H_g^{s,\gamma}}^2 \quad (2.21)$$

and for $|\alpha| = s$,

$$\left| \int_{\mathbb{T}} D^\alpha w^\varepsilon \partial_y D^\alpha w^\varepsilon|_{y=0} dx \right| \leq \begin{cases} \frac{1}{12} \|(1+y)^{\gamma+\alpha_2} \partial_y D^\alpha w^\varepsilon\|_{L^2}^2 + H(t), & \alpha_2 = 2k, k \in \mathbb{N}; \\ \frac{1}{12} \|(1+y)^{\gamma+\alpha_2+1} \partial_x^{\alpha_1-1} \partial_y^{\alpha_2+2} w^\varepsilon\|_{L^2}^2 + H(t), & \alpha_2 = 2k+1, k \in \mathbb{N}; \end{cases} \quad (2.22)$$

where $H(t) = C_s \sum_{l=0}^{s/2} \|\partial_t^l \partial_x p^\varepsilon\|_{H^{s-2l}(\mathbb{T})}^2 + C_{s,\gamma} (1 + \|w^\varepsilon\|_{H_g^{s,\gamma}})^{s-2} \|w^\varepsilon\|_{H_g^{s,\gamma}}^2$. Thus, plugging the estimates (2.21) and (2.22) into (2.20), and summing over α , we complete the proof of this lemma. \square

Next, define $a^\varepsilon := \frac{\partial_y w^\varepsilon}{w^\varepsilon}$ and $g_s^\varepsilon := \partial_x^s w^\varepsilon - a^\varepsilon \partial_x^s (w^\varepsilon - U)$, we are going to derive the L^2 estimate for $(1+y)^\gamma g_s^\varepsilon$ by using the standard energy methods. As mentioned in [10], this quantity g_s^ε will avoid the loss of x -derivative by a nonlinear cancellation. By routine checking, it is easy to justify that the quantity g_s^ε satisfies the evolution equation(cf.[10])

$$\begin{aligned} & (\partial_t + u^\varepsilon \partial_x + v^\varepsilon \partial_y - \varepsilon^2 \partial_x^2 - \partial_y^2) g_s^\varepsilon \\ & = 2\varepsilon^2 \left\{ \partial_x^{s+1} (u^\varepsilon - U) - \frac{\partial_x w^\varepsilon}{w^\varepsilon} \partial_x^s (u^\varepsilon - U) \right\} \partial_x a^\varepsilon + 2g_s^\varepsilon \partial_y a^\varepsilon - g_1^\varepsilon \partial_x^s U \\ & \quad - \sum_{j=1}^{s-1} \binom{s}{j} g_{j+1}^\varepsilon \partial_x^{s-j} u^\varepsilon - \sum_{j=1}^{s-1} \binom{s}{j} \partial_x^{s-j} v^\varepsilon \{ \partial_x^j \partial_y w^\varepsilon - a^\varepsilon \partial_x^j w^\varepsilon \} \\ & \quad + a^\varepsilon \sum_{j=0}^{s-1} \binom{s}{j} \partial_x^j (u^\varepsilon - U) \partial_x^{s-j+1} U. \end{aligned} \quad (2.23)$$

where $g_k^\varepsilon := \partial_x^k w^\varepsilon - a^\varepsilon \partial_x^k (w^\varepsilon - U)$.

Now, we are going to derive the following weighted energy estimate for g_s^ε :

Lemma 2.3. *Under the hypothesis of Theorem 2.1, we have the following estimate:*

$$\begin{aligned} & \frac{d}{dt} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2}^2 + \varepsilon^2 \|(1+y)^\gamma \partial_x g_s^\varepsilon\|_{L^2}^2 + \|(1+y)^\gamma \partial_y g_s^\varepsilon\|_{L^2}^2 \\ & \leq C \|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})}^4 + C_{s,\gamma,\sigma,\delta} \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})}^4 + C_{s,\gamma,\sigma,\delta} (1 + Q^4(t) + \|w^\varepsilon\|_{H_g^{s,\gamma}}^6), \end{aligned}$$

where the quantity $Q(t)$ is defined in (2.8), the positive constants C and $C_{s,\gamma,\sigma,\delta}$ are independent of ε .

Proof. Multiplying the equation (2.23) by $(1+y)^{2\gamma}g_s^\varepsilon$ and integrating over $\mathbb{T} \times \mathbb{R}^+$, we have

$$\frac{1}{2} \frac{d}{dt} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2}^2 + \varepsilon^2 \|(1+y)^\gamma \partial_x g_s^\varepsilon\|_{L^2}^2 = \sum_{i=1}^8 K_i, \quad (2.24)$$

where the terms $K_i (i = 1, \dots, 8)$ are defined by

$$\begin{aligned} K_1 &= \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon \partial_y^2 g_s^\varepsilon dx dy, & K_2 &= - \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon (u^\varepsilon \partial_x g_s^\varepsilon + v^\varepsilon \partial_y g_s^\varepsilon) dx dy, \\ K_3 &= 2\varepsilon^2 \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon \left\{ \partial_x^{s+1} (u^\varepsilon - U) - \frac{\partial_x w^\varepsilon}{w^\varepsilon} \partial_x^s (u^\varepsilon - U) \right\} \partial_x a^\varepsilon dx dy, \\ K_4 &= 2 \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |g_s^\varepsilon|^2 \partial_y a^\varepsilon dx dy, & K_5 &= \sum_{j=1}^{s-1} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_{j+1}^\varepsilon \partial_x^{s-j} u^\varepsilon dx dy, \\ K_6 &= \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_1^\varepsilon \partial_x^s U dx dy, & K_7 &= \sum_{j=0}^{s-1} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon a^\varepsilon \partial_x^j (u^\varepsilon - U) \partial_x^{s-j+1} U dx dy, \\ K_8 &= \sum_{j=1}^{s-1} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} \partial_x^{s-j} v^\varepsilon (\partial_x^j \partial_y w^\varepsilon - a^\varepsilon \partial_x^j w^\varepsilon) g_s^\varepsilon dx dy. \end{aligned}$$

Deal with the term K_1 . Integrating by part in the y -variable, we get

$$\begin{aligned} K_1 &= \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon \partial_y^2 g_s^\varepsilon dx dy \\ &= \int_{\mathbb{T}} g_s^\varepsilon \partial_y g_s^\varepsilon|_{y=0} dx - \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |\partial_y g_s^\varepsilon|^2 dx dy - 2\gamma \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma-1} g_s^\varepsilon \partial_y g_s^\varepsilon dx dy \\ &= -\frac{3}{4} \|(1+y)^\gamma \partial_y g_s^\varepsilon\|_{L^2}^2 - \int_{\mathbb{T}} g_s^\varepsilon \partial_y g_s^\varepsilon|_{y=0} dx + C_\gamma \|(1+y)^{\gamma-1} g_s^\varepsilon\|_{L^2}^2. \end{aligned}$$

Due to the definition of g_s^ε and boundary condition $\partial_y w^\varepsilon|_{y=0} = \partial_x p^\varepsilon$, it follows

$$\partial_y g_s^\varepsilon|_{y=0} = \partial_x^{s+1} p^\varepsilon + \frac{\partial_y^2 w^\varepsilon}{w^\varepsilon} \partial_x^s U|_{y=0} - \frac{\partial_y w^\varepsilon}{w^\varepsilon} g_s^\varepsilon|_{y=0}.$$

Using the Hölder and trace inequalities, we get

$$\begin{aligned} \left| \int_{\mathbb{T}} g_s^\varepsilon \partial_x^{s+1} p^\varepsilon|_{y=0} dx \right| &\leq \|g_s^\varepsilon|_{y=0}\|_{L^2(\mathbb{T})} \|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})} \\ &\leq \sqrt{2} \|g_s^\varepsilon\|_{L^2}^{\frac{1}{2}} \|\partial_y g_s^\varepsilon\|_{L^2}^{\frac{1}{2}} \|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})} \\ &\leq \frac{1}{8} \|\partial_y g_s^\varepsilon\|_{L^2}^2 + C \|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})}^{\frac{4}{3}} \|g_s^\varepsilon\|_{L^2}^{\frac{2}{3}}. \end{aligned}$$

Due to $(1+y)^\sigma w^\varepsilon \geq \delta$, it follows $w^\varepsilon|_{y=0} \geq \delta$, and hence, we get, after using $\partial_y w^\varepsilon|_{y=0} = \partial_x p^\varepsilon$,

$$\left| \int_{\mathbb{T}} g_s^\varepsilon \frac{\partial_y w^\varepsilon}{w^\varepsilon} g_s^\varepsilon|_{y=0} dx \right| \leq \frac{1}{8} \|\partial_y g_s^\varepsilon\|_{L^2}^2 + C_\delta \|\partial_x p^\varepsilon\|_{L^\infty(\mathbb{T})}^2 \|g_s^\varepsilon\|_{L^2}^2.$$

and

$$\begin{aligned} \left| \int_{\mathbb{T}} g_s^\varepsilon \frac{\partial_y^2 w^\varepsilon}{w^\varepsilon} \partial_x^s U|_{y=0} dx \right| &\leq \delta^{-1} \|\partial_x^s U\|_{L^\infty(\mathbb{T})} \|g_s^\varepsilon|_{y=0}\|_{L^2(\mathbb{T})} \|\partial_y^2 w^\varepsilon|_{y=0}\|_{L^2(\mathbb{T})} \\ &\leq 2\delta^{-1} \|\partial_x^s U\|_{L^\infty(\mathbb{T})} \|g_s^\varepsilon\|_{L^2}^{\frac{1}{2}} \|\partial_y g_s^\varepsilon\|_{L^2}^{\frac{1}{2}} \|\partial_y^2 w^\varepsilon\|_{L^2}^{\frac{1}{2}} \|\partial_y^3 w^\varepsilon\|_{L^2}^{\frac{1}{2}} \\ &\leq \frac{1}{8} \|\partial_y g_s^\varepsilon\|_{L^2}^2 + C_\delta \|\partial_x^s U\|_{L^\infty(\mathbb{T})}^{\frac{4}{3}} \|g_s^\varepsilon\|_{L^2}^{\frac{2}{3}} \|(\partial_y^2 w^\varepsilon, \partial_y^3 w^\varepsilon)\|_{L^2}^{\frac{4}{3}}. \end{aligned}$$

Thus, the term K_1 can be estimated as follows

$$K_1 \leq -\frac{1}{2} \|(1+y)^\gamma \partial_y g_s^\varepsilon\|_{L^2}^2 + C_{\gamma,\delta} (1 + \|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})}^2 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})}^2) (1 + \|w^\varepsilon\|_{H_g^{s,\gamma}}^2).$$

Deal with the term K_2 . Integrating by part and applying the divergence-free condition, it follows

$$|K_2| = \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon (u^\varepsilon \partial_x g_s^\varepsilon + v^\varepsilon \partial_y g_s^\varepsilon) dx dy \right| \leq C_\gamma \left\| \frac{v^\varepsilon}{1+y} \right\|_{L^\infty} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2}^2,$$

which, along with inequality (B.13), yields directly

$$|K_2| \leq C_{\gamma,\sigma,\delta} (1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}) \|w^\varepsilon\|_{H_g^{s,\gamma}}^3.$$

Deal with the term K_3 . First of all, using the Hölder inequality, we get

$$\begin{aligned} & \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon \frac{\partial_x w^\varepsilon}{w^\varepsilon} \partial_x^s (u^\varepsilon - U) \partial_x a^\varepsilon dx dy \right| \\ & \leq \left\| \frac{\partial_x w^\varepsilon}{w^\varepsilon} \right\|_{L^\infty} \|(1+y) \partial_x a^\varepsilon\|_{L^\infty} \|(1+y)^{\gamma-1} \partial_x^s (u^\varepsilon - U)\|_{L^2} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2}. \end{aligned} \quad (2.25)$$

Due to the fact $\partial_x a^\varepsilon = \frac{\partial_{xy} w^\varepsilon}{w^\varepsilon} - \frac{\partial_y w^\varepsilon \partial_x w^\varepsilon}{(w^\varepsilon)^2}$, we get, after using the inequality $(1+y)^\sigma w^\varepsilon \geq \delta$,

$$\left\| \frac{\partial_x w^\varepsilon}{w^\varepsilon} \right\|_{L^\infty} \leq \delta^{-1} \|(1+y)^\sigma \partial_x w^\varepsilon\|_{L^\infty} \quad (2.26)$$

and

$$\|(1+y) \partial_x a^\varepsilon\|_{L^\infty} \leq \delta^{-1} \|(1+y)^{\sigma+1} \partial_{xy} w^\varepsilon\|_{L^\infty} + \delta^{-2} \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty} \|(1+y)^\sigma \partial_x w^\varepsilon\|_{L^\infty}. \quad (2.27)$$

Substituting the inequalities (2.26) and (2.27) into (2.25), and applying inequality (B.9), we obtain

$$\left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon \frac{\partial_x w^\varepsilon}{w^\varepsilon} \partial_x^s (u^\varepsilon - U) \partial_x a^\varepsilon dx dy \right| \leq C_{\gamma,\sigma,\delta} (1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})}^2 + Q^2(t)) \|w^\varepsilon\|_{H_g^{s,\gamma}}^2. \quad (2.28)$$

On the other hand, by virtue of the Hölder inequality, we get

$$\begin{aligned} & \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon \partial_x^{s+1} (u^\varepsilon - U) \partial_x a^\varepsilon dx dy \right| \\ & \leq \|(1+y)^{\sigma+1} w^\varepsilon \partial_x a^\varepsilon\|_{L^\infty} \|(1+y)^{\gamma-\sigma-1} \frac{\partial_x^{s+1} (u^\varepsilon - U)}{w^\varepsilon}\|_{L^2} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2}. \end{aligned} \quad (2.29)$$

Due to the fact $(1+y)^\sigma w^\varepsilon \geq \delta$, it follows

$$\|(1+y)^{\sigma+1} w^\varepsilon \partial_x a^\varepsilon\|_{L^\infty} \leq \|(1+y)^{\sigma+1} \partial_{xy} w^\varepsilon\|_{L^\infty} + \delta^{-1} \|(1+y)^\sigma \partial_x w^\varepsilon\|_{L^\infty} \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty} \quad (2.30)$$

By routine checking, it follows

$$\partial_x g_s^\varepsilon = w^\varepsilon \partial_y \left\{ \frac{\partial_x^{s+1} (u^\varepsilon - U)}{w^\varepsilon} \right\} - \partial_x a^\varepsilon \partial_x^s (u^\varepsilon - U),$$

and hence, we get after applying the Hardy inequality

$$\begin{aligned} & \|(1+y)^{\gamma-\sigma-1} \frac{\partial_x^{s+1} (u^\varepsilon - U)}{w^\varepsilon}\|_{L^2} \\ & \leq C_{\gamma,\sigma} \left\{ \left\| \frac{\partial_x^{s+1} U}{w^\varepsilon|_{y=0}} \right\|_{L^2(\mathbb{T})} + \|(1+y)^{\gamma-\sigma} \partial_y \left\{ \frac{\partial_x^{s+1} (u^\varepsilon - U)}{w^\varepsilon} \right\}\|_{L^2} \right\} \\ & \leq C_{\gamma,\sigma,\delta} \left\{ \|\partial_x^{s+1} U\|_{L^2(\mathbb{T})} + \|(1+y)^\gamma \partial_x g_s^\varepsilon\|_{L^2} + \|(1+y) \partial_x a^\varepsilon\|_{L^\infty} \|(1+y)^{\gamma-1} \partial_x^s (u^\varepsilon - U)\|_{L^2} \right\}. \end{aligned} \quad (2.31)$$

Substituting inequalities (2.30) and (2.31) into (2.29), and applying the inequality (B.9), it follows

$$\begin{aligned} & \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} g_s^\varepsilon \partial_x^{s+1} (u^\varepsilon - U) \partial_x a^\varepsilon dx dy \right| \\ & \leq \frac{1}{4} \left\| (1+y)^\gamma \partial_x g_s^\varepsilon \right\|_{L^2}^2 + C_{\gamma, \sigma, \delta} (1 + \|\partial_x^{s+1} U\|_{L^2(\mathbb{T})}^2 + Q^2(t)) (1 + \|w^\varepsilon\|_{H_g^{s, \gamma}}^2). \end{aligned}$$

This and the inequality (2.28) imply directly

$$|K_3| \leq \frac{1}{2} \varepsilon^2 \left\| (1+y)^\gamma \partial_x g_s^\varepsilon \right\|_{L^2}^2 + C_{\gamma, \sigma, \delta} (1 + \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})}^2 + Q^2(t)) (1 + \|w^\varepsilon\|_{H_g^{s, \gamma}}^2).$$

Deal with the term K_4 . Indeed, it is easy to get

$$|K_4| = \left| 2 \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |g_s^\varepsilon|^2 \partial_y a^\varepsilon dx dy \right| \leq C \|\partial_y a^\varepsilon\|_{L^\infty} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2}^2. \quad (2.32)$$

Due to the fact $\partial_y a^\varepsilon = \frac{\partial_y^2 w^\varepsilon}{w^\varepsilon} - \left(\frac{\partial_y w^\varepsilon}{w^\varepsilon}\right)^2$, we obtain

$$\|\partial_y a^\varepsilon\|_{L^\infty} \leq \delta^{-1} \|(1+y)^\sigma \partial_y^2 w^\varepsilon\|_{L^\infty} + \delta^{-2} \|(1+y)^\sigma \partial_y w^\varepsilon\|_{L^\infty}^2.$$

This and the inequality (2.32) give immediately

$$|K_4| \leq C_\delta (1 + Q(t)) \|w^\varepsilon\|_{H_g^{s, \gamma}}^2.$$

Deal with the term K_5 . Using the Hölder inequality, it follows

$$|K_5| \leq C_s \|\partial_x^{s-j} u^\varepsilon\|_{L^\infty} \|(1+y)^\gamma g_{j+1}^\varepsilon\|_{L^2} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2}.$$

This and the inequalities (B.11) and (B.12) imply directly

$$|K_5| \leq C_{s, \gamma, \sigma} (1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})}^2 + Q(t)) (1 + \|w^\varepsilon\|_{H_g^{s, \gamma}}^3).$$

Similarly, it is easy to deduce

$$|K_6| \leq C_{\gamma, \delta} (1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})}^2 + Q(t)) \|w^\varepsilon\|_{H_g^{s, \gamma}}^2.$$

Deal with the term K_7 . Using the Hölder and Hardy inequalities, we get for $j = 0, \dots, s-1$

$$\begin{aligned} |K_7| & \leq C_s \|\partial_x^{s-j+1} U\|_{L^\infty(\mathbb{T})} \|(1+y) a^\varepsilon\|_{L^\infty} \|(1+y)^{\gamma-1} \partial_x^j (u^\varepsilon - U)\|_{L^2} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2} \\ & \leq C_{s, \gamma} \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})} \|(1+y) a^\varepsilon\|_{L^\infty} \|w^\varepsilon\|_{H_g^{s, \gamma}}^2, \end{aligned}$$

and hence, using the fact $\|(1+y) a^\varepsilon\|_{L^\infty} \leq \delta^{-1} \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}$, we get

$$|K_7| \leq C_{s, \gamma, \delta} (\|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})}^2 + Q(t)) \|w^\varepsilon\|_{H_g^{s, \gamma}}^2.$$

Deal with the term K_8 . For the case $j = 1$, it is easy to check that

$$\begin{aligned} & \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} \partial_x^{s-1} v^\varepsilon (\partial_{xy} w^\varepsilon - a^\varepsilon \partial_x w^\varepsilon) g_s^\varepsilon dx dy \\ & = \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+1} \frac{\partial_x^{s-1} v^\varepsilon + y \partial_x^s U}{1+y} (\partial_{xy} w^\varepsilon - a^\varepsilon \partial_x w^\varepsilon) g_s^\varepsilon dx dy \\ & \quad - \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+1} \frac{y \partial_x^s U}{1+y} (\partial_{xy} w^\varepsilon - a^\varepsilon \partial_x w^\varepsilon) g_s^\varepsilon dx dy. \end{aligned}$$

By virtue of the Hölder inequality, it follows

$$\begin{aligned}
 & \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+1} \frac{\partial_x^{s-1} v^\varepsilon + y \partial_x^s U}{1+y} (\partial_{xy} w^\varepsilon - a^\varepsilon \partial_x w^\varepsilon) g_s^\varepsilon dx dy \right| \\
 & \leq (\|(1+y)^{\gamma+1} \partial_{xy} w^\varepsilon\|_{L^\infty} + \|(1+y)a^\varepsilon\|_{L^\infty}) \|(1+y)^\gamma \partial_x w^\varepsilon\|_{L^\infty} \\
 & \quad \times \left\| \frac{\partial_x^{s-1} v^\varepsilon + y \partial_x^s U}{1+y} \right\|_{L^2} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2}
 \end{aligned} \tag{2.33}$$

and

$$\begin{aligned}
 & \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+1} \frac{y \partial_x^s U}{1+y} (\partial_{xy} w^\varepsilon - a^\varepsilon \partial_x w^\varepsilon) g_s^\varepsilon dx dy \right| \\
 & \leq (\|(1+y)^{\gamma+1} \partial_{xy} w^\varepsilon\|_{L^2} + \|(1+y)a^\varepsilon\|_{L^\infty}) \|(1+y)^\gamma \partial_x w^\varepsilon\|_{L^2} \\
 & \quad \times \|\partial_x^s U\|_{L^\infty(T)} \|L^2\| (1+y)^\gamma g_s^\varepsilon\|_{L^2}.
 \end{aligned} \tag{2.34}$$

Due to the fact $\|(1+y)a^\varepsilon\|_{L^\infty} \leq \delta^{-1} \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}$, applying the inequality (B.10) and Sobolev inequality to inequalities (2.33) and (2.34), we get

$$\left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} \partial_x^{s-1} v^\varepsilon (\partial_{xy} w^\varepsilon - a^\varepsilon \partial_x w^\varepsilon) g_s^\varepsilon dx dy \right| \leq C_{\gamma,\delta} (1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})}^2 + Q(t)) (1 + \|w^\varepsilon\|_{H_g^{s,\gamma}}^3). \tag{2.35}$$

On the other hand, by virtue of the Hölder inequality and estimate (B.13), we get for $j = 2, \dots, s-1$,

$$\begin{aligned}
 & \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} \partial_x^{s-j} v^\varepsilon (\partial_x^j \partial_y w^\varepsilon - a^\varepsilon \partial_x^j w^\varepsilon) g_s^\varepsilon dx dy \right| \\
 & \leq (\|(1+y)^{\gamma+1} \partial_x^j \partial_y w^\varepsilon\|_{L^2} + \|(1+y)a^\varepsilon\|_{L^\infty}) \|(1+y)^\gamma \partial_x^j w^\varepsilon\|_{L^2} \\
 & \quad \times \left\| \frac{\partial_x^{s-j} v^\varepsilon}{1+y} \right\|_{L^\infty} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2} \\
 & \leq C_{\gamma,\sigma,\delta} (1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})}^2 + Q(t)) \|w^\varepsilon\|_{H_g^{s,\gamma}}^3.
 \end{aligned}$$

This and the inequality (2.35) imply

$$|K_8| \leq C_{s,\gamma,\sigma,\delta} (1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})}^2 + Q(t)) (1 + \|w^\varepsilon\|_{H_g^{s,\gamma}}^3).$$

Substituting the estimates of K_1 through K_8 into equality (2.24), we complete the proof of lemma. \square

Based on the estimates obtained in Lemmas 2.2 and 2.3, we have the estimate:

$$\begin{aligned}
 & \frac{d}{dt} \|w^\varepsilon\|_{H_g^{s,\gamma}}^2 + \varepsilon^2 \sum_{\substack{|\alpha| \leq s \\ \alpha_1 \leq s-1}} \|(1+y)^{\gamma+\alpha_2} \partial_x D^\alpha w^\varepsilon\|_{L^2}^2 + \varepsilon^2 \|(1+y)^\gamma \partial_x g_s^\varepsilon\|_{L^2}^2 \\
 & \quad + \sum_{\substack{|\alpha| \leq s \\ \alpha_1 \leq s-1}} \|(1+y)^{\gamma+\alpha_2} \partial_y D^\alpha w^\varepsilon\|_{L^2}^2 + \|(1+y)^\gamma \partial_y g_s^\varepsilon\|_{L^2}^2 \\
 & \leq C_s \{ \|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})}^4 + \sum_{k=0}^{s/2} \|\partial_t^k \partial_x p^\varepsilon\|_{H^{s-2k}(\mathbb{T})}^2 \} + C_{s,\gamma,\sigma,\delta} \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})}^4 \\
 & \quad + C_{s,\gamma,\sigma,\delta} \{1 + Q^4(t) + \|w^\varepsilon\|_{H_g^{s,\gamma}}^{s+4}\}.
 \end{aligned} \tag{2.36}$$

Using the regularized Bernoulli's law (1.9), we get

$$\sum_{k=0}^{s/2} \|\partial_t^k \partial_x p^\varepsilon\|_{H^{s-2k}(\mathbb{T})}^2 \leq C_s \{1 + \sum_{k=0}^{s/2+1} \|\partial_t^k U\|_{H^{s-2k+2}(\mathbb{T})}^2\}. \tag{2.37}$$

Using the Sobolev inequality for dimension one, it follows

$$\|\partial_x^{s+1}U\|_{L^\infty(\mathbb{T})} \leq C\|\partial_x^{s+1}\|_{H^1(\mathbb{T})}.$$

This and the inequality (2.37) imply

$$\begin{aligned} & \|\partial_x^{s+1}p^\varepsilon\|_{L^2(\mathbb{T})}^4 + \|\partial_x^{s+1}U\|_{L^\infty(\mathbb{T})}^4 + \sum_{k=0}^{s/2} \|\partial_t^k \partial_x p^\varepsilon\|_{H^{s-2k}(\mathbb{T})}^2 \\ & \leq C_s \left\{ 1 + \sum_{k=0}^{s/2+1} \|\partial_t^k U\|_{H^{s-2k+2}(\mathbb{T})}^2 \right\}^4 \leq C_s (1 + M_U)^4. \end{aligned} \quad (2.38)$$

Substituting estimate (2.38) into (2.36), and integrating the resulting inequality over $[0, t]$, it follows

$$\sup_{0 \leq \tau \leq t} \|w^\varepsilon(\tau)\|_{H_g^{s,\gamma}}^2 \leq \|w_0\|_{H_g^{s,\gamma}}^2 + C_{s,\gamma,\sigma,\delta} (1 + M_U)^4 t + C_{s,\gamma,\sigma,\delta} \int_0^t \{1 + Q^4(\tau) + \|w^\varepsilon(\tau)\|_{H_g^{s,\gamma}}^{s+4}\} d\tau,$$

or equivalently,

$$\sup_{0 \leq \tau \leq t} \|w^\varepsilon(\tau)\|_{H_g^{s,\gamma}}^2 \leq \|w_0\|_{H_g^{s,\gamma}}^2 + C_{s,\gamma,\sigma,\delta} (1 + M_U)^4 t + C_{s,\gamma,\sigma,\delta} \Omega_g(t)^s t, \quad (2.39)$$

for all $s \geq 4$ and $\gamma \geq 1$.

2.2. Weighted L^∞ Estimates for Lower Order Terms

In this subsection, we will establish the estimate for the quantity $Q(t)$ to close the estimate. Since the weight index $\sigma > \gamma + \frac{1}{2}$, we can not close this L^∞ estimate by the Sobolev inequality and weighted energy estimates directly. Similar to [10], we apply the maximum principle of heat equation to control the quantity $Q(t)$ by its initial data, boundary condition and quantity $\Omega_g(t)$. Note that the boundary condition without weight can be controlled by $\Omega_g(t)$ owing to $\gamma \geq 1$.

Lemma 2.4. *Under the hypotheses of Theorem 2.1, we have the following estimates:*

$$\sup_{0 \leq \tau \leq t} Q(\tau) \leq e^{C_{s,\gamma,\sigma,\delta} \{1 + M_U + \Omega_g(t)\} t} \{ \|w_0\|_{\mathcal{B}_g^{s,\gamma,\sigma}}^2 + C_{s,\gamma,\sigma,\delta} (1 + M_U)^4 t + C_{s,\gamma,\sigma,\delta} \Omega_g(t)^s t \}, \quad (2.40)$$

and

$$(1 + y)^\sigma w^\varepsilon(t, x, y) \geq (1 + y)^\sigma w_0(x, y) - Ct(1 + M_U + \Omega(t)). \quad (2.41)$$

Proof. Let us define

$$I(t) := \sum_{1 \leq |\alpha| \leq 2} |(1 + y)^{\sigma + \alpha_2} D^\alpha w^\varepsilon(t)|^2,$$

similar to [10], we may check that the quantity $I(t)$ satisfies:

$$\{\partial_t + u^\varepsilon \partial_x + v^\varepsilon \partial_y - \varepsilon^2 \partial_x^2 - \partial_y^2\} I(t) \leq C_{s,\gamma,\sigma,\delta} \{1 + \|\partial_x^s U\|_{L^\infty}^2 + \|(1 + y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}^2 + \|w^\varepsilon\|_{H_g^{s,\gamma}}^2\} I(t).$$

This and the maximum principle in Lemma A.4 will give directly for all $t \in [0, T^\varepsilon]$

$$\|I(t)\|_{L^\infty(\Omega)} \leq \max\{e^{C_{s,\gamma,\sigma,\delta} (1 + M_U + \Omega_g(t)) t} \|I(0)\|_{L^\infty(\Omega)}, \max_{\tau \in [0, t]} \{e^{C_{s,\gamma,\sigma,\delta} (1 + M_U + \Omega_g(t)) (t-\tau)} \|I(\tau)|_{y=0}\|_{L^\infty(\mathbb{T})}\}\},$$

or equivalently

$$\sup_{0 \leq \tau \leq t} Q(\tau) \leq e^{C_{s,\gamma,\sigma,\delta} \{1 + M_U + \Omega_g(t)\} t} (\|I(0)\|_{L^\infty(\Omega)} + \|I(t)|_{y=0}\|_{L^\infty(\mathbb{T})}). \quad (2.42)$$

By virtue of $s \geq 4$, we apply the Sobolev inequality (A.3) to get

$$\|I(t)|_{y=0}\|_{L^\infty(\mathbb{T})} \leq C\|w^\varepsilon\|_{H_g^{s,\gamma}}^2.$$

This and the inequality (2.42) yield directly

$$\sup_{0 \leq \tau \leq t} Q(\tau) \leq e^{C_{s,\gamma,\sigma,\delta}\{1+M_U+\Omega_g(t)\}t}(Q(0) + \|w^\varepsilon\|_{H_g^{s,\gamma}}^2). \quad (2.43)$$

Substituting the estimate (2.39) into inequality (2.43), we get the inequality (2.40).

Finally, using the first equation of (2.1), it follows

$$\begin{aligned} \|(1+y)^\sigma \partial_t w^\varepsilon\|_{L^\infty} &\leq \varepsilon^2 \|(1+y)^\sigma \partial_x^2 w^\varepsilon\|_{L^\infty} + \|(1+y)^\sigma \partial_y^2 w^\varepsilon\|_{L^\infty} + \|(1+y)^\sigma \partial_x w^\varepsilon\|_{L^\infty} \|u^\varepsilon\|_{L^\infty} \\ &\quad + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty} \left\| \frac{v^\varepsilon}{1+y} \right\|_{L^\infty} \\ &\leq C(1 + \|\partial_x U\|_{L^\infty(\mathbb{T})}^2 + Q(t) + \|w^\varepsilon\|_{H^{s,\gamma}}^2). \end{aligned} \quad (2.44)$$

Due to the basic fact

$$w^\varepsilon(t, x, y) - w_0(x, y) = \int_0^t \partial_\tau w^\varepsilon(\tau, x, y) d\tau,$$

we get after using (2.44)

$$\begin{aligned} (1+y)^\sigma w^\varepsilon(t, x, y) &\geq (1+y)^\sigma w_0(x, y) - \int_0^t \|(1+y)^\sigma \partial_\tau w^\varepsilon(\tau)\|_{L^\infty} d\tau \\ &\geq (1+y)^\sigma w_0(x, y) - Ct(1 + \sup_{0 \leq \tau \leq t} \|\partial_x U(\tau)\|_{L^\infty(\mathbb{T})}^2 + \sup_{0 \leq \tau \leq t} \{\|w^\varepsilon(\tau)\|_{H^{s,\gamma}}^2 + Q(\tau)\}) \\ &\geq (1+y)^\sigma w_0(x, y) - Ct(1 + M_U + \Omega(t)). \end{aligned}$$

Therefore, we complete the proof of this lemma. □

From the estimates (2.39), (2.40) and (2.41), we have the estimates

$$\Omega_g(t) \leq 2e^{C_{s,\gamma,\sigma,\delta}\{1+M_U+\Omega_g(t)\}t} \{\|w_0\|_{\mathcal{B}_g^{s,\gamma,\sigma}}^2 + C_{s,\gamma,\sigma,\delta}(1+M_U)^4 t + C_{s,\gamma,\sigma,\delta} \Omega_g(t)^s t\}, \quad (2.45)$$

and

$$(1+y)^\sigma w^\varepsilon(t, x, y) \geq (1+y)^\sigma w_0(x, y) - Ct(1 + M_U + \Omega(t)). \quad (2.46)$$

The advantage of estimates (2.45) and (2.46) is that the constants C and $C_{s,\gamma,\sigma,\delta}$ are independent of the artificial viscosity ε .

2.3. Proof of Theorem 2.1

Based on the estimates obtained so far, we can complete the proof of Theorem 2.1 in this subsection. First of all, for any $\varepsilon > 0$, we can apply standard energy method to gain the regularity propagates from the initial data(see estimates (C.1) and (C.2) in Lemma C.1), that is to say on $[0, T^\varepsilon]$, we have

$$\Omega(T) = \sup_{0 \leq t \leq T} \|w^\varepsilon(t)\|_{\mathcal{B}^{s,\gamma,\sigma}}^2 < +\infty.$$

Moreover, we can also get from the initial data that (2.6) is valid on $[0, T^\varepsilon]$ (possibly by taking T^ε smaller). An important remark is that if $\Omega(T_{ax}) < +\infty$, the solution can be continued on $[0, T_{bx}]$, $T_{bx} > T_{ax}$ with $\Omega(T_{bx}) < +\infty$. This and the estimates (2.4)-(2.5) can guarantee that the solution can be continued on an interval of time independent of ε . Thus, it suffices to verify the estimates (2.4) and (2.5).

For two constants R and δ , which will be defined later, we define

$$T_*^\varepsilon := \sup\{T \in [0, 1] \mid \Omega(t) \leq R, \quad (1+y)^\sigma w^\varepsilon(t, x, y) \geq \delta, \quad \forall (t, x, y) \in [0, T] \times \mathbb{T} \times \mathbb{R}^+\}. \quad (2.47)$$

Recall the relations (see (B.5) and (B.6))

$$\Omega(t) \leq C_{\gamma, \sigma, \delta} \|\partial_x^s U\|_{L^\infty(T)}^4 + C_{\gamma, \sigma, \delta} (1 + \Omega_g(t)^2),$$

and

$$\Omega_g(t) \leq C_{\gamma, \sigma, \delta} (1 + \Omega(t)^2),$$

and hence, we get for all $t \leq T^\varepsilon$, after using the inequality (2.45),

$$\begin{aligned} \Omega(t) &\leq C_{s, \gamma, \sigma} \{1 + \Omega(0)^4 + M_U^2 + C_{s, \gamma, \sigma, \delta} (1 + M_U)^8 t^2 + C_{s, \gamma, \sigma, \delta} (1 + \Omega(t)^2)^{2s} t^2\} \\ &\quad \times e^{C_{s, \gamma, \sigma, \delta} \{1 + M_U + \Omega(t)^2\} t}. \end{aligned}$$

Then, we may conclude for $T \leq T_*^\varepsilon$

$$\begin{aligned} \Omega(T) &\leq C_{s, \gamma, \sigma} \{1 + \Omega(0)^4 + M_U^4 + C_{s, \gamma, \sigma, \delta} (1 + M_U)^8 T + C_{s, \gamma, \sigma, \delta} (1 + R^2)^{2s} T\} \\ &\quad \times e^{C_{s, \gamma, \sigma, \delta} \{1 + M_U + R^2\} T}. \end{aligned}$$

Choose constants $R = 8C_{s, \gamma, \sigma} \{1 + \Omega(0)^4 + M_U^4\}$ and $\delta = \frac{\delta_0}{2}$, we get

$$\Omega(T_1) \leq 4C_{s, \gamma, \sigma} \{1 + \Omega(0)^4 + M_U^4\} = \frac{R}{2},$$

where $T_1 := \min\left\{\frac{\ln 2}{C_{s, \gamma, \sigma, \delta} (1 + M_U + R^2)}, \frac{1 + \Omega(0)^4 + M_U^4}{2C_{s, \gamma, \sigma, \delta} (1 + M_U)^8}, \frac{1 + \Omega(0)^4 + M_U^4}{2C_{s, \gamma, \sigma, \delta} (1 + R^2)^{2s}}\right\}$. It follows from (2.46)

$$\min_{\mathbb{T} \times \mathbb{R}^+} (1+y)^\sigma w^\varepsilon(t) \geq \delta_0 = 2\delta, \quad t \in [0, T_2].$$

where $T_2 := \min\left\{T_1, \frac{\delta_0}{C(1 + M_U + R)}\right\}$. Obviously, we conclude that there exists a time $T_2 > 0$ depending only on $s, \gamma, \sigma, \delta_0, M_U$ and the initial data $\|w_0\|_{\mathcal{B}^{s, \gamma, \sigma}}$ (hence independent of ε) such that for all $T \leq \min\{T_2, T^\varepsilon\}$, the estimates (2.4) and (2.5) hold on. Of course, it holds that $T_2 \leq T_*^\varepsilon$. Otherwise, our criterion about the continuation of the solution would contradict the definition of T_*^ε in (2.47). Then, taking $T_a = T_2$, we obtain the estimate (2.5) and close the a priori assumption (2.6). Therefore, we complete the proof of Theorem 2.1.

A Calculus Inequalities

In this appendix, we will introduce some basic inequality that be used frequently in this paper. For the proof in detail, the interested readers can refer to [10].

Lemma A.1 (Hardy Type Inequalities). *Let function $f : \mathbb{T} \times \mathbb{R}^+ \rightarrow \mathbb{R}$.*

(i) *if $\lambda > -\frac{1}{2}$ and $\lim_{y \rightarrow +\infty} f(x, y) = 0$, then*

$$\|(1+y)^\lambda f\|_{L^2(\mathbb{T} \times \mathbb{R}^+)} \leq \frac{2}{2\lambda + 1} \|(1+y)^{\lambda+1} \partial_y f\|_{L^2(\mathbb{T} \times \mathbb{R}^+)}. \quad (A.1)$$

(ii) *if $\lambda < -\frac{1}{2}$, then*

$$\|(1+y)^\lambda f\|_{L^2(\mathbb{T} \times \mathbb{R}^+)} \leq \sqrt{-\frac{1}{2\lambda + 1}} \|f|_{y=0}\|_{L^2(\mathbb{T})} - \frac{2}{2\lambda + 1} \|(1+y)^{\lambda+1} \partial_y f\|_{L^2(\mathbb{T} \times \mathbb{R}^+)}. \quad (A.2)$$

Lemma A.2 (Sobolev-Type Inequality). *Let the proper function $f : \mathbb{T} \times \mathbb{R}^+ \rightarrow \mathbb{R}$. Then there exists a universal constant $C > 0$ such that*

$$\|f\|_{L^\infty(\mathbb{T} \times \mathbb{R}^+)} \leq C\{\|f\|_{L^2(\mathbb{T} \times \mathbb{R}^+)} + \|\partial_x f\|_{L^2(\mathbb{T} \times \mathbb{R}^+)} + \|\partial_y^2 f\|_{L^2(\mathbb{T} \times \mathbb{R}^+)}\}. \quad (\text{A.3})$$

Next, we state the Morse type inequality that will be used frequently when we deal with the convective term. For the sake of brevity, we omit the proof for inequality (A.4) since it can be guaranteed by the Sobolev inequality (A.3).

Lemma A.3 (Morse-Type Inequality). *Let f and g be proper functions, $\gamma \in \mathbb{R}$ and an integer $s \geq 3$, we have for all $|\alpha + \tilde{\alpha}| \leq s$*

$$\|(1+y)^{\gamma+\alpha+\tilde{\alpha}}(D^\alpha f \cdot D^{\tilde{\alpha}} g)(t, \cdot)\|_{L^2} \leq C\|f(t)\|_{H^{s,\gamma_1}} \|g(t)\|_{H^{s,\gamma_2}}, \quad (\text{A.4})$$

where $\gamma_1, \gamma_2 \in \mathbb{R}$ with $\gamma_1 + \gamma_2 = \gamma$, and $C > 0$ is a universal constant.

Finally, let us recall the maximum principle for bounded solutions to parabolic equations(cf.[10]).

Lemma A.4 (Maximum Principle for Parabolic Equations). *Let $\varepsilon \geq 0$. If $H \in C([0, T]; C^2(\mathbb{T} \times \mathbb{R}^+)) \cap C^1([0, T]; C^0(\mathbb{T} \times \mathbb{R}^+))$ is a bounded function which satisfies the differential inequality:*

$$\{\partial_t + b_1 \partial_x + b_2 \partial_y - \varepsilon^2 \partial_{xx} - \partial_{yy}\}H \leq fH \quad \text{in } [0, T] \times \mathbb{T} \times \mathbb{R}^+,$$

where the coefficients b_1, b_2 and f are continuous and satisfy

$$\left\| \frac{b_2}{1+y} \right\|_{L^\infty([0, T] \times \mathbb{T} \times \mathbb{R}^+)} < +\infty \quad \text{and} \quad \|f\|_{L^\infty([0, T] \times \mathbb{T} \times \mathbb{R}^+)} \leq \lambda,$$

then for any $t \in [0, T]$,

$$\sup_{\mathbb{T} \times \mathbb{R}^+} H(t) \leq \max\{e^{\lambda t} \|H(0)\|_{L^\infty(\mathbb{T} \times \mathbb{R}^+)}, \max_{\tau \in [0, t]} \{e^{\lambda(t-\tau)} \|H(\tau)|_{y=0}\|_{L^\infty(\mathbb{T})}\}\}.$$

B Almost Equivalence of Weighted Norms

In this section, we will state some estimates that will be used in section 2. First of all, we derive the relation between g_s^ε and $\partial_x^s w^\varepsilon$ as follows.

Lemma B.1. *Let $s \geq 4$ be an even integer, $\gamma \geq 1, \sigma \geq \gamma + \frac{1}{2}$, and $\varepsilon \in (0, 1]$, the smooth solution $(u^\varepsilon, v^\varepsilon, w^\varepsilon)$, defined on $[0, T^\varepsilon]$, to the regularized Prandtl equations (2.1)-(2.3). There exists a small constant $\delta \in (0, 1)$ such that $(1+y)^\sigma w^\varepsilon \geq \delta, \forall (t, x, y) \in [0, T^\varepsilon] \times \mathbb{T} \times \mathbb{R}^+$, then it holds on*

$$\|(1+y)^\gamma g_s^\varepsilon\|_{L^2} \leq C_{\gamma, \delta} (1 + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}) \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2}, \quad (\text{B.1})$$

and

$$\|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2} \leq C_{\gamma, \sigma, \delta} (1 + \|\partial_x^s U\|_{L^\infty(T)} + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}) \|w^\varepsilon\|_{H_g^{s,\gamma}}, \quad (\text{B.2})$$

where $g_s^\varepsilon := \partial_x^s w^\varepsilon - \frac{\partial_y w^\varepsilon}{w^\varepsilon} \partial_x^s (u^\varepsilon - U)$.

Proof. Using the definition $g_s^\varepsilon = \partial_x^s w^\varepsilon - \frac{\partial_y w^\varepsilon}{w^\varepsilon} \partial_x^s (u^\varepsilon - U)$ and Hardy inequality (A.2), it follows

$$\begin{aligned} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2} &\leq \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2} + \|(1+y)^\gamma \frac{\partial_y w^\varepsilon}{w^\varepsilon} \partial_x^s (u^\varepsilon - U)\|_{L^2} \\ &\leq \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2} + \left\| \frac{(1+y)^{\sigma+1} \partial_y w^\varepsilon}{(1+y)^\sigma w^\varepsilon} \right\|_{L^\infty} \|(1+y)^{\gamma-1} \partial_x^s (u^\varepsilon - U)\|_{L^2} \\ &\leq \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2} + C_{\gamma,\delta} \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty} \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2}, \end{aligned}$$

where we have used $(1+y)^\sigma w^\varepsilon \geq \delta$, and hence, we get

$$\|(1+y)^\gamma g_s^\varepsilon\|_{L^2} \leq C_{\gamma,\delta} (1 + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}) \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2}.$$

This implies inequality (B.1). On the other hand, we get from the definition of g_s^ε that

$$\|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2} \leq \|(1+y)^\gamma g_s^\varepsilon\|_{L^2} + \|(1+y)^\gamma \partial_y w^\varepsilon \frac{\partial_x^s (u^\varepsilon - U)}{w^\varepsilon}\|_{L^2} \quad (\text{B.3})$$

By routine checking, it follows the relation $g_s^\varepsilon = w^\varepsilon \partial_y \left\{ \frac{\partial_x^s (u^\varepsilon - U)}{w^\varepsilon} \right\}$, and hence using $u^\varepsilon|_{y=0} = 0$, we get

$$\frac{\partial_x^s (u^\varepsilon - U)}{w^\varepsilon} = -\frac{\partial_x^s U}{w^\varepsilon|_{y=0}} + \int_0^y \frac{g_s^\varepsilon}{w^\varepsilon} d\xi.$$

This and the condition $(1+y)^\sigma w^\varepsilon \geq \delta$ yield directly

$$\begin{aligned} &\|(1+y)^\gamma \partial_y w^\varepsilon \frac{\partial_x^s (u^\varepsilon - U)}{w^\varepsilon}\|_{L^2} \\ &\leq \delta^{-1} \|\partial_x^s U\|_{L^\infty(\mathbb{T})} \|(1+y)^\gamma \partial_y w^\varepsilon\|_{L^2} + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty} \|(1+y)^{\gamma-\sigma-1} \int_0^y \frac{g_s^\varepsilon}{w^\varepsilon} d\xi\|_{L^2} \\ &\leq C_\delta \|\partial_x^s U\|_{L^\infty(\mathbb{T})} \|(1+y)^\gamma \partial_y w^\varepsilon\|_{L^2} + C_{\gamma,\sigma,\delta} \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty} \|(1+y)^\gamma g_s^\varepsilon\|_{L^2}. \end{aligned} \quad (\text{B.4})$$

Plugging inequality (B.4) into (B.3), we obtain the inequality (B.2). \square

Based on the inequalities (B.1)-(B.2), and the definitions of $H^{s,\gamma}$ and $H_g^{s,\gamma}$, we can establish the following estimates, which are important relation for us to obtain the well-posedness for the Prandtl equations in Sobolev space.

Lemma B.2. *Let $s \geq 4$ be an even integer, $\gamma \geq 1, \sigma \geq \gamma + \frac{1}{2}$, and $\varepsilon \in (0, 1]$, the smooth solution $(u^\varepsilon, v^\varepsilon, w^\varepsilon)$, defined on $[0, T^\varepsilon]$, to the regularized Prandtl equations (2.1)-(2.3). There exists a small constant $\delta \in (0, 1)$ such that $(1+y)^\sigma w^\varepsilon \geq \delta$, $\forall (t, x, y) \in [0, T^\varepsilon] \times \mathbb{T} \times \mathbb{R}^+$, then it holds on*

$$\Omega(t) \leq C_{\gamma,\sigma,\delta} \|\partial_x^s U\|_{L^\infty(T)}^4 + C_{\gamma,\sigma,\delta} (1 + \Omega_g(t)^2), \quad (\text{B.5})$$

and

$$\Omega_g(t) \leq C_{\gamma,\sigma,\delta} (1 + \Omega(t)^2). \quad (\text{B.6})$$

Proof. Using the definition of $\|\cdot\|_{H^{s,\gamma}}$ and estimate (B.2), it follows

$$\|w^\varepsilon\|_{H^{s,\gamma}} \leq C_{\gamma,\sigma,\delta} (1 + \|\partial_x^s U\|_{L^\infty(T)} + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}) \|w^\varepsilon\|_{H_g^{s,\gamma}}, \quad (\text{B.7})$$

which along with Cauchy inequality implies directly

$$\begin{aligned} \|w^\varepsilon(t)\|_{\mathcal{B}^{s,\gamma,\sigma}}^2 &\leq C_{\gamma,\sigma,\delta} (1 + \|\partial_x^s U\|_{L^\infty(T)}^2 + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}^2) \|w^\varepsilon\|_{H_g^{s,\gamma}}^2 \\ &\quad + \sum_{1 \leq |\alpha| \leq 2} \|(1+y)^{\sigma+\alpha_2} D^\alpha w^\varepsilon(t)\|_{L^\infty}^2 \\ &\leq C_{\gamma,\sigma,\delta} \|\partial_x^s U\|_{L^\infty(T)}^4 + C_{\gamma,\sigma,\delta} (1 + \|w^\varepsilon(t)\|_{\mathcal{B}^{s,\gamma,\sigma}}^4). \end{aligned}$$

This yields inequality (B.5). Similarly, it follows from inequality (B.1) that

$$\|w^\varepsilon\|_{H_g^{s,\gamma}} \leq C_{\gamma,\sigma,\delta}(1 + \|(1+y)^{\sigma+1}\partial_y w^\varepsilon\|_{L^\infty})\|w^\varepsilon\|_{H^{s,\gamma}}, \quad (\text{B.8})$$

and hence, we obtain

$$\|w^\varepsilon(t)\|_{\mathcal{B}^{s,\gamma,\sigma}}^2 \leq C_{\gamma,\sigma,\delta}(1 + \|w^\varepsilon(t)\|_{\mathcal{B}^{s,\gamma,\sigma}}^4).$$

Then we prove the inequality (B.6). □

Next, we establish some estimates for the quantity $(u^\varepsilon, v^\varepsilon, g_k^\varepsilon)$ in weighted L^2 -norm.

Lemma B.3. *Let $s \geq 4$ be an even integer, $\gamma \geq 1, \sigma \geq \gamma + \frac{1}{2}$, and $\varepsilon \in (0, 1]$, the smooth solution $(u^\varepsilon, v^\varepsilon, w^\varepsilon)$, defined on $[0, T^\varepsilon]$, to the regularized Prandtl equations (2.1)-(2.3). There exists a small constant $\delta \in (0, 1)$ such that $(1+y)^\sigma w^\varepsilon \geq \delta, \forall (t, x, y) \in [0, T^\varepsilon] \times \mathbb{T} \times \mathbb{R}^+$, then it holds on:*

(i) For all $k = 0, 1, \dots, s$,

$$\|(1+y)^{\gamma-1}\partial_x^k(u^\varepsilon - U)\|_{L^2} \leq C_{\gamma,\sigma,\delta}(1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|(1+y)^{\sigma+1}\partial_y w^\varepsilon\|_{L^\infty})\|w^\varepsilon\|_{H_g^{s,\gamma}}, \quad (\text{B.9})$$

(ii) For all $k = 0, 1, \dots, s-1$,

$$\left\| \frac{\partial_x^k v^\varepsilon + y\partial_x^{k+1}U}{1+y} \right\|_{L^2} \leq C_{\gamma,\sigma,\delta}(1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|(1+y)^{\sigma+1}\partial_y w^\varepsilon\|_{L^\infty})\|w^\varepsilon\|_{H_g^{s,\gamma}}, \quad (\text{B.10})$$

(iii) For all $k = 1, 2, \dots, s$

$$\|(1+y)^\gamma g_k^\varepsilon\|_{L^2} \leq C_{\gamma,\delta}(1 + \|(1+y)^{\sigma+1}\partial_y w^\varepsilon\|_{L^\infty})\|w^\varepsilon\|_{H_g^{s,\gamma}}. \quad (\text{B.11})$$

Proof. (i) It follows from Hardy inequality (A.1) that $\|(1+y)^{\gamma-1}\partial_x^k(u^\varepsilon - U)\|_{L^2} \leq C_\gamma\|(1+y)^\gamma\partial_x^k w^\varepsilon\|_{L^2}$, and hence inequality (B.9) is a direct consequence of the inequality (B.7).

(ii) Due to the Hardy inequality (A.2) and divergence-free condition, it follows

$$\left\| \frac{\partial_x^k v^\varepsilon + y\partial_x^{k+1}U}{1+y} \right\|_{L^2} \leq C\|\partial_x^k(\partial_y v^\varepsilon + \partial_x U)\|_{L^2} \leq C\|\partial_x^{k+1}(u^\varepsilon - U)\|_{L^2},$$

and hence, we get inequality (B.10) after using inequality (B.9).

(iii) For the case $s = 1, 2, \dots, s-1$, we get after using $(1+y)^\sigma w^\varepsilon \geq \delta$,

$$\begin{aligned} \|(1+y)^\gamma g_k^\varepsilon\|_{L^2} &\leq \|(1+y)^\gamma\partial_x^k w^\varepsilon\|_{L^2} + \left\| \frac{(1+y)^{\sigma+1}\partial_y w^\varepsilon}{(1+y)^{\sigma} w^\varepsilon} \right\|_{L^\infty} \|(1+y)^{\gamma-1}\partial_x^k(u^\varepsilon - U)\|_{L^2} \\ &\leq \|(1+y)^\gamma\partial_x^k w^\varepsilon\|_{L^2} + C_\delta \|(1+y)^{\sigma+1}\partial_y w^\varepsilon\|_{L^\infty} \|(1+y)^{\gamma-1}\partial_x^k(u^\varepsilon - U)\|_{L^2}. \end{aligned}$$

This and the Hardy inequality yield directly

$$\|(1+y)^\gamma g_k^\varepsilon\|_{L^2} \leq C_{\gamma,\delta}(1 + \|(1+y)^{\sigma+1}\partial_y w^\varepsilon\|_{L^\infty})\|(1+y)^\gamma\partial_x^k w^\varepsilon\|_{L^2}.$$

Therefore, we complete the proof of this lemma. □

Finally, we establish some estimates for the quantity $(u^\varepsilon, v^\varepsilon)$ in L^∞ -norm.

Lemma B.4. *Let $s \geq 4$ be an even integer, $\gamma \geq 1, \sigma \geq \gamma + \frac{1}{2}$, and $\varepsilon \in (0, 1]$, the smooth solution $(u^\varepsilon, v^\varepsilon, w^\varepsilon)$, defined on $[0, T^\varepsilon]$, to the regularized Prandtl equations (2.1)-(2.3). There exists a small constant $\delta \in (0, 1)$ such that $(1+y)^\sigma w^\varepsilon \geq \delta, \forall (t, x, y) \in [0, T^\varepsilon] \times \mathbb{T} \times \mathbb{R}^+$, then it holds on:*

(i) For all $k = 0, 1, \dots, s-1$,

$$\|\partial_x^k u^\varepsilon\|_{L^\infty} \leq C_{\gamma,\sigma,\delta}(1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|(1+y)^{\sigma+1}\partial_y w^\varepsilon\|_{L^\infty})(1 + \|w^\varepsilon\|_{H_g^{s,\gamma}}) \quad (\text{B.12})$$

(ii) For all $k = 0, 1, \dots, s-2$,

$$\left\| \frac{\partial_x^k v^\varepsilon}{1+y} \right\|_{L^\infty} \leq C_{\gamma,\sigma,\delta}(1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|(1+y)^{\sigma+1}\partial_y w^\varepsilon\|_{L^\infty})\|w^\varepsilon\|_{H_g^{s,\gamma}}. \quad (\text{B.13})$$

Proof. By virtue of the Sobolev inequality (A.3) and estimate (B.9), it follows

$$\begin{aligned} \|\partial_x^k u^\varepsilon\|_{L^\infty} &\leq C(\|\partial_x^k(u^\varepsilon - U)\|_{L^2} + \|\partial_x^{k+1}(u^\varepsilon - U)\|_{L^2} + \|\partial_x^k \partial_y w^\varepsilon\|_{L^2}) + \|\partial_x^k U\|_{L^\infty(\mathbb{T})} \\ &\leq C_{\gamma, \sigma, \delta}(1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|(1+y)^{\sigma+1} \partial_y w^\varepsilon\|_{L^\infty}) \|w^\varepsilon\|_{H_g^{s, \gamma}} + \|\partial_x^k U\|_{L^\infty(\mathbb{T})}. \end{aligned} \quad (\text{B.14})$$

Due to the fact $U = \int_0^{+\infty} w^\varepsilon dy$, we get

$$\|U\|_{L^2(\mathbb{T})} \leq C_\gamma \|(1+y)^\gamma w^\varepsilon\|_{L^2},$$

and hence, we get after using the Sobolev and Wirtinger inequalities for $k = 0, 1, \dots, s-1$,

$$\|\partial_x^k U\|_{L^\infty(\mathbb{T})} \leq C(\|\partial_x^k U\|_{L^2(\mathbb{T})} + \|\partial_x^{k+1} U\|_{L^2(\mathbb{T})}) \leq C_\gamma \|(1+y)^\gamma w^\varepsilon\|_{L^2} + \|\partial_x^s U\|_{L^\infty(\mathbb{T})}. \quad (\text{B.15})$$

Submitting inequality (B.15) into (B.14), we obtain the inequality (B.12). Finally, the inequality (B.13) is consequence of Sobolev inequality (A.3), estimates (B.9) and (B.10). Thus, we complete the proof of this lemma. \square

C Existence for the Regularized Prandtl Equations

In this section, we state the local in time well-posedness theory for the regularized Prandtl equations (2.1)-(2.3). More precisely, we have the following results:

Lemma C.1. *Let $s \geq 4$ be an even integer, $\gamma \geq 1, \sigma > \gamma + \frac{1}{2}, \delta_0 \in (0, \frac{1}{2})$ and $\varepsilon \in (0, 1]$. If the vorticity $w_0 \in \tilde{H}_{\sigma, 2\delta_0}^{s, \gamma}$, U and p^ε are given and satisfy the regularized Bernoulli's law (1.9) and the regularity assumption (1.7), then there exist a time*

$$T^\varepsilon := T(s, \gamma, \sigma, \delta_0, \varepsilon, \|w_0\|_{\mathcal{B}^{s, \gamma, \sigma}}, M_U) > 0,$$

and a solution w^ε , to the regularized vorticity system (2.1)-(2.3), satisfying the estimates:

$$\Omega(t) := \sup_{0 \leq \tau \leq t} \|w^\varepsilon(\tau)\|_{\mathcal{B}^{s, \gamma, \sigma}}^2 \leq C_*(1 + \|w_0\|_{\mathcal{B}^{s, \gamma, \sigma}}^2) < +\infty, \quad (\text{C.1})$$

and

$$(1+y)^\sigma w^\varepsilon(t, x, y) \geq c_* \delta_0, \quad (\text{C.2})$$

for all $(t, x, y) \in [0, T^\varepsilon] \times \mathbb{T} \times \mathbb{R}^+$, here C_*, c_* are positive constants and $c_* \in (0, 1)$.

Proof. We only establish the a priori estimates (C.1) and (C.2), and the well-posedness results of the regularized vorticity system (2.1)-(2.3) can be obtained immediately(cf.[10]).

Step 1: $H^{s, \gamma}$ -estimates. Differentiating the regularized vorticity equation (2.1) with differential operator D^α ($|\alpha| \leq s$), we get

$$\{\partial_t + u^\varepsilon \partial_x + v^\varepsilon \partial_y - \varepsilon^2 \partial_x^2 - \partial_y^2\} D^\alpha w^\varepsilon = -[D^\alpha, u^\varepsilon \partial_x] w^\varepsilon - [D^\alpha, v^\varepsilon \partial_y] w^\varepsilon.$$

Multiplying this equation by $(1+y)^{2\gamma+2\alpha_2} D^\alpha w^\varepsilon$ and integrating over $\mathbb{T} \times \mathbb{R}^+$, it follows

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2} |D^\alpha w^\varepsilon|^2 dx dy + \varepsilon^2 \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2} |\partial_x D^\alpha w^\varepsilon|^2 dx dy \\ &= (\gamma + \alpha_2) \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2-1} v^\varepsilon |D^\alpha w^\varepsilon|^2 dx dy + \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2} \partial_y^2 D^\alpha w^\varepsilon \cdot D^\alpha w^\varepsilon dx dy \\ &\quad - \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2} [D^\alpha, u^\varepsilon \partial_x] w^\varepsilon \cdot D^\alpha w^\varepsilon dx dy - \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma+2\alpha_2} [D^\alpha, v^\varepsilon \partial_y] w^\varepsilon \cdot D^\alpha w^\varepsilon dx dy, \end{aligned} \quad (\text{C.3})$$

where we have used the divergence-free condition.

First of all, we deal with the case $|\alpha| \leq s$ and $\alpha_1 \leq s - 1$ in (C.3). Similar to (2.19), we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2}^2 + \varepsilon^2 \|(1+y)^{\gamma+\alpha_2} \partial_x D^\alpha w^\varepsilon\|_{L^2}^2 + \frac{3}{4} \|(1+y)^{\gamma+\alpha_2} \partial_y D^\alpha w^\varepsilon\|_{L^2}^2 \\ & \leq - \int_{\mathbb{T}} D^\alpha w^\varepsilon \partial_y D^\alpha w^\varepsilon|_{y=0} dx + C_{s,\gamma} (1 + \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}) \|w^\varepsilon\|_{H^{s,\gamma}}^2, \end{aligned} \quad (\text{C.4})$$

Note the boundary term above can be estimated as follows(cf.[10]) for $|\alpha| \leq s - 1$,

$$\left| \int_{\mathbb{T}} D^\alpha w^\varepsilon \partial_y D^\alpha w^\varepsilon|_{y=0} dx \right| \leq \frac{1}{12} \|(1+y)^{\gamma+\alpha_2+1} \partial_y^2 D^\alpha w^\varepsilon\|_{L^2}^2 + C \|w^\varepsilon\|_{H^{s,\gamma}}^2 \quad (\text{C.5})$$

and for $|\alpha| = s$,

$$\left| \int_{\mathbb{T}} D^\alpha w^\varepsilon \partial_y D^\alpha w^\varepsilon|_{y=0} dx \right| \leq \begin{cases} \frac{1}{12} \|(1+y)^{\gamma+\alpha_2} \partial_y D^\alpha w^\varepsilon\|_{L^2}^2 + G(t), & \alpha_2 = 2k, k \in \mathbb{N}; \\ \frac{1}{12} \|(1+y)^{\gamma+\alpha_2+1} \partial_x^{\alpha_1-1} \partial_y^{\alpha_2+2} w^\varepsilon\|_{L^2}^2 + G(t), & \alpha_2 = 2k+1, k \in \mathbb{N}; \end{cases} \quad (\text{C.6})$$

where

$$G(t) = C_s \sum_{l=0}^{s/2} \|\partial_t^l \partial_x p^\varepsilon\|_{H^{s-2l}(\mathbb{T})}^2 + C_{s,\gamma} (1 + \|w^\varepsilon\|_{H^{s,\gamma}})^{s-2} \|w^\varepsilon\|_{H^{s,\gamma}}^2.$$

Thus, plugging the estimates (C.5) and (C.6) into (C.4), and summing over α , we get

$$\begin{aligned} & \frac{d}{dt} \sum_{\substack{|\alpha| \leq s \\ \alpha_1 \leq s-1}} \|(1+y)^{\gamma+\alpha_2} D^\alpha w^\varepsilon\|_{L^2}^2 + \sum_{\substack{|\alpha| \leq s \\ \alpha_1 \leq s-1}} \|(1+y)^{\gamma+\alpha_2} (\varepsilon \partial_x D^\alpha w^\varepsilon, \partial_y D^\alpha w^\varepsilon)\|_{L^2}^2 \\ & \leq C_{s,\gamma} \|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})}^2 + C_s \sum_{k=0}^{s/2} \|\partial_t^k \partial_x p^\varepsilon\|_{H^{s-2k}(\mathbb{T})}^2 + C_{s,\gamma} (1 + \|w^\varepsilon\|_{H^{s,\gamma}}^s). \end{aligned} \quad (\text{C.7})$$

Next, we deal with the case $|\alpha| = s$ and $\alpha_1 = s$ in (C.3), and hence, it follows

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |\partial_x^s w^\varepsilon|^2 dx dy + \varepsilon^2 \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |\partial_x^{s+1} w^\varepsilon|^2 dx dy \\ & = (\gamma + \alpha_2) \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma-1} v^\varepsilon |\partial_x^s w^\varepsilon|^2 dx dy + \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} \partial_y^2 \partial_x^s w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy \\ & \quad - \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} [\partial_x^s, u^\varepsilon \partial_x] w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy - \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} [\partial_x^s, v^\varepsilon \partial_y] w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy, \end{aligned} \quad (\text{C.8})$$

Using the inequality (2.10), it follows

$$\begin{aligned} & |(\gamma + \alpha_2) \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma-1} v^\varepsilon |\partial_x^s w^\varepsilon|^2 dx dy| \\ & \leq C_{s,\gamma} \left\| \frac{v^\varepsilon}{1+y} \right\|_{L^\infty} \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2}^2 \\ & \leq C_{s,\gamma} (\|\partial_x U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}) \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2}^2. \end{aligned} \quad (\text{C.9})$$

Integrating by part and applying Cauchy inequality, we get

$$\begin{aligned}
 & \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} \partial_y^2 \partial_x^s w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy \\
 &= \int_{\mathbb{T}} \partial_y \partial_x^s w^\varepsilon \cdot \partial_x^s w^\varepsilon|_{y=0} dx - \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |\partial_y \partial_x^s w^\varepsilon|^2 dx dy \\
 & \quad - \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma-1} \partial_y \partial_x^s w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy \\
 & \leq -\frac{3}{4} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |\partial_y \partial_x^s w^\varepsilon|^2 dx dy + C \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma-2} |\partial_x^s w^\varepsilon|^2 dx dy \\
 & \quad + \int_{\mathbb{T}} \partial_y \partial_x^s w^\varepsilon \cdot \partial_x^s w^\varepsilon|_{y=0} dx.
 \end{aligned} \tag{C.10}$$

Using the boundary condition $\partial_y w^\varepsilon|_{y=0} = p^\varepsilon$ and Sobolev inequality, it follows

$$\begin{aligned}
 \int_{\mathbb{T}} \partial_y \partial_x^s w^\varepsilon \cdot \partial_x^s w^\varepsilon|_{y=0} dx & \leq \sqrt{2} \|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})} \|\partial_y \partial_x^s w^\varepsilon\|_{L^2}^{\frac{1}{2}} \|\partial_x^s w^\varepsilon\|_{L^2}^{\frac{1}{2}} \\
 & \leq \frac{1}{4} \|\partial_y \partial_x^s w^\varepsilon\|_{L^2}^2 + C(\|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})}^2 + \|\partial_x^s w^\varepsilon\|_{L^2}^2).
 \end{aligned}$$

This and the inequality (C.10) yield directly

$$\begin{aligned}
 & \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} \partial_y^2 \partial_x^s w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy \\
 & \leq -\frac{1}{2} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |\partial_y \partial_x^s w^\varepsilon|^2 dx dy + C(\|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})}^2 + \|(1+y)^{\gamma-1} \partial_x^s w^\varepsilon\|_{L^2}^2).
 \end{aligned} \tag{C.11}$$

By virtue of Hölder inequality, we get

$$\begin{aligned}
 & \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} [\partial_x^s, u^\varepsilon \partial_x] w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy \right| \\
 & \leq \sum_{1 \leq k \leq s} C_{s,k} \|(1+y)^\gamma \partial_x^k (u^\varepsilon - U) \partial_x^{s+1-k} w^\varepsilon\|_{L^2} \|(1+y)^\gamma \partial_x w^\varepsilon\|_{L^2} \\
 & \quad + \sum_{1 \leq k \leq s} C_{s,k} \|(1+y)^\gamma \partial_x^k U \partial_x^{s+1-k} w^\varepsilon\|_{L^2} \|(1+y)^\gamma \partial_x w^\varepsilon\|_{L^2}.
 \end{aligned}$$

Using the Hardy and Morse type inequalities, it follows for $1 \leq k \leq s$

$$\|(1+y)^\gamma \partial_x^k (u^\varepsilon - U) \partial_x^{s+1-k} w^\varepsilon\|_{L^2} \leq \|\partial_x (u^\varepsilon - U)\|_{H^{s-1,0}} \|\partial_x w^\varepsilon\|_{H^{s-1,\gamma}} \leq C \|w^\varepsilon\|_{H^{s,\gamma}}^2,$$

and applying Wirtinger inequality, it follows

$$\|(1+y)^\gamma \partial_x^k U \partial_x^{s+1-k} w^\varepsilon\|_{L^2} \leq C_s \|\partial_x^s U\|_{L^\infty(\mathbb{T})} \|w^\varepsilon\|_{H^{s,\gamma}}.$$

Thus, we can conclude the estimate

$$\left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} [\partial_x^s, u^\varepsilon \partial_x] w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy \right| \leq C_s (\|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}) \|w^\varepsilon\|_{H^{s,\gamma}}^2. \tag{C.12}$$

Applying the Hölder inequality, it follows

$$\begin{aligned}
 & \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} [\partial_x^s, v^\varepsilon \partial_y] w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy \\
 & \leq \sum_{1 \leq k \leq s} C_{s,k} \|(1+y)^\gamma \partial_x^k (v^\varepsilon + y \partial_x U) \partial_x^{s-k} \partial_y w^\varepsilon\|_{L^2} \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2} \\
 & \quad + \sum_{1 \leq k \leq s} C_{s,k} \|\partial_x^{k+1} U\|_{L^\infty(\mathbb{T})} \|(1+y)^{\gamma+1} \partial_x^{s-k} \partial_y w^\varepsilon\|_{L^2} \|(1+y)^\gamma \partial_x^s w^\varepsilon\|_{L^2}.
 \end{aligned} \tag{C.13}$$

For $1 \leq k \leq s-1$, using the Hardy and Morse type inequalities, it follows

$$\begin{aligned} & \|(1+y)^\gamma \partial_x^k (v^\varepsilon + y \partial_x U) \partial_x^{s-k} \partial_y w^\varepsilon\|_{L^2} \\ & \leq C \|\partial_x (v^\varepsilon + y \partial_x U)\|_{H^{s-2,-1}} \|(1+y) \partial_{xy} w^\varepsilon\|_{H^{s-2,\gamma}} \\ & \leq C \|w^\varepsilon\|_{H^{s,\gamma}}^2, \end{aligned} \tag{C.14}$$

and

$$\begin{aligned} & \|(1+y)^\gamma \partial_x^s (v^\varepsilon + y \partial_x U) \partial_y w^\varepsilon\|_{L^2} \\ & \leq \|(1+y)^{\gamma-1} \partial_x^s (v^\varepsilon + y \partial_x U)\|_{L^2} \|(1+y) \partial_y w^\varepsilon\|_{L^\infty} \\ & \leq C_\gamma \|(1+y)^\gamma \partial_x^{s+1} w^\varepsilon\|_{L^2} \|w^\varepsilon\|_{H^{3,0}}. \end{aligned} \tag{C.15}$$

Substituting the estimates (C.14) and (C.15) into (C.13), and using the Cauchy inequality, we get

$$\begin{aligned} & \left| \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} [\partial_x^s, v^\varepsilon \partial_y] w^\varepsilon \cdot \partial_x^s w^\varepsilon dx dy \right| \\ & \leq \frac{1}{4} \varepsilon^2 \|(1+y)^\gamma \partial_x^{s+1} w^\varepsilon\|_{L^2}^2 + C_s \|\partial_x^{s+1} U\|_{L^\infty}^2 + C_{s,\gamma,\varepsilon} (1 + \|w^\varepsilon\|_{H^{s,\gamma}}^4). \end{aligned} \tag{C.16}$$

Substituting the estimates (C.9), (C.11), (C.12), (C.16) into (C.8), we obtain

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |\partial_x^s w^\varepsilon|^2 dx dy + \varepsilon^2 \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |\partial_x^{s+1} w^\varepsilon|^2 dx dy + \int_{\mathbb{T} \times \mathbb{R}^+} (1+y)^{2\gamma} |\partial_y \partial_x^s w^\varepsilon|^2 dx dy \\ & \leq C_s (\|\partial_x^{s+1} U\|_{L^\infty(\mathbb{T})}^2 + \|\partial_x^{s+1} p^\varepsilon\|_{L^2(\mathbb{T})}^2) + C_{s,\gamma,\varepsilon} (1 + \|w^\varepsilon\|_{H^{s,\gamma}}^4). \end{aligned}$$

This along with the inequalities (C.7) and (2.38) yield directly

$$\frac{d}{dt} \|w^\varepsilon\|_{H^{s,\gamma}}^2 + \varepsilon^2 \|\partial_x w^\varepsilon\|_{H^{s,\gamma}}^2 + \|\partial_y w^\varepsilon\|_{H^{s,\gamma}}^2 \leq C_{s,\gamma} (1 + M_U)^2 + C_{s,\gamma,\varepsilon} \|w^\varepsilon\|_{H^{s,\gamma}}^s,$$

and hence, we conclude that

$$1 + \|w^\varepsilon(t)\|_{H^{s,\gamma}}^2 \leq \frac{1 + \|w_0\|_{H^{s,\gamma}}^2}{\left\{1 - \frac{2}{s-2} \max\{C_{s,\gamma}(1 + M_U)^2, C_{s,\gamma,\varepsilon}\} \{1 + \|w_0\|_{H^{s,\gamma}}^2\}^{\frac{s-2}{2}} t\right\}^{\frac{2}{s-2}}}, \tag{C.17}$$

as long as $t < \frac{s-2}{2 \max\{C_{s,\gamma}(1 + M_U)^2, C_{s,\gamma,\varepsilon}\} \{1 + \|w_0\|_{H^{s,\gamma}}^2\}^{\frac{s-2}{2}}}$.

Step 2: L^∞ -estimates. Denote $B_\alpha := (1+y)^{\sigma+\alpha_2} D^\alpha w^\varepsilon$ and $I := \sum_{1 \leq |\alpha| \leq 2} |B_\alpha|^2$, it is easy to check that I satisfies the evolution equation

$$\begin{aligned} & (\partial_t + u^\varepsilon \partial_x + v^\varepsilon \partial_y - \varepsilon^2 \partial_x^2 - \partial_y^2) I \\ & = -2 \sum_{1 \leq |\alpha| \leq 2} \{\varepsilon^2 |\partial_x B_\alpha|^2 + |\partial_y B_\alpha|^2\} + 2 \sum_{1 \leq |\alpha| \leq 2} \{Q_\alpha B_\alpha \partial_y B_\alpha + R_\alpha |B_\alpha|^2 + S_\alpha B_\alpha\}, \end{aligned}$$

where the quantities Q_α, R_α and S_α are given explicitly by

$$Q_\alpha := -\frac{2(\sigma + \alpha_2)}{1+y}, \quad R_\alpha := \frac{\sigma + \alpha_2}{1+y} v^\varepsilon + \frac{(\sigma + \alpha_2)(\sigma + \alpha_2 + 1)}{(1+y)^2}$$

and

$$S_\alpha := -\sum_{0 < \beta \leq \alpha} C_{\alpha,\beta} (1+y)^{\beta_2} \left\{ D^\beta u^\varepsilon B_{\alpha-\beta+e_1} + \frac{D^\beta v^\varepsilon B_{\alpha-\beta+e_2}}{1+y} \right\}.$$

By routine checking, we get that

$$|Q_\alpha| \leq C_\sigma, \quad |R_\alpha| \leq C_\sigma (\|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}),$$

and

$$|S_\alpha| \leq C_\sigma (\|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}) \sum_{0 < \beta \leq \alpha} \{|B_{\alpha-\beta+e_1}| + |B_{\alpha-\beta+e_2}|\}.$$

Then, we can verify that the quantity I satisfies

$$(\partial_t + u^\varepsilon \partial_x + v^\varepsilon \partial_y - \varepsilon^2 \partial_x^2 - \partial_y^2)I \leq C_\sigma \{1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}\}I,$$

and hence, we apply the maximum principle in Lemma A.4 to get

$$\sup_{0 \leq \tau \leq t} Q(\tau) \leq e^{C_\sigma \{1 + \|\partial_x^s U\|_{L^\infty(\mathbb{T})} + \|w^\varepsilon\|_{H^{s,\gamma}}\}t} (\|I(0)\|_{L^\infty(\Omega)} + \|I(t)|_{y=0}\|_{L^\infty(\mathbb{T})}). \quad (\text{C.18})$$

By virtue of $s \geq 4$, we apply the Sobolev inequality (A.3) to get

$$\|I(t)|_{y=0}\|_{L^\infty(\mathbb{T})} \leq C \|w^\varepsilon\|_{H^{s,\gamma}}^2.$$

This and the inequality (C.18) yield directly

$$\sup_{0 \leq \tau \leq t} Q(\tau) \leq e^{C_\sigma \{1 + \sup_{0 \leq \tau \leq t} \|\partial_x^s U(\tau)\|_{L^\infty(\mathbb{T})} + \sup_{0 \leq \tau \leq t} \|w^\varepsilon(\tau)\|_{H^{s,\gamma}}\}t} (Q(0) + \|w^\varepsilon\|_{H^{s,\gamma}}^2). \quad (\text{C.19})$$

Step 3: Life span time. Taking $T_1 = \frac{\{1 - (\frac{1}{2})^{\frac{s-2}{2}}\}(s-2)}{2 \max\{C_{s,\gamma}(1+M_U)^2, C_{s,\gamma,\varepsilon}\} \{1 + \|w_0\|_{H^{s,\gamma}}^2\}^{\frac{s-2}{2}}}$, we get by using (C.17)

$$\sup_{0 \leq \tau \leq T_1} \|w^\varepsilon(\tau)\|_{H^{s,\gamma}}^2 \leq 2(1 + \|w_0\|_{H^{s,\gamma}}^2). \quad (\text{C.20})$$

Taking $T_2 = \min\{T_1, \frac{\ln 2}{C_\sigma(3+M_U^{\frac{1}{2}} + \sqrt{2}\|w_0\|_{H^{s,\gamma}})}\}$, it follows from (C.19)

$$\sup_{0 \leq t \leq T_2} Q(t) \leq 4(1 + Q(0) + \|w_0\|_{H^{s,\gamma}}^2). \quad (\text{C.21})$$

Taking $T_3 = \min\{T_1, T_2, \frac{\delta_0}{C(7+M_U+4Q(0)+6\|w_0\|_{H^{s,\gamma}}^2)}\}$, we get by using (2.41)

$$(1+y)^\sigma w^\varepsilon(t, x, y) \geq (1+y)^\sigma w_0(x, y) - \delta_0 \geq 2\delta_0 - \delta_0 = \delta_0. \quad (\text{C.22})$$

Then, we have chosen the life span time $T_a := T_3$ such the estimates (C.20)-(C.22) hold on. Finally, we point out that we can use the local existence results established above to extend the solution (defined on $[0, T_a]$) step by step to the time interval $[0, T^\varepsilon]$ such the estimates (C.1) and (C.2) hold on. Therefore, we complete the proof of this lemma. \square

Acknowledgements

Jincheng Gao's research was partially supported by Fundamental Research Funds for the Central Universities (Grants No.18lgpy66) and NNSF of China (Grants No.11801586). Daiwen Huang's research was partially supported by the NNSF of China (Grants No.11631008) and National Basic Research Program of China 973 Program (Grants No.2007CB814800). Zheng-an Yao's research was partially supported by NNSF of China (Grant No.11431015).

References

- [1] L. Prandtl, *Über flüssigkeits-bewegung bei sehr kleiner reibung*. Verhandlungen des III. Internationalen Mathematiker Kongresses, Heidelberg. Teubner, Leipzig, (1904), 484-491.
- [2] O.A. Oleinik, *On the system of Prandtl equations in boundary-layer theory*, Dokl. Akad. Nauk SSSR 150(1963) 28-31.
- [3] O.A. Oleinik, *On the mathematical theory of boundary layer for an unsteady flow of incompressible fluid*, Prikl. Mat. Meh. 30, 801-821(Russian); translated as J. Appl. Math. Mech. 30(1966)951-974.
- [4] O.A. Oleinik, V.N. Samokhin, *Mathematical Models in Boundary Layers Theory*, Applied Mathematics and Mathematical Computation, 15. Chapman & Hall/CRC, Boca Raton, FL, 1999.
- [5] Z.P.Xin, L.Q.Zhang, *On the global existence of solutions to the Prandtl's system*, Adv. Math. 181 (2004) 88-133.
- [6] R.E. Caflisch, M.Sammartino, *Existence and singularities for the Prandtl boundary layer equations*, Z.Angew.Math.Mech. 80 (2000), no.11-12, 733-744.
- [7] D.Gérard-Varet, E.Dormy, *On the ill-posedness of the Prandtl equation*, J. Amer. Math. Soc. 23 (2010), no. 2, 591-609.
- [8] D.Gérard-Varet, T.Nguyen, *Remarks on the ill-posedness of the Prandtl equation*, Asymptot. Anal. 77 (2012), no. 1-2, 71-88.
- [9] R. Alexander, Y.G. Wang, C.J. Xu, T. Yang, *Well-posedness of the Prandtl equation in Sobolev spaces*, J. Amer. Math. Soc. 28(2015) 745-784.
- [10] N.Masmoudi, T.K.Wong, *Local-in-time existence and uniqueness of solutions to the Prandtl equations by energy methods*, Comm. Pure Appl. Math., 68(2015), 10, 1683-1741.
- [11] M. Sammartino, R.E. Caflisch, *Zero viscosity limit for analytic solutions, of the Navier-Stokes equation on a half-space. I. Existence for Euler and Prandtl equations*, Comm. Math. Phys. 192(1998) 433-461.
- [12] M. Sammartino, R.E. Caflisch, *Zero viscosity limit for analytic solutions of the Navier-Stokes equation on a half-space. II. Construction of the Navier-Stokes solution*, Comm. Math. Phys. 192(1998) 463-491.
- [13] Y.Guo, T.Nguyen, *Prandtl boundary layer expansions of steady Navier-Stokes flows over a moving plate*, Ann. PDE 3 (2017), no. 1, Art. 10, 58 pp.
- [14] Y.Guo, S.Iyer, *Validity of Steady Prandtl Layer Expansions*, arXiv:1805.05891v5.
- [15] Y.Guo, S.Iyer, *Steady Prandtl Layer Expansions with External Forcing*, arXiv:1810.06662v1.
- [16] E.Grenier, Y.Guo, T.Nguyen, *Spectral instability of characteristic boundary layer flows*, Duke Math. J. 165(2016) 3085-3146.
- [17] E.Grenier, Y.Guo, T.Nguyen, *Spectral stability of Prandtl boundary layers: an overview*, Analysis (Berlin) 35 (2015) 343-355.

- [18] G.M.Gie, C.Y.Jung, R.Temam, Recent progresses in boundary layer theory, *Discrete Contin. Dyn. Syst.* 36 (2016) 2521-2583.
- [19] Y.L. Xiao, Z.P. Xin, On the vanishing viscosity limit for the 3D Navier-Stokes equations with a slip boundary condition, *Commun. Pure Appl. Math.* 60 (2007) 1027-1055.
- [20] N. Masmoudi, F. Rousset, Uniform Regularity for the Navier-Stokes equation with Navier boundary condition, *Arch. Ration. Mech. Anal.* 203 (2012) 529-575.
- [21] N. Masmoudi, F. Rousset, Uniform regularity and vanishing viscosity limit for the free surface Navier-Stokes equations, *Arch. Ration. Mech. Anal.* 223 (2017) 301-417.