

# ExtraOrdinary Gauge Mediation Extension of deflected AMSB

---

Fei Wang<sup>1</sup>,

<sup>1</sup> *School of Physics, Zhengzhou University, 450000, Zhengzhou P.R.China*

ABSTRACT: Extraordinary gauge mediation extension of deflected AMSB scenarios can be interesting because it can accommodate together the deflection in the Kahler potential and the superpotential. We revisit the EGM scenario and derive the analytical expressions for soft SUSY breaking parameters in EGM and EGM extension of deflected AMSB scenarios with wavefunction renormalization approach, especially the case with vanishing gauge beta-function at an intermediate energy scale. The Landau pole and proton decay constraints are also discussed.

---

## Contents

<b>1. Introduction</b>	<b>2</b>
<b>2. Extraordinary gauge mediation in deflected AMSB</b>	<b>3</b>
<b>3. Analytical expressions within EGM</b>	<b>5</b>
3.1 Class A: Partition without vanishing gauge beta functions	7
3.2 Class B: Partition with vanishing gauge beta functions	10
3.3 Dependence of $\ln M_a$ on $\ln X$	14
<b>4. Messengers on GUT and Landau Pole</b>	<b>19</b>
<b>5. Conclusions</b>	<b>21</b>
<b>A. Coefficients In the Wavefunction of MSSM Superfields</b>	<b>21</b>

---

## 1. Introduction

Weak scale supersymmetry(SUSY), which is a leading candidate for physics beyond the standard model(SM), can solve elegantly the hierarchy problem of the Higgs boson by introducing various superpartners at TeV scale. Besides, gauge couplings unification, which can not be exact in SM, can be successful in its SUSY extensions. The dark matter(DM) puzzle as well as the BAU(baryon-asymmetric universe) puzzle etc, can also be explained with proper DM candidates and baryogenesis mechanisms in SUSY. It is worth to note that the Higgs scalar, which was discovered by the ATLAS[1] and CMS[2] collaborations of LHC in 2012, lie miraculously in the small '115 – 135' GeV window predicted by the low energy SUSY.

It is well known that the soft SUSY spectrum, including the gaugino and sfermion masses, are determined by the SUSY breaking mechanism. Depending on the way the visible sector '*feels*' the SUSY breaking effects in the hidden sector, the SUSY breaking mechanisms can be classified into gravity mediation[3], gauge mediation[4], anomaly mediation[5] scenarios, etc. Gauge mediated SUSY breaking(GMSB) scenarios, which will not cause flavor and CP problems that bothers gravity mediation models, are calculable, predictive, and phenomenologically distinctive with minimal messenger sector. However, it is difficult for minimal GMSB to explain 125 GeV Higgs with TeV scale soft SUSY breaking parameters because of the vanishing trilinear terms at the messenger scale. An interesting extension is the (extra)ordinary gauge mediation (EGM) scenarios[6, 7], in which the messenger sector can include all renormalizable, gauge invariant couplings between the messengers and any number of singlet fields. In fact, many examples in the literature of OGM deformed by mass terms can fall into this category and their generic properties can be obtained therein.

Gravity, which can couple to everything, can generate the soft masses by the auxiliary field of the gravitational multiplet. Such a '*pure*' gravity mediation scenario with negligible contributions from direct non-renormalizable contact terms is called the anomaly mediated SUSY breaking(AMSB). Pure anomaly mediation is bothered by the tachyonic slepton problem [8]. Its non-trivial extensions with messenger sectors, namely the deflected AMSB[9, 10, 11], can elegantly solve such tachyonic slepton problems through the deflection of the renormalization group equation (RGE) trajectory [12]. There are two types of deflections in the literature, namely, the deflection in the superpotential or deflection in the Kahler potential[13, 14, 15, 16]. However, it is difficult to determined consistently the deflection parameter '*d*' and soft SUSY parameters if both deflections are present. We find that such a scenario can be seen as a special case of the EGM extension of deflected AMSB. Besides, the most generic extension of the messenger sector can also be interesting theoretically.

We revisit the EGM scenario and derive the analytical results for the GMSB contributions in EGM and EGM extension of deflected AMSB, especially the case with a vanishing beta function at an intermediate scale and the case with  $rank(\lambda_{ij}) = rank(m_{ij}) = 0$ . Our result is especially useful for cases with hierarchical messenger scales. The extension of the EGM to deflected AMSB is straightforward. We show that such an extension can indeed

accommodate both types of deflections in AMSB.

This paper is organized as follows. In Sec 2, we discuss the deflected AMSB scenario with EGM extension. In Sec 3, we discuss the analytical expressions of soft SUSY parameters in EGM. In Sec 4, constraints from the Landau pole with multiple messengers are given. Sec 5 contains our conclusions.

## 2. Extraordinary gauge mediation in deflected AMSB

To fully understand the deflected AMSB scenarios with the presence of both the superpotential and Kahler potential deflection, we need the deflection parameter ' $d$ ' to derive the soft SUSY breaking parameters. One can minimize the whole scalar potential to obtain ' $d$ ', but such a case by case study seems tedious. We find that the solution can be obtained in EGM extension of deflected AMSB.

In deflected AMSB, the Kahler potential can have the following types of deflection with holomorphic terms for messengers

$$K \supseteq \kappa_{ij} \frac{\phi^\dagger}{\phi} \tilde{P}_i P_j , \quad (2.1)$$

or the deflection from the couplings in the superpotential

$$W \supseteq \lambda_{ij} \tilde{X} \tilde{Q}_i Q_j + W(\tilde{X}) , \quad (2.2)$$

with a proper form of superpotential  $W(\tilde{X})$  for the pseudo-moduli field  $\tilde{X}$  to determine the deflection parameter  $\tilde{d}$

$$\frac{F_{\tilde{X}}}{\tilde{X}} = (\tilde{d} + 1) F_\phi . \quad (2.3)$$

We should note that the messengers  $\tilde{P}_i, P_i$  can possibly be identified to be the  $\tilde{Q}_i, Q_i$  superfields. Combining both Kahler and superpotential deflection, we have

$$\mathcal{L} \supseteq \int d^4\theta \left[ \kappa_{ij} \frac{\phi^\dagger}{\phi} \tilde{P}_i P_j \right] + \int d^2\theta \left[ \lambda_{ij} \tilde{X} \tilde{Q}_i Q_j + W(\tilde{X}) \right] , \quad (2.4)$$

which can be rewritten as

$$\mathcal{L} \supseteq \int d^2\theta (\lambda_{ij} \tilde{X} + \kappa_{ij} \tilde{T}) \tilde{\phi}_i \phi_j + W(\tilde{X}) , \quad (2.5)$$

with  $\tilde{T}$  the auxiliary field with VEV

$$\langle \tilde{T} \rangle = F_\phi - \theta^2 F_\phi^2 , \quad (2.6)$$

and  $\phi$  the conformal compensator field which carries the SUSY breaking information in the SUSY breaking sector

$$\phi = 1 + \theta^2 F_\phi . \quad (2.7)$$

We can rotate  $\tilde{X}$  and  $\tilde{T}$  so that only one combination  $X$  will acquire F-term VEVs while  $T$  will acquire only the lowest component VEVs

$$\begin{aligned} X &= \frac{1}{\sqrt{F_{\tilde{X}}^2 + F_{\tilde{T}}^2}} \left[ F_{\tilde{X}} \tilde{X} - F_{\tilde{T}}^2 \tilde{T} \right] , \\ T &= \frac{1}{\sqrt{F_{\tilde{X}}^2 + F_{\tilde{T}}^2}} \left[ F_{\tilde{T}}^2 \tilde{X} + F_{\tilde{X}} \tilde{T} \right] , \end{aligned} \quad (2.8)$$

So the superpotential can be rewritten as

$$W \supseteq (\lambda_{ij} X + m_{ij}) \tilde{\phi}_i \phi_j , \quad (2.9)$$

with  $m_{ij} = \kappa_{ij} \langle T \rangle$ .

We will impose a non-trivial R-symmetry

$$\begin{cases} \lambda_{ij} \neq 0 , & \text{only if } R(\tilde{\phi}_i) + R(\phi_j) = 2 - R(X) , \\ \kappa_{ij} \neq 0 , & \text{only if } R(\tilde{\phi}_i) + R(\phi_j) = 2 , \end{cases} \quad (2.10)$$

as well as a symmetry for messengers to prevent destructive D-term contributions to sfermion masses. After integrating out the messenger fields, the messenger determinant is proven by [6] to be a monomial in  $X$

$$\det(\lambda_{ij} X + \kappa_{ij} T) = X^{n_0} G(\lambda, \kappa T) , \quad (2.11)$$

with

$$n_0 = \frac{1}{R(X)} \sum_{i=1}^n \left[ 2 - R(\tilde{\phi}_i) + R(\phi_i) \right] . \quad (2.12)$$

Note that the most general deflection in AMSB is quite similar to that of EGM in GMSB except that the  $X$  is given by

$$\sqrt{F_{\tilde{X}}^2 + F_{\tilde{T}}^2} \langle X \rangle = F_{\tilde{T}} \left[ (\tilde{d} + 1) \tilde{X}^2 - F_{\tilde{T}}^2 \right] + (F_{\tilde{X}}^2 + F_{\tilde{T}}^4) \theta^2 , \quad (2.13)$$

from eqn.(2.8). Thus the effective deflection parameter  $d$  is given by

$$\begin{aligned} d &= \frac{(F_{\tilde{X}}^2 + F_{\tilde{T}}^4)}{F_{\tilde{T}}^2 \left[ (\tilde{d} + 1) \tilde{X}^2 - F_{\tilde{T}}^2 \right]} - 1, \\ &= \frac{\left[ (\tilde{d} + 1)^2 \tilde{X}^2 + F_{\tilde{T}}^2 \right]}{\left[ (\tilde{d} + 1) \tilde{X}^2 - F_{\tilde{T}}^2 \right]} - 1, \end{aligned} \quad (2.14)$$

For later convenience, the VEV of  $X$  is denoted by  $\langle X \rangle = M + \theta^2 F_X$ .

The soft SUSY breaking parameters in EGM extension of deflected AMSB can be obtained by wavefunction renormalization[17] approach

- The gaugino masses are given as

$$\begin{aligned}
M_i &= g_i^2 \left( \frac{F_\phi}{2} \frac{\partial}{\partial \ln \mu} - \frac{dF_\phi}{2} \frac{\partial}{\partial \ln |X|} \right) \frac{1}{g_i^2}(\mu, |X|) , \\
&= g_i^2 \left( \frac{F_\phi}{2} \frac{\partial}{\partial \ln \mu} - \sum_{a=1}^N \frac{dF_\phi}{2} \frac{\partial \ln M_a}{\partial \ln |X|} \frac{\partial}{\partial \ln M_a} \right) \frac{1}{g_i^2}(\mu, M_a) , \quad (2.15)
\end{aligned}$$

- The trilinear couplings are given as

$$A_0^{ijk} \equiv \frac{A_{ijk}}{y_{ijk}} = \sum_i \left( -\frac{F_\phi}{2} \frac{\partial}{\partial \ln \mu} + \frac{dF_\phi}{2} \frac{\partial}{\partial \ln |X|} \right) \ln [Z_i(\mu, M_a)] . \quad (2.16)$$

So we need to calculate

$$\frac{\partial \ln Z_l(\mu, M_a)}{\partial \ln |X|} = \sum_{M_j} \sum_{g_i(\mu')} \frac{\partial \ln M_j}{\partial \ln |X|} \frac{\partial \ln g_i(\mu')}{\partial \ln M_j} \frac{\partial \ln Z_l}{\partial \ln g_i(\mu')} + [g_i(\mu') \rightarrow y_a(\mu')] . \quad (2.17)$$

in which the sum over  $g_i(\mu')$ , which depends on the messenger thresholds  $M_i$ , take the values  $g_i(M_1), g_i(M_2), \dots, g_i(M_N), g_i(\mu)$ . The second term always vanishes because the anomalous dimension is continuous across the messenger thresholds.

- The soft SUSY breaking scalar masses are given as

$$\begin{aligned}
m_{soft}^2 &= - \left| -\frac{F_\phi}{2} \frac{\partial}{\partial \ln \mu} + \frac{dF_\phi}{2} \frac{\partial}{\partial \ln |X|} \right|^2 \ln [Z_i(\mu, M_a)] , \quad (2.18) \\
&= - \left( \frac{F_\phi^2}{4} \frac{\partial^2}{\partial (\ln \mu)^2} + \frac{d^2 F_\phi^2}{4} \frac{\partial^2}{\partial (\ln |X|)^2} - \frac{dF_\phi^2}{2} \frac{\partial^2}{\partial \ln |X| \partial \ln \mu} \right) \ln [Z_i(\mu, M_a)] ,
\end{aligned}$$

We need to calculate

$$\begin{aligned}
\frac{\partial^2}{\partial \ln |X|^2} \ln Z_i(\mu, M_a) &= \left[ \frac{\partial \ln M_j}{\partial \ln |X|} \frac{\partial}{\partial \ln M_j} \right] \left[ \frac{\partial \ln M_i}{\partial \ln |X|} \frac{\partial}{\partial \ln M_i} \right] \ln Z_a(\mu, M_a) , \\
&= \left[ \frac{\partial \ln M_j}{\partial \ln |X|} \frac{\partial \ln M_i}{\partial \ln |X|} \right] \left[ \frac{\partial^2 \ln Z_a(\mu, M_a)}{\partial \ln M_i \partial \ln M_j} \right] . \quad (2.19)
\end{aligned}$$

The dependencies of  $\ln M_a$  on  $\ln |X|$  are in general non-trivial, which depends crucially on the properties of matrices  $\lambda_{ij}$  and  $m_{ij}$ . We will discuss their expressions in the subsequent sections.

It is well known that in the  $d \rightarrow \infty$  limit, the anomaly mediation contributions in the deflected AMSB are sub-leading and the gauge mediation contributions are dominant. So we will derive the EGM contributions first and return to deflected AMSB cases subsequently.

### 3. Analytical expressions within EGM

We assume that the mass thresholds of the  $N$  messengers can be degenerated and separated into ' $p$ ' groups as

$$\underbrace{(M_1, \dots, M_1)}_{n_1}, \underbrace{(M_2, \dots, M_2)}_{n_2}, \dots, \underbrace{(M_p, \dots, M_p)}_{n_p} . \quad (3.1)$$

with  $\sum_{i=1}^p n_p = N$ .

The gauge coupling at a scale  $\mu$  that below all the messenger thresholds are given as

$$\begin{aligned} \frac{1}{g_i^2(\mu, X)} &= \frac{1}{g_i^2(\Lambda)} + \frac{b'_i}{8\pi^2} \ln \frac{\Lambda}{M_1} + \frac{b'_i - n_1}{8\pi^2} \ln \frac{M_1}{M_2} + \frac{b'_i - \sum_{k=1}^2 n_k}{8\pi^2} \ln \frac{M_2}{M_3} + \cdots + \frac{b'_i - N}{8\pi^2} \ln \frac{M_p}{\mu}, \\ &= \frac{1}{g_i^2(\Lambda)} + \frac{b'_i}{8\pi^2} \ln \Lambda - \frac{1}{8\pi^2} \ln \det \mathcal{M} - \frac{b'_i - N}{8\pi^2} \ln \mu. \end{aligned} \quad (3.2)$$

Here we assume that the eigenvalues of the messenger mass matrix are given by  $M_1 \geq M_2 \geq \cdots \geq M_p$ . So we can obtain that gaugino mass

$$\begin{aligned} M_i &= g_i^2 \left( \frac{F_\phi}{2} \frac{\partial}{\partial \ln \mu} - \sum_{a=1}^p \frac{dF_\phi}{2} \frac{\partial \ln M_a}{\partial \ln |X|} \frac{\partial}{\partial \ln M_a} \right) \frac{1}{g_i^2}(\mu, M_a), \\ &= -F_\phi \frac{g_i^2}{16\pi^2} b_i + dF_\phi \frac{g_i^2}{16\pi^2} \sum_{a=1}^p n_a \frac{\partial \ln M_a}{\partial \ln |X|}, \\ &= -F_\phi \frac{g_i^2}{16\pi^2} b_i + dF_\phi \frac{g_i^2}{16\pi^2} \frac{\partial \ln \det \mathcal{M}}{\partial \ln |X|}, \\ &\equiv -F_\phi \frac{g_i^2}{16\pi^2} [b_i - d n_0]. \end{aligned} \quad (3.3)$$

Here  $b_i = b'_i - N$ , which are given by

$$(b_1, b_2, b_3) = (33/5, 1, -3), \quad (3.4)$$

with  $N = N_5 + 3N_{10}$  and  $n_0$  is given in eqn.(2.11).

For the soft sfermion masses and trilinear couplings, we need the dependence of wavefunction  $Z_i$  on the messenger thresholds  $M_a(\sqrt{X^\dagger X})$ . The derivative of  $\ln Z_i$  with respect to  $\ln |X|$  can be obtained via

$$\begin{aligned} &\frac{\partial \ln Z_l[\mu, g_i(\mu'), y_i(\mu'), M_a]}{\partial \ln |X|} \\ &= \sum_{M_j} \sum_{g_i(\mu')} \frac{\partial \ln M_j}{\partial \ln |X|} \frac{\partial \ln g_i(\mu')}{\partial \ln M_j} \frac{\partial \ln Z_l[\mu, g_i(\mu'), y_i(\mu'), M_a]}{\partial \ln g_i(\mu')} + [g_i(\mu') \rightarrow y_a(\mu')] \\ &+ \sum_{M_a} \frac{\partial \ln M_a}{\partial \ln |X|} \frac{\partial \ln Z_l[\mu, g_i(\mu'), y_i(\mu'), M_a]}{\partial \ln M_a}. \end{aligned} \quad (3.5)$$

The sum over  $g_i(\mu')$ , which depends on the messenger thresholds  $M_i$ , take the values  $g_i(M_1), g_i(M_2), \cdots, g_i(M_p)$ . The second term always vanishes because the anomalous dimension is continuous across the messenger thresholds.

To get the expressions for wavefunction  $Z_i$ , we need to the following classification

- In the case  $b'_i - \sum_{r=1}^k n_r \neq 0$  for all  $0 \leq k \leq p$ , the expression will fall into class A.
- In the case  $b'_i - \sum_{r=1}^k n_r = 0$  for some  $0 \leq k \leq p$ , the expression will fall into class B.

### 3.1 Class A: Partition without vanishing gauge beta functions

To obtain  $Z_i$ , we can construct an invariant by surveying the anomalous dimension of  $Z_i$

$$\frac{d \ln Z_i}{dt} = -\frac{1}{8\pi^2} \left[ d_{ij}^k y_{ijk}^2 - 2C(r)g_i^2 \right], \quad (3.6)$$

and solve the differential equation in the basis of  $(y_t^2, y_b^2, y_\tau^2, g_3^2, g_2^2, g_1^2)$  with

$$\frac{d}{dt} \ln Z_i = \sum_{l=g_3, g_2, g_1} 2\tilde{A}_l \frac{d \ln g_l}{dt} + \sum_{l=y_t, y_b, y_\tau} 2B_l \frac{d \ln y_l}{dt}, \quad (3.7)$$

at the scale  $\Lambda$ . The expressions of the wavefunction can be solved<sup>1</sup> as

$$Z_l(\mu, M_a) = Z_l(\Lambda) \left( \frac{y_t^2(\mu)}{y_t^2(\Lambda)} \right)^{B_t} \left( \frac{y_b^2(\mu)}{y_b^2(\Lambda)} \right)^{B_b} \left( \frac{y_\tau^2(\mu)}{y_\tau^2(\Lambda)} \right)^{B_\tau} \quad (3.8)$$

$$\prod_{i=1}^3 \left[ \left( \frac{g_i^2(M_1)}{g_i^2(\Lambda)} \right)^{\frac{A_i}{b'_i}} \left( \frac{g_i^2(M_2)}{g_i^2(M_1)} \right)^{\frac{A_i}{b'_i - n_1}} \left( \frac{g_i^2(M_3)}{g_i^2(M_2)} \right)^{\frac{A_i}{b'_i - \sum_{k=1}^2 b_i}} \dots \left( \frac{g_i^2(\mu)}{g_i^2(M_p)} \right)^{\frac{A_i}{b'_i - N}} \right].$$

with the corresponding coefficients  $\tilde{A}_i$  in given as  $\tilde{A}_i \equiv A_i/b'_i, A_i/(b'_i - n_1), \dots$  at the energy interval  $[M_i, M_{i+1}]$  ( $M_0 \equiv \Lambda$ ), respectively.

If all the Yukawa terms within the wavefunction  $Z_i$  are neglected, the  $A_i$  will take the value  $4C(r)_i$  with  $C(r)_i$  the quadratic Casimir invariant for the superfield  $\Phi_i$ .

So we have

$$\ln Z_l(\mu, M_a) = \ln Z_l(\Lambda) + \sum_i \left\{ -\frac{A_i}{b'_i} \ln g_i^2(\Lambda) + \left( \frac{A_i}{b'_i} - \frac{A_i}{b'_i - n_1} \right) \ln g_i^2(M_1) + \left( \frac{A_i}{b'_i - n_1} - \frac{A_i}{b'_i - \sum_{k=1}^2 n_k} \right) \ln g_i^2(M_2) \right.$$

$$\left. + \left( \frac{A_i}{b'_i - \sum_{k=1}^2 n_k} - \frac{A_i}{b'_i - \sum_{k=1}^3 n_k} \right) \ln g_i^2(M_3) + \dots + \frac{A_i}{b'_i - N} \ln g_i^2(\mu) \right\} + \dots \quad (3.9)$$

From eqn.(3.9), we get

$$\left( \frac{\partial \ln Z_l(\mu, M_a)}{\partial \ln g_i(\mu')} \right) = 2 \left( \frac{A_i}{b'_i} - \frac{A_i}{b'_i - n_1}, \frac{A_i}{b'_i - n_1} - \frac{A_i}{b'_i - \sum_{k=1}^2 n_k}, \dots, \frac{A_i}{b'_i - \sum_{k=1}^{p-1} n_k} - \frac{A_i}{b'_i - N}, \frac{A_i}{b'_i - N} \right), \quad (3.10)$$

with the column indices  $g_i(M_1), g_i(M_2), \dots, g_i(M_p), g_i(\mu)$ . From the expressions of gauge coupling at scale  $\mu$  within each threshold interval, we can obtain an  $(p+1) \times p$  matrix

$$\begin{array}{cccccc} \ln M_1 & \ln M_2 & \ln M_3 & \dots & \ln M_p \end{array}$$

<sup>1</sup>For example, see appendix A in Ref.[18] for details.



$$\left( \frac{\partial \ln g_i(\mu')}{\partial \ln M_j} \right)_{\mu',j} = \frac{1}{16\pi^2} \begin{pmatrix} b'_i g_i^2(M_1) & 0 & 0 & \cdots & 0 \\ n_1 g_i^2(M_2) & (b'_i - n_1) g_i^2(M_2) & 0 & \cdots & 0 \\ n_1 g_i^2(M_3) & n_2 g_i^2(M_3) & (b'_i - \sum_{i=1}^2 n_i) g_i^2(M_3) & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ n_1 g_i^2(M_p) & n_2 g_i^2(M_p) & n_3 g_i^2(M_p) & \cdots & (b'_i - \sum_{i=1}^{p-1} n_i) g_i^2(M_p) \\ n_1 g_i^2(\mu) & n_2 g_i^2(\mu) & n_3 g_i^2(\mu) & \cdots & n_p g_i^2(\mu) \end{pmatrix} \begin{pmatrix} g_i(M_1) \\ g_i(M_2) \\ g_i(M_3) \\ \cdots \\ g_i(M_p) \\ g_i(\mu) \end{pmatrix} \quad (3.11)$$

So we have

$$U_b \equiv \left( \frac{\partial \ln Z_l(\mu, M_a)}{\partial \ln M_b} \right)^T \equiv - \sum_{i=1,2,3} \frac{A_i}{8\pi^2} U_{b;i} , \quad (3.12)$$

$$U_{b;i} = \begin{pmatrix} n_1 (P_i[1] + Q_i[1]) \\ n_2 (P_i[2] + Q_i[2]) \\ n_3 (P_i[3] + Q_i[3]) \\ \cdots \\ n_p (P_i[p] + Q_i[p]) \end{pmatrix} \equiv \begin{pmatrix} \left[ \frac{n_1 g_i^2(M_1)}{b'_i - n_1} - \frac{n_1 g_i^2(\mu)}{b'_i - N} \right] + \sum_{a=2}^p \frac{n_1 n_a g_i^2(M_a)}{(b'_i - \sum_{k=1}^{a-1} n_k)(b'_i - \sum_{k=1}^a n_k)} \\ \left[ \frac{n_2 g_i^2(M_2)}{b'_i - \sum_{k=1}^2 n_k} - \frac{n_2 g_i^2(\mu)}{b'_i - N} \right] + \sum_{a=3}^p \frac{n_2 n_a g_i^2(M_a)}{(b'_i - \sum_{k=1}^{a-1} n_k)(b'_i - \sum_{k=1}^a n_k)} \\ \left[ \frac{n_3 g_i^2(M_3)}{b'_i - \sum_{k=1}^3 n_k} - \frac{n_3 g_i^2(\mu)}{b'_i - N} \right] + \sum_{a=4}^p \frac{n_3 n_a g_i^2(M_a)}{(b'_i - \sum_{k=1}^{a-1} n_k)(b'_i - \sum_{k=1}^a n_k)} \\ \cdots \cdots \cdots \\ \left[ \frac{n_p g_i^2(M_p)}{b'_i - N} - \frac{n_p g_i^2(\mu)}{b'_i - N} \right] \end{pmatrix}$$

with the column indices corresponding to  $M_1, M_2, \dots, M_p$  etc. Here we rewrite our expressions neatly by define

$$P_i[a] = \frac{g_i^2(M_a)}{b'_i - \sum_{k=1}^a n_k} - \frac{g_i^2(\mu)}{b'_i - N} ,$$

$$Q_i[a] = \sum_{c=a+1}^p \frac{n_c g_i^2(M_c)}{(b'_i - \sum_{k=1}^{c-1} n_k)(b'_i - \sum_{k=1}^c n_k)} , \quad (3.13)$$

within which  $b_i \equiv b'_i - N$  is just the beta function coefficient below all the messenger thresholds. From the previous expressions, we can check that the each row will vanish if we neglect the scale dependence of  $g_i^2$ . This observation agrees with the ordinary conclusion that the trilinear couplings of GMSB vanish if no Yukawa deflections are present.

From the expressions in eqn.(3.12), we can obtain the symmetric matrix (for indices  $j$  and  $k$ )

$$\left( \frac{\partial^2 \ln Z_a(\mu, M_a)}{\partial \ln M_b \partial \ln M_a} \right)^T \equiv - \sum_{i=1,2,3} \frac{4A_i}{(16\pi^2)^2} K_{ab;i} , \quad (3.14)$$

with the contributions  $K_{ab;i}$  from each  $i = 1, 2, 3$  gauge fields given by

$$\begin{pmatrix} n_1^2 X_i[1], & n_1 n_2 Y_i[2], & n_1 n_3 Y_i[3], & n_1 n_4 Y_i[4], & n_1 n_5 Y_i[5], & \cdots, & n_1 n_p Y_i[p] \\ n_1 n_2 Y_i[2], & n_2^2 X_i[2], & n_2 n_3 Y_i[3], & n_2 n_4 Y_i[4], & n_2 n_5 Y_i[5], & \cdots, & n_2 n_p Y_i[p] \\ n_1 n_3 Y_i[3], & n_2 n_3 Y_i[3], & n_3^2 X_i[3], & n_3 n_4 Y_i[4], & n_3 n_5 Y_i[5], & \cdots, & n_3 n_p Y_i[p] \\ n_1 n_4 Y_i[4], & n_2 n_4 Y_i[4], & n_3 n_4 Y_i[4], & n_4^2 X_i[4], & n_4 n_5 Y_i[5], & \cdots, & n_4 n_p Y_i[p] \\ n_1 n_5 Y_i[5], & n_2 n_5 Y_i[5], & n_3 n_5 Y_i[5], & n_4 n_5 X_i[5], & n_5^2 X_i[5], & \cdots, & n_5 n_p Y_i[p] \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ n_1 n_p Y_i[p], & n_2 n_p Y_i[p], & n_3 n_p Y_i[p], & n_4 n_p Y_i[p], & n_5 n_p Y_i[p], & \cdots, & n_p^2 X_i[p], \end{pmatrix}. \quad (3.15)$$

The functions within  $K_{ab;i}$  are defined as

$$X_i[a] = G_i^E[a] + K_i^E[a], \quad Y_i[a] = F_i^E[a] + K_i^E[a], \quad (3.16)$$

within which

$$\begin{aligned} F_i^E[a] &= \frac{g_i^4(M_a)}{b'_i - \sum_{i=1}^a n_i} - \frac{g_i^4(\mu)}{b'_i - N}, \\ G_i^E[a] &= \frac{(b'_i - \sum_{i=1}^{a-1} n_i) g_i^4(M_a)}{n_a (b'_i - \sum_{i=1}^a n_i)} - \frac{g_i^4(\mu)}{b'_i - N}, \\ K_i^E[a] &= \sum_{c=a+1}^p \frac{n_c g_i^4(M_c)}{(b'_i - \sum_{i=1}^{c-1} n_i) (b'_i - \sum_{i=1}^c n_i)}, \end{aligned} \quad (3.17)$$

Here we define the summation to vanish if its index lies out of its definition range. For example,  $\sum_{i=1}^0 (\cdots) \equiv \sum_{i=p+1}^p (\cdots) = 0$ . From the previous expressions, we can check that each non-diagonal element of  $K_{ab}$  will vanish if we neglect the scale dependence of  $g_i^2$ . Only the diagonal elements of  $K_{ab}$  can give non-vanishing values of order

$$K_{aa} \sim n_a^2 \frac{g_i^4}{n_a} = n_a g_i^4. \quad (3.18)$$

The inclusion of top-Yukawa coupling is straightforward in the analytical expressions. The scale dependence of top-Yukawa in the simplest case, in which only the leading top Yukawa  $\alpha_t \equiv y_t^2/4\pi$  and  $\alpha_s \equiv g_s^2/4\pi$  are kept in the anomalous dimension, takes the form

$$\frac{d}{dt} \ln \alpha_t = \frac{1}{\pi} \left( 3\alpha_t - \frac{8}{3}\alpha_s \right), \quad \frac{d}{dt} \ln \alpha_s = \frac{1}{2\pi} b_3 \alpha_s, \quad (3.19)$$

Define  $A = \ln \left( \alpha_t \alpha_s^{\frac{16}{3b_3}} \right)$ , the equation can be written as

$$d[e^{-A}] = -\frac{3}{\pi} \alpha_s^{-\frac{16}{3b_3}} dt = -\frac{6}{b_3} \alpha_s^{-\frac{16}{3b_3}-2} d\alpha_s. \quad (3.20)$$

So we can exactly solve the differential equation to get

$$\left[ \frac{\alpha_t(\mu)}{\alpha_t(\Lambda)} \left( \frac{\alpha_s(\mu)}{\alpha_s(\Lambda)} \right)^{\frac{16}{3b_3}} \right]^{-1} = 1 - \frac{6\alpha_t(\Lambda)}{\frac{16}{3} + b_3} \left[ \alpha_s(\Lambda)^{-1} - \left( \frac{\alpha_s(\mu)}{\alpha_s(\Lambda)} \right)^{-\frac{16}{3b_3}} \alpha_s^{-1}(\mu) \right].$$

Expanding the expressions and neglect high order terms, we finally have

$$[\ln \alpha_t(\mu) - \ln \alpha_t(\Lambda)] \approx \left[ -\frac{8}{3\pi} \alpha_s(\mu) + \frac{3}{\pi} \alpha_t(\mu) \right] \ln \left( \frac{\Lambda}{\mu} \right). \quad (3.21)$$

It can be observed that the (leading order approximation) expression within the square bracket is just the beta function of top Yukawa coupling. As there are no Yukawa deflection contributions related to the introduction of messengers, the Yukawa coupling contributions will not enter the expression within the GMSB part of deflection GM.

### 3.2 Class B: Partition with vanishing gauge beta functions

In previous discussions, apparent poles  $A_i/(b'_i - \sum_{a=1}^j n_a)$  may arise in the expressions if the gauge beta function coefficient  $b'_i - \sum_{a=1}^j n_a = 0$  between certain messenger scales. For example, with  $N > 3$  non-degenerate messengers in  $\mathbf{5} \oplus \bar{\mathbf{5}}$  representation, the beta function for  $\alpha_3$ , which is given by  $b'_3 = -3 + N$ , will vanish after decoupling  $N - 3$  family of vector-like messengers at one-loop level. (The beta function for  $i = 1, 2$  gauge fields will not encounter this possibility.) Such an artificial pole can be resolved by revisiting the deduction procedure of wavefunctions.

Assume that all the ' $i$ '-th gauge coupling beta function coefficients  $b'_i - \sum_{a=1}^{k-1} n_a$  are non-vanishing for  $k < j$ . After integrating out the  $n_j$  family of vector-like messengers at  $M_j$  scale, the beta function coefficient is assumed to vanish (so that  $b'_i - \sum_{a=1}^j n_a = 0$ ).

The wavefunction at the  $M_j$  scale takes value

$$Z_l(M_j, M_a) = Z_l(\Lambda) \prod_{i=1}^3 \left[ \left( \frac{g_i^2(M_1)}{g_i^2(\Lambda)} \right)^{A_i/b'_i} \left( \frac{g_i^2(M_2)}{g_i^2(M_1)} \right)^{A_i/(b'_i - n_1)} \cdots \left( \frac{g_i^2(M_j)}{g_i^2(M_{j-1})} \right)^{A_i/(b'_i - \sum_{a=1}^{j-1} n_a)} \right],$$

$$\left( \frac{y_t^2(M_j)}{y_t^2(\Lambda)} \right)^{B_t} \left( \frac{y_b^2(M_j)}{y_b^2(\Lambda)} \right)^{B_b} \left( \frac{y_\tau^2(M_j)}{y_\tau^2(\Lambda)} \right)^{B_\tau}. \quad (3.22)$$

As the ' $i$ '-th beta function vanishes at one-loop level

$$\frac{dg_i}{dt} = 0, \quad M_{j+1} < \mu < M_j \quad (3.23)$$

it can be seen as a constant between  $M_{j+1} < \mu < M_j$ . Within this range, the RGE invariant became

$$\frac{d}{dt} \ln \left[ Z(\mu) \prod_{l=y_t, y_b, y_\tau} [y_l(\mu)]^{-2B_l} \prod_{k \neq i} [g_k(\mu)]^{-2\tilde{A}_k} \right] = D_i g_i^2(\mu) = D_i g_i^2(M_j), \quad (3.24)$$

and we can deduce that

$$\begin{aligned}
& Z(\mu) \prod_{l=y_t, y_b, y_\tau} [y_l(\mu)]^{-2B_l} \prod_{k \neq i} [g_k(\mu)]^{-2\tilde{A}_k} \\
&= \left( \frac{\mu}{M_j} \right)^{D_i g_i^2(M_j)} \left[ Z(M_j) \prod_{l=y_t, y_b, y_\tau} [y_l(M_j)]^{-2B_l} \prod_{k \neq i} [g_k(M_j)]^{-2\tilde{A}_k} \right]. \quad (3.25)
\end{aligned}$$

The value  $D_i \equiv A_i/8\pi^2$  with the value  $A_i$  given in the appendix A. We keep to use  $'D'_i$  in this paper to indicate clearly the consequence of vanishing one-loop beta functions.

So, for  $M_{j+1} < \mu < M_j$

$$\begin{aligned}
Z(\mu) &= Z(M_j) \left( \frac{\mu}{M_j} \right)^{D_i g_i^2(M_j)} \prod_{l=y_t, y_b, y_\tau} \left[ \frac{y_l(M_j)}{y_l(\mu)} \right]^{-2B_l} \prod_{k \neq i} \left[ \frac{g_k(M_j)}{g_k(\mu)} \right]^{-2\tilde{A}_k} \\
&= Z(\Lambda) \prod_{l=y_t, y_b, y_\tau} \left[ \frac{y_l(\Lambda)}{y_l(\mu)} \right]^{-2B_l} \prod_{k \neq i} \left[ \left( \frac{g_k^2(M_1)}{g_k^2(\Lambda)} \right)^{\frac{A_k}{b'_k}} \left( \frac{g_k^2(M_2)}{g_k^2(M_1)} \right)^{\frac{A_k}{b'_k - n_1}} \dots \left( \frac{g_k^2(\mu)}{g_k^2(M_j)} \right)^{\frac{A_k}{b'_k - \sum_{a=1}^j n_a}} \right], \\
&\quad \left[ \left( \frac{g_i^2(M_1)}{g_i^2(\Lambda)} \right)^{\frac{A_i}{b'_i}} \left( \frac{g_i^2(M_2)}{g_i^2(M_1)} \right)^{\frac{A_i}{b'_i - n_1}} \dots \left( \frac{g_i^2(M_j)}{g_i^2(M_{j-1})} \right)^{\frac{A_i}{b'_i - \sum_{a=1}^{j-1} n_a}} \right] \left( \frac{\mu}{M_j} \right)^{D_i g_i^2(M_j)}.
\end{aligned}$$

and for  $\mu < M_p$

$$\begin{aligned}
Z(\mu) &= Z(\Lambda) \prod_{l=y_t, y_b, y_\tau} \left[ \frac{y_l(\Lambda)}{y_l(\mu)} \right]^{-2B_l} \\
&\quad \prod_{k \neq i} \left[ \left( \frac{g_k^2(M_1)}{g_k^2(\Lambda)} \right)^{\frac{A_k}{b'_k}} \left( \frac{g_k^2(M_2)}{g_k^2(M_1)} \right)^{\frac{A_k}{b'_k - n_1}} \dots \left( \frac{g_k^2(M_j)}{g_k^2(M_{j-1})} \right)^{\frac{A_k}{b'_k - \sum_{a=1}^j n_a}} \dots \left( \frac{g_k^2(\mu)}{g_k^2(M_p)} \right)^{\frac{A_k}{b'_k}} \right], \\
&\quad \left[ \left( \frac{g_i^2(M_1)}{g_i^2(\Lambda)} \right)^{\frac{A_i}{b'_i}} \left( \frac{g_i^2(M_2)}{g_i^2(M_1)} \right)^{\frac{A_i}{b'_i - n_1}} \dots \left( \frac{g_i^2(M_j)}{g_i^2(M_{j-1})} \right)^{\frac{A_i}{b'_i - \sum_{a=1}^{j-1} n_a}} \right] \left( \frac{M_{j+1}}{M_j} \right)^{D_i g_i^2(M_j)} \\
&\quad \left( \frac{g_i^2(M_{j+2})}{g_i^2(M_{j+1})} \right)^{\frac{A_i}{b'_i - \sum_{a=1}^{j+1} n_a}} \dots \left( \frac{g_i^2(\mu)}{g_i^2(M_p)} \right)^{\frac{A_i}{b'_i}}. \quad (3.26)
\end{aligned}$$

For  $\mu < M_p$ ,

$$\frac{d \ln Z [\mu, g_i(\mu'), M_j, M_{j+1}]}{d \ln M_a} = \left[ \sum_{g_i(\mu')} \frac{\partial g_i(\mu')}{\partial \ln M_a} \frac{\partial}{\partial g_i(\mu')} + \frac{\partial}{\partial \ln M_a} \right] \ln Z [\mu, g_i(\mu'), M_j, M_{j+1}].$$

with the last term gives non-vanishing contributions only for  $a = j, j + 1$

$$\frac{\partial}{\partial \ln M_a} \ln Z [\mu, g_i(\mu'), M_j, M_{j+1}] = [\delta_{a, j+1} - \delta_{a, j}] D_i g_i^2(M_j), \quad (3.27)$$

From the general expressions, we can see that the ' $j'$ '-th and ' $(j+1)'$ '-th components will change into

$$\left( \frac{\partial \ln Z[\mu; g_i(\mu'), M_a]}{\partial \ln g_i(\mu')} \right)_{j,j+1} = 2 \left( \frac{A_i}{b'_i - \sum_{a=1}^{j-1} n_a} + D_i g_i^2(M_j) \ln \frac{M_{j+1}}{M_j}, -\frac{A_i}{b'_i - \sum_{a=1}^{j+1} n_a} \right),$$

while other columns are unchanged as eqn.(3.10) if  $b_i(\mu) = 0$  for  $M_{j+1} < \mu < M_j$ . The matrix  $\partial \ln g_i(\mu')/\partial M_j$ , which is given by eqn.(3.11), is unchanged. Then the contributions from the ' $i'$ '-th gauge field, which has vanishing beta functions between  $[M_{j+1}, M_j]$ , are given as

$$\begin{aligned} U_{b;i} &\equiv \left. \frac{d \ln Z[g_i(\mu'), M_n]}{d \ln M_b} \right|_i \\ &= \left[ \frac{\partial g_i(\mu')}{\partial \ln M_b} \frac{\partial}{\partial g_i(\mu')} + \frac{\partial}{\partial \ln M_b} \right] \ln Z[g_i(\mu'), M_j, M_{j+1}] \\ &= -\frac{A_i}{8\pi^2} \begin{pmatrix} n_1 \left( P_i^S[1] + Q_i^S[1] - \frac{D_i}{A_i} g_i^4(M_j) \ln \frac{M_{j+1}}{M_j} \right), \\ n_2 \left( P_i^S[2] + Q_i^S[2] - \frac{D_i}{A_i} g_i^4(M_j) \ln \frac{M_{j+1}}{M_j} \right), \\ \dots \\ n_j \left( P_i^S[j] + Q_i^S[j] - \frac{D_i}{A_i} g_i^4(M_j) \ln \frac{M_{j+1}}{M_j} \right) + \frac{8\pi^2}{A_i} D_i g_i^2(M_j), \\ n_{j+1} \left( P_i^S[j+1] + Q_i^S[j+1] \right) - \frac{8\pi^2}{A_i} D_i g_i^2(M_j), \\ n_{j+2} \left( P_i^S[j+2] + Q_i^S[j+2] \right), \\ \dots \\ n_p \left( P_i^S[p] + Q_i^S[p] \right). \end{pmatrix}. \end{aligned} \quad (3.28)$$

Within the deduction, we use the fact that  $g_i(M_j) = g_i(M_{j+1})$  and  $b'_i - \sum_{k=1}^j n_k = 0$ . Besides, we define within the expression

$$\begin{aligned} Q_i^S[c] &= \begin{cases} \sum_{a=c+1; a \neq j, j+1}^p \frac{n_a g_i^2(M_a)}{(b'_i - \sum_{k=1}^{a-1} n_k)(b'_i - \sum_{k=1}^a n_k)} + \frac{(n_j + n_{j+1}) g_i^2(M_j)}{(b'_i - \sum_{k=1}^{j-1} n_k)(b'_i - \sum_{k=1}^{j+1} n_k)}, & 1 \leq c \leq j \\ \sum_{a=c+1}^p \frac{n_a g_i^2(M_a)}{(b'_i - \sum_{k=1}^{a-1} n_k)(b'_i - \sum_{k=1}^a n_k)}, & j+1 \leq c \leq p \end{cases} \\ P_i^S[c] &= \begin{cases} \frac{g_i^2(M_c)}{b'_i - \sum_{k=1}^c n_k} - \frac{g_i^2(\mu)}{b'_i - N}, & c \neq j, j+1 \\ -\frac{g_i^2(\mu)}{b'_i - N}, & c = j, j+1 \end{cases} \end{aligned} \quad (3.29)$$

Within the expression

$$\frac{(n_j + n_{j+1}) g_i^2(M_j)}{(b'_i - \sum_{k=1}^{j-1} n_k)(b'_i - \sum_{k=1}^{j+1} n_k)} = -\frac{n_j + n_{j+1}}{n_j n_{j+1}} g_i^2(M_j). \quad (3.30)$$

We note that when  $c$  takes value  $j-1$  or  $j$  in the summation of  $Q_i^S[c]$ , the sum skip  $j, j+1$  and begins at  $a = j+2$ .

From the previous expressions, we can see that the each row will vanish if we neglect the scale dependence of  $g_i^2$  and higher order  $g_i^6$  terms. In fact, with such an approximation, the  $j$ -th and  $j+1$ -th row is given by

$$\begin{aligned} U_{b;j} &\sim -\frac{g^2}{n_j}n_j + g^2 \sim 0, \\ U_{b;j+1} &\sim \frac{g^2}{n_{j+1}}n_{j+1} - g^2 \sim 0. \end{aligned} \quad (3.31)$$

In the summation

$$\frac{\partial \ln Z[\mu, g_i(\mu'), M_n]}{\partial \ln M_b} = \sum_{i=1}^3 \frac{\partial \ln Z[\mu, g_i(\mu'), M_n]}{\partial \ln M_b} \Big|_i, \quad (3.32)$$

the expressions for  $i = 1, 2$  gauge fields (which have no vanishing beta functions) are still given by

$$\left( \frac{\partial \ln Z[\mu, g_i(\mu')]}{\partial \ln M_b} \right) \Big|_{i=1,2} \equiv -\frac{A_i}{8\pi^2} \begin{pmatrix} n_1 (P_i[1] + Q_i[1]) \\ n_2 (P_i[2] + Q_i[2]) \\ n_3 (P_i[3] + Q_i[3]) \\ \dots \\ n_p (P_i[p] + Q_i[p]) \end{pmatrix}, \quad (3.33)$$

from eqn.(3.12).

With previous results, we can derive the expression of  $\frac{\partial^2}{\partial \ln M_a \partial \ln M_b} Z[\mu; g_i(\mu'), M_n]$  from ' $i$ '-th ( here  $i = 3$ ) gauge fields

$$\begin{aligned} &\frac{\partial}{\partial \ln M_a} \left( \frac{\partial \ln Z[g_i(\mu'), M_n]}{\partial \ln M_b} \Big|_i \right) \equiv -\frac{A_i}{8\pi^2} \left[ \frac{\partial \ln g_j(\mu')}{\partial \ln M_a} \frac{\partial}{\partial \ln g_j(\mu')} + \delta_{a;j,j+1} \frac{\partial}{\partial M_{j,j+1}} \right] V_{b;i} \\ &= -\frac{4A_i}{(16\pi^2)^2} K_{ab;i} \end{aligned} \quad (3.34)$$

with  $K_{ab;i}$  a symmetric matrix given as

$$\begin{pmatrix} n_1^2 J[1] & n_1 n_2 H[2] & n_1 n_3 H[3] & \dots & n_1 n_j H[j] & n_1 n_{j+1} H[j+1] & n_1 n_{j+2} H[j+2] & \dots & n_1 n_p H[p] \\ n_1 n_2 H[2] & n_2^2 J[2] & n_2 n_3 H[3] & \dots & n_2 n_j H[j] & n_2 n_{j+1} H[j+1] & n_2 n_{j+2} H[j+2] & \dots & n_2 n_p H[p] \\ n_1 n_3 H[3] & n_3 n_2 H[3] & n_3^2 J[3] & \dots & n_3 n_j H[j] & n_3 n_{j+1} H[j+1] & n_3 n_{j+2} H[j+2] & \dots & n_3 n_p H[p] \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ n_1 n_j H[j] & n_2 n_j H[j] & n_3 n_j H[j] & \dots & n_j^2 J[j] & n_j n_{j+1} H[j+1] & n_j n_{j+2} H[j+2] & \dots & n_j n_p H[p] \\ n_1 n_{j+1} H[j+1] & n_2 n_{j+1} H[j+1] & n_3 n_{j+1} H[j+1] & \dots & n_j n_{j+1} H[j+1] & n_{j+1}^2 J[j+1] & n_{j+1} n_{j+2} H[j+2] & \dots & n_{j+1} n_p H[p] \\ n_1 n_{j+2} H[j+2] & n_2 n_{j+2} H[j+2] & n_3 n_{j+2} H[j+2] & \dots & n_j n_{j+2} H[j+2] & n_{j+1} n_{j+2} H[j+2] & n_{j+2}^2 J[j+2] & \dots & n_{j+2} n_p H[p] \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ n_1 n_p H[p] & n_2 n_p H[p] & n_3 n_p H[p] & \dots & n_j n_p H[p] & n_{j+1} n_p H[p] & n_{j+2} n_p H[p] & \dots & n_p^2 J[p] \end{pmatrix}, \quad (3.35)$$

The functions within  $K_{ab;i}$  are defined as

$$J[m] = \begin{cases} G_i^S[m] + K_i^S[m] - 2\frac{D_i}{A_i} g_i^6(M_j) \ln \frac{M_{j+1}}{M_j}, & 1 \leq m \leq j-1 \\ G_i^S[j] + K_i^S[j] + 16\pi^2 \frac{D_i}{n_j A_i} g_i^4(M_j) - 2\frac{D_i}{A_i} g_i^6(M_j) \ln \frac{M_{j+1}}{M_j}, & m = j \\ G_i^S[m] + K_i^S[m], & j+1 \leq m \leq p \end{cases}$$

$$H[m] = \begin{cases} F_i^S[m] + K_i^S[m] - 2\frac{D_i}{A_i}g_i^6(M_j) \ln \frac{M_{j+1}}{M_j}, & 1 \leq m \leq j-1 \\ F_i^S[j] + K_i^S[j+1] + 8\pi^2 \frac{D_i}{n_j A_i} g_i^4(M_j) - 2\frac{D_i}{A_i}g_i^6(M_j) \ln \frac{M_{j+1}}{M_j}, & m = j \\ F_i^S[j+1] + K_i^S[j+1] - 8\pi^2 \frac{D_i}{n_{j+1} A_i} g_i^4(M_j), & m = j+1 \\ F_i^S[m] + K_i^S[m], & j+2 \leq m \leq p \end{cases}$$

with

$$K_i^S[c] = \begin{cases} \sum_{a=c+1; a \neq j, j+1}^p \frac{n_a g_i^4(M_a)}{(b'_i - \sum_{k=1}^{a-1} n_k)(b'_i - \sum_{k=1}^a n_k)} + \frac{(n_j + n_{j+1})g_i^4(M_j)}{(b'_i - \sum_{k=1}^{j-1} n_k)(b'_i - \sum_{k=1}^{j+1} n_k)}, & 1 \leq c \leq j \\ \sum_{a=c+1}^p \frac{n_a g_i^4(M_a)}{(b'_i - \sum_{k=1}^{a-1} n_k)(b'_i - \sum_{k=1}^a n_k)}, & j+1 \leq c \leq p \end{cases}$$

$$F_i^S[c] = \begin{cases} \frac{g_i^4(M_c)}{b'_i - \sum_{k=1}^c n_k} - \frac{g_i^4(\mu)}{b'_i - N}, & c \neq j, j+1 \\ \frac{n_j + n_{j+1}}{b'_i - \sum_{k=1}^{j-1} n_k} \frac{g_i^4(M_c)}{b'_i - \sum_{k=1}^{j+1} n_k} - \frac{g_i^4(\mu)}{b'_i - N}, & c = j \\ -\frac{g_i^4(\mu)}{b'_i - N}, & c = j+1 \end{cases} \quad (3.36)$$

$$G_i^S[c] = \begin{cases} \frac{(b'_i - \sum_{k=1}^{c-1} n_k)g_i^4(M_a)}{n_c(b'_i - \sum_{k=1}^c n_k)} - \frac{g_i^4(\mu)}{b'_i - N}, & c \neq j, j+1 \\ -\frac{g_i^4(\mu)}{b'_i - N}, & c = j, j+1 \end{cases} \quad (3.37)$$

From the previous expressions, we can check that each non-diagonal element of  $K_{ab}(a, b \neq j, j+1)$  will vanish if we neglect the scale dependence of  $g_i^2$  and higher order  $g_i^6$  terms. The diagonal elements of  $K_{ab}(a, b \neq j, j+1)$  can give non-vanishing values of order

$$K_{aa} \equiv n_a^2 J[a] \sim n_a^2 \frac{g_i^4}{n_a} = n_a g_i^4. \quad (3.38)$$

The  $J[j]$  and  $H[j]$  term will be given as

$$\begin{aligned} J[j] &\sim -\frac{g_i^4}{n_j} + \frac{2}{n_j} g_i^4, & J[j+1] &\sim \frac{g_i^4}{n_{j+1}}, \\ H[j] &\sim -\frac{g_i^4}{n_j} + \frac{g_i^4}{n_j} \sim 0, & H[j+1] &\sim \frac{g_i^4}{n_{j+1}} - \frac{g_i^4}{n_{j+1}} \sim 0, \end{aligned} \quad (3.39)$$

if we neglect the scale dependence of  $g_i^2$  and higher order  $g_i^6$  terms.

The contributions from  $i = 1, 2$  gauge fields are still given by eqn.(3.15). The total contributions are given by the sum of  $i = 1, 2, 3$  gauge fields.

### 3.3 Dependence of $\ln M_a$ on $\ln X$

As noted before, with non-trivial  $U(1)_R$  symmetry, the messenger determinant is proven by [6] to be a monomial in  $X$

$$\det(\lambda_{ij}X + \kappa_{ij}T) = X^{n_0} G(\lambda, \kappa T), \quad (3.40)$$

with  $\kappa_{ij}\langle T \rangle = m_{ij}$ . We need to replace the lowest component VEV  $\langle X \rangle \equiv M$  of  $X$ , which should be substituted in  $\mathcal{M}$  to get all the eigenvalues, by its superspace extension  $\sqrt{X^\dagger X}$  into  $\mathcal{M}$  to incorporate the SUSY breaking effects.

Knowing the value of the determinant, it is still nontrivial to express the eigenvalues of  $\mathcal{M}$  in terms of  $\langle X \rangle$ . Fortunately, the asymptotic behavior will display a simple form. In large  $\langle X \rangle$  region,  $r_\lambda \equiv \text{rank}(\lambda_{ij})$  messengers acquire masses  $\mathcal{O}(\langle X \rangle)$  while the remaining  $N - r_\lambda$  messengers acquire masses of order

$$M_i \sim \frac{m^{n_i+1}}{\langle X \rangle^{n_i}}, \quad \sum_{i=1}^{N-r_\lambda} n_i = r_\lambda - n_0, \quad (3.41)$$

with  $n_i \geq 0$ . At small  $\langle X \rangle$  region,  $r_m \equiv \text{rank}(m_{ij})$  messengers acquire masses  $\mathcal{O}(m)$  while the remaining  $N - r_m$  messengers acquire masses of order

$$M_i \sim \frac{\langle X \rangle^{\tilde{n}_i+1}}{m^{\tilde{n}_i}}, \quad \sum_{i=1}^{N-r_m} (\tilde{n}_i + 1) = n_0. \quad (3.42)$$

with  $\tilde{n}_i \geq 0$ .

Depending on the singularity properties of the messenger mass matrix, we have the following the discussions

- Type I:  $\det m \neq 0$ .

In the basis in which  $m$  is diagonal, it can easily obtain [6] that the eqn.(3.40) takes the form

$$n = 0, \quad \det(\lambda\langle X \rangle + m) = \det m, \quad (3.43)$$

which necessarily imply  $\det \lambda = 0$ . As the matrix is upper triangular, the eigenvalues are  $m_{ii}$  that do not depend on  $\langle X \rangle$ .

So in this case, we will have

$$\frac{\partial \ln M_i}{\partial \ln |X|} \equiv 0. \quad (3.44)$$

So we can see that the gauginos, the trilinear couplings as well as the sfermions will not receive any gauge mediation contributions.

- Type II:  $\det \lambda \neq 0$ .

Similarly, we can obtain an upper triangular matrix with eigenvalues the diagonal elements of diagonalized matrix  $\lambda'_{ii}$ . The determinant is

$$n = N, \quad \det(\lambda\langle X \rangle + m) = \langle X \rangle^N \det \lambda. \quad (3.45)$$

So the eigenvalues will be  $\lambda'_{ii}\langle X \rangle$  and depends linearly on  $\langle X \rangle$ . We will arrange  $\lambda'_{ii}$  to obtain the eigenvalues  $\tilde{M}_1(\langle X \rangle), \tilde{M}_2(\langle X \rangle), \dots, \tilde{M}_N(\langle X \rangle)$ .



Suppose the  $T_i \equiv \lambda'_{ii}$  are ordered so as that  $T_1 \geq T_2 \geq T_3 \cdots \geq T_N$ , we define

$$V_i \equiv \frac{d \ln M_i}{d \ln |X|} \equiv (1, 1, 1, \dots, 1). \quad (3.46)$$

For degenerate eigenvalues

$$\begin{aligned} \tilde{M}_1 = \tilde{M}_2 = \cdots = \tilde{M}_{n_1} &\equiv M_1 = \lambda'_{n_1 n_1} \langle X \rangle, \quad \tilde{M}_{n_1+1} = \tilde{M}_{n_1+2} = \cdots = \tilde{M}_{n_2} \equiv M_2 = \lambda'_{n_2 n_2} \langle X \rangle, \\ \tilde{M}_{n_2+1} = \tilde{M}_{n_2+2} = \cdots = \tilde{M}_{n_3} &\equiv M_3 = \lambda'_{n_3 n_3} \langle X \rangle, \quad \cdots \\ \tilde{M}_{n_{p-1}+1} = \tilde{M}_{n_{p-1}+2} = \cdots = \tilde{M}_{n_p} &\equiv M_p = \lambda'_{n_p n_p} \langle X \rangle, \end{aligned} \quad (3.47)$$

with  $M_1 \geq M_2 \geq \cdots \geq M_p$ , the matrix  $V_i$  reduces to a  $1 \times p$  matrix

$$V_i \equiv \frac{d \ln M_i}{d \ln |X|} \equiv (1, 1, \dots, 1). \quad (3.48)$$

So the soft SUSY breaking parameters from GMSB

– The gaugino mass:

$$M_i = \frac{F_X}{M} \frac{g_i^2}{16\pi^2} n_0. \quad (3.49)$$

– GMSB contributions to trilinear terms:

$$A_i = \frac{A_{ijk}}{y_{ijk}} = \sum_{i,j,k} \frac{F_X}{2M} \frac{\partial \ln Z_i(\mu, M_a)}{\partial \ln |X|} = \frac{F_X}{2M} \sum_{i,j,k} \sum_{v=1}^N U_b V_b. \quad (3.50)$$

– Pure GMSB contributions to soft sfermion masses:

$$\begin{aligned} m_i^2 &= -\frac{F_X^2}{4M^2} \frac{\partial^2}{\partial \ln |X|^2} \ln Z_i(\mu, M_a) \\ &= \frac{F_X^2}{4M^2} \sum_{i=1,2,3} \frac{4A_i}{(16\pi^2)^2} \sum_{a,b} V_a K_{ab;i} V_b. \end{aligned} \quad (3.51)$$

We should note that if hierarchical structure appears within the diagonal elements of the diagonalized matrix  $\lambda'_{ii} (i = 1, \dots, N)$ , the splitting of the messenger scales will lead to large GMSB contributions.

- Type III:  $\det m = \det \lambda = 0$ .

As the matrix  $\lambda X + m$  is non-singular, its eigenvalues can be written as  $x_1, \dots, x_n$  which should satisfy

$$\prod_i x_i = \det(\lambda X + m) = X^{n_0} G(\lambda, m), \quad (3.52)$$

and

$$\sum_i x_i = -Tr(\lambda X + m) = cX + d. \quad (3.53)$$



So we can obtain the GMSB contributions

$$\begin{aligned}
A_i &= \frac{A_{ijk}}{y_{ijk}} = \sum_{i,j,k} \frac{F_X}{2M} \frac{\partial \ln Z_l(\mu, M_a)}{\partial \ln |X|} = \frac{F_X}{2M} \sum_{i,j,k} \sum_{v=1}^p U_b V_b, \\
m_i^2 &= -\frac{F_X^2}{4M^2} \frac{\partial^2}{\partial \ln |X|^2} \ln Z_i(\mu, M_a) \\
&= -\frac{F_X^2}{4M^2} \sum_{i=1,2,3} \frac{4A_i}{(16\pi^2)^2} \sum_{a,b} V_a K_{ab;i} V_b.
\end{aligned} \tag{3.60}$$

with the  $V_a$  takes the value in eqn.(3.58). The partition of  $N$  can be obtained numerically by diagonalizing  $\lambda\langle X \rangle + m$  to obtain its eigenvalues as functions of  $\langle X \rangle$ .

The inclusion of EGM in deflected AMSB is straightforward. The AMSB type contributions can be given as

$$\begin{aligned}
\frac{\partial}{\partial \ln \mu} \ln [Z_i(\mu, M_a)] &= -\frac{1}{8\pi^2} G_i^- [g_l(\mu), y_l(\mu)], \\
\frac{\partial^2}{\partial (\ln \mu)^2} \ln [Z_i(\mu, M_a)] &= -\frac{1}{8\pi^2} \left[ \frac{\partial g_l(\mu)}{\partial \ln \mu} \frac{\partial}{\partial g_l(\mu)} + \frac{\partial y_l(\mu)}{\partial \ln \mu} \frac{\partial}{\partial y_l(\mu)} \right] G_i^- [g_l(\mu), y_l(\mu)] \\
&= -\frac{2}{(16\pi^2)^2} \left[ \beta_{g_l} \frac{\partial}{\partial g_l(\mu)} + \beta_{y_l} \frac{\partial}{\partial y_l(\mu)} \right] G_i^- [g_l(\mu), y_l(\mu)].
\end{aligned} \tag{3.61}$$

The interference terms between AMSB and GMSB can also easily obtained with eqn.(3.12) and eqn.(3.29).

In ref.(3), the '*effective messenger number*' is defined as

$$N_{eff} \equiv \frac{\Lambda_G^2}{\Lambda_S^2}, \tag{3.62}$$

with

$$M_i = \frac{g_i^2}{16\pi^2} \Lambda_G, \quad m_{\tilde{f}}^2 = 2 \frac{g_i^4}{(16\pi^2)^2} \sum_i C_{\tilde{f}}(r) \Lambda_S^2. \tag{3.63}$$

So the approximate value of  $N_{eff}$  can be given as

$$N_{eff} = \frac{n_0^2 g_i^4}{\sum_{a,b} V_a K_{ab;i} V_b}, \tag{3.64}$$

by neglecting the scale dependence of  $g_i$  and higher order terms in the expressions of soft SUSY parameters. With previous approximation, the value of  $N_{eff}$  in Type II can be calculated to be  $N_{eff} = N$  after simplifications. While in Type III EGM, it can be calculated to be  $N_{eff} = n_0$ . Such a result holds for both Class A and Class B. Taking into account the scale dependence of  $g_i$ ,  $N_{eff}$  can be different to  $n_0$  and  $N$ .

#### 4. Messengers on GUT and Landau Pole

We must ensure that no Landau pole will be reached below the GUT scale. It is obvious that the gauge coupling unification will be preserved because the messengers are fitted into complete SU(5) representations. The presence of (complete GUT representation) messenger fields at an intermediate scale does not modify the value of  $M_{GUT}$ . However, proton decay could possibly set constraints on the gauge couplings at the GUT scale.

We can define the quantity

$$\delta = - \sum_{r=1}^p \frac{n_r}{2\pi} \ln \frac{M_{GUT}}{M_r} , \quad (4.1)$$

which contributes to the inverse gauge coupling strength. Perturbativity at the GUT scale set a bound on this quantity

$$|\delta| \lesssim 24.3 . \quad (4.2)$$

with 24.3 the value of  $\alpha_{GUT}^{-1}$  without additional messengers. The SUSY scale is taken to be 2 TeV.

Proton decay experiments will also constrained the value of  $\delta$ . As the proton decay induced from the triplet Higgs depends on the scale of the triplets, we just take constraints from proton decay induced by heavy gauge bosons. The decay channel  $p \rightarrow \pi^0 e^+$  has the lifetime

$$\tau(p \rightarrow \pi^0 e^+) = \frac{4f_\pi^2 M_X^4}{\pi m_p \alpha_{GUT}^2 (1 + D + F)^2 \alpha_N^2 [A_{R;1}^2 + (1 + |V_{ud}|^2) A_{R;2}^2]} \left[ 1 + 2 \frac{m_\pi^2}{m_p^2} \right] . \quad (4.3)$$

With updated experimental bounds from Super-Kamiokande[19]  $\tau > 1.67 \times 10^{34}$  years, we can constraint the inputs

$$\alpha_{GUT} \lesssim (5.27)^{-1} , \quad (4.4)$$

by taking  $f_\pi = 131$  MeV, chiral Lagrangian factor  $1 + D + F = 2.27$  with  $D = 0.80, F = 0.47$ [20], the hadronic matrix element  $\alpha_N = 0.0112$  GeV<sup>3</sup> (at renormalization scale  $\mu = 2$ GeV) and  $A_{R;1} = A_{R;2} \approx 5$ . This value constrained  $\delta$  to be

$$|\delta| \lesssim 19 . \quad (4.5)$$

We should note that  $A_{R;1}, A_{R;2}$ , which represent the renormalization effects resulting from the anomalous dimensions of the operators, will also be amended by presence of vector-like messengers[21]. They are defined as

$$A_{R;1} = A_L A_{S;1} , \quad A_{R;2} = A_L A_{S;2} , \quad (4.6)$$

with  $A_L, A_{S;i}$  the long and short distance factors, respectively. Here the long-distance contribution  $A_L$  is taken to be 1.25. The short distance factors will be changed into

$$A_{S;i} = \left( \prod_j \frac{\alpha_j(M_{SUSY})}{\alpha_j(M_Z)} \right)^{\frac{\gamma_{ji}^{SM}}{b_j^{SM}}} \left( \prod_j \frac{\alpha_j(M_{mess})}{\alpha_j(M_{SUSY})} \right)^{\frac{\gamma_{ji}^{MSSM}}{b_j^{MSSM}}} \left( \prod_j \frac{\alpha_j(M_{GUT})}{\alpha_j(M_{mess})} \right)^{\frac{\gamma_{ji}^{MSSM}}{b_j^{MSSM}}} ,$$

$$\begin{aligned}
&= \left( \prod_j \frac{\alpha_j(M_{SUSY})}{\alpha_j(M_Z)} \right)^{\frac{\gamma_{j;i}^{SM}}{b_j^{SM}}} \left( \prod_j \frac{\alpha_j(M_{GUT})}{\alpha_j(M_{SUSY})} \right)^{\frac{\gamma_{j;i}^{MSSM}}{b_j^{MSSM}}} \left( \prod_j \frac{\alpha_j(M_{GUT})}{\alpha_j(M_{mess})} \right)^{\frac{\gamma_{j;i}^{MSSM}}{b'_j} - \frac{\gamma_{j;i}^{MSSM}}{b_j^{MSSM}}}, \\
&\equiv A_{S;i}^0 \left( \prod_j \frac{\alpha_j(M_{GUT})}{\alpha_j(M_{mess})} \right)^{\frac{\gamma_{j;i}^{MSSM}}{b'_j} - \frac{\gamma_{j;i}^{MSSM}}{b_j^{MSSM}}}. \tag{4.7}
\end{aligned}$$

in the case that one vector-like family of messengers at scale  $M_{mess}$  are present. Results with multiple messenger thresholds can be trivially extended. The relevant coefficients within the expressions are given[22] as

$$\begin{aligned}
\gamma_{j;1}^{SM} &= (2, 9/4, 11/20), \quad \gamma_{j;2}^{SM} = (2, 9/4, 23/20), \\
\gamma_{j;1}^{MSSM} &= (4/3, 3/2, 11/30), \quad \gamma_{j;2}^{SM} = (4/3, 3/2, 23/30), \tag{4.8}
\end{aligned}$$

with  $b_j$  the relevant gauge beta functions upon each threshold. The multiple factor for  $A_{S;i}^0$  in the presence of messengers is given approximately by

$$\begin{aligned}
F_1 &= \left( \prod_j \frac{\alpha_j(M_{GUT})}{\alpha_j(M_{mess})} \right)^{\frac{\gamma_{j;i}^{MSSM}}{b'_j} - \frac{\gamma_{j;i}^{MSSM}}{b_j^{MSSM}}} \\
&\approx \left[ 1 + \frac{b'_j}{2\pi} \alpha_j(M_{GUT}) \ln \frac{M_{GUT}}{M_{mess}} \right]^{\frac{\gamma_{j;i}^{MSSM}}{b'_j} - \frac{\gamma_{j;i}^{MSSM}}{b_j^{MSSM}}}, \\
&\approx 1 - \frac{\gamma_{j;i}^{MSSM}}{2\pi} \frac{\Delta b_j^m}{b_j^{MSSM}} \alpha_j(M_{GUT}) \ln \frac{M_{GUT}}{M_{mess}}, \tag{4.9}
\end{aligned}$$

in which we define  $\Delta b_j^m = b'_j - b_j^{MSSM} = n_1$ . This multiple factor can be easily extended to include multiple messengers. For example, with additional messenger thresholds at  $M_2$ , the new multiple factor is given by

$$F_2 \approx 1 - \frac{\gamma_{j;i}^{MSSM}}{2\pi} \frac{n_2}{b_j^{MSSM} + n_1} \alpha_j(M_{GUT}) \ln \frac{M_{GUT}}{M_2}, \tag{4.10}$$

with the total multiple factor

$$F = \prod_k F_k \approx 1 - \frac{\gamma_{j;i}^{MSSM}}{2\pi} \alpha_j(M_{GUT}) \sum_{k=1}^p \frac{n_k}{b_j^{MSSM} + \sum_{l=1}^{k-1} n_l} \ln \frac{M_{GUT}}{M_k}. \tag{4.11}$$

As the coefficients  $A_{R;1}, A_{R;2}$  depend on the messenger scales, the proton decay constraints will feed back into the constraints on  $\delta$ . Detailed discussions on constraints for  $\delta$  will be given in our subsequent studies.

## 5. Conclusions

Extraordinary gauge mediation extension of deflected AMSB scenarios can be interesting because it can accommodate together the deflection in the Kahler potential and the superpotential. We revisit the EGM scenario and derive the analytical expressions for soft SUSY breaking parameters in EGM and EGM extension of deflected AMSB scenarios with wavefunction renormalization approach, especially the case with vanishing gauge beta-function at an intermediate energy scale. We find in EGM that large hierarchy among the messenger thresholds may indicate non-negligible contributions to trilinear couplings at the lightest messenger threshold scale. The Landau pole and proton decay constraints are also discussed.

### A. Coefficients In the Wavefunction of MSSM Superfields

From the anomalous dimension

$$\frac{d}{dt} \ln Z_f = \sum_{l=g_3, g_2, g_1} 2\tilde{A}_l \frac{d \ln g_l}{dt} + \sum_{l=y_t, y_b, y_\tau} 2B_l \frac{d \ln y_l}{dt},$$

in the basis of  $(y_t^2, y_b^2, y_\tau^2, g_3^2, g_2^2, g_1^2)$ , the coefficients  $\tilde{A}_l, B_l$  can be solved. Expressions of the coefficients had already been obtained in our previous paper[18]. The coefficient  $A_i$  are listed in Table.1.

**Table 1:** The gauge field coefficients  $A_i (i = 1, 2, 3)$  within the wavefunction for MSSM superfields:

	$A_3(g_3)$	$A_2(g_2)$	$A_1(g_1)$
$Q_3$	$\frac{128}{61}$	$\frac{87}{61}$	$-\frac{11}{61}$
$U_3$	$\frac{144}{61}$	$-\frac{108}{61}$	$\frac{144}{305}$
$D_3$	$\frac{112}{61}$	$-\frac{84}{61}$	$\frac{112}{305}$
$L_3$	$\frac{80}{61}$	$\frac{123}{61}$	$-\frac{103}{305}$
$E_3$	$\frac{160}{61}$	$-\frac{120}{61}$	$\frac{61}{32}$
$H_u$	$-\frac{272}{61}$	$\frac{21}{61}$	$-\frac{89}{305}$
$H_d$	$-\frac{240}{61}$	$-\frac{3}{61}$	$-\frac{57}{305}$
$Q_2$	$\frac{16}{3}$	3	$\frac{1}{15}$
$U_2$	$\frac{16}{3}$	0	$\frac{16}{15}$
$D_2$	$\frac{16}{3}$	0	$\frac{4}{15}$
$L_2$	0	3	$\frac{3}{5}$
$E_2$	0	0	$\frac{12}{5}$

### Acknowledgments

This work was supported by the Natural Science Foundation of China under grant numbers 11675147,11775012.

## References

- [1] G. Aad et al.(ATLAS Collaboration), Phys. Lett. B710, 49 (2012).
- [2] S. Chatrchyan et al.(CMS Collaboration), Phys. Lett.B710, 26 (2012).
- [3] A. H. Chamseddine, R. L. Arnowitt and P. Nath, Phys. Rev. Lett. **49**, 970 (1982);  
H. P. Nilles, Phys. Lett. B **115**, 193 (1982); L. E. Ibanez, Phys. Lett. B **118**, 73 (1982);  
R. Barbieri, S. Ferrara and C. A. Savoy, Phys. Lett. B **119**, 343 (1982); H. P. Nilles,  
M. Srednicki and D. Wyler, Phys. Lett. B **120**, 346 (1983); J. R. Ellis, D. V. Nanopoulos and  
K. Tamvakis, Phys. Lett. B **121**, 123 (1983); J. R. Ellis, J. S. Hagelin, D. V. Nanopoulos and  
K. Tamvakis, Phys. Lett. B **125**, 275 (1983); N. Ohta, Prog. Theor. Phys. 70 (1983) 542;  
L. J. Hall, J. D. Lykken and S. Weinberg, Phys. Rev. D **27**, 2359 (1983); Fei Wang, Kun  
Wang, Jin Min Yang, Jingya Zhu, JHEP12(2018)041;  
Fei Wang, Wenyu Wang, Jin Min Yang, JHEP 06 (2015) 079;  
Fei Wang, Wenyu Wang, Jin Min Yang, JHEP03(2015)050;  
Kun Wang, Fei Wang, Jingya Zhu, Quanlin Jie, Chinese Physics C, 2018, 42(10): 103109.
- [4] M. Dine, W. Fischler and M. Srednicki, Nucl. Phys. B **189**, 575 (1981);  
S. Dimopoulos and S. Raby, Nucl. Phys. B **192**, 353 (1981);  
M. Dine and W. Fischler, Phys. Lett. B **110**, 227 (1982);  
M. Dine and A. E. Nelson, Phys. Rev. **D48**, 1277 (1993);  
M. Dine, A. E. Nelson and Y. Shirman, Phys. Rev. **D51**, 1362 (1995);  
M. Dine, A. E. Nelson, Y. Nir and Y. Shirman, Phys. Rev. **D53**, 2658 (1996);  
G. F. Giudice and R. Rattazzi, Phys. Rept. **322**, 419 (1999).
- [5] L. Randall and R. Sundrum, Nucl. Phys. B **557**, 79 (1999); G. F. Giudice, M. A. Luty,  
H. Murayama and R. Rattazzi, JHEP **9812**, 027 (1998).
- [6] C. Cheung, A. L. Fitzpatrick and D. Shih, JHEP **0807**, 054 (2008)  
doi:10.1088/1126-6708/2008/07/054 [arXiv:0710.3585 [hep-ph]].
- [7] T. T. Dumitrescu, Z. Komargodski, N. Seiberg and D. Shih, JHEP **1005**, 096 (2010)  
doi:10.1007/JHEP05(2010)096 [arXiv:1003.2661 [hep-ph]].
- [8] I. Jack and D. R. T. Jones, Phys. Lett. B 482, 167 (2000);  
E. Katz, Y. Shadmi and Y. Shirman, JHEP 9908, 015 (1999);  
N. ArkaniHamed, D. E. Kaplan, H. Murayama and Y. Nomura, JHEP 0102, 041 (2001);  
R. Sundrum, Phys. Rev. D 71, 085003 (2005);  
K. Hsieh and M. A. Luty, JHEP 0706, 062 (2007);  
Y. Cai and M. A. Luty, JHEP 1012, 037 (2010);  
T. Kobayashi, Y. Nakai and M. Sakai, JHEP 1106, 039 (2011);  
Fei Wang, Lei Wu, Jin Min Yang, Mengchao Zhang, Phys.Lett.B759 (2016)191;  
F. Wang, Phys. Lett.B 751(2015)402;  
Fei Wang, Jin Min Yang, Yang Zhang, JHEP04(2016)177;  
Fei Wang, Wenyu Wang, Jin Min Yang, Phys. Rev. D 96, 075025 (2017);  
X. Ning, F. Wang, JHEP08(2017)089.
- [9] A. Pomarol and R. Rattazzi, JHEP 9905, 013 (1999);  
R. Rattazzi, A. Strumia, James D. Wells, Nucl.Phys.B576:3-28(2000).
- [10] Nobuchika Okada, Phys.Rev. D65 (2002) 115009;  
Nobuchika Okada, Hieu Minh Tran, Phys.Rev. D87 (2013) 3, 035024;

- [11] Fei Wang, Wenyu Wang, Jin Min Yang, Yang Zhang, JHEP07(2015)138;  
Li, Z., Liu, GL., Wang, F. et al. Sci. China Phys. Mech. Astron. (2018) 61: 91011;  
Guo-Li Liu, Fei Wang, Wenyu Wang, Jin Min Yang, Chinese Physics C, Vol. 42, No. 3 (2018)  
035101;
- [12] I. Jack, D.R.T. Jones, Phys.Lett. B465 (1999) 148-154.
- [13] A. E. Nelson and N. J. Weiner, hep-ph/0210288.
- [14] K. Hsieh and M. A. Luty, JHEP 0706, 062 (2007).
- [15] Y. Cai and M. A. Luty, JHEP 1012, 037 (2010).
- [16] Xiaokang Du, Fei Wang, Eur. Phys. J. C (2018) 78:431.
- [17] G. F. Giudice, R. Rattazzi, Nucl. Phys. B511, 25 (1998).
- [18] Fei Wang, JHEP 1811 (2018) 062;  
Zhuang Li, Fei Wang, JHEP05(2019)009;  
Xiao Kang Du, Guo-Li Liu, Fei Wang, Wenyu Wang, Jin Min Yang, Yang Zhang, Eur. Phys.  
J. C (2019) 79: 397.
- [19] K. Abe et al. [Super-Kamiokande Collaboration], Phys. Rev. D95, 012004 (2017)  
[arXiv:1610.03597];  
S. Mine for the Super Kamiokande Collaboration, J. Phys. Conf. Ser. 718, 062044 (2016).
- [20] Y. Aoki, et al., RBC-UKQCD Collaboration, Phys. Rev. D 78 (2008) 054505.
- [21] J.Hisano, D. Kobayashi, N. Nagata, Phys. Lett. B 716(2012)406.
- [22] L.F. Abbott, M.B. Wise, Phys. Rev. D 22 (1980) 2208;  
C. Munoz, Phys. Lett. B 177 (1986) 55.