

PREPARED FOR SUBMISSION TO JHEP

Gravity with antisymmetric components

Chrysoula Markou, Felix J. Rudolph & Angnis Schmidt-May

*Max-Planck-Institut für Physik (Werner-Heisenberg-Institut)
Föhringer Ring 6, 80805 Munich, Germany*

E-mail: cmarkou@mpp.mpg.de, frudolph@mpp.mpg.de, angnisss@mpp.mpg.de

ABSTRACT: This work proposes a new gravitational theory formulated in terms of the vierbein field. The vierbein contains components which can be shifted by local Lorentz transformations and therefore do not show up in the spacetime metric. These components are given dynamics and become physical in our setup. They enter the massless theory in the form of an antisymmetric tensor field which makes the action reminiscent of the bosonic sector of supergravity. We then demonstrate that both the metric and the antisymmetric tensor can be made massive by adding a potential term for the vierbein. The form of this mass potential is inspired by ghost-free massive gravity. We confirm the absence of additional and potentially pathological degrees of freedom in an ADM analysis.

Contents

1	Introduction	1
1.1	Motivation	1
1.2	The standard massless theory for $G_{\mu\nu}$ and $B_{\mu\nu}$	4
2	A new vierbein formulation	5
2.1	Equations of motion	6
2.2	Local symmetries	7
3	The massive theory	9
3.1	Massive gravity in vierbein formulation	9
3.2	The vierbein action with mass term	10
3.3	Linear theory	11
4	ADM analysis	13
5	Discussion	15
A	Independence of $G_{\mu\nu}$ and $B_{\mu\nu}$	16
B	Details of the ADM analysis	18
B.1	Standard massive theory for $B_{\mu\nu}$	18
B.2	Action with new mass term	19
B.2.1	Einstein-Hilbert term	19
B.2.2	Mass term	19
B.2.3	Kinetic term for $B_{\mu\nu}$	20
B.2.4	Full action in ADM variables	21

1 Introduction

1.1 Motivation

General relativity is the standard theory for gravitational interactions. It possesses a mathematically well-defined interpretation in terms of Riemannian geometry and describes the dynamics of a metric $G_{\mu\nu}$ on a Riemannian manifold \mathcal{M} . The theory is well-tested over a large range of different energies and, from an experimental or observational point of view, there may be little motivation to modify or extend its unique and exceptionally beautiful structure. On the other hand, general relativity is not a fundamental quantum field theory

because it is not renormalizable. It can be treated as an effective field theory valid at low energies which requires an ultraviolet completion at the Planck scale the latest. A well-studied and promising candidate for the quantum theory of gravity at high energies is string theory, in which the notion of a point particle is replaced by a one-dimensional string.

The spacetime metric $G_{\mu\nu}$ arises as part of the massless bosonic excitations of the closed string, which in addition contain the scalar dilaton ϕ and the antisymmetric tensor $B_{\mu\nu}$, also known as the Kalb-Ramond field [1]. At low energies, string theory reduces to supergravity in 10 spacetime dimensions whose bosonic field content is precisely given by the massless fields $G_{\mu\nu}$, $B_{\mu\nu}$ and ϕ . For many purposes, one can consistently ignore the effects of the dilaton and work in an approximation where it is constant.¹ We therefore restrict our discussion in the following to the tensor fields.

The geometry of quantum gravity is not known, but string theory can provide us with hints on its nature. For instance, strings are subject to duality symmetries which do not have an obvious interpretation in standard field theory but are crucial for quantum consistency. Some aspects of these dualities can be captured using the mathematical concept of *generalized geometry* [3, 4]. Attempts to incorporate the T-duality symmetry of closed strings into 10-dimensional supergravity inspired the creation of *double field theory (DFT)* [5–7] (see also the reviews [8–10]). DFT is formulated on a doubled space obtained by augmenting the coordinates of regular supergravity with additional spatial coordinates associated with the winding modes of the string. The metric $G_{\mu\nu}$ and Kalb-Ramond field $B_{\mu\nu}$ are combined into a single symmetric tensor corresponding to a “generalized metric” of the doubled space. Thus $G_{\mu\nu}$ and $B_{\mu\nu}$ are treated on an equal footing and are both seen as part of the geometry. For an extensive analysis of the mathematical structure of the doubled geometry and its relation to generalized geometry see for example [11, 12].

String theory also contains other extended objects such as D-branes whose dynamics can be described by the Dirac-Born-Infeld action [13, 14]. This action is another example of containing the massless fields in the form of a single tensor, namely the linear combination $G_{\mu\nu} + B_{\mu\nu}$.

In this work we propose a new way to combine the degrees of freedom of the fields $G_{\mu\nu}$ and $B_{\mu\nu}$ into a single object with a well-known geometrical interpretation. This object is taken to be the vierbein field $e^a{}_\mu$ defined via $G_{\mu\nu} = e^a{}_\mu \eta_{ab} e^b{}_\nu$. Since the metric is invariant under local Lorentz transformations of the vierbein, $e^a{}_\mu \mapsto \Lambda^a{}_b e^b{}_\mu$ with $\Lambda^a{}_b \eta_{ac} \Lambda^c{}_d = \eta_{bd}$, $G_{\mu\nu}$ depends only on 10 of the 16 components in $e^a{}_\mu$.² In general relativity, the remaining 6 vierbein components are unphysical. Our idea is to make them dynamical and identify them with the 6 components of the antisymmetric tensor $B_{\mu\nu}$. This nontrivial way of re-packaging the field components into a single matrix is not directly inspired by string theory but rather follows a bottom-up approach, starting from the simple setup of general relativity coupled to

¹For example, one can assume that the value of the dilaton has been stabilized by means of a flux compactification (for a comprehensive review of the latter see [2]).

²For concreteness, we work in $D = 4$ spacetime dimensions throughout this paper, but our results straightforwardly generalize to arbitrary D , where $e^a{}_\mu$ contains D^2 and $G_{\mu\nu}$ contains $D(D + 1)/2$ components.

an antisymmetric tensor $B_{\mu\nu}$.

In fact, there exists already a well-studied set of theories for which the additional Lorentz components in the vierbein are non-dynamical but not unphysical, at least not initially. These theories are the vierbein formulations of *nonlinear massive gravity* [15], *bimetric gravity* [16] and *multimetric gravity* [17] (for reviews see [18–20]). In these setups, the 6 extra vierbein components are determined by solving a set of constraint equations. Generically, these constraints eventually require the additional degrees of freedom to vanish. This feature is crucial for the absence of ghost instabilities in these theories [21] and therefore the extra components cannot be made to remain physical in a straightforward way. The idea to make the additional Lorentz components dynamical in a way that avoids the ghost was already mentioned in Ref. [22] where, however, the suggested parametrization was slightly different from our setup.

A discussion of a dynamical antisymmetric part of the vierbein fluctuation in the context of linearized teleparallel theories can be found in section 4.6 of the encyclopaedic book by Ortín [23]. It is also worth mentioning that a previous attempt to extend general relativity by additional antisymmetric components was made by introducing Hermitian metrics [24–27]. For more work on gravity in the presence of an independent $B_{\mu\nu}$ field, see e.g. Ref. [28–32].

Summary of results

We extract the additional components of the vierbein that do not appear in the metric $G_{\mu\nu} = e^a{}_\mu \eta_{ab} e^b{}_\nu$ by introducing a second (non-dynamical) vierbein $\tilde{e}^a{}_\mu$ and defining $B_{\mu\nu}$ as the antisymmetric part of the matrix $e^a{}_\mu \eta_{ab} \tilde{e}^b{}_\nu$. These definitions are inserted into the action for the massless fields, which contains an Einstein-Hilbert term for the metric and a standard quadratic kinetic term for the antisymmetric tensor. We demonstrate that the formulation of the massless theory in terms of the vierbein field is entirely equivalent to the tensor formulation with independent $G_{\mu\nu}$ and $B_{\mu\nu}$.

We then extend the massless action by a set of vierbein interactions whose structure is motivated by massive gravity. As a highly nontrivial result, we find that these interactions make both fields $G_{\mu\nu}$ and $B_{\mu\nu}$ massive without introducing any further degrees of freedom into the theory (which usually are pathological). This is verified both at the linear level where the fields decouple and in a fully nonlinear ADM analysis.

The resulting vierbein theory thus yields an entirely new set of ghost-free gravitational interactions, which can both be seen as an extension of massive gravity and as an extension of the massless action for the supergravity fields $G_{\mu\nu}$ and $B_{\mu\nu}$.

Conventions

Throughout this paper we will use the following conventions. We are working in 4 dimensions and the signature of the spacetime metric is $(-, +, +, +)$. The spacetime indices are denoted by Greek letters μ, ν whereas for the Lorentz indices we use Latin letters a, b . These two types of indices are related via the vierbein $e^a{}_\mu$ and its inverse, $e^\mu{}_a$. Indices are raised and lowered by $G_{\mu\nu}$ and η_{ab} and their inverses respectively. Brackets to denote symmetrization and

antisymmetrization over indices are defined as $T_{\mu\nu} = T_{(\mu\nu)} + T_{[\mu\nu]}$ with $T_{(\mu\nu)} = \frac{1}{2}(T_{\mu\nu} + T_{\nu\mu})$ and $T_{[\mu\nu]} = \frac{1}{2}(T_{\mu\nu} - T_{\nu\mu})$.

1.2 The standard massless theory for $G_{\mu\nu}$ and $B_{\mu\nu}$

Our starting point will be the Einstein-Hilbert action for a metric $G_{\mu\nu}$ in the presence of an antisymmetric tensor field $B_{\mu\nu}$, whose basic properties will be briefly reviewed in the following. The action is,

$$S_{GB} = m_{\text{P}}^2 \int d^4x \sqrt{G} [R(G) - 2\Lambda] - \frac{1}{2 \cdot 3!} \int d^4x \sqrt{G} H_{\mu\nu\rho} H^{\mu\nu\rho}, \quad (1.1)$$

where $H_{\mu\nu\rho} = 3\nabla_{[\mu} B_{\nu\rho]}$ are the components of the 3-form field strength. They enter the action in the kinetic term for the massless field $B_{\mu\nu}$ in curved space, i.e. all indices are raised with $G^{\mu\nu}$. The Einstein-Hilbert action including the cosmological constant Λ is the standard nonlinear kinetic term for a massless spin-2 field.

The equations of motion obtained by varying the action with respect to the tensor fields $G_{\mu\nu}$ and $B_{\mu\nu}$ are,

$$\mathcal{G}^{\mu\nu} \equiv R^{\mu\nu} - \frac{1}{2}(R - 2\Lambda)G^{\mu\nu} - \frac{1}{4m_{\text{P}}^2}(H^{\mu\rho\sigma} H^{\nu}{}_{\rho\sigma} - \frac{1}{6}H^2 G^{\mu\nu}) = 0, \quad (1.2a)$$

$$\mathcal{B}^{\mu\nu} \equiv -\frac{1}{2m_{\text{P}}^2} \nabla_{\rho} H^{\rho\mu\nu} = 0. \quad (1.2b)$$

Note that $\mathcal{G}_{\mu\nu}$ consists of the usual Einstein tensor $R_{\mu\nu} - \frac{1}{2}G_{\mu\nu}R$ and the contribution from the ‘‘matter source’’ including $B_{\mu\nu}$ through its field strength. Also, $\mathcal{G}_{\mu\nu}$ is symmetric while $\mathcal{B}_{\mu\nu}$ is antisymmetric in its two indices.

Interpretation in terms of torsion. The Ricci scalar in the action S_{GB} in (1.1) is constructed from the torsionless Levi-Civita connection Γ of the metric $G_{\mu\nu}$, whose components satisfy $\Gamma_{[\mu\nu]}{}^{\rho} = 0$. It is possible to write the entire action in Einstein-Hilbert form with a different metric-compatible connection $\hat{\Gamma}$ which is torsionful, i.e. $\hat{\Gamma}_{[\mu\nu]}{}^{\rho} \neq 0$. The precise form is,

$$S_{GB} = m_{\text{P}}^2 \int d^4x \sqrt{G} [R(\hat{\Gamma}) - 2\Lambda], \quad \hat{\Gamma}_{\mu\nu}{}^{\rho} = \Gamma_{\mu\nu}{}^{\rho} + \frac{1}{\sqrt{12} m_{\text{P}}} H_{\mu\nu}{}^{\rho}, \quad (1.3)$$

where the Ricci scalar is now constructed from $\hat{\Gamma}$ and the contorsion of the new connection is given precisely by the 3-form field strength of $B_{\mu\nu}$. From this point of view, the action S_{GB} is a particular version of general relativity with propagating torsion. An interpretation of this setup in the context of supergravity and string theory can be found in Ref. [33].

Local symmetries. The action in (1.1) has several local symmetries (corresponding to gauge redundancies).

- (i) Diffeomorphisms which infinitesimally transform the coordinates as $\Delta_{\xi} x^{\mu} = \xi^{\mu}$, and the fields as,

$$\Delta_{\xi} G_{\mu\nu} = \nabla_{\mu} \xi_{\nu} + \nabla_{\nu} \xi_{\mu}, \quad \Delta_{\xi} B_{\mu\nu} = \xi^{\rho} \nabla_{\rho} B_{\mu\nu} + B_{\rho\nu} \nabla_{\mu} \xi^{\rho} - B_{\rho\mu} \nabla_{\nu} \xi^{\rho}. \quad (1.4)$$

It is well-known that these remove 4+4 degrees of freedom, leaving 2 propagating modes in $G_{\mu\nu}$. These correspond to the 2 helicity states of the massless spin-2 field.

- (ii) B -gauge transformations under which only the $B_{\mu\nu}$ field transforms,

$$\Delta_\lambda B_{\mu\nu} = \nabla_\mu \lambda_\nu - \nabla_\nu \lambda_\mu, \quad \Delta_\lambda G_{\mu\nu} = 0. \quad (1.5)$$

It can be shown that these remove 3 + 2 degrees of freedom, leaving 1 propagating mode in $B_{\mu\nu}$. This is of course consistent with the fact that a massless $B_{\mu\nu}$ field can be dualized to a scalar in $D = 4$.

- (iii) Local Lorentz invariance, i.e. $\text{SO}(1,3)$ which transforms neither $G_{\mu\nu}$ nor $B_{\mu\nu}$ and therefore is not explicit in the tensor formulation. We see it only when we introduce a vierbein $e^a{}_\mu$ for the metric, $G_{\mu\nu} = e^a{}_\mu \eta_{ab} e^b{}_\nu$. Then the infinitesimal local Lorentz transformations (LLT) read,

$$\Delta_\omega e^a{}_\mu = \eta^{ab} \omega_{bc} e^c{}_\mu \quad \text{with} \quad \omega_{bc} = -\omega_{cb}, \quad \Delta_\omega B_{\mu\nu} = \Delta_\omega G_{\mu\nu} = 0. \quad (1.6)$$

These precisely remove the 6 redundant degrees of freedom contained in the vierbein which do not enter the metric $G_{\mu\nu}$.

As a side remark, we also note that the diffeomorphisms in (1.4) can be written as a transformation of the vierbein $e^a{}_\mu$ instead of the metric,

$$\Delta_\xi e^a{}_\mu = \xi^\rho \nabla_\rho e^a{}_\mu + e^a{}_\rho \nabla_\mu \xi^\rho, \quad \Delta_\xi B_{\mu\nu} = \xi^\rho \nabla_\rho B_{\mu\nu} + B_{\rho\nu} \nabla_\mu \xi^\rho - B_{\rho\mu} \nabla_\nu \xi^\rho. \quad (1.7)$$

The first term in the transformation of $e^a{}_\mu$ drops out from the transformation of the metric but is part of the proper transformation of a 1-form under diffeomorphisms.

2 A new vierbein formulation

Our aim is to package the $10 + 6 = 16$ degrees of freedom of $G_{\mu\nu}$ and $B_{\mu\nu}$ into a single 4×4 matrix and identify the latter with the vierbein field $e^a{}_\mu$. As before, the metric is given by,

$$G_{\mu\nu} = e_\mu{}^a \eta_{ab} e^b{}_\nu. \quad (2.1)$$

In GR, 6 of the 16 components in $e^a{}_\mu$ do not show up in $G_{\mu\nu}$ and are thus unphysical due to local Lorentz invariance. Here we want them to remain physical and define the components of the field $B_{\mu\nu}$. This means that we cannot simply contract the Lorentz index of $e^a{}_\mu$ with $\text{SO}(1,3)$ invariant tensors such as η_{ab} or ϵ_{abcd} everywhere in the action. We are forced to introduce a new object that carries a Lorentz index. Moreover, we need to convert the Lorentz index of $e^a{}_\mu$ into a coordinate index in order to translate its components into those of the antisymmetric tensor. Therefore we choose the additional object to be a second vierbein $\tilde{e}^a{}_\mu$ which, when all its components are specified, introduces a fixed Lorentz frame. For instance, as a matrix, $\tilde{e}^a{}_\mu$ could just be the identity $\delta^a{}_\mu$.³

³The reader familiar with the topic will notice that this is the exact same situation as in massive gravity.

Clearly there are infinitely many possibilities to build the tensor $B_{\mu\nu}$ out of the vierbeine $e^a{}_\mu$ and $\tilde{e}^a{}_\mu$ but here we restrict to expressions that are quadratic in the vierbeine. Then the only antisymmetric rank-2 tensor without any free Lorentz indices is,

$$B_{\mu\nu} \equiv e^a{}_\mu \eta_{ab} \tilde{e}^b{}_\nu - \tilde{e}^a{}_\mu \eta_{ab} e^b{}_\nu. \quad (2.2)$$

Since the definitions of the tensors cannot be directly inverted in a covariant way, it is not clear how exactly the 16 components split into the 10 components of $G_{\mu\nu}$ and 6 components of $B_{\mu\nu}$. Thus, a priori, it seems that now $G_{\mu\nu}$ and $B_{\mu\nu}$ are no longer independent objects.

The next step is to insert the vierbein expressions for $G_{\mu\nu}$ and $B_{\mu\nu}$ into the action (1.1).⁴ It is not clear that this will give a description equivalent to the tensor formulation with independent $G_{\mu\nu}$ and $B_{\mu\nu}$. However, as we will show now, the equations of motion for the vierbein guarantee the equivalence of the two formulations.

2.1 Equations of motion

Using the chain rule it is easy to vary the action (1.1) with $G_{\mu\nu}$ and $B_{\mu\nu}$ given by (2.1) and (2.2) with respect to $e^a{}_\mu$. In this way we obtain the equation of motion for the vierbein,

$$\mathcal{E}_a{}^\mu \equiv \frac{\delta S}{\delta e^a{}_\mu} = \frac{\delta G^{\rho\sigma}}{\delta e^a{}_\mu} \frac{\delta S_{GB}}{\delta G^{\rho\sigma}} \Big|_{B=\text{const.}} + \frac{\delta B_{\rho\sigma}}{\delta e^a{}_\mu} \frac{\delta S_B}{\delta B_{\rho\sigma}} \Big|_{G=\text{const.}} = 0, \quad (2.3)$$

where the variations of the tensors with respect to the vierbein are obtained from,

$$\delta G_{\mu\nu} = \delta e_\mu{}^a \eta_{ab} e^b{}_\nu + e_\mu{}^a \eta_{ab} \delta e^b{}_\nu \quad \delta B_{\mu\nu} = \delta e_\mu{}^a \eta_{ab} \tilde{e}^b{}_\nu - \tilde{e}_\mu{}^a \eta_{ab} \delta e^b{}_\nu. \quad (2.4)$$

We then simply use the results for the tensor variations in (1.2) to see that the equations of motion for the vierbein take the following form,

$$\begin{aligned} \mathcal{E}_a{}^\mu &= 2\eta_{ab} e_\nu{}^b \left[R^{\mu\nu} - \frac{1}{2}(R - 2\Lambda)G^{\mu\nu} - \frac{1}{4m_{\text{P}}^2}(H^{\mu\rho\sigma}H^\nu{}_{\rho\sigma} - \frac{1}{6}H^2G^{\mu\nu}) \right] - \frac{1}{m_{\text{P}}^2}\eta_{ab}\tilde{e}_\nu{}^b \nabla_\rho H^{\rho\mu\nu} \\ &= 0. \end{aligned} \quad (2.5)$$

In terms of the tensors defined in (1.2), we can write these as,

$$\mathcal{E}_a{}^\mu = 2\eta_{ab} e_\nu{}^b \mathcal{G}^{\mu\nu} + 2\eta_{ab} \tilde{e}_\nu{}^b \mathcal{B}^{\mu\nu} = 0. \quad (2.6)$$

A priori, it is now not obvious that $\mathcal{G}^{\mu\nu}$ and $\mathcal{B}^{\mu\nu}$ should vanish separately as they do in the tensor formulation of the theory.

⁴Since, in our parametrization of the action, $B_{\mu\nu}$ has mass dimension 1 while $G_{\mu\nu}$ (and thus $e^a{}_\mu$) have mass dimension 0, we take $\tilde{e}^a{}_\mu$ to have mass dimension 1. These conventions can of course easily be changed by performing a rescaling of the vierbeine.

Separating the equations. Consider the following antisymmetrized linear combination of the vierbein equations,

$$2\eta^{ab}e^{[\mu}{}_a\frac{\delta S}{\delta e^b{}_{\nu]}} = 2\eta^{ab}e^{[\mu}{}_a\mathcal{E}_b^{\nu]} = 0. \quad (2.7)$$

These correspond to a subset of 6 out of the 16 equations. Inserting \mathcal{E}_a^μ from above and using the symmetry of $\mathcal{G}_{\mu\nu}$, they reduce to,

$$4\tilde{e}^a{}_\rho e^{[\mu}{}_a\mathcal{B}^{\nu]\rho} = 4e^{[\mu}{}_a\tilde{e}^{\nu]}{}_b\tilde{e}^b{}_\sigma\tilde{e}^a{}_\rho\mathcal{B}^{\sigma\rho} = -2\tilde{P}^{\mu\nu}{}_{ab}\tilde{e}^a{}_\rho\tilde{e}^b{}_\sigma\mathcal{B}^{\rho\sigma} = 0 \quad (2.8)$$

where we have defined the operator $\tilde{P}^{\mu\nu}{}_{ab} \equiv 2e^{[\mu}{}_a\tilde{e}^{\nu]}{}_b$. As we will argue now, this operator is generically invertible, since on the space of antisymmetric matrices it acts just like a linear map. One way to see this is to introduce a combined index $M = [\mu\nu]$ and $A = [ab]$ for antisymmetrized pairs of normal indices. Then \tilde{P} is like a matrix acting on a vector,

$$\tilde{P}_A^M\mathcal{B}^A = 0. \quad (2.9)$$

Under the restriction to the space of antisymmetric matrices, \tilde{P}_A^M can thus generically be inverted. Then the subset of equations in (2.7) reduces to $\mathcal{B}^{\mu\nu} = 0$, which are the equations for $B_{\mu\nu}$ in the tensor formulation. Upon inserting this back into the full equations (2.6), they reduce to $\mathcal{G}_{\mu\nu} = 0$, which is the equation for $G_{\mu\nu}$ in the tensor formulation. This shows how the $G_{\mu\nu}$ and $B_{\mu\nu}$ equations separate and our vierbein formulation is equivalent to the tensor formulation, provided that the operator $\tilde{P}^{\mu\nu}{}_{ab}$ is invertible. More details on the independence of $G_{\mu\nu}$ and $B_{\mu\nu}$ and the relation to the invertibility of the operator can be found in appendix A.

2.2 Local symmetries

We have explicitly broken local Lorentz invariance by making all 16 components in the vierbein physical.⁵ The remaining local symmetries of the action (1.1) that we discussed in section 1.2 must be recovered in the vierbein formulation. However, it is not immediately obvious how the diffeomorphisms and gauge transformations can be written as a transformation of the vierbein $e^a{}_\mu$ which now appears in both tensor fields. In the following we discuss how exactly the symmetries emerge in this case.

Diffeomorphisms. Let us start from the diffeomorphisms transformation of the metric, $\Delta_\xi G_{\mu\nu} = \nabla_\mu\xi_\nu + \nabla_\nu\xi_\mu$. As we already saw in (1.7), this can be produced by the following transformation of the vierbein,

$$\Delta_\xi e^a{}_\mu = \xi^\rho\nabla_\rho e^a{}_\mu + e^a{}_\rho\nabla_\mu\xi^\rho + \eta^{ac}\omega_{cb}(\xi)e^b{}_\mu. \quad (2.10)$$

⁵In fact, local Lorentz invariance can be restored in two ways. As shown in the previous subsection, $G_{\mu\nu}$ and $B_{\mu\nu}$ become independent dynamically. We can hence express $G_{\mu\nu}$ in terms of a new vierbein whose components do not depend on $B_{\mu\nu}$ and whose Lorentz rotations thus leave the action invariant. The other option is to make $\tilde{e}^a{}_\mu$ dynamical, in which case simultaneous Lorentz rotations of the two vierbeine leave the action invariant. This latter case is particularly interesting and will be discussed in a future publication.

Here we have allowed for a LLT with $\omega_{ab} = -\omega_{ba}$ depending on the diffeo parameter ξ_μ , since this leaves the metric $G_{\mu\nu}$ invariant. Requiring the correct transformation behaviour for $B_{\mu\nu}$ will fix the form of $\omega_{ab}(\xi)$. Namely, under diffeomorphisms, the 2-form should transform as,

$$\begin{aligned}\Delta_\xi B_{\mu\nu} &= \xi^\rho \nabla_\rho B_{\mu\nu} + B_{\rho\nu} \nabla_\mu \xi^\rho - B_{\rho\mu} \nabla_\nu \xi^\rho \\ &= \xi^\rho \nabla_\rho B_{\mu\nu} + 2\eta_{ab} (e^a_{[\rho} \tilde{e}^b_{\nu]} \nabla_\mu \xi^\rho - e^a_{[\rho} \tilde{e}^b_{\mu]} \nabla_\nu \xi^\rho).\end{aligned}\quad (2.11)$$

On the other hand, inserting (2.10) into the definition of $B_{\mu\nu}$, we obtain,

$$\begin{aligned}\Delta_\xi B_{\mu\nu} &= 2\Delta_\xi e^a_{[\mu} \eta_{ab} \tilde{e}^b_{\nu]} = 2\eta_{ab} \xi^\rho \nabla_\rho e^a_{[\mu} \tilde{e}^b_{\nu]} + 2\eta_{ab} e^a_{\rho} \tilde{e}^b_{[\nu} \nabla_{\mu]} \xi^\rho + 2e^a_{[\mu} \tilde{e}^b_{\nu]} \omega_{ba}(\xi) \\ &= \xi^\rho \nabla_\rho B_{\mu\nu} - 2\eta_{ab} \xi^\rho \nabla_\rho \tilde{e}^b_{[\nu} e^a_{\mu]} + \eta_{ab} e^a_{\rho} (\tilde{e}^b_{\nu} \nabla_\mu \xi^\rho - \tilde{e}^b_{\mu} \nabla_\nu \xi^\rho) - 2e^a_{[\mu} \tilde{e}^b_{\nu]} \omega_{ab}(\xi).\end{aligned}\quad (2.12)$$

Comparing (2.12) to (2.11), we see that we must have,

$$-2\eta_{ab} \xi^\rho \nabla_\rho \tilde{e}^b_{[\nu} e^a_{\mu]} + \eta_{ab} \tilde{e}^b_{\rho} (e^a_{\nu} \nabla_\mu \xi^\rho - e^a_{\mu} \nabla_\nu \xi^\rho) = 2e^{[a}_{[\mu} \tilde{e}^b_{\nu]} \omega_{ab}(\xi) \equiv P_{\mu\nu}{}^{ab} \omega_{ab}(\xi) \quad (2.13)$$

The object $P_{\mu\nu}{}^{ab} = 2e^{[a}_{[\mu} \tilde{e}^b_{\nu]}$ is an operator similar to $\tilde{P}^{\mu\nu}{}_{ab}$ which we encountered earlier in eq. (2.8). For the same reason as before, $P_{\mu\nu}{}^{ab}$ is generically invertible on the space of antisymmetric matrices. We thus find the local Lorentz transformation that is needed to generate the diffeomorphism of $B_{\mu\nu}$,

$$\omega_{ab}(\xi) = 2(P^{-1})_{ab}{}^{\mu\nu} \left(\xi^\rho \nabla_\rho \tilde{e}^c_{\mu} + \tilde{e}^c_{\rho} \nabla_\mu \xi^\rho \right) \eta_{cd} e^d_{\nu}, \quad (2.14)$$

and the diffeomorphisms are generated by the following transformation of the vierbein,

$$\Delta_\xi e^a_{\mu} = \xi^\rho \nabla_\rho e^a_{\mu} + e^a_{\rho} \nabla_\mu \xi^\rho + 2\eta^{ac} \eta_{de} e^b_{\mu} e^e_{\nu} (P^{-1})^{cb}{}^{\sigma\nu} \left(\xi^\rho \nabla_\rho \tilde{e}^d_{\sigma} + \tilde{e}^d_{\rho} \nabla_\sigma \xi^\rho \right). \quad (2.15)$$

Gauge transformations. We would also like to obtain the gauge transformation of the form $\Delta_\lambda B = d\lambda$ from a transformation of Δe^a_{μ} . Since such a gauge transformation should leave $G_{\mu\nu}$ invariant, it must be recast as a local Lorentz transformation. At the linearized level it is thus given by,

$$\Delta_\lambda e^a_{\mu} = \omega^a_b(\lambda) e^b_{\mu}, \quad \omega_{ab}(\lambda) = -\omega_{ba}(\lambda). \quad (2.16)$$

The form of $\omega_{ab}(\lambda)$ is now again determined by the correct transformation behaviour of $B_{\mu\nu}$. Namely, the gauge transformation of the B -field gives,

$$\Delta_\lambda B_{\mu\nu} = 2\nabla_{[\mu} \lambda_{\nu]} = 2e_{[\mu}{}^{[a} \tilde{e}_{\nu]}{}^{b]} \omega_{ab}(\lambda) \equiv P_{\mu\nu}{}^{ab} \omega_{ab}(\lambda). \quad (2.17)$$

The object $P_{\mu\nu}{}^{ab}$ is the same operator as in equation (2.13). We can again invert it and find the local Lorentz transformation corresponding to a given gauge transformation of $B_{\mu\nu}$,

$$\omega_{ab}(\lambda) = (P^{-1})_{ab}{}^{\mu\nu} \Delta B_{\mu\nu} = 2(P^{-1})_{ab}{}^{\mu\nu} \nabla_{[\mu} \lambda_{\nu]}. \quad (2.18)$$

The transformation on the vierbein generating B -gauge transformations is thus,

$$\Delta_\lambda e^a{}_\mu = 2\eta^{ac}(P^{-1})_{cb}\rho^\sigma\nabla_{[\rho}\lambda_{\sigma]}e^b{}_\mu. \quad (2.19)$$

Since the equations separate and the gauge symmetries can be reproduced, we conclude that the vierbein formulation for the massless theory is entirely equivalent to the tensor formulation in section 1.2.

3 The massive theory

We would like to add interactions to the massless theory which render the fields massive. In the tensor formulation with independent $G_{\mu\nu}$ and $B_{\mu\nu}$, one option would be to include the standard mass term for the antisymmetric tensor which is of the form,

$$-m^2 \int d^4x \sqrt{G} B_{\mu\nu} B_{\rho\sigma} G^{\mu\rho} G^{\nu\sigma}. \quad (3.1)$$

This gives a consistent theory, describing a massive antisymmetric tensor in curved background given by a metric which remains massless. We perform a 3+1 split and degree of freedom counting for the corresponding action in appendix B.1. A massive $B_{\mu\nu}$ field has 3 propagating degrees of freedom, which is consistent with the fact that in $D = 4$ it is dual to a massive vector (while a massless $B_{\mu\nu}$ is dual to a massless scalar) [34].

Here we are going to take a different approach and show that a mass term that is usually used for $G_{\mu\nu}$ can make both fields $G_{\mu\nu}$ and $B_{\mu\nu}$ massive. This is achieved precisely by our identification of the field $B_{\mu\nu}$ with the 6 additional components in the vierbein $e^a{}_\mu$. In section 4 we show explicitly that the resulting action has 5 + 3 propagating modes, as expected for a massive $G_{\mu\nu}$ and a massive $B_{\mu\nu}$. No additional (potentially ghost-like) degrees of freedom are introduced.⁶

In order to set the stage, we begin by reviewing the mass term for the metric $G_{\mu\nu}$ which usually does not contain the $B_{\mu\nu}$ field, since its consistency requires $B_{\mu\nu} = 0$.

3.1 Massive gravity in vierbein formulation

Massive gravity extends the Einstein-Hilbert term of GR by a mass potential which assumes a rather simple form when written in terms of the vierbein field $e^a{}_\mu$. The only nontrivial term that can be built from this vierbein alone is $\det e$, which corresponds to the gauge-invariant cosmological constant term and is not part of the mass giving potential. Writing down the mass term therefore requires introducing a second vierbein $\tilde{e}^a{}_\mu$. This field can be made dynamical in a consistent way but for now we focus on the case with non-dynamical $\tilde{e}^a{}_\mu$.

The ghost-free action for massive gravity then is,

$$S_{\text{MG}} = m_{\text{P}}^2 \int d^4x \sqrt{G} (R(G) - \Lambda) - m_{\text{P}}^2 m^2 \int \epsilon_{abcd} (b_1 e^a \wedge e^b \wedge e^c \wedge \tilde{e}^d + b_2 e^a \wedge e^b \wedge \tilde{e}^c \wedge \tilde{e}^d + b_3 e^a \wedge \tilde{e}^b \wedge \tilde{e}^c \wedge \tilde{e}^d), \quad (3.2)$$

⁶It turns out that it is then inconsistent to also add the standard mass term for $B_{\mu\nu}$ in (3.1) to this setup.

where the interaction parameters b_n are arbitrary constants. As advertised, the action contains an Einstein-Hilbert kinetic term for the metric $G_{\mu\nu} = e^a{}_\mu \eta_{ab} e^b{}_\nu$ and an interaction potential giving mass to the spin-2 field. The interactions are forced to be of the above totally antisymmetrized form in order to guarantee the absence of the Boulware-Deser ghost [35]. The Hamiltonian analysis for the action (3.2) has been performed in Ref. [17, 36, 37]. The mass term can be consistently added to (1.1) with independent tensor fields, in which case the spin-2 degrees of freedom become massive while $B_{\mu\nu}$ remains massless.

The equations of motion for the vierbein field $e^a{}_\mu$ following from (3.2) contain a set of constraints arising from the potential. These read [17],

$$\eta^{ab} e^{[\mu}{}_a \frac{\delta S_{\text{MG}}}{\delta e^b{}_{\nu]}} = 0. \quad (3.3)$$

The kinetic term does not contribute to this antisymmetric part of the equations due to its invariance under local Lorentz transformations, $\Delta e^a{}_\mu = \eta^{ab} \omega_{bc} e^c{}_\mu$ with $\omega_{bc} = -\omega_{cb}$. The constraints (3.3) have a unique covariant solution which reads,

$$e^a{}_\mu \eta_{ab} \tilde{e}^b{}_\nu - \tilde{e}^a{}_\mu \eta_{ab} e^b{}_\nu = 0. \quad (3.4)$$

These 6 constraints determine the extra components in $e^a{}_\mu$ which do not show up in $G_{\mu\nu}$. In fact, it is crucial that (3.4) holds because it ensures the absence of the Boulware-Deser ghost in the action [21]. It also allows us to express the vierbein potential in terms of the respective metrics $G_{\mu\nu}$ and $f_{\mu\nu} = \tilde{e}^a{}_\mu \eta_{ab} \tilde{e}^b{}_\nu$, in which case the interactions depend on the square-root matrix $\sqrt{G^{-1}f}$.

Massive gravity does therefore not allow configurations with $e^a{}_\mu \eta_{ab} \tilde{e}^b{}_\nu - \tilde{e}^a{}_\mu \eta_{ab} e^b{}_\nu \neq 0$.⁷ However, this result crucially depends on the fact that $B_{\mu\nu} \equiv e^a{}_\mu \eta_{ab} \tilde{e}^b{}_\nu - \tilde{e}^a{}_\mu \eta_{ab} e^b{}_\nu$ does not possess a kinetic term. In section 4 we will demonstrate that by giving dynamics to the antisymmetric components we can have $B_{\mu\nu} \neq 0$ without introducing the Boulware-Deser ghost.

3.2 The vierbein action with mass term

We now add the ghost-free mass term for the vierbein field $e^a{}_\mu$ in (3.2) to the action (1.1), identifying the non-dynamical vierbein $\tilde{e}^a{}_\mu$ in the antisymmetric tensor with the reference field in the mass term.⁸ The full action $S_{\text{m}} = S_{GB} + S_V$ reads,

$$\begin{aligned} S_{\text{m}} = & m_{\text{P}}^2 \int d^4x \sqrt{G} [R(G) - 2\Lambda] - \frac{1}{2 \cdot 3!} \int d^4x \sqrt{G} H_{\mu\nu\rho} H^{\mu\nu\rho} \\ & - m_{\text{P}}^2 m^2 \int \epsilon_{abcd} \left(b_1 e^a \wedge e^b \wedge e^c \wedge \tilde{e}^d + b_2 e^a \wedge e^b \wedge \tilde{e}^c \wedge \tilde{e}^d + b_3 e^a \wedge \tilde{e}^b \wedge \tilde{e}^c \wedge \tilde{e}^d \right), \end{aligned} \quad (3.5)$$

⁷In fact, the antisymmetric combination of vierbeine can be non-vanishing only in theories involving at least 3 vierbein fields and a very particular type of ghost-free coupling [38].

⁸Using two different reference fields would not change the conclusions below but seems like an unnecessary complication of our minimal setup.

with $G_{\mu\nu}$ and $B_{\mu\nu}$ defined in terms of the vierbein as in (2.1) and (2.2).⁹ The mass term breaks both the diffeomorphism and the gauge symmetry of the $B_{\mu\nu}$ field, which were present in the massless theory. We will show that the above action provides a nonlinear description of interacting massive $G_{\mu\nu}$ and massive $B_{\mu\nu}$.

In contrast to massive gravity, since now $B_{\mu\nu} \neq 0$, the interaction potential does not possess a formulation in terms of metrics. Nevertheless, it is still possible to separate the equations of motion into an antisymmetric and symmetric part, which contain exactly the same kinetic structures as in the massless theory and its equivalent tensor formulation.

Separating the equations. The mass term gives an extra contribution to the equation of motion of $e^a{}_\mu$, which we shall denote by $\mathcal{V}_a{}^\mu \equiv -(m_{\text{P}}^2 \det e)^{-1} \frac{\delta S_V}{\delta e^a{}_\mu}$. The full equations – extending (2.6) – then read,

$$\mathcal{E}_a{}^\mu = 2\eta_{ab}e^b{}_\nu \mathcal{G}^{\mu\nu} + 2\eta_{ab}\tilde{e}^b{}_\nu \mathcal{B}^{\mu\nu} + \mathcal{V}_a{}^\mu = 0, \quad (3.6)$$

where $\mathcal{G}^{\mu\nu}$ and $\mathcal{B}^{\mu\nu}$ are still the same as in (1.2). After multiplying by $\frac{1}{2}\eta^{ac}e_c{}^\rho$ and taking the antisymmetric part of the resulting equation, a few simple manipulations lead to,

$$\mathcal{B}^{\mu\nu} - (\tilde{P}^{-1})_{\rho\sigma}{}^{ab}\tilde{e}^\mu{}_a\tilde{e}^\nu{}_b e^\rho{}_c \eta^{cd}\mathcal{V}_d{}^\sigma = 0, \quad (3.7)$$

instead of the earlier result $\mathcal{B}^{\mu\nu} = 0$. Here, $\tilde{P}^{\mu\nu}{}_{ab} \equiv 2e^{[\mu}{}_{[a}\tilde{e}^{\nu]}{}_{b]}$ is the same invertible operator that we encountered in the massless case. Plugging (3.7) back into the full equation (3.6), we find,

$$\mathcal{G}^{\mu\nu} + (\tilde{P}^{-1})_{\rho\sigma}{}^{ab}\tilde{e}^\mu{}_a e^\nu{}_b e^\rho{}_c \eta^{cd}\mathcal{V}_d{}^\sigma + \frac{1}{2}e^\nu{}_a \eta^{ab}\mathcal{V}_b{}^\mu = 0. \quad (3.8)$$

By construction, the antisymmetric part of this expression identically vanishes¹⁰ and we are left with a symmetric equation for $\mathcal{G}^{\mu\nu}$. It captures the contribution from the interaction term to the equation of motion for the components in the metric $G_{\mu\nu}$. From the form of the equations (3.7) and (3.8) we conclude that the kinetic structures have not been altered with respect to the massless theory.

We also observe that the action (3.5) in fact contains ghost-free massive gravity in the form of a particular solution to (3.7). This solution corresponds to $B_{\mu\nu} = 0$, in which case (3.8) reduces to the equation of motion for the vierbein following from varying the massive gravity action in (3.2). This shows that our new theory can be viewed as a generalization of massive gravity.

3.3 Linear theory

In this section we study the spectrum of linear perturbations around the background $\bar{e}^a{}_\mu$ for which the dynamical vierbein is proportional to the reference frame, $\tilde{e}^a{}_\mu = c\bar{e}^a{}_\mu$ with constant

⁹Note that since in our parametrization, the vierbein $\tilde{e}^a{}_\mu$ has mass dimension 1, the b_n parameters have mass dimension $-n$.

¹⁰This is easy to verify directly using $(\tilde{P}^{-1})_{\rho\sigma}{}^{ab}e^{[\nu}{}_{[a}\tilde{e}^{\mu]}{}_{b]} = \frac{1}{2}(\tilde{P}^{-1})_{\rho\sigma}{}^{ab}\tilde{P}_{ab}{}^{\nu\mu} = \frac{1}{2}\delta^{[\nu}{}_{[\rho}\delta^{\mu]}{}_{\sigma]}$.

$c \neq 0$ of mass dimension 1. Clearly, for such a background we have that $\bar{B}_{\mu\nu} = 0$, which solves eq. (3.7). Plugging the proportional ansatz into (3.8), we find that $\bar{G}_{\mu\nu}$ is the metric of a maximally symmetric spacetime with $R_{\mu\nu}(\bar{G}) = \Lambda_G \bar{G}_{\mu\nu}$. The constant background curvature is given by the following combination of parameters,

$$\Lambda_G = \Lambda + 3m^2 \left(3b_1 c + 2b_2 c^2 + b_3 c^3 \right). \quad (3.9)$$

Next, we write the nonlinear vierbein in terms of this background and a small fluctuation,

$$e^a{}_\mu = \bar{e}^a{}_\mu + \delta e^a{}_\mu. \quad (3.10)$$

We split the fluctuation into a symmetric and an antisymmetric component as follows,

$$\eta_{ab} \bar{e}^b{}_\nu \delta e^a{}_\mu \equiv h_{\mu\nu} + b_{\mu\nu}, \quad (3.11)$$

where,

$$h_{\mu\nu} = h_{\nu\mu} = \eta_{ab} \bar{e}^b{}_{(\nu} \delta e^a{}_{\mu)}, \quad b_{\mu\nu} = -b_{\nu\mu} = \eta_{ab} \bar{e}^b{}_{[\nu} \delta e^a{}_{\mu]}. \quad (3.12)$$

Due to this definition, the fluctuation of the metric is $\delta G_{\mu\nu} = 2h_{\mu\nu}$ and the fluctuation of the B -field is $\delta B_{\mu\nu} = 2cb_{\mu\nu}$. By a tedious but straightforward computation one can show that the linearized version of the full equations of motion (3.6) can be separated and diagonalized into the two sets,

$$\mathcal{E}_{\mu\nu}{}^{\rho\sigma} h_{\rho\sigma} - \Lambda_G (h_{\mu\nu} - \frac{1}{2} h_{\rho\sigma} \bar{G}^{\rho\sigma} \bar{G}_{\mu\nu}) - \frac{m_{\text{FP}}^2}{2} (h_{\mu\nu} - h_{\rho\sigma} \bar{G}^{\rho\sigma} \bar{G}_{\mu\nu}) = 0, \quad (3.13a)$$

$$\bar{\nabla}^\rho \bar{\nabla}_{[\rho} b_{\mu\nu]} + m_b^2 b_{\mu\nu} = 0. \quad (3.13b)$$

Here we have defined the linearized Einstein tensor in terms of the covariant derivative $\bar{\nabla}_\mu$ compatible with the background metric $\bar{G}_{\mu\nu}$,

$$\begin{aligned} \mathcal{E}_{\mu\nu}{}^{\rho\sigma} h_{\rho\sigma} = & -\frac{1}{2} \left[\delta_\mu^\rho \delta_\nu^\sigma \bar{\nabla}^2 + \bar{G}^{\rho\sigma} \bar{\nabla}_\mu \bar{\nabla}_\nu - \delta_\mu^\rho \bar{\nabla}^\sigma \bar{\nabla}_\nu - \delta_\nu^\rho \bar{\nabla}^\sigma \bar{\nabla}_\mu \right. \\ & \left. - \bar{G}_{\mu\nu} \bar{G}^{\rho\sigma} \bar{\nabla}^2 + \bar{G}_{\mu\nu} \bar{\nabla}^\rho \bar{\nabla}^\sigma \right] h_{\rho\sigma}. \end{aligned} \quad (3.14)$$

These are the linear equations for a massive spin-2 field and a massive antisymmetric tensor in maximally symmetric backgrounds. Their masses are given by the following parameter combinations,

$$m_{\text{FP}}^2 = m^2 \left(3b_1 c + 4b_2 c^2 + 3b_3 c^3 \right), \quad (3.15a)$$

$$m_b^2 = \frac{m_{\text{P}}^2}{3c^2} m^2 \left(3b_1 c + 4b_2 c^2 + 3b_3 c^3 \right). \quad (3.15b)$$

Recall that c has mass dimension 1 and b_n has mass dimension $-n$, such that m_{FP} and m_b have the correct dimension. We note that these two masses are related by $m_b^2 = \frac{m_{\text{P}}^2}{3c^2} m_{\text{FP}}^2$. Since c is an arbitrary parameter in our setup, the two masses are independent.

4 ADM analysis

In this section we will perform a 3+1 split for the vierbein theory defined by the action (3.5). This will allow us to count the number of propagating degrees of freedom at the nonlinear level and confirm that it agrees with the linear theory.

We parameterize the general vierbein $e^a{}_\mu$ as a Lorentz transformation of a gauge-fixed vierbein $E^a{}_\mu$,

$$e^a{}_\mu = \Lambda^a{}_b E^b{}_\mu = \begin{pmatrix} \Gamma & \Gamma v_\beta \\ \Gamma v^\alpha & \mathcal{V}^\alpha_\beta \end{pmatrix} \begin{pmatrix} N & 0 \\ E_j^\beta N^j & E_i^\beta \end{pmatrix}, \quad (4.1)$$

where

$$\Gamma \equiv \frac{1}{\sqrt{1 - v^\alpha v_\alpha}}, \quad \mathcal{V}^\alpha_\beta \equiv \delta^\alpha_\beta + \frac{\Gamma^2}{1 + \Gamma} v^\alpha v_\beta. \quad (4.2)$$

Here $\alpha, \beta = 1, 2, 3$ are spatial Lorentz and $i, j = 1, 2, 3$ are spatial coordinate indices. The Lorentz rotations sit entirely in E_i^β , that is,

$$E_i^\beta = R^\beta{}_\alpha \bar{E}_i^\alpha, \quad (4.3)$$

for some gauge-fixed \bar{E}_i^α . The gauge-fixed vierbein depends on 10 independent field variables and the Lorentz matrix contains the 6 additional components that do not show up in the kinetic term for $G_{\mu\nu}$, but only appear in the mass potential.

For simplicity, let us take $\tilde{e}^a{}_\mu = \delta^a{}_\mu$.¹¹ The expressions with a general form for $\tilde{e}^a{}_\mu$ can be found in appendix B and all our results derived in the following straightforwardly generalize to that case. With this simple choice for $\tilde{e}^a{}_\mu$ we have that,

$$\tilde{e}^a{}_\mu \eta_{ab} e^b{}_\nu = \begin{pmatrix} -N\Gamma - \Gamma v_\alpha E_k^\alpha N^k & -\Gamma v_\alpha E_j^\alpha \\ N\Gamma \delta_{i\alpha} v^\alpha + \delta_{i\alpha} \mathcal{V}^\alpha_\beta E_k^\beta N^k & \delta_{i\alpha} \mathcal{V}^\alpha_\beta E_j^\beta \end{pmatrix}, \quad (4.4)$$

from where we read off the components of the antisymmetric tensor,

$$B_{ij} = 2\delta_{\alpha[i} E_{j]}^\beta \mathcal{V}^\alpha_\beta = 2\delta_{\alpha[i} \bar{E}_{j]}^\beta \mathcal{V}^\alpha_\beta R^\epsilon{}_\beta, \quad (4.5a)$$

$$B_{0i} = \Gamma v_\alpha R^\alpha{}_\beta \bar{E}_i^\beta + N\Gamma \delta_{i\alpha} v^\alpha + \delta_{i\alpha} \mathcal{V}^\alpha_\beta R^\beta{}_\epsilon E_k^\epsilon N^k. \quad (4.5b)$$

The expressions for the Einstein-Hilbert action in terms of ADM variables are well-known [41] and those for the mass potential were discussed in detail in Ref. [17, 21]. These results are summarized in appendix B, where we also derive the expressions for the $B_{\mu\nu}$ kinetic term. From those results it follows that, before solving any of the constraints, the action assumes the following form,

$$S = \int d^4x \left(\bar{\Pi}_a{}^j \dot{\bar{E}}_j^a + \Pi^{ij} \dot{B}_{ij} + NC + N^i \mathcal{C}_i + \mathcal{L}_{\text{rest}} \right), \quad (4.6)$$

¹¹For notational simplicity, we neglect the fact that $\tilde{e}^a{}_\mu$ is of mass dimension 1, since scales are irrelevant for the ADM analysis.

where $\bar{\Pi}_a^j$ and Π^{ij} are the canonical conjugate momenta of \bar{E}_j^a and B_{ij} , respectively. The canonical momenta for the remaining variables (N , N^i , B_{0j} and the Lorentz fields v^α , R^α_β) all vanish and hence the purely spatial components \bar{E}_j^a and B_{ij} are the only dynamical variables. Crucially, the constraints \mathcal{C} and \mathcal{C}_i as well as the additional terms in $\mathcal{L}_{\text{rest}}$ do not contain the lapse N nor the shift components N^i of the vierbein.

The next step is to perform an invertible field redefinition for the 3 independent components of the Lorentz rotation R^α_β .¹² Namely, we introduce $R'_{ij} \equiv \delta_{\alpha i} \bar{E}_j^\beta \mathcal{V}_\epsilon^\alpha R^\epsilon_\beta$ whose 3 independent components now replace the 3 rotation parameters. Note that this redefinition does not involve the lapse and shift functions. The spatial components B_{ij} are precisely the antisymmetric part of R'_{ij} . We can thus interpret the 3 independent components of $B_{ij} = R'_{[ij]}$ as giving dynamics to the 3 components of R^α_β . This implies that, in contrast to massive gravity, the equations of motion for the rotations R^α_β are now dynamical equations instead of constraints.

The set of nondynamical fields consists of N , N^i and the boosts v^α . Their equations are therefore constraints, which take the following form,

$$\frac{\delta S}{\delta N} = \mathcal{C}(v, \bar{E}, R') = 0, \quad (4.7a)$$

$$\frac{\delta S}{\delta N^i} = \mathcal{C}_i(v, \bar{E}, R') = 0, \quad (4.7b)$$

$$\frac{\delta S}{\delta v^\alpha} = N \frac{\delta \mathcal{C}}{\delta v^\alpha} + N^i \frac{\delta \mathcal{C}_i}{\delta v^\alpha} + \frac{\delta \mathcal{L}_{\text{rest}}}{\delta v^\alpha} = 0. \quad (4.7c)$$

We can solve the 3 equations in (4.7b) for the 3 components of the boost vector, yielding a solution $v^\alpha = v^\alpha(\bar{E}, R')$, which is independent of N and N^i . Then we solve the 3 equations in (4.7c) for the shift components N^i . Since these equations are linear in N and N^i , they result in a solution of the form,

$$N^i = A^i(v(\bar{E}, R'), \bar{E}, R') + N B^i(v(\bar{E}, R'), \bar{E}, R'), \quad (4.8)$$

which is linear in N . Thus, plugging the solutions to (4.7b) and (4.7c) for v^α and N^i back into the action, we arrive at the following form,

$$S = \int d^4x \left(\bar{\Pi}_a^j \dot{\bar{E}}_j^a + \Pi^{ij} R'_{[ij]} + N \mathcal{C}'(\bar{E}, R') + \mathcal{L}'_{\text{rest}}(\bar{E}, R') \right). \quad (4.9)$$

The action now contains 6 + 3 dynamical variables in \bar{E}_j^a and B_{ij} which are subject to one constraint coming from the N equation of motion, $\mathcal{C}' = 0$. This means we have 8 = 5 + 3 propagating degrees of freedom, in agreement with the linear theory for a massive spin-2 and a massive B -field.

Let us emphasize again that for massive gravity (i.e. in the absence of a kinetic term for $B_{\mu\nu}$), it was shown that the only ghost-free configuration is $B_{\mu\nu} = 0$ [21]. More precisely,

¹²We can extract the 3 independent components of R^α_β explicitly, for instance by using the Cayley transform for orthogonal transformations, $R = (\eta + A)^{-1}(\eta - A)$ with $A^T = -A$ which has 3 independent entries [42].

in massive gravity the equations for the non-dynamical rotation matrices R^α_β are constraints which have to be solved in order to determine whether the final action is linear in the lapse N . If the algebraic solution for R^α_β was such that the antisymmetric tensor was non-vanishing, $B_{\mu\nu} \neq 0$, the additional constraint needed to remove the Boulware-Deser ghost would be lost. The reason why our setup does not re-introduce the Boulware-Deser ghost, even though it contains $B_{\mu\nu} \neq 0$, is the dynamical nature of the antisymmetric tensor. We do not need to solve the R^α_β equations in order to see the full set of constraints. As a consequence, our theory contains the 3 propagating modes of the massive $B_{\mu\nu}$ field, which all appear with a standard (ghost-free) kinetic term.

5 Discussion

We have shown that the new classical vierbein theory defined by the action (3.5) is a ghost-free nonlinear theory for a massive spin-2 and a massive antisymmetric tensor field. Let us stress once more that our entire discussion was restricted to $D = 4$ but the results straightforwardly generalize to any spacetime dimension. The physical vierbein e^a_μ of the space time metric also defines the antisymmetric tensor whose kinetic term gives dynamics to the Lorentz components in e^a_μ . Since $B_{\mu\nu} \neq 0$ in general, the action is not expressible in terms of tensor fields.

The new action can be viewed as a deformation of the massless theory for $G_{\mu\nu}$ and $B_{\mu\nu}$. At the same time, it is a non-trivial generalization of ghost-free massive gravity, which is in fact contained in the theory and corresponds to the solution $B_{\mu\nu} = 0$. Our results can be extended to the case where the reference frame \tilde{e}^a_μ is dynamical, or even to a setup with multiple dynamical vierbein fields. This yields generalizations of bi- and multimetric gravity which are manifestly invariant under local Lorentz transformations and diffeomorphisms. We will discuss this in detail in a future publication.

As a ghost-free theory with $B_{\mu\nu} = 0$, our result could also allow for the inclusion of new matter couplings into multi-vierbein theories. Ref. [46] suggested to couple a linear combination of vierbein fields to the matter sector. This construction failed to be ghost-free precisely because the antisymmetric vierbein combination was non-vanishing but also non-dynamical [21]. With a kinetic term for $B_{\mu\nu}$, this problem could be resolved.

Our ADM analysis shows that the action propagates the correct number of degrees of freedom at the nonlinear level. In particular, the Boulware-Deser ghost (which is a notorious problem in theories with massive spin-2 fields) is removed from the physical spectrum by an additional constraint. Nevertheless, in principle there could be more ghosts or other pathologies hidden in the theory, as part of the dynamical modes in $G_{\mu\nu}$ and $B_{\mu\nu}$. We believe that this is rather unlikely because, as we showed in sections 2.1 and 3.2, the kinetic structures of the massless theory for independent tensor fields are not modified by the identification of $B_{\mu\nu}$ with the antisymmetric vierbein components nor by the presence of the mass term.

Another potential issue is the question of causality. In Ref. [47] it was shown that the possibility to address the question of causal propagation in ghost-free bimetric theory

requires precisely the vanishing of the antisymmetric vierbein combination. The presence of our dynamical $B_{\mu\nu}$ could in principle spoil this feature. Once again, the fact that the kinetic structures are the same as in the massless theory may indicate that this is not the case. A different approach to studying the causal structure of massive spin-2 theories in the framework of scattering amplitudes was taken in Ref. [48–52]. A thorough investigation of this issue is beyond the scope of this paper.

It would be very interesting to develop a geometrical interpretation for our setup. The vierbein (or frame field) e^a_μ together with the invariant η_{ab} of the $\text{SO}(1,3)$ Lorentz structure group define the metric $G_{\mu\nu} = e^a_\mu \eta_{ab} e^b_\nu$ of the pseudo-Riemannian geometry. For certain configurations, the presence of the $B_{\mu\nu}$ field can be interpreted as an additional symplectic structure on the manifold with the structure group $\text{Sp}(4)$. This can be seen by introducing a symplectic frame field b^a_μ which, together with the symplectic invariant Ω_{ab} , gives a decomposition $B_{\mu\nu} = b^a_\mu \Omega_{ab} b^b_\nu$. It is clear that then $B_{\mu\nu}$ is invariant under symplectic transformations of its frame field. Well-studied cases with *compatible* metric and symplectic structure are manifolds with (almost) Hermitian or (almost) Kähler geometry. In our setup, $G_{\mu\nu}$ and $B_{\mu\nu}$ are intimately related via the vierbein e^a_μ and it could be interesting to investigate the consequences when their respective structures are made compatible with each other. In spirit this would be similar to recent attempts to construct the geometry of quantum gravity [11, 12]. Moreover our setup may allow to relate the symplectic interpretation of the $B_{\mu\nu}$ field to that of a geometry with torsion, as briefly discussed in section 1.2. The torsion perspective might also be related to teleparallel theories where the antisymmetric part of the linearized vierbein becomes dynamical, see section 4.6 of [23]. Moreover, it would be interesting to see whether there is a relation to the non-commutative setup of Ref. [53].

As outlined in the introduction, in string theory the antisymmetric tensor field appears together with the metric as massless excitation of the closed string. In the low-energy effective theory, together with the dilaton field, they make up the bosonic sector of supergravity. It would thus be interesting to investigate the consequences of adding our mass term to the supergravity action, which could also be used to deform the structure of double field theory. Possibly, this could even help with supersymmetrizing the massive gravity potential.

Acknowledgements. We are grateful to Fawad Hassan and Ignatios Antoniadis for useful discussions. This work is supported by a grant from the Max-Planck-Society.

A Independence of $G_{\mu\nu}$ and $B_{\mu\nu}$

In this appendix we discuss the mild assumptions under which the vierbein combinations $G_{\mu\nu}$ and $B_{\mu\nu}$ can be treated as independent. Consider a gauge-fixed vierbein E^a_μ . The general vierbein is given by $e^a_\mu = \Lambda^a_b E^b_\mu$ where Λ^a_b is a local Lorentz transformation satisfying $\Lambda^T \eta \Lambda = \eta$. As already discussed in Ref. [43], we can parameterize it by its Cayley transform [42],

$$\Lambda^a_b = [(\eta - A)^{-1}]^{ac} [\eta_{cb} + A_{cb}], \quad (\text{A.1})$$

where $A_{ab} = -A_{ba}$ contains the 6 Lorentz parameters. We then have that,

$$G_{\mu\nu} = E^a{}_\mu \eta_{ab} E^b{}_\nu, \quad B_{\mu\nu} = \Lambda^a{}_b E^b{}_\mu \eta_{ac} \tilde{e}^c{}_\nu - \Lambda^a{}_b E^b{}_\nu \eta_{ac} \tilde{e}^c{}_\mu. \quad (\text{A.2})$$

The 6 components in A_{ab} do not show up in the Lorentz invariant metric $G_{\mu\nu}$, whereas $B_{\mu\nu}$ depends on all of them. A field redefinition $B_{\mu\nu} \rightarrow A_{ab}$ would thus render the metric and antisymmetric components independent. But it is not immediately obvious how to invert this dependence and express A_{ab} as a function of $B_{\mu\nu}$ in a covariant way. Nevertheless, at the level of a linearized Lorentz transformation, we can easily derive the condition for the invertibility of the dependence.

To this end, let us assume that the symmetric vierbein condition can be enforced by a local Lorentz transformation $\bar{\Lambda}^a{}_b$ of the general vierbein $e^a{}_\mu$,¹³

$$\bar{\Lambda}^a{}_b e^b{}_\mu \eta_{ac} \tilde{e}^c{}_\nu - \bar{\Lambda}^a{}_b e^b{}_\nu \eta_{ac} \tilde{e}^c{}_\mu = 0. \quad (\text{A.3})$$

Infinitesimally we write the transformation as $\bar{\Lambda}_a{}^b = \delta_a{}^b - \bar{\omega}_{ac} \eta^{cb}$, such that the above equation can be written in the form,

$$B_{\mu\nu} - e^a{}_\mu \tilde{e}^b{}_\nu \bar{\omega}_{ab} + e^a{}_\nu \tilde{e}^b{}_\mu \bar{\omega}_{ab} = 0 \quad (\text{A.4})$$

Now we see that the right-hand side can be written as the invertible operator $P_{\mu\nu}{}^{ab}$ (which we encountered in section 2.2 when reproducing the local symmetry transformations) acting on the infinitesimal Lorentz parameter $\bar{\omega}_{ab}$,

$$P_{\mu\nu}{}^{ab} \bar{\omega}_{ab} = 2e^{[a}{}_{[\mu} \tilde{e}^{b]}{}_{\nu]} \bar{\omega}_{ab} = e^a{}_\mu \tilde{e}^b{}_\nu \bar{\omega}_{ab} - e^a{}_\nu \tilde{e}^b{}_\mu \bar{\omega}_{ab}. \quad (\text{A.5})$$

Comparing to (A.4) we see that,

$$P_{\mu\nu}{}^{ab} \bar{\omega}_{ab} = B_{\mu\nu}. \quad (\text{A.6})$$

In order to obtain the Lorentz transformation which satisfies (A.3), we need to invert this equation and determine $\bar{\omega}_{ab}$ in terms of the components of $B_{\mu\nu}$. In other words, the invertibility of P makes it possible to map the 6 components of $B_{\mu\nu}$ onto the 6 Lorentz parameters $\bar{\omega}_{ab}$ which do not show up in $G_{\mu\nu}$. This implies that $B_{\mu\nu}$ and $G_{\mu\nu}$ can be treated as independent objects precisely when the symmetric vierbein condition can be satisfied by a Lorentz transformed $e^a{}_\mu$.

Analogously, the operator $\tilde{P}_{\mu\nu}{}^{ab}$ (which we encountered in section 2.1 when separating the tensor equations) maps the Lorentz parameters which symmetrize the inverse $e_a{}^\mu \eta^{ab} \tilde{e}_b{}^\nu$ onto the components of its antisymmetric part $\tilde{B}^{\mu\nu} = e_a{}^{[\mu} \eta^{ab} \tilde{e}_b{}^{\nu]}$. Since the symmetry of $e_a{}^\mu \eta^{ab} \tilde{e}_b{}^\nu$ is equivalent to the symmetry of its inverse $e_a{}^\mu \eta^{ab} \tilde{e}_b{}^\nu$, the local Lorentz transformation parametrized by $\bar{\omega}_{ab}$ is the same in both cases. Thus we have that $\tilde{P}_{ab}{}^{\mu\nu} \eta^{ac} \eta^{bd} \bar{\omega}_{cd} = \tilde{B}^{\mu\nu}$.

¹³This assumption is equivalent to the existence of the square-root matrix $\sqrt{G^{-1}f}$ with $f = \tilde{e}^T \eta \tilde{e}$ [39], which appears in the tensor formulation of massive gravity [40].

B Details of the ADM analysis

This appendix contains technical details on the 3+1 split and degree of freedom counting performed in section 4.

B.1 Standard massive theory for $B_{\mu\nu}$

We begin by performing an ADM analysis of the standard massive theory for $B_{\mu\nu}$ in curved space, whose Lagrangian reads,

$$\mathcal{L}_B = -\frac{1}{2 \cdot 3!} \sqrt{G} H_{\mu\nu\rho} H^{\mu\nu\rho} - m^2 \sqrt{G} B_{\mu\nu} B_{\rho\sigma} G^{\mu\rho} G^{\nu\sigma}. \quad (\text{B.1})$$

We introduce ADM variables to parametrize the metric and its inverse by [41],

$$G_{\mu\nu} = \begin{pmatrix} -N^2 + N_k N^k & N_j \\ N_i & \gamma_{ij} \end{pmatrix}, \quad G^{\mu\nu} = \frac{1}{N^2} \begin{pmatrix} -1 & N^j \\ N^i & N^2 \gamma^{ij} - N^i N^j \end{pmatrix} \quad (\text{B.2})$$

where $N^i = \gamma^{ij} N_j$ and for the determinants we now have $\sqrt{G} = N\sqrt{\gamma}$. Moreover, we split the antisymmetric tensor $B_{\mu\nu}$ into its spatial components B_{ij} and its mixed space-time components B_{0j} . The canonical momentum conjugate to B_{mn} is then obtained as,

$$\Pi^{mn} \equiv \frac{\partial \mathcal{L}_B}{\partial \dot{B}_{mn}} = \frac{\sqrt{\gamma}}{2N} \left(\partial_0 B_{ij} - 3N^k \partial_{[k} B_{ij]} - 2\partial_i B_{0j} \right) \gamma^{i[m} \gamma^{n]j}, \quad (\text{B.3})$$

while the components B_{0j} remain nondynamical, i.e. $\Pi^{0j} = 0$. With some manipulations we can now rewrite the Lagrangian in the form,

$$\begin{aligned} \mathcal{L}_B = & \Pi^{mn} \dot{B}_{mn} - \frac{N}{\sqrt{\gamma}} \Pi^{mn} \Pi_{mn} - 3N^{[k} \Pi^{ij]} \partial_k B_{ij} - 2\Pi^{ij} \partial_i B_{0j} - \frac{1}{4} N \sqrt{\gamma} \partial_k B_{ij} \partial_l B_{pq} \Xi^{kl,[ij],[pq]} \\ & + m^2 \frac{\sqrt{\gamma}}{N} [2B_{0i} B_{0j} \gamma^{ij} - 4B_{0j} B_{ik} N^i \gamma^{jk} - N^2 B_{ij} B_{kl} G^{ik} G^{jl}], \end{aligned} \quad (\text{B.4})$$

where we have defined the following spatial tensor combination,

$$\Xi^{kl,[ij],[pq]} \equiv \gamma^{lk} \gamma^{pi} \gamma^{qj} + \gamma^{lj} \gamma^{pk} \gamma^{qi} + \gamma^{li} \gamma^{pj} \gamma^{qk}. \quad (\text{B.5})$$

Due to $\Pi^{0i} = 0$, the B_{0i} equations are constraints which we need to solve explicitly in order to see the linearity of the final action in the lapse N and the shift N^i . The solution is easily obtained and reads,

$$B_{0i} = N^j B_{ji} - \frac{N}{2m^2 \sqrt{\gamma}} \partial_j \Pi^{jk} \gamma_{ki}. \quad (\text{B.6})$$

The final Lagrangian after integrating out the B_{0j} then is,

$$\begin{aligned} \mathcal{L}_B = & \Pi^{mn} \dot{B}_{mn} - N^i \left(3\Pi^{jk} \partial_{[i} B_{jk]} + 2B_{ij} \partial_k \Pi^{jk} \right) - N \left\{ \frac{1}{\sqrt{\gamma}} \left(\Pi^{mn} \Pi_{mn} + \frac{1}{2m^2} \partial_i \Pi^{ik} \partial_j \Pi^{jl} \gamma_{kl} \right) \right. \\ & \left. + \sqrt{\gamma} \left(\frac{1}{4} \partial_k B_{ij} \partial_l B_{pq} [\gamma^{lk} \gamma^{pi} \gamma^{qj} + \gamma^{lj} \gamma^{pk} \gamma^{qi} + \gamma^{li} \gamma^{pj} \gamma^{qk}] + m^2 B_{ij} B_{kl} \gamma^{ik} \gamma^{jl} \right) \right\}. \end{aligned} \quad (\text{B.7})$$

This is manifestly linear in N and N^i , as expected since the Lagrangian in (B.1) is diffeomorphism invariant.

Let us count the number of propagating degrees of freedom. The components B_{0j} have been algebraically determined by their own equations. The dynamical objects are the spatial components B_{mn} , on which there are no further constraints. Thus the massive theory for $B_{\mu\nu}$ defined by (B.1) has 3 propagating degrees of freedom.¹⁴ Adding the Einstein-Hilbert term to the action will introduce the 2 additional degrees of freedom of the massless graviton.

B.2 Action with new mass term

We now focus on our new action defined by (3.5) and show that it propagates the degrees of freedom of a massive $B_{\mu\nu}$ and a massive $G_{\mu\nu}$ field. We parametrize the general vierbein e^a_μ as in equation (4.1) and we allow for a general form of \tilde{e}^a_μ , which we write in 3 + 1 form,

$$\tilde{e}^a_\mu = \begin{pmatrix} L & 0 \\ \varphi^\beta_j L^j & \varphi^\beta_i \end{pmatrix}. \quad (\text{B.8})$$

There is no need to include Lorentz parameters for the second vierbein. They can all be shifted to the Lorentz matrix in e^a_μ , using the fact that a simultaneous Lorentz transformation of both vierbeine leaves the action invariant.

B.2.1 Einstein-Hilbert term

As is well-known, the Einstein-Hilbert term in the Lagrangian written in terms of the vierbein is of the form [44] (see also the review [45]),

$$\mathcal{L}_e = \bar{\Pi}_a^i \dot{\tilde{E}}_i^a - N\mathcal{C}^{(e)} - N^i \mathcal{C}_i^{(e)}, \quad (\text{B.9a})$$

where $\bar{\Pi}_a^i$ is the canonical momenta conjugate to the spatial vierbein components \tilde{E}_i^a . The kinetic term for $G_{\mu\nu}$ is linear in N , and N^i since the constraint contributions $\mathcal{C}^{(e)}$ and $\mathcal{C}_i^{(e)}$ depend only on the remaining ADM variables. Their precise form is irrelevant for the arguments used in this work.

B.2.2 Mass term

We summarize briefly the relevant results of Ref. [17]. The mass term and the cosmological constant term in equation (3.5) can be combined into the potential,

$$\begin{aligned} V &= \sum_{n=0}^3 b_n \epsilon_{a_1 a_2 a_3 a_4} \tilde{e}^{a_1} \wedge \cdots \wedge \tilde{e}^{a_n} \wedge e^{a_{n+1}} \wedge \cdots \wedge e^{a_4} \\ &= \sum_{n=0}^3 b_n \epsilon_{a_1 a_2 a_3 a_4} \epsilon^{\mu_1 \mu_2 \mu_3 \mu_4} \tilde{e}^{a_1}_{\mu_1} \cdots \tilde{e}^{a_n}_{\mu_n} e^{a_{n+1}}_{\mu_{n+1}} \cdots e^{a_4}_{\mu_4}. \end{aligned} \quad (\text{B.10})$$

¹⁴This result is of course in agreement with the covariant counting, which is particularly easy to see in flat space. When $G_{\mu\nu} = \eta_{\mu\nu}$, the divergence of the equations of motion for the massive $B_{\mu\nu}$ field gives the constraints $\partial^\mu B_{\mu\nu} = 0$. Note that these are only 3 independent equations because $\partial^\nu \partial^\mu B_{\mu\nu} = 0$ is identically satisfied. The constraints therefore remove 3 out of the 6 components in $B_{\mu\nu}$, leaving 3 propagating modes.

Note that there are 4 terms since we have included the cosmological constant Λ for $e^a{}_\mu$ by defining $b_0 \equiv 2\Lambda/(m^2 4!)$. The ADM parametrizations of the two vierbeine in (4.1) and (B.8) give,

$$e^0{}_0 = \Gamma(N + v_\alpha E^\alpha{}_i N^i), \quad \tilde{e}^0{}_0 = L, \quad (\text{B.11a})$$

$$e^\alpha{}_0 = \Gamma v^\alpha N + \mathcal{V}^\alpha{}_\beta E^\beta{}_i N^i, \quad \tilde{e}^\alpha{}_0 = \varphi^\alpha{}_i L^i, \quad (\text{B.11b})$$

while all other (purely spatial) components are independent of N , N^i , L and L^i . The anti-symmetric wedge product (i.e. the factor of $\epsilon^{\mu_1\mu_2\mu_3\mu_4}$) in the potential ensures its linearity in the components $e^a{}_0$ and $\tilde{e}^a{}_0$ and thus the linearity in N , N^i , L and L^i . We can therefore write the potential in the form,

$$V = N\mathcal{C}^{(V)} + N^i\mathcal{C}_i^{(V)} + L\tilde{\mathcal{C}}^{(V)} + L^i\tilde{\mathcal{C}}_i^{(V)}, \quad (\text{B.12})$$

where $\mathcal{C}^{(V)}$, $\mathcal{C}_i^{(V)}$, $\tilde{\mathcal{C}}^{(V)}$ and $\tilde{\mathcal{C}}_i^{(V)}$ are all functions of the remaining ADM variables alone.

B.2.3 Kinetic term for $B_{\mu\nu}$

So far we have seen that the Einstein-Hilbert term and the mass potential are linear in the lapse and the shift functions. We now show that this is also the case for the kinetic term of the antisymmetric components $B_{\mu\nu} = e^a{}_\mu \eta_{ab} \tilde{e}^b{}_\nu - \tilde{e}^a{}_\mu \eta_{ab} e^b{}_\nu$.

To this end, we again split the $B_{\mu\nu}$ into its spatial components B_{ij} and its mixed space-time components B_{0j} . The 3 + 1 form of the kinetic term of $B_{\mu\nu}$ in (1.1) is the same as in (B.7). It reads,

$$\begin{aligned} \mathcal{L}_B = & \Pi^{mn} \dot{B}_{mn} - \frac{N}{\sqrt{\gamma}} \Pi^{mn} \Pi_{mn} - 3N^{[k} \Pi^{ij]} \partial_k B_{ij} + 2\partial_i \Pi^{ij} B_{0j} \\ & - \frac{1}{4} N \sqrt{\gamma} \partial_k B_{ij} \partial_l B_{pq} \Xi^{kl,[ij],[pq]}, \end{aligned} \quad (\text{B.13})$$

where $\Xi^{kl,[ij],[pq]}$ was already defined in (B.5) and $\gamma_{ij} = \delta_{\alpha\beta} E^\alpha{}_i E^\beta{}_j = G_{ij}$ is the spatial part of our spacetime metric $G_{\mu\nu}$ as in (B.2) with determinant $\gamma \equiv \det \gamma_{ij}$. The canonical momentum for the spatial components B_{mn} is given by (B.3). We remind the reader that the standard mass term for the $B_{\mu\nu}$ field is not contained in the action.

The definition of $B_{\mu\nu}$ in terms of the vierbeine gives the following ADM expressions,

$$B_{0i} = -\Gamma v_\alpha (L E^\alpha{}_i + N \varphi^\alpha{}_i) + \varphi^\alpha{}_k \mathcal{V}_{\alpha\beta} E^\beta{}_l (L^k \delta_i^l - \delta_i^k N^l), \quad (\text{B.14a})$$

$$B_{ij} = \mathcal{V}_{\alpha\beta} \varphi^\alpha{}_{[i} E^\beta{}_{j]} = \mathcal{V}_{\alpha\beta} R^\beta{}_\gamma \varphi^\alpha{}_{[i} \bar{E}^\gamma{}_{j]}. \quad (\text{B.14b})$$

Note that B_{0i} is linear in the lapses and shifts while B_{ij} is independent of them. Inserting these expressions into (B.13), we obtain,

$$\begin{aligned} \mathcal{L}_B = & \Pi^{mn} \dot{B}_{mn} - \frac{N}{\sqrt{\gamma}} \Pi^{mn} \Pi_{mn} - 3N^{[k} \Pi^{ij]} \partial_k (\varphi^\alpha{}_i \mathcal{V}_{\alpha\beta} E^\beta{}_j) \\ & + \partial_i \Pi^{ij} [-\Gamma v_\alpha (L E^\alpha{}_j + N \varphi^\alpha{}_j) + \varphi^\alpha{}_k \mathcal{V}_{\alpha\beta} E^\beta{}_l (L^k \delta_j^l - \delta_j^k N^l)] \\ & - \frac{1}{4} N \sqrt{\gamma} \partial_k (\varphi^\alpha{}_i \mathcal{V}_{\alpha\beta} E^\beta{}_j) \partial_l (\varphi^\alpha{}_p \mathcal{V}_{\alpha\beta} E^\beta{}_q) \Xi^{kl,[ij],[pq]}. \end{aligned} \quad (\text{B.15})$$

This can be written in the form,

$$\mathcal{L}_B = \Pi^{mn} \dot{B}_{mn} - N\mathcal{C}^{(B)} - L\tilde{\mathcal{C}}^{(B)} - N^i \mathcal{C}_i^{(B)} - L^i \tilde{\mathcal{C}}_i^{(B)}, \quad (\text{B.16})$$

where again none of $\mathcal{C}^{(B)}$, $\mathcal{C}_i^{(B)}$, $\tilde{\mathcal{C}}^{(B)}$ and $\tilde{\mathcal{C}}_i^{(B)}$ depend on N , N^i , L and L^i .

B.2.4 Full action in ADM variables

Putting together the results for the two kinetic terms and the mass potential obtained in the previous subsections, the whole action can be written in the form

$$S = \int d^4x \left(\bar{\Pi}_a^j \dot{E}_j^a + \Pi^{ij} \dot{B}_{ij} - N\mathcal{C} - N^i \mathcal{C}_i - L\tilde{\mathcal{C}} - L^i \tilde{\mathcal{C}}_i \right), \quad (\text{B.17})$$

where the \mathcal{C} and \mathcal{C}_i , $\tilde{\mathcal{C}}$ and $\tilde{\mathcal{C}}_i$ contain the contributions from (B.9), (B.12) and (B.16) which do not depend on any of the shifts or lapses. Note that since \tilde{e}_μ^a is not a dynamical field in our setup, only the terms proportional to N and N^i correspond to the constraints. The remaining terms make up $\mathcal{L}_{\text{rest}}$ in (4.6).

References

- [1] M. Kalb and P. Ramond, Phys. Rev. D **9** (1974) 2273.
- [2] M. Grana, Phys. Rept. **423** (2006) 91 [hep-th/0509003].
- [3] M. Gualtieri, math/0401221 [math-dg].
- [4] N. Hitchin, Quart. J. Math. **54** (2003) 281 [math/0209099 [math-dg]].
- [5] W. Siegel, Phys. Rev. D **47** (1993) 5453 [hep-th/9302036].
- [6] W. Siegel, Phys. Rev. D **48** (1993) 2826 [hep-th/9305073].
- [7] C. Hull and B. Zwiebach, JHEP **0909** (2009) 099 [arXiv:0904.4664 [hep-th]].
- [8] G. Aldazabal, D. Marques and C. Nunez, Class. Quant. Grav. **30** (2013) 163001 [arXiv:1305.1907 [hep-th]].
- [9] D. S. Berman and D. C. Thompson, Phys. Rept. **566** (2014) 1 [arXiv:1306.2643 [hep-th]].
- [10] O. Hohm, D. Lst and B. Zwiebach, Fortsch. Phys. **61** (2013) 926 [arXiv:1309.2977 [hep-th]].
- [11] L. Freidel, F. J. Rudolph and D. Svoboda, JHEP **1711** (2017) 175 [arXiv:1706.07089 [hep-th]].
- [12] L. Freidel, F. J. Rudolph and D. Svoboda, arXiv:1806.05992 [hep-th].
- [13] R. G. Leigh, Mod. Phys. Lett. A **4** (1989) 2767.
- [14] J. Polchinski, Phys. Rev. Lett. **75** (1995) 4724 [hep-th/9510017].
- [15] C. de Rham, G. Gabadadze and A. J. Tolley, Phys. Rev. Lett. **106** (2011) 231101 [arXiv:1011.1232 [hep-th]].
- [16] S. F. Hassan and R. A. Rosen, JHEP **1202** (2012) 126 [arXiv:1109.3515 [hep-th]].
- [17] K. Hinterbichler and R. A. Rosen, JHEP **1207** (2012) 047 [arXiv:1203.5783 [hep-th]].

- [18] K. Hinterbichler, *Rev. Mod. Phys.* **84** (2012) 671 [arXiv:1105.3735 [hep-th]].
- [19] C. de Rham, *Living Rev. Rel.* **17** (2014) 7 [arXiv:1401.4173 [hep-th]].
- [20] A. Schmidt-May and M. von Strauss, *J. Phys. A* **49** (2016) no.18, 183001 [arXiv:1512.00021 [hep-th]].
- [21] C. de Rham and A. J. Tolley, *Phys. Rev. D* **92** (2015) no.2, 024024 [arXiv:1505.01450 [hep-th]].
- [22] G. Gabadadze, K. Hinterbichler, D. Pirtskhalava and Y. Shang, *Phys. Rev. D* **88** (2013) no.8, 084003 [arXiv:1307.2245 [hep-th]].
- [23] T. Ortin, Cambridge Univ. Press (2004)
- [24] J. W. Moffat, *Phys. Rev. D* **19** (1979) 3554.
- [25] A. H. Chamseddine, *Commun. Math. Phys.* **218** (2001) 283 [hep-th/0005222].
- [26] A. H. Chamseddine and V. Mukhanov, *JHEP* **1003** (2010) 033 [arXiv:1002.0541 [hep-th]].
- [27] A. H. Chamseddine and V. Mukhanov, *JHEP* **1208** (2012) 036 [arXiv:1205.5828 [hep-th]].
- [28] B. Altschul, Q. G. Bailey and V. A. Kostelecky, *Phys. Rev. D* **81** (2010) 065028 [arXiv:0912.4852 [gr-qc]].
- [29] K. N. Lau and M. D. Seifert, *Phys. Rev. D* **95** (2017) no.2, 025023 [arXiv:1309.2241 [hep-th]].
- [30] S. Aashish and S. Panda, *Phys. Rev. D* **97** (2018) no.12, 125005 [arXiv:1803.10157 [gr-qc]].
- [31] S. Aashish and S. Panda, arXiv:1806.08194 [gr-qc].
- [32] S. Aashish, A. Padhy, S. Panda and A. Rana, *Eur. Phys. J. C* **78** (2018) no.11, 887 [arXiv:1808.04315 [gr-qc]].
- [33] J. Scherk and J. H. Schwarz, *Phys. Lett.* **52B** (1974) 347.
- [34] A. Smailagic and E. Spallucci, *J. Phys. A* **34** (2001) L435 [hep-th/0106173].
- [35] D. G. Boulware and S. Deser, *Phys. Rev. D* **6** (1972) 3368.
- [36] S. F. Hassan and R. A. Rosen, *Phys. Rev. Lett.* **108** (2012) 041101 [arXiv:1106.3344 [hep-th]].
- [37] S. F. Hassan, R. A. Rosen and A. Schmidt-May, *JHEP* **1202** (2012) 026 [arXiv:1109.3230 [hep-th]].
- [38] S. F. Hassan and A. Schmidt-May, arXiv:1804.09723 [hep-th].
- [39] C. Deffayet, J. Mourad and G. Zahariade, *JHEP* **1303** (2013) 086 [arXiv:1208.4493 [gr-qc]].
- [40] S. F. Hassan and R. A. Rosen, *JHEP* **1107** (2011) 009 [arXiv:1103.6055 [hep-th]].
- [41] R. L. Arnowitt, S. Deser and C. W. Misner, *Gen. Rel. Grav.* **40** (2008) 1997 [gr-qc/0405109].
- [42] A. Cayley, *J. Reine Angew. Math.*, **32**, 1846.
- [43] S. F. Hassan, A. Schmidt-May and M. von Strauss, arXiv:1204.5202 [hep-th].
- [44] S. Deser and C. J. Isham, *Phys. Rev. D* **14** (1976) 2505.
- [45] P. Peldan, *Class. Quant. Grav.* **11** (1994) 1087 [gr-qc/9305011].
- [46] K. Hinterbichler and R. A. Rosen, *Phys. Rev. D* **92** (2015) no.2, 024030 [arXiv:1503.06796 [hep-th]].

- [47] S. F. Hassan and M. Kocic, JHEP **1805** (2018) 099 [arXiv:1706.07806 [hep-th]].
- [48] K. Hinterbichler, A. Joyce and R. A. Rosen, JHEP **1803** (2018) 051 [arXiv:1708.05716 [hep-th]].
- [49] J. Bonifacio, K. Hinterbichler, A. Joyce and R. A. Rosen, JHEP **1806** (2018) 075 [arXiv:1712.10020 [hep-th]].
- [50] C. de Rham, S. Melville, A. J. Tolley and S. Y. Zhou, JHEP **1803** (2018) 011 [arXiv:1706.02712 [hep-th]].
- [51] C. de Rham, S. Melville and A. J. Tolley, JHEP **1804** (2018) 083 [arXiv:1710.09611 [hep-th]].
- [52] C. de Rham, S. Melville, A. J. Tolley and S. Y. Zhou, arXiv:1804.10624 [hep-th].
- [53] M. de Cesare, M. Sakellariadou and P. Vitale, Class. Quant. Grav. **35** (2018) no.21, 215009 [arXiv:1806.04666 [gr-qc]].