

Chapter 1

Relaxation of Periodic and Nonstandard Growth Integrals by means of Two-scale convergence

Joel Fotso Tachago, Hubert Nnang and Elvira Zappale

Abstract An integral representation result is obtained for the variational limit of the family functionals $\int_{\Omega} f(\frac{x}{\varepsilon}, Du)dx$, $\varepsilon > 0$, when the integrand $f = f(x, v)$ is a Carathéodory function, periodic in x , convex in v and with nonstandard growth.

1.1 Introduction

In [TN1], the authors extended the notion of two-scale convergence introduced by [N] (see also [A], [CDG], [FZ], [V] among a wider literature for extensions and related notions) to the Orlicz-Sobolev setting and obtained, under strict convexity assumption on f and suitable boundary conditions, the existence of a unique minimizer for a suitable limit functional as the limit of the minimizers of the original functionals $\int_{\Omega} f(\frac{x}{\varepsilon}, Du)dx$ as $\varepsilon \rightarrow 0$.

In particular they proved (cf. [TN1, Lemma 5.2]) that for every sequence $(u_{\varepsilon})_{\varepsilon} \in W^1L^B(\Omega; \mathbb{R})$ such that $(Du_{\varepsilon})_{\varepsilon}$ weakly $2s$ -converges to $\mathbb{D}u_0 = Du + D_y u_1$, where

J. Fotso Tachago

University of Bamenda , Faculty of Sciences, Department of Mathematics and Computers Sciences, P.O. Box 39, Bambili, Cameroon

and

Dipartimento di Ingegneria Industriale, Università degli Studi di Salerno, Via Giovanni Paolo II, 132, Fisciano (Sa) 84084, Italy

e-mail: fotsotachago@yahoo.fr,

H. Nnang,

University of Yaoundé I, and École Normale Supérieure de Yaoundé, P.O. Box 47 Yaounde, Cameroon,

e-mail: hnnang@u1.uninet.cm;hnnang@yahoo.fr,

E. Zappale

Dipartimento di Ingegneria Industriale, Università degli Studi di Salerno, 84084 Fisciano (SA), Italy

e-mail: ezappale@unisa.it.

$u_0 = (u, u_1) \in W^1 L^B(\Omega) \times L^B(\Omega; W_{\#}^1 L_{\text{per}}^B(Y))$. Then

$$\iint_{\Omega \times Y} f(y, \mathbb{D}u_0) dx dy \leq \liminf_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\varepsilon}\right) dx, \quad (1.1)$$

where $Y := (0, 1)^d$ ($d \in \mathbb{N}$) and $\mathbb{D}u_0 := Du + D_y u_1$ (see Section 2 for the notations adopted in this introduction).

On the other hand, by the very nature of two-scale convergence they obtain, under homogeneous boundary conditions on $\partial\Omega$, (the same proof can be performed for periodic **any** boundary conditions), for u and u_1 regular, the existence of a suitable sequence $(\bar{u}_{\varepsilon})_{\varepsilon} \subseteq W^1 L^B(\Omega)$ such that $\bar{u}_{\varepsilon} \rightharpoonup u$, and the opposite inequality holds:

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, D\bar{u}_{\varepsilon}\right) dx = \iint_{\Omega \times Y} f(y, \mathbb{D}u_0) dx dy.$$

Here, by means of two scale convergence we aim to extend their result to any couple of functions $(u_0 \equiv) (u, u_1) \in W^1 L^B(\Omega) \times L^B(\Omega; W_{\#}^1 L_{\text{per}}^B(Y))$, and also to obtain an integral representation result for

$$\inf \left\{ \liminf_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\varepsilon}\right) dx : u_{\varepsilon} \rightharpoonup u \text{ weakly in } W^1 L^B(\Omega) \right\}.$$

Indeed, after stating preliminary results in Section 1.2 on Orlicz-Sobolev spaces and homogenization theory, in Section 1.3 we will prove the following theorem:

Theorem 1. *Let Ω be a bounded open set with Lipschitz boundary and let $f : \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}$ be a Carathéodory function such that*

$$f(x, \cdot) \text{ is convex for a.e. } x \in \Omega,$$

and there exist constants c, c' and $C \in \mathbb{R}^+$ such that for a.e. $x \in \Omega$ and every $\xi \in \mathbb{R}^d$,

$$cB'(|\xi|) - c' \leq f(x, \xi) \leq C(1 + B(|\xi|)) \quad (1.2)$$

with B, B' equivalent N -functions which satisfy the Δ_2 condition. Then, it results that for every $u \in W^1 L^B(\Omega)$,

$$\begin{aligned} & \inf \left\{ \liminf_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\varepsilon}\right) dx : u_{\varepsilon} \rightharpoonup u \text{ weakly in } W^1 L^B(\Omega) \right\} \\ &= \inf \left\{ \limsup_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\varepsilon}\right) dx : u_{\varepsilon} \rightharpoonup u \text{ weakly in } W^1 L^B(\Omega) \right\} = \int_{\Omega} f_{\text{hom}}(Du) dx, \end{aligned} \quad (1.3)$$

where $f_{\text{hom}} : \mathbb{R}^d \rightarrow \mathbb{R}$ is the density defined by

$$f_{\text{hom}}(\xi) := \inf \left\{ \int_Y f(y, \xi + Du) dy : u \in W^1 L_{\text{per}}^B(Y) \right\}. \quad (1.4)$$

We underline that the analysis presented in this paper, holds also in the vectorial case, i.e. fields $u \in W^1L^B(\Omega; \mathbb{R}^m)$, with the exact same techniques, provided that $f(x, \cdot)$ is convex.

Furthermore, in order to prove (1.3), we also obtain for every $u_0 \in W^1L^B(\Omega) \times L^B(\Omega; W_{\#}^1L_{\text{per}}^B(Y))$, the following two-scale representation:

$$\begin{aligned} & \inf \left\{ \liminf_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\varepsilon}\right) dx : u_{\varepsilon} \xrightarrow{2s} u_0 \right\} \\ &= \inf \left\{ \limsup_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\varepsilon}\right) dx : u_{\varepsilon} \xrightarrow{2s} u_0 \right\} = \iint_{\Omega \times Y} f(y, \mathbb{D}u_0) dx dy. \end{aligned}$$

1.2 Preliminaries

This section is devoted to fix notation adopted in the sequel and state preliminary results on Orlicz-Sobolev spaces and homogenization results that will be exploited in the next section. For more details concerning these latter results, for the sake of brevity, we will refer directly to [TN1].

$\Omega \subset \mathbb{R}^d$ ($d \in \mathbb{N}$) denotes a bounded open set with Lipschitz boundary.

1.2.1 Orlicz-Sobolev spaces

Let $B : [0, +\infty[\rightarrow [0, +\infty[$ be an N -function as in [Ad], i.e., B is continuous, convex, with $B(t) > 0$ for $t > 0$, $\frac{B(t)}{t} \rightarrow 0$ as $t \rightarrow 0$, and $\frac{B(t)}{t} \rightarrow \infty$ as $t \rightarrow \infty$.

Equivalently, B is of the form $B(t) = \int_0^t b(\tau) d\tau$, where $b : [0, +\infty[\rightarrow [0, +\infty[$ is non decreasing, right continuous, with $b(0) = 0$, $b(t) > 0$ if $t > 0$ and $b(t) \rightarrow +\infty$ if $t \rightarrow +\infty$. We denote by \tilde{B} , the Fenchel's conjugate, also called the complementary N -function of B defined by

$$\tilde{B}(t) = \sup_{s \geq 0} \{st - B(s)\}, \quad t \geq 0.$$

It can be proven that (see [TN1, Lemma 2.1]) if B is an N -function and \tilde{B} is its conjugate, then for all $t > 0$, it results

$$\begin{aligned} \frac{tb(t)}{B(t)} &\geq 1 \quad (> 1 \text{ if } b \text{ is strictly increasing}), \\ \tilde{B}(b(t)) &\leq tb(t) \leq B(2t). \end{aligned}$$

An N -function B is of class Δ_2 (denoted $B \in \Delta_2$) if there are $\alpha > 0$ and $t_0 \geq 0$ such that $B(2t) \leq \alpha B(t)$ for all $t \geq t_0$.

In all what follows B and \tilde{B} are conjugates N -functions satisfying the delta-2 (Δ_2) condition and c refers to a constant that may vary from line to line.

The Orlicz-space $L^B(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{C} \text{ measurable, } \lim_{\delta \rightarrow 0^+} \int_{\Omega} B(\delta |u(x)|) dx = 0 \right\}$ is a Banach space for the Luxemburg norm:

$$\|u\|_{B,\Omega} = \inf \left\{ k > 0 : \int_{\Omega} B\left(\frac{|u(x)|}{k}\right) dx \leq 1 \right\} < +\infty.$$

It follows that: $C_c^\infty(\Omega)$ is dense in $L^B(\Omega)$, $L^B(\Omega)$ is separable and reflexive, the dual of $L^B(\Omega)$ is identified with $L^{\tilde{B}}(\Omega)$, and the norm induced on $L^{\tilde{B}}(\Omega)$ as a dual space is equivalent to $\|\cdot\|_{\tilde{B},\Omega}$.

Analogously one can define the Orlicz-Sobolev functional space as follows:

$$W^1 L^B(\Omega) = \left\{ u \in L^B(\Omega) : \frac{\partial u}{\partial x_i} \in L^B(\Omega), 1 \leq i \leq d \right\},$$

where derivatives are taken in the distributional sense on Ω . Endowed with the norm $\|u\|_{W^1 L^B(\Omega)} = \|u\|_{B,\Omega} + \sum_{i=1}^d \left\| \frac{\partial u}{\partial x_i} \right\|_{B,\Omega}$, $W^1 L^B(\Omega)$ is a reflexive Banach space. We denote by $W_0^1 L^B(\Omega)$, the closure of $C_c^\infty(\Omega)$ in $W^1 L^B(\Omega)$ and the semi-norm $u \rightarrow \|u\|_{W_0^1 L^B(\Omega)} = \|Du\|_{B,\Omega} = \sum_{i=1}^d \left\| \frac{\partial u}{\partial x_i} \right\|_{B,\Omega}$ is a norm on $W_0^1 L^B(\Omega)$ equivalent to $\|\cdot\|_{W^1 L^B(\Omega)}$.

1.2.2 Homogenization

In order to deal with periodic integrands we will adopt the following notation.

Let $Y := (0, 1)^d$. The letter ε throughout will denote a family of positive real numbers converging to 0. The set \mathbb{R}_y^d will denote \mathbb{R}^d , but the subscript y emphasizes the fact that this is the set where the space variable y is. We also define

$$C_{\text{per}}(Y) = \{v \in C(\mathbb{R}_y^d) : Y\text{-periodic}\},$$

and

$$L_{\text{per}}^B(Y) := \{v \in L_{\text{loc}}^B(\mathbb{R}_y^N) : Y\text{-periodic}\}.$$

Moreover we observe that $L_{\text{per}}^B(Y)$ is a Banach space under the Luxemburg norm $\|\cdot\|_{B,Y}$, and $C_{\text{per}}(Y)$ is dense in $L_{\text{per}}^B(Y)$ (see [TN1, Lemma 2.1]).

For $v \in L_{\text{per}}^B(Y)$ let

$$v^\varepsilon(x) = v\left(\frac{x}{\varepsilon}\right), x \in \mathbb{R}^d.$$

Given $v \in L_{\text{loc}}^B(\Omega \times \mathbb{R}_y^N)$ and $\varepsilon > 0$, we put

$$v^\varepsilon(x) = v\left(x, \frac{x}{\varepsilon}\right), x \in \mathbb{R}^d \text{ whenever it makes sense.}$$

We define the vector space

$$L^B(\Omega \times Y_{\text{per}}) := \{u \in L^B_{\text{loc}}(\Omega \times \mathbb{R}_y^N) : \text{for a.e. } x \in \Omega, u(x, \cdot) \text{ is } Y\text{-periodic}\}.$$

and observe that the embedding $L^B(\Omega, C_{\text{per}}(Y)) \rightarrow L^B(\Omega \times Y_{\text{per}})$ is continuous.

Moreover we will make use of the space

$$W^1 L^B_{\text{per}}(Y) := \left\{ u \in W^1 L^B_{\text{loc}}(\mathbb{R}_y^N) : u, \frac{\partial u}{\partial x_i}, i = 1, \dots, N, Y\text{-periodic} \right\}$$

where the derivative $\frac{\partial u}{\partial x_i}$ is taken in the distributional sense on \mathbb{R}_y^N , and we endow it with the norm $\|u\|_{W^1 L^B_{\text{per}}} = \|u\|_{B,Y} + \sum_{i=1}^N \left\| \frac{\partial u}{\partial x_i} \right\|_{B,Y}$, which makes it a Banach space.

We also consider the space

$$W^1_{\#} L^B_{\text{per}}(Y) = \left\{ u \in W^1 L^B_{\text{per}}(Y) : \int_Y u(y) dy = 0 \right\},$$

and we endow it with the gradient norm

$$\|u\|_{W^1_{\#} L^B_{\text{per}}(Y)} = \sum_{i=1}^N \left\| \frac{\partial u}{\partial x_i} \right\|_{B,Y}.$$

Denoting by $C_{\text{per}}^\infty(Y) = C_{\text{per}}(Y) \cap C^\infty(\mathbb{R}^N)$, and recalling that the space $C_{\#, \text{per}}^\infty(Y; \mathbb{R}) = \{u \in C_{\text{per}}^\infty(Y; \mathbb{R}) : \int_Y u(y) dy = 0\}$ is dense in $W^1_{\#} L^B_{\text{per}}(Y)$, one can deduce (cf. [TN1]) the density of the embedding

$$C_c^\infty(\Omega; \mathbb{R}) \otimes C_{\#, \text{per}}^\infty(Y; \mathbb{R}) \subseteq L^B(\Omega; W^1_{\#} L^B_{\text{per}}(Y)). \quad (1.5)$$

In [TN1] the notion of two-scale convergence introduced by [N] and developed by [A] (see also, among a wide literature, [CDG], [FZ], [Ne], [V] for further developments and related notions like periodic unfolding method), has been extended to the Orlicz-Sobolev setting.

Definition 1. A sequence of functions $(u_\varepsilon)_\varepsilon$ in $L^B(\Omega)$ is said to be:

-weakly two-scale convergent in $L^B(\Omega)$ to a function $u_0 \in L^B(\Omega \times Y_{\text{per}})$ if for every $\varepsilon \rightarrow 0$, we have

$$\int_\Omega u_\varepsilon f^\varepsilon dx \rightarrow \iint_{\Omega \times Y} u_0 f dx dy, \text{ for all } f \in L^{\tilde{B}}(\Omega; C_{\text{per}}(Y)) \quad (1.6)$$

-strongly two-scale convergent in $L^B(\Omega)$ to $u_0 \in L^B(\Omega \times Y_{\text{per}})$ if for $\eta > 0$ and $f \in L^{\tilde{B}}(\Omega; C_{\text{per}}(Y))$ verifying $\|u_0 - f\|_{L^B(\Omega \times Y)} \leq \frac{\eta}{2}$ there exist $\rho > 0$ such that $\|u_\varepsilon - f^\varepsilon\|_{L^B(\Omega)} \leq \eta$ for all $0 < \varepsilon \leq \rho$.

When (1.6) happens for all $f \in L^{\tilde{B}}(\Omega; C_{per}(Y))$ we denote it by " $u_\varepsilon \xrightarrow{2s} u_0$ in $L^B(\Omega)$ weakly" or simply " $u_\varepsilon \xrightarrow{2s} u_0$ in $L^B(\Omega)$ two-scale weakly" and we will say that u_0 is the weak two-scale limit in $L^B(\Omega)$ of the sequence $(u_\varepsilon)_\varepsilon$. In order to denote strong two scale convergence of $u_\varepsilon \rightarrow u_0$ we adopt the symbol $\|u_\varepsilon - u_0\|_{2s-L^B(\Omega \times Y)} \rightarrow 0$.

The following result, whose proof can be found in [TN1], allows to extend the notion of weak two-scale convergence at Orlicz-Sobolev functions, guaranteeing, at the same time, a compactness result.

Proposition 1. *Let Ω be a bounded open set in \mathbb{R}^d and let $(u_\varepsilon)_\varepsilon$ be bounded in $W^1 L^B(\Omega)$. There exist a subsequence, still denoted in the same way, and $u \in W^1 L^B(\Omega)$, $u_1 \in L^1(\Omega; W_\#^1 L_{per}^B(Y))$ such that:*

- (i) $u_\varepsilon \xrightarrow{2s} u$ in $L^B(\Omega)$,
- (ii) $D_{x_i} u_\varepsilon \xrightarrow{2s} D_{x_i} u + D_{y_i} u_1$ in $L^B(\Omega)$, $1 \leq i \leq d$.

In the sequel we denote by $u_0(x, y)$ the function $u(x) + u_1(x, y)$, and by $\mathbb{D}u_0$ the vector $Du + D_y u_1$.

For the sake of brevity, we cannot explicitly quote all the results used throughout the paper but we will refer to [TN1] for further necessary properties of Orlicz-Sobolev spaces, two-scale convergence and homogenization in the Orlicz setting.

1.3 Proof of Theorem 1

This section is devoted to the proof of Theorem 1. To this aim recall the definition of f_{hom} given by (1.4).

Proof (of Theorem 1). We start observing that the coercivity assumptions on f , the compactness result, given by Proposition 1, and (1.1) guarantee that for every $u_\varepsilon \rightharpoonup u \in W^1 L^B(\Omega)$

$$\liminf_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_\varepsilon\right) dx \geq \iint_{\Omega \times Y} f(y, \mathbb{D}u_0) dx dy, \quad (1.7)$$

where $u_0(x, y) = u(x) + u_1(x, y)$ is the weak two scale limit of $(u_\varepsilon)_\varepsilon$. Clearly passing to the infimum on both sides of the equation (1.7), and recalling that $\mathbb{D}u_0 = Du + D_y u_1$, we obtain

$$\begin{aligned} & \inf \left\{ \liminf_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_\varepsilon\right) dx : u_\varepsilon \rightharpoonup u \text{ in } W^1 L^B(\Omega) \right\} \\ & \geq \inf \left\{ \iint_{\Omega \times Y} f(y, Du + D_y u_1) dx dy : u_1 \in L^1(\Omega; W_\#^1 L_{per}^B(Y)) \right\} = \int_{\Omega} f_{\text{hom}}(Du) dx, \end{aligned}$$

where one can replicate the same proof as [CDDA, Lemma 2.2] replacing t by 1 and f_1 by f_{hom} in (1.4) therein and exploit the convexity of f to replace functions with null boundary datum on ∂Y , with periodic ones (see also end of [Ne, Chapter 3]).

The upper bound exploits an argument very similar to the one presented in [Ne], relying, in the present context, on the density result in (1.5). Indeed we can first observe that, as in [TN1, Corollary 5.1] for any given $u \in C^\infty(\overline{\Omega})$ and $\phi_1 \in C_c^\infty(\Omega) \otimes C_{\text{per}}^\infty(Y)$ it results that, given $\phi_\varepsilon(x) := u(x) + \varepsilon\phi_1\left(x, \frac{x}{\varepsilon}\right)$

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, D\phi_\varepsilon(x)\right) dx = \iint_{\Omega \times Y} f(y, Du + D_y\phi_1(x, y)) dx dy. \quad (1.8)$$

On the other hand, given $u \in W^1L^B(\Omega)$ and $u_1 \in L^B(\Omega; W_{\#}^1L_{\text{per}}^B(Y))$, (1.5) guarantees that for each $\delta > 0$ we can find maps $u_\delta \in C^\infty(\overline{\Omega})$ and $v_\delta \in C_c^\infty(\Omega; C_{\text{per}}^\infty(Y))$ such that

$$\|u - u_\delta\|_{W^1L^B(\Omega)} + \|u_1 - v_\delta\|_{L^B(\Omega; W^1L_{\text{per}}^B(Y))} \leq \delta \quad (1.9)$$

Next defining, for every δ , and for every $x \in \Omega$,

$$u_{\delta, \varepsilon}(x) := u_\delta(x) + \varepsilon v_\delta^\varepsilon(x),$$

one has

$$Du_{\delta, \varepsilon}(x) = Du_\delta(x) + \varepsilon D_x v_\delta\left(x, \frac{x}{\varepsilon}\right) + D_y v_\delta\left(x, \frac{x}{\varepsilon}\right).$$

Clearly, as $\varepsilon \rightarrow 0$, it results

$$\begin{aligned} u_{\delta, \varepsilon} &\rightarrow u \text{ in } L^B(\Omega), \\ Du_{\delta, \varepsilon}(x) &\xrightarrow{2s} Dv_\delta(x) + D_y v_\delta(x, y) \text{ strongly in } L^B(\Omega \times Y_{\text{per}}). \end{aligned}$$

Now, considering the following quantity and the notations introduced at page 6,

$$c_{\delta, \varepsilon} := \|u_{\delta, \varepsilon} - u\|_{W^1L^B(\Omega)} + \|Du_{\delta, \varepsilon} - Du + D_y u_1\|_{2s-L^B(\Omega \times Y)}, \quad (1.10)$$

it is easily seen that

$$\lim_{\delta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} c_{\delta, \varepsilon} = 0,$$

which allows us to apply H. Attouch Diagonalization Lemma, thus detecting a sequence $\delta(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$, such that $c_{\delta(\varepsilon), \varepsilon} \rightarrow 0$ and $u_{\delta(\varepsilon), \varepsilon} \rightarrow u$ in $L^B(\Omega)$, with

$$Du_{\delta(\varepsilon), \varepsilon}(x) \xrightarrow{2s} Du(x) + D_y u_1(x, y) \text{ strongly in } L^B(\Omega \times Y_{\text{per}}).$$

This latter convergence, and [TN1, Remark 4.1] ensure that $Du_{\delta(\varepsilon), \varepsilon} \rightharpoonup Du$ weakly in $L^B(\Omega)$, thus, (1.8), the continuity of f in the second variable, (1.2), guarantee that for every $u_1 \in L^B(\Omega; W_{\#}^1L_{\text{per}}^B(Y))$,

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\delta(\varepsilon), \varepsilon}\right) dx = \iint_{\Omega \times Y} f(y, Du + D_y u_1(x, y)) dx dy.$$

as desired.

Thus we can conclude that

$$\begin{aligned} & \inf \left\{ \limsup_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\varepsilon}\right) dx : u_{\varepsilon} \rightharpoonup u \text{ in } W^1 L^B(\Omega) \right\} \\ & \leq \lim_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\delta(\varepsilon), \varepsilon}\right) dx \leq \iint_{\Omega \times Y} f(y, Du(x) + D_y u_1(x, y)) dx dy. \end{aligned}$$

Hence

$$\begin{aligned} & \inf \left\{ \limsup_{\varepsilon \rightarrow 0} \int_{\Omega} f\left(\frac{x}{\varepsilon}, Du_{\varepsilon}\right) dx : u_{\varepsilon} \rightharpoonup u \text{ in } W^{1,B}(\Omega; \mathbb{R}) \right\} \\ & \leq \inf \left\{ \iint_{\Omega \times Y} f(y, Du + D_y u_1) dx dy : u_1 \in L^B(\Omega; W_{\#}^1 L^B_{\text{per}}(Y)) \right\}. \end{aligned}$$

This, together with the last equality in (1.7) concludes the proof.

Acknowledgements The first and the third author acknowledge the support of the Programme ICTP-INdAM research in pairs 2018. Joel Fotso Tachago thanks Dipartimento di Ingegneria Industriale at University of Salerno for its hospitality. Elvira Zappale is a member of GNAMPA-INdAM, whose support is gratefully acknowledged.

References

- Ad. Adams R. A., Sobolev spaces, Pure and Applied Mathematics, Vol. 65, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975, xviii+268.
- A. Allaire, G., Two-scale convergence: a new method in periodic homogenization, Non-linear partial differential equations and their applications. Collège de France Seminar, Vol. XII (Paris, 1991–1993), Pitman Res. Notes Math. Ser., **302**, 1–14, Longman Sci. Tech., Harlow, 1994.
- CDDA. Cioranescu, D., Damlamian, A., and De Arcangelis, R., Homogenization of quasiconvex integrals via the periodic unfolding method., *SIAM J. Math. Anal.*, **37**, (2006), 5, 1435–1453.
- CDG. Cioranescu, D. Damlamian, A., and Griso G., The periodic unfolding method in homogenization, *SIAM J. Math. Anal.*, **40**, (2008), n. 4, 1585-1620.
- TN1. Fotso Tachago, J., and Nnang, H. Two-scale convergence of integral functionals with convex, periodic and nonstandard growth integrands, *Acta Appl. Math.*, **121**, (2012), 175–196.
- FZ. Fonseca, I, and Zappale E., Multiscale relaxation of convex functionals, *J. Convex Anal.*, **10**, (2003), n. 2, 325–350.
- Ne. Neukamm, S. Ph.d Dissertation
- N. Nguentseng, A general convergence result for a functional related to the theory of homogenization. *SIAM J. Math. Anal.* **20**, (1989), 608-623.
- V. Visintin, A. Towards a two-scale calculus, *ESAIM Control Optim. Calc. Var.*, **12**, (2006), n. 3, 371–397.