

Light vector mesons (ω , ρ and ϕ) in strong magnetic fields: A QCD sum rule approach

Amruta Mishra,^{*} Ankit Kumar,[†] Pallabi Parui,[‡] and Sourodeep De[§]

*Department of Physics, Indian Institute of Technology,
Delhi, Hauz Khas, New Delhi – 110 016, India*

Abstract

The mass modifications of the light vector mesons (ω , ρ and ϕ) are investigated in asymmetric nuclear matter in the presence of strong magnetic fields, using a QCD sum rule approach. These are computed from the medium modifications of the non-strange and strange light quark condensates as well as scalar gluon condensate. The quark and gluon condensates are calculated from the medium changes of the scalar fields (non-strange and strange) and a scalar dilaton field in the magnetized nuclear matter, within a chiral SU(3) model. The scalar dilaton field within the model breaks the scale invariance of QCD and simulates the gluon condensate. The anomalous magnetic moments for the nucleons are taken into account in the present study.

^{*}Electronic address: amruta@physics.iitd.ac.in

[†]Electronic address: ankitchahal17795@gmail.com

[‡]Electronic address: pallabiparui123@gmail.com

[§]Electronic address: sourodeepde2015@gmail.com

I. INTRODUCTION

The study of properties of hadrons under extreme conditions, e.g, high densities and/or temperatures, is an important topic of contemporary research in strong interaction physics. The subject is of relevance to ultrarelativistic heavy ion collision experiments, where the experimental observables of these high energy nuclear collisions are affected by the medium modifications of the hadrons. Furthermore, the heavy colliding nuclei have large isospin asymmetry as the number of the neutrons is much larger than the number of protons of these nuclei. It is thus important to study the effects of isospin asymmetry on the hadron properties. The estimation of huge magnetic fields being created in non-central ultra-relativistic heavy ion collision experiments necessitates the study of magnetic field effects on the properties of the hadrons. The medium modifications of the hadrons have been studied extensively in the literature. The different formalisms for these studies are the effective hadronic models, e.g., Quantum Hadrodynamics (QHD) model [1], the QCD sum rule (QCDSR) approach [2, 3], the quark meson coupling (QMC) model[4], the chiral effective models, as well as using the coupled channel approach. The models like Nambu Jona Lasinio model, which simulate the spontaneous chiral symmetry breaking of QCD (through four fermion interactions), have been extensively used in the literature [5–9], to study the strongly interacting matter. The AdS/CFT correspondence and the conjecture of gravity/gauge duality [10] have also been used to study the hadrons [11].

The in-medium masses of the light vector mesons (ρ , ω and ϕ) are studied in the present work using a QCD sum rule approach [12–23], in asymmetric nuclear matter in the presence of strong magnetic fields. The medium modifications are due to the changes of the light quark condensates and gluon condensate in the magnetized isospin asymmetric hadronic matter, which are calculated within mean field approximation, from the changes in the expectation values of non-strange (σ) and strange (ζ) scalar-isoscalar fields, the third component of a scalar isovector field (δ) and a dilaton field (χ) from their vacuum values, within a chiral SU(3) model [24, 25]. The quark condensates, are obtained from the explicit symmetry breaking term within the chiral SU(3) model, in terms of the scalar fields, σ , ζ and δ , and the gluon condensate is related to the dilaton field, χ , which mimics the scale symmetry of QCD, through a logarithmic potential. The model has been used to describe nuclear

matter [24], finite nuclei [25] and the bulk properties of (proto) neutron stars [26]. Using the chiral $SU(3)$ model, the vector mesons have also been studied [27], accounting for the Dirac polarization effects [28]. The model has been used to study the kaons and antikaons in isospin asymmetric nuclear (hyperonic) matter [29–32]. The model has been generalized to the charm and bottom sectors and the in-medium masses of open charm [33–37], open bottom mesons [38, 39], charmonium [36, 40] and bottomonium states [41]. Using the mass modifications of the charmonium states and the open charm mesons, the partial decay widths of the charmonium states to $D\bar{D}$ pair in the hadronic medium have been studied using 3P_0 model [36, 42, 43]. The in-medium partial decay widths of the charmonium (bottomonium) to $D\bar{D}$ ($B\bar{B}$) [44, 45] have also been studied using a field theoretic model for composite hadrons, from the mass modifications of these heavy flavour mesons calculated within the chiral effective model. The effects of magnetic fields on these heavy flavour mesons (D , B , charmonium and bottomonium states) in asymmetric nuclear matter have also been studied [46–50], by including the coupling terms with the electromagnetic field to the baryons in the Lagrangian density of the chiral effective model. The masses of these heavy flavour mesons have been studied accounting for the effects of the anomalous magnetic moments of the nucleons. The light vector mesons (ω , ρ and ϕ) in strange hadronic matter have been studied using a QCD sum rule approach [51], using the medium dependent quark and gluon condensates, calculated within the chiral $SU(3)$ model. In the present investigation, we compute the medium modifications of these vector mesons in asymmetric nuclear matter in the presence of strong magnetic fields using QCD sum rule approach, with the quark and gluon condensates obtained from the scalar fields and dilaton field within the chiral model.

The outline of the paper is as follows : In section II, we describe briefly the chiral $SU(3)$ model used to calculate the quark and gluon condensates in the nuclear medium in the presence of strong magnetic fields. The in-medium values of these condensates are calculated from the medium changes of the scalar fields of the explicit symmetry breaking term and of the dilaton field, which mimics the gluon condensate of QCD in the chiral $SU(3)$ model. In section III, we present the QCD sum rule approach using which the in-medium masses of the light vector mesons (ω , ρ , ϕ) are studied. Section IV discusses the results of the mass modifications of the light vector mesons (ω , ρ and ϕ) in the magnetized asymmetric

nuclear matter. In section V, we summarize the findings of the present investigation.

II. THE HADRONIC CHIRAL $SU(3) \times SU(3)$ MODEL

We use an effective chiral $SU(3)$ model [25] to obtain the in-medium quark and gluon condensates for the study of modifications of the masses of the light vector mesons using a QCD sum rule approach. The model is based on the nonlinear realization of chiral symmetry [52–54] and broken scale invariance [24, 25, 27]. The concept of broken scale invariance leading to the trace anomaly in QCD, $\theta_\mu^\mu = \frac{\beta_{QCD}}{2g} G_{\mu\nu}^a G^{\mu\nu a}$, where $G_{\mu\nu}^a$ is the gluon field strength tensor of QCD, is simulated in the effective Lagrangian at tree level through the introduction of the scale breaking terms [55, 56]

$$\mathcal{L}_{scalebreaking} = -\frac{1}{4}\chi^4 \ln\left(\frac{\chi^4}{\chi_0^4}\right) + \frac{d}{3}\chi^4 \ln\left(\left(\frac{\sigma^2 \zeta}{\sigma_0^2 \zeta_0}\right)\left(\frac{\chi}{\chi_0}\right)^3\right). \quad (1)$$

The Lagrangian density corresponding to the dilaton field, χ leads to the trace of the energy momentum tensor as [40, 51, 57]

$$\theta_\mu^\mu = \chi \frac{\partial \mathcal{L}}{\partial \chi} - 4\mathcal{L} = -(1-d)\chi^4. \quad (2)$$

Equating the trace of the energy momentum tensor arising from the trace anomaly of QCD with that of the present chiral model given by equation (2), gives the relation of the dilaton field to the scalar gluon condensate. The trace of the energy momentum tensor in QCD is given as [58],

$$T_\mu^\mu = \sum_{q_i=u,d,s} m_{q_i} \bar{q}_i q_i + \left\langle \frac{\beta_{QCD}}{2g} G_{\mu\nu}^a G^{\mu\nu a} \right\rangle \equiv -(1-d)\chi^4. \quad (3)$$

In the above, the first term of the energy-momentum tensor, within the chiral $SU(3)$ model is the negative of the explicit chiral symmetry breaking term. This relates the light quark condensates to the values of the scalar fields σ , ζ and δ in the mean field approximation as [51]

$$\begin{aligned} m_u \langle \bar{u}u \rangle &= \frac{1}{2} m_\pi^2 f_\pi (\sigma + \delta), \\ m_d \langle \bar{d}d \rangle &= \frac{1}{2} m_\pi^2 f_\pi (\sigma - \delta), \\ m_s \langle \bar{s}s \rangle &= \left(\sqrt{2} m_k^2 f_k - \frac{1}{\sqrt{2}} m_\pi^2 f_\pi \right) \zeta. \end{aligned} \quad (4)$$

Using the QCD β function occurring in the right hand side of equation (3) at one loop order, for $N_c = 3$ colors and $N_f = 3$ flavors, one gets the dilaton field related to the scalar gluon condensate as

$$\left\langle \frac{\alpha_s}{\pi} G^a_{\mu\nu} G^{a\mu\nu} \right\rangle = \frac{8}{9} \left[(1-d)\chi^4 + \left(m_\pi^2 f_\pi \sigma + \left(\sqrt{2} m_k^2 f_k - \frac{1}{\sqrt{2}} m_\pi^2 f_\pi \right) \zeta \right) \right]. \quad (5)$$

The coupled equations of motion for the non-strange scalar isoscalar field σ , scalar isovector field, δ , the strange scalar field ζ , and the dilaton field χ , derived from the Lagrangian density of the chiral SU(3) model, are solved to obtain the values of these fields in the asymmetric nuclear medium in the presence of magnetic field.

III. QCD SUM RULE APPROACH

In the present section, we briefly describe the QCD sum rule approach used to study the properties of the light vector mesons (ω , ρ , ϕ) in the nuclear medium in presence of a magnetic field. The study uses the values of quark and gluon condensates in the magnetized nuclear medium obtained in a chiral SU(3) model as described in the previous section. The current current correlation function for the vector meson, $V(=\omega, \rho, \phi)$ is written as

$$\Pi_{\mu\nu}^V(q) = i \int d^4x e^{iq \cdot x} \langle 0 | T j_\mu^V(x) j_\nu^V(0) | 0 \rangle, \quad (6)$$

where T is the time ordered product and J_μ^V is the current for the vector meson, $V = \rho, \omega, \phi$. Current conservation gives the structure of the correlation function as

$$\Pi_{\mu\nu}^V(q) = \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) \Pi^V(q^2). \quad (7)$$

In large space-like region, $Q^2 = -q^2 \gg 1 \text{ GeV}^2$, the scalar correlation function $\Pi^V(q^2)$ for the light vector mesons (ω , ρ and ϕ) can be written in terms of the operator product expansion (OPE) as [17, 19]

$$12\pi^2 \tilde{\Pi}^V(q^2 = -Q^2) = d_V \left[-c_0^V \ln \left(\frac{Q^2}{\mu^2} \right) + \frac{c_1^V}{Q^2} + \frac{c_2^V}{Q^4} + \frac{c_3^V}{Q^6} + \dots \right] \quad (8)$$

where, $\tilde{\Pi}^V(q^2 = -Q^2) = \frac{\Pi^V(q^2 = -Q^2)}{Q^2}$, $d_V = 3/2, 1/6$ and $1/3$ for ρ, ω and ϕ mesons respectively, and, μ is a scale chosen to be 1 GeV in the present investigation [51]. The leading term in the

OPE, given by the first term, is calculated in the perturbative QCD. The coefficients c_i^V ($i = 1, 2, 3$) in the OPE contain the informations of the nonperturbative effects of QCD in terms of the quark and gluon condensates, as well as, of the Wilson coefficients [14, 16]. The Wilson coefficients are taken as medium independent, with all the medium effects incorporated into the quark and gluon condensates [14, 16, 17, 19, 20]. After Borel transformation, the correlator for the vector meson can be written as

$$12\pi^2\tilde{\Pi}^V(M^2) = d_V \left[c_0^V M^2 + c_1^V + \frac{c_2^V}{M^2} + \frac{c_3^V}{2M^4} \right] \quad (9)$$

On the phenomenological side the correlator function, $\tilde{\Pi}^V(Q^2)$ can be written as

$$12\pi^2\tilde{\Pi}_{phen}^V(Q^2) = \int_0^\infty ds \frac{R_{phen}^V(s)}{s + Q^2} \quad (10)$$

where $R_{phen}^V(s)$ is the spectral density proportional to the imaginary part of the correlator

$$R_{phen}^V(s) = 12\pi \text{Im} \Pi_{phen}^V(s). \quad (11)$$

On Borel transformation, equation (10) reduces to

$$12\pi^2\tilde{\Pi}^V(M^2) = \int_0^\infty ds e^{-s/M^2} R_{phen}^V(s) \quad (12)$$

Equating the correlation functions from the phenomenological side given by equation (12) to that from the operator product expansion given by equation (9), we obtain,

$$\int_0^\infty ds e^{-s/M^2} R_{phen}^V(s) + 12\pi^2\Pi^V(0) = d_V \left[c_0^V M^2 + c_1^V + \frac{c_2^V}{M^2} + \frac{c_3^V}{2M^4} \right], \quad (13)$$

where the second term in the left hand side of the above equation is the contribution due to scattering of the vector meson with the baryons in the hadronic medium. In the nuclear medium as is the case of the present work of the study of in-medium masses of vector meson, $\Pi^V(0) = \frac{\rho_B}{4M_N}$ for $V=\omega, \rho$. Further, $\Pi^V(0)$ vanishes for ϕ meson, since the ϕ -meson nucleon coupling is zero in the parametrization for the vector meson–baryon interactions within the chiral SU(3) model [25]. This is a reasonable assumption due to the fact that the ϕ -meson nucleon scattering amplitude is negligibly small as compared to the ω -nucleon as well as ρ -nucleon scattering amplitudes [13, 17, 59, 60]. The spectral density is assumed to be of the form of a resonance part $R_{phen}^{V(res)}(s)$ and a perturbative continuum as [17, 51]

$$R_{phen}^V(s) = R_{phen}^{V(res)}(s)\theta(s_0^V - s) + d_V c_0^V \theta(s - s_0^V) \quad (14)$$

For $M > \sqrt{s_0^V}$, the exponential function in the integral of the left hand side of the equation (13) is expanded in powers of s/M^2 for $s < s_0^V$ and one obtains the Finite energy sum rules (FESR) [17] by equating the powers in $1/M^2$ of both sides of equation (13). In the hadronic medium, the FESRs are obtained as [51]

$$F_V^* = d_V(c_0^V s_0^{*V} + c_1^V) - 12\pi^2 \Pi^V(0) \quad (15)$$

$$F_V^* m_V^{*2} = d_V \left(\frac{(s_0^{*V})^2 c_0^V}{2} - c_2^{*V} \right) \quad (16)$$

$$F_V^* m_V^{*4} = d_V \left(\frac{(s_0^{*V})^3 c_0^V}{3} + c_3^{*V} \right) \quad (17)$$

The coefficient c_2^{*V} contains the quark and gluon condensates in the medium, and c_3^{*V} corresponds to the four quark condensate, which is calculated using a factorization method [3] along with a parameter $\kappa_i, i = u, d, s$ which measures the deviation from exact factorization ($\kappa_i=1$). For the nonstrange vector mesons, ρ and ω , these coefficients are given as

$$c_0^{(\rho,\omega)} = 1 + \frac{\alpha_s(Q^2)}{\pi}, \quad c_1^{(\rho,\omega)} = -3(m_u^2 + m_d^2) \quad (18)$$

$$c_2^{*(\rho,\omega)} = \frac{\pi^2}{3} \left\langle \frac{\alpha_s}{\pi} G^{\mu\nu} G_{\mu\nu} \right\rangle + 4\pi^2 \langle m_u \bar{u}u + m_d \bar{d}d \rangle \quad (19)$$

$$c_3^{*(\rho,\omega)} = -\alpha_s \pi^3 \times \frac{448}{81} \kappa_q (\langle \bar{u}u \rangle^2 + \langle \bar{d}d \rangle^2), \quad (20)$$

where, we take $\kappa_u \simeq \kappa_d = \kappa_q$. For ϕ meson, these coefficients are given as [2, 17]

$$c_0^\phi = 1 + \frac{\alpha_s(Q^2)}{\pi}, \quad c_1^\phi = -6m_s^2 \quad (21)$$

$$c_2^{*\phi} = \frac{\pi^2}{3} \left\langle \frac{\alpha_s}{\pi} G^{\mu\nu} G_{\mu\nu} \right\rangle + 8\pi^2 \langle m_s \bar{s}s \rangle \quad (22)$$

$$c_3^{*\phi} = -8\pi^3 \times \frac{224}{81} \alpha_s \kappa_s \langle \bar{s}s \rangle^2. \quad (23)$$

Solving the FESR for vector meson, V (ρ, ω, ϕ) in vacuum, assuming the vacuum mass of the vector meson, determines the value of the coefficient κ_q (κ_s), of the 4-quark condensate, along with the parameters, F_V and s_0^V in vacuum [51]. The equations (15), (16) and (17) are then solved to obtain the medium dependent mass, m_V^* , the scale s_0^{*V} and F_V^* for the vector meson, V , using the value of κ_i as determined from the FESR in vacuum.

| | | $\eta = 0$ | | $\eta = 0.5$ | |
|----|-----|-------------------|--------------------|-------------------|--------------------|
| | | $\rho_B = \rho_0$ | $\rho_B = 2\rho_0$ | $\rho_B = \rho_0$ | $\rho_B = 2\rho_0$ |
| 0 | | 777 | 953.7 | 788.8 | 965.6 |
| 4 | (a) | 773.4 | 950 | 787.24 | 964.2 |
| | (b) | 775.7 | 953.4 | 792.5 | 967.56 |
| 8 | (a) | 772.8 | 947.8 | 787.24 | 964.2 |
| | (b) | 775.6 | 952.4 | 793.74 | 972 |
| 12 | (a) | 772.34 | 947.3 | 787.24 | 964.2 |
| | (b) | 775.87 | 953.34 | 795.3 | 974.1 |

TABLE I: In-medium masses for ω meson in magnetized nuclear matter for densities of ρ_0 and $2\rho_0$, asymmetric parameter, $\eta=0$ and 0.5 and for magnetic fields, eB/m_π^2 as 4,8 and 12. These masses are compared with the in-medium masses of ω meson for zero magnetic field.

IV. RESULTS AND DISCUSSIONS

In this section, we study the in-medium masses of light vector mesons (ω , ρ and ϕ) in strongly magnetized nuclear matter, using the QCD sum rule approach from the light quark condensates and the scalar gluon condensate calculated within a chiral SU(3) model. The broken scale invariance of QCD is incorporated in the effective hadronic model through a scale breaking logarithmic potential in terms of a scalar dilaton field, χ and the gluon condensate in the magnetized nuclear matter is calculated from the medium modification of this scalar dilaton field. In the chiral effective model, the calculations are done in the mean field approximation. In mean field approximation, the meson fields are treated as classical fields. The nonstrange light quark condensates ($\langle\bar{u}u\rangle$, $\langle\bar{d}d\rangle$) and strange condensate ($\langle\bar{s}s\rangle$) in the asymmetric nuclear matter in the presence of magnetic field are calculated within the model from the values of the nonstrange (σ , δ) and strange (ζ) scalar fields within the chiral SU(3) model. For given values of baryon density, ρ_B , the magnetic field, and, the isospin asymmetry parameter, $\eta = (\rho_n - \rho_p)/(2\rho_B)$ (ρ_p and ρ_n are the proton and neutron number densities, respectively), the mean values of the scalar fields, σ , ζ , δ and χ are obtained by solving the coupled equations of motion of these fields. In the absence of a magnetic field, the

| eB/m_π^2 | | $\eta = 0$ | | $\eta = 0.5$ | |
|--------------|-----|-------------------|--------------------|-------------------|--------------------|
| | | $\rho_B = \rho_0$ | $\rho_B = 4\rho_0$ | $\rho_B = \rho_0$ | $\rho_B = 4\rho_0$ |
| 0 | | 622.2 | 391 | 636.3 | 473.3 |
| 4 | (a) | 618 | 374.9 | 634.5 | 468.5 |
| | (b) | 620.7 | 398 | 640.85 | 477.5 |
| 8 | (a) | 617.26 | 335.8 | 634.5 | 468.5 |
| | (b) | 620.65 | 411.7 | 642.3 | 502.8 |
| 12 | (a) | 616.68 | 315.8 | 634.5 | 468.5 |
| | (b) | 620.9 | 433.7 | 644.2 | 527.6 |

TABLE II: In-medium masses for ρ meson in magnetized nuclear matter for densities of ρ_0 and $4\rho_0$, asymmetric parameter, $\eta=0$ and 0.5 and for magnetic fields, eB/m_π^2 as $4,8$ and 12 . These masses are compared with the in-medium masses of ρ meson for zero magnetic field.

in-medium masses of the light vector mesons in asymmetric hadronic matter were studied using QCD sum rule approach and using the gluon and quark condensates calculated within the chiral SU(3) model [51]. In the present work, the medium modifications of the vector meson masses are studied in the presence of a strong magnetic field.

In the present investigation, the values of the current quark masses are taken as $m_u=4$ MeV, $m_d=7$ MeV and $m_s=150$ MeV. The vacuum masses of the ρ , ω and ϕ mesons are taken to be 770 , 783 and 1020 MeV. The coefficients κ_i ($i = q, s$) are solved from the FESRs in vacuum [51] and are obtained as 7.788 , 7.236 and -1.21 for the ω , ρ and ϕ mesons. The difference in the values of κ_q obtained from solving the FESRs of the ω and ρ mesons are due to the difference in their vacuum masses.

In the presence of a magnetic field, the proton has contributions from the Landau energy levels. The anomalous magnetic moments of the nucleons are also taken into consideration in studying the in-medium masses of light vector mesons in nuclear matter in presence of a magnetic field, using the QCDSR approach. The light quark condensates and the scalar gluon condensate, are calculated from the equations (4) and (5), which are then used to calculate the values of the coefficients, c_2^{*V} and c_3^{*V} for the vector mesons, ρ , ω and ϕ as

| eB/m_π^2 | | $\eta = 0$ | | $\eta = 0.5$ | |
|--------------|-----|-------------------|--------------------|-------------------|--------------------|
| | | $\rho_B = \rho_0$ | $\rho_B = 4\rho_0$ | $\rho_B = \rho_0$ | $\rho_B = 4\rho_0$ |
| 0 | | 1001.49 | 998.79 | 1001.79 | 998.38 |
| 4 | (a) | 1001.11 | 998.4 | 1001.6 | 997.85 |
| | (b) | 1001.28 | 998.08 | 1002.03 | 997.74 |
| 8 | (a) | 1000.9 | 998.9 | 1001.6 | 997.85 |
| | (b) | 1001.28 | 997.87 | 1002.14 | 997.5 |
| 12 | (a) | 1001.09 | 999.2 | 1001.6 | 997.85 |
| | (b) | 1001.22 | 997.63 | 1002.28 | 997.38 |

TABLE III: In-medium masses for ϕ meson in magnetized nuclear matter for densities of ρ_0 and $4\rho_0$, asymmetric parameter, $\eta=0$ and 0.5 and for magnetic fields, eB/m_π^2 as 4,8 and 12. These masses are compared with the in-medium masses of ϕ meson for zero magnetic field.

given by equations (19), (20), (22), and (23). Using the values of these coefficients, the in-medium masses of the light vector mesons are calculated by solving the coupled equations (15), (16) and (17) involving the in-medium values F_V^* , s_0^{*V} and m_V^* .

In figures 1 and 2, the in-medium masses of ω mesons are plotted as functions of baryon density, ρ_B (in units of nuclear matter saturation density, ρ_0) for values of magnetic fields, $eB = 4m_\pi^2$ and $eB = 12m_\pi^2$ respectively, with values of isospin asymmetry parameter, η taken to be 0, 0.3 and 0.5. These masses are plotted including the anomalous magnetic moments (AMM) of the nucleons and are compared to the case when the AMMs are not taken into consideration. The masses of the ω meson are also given in Table 1, for the isospin symmetric ($\eta=0$) as well as for the extreme isospin asymmetric case of $\eta=0.5$, for magnetic fields, $eB = 4m_\pi^2$, $8m_\pi^2$, $12m_\pi^2$, (a) without and (b) with the anomalous magnetic moments (AMM) of nucleons taken into account. The ω meson mass is observed to have an initial drop with increase in density, for sub-nuclear densities. This is due to the fact that the contribution of the scattering term, which is proportional to baryon density, ρ_B , is small at low densities, and, the medium modification of the ω meson mass is dominated by modifications of the the light quark condensates in the medium, which leads to a drop in




FIG. 1: (Color online) The mass of ω meson plotted as a function of the baryon density in units of nuclear matter saturation density, for magnetized nuclear matter (for $\eta=0, 0.3, 0.5$) with $eB = 4m_\pi^2$.

the mass of the ω meson. As the density is further increased, the effects of the scattering of the vector meson ω from the nucleons become appreciable, leading to a rise in the ω meson mass. This observed behaviour can be understood from the first two FESRs. The effective




FIG. 2: (Color online) The mass of ω meson plotted as a function of the baryon density in units of nuclear matter saturation density, for magnetized nuclear matter (for $\eta=0, 0.3, 0.5$) with $eB = 12m_\pi^2$.

mass squared of the vector meson, V , is obtained as dividing equation (16) by (15) as

$$m_V^{*2} = \frac{\left(\frac{(s_0^{*V})^2 c_0^V}{2} - c_2^{*V}\right)}{(c_0^V s_0^{*V} + c_1^V) - (1/d_V) \cdot 12\pi^2 \Pi^V(0)}, \quad (24)$$

where, as has already been mentioned, $\Pi^V(0) = \rho_B/(4M_N)$, for $V = \omega, \rho$ and $\Pi^V(0)$ vanishes




FIG. 3: (Color online) The mass of ρ meson plotted as a function of the baryon density in units of nuclear matter saturation density, for magnetized nuclear matter (for $\eta=0, 0.3, 0.5$) with $eB = 4m_\pi^2$.

for ϕ meson as ϕ -nucleon coupling is assumed to be zero in the present work. The scattering term for the non-strange vector mesons, being proportional to the density, becomes appreciable at higher densities. This leads to a smaller value for the denominator and hence to an increase in the mass of ω meson at high densities. This behaviour of the ω mass with density was also observed for the case of zero magnetic field in Ref. [51]. The effects of the




FIG. 4: (Color online) The mass of ρ meson plotted as a function of the baryon density in units of nuclear matter saturation density, for magnetized nuclear matter (for $\eta=0, 0.3, 0.5$) with $eB = 12m_\pi^2$.

magnetic field, which are through the light quark and gluon condensates, are observed to be much smaller as compared to the density effects. The values of the ω meson mass (in MeV) at densities $\rho_0(2\rho_0)$, for the values of the isospin asymmetric parameter, $\eta=0, 0.3$ and 0.5 , are observed to be 773.4 (950), 781.55 (956.86) and 787.24 (964.2) at the value of the




FIG. 5: (Color online) The mass of ϕ meson plotted as a function of the baryon density in units of nuclear matter saturation density, for magnetized nuclear matter (for $\eta=0, 0.3, 0.5$) with $eB = 4m_\pi^2$.

magnetic field as $eB = 4m_\pi^2$, when the anomalous magnetic moments (AMM) of nucleons are not considered, and, 775.7 (953.4), 785.2 (960) and 792.5 (967.56), when AMMs are taken into consideration. The isospin asymmetry effects are observed to be larger for the higher value of the magnetic field, $eB = 12m_\pi^2$, plotted in figure 2. The contribution of the scattering term which is the dominant contribution to the mass of ω meson at high densities,




FIG. 6: (Color online) The mass of ϕ meson plotted as a function of the baryon density in units of nuclear matter saturation density, for magnetized nuclear matter (for $\eta=0, 0.3, 0.5$) with $eB = 12m_\pi^2$.

being proportional to ρ_B , is independent of the isospin asymmetry. This leads to lessening of the isospin asymmetry effects at higher densities, as is evident from the figures 1 and 2. The isospin asymmetry effects on the ω mesons mass are observed only at densities, of around $0.5\rho_0$ to about $2\rho_0$, above which these effects are seen to be diminished as the scattering

effects start becoming important. The isospin asymmetry effects are observed to be more appreciable for the higher value of the magnetic field, $eB = 12m_\pi^2$, as can be seen from figure 2.

The density, isospin asymmetry effects on the masses of the ρ meson are illustrated in figures 3 and 4 for the values of eB as $4m_\pi^2$ and $12m_\pi^2$ respectively. The in-medium ρ meson masses are given in Table 2, for typical values of the magnetic field, density and isospin asymmetry parameters, which are calculated (a) without and (b) with the anomalous magnetic moments (AMM) of the nucleons. The ρ meson mass in the presence of a magnetic field is observed to drop with baryon density, similar to the case of zero magnetic field studied previously [51]. The contribution of the scattering of the ρ meson due to the nucleons in the nuclear matter is small, due to the factor $(1/d_V)$ in this term, which makes the contribution of the Landau scattering term to be 9 times smaller than that of the ω meson, as $(1/d_\rho)/(1/d_\omega) = 1/9$. The effects of the isospin asymmetry are observed to be large at high densities, as was seen for the case of zero magnetic field [51]. The effects of magnetic fields as well as of anomalous magnetic moments are observed to be appreciable at high densities, as can be seen in Table 2. Accounting for the anomalous magnetic moments (AMM) of the nucleons, the values of the ρ meson mass for symmetric nuclear matter ($\eta=0$) at densities $\rho_0(3\rho_0)$, are 620.7 (445.4) and 620.9 (455.9) for $eB = 4m_\pi^2$, and $eB = 12m_\pi^2$ respectively. These values are modified to 632.1 (468.9) and 634.85 (500) at $\eta = 0.3$ and 640.85 (508.7) and 644.2 (545.36) at $\eta=0.5$, for the same magnetic fields. The drop in the ρ meson mass is observed to be smaller when the AMMs of the nucleons are taken into consideration. The isospin asymmetry effect is observed to be much larger for the larger magnetic field, $eB = 12m_\pi^2$, especially at higher densities, as can be seen in figure 4. In Table 2, the values of the ρ meson mass in the presence of magnetic field for the symmetric nuclear matter as well as for asymmetric nuclear matter with $\eta=0.5$, at densities ρ_0 and $4\rho_0$, are compared with the values obtained for zero magnetic field in Ref. [51]. As can be seen from Table 2, in the absence of a magnetic field, the in-medium ρ mass calculated at the nuclear matter saturation density, ρ_0 , in symmetric nuclear matter, is 622.2 MeV [51], which is similar to the value obtained in Ref. [12], using the linear density approximation. The mass of ρ meson at ρ_0 at zero magnetic field, may be compared with the value of 670

MeV obtained using QCD sum rule approach in Ref. [19], and, of around 530 MeV, in an improved QCD sum rule calculation [15]. In the present work, the effects of magnetic field on the ρ meson mass in isospin asymmetric nuclear matter have been investigated.

The in-medium masses of the ϕ meson, plotted in figures 5 and 6 for the values of eB as $4m_\pi^2$ and $12m_\pi^2$ respectively, are observed to decrease sharply with density upto a density of around ρ_0 , above which there is observed to be very less modification in the mass of the ϕ meson. There is no contribution to the ϕ mass from the scattering term. This is due to the fact that the nucleon- ϕ meson coupling is zero in the parameter set chosen for the vector meson-baryon interactions within the chiral SU(3) model. It might be noted here that $\phi \rightarrow K\bar{K}$ is OZI allowed, whereas OZI rule forbids the decay of ϕ meson to pions. There is, however, observed to be violation of OZI rule by about 5% in the $\phi \rightarrow 3\pi$ channel [17, 18]. The decay $\phi \rightarrow 3\pi$ and the decay of $\omega \rightarrow 3\pi$ (through direct decay as well as the two step decay, $\omega \rightarrow \rho\pi$ followed by $\rho \rightarrow 2\pi$) imply that there is mixing between the ϕ meson and the non-strange (ρ, ω) vector mesons. However, the decay of ϕ is dominated by the OZI allowed $\phi \rightarrow K\bar{K}$ and the mass modification of the ϕ meson is predominantly due to the change of the strange condensate in the hadronic medium. The drop in ϕ mass is much smaller than the mass drop of the ρ meson (where the scattering term has negligible contribution) due to the large drop of the nonstrange quark condensate as compared to the drop of the strange quark condensate in the nuclear medium. At higher densities, the strange condensate remains almost constant and hence the ϕ meson mass changes very little with density, whereas the nonstrange quark condensate continues to drop as density is increased, leading to a monotonic drop of the ρ meson mass with density. These behaviours of the ρ and ϕ vector meson masses were also observed for the case of zero magnetic field [51]. The effects of anomalous magnetic moments of the nucleons are observed to be larger at higher densities, even though the magnitudes of these still remain small. The strange quark condensate as well as the scalar gluon condensate have very small effects from isospin asymmetry, leading to the modifications of the ϕ meson mass to be very similar in the isospin symmetric and asymmetric nuclear matter.

V. SUMMARY

In summary, in the present investigation, we have calculated the masses of the light vector mesons (ω , ρ and ϕ) in the nuclear matter in the presence of a strong magnetic field, using QCD sum rule approach, due to the modifications of the light quark and scalar gluon condensates, calculated within a chiral effective model. The medium modifications are due to the changes of the light quark condensates and the scalar gluon condensate in the magnetized isospin asymmetric hadronic matter, which are calculated within mean field approximation, from the changes in the expectation values of the scalar fields (σ , ζ and δ) and a dilaton field (χ) from their vacuum values, respectively, within a chiral SU(3) model. The masses of the ρ mesons are dominantly governed by the nonstrange quark condensates which lead to decrease in these masses with increase in density. For ω meson, the drop in the nonstrange quark condensates leads to a drop in the mass at sub-nuclear matter densities, which however, is overcome by the ω -nucleon scattering term, leading to an increase in the ω mass as the density is further increased. The scattering term which dominates for the ω meson mass at high densities, is of the form $(\rho_p + \rho_n)$ and is thus independent of the isospin asymmetry of the nuclear medium. The mass of ϕ meson is dominated by the behaviour of the strange condensate, $\langle \bar{s}s \rangle$ in the medium, leading to an initial drop with density and the mass changes slightly as the density is further increased. The isospin asymmetry effects are observed to be more for ρ meson as compared to the ω and ϕ mesons. The effects of anomalous magnetic moments of the nucleons are seen to be appreciable at higher densities and higher magnetic fields for the ρ meson mass. The density effects are observed to be the dominant medium effects on the masses of these light vector mesons.

Acknowledgments

One of the authors (AM) is grateful to ITP, University of Frankfurt, for warm hospitality and acknowledges financial support from Alexander von Humboldt foundation when this

work was initiated.

-
- [1] B. D. Serot and J. D. Walecka, *Adv. Nucl. Phys.* **16**, 1 (1986); S. A. Chin, *Ann. Phys. (N.Y.)* **108**, 301 (1977).
 - [2] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, *Nucl. Phys. B* **147**, 385 (1979).
 - [3] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, *Nucl. Phys. B* **147**, 448 (1979).
 - [4] K. Tsushima, D.H. Lu, A.W. Thomas, K. Saito, R.H. Landau, *Phys. Rev. C* **59**, 2824 (1999); A. Sibirtsev, K. Tsushima, A.W. Thomas, *Eur. Phys. J. A* **6**, 351 (1999).
 - [5] S. P. Klevansky, *Rev. Mod. Phys.* **64** 649 (1992).
 - [6] M. Buballa, *Phys. Rep.* **407**, 205 (2004).
 - [7] T. Hatsuda and T. Kunihiro, *Phys. Rep.* **247**, 221 (1994).
 - [8] H. Mishra and S. P. Misra, *Phys. Rev. D* **48**, 5376 (1993); H. Mishra and J. C. Parikh, *Nucl. Phys. A* **679**, 597 (2001).
 - [9] A. Mishra and H. Mishra, *Phys. Rev. D* **69**, 014014 (2004); *ibid*, *Phys. Rev. D* **71**, 074023 (2005).
 - [10] J.M. Maldacena, *Int. J. Theor. Phys.* **38**, 1113 (1999); S. Gubser, I.R. Klebanov and A.M. Polyakov, *Phys. Lett. B* **428**, 105 (1998); E. Witten, *Adv. Theor. Math. Phys.* **2**, 253 (1998).
 - [11] J. Erlich, E. Katz, D.T. Son and M.A. Stephanov, *Phys. Rev. Lett.* **95**, 261602 (2005); A. Karch, E. Katz, D. T. Son and M. A. Stephanov, *Phys. Rev. D* **74** 015005 (2006); T. Sakai and S. Sugimoto, *Prog. Theor. Phys.* **113**, 843 (2005); G.F. de Teramond and S.J. Brodsky, *Phys. Rev. Lett.* **94**, 201601 (2005); D.K. Hong, T. Inami and H.U. Yee, *Phys. Lett. B* **646**, 165 (2007); K. Nawa, H. Suganuma and T. Kojo, *Phys. Rev. D* **75**, 086003 (2007); Danning Li and Mei Huang, *Jour. High Energy Phys.* **11**, 088 (2013).
 - [12] T. Hatsuda, S.H. Lee, *Phys. Rev. C* **46**, R34 (1992).
 - [13] T. Hatsuda, S.H. Lee, H. Shiomi, *Phys. Rev. C* **52**, 3364 (1995).
 - [14] T. Hatsuda, Y. Koike and S. H. Lee, *Nucl. Phys. B* **394**, 221 (1993).
 - [15] M. Asakawa and C. M. Ko, *Nucl. Phys. A* **560**, 399 (1993).
 - [16] S. Zschocke, O. P. Pavlenko and B. Kämpfer, *Eur. Phys. Jour. A* **15**, 529 (2002).

- [17] F.Klingl, N. Kaiser, W. Weise, Nucl. Phys. **A 624**, 527 (1997).
- [18] F.Klingl, N. Kaiser, W. Weise, Z. Phys. **A 356**, 193 (1996).
- [19] Y. Kwon, M. Procura and W. Weise, Phys. Rev. **C 78**, 055203 (2008).
- [20] Y. Kwon, C. Sasaki and W. Weise, Phys. Rev. **C 81**, 065203 (2010).
- [21] C. Adami, T. Hatsuda and I. Zahed, Phys. Rev. **D 43**, 921 (1991).
- [22] R. Thomas, T. Hilger and B. Kaempfer, Nucl. Phys. **A 795**, 19 (2007).
- [23] A.K. Dutt-Mazumder, R. Hofmann, M. Pospelov, Phys. Rev. **C 63**, 015204 (2000).
- [24] A. Mishra, K. Balazs, D. Zschesche, S. Schramm, H. Stöcker and W. Greiner, Phys. Rev. **C 69**, 024903 (2004).
- [25] P. Papazoglou, D. Zschesche, S. Schramm, J. Schaffner-Bielich, H. Stöcker, and W. Greiner, Phys. Rev. **C 59**, 411 (1999).
- [26] A. Mishra, A. Kumar, S. Sanyal, V. Dexheimer, S. Schramm, Eur. Phys. Jour. **A 45**, 169 (2010).
- [27] D. Zschesche, A. Mishra, S. Schramm, H. Stöcker and W. Greiner, Phys. Rev. **C 70**, 045202 (2004).
- [28] H. Shiomi and T. Hatsuda, Phys. Lett. **B 334**, 281 (1994); T. Hatsuda, H. Shiomi, and H. Kuwabara, Prog. Theor. Phys. **95**, 1009 (1996); H.-C. Jean, J. Piekarewicz, and A. G. Williams, Phys. Rev. **C 49**, 1981 (1994); K. Saito, K. Tsushima, A. W. Thomas, and A. G. Williams, Phys. Lett. **B 433**, 243 (1998).
- [29] A. Mishra, E. L. Bratkovskaya, J. Schaffner-Bielich, S. Schramm and H. Stöcker, Phys. Rev. **C 70**, 044904 (2004).
- [30] A. Mishra and S. Schramm, Phys. Rev. **C 74**, 064904 (2006).
- [31] A. Mishra, S. Schramm and W. Greiner, Phys. Rev. **C 78**, 024901 (2008).
- [32] Amruta Mishra, Arvind Kumar, Sambuddha Sanyal, S. Schramm, Eur. Phys. **J. A 41**, 205 (2009).
- [33] A. Mishra, E. L. Bratkovskaya, J. Schaffner-Bielich, S.Schramm and H. Stöcker, Phys. Rev. **C 69**, 015202 (2004).
- [34] Amruta Mishra and Arindam Mazumdar, Phys. Rev. **C 79**, 024908 (2009).
- [35] Arvind Kumar and Amruta Mishra, Phys. Rev. **C 81**, 065204 (2010).

- [36] Arvind Kumar and Amruta Mishra, *Eur. Phys. J. A* **47**, 164 (2011).
- [37] Divakar Pathak and Amruta Mishra, *Adv. High Energy Phys.* 2015, 697514 (2015).
- [38] Divakar Pathak and Amruta Mishra, *Phys. Rev. C* **91**, 045206 (2015).
- [39] Divakar Pathak and Amruta Mishra, *Int. J. Mod. Phys. E* **23**, 1450073 (2014).
- [40] Arvind Kumar and Amruta Mishra, *Phys. Rev. C* **82**, 045207 (2010).
- [41] Amruta Mishra and Divakar Pathak, *Phys. Rev. C* **90**, 025201 (2014).
- [42] A. Le Yaouanc, L. Oliver, O. Pene and J. C. Raynal, *Phys. Rev. D* **8**, 2223 (1973); *ibid*, *Phys. Rev. D* **9**, 1415 (1974); *ibid*, *Phys. Rev. D* **11**, 1272 (1975); T. Barnes, F. E. Close, P. R. Page and E. S. Swanson, *Phys. Rev. D* **55**, 4157 (1997). (2002).
- [43] B. Friman, S. H. Lee and T. Song, *Phys. Lett, B* **548**, 153 (2002).
- [44] Amruta Mishra, S. P. Misra and W. Greiner, *Int. J. Mod. Phys. E* **24**, 155053 (2015).
- [45] Amruta Mishra and S. P. Misra, *Phys. Rev. C* **95**, 065206 (2017).
- [46] Sushruth Reddy P., Amal Jahan C.S., Nikhil Dhale, Amruta Mishra, J. Schaffner-Bielich, *Phys. Rev. C* **97**, 065208 (2018).
- [47] Nikhil Dhale, Sushruth Reddy P., Amal Jahan C.S., Amruta Mishra, *Phys. Rev. C* **98**, 015202 (2018).
- [48] Amal Jahan C.S., Nikhil Dhale, Sushruth Reddy P., Shivam Kesarwani and Amruta Mishra, *Phys. Rev. C* **98**, 065202 (2018).
- [49] Amal Jahan C.S., Shivam Kesarwani, Sushruth Reddy P., Nikhil Dhale, and Amruta Mishra, [arXiv:1807.07572](https://arxiv.org/abs/1807.07572) (nucl-th).
- [50] Pallabi Parui, Ankit Kumar, Sourodeep De, Amruta Mishra, [arXiv: 1811.04622](https://arxiv.org/abs/1811.04622) (nucl-th).
- [51] Amruta Mishra, *Phys. Rev. C* **91**, 035201 (2015).
- [52] S. Weinberg, *Phys. Rev.* **166** 1568 (1968).
- [53] S. Coleman, J. Wess, B. Zumino, *Phys. Rev.* **177** 2239 (1969); C.G. Callan, S. Coleman, J. Wess, B. Zumino, *Phys. Rev.* **177** 2247 (1969).
- [54] W. A. Bardeen and B. W. Lee, *Phys. Rev.* **177** 2389 (1969).
- [55] J. Schechter, *Phys. Rev. D* **21**, 3393 (1980).
- [56] J. Ellis, *Nucl. Phys.* **B 22**, 478 (1970); B. A. Campbell, J. Ellis and K. A. Olive, *Nucl. Phys.* **B 345**, 57 (1990).

- [57] Erik K. Heide, Serge Rudaz and Paul J. Ellis, Nucl. Phys. A **571**, 713 (2001).
- [58] Thomas D. Cohen, R. J. Furnstahl and David K. Griegel, Phys. Rev. C **45**, 1881 (1992).
- [59] A. I. Bochkarev and M. E. Shaposhnikov, Phys. Lett. B **145**, 276 (1984); *ibid*, Nucl. Phys. B **268**, 220 (1986).
- [60] W. Florkowski, W. Broniowski, Nucl. Phys. A **651**, 397 (1999).