Abstract

We develop simple diffusion-advection models to estimate the average time it takes fish to reach one of the boundaries of an enclosure and the population distribution over time moving in the enclosure (such as a lake or slough). We start with a combination of random walks and directed movement and then, from these, proceeding to the associated Partial Differential Equations and their solution. We also find the evolution of the population distribution and communities composition over time moving in the enclosure. Although this model was developed with fish movements in mind it has wide ranging applicability scaling from the molecular to human and action from inert to deliberate.

Thinking Inside the Box: An Advection-Diffusion Model of Animal Movement in an Enclosed Region

Stephen Tennenbaum*, John Gatto and Joel Trexler

Department of Biology, Florida International University

*set1@fiu.edu (corresponding author)

September 18, 2019

1 Introduction

There is a wide range of movement phenomena occurring at scales from the molecular to the human that share certain common features. There is the dispersion of smoke from a chimney in the breeze, the spread of an effluent in a stream, the transport of proteins in the cytoplasm of a cell, taxis of protozoa in pond water responding to dissolved nutrients, movement of animals, the spread of a tree species across a continent, these movements have both directed components and random components. Both of these aspects can occur, at one extreme, as completely passive or at the other, as an act of individual volition. Nevertheless, if these two aspects of movement act at similar scales in space and time then they can be described mathematically at any of these scales using the similar principles. In this paper we derive an advection-diffusion partial differential equation (PDE) from a mixed model based on the hypothetical movements of fish in an enclosed body of water. This model includes a random walk component corresponding to the fish searching its immediate area and a directed movement component corresponding to the fish's response to some environmental cue. The solution of this PDE allows us to relate; (1) the parameters of movement to the average time to arrive at a particular location from any starting point, (2) the probability of a fish being at a particular location at a given time, and comparisons of the movement characteristics of different species of fish (or different individuals) with differing parameter values. This approach has been used in studies fish movements in rivers, streams, and open water situations (see for example [Skalski and Gilliam(2000)], [Sparrevohn et al.(2002)Sparrevohn, Nielsen, and Støttrup], [Faugeras and Maury(2007)], [Kerckhove et al.(2015)Kerckhove, Milne, Shuter, and Abrams].) To our knowledge this is the first time this method is applied to a completely inclosed situation such as a lake, pond, marsh or slough.

We calculate first the average time to arrive at the designated goal starting from any point in the enclosure. This will give us a general idea of how the average individual performs given the particular aspects of its behavior. Next we calculate the probability of being at any point in the enclosure at a particular time. This will be useful for looking at the distribution of individuals and any steady-state. We can also use this information to determine from among different species the probabilities of first arrival at the designated goal (or for that matter, any order of arrival). Finally, we can determine the relative species composition of arrivals and distribution.

Our approach in both of the above cases will be to start with a random walk alternated with directed movement. The derivations of diffusion type equations from random walks can be found in many sources, but the one that we used here is found in Random Walks in Biology [Berg(1993)].



Figure 1: Mixed random walk and directed movement. Movement is directed a fraction of the time p, and random movement a fraction of time 1 - p = q. The average length of time to reach the absorbing boundary from a given point is W(x, y).

2.1 Derivation of equations

We start with the assumptions that movement will be either completely random in two dimensions or completely deterministic in one dimension. The directed movement will occur a fraction of time p and the random movement will occur a fraction of time (1 - p). Random movement will involve steps in only one of the four directions parallel to the boundaries of the (rectangular) enclosure, each occurring with equal probability. Each of the steps will move the individual a fixed distance Δ in a time τ (Figure 1). When the movement is directed, it will be exclusively in the positive x direction toward the designated goal. These steps shall transport the individual a distance $v\tau$ toward the right, where v is the speed of directed movement and τ is our given time increment. Starting at location (x, y), we can write the average length of time to get to the goal from that point as W(x, y). In the next time increment τ the individual will have moved either randomly or directed. If the movement was directed, it will now be at the location $(x + v\tau, y)$ and the average time it takes to get to goal from there will be $W(x + v\tau, y)$. Note that if there were only directed movement then the time from (x, y) would be τ units of time more than from $(x + v\tau, y)$, which is closer to goal, so that $W(x, y) = \tau + W(x + v\tau, y)$. However, these steps only occur with frequency p, the rest of the time the individual is moving in a random direction that will put it either closer $(x + \Delta, y)$ or further $(x - \Delta, y)$ or the same distance but to one side or the other $(x, y \pm \Delta)$ in any event the individual will be at these new locations τ units of time later. So the average time from (x, y) will be the average of all the times from the new locations plus the time τ it took to get there. Additionally, we assume that a fraction s of the population does not move at all in the interval, or alternatively, a given individual rests s fraction of the time. This is written explicitly as

$$W(x,y) = \tau + (1-s)\frac{1-p}{4} [W(x+\Delta,y) + W(x-\Delta,y) + W(x,y+\Delta) + W(x,y-\Delta)] + (1-s)pW(x+v\tau,y) + sW(x,y).$$
(1)

Subtracting W(x, y) from both sides, we obtain

$$0 = \tau + (1 - s) \frac{1 - p}{4} \left[(W(x + \Delta, y) - W(x, y)) - (W(x, y) - W(x - \Delta, y)) + (W(x, y + \Delta) - W(x, y)) - (W(x, y) - W(x, y - \Delta)) \right] + (1 - s) p (W(x + v\tau, y) - W(x, y)).$$

Then dividing through by $\frac{1}{2}\Delta$, we get

$$0 = \frac{2\tau}{(1-s)\Delta} + \frac{1-p}{2} \left[\left(\frac{W(x+\Delta,y) - W(x,y)}{\Delta} \right) - \left(\frac{W(x,y) - W(x-\Delta,y)}{\Delta} \right) \right] \\ + \left(\frac{W(x,y+\Delta) - W(x,y)}{\Delta} \right) - \left(\frac{W(x,y) - W(x,y-\Delta)}{\Delta} \right) \right] \\ + \frac{2pv\tau}{\Delta} \left(\frac{W(x+v\tau,y) - W(x,y)}{v\tau} \right).$$
(2)

Noting that as Δ and τ approach 0, we get (by the definition of the derivative)

$$\lim_{\Delta \to 0} \frac{W\left(x + \Delta, y\right) - W\left(x, y\right)}{\Delta} = \frac{\partial}{\partial x} W\left(x + \Delta, y\right)$$

and
$$\lim_{v\tau \to 0} \frac{W\left(x + v\tau, y\right) - W\left(x, y\right)}{v\tau} = \frac{\partial}{\partial x} W\left(x + v\tau, y\right).$$

Hence,

$$0 = \frac{2\tau}{(1-s)\Delta} + \frac{1-p}{2} \left[\left(\frac{\partial}{\partial x} W(x+\Delta,y) - \frac{\partial}{\partial x} W(x,y) \right) + \left(\frac{\partial}{\partial y} W(x,y+\Delta) - \frac{\partial}{\partial y} W(x,y) \right) \right] + \frac{2pv\tau}{\Delta} \frac{\partial}{\partial x} W(x+\tau v,y).$$

Dividing the terms in the bracket again by Δ , we obtain the second derivative with

$$\lim_{\Delta \to 0} \frac{\frac{\partial}{\partial x} W\left(x + \Delta, y\right) - \frac{\partial}{\partial x} W\left(x, y\right)}{\Delta} = \frac{\partial^2}{\partial x^2} W\left(x, y\right)$$

and
$$\lim_{v\tau \to 0} \frac{\partial}{\partial x} W\left(x + v\tau, y\right) = \frac{\partial}{\partial x} W\left(x, y\right).$$

Thus, we continue as

$$0 = \frac{2\tau}{(1-s)\,\Delta^2} + \frac{1-p}{2}\left(\frac{\partial^2 W}{\partial x^2} + \frac{\partial^2 W}{\partial y^2}\right) + \frac{2pv\tau}{\Delta^2}\frac{\partial W}{\partial x}.$$

The diffusion coefficient is assumed to be a constant, for a 2d random walk $D = \frac{\Delta^2}{4\tau}$ and q = 1 - p, yielding

$$0 = \frac{1}{(1-s)2D} + \frac{q}{2} \left(\frac{\partial^2 W}{\partial x^2} + \frac{\partial^2 W}{\partial y^2} \right) + \frac{pv}{2D} \frac{\partial W}{\partial x}.$$

$$0 = \frac{1}{(1-s)D} + q \left(\frac{\partial^2 W}{\partial x^2} + \frac{\partial^2 W}{\partial y^2} \right) + \frac{pv}{D} \frac{\partial W}{\partial x}.$$
(3)

Rearranging parameters and defining $\phi = \frac{pv}{qD}$ and $\eta = \frac{1}{(1-s)qD}$, we obtain

$$0 = \left(\frac{\partial^2 W}{\partial x^2} + \frac{\partial^2 W}{\partial y^2}\right) + \phi \frac{\partial W}{\partial x} + \eta, \qquad (4)$$

The above equation does not lend itself to a solution by separation of variables due to the presence of the constant η (it is non-homogeneous), so we will take as our ansatz a new function U defined as,

$$U(x,y) = \phi W(x,y) + \eta x + k \tag{5}$$

where k is an arbitrary constant. We then have

$$\frac{\partial}{\partial x}U(x,y) = \phi \frac{\partial}{\partial x}W(x,y) + \eta \quad \text{and} \quad \frac{\partial^2}{\partial x^2}U(x,y) = \phi \frac{\partial^2}{\partial x^2}W(x,y)$$
$$\frac{\partial}{\partial y}U(x,y) = \phi \frac{\partial}{\partial y}W(x,y) \quad \text{and} \quad \frac{\partial^2}{\partial y^2}U(x,y) = \phi \frac{\partial^2}{\partial y^2}W(x,y)$$

Substituting the above into equation (4), we have

$$0 = \frac{1}{\phi} \left(\frac{\partial^2 U(x,y)}{\partial x^2} + \frac{\partial^2 U(x,y)}{\partial y^2} \right) + \frac{\partial U(x,y)}{\partial x}.$$
 (6)

The next step is to try a separable solution for U, that is, U(x, y) = X(x) Y(y) yields

$$\frac{\partial U(x,y)}{\partial x} = Y(y) \frac{\partial X(x)}{\partial x} \quad \text{and} \quad \frac{\partial^2 U(x,y)}{\partial x^2} = Y(y) \frac{\partial^2 X(x)}{\partial x^2}$$
$$\frac{\partial U(x,y)}{\partial y} = X(x) \frac{\partial (y)}{\partial y} Y \quad \text{and} \quad \frac{\partial^2 U(x,y)}{\partial y^2} = X(x) \frac{\partial^2 Y(y)}{\partial y^2}$$
$$0 = \frac{1}{\phi} \left(Y \frac{\partial^2 X}{\partial x^2} + X \frac{\partial^2 Y}{\partial y^2} \right) + Y \frac{\partial X}{\partial x}.$$

Multiplying by ϕ and dividing by XY, we get

$$0 = \left(\frac{1}{X}\frac{\partial^2 X}{\partial x^2} + \frac{1}{Y}\frac{\partial^2 Y}{\partial y^2}\right) + \phi \frac{1}{X}\frac{\partial X}{\partial x},\tag{7}$$

and moving the Y term to the left hand side gives

$$-\frac{1}{Y}\frac{\partial^2 Y}{\partial y^2} = \frac{1}{X}\left(\frac{\partial^2 X}{\partial x^2} + \phi\frac{\partial X}{\partial x}\right).$$
(8)

Since each side of the equality in (8) is independent of the the other side, both sides must be equal to some constant (a standard PDE approach)

$$-\frac{1}{Y}\frac{\partial^2 Y}{\partial y^2} = \lambda = \frac{1}{X}\frac{\partial^2 X}{\partial x^2} + \phi \frac{1}{X}\frac{\partial X}{\partial x}.$$
(9)

Before attacking this equation, we discuss the boundary conditions. We have three reflecting boundaries at x = 0, y = -b/2, and y = b/2. For these we write $\frac{\partial}{\partial x}W(0, y) = 0$, and $\frac{\partial}{\partial x}W(x, \pm b/2) = 0$. At the goal, x = a the boundary condition is "absorbing", that is, the average time to reach the goal is zero. So W(a, y) = 0. Going back to the definition of U, we have

$$U(a, y) = \phi W(a, y) - \eta a + k \quad \text{from equation (5)}$$
(10)
 $X(a) Y(y) = -\eta a + k, \quad \text{since } U(a, y) = X(a) Y(y) \text{ and } W(a, y) = 0.$

Solving for Y, we obtain

$$Y(y) = \frac{-\eta a + k}{X(a)} = C_0, \quad \text{a constant.}$$
(11)

Since the right hand side is composed entirely of constants and the function Y(y) is constant for all x and y, we have $\frac{\partial Y}{\partial y} = 0$ and $\frac{\partial^2 Y}{\partial y^2} = 0$. Therefore, $\lambda = 0$ and we can now write

$$\frac{\partial^2 X}{\partial x^2} + \phi \frac{\partial X}{\partial x} = 0. \tag{12}$$

The associated characteristic equation is $r^2 + \phi r = 0$, which has roots $r_1 = 0$ and $r_2 = -\phi$. This gives the general solution

$$X(x) = C_1 e^{r_1 x} + C_2 e^{r_2 x} = C_1 + C_2 e^{-\phi x}$$
(13)

Rearranging equation (5) to solve for W, we get

$$W(x, y) = \phi^{-1} (U(x, y) - \eta x - k) = \phi^{-1} (X(x) Y(y) - \eta x - k)$$

Substituting for X from (13) and Y from (11), we get

$$W(x,y) = \phi^{-1} \left(\left(C_1 + C_2 e^{-\phi x} \right) C_0 - \eta x - k \right).$$
(14)

Taking the derivative with respect to x, we have

$$\frac{\partial}{\partial x}W(x,y) = \frac{\partial}{\partial x}\left(\phi^{-1}\left(\left(C_1 + C_2 e^{-\phi x}\right)C_0 - \eta x - k\right)\right)$$
$$= -C_0 C_2 e^{-\phi x} - \eta \phi^{-1}.$$
(15)

From our boundary condition at x = 0, we have $\frac{\partial}{\partial x}W(0, y) = 0$. Plugging this into (15) yields

$$\frac{\partial}{\partial x}W(0,y) = 0 = -\eta\phi^{-1} - C_0C_2e^{-\phi_0}$$

$$C_0C_2 = -\eta\phi^{-1}.$$
(16)

Equation (14) then becomes

$$W(x,y) = \phi^{-1} \left(\left(C_1 C_0 + C_2 C_0 e^{-\phi x} \right) - \eta x - k \right) = \phi^{-1} \left(\left(C_1 C_0 - \eta \phi^{-1} e^{-\phi x} \right) - \eta x - k \right) = \phi^{-1} \left(\left(C_1 C_0 - k \right) - \eta \left(\phi^{-1} e^{-\phi x} + x \right) \right).$$
(17)

At x = a we have W(a, y) = 0, i.e.,

$$W(a,y) = 0 = \phi^{-1} \left((C_1 C_0 - k) - \eta \left(\phi^{-1} e^{-\phi a} + a \right) \right)$$

Solving for the constants, we obtain

$$C_0 C_1 - k = \eta \left(\phi^{-1} e^{-\phi a} + a \right).$$
(18)

Finally, substituting (18) into the remaining constants in (17), we get

$$W(x,y) = \phi^{-1} \left((C_0 C_1 - k) - \eta \left(\phi^{-1} e^{-\phi x} + x \right) \right)$$

= $\phi^{-1} \left(\eta \left(\phi^{-1} e^{-\phi a} + a \right) - \eta \left(\phi^{-1} e^{-\phi x} + x \right) \right)$
= $\eta \phi^{-1} \left(a - x + \phi^{-1} e^{-\phi a} - \phi^{-1} e^{-\phi x} \right)$, or

$$W(x,y) = \eta \phi^{-1} (a-x) - \eta \phi^{-2} \left(e^{-\phi x} - e^{-\phi a} \right).$$
(19)

Recalling that $\phi = \frac{pv}{qD}$ and $\eta = \frac{1}{(1-s)qD}$, we obtain

$$W(x,y) = \frac{1}{(1-s)} \left(\frac{(a-x)}{pv} - \frac{qD}{p^2 v^2} \left(e^{-\frac{pvx}{qD}} - e^{-\frac{pva}{qD}} \right) \right).$$
(20)

Note that the solution is independent of the y coordinate. On reflection this makes perfect sense, since the back and side boundaries are reflecting, the time to goal from any point on a line parallel to the goal should be the same. This observation allows us to remark, in hindsight, that since W(x, y) = W(x), then $\partial^2 W(x, y) / \partial y^2 = 0$ with the result that the PDE could have been solved by directly integrating twice and the exact same solution obtained.

Using equation (20) we have the following special cases.

- 1. If the fish search and then move in a directed fashion very slowly then $v \to 0$ and $\lim_{v \to 0} W(x, y) = \frac{a^2 x^2}{(1 s)qD}$.
- 2. If they search all the time q = 1 and don't rest (s = 0), then $W(x, y) = \frac{a^2 x^2}{2D}$.
- 3. If the fish move forward and then search very slowly for q fraction of the time, then $D \to 0$ and $\lim_{D\to 0} W(x, y) = \frac{a-x}{(1-s)pv}$.
- 4. If they move forward without resting or searching, then p = 1, s = 0, and $W(x, y) = \frac{a-x}{v}$

This is not the only approach to this approach to this problem. We will see later that the average time to arrive calculated here is slightly different than the *median* time to arrive calculated from the probability distribution but has the advantage of being relatively simple to calculate. It also allows us to characterize the system as dominated by random or directed moment with a dimensionless index and can even provide a rough estimate of the diffusion coefficient.

For example, the relative importance of advection versus diffusion can be expressed with the dimensionless quantity the *Péclet number* [Cushman-Roisin(2012)]. The Péclet number is defined as follows:

$$P\acute{e} = \frac{\text{advection transport rate}}{\text{diffusive transport rate}} = \frac{v}{D/L} = \frac{vL}{D},$$

where L is the characteristic length of the system (for the above process we will take it as the distance between the starting point and the goal on the x-axis, i.e. L = a - x). For strictly physical processes this is fine, however, in our formulation, we have included a parameter to indicate the relative amount of time individuals do one thing or the other. Thus, we need to include these modifications in the definition. We will do this by defining an effective advection rate, pv (directed speed), and effective diffusion coefficient, qD, so that

$$\mathbf{P}\acute{\mathbf{e}} = \frac{pv\left(a-x\right)}{qD}.$$

Substituting this in the expression for the average time to goal allows us to see the general behavior of the solution without having to manipulate multiple variables or worrying about units of measurement. Rules of thumb are for values of Pé < 0.1 diffusion dominates and for Pé > 10 advection dominates. Thus, we can rewrite (20) as

$$W(x,y) = \frac{a-x}{(1-s)\,pv} \left[1 - \frac{1}{P\acute{e}} \left(e^{-\frac{x}{(a-x)}P\acute{e}} - e^{-\frac{a}{(a-x)}P\acute{e}} \right) \right].$$
 (21)

Comparing W to the time it takes to get to goal by directed movement exclusively $\left(\frac{a-x}{v}\right)$ and letting $r = 1 - \frac{x}{a}$ (the fraction of the distance to go to reach the goal) and s = 0 (no rest), we have

$$\Omega = \frac{W(x,y)}{\left(\frac{a-x}{v}\right)} = \frac{v}{\left(\frac{a-x}{W(x,y)}\right)} = \frac{1}{p} \left(1 - \frac{1}{P\acute{e}} \left(e^{-\frac{(1-r)}{r}P\acute{e}} - e^{-\frac{1}{r}P\acute{e}}\right)\right).$$
(22)

This is the average time it takes to get to the goal relative to traveling the entire distance at the advection rate. Alternatively, it is the speed of advection exclusively relative to the average speed. A graph of this for various values of p is shown in Figure 2.

The relationship in Equation (22), relating the *measurement* of average time to goal relative to time required for direct travel and the Péclet number, provides a possible means of estimating the diffusion coefficient. For example, starting at the far end of the enclosure (r = 1), we have

$$\Omega = \frac{1}{p} \left(1 - \frac{1}{\mathrm{P}\acute{\mathrm{e}}} \left(1 - e^{-\mathrm{P}\acute{\mathrm{e}}} \right) \right).$$
(23)

Graphing Pé as a function of Ω for various values of p, we have the following Figure 3.

We now look at how the distribution of individuals actually evolves over time.

3 Progression of the probability distribution.

In order to derive the 2-d probability distribution of individuals, we start by looking at the change in numbers of individuals at a particular location. The approach is similar to the above: we begin with the assumptions of a semi-random walk in discrete time increments τ , moving in a directed manner a fraction of the time p and in a random manner a fraction of



Figure 2: The time to reach the goal with random and constant directed movement relative to the time to goal with just constant directed movement, $\Omega = W(x, y)/((a - x)/v)$. Here, p is the fraction of time spent in directed movement, r is the fraction of the distance from start to goal. Note that for a fixed Péclet number the greater the percent time searching (i.e., diffusion), the longer the relative time to reach the goal on average. However even if p = 1the time relative to advection exclusively is not 1, since the Péclet number includes divisor by q. Thus, in order for Pé = 1, the parameter $D \to \infty$.

the time (1-p) = q. On the right hand side we write the number of individuals at location (x, y) and at time $t + \tau$. On the left hand side is the number of individuals that have moved in the time interval τ , from one location to another due to random or directed movement. We assume that a fraction s of all individuals everywhere will *not* have moved in the time interval τ . With reference to Figure 3, we have*

*Alternatively we could formulate the problem in the following way,

$$\begin{split} N\left(x, y, t + \tau\right) &= (1 - s) \left[q' N\left(x + \Delta, y, t\right) + p' N\left(x - \Delta, y, t\right) \right. \\ &+ q' N\left(x, y + \Delta, t\right) + q' N\left(x, y - \Delta, t\right)\right] + s N\left(x, y, t\right) \end{split}$$

In this approach we have just a random walk but with a bias in the north direction p' > q' and 3q' + p' = 1. The results of the derivation would be the same with 4q' = q and p' - q' = p. The main difference is in the interpretation, in the main text there is a random movement and a deterministic movement, in the alternate derivation all movement is random but with a bias like dust in a gentle breeze.



Figure 3: The Péclet number as a function of time to goal relative to time to goal with constant directed movement, plotted as the log base 2 of both variables $log_2(P\acute{e})$ vs. $log_2(\Omega)$

$$N(x, y, t + \tau) = (1 - s) \left[\frac{q}{4} N(x + \Delta, y, t) + \frac{q}{4} N(x - \Delta, y, t) + \frac{q}{4} N(x, y + \Delta, t) + \frac{q}{4} N(x, y - \Delta, t) + p N(x - v\tau, y, t) \right] + s N(x, y, t)$$
(24)

The fraction $\frac{1}{4}$ in the above equation is because, for example, only a quarter of the number moving from $(x + \Delta, y)$ go to (x, y), the other $\frac{3}{4}$ go to one of the squares centered at $(x + \Delta, y + \Delta), (x + \Delta, y - \Delta)$, or $(x + 2\Delta, y)$. There is no analogous (fraction) term on the directed movement, since the movement is in one direction only. Subtracting N(x, y, t) from both sides, we have the change in numbers in the interval τ :

$$N(x, y, t + \tau) - N(x, y, t) = (1 - s) \left[\frac{q}{4} \left(N(x + \Delta, y, t) + N(x - \Delta, y, t) + N(x, y + \Delta, t) + N(x, y - \Delta, t) \right) + pN(x - v\tau, y, t) \right] - (1 - s) N(x, y, t).$$
(25)

Regrouping terms, we rewrite as

$$N(x, y, t + \tau) - N(x, y, t) = (1 - s) \frac{q}{4} [(N(x + \Delta, y, t) - N(x, y, t)) + (N(x - \Delta, y, t) - N(x, y, t)) + (N(x, y + \Delta, t) - N(x, y, t)) + (N(x, y - \Delta, t) - N(x, y, t))] + (1 - s) p (N(x - v\tau, y, t) - N(x, y, t)).$$
(26)



Figure 4: The number of individuals or particles relative to a point (x,y) at time t.

Multiplying and dividing the random movement terms by the step increment Δ , the directed movement term by the step increment $v\tau$, and dividing both sides of the equation by the time increment τ , we have

$$\frac{N(x, y, t+\tau) - N(x, y, t)}{\tau} = \frac{(1-s) q \Delta}{4\tau} \left[\left(\frac{N(x+\Delta, y, t) - N(x, y, t)}{\Delta} \right) - \left(\frac{N(x, y, t) - N(x-\Delta, y, t)}{\Delta} \right) \right] \\
+ \left(\frac{N(x, y+\Delta, t) - N(x, y, t)}{\Delta} \right) - \left(\frac{N(x, y, t) - N(x, y-\Delta, t)}{\Delta} \right) \right] \\
- \frac{(1-s) pv\tau}{\tau} \left(\frac{N(x, y, t) - N(x-v\tau, y, t)}{v\tau} \right).$$
(27)

Taking the limit as $\tau, \Delta \to 0$

$$\frac{\partial}{\partial t}N(x,y,t+\tau) = \frac{(1-s)\,q\Delta}{4\tau} \left[\left(\frac{\partial}{\partial x}N(x+\Delta,y,t) - \frac{\partial}{\partial x}N(x,y,t) \right) + \left(\frac{\partial}{\partial y}N(x,y+\Delta,t) - \frac{\partial}{\partial y}N(x,y,t) \right) \right] - (1-s)\,pv\frac{\partial}{\partial x}N(x,y,t) \,.$$
(28)

Multiplying and dividing the random movement term by Δ again, we have

$$\frac{\partial}{\partial t}N(x,y,t+\tau) = \frac{(1-s)q\Delta^2}{4\tau} \left[\left(\frac{\frac{\partial}{\partial x}N(x+\Delta,y,t) - \frac{\partial}{\partial x}N(x,y,t)}{\Delta} \right) + \left(\frac{\frac{\partial}{\partial y}N(x,y+\Delta,t) - \frac{\partial}{\partial y}N(x,y,t)}{\Delta} \right) \right] - (1-s)pv\frac{\partial}{\partial x}N(x,y,t).$$
(29)

Taking the limit as $\tau \to 0, \Delta \to 0$ again

$$\frac{\partial}{\partial t}N\left(x,y,t\right) = \frac{(1-s)\,q\Delta^2}{4\tau} \left(\frac{\partial^2}{\partial x^2}N\left(x,y,t\right) + \frac{\partial^2}{\partial y^2}N\left(x,y,t\right)\right) - (1-s)\,pv\frac{\partial}{\partial x}N\left(x,y,t\right). \tag{30}$$

Recalling that the diffusion coefficient is $D = \frac{\Delta^2}{4\tau}$, we get

$$\frac{1}{(1-s)}\frac{\partial}{\partial t}N(x,y,t) = qD\left(\frac{\partial^2}{\partial x^2}N(x,y,t) + \frac{\partial^2}{\partial y^2}N(x,y,t)\right) - pv\frac{\partial}{\partial x}N(x,y,t).$$
(31)

If we assume that the total population over the entire area in question is constant and equal to N_T , then dividing through by this amount we have the "fraction" of the population at every point. The probability density of finding an individual at a point is therefore $P(x, y, t) = N(x, y, t)/N_T$. Recall that in the average time to capture calculation in the first section, the average time was scaled by the fraction of time spent resting or staying put at a single location. Observe that the same thing is occuring in (31), rescaling time as $(1-s) t \to t$ (so that henceforth t will represent the "active" time, or the time for the active fraction of the population). If at some point in the future we require the total time (say, for instance, we are measuring over the course of a few days, or individuals are resting at regular intervals), then s becomes a significant fraction and can simply divide all t's by (1-s) to get the overall time. Equation (31) simplifies to

$$\frac{\partial}{\partial t}P(x,y,t) = qD\left(\frac{\partial^2}{\partial x^2}P(x,y,t) + \frac{\partial^2}{\partial y^2}P(x,y,t)\right) - pv\frac{\partial}{\partial x}P(x,y,t), \quad (32)$$

where t now represents movement during active times.

3.1 Solution to diffusion-advection equation in a closed area.

In order to solve this equation for P, we will change our frame of reference from the stationary coordinates (x, y, t) to moving coordinates (x', y', t'). The new reference frame is moving at the speed pv to the right (parallel to the stationary x-axis) so that we have $x' = x - (x_0 + pvt)$, $y' = y - y_0$, and t' = t. At t = 0, we have $(x, y) = (x_0, y_0)$ and (x', y') = (0, 0). From these, we have the following derivatives

$$\frac{\partial x'}{\partial t} = -pv, \quad \frac{\partial x'}{\partial x} = 1,$$

$$\frac{\partial y'}{\partial t} = 0, \quad \frac{\partial y'}{\partial y} = 1,$$

$$\frac{\partial t'}{\partial t} = 1, \quad \frac{\partial t'}{\partial x} = 0, \quad \text{and} \quad \frac{\partial t'}{\partial y} = 0.$$
(33)

Using these and applying the chain rule we obtain the following change of variables,

$$\frac{\partial}{\partial t}P(x,y,t) = \frac{\partial}{\partial x'}P(x',y',t')\frac{\partial x'}{\partial t} + \frac{\partial}{\partial y'}P(x',y',t')\frac{\partial y'}{\partial t} + \frac{\partial}{\partial t'}P(x',y',t')\frac{\partial t'}{\partial t} = -pv\frac{\partial}{\partial x'}P(x',y',t') + \frac{\partial}{\partial t'}P(x',y',t'),$$
(34)

$$\frac{\partial}{\partial x}P(x,y,t) = \frac{\partial}{\partial x'}P(x',y',t')\frac{\partial x'}{\partial x} + \frac{\partial}{\partial t'}P(x',y',t')\frac{\partial t'}{\partial x} = \frac{\partial}{\partial x'}P(x',y',t')$$
(35)

$$\frac{\partial^2}{\partial x^2} P\left(x, y, t\right) = \frac{\partial}{\partial x'} \left(\frac{\partial}{\partial x'} P\left(x', y', t'\right)\right) \frac{\partial x'}{\partial x} = \frac{\partial^2}{\left(\partial x'\right)^2} P\left(x', y', t'\right).$$
(36)

Similarly,

$$\frac{\partial}{\partial y}P(x,y,t) = \frac{\partial}{\partial y'}P(x',y',t')$$
(37)

$$\frac{\partial^2}{\partial y^2} P\left(x, y, t\right) = \frac{\partial^2}{\left(\partial y'\right)^2} P\left(x', y', t'\right).$$
(38)

Substituting the appropriate expressions from lines (33) to (38) into the derivatives in equation (32), we have

$$- pv\frac{\partial}{\partial x'}P(x',y',t') + \frac{\partial}{\partial t'}P(x',y',t')$$

$$= qD\left(\frac{\partial^2}{(\partial x')^2}P(x',y',t') + \frac{\partial^2}{(\partial y')^2}P(x',y',t')\right) - pv\frac{\partial}{\partial x'}P(x',y',t')$$

Canceling the $-pv\frac{\partial}{\partial x'}P(x',y',t')$ term on either side, we reduce to

$$\frac{\partial}{\partial t'}P\left(x',y',t'\right) = qD\left(\frac{\partial^2}{\left(\partial x'\right)^2}P\left(x',y',t'\right) + \frac{\partial^2}{\left(\partial y'\right)^2}P\left(x',y',t'\right)\right).$$
(39)

This is just the form of an equation for diffusion in 2-d. We can try separation of variables, where P(x', y', t') = X(x', t') Y(y', t'), to get

$$\frac{\partial}{\partial t'} P\left(x', y', t'\right) = \frac{\partial}{\partial t'} \left[X\left(x', t'\right) Y\left(y', t'\right)\right]
= Y\left(y', t'\right) \frac{\partial}{\partial t'} X\left(x', t'\right) + X\left(x', t'\right) \frac{\partial}{\partial t'} Y\left(y', t'\right),$$
(40)

$$\frac{\partial^2}{\left(\partial x'\right)^2} P\left(x', y', t'\right) = Y\left(y', t'\right) \frac{\partial^2}{\left(\partial x'\right)^2} X\left(x', t'\right),\tag{41}$$

and
$$\frac{\partial^2}{\left(\partial y'\right)^2} P\left(x', y', t'\right) = X\left(x', t'\right) \frac{\partial^2}{\left(\partial y'\right)^2} Y\left(y', t'\right).$$
 (42)

Substituting these in (39) (and leaving off the function arguments for brevity) gives us

$$Y\frac{\partial X}{\partial t'} + X\frac{\partial Y}{\partial t'} = qD\left(Y\frac{\partial^2 X}{\left(\partial x'\right)^2} + X\frac{\partial^2 Y}{\left(\partial y'\right)^2}\right).$$
(43)

Putting all the Y terms together and all the X terms together we have

$$\frac{1}{Y}\left(\frac{\partial Y}{\partial t'} - qD\frac{\partial^2 Y}{\left(\partial y'\right)^2}\right) + \frac{1}{X}\left(\frac{\partial X}{\partial t'} - qD\frac{\partial^2 X}{\left(\partial x'\right)^2}\right) = 0.$$
(44)

Since both X and Y are positive functions and recalling that, from the original construction of the problem, movement in the x direction is independent of movement in the y direction, we determine that

$$\frac{\partial Y}{\partial t'} - qD \frac{\partial^2 Y}{\left(\partial y'\right)^2} = 0 \tag{45}$$

and
$$\frac{\partial X}{\partial t'} - qD \frac{\partial^2 X}{\left(\partial x'\right)^2} = 0.$$
 (46)

We will start with a solution on an infinite domain[†], i.e. $-\infty < x' < \infty$ and $-\infty < y' < \infty$ and with the initial condition as a instantaneous point source

$$\left. \begin{array}{l}
Y\left(y',0\right) = \delta\left(y'\right) \\
X\left(x',0\right) = \delta\left(x'\right)
\end{array} \right\} \quad \text{Dirac delta functions.}$$
(47)

Since the equations for X and Y have identical forms we will go through the solution only for one of them. Applying the Fourier transform to the X equation, gives

$$\mathcal{F}\left(\frac{\partial}{\partial t'}X\left(x',t'\right)\right) = qD\mathcal{F}\left(\frac{\partial^2 X}{\left(\partial x'\right)^2}\right).$$
(48)

Which yields the ordinary differential equation (the circumflex over the variable name indicates the variable is transformed from the time domain to the frequency domain),

$$\frac{d}{dt'}\hat{X}(\omega,t') = -qD\omega^2\hat{X}(\omega,t').$$
(49)

This has the solution

$$\hat{X}(\omega, t') = \hat{X}(\omega, 0) e^{-qDt'\omega^2}.$$
(50)

We now need to return to the time domain. To do this we note that, *in general*, the Fourier transform of the convolution of two functions is the product of their Fourier transforms. Therefore, the inverse Fourier transform of a product of two functions in the frequency domain is the convolution of the two functions in the time domain. For example,

$$\mathcal{F}\left(f\left(x\right)*g\left(x\right)\right) = \mathcal{F}\left(f\left(x\right)\right)\mathcal{F}\left(g\left(x\right)\right).$$
(51)

Switching right and left-hand sides and taking the inverse transform of both sides, we have

[†]We could also seek a solution directly on a finite domain for diffusion along the y axis, however, the method used here is more succinct and provides a parallel approach in both directions.

$$\mathcal{F}^{-1}\left(\mathcal{F}\left(f\left(x\right)\right)\mathcal{F}\left(g\left(x\right)\right)\right) = \mathcal{F}^{-1}\left(\mathcal{F}\left(f\left(x\right)\ast g\left(x\right)\right)\right) = f\left(x\right)\ast g\left(x\right),\tag{52}$$

or, using the "hat" notation for the functions in the frequency domain $(\hat{f}(\omega) = \mathcal{F}(f(x)))$ and $\hat{g}(\omega) = \mathcal{F}(g(x)))$, we can rewrite the inverse transform of the product of two transforms as

$$\mathcal{F}^{-1}\left(\hat{f}\left(\omega\right)\hat{g}\left(\omega\right)\right) = f\left(x\right) * g\left(x\right).$$
(53)

Employing the above device, we obtain the inverse transform in the following steps.

$$X(x',t') = \mathcal{F}^{-1}\left(\hat{X}(\omega,t')\right)$$
$$= \mathcal{F}^{-1}\left(\hat{X}(\omega,0)e^{-qDt'\omega^2}\right).$$
(54)

Note that

$$\mathcal{F}^{-1}\left(\hat{X}\left(\omega,0\right)\right) = X\left(x',0\right) \Leftrightarrow \hat{X}\left(\omega,0\right) = \mathcal{F}\left(X\left(x',0\right)\right) = \mathcal{F}\left(\delta\left(x'\right)\right) \tag{55}$$

and
$$\mathcal{F}^{-1}\left(e^{-qDt'\omega^2}\right) = \frac{1}{\sqrt{2qDt'}}e^{-\frac{(x')^2}{4qDt'}} \Leftrightarrow e^{-\frac{qD}{2}t'\omega^2} = \mathcal{F}\left(\frac{1}{\sqrt{2qDt'}}e^{-\frac{(x')^2}{4qDt'}}\right).$$
 (56)

Applying the convolution theorem for Fourier transforms and the properties of the Dirac delta function, we get

$$X(x',t') = \mathcal{F}^{-1}\left(\hat{X}(\omega,0)e^{-qDt'\omega^{2}}\right)$$
$$= \mathcal{F}^{-1}\left(\mathcal{F}\left(\delta(x')\right)\mathcal{F}\left(\frac{1}{\sqrt{2qDt'}}e^{-\frac{(x')^{2}}{4qDt'}}\right)\right)$$
$$= \mathcal{F}^{-1}\left(\mathcal{F}\left(\delta(x')*\frac{1}{\sqrt{2qDt'}}e^{-\frac{(x')^{2}}{4qDt'}}\right)\right).$$
(57)

$$X(x',t') = \delta(x') * \frac{1}{\sqrt{2qDt'}} e^{-\frac{(x')^2}{4qDt'}} = \frac{1}{\sqrt{4\pi qDt'}} \int_{-\infty}^{\infty} \delta(s) e^{-\frac{(x'-s)^2}{4qDt'}} ds = \frac{1}{\sqrt{4\pi qDt'}} e^{-\frac{(x')^2}{4qDt'}}.$$
(58)

With a similar result for the Y function, we obtain

$$Y(y',t') = \frac{1}{\sqrt{4\pi q D t'}} e^{-\frac{(y')^2}{4q D t'}}.$$
(59)

Switching back to the original reference frame we have

$$X(x,t) = \frac{1}{\sqrt{4\pi q D t}} e^{-\frac{(x-x_0 - pvt)^2}{4q D t}},$$
(60)

$$Y(y,t) = \frac{1}{\sqrt{4\pi q D t}} e^{-\frac{(y-y_0)^2}{4q D t}},$$
(61)

While we get solutions, this is not the end of the story. This result is only valid on an unbounded plane but that is not the situation which we have here. We have impenetrable boundaries at $y = \pm b/2$, x = 0 and x = a. The above solution does not satisfy these conditions. To fulfill the desired initial and boundary conditions we can add together multiple solutions to create a new one by superposition given that the original differential equation is linear. Three of the four boundaries $(y = \pm b/2 \text{ and } x = 0)$ have no flux $\partial P(0, y, t) / \partial x = 0$, $\partial P(x, b, t) / \partial y = 0$, and $\partial P(x, b, t) / \partial x = 0$, they just reflect anything coming at them. The fourth boundary, at x = a, requires a bit more consideration, which we will get to in a minute. For diffusion along the y-axis the fix is straightforward, treat the two boundaries as if they were two mirrors facing each other. We then take the part of each reflected image that lies between y = -b/2 and y = b/2 and add it to the original distribution. If the "y" distribution is centered at y_0 then the images are centered at $nb + (-1)^n y_0$ for $n = \{0, \pm 1, \pm 2, \pm 3, \ldots\}$. Or letting $n_{\text{even}} = 2k$ for positive images and $n_{\text{odd}} = 2k + 1$ for negative images, the Y(y, t) factor is then

$$\sum_{k=-\infty}^{\infty} Y_k(y,t) = \frac{1}{2\sqrt{\pi q D t}} \sum_{k=-\infty}^{\infty} \left(e^{-\frac{(y-y_0+2kb)^2}{4q D t}} + e^{-\frac{(y+y_0+(2k+1)b)^2}{4q D t}} \right) \quad \text{for} \quad -b/2 \le y \le b/2$$
(62)

This spreads and becomes uniform distribution as time progresses, and

$$\lim_{t \to \infty} \sum_{k=-\infty}^{\infty} Y_k(y,t) = \frac{1}{b}.$$
(63)



Figure 5: The primary distribution crosses the boundary at y = -b/2 and y = b/2. These tails of the pdf can be exactly replaced within the boundaries of the problem by adding a new "reflected" distributions centered at $y = -b - y_0$ on the left and $y = b - y_0$ on the right. Each of these have their own reflections at $y = \pm 2b + y_0$ and so on ad infinitum. The superposition of the imaged distributions (colored at bottom) in the practical boundary is shown as blended colors over the primary distribution.

In practice only a few terms are needed since the tails of the distributions coming from images that are far off from the original are very small. Consequently the series converges quickly.

This approach does not work for movement parallel to the x-axis. If we treat both of the boundaries at x = 0 and x = a as reflecting, the distribution spreads with time as it does in diffusion only, but also the entire distribution is translated to the right due to the advection term. This looks alright until the center of the original distribution passes x = a. At this point, since the distribution's reflection is traveling in the opposite direction, more of the image distribution is between the boundaries traveling to the left than the original still traveling to the right. Then as the first image reaches the back wall it's image pokes it's head out and moves to the right again. The entire affect is to make a superposed PDE that spreads as it sloshes back and forth between the fore and aft walls!

What is to be done then? If we think about the situation of a chimney emitting smoke into a very calm breeze the analogy is apt (see examples in Chapter 2 sections 6 & 7 of [Cushman-Roisin(2012)]): on the downwind side of the chimney the breeze carries off the smoke, however, on the upwind side just enough smoke diffuses in that direction to replace the amount carried back downwind. If the amount of smoke and wind were perfectly unwavering, the plume would quickly reach a steady state.

Now, although this analogy is far from perfect, it does give us a starting point for the analysis. At the upstream end, x = 0, we have a reflective boundary $\frac{\partial}{\partial x}P(0, y, t) = 0$ and at the downstream end the barrier, instead of behaving as a boundary, acts like a continuous source at x = a, emitting particles into a steady flow, back into the container by diffusion equal to the amount (Q) to that would have moved out if unimpeded. In other words, as the downstream end of the original distribution moves mathematically past the boundary by a certain amount, that amount actually diffuses back into the container. Again we start with our advection-diffusion equation,

$$\frac{\partial}{\partial t}P(x,y,t) = qD\left(\frac{\partial^2}{\partial x^2}P(x,y,t) + \frac{\partial^2}{\partial y^2}P(x,y,t)\right) - pv\frac{\partial}{\partial x}P(x,y,t).$$
(64)

Since the flow is hitting a stationary wall, we apply separation of variables without changing reference frame: $P(x, y, t) = \Psi(x, t) Y(y, t)$. We use Ψ instead of X to emphasize that this factor is a distribution due to the reflux parallel to the x axis and not the original flow. Thus we have

$$\frac{\partial}{\partial t}P(x, y, t) = \frac{\partial}{\partial t} \left[\Psi(x, t) Y(y, t)\right]$$
$$= Y(y, t) \frac{\partial}{\partial t}\Psi(x, t) + \Psi(x, t) \frac{\partial}{\partial t}Y(y, t),$$
(65)

$$\frac{\partial}{\partial x}P(x,y,t) = Y(y,t)\frac{\partial}{\partial x}\Psi(x,t), \qquad (66)$$

$$\frac{\partial^2}{\partial x^2} P\left(x, y, t\right) = Y\left(y, t\right) \frac{\partial^2}{\partial x^2} \Psi\left(x, t\right),\tag{67}$$

and
$$\frac{\partial^2}{\partial y^2} P(x, y, t) = \Psi(x, t) \frac{\partial^2}{\partial y^2} Y(y, t).$$
 (68)

Substituting and regrouping like terms, we get

$$Y(y,t)\frac{\partial}{\partial t}\Psi(x,t) + \Psi(x,t)\frac{\partial}{\partial t}Y(y,t)$$

$$= qD\left(Y(y,t)\frac{\partial^{2}}{\partial x^{2}}\Psi(x,t) + \Psi(x,t)\frac{\partial^{2}}{\partial y^{2}}Y(y,t)\right) - pvY(y,t)\frac{\partial}{\partial x}\Psi(x,t),$$

$$0 = \frac{1}{\Psi(x,t)}\left(\frac{\partial}{\partial t}\Psi(x,t) - qD\frac{\partial^{2}}{\partial x^{2}}\Psi(x,t) + pv\frac{\partial}{\partial x}\Psi(x,t)\right)$$

$$+ \frac{1}{Y(y,t)}\left(\frac{\partial}{\partial t}Y(y,t) - qD\frac{\partial^{2}}{\partial y^{2}}Y(y,t)\right).$$
(69)

The Y terms are diffusion parallel to the y axis, which we have solved for above. The Ψ term is advection and diffusion parallel to the x axis but we now have a source term at the right hand boundary:

$$\frac{\partial}{\partial t}\Psi(x,t) = qD\frac{\partial^2}{\partial x^2}\Psi(x,t) - pv\frac{\partial}{\partial x}\Psi(x,t)$$
(70)

with initial and boundary conditions;

$$\Psi(x,0) = 0 \quad \text{initial conditions,} \tag{71}$$

$$\Psi(-\infty,t) = 0$$
, left-hand side (momentarily ignoring the boundary), (72)

$$\frac{\partial}{\partial t}\Psi(a,t) = \frac{d}{dt}Q(t) \quad \text{right-hand boundary (a variable source).}$$
(73)

The Laplace transform in most useful in dealing with functions defined on a semi-infinite domain. In preparation for applying a Laplace transform we change coordinates (still fixed) from $x \in (-\infty, a)$ to $x' \in (0, \infty)$,

$$x = a - x' \quad \Leftrightarrow \quad x' = a - x \tag{74}$$

$$y = y'$$
 and $t = t'$ (75)

$$\frac{\partial t'}{\partial t} = 1, \quad \frac{\partial x'}{\partial x} = -1, \quad \frac{\partial^2 x'}{\partial x^2} = 0, \tag{76}$$

resulting in

$$\frac{\partial}{\partial t'}\Psi\left(x',t'\right) = \frac{qD}{2}\frac{\partial^2}{\left(\partial x'\right)^2}\Psi\left(x',t'\right) + pv\frac{\partial}{\partial x'}\Psi\left(x',t'\right).$$
(77)

Applying the Laplace transform (here we use a tilde to indicate the transformed function, i.e., $\mathcal{L}(\Psi(x',t')) = \int_0^\infty \Psi(x',t') e^{-st} dt = \tilde{\Psi}(x',s))^{\ddagger}$,

$$\mathcal{L}\left(\frac{\partial}{\partial t'}\Psi\left(x',t'\right)\right) = qD\mathcal{L}\left(\frac{\partial^2}{\left(\partial x'\right)^2}\Psi\left(x',t'\right)\right) + pv\mathcal{L}\left(\frac{\partial}{\partial x'}\Psi\left(x',t'\right)\right),\tag{78}$$

$$s\tilde{\Psi}(x',s) - \Psi(x',0) = qD\frac{d^2}{(dx')^2}\tilde{\Psi}(x',s) + pv\frac{d}{dx'}\tilde{\Psi}(x',s).$$

$$\tag{79}$$

^{\ddagger}Here the "s" is the frequency variable of the Laplace transform and not the fraction resting that we used in the random walk example.

Substituting the initial condition from (71), we get

$$s\tilde{\Psi}(x',s) = qD\frac{d^2}{\left(dx'\right)^2}\tilde{\Psi}(x',s) + pv\frac{d}{dx'}\tilde{\Psi}(x',s), \qquad (80)$$

$$0 = qD \frac{d^2}{\left(dx'\right)^2} \tilde{\Psi}\left(x',s\right) + pv \frac{d}{dx'} \tilde{\Psi}\left(x',s\right) - s\tilde{\Psi}\left(x',s\right).$$
(81)

The transform of the boundary condition (73) gives

$$\Psi(x',0) = 0 \quad \text{and} \quad \frac{\partial}{\partial t'}\Psi(0,t') = \frac{d}{dt'}Q(t'),$$

$$s\tilde{\Psi}(0,s) - \Psi(0,0) = s\tilde{Q}(s) - Q(0) \Rightarrow \tilde{\Psi}(0,s) = \tilde{Q}(s).$$
(82)

From (81) we have the characteristic equation

$$0 = qDr^2 + pvr - s,$$

which has roots

$$r_1 = -\frac{pv}{2qD} \left(1 + \sqrt{1 + \frac{4qDs}{p^2v^2}} \right), \text{ and } r_2 = -\frac{pv}{2qD} \left(1 - \sqrt{1 + \frac{4qDs}{p^2v^2}} \right).$$
(83)

So the solution has the form

$$\tilde{\Psi}(x',s) = c_1 e^{-\frac{pvx'}{2qD}\left(\sqrt{1+\frac{4qDs}{p^2v^2}}+1\right)} + c_2 e^{\frac{pvx'}{2qD}\left(\sqrt{1+\frac{4qDs}{p^2v^2}}-1\right)}.$$
(84)

Now as $s \to \infty$, $\tilde{\Psi}(x',s) < \infty \Longrightarrow c_2 = 0$ so that

$$\tilde{\Psi}(x',s) = c_1 e^{-\frac{pvx'}{2qD} \left(\sqrt{1 + \frac{4qDs}{p^2v^2}} + 1\right)}.$$
(85)

However, at x' = 0, $\tilde{\Psi}(0, s) = c_1$, hence,

$$\tilde{\Psi}(x',s) = \tilde{\Psi}(0,s) e^{-\frac{pvx'}{2qD} \left(\sqrt{1 + \frac{4qDs}{p^2v^2}} + 1\right)}.$$
(86)

From (82), we have

$$\tilde{\Psi}(x',s) = \tilde{Q}(s) e^{-\frac{pvx'}{2qD} \left(\sqrt{1 + \frac{4qDs}{p^2v^2}} + 1\right)}.$$
(87)

We again need to return to the time domain, and as it was in the case of the Fourier transform, the inverse Laplace transform of a product of two functions in the frequency domain is the convolution of the two functions in the time domain (or, equivalently, the convolution of the inverse Laplace transforms of the functions in the frequency domain) [§]. Applying an inverse Laplace transform to both sides of (87), we have

$$\Psi(x',t') = \mathcal{L}^{-1}\left(\tilde{Q}(s) e^{-\frac{pvx'}{2qD}\left(\sqrt{1+\frac{4qDs}{p^2v^2}}+1\right)}\right).$$
(88)

In (88) the argument of the inverse Laplace transform is the product of two Laplace transforms. Therefore the inverse transform is given by

$$\Psi(x',t') = \mathcal{L}^{-1}\tilde{Q}(s) * \mathcal{L}^{-1}\left(e^{-\frac{pvx'}{2qD}\left(\sqrt{1+\frac{4sqD}{p^2v^2}}+1\right)}\right).$$
(89)

Simplifying (89) a bit, we arrive at

$$\Psi\left(x',t'\right) = Q\left(t'\right) * e^{-\frac{pvx'}{2qD}} \mathcal{L}^{-1}\left(e^{-\frac{pvx'}{2qD}\sqrt{1 + \frac{4qD}{p^2v^2}s}}\right).$$
(90)

In order to use Laplace transform tables to deal with the remaining inverse transform, we need to simplify notation, which we can do by using the following change of variables.

$$\xi = \frac{pvx'}{2qD}, \quad \text{and} \quad \gamma = 1 + \frac{4qD}{p^2v^2}s, \tag{91}$$

$$s = \frac{p^2 v^2}{4qD} \left(\gamma - 1\right) \Rightarrow ds = \frac{p^2 v^2}{4qD} d\gamma.$$
(92)

[§]The general approach, starting with the Laplace transform of a convolution of two functions, is

$$\mathcal{L}\left(f\left(t\right)\ast g\left(t\right)\right)=\mathcal{L}\left(f\left(t\right)\right)\mathcal{L}\left(g\left(t\right)\right).$$

Switching the sides and applying the inverse transform, we have

$$\mathcal{L}^{-1}\left(\mathcal{L}\left(f\left(t\right)\right)\mathcal{L}\left(g\left(t\right)\right)\right) = \mathcal{L}^{-1}\mathcal{L}\left(f\left(t\right)*g\left(t\right)\right) = f\left(t\right)*g\left(t\right),$$

or using the "tilda" notation for the functions in the frequency domain we have

$$\mathcal{L}^{-1}\left(\tilde{f}\left(s\right)\tilde{g}\left(s\right)\right) = f\left(t\right) * g\left(t\right).$$

We write out the inverse transform explicitly to sort out the re-scaling necessary when changing variables:

$$\mathcal{L}^{-1}\left(e^{-\frac{pvx'}{2qD}\sqrt{1+\frac{4qD}{p^2v^2}s}}\right) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{-\frac{pvx'}{2qD}\sqrt{1+\frac{4qD}{p^2v^2}s}} e^{st'} ds.$$
(93)

Now substituting $x' \to \xi$ and $s \to \gamma$ terms, we get

$$\mathcal{L}^{-1}\left(e^{-\frac{pvx'}{2qD}\sqrt{1+\frac{4qD}{p^2v^2}s}}\right) = \frac{p^2v^2}{4qD}\frac{1}{2\pi i}\int_{\left(1+\frac{4qD}{p^2v^2}c\right)-i\infty}^{\left(1+\frac{4qD}{p^2v^2}c\right)+i\infty}e^{-\xi\sqrt{\gamma}}e^{\frac{p^2v^2}{4qD}(\gamma-1)t'}d\gamma$$
$$= \frac{p^2v^2}{4qD}e^{-\theta}\frac{1}{2\pi i}\int_{g-i\infty}^{g+i\infty}e^{-\xi\sqrt{\gamma}}e^{\theta\gamma}d\gamma,$$
(94)

where
$$g = \left(1 + \frac{4qD}{p^2v^2}c\right)$$
 and $\theta = \frac{p^2v^2}{4qD}t'.$ (95)

Rewriting the integral in function notation, we obtain

$$\mathcal{L}^{-1}\left(e^{-\frac{pvx'}{2qD}\sqrt{1+\frac{4qD}{p^2v^2}s}}\right) = \frac{p^2v^2}{4qD}e^{-\theta}\mathcal{L}^{-1}\left(e^{-\xi\sqrt{\gamma}}\right).$$
(96)

From the tables [Zwillinger(1996)] (p. 563, #74) the corresponding inverse Laplace transform is

$$\mathcal{L}^{-1}\left(e^{-\xi\sqrt{\gamma}}\right) = \frac{\xi}{2\theta\sqrt{\pi\theta}}e^{-\frac{\xi^2}{4\theta}},\tag{97}$$

$$\mathcal{L}^{-1}\left(e^{-\frac{pvx'}{2qD}\sqrt{1+\frac{4qD}{p^2v^2}s}}\right) = \left(\frac{p^2v^2}{4qD}e^{-\theta}\right)\left(\frac{\xi}{2\theta\sqrt{\pi\theta}}e^{-\frac{\xi^2}{4\theta}}\right).$$
(98)

Substituting this result in (90) and replacing $e^{-\frac{pvx'}{2qD}}$ with $e^{-\xi}$, we obtain

$$\Psi\left(x',t'\right) = Q\left(t'\right) * e^{-\xi} \left(\frac{p^2 v^2}{4qD} e^{-\theta}\right) \left(\frac{\xi}{2\theta\sqrt{\pi\theta}} e^{-\frac{\xi^2}{4\theta}}\right).$$
(99)

Simplifying and then returning to our unscaled space and time variables, we get

$$\Psi(x',t') = Q(t') * \frac{p^2 v^2}{2qD} \frac{\xi}{4\theta \sqrt{\pi\theta}} e^{-\frac{(2\theta+\xi)^2}{4\theta}}$$
(100)

$$= Q(t') * \frac{x'}{t'\sqrt{4\pi q D t'}} e^{-\frac{(x'+pvt')^2}{4q D t'}}.$$
 (101)

Finally, writing the convolution as an integral, we obtain

$$\Psi(x',t') = \int_0^{t'} \frac{x'}{\tau\sqrt{4\pi q D\tau}} e^{-\frac{(x'+pv\tau)^2}{4qD\tau}} Q(t'-\tau) d\tau.$$
 (102)

(Here τ is just a dummy variable for the integration.)

By the mean value theorem there is some value of time t'_* , lying in the interval (0, t'), such that

$$\Psi(x',t') = Q(t'_{*}) \int_{0}^{t'} \frac{x'}{\tau\sqrt{4\pi q D\tau}} e^{-\frac{(x'+pv\tau)^{2}}{4q D\tau}} d\tau.$$
(103)

Using definitions (91) and (95), we switch back to the unitless variables to make the integration simpler $\left(\frac{2qD}{pv}\xi = x', \frac{4qD}{p^2v^2}\theta = t'\right)$, and $\frac{4qD}{p^2v^2}u = \tau$:

$$\int_{0}^{t'} \frac{x'}{\tau\sqrt{4\pi q D\tau}} e^{-\frac{(x'+pv\tau)^{2}}{4q D\tau}} d\tau = \frac{\xi}{2\sqrt{\pi}} \int_{0}^{\theta} \frac{1}{\sqrt{u^{3}}} e^{-\frac{(2u+\xi)^{2}}{4u}} du$$
$$= \frac{1}{2} \left[e^{-2\xi} \left(1 + \operatorname{erf}\left(\frac{2\theta - \xi}{2\sqrt{\theta}}\right) \right) + \operatorname{erfc}\left(\frac{2\theta + \xi}{2\sqrt{\theta}}\right) \right],$$
$$\Psi\left(x',t'\right) = Q\left(t'_{*}\right) \frac{1}{2} \left[e^{-\frac{pvx'}{qD}} \left(1 + \operatorname{erf}\left(\frac{pvt' - x'}{\sqrt{4qt'D}}\right) \right) + \operatorname{erfc}\left(\frac{pvt' + x'}{\sqrt{4qDt'}}\right) \right].$$
(104)

However, from the way we defined the problem, the total integral of $\Psi(x', t')$ is the total probability of an uninhibited flow passing x = a over all x and is to equal Q(t'). Integrating (103) over all x', we have

$$\int_{0}^{\infty} \Psi(x',t') \, dx' = Q(t') \,. \tag{105}$$

Therefore,

$$Q(t') = Q(t'_{*}) \int_{0}^{\infty} \int_{0}^{t'} \frac{x'}{\tau \sqrt{4\pi q D \tau}} e^{-\frac{(x'+pv\tau)^{2}}{4q D \tau}} d\tau dx'.$$
 (106)

If we take the integral in (106) as the probability distribution density for Q(t'). then we need a normalizing factor (h(t')) to make it integrate to unity.

$$1 = h(t') \int_0^\infty \int_0^{t'} \frac{x'}{\tau \sqrt{4\pi q D \tau}} e^{-\frac{(x' + pv\tau)^2}{4q D \tau}} d\tau dx'$$
(107)

Again we switch to the non-dimensional variables before integrating $\left(\frac{2qD}{pv}\xi = x', \frac{4qD}{p^2v^2}\theta = t'\right)$, and $\frac{4qD}{p^2v^2}u = \tau$, and $\frac{4qD}{p^2v^2}u = \tau$, and proceed as follows[¶]

$$1 = h(t') \frac{qD}{\sqrt{\pi}pv} \int_{0}^{\infty} \int_{0}^{\theta} \xi u^{-\frac{3}{2}} e^{-\frac{1}{4u}(2u+\xi)^{2}} dud\xi$$

$$= h(t') \frac{qD}{\sqrt{\pi}pv} \int_{0}^{\theta} u^{-\frac{3}{2}} \left(\int_{0}^{\infty} \xi e^{-\frac{1}{4u}(2u+\xi)^{2}} d\xi \right) du$$

$$= h(t') \frac{2qD}{pv} \int_{0}^{\theta} \left(\frac{1}{\sqrt{\pi u}} e^{-u} - \operatorname{erfc}(\sqrt{u}) \right) du$$

$$= h(t') \frac{2qD}{pv} \left(\frac{1}{2} \operatorname{erf}(\sqrt{\theta}) - \theta \operatorname{erfc}(\sqrt{\theta}) + \sqrt{\frac{\theta}{\pi}} e^{-\theta} \right).$$

$$1 = h(t') \left[\frac{qD}{pv} \operatorname{erf}\left(\sqrt{\frac{p^{2}v^{2}t'}{4qD}} \right) - \frac{pvt'}{2} \operatorname{erfc}\left(\sqrt{\frac{p^{2}v^{2}t'}{4qD}} \right) + \sqrt{\frac{qDt'}{\pi}} e^{-\frac{p^{2}v^{2}}{4qD}t'} \right].$$

Solving for h(t'), we have

$$h(t') = \left(\frac{qD}{pv}\operatorname{erf}\left(\frac{pvt'}{\sqrt{4qDt'}}\right) - \frac{pvt'}{2}\operatorname{erfc}\left(\frac{pvt'}{\sqrt{4qDt'}}\right) + \sqrt{\frac{qDt'}{\pi}}e^{-\frac{p^2v^2t'}{4qD}}\right)^{-1}.$$
 (108)

Note that from (106) and (107), we have

$$Q(t'_{*}) = h(t') Q(t').$$
(109)

Putting this together with (104) and (108), we get

$$\Psi\left(x',t'\right) = Q\left(t'\right) \left(\frac{e^{-\frac{pvx'}{qD}}\left(1 + \operatorname{erf}\left(\frac{pvt'-x'}{\sqrt{4qDt'}}\right)\right) + \operatorname{erfc}\left(\frac{pvt'+x'}{\sqrt{4qDt'}}\right)}{\frac{2qD}{pv}\operatorname{erf}\left(\frac{pvt'}{\sqrt{4qDt'}}\right) - pvt'\operatorname{erfc}\left(\frac{pvt'}{\sqrt{4qDt'}}\right) + \sqrt{\frac{4qDt'}{\pi}}e^{-\frac{p^2v^2t'}{4qD}}\right)$$
(110)

As we did with the distribution parallel to the y-axis, we will need to add reflections of this distribution to get the correct image within the boundaries of the enclosure. The presence

[¶]Note that $\operatorname{erf}(x) = \frac{2}{\pi} \int_0^x e^{-t^2} dt$ and $\operatorname{erfc}(x) = \frac{2}{\pi} \int_x^\infty e^{-t^2} dt = 1 - \operatorname{erf}(x)$.

of x' terms in the argument of the error functions will make for very messy expressions. To simplify (a little bit), we can approximate this expression as just an exponential. We obtain the approximation by evaluating the above at x' = 0 and using the result for the parametrization of the exponential. As it turns out this is just the normalizing factor, h, that we obtained above:

$$\Psi(x',t') \approx Q(t') h(t') e^{-h(t')x'},$$
(111a)

where
$$h(t') = \left(\frac{qD}{pv}\operatorname{erf}\left(\frac{pvt'}{\sqrt{4qDt'}}\right) - \frac{pvt'}{2}\operatorname{erfc}\left(\frac{pvt'}{\sqrt{4qDt'}}\right) + \sqrt{\frac{qDt'}{\pi}}e^{-\frac{p^2v^2t'}{4qD}}\right)^{-1}$$
. (111b)

Finally, returning to the original coordinate system, we have

$$\Psi(x,t) = Q(t) \left(\frac{e^{-\frac{pv(a-x)}{qD}} \left(1 + \operatorname{erf}\left(\frac{pvt-(a-x)}{\sqrt{4qDt}}\right)\right) + \operatorname{erfc}\left(\frac{pvt+(a-x)}{\sqrt{4qDt}}\right)}{\frac{2qD}{pv} \operatorname{erf}\left(\frac{pvt}{\sqrt{4qDt}}\right) - pvt \operatorname{erfc}\left(\frac{pvt}{\sqrt{4qDt}}\right) + \sqrt{\frac{4qDt}{\pi}}e^{-\frac{p^2v^2t}{4qD}}} \right).$$
(112)

The approximation we use is

$$\Psi(x,t) \approx Q(t) h(t) e^{h(t)(x-a)}.$$
(113)

To find the area redistributed, we scale Ψ by the total area of the original PDE X_0 and its image X_{-1} , that is to the right of the right most boundary at x = a. The image Ψ_{-1} is the only image, since we do not treat the right boundary as reflecting the original (traveling) distribution (see 6). The original distribution and its image is given by

$$X_{0}(x,t) = \frac{1}{\sqrt{4\pi q D t}} e^{-\frac{(x - (x_{0} + pvt))^{2}}{4q D t}},$$

$$X_{-1}(x,t) = \frac{1}{\sqrt{4\pi q D t}} e^{-\frac{(x + (x_{0} + pvt))^{2}}{4q D t}}.$$
(114)

The total area redistributed is then

$$Q(t) = \int_{a}^{\infty} \left(X_{0}(x,t) + X_{-1}(x,t) \right) dx$$

= $\frac{1}{\sqrt{4\pi q D t}} \int_{a}^{\infty} \left(e^{-\frac{(x - (x_{0} + pvt))^{2}}{4q D t}} + e^{-\frac{(x + (x_{0} + pvt))^{2}}{4q D t}} \right) dx$
= $1 + \frac{1}{2} \left(\operatorname{erf} \left(\frac{x_{0} - a + pvt}{\sqrt{4q D t}} \right) - \operatorname{erf} \left(\frac{(x_{0} + a + pvt)}{\sqrt{4q D t}} \right) \right).$ (115)



Figure 6: The primary distribution crosses the boundary at x = 0 and x = a. The left tail of the pdf can be replaced within the boundaries of the problem by adding the tail of its "reflected" distribution centered at $x = -x_0 - vt$ on the left. To the right of a we integrate the area of the tails of the original distribution and its image. This area (colored yellow) is distributed within the boundaries as an exponential pdf with its maximum value at x = a. This distribution is fixed in location and has infinite images as in the case for the y-axis. Each of these is superposed, as before, within the practical boundary and is shown as blended colors over the primary distribution.

$$\sum_{m=0}^{\infty} \Psi_m(x,t) = \sum_{m=0}^{\infty} \left(Q(t) h(t) e^{h(t)(x-(2m+1)a)} + Q(t) h(t) e^{h(t)(-x-(2m+1)a)} \right) \quad \text{for} \quad 0 \le x \le a$$
$$= Q(t) h(t) \sum_{m=0}^{\infty} \left(e^{h(t)x} e^{-h(t)a} e^{-2mh(t)a} + e^{-h(t)x} e^{-h(t)a} e^{-2mh(t)a} \right)$$
$$= Q(t) h(t) e^{-h(t)a} \left(e^{h(t)x} + e^{-h(t)x} \right) \sum_{m=0}^{\infty} e^{-2mh(t)a} v \tag{117}$$

Note that $(e^{h(t)x} + e^{-h(t)x}) = 2\cosh[h(t)x]$, so then

$$\sum_{m=0}^{\infty} \Psi_m(x,t) = Q(t) 2h(t) e^{-h(t)a} \cosh[h(t)x] \sum_{m=0}^{\infty} \left(e^{-2h(t)a}\right)^m.$$
(118)

Note that $\sum_{m=0}^{\infty} \left(e^{-2h(t)a}\right)^m$ is a geometric series, equal to $\left(1 - e^{-2h(t)a}\right)^{-1}$, hence,

$$\sum_{m=0}^{\infty} \Psi_m(x,t) = Q(t) 2h(t) \frac{e^{-h(t)a}}{1 - e^{-2h(t)a}} \cosh[h(t)x] = Q(t) h(t) \frac{\cosh[h(t)x]}{\sinh[h(t)a]},$$
(119)

where the sinh term came from the fact that

$$\frac{e^{-h(t)a}}{1 - e^{-2h(t)a}} = \frac{1}{e^{h(t)a} - e^{-h(t)a}} = \frac{1}{2\sinh(h(t)a)}.$$

Putting the pieces together, we get

$$(X_0(x,t) + X_{-1}(x,t)) + \sum_{m=0}^{\infty} \Psi_m(x,t)$$

= $\frac{1}{\sqrt{4\pi q D t}} \left(e^{-\frac{(x-x_0-pvt)^2}{4q D t}} + e^{-\frac{(x+x_0+pvt)^2}{4q D t}} \right) + Q(t) h(t) \frac{\cosh(h(t)x)}{\sinh(h(t)a)}.$ (120)

Finally, multiplying the x and y oriented distributions, we have the following solution on 0 < x < a

$$P(x, y, t) = \left(\frac{1}{\sqrt{4\pi q D t}} \left(e^{-\frac{(x-x_0-pvt)^2}{4q D t}} + e^{-\frac{(x+x_0+pvt)^2}{4q D t}}\right) + Q(t) h(t) \frac{\cosh(h(t)x)}{\sinh(h(t)a)}\right) \\ \times \frac{1}{\sqrt{4\pi q D t}} \sum_{k=-\infty}^{\infty} \left(e^{-\frac{(y-y_0+2kb)^2}{4q D t}} + e^{-\frac{(y+y_0+(2k+1)b)^2}{4q D t}}\right),$$
(121a)

where
$$Q(t) = 1 + \frac{1}{2} \left(\operatorname{erf} \left(\frac{x_0 - a + pvt}{\sqrt{4qDt}} \right) - \operatorname{erf} \left(\frac{(x_0 + a + pvt)}{\sqrt{4qDt}} \right) \right)$$
 (121b)

and
$$h(t) = \left(\frac{qD}{pv}\operatorname{erf}\left(\frac{pvt}{\sqrt{4qDt}}\right) - \frac{pvt}{2}\operatorname{erfc}\left(\frac{pvt}{\sqrt{4qDt}}\right) + \sqrt{\frac{qDt}{\pi}}e^{-\frac{p^2v^2t}{4qD}}\right)^{-1}$$
. (121c)

As t gets very large, we have the following limits

$$\lim_{t \to \infty} \frac{1}{\sqrt{4\pi qDt}} \left(e^{-\frac{(x-x_0-pvt)^2}{4qDt}} + e^{-\frac{(x+x_0+pvt)^2}{4qDt}} \right) = 0.$$

$$\lim_{t \to \infty} \left(\frac{1}{2} \left(2 + \operatorname{erf} \left(\frac{pvt-a+x_0}{\sqrt{4qDt}} \right) - \operatorname{erf} \left(\frac{pvt+a+x_0}{\sqrt{4qDt}} \right) \right) \right) = 1.$$

$$\lim_{t \to \infty} \frac{1}{\sqrt{4\pi qDt}} \sum_{k=-\infty}^{\infty} \left(e^{-\frac{(y-y_0+2kb)^2}{4qDt}} + e^{-\frac{(y+y_0+(2k+1)b)^2}{4qDt}} \right) = \frac{1}{b}.$$
(122)
$$\lim_{t \to \infty} \frac{1}{qDt} \left(1 + \frac{1}{2} \right) = \frac{1}{b}.$$

$$\lim_{t \to \infty} \frac{\cosh(h(t)x)}{\sinh(h(t)a)} = \frac{\cosh\left(\frac{pvx}{qD}\right)}{\sinh\left(\frac{pva}{qD}\right)}.$$

$$P(x, y, \infty) = \frac{pv}{qDb} \frac{\cosh\left(\frac{pvx}{qD}\right)}{\sinh\left(\frac{pva}{qD}\right)} \quad \left\{ \begin{array}{c} 0 < x < a \\ -\frac{b}{2} < y < -\frac{b}{2} \end{array} \right\}.$$
(123)

This is the steady-state solution for the probability distribution.

3.2 Median time to first arrival at goal.

We can also use the expression for Q(t) to obtain the median time to reach the goal. This is the time, t_M , defined when $Q(t_M) = 1/2$.

$$Q(t_M) = \frac{1}{2} = 1 + \frac{1}{2} \left(\operatorname{erf} \left(\frac{x_0 - a + pvt_M}{\sqrt{4qDt_M}} \right) - \operatorname{erf} \left(\frac{(x_0 + a + pvt_M)}{\sqrt{4qDt_M}} \right) \right)$$
$$1 = \operatorname{erf} \left(\frac{(x_0 + a + pvt_M)}{\sqrt{4qDt_M}} \right) - \operatorname{erf} \left(\frac{x_0 - a + pvt_M}{\sqrt{4qDt_M}} \right)$$
(124)

This, of course, has to be solved numerically.

3.3 Summary

Summarizing our results, the probability distribution with the original parametrization is

$$P(x, y, t) dxdy = \left(\frac{1}{\sqrt{4\pi qDt}} \left(e^{-\frac{(x-x_0-pvt)^2}{4qDt}} + e^{-\frac{(x+x_0+pvt)^2}{4qDt}}\right) + Q(t) h(t) \frac{\cosh\left(h\left(t\right)x\right)}{\sinh\left(h\left(t\right)a\right)}\right)$$

$$\times \frac{1}{\sqrt{4\pi qDt}} \sum_{k=-\infty}^{\infty} \left(e^{-\frac{(y-y_0+2kb)^2}{4qDt}} + e^{-\frac{(y+y_0+(2k+1)b)^2}{4qDt}}\right) dxdy \tag{125}$$
where $Q(t) = 1 + \frac{1}{2} \left(\operatorname{erf} \left(\frac{x_0 - a + pvt}{\sqrt{4qDt}}\right) - \operatorname{erf} \left(\frac{(x_0 + a + pvt)}{\sqrt{4qDt}}\right) \right)$
and $h(t) = \left(\frac{qD}{pv} \operatorname{erf} \left(\frac{pvt}{\sqrt{4qDt}}\right) - \frac{pvt}{2} \operatorname{erfc} \left(\frac{pvt}{\sqrt{4qDt}}\right) + \sqrt{\frac{qDt}{\pi}} e^{-\frac{p^2v^2t}{2qD}}\right)^{-1},$

or rescaling time and distance, using the following transformations

$$\begin{aligned} x &= \frac{2qD}{pv}\xi & x_0 = \frac{2qD}{pv}\xi_0 & a = \frac{2qD}{pv}\alpha\\ y &= \frac{2qD}{pv}\zeta & y_0 = \frac{2qD}{pv}\zeta_0 & b = \frac{2qD}{pv}\beta\\ & t = \frac{4qD}{p^2v^2}\theta. \end{aligned}$$

we also have,

$$P\left(\xi,\zeta,\theta\right)d\xi d\zeta = \left(\frac{1}{2\sqrt{\pi\theta}} \left(e^{-\frac{\left(\xi-\xi_{0}-2\theta\right)^{2}}{4\theta}} + e^{-\frac{\left(\xi+\xi_{0}+2\theta\right)^{2}}{4\theta}}\right) + Q\left(\theta\right)h\left(\theta\right)\frac{\cosh\left(h\left(\theta\right)\xi\right)}{\sinh\left(h\left(\theta\right)\alpha\right)}\right)$$
$$\times \frac{1}{2\sqrt{\pi\theta}} \sum_{k=-\infty}^{\infty} \left(e^{-\frac{\left(\zeta-\zeta_{0}+2k\beta\right)^{2}}{4\theta}} + e^{-\frac{\left(\zeta+\zeta_{0}+\left(2k+1\right)\beta\right)^{2}}{4\theta}}\right)d\xi d\zeta$$
where $Q\left(\theta\right) = 1 + \frac{1}{2}\operatorname{erf}\left(\frac{\xi_{0}-\alpha+2\theta}{2\sqrt{\theta}}\right) - \frac{1}{2}\operatorname{erf}\left(\frac{\xi_{0}+\alpha+2\theta}{2\sqrt{\theta}}\right)$
and $h\left(\theta\right) = 2\left(\operatorname{erf}\left(\sqrt{\theta}\right) - 2\theta\operatorname{erfc}\left(\sqrt{\theta}\right) + 2\sqrt{\frac{\theta}{\pi}}e^{-\theta}\right)^{-1}.$

4 Calculation of the probability of reaching the goal by a given point in time, and the probability of placing at that time.



Figure 7: Evolution of the probability distribution at three points in time.

The probability of fish species 1 reaching a point along the right-hand boundary (located at x = a) by time t is given by $Q_1(t)$ (the subscript indicating species 1)

$$Q_{1}(t) = \int_{a}^{\infty} \frac{1}{\sqrt{4\pi q_{1}D_{1}t}} \left(e^{-\frac{(x-x_{0}-p_{1}v_{1}t)^{2}}{4q_{1}D_{1}t}} + e^{-\frac{(x+x_{0}+p_{1}v_{1}t)^{2}}{4q_{1}D_{1}t}} \right) dx$$

= $1 + \frac{1}{2} \left(\operatorname{erf} \left(\frac{p_{1}v_{1}t + x_{0} - a}{\sqrt{4q_{1}D_{1}t}} \right) - \operatorname{erf} \left(\frac{p_{1}v_{1}t + x_{0} + a}{\sqrt{4q_{1}D_{1}t}} \right) \right).$ (126)

The probability of fish species 2 (subscript 2) not reaching a point along the right hand boundary (located at x = a) by time t is given by $1 - Q_2(t)$

$$1 - Q_2(t) = \frac{1}{2} \left(\operatorname{erf}\left(\frac{p_2 v_2 t + x_0 + a}{\sqrt{4q_2 D_2 t}}\right) - \operatorname{erf}\left(\frac{p_2 v_2 t + x_0 - a}{\sqrt{4q_2 D_2 t}}\right) \right).$$
(127)

Thus, the probability of fish species 1, beating fish species 2 by time t, is given by the product $Q_1(t)(1-Q_2(t))$:

$$\left[1 + \frac{1}{2} \left(\operatorname{erf} \left(\frac{p_1 v_1 t + x_0 - a}{\sqrt{4q_1 D_1 t}} \right) - \operatorname{erf} \left(\frac{p_1 v_1 t + x_0 + a}{\sqrt{4q_1 D_1 t}} \right) \right) \right] \times \frac{1}{2} \left(\operatorname{erf} \left(\frac{p_2 v_2 t + x_0 + a}{\sqrt{4q_2 D_2 t}} \right) - \operatorname{erf} \left(\frac{p_2 v_2 t + x_0 - a}{\sqrt{4q_2 D_2 t}} \right) \right).$$
(128)

In general, he probability of fish species k arriving first of n fish species is given by

$$\frac{Q_k(t)}{(1 - Q_k(t))} \prod_{i=1}^n (1 - Q_i(t)).$$
(129)



Figure 8: Competition between two species. Probability that a given species will arrive first (left axis, solid curves). Probability that neither species has arrived or that both have arrived (pink and brown dashed lines respectively, scale on right axis).

The relative proportion of arrivals of species k by time t out of all n species is

$$F(t, n, k) = \frac{N_k Q_k(t)}{\sum_{i=1}^n N_i Q_i(t)},$$
(130)

where N_i is the total number of individuals in species i.

We need to be a bit more specific with function arguments so that we can specify order of arrival of distributions for placing second, third, and so on. For the marginal distribution on the x axis, we let

$$P_{i}(x,t) = X_{0}(x,t) + X_{-1}(x,t), \qquad (131)$$

and the probability of passing point x is given by

$$Q_i(x,t) = \int_x^\infty P_i(\xi,t)d\xi$$

= $1 + \frac{1}{2} \left(\operatorname{erf} \left(\frac{x_0 - x + p_i v_i t}{\sqrt{4q_i D_i t}} \right) - \operatorname{erf} \left(\frac{x_0 + x + p_i v_i t}{\sqrt{4q_i D_i t}} \right) \right).$ (132)

Second Place is

$$\sum_{i \neq k} \frac{\int_{a}^{\infty} P_{k}(x,t) Q_{i}(x,t) dx}{(1 - Q_{i}(a,t))} \frac{\prod_{j=1}^{n} (1 - Q_{j}(t))}{Q_{k}(a,t) (1 - Q_{k}(a,t))}.$$
(133)



Figure 9: Competition between three species. Probability that a given species will arrive first.

Third place is

$$\sum_{i \neq k} \sum_{j \neq k} \frac{\int_{a}^{\infty} P_{k}(x,t) Q_{i}(x,t) Q_{j}(x,t) dx}{(1 - Q_{i}(a,t)) (1 - Q_{j}(a,t))} \frac{\prod_{h=1}^{n} (1 - Q_{h}(t))}{Q_{k}(a,t) (1 - Q_{k}(a,t))}.$$
(134)

And so on.

5 Discussion

Though we have used examples of an individual's movements to illustrate the rationale behind the development of equations, in truth the distributions and movements described are a picture of the actions of an infinite number of individuals each starting from the same conditions and acting under the same rules. Thus, though an individual fish may swim here or there and never appear to have any goal in mind, it may be found that, over time, the whole school moves toward one destination as if the school itself had a goal and self determination. Yet, if we knew ahead of time the arrow of that mass movement and we broke down the components of the movements of each individual along that axis and its perpendicular, we would find that the whole of the action would be just an infinitesimal excess of desire for a slight majority of individuals of movement in that one direction. It is thus for creatures as it is for molecules of air in a breeze, though each moves in any direction, on average they all move toward one direction. When confined to an enclosure, they eventually pile up at the boundary toward



Figure 10: Competition between two species. Fraction of the total number of individuals that members of a given species comprise. (If the probability of a individual arriving is less than 10^{-4} then probability is set to 10^{-4} , this prevents division by very small numbers.)

which they tend. This changes their distribution in space from a dispersing bell shaped curve to a steady state exponential shaped curve, as the forces of diffusion and drift balance one another.



Figure 11: Competition between three species. Fraction of the total number of individuals that members of a given species comprise. (If the probability of a individual arriving is less than 10^{-4} then probability is set to 10^{-4} , this prevents division by very small numbers.)

References

- [Skalski and Gilliam(2000)] Garrick T. Skalski and James F. Gilliam. Modeling diffusive spread in a heterogeneous population: A movement study with stream fish. *Ecology*, 81 (6), 2000.
- [Sparrevohn et al.(2002)Sparrevohn, Nielsen, and Støttrup] Claus R. Sparrevohn, Anders Nielsen, and Josianne G. Støttrup. Diffusion of fish from a single release point. Can. J. Fish. Aquat. Sci., 59:844–853, 2002.
- [Faugeras and Maury(2007)] Blaise Faugeras and Olivier Maury. Modeling fish population movements: From an individual-based representation to an advection-diffusion equation. J. Theoretical Biology, 247:837–848, 2007.
- [Kerckhove et al.(2015)Kerckhove, Milne, Shuter, and Abrams] Derrick T. de Kerckhove, Scott Milne, Brian J. Shuter, and Peter A. Abrams. Ideal gas model adequately describes movement and school formation in a pelagic freshwater fish. *Behavioral Ecology*, 26(4), 2015.
- [Berg(1993)] Howard C. Berg. Random Walks in Biology. Princeton University Press, Princeton, NJ, 2nd edition, 1993. ISBN 0691000646.
- [Cushman-Roisin(2012)] Benoit Cushman-Roisin. Environmental Transport and Fate (ENGS 43: course notes). Thayer School of Engineering, Dartmouth College, Hanover, N.H., 2012. URL http://www.dartmouth.edu/~cushman/courses/engs43.html.
- [Zwillinger(1996)] Daniel Zwillinger, editor. CRC Standard Mathematical Tables and Formulae. CRC Press, Boca Raton, FL, 30th edition, 1996. ISBN 0849324793.