EXISTENCE OF η -QUOTIENTS FOR SQUAREFREE LEVELS

MICHAEL ALLEN

ABSTRACT. In this paper, we investigate which modular form spaces can contain η quotients, functions of the form $f(\tau) = \prod_{\delta \mid N} \eta(\delta \tau)^{r_{\delta}}$. For $k \geq 2$ even and N coprime to 6, we give necessary conditions for the space $M_k(\Gamma_1(N))$ to contain η -quotients. We then show that these conditions are sufficient as well for a large family of squarefree levels N coprime to 6.

1. INTRODUCTION AND STATEMENT OF RESULTS

Dedekind's η function, defined by

$$\eta(\tau) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1-q^n),$$

where τ is in the upper-half of the complex plane and $q := e^{2\pi i \tau}$, is one of the most famous and classical examples of a weight $\frac{1}{2}$ modular form. The infinite product formula makes $\eta(\tau)$ particularly nice to work with for many computations. As such, it is interesting to know when a modular form can be written as a linear combination of products and quotients of η -functions. An η -quotient is a function of the form

$$f(\tau) = \prod_{\delta \mid N} \eta(\delta \tau)^{r_{\delta}},$$

where each $r_{\delta} \in \mathbb{Z}$.

Due to the nice form of $\eta(\tau)$, a question of interest is which modular objects can be expressed as an η -quotient or a linear combination of η -quotients. In [11], Ono shows that every holomorphic modular form for the entire modular group $\operatorname{SL}_2(\mathbb{Z})$ can be written as a rational function of $\eta(z)$, $\eta(2z)$, and $\eta(4z)$. Martin and Ono [7] have classified all elliptic curves which are represented by η -quotients via the modularity theorem. Lemke Oliver [10] has classified all θ -functions which are η -quotients. Additional work on spaces of modular forms spanned by η -quotients has been done by Arnold-Roksandich, James, and Keaton [2] and Rouse and Webb [12], for example. In [1], the author et al. investigate the question of which modular form spaces cannot contain η -quotients; these results are extended in this paper.

Throughout the paper we use the following notation. Given a subgroup $\Gamma \subset \mathrm{SL}_2(\mathbb{Z})$, we use $S_k(\Gamma)$, $M_k(\Gamma)$, and $M_k^!(\Gamma)$ to denote the complex vector spaces of weight k cusp forms, holomorphic modular forms, and weakly holomorphic forms, respectively, for the group Γ . We use $M(\Gamma)$ to denote the graded ring of all holomorphic modular forms for Γ . In the case where $\Gamma = \Gamma_0(N)$, we use $S_k(\Gamma_0(N), \chi)$, $M_k(\Gamma_0(N), \chi)$, and $M_k^!(\Gamma_0(N), \chi)$ to denote the spaces of weight k cusp forms, holomorphic modular forms, and weakly

Date: November 4, 2019.

holomorphic forms, respectively, for the group $\Gamma_0(N)$ with Nebentypus χ .

The following theorem, originally appearing in work of Newman [8, 9] as well as Gordon and Hughes [4], is useful for determining when a given η -quotient is a weakly holomorpic modular form.

Theorem 1.1 (Theorem 1.64, [11]). Let f be the η -quotient of level N given by

$$f(\tau) = \prod_{\delta \mid N} \eta(\delta \tau)^{r_{\delta}}.$$

If f satisfies both

$$\sum_{\delta|N} \delta r_{\delta} \equiv 0 \pmod{24}$$
$$\sum_{\delta|N} \frac{N}{\delta} r_{\delta} \equiv 0 \pmod{24},$$

then for $k = \frac{1}{2} \sum_{\delta \mid N} r_{\delta}$ and $\chi(d) = \left(\frac{(-1)^{k_s}}{d}\right)$, where $s = \prod_{\delta \mid N} \delta^{r_{\delta}}$, we have $f \in M_k^!(\Gamma_0(N), \chi)$.

Remark 1. In the case where N is coprime to 6, the two congruences in Theorem 1.1 are equivalent, as any number coprime to 6 is its own inverse modulo 24. Additionally, the converse to Theorem 1.1 holds when N is coprime to 6. Moreover, any weakly holomorphic η -quotient on $\Gamma_1(N)$ must also satisfy the hypotheses of Theorem 1.1 (see Corollary 1.8 of [1]). That is, given $f(\tau) = \prod_{\delta \mid N} \eta(\delta \tau)^{r_{\delta}} \in M_k^!(\Gamma_1(N))$, we have

$$\sum_{\delta|N} \delta r_{\delta} \equiv 0 \pmod{24},$$

and hence $f(\tau) \in M_k^!(\Gamma_0(N), \chi)$ for some Nebentypus χ . Thus, every η -quotient which is modular for $\Gamma_1(N)$ is in fact modular for $\Gamma_0(N)$ whenever gcd(N, 6) = 1. Because of this, in working with η -quotients on $\Gamma_1(N)$, we only need to consider the behavior at the cusps of $\Gamma_0(N)$.

In work of Martin [6], the following is shown to be a complete set of representatives for the cusps of $\Gamma_0(N)$:

$$C_{\Gamma_0(N)} = \left\{ \frac{a_c}{c} \in \mathbb{Q} \mid c \mid N, 1 \le a_c \le N, \gcd(a_c, N) = 1 \\ \text{and } a_c \equiv a'_c \pmod{\gcd(c, \frac{N}{c})} \iff a_c = a'_c \right\}$$

When N is squarefree, as will be the case throughout this paper, this can be reduced to

$$C_{\Gamma_0(N)} = \left\{ \frac{1}{d} : d \mid N \right\}.$$
(1)

In order to determine when η -quotients are fully holomorphic modular forms, we need to compute orders of vanishing at the cusps. We do so using a theorem of Ligozat [5], also appearing in work of Biagioli [3] and Martin [6].

Theorem 1.2 (Theorem 1.65, [11]). N Let c, d, and N be positive integers with d | N and gcd(c, d) = 1. Then if $f(\tau)$ is an η -quotient satisfying the conditions given in Theorem 1.1 for N, then the order of vanishing for $f(\tau)$ at the cusp c/d relative to $\Gamma_0(N)$ is

$$v_{\frac{c}{d}} := v_{\Gamma_0(N)}\left(f, \frac{c}{d}\right) = \frac{N}{24} \sum_{\delta \mid N} \frac{\gcd(d, \delta)^2 r_{\delta}}{\gcd(d, \frac{N}{d}) d\delta}$$

The next theorem, first appearing in work of Rouse and Webb [12], says that for $M(\Gamma_0(N))$ to be generated by η -quotients, N must be sufficiently composite relative to its size.

Theorem 1.3 ([12]). Suppose that $f(\tau) = \prod_{\delta \mid N} \eta(\delta \tau)^{r_{\delta}} \in M_k(\Gamma_0(N), \chi)$. Then we have

$$\sum_{\delta|N} |r_{\delta}| \le 2k \prod_{p|N} \left(\frac{p+1}{p-1}\right)^{\min\{2, \operatorname{ord}_p(N)\}}$$

This bound implies the existence of an upper bound on the number of η -quotients in any space $M_k(\Gamma_0(N), \chi)$ which depends only on the number of prime divisors of N. In particular, as dim $(M_k(\Gamma_0(N), \chi))$ tends towards infinity as N does, there can only exist a finite number of levels with a given number of prime divisors such that the ring $M(\Gamma_0(N))$ is generated by η -quotients.

In [1], the author et al. investigate other factors in addition to Theorem 1.3 which inhibit the existence of η -quotients in certain modular form spaces. The first result from [1] gives a complete categorization of which space $M_k(p)$ contain η -quotients when $k \ge 2$ is even and p > 5 is prime.

Theorem 1.4 (Theorem 1.9 [1]). Let p be prime, set $h_p = \frac{1}{2} \operatorname{gcd}(p-1, 24)$, and let k be an even integer. Then there exists $f(\tau) = \eta^{r_1}(\tau)\eta^{r_p}(p\tau) \in M_k(\Gamma_1(p))$ if and only if both of the following conditions hold.

- (i) $h_p \mid k$
- (ii) It is not the case that $k = 2, p \equiv 5 \pmod{24}$, and $p \neq 5$.

The next theorem gives a similar classification for spaces $M_k(\Gamma_1(pq))$, where again $k \ge 2$ is even and p, q > 5 are distinct primes.

Theorem 1.5 (Theorem 1.10 [1]). Let p, q be distinct primes and let N = pq. Let $k \ge 2$ be even, and define $h_N := \frac{1}{2} \operatorname{gcd} \{p-1, q-1, 24\}$. There exists an η -quotient $f(\tau) \in M_k(\Gamma_1(N))$ if and only if

- (i) $h_N \mid k$
- (ii) It is not the case that k = 2, $(p,q) \equiv (1,5)$, (5,1), or $(5,5) \pmod{24}$, and neither p nor q is equal to 5.

These results generalize to squarefree levels as follows. First, define

$$S := \left\{ \prod_{i=1}^{\ell} p_i^{e_i} \middle| \begin{array}{c} \text{For all } i, p_i > 5 \text{ prime, } p_i \equiv 1 \text{ or } 5 \pmod{24} \\ \text{and at least one } p_i \equiv 5 \pmod{24} \end{array} \right\}$$

Note that with this definition, condition (ii) in Theorem 1.4 (resp. Theorem 1.5) can be rephrased as "It is not the case that k = 2 and $p \in S$ (resp. $pq \in S$)". Our first result generalizes the forwards direction of Theorem 1.5 to arbitrary levels coprime to 6.

Theorem 1.6. Let $N = p_1^{e_1} p_2^{e_2} \cdots p_{\ell}^{e_{\ell}}$ be an integer coprime to 6, let $k \ge 2$ be even, and define $h_N = \frac{1}{2} \operatorname{gcd} \{p_1 - 1, p_2 - 1, \dots, p_{\ell} - 1, 24\}$. There exists an η -quotient in $M_k(\Gamma_1(N))$ if

(i) $h_N \mid k$ (ii) It is not the case that k = 2 and $N \in S$.

MICHAEL ALLEN

Theorems 1.4 and 1.5 suggest that $M_2(\Gamma_1(N))$ cannot contain η -quotients if the prime divisors of N fall into certain residue classes modulo 24. On the other hand, Theorem 1.3 tells us that the more composite a number is, the easier it is to have η -quotients of that level. This complicates proving the converse of Theorem 1.6, as the bounding term in Theorem 1.3 can be made arbitrarily large as N ranges through all squarefree integers. However, we are able to give the following partial converse to Theorem 1.6.

Theorem 1.7. Let $N \in S$ be squarefree and coprime to 6, and let p_1 denote the smallest prime dividing N. If

$$4\prod_{p|N} \frac{p+1}{p-1} < p_1 + 1, \tag{2}$$

then there is no η -quotient in $M_2(\Gamma_1(N))$.

The product in the left-hand side of (2) is made larger the smaller the prime divisors of N are. Thus, the smallest number N for which this extra hypothesis fails is

$$N = \prod_{\substack{\text{prime } p \le M \\ p \equiv 1 \text{ or } 5 \pmod{24}}} p,$$

where

$$M := \inf \left\{ x \quad \middle| \begin{array}{c} 4 \times \prod_{\substack{\text{prime } p \le x \\ p \equiv 1 \text{ or } 5 \pmod{24}}} \left(\frac{p+1}{p-1} \right) \ge 30 \right\}.$$

The product

$$\prod_{\substack{\text{prime } p \le x \\ p \equiv 1 \text{ or } 5 \pmod{24}}} \left(\frac{p+1}{p-1}\right)$$

grows numerically like $\log \log \log(x)$, and so M will be extremely large. As N is then the product of all primes congruent to 1 or 5 (mod 24) up to M, N must be astronomically large before this hypothesis can fail. We do not claim that the converse is false when N becomes large enough for (2) to fail, only that different techniques from ours would be needed to address this case.

The remainder of this paper is devoted to proving these two theorems. Theorem 1.6 is handled in Section 2, and Section 3 deals with the proof of Theorem 1.7.

2. Modular form spaces which contain η -quotients

In order for there to exist η -quotients which are fully holomorphic modular forms for $\Gamma_1(N)$ and weight k, there certainly must at least exist weakly holomorphic η -quotients. In [1], the authors et al. obtained the following result for existence of weakly holomorphic η -quotients.

Theorem 2.1 (Theorem 4.1, [1]). Let $N = p_1^{e_1} p_2^{e_2} \cdots p_{\ell}^{e_{\ell}}$ with each $p_i \geq 5$, and define $h_N = \frac{1}{2} \operatorname{gcd}(p_1 - 1, \cdots, p_{\ell} - 1, 24)$. There exists an η -quotient $f(\tau) = \prod_{\delta \mid N} \eta^{r_{\delta}}(\delta \tau) \in M_K^!(\Gamma_1(N))$ if and only if $h_N \mid k$.

Remark 2. In [1], the above theorem is only stated to hold for N squarefree. However, the proof makes no use of this assumption, and so still holds for this generalized version.

Before proving Theorem 1.6 in full generality, we will prove it for the specialized case where N is squarefree. We begin with a lemma which allows us to reduce this case to Theorem 1.5.

Lemma 2.2. Let $N = p_1 p_2 \cdots p_\ell$ with $\ell \geq 2$. Then there exist distinct primes p and q dividing N such that $h_N = h_{pq}$. Moreover, if $N \notin S$, then p and q can be chosen such that $pq \notin S$.

Proof. We begin by setting some notation. For an integer a, let \tilde{a} denote the least residue of $a \mod 24$. Let R_N be the set of least residues of N's prime divisors mod 24. That is,

$$R_N = \{\tilde{p_1}, \tilde{p_2}, \dots, \tilde{p_\ell}\}.$$

By definition, $p_1 = \tilde{p_1} + 24m$ for some integer m. Observe that if $d \mid p_1 - 1$ and $d \mid 24$, then $d \mid (p_1 - 1 - 24m) = \tilde{p_1} - 1$. Similarly, any divisor of both $\tilde{p_1} - 1$ and 24 must also divide $p_1 - 1$, and so in particular,

$$h_N = \frac{1}{2} \operatorname{gcd} \left\{ p_1 - 1, p_2 - 1, \cdots, p_{\ell} - 1, 24 \right\} = \frac{1}{2} \operatorname{gcd} \left\{ \tilde{p}_1 - 1, \tilde{p}_2 - 1, \cdots, \tilde{p}_{\ell} - 1, 24 \right\}.$$

That is, h_N depends only on R_N , the residues of the prime divisors of N modulo 24. We proceed to the proof by cases on the value of h_N . As h_N is one half an even divisor of 24, the possible values of h_N are 12, 6, 4, 3, 2, and 1. Note that if $N \in S$, then $R_N = \{5\}$ or $\{1, 5\}$, and so $h_N = 2$. Thus, in all cases excluding $h_N = 2$, we may assume that $N \notin S$.

If $h_N = 12$, we equivalently have that every prime divisor of N is 1 (mod 24), and so $R_N = \{1\}$. Letting p and q be any divisors of N, we again have $R_{pq} = \{1\}$, and so $h_{pq} = h_N$. Additionally, as $p \equiv q \equiv 1 \pmod{24}$, $pq \notin S$.

Next, if $h_N = 6$, then every prime divisor is congruent to 1 (mod 12), and so $R_N \subseteq \{1, 13\}$. However, R_N must contain 13, as otherwise we would have $h_N = 12$. Let $p \mid N$ be such that $p \equiv 13 \pmod{24}$, and let q be any other prime divisor of N. Then R_{pq} is either $\{1, 13\}$ or $\{13\}$ depending on the residue class of q, but in either case $h_{pq} = 6 = h_N$. As $p \equiv 13 \pmod{24}$, we also have that $pq \notin S$, and so gives the desired divisor. The case $h_N = 4$ is very similar. For this to hold, we must have $R_N = \{1, 17\}$ or $\{17\}$, and in either case so long as we choose $p \equiv 17 \pmod{24}$ then any choice of q will suffice.

If $h_N = 3$, then every prime divisor of N is 1 (mod 6), and so $R_N \subseteq \{1, 7, 13, 19\}$. But R_N must contain at least one of 7 or 19, as otherwise we would have $h_N = 6$ or 12. We may thus pick some prime divisor p of N which belongs to one of these two residue classes. Let q be any other prime divisor of N. The possible residue sets for pq are

$$\{7\},\{19\},\{1,7\},\{7,13\},\{7,19\},\{1,19\}, \text{ and } \{13,19\}$$

One can quickly check that for each of these possibilities, $h_{pq} = 3$. Additionally, as p is not congruent to 1 or 5 (mod 24), it follows that $pq \notin S$.

For $h_N = 2$, we have a similar situation. The residue set must satisfy $R_N \subseteq \{1, 5, 13, 17\}$, and either 5 or 17 must belong to R_N , otherwise h_N will be larger than 2. We first consider the case where $R_N = \{1, 5\}$ or $\{5\}$. First, if $N \in S$, then we may choose $p \equiv 5$ (mod 24) and take q to be any other prime divisor, and we have $h_{pq} = 2 = h_N$. On the other hand, if $N \notin S$ then we must have $5 \mid N$. Letting q be any divisor of N distinct from 5, $5q \notin S$ and $h_{5q} = 2 = h_N$. Next suppose $5 \notin R_N$. This implies $13 \in R_N$, as if it

MICHAEL ALLEN

were not we would have $h_N = 4$. Hence, we may choose divisors $p \equiv 17$ and $q \equiv 13 \pmod{24}$, which will suffice. Finally, if $5 \in R_N$ and $R_N \neq \{5\}$ or $\{1, 5\}$, then taking $p \equiv 5$ and $q \equiv 13$ or 17 will work.

Lastly, we consider the case where $h_N = 1$. As $2 \nmid h_N$, there must exist a divisor p which is congruent to 3 (mod 4). Additionally, as $3 \nmid h_N$, there must exist some prime divisor q which is congruent to either 3 or 5 (mod 6). Our choice of p guarantees that $2 \nmid h_{pq}$, and our choice of q is such that $3 \nmid h_{pq}$. Therefore, as h_{pq} is a divisor of 24 which is divisible by neither 2 nor 3, it follows that $h_{pq} = 1$. Additionally, since $p \equiv 3 \pmod{4}$, $pq \notin S$.

We now have enough to prove the specialization of Theorem 1.6 to the case where N is squarefree.

Proposition 2.3. Let $N = p_1 p_2 \cdots p_\ell$ be squarefree, and let $k \ge 2$ be even. Define $h_N = \frac{1}{2} \operatorname{gcd} \{p_1 - 1, p_2 - 1, \cdots, p_\ell - 1, 24\}$. Then there exists an η -quotient in $M_k(\Gamma_1(N))$ if

(i) $h_N \mid k$

(ii) It is not the case that k = 2 and $N \in S$.

Proof. Let N be as in the hypotheses and let $k \ge 2$. We wish to show that there exists an η -quotient in $M_k(\Gamma_1(N))$ whenever $h_N \mid k$ and either $N \notin S$ or $N \in S$ and $k \ge 4$. If $\ell = 1$ or 2, this proposition reduces to Theorem 1.4 or 1.5 respectively, and so we need only consider the case where $\ell \ge 3$. By Lemma 2.2, there exists a product of two distinct primes $pq \mid N$ such that $h_{pq} = h_N$, and such that if $N \notin S$ then neither is pq. But then, as $h_{pq} = h_N \mid k$, it follows from Theorem 1.5 that there exists an η quotient $f(\tau) \in M_k(\Gamma_1(pq))$. As $pq \mid N, M_k(\Gamma_1(pq)) \subset M_k(\Gamma_1(N))$, and so in particular $f(\tau) \in M_k(\Gamma_1(N))$.

Theorem 1.6 now follows easily.

Proof of Theorem 1.6. Let $N = p_1^{e_1} p_2^{e_2} \cdots p_\ell^{e_\ell}$, and define h_N as above. We want to show that there exists an η -quotient $f(\tau) \in M_k(\Gamma_1(N))$ whenever $h_N \mid k$ and it is not the case that k = 2 and $N \in S$. Define $\widehat{N} = p_1 p_2 \cdots p_\ell$. As membership in S is determined entirely by the set of prime divisors of N, we have that $\widehat{N} \in S$ if and only if $N \in S$. Additionally, $h_N = h_{\widehat{N}}$. By Proposition 2.3, it then follows that there exists an η -quotient $f(\tau) \in M_k(\Gamma_1(\widehat{N}))$ whenever $h_N \mid k$ and it is not the case that k = 2 and $N \in S$. The fact that $M_k(\Gamma_1(\widehat{N})) \subset M_k(\Gamma_1(N))$ completes the proof. \Box

3. Modular form spaces which do not contain η -quotients

In [1], the author et al. use the bound from Theorem 1.3 to show the converse to Theorem 1.6 in the case where N is a prime or a product of two distinct primes. We now investigate how well this converse generalizes to arbitrary squarefree levels. Since holomorphic η -quotients are also weakly holomorphic, Theorem 2.1 shows that there can be no η -quotient in $M_k(\Gamma_1(N))$ whenever $h_N \nmid k$. The impact that membership in S has on existence of η -quotients in modular form spaces will be partially answered by Theorem 1.7. In order to prove Theorem 1.7, we must first make some observations about the structure of η -quotients and their orders of vanishing using Theorem 1.2.

Definition 1. Let $N = p_1 p_2 \cdots p_\ell$ be coprime to 6, and let $f(\tau) \in M_k(\Gamma_1(N))$ be an η -quotient. Note that N has 2^ℓ divisors, given by the 2^ℓ ways we can choose to include or exclude each p_i . Using Theorem 1.2 and (1), we have the following system of 2^ℓ equations associated to the orders of vanishing of f at each cusp of $\Gamma_0(N)$:

$$\left\{24v_{\frac{1}{d}} = \sum_{\delta \mid N} \frac{N \gcd(d, \delta)^2}{d\delta} r_{\delta}\right\}_{d \mid N}$$

Let A_N be the coefficient matrix for the right-hand side of the above system of equations. That is, given some ordering $d_1, d_2, \ldots, d_{2^{\ell}}$ of the divisors of N,

$$A_N = \left(\frac{N \operatorname{gcd}(d_i, d_j)^2}{d_i d_j}\right)_{1 \le i, j \le 2^\ell}$$

Lemma 3.1. Let N be squarefree and coprime to 6, and define A_N as above. Every $c \mid N$ appears exactly once in each column and row of A_N .

Proof. As the expression $\frac{N \operatorname{gcd}(d_i d_j)^2}{d_i d_j}$ is symmetric in d_i and d_j , the matrix A_N is symmetric. Therefore, it suffices to show that c appears exactly once in every row of A_N , as we will then know that c appears exactly once in every column of $A_N^T = A_N$. Additionally, as there are as many rows of A_N as there are divisors or N, we need only show that every divisor appears in each row of A_N to know that it appears exactly once. Hence, we wish to show that, for any fixed $1 \leq i \leq 2^{\ell}$, there exists d_j such that

$$c = \frac{N \operatorname{gcd}(d_i, d_j)^2}{d_i d_j}.$$
(3)

Observe that $gcd(c, d_i) gcd(N/c, N/d_i)$ is a divisor of N, so there exists $1 \le j \le 2^{\ell}$ such that

$$d_i = \gcd(c, d_i) \gcd(N/c, N/d_i).$$

We will show that (3) is satisfied by this choice of j, which will finish the proof.

Because N is squarefree, so too is any divisor, and so every divisor is a product of distinct primes. Thus, in checking that various divisors of N are equal, we need only check that they share the same prime divisors. For example, our choice of d_j can be expressed as the product of the primes dividing both c and d_i and of primes dividing N but dividing neither c nor d_i . That is,

$$d_j = \prod_{\substack{p \text{ prime} \\ p|c \text{ and } p|d_i}} p \times \prod_{\substack{p|N \text{ prime} \\ p\nmid c \text{ and } p\nmid d_i}} p.$$

The primes dividing both d_i and d_j are exactly those appearing in the first term of the above representation of d_j . Thus,

$$gcd(d_i, d_j) = \prod_{\substack{p \text{ prime} \\ p \mid c \text{ and } p \mid d_i}} p = gcd(c, d_i).$$

We can hence reduce the right-hand side of (3) by

$$\frac{N \operatorname{gcd}(d_i, d_j)^2}{d_i d_j} = \frac{N \operatorname{gcd}(c, d_i)}{d_i \operatorname{gcd}(N/c, N/d_i)}.$$
(4)

In the denominator of the right-hand side of (4), we have the term $d_i \operatorname{gcd}(N/c, N/d_i)$, which is the product of all primes which divide d_i and all primes dividing N but dividing neither c nor d_i . That is,

$$d_i \operatorname{gcd}(N/c, N/d_i) = \prod_{\substack{\text{prime } p \mid d_i}} p \times \prod_{\substack{p \mid N \text{ prime } \\ p \nmid c \text{ and } p \nmid d_i}} p.$$
(5)

So, $N/(d_i \operatorname{gcd}(N/c, N/d_i))$ is the product of all prime divisors of N which do not appear in (5). As we are removing every prime which divides d_i and every prime which divides neither c nor d_i , we are left only with the primes dividing c but not d_i . That is,

$$\frac{N}{d_i \operatorname{gcd}(N/c, N/d_i)} = \prod_{\substack{p \mid N \text{ prime} \\ p \mid c \text{ and } p \nmid d_i}} p.$$

On the other hand, $gcd(c, d_i)$ is the product of all primes dividing both c and d_i . Combining these, we have

$$\frac{N \operatorname{gcd}(c, d_i)}{d_i \operatorname{gcd}(N/c, N/d_i)} = \prod_{\substack{p \text{ prime} \\ p|c \text{ and } p \nmid d_i}} p \times \prod_{\substack{p \text{ prime} \\ p|c \text{ and } p|d_i}} p = \prod_{\substack{p \text{ prime} \\ p|c \text{ and } p|d_i}} p = c,$$

showing that (3) holds.

Using this result, we are able to show that in order for $M_k(\Gamma_1(N))$ to contain η -quotients which are truly quotients and not η -products, then we must have k large enough or N sufficiently composite relative to its size in the sense of Theorem 1.3. We make these ideas precise in the following Lemma.

Lemma 3.2. Let N be squarefree and coprime to 6, and let p_1 denote the smallest prime dividing N. If

$$2k \prod_{\substack{p \mid N \\ p \text{ prime}}} \frac{p+1}{p-1} < p_1 + 1,$$

then for any $f(\tau) = \prod_{\delta \mid N} \eta^{r_{\delta}}(\delta \tau) \in M_k(\Gamma_1(N))$, we must have $r_{\delta} \ge 0$ for all $\delta \mid N$, so $f(\tau)$ is an η -product.

Proof. Towards contradiction, suppose $r_d < 0$ for some $d \mid N$. By Theorem 1.2 and Lemma 3.1, the order of vanishing at the cusp of $\Gamma_0(N)$ represented by 1/d takes the form

$$24v_{\frac{1}{d}} = Nr_d + \sum_{\substack{\delta \mid N\\ \delta \neq d}} c_{\delta} r_{\delta},$$

where each $c_{\delta} \mid N, c_{\delta} \neq N$. As $r_d < 0$ is an integer, $r_d \leq -1$, and so

$$Nr_d \le -N.$$
 (6)

As each $c_{\delta} \leq \frac{N}{p_1}$, we also have, for each $d \neq \delta \mid N$,

$$c_{\delta}r_{\delta} \le \frac{N}{p_1}|r_{\delta}|.\tag{7}$$

Together (6) and (7) yield the inequality

$$24v_{\frac{1}{d}} \le -N + \frac{N}{p_1} \sum_{\substack{\delta \mid N \\ \delta \neq d}} |r_{\delta}|.$$

$$\tag{8}$$

By Theorem 1.3 and the hypothesis of the Lemma,

$$|r_d| + \sum_{\substack{\delta \mid N \\ \delta \neq d}} |r_\delta| \le 2k \prod_{p \mid N} \left(\frac{p+1}{p-1}\right) < p_1 + 1.$$

In particular, as $|r_d| \ge 1$ and each $|r_\delta|$ is an integer,

$$\sum_{\substack{\delta \mid N\\ \delta \neq d}} |r_{\delta}| \le p_1 - 1.$$
(9)

Combining (8) and (9) we have

$$24v_{\frac{1}{d}} \le -N + \frac{N}{p_1}(p_1 - 1) = N\left(\frac{p_1 - 1}{p_1} - 1\right) < 0.$$

But this would imply that $f(\tau)$ has a pole at the cusp represented by $\frac{1}{d}$, contradicting the assumption that $f(\tau) \in M_k(\Gamma_1(N))$. Thus, $r_d \ge 0$ for all $d \mid N$.

We now prove Theorem 1.7

Proof of Theorem 1.7. Let N be as in the statement of the theorem. Suppose we had an η -quotient $f(\tau) = \prod_{\delta \mid N} \eta^{r_{\delta}}(\delta \tau) \in M_2(\Gamma_1(N))$. By Lemma 3.2, each $r_{\delta} \geq 0$. As N is coprime to 6, as noted in Remark 1, $f(\tau)$ must satisfy the congruence conditions in Theorem 1.1. That is,

$$\sum_{\substack{\delta \mid N \\ \delta \equiv 1 \pmod{24}}} r_{\delta} + \sum_{\substack{\delta \mid N \\ \delta \equiv 5 \pmod{24}}} 5r_{\delta} \equiv 0 \pmod{24}.$$
(10)

Additionally, as the weight k is equal to 1/2 times the sum of the r_{δ} ,

$$\sum_{\delta|N} r_{\delta} = 4. \tag{11}$$

However, the largest the left-hand side of (10) can be for non-negative r_{δ} satisfying (11) is 20, and so these have no simultaneous solutions in non-negative integers. Hence, there is no η -quotient in $M_2(\Gamma_1(N))$.

MICHAEL ALLEN

References

- [1] Michael Allen et al. "Eta-quotients of Prime or Semiprime Level and Elliptic Curves". In: *arXiv preprint arXiv:1901.10511* (2019).
- [2] Allison Arnold-Roksandich, Kevin James, and Rodney Keaton. "Counting etaquotients of prime level". In: *Involve* 11.5 (2018), pp. 827–844. ISSN: 1944-4176. DOI: 10.2140/involve.2018.11.827. URL: https://doi.org/10.2140/involve.2018.11.827.
- [3] Anthony J. F. Biagioli. "The construction of modular forms as products of transforms of the Dedekind eta function". In: Acta Arith. 54.4 (1990), pp. 273–300. ISSN: 0065-1036. DOI: 10.4064/aa-54-4-273-300. URL: https://doi.org/10.4064/aa-54-4-273-300.
- [4] Basil Gordon and Kim Hughes. "Multiplicative properties of η-products. II". In: A tribute to Emil Grosswald: number theory and related analysis. Vol. 143. Contemp. Math. Amer. Math. Soc., Providence, RI, 1993, pp. 415–430. DOI: 10.1090/conm/143/01008. URL: https://doi.org/10.1090/conm/143/01008.
- [5] G. Ligozat. Courbes modulaires de genre 1. Publication Mathématique d'Orsay, No. 75 7411. U.E.R. Mathématique, Université Paris XI, Orsay, 1974, p. 102.
- [6] Yves Martin. "Multiplicative η-quotients". In: Trans. Amer. Math. Soc. 348.12 (1996), pp. 4825–4856. ISSN: 0002-9947. DOI: 10.1090/S0002-9947-96-01743-6. URL: https://doi.org/10.1090/S0002-9947-96-01743-6.
- [7] Yves Martin and Ken Ono. "Eta-quotients and elliptic curves". In: Proc. Amer. Math. Soc. 125.11 (1997), pp. 3169–3176. ISSN: 0002-9939. DOI: 10.1090/S0002-9939-97-03928-2
 URL: https://doi.org/10.1090/S0002-9939-97-03928-2.
- [8] Morris Newman. "Construction and application of a class of modular functions". In: Proc. London. Math. Soc. (3) 7 (1957), pp. 334-350. ISSN: 0024-6115. DOI: 10.1112/plms/s3-7.1.334. URL: https://doi.org/10.1112/plms/s3-7.1.334.
- [9] Morris Newman. "Construction and application of a class of modular functions. II". In: Proc. London Math. Soc. (3) 9 (1959), pp. 373-387. ISSN: 0024-6115. DOI: 10.1112/plms/s3-9.3.373. URL: https://doi.org/10.1112/plms/s3-9.3.373.
- [10] Robert J. Lemke Oliver. "Eta-quotients and theta functions". In: Advances in Mathematics 241 (2013), pp. 1–17. ISSN: 0001-8708. DOI: https://doi.org/10.1016/j.aim.2013.03.0 URL: http://www.sciencedirect.com/science/article/pii/S0001870813001102.
- [11] Ken Ono. The web of modularity: arithmetic of the coefficients of modular forms and q-series. Vol. 102. CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2004, pp. viii+216. ISBN: 0-8218-3368-5.
- [12] Jeremy Rouse and John J. Webb. "On spaces of modular forms spanned by etaquotients". In: Adv. Math. 272 (2015), pp. 200-224. ISSN: 0001-8708. DOI: 10.1016/j.aim.2014.12
 URL: https://doi.org/10.1016/j.aim.2014.12.002.