

MEAN-FIELD REFLECTED BACKWARD STOCHASTIC DIFFERENTIAL EQUATIONS

BOUALEM DJEHICHE, ROMUALD ELIE AND SAID HAMADÈNE

ABSTRACT. In this paper, we study a class of reflected backward stochastic differential equations (BSDEs) of mean-field type, where the mean-field interaction in terms of the distribution of the Y -component of the solution enters in both the driver and the lower obstacle. We consider in details the case where the lower obstacle is a deterministic function of $(Y, \mathbb{E}[Y])$ and discuss the more general dependence on the distribution of Y . Under mild Lipschitz and integrability conditions on the coefficients, we obtain the well-posedness of such a class of equations. Under further monotonicity conditions, we show convergence of the standard penalization scheme to the solution of the equation, which hence satisfies a minimality property. This class of equations is motivated by applications in pricing life insurance contracts with surrender options.

1. INTRODUCTION

Backward stochastic differential equations (BSDEs) have been extensively studied in a variety of context since the seminal paper by Pardoux and Peng [PP90]. Much of the interest in BSDEs is due to the induced probabilistic representation of solutions to a large class of semilinear PDEs and stochastic control problems. Hereby, it constitutes a powerful tool for investigating several meaningful applications in engineering, investment science including mathematical finance, game theory and insurance, among many other areas.

Given a square integrable terminal condition ξ and a Lipschitz continuous driver f , a solution to a typical BSDE consists of a pair of adapted processes (Y, Z) defined on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_t, \mathbb{P})$, on which is defined a Brownian motion $(B_t)_t$, which satisfy

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dB_s, \quad 0 \leq t \leq T,$$

where Z is a control process which ensures that Y may be expressed as a recursive utility function

$$Y_t = E \left[\xi + \int_t^T f(s, Y_s, Z_s) ds \mid \mathcal{F}_t \right].$$

In investment science (see e.g. [DE92], [DE92b], [EKPQ97]), this formulation has the plausible interpretation of Y as the yield of an investment under uncertainty, and Z characterizes the optimal investment strategy. In Norberg [Nor91], [Nor92], Y is interpreted as the prospective reserve of a life insurance contract for which the driver f represents the payment process of contractual benefits less premiums payable during the time interval dt and ξ identifies to the terminal payment at the horizon T .

Given some constraint such as a minimum required amount L_t , a.k.a. solvency constraint, at each time t , one would like to find an optimal time D_t after t at which one can

Date: November 15, 2019.

2010 Mathematics Subject Classification. 60H10, 60H07, 49N90.

Key words and phrases. Mean-field, Backward SDEs, Snell envelope, penalization.

Acknowledgements. The first author gratefully acknowledges the financial support provided by the Swedish Research Council grant (2016-04086).

exit the investment so that at any date, the yield Y is always kept larger than the constraint L :

$$(1.1) \quad Y_t \geq L_t, \quad 0 \leq t \leq T.$$

This amounts to express the yield Y as a solution of an optimal stopping problem:

$$Y_t = \operatorname{ess\,sup}_{\tau \text{ stopping time } \geq t} \mathbb{E} \left[\int_t^\tau f(s, Y_s, Z_s) ds + L_\tau 1_{\{\tau < T\}} + \zeta 1_{\{\tau = T\}} \mid F_t \right], \quad t \leq T.$$

Intuitively, the smallest optimal time D_t should be the first instant s after t where Y_s reaches the constraint L_s :

$$D_t = \inf\{s \geq t, Y_s = L_s\} \wedge T.$$

El Karoui *et al.* [EKKP⁺97] were the first to show that such a pair (Y, Z) of adapted processes satisfies

$$(1.2) \quad Y_t = \zeta + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dB_s + K_T - K_t, \quad 0 \leq t \leq T,$$

where the extra term K is an adapted increasing process for which $K_T - K$ is the running cost for keeping Y above L at all times. In connection to optimal stopping problems, this cost must be minimal in the sense that K is only required to push the investment yield Y above L , whenever it may cross it. Namely, whenever $Y_t > L_t$, there is no reason to stop at time t so that $dK_t = 0$. Hereby, the minimal solution of interest to (1.2) is uniquely characterized by the famous Skorohod flatness condition for Snell envelopes

$$(1.3) \quad \int_0^T (Y_t - L_t)^+ dK_t = 0.$$

The dynamics (1.2)-(1.1)-(1.5) is called reflected BSDE whose solution is the adapted process (Y, Z, K) . This class of constrained BSDEs has been widely extended in different directions, in relation to their possible connections to zero-sum games [CK96], investment strategies with portfolio constraints [CKS98], switching problems [HJ07, EK14], robust optimal stopping [MPZ13] and many others. We refer to the recent paper [BEH18] for further discussions and references.

Motivated by considerations related to partial hedging of financial derivatives in mathematical finance, Briand *et al.* [BEH18] built on the weak hedging approach considered in [BER15] and introduce a class of BSDEs where the pathwise running constraint $Y_t \geq L_t$ is replaced by a weaker constraint on the distribution of the yield Y of the form

$$(1.4) \quad \mathbb{E}[\ell(Y_t - L_t)] \geq 0, \quad 0 \leq t \leq T,$$

for a given loss function ℓ . A typical situation is when the process Y is required to remain above a certain benchmark (or solvency level) u with a probability higher than some given level v in which case $\ell : (t, x) \mapsto \mathbb{1}_{\{x \geq u_t\}} - v_t$. For that reason, it is known in the literature as BSDE with mean reflection. Under Lipschitz and integrability conditions on f and ζ , as well as regularity and monotonicity of ℓ , they obtain a unique solution (Y, Z, K) with *deterministic* K to the BSDE with mean reflection with the Skorohod type condition

$$(1.5) \quad \int_0^T \mathbb{E}[\ell(Y_t - L_t)]^+ dK_t = 0.$$

Observe that the constraint (1.4) is much weaker than the one considered in [DESZ17], where the constraint in expectation (1.4) must be satisfied for any stopping time, although such distinction is not relevant for strong reflections of the form (1.1).

In life insurance and pension (see the example in Section 2.1 below), the lower barrier L , usually interpreted as a dynamical solvency level, is typically of the form

$$L_t = u - c(Y_t) + \lambda(\mathbb{E}[Y_t] - u)^+, \quad 0 \leq t \leq T,$$

where u is a required minimum or a benchmark return, $c(Y_t)$ is a reserve dependent management fee and $\lambda(\mathbb{E}[Y_t] - u)^+$ is a bonus option i.e. a fraction λ of the possible surplus realized by the mean value of Y_t . This extra term typically reflects the cooperative aspects induced by the pooling principles of insurance policies.

Motivated by this example, the purpose of this paper is to study of the following fairly general class of reflected BSDEs of mean-field type

$$\begin{cases} Y_t = \xi + \int_t^T f(s, Y_s, \mathbb{P}_{Y_s}, Z_s) ds - \int_t^T Z_s dB_s + K_T - K_t, & 0 \leq t \leq T, \\ Y_t \geq h(Y_t, \mathbb{P}_{Y_t}), \quad \forall t \in [0, T] \text{ and } \int_0^T (Y_t - h(Y_t, \mathbb{P}_{Y_t})) dK_t = 0. \end{cases}$$

where \mathbb{P}_{Y_t} is the marginal probability distribution of the process Y at time t . Mean Field BSDEs have been introduced in [BLP09] and motivated by their connection to control of McKean Vlasov equation or Mean Field games. While the addition of a reflection to these BSDEs has already been considered in [Li14], our point of interest here is to allow the obstacle to depend on the distribution of the Y -component of the solution. Observe also that this class of Mean Field RBSDE shares some possibly fruitful proximity with the notion of averaged obliquely reflected BSDE discussed in [CR17], whenever \mathbb{P}_Y reduces to $\mathbb{E}[Y]$.

In Section 2, we provide a clear formulation of the problem and introduce the required assumptions on the coefficients, while focusing for ease of presentation on the simpler case where the distributional dependence with respect to \mathbb{P}_Y boils down to $\mathbb{E}[Y]$ and the driver does not depend on Z . In Section 3 we state and prove the main results of the paper, namely Theorem 3.1 and 3.2, dedicated to the solvability of mean-field reflected BSDEs. The existence of a unique solution is derived in terms of a fixed point argument for the associated Snell envelope of processes, and the extension to more general dependence with respect to \mathbb{P}_Y is provided in Remark 3.6. In Section 4, we use an alternative approach and show convergence of the classical penalization scheme for BSDEs to the solution of mean-field reflected BSDEs, under further monotone assumptions on the driver f and the solvency constraint h . Such monotone property also ensures that the Skorohod type condition indeed induces the minimality property of the solution. The more involved case where the driver also depends on Z is finally discussed in Section 5.

Notation. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space on which is defined a standard d -dimensional Brownian motion $B = (B_t)_{0 \leq t \leq T}$. We denote by $(\mathcal{F}_t^0 := \sigma\{B_s, s \leq t\})_{0 \leq t \leq T}$ the natural filtration of B and $\mathbb{F} := (\mathcal{F}_t)_{0 \leq t \leq T}$ its completion with the \mathbb{P} -null sets of \mathcal{F} . Let \mathcal{P} be the σ -algebra on $\Omega \times [0, T]$ of \mathcal{F}_t -progressively measurable sets. Next, we introduce the following spaces. For $p \geq 1$, we let

- (i) $L^p = \{\eta : \mathcal{F}_T\text{-measurable random variable such that } \mathbb{E}[|\eta|^p] < \infty\}$;
- (ii) $\mathcal{H}^{p,m} = \{(v_t)_{0 \leq t \leq T} : \mathcal{P}\text{-measurable, } \mathbb{R}^m\text{-valued process such that } \mathbb{E}[(\int_0^T |v_s|^2 ds)^{\frac{p}{2}}] < \infty\} (m \geq 1)$;
- (iii) $\mathcal{S}^p = \{(y_t)_{0 \leq t \leq T} : \mathcal{P}\text{-measurable process such that } \mathbb{E}[\sup_{0 \leq t \leq T} |y_t|^p] < \infty\}$;

- (iv) $\mathcal{S}_c^p = \{(y_t)_{0 \leq t \leq T} : \text{continuous process of } \mathcal{S}^p\}$ which is a complete separable space.
For $0 \leq s \leq t \leq T$, we set $\|y\|_{\mathcal{S}_c^p([s,t])} := (\mathbb{E}[\sup_{s \leq u \leq t} |y_u|^p])^{\frac{1}{p}}$.
- (v) $\mathcal{S}_i^p = \{(k_t)_{0 \leq t \leq T} : \text{continuous non-decreasing process of } \mathcal{S}^p \text{ such that } k_0 = 0\}$;
- (vi) $\mathcal{T}_t = \{v, \mathcal{F}_t\text{-stopping time such that } \mathbb{P}\text{-a.s. } v \geq t\}$;
- (vii) $\mathcal{D} := \{(y_t)_{0 \leq t \leq T} : \mathbb{F}\text{-adapted } \mathbb{R}\text{-valued continuous process such that } \|y\|_1 = \sup_{\tau \in \mathcal{T}_0} \mathbb{E}[|y_\tau|] < \infty\}$.
The normed space $(\mathcal{D}, \|\cdot\|_1)$ is complete (see e.g. [DM82], pp.90). For $0 \leq s \leq t \leq T$, we let $(\mathcal{D}([s,t]), \|\cdot\|_1)$ denote the complete metric space endowed with the norm $\|\cdot\|_1$ restricted to $[s,t]$:

$$\|X\|_1 := \sup_{\tau \in \mathcal{T}_0, s \leq \tau \leq t} \mathbb{E}[|X_\tau|] < \infty.$$

- (viii) For $q \in (0, 1)$, we define

$\mathcal{M}^q = \{(v_t)_{0 \leq t \leq T} : \mathcal{P}\text{-measurable, } \mathbb{R}^m\text{-valued process such that}$

$$\|v\|_{\mathcal{M}^q} := \mathbb{E} \left[\left(\int_0^T |v_s|^2 ds \right)^{\frac{q}{2}} \right] < \infty \}.$$

The space $(\mathcal{M}^q, \|\cdot\|_{\mathcal{M}^q})$ is a complete metric space. The restriction of \mathcal{M}^q to $[s,t]$ is denoted by $\mathcal{M}^q([s,t])$.

2. FORMULATION OF THE PROBLEM

2.1. The class of reflected mean-field BSDEs. In this paper we propose to find solutions (Y, Z, K) to the following class of reflected BSDE of mean-field type associated with the driver f , the terminal condition ξ and the lower barrier h , in the cases $p > 1$ and $p = 1$ respectively, that we make precise in the following

Definition 2.1. *We say that the triple of progressively measurable processes $(Y_t, Z_t, K_t)_{t \leq T}$ is a solution of the mean-field reflected BSDE associated with (f, ξ, h) if,*

- (1) *when $p > 1$,*

$$(2.1) \quad \begin{cases} Y \in \mathcal{S}_c^p, Z \in \mathcal{H}^{p,d} \text{ and } K \in \mathcal{S}_i^p, \\ Y_t = \xi + \int_t^T f(s, Y_s, \mathbb{E}[Y_s]) ds + K_T - K_t - \int_t^T Z_s dB_s, \quad 0 \leq t \leq T, \\ Y_t \geq h(Y_t, \mathbb{E}[Y_t]), \quad \forall t \in [0, T] \text{ and } \int_0^T (Y_t - h(Y_t, \mathbb{E}[Y_t])) dK_t = 0. \end{cases}$$

- (2) *when $p = 1$,*

$$(2.2) \quad \begin{cases} Y \in \mathcal{D}, Z \in \bigcup_{q \in (0,1)} \mathcal{M}^q \text{ and } K \in \mathcal{S}_i^1, \\ Y_t = \xi + \int_t^T f(s, Y_s, \mathbb{E}[Y_s]) ds + K_T - K_t - \int_t^T Z_s dB_s, \quad 0 \leq t \leq T, \\ Y_t \geq h(Y_t, \mathbb{E}[Y_t]), \quad \forall t \in [0, T] \text{ and } \int_0^T (Y_t - h(Y_t, \mathbb{E}[Y_t])) dK_t = 0. \end{cases}$$

In order to alleviate the presentation, we focus in this Section on the particular case where the driver does not depend on Z . We present the general study in Section 5 below as this case needs more involved techniques and more restrictive conditions on the coefficients. Similarly, the main results of the paper, Theorems 3.1 and 3.2 below, which establish the solvability of the systems (2.1) and (2.2) respectively, remain valid if we replace the mean-field coupling $\mathbb{E}[Y_t]$ with the more general marginal law coupling \mathbb{P}_{Y_t} of Y_t , i.e. when the driver and the obstacle are of the form $f(Y_t, \mathbb{P}_{Y_t})$ and $h(Y_t, \mathbb{P}_{Y_t})$ (see Remark 3.6 for further

details). In particular, by taking the barrier h of the form $h(Y_t, \mathbb{P}_{Y_t}) := Y_t - \mathbb{E}[\ell(t, Y_t)]$, observe that we obtain the class of BSDE with mean reflection with loss function ℓ considered in [BEH18] for which the solution Y satisfies $\mathbb{E}[\ell(t, Y_t)] \geq 0$.

Following [EKKP⁺97], the solution Y of the mean-field reflected BSDE in Definition 2.1, if it exists, is the Snell envelope of the process $L := (L_t)_{t \leq T}$ where

$$L_t := \int_0^t f(s, Y_s, \mathbb{E}[Y_s]) ds + h(Y_t, \mathbb{E}[Y_s]_{s=t}) 1_{\{t < T\}} + \xi 1_{\{t=T\}},$$

given by

$$(2.3) \quad Y_t = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[\int_t^\tau f(s, Y_s, \mathbb{E}[Y_s]) ds + h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) 1_{\{\tau < T\}} + \xi 1_{\{\tau=T\}} \mid F_t \right], \quad t \leq T.$$

Driver and obstacles of mean-field type are common in life insurance contracts. A typical example is the case where the benefits less premiums include a cost of capital fee which is proportional to the reserve. Considering payments which involve a mean-field coupling such as $\mathbb{E}[Y]$ comes from a very common practice among actuaries to consider the so-called ‘model points’ method which is some sort of averaging of large homogeneous portfolios when computing reserves and designing policies. We refer to [CDD14] and [DL14] for further details on reserve-dependent policies in life insurance and pensions. Here is an example of such a class of contracts.

• **Guaranteed life endowment with a surrender/withdrawal option.** Consider a portfolio of a large number N of homogeneous life insurance policies ℓ . Denote by $(Y^{\ell, N}, Z^{\ell, N})$ the characteristics of the prospective reserve of each policy $\ell = 1, \dots, N$. We consider non-linear reserving where the driver f depends on the reserve for the particular contract and on the average reserve characteristics over the N contracts (since N is very large, averaging over the remaining $N - 1$ policies has roughly the same effect as averaging over all N policies): For each $\ell = 1, \dots, N$,

$$(2.4) \quad \begin{aligned} f(t, Y_t^{\ell, N}, (Y_t^{m, N})_{m \neq \ell}) &:= \alpha_t - \delta_t Y_t + \beta_t \max(\theta_t, Y_t^{\ell, N} - \frac{1}{N} \sum_{k=1}^N Y_t^{k, N}), \\ h(Y_t^{\ell, N}, (Y_t^{m, N})_{m \neq \ell}) &= u - c(Y_t^{\ell, N}) + \mu (\frac{1}{N} \sum_{k=1}^N Y_t^{k, N} - u)^+, \end{aligned}$$

where $0 < \mu < 1$. The driver includes the discount rate δ_t and deterministic positive functions α_t, β_t and θ_t which constitute the elements of withdrawal option. The solvency level h is constituted of a required minimum or a benchmark return u , a reserve dependent management fee $c(Y_t^{\ell, N})$ (usually much smaller than u) and a bonus option $(\frac{1}{N} \sum_{k=1}^N Y_t^{k, N} - u)^+$ which is the possible surplus realized by the average of all involved contracts, which reflects the cooperative aspect of the pool of insurance contracts.

Sending N to infinity in (2.4), yields the following forms of the driver and the obstacle:

$$\begin{aligned} f(t, Y_t, \mathbb{E}[Y_t]) &:= \alpha_t - \delta_t Y_t + \beta_t \max(\theta_t, Y_t - \mathbb{E}[Y_t]), \\ h(Y_t, \mathbb{E}[Y_t]) &= u - c(Y_t) + \mu (\mathbb{E}[Y_t] - u)^+, \end{aligned}$$

of the prospective reserve of a representative life insurance contract, a.k.a. the model-point among actuaries.

Pricing this type of contracts is one of the main motivations of studying the class (2.1) of MF-reflected BSDEs, while clarifying the connection between such BSDE and Mean Field Games of timing problems [Ber18] is left for further research.

2.2. Assumptions on (f, h, ξ) . We make the following assumption on (f, h, ξ) .

Assumption (H1). The coefficients f, h and ξ satisfy

- (i) f is a mapping from $[0, T] \times \Omega \times \mathbb{R}^2$ into \mathbb{R} such that
 - (a) the process $(f(t, 0, 0))_{t \leq T}$ is \mathcal{P} -measurable and belongs to $\mathcal{H}^{p,1}$;

- (b) f is Lipschitz w.r.t. (y, y') uniformly in (t, ω) , i.e., there exists a positive constant C_f such that \mathbb{P} -a.s. for all $t \in [0, T]$,

$$|f(t, y_1, y'_1) - f(t, y_2, y'_2)| \leq C_f(|y_1 - y_2| + |y'_1 - y'_2|).$$

for any $y_1, y'_1, y_2, y'_2 \in \mathbb{R}$.

- (ii) h is a mapping from \mathbb{R}^2 into \mathbb{R} which is Lipschitz w.r.t. (y, y') , i.e., there exist two positive constants γ_1 and γ_2 such that for any x, y, x' and y'

$$|h(x, x') - h(y, y')| \leq \gamma_1|x - y| + \gamma_2|x' - y'|$$

where γ_1 and γ_2 are two positive constants.

- (iii) ξ is an \mathcal{F}_T -measurable, \mathbb{R} -valued r.v., $\mathbb{E}[\xi^p] < \infty$ and satisfies $\xi \geq h(\xi, \mathbb{E}[\xi])$.

Remark 2.1. *Observe that Assumption (H1) only contains classical Lipschitz and integrability conditions on the coefficients together with a natural condition ensuring that the terminal condition satisfies the constraint of interest.*

Under Assumption (H1), we first derive existence and uniqueness of the solution of (2.1) (for $p > 1$) and (2.2) (for $p = 1$) based on a fixed point argument, using the notion of Snell envelope of processes. Then, under further monotonicity assumptions on (f, h) , we show that the classical penalization scheme converges to that solution together with a minimality property of the solution satisfying the Skorohod type condition.

3. EXISTENCE OF A UNIQUE SOLUTION VIA THE SNELL ENVELOPE METHOD

This Section is dedicated to the construction of a unique solution to the MF-reflected BSDE (2.1) under Assumption (H1), studying successively the cases where $p > 1$ and $p = 1$. Such result will require an additional smallness condition on the Lipschitz regularity of the constraint, see Relation (3.8) on the coefficients γ_1 and γ_2 below together with the discussion in Remark 3.1. In both cases, our argumentation follows from a fixed point property of the Snell Enveloppe representation of the solution on small time intervals, combined with a proper pasting of the solutions on small intervals into a global solution on $[0, T]$. These results are provided below in Theorem 3.1 for $p > 1$ and Theorem 3.2 for $p = 1$, while the more general case where the driver and the constraints depend at time t on the marginal distribution of Y_t is treated in Remark 3.6.

Let Φ be the mapping that associates to a process Y another process $\Phi(Y)$ defined by

$$(3.1) \quad \Phi(Y)_t := \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[\int_t^\tau f(s, Y_s, \mathbb{E}[Y_s]) ds + h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau = T\}} | F_t \right], \quad t \leq T.$$

We will show that the map Φ is well defined and admits a fixed point, first considering the situation where $p > 1$ and then turning to the case where $p = 1$.

3.1. The case $p > 1$.

In this case, we will establish in this Section that the map Φ has a unique fixed point on the complete metric space \mathcal{S}_c^p .

Let first observe in the next Lemma that Φ is a well-defined map from \mathcal{S}_c^p to itself.

Lemma 3.1. *Let f, h and ξ satisfy Assumption (H1) for some $p > 1$. If $Y \in \mathcal{S}_c^p$ then $\Phi(Y) \in \mathcal{S}_c^p$.*

Proof. Set, for $t \leq T$,

$$L_t := \int_0^t f(s, Y_s, \mathbb{E}[Y_s]) ds + h(Y_t, \mathbb{E}[Y_s]_{s=t}) \mathbf{1}_{\{t < T\}} + \xi \mathbf{1}_{\{t=T\}}.$$

By construction, observe that

$$(3.2) \quad \Phi(Y)_t + \int_0^t f(s, Y_s, \mathbb{E}[Y_s]) ds = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E}[L_\tau | \mathcal{F}_t], \quad 0 \leq t \leq T.$$

For $s \leq T$, we linearize the mappings f and h as follows:

$$\begin{aligned} f(s, Y_s, \mathbb{E}[Y_s]) &= f(s, 0, 0) + a_f(s) Y_s + b_f(s) \mathbb{E}[Y_s], \\ h(Y_s, \mathbb{E}[Y_s]) &= h(0, 0) + a_h(s) Y_s + b_h(s) \mathbb{E}[Y_s] \end{aligned}$$

where $a_f(\cdot), b_f(\cdot), a_h(\cdot)$ and $b_h(\cdot)$ are the adapted processes defined, for any $s \leq T$, by

$$(3.3) \quad \begin{cases} a_f(s) := \frac{f(s, Y_s, \mathbb{E}[Y_s]) - f(s, 0, \mathbb{E}[Y_s])}{Y_s} \mathbf{1}_{\{Y_s \neq 0\}}, & a_h(s) := \frac{h(Y_s, \mathbb{E}[Y_s]) - h(0, \mathbb{E}[Y_s])}{Y_s} \mathbf{1}_{\{Y_s \neq 0\}}. \\ b_f(s) := \frac{f(s, 0, \mathbb{E}[Y_s]) - f(s, 0, 0)}{\mathbb{E}[Y_s]} \mathbf{1}_{\{\mathbb{E}[Y_s] \neq 0\}}, & b_h(s) := \frac{h(0, \mathbb{E}[Y_s]) - h(0, 0)}{\mathbb{E}[Y_s]} \mathbf{1}_{\{\mathbb{E}[Y_s] \neq 0\}}. \end{cases}$$

In view of the Lipschitz continuity of f and h provided by Assumption (H1), we have

$$\max(|a_f(\cdot)|, |b_f(\cdot)|) \leq C_f, \quad |a_h(\cdot)| \leq \gamma_1, \quad |b_h(\cdot)| \leq \gamma_2.$$

We have, for any (\mathcal{F}_t) -stopping time ν ,

$$\begin{aligned} L_\nu &= \int_0^\nu (f(s, 0, 0) + a_f(s) Y_s + b_f(s) \mathbb{E}[Y_s]) ds \\ &\quad + (h(0, 0) + a_h(\nu) Y_\nu + b_h(\nu) \mathbb{E}[Y_s]_{s=\nu}) \mathbf{1}_{\{\nu < T\}} + \xi \mathbf{1}_{\{\nu=T\}}. \end{aligned}$$

Moreover, since $Y \in \mathcal{S}_c^p$, the process $(M_t)_{0 \leq t \leq T}$ defined by

$$\begin{aligned} M_t &:= \mathbb{E} \left[\int_0^T |f(s, 0, 0)| + C_f (|Y_s| + \mathbb{E}[|Y_s|]) ds + |h(0, 0)| + \gamma_1 \sup_{s \leq T} |Y_s| \right. \\ &\quad \left. + \gamma_2 \sup_{s \leq T} \mathbb{E}[|Y_s|] + |\xi| |F_t| \right], \end{aligned}$$

is a continuous martingale which, by Doob's inequality, belongs to \mathcal{S}_c^p . We deduce that

$$|\mathbb{E}[L_\tau | F_t]| \leq M_t.$$

for any $t \leq T$ and $\tau \in \mathcal{T}_t$ and, in view of (3.2), obtain

$$|\Phi(Y)_t + \int_0^t f(s, Y_s, \mathbb{E}[Y_s]) ds| \leq M_t.$$

Therefore,

$$\mathbb{E}[\sup_{t \leq T} |\Phi(Y)_t|^p] \leq C_p \left\{ \mathbb{E} \left[\left(\int_0^T |f(s, Y_s, \mathbb{E}[Y_s])| ds \right)^p \right] + \mathbb{E}[\sup_{t \leq T} |M_t|^p] \right\},$$

where C_p is a positive constant that only depends on p and T . Using once more the fact that $Y \in \mathcal{S}_c^p$ and the Lipschitz property of the driver f , it finally holds that $\Phi(Y) \in \mathcal{S}_c^p$. \square

We are now in position to obtain a contraction property of the map Φ on \mathcal{S}_c^p and first derive such property on a small time horizon.

Proposition 3.2. *Let Assumption (H1) hold for some $p > 1$. If γ_1 and γ_2 satisfy*

$$(\gamma_1 + \gamma_2)^{\frac{p-1}{p}} \left\{ \left(\frac{p}{p-1} \right)^p \gamma_1 + \gamma_2 \right\}^{\frac{1}{p}} < 1,$$

then there exists $\delta > 0$ depending only on $p, C_f, \gamma_1, \gamma_2$ and T such that Φ is a contraction on the time interval $[T - \delta, T]$.

Proof. Let $Y, Y' \in \mathcal{S}_c^p$. Then, for any $t \leq T$, we have

$$\begin{aligned} |\Phi(Y)_t - \Phi(Y')_t| &= \left| \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[\int_t^\tau f(s, Y_s, \mathbb{E}[Y_s]) ds + h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau=T\}} \right] \middle| \mathcal{F}_t \right| \\ &\quad - \left| \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[\int_t^\tau f(s, Y'_s, \mathbb{E}[Y'_s]) ds + h(Y'_\tau, \mathbb{E}[Y'_s]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau=T\}} \right] \middle| \mathcal{F}_t \right| \\ &\leq \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[\int_t^\tau |f(s, Y_s, \mathbb{E}[Y_s]) - f(s, Y'_s, \mathbb{E}[Y'_s])| ds \right. \\ &\quad \left. + |h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) - h(Y'_\tau, \mathbb{E}[Y'_s]_{s=\tau})| \middle| \mathcal{F}_t \right]. \end{aligned}$$

Therefore, for any $t \leq T$,

$$(3.4) \quad |\Phi(Y)_t - \Phi(Y')_t| \leq \mathbb{E} \left[\int_t^T |f(s, Y_s, \mathbb{E}[Y_s]) - f(s, Y'_s, \mathbb{E}[Y'_s])| ds \right. \\ \left. + \gamma_1 \sup_{t \leq s \leq T} |Y_s - Y'_s| + \gamma_2 \sup_{t \leq s \leq T} \mathbb{E}[|Y_s - Y'_s|] \middle| \mathcal{F}_t \right].$$

Next, let $\delta > 0$ and $t \in [T - \delta, T]$. From (3.4) and the Lipschitz property of f we obtain

$$|\Phi(Y)_t - \Phi(Y')_t| \leq (\delta C_f + \gamma_1) \mathbb{E} \left[\sup_{T - \delta \leq s \leq T} |Y_s - Y'_s| \middle| \mathcal{F}_t \right] + (\delta C_f + \gamma_2) \sup_{T - \delta \leq s \leq T} \mathbb{E}[|Y_s - Y'_s|]$$

which implies that

$$\begin{aligned} |\Phi(Y)_t - \Phi(Y')_t|^p &\leq (2\delta C_f + \gamma_1 + \gamma_2)^{p-1} \{ (\delta C_f + \gamma_1) (\mathbb{E} \left[\sup_{T - \delta \leq s \leq T} |Y_s - Y'_s| \middle| \mathcal{F}_t \right])^p \\ &\quad + (\delta C_f + \gamma_2) (\sup_{T - \delta \leq s \leq T} \mathbb{E}[|Y_s - Y'_s|])^p \}, \end{aligned}$$

since the convexity relation $(ax_1 + bx_2)^p \leq (a+b)^{p-1}(ax_1^p + bx_2^p)$ holds for any non negative real constants a, b, x_1 and x_2 . Taking the expectation of the supremum over $t \in [T - \delta, T]$ on both sides, and using Doob's inequality (only with one term, the other one is deterministic) together with Jensen's inequality, we obtain

$$\begin{aligned} \mathbb{E} \left[\sup_{T - \delta \leq s \leq T} |\Phi(Y)_s - \Phi(Y')_s|^p \right] &\leq (2\delta C_f + \gamma_1 + \gamma_2)^{p-1} \left\{ \left(\frac{p}{p-1} \right)^p (\delta C_f + \gamma_1) \mathbb{E} \left[\sup_{T - \delta \leq s \leq T} |Y_s - Y'_s|^p \right] + \right. \\ &\quad \left. + (\delta C_f + \gamma_2) \mathbb{E} \left[\sup_{T - \delta \leq s \leq T} |Y_s - Y'_s|^p \right] \right\}. \end{aligned}$$

Hence,

$$(3.5) \quad \|\Phi(Y) - \Phi(Y')\|_{\mathcal{S}_c^p(\delta, T)} \leq \Lambda(C_f, p, \gamma_1, \gamma_2) \times \|Y - Y'\|_{\mathcal{S}_c^p(\delta, T)},$$

where $\mathcal{S}_c^p(\delta, T) := \mathcal{S}_c^p([T - \delta, T])$ is the space \mathcal{S}_c^p restricted to $[T - \delta, T]$ and

$$\Lambda(C_f, p, \gamma_1, \gamma_2)(\delta) := (2\delta C_f + \gamma_1 + \gamma_2)^{\frac{p-1}{p}} \left\{ \left(\frac{p}{p-1} \right)^p (\delta C_f + \gamma_1) + (\delta C_f + \gamma_2) \right\}^{\frac{1}{p}}.$$

Now, it is easy to see that if

$$(\gamma_1 + \gamma_2)^{\frac{p-1}{p}} \left\{ \left(\frac{p}{p-1} \right)^p \gamma_1 + \gamma_2 \right\}^{\frac{1}{p}} < 1,$$

we can find $\delta > 0$ which only depends on C_f, p, γ_1 and γ_2 and more specifically not on the terminal value ξ such that

$$(3.6) \quad \Lambda(C_f, p, \gamma_1, \gamma_2)(\delta) < 1.$$

This directly implies that Φ is a contraction on $\mathcal{S}_c^p([T - \delta, T])$ and hereby has a fixed point Y which satisfies, for any $t \in [T - \delta, T]$,

$$(3.7) \quad Y_t = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[\int_t^\tau f(s, Y_s, \mathbb{E}[Y_s]) ds + h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau=T\}} \right] \middle| \mathcal{F}_t.$$

□

By repeatedly applying the fixed point argument of Proposition (3.2) over adjacent time intervals of fixed length δ and pasting the solutions, we finally obtain existence of a unique solution in $\mathcal{S}_c^p \times \mathcal{H}^{p,d} \times \mathcal{S}_i^p$ to the mean-field reflected BSDE (2.1) over the whole time interval $[0, T]$, as shown in the following theorem.

Theorem 3.1. *Suppose that Assumption (H1) holds for some $p > 1$. If γ_1 and γ_2 satisfy*

$$(3.8) \quad (\gamma_1 + \gamma_2)^{\frac{p-1}{p}} \left\{ \left(\frac{p}{p-1} \right)^p \gamma_1 + \gamma_2 \right\}^{\frac{1}{p}} < 1,$$

then the mean-field reflected BSDE (2.1) has a unique solution $(Y, Z, K) \in \mathcal{S}_c^p \times \mathcal{H}^{p,d} \times \mathcal{S}_i^p$.

Proof. According to the previous argumentation, recall that the constant δ satisfying the inequality (3.6) only depends on p, C_f, γ_1 and γ_2 . Hence, the fixed point argument in Proposition 3.2 can be applied for any terminal condition ξ on a time interval of size δ .

Taking Y^0 the unique fixed point of Φ in $\mathcal{S}_c^p([T - \delta, T])$, observe that Y satisfies (3.7) on $[T - \delta, T]$. In view of the link between the Snell envelope of processes and solutions of reflected BSDEs (see [EKKP⁺97] for more details), there exist processes $(Z^0, K^0) \in \mathcal{H}^{2,d} \times \mathcal{S}_i^2([T - \delta, T])$ such that $K_{T-\delta}^0 = 0$ and (Y^0, Z^0, K^0) solves (2.1) on $[T - \delta, T]$.

Applying the same reasoning on each time interval $[T - (i + 1)\delta, T - i\delta]$ with a similar dynamics but terminal condition $Y_{T-i\delta}^{i-1}$ at time $T - i\delta$, we build recursively for $i = 1$ to any n a solution $(Y^i, Z^i, K^i) \in \mathcal{S}_c^p \times \mathcal{H}^{2,d} \times \mathcal{S}_i^2$ of on each time interval $[T - (i + 1)\delta, T - i\delta]$. Pasting properly these processes, we naturally derive a solution (Y, Z, K) satisfying (2.1) on the full time interval $[0, T]$.

As for the uniqueness of solution to (2.1) on the full time interval $[0, T]$, observe that Y is recursively uniquely defined on each time interval by the fixed point contraction, thanks to the Snell Envelope representation (3.7). Hence, Y and hereby Z are uniquely defined on the full time interval $[0, T]$ while K simply identifies to

$$K_t = Y_0 - Y_t - \int_t^T f(s, Y_s, \mathbb{E}[Y_s]) ds + \int_t^T Z_s dB_s, 0 \leq t \leq T.$$

Hence, (2.1) admits a unique solution in $\mathcal{S}_c^p \times \mathcal{H}^{2,d} \times \mathcal{S}_i^2$ on $[0, T]$. \square

Remark 3.1. *We make the following observations on the sufficient condition (3.8).*

(a) *Since $(\frac{p}{p-1})^p > 1$, the inequality (3.8) implies that $\gamma_1 + \gamma_2 < 1$. The term $(\frac{p}{p-1})^p$ which inflates the Lipschitz constant γ_1 , due to the use of Doob's inequality, makes the condition (3.8) a bit heavy. We will see in Theorem 3.2 (for the case $p = 1$) that the solvability of the MF-BSDE (2.2) only requires $\gamma_1 + \gamma_2 < 1$, as Doob's inequality is not required for the proof.*

(b) *Noting that (3.8) also reads*

$$(3.9) \quad \left(\frac{p}{p-1} \right)^p \gamma_1 + \gamma_2 < \frac{1}{(\gamma_1 + \gamma_2)^{p-1}},$$

using the fact that $\lim_{p \rightarrow \infty} (\frac{p}{p-1})^p = e$, if $\gamma_1 + \gamma_2 < 1$ there exists p for which (3.8) is satisfied, since, when $p \rightarrow \infty$, the left-hand side converges to $e\gamma_1 + \gamma_2$ while the right-hand side diverges to $+\infty$.

On the other hand, since $\lim_{p \rightarrow 1} (\frac{p}{p-1})^p = +\infty$, then (3.8) is satisfied only if γ_1 very small which means that h varies very slowly w.r.t the component y .

(c) *When the barrier h does not depend on the mean-field term $\mathbb{E}[Y]$ (i.e. $\gamma_2 = 0$), the fixed point argument through the Snell envelope may not be appropriate to obtain the solvability of the BSDE unless $\gamma_1 < \frac{p-1}{p}$. For example, if $\frac{1}{2} < \gamma_1 < 1$, the BSDE with a standard driver f i.e. independent of $\mathbb{E}[Y]$ and barrier $h(y) = \gamma_1 y$ does not satisfy Theorem 3.1 for $p = 2$. But if $\xi \geq 0$ and $f(t, \omega, y) \geq 0$, then any solution Y of equation (3.7) is non-negative. Then the condition*

$Y_t \geq \gamma_1 Y_t$ is equivalent to $Y_t \geq 0$, and by Corollary 3.7 in [EKKP⁺97], the corresponding BSDE has a unique solution.

(d) Whenever the solvency constraint only depends on $\mathbb{E}[Y_t]$, i.e. when $\gamma_1 = 0$, observe that (3.8) simply reduces to the condition $\gamma_2 < 1$. This is quite natural as for example a linear constraint $Y_t \geq \gamma_2 \mathbb{E}[Y_t]$ would be automatically violated as soon as $\gamma_2 > 1$.

(e) In the particular case of linear constraint of the form $Y_t \geq \gamma_1 Y_t + h(\mathbb{E}[Y_t])$ with $\gamma_1 \geq 0$, the condition (3.8) simply rewrites $\gamma_1 + \gamma_2 < 1$ as the condition is simply equivalent to $Y_t \geq \frac{h(\mathbb{E}[Y_t])}{1-\gamma_1}$.

3.2. The case $p = 1$.

In this case we establish that the map Φ has a unique fixed point on the complete metric space \mathcal{D} .

Following the same line of proof as in Lemma 3.1, we can verify that Φ is a well-defined map from \mathcal{D} to itself.

Lemma 3.3. *Let f, h and ξ satisfy Assumption (H1) for $p = 1$. If $Y \in \mathcal{D}$ then $\Phi(Y) \in \mathcal{D}$.*

In the next proposition we show that Φ admits a local fixed point.

Proposition 3.4. *Let f, h and ξ satisfy Assumption (H1) for $p = 1$ and suppose that*

$$\gamma_1 + \gamma_2 < 1.$$

Then there exists $\delta > 0$, depending only on C_f, γ_1 and γ_2 and a process Y which belongs to $\mathcal{D}([T - \delta, T])$, such that for any $t \in [T - \delta, T]$,

$$\Phi(Y)_t = Y_t.$$

Proof. We are going to show existence of such a δ such that Φ is a contraction on $(\mathcal{D}([T - \delta, T]), \|\cdot\|_1)$. Indeed, let Y and Y' be two processes of \mathcal{D} and let δ be a positive constant to be determined later on. For any stopping time θ such that \mathbb{P} -a.s. $T - \delta \leq \theta \leq T$, we have

$$\begin{aligned} |\Phi(Y)_\theta - \Phi(Y')_\theta| &= \left| \operatorname{ess\,sup}_{\tau \in \mathcal{T}_\theta} \mathbb{E} \left[\int_\theta^\tau f(s, Y_s, \mathbb{E}[Y_s]) ds + h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau = T\}} \mid \mathcal{F}_\theta \right] \right. \\ &\quad \left. - \operatorname{ess\,sup}_{\tau \in \mathcal{T}_\theta} \mathbb{E} \left[\int_\theta^\tau f(s, Y'_s, \mathbb{E}[Y'_s]) ds + h(Y'_\tau, \mathbb{E}[Y'_s]_{s=\tau}) \mathbf{1}_{\{\tau < T\}} + \xi \mathbf{1}_{\{\tau = T\}} \mid \mathcal{F}_\theta \right] \right| \\ &\leq \operatorname{ess\,sup}_{\tau \in \mathcal{T}_\theta} \mathbb{E} \left[\int_\theta^\tau |f(s, Y_s, \mathbb{E}[Y_s]) - f(s, Y'_s, \mathbb{E}[Y'_s])| ds + |h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) - h(Y'_\tau, \mathbb{E}[Y'_s]_{s=\tau})| \mid \mathcal{F}_\theta \right]. \end{aligned}$$

In view of the Lipschitz continuity of f and h , we have

$$\begin{aligned} &\mathbb{E}[|\Phi(Y)_\theta - \Phi(Y')_\theta|] \\ &\leq \sup_{\tau \in \mathcal{T}_\theta} \mathbb{E} \left[\int_\theta^\tau |f(s, Y_s, \mathbb{E}[Y_s]) - f(s, Y'_s, \mathbb{E}[Y'_s])| ds + |h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) - h(Y'_\tau, \mathbb{E}[Y'_s]_{s=\tau})| \right] \\ &\leq \sup_{\tau \in \mathcal{T}_\theta} \mathbb{E} \left[\int_{T-\delta}^T |f(s, Y_s, \mathbb{E}[Y_s]) - f(s, Y'_s, \mathbb{E}[Y'_s])| ds + |h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) - h(Y'_\tau, \mathbb{E}[Y'_s]_{s=\tau})| \right] \\ &= (2\delta C_f + \gamma_1 + \gamma_2) \sup_{T-\delta \leq \tau \leq T} \mathbb{E}[|Y_\tau - Y'_\tau|]. \end{aligned}$$

Now, since θ is an arbitrary stopping time in $[T - \delta, T]$, it holds that

$$\sup_{T-\delta \leq \tau \leq T} \mathbb{E}[|\Phi(Y)_\tau - \Phi(Y')_\tau|] \leq (2\delta C_f + \gamma_1 + \gamma_2) \sup_{T-\delta \leq \tau \leq T} \mathbb{E}[|Y_\tau - Y'_\tau|].$$

Now, it is immediate to see that, whenever $\gamma_1 + \gamma_2 < 1$, there exists $\delta > 0$ only depending on C_f, γ_1 and γ_2 such that $2\delta C_f + \gamma_1 + \gamma_2 < 1$. This yields that the map Φ is a contraction on the complete metric space $(\mathcal{D}[T - \delta, T], \|\cdot\|_1)$. Hence, Φ has a unique fixed point, i.e., there exists $Y \in \mathcal{D}[T - \delta, T]$, such that for any $t \in [T - \delta, T]$, $\Phi(Y)_t = Y_t$. \square

In the next proposition we show that the fixed point Y of Φ on $(\mathcal{D}([T - \delta, T]), \|\cdot\|_1)$ yields the existence of a unique local solution of (2.1) in $(\mathcal{D}([T - \delta, T]))$.

Proposition 3.5. *Let f, h and ξ satisfy Assumption (H1) for $p = 1$ and suppose that*

$$(3.10) \quad \gamma_1 + \gamma_2 < 1.$$

Then, there exist $\delta > 0$ only depending on C_f, γ_1 and γ_2 , and a triplet of processes (Y, Z, K) such that

$$(3.11) \quad \begin{cases} Y \in \mathcal{D}([T - \delta, T]), Z \in \bigcup_{q \in (0,1)} \mathcal{M}^q([T - \delta, T]), \\ K \text{ continuous nondecreasing such that } K_{T-\delta} = 0, \mathbb{E}[K_T] < \infty, \\ Y_t = \xi + \int_t^T f(s, Y_s, \mathbb{E}[Y_s]) ds + K_T - K_t - \int_t^T Z_s dB_s, \quad T - \delta \leq t \leq T, \\ Y_t \geq h(Y_t, \mathbb{E}[Y_t]), \quad \forall t \in [T - \delta, T] \text{ and } \int_{T-\delta}^T (Y_t - h(Y_t, \mathbb{E}[Y_t])) dK_t = 0. \end{cases}$$

Moreover, if $(\bar{Y}, \bar{Z}, \bar{K})$ is another triple which satisfies (3.11), then for $t \in [T - \delta, T]$, $Y_t = \bar{Y}_t$, $K_t = \bar{K}_t$ and $Z_t 1_{\{T-\delta \leq t \leq T\}} = \bar{Z}_t 1_{\{T-\delta \leq t \leq T\}}$, $dt \otimes d\mathbb{P}$ -a.e.

Proof. Let δ be the positive constant and Y the process defined in Proposition 3.4 above. Then, for all $t \in [T - \delta, T]$,

$$(3.12) \quad Y_t = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[\int_t^\tau f(s, Y_s, \mathbb{E}[Y_s]) ds + h(Y_\tau, \mathbb{E}[Y_s]_{s=\tau}) 1_{\{\tau < T\}} + \xi 1_{\{\tau=T\}} \middle| \mathcal{F}_t \right].$$

Therefore, $(Y_t + \int_{T-\delta}^t f(s, Y_s, \mathbb{E}[Y_s]) ds)_{t \in [T-\delta, T]}$ belongs to \mathcal{D} since $Y \in \mathcal{D}$ and it is the Snell envelope of the process $(\int_{T-\delta}^t f(s, Y_s, \mathbb{E}[Y_s]) ds + h(Y_t, \mathbb{E}[Y_t]) 1_{\{t < T\}} + \xi 1_{\{t=T\}})_{t \in [T-\delta, T]}$. Then, by applying the Doob-Meyer decomposition ([DM82], pp. 211), there exists a continuous martingale $(M_t)_{t \in [T-\delta, T]}$ and a non-decreasing process $(K_t)_{t \in [T-\delta, T]}$ ($K_{T-\delta} = 0$) such that

$$\forall t \in [T - \delta, T], \quad Y_t + \int_{T-\delta}^t f(s, Y_s, \mathbb{E}[Y_s]) ds = M_t - K_t.$$

Moreover, $\mathbb{E}[K_T] + \mathbb{E}[|M_T|] < \infty$. Next, for $t \leq T - \delta$, we set $M_t := Y_t = \mathbb{E}[M_{T-\delta} | \mathcal{F}_t]$ and $K_t = 0$. Then, by the Martingale Representation Theorem there exists a \mathcal{P} -measurable process $(Z_t)_{t \in [T-\delta, T]}$ valued in \mathbb{R}^d , such that for any $t \in [T - \delta, T]$,

$$M_t = M_{T-\delta} + \int_{T-\delta}^t Z_s dB_s.$$

Thus, for any $t \in [T - \delta, T]$,

$$Y_t = \xi + \int_t^T f(s, Y_s, \mathbb{E}[Y_s]) ds + K_T - K_t - \int_t^T Z_s dB_s \text{ and } Y_t \geq h(Y_t, \mathbb{E}[Y_t]).$$

On the other hand, since $\mathbb{E}[|M_T|] < \infty$, we observe that $\mathbb{E}[\sup_{t \in [T-\delta, T]} |M_t|^q] < \infty$, for any $q \in (0, 1)$ (see [BDH⁺03], pp.125). Therefore, the Burkholder-Davis-Gundy inequality ([RY13], pp. 161) implies that $Z \in \mathcal{M}^q([T - \delta, T])$ for any $q \in (0, 1)$ and we deduce that $Z \in \bigcup_{q \in (0,1)} \mathcal{M}^q([T - \delta, T])$. Finally, for $t \in [T - \delta, T]$, the random time defined by

$$D_t := \inf\{s \in [T - \delta, T], Y_s = h(Y_s, \mathbb{E}[Y_s])\} \wedge T,$$

is a stopping time which realizes the essential supremum in (3.12), i.e., it is optimal. Thus, by the optimal stopping properties we have that $\int_t^{D_t} (Y_s - h(Y_s, \mathbb{E}[Y_s])) dK_s = 0$ for any $t \in [T - \delta, T]$. Consequently we have $\int_{T-\delta}^T (Y_s - h(Y_s, \mathbb{E}[Y_s])) dK_s = 0$ (see[EKKP⁺97], pp. 717).

We now prove uniqueness. Let $(\bar{Y}, \bar{Z}, \bar{K})$ be another triple that satisfies (3.11). Therefore, \bar{Y} satisfies, for all $t \in [T - \delta, T]$,

$$(3.13) \quad \bar{Y}_t = \operatorname{ess\,sup}_{\tau \in \mathcal{T}_t} \mathbb{E} \left[\int_t^\tau f(s, \bar{Y}_s, \mathbb{E}[\bar{Y}_s]) ds + h(\bar{Y}_\tau, \mathbb{E}[\bar{Y}_s]_{s=\tau}) 1_{\{\tau < T\}} + \xi 1_{\{\tau=T\}} \middle| \mathcal{F}_t \right].$$

As \bar{Y} belongs $\mathcal{D}([T - \delta, T])$, then it is the fixed point of the mapping Φ on $[T - \delta, T]$, thus $\bar{Y}_t = Y_t$ for any $t \in [T - \delta, T]$. Now, the equations satisfied by Y and \bar{Y} imply that for any $t \in [T - \delta, T]$ $K_t = \bar{K}_t$ and $Z_t \mathbb{1}_{\{T - \delta \leq t \leq T\}} = \bar{Z}_t \mathbb{1}_{\{T - \delta \leq t \leq T\}}$, as claimed. \square

Arguing now as in Theorem 3.1, we are able to paste solutions on small intervals and derive a global solution on any time interval $[0, T]$, which is the main result of this section, together with Remark 3.6 below:

Theorem 3.2. *Let f, h and ξ satisfy Assumption (H1) for $p = 1$ and suppose that $\gamma_1 + \gamma_2 < 1$. Then, there is a unique triple of processes (Y, Z, K) such that*

$$(3.14) \quad \begin{cases} Y \in \mathcal{D}, Z \in \bigcup_{q \in (0,1)} \mathcal{M}^q, K \in \mathcal{S}_t^1, \\ Y_t = \xi + \int_t^T f(s, Y_s, \mathbb{E}[Y_s]) ds + K_T - K_t - \int_t^T Z_s dB_s, \quad \forall t \in [0, T], \\ Y_t \geq h(Y_t, \mathbb{E}[Y_t]), \quad \forall t \in [0, T] \text{ and } \int_0^T (Y_t - h(Y_t, \mathbb{E}[Y_t])) dK_t = 0. \end{cases}$$

Remark 3.2. *Recall that the process Y which satisfies (3.14) also admits the representation (2.3).*

Remark 3.6. *Theorems 3.1 and 3.2 are still valid if we replace the mean-field coupling $\mathbb{E}[Y_t]$ with the marginal law \mathbb{P}_{Y_t} of Y at time t in both the driver f and the obstacle h , i.e. consider a MF-BSDE driven by $f(Y_t, \mathbb{P}_{Y_t})$ reflected on $h(Y_t, \mathbb{P}_{Y_t})$ and with terminal condition ξ , provided that f and h are Lipschitz continuous w.r.t. the Wasserstein distance i.e. Assumptions (i) (b) and (ii) are replaced with the following. For any $y_1, y_2 \in \mathbb{R}$ and $\mu, \nu \in \mathcal{P}_p(\mathbb{R}^d)$,*

$$|f(t, y_1, \mu) - f(t, y_2, \nu)| \leq C_f(|y_1 - y_2| + d_p(\mu, \nu)),$$

$$|h(y_1, \mu) - h(y_2, \nu)| \leq \gamma_1|x - y| + \gamma_2 d_p(\mu, \nu),$$

where $d_p(\cdot, \cdot)$ is the p -Wasserstein distance between two probability measures on the subset $\mathcal{P}_p(\mathbb{R}^d)$ of probability measures with finite p -th moment, formulated in terms of a coupling between two random variables X and Y defined on the same probability space:

$$d_p(\mu, \nu) := \inf \left\{ (\mathbb{E}[|X - Y|^p])^{1/p}, \text{law}(X) = \mu, \text{law}(Y) = \nu \right\}.$$

In particular, we have, for any $0 \leq s \leq u \leq t \leq T$,

$$(3.15) \quad \sup_{u \in [s, t]} d_p(\mathbb{P}_{Y_u}, \mathbb{P}_{Y'_u}) \leq \sup_{u \in [s, t]} (\mathbb{E}[|Y_u - Y'_u|^p])^{1/p}$$

from which we derive the following useful inequality

$$(3.16) \quad \sup_{u \in [s, t]} d_p(\mathbb{P}_{Y_u}, \delta_0) \leq \sup_{u \in [s, t]} (\mathbb{E}[|Y_u|^p])^{1/p}.$$

Moreover, the linearization (3.3) of the mappings f and h used in the proof above is still valid. Indeed, for $s \leq T$, we have

$$\begin{aligned} f(s, Y_s, \mathbb{P}_{Y_s}) &= f(s, 0, \delta_0) + a_f(s)Y_s + b_f(s)d_p(\mathbb{P}_{Y_s}, \delta_0) \\ h(Y_s, \mathbb{P}_{Y_s}) &= h(0, \delta_0) + a_h(s)Y_s + b_h(s)d_p(\mathbb{P}_{Y_s}, \delta_0), \end{aligned}$$

where, in view of the Lipschitz continuity of f and h w.r.t. (y, μ) , the coefficients $b_f(\cdot)$ and $b_h(\cdot)$ become, for any $s \leq T$,

$$b_f(s) := \frac{f(s, 0, \mathbb{P}_{Y_s}) - f(s, 0, \delta_0)}{d_p(\mathbb{P}_{Y_s}, \delta_0)} \mathbb{1}_{\{d_p(\mathbb{P}_{Y_s}, \delta_0) \neq 0\}}, \quad b_h(s) := \frac{h(0, \mathbb{P}_{Y_s}) - h(0, \delta_0)}{d_p(\mathbb{P}_{Y_s}, \delta_0)} \mathbb{1}_{\{d_p(\mathbb{P}_{Y_s}, \delta_0) \neq 0\}}.$$

and satisfy together with $a_f(\cdot)$ and $a_h(\cdot)$

$$\max(|a_f(\cdot)|, |b_f(\cdot)|) \leq C_f, \quad |a_h(\cdot)| \leq \gamma_1, \quad |b_h(\cdot)| \leq \gamma_2.$$

Thanks to this linearization and the inequality (3.16), Propositions 3.1 and 3.3 extend to the general case under consideration. Moreover, the inequality (3.15) makes the above fixed point argument used in Theorems 3.1 and 3.2 extend to this general case as well.

4. CONVERGENCE OF THE PENALIZATION SCHEME

We now present an alternative approach for the derivation of a unique solution to (2.1), using a penalization type constructive argumentation, which may also reinterpret as a recursive scheme reflecting at step n the solution to the BSDE Y^n on the obstacle generated by the solution Y^{n-1} . The monotonic convergence of the penalized BSDEs naturally relies on a comparison argumentation, which requires as for non-reflected MF BSDE [BLP09] additional monotone properties on the driver and the constraint, see Assumption (H2) right below. Under such assumption, we verify that the Skorohod type condition indeed characterizes the minimality property of the solution to the constrained BSDE. The result is besides derived in Theorem 4.3 under an additional monotony property of the driver, which is in fact not necessary as explained in Corrolary 4.1. For the sake of simplicity, we choose to restrict to the case $p = 2$.

The following assumptions on the mappings f , h and terminal value ξ will be in force throughout this Section.

Assumption (H2) f , h and ξ satisfy

- (a) Assumption (H1) for $p = 2$.
- (b) For any fixed y (resp. y'), the mapping $y' \in \mathbb{R} \mapsto h(y, y')$ (resp. $y \in \mathbb{R} \mapsto h(y, y')$) is non-decreasing.
- (c) the mapping $y' \in \mathbb{R} \mapsto f(t, y, y')$ is non-decreasing, for t, y fixed.

Remark 4.3. (i) The monotonicity conditions imposed on f and h in (H2)(b,c) are needed to be able to apply the comparison theorem for solutions of mean-field BSDEs (cf. [BLP09]).

(ii) Since h is Lipschitz, by (H2)-b), linearizing h , we obtain, for all y and y' ,

$$(4.1) \quad (h(y, y'))^+ \leq h(0, 0)^+ + \gamma_1 y^+ + \gamma_2 (y')^+.$$

Let us now introduce the following penalization scheme of the equation (2.1). For $n \geq 0$, let (Y^n, Z^n) be the pair of processes defined as follows: $Y^0 = \underline{Y}$ where $(\underline{Y}, \underline{Z})$ is the solution of the following BSDE of mean field type (which exists according to Theorem 3.1. in [BLP09]):

$$(4.2) \quad \begin{cases} \underline{Y} \in \mathcal{S}_c^2, & \underline{Z} \in \mathcal{H}^{2,d}; \\ \underline{Y}_t = \xi + \int_t^T f(s, \underline{Y}_s, \mathbb{E}[\underline{Y}_s]) ds - \int_t^T \underline{Z}_s dB_s, & t \leq T. \end{cases}$$

Next, for $n \geq 1$, we define (Y^n, Z^n) as the solution of the following standard BSDE:

$$(4.3) \quad \begin{cases} Y^n \in \mathcal{S}_c^2, Z^n \in \mathcal{H}^{2,d}; \\ Y_t^n = \xi + \int_t^T f(s, Y_s^n, \mathbb{E}[Y_s^{n-1}]) ds + K_T^n - K_t^n - \int_t^T Z_s^n dB_s, & t \leq T, \end{cases}$$

where, for $n \geq 1$, K^n denotes the process

$$K_t^n := n \int_0^t (Y_s^n - h(Y_s^{n-1}, \mathbb{E}[Y_s^{n-1}]))^- ds, \quad t \leq T.$$

Theorem 4.3. Suppose that (γ_1, γ_2) satisfies (3.8) with $p = 2$ and that Assumption (H2) is fulfilled. Suppose also that $y \in \mathbb{R} \mapsto f(t, y, y')$ is non-decreasing, for any t, y' fixed.

Then, the sequence $(Y^n, Z^n, K^n)_{n \geq 0}$ converges in $\mathcal{S}_c^2 \times \mathcal{H}^{2,d} \times \mathcal{S}^2$ to the unique solution (Y, Z, K) of (2.1) i.e.

$$(4.4) \quad \begin{cases} Y \in \mathcal{S}_c^2, Z \in \mathcal{H}^{2,d} \text{ and } K \in \mathcal{S}_{ci}^2; \\ Y_t = \xi + \int_t^T f(s, Y_s, \mathbb{E}[Y_s]) ds + K_T - K_t - \int_t^T Z_s dB_s, \quad t \leq T; \\ Y_t \geq h(Y_t, \mathbb{E}[Y_t]), \quad \forall t \geq T \text{ and } \int_0^T (Y_t - h(Y_t, \mathbb{E}[Y_t])) dK_t = 0. \end{cases}$$

Besides, the Skorohod condition ensures the minimality property of the solution.

Remark 4.4. The additional assumption on the monotone property of $f(t, \cdot, y')$ for any (t, y') is in fact not necessary for such result, as detailed in Corrolary 4.1 below

Proof. : Observe first, that, since the solution $(\underline{Y}, \underline{Z})$ of (4.2) exists, then, by induction, it follows that, for any $n \geq 1$, there is a unique solution (Y^n, Z^n) of (4.3). We proceed to the proof of the convergence in the following five steps. We first verify that the sequence $(Y^n)_n$ is monotone and bounded, leading to a limit \hat{Y} which is proved to satisfy the constraint in the second step. Next, we derive the continuity of \hat{Y} and finally conclude on the dynamics of the limit of (Y^n, Z^n, K^n) as well as on the minimality property.

Step 1. We have, for any $n \geq 0$, $Y^n \leq Y^{n+1} \leq Y$.

We will show by induction that, for any $n \geq 0$, $Y^0 \leq Y^1 \leq Y^n \leq Y^{n+1} \leq Y$. To begin with, recall that (Y, Z, K) satisfies

$$(4.5) \quad \begin{cases} Y_t = \xi + \int_t^T f(s, Y_s, \mathbb{E}[Y_s]) ds + K_T - K_t - \int_t^T Z_s dB_s, \quad t \leq T; \\ Y_t \geq h(Y_t, \mathbb{E}[Y_t]), \quad \forall t \geq T \text{ and } \int_0^T (Y_t - h(Y_t, \mathbb{E}[Y_t])) dK_t = 0. \end{cases}$$

Now, consider the following standard reflected BSDE satisfied by $(\bar{Y}, \bar{Z}, \bar{K})$:

$$(4.6) \quad \begin{cases} \bar{Y}_t = \xi + \int_t^T f(s, \bar{Y}_s, \mathbb{E}[\bar{Y}_s]) ds + \bar{K}_T - \bar{K}_t - \int_t^T \bar{Z}_s dB_s, \quad t \leq T; \\ \bar{Y}_t \geq h(\bar{Y}_t, \mathbb{E}[\bar{Y}_t]), \quad \forall t \geq T \text{ and } \int_0^T (\bar{Y}_t - h(\bar{Y}_t, \mathbb{E}[\bar{Y}_t])) d\bar{K}_t = 0. \end{cases}$$

Since the solution of (4.6) is unique, then we obviously have $\bar{Y} = Y$.

The fact that $Y^0 \leq Y^1$ follows from the standard comparison theorem for solutions of BSDEs (see e.g. [EKPQ97]) since the penalization term is non negative. Next, if for some $n \geq 1$ we have $Y^{n-1} \leq Y^n$, then using once more the comparison theorem for standard BSDEs we get that $Y^n \leq Y^{n+1}$ since the mappings $y' \mapsto f(y, y')$, $y \mapsto h(y, y')$ and $y' \mapsto h(y, y')$ are non-decreasing. Thus, for any $n \geq 0$, we have $Y^n \leq Y^{n+1}$.

Since the mapping $y' \mapsto f(y, y')$ is non-decreasing, by the comparison theorem for solutions of mean-field BSDEs in [BLP09] (see Theorem 6.5 in Appendix), we have also $Y^0 = \underline{Y} \leq Y = \bar{Y}$.

The mappings $y' \mapsto f(y, y')$, $y \mapsto h(y, y')$ and $y' \mapsto h(y, y')$ being nondecreasing, we have

$$\{f(s, y, \mathbb{E}[Y_s^0]) + n(y - h(Y_s^0, \mathbb{E}[Y_s^0]))^-\}_{|y=\bar{Y}_s} = f(s, y, \mathbb{E}[Y_s^0])_{|y=\bar{Y}_s} \leq f(s, y, \mathbb{E}[Y_s])_{|y=\bar{Y}_s}.$$

In view of the comparison theorem for solutions of standard BSDEs we obtain that $Y^1 \leq \bar{Y}$. But, \bar{Y} is nothing else but Y . Therefore, $Y^1 \leq Y$. Next, if for some $n \geq 1$, we have $Y^n \leq \bar{Y}$, we use the same argument to show that $Y^{n+1} \leq \bar{Y} = Y$. Therefore, for any $n \geq 0$, $Y^n \leq Y$.

Summing up, for any $n \geq 0$, $Y^n \leq Y^{n+1} \leq Y$.

Step 2. The limit \hat{Y} satisfies the solvency constraint.

Set, for $t \in [0, T]$,

$$\hat{Y}_t := \lim_{n \rightarrow \infty} Y_t^n.$$

Then the process \hat{Y} is rcll and satisfies

$$(4.7) \quad \mathbb{E}[\sup_{t \leq T} |\hat{Y}_t|^2] < \infty \quad \text{and} \quad \forall t \leq T, \quad \hat{Y}_t \geq h(\hat{Y}_t, \mathbb{E}[\hat{Y}_t]).$$

Indeed, the existence of the process \hat{Y} is due to the fact that the sequence $(Y^n)_{n \geq 0}$ is non decreasing. Moreover, since $Y^0 \leq \hat{Y} \leq Y$, the square integrability of \hat{Y} is due to the square-integrability of Y^0 and Y .

Next, for $n \geq 1$, let us set $\bar{Y}^n = Y^n - Y^0$. Therefore, \bar{Y}^n satisfies, for all $t \leq T$,

$$\begin{aligned} \bar{Y}_t^n &= \int_t^T \{f(s, Y_s^n, \mathbb{E}[Y_s^{n-1}]) - f(s, \underline{Y}_s, \mathbb{E}[\underline{Y}_s])\} ds \\ &\quad + n \int_t^T (Y_s^n - h(Y_s^{n-1}, \mathbb{E}[Y_s^{n-1}]))^- ds - \int_t^T (Z_s^n - \underline{Z}_s) dB_s. \end{aligned}$$

Since f is nondecreasing w.r.t y and y' , and we have $Y^n \geq \underline{Y}$, then for any $s \leq T$, $f(s, Y_s^n, \mathbb{E}[Y_s^{n-1}]) - f(s, \underline{Y}_s, \mathbb{E}[\underline{Y}_s]) \geq 0$. This in turn implies that for any $n \geq 0$, \bar{Y}^n is a continuous supermartingale which converges increasingly to $\hat{Y} - Y^0$, therefore $\hat{Y} - Y^0$ is rcll ([DM82], pp. 86). Consequently, \hat{Y} is rcll, since Y^0 is continuous. Next, by (4.3), we have

$$\mathbb{E}[\int_t^T (Y_s^n - h(Y_s^{n-1}, \mathbb{E}[Y_s^{n-1}]))^- ds] = \frac{1}{n} \mathbb{E}[Y_0^n - \zeta - \int_t^T f(s, Y_s^n, \mathbb{E}[Y_s^{n-1}]) ds].$$

By Fatou's Lemma, we have

$$\mathbb{E}[\int_t^T (\hat{Y}_s - h(\hat{Y}_s, \mathbb{E}[\hat{Y}_s]))^- ds] \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[Y_0^n - \zeta - \int_t^T f(s, Y_s^n, \mathbb{E}[Y_s^{n-1}]) ds] = 0$$

since $Y^0 \leq Y^n \leq Y$ and the processes Y^0, Y belong to \mathcal{S}_c^2 . As \hat{Y} and $h(\hat{Y}, \mathbb{E}[\hat{Y}])$ are rcll, then for any $t < T$, $\hat{Y}_t \geq h(\hat{Y}_t, \mathbb{E}[\hat{Y}_t])$. Finally, the inequality holds also on T since $\hat{Y}_T = \zeta$ and $\zeta \geq h(\zeta, \mathbb{E}[\zeta])$ by (H1)-iii).

Step 3. \hat{Y} is continuous.

First note that by the uniform square integrability of \hat{Y} , the process $(h(\hat{Y}_t, \mathbb{E}[\hat{Y}_t]))_{t \leq T}$ is uniformly square integrable since h is Lipschitz w.r.t (y, y') . On the other hand the process Y^n has the following representation as a Snell Envelope of processes: for all $t \in [0, T]$,

$$Y_t^n = \operatorname{ess\,sup}_{\tau \geq t} \mathbb{E}[\int_t^\tau f(s, Y_s^n, \mathbb{E}[Y_s^{n-1}]) ds + Y_\tau^n \wedge h(Y_\tau^{n-1}, \mathbb{E}[Y_\tau^{n-1}]) \mathbf{1}_{\{\tau < T\}} + \zeta \mathbf{1}_{\{\tau = T\}} | \mathcal{F}_t],$$

since

$$Y^n \geq Y^n \wedge h(Y^{n-1}, \mathbb{E}[Y^{n-1}]) \quad \text{and} \quad \int_0^T (Y_s^n - Y_s^n \wedge h(Y_s^{n-1}, \mathbb{E}[Y_s^{n-1}])) dK_s^n = 0.$$

One can see the paper by El-Karoui et al. [EKKP⁺97] for more details. Now, if for any $n \geq 0$, U^n and U are rcll and $U^n \nearrow U$ then $\operatorname{SN}(U^n) \nearrow \operatorname{SN}(U)$ (see e.g. [DHP09] for more details). It follows that, for any $t \leq T$,

$$(4.8) \quad \hat{Y}_t = \operatorname{ess\,sup}_{\tau \geq t} \mathbb{E}[\int_t^\tau f(s, \hat{Y}_s, \mathbb{E}[\hat{Y}_s]) ds + h(\hat{Y}_\tau, \mathbb{E}[\hat{Y}_\tau]) \mathbf{1}_{\{\tau < T\}} + \zeta \mathbf{1}_{\{\tau = T\}} | \mathcal{F}_t]$$

since $Y^n \nearrow \hat{Y}$, f and h are increasing in their arguments and finally $\hat{Y} \geq h(\hat{Y}, \mathbb{E}[\hat{Y}])$.

Next, for $t \leq T$, let us set $\Phi(t) := \mathbb{E}[\hat{Y}_t]$ and show that Φ is continuous.

Indeed, first note that since the process $(\hat{Y}_t + \int_0^t f(s, \hat{Y}_s, \mathbb{E}[\hat{Y}_s]) ds)_{t \leq T}$ is a supermartingale, the possible jumps of the process \hat{Y} are only negative. Suppose now that there exists t such that $\Delta\Phi(t) < 0$. Then $\mathbb{P}\{\omega, \Delta\hat{Y}_t(\omega) < 0\} > 0$ and $\Delta\Phi(t) = \mathbb{E}[\Delta\hat{Y}_t]$. But (4.8) and the characterization of the jumps of the Snell envelope (see e.g. [Ham02]) imply that

$$-\Delta\hat{Y}_t = \lim_{s \nearrow t} (h(\hat{Y}_s, \mathbb{E}[\hat{Y}_s]) - h(\hat{Y}_t, \mathbb{E}[\hat{Y}_t])) \leq \gamma_1(-\Delta\hat{Y}_t) + \gamma_2(-\Delta\Phi(t)).$$

Take now the expectation in both hand-sides to obtain that

$$-\Delta\Phi(t) \leq \gamma_1(-\Delta\Phi(t)) + \gamma_2(-\Delta\Phi(t)).$$

But, this contradicts the inequality $\gamma_1 + \gamma_2 < 1$ (see (3.8) for $p = 2$ and Remark 3.1). Therefore, Φ is continuous.

We now show that \hat{Y} is continuous. Suppose there exists a stopping time σ such that $\Delta\hat{Y}_\sigma < 0$. Then the characterization of jumps of a Snell envelope of processes imply that

$$-\Delta\hat{Y}_\sigma \leq h(\hat{Y}_{\sigma-}, \mathbb{E}[\hat{Y}_t]_{t=\sigma}) - h(\hat{Y}_\sigma, \mathbb{E}[\hat{Y}_t]_{t=\sigma}) \leq \gamma_1(-\Delta\hat{Y}_\sigma)$$

since $\Phi(t) = \mathbb{E}[\hat{Y}_t]$ is continuous. The last inequality is absurd since $\gamma_1 < 1$. Thus, for any stopping time σ , $\Delta\hat{Y}_\sigma = 0$ i.e. \hat{Y} is continuous.

Step 4. The sequence $(Y^n, Z^n, K^n)_{n \geq 0}$ converges to $(\hat{Y}, \hat{Z}, \hat{K})$ solution of the reflected BSDE (4.4).

First note that since \hat{Y} is continuous then, by Dini's theorem, the convergence of $(Y^n)_{n \geq 0}$ to \hat{Y} is uniform on $[0, T]$, i.e.,

$$\mathbb{P}\text{-a.s.}, \quad \lim_{n \rightarrow \infty} \sup_{t \leq T} |Y_t^n - \hat{Y}_t| = 0.$$

Next, the characterization (4.8) implies the existence of a martingale \hat{M} and a continuous increasing process \hat{K} ($\hat{K}_0 = 0$) such that

$$\hat{Y}_t + \int_0^t f(s, \hat{Y}_s, \mathbb{E}[\hat{Y}_s]) ds = \hat{M}_t - \hat{K}_t, \quad t \leq T.$$

Moreover, in view of (4.7), we have

$$\mathbb{E}[(\hat{K}_T)^2] < \infty \quad \text{and then} \quad \mathbb{E}[\sup_{t \leq T} |\hat{M}_t|^2] < \infty.$$

The rest of the proof is classical i.e the martingale representation provides the process \hat{Z} and \hat{K} satisfies the Skorohod condition, i.e., $(\hat{Y}, \hat{Z}, \hat{K})$ is a solution of the mean-field reflected BSDE (2.1) (see [EKKP⁺97] for more details).

Step 5. The minimality property

Lastly, let us show that this solution is minimal. Let (Y', Z', K') be another solution of (2.1) without imposing the Skorohod type condition. Then, by comparison of solutions of mean-field BSDEs (see Theorem 6.5 below) we obtain $Y^0 \leq Y'$. Next, assume that for some $n \geq 0$, we have $Y^n \leq Y'$. Once more by comparison we obtain $Y^{n+1} \leq Y'$ since

$$\begin{aligned} & f(t, y, \mathbb{E}[Y_t^n]) + (n+1)(y - h(Y_t^n, \mathbb{E}[Y_t^n]))^- \Big|_{y=Y_t'} \\ &= f(t, Y_t', \mathbb{E}[Y_t^n]) \leq f(t, Y_t', \mathbb{E}[Y_t']) \leq f(t, Y_t', \mathbb{E}[Y_t']) + dK_t'. \end{aligned}$$

Thus, for any $n \geq 0$, $Y^n \leq Y'$ which implies, in taking the limit, that $Y \leq Y'$ and then Y is the minimal solution. \square

Corollary 4.1. *Assume that (γ_1, γ_2) satisfies (3.8) and f, h and ξ satisfy Assumption (H2). Then, the sequence $(Y^n, Z^n, K^n)_{n \geq 0}$ converges in $\mathcal{S}_c^2 \times \mathcal{H}^{2,d} \times \mathcal{S}_c^2$ to the unique solution of the mean-field RBSDE (2.1) and the Skorohod condition induces the minimality of the solution.*

Proof. Let θ be a real constant and (Y, Z, K) be a solution of (2.1). For $t \leq T$, set

$$\underline{Y}_t = e^{\theta t} Y_t, \quad \underline{Z}_t = e^{\theta t} Z_t \quad \text{and} \quad \underline{K}_t = e^{\theta t} K_t.$$

Then, $(\underline{Y}, \underline{Z}, \underline{K})$ is a solution of the following mean-field RBSDE:

$$(4.9) \quad \begin{cases} \underline{Y} \in \mathcal{S}_c^2, \quad \underline{Z} \in \mathcal{H}^{2,d}, \quad \underline{K} \in \mathcal{S}_c^2; \\ \underline{Y}_t = e^{\theta T} \xi + \int_t^T \underline{F}(s, \underline{Y}_s, \mathbb{E}[\underline{Y}_s]) ds + \underline{K}_T - \underline{K}_t - \int_t^T \underline{Z}_s dB_s, \quad t \geq T; \\ \underline{Y}_t \geq e^{\theta t} h(e^{-\theta t} \underline{Y}_t, e^{-\theta t} \mathbb{E}[\underline{Y}_t]), \quad \forall t \leq T, \\ \int_0^T (\underline{Y}_t - e^{\theta t} h(e^{-\theta t} \underline{Y}_t, e^{-\theta t} \mathbb{E}[\underline{Y}_t])) d\underline{K}_t = 0, \end{cases}$$

where

$$\underline{F} : (t, y, y') \mapsto e^{\theta t} f(t, e^{-\theta t} y, e^{-\theta t} y') - \theta y.$$

But, by choosing θ appropriately we make that the function \underline{F} satisfy all the assumptions (H2). Therefore, by Theorem 4.3, its penalization scheme $(\underline{Y}^n, \underline{Z}^n, \underline{K}^n)$, $n \geq 0$, defined similarly as in (4.3), converges in $\mathcal{S}_c^2 \times \mathcal{H}^{2,d} \times \mathcal{S}_c^2$ to $(\underline{Y}, \underline{Z}, \underline{K})$ the unique solution of (4.9). But, by uniqueness, for any $t \leq T$,

$$\underline{Y}_t^n = e^{\theta t} Y_t^n, \quad \underline{Z}_t^n = e^{\theta t} Z_t^n \quad \text{and} \quad d\underline{K}_t^n = e^{\theta t} dK_t^n.$$

Therefore $(Y^n, Z^n, K^n)_{n \geq 0}$ converges in $\mathcal{S}_c^2 \times \mathcal{H}^{2,d} \times \mathcal{S}_c^2$ to (Y, Z, K) , the unique solution of (2.1). \square

Remark 4.2. *The constraint in (H2)-b) related to the monotonicity of $y \mapsto h(y, y')$ can also be relaxed substantially. Indeed, for $\kappa > 0$, let us set*

$$\Psi_\kappa : (y, y') \mapsto \frac{1}{1 + \kappa \gamma_1} h(y, y') + \frac{\kappa \gamma_1}{1 + \kappa \gamma_1} y.$$

Observe, that, for any $\kappa > 1$, the mapping $y \mapsto \Psi_\kappa(y, y')$ is non decreasing w.r.t y even when $y \mapsto h(y, y')$ does not enjoy any specific property of monotonicity. On the other hand, the condition $y \geq h(y, y')$ is equivalent to $y \geq \Psi_\kappa(y, y')$, and the corresponding Skorohod conditions also coincide. Therefore as soon as the Lipschitz constants of Ψ_κ verify the condition (4.10), i.e.,

$$(4.10) \quad \left(\frac{\gamma_1 + \kappa \gamma_1}{1 + \kappa \gamma_1} + \frac{\gamma_2}{1 + \kappa \gamma_1} \right)^{\frac{p-1}{p}} \left\{ \left(\frac{p}{p-1} \right)^p \frac{\gamma_1 + \kappa \gamma_1}{1 + \kappa \gamma_1} + \frac{\gamma_2}{1 + \kappa \gamma_1} \right\}^{\frac{1}{p}} < 1,$$

the penalization scheme of the MFBSDE associated with coefficients (f, ξ, Ψ_κ) converges to the solution of the MFBSDE of interest associated with (f, ξ, h) .

5. A MORE GENERAL CASE: f FURTHER DEPENDS ON z

The last Section of the paper is dedicated to the more involved framework, where the driver f may also depend on the Z process and, once again, in this Section we restrict to the case where $p = 2$, for the sake of simplicity.

The following assumptions on the mappings f , h and terminal value ξ will be assumed hereafter.

Assumption (H3). The coefficients f , h and ξ satisfy the following set of assumptions.

- (i) f is a mapping from $[0, T] \times \Omega \times \mathbb{R}^{2+d}$ into \mathbb{R} such that
 - (a) the process $(f(t, 0, 0, 0))_{t \leq T}$ is \mathcal{P} -measurable and belongs to $\mathcal{H}^{2,1}$;
 - (b) f is Lipschitz w.r.t. (y, z, y') uniformly in (t, ω) , i.e., there exists a positive constant C_f such that \mathbb{P} -a.s. for all $t \in [0, T]$,

$$|f(t, y_1, z_1, y'_1) - f(t, y_2, z_2, y'_2)| \leq C_f (|y_1 - y_2| + |z_1 - z_2| + |y'_1 - y'_2|).$$

for any $y_1, y'_1, y_2, y'_2 \in \mathbb{R}$ and z_1, z_2 in \mathbb{R}^d .

- (c) the mapping $y' \in \mathbb{R} \mapsto f(t, y, z, y')$ is non-decreasing, for t, y, z fixed.

- (d) The domination condition: There exists a measurable function $\Phi(t, \omega, y, y')$ from $[0, T] \times \Omega \times \mathbb{R}^2$ into \mathbb{R}^+ such that: i) Φ is Lipschitz in (y, y') uniformly w.r.t (t, ω) ; ii) the process $(\Phi(t, \omega, 0, 0))_{t \leq T}$ belongs to $\mathcal{H}^{2,1}$; iii) \mathbb{P} -a.s. for any t, y, y', z we have

$$f(t, \omega, y, z, y') \leq \Phi(t, \omega, y, y').$$

- (ii) (a) h is a mapping from \mathbb{R}^2 into \mathbb{R} which is Lipschitz w.r.t. (y, y') , i.e., there exist two positive constants γ_1 and γ_2 such that for any x, y, x' and y'

$$|h(x, x') - h(y, y')| \leq \gamma_1|x - y| + \gamma_2|x' - y'|,$$

where γ_1 and γ_2 are two positive constants.

- (b) For any fixed y (resp. y'), the mapping $y' \in \mathbb{R} \mapsto h(y, y')$ (resp. $y \in \mathbb{R} \mapsto h(y, y')$) is non-decreasing.

- (iii) ξ is an \mathcal{F}_T -measurable, \mathbb{R} -valued r.v., $\mathbb{E}[\xi^2] < \infty$ and satisfies $\xi \geq h(\xi, \mathbb{E}[\xi])$.

We have the following result.

Theorem 5.4. *Suppose that (γ_1, γ_2) satisfies (3.8) with $p = 2$ and that Assumption (H3) is fulfilled. Then there exists a triplet of processes (Y, Z, K) which is the minimal solution of the following mean-field reflected BSDE:*

$$(5.1) \quad \begin{cases} Y \in \mathcal{S}_c^2, Z \in \mathcal{H}^{2,d} \text{ and } K \in \mathcal{S}_{ci}^2; \\ Y_t = \xi + \int_t^T f(s, Y_s, Z_s, \mathbb{E}[Y_s]) ds + K_T - K_t - \int_t^T Z_s dB_s, \quad t \leq T; \\ Y_t \geq h(Y_t, \mathbb{E}[Y_t]), \quad \forall t \geq T \text{ and } \int_0^T (Y_t - h(Y_t, \mathbb{E}[Y_t])) dK_t = 0. \end{cases}$$

Proof. Let $Y^0 = \underline{Y}$ where $(\underline{Y}, \underline{Z})$ is the solution of the following BSDE of mean field type (which exists according to Theorem 3.1. in [BLP09]):

$$(5.2) \quad \begin{cases} \underline{Y} \in \mathcal{S}_c^2, \quad \underline{Z} \in \mathcal{H}^{2,d}; \\ \underline{Y}_t = \xi + \int_t^T f(s, \underline{Y}_s, \underline{Z}_s, \mathbb{E}[\underline{Y}_s]) ds - \int_t^T \underline{Z}_s dB_s, \quad t \leq T. \end{cases}$$

For $n \geq 1$, we define (Y^n, Z^n) as the solution of the following standard BSDE:

$$(5.3) \quad \begin{cases} Y^n \in \mathcal{S}_c^2, Z^n \in \mathcal{H}^{2,d}; \\ Y_t^n = \xi + \int_t^T f(s, Y_s^n, Z_s^n, \mathbb{E}[Y_s^{n-1}]) ds + K_T^n - K_t^n - \int_t^T Z_s^n dB_s, \quad t \leq T, \end{cases}$$

where, for $n \geq 1$, K^n denotes the process

$$K_t^n := n \int_0^t (Y_s^n - h(Y_s^{n-1}, \mathbb{E}[Y_s^{n-1}]))^- ds, \quad t \leq T.$$

Finally, let $(\bar{Y}, \bar{Z}, \bar{K})$ be the unique solution of the following mean-field reflected BSDE:

$$(5.4) \quad \begin{cases} \bar{Y} \in \mathcal{S}_c^2, \bar{Z} \in \mathcal{H}^{2,d} \text{ and } \bar{K} \in \mathcal{S}_{ci}^2; \\ \bar{Y}_t = \xi + \int_t^T \Phi(s, \bar{Y}_s, \mathbb{E}[\bar{Y}_s]) ds + \bar{K}_T - \bar{K}_t - \int_t^T \bar{Z}_s dB_s, \quad t \leq T; \\ \bar{Y}_t \geq h(\bar{Y}_t, \mathbb{E}[\bar{Y}_t]), \quad \forall t \geq T \text{ and } \int_0^T (\bar{Y}_t - h(\bar{Y}_t, \mathbb{E}[\bar{Y}_t])) d\bar{K}_t = 0. \end{cases}$$

The triplet of processes $(\bar{Y}, \bar{Z}, \bar{K})$ solution of (5.4) exists by Theorem 3.1.

The proof of the theorem will be obtained in the following steps.

Step 1.

$$(5.5) \quad \text{For any } n \geq 0, \quad Y^0 \leq Y^n \leq Y^{n+1} \leq \bar{Y}.$$

The fact that $Y^0 \leq Y^1$ is obtained by the comparison result by using the standard argument of Itô's formula with $((Y^0 - Y^1)^+)^2$ since the driver of Y^1 is $f(t, y, z, \mathbb{E}[Y_t^0]) + (y - \mathbb{E}[Y_t^0])^+$ which is greater than $f(t, y, z, \mathbb{E}[Y_t^0])$. Actually, we can easily show that $\mathbb{E}[(Y^0 - Y^1)^+] = 0$, which is the desired result. Suppose that, for some $n \geq 0$, we have $Y^n \leq Y^{n+1}$. Then we also have $Y^{n+1} \leq Y^{n+2}$ by the comparison theorem of solutions

since the mappings $y' \in \mathbb{R} \mapsto f(t, y, z, y')$, $y' \in \mathbb{R} \mapsto h(y, y')$ and $y \in \mathbb{R} \mapsto h(y, y')$ are nondecreasing. Finally, let us show by induction that for any $n \geq 0$, $Y^n \leq \tilde{Y}$. For $n = 0$, it holds true by the comparison theorem of solutions of mean-field BSDEs since the function $y' \in \mathbb{R} \mapsto f(t, y, z, y')$ is non-decreasing and $d\tilde{K} \geq 0$ (see Theorem 6.5 in the appendix). Next, assume that the property holds true for some $n \geq 0$, i.e. $Y^n \leq \tilde{Y}$. But the solution of (5.4) is unique, then $(\tilde{Y}, \tilde{Z}, \tilde{K}) = (\tilde{Y}, \tilde{Z}, \tilde{K})$ where $(\tilde{Y}, \tilde{Z}, \tilde{K})$ is the unique solution of the following standard BSDE:

$$(5.6) \quad \begin{cases} \tilde{Y} \in \mathcal{S}_c^2, \tilde{Z} \in \mathcal{H}^{2,d} \text{ and } \tilde{K} \in \mathcal{S}_{ci}^2; \\ \tilde{Y}_t = \xi + \int_t^T \Phi(s, \tilde{Y}_s, \mathbb{E}[\tilde{Y}_s]) ds + \tilde{K}_T - \tilde{K}_t - \int_t^T \tilde{Z}_s dB_s, \quad t \leq T; \\ \tilde{Y}_t \geq h(\tilde{Y}_t, \mathbb{E}[\tilde{Y}_t]), \quad \forall t \leq T \text{ and } \int_0^T (\tilde{Y}_t - h(\tilde{Y}_t, \mathbb{E}[\tilde{Y}_t])) d\tilde{K}_t = 0. \end{cases}$$

Now, by the induction hypothesis and the fact that the mappings, $y' \in \mathbb{R} \mapsto \Phi(t, y, y')$, $y' \in \mathbb{R} \mapsto f(t, y, z, y')$, $y' \in \mathbb{R} \mapsto h(y, y')$ and $y \in \mathbb{R} \mapsto h(y, y')$ being nondecreasing, we obtain

$$\begin{aligned} f(t, y, z, \mathbb{E}[Y_t^n]) + (n+1)(y - h(Y_t^n, \mathbb{E}[Y_t^n]))^- \Big|_{(y,z)=(\tilde{Y}_t, \tilde{Z}_t)} \\ = f(t, \tilde{Y}_t, \tilde{Z}_t, \mathbb{E}[Y_t^n]) \leq f(t, \tilde{Y}_t, \tilde{Z}_t, \mathbb{E}[\tilde{Y}_t]) \leq \Phi(s, \tilde{Y}_s, \mathbb{E}[\tilde{Y}_s]) \leq \Phi(s, \tilde{Y}_s, \mathbb{E}[\tilde{Y}_s]) + d\tilde{K}_t. \end{aligned}$$

Therefore, by the comparison theorem of the solutions of standard BSDEs, we obtain $Y^{n+1} \leq \tilde{Y} = \tilde{Y}$, which completes the proof.

Step 2. For any $t \leq T$, we set $Y_t = \lim_{n \rightarrow \infty} Y_t^n$. The process Y is rcll and there exist a \mathcal{P} -measurable process Z such that for any $p \in [1, 2)$, $(Z^n)_{n \geq 0}$ converge to Z in $L^p(dt \otimes d\mathbb{P})$.

First, note that the process Y exists since by Step 1 the sequence of processes $(Y^n)_{n \geq 0}$ is non-decreasing and $Y^n \leq \tilde{Y}$. On the other hand since Y^0 and \tilde{Y} belongs to \mathcal{S}_c^2 then

$$(5.7) \quad \mathbb{E}[\sup_{t \leq T} |Y_t|^2] \leq C.$$

Furthermore, using Itô's formula and taking into account the representation of Y^n similar to (4) we have, for any $t \leq T$ and $\varepsilon > 0$,

$$(5.8) \quad \begin{aligned} (Y_t^n)^2 + \int_t^T |Z_s^n|^2 ds &= \xi^2 + 2 \int_t^T Y_s^n f(s, Y_s^n, Z_s^n, \mathbb{E}[Y_s^{n-1}]) ds + 2 \int_t^T Y_s^n dK_s^n - 2 \int_t^T Y_s^n Z_s^n dB_s \\ &\leq \xi^2 + 2 \int_t^T Y_s^n \{f(s, 0, 0, 0) + a(s)Y_s^n + b(s)Z_s^n + c(s)\mathbb{E}[Y_s^{n-1}]\} ds \\ &\quad + 2K_T^n \sup_{s \leq T} \{Y_s^n \wedge h(Y_s^{n-1}, \mathbb{E}[Y_s^{n-1}])\} - 2 \int_t^T Y_s^n Z_s^n dB_s \end{aligned}$$

$$(5.9) \quad \begin{aligned} &\leq \xi^2 + 2 \int_t^T Y_s^n \{f(s, 0, 0, 0) + a(s)Y_s^n + b(s)Z_s^n + c(s)\mathbb{E}[Y_s^{n-1}]\} ds \\ &\quad + \varepsilon(K_T^n)^2 + \varepsilon^{-1}(\sup_{s \leq T} \{Y_s^n \wedge h(Y_s^{n-1}, \mathbb{E}[Y_s^{n-1}])\})^2 - 2 \int_t^T Y_s^n Z_s^n dB_s \end{aligned}$$

where $a(\cdot)$, $b(\cdot)$ and $c(\cdot)$ are measurable processes bounded by the Lipschitz constant of f . Now, taking into account of the inequalities of Step 1 and the monotonicity property of h to deduce that, for any $n \geq 1$,

$$\mathbb{E}[(\sup_{s \leq T} \{Y_s^n \wedge h(Y_s^{n-1}, \mathbb{E}[Y_s^{n-1}])\})^2] \leq \mathbb{E}[(\sup_{s \leq T} h(\tilde{Y}_s, \mathbb{E}[\tilde{Y}_s]))^2] + \mathbb{E}[(\sup_{s \leq T} \{Y_s^1 \wedge h(Y_s^0, \mathbb{E}[Y_s^0])\})^2].$$

As

$$K_T^n = Y_0^n - \xi - \int_0^T f(s, Y_s^n, Z_s^n, \mathbb{E}[Y_s^{n-1}]) ds + \int_0^T Z_s^n dB_s, \quad t \leq T,$$

then, by using (5.5), the inequality $(u + v + w + z)^2 \leq 4(u^2 + v^2 + w^2 + z^2)$, the Burkholder-Davis-Gundy inequality and finally by choosing ε appropriately we obtain

$$\mathbb{E}\left[\int_0^T |Z_s^n|^2 ds\right] \leq C \text{ and } \mathbb{E}\left[\int_0^T |f(s, Y_s^n, Z_s^n, \mathbb{E}[Y_s^{n-1}])|^2 ds\right] \leq C$$

where C is a constant independent of n . Consequently, in view of Theorem 2.1 in [Pen99], we obtain that

- (a) Y is rcll;
- (b) there exist a \mathcal{P} -measurable process Z such that for any $p \in [1, 2)$,

$$\lim_{n \rightarrow \infty} \mathbb{E}\left[\int_0^T |Z_s^n - Z_s|^p ds\right] = 0.$$

Step 3. The process Y is continuous and satisfies

$$(5.10) \quad Y_t = \operatorname{ess\,sup}_{\tau \geq t} \mathbb{E}\left[\int_t^\tau f(s, Y_s, Z_s, \mathbb{E}[Y_s]) ds + h(Y_\tau, \mathbb{E}[Y_t]_{|t=\tau}) 1_{\{\tau < T\}} + \xi 1_{\{\tau = T\}} \mid \mathcal{F}_t\right].$$

For any $n \geq 1$, in view of the representation (4), we have, for all $t \in [0, T]$,

$$(5.11) \quad \begin{aligned} Y_t^n &= \operatorname{ess\,sup}_{\tau \geq t} \mathbb{E}\left[\int_t^\tau f(s, Y_s^n, Z_s^n, \mathbb{E}[Y_s^{n-1}]) ds + Y_\tau^n \wedge h(Y_\tau^{n-1}, \mathbb{E}[Y_t^{n-1}]_{|t=\tau}) 1_{\{\tau < T\}} + \xi 1_{\{\tau = T\}} \mid \mathcal{F}_t\right] \\ &= G^n(t) + H^n(t) \end{aligned}$$

where, for all $t \leq T$,

$$\begin{aligned} G^n(t) &= \operatorname{ess\,sup}_{\tau \geq t} \mathbb{E}\left[\int_t^\tau f(s, Y_s^n, Z_s^n, \mathbb{E}[Y_s^{n-1}]) ds + Y_\tau^n \wedge h(Y_\tau^{n-1}, \mathbb{E}[Y_t^{n-1}]_{|t=\tau}) 1_{\{\tau < T\}} + \xi 1_{\{\tau = T\}} \mid \mathcal{F}_t\right] \\ &\quad - \operatorname{ess\,sup}_{\tau \geq t} \mathbb{E}\left[\int_t^\tau f(s, Y_s, Z_s, \mathbb{E}[Y_s]) ds + Y_\tau^n \wedge h(Y_\tau^{n-1}, \mathbb{E}[Y_t^{n-1}]_{|t=\tau}) 1_{\{\tau < T\}} + \xi 1_{\{\tau = T\}} \mid \mathcal{F}_t\right] \end{aligned}$$

and

$$H^n(t) = \operatorname{ess\,sup}_{\tau \geq t} \mathbb{E}\left[\int_t^\tau f(s, Y_s, Z_s, \mathbb{E}[Y_s]) ds + Y_\tau^n \wedge h(Y_\tau^{n-1}, \mathbb{E}[Y_t^{n-1}]_{|t=\tau}) 1_{\{\tau < T\}} + \xi 1_{\{\tau = T\}} \mid \mathcal{F}_t\right].$$

But, for any $t \leq T$,

$$|G^n(t)| \leq \mathbb{E}\left[\int_0^T |f(s, Y_s^n, Z_s^n, \mathbb{E}[Y_s^{n-1}]) - f(s, Y_s, Z_s, \mathbb{E}[Y_s])| ds \mid \mathcal{F}_t\right]$$

since $|\operatorname{ess\,sup}_{\tau \geq t} \Sigma_{t,\tau}^1 - \operatorname{ess\,sup}_{\tau \geq t} \Sigma_{t,\tau}^2| \leq \operatorname{ess\,sup}_{\tau \geq t} |\Sigma_{t,\tau}^1 - \Sigma_{t,\tau}^2|$. Therefore, by the convergence results stated in Step 2 for $(Y^n)_n$ and $(Z^n)_n$ and using Doob's inequality, the sequence of processes $(G^n)_{n \geq 0}$ converges to 0 in \mathcal{S}_c^p for any $p \in [1, 2)$. On the other hand, using the same argument as in (4.8), the sequence $(H^n)_{n \geq 1}$ converges increasingly to the following rcll process H defined, for any $t \leq T$, by

$$H(t) = \operatorname{ess\,sup}_{\tau \geq t} \mathbb{E}\left[\int_t^\tau f(s, Y_s, Z_s, \mathbb{E}[Y_s]) ds + Y_\tau \wedge h(Y_\tau, \mathbb{E}[Y_t]_{|t=\tau}) 1_{\{\tau < T\}} + \xi 1_{\{\tau = T\}} \mid \mathcal{F}_t\right].$$

Thus, for any $t \leq T$, $Y_t = H(t)$. Finally, the continuity of Y is obtained in the same way as in Step 3 of the previous subsection, since (γ_1, γ_2) satisfies (3.8). The proof of the claim is now complete.

Step 4. For any $t \leq T$, we define the continuous process K by

$$K_t = Y_0 - Y_t - \int_0^t f(s, Y_s, Z_s, \mathbb{E}[Y_s]) ds + \int_0^t Z_s dB_s.$$

The triple of processes (Y, Z, K) is a solution of the mean-field reflected BSDE (5.1) and it is minimal.

The process K is nothing but the limit of the sequence $(K^n)_n$ and since for any n , K^n is non-decreasing then K is also non-decreasing. On the other hand, by Dini's Theorem $(Y^n)_n$ converges to Y in \mathcal{S}_c^2 and $(Z^n)_n$ converges to Z in $L^p(dt \otimes d\mathbb{P})$ for any $p \in [1, 2)$, then, by Burkholder-Davis-Gundy inequality and Jensen's inequality, $(K^n)_n$ converges in \mathcal{S}_c^p ($p \in [1, 2)$) to K .

Finally, let us show that the Skorohod condition $\int_0^T (Y_s - h(Y_s, \mathbb{E}[Y_s])) dK_s = 0$ holds. Consider a subsequence which we still denote by $\{n\}$ such that

$$\mathbb{P}\text{-a.s.}, \quad \limsup_{n \rightarrow \infty} \sup_{s \leq T} |K_s^n - K_s| \rightarrow 0.$$

As pointed out previously, for any $n \geq 0$, we have

$$\int_0^T (Y_s^n - Y_s^n \wedge h(Y_s^n, \mathbb{E}[Y_s^n])) dK_s^n = 0.$$

On the other hand

$$\begin{aligned} & \int_0^T (Y_s^n - Y_s^n \wedge h(Y_s^n, \mathbb{E}[Y_s^n])) dK_s^n \\ &= \int_0^T \{ (Y_s^n - Y_s^n \wedge h(Y_s^n, \mathbb{E}[Y_s^n])) - (Y_s - h(Y_s, \mathbb{E}[Y_s])) \} dK_s^n \\ & \quad + \int_0^T (Y_s - h(Y_s, \mathbb{E}[Y_s])) dK_s^n = A_1^n + A_2^n. \end{aligned}$$

But,

$$|A_1^n| \leq \sup_{s \leq T} | (Y_s^n - Y_s^n \wedge h(Y_s^n, \mathbb{E}[Y_s^n])) - (Y_s - h(Y_s, \mathbb{E}[Y_s])) | \times K_T^n.$$

Thus, \mathbb{P} -a.s. $\lim_{n \rightarrow \infty} A_1^n \rightarrow 0$, since K_T^n is bounded thanks to the uniform convergence of $(K^n)_n$ to K , the uniform convergence of $(Y^n)_n$ to Y , \mathbb{P} -a.s. and in \mathcal{S}_c^2 , and finally the fact that h is Lipschitz. On the other hand, in view of Helly's Convergence Theorem (see [KF70], pp. 370), we have

$$\mathbb{P}\text{-a.s.}, \quad \lim_{n \rightarrow \infty} A_2^n = \int_0^T (Y_s - h(Y_s, \mathbb{E}[Y_s])) dK_s.$$

Therefore, $\int_0^T (Y_s - h(Y_s, \mathbb{E}[Y_s])) dK_s = 0$ which means that the Skorohod condition is satisfied and then (Y, Z, K) is a solution of (5.1).

Lastly, the minimality property follows in a similar manner as in Theorem 4.3 above. \square

Remark 5.1. Since $(Y^n)_n$ converges to Y in \mathcal{S}_c^2 , this entails the convergence of $(Z^n)_n$ to Z in $\mathcal{H}^{2,d}$.

6. APPENDIX: COMPARISON OF SOLUTIONS OF BSDEs OF MEAN-FIELD TYPE

For sake of completeness, we recall the following result related to comparison of solutions of mean-field BSDEs given in ([BLP09], Theorem 3.2) in a more general setting. We present here the form which we need it throughout this paper.

Theorem 6.5 ([BLP09], Theorem 3.2). *Let us consider the two following two BSDEs of mean-field type: For any $t \leq T$,*

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s, \mathbb{E}[Y_s]) ds + \int_t^T Z_s dB_s$$

and

$$(6.1) \quad Y'_t = \xi' + \int_t^T f'(s, Y'_s, Z'_s, \mathbb{E}[Y'_s]) ds + \int_t^T Z'_s dB_s.$$

We assume that the mappings f and f' satisfy (H3)-(i) (a)-(b) and ξ, ξ' are square integrable. If

- (a) $\xi \leq \xi'$,
- (b) $f(t, Y'_t, Z'_t, \mathbb{E}[Y'_t]) \leq f(t, Y_t, Z_t, \mathbb{E}[Y_t])$,
- (c) f is non-decreasing w.r.t. y' .

Then $Y \leq Y'$.

Remark 6.5. *In this theorem, we do not need that f' satisfies (H3)-(i) (a)-(b) but only the existence of the solution (Y', Z') of (6.1).*

REFERENCES

- [BEH18] P. Briand, R. Elie, and Y. Hu, *BSDEs with mean reflection*, The Annals of Applied Probability **28** (2018), no. 1, 482–510.
- [BDH⁺03] P. Briand, B. Delyon, Y. Hu, E. Paroux, and L. Stoica, *BSDEs with mean reflection*, Stochastic Processes and their Applications **108** (2003), no. 1, 109–129.
- [BLP09] R. Buckdahn, J. Li, and S. Peng, *Mean-field backward stochastic differential equations and related partial differential equations*, Stochastic Processes and their Applications **119** (2009), no. 10, 3133–3154.
- [CDD14] M. Christiansen, M. M Denuit, and J. Dhaene, *Reserve-dependent benefits and costs in life and health insurance contracts*, Insurance: Mathematics and Economics **57** (2014), 132–137.
- [Li14] J. Li, *Reflected mean-field backward stochastic differential equations. Approximation and associated non-linear PDEs*, Journal of Mathematical Analysis and Applications **413** (2014), 47–68.
- [DM82] C Dellacherie and P A Meyer, *Probabilities and potential B, Chapter V to VIII*, North-Holland Amsterdam, 1982.
- [RY13] D. Revuz and M. Yor, *Continuous martingales and Brownian motion*, Springer Science & Business Media, 2013.
- [DL14] B. Djehiche and B. Löfdahl, *Risk aggregation and stochastic claims reserving in disability insurance*, Insurance: Mathematics and Economics **59** (2014), 100–108.
- [DHP09] B. Djehiche, S. Hamadene, and A. Popier, *A finite horizon optimal multiple switching problem*, SIAM Journal on Control and Optimization **48** (2009), no. 4, 2751–2770.
- [DE92a] D. Duffie and L. G Epstein, *Stochastic differential utility*, Econometrica: Journal of the Econometric Society (1992), 353–394.
- [DE92b] D. Duffie and L. G Epstein, *Asset pricing with stochastic differential utility*, The Review of Financial Studies **5** (1992), no. 3, 411–436.
- [EKQP97] N. El Karoui, S. Peng, and M. C. Quenez, *Backward stochastic differential equations in finance*, Mathematical finance **7** (1997), no. 1, 1–71.
- [EKKP⁺97] N. El Karoui, C. Kapoudjian, É. Pardoux, S. Peng, and M.-C. Quenez, *Reflected solutions of backward SDE's, and related obstacle problems for PDE's*, the Annals of Probability **25** (1997), no. 2, 702–737.
- [Ham02] S. Hamadene, *Reflected BSDE's with discontinuous barrier and application*, Stochastics: An International Journal of Probability and Stochastic Processes **74** (2002), no. 3-4, 571–596.
- [KF70] A. Kolmogorov and S. Fomin, *Introductory Real Analysis (translated and edited by Richard A. Silverman)* Prentice-Hall, 1970.
- [Nor91] R. Norberg, *Reserves in life and pension insurance*, Scandinavian Actuarial Journal **1991** (1991), no. 1, 3–24.
- [Nor92] R. Norberg, *Hattendorff's theorem and Thiele's differential equation generalized*, Scandinavian Actuarial Journal **1992** (1992), no. 1, 2–14.
- [PP90] E. Pardoux and S. Peng, *Adapted solution of a backward stochastic differential equation*, Systems & Control Letters **14** (1990), no. 1, 55–61.
- [Pen99] S. Peng, *Monotonic limit theorem of BSDE and nonlinear decomposition theorem of Doob–Meyers type*, Probability theory and related fields **113** (1999), no. 4, 473–499.
- [HJ07] S. Hamadene and M. Jeanblanc, *On the starting and stopping problem: application in reversible investments*, Math. Oper. Res. **32** (2007), no. 1, 182–192.
- [Ber18] C. Bertucci, *Optimal stopping in Mean Field Games, an obstacle problem approach*, J; Math. pures et appliquees **120** (2018), 165–194.
- [EK14] R. Elie and I. Kharroubi, *Adding constraints to BSDEs with Jumps : an alternative to multidimensional reflections*, ESAIM probability and statistics **18** (2014), 233–250.
- [CK96] J. Cvitanic and I. Karatzas, *Backward stochastic differential equations with reflection and Dynkin games*, Ann. Probab. **24** (1996), no. 4, 2024–2056.
- [CKS98] J. Cvitanic, I. Karatzas, and M. Soner, *Backward stochastic differential equations with constraints on the gains-process*, Ann. Probab. **26** (1998), no. 4, 1522–1551.
- [BER15] B. Bouchard, R. Elie, and A. Reveillac, *BSDEs with weak terminal condition*, Ann. Probab. **43** (2015), no. 2, 572–604.
- [DESZ17] R. Dumitrescu, R. Elie, W. Sabbagh, and C. Zhou, *BSDEs with weak reflections and partial hedging of American options*, Arxiv:1708.05957 (2017).
- [CR17] J.-F. Chassagneux and A. Richou, *Obliquely reflected BSDEs*, Arxiv:1710.08989 (2017).

- [MPZ13] A. Matoussi, D. Possamai, and C. Zhou, *BSDEs with weak terminal condition*, Ann. Applied Probab. **12** (2013), no. 6, 2420-2457.
- [RY13] D. Revuz and M. Yor, *Continuous martingales and Brownian motion*, Springer Science & Business Media, 2013.

DEPARTMENT OF MATHEMATICS, KTH ROYAL INSTITUTE OF TECHNOLOGY, 100 44, STOCKHOLM, SWEDEN

E-mail address: boualem@kth.se

LAMA UMR CNRS 8050, UNIVERSITÉ PARIS-EST, CHAMPS-SUR-MARNE, FRANCE

E-mail address: romuald.elie@univ-mlv.fr

LMM, LE MANS UNIVERSITY, AVENUE OLIVIER MESSIAEN, 72085 LE MANS, CEDEX 9, FRANCE

E-mail address: hamadene@univ-lemans.fr