

In the name of Allah, the Beneficent, the Merciful

ON IDENTITIES OF 2-DIMENSIONAL ALGEBRAS

H. AHMED¹, U.BEKBAEV², I.RAKHIMOV³

¹Depart. of Math., Faculty of Science, Taiz University, Taiz, Yemen

²Depart. of Mathematical and Natural Sciences, TTPU, Tashkent, Uzbekistan

³Depart. of Math., Faculty of Computer and Mathematical Sciences, UiTM, Shah Alam, Malaysia

& V.I.Romanovski Institute of Mathematics, Uzbekistan Academy of Sciences

ABSTRACT. In the paper we provide some polynomial identities for finite-dimensional algebras. A list of well known single polynomial identities is exposed and the classification of all 2-dimensional algebras with respect to these identities is given.

1. INTRODUCTION

It is known that many important algebras in use are so called PI-algebras, that is, algebras satisfying a certain set of polynomial identities. Therefore, the classification of such algebras, up to isomorphism, is of a great interest. Earlier we have given classification results for some important classes of two-dimensional PI-algebras [1, 3, 4, 5]. In this paper we consider a list of some important polynomial identities which have appeared earlier in the theory of algebras and present a classification of two-dimensional algebras with respect to these identities. For other results related to the classification problem and the problems raised in this paper we refer the reader to [6, 7, 8, 9, 10, 11].

The organization of the paper is as follows. In the next section we introduce definitions, notations and results needed in the course of the study followed by two section where we present main results of the paper. In Section 3 we provide some polynomial identities for finite-dimensional algebras. The last section is devoted to the classification of two-dimensional algebras with respect to the identities specified.

2. PRELIMINARIES

In this paper an algebra (\mathbb{A}, \cdot) means a vector space \mathbb{A} over a field \mathbb{F} with a given bilinear map

$$\cdot : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{A}, (\mathbf{u}, \mathbf{v}) \mapsto \mathbf{u} \cdot \mathbf{v}$$

and often we drop \cdot in the writing.

If \mathbb{A} such a two-dimensional algebra and $e = (e_1, e_2)$ is a fixed basis by

$A = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \beta_1 & \beta_2 & \beta_3 & \beta_4 \end{pmatrix}$ we denote its matrix of structural constants (MSC) with respect to this basis, i.e.,

$$e_1 e_1 = \alpha_1 e_1 + \beta_1 e_2, e_1 e_2 = \alpha_2 e_1 + \beta_2 e_2, e_2 e_1 = \alpha_3 e_1 + \beta_3 e_2, e_2 e_2 = \alpha_4 e_1 + \beta_4 e_2.$$

Further it is assumed that the basis e is fixed and we don't make difference between an algebra \mathbb{A} and its MSC A with respect to this basis.

The classification problem of all two dimensional algebras over any field \mathbb{F} , over that any the second and third degree polynomial has a root, has been solved in [2]. The classification there was done via providing the canonical MSCs for such algebras. In this paper we rely on the result of [2], follow its notations and for a convenience we present here the corresponding canonical representatives according to $\text{Char}(\mathbb{F}) \neq 2, 3$, $\text{Char}(\mathbb{F}) = 2$ and $\text{Char}(\mathbb{F}) = 3$ cases. Note that the parameters given in the canonical representatives may take any values in \mathbb{F} .

emails: ¹houida_m7@yahoo.com; ²uralbekbaev@gmail.com; ³isamiddin@uitm.edu.my.

Algebra	Structure constants							
	α_1	α_2	α_3	α_4	β_1	β_2	β_3	β_4
$A_1(\mathbf{c})$	α_1	α_2	$\alpha_2 + 1$	α_4	β_1	$-\alpha_1$	$-\alpha_1 + 1$	$-\alpha_2$
$A_2(\mathbf{c})$	α_1	0	0	1	β_1	β_2	$1 - \alpha_1$	0
$A_3(\mathbf{c})$	0	1	1	0	β_1	β_2	1	-1
$A_4(\mathbf{c})$	α_1	0	0	0	0	β_2	$1 - \alpha_1$	0
$A_5(\mathbf{c})$	α_1	0	0	0	1	$2\alpha_1 - 1$	$1 - \alpha_1$	0
$A_6(\mathbf{c})$	α_1	0	0	1	β_1	$1 - \alpha_1$	$-\alpha_1$	0
$A_7(\mathbf{c})$	0	1	1	0	β_1	1	0	-1
$A_8(\mathbf{c})$	α_1	0	0	0	0	$1 - \alpha_1$	$-\alpha_1$	0
A_9	$\frac{1}{3}$	0	0	0	1	$\frac{2}{3}$	$-\frac{1}{3}$	0
A_{10}	0	1	1	0	0	0	0	-1
A_{11}	0	1	1	0	1	0	0	-1
A_{12}	0	0	0	0	1	0	0	0

$$A_2(\mathbf{c}) = \begin{pmatrix} \alpha_1 & 0 & 0 & 1 \\ \beta_1 & \beta_2 & 1 - \alpha_1 & 0 \end{pmatrix} \cong \begin{pmatrix} \alpha_1 & 0 & 0 & 1 \\ -\beta_1 & \beta_2 & 1 - \alpha_1 & 0 \end{pmatrix}, \text{ where } \mathbf{c} = (\alpha_1, \beta_1, \beta_2) \in \mathbb{F}^3,$$

$$A_6(\mathbf{c}) = \begin{pmatrix} \alpha_1 & 0 & 0 & 1 \\ \beta_1 & 1 - \alpha_1 & -\alpha_1 & 0 \end{pmatrix} \cong \begin{pmatrix} \alpha_1 & 0 & 0 & 1 \\ -\beta_1 & 1 - \alpha_1 & -\alpha_1 & 0 \end{pmatrix}, \text{ where } \mathbf{c} = (\alpha_1, \beta_1) \in \mathbb{F}^2,$$

Algebra	The structure constants							
	α_1	α_2	α_3	α_4	β_1	β_2	β_3	β_4
$A_{1,2}(\mathbf{c})$	α_1	α_2	$1 + \alpha_2$	α_4	β_1	α_1	$1 + \alpha_1$	α_2
$A_{2,2}(\mathbf{c})$	α_1	0	0	1	β_1	β_2	$1 + \alpha_1$	0
$A_{3,2}(\mathbf{c})$	α_1	1	1	0	0	β_2	$1 + \alpha_1$	1
$A_{4,2}(\mathbf{c})$	α_1	0	0	0	0	β_2	$1 + \alpha_1$	0
$A_{5,2}(\mathbf{c})$	α_1	0	0	0	1	1	$1 + \alpha_1$	0
$A_{6,2}(\mathbf{c})$	α_1	0	0	1	β_1	$1 + \alpha_1$	α_1	0
$A_{7,2}(\mathbf{c})$	α_1	1	1	0	0	$1 + \alpha_1$	α_1	1
$A_{8,2}(\mathbf{c})$	α_1	0	0	0	0	$1 + \alpha_1$	α_1	0
$A_{9,2}$	1	0	0	0	1	0	1	0
$A_{10,2}$	0	1	1	0	0	0	0	1
$A_{11,2}$	1	1	1	0	0	1	1	1
$A_{12,2}$	0	0	0	0	1	0	0	0

Algebra	The structure constants							
	α_1	α_2	α_3	α_4	β_1	β_2	β_3	β_4
$A_{1,3}(\mathbf{c})$	α_1	α_2	$\alpha_2 + 1$	α_4	β_1	$-\alpha_1$	$-\alpha_1 + 1$	$-\alpha_2$
$A_{2,3}(\mathbf{c})$	α_1	0	0	1	β_1	β_2	$1 - \alpha_1$	0
$A_{3,3}(\mathbf{c})$	0	1	1	0	β_1	β_2	1	-1
$A_{4,3}(\mathbf{c})$	α_1	0	0	0	0	β_2	$1 - \alpha_1$	0
$A_{5,3}(\mathbf{c})$	α_1	0	0	0	1	$-\alpha_1 - 1$	$1 - \alpha_1$	0
$A_{6,3}(\mathbf{c})$	α_1	0	0	1	β_1	$1 - \alpha_1$	$-\alpha_1$	0
$A_{7,3}(\mathbf{c})$	0	1	1	0	β_1	1	0	-1
$A_{8,3}(\mathbf{c})$	α_1	0	0	0	0	$1 - \alpha_1$	$-\alpha_1$	0
$A_{9,3}$	0	1	1	0	1	0	0	-1
$A_{10,3}$	0	1	1	0	0	0	0	-1
$A_{11,3}$	1	0	0	0	1	-1	-1	0
$A_{12,3}$	0	0	0	0	1	0	0	0

$$A_{2,3}(\mathbf{c}) = \begin{pmatrix} \alpha_1 & 0 & 0 & 1 \\ \beta_1 & \beta_2 & 1 - \alpha_1 & 0 \end{pmatrix} \cong \begin{pmatrix} \alpha_1 & 0 & 0 & 1 \\ -\beta_1 & \beta_2 & 1 - \alpha_1 & 0 \end{pmatrix}, \text{ where } \mathbf{c} = (\alpha_1, \beta_1, \beta_2) \in \mathbb{F}^3,$$

$$A_{6,3}(\mathbf{c}) = \begin{pmatrix} \alpha_1 & 0 & 0 & 1 \\ \beta_1 & 1 - \alpha_1 & -\alpha_1 & 0 \end{pmatrix} \cong \begin{pmatrix} \alpha_1 & 0 & 0 & 1 \\ -\beta_1 & 1 - \alpha_1 & -\alpha_1 & 0 \end{pmatrix}, \text{ where } \mathbf{c} = (\alpha_1, \beta_1) \in \mathbb{F}^2.$$

3. POLYNOMIAL IDENTITIES OF TWO-DIMENSIONAL ALGEBRAS

Let \mathbb{F} be a field, $m \leq n \leq l$ be fixed natural numbers, S_n be the symmetric group and x^1, x^2, \dots, x^l be non-commutative, non-associative variables. By the use of these variables and parenthesis $(,)$ one can construct different non-associative monomials(words) containing each of the variables x^1, x^2, \dots, x^l only once and they may occur in the monomial in any order. For example, in the case of $l = 2$ one has only two such monomials x^1x^2 and x^2x^1 , whereas in the case of $l = 3$ there are the following twelve possibilities

$$\{(x^{\sigma(1)}x^{\sigma(2)})x^{\sigma(3)} : \sigma \in S_3\} \cup \{x^{\sigma(1)}(x^{\sigma(2)}x^{\sigma(3)}) : \sigma \in S_3\}.$$

Further $w(x^1, x^2, \dots, x^l)$ stands for such a monomial and we associate with $w(x^1, x^2, \dots, x^l)$ the following multi-linear polynomial

$$w_{n,d}(x^1, x^2, \dots, x^l) = \sum_{\sigma \in S_n} sgn(\sigma) w(x^{\sigma(1)}, x^{\sigma(2)}, \dots, x^{\sigma(n)}, x^{n+1}, x^{n+2}, \dots, x^l).$$

Let \mathbb{A} be any m -dimensional algebra over \mathbb{F} . We use the notations $[\mathbf{u}, \mathbf{v}] = \mathbf{u} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{u}$ and $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = (\mathbf{u} \cdot \mathbf{v}) \cdot \mathbf{w} - \mathbf{u} \cdot (\mathbf{v} \cdot \mathbf{w})$ for the commutator and the associator of \mathbf{u}, \mathbf{v} and $\mathbf{u}, \mathbf{v}, \mathbf{w}$, respectively. It is assumed that a basis $\mathbf{e} = \{e_j\}_{j=1,2,3,\dots,m}$ of \mathbb{A} over \mathbb{F} is fixed and $\mathbf{u}^i = \sum_{j=1}^m x_j^i e_j \in \mathbb{A}$, where $i = 1, 2, 3, \dots, l$, stand for any elements of \mathbb{A} . Define

$$|\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^m| = \begin{vmatrix} x_1^1 & x_2^1 & \dots & x_m^1 \\ x_1^2 & x_2^2 & \dots & x_m^2 \\ \vdots & \vdots & \dots & \vdots \\ x_1^m & x_2^m & \dots & x_m^m \end{vmatrix}.$$

Theorem 3.1.

- If $m = n = l$ then there exists such an element $\mathbf{u}_0 \in \mathbb{A}$ that for any $\mathbf{u}^i = \sum_{j=1}^m x_j^i e_j \in \mathbb{A}$, where $i = 1, 2, 3, \dots, m$, the equality

$$(3.1) \quad w_{m,d}(\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^m) = |\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^m| \mathbf{u}_0$$

- holds true;
- if $n > m$ then $w_d(\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^l) = 0$.

Proof. It is clear that

$$w(\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^l) = \sum_{\substack{i_j = 1, j=1, \dots, l, \\ s=1}}^m x_{i_1}^1 x_{i_2}^2 \dots x_{i_l}^l c_s^{(i_1, i_2, \dots, i_l)} e_s,$$

where $c_s^{(i_1, i_2, \dots, i_l)} \in \mathbb{F}$ and therefore,

$$\begin{aligned} w_{n,d}(\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^l) &= \sum_{\substack{i_j = 1, j=1,\dots,l, \\ s=1}}^m \sum_{\sigma \in S_n} sgn(\sigma) x_{i_1}^{\sigma(1)} x_{i_2}^{\sigma(2)} \dots x_{i_n}^{\sigma(n)} x_{i_{n+1}}^{n+1} x_{i_{n+2}}^{n+2} \dots x_{i_l}^l c_s^{(i_1, i_2, \dots, i_l)} e_s \\ &= \sum_{\substack{i_j = 1, j=1,\dots,l, \\ s=1}}^m \begin{vmatrix} x_{i_1}^1 & x_{i_2}^1 & \dots & x_{i_n}^1 \\ x_{i_1}^2 & x_{i_2}^2 & \dots & x_{i_n}^2 \\ \vdots & \vdots & \dots & \vdots \\ x_{i_1}^n & x_{i_2}^n & \dots & x_{i_n}^n \end{vmatrix} x_{i_{n+1}}^{n+1} x_{i_{n+2}}^{n+2} \dots x_{i_l}^l c_s^{(i_1, i_2, \dots, i_l)} e_s. \end{aligned}$$

The determinant in the expression above is zero whenever either two of the numbers i_1, i_2, \dots, i_n are the same and hence, in this case we have $w_{n,d}(\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^l) = 0$ whenever $n > m$, whereas if $m = n = l$ one has

$$w_{m,d}(\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^m) = |\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^m| \mathbf{u}_0, \text{ where } \mathbf{u}_0 = \sum_{s=1}^m \sum_{\sigma \in S_m} c_s^{(\sigma(1), \sigma(2), \dots, \sigma(m))} e_s.$$

□

Corollary 3.2. *For any m -dimensional algebra \mathbb{A} the following identity*

$$[w_{m,d}(\mathbf{u}^1, \mathbf{u}^2, \dots, \mathbf{u}^m), w_{m,d}(\mathbf{v}^1, \mathbf{v}^2, \dots, \mathbf{v}^m)] = 0$$

holds true.

Further we deal only with two-dimensional algebras. Due to the theorem above in any two-dimensional algebra \mathbb{A} the following identities hold true

$$(3.2) \quad [[\mathbf{u}, \mathbf{v}], [\mathbf{u}', \mathbf{v}']] = 0, \quad [\mathbf{u}, \mathbf{v}]\mathbf{w} + [\mathbf{v}, \mathbf{w}]\mathbf{u} + [\mathbf{w}, \mathbf{u}]\mathbf{v} = 0, \quad \mathbf{w}[\mathbf{u}, \mathbf{v}] + \mathbf{u}[\mathbf{v}, \mathbf{w}] + \mathbf{v}[\mathbf{w}, \mathbf{u}] = 0$$

for all $\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{u}', \mathbf{v}' \in \mathbb{A}$. In particular, for any such algebra (\mathbb{A}, \cdot) the corresponding $(\mathbb{A}, [\cdot, \cdot])$ is a Lie algebra. Note that any three-linear identity in a two-dimensional algebra is a linear combination of the identities (3.2).

Let us now observe a few simple identities for some classes of two-dimensional algebras presented above in terms of their MSC (note that these identities also hold true for the corresponding isomorphic algebras).

In the cases of A_4 , A_5 , $A_8(\alpha_1)$ and A_9 one has

$$\begin{aligned} [\mathbf{u}, \mathbf{v}] &= |\mathbf{u}, \mathbf{v}|(\beta_2 + \alpha_1 - 1)e_2, \\ [\mathbf{u}, \mathbf{v}] &= (3\alpha_1 - 2)|\mathbf{u}, \mathbf{v}|e_2, \\ [\mathbf{u}, \mathbf{v}] &= (x_1y_2 - x_2y_1)e_2 \end{aligned}$$

and

$$[\mathbf{u}, \mathbf{v}] = (x_1y_2 - x_2y_1)e_2,$$

respectively.

Due to $e_2^2 = 0$ these imply the identities

$$[\mathbf{u}, \mathbf{v}][\mathbf{u}', \mathbf{v}'] = 0 \text{ and } [\mathbf{u}, \mathbf{v}, \mathbf{w}][\mathbf{u}', \mathbf{v}', \mathbf{w}'] = 0.$$

Moreover, for A_9 the identity

$$2[\mathbf{u}, \mathbf{v}]\mathbf{w} + \mathbf{w}[\mathbf{u}, \mathbf{v}] = 0$$

also holds true. Indeed,

$$\begin{aligned} \mathbf{u}\mathbf{v} &= (x_1e_1 + x_2e_2)(y_1e_1 + y_2e_2) = \frac{x_1y_1}{3}e_1 + \frac{3x_1y_1 + 2x_1y_2 - x_2y_1}{3}e_2, \\ [\mathbf{u}, \mathbf{v}] &= (x_1y_2 - x_2y_1)e_2, \\ [\mathbf{u}, \mathbf{v}]\mathbf{w} &= -\frac{z_1}{3}(x_1y_2 - x_2y_1)e_2, \\ \mathbf{w}[\mathbf{u}, \mathbf{v}] &= \frac{2z_1}{3}(x_1y_2 - x_2y_1)e_2. \end{aligned}$$

Therefore, we have

$$2[\mathbf{u}, \mathbf{v}]\mathbf{w} + \mathbf{w}[\mathbf{u}, \mathbf{v}] = 0.$$

For algebras A_{10} , A_{11} and A_{12} one has $\mathbf{uv} = \mathbf{vu}$. Moreover, in the case of A_{10} we have

$$\begin{aligned}\mathbf{uv} &= (x_1e_1 + x_2e_2)(y_1e_1 + y_2e_2) = (x_1y_2 + x_2y_1)e_1 - x_2y_2e_2, \\ (\mathbf{uv})\mathbf{w} &= ((x_1y_2 + x_2y_1)z_2 - x_2y_2z_1)e_1 + x_2y_2z_2e_2, \\ \mathbf{u}(\mathbf{vw}) &= (-x_1y_2z_2 + x_2(y_1z_2 + y_2z_1))e_1 + x_2y_2z_2e_2, \\ [\mathbf{u}, \mathbf{v}, \mathbf{w}] &= (\mathbf{uv})\mathbf{w} - \mathbf{u}(\mathbf{vw}) = 2y_2(x_1z_2 - x_2z_1)e_1\end{aligned}$$

and hence, the identities

$$[\mathbf{u}, \mathbf{v}, \mathbf{w}][\mathbf{u}', \mathbf{v}', \mathbf{w}'] = 0, \quad [\mathbf{u}, \mathbf{v}, \mathbf{w}] + [\mathbf{v}, \mathbf{w}, \mathbf{u}] - [\mathbf{w}, \mathbf{u}, \mathbf{v}] = 0$$

hold true.

In A_{11} we have $\mathbf{uv} = (x_1e_1 + x_2e_2)(y_1e_1 + y_2e_2) = (x_1y_2 + x_2y_1)e_1 + (x_1y_1 - x_2y_2)e_2$,

$$\begin{aligned}(\mathbf{uv})\mathbf{w} &= ((x_1y_2 + x_2y_1)z_2 + (x_1y_1 - x_2y_2)z_1)e_1 + ((x_1y_2 + x_2y_1)z_1 - (x_1y_1 - x_2y_2)z_2)e_2, \\ \mathbf{u}(\mathbf{vw}) &= (x_1(y_1z_1 - y_2z_2)z_2 + x_2(y_1z_2 + y_2z_1))e_1 + (x_1(y_1z_2 + y_2z_1) - x_2(y_1z_1 - y_2z_2))e_2, \\ [\mathbf{u}, \mathbf{v}, \mathbf{w}] &= (\mathbf{uv})\mathbf{w} - \mathbf{u}(\mathbf{vw}) = 2(x_1z_2 - x_2z_1)(y_2e_1 - y_1e_2)\end{aligned}$$

and therefore,

$$[\mathbf{u}, \mathbf{v}, \mathbf{w}] = -[\mathbf{w}, \mathbf{v}, \mathbf{u}], \quad [\mathbf{u}, \mathbf{v}, \mathbf{w}] + [\mathbf{v}, \mathbf{w}, \mathbf{u}] + [\mathbf{w}, \mathbf{u}, \mathbf{v}] = 0$$

hold true for the algebra A_{11} .

In the case of A_{12} the identities $(\mathbf{uv})\mathbf{w} = \mathbf{u}(\mathbf{vw}) = 0$ are valid.

4. CLASSIFICATION OF TWO-DIMENSIONAL ALGEBRAS WITH RESPECT TO IDENTITIES

Recall that in the opposite algebra \mathbb{A}^{op} to an algebra (\mathbb{A}, \cdot) the product $*$ is defined by $\mathbf{u} * \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$. Observe that an algebra (\mathbb{A}, \cdot) satisfies a polynomial identity if and only if its “opposite” $(\mathbb{A}^{op}, *)$ satisfies the corresponding “opposite” polynomial identity, i.e., the polynomial identities of an “opposite” algebra can be derived from the polynomial identities of the original algebra.

For an algebra \mathbb{A} with MSC A the MSC A^{op} of \mathbb{A}^{op} can be easily found. In two-dimensional case it is evident that MSC of \mathbb{A}^{op} with respect to the basis e is

$$A^{op} = \begin{pmatrix} \alpha_1 & \alpha_3 & \alpha_2 & \alpha_4 \\ \beta_1 & \beta_3 & \beta_2 & \beta_4 \end{pmatrix}, \text{ provided that } A = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \beta_1 & \beta_2 & \beta_3 & \beta_4 \end{pmatrix}$$

is MSC of \mathbb{A} with respect to e .

The main motivation to raise the question on the opposite algebras is the fact that the list of canonical representatives presented in Section 2 is not invariant with respect to the “opposite” operation and therefore the study of their behavior with respect to this operation is important, moreover, later on we use the result in the classification of two-dimensional algebras with respect to identities. Here are the results on “opposite”s of two-dimensional algebras.

Theorem 4.1. *Let $\text{Char}(\mathbb{F}) \neq 2, 3$. The following hold true:*

- $(A_1(\alpha_1, \alpha_2, \alpha_4, \beta_1))^{op} \cong A_1(-\alpha_2, -\alpha_1, \beta_1, \alpha_4)$,
- $(A_2(\alpha_1, \beta_1, \beta_2))^{op} \cong A_2\left(\frac{\alpha_1}{\alpha_1+\beta_2}, \frac{\beta_1}{(\alpha_1+\beta_2)\sqrt{\alpha_1+\beta_2}}, \frac{1-\alpha_1}{\alpha_1+\beta_2}\right)$, whenever $\alpha_1 + \beta_2 \neq 0$,
 $(A_2(\alpha_1, \beta_1, -\alpha_1))^{op} = A_6(\alpha_1, \beta_1)$,
- $(A_3(\beta_1, \beta_2))^{op} \cong A_3\left(\frac{\beta_1}{\beta_2^2}, \frac{1}{\beta_2}\right)$, whenever $\beta_2 \neq 0$,
 $(A_3(\beta_1, 0))^{op} = A_7(\beta_1)$,
- $(A_4(\alpha_1, \beta_2))^{op} \cong A_4\left(\frac{\alpha_1}{\alpha_1+\beta_2}, \frac{1-\alpha_1}{\alpha_1+\beta_2}\right)$, whenever $\alpha_1 + \beta_2 \neq 0$,
 $(A_4(\alpha_1, -\alpha_1))^{op} = A_8(\alpha_1)$,
- $(A_5(\alpha_1))^{op} \cong A_5\left(\frac{\alpha_1}{3\alpha_1-1}\right)$, whenever $\alpha_1 \neq \frac{1}{3}$,
 $(A_5\left(\frac{1}{3}\right))^{op} = A_9$,
- $(A_6(\alpha_1, \beta_1))^{op} = A_2(\alpha_1, \beta_1, -\alpha_1)$,

- $(A_7(\beta_1))^{op} = A_3(\beta_1, 0)$,
- $(A_8(\alpha_1))^{op} = A_4(\alpha_1, -\alpha_1)$,
- $(A_9)^{op} = A_5 \left(\frac{1}{3} \right)$,
- $(A_{10})^{op} = A_{10}$,
- $(A_{11})^{op} = A_{11}$,
- $(A_{12})^{op} = A_{12}$.

Proof. We provide non-trivial base change that brings an algebra to its opposite unless they are equal.

Indeed, $g(A_1(\alpha_1, \alpha_2, \alpha_4, \beta_1))^{op}(g^{-1} \otimes g^{-1}) = \begin{pmatrix} -\alpha_2 & -\alpha_1 & 1-\alpha_1 & \beta_1 \\ \alpha_4 & \alpha_2 & \alpha_2+1 & \alpha_1 \end{pmatrix} = A_1(-\alpha_2, -\alpha_1, \beta_1, \alpha_4)$, where
 $g = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

- If $\alpha_1 + \beta_2 \neq 0$ then $g(A_2(\alpha_1, \beta_1, \beta_2))^{op}(g^{-1} \otimes g^{-1}) = A_2 \left(\frac{\alpha_1}{\alpha_1+\beta_2}, \frac{\beta_1}{(\alpha_1+\beta_2)\sqrt{\alpha_1+\beta_2}}, \frac{1-\alpha_1}{\alpha_1+\beta_2} \right)$, where $g = \begin{pmatrix} \alpha_1 + \beta_2 & 0 \\ 0 & \sqrt{\alpha_1 + \beta_2} \end{pmatrix}$;
- If $\alpha_1 + \beta_2 = 0$ then $(A_2(\alpha_1, \beta_1, -\alpha_1))^{op} = A_6(\alpha_1, \beta_1)$.
- If $\beta_2 \neq 0$ then

$$g(A_3(\alpha_1, \beta_1, \beta_2))^{op}(g^{-1} \otimes g^{-1}) = \begin{pmatrix} 0 & 1 & 1 & 0 \\ \frac{\beta_1}{\beta_2^2} & \frac{1}{\beta_2} & 1 & -1 \end{pmatrix} = A_3 \left(\frac{\beta_1}{\beta_2^2}, \frac{1}{\beta_2} \right),$$

where $g = \begin{pmatrix} \beta_2 & 0 \\ 0 & 1 \end{pmatrix}$;

- $(A_3(\beta_1, 0))^{op} = A_7(\beta_1)$.
- If $\alpha_1 + \beta_2 \neq 0$ then

$$g(A_4(\alpha_1, \beta_2))^{op}(g^{-1} \otimes g^{-1}) = \begin{pmatrix} \frac{\alpha_1}{\alpha_1+\beta_2} & 0 & 0 & 0 \\ 0 & \frac{1-\alpha_1}{\alpha_1+\beta_2} & 1 - \frac{\alpha_1}{\alpha_1+\beta_2} & 0 \end{pmatrix} = A_4 \left(\frac{\alpha_1}{\alpha_1+\beta_2}, \frac{1-\alpha_1}{\alpha_1+\beta_2} \right),$$

where $g = \begin{pmatrix} \alpha_1 + \beta_2 & 0 \\ 0 & 1 \end{pmatrix}$;

- $(A_4(\alpha_1, -\alpha_1))^{op} = A_8(\alpha_1)$.
- If $3\alpha_1 - 1 \neq 0$ and $g = \begin{pmatrix} 3\alpha_1 - 1 & 0 \\ 0 & (3\alpha_1 - 1)^2 \end{pmatrix}$ then

$$g(A_5(\alpha_1))^{op}(g^{-1} \otimes g^{-1}) = \begin{pmatrix} \frac{\alpha_1}{3\alpha_1+1} & 0 & 0 & 0 \\ 1 & 2\frac{\alpha_1}{3\alpha_1-1} - 1 & 1 - \frac{\alpha_1}{3\alpha_1+1} & 0 \end{pmatrix} = A_5 \left(\frac{\alpha_1}{3\alpha_1-1} \right);$$

- $(A_5 \left(\frac{1}{3} \right))^{op} = A_9$.

Now due to the equality $(A^{op})^{op} = A$ the above obtained isomorphisms and equalities imply that

$$(A_6(\alpha_1, \beta_1))^{op} = A_2(\alpha_1, \beta_1, -\alpha_1), \quad (A_7(\beta_1))^{op} = A_3(\beta_1, 0),$$

$$(A_8(\alpha_1))^{op} = A_4(\alpha_1, -\alpha_1), \quad (A_9)^{op} = A_5 \left(\frac{1}{3} \right).$$

The equalities

$$(A_{10})^{op} = A_{10}, \quad (A_{11})^{op} = A_{11}, \quad (A_{12})^{op} = A_{12}$$

are evident. \square

Corollary 4.2. Let $\text{Char}(\mathbb{F}) \neq 2, 3$. Then, up to isomorphism, there exist only the following nontrivial two-dimensional algebras \mathbb{A} with $\mathbb{A}^{op} \cong \mathbb{A}$:

$$\begin{aligned} A_1(\alpha_1, -\alpha_1, \alpha_2, \alpha_2), \quad & A_2(\alpha_1, \beta_1, 1 - \alpha_1), \quad A_2(0, 0, -1), \quad A_3(\beta_1, -1), \quad A_3(\beta_1, 1), \quad A_4(\alpha_1, 1 - \alpha_1), \\ & A_4(0, -1), \quad A_5\left(\frac{2}{3}\right), \quad A_5(0), \quad A_{10}, \quad A_{11}, \quad A_{12}. \end{aligned}$$

Theorem 4.3. Let $\text{Char}(\mathbb{F}) = 2$. Then the following isomorphisms and equalities hold true

- $(A_{1,2}(\alpha_1, \alpha_2, \alpha_4, \beta_1))^{op} \cong A_{1,2}(\alpha_2, \alpha_1, \beta_1, \alpha_4)$,
- $(A_{2,2}(\alpha_1, \beta_1, \beta_2))^{op} \cong A_{2,2}\left(\frac{\alpha_1}{\alpha_1 + \beta_2}, \frac{\beta_1}{(\alpha_1 + \beta_2)\sqrt{\alpha_1 + \beta_2}}, \frac{1 + \alpha_1}{\alpha_1 + \beta_2}\right)$, whenever $\alpha_1 + \beta_2 \neq 0$,
 $(A_{2,2}(\alpha_1, \beta_1, \alpha_1))^{op} = A_{6,2}(\alpha_1, \beta_1)$,
- $(A_{3,2}(\alpha_1, \beta_2))^{op} \cong A_{3,2}\left(\frac{\alpha_1}{\alpha_1 + \beta_2}, \frac{1 + \alpha_1}{\alpha_1 + \beta_2}\right)$, whenever $\alpha_1 + \beta_2 \neq 0$,
 $(A_{3,2}(\alpha_1, \alpha_1))^{op} = A_{7,2}(\alpha_1)$,
- $(A_{4,2}(\alpha_1, \beta_2))^{op} \cong A_{4,2}\left(\frac{\alpha_1}{\alpha_1 + \beta_2}, \frac{1 + \alpha_1}{\alpha_1 + \beta_2}\right)$, whenever $\alpha_1 + \beta_2 \neq 0$,
 $(A_{4,2}(\alpha_1, \alpha_1))^{op} = A_{8,2}(\alpha_1)$,
- $(A_{5,2}(\alpha_1))^{op} \cong A_{5,2}\left(\frac{\alpha_1}{\alpha_1 + 1}\right)$, whenever $\alpha_1 \neq 1$,
 $(A_{5,2}(1))^{op} = A_{9,2}$,
- $(A_{6,2}(\alpha_1, \beta_1))^{op} = A_{2,2}(\alpha_1, \beta_1, \alpha_1)$,
- $(A_{7,2}(\alpha_1))^{op} = A_{3,2}(\alpha_1, \alpha_1)$,
- $(A_{8,2}(\alpha_1))^{op} = A_{4,2}(\alpha_1, \alpha_1)$,
- $(A_{9,2})^{op} = A_{5,2}(1)$,
- $(A_{10,2})^{op} = A_{10,2}$,
- $(A_{11,2})^{op} = A_{11,2}$, $(A_{12,2})^{op} = A_{12,2}$.

Proof. The proof is similar to that of previous theorem, for example,

$$\begin{aligned} g(A_{3,2}(\alpha_1, \beta_2))^{op}(g^{-1})^{\otimes 2} &= A_{3,2}\left(\frac{\alpha_1}{\alpha_1 + \beta_2}, \frac{1 + \alpha_1}{\alpha_1 + \beta_2}\right), \\ \text{where } g &= \begin{pmatrix} \alpha_1 + \beta_2 & 0 \\ 0 & 1 \end{pmatrix} \text{ and } \alpha_1 + \beta_2 \neq 0. \end{aligned} \quad \square$$

Corollary 4.4. Let $\text{Char}(\mathbb{F}) = 2$. Then, up to isomorphism, there exist only the following nontrivial two-dimensional algebras with $\mathbb{A}^{op} \cong \mathbb{A}$:

$$A_{1,2}(\alpha_1, \alpha_1, \alpha_2, \alpha_2), \quad A_{2,2}(\alpha_1, \beta_1, 1 + \alpha_1), \quad A_{3,2}(\alpha_1, 1 + \alpha_1), \quad A_{4,2}(\alpha_1, 1 + \alpha_1), \quad A_{5,2}(0), \quad A_{10,2}, \quad A_{11,2}, \quad A_{12,2}.$$

Theorem 4.5. Let $\text{Char}(\mathbb{F}) = 3$. The following isomorphisms and equalities are true

- $(A_{1,3}(\alpha_1, \alpha_2, \alpha_4, \beta_1))^{op} \cong A_{1,3}(-\alpha_2, -\alpha_1, \beta_1, \alpha_4)$,
- $(A_{2,3}(\alpha_1, \beta_1, \beta_2))^{op} \cong A_{2,3}\left(\frac{\alpha_1}{\alpha_1 + \beta_2}, \frac{\beta_1}{(\alpha_1 + \beta_2)\sqrt{\alpha_1 + \beta_2}}, \frac{1 - \alpha_1}{\alpha_1 + \beta_2}\right)$, whenever $\alpha_1 + \beta_2 \neq 0$,
 $(A_{2,3}(\alpha_1, \beta_1, -\alpha_1))^{op} = A_{6,3}(\alpha_1, \beta_1)$,
- $(A_{3,3}(\beta_1, \beta_2))^{op} \cong A_{3,3}\left(\frac{\beta_1}{\beta_2^2}, \frac{1}{\beta_2}\right)$, whenever $\beta_2 \neq 0$,
 $(A_{3,3}(\beta_1, 0))^{op} = A_{7,3}(\beta_1)$,
- $(A_{4,3}(\alpha_1, \beta_2))^{op} \cong A_{4,3}\left(\frac{\alpha_1}{\alpha_1 + \beta_2}, \frac{1 - \alpha_1}{\alpha_1 + \beta_2}\right)$, whenever $\alpha_1 + \beta_2 \neq 0$,
 $(A_{4,3}(\alpha_1, -\alpha_1))^{op} = A_{8,3}(\alpha_1)$,
- $(A_{5,3}(\alpha_1))^{op} \cong A_{5,3}(-\alpha_1)$,
- $(A_{6,3}(\alpha_1, \beta_1))^{op} = A_{2,3}(\alpha_1, \beta_1, -\alpha_1)$,
- $(A_{7,3}(\beta_1))^{op} = A_{3,3}(\beta_1, 0)$,
- $(A_{8,3}(\alpha_1))^{op} = A_{4,3}(\alpha_1, -\alpha_1)$,
- $(A_{9,3})^{op} = A_{9,3}$,
- $(A_{10,3})^{op} = A_{10,3}$,

- $(A_{11,3})^{op} = A_{11,3}$,
- $(A_{12,3})^{op} = A_{12,3}$.

Corollary 4.6. *In the case of $\text{Char}(\mathbb{F}) = 3$, up to isomorphism, there exist only the following nontrivial two-dimensional algebras with $\mathbb{A}^{op} \cong \mathbb{A}$:*

$$\begin{aligned} & A_{1,3}(\alpha_1, -\alpha_1, \alpha_2, \alpha_2), \quad A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1), \quad A_{2,3}(0, 0, -1), \quad A_{3,3}(\beta_1, -1), \quad A_{3,3}(\beta_1, 1), \\ & A_{4,3}(\alpha_1, 1 - \alpha_1), \quad A_{4,3}(0, -1), \quad A_{5,3}(0), \quad A_{9,3}, \quad A_{10,3}, \quad A_{11,3}, \quad A_{12,3}. \end{aligned}$$

Now we consider a set of the most important identities and classify all nontrivial two-dimensional algebras with respect to this set of identities. The set of identities mainly consists of identities which have appeared before in definitions of different classes of algebras and their “anti” variants. We tried not include both of an identity and its “opposite” versions into this set except for the case when an identity and its opposite are the same. Therefore, in calling identities the term “Left” is used. Of course, the use of terms “Left” and “Anti” seems to be a questionable, but we have tried to follow the tradition given in the literature earlier. From this classification one can easily derive the classification of two-dimensional algebras with a given subset of identities and due to Theorems 4.1, 4.3 and 4.5, the classification of two-dimensional algebras satisfying the corresponding “Right” (“Opposite”) identities also can easily be obtained.

In the theorems below we give the polynomial identities denoted by I_1, I_2, \dots, I_{30} followed by the classification of two-dimensional algebras satisfying these identities in terms of canonical representatives $A_1 - A_{12}$.

The following notations are used:

I stands for the second order identity matrix,

i stands for a fixed element of \mathbb{F} such that $i^2 = -1$.

Elements of \mathbb{A} and their position vectors are denoted by $\mathbf{u}, \mathbf{v}, \mathbf{w}$ and u, v, w , respectively.

Theorem 4.7. *Let $\text{Char}(\mathbb{F}) \neq 2, 3$. The following classification of two-dimensional algebras with respect to $I_1 - I_{30}$ holds true:*

I_1 . Commutativity identity $\mathbf{uv} = \mathbf{vu}$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_5(\frac{2}{3}), A_{10}, A_{11}, A_{12}.$$

I_2 . Anti-commutativity identity $\mathbf{uv} = -\mathbf{vu}$. $A_4(0, -1)$, A_{12} .

I_3 . Associativity identity $(\mathbf{uv})\mathbf{w} = \mathbf{u}(\mathbf{vw})$.

$$A_2(\frac{1}{2}, 0, \frac{1}{2}), A_4(\frac{1}{2}, \frac{1}{2}), A_4(1, 0), A_4(1, 1), A_4(\frac{1}{2}, 0), A_{12}.$$

I_4 . Anti-associativity identity $(\mathbf{uv})\mathbf{w} = -\mathbf{u}(\mathbf{vw})$. A_{12} .

I_5 . Well defined cube identity $\mathbf{u}^2\mathbf{u} = \mathbf{uu}^2$.

$$A_1(\frac{1}{3}, -\frac{1}{3}, 0, 0), A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1),$$

$$\text{where } \alpha_1 \neq \frac{2}{3}, A_4(\alpha_1, 2\alpha_1 - 1), A_5(\frac{2}{3}), A_8(\frac{1}{3}), A_{10}, A_{11}, A_{12}.$$

I_6 . Half-commutativity identity $[\mathbf{u}, \mathbf{v}]\mathbf{w} = \mathbf{w}[\mathbf{u}, \mathbf{v}]$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_5(\frac{2}{3}), A_{10}, A_{11}, A_{12}.$$

I_7 . Anti-half-commutativity identity $[\mathbf{u}, \mathbf{v}]\mathbf{w} = -\mathbf{w}[\mathbf{u}, \mathbf{v}]$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_4(\alpha_1, \alpha_1 - 1), A_5(0), A_5(\frac{2}{3}),$$

$$A_8(\frac{1}{2}), A_{10}, A_{11}, A_{12}.$$

I_8 . Mixed associativity identity $[\mathbf{u}, \mathbf{v}]\mathbf{w} = \mathbf{u}[\mathbf{v}, \mathbf{w}]$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_5(\frac{2}{3}), A_{10}, A_{11}, A_{12}.$$

I_9 . Anti-mixed-associativity identity $[\mathbf{u}, \mathbf{v}]\mathbf{w} = -\mathbf{u}[\mathbf{v}, \mathbf{w}]$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_5(\frac{2}{3}), A_{10}, A_{11}, A_{12}.$$

I_{10} . Flexibility identity $\mathbf{u}(\mathbf{vu}) = (\mathbf{uv})\mathbf{u}$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_4(\alpha_1, 2\alpha_1 - 1),$$

$$\text{where } \alpha_1 \neq \frac{2}{3}, A_5(\frac{2}{3}), A_8(\frac{1}{3}), A_{10}, A_{11}, A_{12}.$$

I_{11} . Anti-flexibility identity $\mathbf{u}(\mathbf{vu}) = -(\mathbf{uv})\mathbf{u}$. A_{12} .

I_{12} . Mixed flexibility identity $\mathbf{u}[\mathbf{v}, \mathbf{u}] = [\mathbf{u}, \mathbf{v}]\mathbf{u}$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_4(\alpha_1, \alpha_1 - 1), \\ A_5(0), A_5(\frac{2}{3}), A_8(\frac{1}{2}), A_{10}, A_{11}, A_{12}.$$

I₁₃. Mixed anti-flexibility identity $\mathbf{u}[\mathbf{v}, \mathbf{u}] = -[\mathbf{u}, \mathbf{v}]\mathbf{u}$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_5(\frac{2}{3}), A_{10}, A_{11}, A_{12}.$$

I₁₄. Left Leibniz identity $\mathbf{u}(\mathbf{vw}) = (\mathbf{uv})\mathbf{w} + \mathbf{v}(\mathbf{uw})$. $A_4(0, -1), A_8(0), A_{12}$.

I₁₅. Left anti-Leibniz identity $\mathbf{u}(\mathbf{vw}) = -(\mathbf{uv})\mathbf{w} - \mathbf{v}(\mathbf{uw})$. A_{12} .

I₁₆. Mixed left Leibniz identity $\mathbf{u}[\mathbf{v}, \mathbf{w}] = [\mathbf{u}, \mathbf{v}]\mathbf{w} + \mathbf{v}[\mathbf{u}, \mathbf{w}]$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_4(\alpha_1, \alpha_1 - 1), \\ A_5(0), A_5(\frac{2}{3}), A_8(\frac{1}{2}), A_{10}, A_{11}, A_{12}.$$

I₁₇. Mixed anti-left Leibniz identity $\mathbf{u}[\mathbf{v}, \mathbf{w}] = -[\mathbf{u}, \mathbf{v}]\mathbf{w} - \mathbf{v}[\mathbf{u}, \mathbf{w}]$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_5(\frac{2}{3}), A_{10}, A_{11}, A_{12}.$$

I₁₈. Left Poisson identity $(\mathbf{uv})\mathbf{w} + (\mathbf{vw})\mathbf{u} + (\mathbf{wu})\mathbf{v} = 0$. $A_4(0, -1), A_8(0), A_{12}$.

I₁₉. Left Jordan identity $(\mathbf{uv})\mathbf{u}^2 = \mathbf{u}(\mathbf{vu}^2)$.

$$\begin{aligned} & - \text{If } \text{Char}(\mathbb{F}) \neq 5 \text{ then } A_2(\frac{1}{2}, 0, \frac{1}{2}), A_2(\frac{1}{2}, 0, -\frac{1}{2}), A_4(\alpha_1, -1 + 2\alpha_1), \text{ where } \alpha_1 \neq \frac{1}{10}(5 \pm \sqrt{5}), \\ & \quad A_4(\alpha_1, \sqrt{\alpha_1 - \alpha_1^2}), A_4(\alpha_1, -\sqrt{\alpha_1 - \alpha_1^2}), \text{ where } \alpha_1 \neq 0, 1, A_5(\frac{1}{10}(5 - \sqrt{5})), \\ & \quad A_5(\frac{1}{10}(5 + \sqrt{5})), A_8(\frac{1}{3}), A_8(\frac{1}{2} - \frac{i}{2}), A_8(\frac{1}{2} + \frac{i}{2}), A_{12}. \\ & - \text{If } \text{Char}(\mathbb{F}) = 5 \text{ then } A_2(\frac{1}{2}, 0, \frac{1}{2}), A_2(\frac{1}{2}, 0, -\frac{1}{2}), A_4(\alpha_1, -1 + 2\alpha_1), A_4(\alpha_1, \sqrt{\alpha_1 - \alpha_1^2}), A_4(\alpha_1, -\sqrt{\alpha_1 - \alpha_1^2}), \\ & \quad \text{where } \alpha_1 \neq 0, 1, A_8(\frac{1}{3}), A_8(\frac{3}{2}), A_9, A_{12}. \end{aligned}$$

I₂₀. Left anti-Jordan identity $(\mathbf{uv})\mathbf{u}^2 = -\mathbf{u}(\mathbf{vu}^2)$. $A_4(0, -1), A_4(0, 0), A_{12}$.

I₂₁. Mixed left Jordan identity $[\mathbf{u}, \mathbf{v}]\mathbf{u}^2 = \mathbf{u}[\mathbf{v}, \mathbf{u}^2]$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(0, -1), A_4(0, 0), A_4(\alpha_1, 1 - \alpha_1), \\ A_5(\frac{2}{3}), A_{10}, A_{11}, A_{12}.$$

I₂₂. Mixed anti-left Jordan identity $[\mathbf{u}, \mathbf{v}]\mathbf{u}^2 = -\mathbf{u}[\mathbf{v}, \mathbf{u}^2]$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_2(\alpha_1, -\beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(0, -1), A_4(0, 0), A_4(\alpha_1, 1 - \alpha_1), \\ A_5(\frac{2}{3}), A_{10}, A_{11}, A_{12}.$$

I₂₃. Left Malcev identity $((\mathbf{uv})\mathbf{w} + (\mathbf{vw})\mathbf{u} + (\mathbf{wu})\mathbf{v})\mathbf{u} = (\mathbf{uv})(\mathbf{uw}) + (\mathbf{v}(\mathbf{uw}))\mathbf{u} + ((\mathbf{uw})\mathbf{u})\mathbf{v}$.

$$A_2(\frac{1}{2}, 0, \frac{1}{2}), A_4(0, -1), A_4(\frac{1}{2}, \frac{1}{2}), A_4(1, 0), A_8(0), A_{12}.$$

I₂₄. Left anti-Malcev identity $((\mathbf{uv})\mathbf{w} + (\mathbf{vw})\mathbf{u} + (\mathbf{wu})\mathbf{v})\mathbf{u} = -(\mathbf{uv})(\mathbf{uw}) - (\mathbf{v}(\mathbf{uw}))\mathbf{u} - ((\mathbf{uw})\mathbf{u})\mathbf{v}$.

$$A_4(0, -1), A_8(0), A_{12}.$$

I₂₅. Left Zinbiel identity $(\mathbf{uv})\mathbf{w} = \mathbf{u}(\mathbf{vw} + \mathbf{wv})$. A_{12} .

I₂₆. Left anti-Zinbiel identity $(\mathbf{uv})\mathbf{w} = -\mathbf{u}(\mathbf{vw} + \mathbf{wv})$. A_{12} .

I₂₇. Left symmetric identity $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = [\mathbf{v}, \mathbf{u}, \mathbf{w}]$.

$$A_2(\frac{1}{2}, 0, \frac{1}{2}), A_2(1, 0, \frac{1}{2}), A_4(1, \beta_2), A_4(\frac{1}{2}, \beta_2), A_5(1), A_5(\frac{1}{2}), A_8(0), A_{12}.$$

I₂₈. Left anti-symmetric identity $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = -[\mathbf{v}, \mathbf{u}, \mathbf{w}]$.

$$A_2(\frac{1}{2}, 0, \frac{1}{2}), A_4(\frac{1}{2}, \frac{1}{2}), A_4(\frac{1}{2}, 0), A_4(1, 0), A_4(1, 1), A_{12}.$$

I₂₉. Centro-symmetric identity $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = [\mathbf{w}, \mathbf{v}, \mathbf{u}]$.

$$A_1(\alpha_1, \alpha_2, -\alpha_1 - 2\alpha_2, 2\alpha_1 + \alpha_2), \text{ where } \alpha_2 = \frac{1}{8}(\pm\sqrt{32\alpha_1 - 15} - 8\alpha_1 - 1), A_2(\frac{1}{2}, 0, \frac{1}{2}),$$

$$A_4\left(\alpha_1, \frac{\alpha_1 - \sqrt{12\alpha_1 - 7\alpha_1^2 - 4}}{2}\right),$$

$$A_4\left(\alpha_1, \frac{\alpha_1 + \sqrt{12\alpha_1 - 7\alpha_1^2 - 4}}{2}\right), A_5(\frac{1}{2}), A_5(1), A_8\left(\frac{3 - \sqrt{7}i}{8}\right), A_8\left(\frac{3 + \sqrt{7}i}{8}\right), A_{12}.$$

I₃₀. Centro-anti-symmetric identity $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = -[\mathbf{w}, \mathbf{v}, \mathbf{u}]$.

$$A_2(\alpha_1, \beta_1, 1 - \alpha_1), A_3(\beta_1, 1), A_4(\alpha_1, 1 - \alpha_1), A_4(\alpha_1, 2\alpha_1 - 1), A_5(\frac{2}{3}), A_8(\frac{1}{3}), A_{10}, A_{11}, A_{12}.$$

Proof. Proofs of the cases $I_1 - I_{30}$ are similar. The basic idea behind the proof of the theorem is just to convert the given identity in terms of MSC A as a matrix equation and solve the corresponding system of polynomials equations for each of $A_1 - A_{12}$ canonical representatives. Here we provide the systems of equations. The solutions to them are a bit technical, but once we substitute the structure constants into the system it is much

simplified and can be managed combining manual and computer calculations. We illustrate the technique in two specific cases (the cases I_{18} and I_{23}).

The cases of commutativity and anticommutativity are clear where the identities are equivalent to the fact that the second and third columns of A to be the same and differ in sign as vectors, respectively. The proof of the cases I_5 and I_{19} are given in [4] and [3], respectively.

The identities I_3 and I_4 in terms of matrices are expressed by $A(A \otimes I) = A(I \otimes A)$ and $A(A \otimes I) = -A(I \otimes A)$, respectively.

- I_6 case. $A(A \otimes I)((u \otimes v - v \otimes u) \otimes w) = A(I \otimes A)(w \otimes (u \otimes v - v \otimes u))$.

The corresponding system is given as below

$$\begin{aligned} \alpha_2\beta_2 - \alpha_3\beta_2 - \alpha_2\beta_3 + \alpha_3\beta_3 &= 0, \\ \alpha_2^2 - 2\alpha_3\alpha_2 + \alpha_3^2 &= 0, \\ \beta_2^2 - 2\beta_3\beta_2 + \beta_3^2 &= 0. \end{aligned}$$

- I_7 case. $A(A \otimes I)((u \otimes v - v \otimes u) \otimes w) = -A(I \otimes A)(w \otimes (u \otimes v - v \otimes u))$.

In this case the system is as follows

$$\begin{aligned} 2\alpha_1\alpha_2 + \beta_2\alpha_2 - \beta_3\alpha_2 - 2\alpha_1\alpha_3 + \alpha_3\beta_2 - \alpha_3\beta_3 &= 0, \\ \alpha_2\beta_2 - \alpha_3\beta_2 + 2\beta_4\beta_2 + \alpha_2\beta_3 - \alpha_3\beta_3 - 2\beta_3\beta_4 &= 0, \\ \alpha_2^2 - \alpha_3^2 + 2\alpha_4\beta_2 - 2\alpha_4\beta_3 &= 0, \\ \beta_2^2 - \beta_3^2 + 2\alpha_2\beta_1 - 2\alpha_3\beta_1 &= 0. \end{aligned}$$

- I_8 case. $A(A \otimes I)((u \otimes v - v \otimes u) \otimes w) = A(I \otimes A)(u \otimes (v \otimes w - w \otimes v))$.

The identity is written in terms of structure constants as follows

$$\begin{aligned} 2\alpha_1\alpha_2 + \beta_2\alpha_2 - \beta_3\alpha_2 - 2\alpha_1\alpha_3 + \alpha_3\beta_2 - \alpha_3\beta_3 &= 0, \\ \alpha_2\beta_2 - \alpha_3\beta_2 + 2\beta_4\beta_2 + \alpha_2\beta_3 - \alpha_3\beta_3 - 2\beta_3\beta_4 &= 0, \\ \alpha_1\alpha_2 + \beta_2\alpha_2 - \beta_3\alpha_2 - \alpha_1\alpha_3 &= 0, \\ \alpha_2^2 - \alpha_3\alpha_2 + \alpha_4\beta_2 - \alpha_4\beta_3 &= 0, \\ \alpha_1\alpha_2 - \alpha_1\alpha_3 + \alpha_3\beta_2 - \alpha_3\beta_3 &= 0, \\ \alpha_2^2 - \alpha_3^2 + 2\alpha_4\beta_2 - 2\alpha_4\beta_3 &= 0, \\ \alpha_3^2 - \alpha_2\alpha_3 - \alpha_4\beta_2 + \alpha_4\beta_3 &= 0, \\ \beta_2^2 - \beta_3\beta_2 - \alpha_2\beta_1 - \alpha_3\beta_1 &= 0, \\ \beta_2^2 - \beta_3^2 + 2\alpha_2\beta_1 - 2\alpha_3\beta_1 &= 0, \\ \alpha_2\beta_2 - \alpha_3\beta_2 + \beta_4\beta_2 - \beta_3\beta_4 &= 0, \\ \beta_3^2 - \beta_2\beta_3 - \alpha_2\beta_1 + \alpha_3\beta_1 &= 0, \\ \alpha_2\beta_3 - \alpha_3\beta_3 - \beta_4\beta_3 + \beta_2\beta_4 &= 0. \end{aligned}$$

- I_9 case. $A(A \otimes I)((u \otimes v - v \otimes u) \otimes w) = -A(I \otimes A)(u \otimes (v \otimes w - w \otimes v))$.

Here is the corresponding system of equations

$$\begin{aligned} \alpha_1\alpha_2 + \beta_2\alpha_2 - \beta_3\alpha_2 - \alpha_1\alpha_3 &= 0, & \alpha_2\beta_2 - \alpha_3\beta_2 - \alpha_2\beta_3 + \alpha_3\beta_3 &= 0, \\ \alpha_2^2 - \alpha_3\alpha_2 + \alpha_4\beta_2 - \alpha_4\beta_3 &= 0, & \alpha_1\alpha_2 - \alpha_1\alpha_3 + \alpha_3\beta_2 - \alpha_3\beta_3 &= 0, \\ \beta_2^2 - \beta_3\beta_2 + \alpha_2\beta_1 - \alpha_3\beta_1 &= 0, & \alpha_3^2 - \alpha_2\alpha_3 - \alpha_4\beta_2 + \alpha_4\beta_3 &= 0, \\ \alpha_2\beta_2 - \alpha_3\beta_2 + \beta_4\beta_2 - \beta_3\beta_4 &= 0, & \beta_3^2 - \beta_2\beta_3 - \alpha_2\beta_1 + \alpha_3\beta_1 &= 0, \\ \alpha_2\beta_2 - \alpha_3\beta_2 - \alpha_2\beta_3 + \alpha_3\beta_3 &= 0, & \alpha_2\beta_3 - \alpha_3\beta_3 - \beta_4\beta_3 + \beta_2\beta_4 &= 0, \\ \alpha_2^2 - 2\alpha_3\alpha_2 + \alpha_3^2 &= 0, & \beta_2^2 + 2\beta_3\beta_2 + \beta_3^2 &= 0. \end{aligned}$$

- I_{10} case. $(A(I \otimes A) - A(A \otimes I))(u \otimes v \otimes u) = 0$.

The system of equations is

$$\begin{aligned} \alpha_3\beta_2 - \beta_4\beta_2 - \alpha_2\beta_3 + \beta_3\beta_4 &= 0, & \beta_2^2 - \alpha_1\beta_2 - \beta_3^2 + \alpha_2\beta_1 - \alpha_3\beta_1 + \alpha_1\beta_3 &= 0, \\ \alpha_2\beta_2 - \alpha_3\beta_3 &= 0, & \alpha_2^2 - \beta_4\alpha_2 - \alpha_3^2 + \alpha_4\beta_2 - \alpha_4\beta_3 + \alpha_3\beta_4 &= 0, \\ \alpha_4\beta_2 - \alpha_4\beta_3 &= 0, & \alpha_1\alpha_2 - \beta_3\alpha_2 - \alpha_1\alpha_3 + \alpha_3\beta_2 &= 0, \\ \beta_1\beta_2 - \beta_1\beta_3 &= 0, & \alpha_3\alpha_4 - \alpha_2\alpha_4 &= 0, \\ \alpha_2\beta_1 - \alpha_3\beta_1 &= 0. & & \end{aligned}$$

- I_{11} case. $(A(I \otimes A) + A(A \otimes I))(u \otimes v \otimes u) = 0$.

In this case the system of equations is given as follows

$$\begin{aligned} \alpha_2^2 + \beta_4\alpha_2 + \alpha_3^2 + 2\alpha_1\alpha_4 + \alpha_4\beta_2 + \alpha_4\beta_3 + \alpha_3\beta_4 &= 0, \\ \beta_2^2 + \alpha_1\beta_2 + \beta_3^2 + \alpha_2\beta_1 + \alpha_3\beta_1 + \alpha_1\beta_3 + 2\beta_1\beta_4 &= 0, \\ 2\alpha_1\alpha_2 + \beta_2\alpha_2 + 2\alpha_1\alpha_3 + 2\alpha_4\beta_1 + \alpha_3\beta_3 &= 0, \\ 2\alpha_4\beta_1 + \alpha_2\beta_2 + \alpha_3\beta_3 + 2\beta_2\beta_4 + 2\beta_3\beta_4 &= 0, \\ \alpha_1\alpha_2 + \beta_3\alpha_2 + \alpha_1\alpha_3 + \alpha_3\beta_2 &= 0, \\ \alpha_3\beta_2 + \beta_4\beta_2 + \alpha_2\beta_3 + \beta_3\beta_4 &= 0, \\ 2\alpha_2\alpha_3 + \alpha_4\beta_2 + \alpha_4\beta_3 &= 0, \\ \alpha_2\alpha_4 + \alpha_3\alpha_4 + 2\beta_4\alpha_4 &= 0, \\ 2\alpha_1\beta_1 + \beta_2\beta_1 + \beta_3\beta_1 &= 0, \\ \alpha_2\beta_1 + \alpha_3\beta_1 + 2\beta_2\beta_3 &= 0, \\ 2\alpha_1^2 + \alpha_2\beta_1 + \alpha_3\beta_1 &= 0, \\ 2\beta_4^2 + \alpha_4\beta_2 + \alpha_4\beta_3 &= 0. \end{aligned}$$

- I_{12} case. $A(A \otimes I)((u \otimes v - v \otimes u) \otimes u) = A(I \otimes A)(u \otimes (v \otimes u - u \otimes v))$.

The identity can be written in terms of structure constant as the following system of equations

$$\begin{aligned} 2\alpha_1\alpha_2 + \beta_2\alpha_2 - \beta_3\alpha_2 - 2\alpha_1\alpha_3 + \alpha_3\beta_2 - \alpha_3\beta_3 &= 0, \\ \alpha_2\beta_2 - \alpha_3\beta_2 + 2\beta_4\beta_2 + \alpha_2\beta_3 - \alpha_3\beta_3 - 2\beta_3\beta_4 &= 0, \\ \alpha_2^2 - \alpha_3^2 + 2\alpha_4\beta_2 - 2\alpha_4\beta_3 &= 0, \\ \beta_2^2 - \beta_3^2 + 2\alpha_2\beta_1 - 2\alpha_3\beta_1 &= 0, \end{aligned}$$

- I_{13} case. $A(A \otimes I)((u \otimes v - v \otimes u) \otimes u) = -A(I \otimes A)(u \otimes (v \otimes u - u \otimes v))$.

The system of equations is

$$\alpha_2\beta_2 - \alpha_3\beta_2 - \alpha_2\beta_3 + \alpha_3\beta_3 = 0, \quad \alpha_2^2 - 2\alpha_3\alpha_2 + \alpha_3^2 = 0, \quad \beta_2^2 - 2\beta_3\beta_2 + \beta_3^2 = 0.$$

- I_{14} case. $(A(I \otimes A) - A(A \otimes I))(u \otimes v \otimes w) = A(I \otimes A)(v \otimes u \otimes w)$.

Here is the corresponding system of equations

$$\begin{aligned} \beta_3^2 - \alpha_1\beta_3 + \beta_2\beta_3 + 2\alpha_3\beta_1 - \beta_1\beta_4 &= 0, \quad \alpha_1^2 + \alpha_3\beta_1 = 0, \\ \alpha_2^2 + \alpha_3\alpha_2 - \beta_4\alpha_2 - \alpha_1\alpha_4 + 2\alpha_4\beta_2 &= 0, \quad \alpha_4\beta_1 + \beta_3\beta_4 = 0, \\ \alpha_1\alpha_3 + \beta_3\alpha_3 - \alpha_4\beta_1 + \alpha_2\beta_3 &= 0, \quad \alpha_1\beta_1 + \beta_3\beta_1 = 0, \\ \alpha_1\alpha_2 - \beta_3\alpha_2 + \alpha_4\beta_1 + \alpha_3\beta_2 &= 0, \quad \alpha_2\alpha_4 + \beta_4\alpha_4 = 0, \\ \alpha_2\beta_1 - \alpha_3\beta_1 + \beta_4\beta_1 + \alpha_1\beta_3 &= 0, \quad \alpha_1\beta_2 + \beta_1\beta_4 = 0, \\ \alpha_4\beta_1 - \alpha_2\beta_2 - \alpha_2\beta_3 - \beta_2\beta_4 &= 0, \quad \alpha_1\alpha_2 + \alpha_4\beta_1 = 0, \\ \alpha_4\beta_1 + \alpha_3\beta_2 - \alpha_2\beta_3 + \beta_3\beta_4 &= 0, \quad \alpha_1\alpha_4 + \alpha_3\beta_4 = 0, \\ \alpha_1\alpha_4 - \beta_2\alpha_4 + \beta_3\alpha_4 + \alpha_2\beta_4 &= 0, \quad \beta_4^2 + \alpha_4\beta_2 = 0. \end{aligned}$$

- I_{15} case. $(A(I \otimes A) + A(A \otimes I))(u \otimes v \otimes w) = -A(I \otimes A)(v \otimes u \otimes w)$.

The system of equations is given below

$$\begin{aligned} \alpha_1\alpha_2 + \beta_3\alpha_2 + 2\alpha_1\alpha_3 + \alpha_4\beta_1 + \alpha_3\beta_2 &= 0, \quad 2\alpha_2\alpha_3 + \alpha_1\alpha_4 + \alpha_4\beta_2 + \alpha_4\beta_3 + \alpha_2\beta_4 = 0, \\ \alpha_2\beta_1 + \alpha_3\beta_1 + \beta_4\beta_1 + \alpha_1\beta_3 + 2\beta_2\beta_3 &= 0, \quad \alpha_2^2 + \alpha_3\alpha_2 + \beta_4\alpha_2 + \alpha_1\alpha_4 + 2\alpha_4\beta_2 = 0, \\ \beta_3^2 + \alpha_1\beta_3 + \beta_2\beta_3 + 2\alpha_3\beta_1 + \beta_1\beta_4 &= 0, \quad \alpha_4\beta_1 + \alpha_3\beta_2 + \alpha_2\beta_3 + 2\beta_2\beta_4 + \beta_3\beta_4 = 0, \\ 3\alpha_1\alpha_3 + \beta_3\alpha_3 + \alpha_4\beta_1 + \alpha_2\beta_3 &= 0, \quad \alpha_2\alpha_4 + 2\alpha_3\alpha_4 + 3\beta_4\alpha_4 = 0, \\ 2\alpha_3^2 + \beta_4\alpha_3 + \alpha_1\alpha_4 + 2\alpha_4\beta_3 &= 0, \quad 2\beta_2^2 + \alpha_1\beta_2 + 2\alpha_2\beta_1 + \beta_1\beta_4 = 0, \\ 3\alpha_1^2 + 2\alpha_2\beta_1 + \alpha_3\beta_1 &= 0, \quad \alpha_4\beta_1 + \alpha_2\beta_2 + \alpha_2\beta_3 + 3\beta_2\beta_4 = 0, \\ 3\alpha_1\beta_1 + 2\beta_2\beta_1 + \beta_3\beta_1 &= 0, \quad 3\alpha_1\alpha_2 + 2\beta_2\alpha_2 + \alpha_4\beta_1 = 0, \\ \alpha_4\beta_1 + 2\alpha_3\beta_3 + 3\beta_3\beta_4 &= 0, \quad 3\beta_4^2 + \alpha_4\beta_2 + 2\alpha_4\beta_3 = 0. \end{aligned}$$

- I_{16} case. $A(I \otimes A)(u \otimes (v \otimes w - w \otimes v)) = A(A \otimes I)((u \otimes v - v \otimes u) \otimes w) + A(I \otimes A)(v \otimes (u \otimes w - w \otimes u))$.

The system of equations is

$$\begin{aligned} 2\alpha_1\alpha_2 + \beta_2\alpha_2 - \beta_3\alpha_2 - 2\alpha_1\alpha_3 + \alpha_3\beta_2 - \alpha_3\beta_3 &= 0, \\ \alpha_2\beta_2 - \alpha_3\beta_2 + 2\beta_4\beta_2 + \alpha_2\beta_3 - \alpha_3\beta_3 - 2\beta_3\beta_4 &= 0, \\ \alpha_2^2 - \alpha_3^2 + 2\alpha_4\beta_2 - 2\alpha_4\beta_3 &= 0, \\ \beta_2^2 - \beta_3^2 + 2\alpha_2\beta_1 - 2\alpha_3\beta_1 &= 0. \end{aligned}$$

- I_{17} case. $A(I \otimes A)(u \otimes (v \otimes w - w \otimes v))$

$$= -A(A \otimes I)((u \otimes v - v \otimes u) \otimes w) - A(I \otimes A)(v \otimes (u \otimes w - w \otimes u)).$$

In this case we have the system of equations

$$\begin{aligned} \alpha_2\beta_2 - \alpha_3\beta_2 + 2\beta_4\beta_2 + \alpha_2\beta_3 - \alpha_3\beta_3 - 2\beta_3\beta_4 &= 0, & \alpha_2\beta_2 - \alpha_3\beta_2 - \alpha_2\beta_3 + \alpha_3\beta_3 &= 0, \\ 2\alpha_1\alpha_2 + \beta_2\alpha_2 - \beta_3\alpha_2 - 2\alpha_1\alpha_3 + \alpha_3\beta_2 - \alpha_3\beta_3 &= 0, & \alpha_2^2 - \alpha_3^2 + 2\alpha_4\beta_2 - 2\alpha_4\beta_3 &= 0, \\ \alpha_3^2 - \alpha_2\alpha_3 - \alpha_4\beta_2 + \alpha_4\beta_3 &= 0, & \beta_2^2 - \beta_3^2 + 2\alpha_2\beta_1 - 2\alpha_3\beta_1 &= 0, \\ \beta_2^2 - \beta_3\beta_2 + \alpha_2\beta_1 - \alpha_3\beta_1 &= 0, & \beta_2^2 - 2\beta_3\beta_2 + \beta_3^2 &= 0, \\ \alpha_1\alpha_2 + \beta_2\alpha_2 - \beta_3\alpha_2 - \alpha_1\alpha_3 &= 0, & \alpha_2^2 - 2\alpha_3\alpha_2 + \alpha_3^2 &= 0, \\ \alpha_2\beta_3 - \alpha_3\beta_3 - \beta_4\beta_3 + \beta_2\beta_4 &= 0. \end{aligned}$$

- I_{18} case. The left Poisson identity $(\mathbf{u}\mathbf{v})\mathbf{w} + (\mathbf{v}\mathbf{w})\mathbf{u} + (\mathbf{w}\mathbf{u})\mathbf{v} = 0$ is equivalent to $A(A \otimes I)(u \otimes v \otimes w + v \otimes w \otimes u + w \otimes u \otimes v) = 0$ that produces the system of equations

$$(4.1) \quad \begin{aligned} 2\alpha_1\alpha_2 + \alpha_1\alpha_3 + \alpha_4\beta_1 + \alpha_3\beta_2 + \alpha_3\beta_3 &= 0, \\ \alpha_2^2 + \alpha_3\alpha_2 + \alpha_1\alpha_4 + \alpha_4\beta_2 + \alpha_4\beta_3 + \alpha_3\beta_4 &= 0, \\ \beta_3^2 + \beta_2\beta_3 + \alpha_2\beta_1 + \alpha_3\beta_1 + \alpha_1\beta_2 + \beta_1\beta_4 &= 0, \\ \alpha_4\beta_1 + \alpha_2\beta_2 + \alpha_3\beta_2 + \beta_2\beta_4 + 2\beta_3\beta_4 &= 0, \\ 3\beta_4^2 + 3\alpha_4\beta_2 &= 0, \\ 3\alpha_1^2 + 3\alpha_3\beta_1 &= 0, \\ 3\alpha_2\alpha_4 + 3\beta_4\alpha_4 &= 0, \\ 3\alpha_1\beta_1 + 3\beta_3\beta_1 &= 0, \end{aligned}$$

For A_1 the system (4.1) becomes

$$\begin{aligned} 3\alpha_1^2 + 3\alpha_2\beta_1 + 3\beta_1 &= 0, \\ \alpha_2\alpha_1 - \alpha_1 + \alpha_2 + \alpha_4\beta_1 + 1 &= 0, \\ \alpha_1^2 - 3\alpha_1 + \alpha_2\beta_1 + \beta_1 + 1 &= 0, \\ \alpha_2\alpha_1 - \alpha_1 - 2\alpha_2 + \alpha_4\beta_1 &= 0, \\ 3\alpha_2^2 - 3\alpha_1\alpha_4 &= 0, \\ \alpha_2^2 - \alpha_1\alpha_4 + \alpha_4 &= 0, \\ 3\beta_1 &= 0. \end{aligned}$$

The last equation gives $\beta_1 = 0$ that produces $\alpha_1 = 0$ due to the first equation and we get a contradiction in the equation 3.

For A_2 the system (4.1) has the form

$$\begin{aligned} \alpha_1^2 - 2\alpha_1 + \beta_2 + 1 &= 0, \\ \beta_2 + 1 &= 0, \\ 3\beta_2 &= 0, \\ 3\alpha_1^2 &= 0, \\ \beta_1 &= 0. \end{aligned}$$

The equations 2 and 3 immediately lead to a contradiction.

For A_3 due to the system (4.1) we have

$$\begin{aligned} \beta_1 + \beta_2 + 1 &= 0, \\ \beta_2 + 1 &= 0, \\ \beta_2 - 2 &= 0, \\ 3\beta_1 &= 0, \\ 1 &= 0. \end{aligned}$$

which is a contradiction.

For A_4 the system (4.1) becomes

$$\begin{aligned} \alpha_1^2 - 2\alpha_1 + \beta_2 + 1 &= 0, \\ 3\alpha_1^2 &= 0. \end{aligned}$$

from where we get $\alpha_1 = 0$ and $\beta_2 = -1$.

For A_5 , A_6 and A_7 the system (4.1) gives

$$\begin{aligned} 3\alpha_1^2 &= 0, & 1 - \alpha_1 &= 0, \\ \alpha_1^2 &= 0, & \alpha_1^2 &= 0, \quad \text{and} \quad \beta_1 &= 0, \\ 3 &= 0. & \beta_1 &= 0. \end{aligned}$$

, respectively. Which are contradictions.

Substituting the structure constants A_8 into the system (4.1) we get

$$\alpha_1^2 = 0.$$

hence, $\alpha_1 = 0$.

For the structure constants of A_9 , A_{10} , A_{11} the system (4.1) gives contradictions.

The structure constants of A_{12} satisfy the system of equations (4.1).

- I_{19} case. $(A(A \otimes A) - A(I \otimes A(I \otimes A)))(u \otimes v \otimes u \otimes u) = 0$.

More detailed proof for this case can be seen in [3]. In this case the system of equations has the following form

$$\begin{aligned} &\alpha_4\alpha_1^2 + \alpha_2^2\alpha_1 + 2\alpha_3^2\alpha_1 + 2\alpha_2\alpha_3\alpha_1 + \alpha_4\beta_2\alpha_1 + 2\alpha_4\beta_3\alpha_1 + \alpha_3\beta_4\alpha_1 + \alpha_4\beta_2^2 \\ &+ \alpha_2\alpha_4\beta_1 + \alpha_3\alpha_4\beta_1 + \alpha_2^2\beta_2 + \alpha_3^2\beta_2 + \alpha_2\alpha_3\beta_2 + 2\alpha_2^2\beta_3 + \alpha_2\alpha_3\beta_3 + \alpha_4\beta_2\beta_3 \\ &\quad + 2\alpha_4\beta_1\beta_4 + \alpha_2\beta_2\beta_4 + \alpha_2\beta_3\beta_4 = 0, \\ &\beta_1\alpha_3^2 + \beta_2^2\alpha_3 + \beta_3^2\alpha_3 + \alpha_2\beta_1\alpha_3 + \alpha_1\beta_3\alpha_3 + \beta_2\beta_3\alpha_3 + \beta_1\beta_4\alpha_3 + 2\alpha_2\beta_3^2 \\ &\quad + \beta_1\beta_4^2 + 2\alpha_1\alpha_4\beta_1 + \alpha_4\beta_1\beta_2 + \alpha_1\alpha_2\beta_3 + \alpha_4\beta_1\beta_3 + \alpha_2\beta_2\beta_3 + 2\beta_2^2\beta_4 \\ &\quad + \beta_3^2\beta_4 + 2\alpha_2\beta_1\beta_4 + \alpha_1\beta_2\beta_4 + 2\beta_2\beta_3\beta_4 = 0, \\ &\beta_2^3 + \alpha_1\beta_2^2 + \beta_3\beta_2^2 + 2\alpha_2\beta_1\beta_2 + 2\alpha_3\beta_1\beta_2 + \alpha_1\beta_3\beta_2 + 2\beta_1\beta_4\beta_2 + \alpha_1\beta_3^2 \\ &\quad + 2\alpha_1\alpha_2\beta_1 + 3\alpha_1\alpha_3\beta_1 + \alpha_1^2\beta_3 + 3\alpha_2\beta_1\beta_3 + \alpha_3\beta_1\beta_3 + \alpha_1\beta_1\beta_4 + 2\beta_1\beta_3\beta_4 = 0, \\ &\alpha_3^3 + \alpha_2\alpha_3^2 + \beta_4\alpha_3^2 + 2\alpha_1\alpha_4\alpha_3 + 2\alpha_4\beta_2\alpha_3 + 2\alpha_4\beta_3\alpha_3 + \alpha_2\beta_4\alpha_3 + \alpha_2\beta_4^2 \\ &\quad + 2\alpha_1\alpha_2\alpha_4 + \alpha_2\alpha_4\beta_2 + 3\alpha_2\alpha_4\beta_3 + \alpha_2^2\beta_4 + \alpha_1\alpha_4\beta_4 + 3\alpha_4\beta_2\beta_4 + 2\alpha_4\beta_3\beta_4 = 0, \\ &\beta_1\alpha_2^2 + \beta_2^2\alpha_2 + 2\alpha_3\beta_1\alpha_2 + 3\beta_2\beta_3\alpha_2 + 2\beta_1\beta_2^2 + \alpha_2^2\beta_1 + \alpha_1\alpha_4\beta_1 + 2\alpha_4\beta_1\beta_2 \\ &\quad + \alpha_1\alpha_3\beta_3 + 2\alpha_4\beta_1\beta_3 + 2\alpha_3\beta_2\beta_3 + 2\beta_2^2\beta_4 + 2\alpha_1\beta_3\beta_4 + 2\beta_2\beta_3\beta_4 = 0, \\ &2\alpha_4\alpha_1^2 + 2\alpha_3^2\alpha_1 + 2\alpha_2\alpha_3\alpha_1 + 2\alpha_2\beta_4\alpha_1 + \alpha_4\beta_2^2 + \alpha_4\beta_3^2 + 2\alpha_2\alpha_4\beta_1 + 2\alpha_3\alpha_4\beta_1 \\ &\quad + 2\alpha_2\alpha_3\beta_2 + \alpha_3^2\beta_3 + 3\alpha_2\alpha_3\beta_3 + 2\alpha_4\beta_2\beta_3 + \alpha_4\beta_1\beta_4 + \alpha_2\beta_2\beta_4 = 0, \\ &\beta_3\alpha_3^2 + 2\alpha_4\beta_1\alpha_3 + \alpha_2\beta_3\alpha_3 + 2\beta_3\beta_4\alpha_3 + \alpha_4\beta_2^2 + \alpha_4\beta_3^2 + 4\beta_2\beta_4^2 + 2\beta_3\beta_4^2 \\ &\quad + 2\alpha_2\alpha_4\beta_1 + 4\alpha_4\beta_2\beta_3 + \alpha_4\beta_1\beta_4 + \alpha_2\beta_2\beta_4 + 2\alpha_2\beta_3\beta_4 = 0, \\ &2\alpha_2\alpha_1^2 + 4\alpha_3\alpha_1^2 + \alpha_4\beta_1\alpha_1 + 2\alpha_2\beta_2\alpha_1 + 2\alpha_2\beta_3\alpha_1 + \alpha_3\beta_3\alpha_1 + \alpha_2\beta_2^2 + \alpha_2^2\beta_1 \\ &\quad + \alpha_3^2\beta_1 + 4\alpha_2\alpha_3\beta_1 + 2\alpha_4\beta_1\beta_2 + 2\alpha_4\beta_1\beta_3 + \alpha_2\beta_2\beta_3 = 0, \\ &\alpha_4\beta_3^2 + \beta_4^2\beta_3 + \alpha_1\alpha_4\beta_3 + \alpha_2\beta_4\beta_3 + \beta_2\beta_4^2 + \alpha_3\alpha_4\beta_1 + \alpha_4\beta_1\beta_4 + \alpha_3\beta_2\beta_4 = 0, \\ &\alpha_2\alpha_1^2 + \alpha_3\alpha_1^2 + \alpha_4\beta_1\alpha_1 + \alpha_3\beta_2\alpha_1 + \alpha_2\beta_3\alpha_1 + \alpha_2^2\beta_1 + \alpha_4\beta_1\beta_2 + \alpha_2\beta_1\beta_4 = 0, \\ &\beta_1\alpha_4^2 + 2\alpha_1\alpha_3\alpha_4 + \alpha_3\beta_3\alpha_4 + \beta_2\beta_4\alpha_4 + \beta_3\beta_4\alpha_4 + 2\alpha_2\alpha_3\beta_4 = 0, \\ &\alpha_4\beta_1^2 + \alpha_1\alpha_2\beta_1 + \alpha_1\alpha_3\beta_1 + \alpha_2\beta_2\beta_1 + 2\beta_2\beta_4\beta_1 + 2\alpha_1\beta_2\beta_3 = 0, \\ &2\beta_1\alpha_1^2 + 2\beta_1\beta_2\alpha_1 + \beta_1\beta_3\alpha_1 + \alpha_2\beta_1^2 + \beta_1\beta_2^2 + \beta_1^2\beta_4 = 0, \\ &\alpha_4\alpha_3^2 + 2\alpha_4\beta_4\alpha_3 + \alpha_1\alpha_4^2 + 2\alpha_4\beta_4^2 + \alpha_4^2\beta_3 + \alpha_2\alpha_4\beta_4 = 0, \\ &2\beta_4^3 + \alpha_4\beta_2\beta_4 + 3\alpha_4\beta_3\beta_4 + \alpha_4^2\beta_1 + \alpha_3\alpha_4\beta_3 = 0, \\ &2\alpha_1^3 + 3\alpha_2\beta_1\alpha_1 + \alpha_3\beta_1\alpha_1 + \alpha_4\beta_1^2 + \alpha_2\beta_1\beta_2 = 0. \end{aligned}$$

- I_{20} case. $(A(A \otimes A) + A(I \otimes A(I \otimes A)))(u \otimes v \otimes u \otimes u) = 0$.

Here is the system of equations

$$\begin{aligned}
& \alpha_4\alpha_1^2 + \alpha_2^2\alpha_1 + 2\alpha_3^2\alpha_1 + 2\alpha_2\alpha_3\alpha_1 + \alpha_4\beta_2\alpha_1 + 2\alpha_4\beta_3\alpha_1 + \alpha_3\beta_4\alpha_1 + \alpha_4\beta_2^2 \\
& + \alpha_2\alpha_4\beta_1 + \alpha_3\alpha_4\beta_1 + \alpha_2^2\beta_2 + \alpha_3^2\beta_2 + \alpha_2\alpha_3\beta_2 + 2\alpha_2^2\beta_3 + \alpha_2\alpha_3\beta_3 + \alpha_4\beta_2\beta_3 \\
& \quad + 2\alpha_4\beta_1\beta_4 + \alpha_2\beta_2\beta_4 + \alpha_2\beta_3\beta_4 = 0, \\
& \beta_1\alpha_3^2 + \beta_2^2\alpha_3 + \beta_3^2\alpha_3 + \alpha_2\beta_1\alpha_3 + \alpha_1\beta_3\alpha_3 + \beta_2\beta_3\alpha_3 + \beta_1\beta_4\alpha_3 + 2\alpha_2\beta_3^2 \\
& + \beta_1\beta_4^2 + 2\alpha_1\alpha_4\beta_1 + \alpha_4\beta_1\beta_2 + \alpha_1\alpha_2\beta_3 + \alpha_4\beta_1\beta_3 + \alpha_2\beta_2\beta_3 + 2\beta_2^2\beta_4 \\
& \quad + \beta_3^2\beta_4 + 2\alpha_2\beta_1\beta_4 + \alpha_1\beta_2\beta_4 + 2\beta_2\beta_3\beta_4 = 0, \\
& \alpha_3^3 + \alpha_2\alpha_3^2 + \beta_4\alpha_3^2 + 2\alpha_1\alpha_4\alpha_3 + 2\alpha_4\beta_2\alpha_3 + 2\alpha_4\beta_3\alpha_3 + \alpha_2\beta_4\alpha_3 + \alpha_2\beta_4^2 \\
& + 2\alpha_1\alpha_2\alpha_4 + \alpha_2\alpha_4\beta_2 + 3\alpha_2\alpha_4\beta_3 + \alpha_2^2\beta_4 + \alpha_1\alpha_4\beta_4 + 3\alpha_4\beta_2\beta_4 + 2\alpha_4\beta_3\beta_4 = 0, \\
& \beta_2^3 + \alpha_1\beta_2^2 + \beta_3\beta_2^2 + 2\alpha_2\beta_1\beta_2 + 2\alpha_3\beta_1\beta_2 + \alpha_1\beta_3\beta_2 + 2\beta_1\beta_4\beta_2 + \alpha_1\beta_3^2 \\
& + 2\alpha_1\alpha_2\beta_1 + 3\alpha_1\alpha_3\beta_1 + \alpha_1^2\beta_3 + 3\alpha_2\beta_1\beta_3 + \alpha_3\beta_1\beta_3 + \alpha_1\beta_1\beta_4 + 2\beta_1\beta_3\beta_4 = 0, \\
& 2\alpha_4\alpha_1^2 + 2\alpha_3^2\alpha_1 + 2\alpha_2\alpha_3\alpha_1 + 2\alpha_2\beta_4\alpha_1 + \alpha_4\beta_2^2 + \alpha_4\beta_3^2 + 2\alpha_2\alpha_4\beta_1 + 2\alpha_3\alpha_4\beta_1 \\
& \quad + 2\alpha_2\alpha_3\beta_2 + \alpha_3^2\beta_3 + 3\alpha_2\alpha_3\beta_3 + 2\alpha_4\beta_2\beta_3 + \alpha_4\beta_1\beta_4 + \alpha_2\beta_2\beta_4 = 0, \\
& \beta_1\alpha_2^2 + \beta_2^2\alpha_2 + 2\alpha_3\beta_1\alpha_2 + 3\beta_2\beta_3\alpha_2 + 2\beta_1\beta_4^2 + \alpha_3^2\beta_1 + \alpha_1\alpha_4\beta_1 + 2\alpha_4\beta_1\beta_2 \\
& \quad + \alpha_1\alpha_3\beta_3 + 2\alpha_4\beta_1\beta_3 + 2\alpha_3\beta_2\beta_3 + 2\beta_2^2\beta_4 + 2\alpha_1\beta_3\beta_4 + 2\beta_2\beta_3\beta_4 = 0, \\
& \beta_3\alpha_3^2 + 2\alpha_4\beta_1\alpha_3 + \alpha_2\beta_3\alpha_3 + 2\beta_3\beta_4\alpha_3 + \alpha_4\beta_2^2 + \alpha_4\beta_3^2 + 4\beta_2\beta_4^2 + 2\beta_3\beta_4^2 \\
& \quad + 2\alpha_2\alpha_4\beta_1 + 4\alpha_4\beta_2\beta_3 + \alpha_4\beta_1\beta_4 + \alpha_2\beta_2\beta_4 + 2\alpha_2\beta_3\beta_4 = 0, \\
& 2\alpha_2\alpha_1^2 + 4\alpha_3\alpha_1^2 + \alpha_4\beta_1\alpha_1 + 2\alpha_2\beta_2\alpha_1 + 2\alpha_2\beta_3\alpha_1 + \alpha_3\beta_3\alpha_1 + \alpha_2\beta_2^2 + \alpha_2^2\beta_1 \\
& \quad + \alpha_3^2\beta_1 + 4\alpha_2\alpha_3\beta_1 + 2\alpha_4\beta_1\beta_2 + 2\alpha_4\beta_1\beta_3 + \alpha_2\beta_2\beta_3 = 0, \\
& \alpha_2\alpha_1^2 + \alpha_3\alpha_1^2 + \alpha_4\beta_1\alpha_1 + \alpha_3\beta_2\alpha_1 + \alpha_2\beta_3\alpha_1 + \alpha_2^2\beta_1 + \alpha_4\beta_1\beta_2 + \alpha_2\beta_1\beta_4 = 0, \\
& \alpha_4\beta_3^2 + \beta_4^2\beta_3 + \alpha_1\alpha_4\beta_3 + \alpha_2\beta_4\beta_3 + \beta_2\beta_4^2 + \alpha_3\alpha_4\beta_1 + \alpha_4\beta_1\beta_4 + \alpha_3\beta_2\beta_4 = 0, \\
& \beta_1\alpha_4^2 + 2\alpha_1\alpha_3\alpha_4 + \alpha_3\beta_3\alpha_4 + \beta_2\beta_4\alpha_4 + \beta_3\beta_4\alpha_4 + 2\alpha_2\alpha_3\beta_4 = 0, \\
& \alpha_4\alpha_3^2 + 2\alpha_4\beta_4\alpha_3 + \alpha_1\alpha_4^2 + 2\alpha_4\beta_4^2 + \alpha_4^2\beta_3 + \alpha_2\alpha_4\beta_4 = 0, \\
& 2\beta_1\alpha_1^2 + 2\beta_1\beta_2\alpha_1 + \beta_1\beta_3\alpha_1 + \alpha_2\beta_1^2 + \beta_1\beta_2^2 + \beta_1^2\beta_4 = 0, \\
& \alpha_4\beta_1^2 + \alpha_1\alpha_2\beta_1 + \alpha_1\alpha_3\beta_1 + \alpha_2\beta_2\beta_1 + 2\beta_2\beta_4\beta_1 + 2\alpha_1\beta_2\beta_3 = 0, \\
& 2\alpha_1^3 + 3\alpha_2\beta_1\alpha_1 + \alpha_3\beta_1\alpha_1 + \alpha_4\beta_1^2 + \alpha_2\beta_1\beta_2 = 0, \\
& 2\beta_4^3 + \alpha_4\beta_2\beta_4 + 3\alpha_4\beta_3\beta_4 + \alpha_4^2\beta_1 + \alpha_3\alpha_4\beta_3 = 0.
\end{aligned}$$

- I_{21} case. $A(A \otimes A)((u \otimes v - v \otimes u) \otimes u^{\otimes 2}) = A(I \otimes A(I \otimes A))(u \otimes v \otimes u^{\otimes 2}) - A(I \otimes A(A \otimes I))(u^{\otimes 3} \otimes v)$.

The system of equations is given as follows

$$\begin{aligned}
& \alpha_1\alpha_2^2 + \beta_2\alpha_2^2 + \beta_3\alpha_2^2 + \alpha_3\beta_2\alpha_2 - \alpha_3\beta_3\alpha_2 + \alpha_1\beta_4\alpha_2 + \beta_2\beta_4\alpha_2 - \beta_3\beta_4\alpha_2 - \alpha_1\alpha_3^2 \\
& \quad + 2\alpha_4\beta_2^2 - 2\alpha_4\beta_3^2 - 2\alpha_3^2\beta_3 - \alpha_1\alpha_3\beta_4 = 0, \\
& 2\beta_1\alpha_2^2 + 2\beta_2^2\alpha_2 - \beta_3^2\alpha_2 + \alpha_1\beta_3\alpha_2 + \beta_2\beta_3\alpha_2 - \alpha_3\beta_3^2 - 2\alpha_3^2\beta_1 - \alpha_1\alpha_3\beta_3 - \alpha_3\beta_2\beta_3 \\
& \quad + \beta_2^2\beta_4 - \beta_3^2\beta_4 + \alpha_1\beta_2\beta_4 - \alpha_1\beta_3\beta_4 = 0, \\
& 2\alpha_1\alpha_2^2 + 2\beta_2\alpha_2^2 + \alpha_1\alpha_3\alpha_2 + \alpha_3\beta_2\alpha_2 - 3\alpha_3\beta_3\alpha_2 - 3\alpha_1\alpha_3^2 + \alpha_4\beta_2^2 - \alpha_4\beta_3^2 + \alpha_3^2\beta_2 \\
& \quad + \alpha_1\alpha_4\beta_2 - \alpha_3^2\beta_3 - \alpha_1\alpha_4\beta_3 = 0, \\
& \alpha_3^3 - \alpha_2^2\alpha_3 + 2\alpha_1\alpha_4\alpha_3 - 2\alpha_4\beta_2\alpha_3 + 2\alpha_4\beta_3\alpha_3 + \alpha_2\beta_4\alpha_3 - 2\alpha_1\alpha_2\alpha_4 - 2\alpha_2\alpha_4\beta_2 \\
& \quad + 2\alpha_2\alpha_4\beta_3 - \alpha_2^2\beta_4 - \alpha_4\beta_2\beta_4 + \alpha_4\beta_3\beta_4 = 0, \\
& \beta_2^3 - \beta_3^2\beta_2 + 2\alpha_2\beta_1\beta_2 - 2\alpha_3\beta_1\beta_2 + \alpha_1\beta_3\beta_2 + 2\beta_1\beta_4\beta_2 - \alpha_1\beta_3^2 + \alpha_1\alpha_2\beta_1 \\
& \quad - \alpha_1\alpha_3\beta_1 + 2\alpha_2\beta_1\beta_3 - 2\alpha_3\beta_1\beta_3 - 2\beta_1\beta_3\beta_4 = 0, \\
& \beta_2^2\alpha_2 + 3\beta_2\beta_3\alpha_2 + \beta_1\beta_4\alpha_2 - \alpha_3\beta_2^2 - 2\alpha_3\beta_3^2 - \alpha_3^2\beta_1 + \alpha_2^2\beta_1 - \alpha_3\beta_2\beta_3 \\
& \quad + 3\beta_2^2\beta_4 - 2\beta_3^2\beta_4 - \alpha_3\beta_1\beta_4 - \beta_2\beta_3\beta_4 = 0, \\
& \alpha_3\alpha_1^2 - \alpha_2\beta_2\alpha_1 - \alpha_2\beta_3\alpha_1 + 2\alpha_3\beta_3\alpha_1 - \alpha_2\beta_2^2 + \alpha_2\beta_3^2 - \alpha_1^2\alpha_2 - \alpha_2^2\beta_1 \\
& \quad + \alpha_3^2\beta_1 - 2\alpha_4\beta_1\beta_2 + 2\alpha_4\beta_1\beta_3 = 0, \\
& \beta_3\alpha_2^2 + 2\alpha_4\beta_1\alpha_2 + 2\beta_2\beta_4\alpha_2 - \beta_3\beta_4\alpha_2 + \alpha_4\beta_2^2 - \alpha_4\beta_3^2 + \beta_2\beta_4^2 - \beta_3\beta_4^2 \\
& \quad - 2\alpha_3\alpha_4\beta_1 - \alpha_3^2\beta_3 - \alpha_3\beta_3\beta_4 = 0, \\
& 2\alpha_2\alpha_1^2 - 2\alpha_3\alpha_1^2 + \alpha_2\beta_2\alpha_1 + \alpha_3\beta_2\alpha_1 - \alpha_2\beta_3\alpha_1 - \alpha_3\beta_3\alpha_1 + \alpha_2^2\beta_1 - \alpha_2\alpha_3\beta_1 \\
& \quad + \alpha_4\beta_1\beta_2 - \alpha_4\beta_1\beta_3 = 0, \\
& \alpha_4\beta_3^2 + 2\beta_4^2\beta_3 - \alpha_4\beta_2\beta_3 - \alpha_2\beta_4\beta_3 + \alpha_3\beta_4\beta_3 - 2\beta_2\beta_4^2 - \alpha_2\alpha_4\beta_1 + \alpha_3\alpha_4\beta_1 \\
& \quad - \alpha_2\beta_2\beta_4 + \alpha_3\beta_2\beta_4 = 0, \\
& \beta_4\alpha_2^2 + \alpha_1\alpha_4\alpha_2 - \alpha_1\alpha_3\alpha_4 + \alpha_3\alpha_4\beta_2 - \alpha_3\alpha_4\beta_3 - \alpha_3^2\beta_4 + 2\alpha_4\beta_2\beta_4 - 2\alpha_4\beta_3\beta_4 = 0, \\
& \alpha_1\beta_2^2 + \alpha_2\beta_1\beta_2 - \alpha_3\beta_1\beta_2 + \beta_1\beta_4\beta_2 - \alpha_1\beta_3^2 + 2\alpha_1\alpha_2\beta_1 - 2\alpha_1\alpha_3\beta_1 - \beta_1\beta_3\beta_4 = 0, \\
& \quad \alpha_2\alpha_4\beta_3 - \alpha_3\alpha_4\beta_3 - \alpha_4\beta_4\beta_3 + \alpha_4\beta_2\beta_4 = 0. \\
& \alpha_1\alpha_2\beta_1 - \alpha_1\alpha_3\beta_1 + \alpha_2\beta_2\beta_1 - \alpha_2\beta_3\beta_1 = 0, \\
& \alpha_4\alpha_3^2 - \alpha_2\alpha_4\alpha_3 - \alpha_4^2\beta_2 + \alpha_4^2\beta_3 = 0, \\
& \alpha_2\beta_1^2 - \alpha_3\beta_1^2 + \beta_2^2\beta_1 - \beta_2\beta_3\beta_1 = 0.
\end{aligned}$$

- I_{22} case. $A(A \otimes A)((u \otimes v - v \otimes u) \otimes u^{\otimes 2})$

$$= -A(I \otimes A(I \otimes A))(u \otimes v \otimes u^{\otimes 2}) + A(I \otimes A(A \otimes I))(u^{\otimes 3} \otimes v).$$

In this case the system of equations is

$$\begin{aligned}
 & \alpha_1\alpha_2^2 + \beta_2\alpha_2^2 + \beta_3\alpha_2^2 - \alpha_3\beta_2\alpha_2 - 3\alpha_3\beta_3\alpha_2 - \alpha_1\beta_4\alpha_2 - \beta_2\beta_4\alpha_2 + \beta_3\beta_4\alpha_2 - \alpha_1\alpha_3^2 \\
 & \quad + 2\alpha_3^2\beta_2 + \alpha_1\alpha_3\beta_4 = 0, \\
 & 2\alpha_3\beta_2^2 - \beta_4\beta_2^2 - 3\alpha_2\beta_3\beta_2 - \alpha_3\beta_3\beta_2 + \alpha_1\beta_4\beta_2 + \alpha_2\beta_3^2 + \alpha_3\beta_3^2 + \alpha_1\alpha_2\beta_3 - \alpha_1\alpha_3\beta_3 \\
 & \quad + \beta_3^2\beta_4 - \alpha_1\beta_3\beta_4 = 0, \\
 & \beta_2^2\alpha_2 - 2\beta_3^2\alpha_2 + \beta_2\beta_3\alpha_2 - \beta_1\beta_4\alpha_2 - \alpha_3\beta_2^2 - \alpha_3^2\beta_1 + \alpha_2^2\beta_1 + \alpha_3\beta_2\beta_3 - \beta_2^2\beta_4 \\
 & \quad + \alpha_3\beta_1\beta_4 + \beta_2\beta_3\beta_4 = 0, \\
 & 2\beta_3\alpha_2^2 - \alpha_1\alpha_3\alpha_2 - \alpha_3\beta_2\alpha_2 - \alpha_3\beta_3\alpha_2 + \alpha_1\alpha_3^2 + \alpha_4\beta_2^2 - \alpha_4\beta_3^2 + \alpha_3^2\beta_2 - \alpha_1\alpha_4\beta_2 \\
 & \quad - \alpha_3^2\beta_3 + \alpha_1\alpha_4\beta_3 = 0, \\
 & \beta_3\alpha_2^2 - \beta_3\beta_4\alpha_2 + \alpha_4\beta_2^2 + \alpha_4\beta_3^2 - \beta_2\beta_4^2 + \beta_3\beta_4^2 - \alpha_3^2\beta_3 - 2\alpha_4\beta_2\beta_3 \\
 & \quad + 2\alpha_3\beta_2\beta_4 - \alpha_3\beta_3\beta_4 = 0, \\
 & \alpha_3\alpha_1^2 + \alpha_2\beta_2\alpha_1 - 2\alpha_3\beta_2\alpha_1 + \alpha_2\beta_3\alpha_1 + \alpha_2\beta_2^2 - \alpha_2\beta_3^2 - \alpha_1^2\alpha_2 - \alpha_2^2\beta_1 \\
 & \quad - \alpha_3^2\beta_1 + 2\alpha_2\alpha_3\beta_1 = 0, \\
 & \alpha_4\beta_3^2 - \alpha_4\beta_2\beta_3 + \alpha_2\beta_4\beta_3 - \alpha_3\beta_4\beta_3 - \alpha_2\alpha_4\beta_1 + \alpha_3\alpha_4\beta_1 - \alpha_2\beta_2\beta_4 + \alpha_3\beta_2\beta_4 = 0, \\
 & \beta_1\alpha_2^2 - \alpha_3\beta_1\alpha_2 - \alpha_1\beta_2\alpha_2 + \alpha_1\beta_3\alpha_2 + \alpha_1\alpha_3\beta_2 + \alpha_4\beta_1\beta_2 - \alpha_1\alpha_3\beta_3 - \alpha_4\beta_1\beta_3 = 0, \\
 & \alpha_3^3 - \alpha_2^2\alpha_3 - \alpha_2\beta_4\alpha_3 - 2\alpha_2\alpha_4\beta_2 + 2\alpha_2\alpha_4\beta_3 + \alpha_2^2\beta_4 + \alpha_4\beta_2\beta_4 - \alpha_4\beta_3\beta_4 = 0, \\
 & \beta_2^3 - \beta_3^2\beta_2 - \alpha_1\beta_3\beta_2 + \alpha_1\beta_3^2 - \alpha_1\alpha_2\beta_1 + \alpha_1\alpha_3\beta_1 + 2\alpha_2\beta_1\beta_3 - 2\alpha_3\beta_1\beta_3 = 0, \\
 & \beta_4\alpha_2^2 + \alpha_1\alpha_4\alpha_2 - 2\alpha_3\beta_4\alpha_2 - \alpha_1\alpha_3\alpha_4 + \alpha_3\alpha_4\beta_2 - \alpha_3\alpha_4\beta_3 + \alpha_3^2\beta_4 = 0, \\
 & \alpha_1\beta_2^2 - \alpha_2\beta_1\beta_2 + \alpha_3\beta_1\beta_2 - 2\alpha_1\beta_3\beta_2 - \beta_1\beta_4\beta_2 + \alpha_1\beta_3^2 + \beta_1\beta_3\beta_4 = 0, \\
 & \alpha_1\alpha_2\beta_1 - \alpha_1\alpha_3\beta_1 + \alpha_2\beta_2\beta_1 - \alpha_2\beta_3\beta_1 = 0, \\
 & \alpha_2\alpha_4\beta_3 - \alpha_3\alpha_4\beta_3 - \alpha_4\beta_4\beta_3 + \alpha_4\beta_2\beta_4 = 0, \\
 & \alpha_2\beta_1^2 - \alpha_3\beta_1^2 + \beta_2^2\beta_1 - \beta_2\beta_3\beta_1 = 0, \\
 & \alpha_4\alpha_3^2 - \alpha_2\alpha_4\alpha_3 - \alpha_4^2\beta_2 + \alpha_4^2\beta_3 = 0,
 \end{aligned}$$

- I_{23} case. $A(A(A \otimes I) \otimes I)((u \otimes v \otimes w + v \otimes w \otimes u + w \otimes u \otimes v) \otimes u)$
 $= A(A \otimes A)(u \otimes v \otimes u \otimes w) + A(A(I \otimes A) \otimes I)(v \otimes u \otimes w \otimes u) + A(A(A \otimes I) \otimes I)(u \otimes w \otimes u \otimes v)$. The identity is equivalent to the following system of equations

$$\begin{aligned}
 & \beta_1\alpha_2^2 + \alpha_1\beta_2\alpha_2 + \beta_1\beta_4\alpha_2 + \alpha_3\beta_2^2 - \alpha_3\beta_3^2 - 2\beta_1\beta_4^2 - \alpha_3^2\beta_1 - \alpha_1\alpha_3\beta_2 - \alpha_4\beta_1\beta_2 \\
 & \quad + \alpha_4\beta_1\beta_3 + \beta_3^2\beta_4 + \alpha_3\beta_1\beta_4 - 2\alpha_1\beta_3\beta_4 + \beta_2\beta_3\beta_4 = 0, \\
 & 2\alpha_4\alpha_1^2 - \alpha_2^2\alpha_1 - \alpha_2\alpha_3\alpha_1 - \alpha_4\beta_2\alpha_1 - \alpha_4\beta_3\alpha_1 + 2\alpha_2\beta_4\alpha_1 + \alpha_4\beta_2^2 - \alpha_4\beta_3^2 \\
 & \quad - \alpha_2\alpha_4\beta_1 + \alpha_3\alpha_4\beta_1 + \alpha_2^2\beta_2 - \alpha_3^2\beta_2 + \alpha_3\beta_2\beta_4 - \alpha_3\beta_3\beta_4 = 0, \\
 & \alpha_2\beta_2^2 + \beta_4\beta_2^2 - \alpha_1\alpha_2\beta_2 - \alpha_1\alpha_3\beta_2 - \alpha_4\beta_1\beta_2 - 2\alpha_3\beta_3\beta_2 + \beta_3\beta_4\beta_2 + \alpha_2\beta_3^2 + 2\alpha_1\alpha_4\beta_1 \\
 & \quad + \alpha_4\beta_1\beta_3 - 2\beta_3^2\beta_4 + \alpha_2\beta_1\beta_4 - \alpha_3\beta_1\beta_4 = 0, \\
 & \beta_1\alpha_3^2 - \beta_2^2\alpha_3 + \beta_3^2\alpha_3 + \alpha_2\beta_1\alpha_3 + \beta_2\beta_3\alpha_3 - \beta_1\beta_4^2 - 2\alpha_1\alpha_2\beta_2 - \alpha_4\beta_1\beta_2 + \alpha_4\beta_1\beta_3 \\
 & \quad + \alpha_2\beta_2\beta_3 + \beta_3^2\beta_4 - \alpha_2\beta_1\beta_4 - \alpha_1\beta_2\beta_4 = 0, \\
 & 2\alpha_1\alpha_2^2 - \beta_3\alpha_2^2 - \alpha_1\alpha_3\alpha_2 - \alpha_4\beta_1\alpha_2 + 2\alpha_3\beta_2\alpha_2 - \alpha_1\alpha_3^2 + \alpha_3\alpha_4\beta_1 + \alpha_1\alpha_4\beta_2 - \alpha_3^2\beta_3 \\
 & \quad - \alpha_1\alpha_4\beta_3 - 2\alpha_4\beta_1\beta_4 + \alpha_3\beta_2\beta_4 + \alpha_3\beta_3\beta_4 = 0, \\
 & \alpha_4\alpha_1^2 + \alpha_4\beta_3\alpha_1 + \alpha_3\beta_4\alpha_1 - \alpha_1\alpha_2^2 - \alpha_4\beta_2^2 - \alpha_2\alpha_4\beta_1 + \alpha_3\alpha_4\beta_1 - \alpha_2^2\beta_2 + \alpha_3^2\beta_2 \\
 & \quad - \alpha_2\alpha_3\beta_2 - \alpha_2\alpha_3\beta_3 - \alpha_4\beta_2\beta_3 + 2\alpha_3\beta_3\beta_4 = 0, \\
 & \alpha_3\alpha_1^2 + \alpha_4\beta_1\alpha_1 - 2\alpha_2\beta_2\alpha_1 + \alpha_3\beta_2\alpha_1 + \alpha_3\beta_3\alpha_1 - \alpha_3\beta_2^2 + \alpha_3\beta_3^2 - \alpha_1^2\alpha_2 + \alpha_3^2\beta_1 \\
 & \quad - 2\alpha_2\alpha_3\beta_1 - \alpha_4\beta_1\beta_2 + \alpha_3\beta_1\beta_4 = 0,
 \end{aligned}$$

$$\begin{aligned}
& 2\alpha_1\alpha_2^2 - \beta_3\alpha_2^2 - \alpha_1\alpha_3\alpha_2 + 2\alpha_4\beta_1\alpha_2 + \alpha_3\beta_2\alpha_2 - \alpha_1\alpha_3^2 - \alpha_2^2\beta_2 + \alpha_1\alpha_4\beta_2 - \alpha_3^2\beta_3 \\
& \quad - \alpha_1\alpha_4\beta_3 + \alpha_4\beta_1\beta_4 - \alpha_3\beta_3\beta_4 = 0, \\
& \alpha_2^3 - \alpha_3^2\alpha_2 + \alpha_1\alpha_4\alpha_2 + 2\alpha_4\beta_2\alpha_2 - 2\alpha_4\beta_3\alpha_2 + \alpha_3\beta_4\alpha_2 - \alpha_1\alpha_3\alpha_4 + \alpha_4^2\beta_1 - \alpha_3\alpha_4\beta_3 \\
& \quad - \alpha_3^2\beta_4 + \alpha_4\beta_2\beta_4 - \alpha_4\beta_3\beta_4 = 0, \\
& \beta_3^3 - \beta_2^2\beta_3 - 2\alpha_2\beta_1\beta_3 + 2\alpha_3\beta_1\beta_3 + \alpha_1\beta_2\beta_3 + \beta_1\beta_4\beta_3 + \alpha_4\beta_1^2 - \alpha_1\beta_2^2 - \alpha_1\alpha_2\beta_1 \\
& \quad + \alpha_1\alpha_3\beta_1 - \alpha_2\beta_1\beta_2 - \beta_1\beta_2\beta_4 = 0, \\
& \alpha_2\beta_2^2 + \alpha_3\beta_2^2 + \beta_4\beta_2^2 + \alpha_1\alpha_2\beta_2 - \alpha_3\beta_3\beta_2 + \beta_3\beta_4\beta_2 + \alpha_2\beta_3^2 - \alpha_1\alpha_4\beta_1 - 2\alpha_4\beta_1\beta_3 \\
& \quad - 2\beta_3^2\beta_4 + \alpha_2\beta_1\beta_4 - \alpha_3\beta_1\beta_4 = 0, \\
& \beta_2\alpha_2^2 + \beta_2\beta_4\alpha_2 + \alpha_4\beta_2^2 + \beta_2\beta_4^2 - \beta_3\beta_4^2 - \alpha_3\alpha_4\beta_1 - \alpha_3^2\beta_2 + \alpha_1\alpha_4\beta_2 - 2\alpha_4\beta_2\beta_3 \\
& \quad + \alpha_4\beta_1\beta_4 + \alpha_3\beta_2\beta_4 - 2\alpha_3\beta_3\beta_4 = 0, \\
& \beta_4\alpha_2^2 - \alpha_1\alpha_4\alpha_2 + 2\alpha_4\beta_2\alpha_2 + \alpha_3\beta_4\alpha_2 - 2\alpha_3\beta_4^2 + \alpha_1\alpha_3\alpha_4 - 3\alpha_3\alpha_4\beta_2 + \alpha_3\alpha_4\beta_3 \\
& \quad - 2\alpha_1\alpha_4\beta_4 + 3\alpha_4\beta_2\beta_4 - \alpha_4\beta_3\beta_4 = 0, \\
& 3\alpha_2\alpha_1^2 - 3\alpha_3\alpha_1^2 + 2\alpha_4\beta_1\alpha_1 + \alpha_3\beta_2\alpha_1 - 2\alpha_2\beta_3\alpha_1 - \alpha_3\beta_3\alpha_1 - \alpha_2^2\beta_1 - 2\alpha_3^2\beta_1 \\
& \quad + 3\alpha_2\alpha_3\beta_1 - \alpha_4\beta_1\beta_2 + \alpha_4\beta_1\beta_3 = 0, \\
& \alpha_2^3 + \alpha_3\alpha_2^2 - \beta_4\alpha_2^2 - \alpha_1\alpha_4\alpha_2 + 2\alpha_4\beta_2\alpha_2 + \alpha_4\beta_3\alpha_2 - \alpha_3\beta_4^2 - \alpha_1\alpha_3\alpha_4 - \alpha_3\alpha_4\beta_3 \\
& \quad - \alpha_1\alpha_4\beta_4 + \alpha_4\beta_3\beta_4 = 0, \\
& \beta_3^3 - \alpha_1\beta_2^2 + \beta_2\beta_3^2 + \alpha_2\beta_1\beta_3 + 2\alpha_3\beta_1\beta_3 - \beta_1\beta_4\beta_3 + \alpha_1\alpha_2\beta_1 - \alpha_1^2\beta_2 \\
& \quad - \alpha_2\beta_1\beta_2 - \alpha_1\beta_1\beta_4 - \beta_1\beta_2\beta_4 = 0, \\
& 2\beta_2\alpha_1^2 - \beta_3^2\alpha_1 + \alpha_2\beta_1\alpha_1 - 3\alpha_3\beta_1\alpha_1 - \beta_2\beta_3\alpha_1 + 2\beta_1\beta_4\alpha_1 - \alpha_2\beta_1\beta_2 + 3\alpha_3\beta_1\beta_2 \\
& \quad - 2\alpha_3\beta_1\beta_3 - \beta_1\beta_2\beta_4 + \beta_1\beta_3\beta_4 = 0, \\
& 2\alpha_4\beta_2^2 + 3\beta_4^2\beta_2 - 3\alpha_4\beta_3\beta_2 + \alpha_2\beta_4\beta_2 - \alpha_3\beta_4\beta_2 + \alpha_4\beta_3^2 - 3\beta_3\beta_4^2 - \alpha_2\alpha_4\beta_1 \\
& \quad + \alpha_3\alpha_4\beta_1 - 2\alpha_4\beta_1\beta_4 + 2\alpha_2\beta_3\beta_4 = 0, \\
& \beta_2\alpha_2^2 + \alpha_3\beta_2\alpha_2 + \alpha_4\beta_2^2 - \alpha_4\beta_3^2 - \alpha_2\alpha_4\beta_1 - \alpha_3\alpha_4\beta_1 - \alpha_4\beta_1\beta_4 + \alpha_3\beta_2\beta_4 = 0, \\
& \alpha_3\beta_3^2 + \alpha_3^2\beta_1 - \alpha_2^2\beta_1 - \alpha_1\alpha_4\beta_1 + \alpha_1\alpha_3\beta_2 - \alpha_4\beta_1\beta_2 - \alpha_4\beta_1\beta_3 + \alpha_3\beta_2\beta_3 = 0, \\
& 2\alpha_4\alpha_2^2 - \alpha_3\alpha_4\alpha_2 - \alpha_1\alpha_4^2 + 2\alpha_4^2\beta_2 - \alpha_4^2\beta_3 - \alpha_3\alpha_4\beta_4 = 0, \\
& \alpha_2\beta_1^2 - 2\alpha_3\beta_1^2 + \beta_4\beta_1^2 - 2\beta_3^2\beta_1 + \alpha_1\beta_2\beta_1 + \beta_2\beta_3\beta_1 = 0, \\
& \alpha_4\beta_1^2 + 2\alpha_1\alpha_2\beta_1 - 2\alpha_1\alpha_3\beta_1 + \alpha_3\beta_2\beta_1 - 2\alpha_3\beta_3\beta_1 = 0, \\
& 2\alpha_2\beta_2\alpha_4 - \alpha_3\beta_2\alpha_4 + 2\beta_2\beta_4\alpha_4 - 2\beta_3\beta_4\alpha_4 - \alpha_4^2\beta_1 = 0.
\end{aligned}$$

For A_1 the system above has the form

$$\begin{aligned}
& \alpha_4\beta_1^2 - \alpha_1\beta_1 + \alpha_1\alpha_2\beta_1 - 2\alpha_2\beta_1 - 2\beta_1 = 0, \\
& \alpha_1^2 + \alpha_2\alpha_1 - 2\alpha_4\beta_1\alpha_1 + \alpha_1 - \alpha_2 + 2\alpha_2^2\beta_1 + \alpha_2\beta_1 - \beta_1 - 1 = 0, \\
& \alpha_2\alpha_1^2 + \alpha_1^2 - 3\alpha_2\alpha_1 + \alpha_4\beta_1\alpha_1 - 3\alpha_1 + \alpha_2 + 2\alpha_2\beta_1 - \alpha_4\beta_1 + \beta_1 + 1 = 0, \\
& 2\alpha_4\alpha_1^2 - 2\alpha_2^2\alpha_1 - \alpha_2\alpha_1 - 2\alpha_4\alpha_1 + \alpha_1 + 3\alpha_2^2 + 3\alpha_2 - \alpha_4\beta_1 = 0, \\
& 2\alpha_2\alpha_1^2 - 3\alpha_1^2 - 3\alpha_2\alpha_1 + 2\alpha_4\beta_1\alpha_1 - \alpha_1 - \alpha_2\beta_1 + \alpha_4\beta_1 - 2\beta_1 = 0, \\
& 4\alpha_4\alpha_1^2 - 4\alpha_2^2\alpha_1 + \alpha_2\alpha_1 + \alpha_4\alpha_1 + \alpha_1 + \alpha_2^2 + \alpha_2 - \alpha_4 + \alpha_4\beta_1 = 0, \\
& 2\alpha_1\alpha_2^2 - 3\alpha_2^2 - \alpha_1\alpha_2 + 2\alpha_4\beta_1\alpha_2 - 3\alpha_2 - \alpha_1\alpha_4 + \alpha_4\beta_1 - 1 = 0, \\
& 4\alpha_2^3 + 3\alpha_2^2 - 4\alpha_1\alpha_4\alpha_2 - 2\alpha_4\alpha_2 - 3\alpha_1\alpha_4 - \alpha_4 = 0, \\
& \alpha_1\alpha_2^2 - \alpha_2^2 - \alpha_1\alpha_2 + \alpha_4\beta_1\alpha_2 - \alpha_2 + \alpha_1 - \alpha_1\alpha_4 - 1 = 0, \\
& 2\alpha_2^3 - 2\alpha_1\alpha_4\alpha_2 - \alpha_4\alpha_2 - \alpha_4 = 0, \\
& \alpha_2^2 - \alpha_1\alpha_4\alpha_2 + 2\alpha_4\alpha_2 + \alpha_4 - \alpha_4^2\beta_1 = 0,
\end{aligned}$$

$$\begin{aligned}
2\alpha_4\alpha_2^2 - 2\alpha_1\alpha_4^2 - \alpha_4^2 &= 0, \\
2\beta_1\alpha_1^2 - 3\beta_1\alpha_1 + 2\alpha_2\beta_1^2 + 2\beta_1^2 + 2\beta_1 &= 0, \\
\alpha_1^2 + \alpha_2\beta_1\alpha_1 - \beta_1\alpha_1 - 3\alpha_1 + \alpha_4\beta_1^2 - \alpha_2\beta_1 + 2\beta_1 + 1 &= 0, \\
2\alpha_1^3 - 7\alpha_1^2 + 2\alpha_2\beta_1\alpha_1 + 2\beta_1\alpha_1 + 5\alpha_1 - 4\alpha_2\beta_1 - 2\beta_1 - 1 &= 0, \\
\alpha_2\alpha_1^2 - 4\alpha_2\alpha_1 + \alpha_4\beta_1\alpha_1 + \alpha_1 + 3\alpha_2 + \alpha_2\beta_1 - 2\alpha_4\beta_1 &= 0, \\
4\alpha_1^3 - 3\alpha_1^2 + 4\alpha_2\beta_1\alpha_1 + 4\beta_1\alpha_1 + \alpha_1 + 3\alpha_2\beta_1 + 2\beta_1 &= 0, \\
2\alpha_2\alpha_1^2 - \alpha_1^2 - 3\alpha_2\alpha_1 + 2\alpha_4\beta_1\alpha_1 + 2\alpha_1 + 3\alpha_2 + \alpha_2\beta_1 + \alpha_4\beta_1 &= 0, \\
4\alpha_2\alpha_1^2 - \alpha_1^2 - 7\alpha_2\alpha_1 - 2\alpha_1 + 2\alpha_2 + 4\alpha_2^2\beta_1 + 3\alpha_2\beta_1 - \alpha_4\beta_1 + \beta_1 + 1 &= 0, \\
2\alpha_1\alpha_2^2 - 5\alpha_2^2 - \alpha_1\alpha_2 + 2\alpha_4\beta_1\alpha_2 + \alpha_1\alpha_4 + \alpha_4 + \alpha_4\beta_1 &= 0, \\
2\alpha_2\alpha_1^2 + \alpha_1^2 - 2\alpha_2\alpha_1 - 3\alpha_1 + 2\alpha_2^2\beta_1 + 3\alpha_2\beta_1 + \alpha_4\beta_1 + \beta_1 + 1 &= 0, \\
\alpha_1\alpha_2^2 + \alpha_4\beta_1\alpha_2 - 2\alpha_1\alpha_4 + \alpha_4 + \alpha_4\beta_1 &= 0, \\
2\alpha_4\alpha_1^2 - \alpha_2\alpha_1 - 2\alpha_4\alpha_1 - \alpha_1 - \alpha_2^2 - 2\alpha_2 + 2\alpha_2\alpha_4\beta_1 + \alpha_4\beta_1 &= 0, \\
\beta_1\alpha_4^2 - \alpha_1\alpha_4 + \alpha_1\alpha_2\alpha_4 - 2\alpha_2\alpha_4 &= 0.
\end{aligned}$$

The last equation of the system is written $\alpha_4(\beta_1\alpha_4 - \alpha_1 + \alpha_1\alpha_2 - 2\alpha_2) = 0$ from that we get the following options:

Case 1 If $\alpha_4 = 0$ then from the equation 4 we have $\alpha_2 = 0$ or $\alpha_2 = -\frac{3}{4}$.

- * If $\alpha_2 = 0$ then from the equations 16 and 1 we get $\alpha_1 = 0$, $\beta_1 = 0$ which implies a contradiction with the equation 2.
- * If $\alpha_2 = -\frac{3}{4}$ then applying the equations 22 and 1 we get $\alpha_1 = 0$, $\beta_1 = 0$ which implies a contradiction with the equation 2.

Case 2 If $\alpha_4 \neq 0$ then $-\beta_1\alpha_4 + \alpha_1 - \alpha_1\alpha_2 + 2\alpha_2 = 0$ and the equation 1 leads to $\beta_1 = 0$. Then the equation 14 implies $\alpha_1^2 - 3\alpha_1 + 1 = 0$ (so α_1 cannot be 0). Now using the equation 12 one has $4\alpha_1^3 - 3\alpha_1^2 + \alpha_1 = \alpha_1(4\alpha_1^2 - 3\alpha_1 + 1) = \alpha_1(3\alpha_1^2) = 0$ hence, $\alpha_1 = 0$ which is a contradiction, therefore A_1 is not a Malcev algebra.

Substituting the structure constants of A_2 into the general system we derive the system equations

$$\begin{aligned}
\alpha_1^3 + \beta_2\alpha_1^2 - 3\alpha_1^2 - \beta_2\alpha_1 + 3\alpha_1 - \beta_1^2 + \beta_2^2 - 1 &= 0, \\
2\alpha_1^3 - 5\alpha_1^2 + 2\beta_2\alpha_1 + 4\alpha_1 - \beta_2 - 1 &= 0, \\
\alpha_1^2 + 3\beta_2\alpha_1 - 2\alpha_1 + 2\beta_2^2 - 3\beta_2 + 1 &= 0, \\
\alpha_1^3 - 3\beta_2\alpha_1^2 - 2\alpha_1^2 + \beta_2\alpha_1 + \alpha_1 &= 0, \\
2\beta_1\alpha_1^2 - 4\beta_1\alpha_1 + 2\beta_1 - \beta_1\beta_2 &= 0, \\
2\alpha_1^2 - \beta_2\alpha_1 + \alpha_1 + \beta_2^2 - 1 &= 0, \\
\beta_2^2 - \alpha_1\beta_2 + \beta_2 - \alpha_1 &= 0, \\
\alpha_1^2 - 2\alpha_1 - \beta_2^2 + 1 &= 0, \\
\beta_2^2 + 3\alpha_1\beta_2 - 2\beta_2 &= 0, \\
\alpha_1\beta_1 - \beta_2\beta_1 + \beta_1 &= 0, \\
\alpha_1\beta_1 + \beta_2\beta_1 - \beta_1 &= 0, \\
\alpha_1^2 + \beta_2\alpha_1 - \alpha_1 &= 0, \\
2\beta_1 - \alpha_1\beta_1 &= 0, \\
\alpha_1\beta_1 - \beta_1\beta_2 &= 0, \\
\beta_2\beta_1 + \beta_1 &= 0, \\
2\beta_2 - 1 &= 0, \\
\beta_1 &= 0,
\end{aligned}$$

It is easy to see the solution $\beta_1 = 0$, $\beta_2 = \frac{1}{2}$ and $\alpha_1 = \frac{1}{2}$ to the system.

For A_3 the general system has the form

$$\begin{aligned}
 2\beta_1 - \beta_1\beta_2 &= 0, & \beta_2^2 + 2\beta_1 - 1 &= 0, \\
 \beta_2 + 1 &= 0, & \beta_2 + 3 &= 0, \\
 \beta_2 - 1 &= 0, & \beta_2 - 3 &= 0, \\
 4 &= 0, & -1 &= 0, \\
 2 &= 0, & 2\beta_1^2 - \beta_2\beta_1 + 2\beta_1 &= 0, \\
 \beta_2^2 + \beta_1 - 1 &= 0, & 4\beta_1 + \beta_2 + 1 &= 0, \\
 \beta_2^2 - 2\beta_2 + 3 &= 0, & 3\beta_1\beta_2 - 3\beta_1 &= 0, \\
 \beta_2^2 - \beta_2 - 4\beta_1 - 2 &= 0, & 5 - 3\beta_2 &= 0, \\
 \beta_2^2 - 2\beta_2 - 2\beta_1 &= 0, & \beta_2 &= 0.
 \end{aligned}$$

which is a contradiction.

For A_4 the system has the form

$$\begin{aligned}
 \alpha_1^3 + \beta_2\alpha_1^2 - 3\alpha_1^2 - \beta_2\alpha_1 + 3\alpha_1 + \beta_2^2 - 1 &= 0, \\
 2\alpha_1^3 - 5\alpha_1^2 + 2\beta_2\alpha_1 + 4\alpha_1 - \beta_2 - 1 &= 0, \\
 \alpha_1^3 - 3\beta_2\alpha_1^2 - 2\alpha_1^2 + \beta_2\alpha_1 + \alpha_1 &= 0.
 \end{aligned}$$

The equation 3 is written as follows

$$\alpha_1(\alpha_1^2 - 3\alpha_1\beta_2 - 2\alpha_1 + \beta_2 + 1) = 0$$

Case 1: If $\alpha_1 = 0$ the equation 2 implies $\beta_2 = -1$ therefore, $A_4(0, -1)$ is a Malcev algebra.

Case 2: If $\alpha_1 \neq 0$ then $\alpha_1^2 - 3\alpha_1\beta_2 - 2\alpha_1 + \beta_2 + 1 = 0$ and the equation 2 gives

$$2\alpha_1^3 - 5\alpha_1^2 + 2\alpha_1(\beta_2 + 1) + 2\alpha_1 - (\beta_2 + 1) = 0$$

from where we get $2\alpha_1^3 - 5\alpha_1^2 + 2\alpha_1 - (\beta_2 + 1)(2\alpha_1 - 1) = 0$. Hence, $3\beta_2\alpha_1(-2\alpha_1 + 1) = 0$ and we obtain $\beta_2 = 0$, $\alpha_1 = 1$ or $\alpha_1 = \frac{1}{2}$, $\beta_2 = \frac{1}{2}$.

Substituting the structure constants of algebras A_5 , A_6 , A_7 , A_9 , A_{10} and A_{11} we can easily see that the systems obtained are contradictions.

As for the algebras A_8 the system has the solution $\alpha_1 = 0$. Finally, the set of structure constants of A_{12} satisfies the general system of equations.

- I_{24} case. $A(A(A \otimes I) \otimes I)((u \otimes v \otimes w + v \otimes w \otimes u + w \otimes u \otimes v) \otimes u)$
 $= -A(A \otimes A)(u \otimes v \otimes u \otimes w) - A(A(I \otimes A) \otimes I)(v \otimes u \otimes w \otimes u) - A(A(A \otimes I) \otimes I)(u \otimes w \otimes u \otimes v)$.

Here is the system of equations in terms of structure constants

$$\begin{aligned}
 4\alpha_4\alpha_1^2 + 5\alpha_2^2\alpha_1 + 3\alpha_2\alpha_3\alpha_1 + 3\alpha_4\beta_2\alpha_1 + \alpha_4\beta_3\alpha_1 + 2\alpha_2\beta_4\alpha_1 + 2\alpha_3\beta_4\alpha_1 \\
 + \alpha_4\beta_2^2 + \alpha_4\beta_3^2 + 3\alpha_2\alpha_4\beta_1 + 5\alpha_3\alpha_4\beta_1 + \alpha_2^2\beta_2 + \alpha_3^2\beta_2 + 4\alpha_2\alpha_3\beta_2 \\
 + 2\alpha_2\alpha_3\beta_3 + 2\alpha_4\beta_2\beta_3 + 2\alpha_4\beta_1\beta_4 + 3\alpha_3\beta_2\beta_4 + 3\alpha_3\beta_3\beta_4 &= 0, \\
 2\alpha_4\alpha_1^2 + 4\alpha_2^2\alpha_1 + \alpha_3^2\alpha_1 + 5\alpha_2\alpha_3\alpha_1 + 3\alpha_4\beta_2\alpha_1 + 3\alpha_4\beta_3\alpha_1 + 2\alpha_3\beta_4\alpha_1 \\
 + 2\alpha_4\beta_3^2 + 5\alpha_2\alpha_4\beta_1 + 3\alpha_3\alpha_4\beta_1 + 2\alpha_3^2\beta_2 + 2\alpha_2\alpha_3\beta_2 + \alpha_2^2\beta_3 + \alpha_3^2\beta_3 \\
 + 2\alpha_2\alpha_3\beta_3 + 2\alpha_4\beta_2\beta_3 + 4\alpha_4\beta_1\beta_4 + \alpha_3\beta_2\beta_4 + 3\alpha_3\beta_3\beta_4 &= 0, \\
 2\beta_1\alpha_2^2 + \beta_2^2\alpha_2 + \beta_3^2\alpha_2 + 2\alpha_3\beta_1\alpha_2 + 3\alpha_1\beta_2\alpha_2 + 2\beta_2\beta_3\alpha_2 + 3\beta_1\beta_4\alpha_2 + 2\alpha_3\beta_2^2 \\
 + 2\beta_1\beta_4^2 + 4\alpha_1\alpha_4\beta_1 + \alpha_1\alpha_3\beta_2 + 3\alpha_4\beta_1\beta_2 + 5\alpha_4\beta_1\beta_3 + 2\alpha_3\beta_2\beta_3 + \beta_2^2\beta_4 \\
 + 4\beta_3^2\beta_4 + 3\alpha_3\beta_1\beta_4 + 2\alpha_1\beta_2\beta_4 + 5\beta_2\beta_3\beta_4 &= 0, \\
 \beta_1\alpha_2^2 + 2\alpha_3\beta_1\alpha_2 + 3\alpha_1\beta_2\alpha_2 + 2\beta_2\beta_3\alpha_2 + \beta_1\beta_4\alpha_2 + \alpha_3\beta_2^2 + \alpha_3\beta_3^2 + 4\beta_1\beta_4^2 \\
 + \alpha_3^2\beta_1 + 2\alpha_1\alpha_4\beta_1 + 3\alpha_1\alpha_3\beta_2 + 5\alpha_4\beta_1\beta_2 + 3\alpha_4\beta_1\beta_3 + 4\alpha_3\beta_2\beta_3 \\
 + 5\beta_3^2\beta_4 + 3\alpha_3\beta_1\beta_4 + 2\alpha_1\beta_2\beta_4 + 2\alpha_1\beta_3\beta_4 + 3\beta_2\beta_3\beta_4 &= 0,
 \end{aligned}$$

$$\begin{aligned}
& \beta_1\alpha_3^2 + \beta_2^2\alpha_3 + \beta_3^2\alpha_3 + \alpha_2\beta_1\alpha_3 + 2\alpha_1\beta_2\alpha_3 + 3\beta_2\beta_3\alpha_3 + 2\beta_1\beta_4\alpha_3 + \beta_1\beta_4^2 \\
& \quad + 2\alpha_1\alpha_2\beta_2 + \alpha_4\beta_1\beta_2 + \alpha_4\beta_1\beta_3 + \alpha_2\beta_2\beta_3 + 3\beta_3^2\beta_4 + \alpha_2\beta_1\beta_4 \\
& \quad \quad \quad + \alpha_1\beta_2\beta_4 + 2\beta_2\beta_3\beta_4 = 0, \\
& 2\alpha_1\alpha_2^2 + \beta_3\alpha_2^2 + 3\alpha_1\alpha_3\alpha_2 + 2\alpha_4\beta_1\alpha_2 + \alpha_3\beta_2\alpha_2 + 2\alpha_3\beta_3\alpha_2 + \alpha_1\alpha_3^2 + 2\alpha_4\beta_3^2 \\
& \quad + 2\alpha_3\alpha_4\beta_1 + \alpha_3^2\beta_2 + \alpha_1\alpha_4\beta_2 + \alpha_3^2\beta_3 + \alpha_1\alpha_4\beta_3 + 2\alpha_4\beta_2\beta_3 \\
& \quad \quad \quad + \alpha_4\beta_1\beta_4 + \alpha_3\beta_3\beta_4 = 0, \\
& 2\beta_1\alpha_2^2 + \beta_2^2\alpha_2 + \beta_3^2\alpha_2 + 2\alpha_3\beta_1\alpha_2 + \alpha_1\beta_2\alpha_2 + 2\beta_2\beta_3\alpha_2 + \beta_1\beta_4\alpha_2 + \alpha_3\beta_2^2 \\
& \quad + \alpha_1\alpha_4\beta_1 + 2\alpha_4\beta_1\beta_2 + 2\alpha_4\beta_1\beta_3 + \alpha_3\beta_2\beta_3 + \beta_2^2\beta_4 + 2\beta_3^2\beta_4 \\
& \quad \quad \quad + \alpha_3\beta_1\beta_4 + 3\beta_2\beta_3\beta_4 = 0, \\
& \alpha_4\alpha_1^2 + 3\alpha_2^2\alpha_1 + 2\alpha_2\alpha_3\alpha_1 + 2\alpha_4\beta_2\alpha_1 + \alpha_4\beta_3\alpha_1 + \alpha_3\beta_4\alpha_1 + \alpha_4\beta_2^2 + \alpha_2\alpha_4\beta_1 \\
& \quad + \alpha_3\alpha_4\beta_1 + \alpha_2^2\beta_2 + \alpha_3^2\beta_2 + 3\alpha_2\alpha_3\beta_2 + \alpha_2\alpha_3\beta_3 + \alpha_4\beta_2\beta_3 \\
& \quad \quad \quad + 2\alpha_3\beta_2\beta_4 + 2\alpha_3\beta_3\beta_4 = 0, \\
& 2\alpha_2^3 + 2\alpha_3\alpha_2^2 + \beta_4\alpha_2^2 + 7\alpha_1\alpha_4\alpha_2 + 6\alpha_4\beta_2\alpha_2 + 2\alpha_4\beta_3\alpha_2 + 3\alpha_3\beta_4\alpha_2 + 4\alpha_3\beta_4^2 \\
& \quad + \alpha_1\alpha_3\alpha_4 + 2\alpha_4^2\beta_1 + 5\alpha_3\alpha_4\beta_2 + \alpha_3\alpha_4\beta_3 + 4\alpha_1\alpha_4\beta_4 \\
& \quad \quad \quad + 5\alpha_4\beta_2\beta_4 + 3\alpha_4\beta_3\beta_4 = 0, \\
& \beta_3^3 + \alpha_1\beta_3^2 + \beta_2\beta_3^2 + \alpha_2\beta_1\beta_3 + 2\alpha_3\beta_1\beta_3 + 2\alpha_1\beta_2\beta_3 + 3\beta_1\beta_4\beta_3 + 2\alpha_4\beta_1^2 \\
& \quad + 3\alpha_1\alpha_2\beta_1 + 2\alpha_1\alpha_3\beta_1 + \alpha_1^2\beta_2 + \alpha_2\beta_1\beta_2 + 2\alpha_3\beta_1\beta_2 \\
& \quad \quad \quad + \alpha_1\beta_1\beta_4 + \beta_1\beta_2\beta_4 = 0, \\
& \alpha_2^3 + \alpha_3\alpha_2^2 + \beta_4\alpha_2^2 + 3\alpha_1\alpha_4\alpha_2 + 2\alpha_4\beta_2\alpha_2 + \alpha_4\beta_3\alpha_2 + 2\alpha_3\beta_4\alpha_2 + \alpha_3\beta_4^2 \\
& \quad + \alpha_1\alpha_3\alpha_4 + 2\alpha_4^2\beta_1 + 2\alpha_3\alpha_4\beta_2 + \alpha_3\alpha_4\beta_3 + \alpha_1\alpha_4\beta_4 \\
& \quad \quad \quad + 2\alpha_4\beta_2\beta_4 + 3\alpha_4\beta_3\beta_4 = 0, \\
& 2\beta_3^3 + \alpha_1\beta_3^2 + 2\beta_2\beta_3^2 + 2\alpha_2\beta_1\beta_3 + 6\alpha_3\beta_1\beta_3 + 3\alpha_1\beta_2\beta_3 + 7\beta_1\beta_4\beta_3 + 2\alpha_4\beta_1^2 \\
& \quad + 3\alpha_1\alpha_2\beta_1 + 5\alpha_1\alpha_3\beta_1 + 4\alpha_1^2\beta_2 + \alpha_2\beta_1\beta_2 + 5\alpha_3\beta_1\beta_2 \\
& \quad \quad \quad + 4\alpha_1\beta_1\beta_4 + \beta_1\beta_2\beta_4 = 0, \\
& 2\beta_2\alpha_2^2 + 5\alpha_4\beta_1\alpha_2 + 2\alpha_3\beta_2\alpha_2 + 3\beta_2\beta_4\alpha_2 + 2\beta_3\beta_4\alpha_2 + 4\alpha_4\beta_2^2 + \alpha_4\beta_3^2 + 5\beta_2\beta_4^2 \\
& \quad + 7\beta_3\beta_4^2 + \alpha_3\alpha_4\beta_1 + 2\alpha_1\alpha_4\beta_2 + 5\alpha_4\beta_2\beta_3 + 6\alpha_4\beta_1\beta_4 + 3\alpha_3\beta_2\beta_4 = 0, \\
& 7\alpha_2\alpha_1^2 + 5\alpha_3\alpha_1^2 + 6\alpha_4\beta_1\alpha_1 + 3\alpha_3\beta_2\alpha_1 + 2\alpha_2\beta_3\alpha_1 + 3\alpha_3\beta_3\alpha_1 + 2\alpha_3\beta_3^2 + \alpha_2^2\beta_1 \\
& \quad + 4\alpha_3^2\beta_1 + 5\alpha_2\alpha_3\beta_1 + \alpha_4\beta_1\beta_2 + 5\alpha_4\beta_1\beta_3 + 2\alpha_3\beta_2\beta_3 + 2\alpha_3\beta_1\beta_4 = 0, \\
& \alpha_2^3 + 2\alpha_3\alpha_2^2 + \alpha_3^2\alpha_2 + \alpha_1\alpha_4\alpha_2 + 2\alpha_4\beta_2\alpha_2 + 4\alpha_4\beta_3\alpha_2 + \alpha_3\beta_4\alpha_2 + \alpha_1\alpha_3\alpha_4 \\
& \quad + \alpha_4^2\beta_1 + 2\alpha_3\alpha_4\beta_2 + \alpha_3\alpha_4\beta_3 + \alpha_3^2\beta_4 + \alpha_4\beta_2\beta_4 + 5\alpha_4\beta_3\beta_4 = 0, \\
& \beta_3^3 + 2\beta_2\beta_3^2 + \beta_2^2\beta_3 + 4\alpha_2\beta_1\beta_3 + 2\alpha_3\beta_1\beta_3 + \alpha_1\beta_2\beta_3 + \beta_1\beta_4\beta_3 + \alpha_4\beta_1^2 + \alpha_1\beta_2^2 \\
& \quad + 5\alpha_1\alpha_2\beta_1 + \alpha_1\alpha_3\beta_1 + \alpha_2\beta_1\beta_2 + 2\alpha_3\beta_1\beta_2 + \beta_1\beta_2\beta_4 = 0, \\
& \beta_2\alpha_2^2 + \alpha_4\beta_1\alpha_2 + \alpha_3\beta_2\alpha_2 + 2\beta_2\beta_4\alpha_2 + \alpha_4\beta_2^2 + \alpha_4\beta_3^2 + 2\beta_2\beta_4^2 + 4\beta_3\beta_4^2 + \alpha_3\alpha_4\beta_1 \\
& \quad + 2\alpha_1\alpha_4\beta_2 + 2\alpha_4\beta_2\beta_3 + 3\alpha_4\beta_1\beta_4 + 3\alpha_3\beta_2\beta_4 = 0, \\
& \beta_2\alpha_2^2 + 2\alpha_3\beta_2\alpha_2 + \beta_2\beta_4\alpha_2 + \alpha_4\beta_2^2 + \beta_2\beta_4^2 + 5\beta_3\beta_4^2 + \alpha_3\alpha_4\beta_1 + \alpha_3^2\beta_2 + \alpha_1\alpha_4\beta_2 \\
& \quad + 4\alpha_4\beta_2\beta_3 + \alpha_4\beta_1\beta_4 + 3\alpha_3\beta_2\beta_4 + 2\alpha_3\beta_3\beta_4 = 0, \\
& 5\alpha_2\alpha_1^2 + \alpha_3\alpha_1^2 + \alpha_4\beta_1\alpha_1 + 2\alpha_2\beta_2\alpha_1 + 3\alpha_3\beta_2\alpha_1 + \alpha_3\beta_3\alpha_1 + \alpha_3\beta_2^2 + \alpha_3\beta_3^2 + \alpha_3^2\beta_1 \\
& \quad + 4\alpha_2\alpha_3\beta_1 + \alpha_4\beta_1\beta_2 + 2\alpha_3\beta_2\beta_3 + \alpha_3\beta_1\beta_4 = 0, \\
& 4\alpha_2\alpha_1^2 + 2\alpha_3\alpha_1^2 + 3\alpha_4\beta_1\alpha_1 + 3\alpha_3\beta_2\alpha_1 + 2\alpha_3\beta_3\alpha_1 + \alpha_3\beta_2^2 + \alpha_2^2\beta_1 + \alpha_3^2\beta_1 \\
& \quad + 2\alpha_2\alpha_3\beta_1 + \alpha_4\beta_1\beta_2 + \alpha_4\beta_1\beta_3 + \alpha_3\beta_2\beta_3 + 2\alpha_3\beta_1\beta_4 = 0, \\
& 4\alpha_4\alpha_2^2 + \alpha_3\alpha_4\alpha_2 + 6\alpha_4\beta_4\alpha_2 + \alpha_1\alpha_4^2 + 6\alpha_4\beta_4^2 + 4\alpha_4^2\beta_2 + \alpha_4^2\beta_3 + \alpha_3\alpha_4\beta_4 = 0, \\
& 6\beta_1\alpha_1^2 + \beta_1\beta_2\alpha_1 + 6\beta_1\beta_3\alpha_1 + \alpha_2\beta_1^2 + 4\alpha_3\beta_1^2 + 4\beta_1\beta_3^2 + \beta_1\beta_2\beta_3 + \beta_1^2\beta_4 = 0, \\
& 6\beta_4^3 + 10\alpha_4\beta_2\beta_4 + 2\alpha_4\beta_3\beta_4 + \alpha_4^2\beta_1 + 4\alpha_2\alpha_4\beta_2 + \alpha_3\alpha_4\beta_2 = 0, \\
& 6\alpha_1^3 + 2\alpha_2\beta_1\alpha_1 + 10\alpha_3\beta_1\alpha_1 + \alpha_4\beta_1^2 + \alpha_3\beta_1\beta_2 + 4\alpha_3\beta_1\beta_3 = 0.
\end{aligned}$$

- I_{25} case. $A(A \otimes I)(u \otimes v \otimes w) = A(I \otimes A)(u \otimes v \otimes w) + A(I \otimes A)(u \otimes w \otimes v)$.

The system of equations is

$$\begin{aligned}
\alpha_1\alpha_3 - \alpha_4\beta_1 + \alpha_2\beta_2 + \alpha_2\beta_3 &= 0, & \alpha_1^2 + 2\alpha_2\beta_1 - \alpha_3\beta_1 &= 0, \\
\alpha_1\alpha_3 - \beta_2\alpha_3 + \alpha_2\beta_2 + \alpha_2\beta_3 &= 0, & \beta_2^2 + \alpha_3\beta_1 &= 0, \\
\alpha_3\beta_2 - \beta_4\beta_2 - \alpha_2\beta_3 - \alpha_3\beta_3 &= 0, & \alpha_3^2 + \alpha_4\beta_2 &= 0, \\
\alpha_2\alpha_4 - 2\alpha_3\alpha_4 - \beta_4\alpha_4 &= 0, & \alpha_1\beta_1 + 2\beta_2\beta_1 - \beta_3\beta_1 &= 0, \\
\alpha_2^2 - 2\beta_4\alpha_2 - 2\alpha_1\alpha_4 + \alpha_4\beta_2 &= 0, & 2\alpha_4\beta_1 - \alpha_2\beta_2 + \beta_2\beta_4 &= 0, \\
\beta_3^2 - 2\alpha_1\beta_3 + \alpha_3\beta_1 - 2\beta_1\beta_4 &= 0, & \alpha_1\alpha_3 - \beta_3\alpha_3 + 2\alpha_4\beta_1 &= 0, \\
\alpha_4\beta_1 - \alpha_2\beta_3 - \alpha_3\beta_3 - \beta_2\beta_4 &= 0, & \beta_4^2 - \alpha_4\beta_2 + 2\alpha_4\beta_3 &= 0, \\
\alpha_3^2 + \alpha_2\alpha_3 - \beta_4\alpha_3 - \alpha_1\alpha_4 & & \beta_2^2 - \alpha_1\beta_2 + \beta_3\beta_2 + \alpha_2\beta_1 & \\
& & + \alpha_4\beta_2 + \alpha_4\beta_3 & = 0, & + \alpha_3\beta_1 - \beta_1\beta_4 & = 0.
\end{aligned}$$

- I_{26} case. $A(A \otimes I)(u \otimes v \otimes w) = -A(I \otimes A)(u \otimes v \otimes w) - A(I \otimes A)(u \otimes w \otimes v)$.

The identity is equivalent to the system of equations

$$\begin{aligned}
2\alpha_1\alpha_2 + \beta_2\alpha_2 + \beta_3\alpha_2 + \alpha_1\alpha_3 + \alpha_4\beta_1 &= 0, & \alpha_2^2 + 2\beta_4\alpha_2 + 2\alpha_1\alpha_4 + \alpha_4\beta_2 &= 0, \\
2\alpha_1\alpha_2 + \beta_2\alpha_2 + \beta_3\alpha_2 + \alpha_1\alpha_3 + \alpha_3\beta_2 &= 0, & \beta_3^2 + 2\alpha_1\beta_3 + \alpha_3\beta_1 + 2\beta_1\beta_4 &= 0, \\
\alpha_3^2 + 2\alpha_2\alpha_3 + \alpha_4\beta_2 + 2\alpha_4\beta_3 &= 0, & 3\alpha_1^2 + 2\alpha_2\beta_1 + \alpha_3\beta_1 &= 0, \\
\alpha_3^2 + \alpha_2\alpha_3 + \beta_4\alpha_3 + \alpha_1\alpha_4 + \alpha_4\beta_2 + \alpha_4\beta_3 &= 0, & \alpha_2\alpha_4 + 2\alpha_3\alpha_4 + 3\beta_4\alpha_4 &= 0, \\
\beta_2^2 + \alpha_1\beta_2 + \beta_3\beta_2 + \alpha_2\beta_1 + \alpha_3\beta_1 + \beta_1\beta_4 &= 0, & 3\alpha_1\beta_1 + 2\beta_2\beta_1 + \beta_3\beta_1 &= 0, \\
\beta_2^2 + 2\beta_3\beta_2 + 2\alpha_2\beta_1 + \alpha_3\beta_1 &= 0, & 2\alpha_4\beta_1 + \alpha_2\beta_2 + 3\beta_2\beta_4 &= 0, \\
\alpha_3\beta_2 + \beta_4\beta_2 + \alpha_2\beta_3 + \alpha_3\beta_3 + 2\beta_3\beta_4 &= 0, & 3\alpha_1\alpha_3 + \beta_3\alpha_3 + 2\alpha_4\beta_1 &= 0, \\
\alpha_4\beta_1 + \alpha_2\beta_3 + \alpha_3\beta_3 + \beta_2\beta_4 + 2\beta_3\beta_4 &= 0, & 3\beta_4^2 + \alpha_4\beta_2 + 2\alpha_4\beta_3 &= 0.
\end{aligned}$$

- I_{27} case. $A((A \otimes I) - (I \otimes A))(u \otimes v \otimes w) = A((A \otimes I) - (I \otimes A))(v \otimes u \otimes w)$.

The identity can be written in terms of structure constants as a system of equations as follows

$$\begin{aligned}
\alpha_1\alpha_2 - \beta_3\alpha_2 - \alpha_1\alpha_3 + \alpha_4\beta_1 + \alpha_3\beta_2 - \alpha_3\beta_3 &= 0, & \alpha_2^2 - \beta_4\alpha_2 - \alpha_1\alpha_4 + 2\alpha_4\beta_2 - \alpha_4\beta_3 &= 0, \\
\alpha_4\beta_1 - \alpha_2\beta_2 + \alpha_3\beta_2 - \alpha_2\beta_3 - \beta_2\beta_4 + \beta_3\beta_4 &= 0, & \beta_3^2 - \alpha_1\beta_3 - \alpha_2\beta_1 + 2\alpha_3\beta_1 - \beta_1\beta_4 &= 0.
\end{aligned}$$

- I_{28} case. $A((A \otimes I) - (I \otimes A))(u \otimes v \otimes w) = -A((A \otimes I) - (I \otimes A))(v \otimes u \otimes w)$.

Here is the the system of equations

$$\begin{aligned}
\alpha_4\beta_1 - \alpha_2\beta_2 - \alpha_3\beta_2 + \alpha_2\beta_3 + \beta_2\beta_4 - \beta_3\beta_4 &= 0, & \alpha_4\beta_1 - \alpha_2\beta_2 &= 0, \\
\alpha_1\alpha_2 - \beta_3\alpha_2 - \alpha_1\alpha_3 - \alpha_4\beta_1 + \alpha_3\beta_2 + \alpha_3\beta_3 &= 0, & \alpha_4\beta_1 - \alpha_3\beta_3 &= 0, \\
\alpha_3^2 - \beta_4\alpha_3 - \alpha_1\alpha_4 + \alpha_4\beta_3 &= 0, & \alpha_2\alpha_4 - \alpha_3\alpha_4 &= 0, \\
\beta_2^2 - \alpha_1\beta_2 + \alpha_2\beta_1 - \beta_1\beta_4 &= 0, & \beta_1\beta_3 - \beta_1\beta_2 &= 0, \\
\beta_3^2 - \alpha_1\beta_3 + \alpha_2\beta_1 - \beta_1\beta_4 &= 0, & \alpha_3\beta_1 - \alpha_2\beta_1 &= 0, \\
\alpha_2^2 - \beta_4\alpha_2 - \alpha_1\alpha_4 + \alpha_4\beta_3 &= 0, & \alpha_4\beta_2 - \alpha_4\beta_3 &= 0.
\end{aligned}$$

- I_{29} case. $A((A \otimes I) - (I \otimes A))(u \otimes v \otimes w) = A((A \otimes I) - (I \otimes A))(w \otimes v \otimes u)$.

The system of equations is

$$\begin{aligned}
\alpha_2^2 - \beta_4\alpha_2 + \alpha_3^2 - 2\alpha_1\alpha_4 + \alpha_4\beta_2 + \alpha_4\beta_3 - \alpha_3\beta_4 &= 0, & 2\alpha_4\beta_1 - \alpha_2\beta_2 - \alpha_3\beta_3 &= 0, \\
\beta_2^2 - \alpha_1\beta_2 + \beta_3^2 + \alpha_2\beta_1 + \alpha_3\beta_1 - \alpha_1\beta_3 - 2\beta_1\beta_4 &= 0.
\end{aligned}$$

- I_{30} case. $A((A \otimes I) - (I \otimes A))(u \otimes v \otimes w) = -A((A \otimes I) - (I \otimes A))(w \otimes v \otimes u)$.

One can write the identity above as a system of equations as follows

$$\begin{aligned}
\alpha_2^2 - \beta_4\alpha_2 - \alpha_3^2 + \alpha_4\beta_2 - \alpha_4\beta_3 + \alpha_3\beta_4 &= 0, & \alpha_3\beta_3 - \alpha_2\beta_2 &= 0, \\
\beta_2^2 - \alpha_1\beta_2 - \beta_3^2 + \alpha_2\beta_1 - \alpha_3\beta_1 + \alpha_1\beta_3 &= 0, & \alpha_3\beta_1 - \alpha_2\beta_1 &= 0, \\
\alpha_1\alpha_2 - \beta_3\alpha_2 - \alpha_1\alpha_3 + \alpha_3\beta_2 &= 0, & \alpha_2\alpha_4 - \alpha_3\alpha_4 &= 0, \\
\alpha_3\beta_2 - 2\beta_4\beta_2 - \alpha_2\beta_3 + \beta_3\beta_4 &= 0, & \alpha_4\beta_3 - \alpha_4\beta_2 &= 0, \\
\beta_1\beta_3 - \beta_1\beta_2 &= 0.
\end{aligned}$$

□

Remark 4.8. Note that in the case of I_{19} if $\text{Char}(\mathbb{F}) = 5$ then $i = 2$ and thus here we correct an inaccuracy admitted in Theorem 7 from [3].

The analogues of the result above for the fields of characteristic 2 and 3 can be easily proved following the same manner. Here we give final results without proof as the following two theorems below. In the case of the characteristic 2 some of the identities coincide, this is also denoted by \cong .

Theorem 4.9. Let $\text{Char}(\mathbb{F}) = 2$. Then the following classification of two-dimensional algebras over \mathbb{F} with respect to the identities $I_1 - I_{30}$ is valid.

$I_1 \cong I_2$. Commutativity identity $\mathbf{uv} = \mathbf{vu}$.

$$A_{2,2}(\alpha_1, \beta_1, 1 + \alpha_1), A_{3,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1 + \alpha_1), A_{5,2}(0), A_{10,2}, A_{11,2}, A_{12,2}.$$

$I_3 \cong I_4$. Associativity identity

$$A_{3,2}(1, 0), A_{4,2}(1, 0), A_{4,2}(1, 1), A_{8,2}(1), A_{10,2}, A_{12,2}.$$

I_5 . Well defined cube identity $\mathbf{u}^2\mathbf{u} = \mathbf{uu}^2$.

$$A_{1,2}(1, 1, 0, 0), A_{2,2}(\alpha_1, \beta_1, 1 + \alpha_1), A_{3,2}(\alpha_1, 1 - \alpha_1), A_{4,2}(\alpha_1, 1 + \alpha_1), \text{ where } \alpha_1 \neq 0, \\ A_{4,2}(\alpha_1, 1), A_{5,2}(0), A_{8,2}(1), A_{10,2}, A_{11,2}, A_{12,2}.$$

$I_6 \cong I_7$. Half-commutativity identity $[\mathbf{u}, \mathbf{v}]\mathbf{w} = \mathbf{w}[\mathbf{u}, \mathbf{v}]$.

$$A_{2,2}(\alpha_1, \beta_1, 1 - \alpha_1), A_{3,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1 - \alpha_1), A_{5,2}(0), A_{10,2}, A_{11,2}, A_{12,2}.$$

$I_8 \cong I_9$. Mixed associativity identity $[\mathbf{u}, \mathbf{v}]\mathbf{w} = \mathbf{u}[\mathbf{v}, \mathbf{w}]$.

$$A_{2,2}(\alpha_1, \beta_1, 1 + \alpha_1), A_{3,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1 + \alpha_1), A_{5,2}(0), A_{10,2}, A_{11,2}, A_{12,2}.$$

$I_{10} \cong I_{11}$. Flexibility identity $\mathbf{u}(\mathbf{vu}) = (\mathbf{uv})\mathbf{u}$.

$$A_{2,2}(\alpha_1, \beta_1, 1 + \alpha_1), A_{3,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1), \text{ where } \alpha_1 \neq 0, A_{5,2}(0), \\ A_{8,2}(1), A_{10,2}, A_{11,2}, A_{12,2}.$$

$I_{12} \cong I_{13}$. Mixed flexibility identity $\mathbf{u}[\mathbf{v}, \mathbf{u}] = [\mathbf{u}, \mathbf{v}]\mathbf{u}$.

$$A_{2,2}(\alpha_1, \beta_1, 1 - \alpha_1), A_{3,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1 + \alpha_1), A_{5,2}(0), A_{10,2}, A_{11,2}, A_{12,2}.$$

$I_{14} \cong I_{15}$. Left Leibniz identity $\mathbf{u}(\mathbf{vw}) = (\mathbf{uv})\mathbf{w} + \mathbf{v}(\mathbf{uw})$.

$$A_{4,2}(0, 1), A_{8,2}(0), A_{12,2}.$$

$I_{16} \cong I_{17}$. Mixed left Leibniz identity $\mathbf{u}[\mathbf{v}, \mathbf{w}] = [\mathbf{u}, \mathbf{v}]\mathbf{w} + \mathbf{v}[\mathbf{u}, \mathbf{w}]$.

$$A_{2,2}(\alpha_1, \beta_1, 1 + \alpha_1), A_{3,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1 + \alpha_1), A_{5,2}(0), A_{10,2}, A_{11,2}, A_{12,2}.$$

I_{18} . Left Poisson identity $(\mathbf{uv})\mathbf{w} + (\mathbf{vw})\mathbf{u} + (\mathbf{wu})\mathbf{v} = 0$.

$$A_{4,2}(0, 1), A_{5,2}(0), A_{8,2}(0), A_{12,2}.$$

$I_{19} \cong I_{20}$. Left Jordan identity $(\mathbf{uv})\mathbf{u}^2 = \mathbf{u}(\mathbf{vu}^2)$.

$$A_{3,2}(0, 0), A_{3,2}(1, 0), A_{4,2}(\alpha_1, 1), A_{4,2}\left(\alpha_1, \sqrt{\alpha_1 + \alpha_1^2}\right), \text{ where } \alpha_1^2 + \alpha_1 + 1 \neq 0, A_{5,2}(\alpha_1), \text{ where } \\ \alpha_1^2 + \alpha_1 + 1 = 0, A_{8,2}(1), A_{10,2}, A_{12,2}.$$

$I_{21} \cong I_{22}$. Mixed left Jordan identity $[\mathbf{u}, \mathbf{v}]\mathbf{u}^2 = \mathbf{u}[\mathbf{v}, \mathbf{u}^2]$.

$$A_{2,2}(\alpha_1, \beta_1, 1 + \alpha_1), A_{3,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(0, 0), A_{5,2}(0), A_{10,2}, A_{11,2}, A_{12,2}.$$

$I_{23} \cong I_{24}$. Left Malcev identity $((\mathbf{uv})\mathbf{w} + (\mathbf{vw})\mathbf{u} + (\mathbf{wu})\mathbf{v})\mathbf{u} = (\mathbf{uv})(\mathbf{uw}) + (\mathbf{v}(\mathbf{uw}))\mathbf{u} + ((\mathbf{uw})\mathbf{u})\mathbf{v}$.

$$A_{3,2}(1, 0), A_{4,2}(1, 0), A_{4,2}(0, 1), A_{8,2}(0), A_{10,2}, A_{12,2}.$$

$I_{25} \cong I_{26}$. Left Zinbiel identity $(\mathbf{uv})\mathbf{w} = \mathbf{u}(\mathbf{vw} + \mathbf{wv})$. $A_{12,2}$.

$I_{27} \cong I_{28}$. Left symmetric identity $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = [\mathbf{v}, \mathbf{u}, \mathbf{w}]$.

$$A_{3,2}(1, 0), A_{4,2}(1, \beta_2), A_{5,2}(1), A_{6,2}(\alpha_1, 0), A_{7,2}(1), A_{8,2}(\alpha_1), A_{9,2}, A_{10,2}, A_{12,2}.$$

$I_{29} \cong I_{30}$. Centro-symmetric identity $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = [\mathbf{w}, \mathbf{v}, \mathbf{u}]$.

$$A_{1,2}(\alpha_1, 1 + \alpha_1, \alpha_1, 1 + \alpha_1), A_{2,2}(\alpha_1, \beta_1, 1 + \alpha_1), A_{3,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1 + \alpha_1), A_{4,2}(\alpha_1, 1), \\ \text{ where } \alpha_1 \neq 0, A_{5,2}(\alpha_1), A_{8,2}(1), A_{9,2}, A_{10,2}, A_{11,2}, A_{12,2}.$$

Theorem 4.10. In the case of $\text{Char}(\mathbb{F}) = 3$ the following classification result of two-dimensional algebras over \mathbb{F} with respect to the identities $I_1 - I_{30}$ holds true.

I_1 . Commutativity identity $\mathbf{uv} = \mathbf{vu}$.

$$A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, , 1 - \alpha_1), A_{5,3}(\alpha_1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}.$$

I_2 . Anti-commutativity identity $\mathbf{uv} = -\mathbf{vu}$. $A_{4,3}(0, -1)$.

I_3 . Associativity identity

$$A_{2,3}(-1, 0, -1), A_{4,3}(-1, -1), A_{4,3}(1, 0), A_{4,3}(1, 1), A_{4,3}(-1, 0), A_{12,3}.$$

I_4 . Anti-associativity identity $(\mathbf{uv})\mathbf{w} = -\mathbf{u}(\mathbf{vw})$. $A_{12,3}$.

I_5 . Well defined cube identity

$$A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{3,3}(0, -1), A_{4,3}(\alpha_1, 1 - \alpha_1), \\ A_{4,3}(\alpha_1, 2\alpha_1 - 1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}.$$

- I₆. Half-commutativity identity* $[\mathbf{u}, \mathbf{v}]\mathbf{w} = \mathbf{w}[\mathbf{u}, \mathbf{v}]$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₇. Anti-half-commutativity identity* $[\mathbf{u}, \mathbf{v}]\mathbf{w} = -\mathbf{w}[\mathbf{u}, \mathbf{v}]$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{4,3}(\alpha_1, \alpha_1 - 1), A_{5,3}(0), A_{8,3}(-1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₈. Mixed associativity identity* $[\mathbf{u}, \mathbf{v}]\mathbf{w} = \mathbf{u}[\mathbf{v}, \mathbf{w}]$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₉. Anti-mixed-associativity identity* $[\mathbf{u}, \mathbf{v}]\mathbf{w} = -\mathbf{u}[\mathbf{v}, \mathbf{w}]$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₁₀. Flexibility identity* $\mathbf{u}(\mathbf{v}\mathbf{u}) = (\mathbf{u}\mathbf{v})\mathbf{u}$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{4,3}(\alpha_1, 2\alpha_1 - 1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₁₁. Anti-flexibility identity* $\mathbf{u}(\mathbf{v}\mathbf{u}) = -(\mathbf{u}\mathbf{v})\mathbf{u}$. $A_{12,3}$.
- I₁₂. Mixed flexibility identity* $\mathbf{u}[\mathbf{v}, \mathbf{u}] = [\mathbf{u}, \mathbf{v}]\mathbf{u}$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{4,3}(\alpha_1, \alpha_1 - 1), A_{5,3}(0), A_{8,3}(-1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₁₃. Mixed anti-flexibility identity* $\mathbf{u}[\mathbf{v}, \mathbf{u}] = -[\mathbf{u}, \mathbf{v}]\mathbf{u}$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₁₄. Left Leibniz identity* $\mathbf{u}(\mathbf{v}\mathbf{w}) = (\mathbf{u}\mathbf{v})\mathbf{w} + \mathbf{v}(\mathbf{u}\mathbf{w})$. $A_{4,3}(0, -1), A_{8,3}(0), A_{12,3}$.
- I₁₅. Left anti-Leibniz identity* $\mathbf{u}(\mathbf{v}\mathbf{w}) = -(\mathbf{u}\mathbf{v})\mathbf{w} - \mathbf{v}(\mathbf{u}\mathbf{w})$.
 $A_{2,3}(2, 0, -1), A_{4,3}(1, 0), A_{4,3}(1, 1), A_{4,3}(-1, -1), A_{12,3}$.
- I₁₆. Mixed left Leibniz identity* $\mathbf{u}[\mathbf{v}, \mathbf{w}] = [\mathbf{u}, \mathbf{v}]\mathbf{w} + \mathbf{v}[\mathbf{u}, \mathbf{w}]$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{4,3}(\alpha_1, \alpha_1 - 1), A_{5,3}(0), A_{8,3}(-1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₁₇. Mixed anti-left Leibniz identity* $\mathbf{u}[\mathbf{v}, \mathbf{w}] = -[\mathbf{u}, \mathbf{v}]\mathbf{w} - \mathbf{v}[\mathbf{u}, \mathbf{w}]$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₁₈. Left Poisson identity* $(\mathbf{u}\mathbf{v})\mathbf{w} + (\mathbf{v}\mathbf{w})\mathbf{u} + (\mathbf{w}\mathbf{u})\mathbf{v} = 0$.
 $A_{2,3}(0, 0, -1), A_{2,3}(2, 0, -1), A_{4,3}(\alpha_1, -(1 - \alpha_1)^2), A_{5,3}(0), A_{8,3}(0), A_{12,3}$.
- I₁₉. Left Jordan identity* $(\mathbf{u}\mathbf{v})\mathbf{u}^2 = \mathbf{u}(\mathbf{v}\mathbf{u}^2)$.
 $A_{2,3}(-1, 0, 1), A_{2,3}(-1, 0, -1), A_{4,3}(\alpha_1, -1 - \alpha_1), \text{ where } \alpha_1 \neq -1 \pm i, A_{4,3}(\alpha_1, \sqrt{\alpha_1 - \alpha_1^2}), A_{4,3}(\alpha_1, -\sqrt{\alpha_1 - \alpha_1^2}), \text{ where } \alpha_1 \neq 0, 1, A_{5,3}(-1 + i), A_{5,3}(-1 - i), A_{8,3}(-1 + i), A_{8,3}(-1 - i), A_{10,3}, A_{12,3}$.
- I₂₀. Left anti-Jordan identity* $(\mathbf{u}\mathbf{v})\mathbf{u}^2 = -\mathbf{u}(\mathbf{v}\mathbf{u}^2)$. $A_{4,3}(0, -1), A_{4,3}(0, 0), A_{12,3}$.
- I₂₁. Mixed left Jordan identity* $[\mathbf{u}, \mathbf{v}]\mathbf{u}^2 = \mathbf{u}[\mathbf{v}, \mathbf{u}^2]$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(0, -1), A_{4,3}(0, 0), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₂₂. Mixed anti-left Jordan identity* $[\mathbf{u}, \mathbf{v}]\mathbf{u}^2 = -\mathbf{u}[\mathbf{v}, \mathbf{u}^2]$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(0, -1), A_{4,3}(0, 0), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{9,3}, A_{10,3}, A_{11,3}, A_{12,3}$.
- I₂₃. Left Malcev identity* $((\mathbf{u}\mathbf{v})\mathbf{w} + (\mathbf{v}\mathbf{w})\mathbf{u} + (\mathbf{w}\mathbf{u})\mathbf{v})\mathbf{u} = (\mathbf{u}\mathbf{v})(\mathbf{u}\mathbf{w}) + (\mathbf{v}(\mathbf{u}\mathbf{w}))\mathbf{u} + ((\mathbf{u}\mathbf{w})\mathbf{u})\mathbf{v}$.
 $A_{2,3}(-1, 0, -1), A_{2,3}(0, 0, -1), A_{4,3}(\alpha_1, -(1 - \alpha_1)^2), A_{5,3}(0), A_{8,3}(0), A_{12,3}$.
- I₂₄. Left anti-Malcev identity* $((\mathbf{u}\mathbf{v})\mathbf{w} + (\mathbf{v}\mathbf{w})\mathbf{u} + (\mathbf{w}\mathbf{u})\mathbf{v})\mathbf{u} = -(\mathbf{u}\mathbf{v})(\mathbf{u}\mathbf{w}) - (\mathbf{v}(\mathbf{u}\mathbf{w}))\mathbf{u} - ((\mathbf{u}\mathbf{w})\mathbf{u})\mathbf{v}$.
 $A_{2,3}(-1, 0, -1), A_{2,3}(0, 0, -1), A_{4,3}(\alpha_1, -(1 - \alpha_1)^2), A_{5,3}(0), A_{8,3}(0), A_{12,3}$.
- I₂₅. Left Zinbiel identity* $(\mathbf{u}\mathbf{v})\mathbf{w} = \mathbf{u}(\mathbf{v}\mathbf{w} + \mathbf{w}\mathbf{v})$. $A_{12,3}$.
- I₂₆. Left anti-Zinbiel identity* $(\mathbf{u}\mathbf{v})\mathbf{w} = -\mathbf{u}(\mathbf{v}\mathbf{w} + \mathbf{w}\mathbf{v})$.
 $A_{2,3}(-1, 0, -1), A_{4,3}(-1, 0), A_{4,3}(-1, -1), A_{4,3}(1, 0), A_{12,3}$.
- I₂₇. Left symmetric identity* $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = [\mathbf{v}, \mathbf{u}, \mathbf{w}]$.
 $A_{2,3}(-1, 0, -1), A_{2,3}(1, 0, -1), A_{4,3}(1, \beta_2), A_{4,3}(-1, \beta_2), A_{5,3}(-1), A_{5,3}(1), A_{8,3}(0), A_{12,3}$.

- I₂₈. Left anti-symmetric identity* $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = -[\mathbf{v}, \mathbf{u}, \mathbf{w}]$.
 $A_{2,3}(-1, 0, -1), A_{4,3}(-1, -1), A_{4,3}(-1, 0), A_{4,3}(1, 0), A_{4,3}(1, 1), A_{12,3}$.
- I₂₉. Centro-symmetric identity* $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = [\mathbf{w}, \mathbf{v}, \mathbf{u}]$.
 $A_{1,3}(\alpha_1, \alpha_2, -\alpha_1 - 2\alpha_2, 2\alpha_1 + \alpha_2)$, where $\alpha_2 = (\pm\sqrt{2\alpha_1} - \alpha_1 + 1)$,
 $A_{2,3}(-1, 0, -1), A_{4,3}(\alpha_1, -\alpha_1 + \sqrt{-\alpha_1^2 - 1}), A_{4,3}(\alpha_1, -\alpha_1 - \sqrt{-\alpha_1^2 - 1})$, $A_{5,3}(-1), A_5(1), A_{8,3}(i), A_{8,3}(-i)$,
 $A_{12,3}$.
- I₃₀. Centro-anti-symmetric identity* $[\mathbf{u}, \mathbf{v}, \mathbf{w}] = -[\mathbf{w}, \mathbf{v}, \mathbf{u}]$.
 $A_{2,3}(\alpha_1, \beta_1, 1 - \alpha_1) \cong A_{2,3}(\alpha_1, -\beta_1, 1 - \alpha_1), A_{3,3}(\beta_1, 1), A_{4,3}(\alpha_1, 1 - \alpha_1), A_{4,3}(\alpha_1, 2\alpha_1 - 1), A_{9,3}, A_{10,3}$,
 $A_{11,3}, A_{12,3}$.

REFERENCES

- [1] H. Ahmed, U. Bekbaev, I. Rakhimov, Classification of 2-dimensional evolution algebras, their groups of automorphisms and derivation algebras, *arXiv*: 17. (2017).
- [2] H. Ahmed, U. Bekbaev, I. Rakhimov, Complete classification of two-dimensional algebras, 12 pages, *AIP Conf. Proc.*, 1830, 070016, 2017, doi 10.1063/1.4980965.
- [3] H. Ahmed, U. Bekbaev, I. Rakhimov, Classification of two-dimensional Jordan algebras, *AIP Conf. Proc.*, 1905, 030003-1030003-8; <https://doi.org/10.1063/1.5012149>
- [4] H. Ahmed, U. Bekbaev, I. Rakhimov, On Two-Dimensional Power Associative Algebras Over Algebraically Closed Fields and \mathbb{R} , *Lobachevskii Journal of Mathematics*, 2019, 40(1), 1–13.
- [5] H. Ahmed, U. Bekbaev, I. Rakhimov, Classification of two-dimensional Jordan algebras over \mathbb{R} , *Malaysian Journal of Mathematical Sciences*, 2018, 12(3), 287–303
- [6] R. Durán Díaz, J.M. Masqué, A.P. Domínguez, Classifying quadratic maps from plane to plane, *Linear Algebra and Appl.*, 2003, 364, 1–12.
- [7] M. Goze and E. Remm, 2-dimensional algebras, *African Journal of Mathematical Physics*, 10, 2011, 81–91.
- [8] H.P. Petersson, The classification of two-dimensional nonassociative algebras, *Result. Math.*, 3, 2000, 120–154.
- [9] Y.C. Casado, *Evolution algebras*, PhD thesis, Universidad de Mlaga (2016) <http://orcid.org/0000-0003-4299-4392>.
- [10] J.M. Casas, M. Ladra, B.A. Omirov, U.A. Rozikov, On evolution Algebras, *Algebra Colloq.*, 2014, 21, 331–342.
- [11] I. Kaygorodov, Yu. Volkov, The variety of 2-dimensional algebras over an algebraically closed field, *Canadian Journal of Mathematics*, 2019, 71(4), 819–842
- [12] A. Giambruno, S. Mishchenko, M. Zaicev, Codimension growth of two-dimensional non-associative algebras, *Proceedings of the American Mathematical Society*, 2007, 135(11) 3405–3415.