MULTIPLICATIVE LIE DERIVATION OF TRIANGULAR 3-MATRIX RINGS

ZHENHUI CHEN, JINCHUAN HOU

ABSTRACT. A map ϕ on an associative ring is called a multiplicative Lie derivation if $\phi([x,y]) = [\phi(x),y] + [x,\phi(y)]$ holds for any elements x, y, where [x,y] = xy - yx is the Lie product. In the paper, we discuss the multiplicative Lie derivations on the triangular 3-matrix rings $\mathcal{T} = \mathcal{T}_3(\mathcal{R}_i; \mathcal{M}_{ij})$. Under the standard assumption $Q_i \mathcal{Z}(\mathcal{T})Q_i = \mathcal{Z}(Q_i \mathcal{T}Q_i)$, i = 1, 2, 3, we show that every multiplicative Lie derivation $\varphi : \mathcal{T} \to \mathcal{T}$ has the standard form $\varphi = \delta + \gamma$ with δ a derivation and γ a center valued map vanishing each commutator.

1. INTRODUCTION

Let R be an associative ring. Let [x, y] = xy - yx denote the Lie product of $x, y \in R$. An additive map φ of R is called a derivation if $\varphi(xy) = \varphi(x)y + x\varphi(y)$ for all $x, y \in R$; $\varphi : R \to R$ is called a Lie derivation if $\varphi([x,y]) = [\varphi(x),y] + [x,\varphi(y)]$ for all $x,y \in R$. Clearly, each derivation is a Lie derivation. The Lie derivations on various kinds of rings and Algebras have been studied intensively (see [6, 12, 1, 4, 13, 15, 10, 5]). Cheung in [6]described the form of Lie derivation on the triangular algebra. In [13], Lu and Jing proved that if $\delta: B(X) \to B(X)$ is a linear map satisfying $\delta([A, B]) = [\delta(A), B] + [A, \delta(B)]$ for any $A, B \in B(X)$ with AB = 0, then δ can be decomposed as $d + \tau$, where d is a derivation of B(X) and $\tau: B(X) \to \mathbb{C}I$ is a linear map vanishing at commutators [A, B] with AB = 0. In paper [7], Du and Wang gave a description of Lie derivations of generalized matrix algebras. In [14], Qi and Hou characterized Lie derivations on a von Neumann algebra M without central summands of type I_1 . More generally, it was shown in [14] that, for any scalar ξ , a additive map L on M satisfies $L(AB - \xi BA) = L(A)B - \xi BL(A) + AL(B) - \xi L(B)A$ whenever AB = 0if and only if there exists an additive derivation φ such that, (1) if $\xi = 1$, then $L = \varphi + f$, where f is an additive map from M into its center vanishing on [A, B] with AB = 0; (2) if $\xi = 0$, then $L(I) \in Z(M)$ and $L(A) = \varphi(A) + L(I)A$ for all A; (3) if ξ is rational and $\xi \neq 0, 1, \xi$ then $L = \varphi$; (4) if ξ is not rational, then $\varphi(\xi I) = \xi L(I)$ and $L(A) = \varphi(A) + L(I)A$. Benkovič in [2] proved that, under certain conditions, each generalized Lie derivation of a triangular algebra \mathcal{A} is the sum of a generalized Lie derivation and a central map which vanishes on all commutators of \mathcal{A} .

In some situation, the assumption of the linearity or additivity on maps may be omitted. Thus the notion of Lie derivations was generalized to that of multiplicative Lie derivations, and more generally, multiplicative Lie *n*-derivations. Let us recall the following sequence of polynomials: $P_1(x) = x$ and $P_n(x_1, x_2, \dots, x_n) = [P_{n-1}(x_1, x_2, \dots, x_{n-1}), x_n]$ for all integers

²⁰¹⁰ Mathematics Subject Classification. 16W25; 15A78; 47B47.

Key words and phrases. Rings; triangular 3-matrix rings; derivations; multiplicative Lie derivations.

This work is partially supported by National Natural Science Foundation of China (11671294).

ZHENHUI CHEN, JINCHUAN HOU

 $n \ge 2$. Thus, $P_2(x_1, x_2) = [x_1, x_2], P_3(x_1, x_2, x_3) = [[x_1, x_2], x_3], \dots$ Let $n \ge 2$ be an integer. A map (no additivity is assumed) $\varphi : R \to R$ is called a multiplicative Lie *n*-derivation if

$$\varphi(P_n(x_1, x_2, \cdots, x_n)) = \sum_{i=1}^n P_n(x_1, \cdots, x_{i-1}, \varphi(x_i), x_{i+1}, \cdots, x_n)$$

holds for all $x_1, x_2, \ldots, x_n \in \mathcal{R}$. Particularly, a multiplicative Lie 2-derivation is called a multiplicative Lie derivation.

Let \mathcal{A}, \mathcal{B} be unital rings (res. algebras over a commutative unital ring \mathcal{R}) and \mathcal{M} be a $(\mathcal{A}, \mathcal{B})$ -bimodule, which is faithful as a left \mathcal{A} -module and also as a right \mathcal{B} -module; that is, for any $A \in \mathcal{A}$ and $B \in \mathcal{B}, A\mathcal{M} = \mathcal{M}B = \{0\}$ imply A = 0 and B = 0. Recall that the associative ring (res. algebra over \mathcal{R})

$$\mathcal{U} = \operatorname{Tri}(\mathcal{A}, \mathcal{M}, \mathcal{B}) = \left\{ \left(\begin{array}{cc} A & M \\ 0 & B \end{array} \right) : A \in \mathcal{A}, M \in \mathcal{M}, B \in \mathcal{B} \right\}$$

under the usual matrix operations will be called a triangular ring (res. algebra). In [17], Yu and Zhang discussed the multiplicative Lie derivations on a triangular algebra $\mathcal{U} = \text{Tri}(\mathcal{A}, \mathcal{M}, \mathcal{B})$ over a commutative ring. Under a standard assumption

$$\pi_{\mathcal{A}}(\mathcal{Z}(\mathcal{U})) = \mathcal{Z}(\mathcal{A}) \text{ and } \pi_{\mathcal{B}}(\mathcal{Z}(\mathcal{U})) = \mathcal{Z}(\mathcal{B}),$$
 (1.1)

they showed that every multiplicative Lie derivation $\varphi : \mathcal{U} \to \mathcal{U}$ has the standard form $\varphi = \delta + \gamma$, where δ is a derivation and γ is a center valuated map sending each commutator to zero. Here $\pi_{\mathcal{A}}$ and $\pi_{\mathcal{B}}$ are projections defined respectively by $\pi_{\mathcal{A}}\begin{pmatrix} A & M \\ 0 & B \end{pmatrix} = A$ and

$$\pi_{\mathcal{B}}\begin{pmatrix} A & M \\ 0 & B \end{pmatrix}) = B.$$

More generally, under the assumptions that $\mathcal{U} = \operatorname{Tri}(A, M, B)$ is (n-1)-torsion free satisfying Eq.(1.1), and, in addition, for the case $n \geq 3$, \mathcal{A} or \mathcal{B} has the property that $[a, \mathcal{R}] \in \mathcal{Z}(\mathcal{R}) \Rightarrow a \in \mathcal{Z}(\mathcal{R})$, Benkovič and Eremita [3] proved that every multiplicative Lie *n*-derivation $\varphi : \mathcal{T} \to \mathcal{U}$ has the standard form $\varphi = \delta + \gamma$. In [16], Yao Wang and Yu Wang discussed the same problem for multiplicative Lie *n*-derivations on a certain class of generalized matrix algebras.

As a generalization of the notion of triangular rings, for each positive integer $k \ge 2$, in [8], Ferreira introduced a new class of rings which called triangular k-matrix rings. The notion of triangular rings coincides with the notion of triangular 2-matrix rings. The works of [17, 3, 16] motivate us to discuss the problem: for $m \ge 3$, under what conditions, every multiplicative Lie n-derivation on a triangular m-matrix rings will have the standard form? We are not able to solve the problem for general m in the present paper. As a start, we show that, under a standard assumption, every multiplicative Lie derivation on a triangular 3-matrix ring has the standard form. But our method used in this paper is not valid for the case $m \ge 4$.

The paper is organized as follows. In section 2 we recall the notion of triangular 3-matrix rings, provide a kind of the triangular 3-matrix rings that are not triangular rings. The main result, including a corollary, is also given in this section. Section 3 is devoted to discussing the general structural properties of the multiplicative Lie derivations. We show that, under certain more relaxed assumption on the triangular 3-matrix ring, every multiplicative Lie

derivation φ has the form $\varphi = \delta + \gamma + \xi$, where δ is a derivation, γ is a center valued map vanishing each commutator and ξ is a multiplicative Lie derivation with a very small range. Based on the results in Section 3, we give our proof of the main result in Section 4.

2. Main result

Though, for each $k \ge 2$, triangular k-matrix rings was introduced in [8], we only recall the definition of triangular 3-matrix rings here.

Let $\mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3$ be unital rings and \mathcal{M}_{ij} be $(\mathcal{R}_i, \mathcal{R}_j)$ -bimodules with $\mathcal{M}_{ii} = \mathcal{R}_i$ for all $1 \leq i \leq j \leq 3$. Let $\varphi_{ijk} : \mathcal{M}_{ij \bigotimes \mathcal{R}_j} \mathcal{M}_{jk} \to \mathcal{M}_{ik}$ be $(\mathcal{R}_i, \mathcal{R}_k)$ -bimodules homomorphisms with $\varphi_{iij} : \mathcal{R}_i \otimes_{\mathcal{R}_i} \mathcal{M}_{ij} \to \mathcal{M}_{ij}$ and $\varphi_{ijj} : \mathcal{M}_{ij} \otimes_{\mathcal{R}_j} \mathcal{R}_j \to \mathcal{M}_{ij}$ the canonical multiplicative maps for all $1 \leq i \leq j \leq 3$. Write $ab = \varphi_{ijk}(a \otimes b)$ for $a \in \mathcal{M}_{ij}, b \in \mathcal{M}_{jk}$. Assume that \mathcal{M}_{ij} is faithful as a left \mathcal{R}_i -module and faithful as a right \mathcal{R}_j -module for all $1 \leq i < j \leq 3$. Let $\mathcal{T} = \mathcal{T}_3(\mathcal{R}_i; \mathcal{M}_{ij})$ be the set

$$\mathcal{T} = \mathcal{T}_3(\mathcal{R}_i; \mathcal{M}_{ij}) = \{ \begin{pmatrix} r_{11} & m_{12} & m_{13} \\ 0 & r_{22} & m_{23} \\ 0 & 0 & r_{33} \end{pmatrix} : r_{ii} \in R_i, m_{ij} \in M_{ij}, 1 \le i < j \le 3 \}.$$
(2.1)

Clearly, with the matrix operations of addition and multiplication, \mathcal{T} is a ring which is called a triangular 3-matrix ring. Note that, for m > 3, the additional assumption is needed so that $\mathcal{T}_m(\mathcal{R}_i : \mathcal{M}_{i,j})$ to be a ring: a(bc) = (ab)c for all $a \in \mathcal{M}_{ik}, b \in \mathcal{M}_{kl}$ and $c \in \mathcal{M}_{lj}$ with $1 \le i \le k \le l \le j \le m$ (Ref. [8]).

Obviously, setting

$$\mathcal{T}_{ij} = \{ (m_{kt}) : m_{kt} = \begin{cases} m_{ij}, & \text{if } (k,t) = (i,j) \\ 0 & \text{if } (k,t) \neq (i,j) \end{cases}, \ 1 \le i \le j \le 3 \} \subset \mathcal{T},$$

we can write $\mathcal{T} = \bigoplus_{1 \leq i \leq j \leq 3} \mathcal{T}_{ij}$. We also can identify \mathcal{T}_{ij} with \mathcal{M}_{ij} . Obviously, for any element $a_{ij} \in \mathcal{T}_{ij}$, we have $a_{ij}a_{kj} = 0$ whenever $j \neq k$.

Some canonical examples of triangular 3-matrix ring are upper triangular matrix rings $\mathcal{T}_n(\mathcal{R})$ with $n \geq 3$ over a unital associative ring \mathcal{R} and the nest algebras with the nest containing more than 2 nontrivial elements.

It is easily seen that a triangular ring may not be a triangular 3-matrix ring. Conversely, the following example shows that a triangular 3-matrix ring may not be a triangular ring.

Example 2.1 Let \mathcal{R} be a unital associative ring with the unit 1 and $M_6(\mathcal{R})$ be the ring of all 6×6 matrices over \mathcal{R} . Let $\mathcal{A} \subseteq M_6(\mathcal{R})$ be the subset

$$\mathcal{A} = \left\{ \begin{pmatrix} a_{11} & a_{12} & 0 & a_{14} & 0 & a_{16} \\ a_{21} & a_{22} & 0 & a_{24} & 0 & a_{26} \\ 0 & 0 & a & 0 & 0 & a_{36} \\ 0 & 0 & 0 & a & 0 & 0 \\ 0 & 0 & 0 & 0 & b & 0 \\ 0 & 0 & 0 & 0 & b & 0 \end{pmatrix} : a, b, a_{ij} \in \mathcal{R} \right\}.$$

It is easily checked that \mathcal{A} is a subring of $M_6(\mathcal{R})$.

We claim that \mathcal{A} is a triangular 3-matrix ring. To see this, let $\mathcal{A}_1 = \mathcal{M}_2(\mathcal{R}), \ \mathcal{A}_2 = \mathcal{A}_3 = \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} : a \in \mathcal{R} \right\}, \ \mathcal{M}_{12} = \mathcal{M}_{13} = \left\{ \begin{pmatrix} 0 & c \\ 0 & d \end{pmatrix} : c, d \in \mathcal{R} \right\} \text{ and } \mathcal{M}_{23} = \left\{ \begin{pmatrix} 0 & c \\ 0 & 0 \end{pmatrix} : c \in \mathcal{R} \right\};$ then $\mathcal{A} = \mathcal{T}_3(\mathcal{A}_i; \mathcal{M}_{ij})$. In fact, it is obvious that \mathcal{A}_i s are unital rings and \mathcal{M}_{ij} s are $(\mathcal{A}_i, \mathcal{A}_j)$ bimodules. We check that \mathcal{M}_{ij} is faithful as a left \mathcal{A}_i -module and as a right \mathcal{A}_j -module. As $\mathcal{A}_2 = \mathcal{A}_3$ and $\mathcal{M}_{12} = \mathcal{M}_{13}$, we need only to check this for (i, j) = (1, 2) and (i, j) = (2, 3).

Given any
$$A_1 = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \in \mathcal{A}_1$$
. If $A_1 M_{12} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} 0 & c \\ 0 & d \end{pmatrix} = 0$ holds for
every $M_{12} = \begin{pmatrix} 0 & c \\ 0 & d \end{pmatrix} \in \mathcal{M}_{12}$, then $a_{ij} = 0$ holds for all $1 \le i, j \le 2$ by taking $(c, d) = (1, 0)$
or $(0, 1)$, which gives $A_1 = 0$. So \mathcal{M}_{12} is faithful as a left \mathcal{A}_1 -module. It is obvious that \mathcal{M}_{12}
is faithful as a right \mathcal{A}_2 -module, and \mathcal{M}_{23} is faithful. Hence, \mathcal{A} is a triangular 3-matrix ring.

We assert that \mathcal{A} is not a triangular ring.

Clearly, one can not regard \mathcal{A} as a triangular ring $\operatorname{Tri}(\mathcal{B}_1, \mathcal{M}, \mathcal{B}_2)$ by (3+3)-partition, that is, by letting

$$\mathcal{B}_{1} = \left\{ \begin{pmatrix} a_{11} & a_{12} & 0\\ a_{21} & a_{22} & 0\\ 0 & 0 & a \end{pmatrix} : a, a_{ij} \in \mathcal{R} \right\}$$
$$\mathcal{M} = \left\{ \begin{pmatrix} a_{14} & 0 & a_{16}\\ a_{24} & 0 & a_{26}\\ 0 & 0 & a_{36} \end{pmatrix} : a_{ij} \in \mathcal{R} \right\}$$

and

$$\mathcal{B}_2 = \{ \left(egin{array}{ccc} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & b \end{array}
ight) : a, b \in \mathcal{R} \}.$$

In fact, \mathcal{A} is a proper subring of $\text{Tri}(\mathcal{B}_1, \mathcal{M}, \mathcal{B}_2)$ which contains elements $x = (a_{ij})$ so that $a_{33} \neq a_{44}$. For the same reason, \mathcal{A} is not a triangular ring by (5+1)-partition.

Thus there are two possible ways, that is, (2+4)-partition and (4+2)-partition, that might make \mathcal{A} into a triangular ring.

Way 1. $\mathcal{A} = \text{Tri}(\mathcal{B}, \mathcal{M}, \mathcal{A}_3)$, where $\mathcal{B} = \text{Tri}(\mathcal{A}_1, \mathcal{M}_{12}, \mathcal{A}_2)$ and

$$\mathcal{M} = \{ \begin{pmatrix} 0 & b \\ 0 & c \\ 0 & d \\ 0 & 0 \end{pmatrix} : b, c, d \in \mathcal{R} \}.$$

However, it is easily seen that \mathcal{M} is not faithful as a left \mathcal{B} -module by taking

$$B = \begin{pmatrix} a_{11} & a_{12} & 0 & 1\\ a_{21} & a_{22} & 0 & a_{24}\\ 0 & 0 & a & 0\\ 0 & 0 & 0 & a \end{pmatrix} \in \mathcal{B}.$$

Way 2. $\mathcal{A} = \text{Tri}(\mathcal{A}_1, \mathcal{N}, \mathcal{C})$, where $\mathcal{C} = \text{Tri}(\mathcal{A}_2, \mathcal{M}_{23}, \mathcal{A}_3)$ and

$$\mathcal{N} = \left\{ \left(\begin{array}{ccc} 0 & b & 0 & d \\ 0 & c & 0 & e \end{array} \right) : b, c, d, e \in \mathcal{R} \right\}.$$

But, by taking

$$C = \begin{pmatrix} a & 0 & 0 & 1 \\ 0 & a & 0 & 0 \\ 0 & 0 & a' & 0 \\ 0 & 0 & 0 & a' \end{pmatrix} \in \mathcal{C}$$

we see that \mathcal{N} is not faithful as a right \mathcal{C} -module.

Therefore, \mathcal{A} is a triangular 3-matrix ring but not a triangular ring.

Before the statement of the main result, we introduce more notations.

For any ring \mathcal{R} , $\mathcal{Z}(\mathcal{R})$ stands for the center of \mathcal{R} ; that is, $\mathcal{Z}(\mathcal{R}) = \{Z : Z \in \mathcal{R} \text{ and } [Z, \mathcal{R}] = 0\}$. Let $\mathcal{T} = \mathcal{T}_3(\mathcal{R}_i; \mathcal{M}_{ij})$ be a triangular 3-matrix ring as defined in Eq.(2.1). There are three standard idempotent elements in \mathcal{T} :

$$Q_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad Q_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad Q_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

 \mathcal{T} is unital with the unit $I = Q_1 + Q_2 + Q_3$. Denote by $\mathcal{T}_{ij} = Q_i \mathcal{T} Q_j$, i, j = 1, 2, 3. Clearly, $\mathcal{T}_{ij} = 0$ if i > j, $\mathcal{T}_i = \mathcal{T}_{ii} \cong \mathcal{R}_i$ and $\mathcal{T}_{ij} \cong \mathcal{M}_{ij}$ if i < j. Moreover, $\mathcal{T} = \sum_{1 \le i \le j \le 3} \mathcal{T}_{ij}$. In this paper, if no confusion occurs, we identify \mathcal{T}_i with \mathcal{R}_i and \mathcal{T}_{ij} with \mathcal{M}_{ij} .

Theorem 2.2. Let $\mathcal{T} = \mathcal{T}_3(\mathcal{R}_i; \mathcal{M}_{ij})$ be a triangular 3-matrix ring. Assume that $Q_i \mathcal{Z}(\mathcal{T})Q_i = \mathcal{Z}(\mathcal{T}_i)$, i = 1, 2, 3. Then each multiplicative Lie derivation $\varphi : \mathcal{T} \to \mathcal{T}$ has the standard form

$$\varphi = \delta + \gamma,$$

where $\delta: \mathcal{T} \to \mathcal{T}$ is a derivation, $\gamma: \mathcal{T} \to \mathcal{Z}(\mathcal{T})$ is a center valued map such that $\gamma([\mathcal{T}, \mathcal{T}]) = 0$.

A proofs of Theorem 2.2 will be given in Section 4.

Corollary 2.3 Let $\mathcal{A} = T_3(\mathcal{A}_i; \mathcal{M}_{ij})$ be the triangular 3-matrix ring constructed from a ring \mathcal{R} as in Example 2.1. Then every multiplicative Lie derivation $\varphi : \mathcal{A} \to \mathcal{A}$ has the standard form.

Proof. It is obvious that $\mathcal{Z}(\mathcal{A}_i) = \{ \begin{pmatrix} z & 0 \\ 0 & z \end{pmatrix} : z \in \mathcal{Z}(\mathcal{R}) \}$ for i = 1, 2, 3. It is also easily checked that

$$\mathcal{Z}(\mathcal{A}) = \{ \operatorname{diag}(z, z, z, z, z, z) : z \in \mathcal{Z}(\mathcal{R}) \}.$$

As $Q_1 = \text{diag}(1, 1, 0, 0, 0, 0)$, $Q_2 = \text{diag}(0, 0, 1, 1, 0, 0)$ and $Q_3 = \text{diag}(0, 0, 0, 0, 1, 1)$, we see that $Q_i \mathcal{Z}(\mathcal{A})Q_i = \mathcal{Z}(Q_i \mathcal{A}Q_i)$, i = 1, 2, 3. Hence \mathcal{A} satisfies the hypotheses of Theorem 2.2 and the corollary follows.

3. Structure of multiplicative Lie derivations on triangular 3-matrix rings

To prove Theorem 2.2, in this section, we consider the structure of multiplicative Lie derivations on triangular 3-matrix rings in some more general situation.

Denote by \mathcal{T} the triangular 3-matrix ring $\mathcal{T}_3(\mathcal{R}_i; \mathcal{M}_{ij})$. It is easily checked that the center of \mathcal{T} is the set

$$\mathcal{Z}(\mathcal{T}) = \{ \text{diag}(r_{11}, r_{22}, r_{33}) : r_{ii} \in \mathcal{R}_i, \ r_{ii}m_{ij} = m_{ij}r_{jj} \ \forall m_{ij} \in \mathcal{M}_{ij}, 1 \le i < j \le 3 \}.$$
(3.1)

Let \mathcal{T} be the triangular 3-matrix ring. It is clear from Eq.(3.1) that, for any $x \in \mathcal{T}$,

$$[x,\mathcal{T}] \subseteq \mathcal{Z}(\mathcal{T}) \Rightarrow x \in \mathcal{Z}(\mathcal{T}).$$

Denote by $Q'_i = I - Q_i$, i = 1, 2, 3. Write $\mathcal{T}_i = Q_i \mathcal{T} Q_i$, $\mathcal{T}'_i = Q'_i \mathcal{T} Q'_i$, i = 1, 2, 3. Also write $\mathcal{M}_1 = Q_1 \mathcal{T} Q'_1$, $\mathcal{M}_3 = Q'_3 \mathcal{T} Q_3$. Obviously, $\mathcal{T} = \mathcal{T}_1 + \mathcal{M}_1 + \mathcal{T}'_1 = \mathcal{T}'_3 + \mathcal{M}_3 + \mathcal{T}_3$.

Denote by $\mathcal{V}_{23} = \{w_{23} \in \mathcal{M}_{23} : m_{12}w_{23} = 0 \ \forall m_{12} \in \mathcal{M}_{12}\}$ and $\mathcal{V}_{12} = \{w_{12} \in \mathcal{M}_{12} : w_{12}m_{23} = 0 \ \forall m_{23} \in \mathcal{M}_{23}\}$. Also, denote by

$$S_1 = \left\{ \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & w_{23} \\ 0 & 0 & 0 \end{pmatrix} : w_{23} \in \mathcal{V}_{23} \right\}$$

and

$$\mathcal{S}_3 = \{ \begin{pmatrix} 0 & w_{12} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} : w_{12} \in \mathcal{V}_{12} \}.$$

The following is our main result in this section which is needed to prove Theorem 2.2.

Theorem 3.1. Let $\mathcal{T} = \mathcal{T}_3(\mathcal{R}_i; \mathcal{M}_{ij})$ be a triangular 3-matrix ring and let $k \in \{1, 3\}$. Assume that $Q_k \mathcal{Z}(\mathcal{T})Q_k = \mathcal{Z}(\mathcal{T}_k)$ and $Q'_k \mathcal{Z}(\mathcal{T})Q'_k = \mathcal{Z}(\mathcal{T}'_k)$. If every multiplicative Lie derivation from \mathcal{T}'_k into itself has the standard form, then each multiplicative Lie derivation $\varphi : \mathcal{T} \to \mathcal{T}$ has the form

$$\varphi = \delta + \gamma + \xi,$$

where $\delta : \mathcal{T} \to \mathcal{T}$ is a derivation, $\gamma : \mathcal{T} \to \mathcal{Z}(\mathcal{T})$ is a center valued map such that $\gamma([\mathcal{T}, \mathcal{T}]) = 0$ and $\xi : \mathcal{T} \to S_k$ is a multiplicative Lie derivation.

We will give the details of the proof of Theorem 3.1 for the case k = 1. The case k = 3 is treated with similarly.

3.1. Proof of Theorem 3.1: Case k = 1. In this case, $\mathcal{T} = \mathcal{T}_1 + \mathcal{M}_1 + \mathcal{T}'_1$ and we have $Q_1 \mathcal{Z}(\mathcal{T}) Q_1 = \mathcal{Z}(\mathcal{T}_1)$ and $Q'_1 \mathcal{Z}(\mathcal{T}) Q'_1 = \mathcal{Z}(\mathcal{T}'_1)$.

For an arbitrary element $x \in \mathcal{T}$, one can then write x = a + m + b with $a \in \mathcal{T}_1, m \in \mathcal{M}_1$ and $b \in \mathcal{T}'_1$. Let us define projections $\pi_{\mathcal{T}_1} : \mathcal{T} \to \mathcal{T}_1$ and $\pi_{\mathcal{T}'_1} : \mathcal{T} \to \mathcal{T}'_1$ by $\pi_{\mathcal{T}_1}(a + m + b) = a$ and $\pi_{\mathcal{T}'_1}(a + m + b) = b$. Obviously, $\pi_{\mathcal{T}_1}(\mathcal{Z}(\mathcal{T})) \subseteq \mathcal{Z}(\mathcal{T}_1)$ and $\pi_{\mathcal{T}'_1}(\mathcal{Z}(\mathcal{T})) \subseteq \mathcal{Z}(\mathcal{T}'_1)$, and, there exists a unique ring isomorphism $\tau : \pi_{\mathcal{T}_1}(\mathcal{Z}(\mathcal{T})) \to \pi_{\mathcal{T}'_1}(\mathcal{Z}(\mathcal{T}))$ such that $am = m\tau(a)$ for all $m \in \mathcal{M}_1, a \in \pi_{\mathcal{T}_1}(\mathcal{Z}(\mathcal{T}))$.

In the sequel we assume that $\varphi : \mathcal{T} \to \mathcal{T}$ is a multiplicative Lie derivation. We prove Theorem 3.1 for the case k = 1 by a series lemmas.

The first lemma is an analogue of [3, Lemma 3.1].

Lemma 3.2. There exist an inner derivation $d_1 : \mathcal{T} - \mathcal{T}$ and a multiplicative Lie derivation $\varphi_1 : \mathcal{T} \to \mathcal{T}$ such that

$$\varphi = d_1 + \varphi_1$$
 and $Q_1 \varphi_1(Q'_1) Q'_1 = 0.$

Proof. We define maps $d_1, \varphi_1 : \mathcal{T} - \mathcal{T}$ by $d_1(x) = [\varphi(Q'_1), x]$ and $\varphi_1(x) = \varphi(x) - d_1(x)$. Obviously, d_1 is an inner derivation and φ_1 is a multiplicative Lie derivation. Since

$$\begin{aligned} \varphi_1(Q'_1) &= \varphi(Q'_1) - d_1(Q'_1) \\ &= \varphi(Q'_1) - [\varphi(Q'_1), Q'_1] \\ &= \varphi(Q'_1) - \varphi(Q'_1)Q'_1 + Q'_1\varphi(Q'_1), \end{aligned}$$

multiplying by Q_1 from the left and by Q'_1 from the right of the equation, we get $Q_1\varphi_1(Q'_1)Q'_1 = 0$.

Thus, in Lemmas 3.3-3.5 we assume that $\varphi_1 : \mathcal{T} \to \mathcal{T}$ is a multiplicative Lie derivations satisfying $Q_1 \varphi_1(Q'_1) Q'_1 = 0$

Lemma 3.3. For any $a \in \mathcal{T}_1, b \in \mathcal{T}'_1$ and $m \in \mathcal{M}_1$, the following statements are true: (a) $\varphi_1(Q_1), \ \varphi_1(Q'_1) \in \mathcal{R}_1 \oplus \mathcal{R}_2 \oplus \mathcal{R}_3$. (b) $Q_1\varphi_1(a)Q'_1 = Q_1\varphi_1(b)Q'_1 = 0$.

(c) $\varphi_1(a) = Q_1\varphi_1(a)Q_1 + Q'_1\varphi_1(a)Q'_1$, $\varphi_1(b) = Q_1\varphi_1(b)Q_1 + Q'_1\varphi_1(b)Q'_1$ and $\varphi_1(m) = Q_1\varphi_1(m)Q'_1$.

(d) $Q_1\varphi_1(\mathcal{T}'_1)Q_1 \subseteq \mathcal{Z}(\mathcal{T}_1)$ and $Q'_1\varphi_1(\mathcal{T}_1)Q'_1 \subseteq \mathcal{Z}(\mathcal{T}'_1)$.

Proof. It is obvious that $\varphi(0) = 0$ and $\varphi(I) \in \mathcal{Z}(\mathcal{T})$. (a) Since $Q_1\varphi_1(Q'_1)Q'_1 = 0$,

$$\varphi_1(Q_1') \in \left\{ \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} : a_{ii} \in \mathcal{R}_i (i = 1, 2, 3), a_{23} \in \mathcal{M}_{23} \right\}$$

Note that

$$0 = \varphi([Q_1, Q'_1]) = \varphi(Q_1)Q'_1 - Q'_1\varphi(Q_1) + Q_1\varphi(Q'_1) - \varphi(Q'_1)Q_1.$$
(3.2)

By Lemma 3.2, we have

$$\begin{aligned}
\varphi_1(Q_1) &= \varphi(Q_1) - d_1(Q_1) \\
&= \varphi(Q_1) - [\varphi(Q'_1), Q_1] \\
&= \varphi(Q_1) - (\varphi(Q'_1)Q_1 - Q_1\varphi(Q'_1)).
\end{aligned}$$
(3.3)

Thus Eq.(3.2) together with Eq.(3.3) gives

$$\varphi_1(Q_1) = \varphi(Q_1) - (\varphi(Q_1)Q_1' - Q_1'\varphi(Q_1)).$$

Writing

$$\varphi(Q_1) = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix},$$

the above formula leads to

$$\begin{split} \varphi_{1}(Q_{1}) &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} - \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &+ \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} \\ &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} - \begin{pmatrix} 0 & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} \\ &= \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} . \end{split}$$

Therefore

$$\varphi_1(Q_1) \in \left\{ \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} : a_{ii} \in \mathcal{R}_i (i = 1, 2, 3), a_{23} \in \mathcal{M}_{23} \right\}.$$

Similarly one can check that

$$\varphi_1(Q_2) \in \{ \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} : a_{ii} \in \mathcal{R}_i (i = 1, 2, 3), a_{23} \in \mathcal{M}_{23} \}.$$

So we can write

$$\varphi_1(Q_1) = \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix}, \quad \varphi_1(Q_2) = \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix}$$

and

$$\varphi_1(Q_1') = \begin{pmatrix} c_{11} & 0 & 0\\ 0 & c_{22} & c_{23}\\ 0 & 0 & c_{33} \end{pmatrix}.$$

Since

$$\begin{array}{l} 0 = & \varphi_1([Q_1,Q_2]) \\ = & \varphi_1(Q_1)Q_2 - Q_2\varphi_1(Q_1) + Q_1\varphi_1(Q_2) - \varphi_1(Q_2)Q_1 \\ = & \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & b_{33} \end{pmatrix} \\ & + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix} - \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ = & \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ = & \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -a_{23} \\ 0 & 0 & 0 \end{pmatrix}, \end{array}$$

we have $a_{23} = 0$ and thus

$$\varphi_1(Q_1) = \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix} \in \mathcal{R}_1 \oplus \mathcal{R}_2 \oplus \mathcal{R}_3.$$

Similarly

$$\begin{array}{ll} 0 = & \varphi_1([Q_1',Q_2]) \\ = & \varphi_1(Q_1')Q_2 - Q_2\varphi_1(Q_1') + Q_1'\varphi_1(Q_2) - \varphi_1(Q_2)Q_1' \\ = & \begin{pmatrix} c_{11} & 0 & 0 \\ 0 & c_{22} & c_{23} \\ 0 & 0 & c_{33} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} c_{11} & 0 & 0 \\ 0 & c_{22} & c_{23} \\ 0 & 0 & c_{33} \end{pmatrix} \\ & + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix} - \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ & = & \begin{pmatrix} c_{11} & 0 & 0 \\ 0 & c_{22} & 0 \\ 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} c_{11} & 0 & 0 \\ 0 & c_{22} & c_{23} \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix} - \begin{pmatrix} 0 & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix} \end{pmatrix} \\ & = & \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -c_{23} \\ 0 & 0 & 0 \end{pmatrix} \end{array}$$

entails $c_{23} = 0$. Hence

$$\varphi_1(Q_1') = \begin{pmatrix} c_{11} & 0 & 0 \\ 0 & c_{22} & 0 \\ 0 & 0 & c_{33} \end{pmatrix} \in \mathcal{R}_1 \oplus \mathcal{R}_2 \oplus \mathcal{R}_3,$$

too.

(b) Let $x \in \mathcal{T}_1 \bigcup \mathcal{T}'_1$. Then $[x, Q'_1] = Q_1 x Q'_1 = 0$. Since $\varphi_1(0) = 0$, we have

$$0 = \varphi_1(Q_1 x Q'_1) = \varphi_1([x, Q'_1]) = [\varphi_1(x), Q'_1] + [x, \varphi_1(Q'_1)] = Q_1 \varphi_1(x) Q'_1 + Q_1[x, \varphi_1(Q'_1)] Q'_1.$$
(3.4)

However the property $Q_1\varphi_1(Q_1')Q_1'=0$ implies that

$$Q_{1}[x, \varphi_{1}(Q'_{1})]Q'_{1}$$

$$= Q_{1}(x\varphi_{1}(Q'_{1}) - \varphi_{1}(Q'_{1})x)Q'_{1}$$

$$= Q_{1}(Q_{1}xQ_{1}\varphi_{1}(Q'_{1}) - \varphi_{1}(Q'_{1})Q'_{1}xQ_{1})Q'_{1}$$

$$= 0$$

holds for every $x \in \mathcal{T}_1$. So $Q_1[\mathcal{T}_1, \varphi_1(Q'_1)]Q'_1 = 0$. Similarly, we have $Q_1[\mathcal{T}'_1, \varphi_1(Q'_1)]Q'_1 = 0$. So by Eq.(3.4) we see that $Q_1\varphi_1(x)Q'_1 = 0$ for all $x \in \mathcal{T}_1 \bigcup \mathcal{T}'_1$; That is, (b) is true.

(c) By (b) it is clear that $\varphi_1(a) = Q_1\varphi_1(a)Q_1 + Q'_1\varphi_1(a)Q'_1$ and $\varphi_1(b) = Q_1\varphi_1(b)Q_1 + Q'_1\varphi_1(b)Q'_1$ are true respectively for every $a \in \mathcal{T}_1$ and $b \in \mathcal{T}'_1$. Now, for each $m \in \mathcal{M}_1$, we have

$$\begin{aligned} \varphi_1(m) &= \varphi_1[m, Q_1'] \\ &= [\varphi_1(m), Q_1'] + [m, \varphi_1(Q_1')] \\ &= \varphi_1(m)Q_1' - Q_1'\varphi_1(m) + [m, \varphi_1(Q_1')]. \end{aligned}$$

Multiplying by Q_1 from the left hand side and by Q'_1 from the right hand side of the equation, we see that

$$[m, \varphi_1(Q'_1)] = Q_1[m, \varphi_1(Q'_1)]Q'_1 = 0.$$

This implies that $\varphi_1(m) = \varphi_1(m)Q'_1 - Q'_1\varphi_1(m) = Q_1\varphi_1(m)Q'_1$. So (c) is true. (d) Let $a \in \mathcal{T}_1, b \in \mathcal{T}'_1$ and $m \in \mathcal{M}_1$. Since [a, b] = 0, we have

$$[\varphi_1(a), b] + [a, \varphi_1(b)] = \varphi_1([a, b]) = 0.$$

Applying (c), we see that

$$[a, Q_1\varphi_1(b)Q_1] = [a, \varphi_1(b)] = 0$$

and

$$[Q_1'\varphi(a)Q_1',b] = [\varphi_1(a),b] = 0$$

hold for all $a \in \mathcal{T}_1$ and all $b \in \mathcal{T}'_1$. Therefore, $Q_1\varphi(\mathcal{T}'_1)Q_1 \subseteq Z(\mathcal{T}_1)$ and $Q'_1\varphi(\mathcal{T}_1)Q'_1 \subseteq Z(\mathcal{T}'_1)$. This completes the proof of (d).

Furthermore, we have

Lemma 3.4. $\varphi_1(Q_1), \varphi_1(Q'_1) \in \mathcal{Z}(\mathcal{T})$ and $\varphi_1(Q_1xQ'_1) = Q_1\varphi_1(x)Q'_1$ for all $x \in \mathcal{T}$. **Proof.** For any $x \in \mathcal{T}$, we have

$$\varphi_1(Q_1 x Q'_1) = \varphi_1([Q_1, x])
= [\varphi_1(Q_1), x] + [Q_1, \varphi_1(x)].$$
(3.5)

Replacing x by $Q_1 x Q'_1$ in (3.5), we obtain

$$\varphi_1(Q_1 x Q_1') = [\varphi_1(Q_1), Q_1 x Q_1'] + [Q_1, \varphi_1(Q_1 x Q_1')].$$

By (c) one gets $[\varphi_1(Q_1), Q_1 x Q'_1] = 0$, which in turn implies that

$$[\varphi_1(Q_1), \mathcal{M}_1] = 0. \tag{3.6}$$

Meanwhile, by (c), it is clear that

$$[\varphi_1(Q_1), \mathcal{T}_1] = 0, \text{ and } [\varphi_1(Q_1), \mathcal{T}'_1] = 0.$$
 (3.7)

Now (3.6) and (3.7) together ensures that

$$\varphi_1(Q_1) \in \mathcal{Z}(\mathcal{T}).$$

An analogous argument shows that

$$\varphi_1(Q_1') \in \mathcal{Z}(\mathcal{T}),$$

too. Consequently,

$$\begin{aligned}
\varphi_1(Q_1 x Q'_1) &= \varphi_1([Q_1, x]) \\
&= [\varphi_1(Q_1), x] + [Q_1, \varphi_1(x)] \\
&= Q_1 \varphi_1(x) Q'_1
\end{aligned}$$

holds for all $x \in \mathcal{T}$. This establishes the lemma.

Lemma 3.5. For any $a \in \mathcal{T}_1, b \in \mathcal{T}'_1$ and $m \in \mathcal{M}_1$, we have

$$\varphi_1(am) = \varphi_1(a)m + a\varphi_1(m) - m\varphi_1(a)$$

and

$$\varphi_1(mb) = \varphi_1(m)b + m\varphi_1(b) - \varphi_1(b)m.$$

Proof. For any $a \in \mathcal{T}_1, b \in \mathcal{T}'_1$ and $m \in \mathcal{M}_1$, by applying Lemma 3.3, we have

$$\begin{aligned} \varphi_1(am) &= \varphi_1([a,m]) \\ &= [\varphi_1(a),m] + [a,\varphi_1(m)] \\ &= \varphi_1(a)m - m\varphi_1(a) + a\varphi_1(m) - \varphi_1(m)a \\ &= \varphi_1(a)m + a\varphi_1(m) - m\varphi_1(a) \end{aligned}$$

and

$$\varphi_1(mb) = \varphi_1([m, b])$$

= $[\varphi_1(m), b] + [m, \varphi_1(b)]$
= $\varphi_1(m)b - b\varphi_1(m) + m\varphi_1(b) - \varphi_1(b)m$
= $\varphi_1(m)b + m\varphi_1(b) - \varphi_1(b)m$.

So, the lemma is true.

Lemma 3.6. For any $a \in \mathcal{T}_1$, $b \in \mathcal{T}'_1$ and $m \in \mathcal{M}_1$, we have (1) Both $\varphi_1(a+m) - \varphi_1(a) - \varphi_1(m)$ and $\varphi_1(a+b) - \varphi_1(a) - \varphi_1(b) \in \mathcal{Z}(\mathcal{T})$. (2) φ_1 is additive on \mathcal{M}_1 . (3) With $\mathcal{V}_{23} = \{w_{23} \in \mathcal{M}_{23} : \mathcal{M}_{12}w_{23} = 0\}$, we have $\varphi_1(a+m+b) - \varphi_1(a) - \varphi_1(b) - \varphi_1(m)$ $\in \left\{ \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & w_{23} \\ 0 & 0 & a_{33} \end{pmatrix} : w_{23} \in \mathcal{V}_{23}, a_{11} \in \mathcal{Z}(\mathcal{T}_1), a_{11}m_{1j} = m_{1j}a_{jj} \ \forall m_{1j} \in \mathcal{M}_{1j}, j = 2, 3 \}.$

Proof. (1) Let $a \in \mathcal{T}_1$ and $m \in \mathcal{M}_1$. For any $m' \in \mathcal{M}_1$, since [a+m,m'] = [a,m'], we have

$$\varphi_1([a,m']) = \varphi_1([a+m,m']).$$

Due to the fact that $\varphi_1(m) \in \mathcal{M}_1$, we have $[\varphi_1(m), m'] = 0$, and then

$$[\varphi_1(a+m), m'] = [\varphi_1(a), m']$$

 So

$$\varphi_1(a+m) - \varphi_1(a) - \varphi_1(m), m'] = 0 \tag{3.8}$$

holds for all $m' \in \mathcal{M}_1$. Notice that, for any $b' \in \mathcal{T}'_1$,

$$\begin{aligned}
\varphi_1(mb') &= \varphi_1([a+m,b']) \\
&= [\varphi_1(a+m),b'] + [a+m,\varphi_1(b')],
\end{aligned}$$

and, by Lemma 3.5,

$$\varphi_1(mb') = \varphi_1(m)b' + m\varphi_1(b') - \varphi_1(b')m$$

= $[\varphi_1(m), b'] + [m, \varphi_1(b')].$

The above two equations, together with the facts $[a, \varphi_1(b')] = 0$ and $[\varphi_1(a), b'] = 0$ (see Lemma 3.3 (d)), entail that

$$[\varphi_1(a+m) - \varphi_1(m) - \varphi_1(a), b'] = 0$$
(3.9)

holds for all $b' \in \mathcal{T}'_1$.

Next, by Lemma 3.4, we have $Q_1\varphi_1(a+m)Q_1'=\varphi_1(m)$. Thus we may write

$$\varphi_1(a+m) - \varphi_1(a) - \varphi_1(m) = \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix}.$$

Then, by Eq.(3.9),

$$\left(\begin{array}{cc}a_{22}&a_{23}\\0&a_{33}\end{array}\right)\in\mathcal{Z}(\mathcal{T}_1'),$$

which implies that

$$a_{23} = 0 \quad \text{and} \quad a_{22}m_{23} = m_{23}a_{33} \tag{3.10}$$

holds for all $m_{23} \in \mathcal{M}_{23}$ as \mathcal{T}'_1 is a triangular ring.

Now, with

$$m' = \begin{pmatrix} 0 & m_{12} & m_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

by Eq.(3.8), we have

$$0 = \left[\varphi_1(a+m) - \varphi_1(a) - \varphi_1(m), m' \right] \\ = \left(\begin{array}{ccc} 0 & a_{11}m_{12} - m_{12}a_{22} & a_{11}m_{13} - m_{13}a_{33} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right)$$

 So

$$a_{11}m_{12} - m_{12}a_{22} = 0$$
 and $a_{11}m_{13} - m_{13}a_{33} = 0$ (3.11)

hold for all $m_{12} \in \mathcal{M}_{12}$ and all $m_{13} \in \mathcal{M}_{13}$. This, together with Eq.(3.10) and Eq.(3.11) entails that

$$\varphi_1(a+m) - \varphi_1(a) - \varphi_1(m) \in \mathcal{Z}(\mathcal{T})$$

We remark here that, for $b \in \mathcal{T}'_1$ and $m \in \mathcal{M}_1$, though a similar argument by using Lemmas 3.3-3.5 as above gives

$$[\varphi_1(b+m) - \varphi_1(b) - \varphi_1(m), m'] = 0 \quad \forall m' \in \mathcal{M}_1$$

and

$$[\varphi_1(b+m) - \varphi_1(b) - \varphi_1(m), a'] = 0 \quad \forall a' \in \mathcal{T}_1.$$

we cannot drive out $\varphi_1(b+m) - \varphi_1(b) - \varphi_1(m) \in \mathcal{Z}(\mathcal{T})$ at this point.

To show that $\varphi_1(a+b) - \varphi_1(a) - \varphi_1(b) \in \mathcal{Z}(\mathcal{T})$, we need prove (2) first.

(2) By Lemma 3.4, $\varphi_1(Q_1), \varphi_1(Q'_1) \in \mathcal{Z}(\mathcal{T})$. As we just have proved in (1), we have $\varphi_1(Q_1 + m) - \varphi_1(Q_1) - \varphi_1(m) \in \mathcal{Z}(\mathcal{T})$. Therefore,

$$[\varphi_1(Q_1+m), m'+Q_1'] = [\varphi_1(m), m'+Q_1'] = \varphi_1(m)$$
(3.12)

and

$$[Q_1 + m, \varphi_1(m' + Q'_1)] = [Q_1 + m, \varphi_1(m')] = \varphi_1(m')$$
(3.13)

hold for all $m, m' \in \mathcal{M}_1$.

Since $m + m' = [Q_1 + m, m' + Q'_1]$, it follows from Eqs.(3.12) and (3.13) that

$$\varphi_1(m+m') = \varphi_1([Q_1+m,m'+Q'_1]) = [\varphi_1(Q_1+m),m'+Q'_1] + [Q_1+m,\varphi_1(m'+Q'_1)] = \varphi_1(m) + \varphi_1(m')$$

for all $m, m' \in \mathcal{M}_1$. Hence, φ_1 is additive on \mathcal{M}_1 .

Now let us go back to the proof of $\varphi_1(a+b) - \varphi_1(a) - \varphi_1(b) \in \mathcal{Z}(\mathcal{T})$. Let $a \in \mathcal{T}_1, b \in \mathcal{T}'_1$. For any $m \in \mathcal{M}_1$, as [a+b,m] = am - mb, by (2), we have

$$\begin{split} & [\varphi_1(a+b),m] + [a+b,\varphi_1(m)] = \varphi_1([a+b,m]) = \varphi_1(am-mb) \\ & = \varphi_1(am) - \varphi_1(mb) = \varphi_1([a,m]) + \varphi_1([b,m]) \\ & = [\varphi_1(a),m] + [a,\varphi_1(m)] + [\varphi_1(b),m] + [b,\varphi_1(m)]. \end{split}$$

 So

$$[\varphi_1(a+b) - \varphi_1(a) - \varphi_1(b), m] = 0$$
(3.14)

holds for all $m \in \mathcal{M}_1$. Also, for any $b' \in \mathcal{T}'_1$,

_

$$[\varphi_1(a+b), b'] + [a+b, \varphi_1(b')] = \varphi_1([a+b, b'])$$

= $\varphi_1([b, b']) = [\varphi_1(b), b])] + [b, \varphi_1(b')].$

This entails that

$$[\varphi_1(a+b) - \varphi_1(a) - \varphi_1(b), b'] = 0$$
(3.15)

holds for all $b' \in \mathcal{T}'_1$. Similarly one can check that

$$[\varphi_1(a+b) - \varphi_1(a) - \varphi_1(b), a'] = 0$$
(3.16)

holds for all $a' \in \mathcal{T}_1$. Eqs.(3.14)-(3.16) together imply that

$$[\varphi_1(a+b) - \varphi_1(a) - \varphi_1(b), \mathcal{T}] = 0,$$

and hence,

$$\varphi_1(a+b) - \varphi_1(a) - \varphi_1(b) \in \mathcal{Z}(\mathcal{T})$$
(3.17)

holds for all $a \in \mathcal{T}_1$ and $b \in \mathcal{T}'_1$.

(3) Now consider the general case of a + m + b. Note that $Q_1\varphi_1(a + b + m)Q'_1 = \varphi_1(m)$. For any $a' \in \mathcal{T}_1$, since

$$\varphi_1([a+b+m,a']) = [\varphi_1(a+b+m),a'] + [a+b+m,\varphi_1(a')]$$

and

$$\varphi_1([a+b+m,a']) = \varphi_1([a+m,a']) = [\varphi_1(a+m),a'] + [a+m,\varphi_1(a')].$$

we get

$$[\varphi_1(a+b+m) - \varphi_1(a+m), a'] = 0 \quad \forall a' \in \mathcal{T}_1.$$

This, together with $\varphi_1(a+m) - \varphi_1(a) - \varphi_1(m) \in \mathcal{Z}(\mathcal{T})$ by (1) and an obvious fact $[\varphi(b), a'] = 0$ by Lemma 3.3, gives

$$[\varphi_1(a+b+m) - \varphi_1(a) - \varphi_1(b) - \varphi_1(m), a'] = 0, \quad \forall a' \in \mathcal{T}_1.$$
(3.18)

Similarly, for any $m' \in \mathcal{M}_1$,

$$\varphi_1([a+b+m,m']) = [\varphi_1(a+b+m),m'] + [a+b+m,\varphi_1(m')]$$

and

$$\varphi_1([a+b+m,m']) = \varphi_1([a+b,m']) = [\varphi_1(a+b),m'] + [a+b,\varphi_1(m')]$$

leads to

$$[\varphi_1(a+b+m) - \varphi_1(a+b), m'] = 0.$$

Then by Eq.(3.17) proved before we obtain that

$$[\varphi_1(a+b+m) - \varphi_1(a) - \varphi_1(b) - \varphi_1(m), m'] = 0 \quad \forall m' \in \mathcal{M}_1.$$
(3.19)

Write $\varphi_1(a+b+m)$, $\varphi_1(a)$, $\varphi_1(b)$, $\varphi_1(m)$ and m' in the form

$$\varphi_1(a+b+m) = \begin{pmatrix} c_{11} & c_{12} & c_{13} \\ 0 & c_{22} & c_{23} \\ 0 & 0 & c_{33} \end{pmatrix}, \quad \varphi_1(b) = \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix},$$
$$\varphi_1(a) = \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix}, \quad \varphi_1(m) = \begin{pmatrix} 0 & m_{12} & m_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

and

$$m' = \left(\begin{array}{ccc} 0 & m'_{12} & m'_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right).$$

Then

$$\begin{array}{l} \left[\varphi_{1}(a+b+m),m'\right] \\ = & \left(\begin{array}{ccc} c_{11} & c_{12} & c_{13} \\ 0 & c_{22} & c_{23} \\ 0 & 0 & c_{33} \end{array}\right) \left(\begin{array}{ccc} 0 & m'_{12} & m'_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right) - \left(\begin{array}{ccc} 0 & m'_{12} & m'_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right) \left(\begin{array}{ccc} c_{11} & c_{12} & c_{13} \\ 0 & c_{22} & c_{23} \\ 0 & 0 & c_{33} \end{array}\right) \\ = & \left(\begin{array}{ccc} 0 & c_{11}m'_{12} - m'_{12}a_{22} & c_{11}m'_{13} - m'_{13}c_{33} - m'_{12}c_{23} \\ 0 & 0 & 0 \end{array}\right), \end{array}$$

$$\begin{bmatrix} \varphi_1(b), m' \end{bmatrix} \\ = \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix} \begin{pmatrix} 0 & m'_{12} & m'_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & m'_{12} & m'_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} b_{11} & 0 & 0 \\ 0 & b_{22} & b_{23} \\ 0 & 0 & b_{33} \end{pmatrix} \\ = \begin{pmatrix} 0 & b_{11}m'_{12} - m'_{12}b_{22} & b_{11}m'_{13} - m'_{13}b_{33} - m'_{12}b_{23} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

and

$$\begin{bmatrix} \varphi_1(a), m' \end{bmatrix} = \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} \begin{pmatrix} 0 & m'_{12} & m'_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & m'_{12} & m'_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{pmatrix} \\ = \begin{pmatrix} 0 & a_{11}m'_{12} - m'_{12}a_{22} & a_{11}m'_{13} - m'_{13}a_{33} - m'_{12}a_{23} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} .$$

It follows that

$$(c_{11} - a_{11} - b_{11})m'_{12} - m'_{12}(c_{22} - a_{22} - b_{22}) = 0$$

and

$$(c_{11} - a_{11} - b_{11})m'_{13} - m'_{13}(c_{33} - a_{33} - b_{33}) - m'_{12}(c_{23} - a_{23} - b_{23}) = 0$$

hold for all $m'_{12} \in \mathcal{M}_{12}$ and $m'_{13} \in \mathcal{M}_{13}$. Consequently,

$$(c_{11} - a_{11} - b_{11})m'_{12} = m'_{12}(c_{22} - a_{22} - b_{22}),$$
$$(c_{11} - a_{11} - b_{11})m'_{13} = m'_{13}(c_{33} - a_{33} - b_{33})$$

and

 $m_{12}'(c_{23} - a_{23} - b_{23}) = 0$

hold for all $m'_{12}, m'_{13} \in \mathcal{M}_{12}$. Particularly, we have

$$(c_{11}-a_{11}-b_{11})\in \mathcal{Z}(\mathcal{T}_1).$$

Also, by Eq.(3.18) we see that

$$m_{12} = c_{12}$$
 and $m_{13} = c_{13}$

Let $\mathcal{V}_{23} = \{ w_{23} \in \mathcal{M}_{23} : \mathcal{M}_{12}w_{23} = 0 \}$. Therefore,

$$\varphi_1(a+m+b) - \varphi_1(a) - \varphi_1(b) - \varphi_1(m)$$

$$\in \left\{ \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & w_{23} \\ 0 & 0 & a_{33} \end{pmatrix} : w_{23} \in \mathcal{V}_{23}, a_{11} \in \mathcal{Z}(\mathcal{T}_1), a_{11}m_{1j} = m_{1j}a_{jj} \forall m_{1j} \in \mathcal{M}_{1j}, j = 2, 3 \right\}.$$

We remark that, Lemmas 3.2-3.6 hold for all multiplicative Lie derivations φ_1 on \mathcal{T} satisfying $Q_1\varphi(Q'_1)Q'_1 = 0$, and the assumption " $Q_1\mathcal{Z}(\mathcal{T})Q_1 = \mathcal{Z}(\mathcal{T}_1)$ and $Q'_1\mathcal{Z}(\mathcal{T})Q'_1 = \mathcal{Z}(\mathcal{T}'_1)$ " as well as the assumption "every multiplicative Lie derivation on \mathcal{T}'_1 has the standard form" are not needed.

Proof of Theorem 3.1: k = 1. Note that, by Lemma 3.2 and Lemma 3.3, we have $Q_1\varphi_1(Q'_1)Q'_1 = 0, Q_1\varphi_1(\mathcal{T}'_1)Q_1 \subseteq Q_1\mathcal{Z}(\mathcal{T})Q_1, Q'_1\varphi_1(\mathcal{T}_1)Q'_1 \subseteq Q'_1\mathcal{Z}(\mathcal{T})Q'_1 \text{ and } \varphi_1(Q'_1), \varphi_1(Q_1) \in \mathcal{Z}(\mathcal{T}).$

Let $\tau : \pi_{\mathcal{T}_1}(\mathcal{Z}(\mathcal{T})) \to \pi_{\mathcal{T}'_1}(\mathcal{Z}(\mathcal{T}))$ be the unique ring isomorphism so that $z \oplus \tau(z) \in \mathcal{Z}(\mathcal{T})$. According to the hypotheses on \mathcal{T} in Theorem 3.1, we have $\pi_{\mathcal{T}_1}(\mathcal{Z}(\mathcal{T})) = \mathcal{Z}(\mathcal{T}_1)$ and $\pi_{\mathcal{T}'_1}(\mathcal{Z}(\mathcal{T})) = \mathcal{Z}(\mathcal{T}'_1)$. Thus, for each $a \in \mathcal{Z}(\mathcal{T}_1)$, $am = m\tau(a)$ holds for all $m \in \mathcal{M}_1$. Clearly, by identifying $\pi_{\mathcal{T}_1}(\mathcal{T})$ with $Q_1\mathcal{T}Q_1$ and $\pi_{\mathcal{T}'_1}(\mathcal{T})$ with $Q'_1\mathcal{T}Q'_1$, we have $Q_1\varphi_1(b)Q_1 + \tau(Q_1\varphi_1(b)Q_1) \in \mathcal{Z}(\mathcal{T})$ and $\tau^{-1}(Q'_1\varphi_1(a)Q'_1) + Q'_1\varphi_1(a)Q'_1 \in \mathcal{Z}(\mathcal{T})$ for all $a \in \mathcal{T}_1$ and $b \in \mathcal{T}'_1$. Let us define maps $\gamma_1 : \mathcal{T} \to \mathcal{Z}(\mathcal{T})$ and $\delta_0 : \mathcal{T} \to \mathcal{T}$ by

$$\gamma_1(x) = Q_1 \varphi_1 (Q'_1 x Q'_1) Q_1 + \tau (Q_1 \varphi_1 (Q'_1 x Q'_1) Q_1) + \tau^{-1} (Q'_1 \varphi_1 (Q_1 x Q_1) Q'_1) + Q'_1 \varphi_1 (Q_1 x Q_1) Q'_1$$

and

$$\delta_0(x) = \varphi_1(x) - \gamma_1(x).$$

Obviously, $\gamma_1(\mathcal{M}_1) = 0$. Hence, according to Lemma 3.3 (c), we have

$$\delta_0(m) = \varphi_1(m) \in \mathcal{M}_1$$

for all $m \in \mathcal{M}_1$.

We claim that $\gamma_1([\mathcal{T},\mathcal{T}]) = 0$ and that δ_0 is a multiplicative Lie derivation. In fact, for arbitrary $x_1, x_2 \in \mathcal{T}$,

$$\begin{split} \gamma_1([x_1, x_2]) &= Q_1 \varphi_1(Q_1'([x_1, x_2])Q_1')Q_1 + \tau^{-1}(Q_1'\varphi_1(Q_1([x_1, x_2])Q_1)Q_1') \\ &+ Q_1'\varphi_1(Q_1([x_1, x_2])Q_1)Q_1' + \tau(Q_1\varphi_1(Q_1'([x_1, x_2]Q_1')Q_1)) \\ &= Q_1\varphi_1([Q_1'x_1Q_1', Q_1'x_2Q_1'])Q_1 + \tau^{-1}(Q_1'\varphi_1([Q_1x_1Q_1, Q_1x_2Q_1])Q_1') \\ &+ Q_1'\varphi_1([Q_1x_1Q_1, Q_1x_2Q_1])Q_1' + \tau(Q_1\varphi([Q_1'x_1Q_1', Q_1'x_2Q_1'])Q_1). \end{split}$$

Since

$$Q_{1}\varphi_{1}([Q'_{1}x_{1}Q'_{1},Q'_{1}x_{2}Q'_{1}])Q_{1}$$

$$= Q_{1}([\varphi_{1}(Q'_{1}x_{1}Q'_{1}),Q'_{1}x_{2}Q'_{1}])Q_{1} + Q_{1}([Q'_{1}x_{1}Q'_{1},\varphi_{1}(Q'_{1}x_{2}Q'_{1}]))Q_{1}$$

$$= 0$$

and

$$Q_{1}'\varphi_{1}([Q_{1}x_{1}Q_{1},Q_{1}x_{2}Q_{1}])Q_{1}'$$

$$= Q_{1}'([\varphi_{1}(Q_{1}x_{1}Q_{1}),Q_{1}x_{2}Q_{1}])Q_{1}' + Q_{1}'([Q_{1}x_{1}Q_{1},\varphi_{1}(Q_{1}x_{2}Q_{1}]))Q_{1}'$$

$$= 0,$$

we see that $\gamma_1([x_1, x_2]) = 0$. So $\gamma_1([\mathcal{T}, \mathcal{T}]) = 0$. Consequently

$$\begin{split} \delta_0([x_1, x_2]) &= \varphi_1([x_1, x_2]) \\ &= [\varphi_1(x_1), x_2] + [x_1, \varphi_1(x_2)] \\ &= [\varphi_1(x_1) - \gamma_1(x_1), x_2] + [x_1, \varphi_1(x_2) - \gamma_1(x_2)] \\ &= [\delta_0(x_1), x_2] + [x_1, \delta_0(x_2)] \end{split}$$

for all $x_1, x_2 \in \mathcal{T}$; that is, δ_0 is a multiplicative Lie derivation which obviously satisfies $Q_1 \delta_0(Q'_1) Q_1 = 0$. Furthermore, we have

$$\delta_0(\mathcal{M}_1) \subseteq \mathcal{M}_1, \ \delta_0(\mathcal{T}_1) \subseteq \mathcal{T}_1 \text{ and } \delta_0(\mathcal{T}_1') \subseteq \mathcal{T}_1'.$$
 (3.20)

Indeed, $\delta_0(\mathcal{M}_1) \subseteq \mathcal{M}_1$ is clear. Using Lemma 3.3 (c) and the definitions of δ_0 , γ_1 , for each $a \in \mathcal{T}$, one gets

$$\begin{split} \delta_0(a) &= \varphi_1(a) - \gamma_1(a) \\ &= Q_1\varphi_1(a)(a)Q_1 + Q_1'\varphi_1(a)Q_1' - \tau^{-1}(Q_1'\varphi_1(a)Q_1') - Q_1'\varphi_1(a)Q_1' \\ &= Q_1\varphi_1(a)Q_1 - \tau^{-1}(Q_1'\varphi_1(a)Q_1') \in \mathcal{T}_1. \end{split}$$

Analogously, one has $\delta_0(b) = Q'_1 \delta_0(b) Q'_1 \in \mathcal{T}'_1$ for each $b \in \mathcal{T}'_1$.

Next, let us introduce two more maps $\delta_1 : \mathcal{T} \to \mathcal{T}$ and $\xi_1 : \mathcal{T} \to \mathcal{T}$ defined respectively by

$$\delta_1(x) = \delta_0(Q_1 x Q_1) + \delta_0(Q_1 x Q_1') + \delta_0(Q_1' x Q_1')$$

and

$$\xi_1(x) = \delta_0(x) - \delta_1(x)$$

for each $x \in \mathcal{T}$. Note that δ_0 is a multiplicative Lie derivation. We claim that δ_1 is also a multiplicative Lie derivation. To see this, letting $a, a' \in \mathcal{T}_1, b, b' \in \mathcal{T}'_1, m, m' \in \mathcal{M}_1$, we have

$$\begin{split} &\delta_1([a+b+m,a'+b'+m']) \\ &= &\delta_1(\begin{pmatrix} a & m \\ 0 & b \end{pmatrix} \begin{pmatrix} a' & m' \\ 0 & b' \end{pmatrix} - \begin{pmatrix} a' & m' \\ 0 & b' \end{pmatrix} \begin{pmatrix} a & m \\ 0 & b \end{pmatrix}) \\ &= &\delta_1(\begin{pmatrix} aa'-a'a & am'+mb'-a'm-m'b \\ 0 & bb'-b'b \end{pmatrix}) \\ &= &\delta_0(aa'-a'a) + \delta_0(am'+mb'-a'm-m'b) + \delta_0(bb'-b'b) \\ &= &\delta_0([a,a']) + [\delta_0(a),m'] + [a,\delta_0(m')] + [\delta_0(m),b'] + [m,\delta_0(b')] \\ &+ [\delta_0(m),a'] + [m,\delta_0(a')] + [\delta_0(b),m'] + [b,\delta_0(m')] + \delta_0([b,b']) \end{split}$$

and

$$\begin{split} & [\delta_1(a+b+m),a'+b'+m'] + [a+b+m,\delta_1(a'+b'+m')] \\ = & [\delta_0(a)+\delta_0(b)+\delta_0(m),a'+b'+m'] + [a+b+m,\delta_0(a')+\delta_0(b')+\delta_0(m')] \\ = & [\delta_0(a),a'] + [\delta_0(a),m'] + [\delta_0(b),b'] + [\delta_0(b),m'] + [\delta_0(m),a'] + [\delta_0(m),b'] \\ & + [a,\delta_0(a')] + [m,\delta_0(a')] + [b,\delta_0(b')] + [m,\delta_0(b')] + [a,\delta_0(m')] + [b,\delta_0(m')]. \end{split}$$

 \mathbf{So}

$$\delta_1([a+b+m,a'+b'+m']) = [\delta_1(a+b+m),a'+b'+m'] + [a+b+m,\delta_1(a'+b'+m')].$$

It follows that δ_1 is a multiplicative Lie derivation with $Q_1\delta_1(Q'_1)Q'_1 = 0$. Consequently, $\xi_1(x) = \delta_0(x) - \delta_1(x)$ is also a multiplicative Lie derivation with $Q_1\xi_1(Q'_1)Q'_1 = 0$. This ensures that Lemma 3.3-Lemma 3.5 are applicable to both δ_1 and ξ_1 .

Our next aim is to prove that δ_1 is a derivation of \mathcal{T} .

Let $a, a' \in \mathcal{T}_1$. For each $m \in \mathcal{M}_1$. Using Lemmas 3.4-3.6 and the fact that $\delta_1(\mathcal{T}_1) \subseteq \mathcal{T}_1$ (see Eq.(3.20)), we get

$$\begin{split} \delta_1((a+a')m) &= \delta_1(am+a'm) \\ &= \delta_1(am) + \delta_1(a'm) \\ &= \delta_1(a)m + a\delta_1(m) - m\delta_1(a) + \delta_1(a')m + a'\delta_1(m) - m\delta_1(a') \\ &= \delta_1(a)m + a\delta_1(m) + \delta_1(a')m + a'\delta_1(m). \end{split}$$

On the other hand, by using Lemma 3.5 we have

$$\begin{split} \delta_1((a+a')m) &= \delta_1(a+a')m + (a+a')\delta_1(m) - m\delta_1(a+a') \\ &= \delta_1(a+a')m + a\delta_1(m) + a'\delta_1(m). \end{split}$$

The above two equations ensure that

$$(\delta_1(a+a') - \delta_1(a) - \delta_1(a'))m = 0 \tag{3.21}$$

holds for all $a, a' \in \mathcal{T}_1$ and $m \in \mathcal{M}_1$. By the definition of triangular 3-matrix ring, it is clear that \mathcal{M}_1 is faithful as a left \mathcal{T}_1 -module. So Eq.(3.21) entails that $\delta_1(a + a') = \delta_1(a) + \delta_1(a')$ holds for all $a, a' \in \mathcal{T}_1$; that is, δ_1 is additive on \mathcal{T}_1 .

In addition, using Lemma 3.5 on $\delta_1(aa'm)$, we see that

$$\delta_1(aa'm) = \delta_1(aa')m + aa'\delta_1(m) - m\delta_1(aa')$$

= $\delta_1(aa')m + aa'\delta_1(m).$

While on the other hand

$$\delta_{1}(aa'm) = \delta_{1}(a(a'm)) = \delta_{1}(a)a'm + a\delta_{1}(a'm) - a'm\delta_{1}(a) = \delta_{1}(a)a'm + a\delta_{1}(a')m + aa'\delta_{1}(m) - am\delta_{1}(a') - a'm\delta_{1}(a) = \delta_{1}(a)a'm + a\delta_{1}(a')m + aa'\delta_{1}(m).$$

These force that $\delta_1(aa') = \delta_1(a)a' + a\delta_1(a')$ and hence δ_1 is a derivation when restricted to \mathcal{T}_1 .

In the sequel we denote $\delta_1|_{\mathcal{T}_1}$ by $\delta_{\mathcal{T}_1}$ and $\delta_1|_{\mathcal{M}_1}$ by δ_{M_1} .

Note that \mathcal{T}'_1 is a triangular ring. By the hypotheses of Theorem 3.1, the multiplicative Lie derivation $\delta_1|_{\mathcal{T}'_1}: \mathcal{T}'_1 \to \mathcal{T}'_1$ has a standard form $\delta_1|_{\mathcal{T}'_1} = \delta_{\mathcal{T}'_1} + \eta$, where $\delta_{\mathcal{T}'_1}$ is an additive derivation on \mathcal{T}'_1 and $\eta: \mathcal{T}'_1 \to \mathcal{Z}(\mathcal{T}'_1)$ is a map such that $\eta(\mathcal{T}'_1, \mathcal{T}'_1) = 0$. We assert that $\eta = 0$. In fact, for any $x = \begin{pmatrix} a_1 & m \\ 0 & a_2 \end{pmatrix}, y = \begin{pmatrix} b_1 & n \\ 0 & b_2 \end{pmatrix} \in \mathcal{T}$, where $a_1, b_1 \in \mathcal{T}_1, m, n \in \mathcal{M}_1$, $a_2, b_2 \in \mathcal{T}'_1$, we have

$$\delta_{1}([x.y]) = \delta_{1}(\begin{pmatrix} a_{1}b_{1} - b_{1}a_{1} & a_{1}n + mb_{2} - b_{1}m - na_{2} \\ 0 & a_{2}b_{2} - b_{2}a_{2} \end{pmatrix})$$

$$= \delta_{\mathcal{T}_{1}}(a_{1}b_{1} - b_{1}a_{1}) + \delta_{M_{1}}(a_{1}n + mb_{2} - b_{1}m - na_{2})$$

$$+ \delta_{\mathcal{T}_{1}'}(a_{2}b_{2} - b_{2}a_{2}) + \eta(a_{2}b_{2} - b_{2}a_{2}).$$
(3.22)

On the other hand

$$= \begin{pmatrix} \delta_{\tau_{1}}(a_{1}) & \delta_{M_{1}}(m) \\ 0 & \delta_{\tau_{1}'}(a_{2}) + \eta(a_{2}) \end{pmatrix} \begin{pmatrix} b_{1} & n \\ 0 & b_{2} \end{pmatrix} - \begin{pmatrix} b_{1} & n \\ 0 & b_{2} \end{pmatrix} \begin{pmatrix} \delta_{\tau_{1}}(a_{1}) & \delta_{M_{1}}(m) \\ 0 & \delta_{\tau_{1}'}(a_{2}) + \eta(a_{2}) \end{pmatrix} \\ + \begin{pmatrix} a_{1} & m \\ 0 & a_{2} \end{pmatrix} \begin{pmatrix} \delta_{\tau_{1}}(b_{1}) & \delta_{M_{1}}(n) \\ 0 & \delta_{\tau_{1}'}(b_{2}) + \eta(b_{2}) \end{pmatrix} - \begin{pmatrix} \delta_{\tau_{1}}(b_{1}) & \delta_{M_{1}}(n) \\ 0 & \delta_{\tau_{1}'}(b_{2}) + \eta(b_{2}) \end{pmatrix} \begin{pmatrix} a_{1} & m \\ 0 & a_{2} \end{pmatrix} \\ = \begin{pmatrix} \delta_{\tau_{1}}(a_{1})b_{1} - b_{1}\delta_{\tau_{1}}(a_{1}) & \delta_{\tau_{1}}(a_{1})n + \delta_{M_{1}}(m)b_{2} - b_{1}\delta_{M_{1}}(m) - n\delta_{\tau_{1}'}(a_{2}) - n\eta(a_{2}) \\ 0 & \delta_{\tau_{1}'}(a_{2})b_{2} + \eta(a_{2})b_{2} - b_{2}\delta_{\tau_{1}'}(a_{2}) - b_{2}\eta(a_{2}) \end{pmatrix} \\ + \begin{pmatrix} a_{1}\delta_{\tau_{1}}(b_{1}) - \delta_{\tau_{1}}(b_{1})a_{1} & a_{1}\delta_{M_{1}}(n) + m\delta_{\tau_{1}'}(b_{2}) + m\eta(b_{2}) - \delta_{\tau_{1}}(b_{1})m - \delta_{M_{1}}(n)a_{2} \\ 0 & a_{2}\delta_{\tau_{1}'}(b_{2}) + a_{2}\eta(b_{2}) - \delta_{\tau_{1}'}(b_{2})a_{2} - a_{2}\eta(b_{2}) \end{pmatrix}.$$

$$(3.23)$$

As $\delta_1([x,y]) = [\delta_1(x), y] + [x, \delta_1(y)]$, by Eq(3.22), Eq(3.23) and Lemmas 3.4-3.5, we obtain that

$$\begin{split} \delta_{\mathcal{T}_{1}}(a_{1})n + a_{1}\delta_{M_{1}}(n) &- n\delta_{\mathcal{T}_{1}}(a_{1}) + \delta_{M_{1}}(m)b_{2} + m\delta_{\mathcal{T}_{1}'}(b_{2}) - \delta_{\mathcal{T}_{1}'}(b_{2})m \\ &- \delta_{\mathcal{T}_{1}}(b_{1})m - b_{1}\delta_{M_{1}}(m) + m\delta_{\mathcal{T}_{1}}(b_{1}) - \delta_{M_{1}}(n)a_{2} - n\delta_{\mathcal{T}_{1}'}(a_{2}) + \delta_{\mathcal{T}_{1}'}(a_{2})n \\ &= \delta_{M_{1}}(a_{1}n) + \delta_{M_{1}}(mb_{2}) - \delta_{M_{1}}(b_{1}m) - \delta_{M_{1}}(na_{2}) \\ &= \delta_{M_{1}}(a_{1}n + mb_{2} - b_{1}m - na_{2}) \\ &= \delta_{\mathcal{T}_{1}}(a_{1})n + \delta_{M_{1}}(m)b_{2} - b_{1}\delta_{M_{1}}(m) - n\delta_{\mathcal{T}_{1}'}(a_{2}) - n\eta(a_{2}) \\ &+ a_{1}\delta_{M_{1}}(n) + m\delta_{\mathcal{T}_{1}'}(b_{2}) + m\eta(b_{2}) - \delta_{\mathcal{T}_{1}}(b_{1})m - \delta_{M_{1}}(n)a_{2}. \end{split}$$

Since $n\delta_{\mathcal{T}_1}(a_1) = m\delta_{\mathcal{T}_1}(b_1) = \delta_{\mathcal{T}'_1}(a_2)n = \delta_{\mathcal{T}'_1}(b_2)m = 0$, we see that

$$m\eta(b_2) - n\eta(a_2) = 0$$
 for all $n, m \in \mathcal{M}_1$,

which implies that $\eta = 0$. To see this, for any $b_2 \in \mathcal{T}'_1$, picking n = 0 in the above equation gives that $m\eta(b_2) = 0$ holds for all $m \in \mathcal{M}_1$. Write $m = (m_{12}, m_{13}), \eta(b_2) = \begin{pmatrix} z_2 & 0 \\ 0 & z_3 \end{pmatrix} \in \mathcal{Z}(\mathcal{T}'_1).$ Then $m\eta(b_2) = (m_{12}z_2, m_{13}z_3) = (0, 0)$. As $m_{12} \in \mathcal{M}_{12}$ and $m_{13} \in \mathcal{M}_{13}$ are arbitrary, we must have $z_2 = 0$ and $z_3 = 0$. So $\eta(b_2) = 0$. Thus δ_1 is in fact an additive derivation on \mathcal{T}'_1 .

By now, we have proved that δ_1 is additive respectively on $\mathcal{T}_1, \mathcal{T}'_1$, and \mathcal{M}_1 (see Lemma 3.6)

(2)). As $\delta_1(\mathcal{T}_1) \subseteq \mathcal{T}_1, \delta_1(\mathcal{T}_1') \subseteq \mathcal{T}_1', \delta_1(M_1) \subseteq M_1$, it is easily checked the δ_1 is additive. For any $x = \begin{pmatrix} a_1 & m \\ 0 & a_2 \end{pmatrix}$ and $y = \begin{pmatrix} b_1 & n \\ 0 & b_2 \end{pmatrix} \in \mathcal{T}$, using Lemmas 3.3-3.5 and the fact that δ_1 is derivation on \mathcal{T}_1 and \mathcal{T}'_1 , we see that

$$\begin{split} \delta_1(xy) &= & \delta_1(a_1b_1 + (a_1n + mb_2) + a_2b_2) \\ &= & \delta_1(a_1b_1) + \delta_1(a_1n + mb_2) + \delta_1(a_2b_2) \\ &= & \delta_1(a_1)b_1 + a_1\delta_1(b_1) + \delta_1(a_1n) + \delta_1(mb_2) + \delta_1(a_2)b_2 + a_2\delta_1(b_2) \\ &= & \delta_1(a_1)b_1 + a_1\delta_1(b_1) + \delta_1(a_1)n + a_1\delta_1(n) \\ &+ \delta_1(m)b_2 + m\delta_1(b_2) + \delta_1(a_2)b_2 + a_2\delta_1(b_2). \end{split}$$

While on the other hand

$$\begin{split} \delta_1(x)y + x\delta_1(y) &= (\delta_1(a_1) + \delta_1(m) + \delta_1(a_2))(b_1 + n + b_2) \\ &+ (a_1 + m + a_2)(\delta_1(b_1) + \delta_1(n) + \delta_1(b_2)) \\ &= \delta_1(a_1)b_1 + \delta_1(a_1)n + \delta_1(m)b_2 + \delta_1(a_2)b_2 \\ &+ a_1\delta_1(b_1) + a_1\delta_1(n) + m\delta_1(b_2) + a_2\delta_1(b_2). \end{split}$$

Consequently, $\delta_1(xy) = \delta_1(x)y + x\delta_1(y)$. Hence δ_1 is a derivation of \mathcal{T} .

Finally, let us consider the structure of ξ_1 . Since $\xi_1(x) = \delta_0(x) - \delta_1(x)$ and $\varphi_1(x) = \delta_0(x) + \delta_0(x) - \delta_1(x)$ $\gamma_1(x)$, by the definition of γ_1 , we get $\xi_1(x) = \varphi_1(x) - \varphi_1(Q_1xQ_1) - \varphi_1(Q_1xQ_1') - \varphi_1(Q_1xQ_1')$. It follows from Lemma 3.6 (3) that, for each $x \in \mathcal{T}, \xi_1(x) \in \mathcal{S}_{23}$, where

$$S_{23} = \left\{ \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & w_{23} \\ 0 & 0 & a_{33} \end{pmatrix} : \\ w_{23} \in \mathcal{V}_{23}, a_{11} \in \mathcal{Z}(\mathcal{T}_1), a_{11}m_{1j} = m_{1j}a_{jj} \ \forall m_{1j} \in \mathcal{M}_{1j}, j = 2, 3 \right\}$$

Define maps $\gamma_2 : \mathcal{T} \to \mathcal{Z}(\mathcal{T})$ and $\xi_2 : \mathcal{T} \to \mathcal{T}$ by

$$\gamma_2(x) = Q_1 \xi_1(x) Q_1 + \tau(Q_1 \xi_1(x) Q_1)$$

and

$$\xi_2 = \xi_1 - \gamma_2.$$

We claim that $\xi_2(x) \in S_1$ for all $x \in \mathcal{T}$. To see this, write

$$\xi_1(x) = \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & a_{22} & w_{23} \\ 0 & 0 & a_{33} \end{pmatrix} \in \mathcal{S}_{23} \text{ and } \gamma_2(x) = \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & b_{22} & 0 \\ 0 & 0 & b_{33} \end{pmatrix} \in \mathcal{Z}(\mathcal{T}).$$

As $\xi_1(x) - \gamma_2(x) \in S_{23}$, we get $(a_{11} - a_{11})m_{1j} = m_{1j}(a_{jj} - b_{jj})$ holds for all $m_{1j} \in \mathcal{M}_{1,j}$, which forces that $b_{jj} = a_{jj}, j = 2, 3$.

We show that $\gamma_2([\mathcal{T}, \mathcal{T}]) = 0$ and hence ξ_2 is a multiplicative Lie derivation. In fact, for arbitrary $x_1, x_2 \in \mathcal{T}$,

$$\gamma_2([x_1, x_2]) = Q_1 \xi_1([x_1, x_2]) Q_1 + \tau(Q_1 \xi_1([x_1, x_2]) Q_1).$$

Since $Q_1\xi_1(x_i)Q_1 \in \mathcal{Z}(\mathcal{T}_1)$ for i = 1, 2, we see that

$$Q_{1}\xi_{1}([x_{1}, x_{2}])Q_{1}$$

$$= Q_{1}[\xi_{1}(x_{1}), x_{2}]Q_{1} + Q_{1}[x_{1}, \xi_{1}(x_{2})]Q_{1}$$

$$= Q_{1}\xi_{1}(x_{1})x_{2}Q_{1} - Q_{1}x_{2}\xi_{1}(x_{1})Q_{1} + Q_{1}x_{1}\xi_{1}(x_{2})Q_{1} - Q_{1}\xi_{1}(x_{2})x_{1}Q_{1}$$

$$= 0,$$

which entails $\gamma_2([x_1, x_2]) = 0$. So $\gamma_2([\mathcal{T}, \mathcal{T}]) = 0$. Consequencely ξ_2 is a multiplicative Lie derivation which obviously satisfies $Q_1\xi_2(Q'_1)Q'_1 = 0$ and

$$\xi_2 \in \left\{ \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & w_{23} \\ 0 & 0 & 0 \end{pmatrix} : w_{23} \in \mathcal{V}_{23} \right\} = \mathcal{S}_1.$$
(3.24)

So far we have shown that $\varphi_1 = \delta_0 + \gamma_1 = \delta_1 + \xi_1 + \gamma_1 = \delta_1 + \xi_2 + \gamma_2 + \gamma_1$. Let $\delta = d_1 + \delta_1$, $\gamma = \gamma_1 + \gamma_2$ and $\xi = \xi_2$. Then δ is a derivation, γ is a center valued map vanishing each commutator, ξ is a multiplicative Lie derivation with range in S_1 and $\varphi = \delta + \gamma + \xi$, as desired.

3.2. Proof of Theorem 3.1: k = 3. In this case $\mathcal{T} = \mathcal{T}'_3 + \mathcal{M}_3 + \mathcal{T}_3$ with $\mathcal{T}_3 = Q_3 \mathcal{T} Q_3$, $\mathcal{T}'_3 = Q'_3 \mathcal{T} Q'_3$ and $\mathcal{M}_3 = Q'_3 \mathcal{T} Q_3$.

Let $\varphi : \mathcal{T} \to \mathcal{T}$ be a multiplicative Lie derivation. One can prove in a similar way that the analogues of Lemmas 3.2-3.5 are valid. We list them here by omitting the proofs for application later.

Lemma 3.2'. There exists an inner derivation $d'_1 : \mathcal{T} - \mathcal{T}$ and a multiplicative Lie derivation $\varphi'_1 : \mathcal{T} \to \mathcal{T}$ such that

$$\varphi = d'_1 + \varphi'_1$$
 and $Q'_3 \varphi'_1(Q'_3) Q_3 = 0.$

Note that we take $d'_1(x) = [x, \varphi(Q'_3)].$

Lemma 3.3'. For any $f \in \mathcal{T}'_3, g \in \mathcal{T}_3$ and $h \in \mathcal{M}_3$, the following statements are true: (a) $\varphi'_1(Q_3), \varphi'_1(Q'_3) \in \mathcal{R}_1 \oplus \mathcal{R}_2 \oplus \mathcal{R}_3$.

(b) $Q'_3\varphi_1(f)Q_3 = Q'_3\varphi'_1(g)Q_3 = 0.$

(c) $\varphi'_{1}(f) = Q'_{3}\varphi'_{1}(f)Q'_{3} + Q_{3}\varphi'_{1}(f)Q_{3}, \ \varphi'_{1}(g) = Q'_{3}\varphi'_{1}(g)Q'_{3} + Q_{3}\varphi'_{1}(g)Q_{3} \text{ and } \varphi'_{1}(h) = Q'_{3}\varphi'_{1}(h)Q_{3}.$ (d) $Q_{3}\varphi'_{1}(\mathcal{T}'_{3})Q_{3} \subseteq \mathcal{Z}(\mathcal{T}_{3}) \text{ and } Q'_{3}\varphi'_{1}(\mathcal{T}_{3})Q'_{3} \subseteq \mathcal{Z}(\mathcal{T}'_{3}).$ Lemma 3.4'. $\varphi'_{1}(Q_{3}), \varphi'_{1}(Q'_{3}) \in \mathcal{Z}(\mathcal{T}) \text{ and } \varphi'_{1}(Q'_{3}xQ_{3}) = Q'_{3}\varphi'_{1}(x)Q_{3} \text{ for all } x \in \mathcal{T}.$ Lemma 3.5'. For any $f \in \mathcal{T}'_{3}, g \in \mathcal{T}_{3} \text{ and } h \in \mathcal{M}_{3}, we have$

$$\varphi_1'(fh) = \varphi_1'(f)h + f\varphi_1'(h) - h\varphi_1'(f)$$

and

$$\varphi_1'(hg) = \varphi_1'(h)g + h\varphi_1'(g) - \varphi_1'(g)h$$

Lemma 3.6'. For any $f \in \mathcal{T}'_3$, $g \in \mathcal{T}_3$ and $h \in \mathcal{M}_3$, we have (1) Both $\varphi'_1(f+h) - \varphi'_1(f) - \varphi'_1(h)$ and $\varphi'_1(f+g) - \varphi'_1(f) - \varphi'_1(g) \in \mathcal{Z}(\mathcal{T})$. (2) φ'_1 is additive on \mathcal{M}_3 . (3) With $\mathcal{V}_{12} = \{w_{12} \in \mathcal{M}_{12} : w_{12}\mathcal{M}_{23} = 0\}$, we have $\varphi'_1(f+h+g) - \varphi'_1(f) - \varphi'_1(g) - \varphi'_1(h)$

$$\in \{ \begin{pmatrix} a_{11} & w_{12} & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix} : w_{12} \in \mathcal{V}_{12}, a_{ii}m_{i3} = m_{i3}a_{33} \forall m_{i3} \in \mathcal{M}_{i3}, a_{33} \in \mathcal{Z}(\mathcal{T}_3) \}.$$

Proof of Theorem 3.1: k = 3. As $Q_1 \mathcal{Z}(\mathcal{T})Q_1 = \mathcal{Z}(\mathcal{T}_1)$ and $Q'_1 \mathcal{Z}(\mathcal{T})Q'_1 = \mathcal{Z}(\mathcal{T}'_1)$, there is a unique isomorphism $\tau : \mathcal{Z}(\mathcal{T}_3) \to \mathcal{Z}(\mathcal{T}'_3)$ so that $\tau(z) + z \in \mathcal{Z}(\mathcal{T})$. Then the map γ'_1 defined by

$$\begin{aligned} \gamma_1'(x) &= Q_3 \varphi_1'(Q_3' x Q_3') Q_3 + \tau'(Q_3 \varphi_1(Q_3' x Q_3') Q_3) \\ &+ (\tau')^{-1} (Q_3' \varphi_1'(Q_3 x Q_3) Q_3') + Q_3' \varphi_1'(Q_3 x Q_3) Q_3' \end{aligned}$$

is a center valued map vanishing each commutator. Consequently,

$$\delta_0' = \varphi_1' - \gamma_1'.$$

is still a multiplicative Lie derivation satisfying $Q'_3\delta'_0(Q'_3)Q_3 = 0$ and Lemmas 3.2'-3.6' are applicable. By using the hypothesis that every multiplicative Lie derivation on \mathcal{T}'_3 has the standard form, one can prove that the map δ'_1 defined by

$$\delta_1'(f+g+h)=\delta_0'(f)+\delta_0'(g)+\delta_0'(h)$$

for any $f \in \mathcal{T}'_3$, $g \in \mathcal{T}_3$ and $h \in \mathcal{M}_3$ is a derivation. Thus

$$\xi_1' = \delta_0' - \delta_1'$$

is still a multiplicative Lie derivation satisfying $Q'_3\xi'_1(Q'_3)Q_3 = 0$. Moreover, the range of ξ'_1 is contained in the set

$$\mathcal{S}_{12} = \{ \begin{pmatrix} a_{11} & w_{12} & 0 \\ 0 & a_{22} & 0 \\ 0 & 0 & a_{33} \end{pmatrix} : w_{12} \in \mathcal{V}_{12}, a_{ii}m_{i3} = m_{i3}a_{33} \ \forall m_{i3} \in \mathcal{M}_{i3}, i = 1, 2, a_{33} \in \mathcal{Z}(\mathcal{T}_3) \}.$$

Let γ'_2 be the map defined by

$$\gamma_2'(x) = (\tau')^{-1}(Q_3\xi_1'(x)Q_3) + Q_3\xi_1'(x)Q_3.$$

 γ'_2 is a center valued map vanishing each commutator. Hence $\xi'_2 = \xi'_1 - \gamma'_2$ is a multiplicative γ_2 is a center valued map valued map value $\mathcal{S}_3 = \left\{ \begin{pmatrix} 0 & w_{12} & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} : w_{12} \in \mathcal{V}_{12} \right\}$ and $\varphi'_1 = \delta'_1 + \gamma'_1 + \gamma'_2 + \xi'_2$. With $\delta = d'_1 + \delta'_1$, $\gamma = \gamma'_1 + \gamma'_2$ and $\xi = \xi'_2$, we see that $\varphi = \delta + \gamma + \xi$ has the desired form. \Box

4. Proof of the main result

We are ready to prove our main result Theorem 2.2.

Proof of Theorem 2.2. Assume that $Q_i \mathcal{Z}(\mathcal{T})Q_i = \mathcal{Z}(\mathcal{T}_i), i = 1, 2, 3$.

Claim 1. For k = 1, 3, we have $Q'_k \mathcal{Z}(\mathcal{T})Q'_k = \mathcal{Z}(\mathcal{T}'_k)$. Thus \mathcal{T} satisfies the hypotheses of Theorem 3.1.

We check $Q'_1 \mathcal{Z}(\mathcal{T})Q'_1 = \mathcal{Z}(\mathcal{T}'_1)$ and the case k = 3 is dealt with similarly. Note that $Q_1' = Q_2 + Q_3 \text{ and obviously } Q_1' \mathcal{Z}(\mathcal{J}) Q_1' \subseteq \mathcal{Z}(\mathcal{T}_1'). \text{ Let } z' = \begin{pmatrix} 0 & 0 & 0 \\ 0 & z_2' & 0 \\ 0 & 0 & z_3' \end{pmatrix} \in \mathcal{Z}(\mathcal{T}_1'). \text{ Then,}$

for any $m_{23} \in \mathcal{M}_{23}$ we have $z'_2 m_{23} = m_{23} z'_3$. By the hypothesis of Theorem 2.2, there exists

 $z = \begin{pmatrix} z_1 & 0 & 0 \\ 0 & z_2 & 0 \\ 0 & 0 & z_3 \end{pmatrix} \in \mathcal{Z}(\mathcal{T}) \text{ such that } z_3 = z'_3. \text{ It follows that } z_2m_{23} = m_{23}z_3 = m_{23}z'_3 =$

 z'_2m_{23} holds for all $m_{23} \in \mathcal{M}_{23}$ which forces $z_2 = z'_2$. So, $Q'_1zQ'_1 = z'$. Hence we have $Q_1'\mathcal{Z}(\mathcal{T})Q_1' = \mathcal{Z}(\mathcal{T}_1').$

Claim 2. If $\xi : \mathcal{T} \to \mathcal{T}$ be a multiplicative Lie derivation of which the range is contained in S_1 , then ξ is a derivation.

Let $d_2: \mathcal{T} \to \mathcal{T}$ and $\varphi_2: \mathcal{T} \to \mathcal{T}$ be the maps defined respectively by

$$d_2(x) = [x, \xi(Q'_3)]$$

and

$$\varphi_2(x) = \xi(x) - d_2(x)$$

for all $x \in \mathcal{T}$. Obviously, d_2 is an inner derivation and φ_2 is a multiplicative Lie derivation satisfying $Q'_3\varphi_2(Q'_3)Q_3 = 0$. Note that $d_2(x) \in S_1$ and hence $\varphi_2(x) \in S_1$ for each $x \in \mathcal{T}$. Consequently, $\varphi_2(Q'_3) = 0$.

We assert that $\varphi_2 = 0$.

By Claim 1, we may apply Theorem 3.1 for the case k = 3 to φ_2 . Thus there exists a derivation δ' , a center valued map γ' and a multiplicative Lie derivation $\xi' : \mathcal{T} \to \mathcal{S}_3$ such that $\varphi_2 = \delta' + \gamma' + \xi'$. However, by checking the proof of Theorem 3.1 for the case k = 1, we must have $\delta'(x) \in S_1$ and $\gamma' = 0$. So, for each $x \in \mathcal{T}$, $\varphi_2(x) \in S_1 \cap S_3 = \{0\}$; that is, $\varphi_2 = 0$.

Claim 3. Every multiplicative Lie derivation on \mathcal{T} has the standard form.

Let $\varphi : \mathcal{T} \to \mathcal{T}$ be a multiplicative Lie derivation. By Claim 1, under the standard assumption $Q_i \mathcal{Z}(\mathcal{T}) Q_i = \mathcal{Z}(\mathcal{T}_i)$, \mathcal{T} meets the assumptions in Theorem 3.1 for both cases k = 1 and k = 3. Then, there exists a derivation, a center valued map γ which vanishes each commutator and a multiplicative Lie derivation $\xi : \mathcal{T} \to \mathcal{S}_1$ such that $\varphi = \delta + \gamma + \xi$. However, by Claim 2, ξ is in fact a derivation. Therefore, φ has the standard form.

MULTIPLICATIVE LIE DERIVATIONS

References

- K. I. Beider, M. A. Chebotar, On Lie derivation of Lie ideals of prime rings, Israel J. Math. 123 (2001) 131-148.
- [2] D Benkovič, Generalized Lie derivations on triangular algebras, Linear Algebra Appl. 434 (2011) 1532-1544.
- [3] D Benkovič, D Eremita, Multiplicative Lie n-derivation of triangular rings, Linear Algebra Appl. 436 (2012) 4223-4240.
- [4] M. Brešar, M.A. Chebotar, A. Mikhail, W.S. Martindale 3rd, Functional Identities, Birkhäuser Verlag, 2007.
- [5] W.S. Cheung, Commuting maps of triangular algebras, J. London Math. Soc. 63 (2001) 117-127.
- [6] W. S. Cheung, Lie derivations of triangular algebras, Linear and Multilinear Algebra, 51 (2003) 299-310.
- [7] Y. Du, Y. Wang, Lie derivations of generalized matrix algebras, Linear Algebra Appl. 437 (2012) 2719-2726.
- [8] B.L.M. Ferreira, Multiplicative maps of triangular n-matrix rings, International Journal of Mathematics, Game Theory and Algebra, 23 (2014) 1-14.
- [9] P.-S. Ji, R.-R. Liu, Y.-Z. Zhao, Nonlinear Lie triple derivations of triangular algebras, Linear and Multilinear Algebra 60 (2012) 1155-1164.
- [10] P.-S. Ji, W.-Q. Qi, Charactrizations of Lie derivations of triangular algebras, Linear Algebra Appl. 435 (2011) 1137C1146.
- [11] M.-Y. Jiao, J.-C. Hou, Additive maps derivable or Jordan derivable at zero point on nest algebras, Linear Algebra Appl. 432 (2010) 2984-2994
- [12] F.-Y. Lu, Lie triple derivations on nest algebras, Math. Nachr., 280 (2007) 882-887.
- [13] F.-Y. Lu, W. Jing, Characterizations of Lie derivations of B(X), Linear Algebra Appl., 432 (2010), 89C99.
- [14] X.-F. Qi, J.-C. Hou, Characterization of Lie derivations on von Neumann algebras, Linear Algebra Appl. 438 (2013) 533-548.
- [15] A.R. Villena, Lie derivations on Banach algebras, J. Algebra. 226 (2000) 390-409.
- [16] Yao Wang, Yu Wang, Multiplicative Lie n-derivation of generalized matrix algebras, Mathematical Reports. 17 (2015) 391-405.
- [17] W.-Y. Yu, J.-H. Zhang, Nonlinear Lie derivations of triangular algebras, Linear Algebra Appl., 432 (2010) 2953-2960.
- [18] J.-H. Zhang, B.-W. Wu, H.-X. Cao, Lie triple derivations of nest algebra, Linear Algebra Appl., 416 (2006) 559-567.

SCHOOL OF MATHEMATICS, TAIYUAN UNIVERSITY OF TECHNOLOGY, TAIYUAN, 030024, P. R. CHINA *E-mail address*, Jinchuan Hou: jinchuanhou@aliyun.com *E-mail address*, Zhenhui Chen: 2691324902@qq.com