

From Best Hits to Best Matches

Peter F. Stadler^{1,6-12}, Manuela Geiß¹, David Schaller¹, Alitzel López Sánchez², Marcos E. González², Dulce I. Valdivia³, Marc Hellmuth^{4,5}, and Maribel Hernández Rosales²

¹Bioinformatics Group, Department of Computer Science; and Interdisciplinary Center of Bioinformatics, University of Leipzig, Härtelstraße 16-18, D-04107 Leipzig, Germany

²CONACYT-Instituto de Matemáticas, UNAM Juriquilla, Blvd. Juriquilla 3001, 76230 Juriquilla, Querétaro, QRO, México

³Centro de Investigación y de Estudios Avanzados del IPN, CINVESTAV-Irapuato, Km. 9.6 Libramiento Norte Carretera Irapuato-León, 36821 Irapuato, GTO, México

⁴Institute of Mathematics and Computer Science, University of Greifswald, Walther-Rathenau-Straße 47, D-17487 Greifswald, Germany

⁵Center for Bioinformatics, Saarland University, Building E 2.1, P.O. Box 151150, D-66041 Saarbrücken, Germany

⁶German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig

⁷Competence Center for Scalable Data Services and Solutions

⁸Leipzig Research Center for Civilization Diseases, Leipzig University, Härtelstraße 16-18, D-04107 Leipzig

⁹Max-Planck-Institute for Mathematics in the Sciences, Inselstraße 22, D-04103 Leipzig

¹⁰Inst. f. Theoretical Chemistry, University of Vienna, Währingerstraße 17, A-1090 Wien, Austria

¹¹Facultad de Ciencias, Universidad Nacional de Colombia, Sede Bogotá, Ciudad Universitaria, 111321, Bogotá, D.C., Colombia

¹²Santa Fe Institute, 1399 Hyde Park Rd., Santa Fe, NM 87501, USA

Abstract

Background: Many of the commonly used methods for orthology detection start from mutually most similar pairs of genes (reciprocal best hits) as an approximation for evolutionary most closely related pairs of genes (reciprocal best matches). This approximation of best matches by best hits becomes exact for ultrametric dissimilarities, i.e., under the Molecular Clock Hypothesis. It fails, however, whenever there are large lineage specific rate variations among paralogous genes. In practice, this introduces a high level of noise into the input data for best-hit-based orthology detection methods.

Results: If additive distances between genes are known, then evolutionary most closely related pairs can be identified by considering certain quartets of genes provided that in each quartet the outgroup relative to the remaining three genes is known. *A priori* knowledge of underlying species phylogeny greatly facilitates the identification of the required outgroup. Although the workflow remains a heuristic since the correct outgroup cannot be determined reliably in all cases, simulations with lineage specific biases and rate asymmetries show that nearly perfect results can be achieved. In a realistic setting, where distances data have to be estimated from sequence data and hence are noisy, it is still possible to obtain highly accurate sets of best matches.

Conclusion: Improvements of tree-free orthology assessment methods can be expected from a combination of the accurate inference of best matches reported here and recent mathematical advances in the understanding of (reciprocal) best match graphs and orthology relations.

Keywords: Best Matches, Gene Tree, Species Tree, Reconciliation, Orthology




Figure 1: Lineage-specific rate variation between paralogs. The gene tree, with branch length indicating an additive evolutionary distances, pre-dates the speciation (\bullet) of the red and blue species. We have $\text{lca}(x, y) \prec \text{lca}(x, y')$ but $d(x, y') < d(x, y)$.

1 Background

The distinction of orthologous and paralogous pairs of genes, respectively, is of key importance in evolutionary biology as well as genome annotation. As defined by Walter Fitch [Fitch \[1970, 2000\]](#), two genes are orthologs if their last common ancestor (in the gene tree) corresponds to a speciation event, and they are paralogs if they arose through a duplication event. In general, orthologs are expected to have the same function in different organisms, while the functions of paralogs are usually similar but clearly distinct [Koonin \[2005\]](#), [Gabaldón and Koonin \[2013\]](#).

A large class of computational approaches to orthology assessment [Altenhoff and Dessimoz \[2009\]](#), [Altenhoff et al. \[2016\]](#) uses symmetric best matches (SBM) [Tatusov et al. \[1997\]](#), also known as bidirectional best hits (BBH) [Overbeek et al. \[1999\]](#), reciprocal best hits (RBH) [Bork et al. \[1998\]](#), or reciprocal smallest distance (RSD) [Wall et al. \[2003\]](#). The intuitive justification for these approaches is that symmetric best matches (in the sense of sequence similarity) approximate the idea of evolutionarily closest relatedness. These two concepts are not the same, however. The notion of evolutionary relatedness depends on the underlying phylogenetic tree T and is naturally expressed by comparing last common ancestors: a gene x is more closely related to a gene y than to y' if the last common ancestor $\text{lca}(x, y)$ is a successor of $\text{lca}(x, y')$ in T .

From an evolutionary point of view, therefore, one is interested in *reciprocal best matches* (defined in terms of the gene tree T) rather than in *reciprocal best hits* (defined in terms of some distance of similarity measure between sequences). Best matches and best hits are equivalent if and only if the Molecular Clock Hypothesis is satisfied [Zuckerandl and Pauling \[1962\]](#), [Kumar \[2005\]](#). In general this is not the case. In particular, paralogous members of a gene family often differ in their evolutionary rates due to (adaptive) changes in the function [Kawahara and Imanishi \[2007\]](#), [Soria et al. \[2014\]](#). Both the ‘‘Duplication-Degeneration-Complementation’’ (DDC) model [Force et al. \[1999\]](#) and the ‘‘Escape from Adaptive Conflict’’ (EAC) model [Hittinger and Carroll \[2007\]](#) predict that the fate of paralogs, including their evolutionary rate, may differ substantially between lineages that diverge soon after the duplication event due to different selective pressures. The simplest case is shown in Fig. 1: an ancestral gene is duplicated before the speciation event leading to two species (indicated by colors), each containing two paralogs (denoted by x and x' in the red species and y and y' in the blue species). The two paralogs evolve with very different rates in the two species. Although x and y as well x' and y' are orthologs, the evolutionary rates are more similar between x and y' , and x' and y , respectively. This situation is not at all uncommon. The asymmetric divergence of the genes in the HOXA cluster following the teleost-specific (3R) genome duplication may serve as a paradigmatic example [Wagner et al. \[2005\]](#). While in fugu (*Takifugu rubripes*) and other percomorphs the HOXA_b paralogs diverge faster, it is the HOXA13_b paralog that evolves at a faster rate in zebrafish (*Danio rerio*), which diverged early from percomorphs within the Teleostei clade.

In Fig. 1 the pair x, y' shows the largest sequence similarity and hence will appear as reciprocal best hit, while the closest evolutionary relative of x is the gene y . This discrepancy is **not** a consequence of inaccurate measurements but an intrinsic feature of the evolutionary process: more evolutionary events have accumulated on the path from x to y than on the path from x to y' . The correct *reciprocal best hit* therefore does not coincide with the correct *reciprocal best match*. This immediately begs the question whether such cases can be detected from sequence comparisons. We consider this issue at two levels: (i) Can (reciprocal) best matches be identified *in principle*, i.e., from perfectly accurate data, and (ii) how well can this be done in practice? To address the first question we will assume that

we can determine an additive distance between any two genes and investigate the consequences of this assumption. To investigate the accuracy that can be achieved from sequence data we will devise a simulation system to generate evolutionary scenarios with complex rate variations.

The focus on additive metrics is motivated by the close connection between additive metrics and evolutionary trees. More precisely, an additive metric determines a unique *unrooted* phylogenetic tree \bar{T} as well as its branch length [Simões-Pereira \[1969\]](#), [Buneman \[1974\]](#), and *vice versa*. The determination of best matches, which are defined in terms of last common ancestors, however, requires a rooted phylogenetic tree T . From a theoretical point of view, therefore, the missing information is the placement of the root of T in the underlying unrooted phylogenetic tree \bar{T} .

The problem of determining the position of the root in an unrooted tree \bar{T} has been well studied in the phylogenetic literature [Kinene et al. \[2016\]](#). The most common approach is the inclusion of an outgroup, i.e., a taxon z known to branch earlier than the taxa of interest. The root is then located in the branch leading to z . Outgroup rooting can be unreliable in the presence of rapid radiations or when only a few very distant outgroups are available [Holland et al. \[2003\]](#), [Shavit et al. \[2007\]](#). The simplest method is midpoint rooting [Swofford et al. \[1996\]](#), which places the root at the midpoint on the longest path in the tree. Despite its simplicity it often works remarkably well [Hess and de Moraes Russo \[2007\]](#). An interesting variation on this theme is minimum variance rooting [Mai et al.](#). The estimation of dated phylogenies using a relaxed clock assumption yields an estimate for the position of the root as a by-product [Drummond et al. \[2006\]](#). A related Bayesian method was introduced in [Huelsenbeck et al. \[2002\]](#). In a phylogenomics setting, the root of the species tree can also be obtained by minimizing the number of inferred gene duplications [Katz et al. \[2012\]](#). Most recently, non-reversible substitution models have been employed for estimating rooted phylogenetic trees [Williams et al. \[2015\]](#), [Cherlin et al. \[2018\]](#).

From a practical point of view, furthermore, we wish to avoid the explicit construction of a (rooted or unrooted) gene tree T since reconstructing accurate evolutionary trees from individual gene sequences is a notoriously difficult problem. Instead we aim to stay as close as possible to the idea of reciprocal best hit methods and thus we will attempt to use only “local” comparisons of as few as possible measurements of evolutionary distances. This idea naturally leads us to considering quartets, i.e., unrooted trees describing four taxa, and the corresponding rooted triples. It is well known that the rooted triples are sufficient to determine the rooted tree in which they reside. Moreover, there is a polynomial-time algorithm that either constructs a rooted tree T from a set of rooted triples or determines that no such tree exists [Aho et al. \[1981\]](#). By Buneman’s Theorem [Simões-Pereira \[1969\]](#), [Buneman \[1974\]](#), an unrooted tree can be uniquely recovered from all its quartets. However, the problem of determining whether a given set is compatible (i.e., whether there is an unrooted tree \bar{T} that contains all quartets) is NP-complete [Steel \[1992\]](#), a fact that reinforces the desire to avoid the explicit reconstruction of \bar{T} . Nevertheless, these classical results ensure that the relevant information is contained in quartets. More directly, we will show in this contribution that if we can reliably determine a suitable outgroup, best matches can be extracted from a small set of quartets.

Although much of this work is based on the assumption that an additive distance between taxa is available, one has to keep in mind that additive evolutionary distances, like divergence times, cannot be measured directly. While it is common practice to determine a dissimilarity $d'(x, y)$ of two taxa (genes) x and y from pairwise alignments, d' is a systematic under-estimate of the number of events d due to back-mutations, and thus not additive. In practice, the conversion of measurements of d' into an additive distance d that quantifies the number of evolutionary events is based on a Markov model of the evolutionary process. For sequence data, this may be the Jukes-Cantor model [Jukes and Cantor \[1969\]](#) or one of its more elaborate variants [Kimura \[1980\]](#), [Hasegawa et al. \[1985\]](#), [Tamura \[1992\]](#). In the most benign setting, d and d' are related by a monotone transformation, which in particular implies that the measured distances d' correctly identify the best hits. It can also be shown, however, that non-additive distances in general cannot identify the correct topology of quartets [Retzlaff and Stadler \[2018\]](#). Hence, we have no hope of computing correct best matches from the directly measured non-additive (dis)similarities.

This contribution is organized as follows: in the following section we give a rigorous mathematical

rendering of the background outlined above and use it to show that, given an additive distance measure, it is indeed possible to perfectly identify all best matches of a gene x of species s among its homologs $\{y_1, \dots, y_k\}$ in species t provided a suitable outgroup z can be found for every set $\{x, y_i, y_j, z\}$ of four genes. As a consequence, the practical problem becomes the reliable identification of correct “relative outgroups”. Assuming knowledge on the phylogeny of species from which the genes are taken, we proceed to derive several conditions under which z cannot be a correct choice and use this insights to devise a heuristic approach that works nearly perfect given additive distance data. We then introduce (in the Methods section) a simulation environment for generating gene family histories with complex rate variations and show that it is possible to recover best matches more accurately than approximating them by best hits.

2 Theory

2.1 Notation and basic definitions

Let T be a phylogenetic (gene) tree with leaf set L . For each gene $x \in L$ we denote by $\sigma(x)$ the species within which it resides. We write $L[s] = \{y \in L \mid \sigma(y) = s\}$ for the set of genes in species s . Given a rooted tree T with vertex set V , we write $\text{lca}(x, y)$ for the *last common ancestor* of $x, y \in V$, i.e., the vertex most distant from the root ρ that is shared by the paths from ρ to x and y , respectively. For a leaf set $L' \subseteq L$ we define the rooted tree $T[L']$ as the tree obtained from T by retaining only the vertices and edges along paths from the root to a leaf in L' and suppressing all vertices with degree 2.

Consider a gene x in species s . Among all genes in species $t \neq s$, the best matches of x are all those genes y in species t that have the lowest $\text{lca}(x, y)$. These y are the closest relatives of x in species t . This concept is made precise in

Definition 1. *Geiß et al. [2019b]* Let T be a phylogenetic tree with leaf set L (denoting genes) and $\sigma : L \rightarrow \mathcal{S}$ identifying the species $\sigma(x) \in \mathcal{S}$ in which a gene x resides. Then $y \in L$ is a best match of $x \in L$, in symbols $x \rightarrow y$, if $\text{lca}(x, y) \preceq \text{lca}(x, y')$ holds for all leaves y' from species $\sigma(y') = \sigma(y)$.

The best match relation \rightarrow is reflexive (since $\text{lca}(x, x) = x$), but it is neither transitive nor symmetric. Its mathematical properties are discussed in detail in Geiß et al. [2019b,a]. In particular, all orthologs of x are among its best matches Geiß et al. [2019c].

The evolutionary relatedness of two taxa x and y is most directly expressed by the divergence time $\tau(x, y)$, which is the total time elapsed in both lineages since the last common ancestor of x and y . Here, we consider only the case that all leaves refer to extant genes or taxa, i.e., $\tau(x, y) = 2\hat{\tau}(\text{lca}(x, y))$, where $\hat{\tau}$ is the age of $\text{lca}(x, y)$. Divergence times are ultrametric by definition. Furthermore, there is a well-known one-to-one correspondence between isomorphism classes of dated, rooted, phylogenetic trees and ultrametrics, cf. Böcker and Dress [1998], Semple and Steel [2003]. The best match relation \rightarrow can thus also be defined in terms of divergence time: $x \rightarrow y$ if and only if

$$y \in \arg \min_{y' \in L[t]} \tau(x, y') \tag{1}$$

The distinction between best hits and best matches thus is simply the distance function: best matches require divergence times, while best hits use one of several (dis)similarity measures for sequence data. They are equivalent under the Molecular Clock Hypothesis, which however fails for most real life data sets.

2.2 Reconciliation of gene tree and species tree

Since genes evolve as part of species, we can expect that *a priori* knowledge of the species phylogeny can be helpful for understanding the phylogeny of a gene family. This link is made precise by considering the embedding of a gene tree T into a species tree S .

We model the species tree S as a planted tree with leaf set \mathcal{S} , where 0_S is the planted root of S with its only child ρ_S . The embedding of the gene tree into the species tree is formalized by

the *reconciliation map* $\mu : V(T) \rightarrow V(S) \cup E(S)$, which maps duplications to the edges of S and speciations to the inner vertices $V^0(S)$ of the species tree. We follow here the notation of [Geiß et al. \[2019c\]](#). Restricting ourselves to duplication/loss scenarios, i.e., disregarding horizontal gene transfer, the reconciliation map satisfies the root constraint (R0) $\mu(x) = 0_S$ if and only if $x = 0_T$; the leaf constraint (R1) $\mu(x) = \sigma(x)$ for $x \in L(T)$, the ancestor preservation (R2), i.e., $x \prec_T y \implies \mu(x) \preceq_S \mu(y)$, and the following two speciation constraints for all speciation vertices $\mu(x) \in V^0(S)$: (R3.i) $\mu(x) = \text{lca}_S(\mu(v'), \mu(v''))$ for at least two distinct children v', v'' of x in T . (R3.ii) $\mu(v')$ and $\mu(v'')$ are incomparable in S for any two distinct children v' and v'' of x in T [Geiß et al. \[2019c\]](#). Equivalent axiom systems are considered e.g. in [Doyon et al. \[2011\]](#), [Rusin et al. \[2014\]](#), [Hellmuth \[2017a\]](#). Such reconciliation maps satisfy

$$\mu(x) \succeq_S \text{lca}_S(\sigma(L(T(x))))), \quad (2)$$

i.e., an event $x \in V(T)$ in the gene tree cannot be mapped to a node in the species tree below the last common ancestor of all the species. In this contribution we assume that μ in addition satisfies (R4): If $\mu(\text{lca}_T(x, y)) = \mu(\text{lca}_T(x, z)) \in V^0(S)$, then $\text{lca}_S(\sigma(x), \sigma(y)) = \text{lca}_S(\sigma(x), \sigma(z))$. In essence, (R4) ensures that a single node in T cannot represent two distinct speciation events, i.e., that the gene tree T is not “less resolved” than the species tree S into which it is embedded.

The reconciliation map μ defines *event labels* on the inner nodes of the gene tree T , identifying u as a duplication node if $\mu(u) \in E(S)$ and as speciation if $\mu(u) \in V(S)$. While it is possible to find a reconciliation map μ for every pair of gene and species tree [Górecki and Tiuryn \[2006\]](#), this is no longer true when event labels on T are given [Hernandez-Rosales et al. \[2012\]](#), [Hellmuth \[2017b\]](#). Conversely, T and S imply constraints on the event labels, identifying nodes that have to be duplications under *any* reconciliation map [Geiß et al. \[2019c\]](#). Here, we characterize these nodes further. We start from the following technical results:

Lemma 2 ([\[Geiß et al., 2019c, Lemma 2\]](#)). *Let T be a gene tree, S be a species tree and $\mu : V(T) \rightarrow V(S) \cup E(S)$ be a reconciliation map without horizontal gene transfer that does not necessarily satisfy (R4). Let $x \in V(T)$ be a vertex with $\mu(x) \in V^0(S)$. Then, $\sigma(L(T(v'))) \cap \sigma(L(T(v''))) = \emptyset$ for all distinct $v', v'' \in \text{child}(x)$.*

Let us first consider the case of binary gene trees:

Lemma 3. *Let T be a binary gene tree, S be a species tree, and $\mu : V(T) \rightarrow V(S) \cup E(S)$ be a reconciliation map without horizontal gene transfer that does not necessarily satisfy (R4). Let $x, y \in L(T)$ be two genes with $\sigma(x) \neq \sigma(y)$. If $\text{lca}_S(\sigma(x), \sigma(y)) \prec \mu(\text{lca}_T(x, y))$, then $\text{lca}_T(x, y)$ is a duplication event.*

Proof. Assume for contradiction $u := \text{lca}_T(x, y)$ is a speciation event, i.e., $\mu(u) \in V^0(S)$. Let v' and v'' be the two children of u in T . Observe that $u := \text{lca}_T(x, y)$ implies that $x \in L(T(v'))$ and $y \in L(T(v''))$ or *vice versa*. W.l.o.g. we assume that $x \in L(T(v'))$ and $y \in L(T(v''))$. By (R3.i) and (R3.ii), $\mu(u) = \text{lca}_S(\mu(v'), \mu(v''))$ and, in particular, $\mu(v')$ and $\mu(v'')$ are incomparable in S . Then by Lemma 2, we have $\sigma(L(T(v'))) \cap \sigma(L(T(v''))) = \emptyset$. This and R2 implies that $\mu(v') \succeq_S \sigma(x)$ and $\mu(v'') \succeq_S \sigma(y)$. The latter two arguments imply that $\text{lca}_S(\sigma(x), \sigma(y)) = \mu(u)$; a contradiction. \square

The assumption that T is binary is necessary here as the example in [Fig. 2](#) shows. Such reconciliations, however, cannot be meaningfully interpreted in terms of evolutionary events. Instead, the root of T confounds the duplication leading to x and y and the speciation separating $\text{lca}_S(\sigma(x), \sigma(y))$ from $\sigma(z)$. To suppress such undesirable cases, we in addition require that μ satisfies axiom (R4). In essence, (R4) forbids to map two distinct speciation events to the same vertex of S .

Lemma 4. *Let $\mu : V(T) \rightarrow V(S) \cup E(S)$ be a reconciliation map without horizontal gene transfer that satisfies (R4) and let $x, y \in L(T)$ be two genes with $\sigma(x) \neq \sigma(y)$. If $\text{lca}_S(\sigma(x), \sigma(y)) \prec \mu(\text{lca}_T(x, y))$, then $\text{lca}_T(x, y)$ is a duplication event.*

Proof. We assume that T is non-binary since the binary case is covered already by Lemma 3. Moreover, we assume, for contradiction, that $u := \text{lca}_T(x, y)$ is a speciation event, i.e., $\mu(u) \in V^0(S)$. Let v_x




Figure 2: The reconciliation μ with $\mu(\text{lca}_T(x, y, z)) = \text{lca}_S(X, Y, Z)$ satisfied (R1), (R2), (R3.i), and (R3.ii) but does not admit an unambiguous interpretation of $\text{lca}_T(x, y, z)$ as single event: it confounds the speciation separating Z and $\text{lca}_S(Y, X)$ with a gene duplication leading the ancestor of x and y or with the speciation separating X and Y . In either interpretation, the reconciliation map μ not correspond to a mechanistic explanation of the gene family history.

and v_y be the children of u with $x \preceq_T v_x$ and $y \preceq_T v_y$; thus we have $\sigma(x) \in \sigma(L(T(v_x)))$ and $\sigma(y) \in \sigma(L(T(v_y)))$. Since $u = \text{lca}_T(x, y)$, v_x and v_y are incomparable in T and hence $v_x \neq v_y$. By (R3.i), $\mu(v_x)$ and $\mu(v_y)$ are incomparable in S . Lemma 2 implies $\sigma(L(T(v'))) \cap \sigma(L(T(v''))) = \emptyset$ for all distinct children v' and v'' of u . The latter two fact together with (R2) imply $\text{lca}_S(\sigma(x), \sigma(y)) = \text{lca}_S(\mu(v_x), \mu(v_y)) \prec \mu(u)$. By (R3.i), $\mu(u) = \text{lca}_S(\mu(v'), \mu(v''))$ for some children v' and v'' of u , and thus $\text{lca}_S(\mu(v'), \mu(v'')) = \text{lca}_S(\sigma(z'), \sigma(z''))$ for some leaves $z' \in L(T(v'))$ and $z'' \in L(T(v''))$ from different species $\sigma(z') \neq \sigma(z'')$.

We proceed by showing that for at least one of $\sigma(z')$ and $\sigma(z'')$ we have $\text{lca}_S(\sigma(x), \sigma(z')) = \text{lca}_S(\sigma(z'), \sigma(z''))$ or $\text{lca}_S(\sigma(x), \sigma(z'')) = \text{lca}_S(\sigma(z'), \sigma(z''))$. Suppose that $\text{lca}_S(\sigma(x), \sigma(z')) \neq \text{lca}_S(\sigma(z'), \sigma(z''))$. Hence, $\text{lca}_S(\sigma(x), \sigma(z')) \prec_S \text{lca}_S(\sigma(z'), \sigma(z'')) = \mu(u)$. Therefore, $\text{lca}_S(\sigma(x), \sigma(z'')) = \text{lca}_S(\sigma(z'), \sigma(z''))$. Similarly, if $\text{lca}_S(\sigma(x), \sigma(z'')) \neq \text{lca}_S(\sigma(z'), \sigma(z''))$, then $\text{lca}_S(\sigma(x), \sigma(z')) = \text{lca}_S(\sigma(z'), \sigma(z''))$. Hence, assume w.l.o.g. that $\text{lca}_S(\sigma(x), \sigma(z')) = \text{lca}_S(\sigma(z'), \sigma(z'')) \neq \text{lca}_S(\sigma(x), \sigma(y))$. Now, by contraposition of (R4), we have $\mu(u) = \mu(\text{lca}_T(x, y)) \neq \mu(\text{lca}_T(x, z')) = \mu(u)$; a contradiction. \square

Lemma 4 conveniently generalizes to sets of genes:

Corollary 5. *Let $\mu : V(T) \rightarrow V(S) \cup E(S)$ be a reconciliation map without horizontal gene transfer that satisfies (R4) and let $A \subseteq L(T)$ with $|\sigma(A)| \geq 2$. If $\text{lca}_S(\sigma(A)) \prec \mu(\text{lca}_T(A))$, then $\text{lca}_T(A)$ is a duplication event.*

Proof. Note first that $\text{lca}_T(A) = \text{lca}_T(x, y)$ for some $x, y \in A$. Assume first $\sigma(x) \neq \sigma(y)$. Thus, $\text{lca}_S(\sigma(A)) \prec \mu(\text{lca}_T(A))$ implies $\text{lca}_S(\sigma(x), \sigma(y)) \preceq \text{lca}_S(\sigma(A)) \prec \mu(\text{lca}_T(A)) = \mu(\text{lca}_T(x, y))$. Hence, the statement follows from Lemma 4. If $\sigma(x) = \sigma(y)$, then $\text{lca}_T(A) = \text{lca}_T(x, y)$ implies that there are distinct children v_x and v_y of $\text{lca}_T(A)$ with $v_x \succeq x$ and $v_y \succeq y$. Thus, $\text{lca}_T(A) = \text{lca}_T(v_x, v_y)$. However, since $\sigma(x) = \sigma(y)$ we have $\sigma(L(T(v_x))) \cap \sigma(L(T(v_y))) \neq \emptyset$. Thus, Lemma 2 implies that $\mu(\text{lca}_T(A)) \notin V^0(S)$ and hence, $\text{lca}_T(A)$ is duplication. \square

2.3 Trees and (dis)similarities

Neither the divergence times nor the lca function of the phylogenetic tree T can be measured directly. The next-best choice is to work with an evolutionary distance, which measures the number of evolutionary events that have taken place to separate two taxa. For each edge $e = uv$ in T it is given by $\ell(e) = \int_{\hat{\tau}(u)}^{\hat{\tau}(v)} \mu_e(t) dt$, where $\mu_e(t)$ is the rate of evolution. In general $\mu_e(t)$ depends both on the lineage, and thus the individual edges in T , as well as on the exact point in time along e . It associates with each edge e a measure $\ell(e)$ of changes incurred, and thus an additive distance. If $\mu_e(t) = \mu_0$ is constant, we simply have $d_{\ell, T}(x, y) = \mu_0 \tau(x, y)$. This is the well-known Molecular Clock Hypothesis [Zuckerandl and Pauling \[1962\]](#), [Kumar \[2005\]](#).

In general, we consider $\ell : E(T) \rightarrow \mathbb{R}^+$ simply as an assignment of positive lengths to the edges of T , which we interpret as a measure proportional to the number of evolutionary events. This gives rise to

a metric distance function $d_{T,\ell}(x, y)$ on L defined as the sum of the lengths $\ell(e)$ of the edges e along the unique path connecting x and y in T . From T we obtain an associated *unrooted* tree \bar{T} by (i) omitting the planted root 0_T and its incident edge, and (ii), in case the root ρ in T has exactly two children u_1 and u_2 , by replacing the path $u_1\rho u_2$ by a single edge u_1u_2 with length $\ell(u_1u_2) := \ell(u_1\rho) + \ell(\rho u_2)$. Note that the dissimilarity function ℓ is by construction the same on T and \bar{T} . Thus \bar{T} determines T up to the position of the root, i.e., T is obtained from \bar{T} by inserting the root into an edge of \bar{T} or declaring an inner vertex of \bar{T} as the root. As for rooted trees, we define the restriction $\bar{T}[L']$ for some subset $L' \subseteq L$ by retaining only the vertices and edges along the paths between pairs of vertices in L' and then suppressing all vertices of degree 2. We note that $\bar{T}[L'] = \bar{T}[L']$.

A dissimilarity d on L is called *additive* if there is an unrooted tree \bar{T} with edge lengths ℓ such that $d = d_{\ell, \bar{T}}$. A key result in mathematical phylogenetics [Simões-Pereira \[1969\]](#), [Buneman \[1974\]](#) characterizes additive (pseudo)metrics as those that satisfy the *four point condition*. It states that d is additive if and only if the restriction of d to each subset L' of L with $|L'| = 4$, usually called a *quartet*, is additive and thus determines a tree on four leaves. Furthermore, the unrooted tree \bar{T} is uniquely defined by d . In principle, therefore, distance data completely determines a phylogenetic tree up to the position of the root.

The results of [Simões-Pereira \[1969\]](#), [Buneman \[1974\]](#) furthermore imply that \bar{T} can be expressed in terms of its four-taxa subtrees. This provides us with a natural possibility to consider only “local” topologies instead of the having to construct the unrooted tree \bar{T} explicitly. To this end, we consider the restrictions $\bar{T}[p, q, r, s]$ of \bar{T} to four distinct leaves $p, q, r, s \in L$ and define the *quartet relation* [Sattah and Tversky \[1977\]](#), [Fitch \[1981\]](#) $(pq|rs)$ if there is an edge e in \bar{T} , and thus in $\bar{T}[p, q, r, s]$, such that $\{p, q\}$ and $\{r, s\}$ are in different connected components of the forest obtained by removing e from \bar{T} or $\bar{T}[p, q, r, s]$. Equivalently, we have [Sattah and Tversky \[1977\]](#), [Fitch \[1981\]](#)

$$(pq|rs) \iff d(p, q) + d(r, s) < d(p, r) + d(q, s), d(p, s) + d(q, r). \quad (3)$$

In fact, for additive metrics, the two distance sums on the r.h.s. are equal [Simões-Pereira \[1969\]](#), [Buneman \[1974\]](#). All three terms are equal if and only if the four points form a star, whence the existence of a separating edge requires the strict inequality. By a slight abuse of notation we write $\bar{T}[p, q, r, s] = (pq|rs)$ if Equ. (3) holds, and $\bar{T}[p, q, r, s] = \times$ if no quartet exists on these four leaves, i.e., if $\bar{T}[p, q, r, s]$ is the star tree.

2.4 From Quartets to rooted triples

In a *planted phylogenetic tree* T with leaf set $L \cup \{0_T\}$ all inner vertices have degree at least 3. The special leaf 0_T identifies the ancestral state. Its only neighbor is the root ρ_T . The vertex set of T is endowed with a partial order \preceq such that $x \preceq y$ whenever y lies along the unique path connecting x and the planted root 0_T and thus also the root ρ_T . Thus the leaves are the minimal elements w.r.t. \preceq . Furthermore, for $A \subseteq L$ we define $\text{lca}(A) = \min\{z \mid x \preceq z \text{ for all } x \in A\}$, where the minimum is taken w.r.t. the partial order \preceq . For every $u \in V(T)$, we denote by $T(u)$ the subtree of T rooted in u . It will sometimes be useful to consider $T(u)$ as planted tree by including the unique parent v of u and the edge vu . The leaf set of $T(u)$ will be denoted by $L(T(u))$.

The most common method to specify the root of a phylogenetic tree is the use of so-called outgroups, that is, additional taxa that are known *a priori* to be outside a monophyletic group of interest. Given a planted (or rooted) phylogenetic tree, on the other hand, monophyletic groups are the leaf sets of a subtree, i.e., L' is a monophyletic group if and only if there is a vertex $u \in V(T)$ such that $L' = L(T(u))$. Every leaf $x \in L \setminus L'$ is an outgroup for L' .

Every edge in an unrooted tree \bar{T} defines a split $L'|L''$ of L , where L' and L'' are the leaves in the connected components of $\bar{T} \setminus e = \bar{T}' \cup \bar{T}''$. At most one of the two subtrees \bar{T}' and \bar{T}'' contains the root of the underlying phylogenetic tree T . If the root is not contained in \bar{T}' , then the tree $T' \cup \{e\}$ planted at the endpoint of e describes a monophyletic group. In this case all $x \in L''$ are outgroups for T' . Which subtrees of \bar{T} correspond to monophyletic groups is determined by the position of the root, and therefore requires external information.




Figure 3: Relation of last common ancestors $\text{lca}(x, y')$ and $\text{lca}(x, y'')$, resp., with quadruples on $\{x, y', y'', z\}$ with a trusted outgroup z .

It will be convenient in the following to define outgroups not only for monophyletic groups.

Definition 6. For a phylogenetic tree T with leaf set L , consider a subset $L' \subseteq L$ and a leaf $z \in L \setminus L'$. We say that z is an outgroup for L' if $\text{lca}(L') \prec \text{lca}(L', z)$.

Let us now return to the quartets of \bar{T} . The following simple result, illustrated in Fig. 3, shows that quartets can be used to infer inequalities between lca vertices in T provided one of the four leaves is known to be an outgroup for the other three:

Lemma 7. Suppose z is an outgroup for $\{x, y', y''\}$ in T . If $\bar{T}[x, y', y'', z]$ is fully resolved, then

(i) $\text{lca}(x, y') = \text{lca}(x, y'')$ iff $\bar{T}[x, y', y'', z] = (xz|y'y'')$,

(ii) $\text{lca}(x, y') \prec \text{lca}(x, y'')$ iff $\bar{T}[x, y', y'', z] = (xy'|y''z)$,
and

(iii) $\text{lca}(x, y') \succ \text{lca}(x, y'')$ iff $\bar{T}[x, y', y'', z] = (xy''|y'z)$.

Otherwise, $\bar{T}[x, y', y'', z] = \times$ and $\text{lca}(x, y') = \text{lca}(x, y'')$.

Proof. Since z is an outgroup by assumption, there are only three possible fully resolved rooted trees with $L = \{x, y', y'', z\}$, see Fig. 3. Each of these trees corresponds to a unique quadruple (annotated at the top). The relationship between $\text{lca}(x, y')$ and $\text{lca}(x, y'')$ is determined by the tree topology. The statement follows by inspecting the three cases. If $\bar{T}[x, y', y'', z]$ is not fully resolved, no quartet is defined on $\{x, y', y'', z\}$, i.e., \bar{T} is the star tree and thus $\text{lca}(x, y') = \text{lca}(x, y'') = \text{lca}(y', y'')$. \square

Observation 8. If $u' = \text{lca}(x, y')$ and $v' = \text{lca}(x, y'')$ for $x, y', y'' \in L$, then u' and v' are comparable w.r.t. \preceq in T .

Lemma 7 together with Obs. 8 implies that quadruples with known outgroups can be used to identify best matches. More precisely, in order to determine the set $\{y \in L[s] \mid x \rightarrow y\}$ it suffices to consider leaf sets $\{x, y', y'', z\}$ with $y', y'' \in L[s]$ such that z is an outgroup for $\{x, y', y''\}$. By Lemma 7, any set of this type implies an (in)equality between $\text{lca}(x, y')$ and $\text{lca}(x, y'')$. It may not be necessary to consider all quadruples. To explore ways to reduce the computational efforts, let us assume that for given $x \in L$ and $s \in S$, $s \neq \sigma(x)$, we can identify sets $Y \subseteq L[s]$ and $Z \subseteq L$ such that the following three assumptions are satisfied:

- (A0) The noise in the data is small enough so that for any four taxa $\{x, y', y'', z\}$ with $y', y'' \in Y$ and $z \in Z$ one of the three possible quartets or the star topology is inferred correctly.
- (A1) The candidate set $Y \subseteq L[s]$ contains all best matches of x in species s (but usually also additional leaves).
- (A2) Z is a non-empty set outgroups for $Y \cup \{x\}$.

Before we proceed, let us consider these three assumptions in some more detail. (A0) is satisfied by construction for additive distance data. In real-life applications it is often possible to obtain at least a very good approximation using explicit models of sequence evolution. In addition, several computational approaches have been proposed to estimate the quartet relation directly from sequence data. It is also worth noting that (A0) does not require precise distance data, it only asks for correct categorical data on the quartet relation.

Condition (A1) can always be enforced by setting $Y = L[s]$. We make this assumption explicit because in practice it will be desirable to work with small subsets $Y \subseteq L[s]$ as using $L[s]$ may be too expensive for large gene families. The inclusion of very distant relatives may be problematic for the construction of good multiple sequence alignments and thus the extraction of the quartet relation. Furthermore, it may be difficult to find suitable outgroup data in this case. Thus we will limit Y to a manageable size and sufficient sequence similarity. In `ProteinOrtho` Lechner et al. [2011], for example, $Y \subseteq L[s]$ is defined as the set sequence with `blast` bit-scores exceeding a certain fraction of the best hit for x in species s .

Condition (A2), i.e., the knowledge of appropriate outgroups, is the only problematic assumption. As discussed above, distance-based methods by construction do not convey information on the root of the phylogenetic tree T but only determine its unrooted version \bar{T} . As a consequence, additional information, not contained in the pairwise distance measurements, is necessary to determine the edge in \bar{T} that harbors the position of the root ρ of T Penny [1976]. In general, Z will be chosen from one or more species that are outgroups to $\sigma(x)$ and s in S . Even if outgroup species are given, gene duplications may pre-date the divergence of the available species set, so that a given data set will usually violate (A2) for some pairs of leaves. We will return to these issues in more detail in the following sections.

Algorithm 1 Overall Workflow

Require: reference vertex x

- 1: retrieve a sufficient set $Y \subseteq L[s]$ of candidate best matches for x with color s
 - 2: determine a set Z of outgroup vertices for $Y \cup \{x\}$
 - 3: initialize an edgeless digraph Γ with vertex set Y
 - 4: **for all** pairs $y', y'' \in Y$ **do**
 - 5: **for all** $z \in Z$ **do**
 - 6: determine significantly supported quartet on $\{x, y', y'', z\}$
 - 7: determine consensus quartet over all choices of $z \in Z$
 - 8: **if** consensus quartet implies $\text{lca}(x, y_1) \preceq \text{lca}(x, y_2)$ **then**
 - 9: insert the directed edge (y_2, y_1) into Γ
 - 10: compute the strongly connected components of Γ
 - 11: report strongly connected components without out-edges as the set of best matches $\{y \in Y \mid x \rightarrow y\}$
-

The discussion so far suggests to use the quadruple relation for sets of the form $\{x, y', y'', z\}$ with $y', y'' \in Y$ and $z \in Z$ to determine the best matches of x in the species containing the homolog set Y . The procedure is summarized in Alg. 1. The main result of this section establishes its correctness.

Theorem 9. *Algorithm 1 correctly identifies the set of best matches of x with color s as the unique strongly connected component of Γ without out-edges provided assumptions (A0), (A1), and (A2) are satisfied.*

Proof. Assumptions (A1) and (A2) imply that comparison of the last common ancestors can be performed in terms of the quartets according to Lemma 7, which by assumption (A0) are all inferred correctly. Therefore, lines 4-7 compute all quartets correctly, and thus the inequality between $\text{lca}(x, y_1)$ and $\text{lca}(x, y_2)$ is inferred correctly. The auxiliary graphs Γ therefore contains at least one arc between any two vertices $y', y'' \in Y$ and both the arc (y', y'') and (y'', y') if and only if $\text{lca}(x, y') = \text{lca}(x, y'')$, i.e., the strongly connected components are cliques. Since the $\text{lca}(x, y)$ are interior vertices of T that are totally ordered along the path from x to the root of T (Observation 8), there is a unique strongly

connected component B in Γ that has no out-edges, whose vertices are those $y \in B$ for which $\text{lca}(x, y)$ is minimal. Thus B is the set of best matches of x with color s . \square

Algorithm 1 therefore works correctly at least under idealized assumptions. It also serves as a heuristic in cases where one of the assumptions (usually (A2)) is violated.

2.5 Identification of outgroups

In many practical applications, the phylogenetic relationships between the *species* under consideration are known. We therefore investigate here to what extent knowledge of the species tree S can help to identify good outgroup sets Z . Ideally, the genes chosen as outgroups Z are co-orthologs of the focal gene set Y , i.e., the duplication event that produced y' and y'' occurred after the speciation event that separates $\sigma(z)$ for all $z \in Z$ from $\sigma(X)$ and $\sigma(Y)$. As we shall see, it is not possible to identify outgroups with complete certainty. It is possible, however, to identify incorrect choices in many situations.

In the following we consider three species $\sigma(X)$, $\sigma(Y)$, and $\sigma(z)$ for $z \in Z$ such that

$$\text{lca}_S(\sigma(X), \sigma(Y)) \prec_S \text{lca}_S(\sigma(X), \sigma(Y), \sigma(z)), \quad (4)$$

i.e., $\sigma(z)$ is an outgroup in the species tree for $\sigma(X)$ and $\sigma(Y)$. Problematic cases in which quartets are interpreted incorrectly may appear whenever the duplication event $\text{lca}_T(y', y'')$ separating two paralogs $y', y'' \in Y$ pre-dates the speciation event separating $\sigma(z)$ from $\text{lca}_S(\sigma(X), \sigma(Y))$. We capture this situation in

Definition 10. *Let u be an inner node of the species tree S , let $y', y'' \in Y$ be paralogs in a species $\sigma(Y) \in L(S(u))$. Then $\text{lca}_T(y', y'')$ is an ancient duplication relative to $u \in V(S)$ for the reconciliation map $\mu : V(T) \rightarrow V(S) \cup E(T)$ if $u \prec_S \mu(\text{lca}_T(y', y''))$.*

Clearly, if $\text{lca}_T(y', y'')$ is an ancient duplication relative to $\text{lca}_S(\sigma(X), \sigma(Y), \sigma(z))$, then genes in $z \in Z$ are bad choices as outgroups $\{x, y', y'', z\}$. The difficulty is that we do not know the reconciliation map μ in our setting. In some cases, however, it is possible to identify vertices in T that are ancient duplications relative to some speciation for *any* reconciliation. Such cases can then be avoided.

Before we investigate possibilities to identify some ancient duplications in distance data, we prove a rather technical result that shows that in cases without too many ancient duplications, Algorithm 1 produces correct results. For the proof we will need to consider the reconciliation map μ for *complete* gene family histories, i.e., gene trees T containing extant genes as well as all branches leading to loss events. As above, we do not consider HGT. The leaf set of T is thus $L := L_e \cup L_0$, where L_e represents the extant genes and L_0 denotes loss events. Since the species map is naturally restricted to extant genes (i.e., $\sigma : L_e(T) \rightarrow L(S)$), we need to restrict (R1): If $x \in L_e(T)$, then $\mu(x) = \sigma(x)$. We will refer to such gene trees and reconciliation maps as *extended* gene trees and *extended* reconciliation maps, respectively. Correspondingly, Lemma 2 only holds for L_e , i.e., we can conclude that $\sigma(L_e(T(w_1))) \cap \sigma(L_e(T(w_2))) = \emptyset$. This can easily be seen by reusing the contradiction argument in Geiß et al. [2019c][Lemma 2]. As a consequence of loss events we now may have $\sigma(L_e(T(v))) = \emptyset$ for some nodes $v \in V(T)$.

Lemma 11. *Let (T, σ) be an extended gene tree with a non-empty set of extant genes $L_e = X \cup Y \cup Z$ with $|\sigma(Z)| = 1$, let S be a species tree on $S = \{\sigma(X), \sigma(Y), \sigma(Z)\}$ such that $\text{lca}_S(\sigma(X), \sigma(Y)) \prec \text{lca}_S(\sigma(X), \sigma(Y), \sigma(Z)) = \rho_S$, and let μ be an extended reconciliation map from (T, σ) to S . If (A0) holds and $|\mu^{-1}(\rho_S)| \leq 2$, then Algorithm 1, using Y as the candidate best match set and Z as outgroup set, correctly determines, for every gene $x \in X$, all best matches in species $\sigma(Y)$.*

Proof. First note that the statement is trivial if there exists only one gene in Y . Hence, we can assume that Y contains more than one gene. Moreover, Condition (A1) is trivially satisfied since, by assumption, the candidate set of best matches of x in Y is exactly Y . Since L_e is non-empty, we have to consider the two cases $|\mu^{-1}(\rho_S)| = 1$ and $|\mu^{-1}(\rho_S)| = 2$.




Figure 4: Minimal examples in which ancient duplications lead to false positives (FP) or false negatives (FN) when choosing outgroups as described in the text. (A) The species tree S displaying the triple $(\sigma(X)\sigma(Y)|\sigma(Z))$. (B-D) Gene trees T with two ancient duplications. We assume that y' and y'' are the only extant genes of color $\sigma(Y)$, i.e. color $\sigma(Y)$ is extinct in the subtree of x in each of the shown cases. The asterisk marks the discriminating edge for the quartet inference. (B) A quartet $(xy'|y''z)$ is inferred so that only y' but not y'' is a best match, (x, y'') is a FN. (C) A quartet $(xz|y'y'')$ is inferred so that y' is a false best match, (x, y') is a FP. (D) A quartet $(xy'|y''z)$ is inferred so that (x, y') is a FP and (x, y'') is a FN.

Assume first $|\mu^{-1}(\rho_S)| = 1$, i.e., there exists exactly one $v \in V(T)$ such that $\mu(v) = \rho_S$. We then have $\sigma(L_e(T(w_1))) \cap \sigma(L_e(T(w_2))) = \emptyset$ for any distinct $w_1, w_2 \in \text{child}_T(v)$ (cf. Geiß et al. [2019c][Lemma 2], Lemma 2), which, by construction of the species tree S , immediately implies $\sigma(L_e(T(w))) \in \{\{\sigma(X), \sigma(Y)\}, \{\sigma(Z)\}\}$ for any $w \in \text{child}_T(v)$. Hence, Z is an outgroup set for $Y \cup \{x\}$, i.e., Condition (A2) is satisfied, and the statement thus follows directly from Theorem 9.

Now suppose $|\mu^{-1}(\rho_S)| = 2$, i.e., there are exactly two distinct $v_1, v_2 \in V(T)$ with $\mu(v_1) = \mu(v_2) = \rho_S$. Let $T_1 := T(v_1)$ and $T_2 := T(v_2)$ be the subtrees of T rooted at v_1 and v_2 , resp., and assume w.l.o.g. $x \in L_e(T_1)$. Note that $L_e(T_1) \cup L_e(T_2) = L_e$. Let $w_1 \in \text{child}_T(v_1)$ such that $x \preceq_T w_1 \prec_T v_1$. If w_1 were mapped to an edge or vertex along the path from ρ_S to $\sigma(Z)$, then $\text{lca}_S(\sigma(X), \sigma(Y)) \prec \text{lca}_S(\sigma(X), \sigma(Y), \sigma(Z)) = \rho_S$ would imply $\sigma(X) \not\preceq_S \mu(w_1)$; a contradiction to (R2). Thus, $\sigma(Z) \notin \sigma(L_e(T(w_1)))$. Since $\mu(v_1) \in V^0(S)$, Condition (R3.i) implies that there exists $w_2 \in \text{child}_T(v_1)$, $w_2 \neq w_1$, such that $\mu(v_1) = \text{lca}(\mu(w_1), \mu(w_2))$. Since $\sigma(L_e(T(w_1))) \cap \sigma(L_e(T(w_2))) = \emptyset$ by Lemma 2, we obtain $\sigma(L_e(T(w_1))) \subseteq \{\sigma(X), \sigma(Y)\}$ and $\sigma(L_e(T(w_2))) \subseteq \{\sigma(Z)\}$. We distinguish the two cases (a) $\sigma(Y) \notin \sigma(L_e(T_1))$ and (b) $\sigma(Y) \in \sigma(L_e(T_1))$.

Case (a): If $\sigma(Y) \notin \sigma(L_e(T_1))$, any leaf $y \in Y$ must reside within a subtree $T(w')$ with $w' \in \text{child}_T(v_2)$, thus all genes in Y are best matches of x . Since the speciation node v_2 separates $\sigma(Z)$ from $\sigma(X)$ and $\sigma(Y)$, we have $\sigma(Z) \notin \sigma(L_e(T(w')))$ for any such w' (cf. Lemma 2). Moreover, reusing the same arguments as for v_1 , we conclude that there exists exactly one such $w' \in \text{child}_T(v_2)$ such that




Figure 5: Incongruence of gene and species quadruples implies the existence of an ancient duplication. Consider four pairwise distinct species A , B , C , and D whose species tree is given on the l.h.s., and let four genes a , b , c , and d be chosen such that $\sigma(a) = A$, $\sigma(b) = B$, $\sigma(c) = C$, and $\sigma(d) = D$. The two speciation events separating A from B and C from D are indicated by \circ . The root of this tree is indicated by \blacksquare . Of the three possible gene quartets, one is congruent with the species tree. The other two are incongruent. In each of these, Equ. (2) implies that the two interior vertices in these quartets cannot be mapped to the species tree below the root. The root of the gene tree must thus be mapped above the root of the species tree.

$\sigma(Y) \in \sigma(L_e(T(w')))$. Hence, any two distinct $y, y' \in Y$ reside within the same subtree $T(w')$ and thus $\text{lca}_T(x, y) = \text{lca}_T(x, y')$. Since $\sigma(Z) \notin \sigma(L_e(T(w')))$, this immediately implies $\bar{T}[x, y, y', z] = (xz|yy')$ for any $z \in Z$. Hence, Γ is the complete graph, i.e., any gene of species $\sigma(Y)$ is correctly inferred as a best match of x .

Case (b): Assume, for contradiction, that there exists $w_3 \in \text{child}_T(v_1) \setminus \{w_1\}$ such that $\sigma(Y) \in \sigma(L_e(T(w_3)))$. Clearly, $w_3 \neq w_2$. Since it must hold $\sigma(L_e(T(w_1))) \cap \sigma(L_e(T(w_3))) = \emptyset$ as well as $\sigma(L_e(T(w_2))) \cap \sigma(L_e(T(w_3))) = \emptyset$ by Lemma 2, we conclude $\sigma(L_e(T(w_1))) = \{\sigma(X)\}$ and $\sigma(L_e(T(w_3))) = \{\sigma(Y)\}$. However, (R4) then implies $\text{lca}_S(\sigma(X), \sigma(Y)) = \text{lca}_S(\sigma(Y), \sigma(Z))$; a contradiction. Hence, there exists an extant gene $y \preceq_T w_1$ in Y . Then, as $\sigma(Z) \notin \sigma(L_e(T(w_1)))$, any $z \in Z$ infers the same quartet on $\{x, y, y', z\}$, $y' \in Y \setminus \{y\}$. We therefore conclude that the auxiliary graph Γ contains a unique strongly connected component without out-edges, which represents the set of best matches of x in Y . Note that in these cases Condition (A2) is not necessarily satisfied, but Algorithm 1 still provides the exact solution. \square

The condition $|\mu^{-1}(\rho_S)| \leq 2$ makes an explicit assumption on the true history of the gene family by limiting the scenario to at most one ancient duplication on $X \cup Y \cup Z$. Fig. 4 shows that this condition cannot be dropped: if there are two or more ancient duplications affecting X , Y , and Z , then the correct inference of best matches from quartets can no longer be guaranteed. It is important to note that the condition $|\mu^{-1}(\rho_S)| \leq 2$ cannot be checked in real data since μ is unknown. In the simulated data, however, it is easy to validate and we observed empirically that it is rarely violated in our data (see Simulation Results section).

In some situations ancient duplications can be inferred unambiguously, independent of the reconciliation map μ . This is in particular the case if there are incongruences between quartets of genes and species. Consider four genes a, b, c, d residing in four pairwise distinct species $\sigma(a)$, $\sigma(b)$, $\sigma(c)$, and $\sigma(d)$, and assume that these four species form the quartet $(\sigma(a)\sigma(b)|\sigma(c)\sigma(d))$. Then we say that the gene and species quartets are *congruent* if $\bar{T}[a, b, c, d] = (ab|cd)$ or \times . Otherwise, i.e., for $\bar{T}[a, b, c, d] \in \{(ac|bd), (ad|bc)\}$, we say they are *incongruent*, see Fig. 5. In the following we show that the incongruence of gene and species quartets implies ancient duplications. More precisely:

Theorem 12. *Let (T, σ) and S be gene and species trees, respectively, and $a, b, c, d \in L(T)$. Moreover, let $\sigma(a)$, $\sigma(b)$, $\sigma(c)$, and $\sigma(d)$ be pairwise distinct species, set $u := \text{lca}_S(\sigma(a), \sigma(b), \sigma(c), \sigma(d))$, $v_1 := \text{lca}_S(\sigma(a), \sigma(b))$, and $v_2 := \text{lca}_S(\sigma(c), \sigma(d))$. If $v_1 \prec_S u$, $v_2 \prec_S u$ and $\bar{T}[a, b, c, d] = (ac|bd)$ or $\bar{T}[a, b, c, d] = (ad|bc)$, then $u \prec_S \mu(\text{lca}_T(a, b, c, d))$ for every reconciliation map $\mu : V(T) \rightarrow V(S) \cup E(S)$ without HGT events. In particular, $\text{lca}_T(a, b, c, d)$ is a duplication event.*




Figure 6: Recall, precision, and F-measure of the true best matches as a function of ϵ for simulated data (2000 scenarios).

Proof. By assumption, $S[\sigma(a), \sigma(b), \sigma(c), \sigma(d)]$ has the topology shown in Fig. 5. Assuming $(ac|bd)$, Equ. (2) implies $\mu(\text{lca}_T(a, c)) \succeq \text{lca}_S(\sigma(a), \sigma(c)) = u$ and $\mu(\text{lca}_T(b, d)) \succeq \text{lca}_S(\sigma(a), \sigma(c)) = u$. Thus both inner nodes p and q of the quartet are mapped no lower than u . The edge between them therefore must be mapped to an edge pre-dating u , since the speciation constraint (R3) implies that two \prec_T -comparable events in T of which one is a speciation cannot be mapped to the same vertex of S . Thus $u \prec_S \mu(\text{lca}_T(a, b, c, d))$. The case $(ad|bc)$ is handled by an analogous argument exchanging c and d . The fact that $\text{lca}_T(a, b, c, d)$ is a duplication event now follows from Lemma 4. \square

This theorem can be used to discard suspicious outgroups: If $\bar{T}[x, y, z_1, z_2]$ is incongruent with the known species tree, then $\sigma(z_1) \neq \sigma(z_2)$ should be replaced by outgroup candidates from earlier-branching species. The downside of using Theorem 12 is that it requires the systematic investigation of a possibly large numbers of quartets.

We suspect that it is possible in most cases to unambiguously identify pairs whose last common ancestor in the gene tree pre-dates the last common ancestor of the species tree under consideration. While it may be difficult to determine the relative order of such duplications, we suspect that clustering methods used to extract groups of co-orthologs (COGs) can be adapted to disentangle such ancient “paralog groups”.

3 Simulation Results

Well curated gene family histories are not available at large scale. We therefore use simulated data to evaluate how well best matches (in the sense of evolutionary relatedness) can be estimated from both perfect and noisy evolutionary distance measurements. For this purpose, it is important to have data sets that emphasize asymmetric rate variations among paralogs, i.e., the situations in which sequence dissimilarities and divergence times are not well correlated. We therefore developed a simulation system (see Methods) that can produce this type of test data and explicitly records the gene family history.

We compare three strategies to estimate best matches:

1. Reciprocal best hits are inferred directly from the distance data. In order to account for rate variations among paralogs, we follow the strategy of `ProteinOrtho` Lechner et al. [2011] and consider nearly co-optimal best hits by considering for a given gene x in species $\sigma(x)$ all those $y \in Y$ as almost best hits that have distance not worse than a factor $1 + \epsilon$ than the most similar gene in Y . In symbols: $H(Y|x) := \{y \in Y \mid d(x, y) \leq (1 + \epsilon) \min_{y' \in Y} d(x, y')\}$

For further comparison we then chose the value of ϵ^* that maximizes the F-measure ($\epsilon^* = 0.5$, see Fig. 6). Still this approach produces a substantial number of both false positives and false negatives in data sets with substantial rate variations. We expect that the optimal value of ϵ^* will depend on the details of the data set, in particular on the extent of evolution rate asymmetries. In general these will have to be estimated from the gene family history. We refer to this approach as the “ ϵ -method”.




Figure 7: Performance comparison of the best match inference methods for simulated data (2000 scenarios). Top panel: Median (solid) and 10th percentile (dashed) of recall and precision as a function of noise level. Lower panel: Boxplots of F-measure for different levels of noise superimposed on the additive distance; $\sigma = 0$ refers to perfect data. Blue: ϵ method, orange: explicit construction of the unrooted tree \bar{T} and midpoint rooting, green: inference of quartets with outgroups chosen in another branch of the root.

Since we chose the cut-off ϵ^* to maximize the F -measure, we effectively determine an upper bound on the performance of the Best Hit approach.

2. Explicit reconstruction of \bar{T} . Since the additive distances completely determine \bar{T} , the only source of errors for perfect data is an incorrect position of the root of T . For additive distance data, the Neighbor Joining algorithm [Saitou and Nei \[1987\]](#) is guaranteed to produce the correct \bar{T} [Atteson \[1999\]](#). We then use midpoint rooting [Hess and de Moraes Russo \[2007\]](#) to pass from \bar{T} to T and compute the best matches in T . We refer to this method as “NJ+midpoint rooting”. This method is not intended as a viable means of analysis for real-life sequence data. It serves, however, as a convenient way to assess the effects of rate imbalances because it isolates the errors that are introduced by the choice of the root alone i.e., by rate imbalances.

3. The “Quartet” approach starts from a known species tree S . For $x \in X$, and $y', y'' \in Y$ we pick the outgroup z from a species $\sigma(z)$ such that $\text{lca}_S(\sigma(X), \sigma(Y)) \prec \text{lca}_S(\sigma(X), \sigma(Y), \sigma(z))$ and then use Algorithm 1. We explored two different ways of determining the quartet relation: (a) After transforming the measured sequences distances into an approximately additive evolutionary distance (see Methods for details), we used Equ. (3), and (b) for sequence data we used the direct method described in the Methods section, thus bypassing the computation of distances altogether.

In order to benchmark the inference of best matches we compute recall and precision w.r.t. the true best matches restricted to pairs of gene sets X and Y for which such outgroups are available. The comparison in Fig. 7 (bottom panel) shows that the quartet method outperforms the alternatives for different levels of the simulated measurement error. The results are robust over a wide range of simulated measurement error. Not surprisingly, the reconstruction of Neighborjoining trees already provides better results than the ϵ -method. The simple midpoint rooting strategy however still incurs noticeable level of error. For the quartet method operating on noiseless data the only source of errors are bad choices of outgroups, which are the consequence of ancient duplications. The number of ancient duplication exceeds 1 in 5.15% of the simulated gene family scenarios. Due to loss events predating the root of the species tree, the condition in Lemma 11 is only violated in 3.7% of the gene trees. Out of these problematic cases, little more than half (2.25%) actually result in a non-perfect




Figure 8: Inference of best matches from simulated sequence data. Heat map of the fraction of false positive best matches inferred by QM as a function of the number of duplication and loss events in the simulated scenario. The false positive rate is computed relative to the number of true best matches.




Figure 9: Performance comparison of two different outgroup choice methods: (1) outgroups chosen randomly from species in another branch of the root (green, same as in Fig. 7), (2) closest outgroups in all relative outgroup species corrected with Theorem 12 (red). Left and middle: Median (solid) and 10th percentile (dashed) of recall and precision as a function of noise level. Right: Boxplots of the F-measure differences for selected noise levels (method 2 – method 1).

inference accuracy.

Moreover, we investigated how the number of duplication and losses influences the inference of false positives inferred by QM (see Fig. 8). As expected, the number of false positives increases with increasing duplication and loss events.

Restricting the choice of outgroup genes z to species that are separated from X and Y by the root of the species tree, i.e., such that $\text{lca}_S(\sigma(z), \text{lca}_S(\sigma(X), \sigma(Y))) = \rho_S$, is likely to be problematic whenever S is skewed in a way that leaves very few choices for $\sigma(z)$ and whenever the divergence between $\sigma(z)$ and $\text{lca}_S(\sigma(X), \sigma(Y))$ is large. In the latter case, saturation effects may impair the quartet inference in fast evolving gene families. Hence, it would be advantageous to consider also genes from closer species. In principle, every relative outgroup w.r.t. species $\sigma(X)$ and $\sigma(Y)$ is a viable candidate. These can then be filtered by applying Theorem 12 to reduce the number of bad choices of z . We find that filtering for outgroups with identifiable ancient duplications and giving preferences to the closest outgroup genes, i.e., those with the lowest $\text{lca}_S(\sigma(z), \text{lca}_S(\sigma(X), \sigma(Y)))$ indeed yields a further moderate improvement of the estimated best matches (see Fig. 9). However, the performance is slightly reduced for perfectly additive data due to ancient duplications that were not detected by the currently available filtering heuristics.

In applications to real-life data sets, additional uncertainties arise through the reconstruction of distances from sequences. We therefore simulated sequences with and without in/dels from the gene




Figure 10: Comparison of best match estimates for short nucleic acid sequences (200 nt), long nucleic acid sequences (2000 nt), and amino acid sequences (500 aa). As expected the F-measure improves when saturation is decreased by down-scaling the total number of events, and with increasing sequence length. The disappointing performance of the QM method is probably the consequence of a majority voting procedure that is too simple-minded. So far, the best results are obtained by estimating additive distances from pairwise sequence alignments using the Jukes-Cantor transform (for nucleic acid sequences) or a BLOSUM-based transformation (for amino acid sequences).

tree/species tree scenarios and inferred the best matches from the sequence data. We compared the results obtained (a) by transforming the Hamming distances using the Jukes-Cantor transform and (b) by direct inference of the quadruples using quartet mapping as outlined in the Methods section.

Fig. 10 summarized the results for simulated nucleic acid and amino acid sequences of different lengths and different scaling of the evolutionary rates. As expected, the short sequences incur a relative large noise level compared to the perfect additive distances. Nevertheless, the overwhelming majority of best matches is still estimated correctly (F -measures well above 0.9 for the vast majority of scenarios even for nucleic acid sequences as short as 200 nt). Larger false positive rates are observed only in a small number of scenarios with many duplications and losses. This is not surprising since our relatively simple rule for outgroup choice tends to fail if there are many ancient duplications. As expected, the F-measure improves with increasing sequence length due to the increased amount of information from which the distances are estimated. The same trends were observed for simulated protein sequences

The performance of the Quartet Mapping strongly depends on how those quartets are handled for which none of the three possible splits dominates. In the default setting, these are interpreted as unresolved tree (\times) and inserted as bi-directional edges into the auxiliary graph Γ . This, however, leads to a moderate overprediction of best matches. Alternatively, a consensus can be taken over multiple choices of the outgroup z . Finally, the unresolved quadruples can be omitted altogether in the construction of the auxiliary graph Γ . Both alternatives perform worse than the default method, see Additional File 1.

4 Discussion and Conclusions

The idea to use quartet structures for improvement of orthology estimates is not new; it was used e.g. in `QuartetS` Yu et al. [2011]. Quartets are also used as witnesses of non-orthology in OMA Train et al. to avoid some types of false-positives. Here, we systematically investigate how and when quartets help to improve and/or correct empirical best-hit data to identify best matches in the sense of closest evolutionary relatives. We propose that reciprocal best matches, rather than the uncorrected reciprocal best hits, should then be used to infer orthology relationships. This second step is the topic of an independent manuscript Geiß et al. [2019c] in which the mathematical connections between (reciprocal) best matches and orthology are elucidated in detail.

The key observation of the present contribution is that the best matches of a gene x in the set Y of genes from a different species can be computed correctly if for every $y', y'' \in Y$ one can find a gene z from a third species that is an outgroup for $\{x, y', y''\}$. From a theoretical point of view, this condition is closely related to rooting the gene tree. The second necessary ingredient is an estimate of an additive evolutionary distance between the genes that is accurate enough to correctly identify the topology of a certain subset of quadruples. We emphasize that this is a much less stringent condition compared to the ability of reconstructing the complete gene tree T .

Empirically, we observe that (partial) knowledge of the species tree (more precisely: reliable monophyletic groups) is very useful for the choice of outgroup genes z : excellent results are obtained by choosing a candidate z from a species that is an outgroup for $\sigma(x)$ and $\sigma(Y)$. The results can be further improved by using filtering criteria that identify ancient duplication events and by computing a consensus over several choices of z . In data sets with little measurement noise, we indeed obtain nearly perfect best match estimates. The theoretical considerations outlined here also suggest additional in-roads for further improvements by means of identifying ancient duplications, which not only serve as “witnesses of non-orthology” but can also be used to prune the set of candidate outgroups.

In order to make the methods described here applicable to very large real-life data sets, it will be necessary to optimize the computational performance. To this end, we will develop heuristic rules to prune the set Y in the case of large gene families. An obvious candidate is to use the ϵ -method as an initial filter, where ϵ is now chosen to optimize the tradeoff between $|Y|$ and false negative predictions of best matches. We expect that the heuristic rules for choosing the set Z of candidate outgroups can also be improved substantially. We expect that methods for orthology assessment can be improved in both reliability and computational performance by combining the accurate estimation of best matches described here with a better understanding of (reciprocal) best match graphs Geiß et al. [2019b,a] and their connection with the orthology relation Geiß et al. [2019c]. Since tree-free methods for orthology detection rely on (pairwise) best hits as proxy for reciprocal best matches, we expect that the accuracy of most tools would improve if best matches are supplied as input data. This is not easy to test, however, since the best hit computation is usually an integral part of the the software. Such a benchmark study is hence beyond the scope of this contribution.

The work reported here is primarily intended to provide a solid theoretical foundation for the construction of improved best match heuristics. The theoretical results give some guarantees for obtaining the correct best matches and highlight some limitations that cannot be overcome with certainty as long as only distance data are available.

5 Methods

5.1 Simulations of *dated* Species Trees

As in previous work Geiß et al. [2019c], we use the Innovation Model Keller-Schmidt and Klemm [2012] to produce realistic topologies for the planted species tree S . We then construct a dating function $\tau : V(S) \rightarrow [0, 1]$ such that $\tau(0_S) = 1$ and $\tau(x) = 0$ for $x \in L(S)$. In order to assign a date to an interior vertex, we traverse S top-down, more precisely for the current node u at time $\tau(u)$ we proceed as follows:

- (1) We pick a child $v \in \text{child}(u)$ and a leaf $x \in L(S(v))$ in the subtree below v . If v is already a leaf,

we set $\tau(v) = 0$ and proceed to the next child of u .

(2) Otherwise, we determine the number k of speciations on the path between v and x . Hence, the path from u to x comprises $k + 2$ edges.

(3) We pick a random number r with mean 1 and range $(0, 2)$ from a uniform distribution and set $\tau(v) = \tau(u)(1 - r/(k + 2))$. This rule is chosen so that the expected time elapsed along the edge uv equals $\tau(u)$ divided by the number $(k + 2)$ of edges along the path to the root and ensures that $\tau(v) > 0$.

The result is a dated species tree in which each edge uv has length $\tau(u) - \tau(v)$.

5.2 Simulation of gene trees in the dated species tree

We use the Gillespie algorithm Gillespie [1977] to simulate the duplication, loss and horizontal gene transfer events (HGT) occurring in S . The branches of the species tree S are independent in Duplication/Loss scenarios. However, horizontal gene transfer introduces dependencies between them. We therefore have to simulate the evolution process in such a way that at each time point τ the possible reactions are given by the Cartesian product $G(\tau) \times \{D, L, H\}$, where $g \in G(\tau)$ is a gene that is present at time τ in any one of the branches of the dated species tree, and $q \in \{D, L, H\}$ is one of the three possible events (Duplication, Loss, HGT). Every possible simulation event $\xi := (g, q)$ is associated with a rate $r_\xi(\tau)$ that may depend explicitly on the point in time. Rate constants are described below.

In each step, two random numbers r_1 and r_2 are drawn independently from the uniform distribution on $[0, 1]$. The first random number r_1 is used to select ξ with probability $r_\xi(\tau)/R(\tau)$, where $R(\tau)$ is the sum of the rates of all reactions available at time τ . We refer to Gillespie [1977] for a convenient way to implement the rate-proportional choice of the “reaction channel”. Depending on the selected event type, the following actions are performed:

($q = L$) Gene loss is modeled by removing g from the list of active genes.

($q = D$) Gene duplications are modeled by placing a copy g' of g into the same branch of S at time τ .

($q = H$) For HGT the copy of g' is placed into a different branch of S . The “landing site” for the HGT copy is chosen uniformly from the branches of S available at time τ with the exception of the branch harboring the parental gene g .

The rules determining the rate parameters for gene copies g' and the optional adjustment of rates for the genes g are discussed below. The second random variable r_2 is used to update the clock according to $\tau \leftarrow \tau - \Delta\tau$ with $\Delta\tau = \ln(1/r_2)/R$. The simulation terminates as soon as $\tau - \Delta\tau \leq 0$.

A complication arises from the fact that the time interval $[\tau, \tau - \Delta\tau]$ may contain a speciation event at time τ_s . At a speciation, the gene content is copied into the daughter-lineages, and the rates are modified in a lineage-specific manner. As a consequence, the waiting time $\Delta\tau$ has to be re-estimated since the set of reaction channels has changed. More precisely, we need to determine the distribution of waiting times from a time point t_0 until the next event conditioned on the fact that no event occurred between t_0 and t_1 , where t_1 designates the time point of the speciation. For the complementary cumulative distribution function and $s := t_0 - t_1$ we have

$$\begin{aligned} \mathbb{P}(T \geq s + t | T \geq s) &= \mathbb{P}(T \geq s + t \wedge T \geq s) / \mathbb{P}(T \geq s) \\ &= \mathbb{P}(T \geq s + t) / \mathbb{P}(T \geq s) \end{aligned}$$

Since the waiting time distributions are exponential with rate r_1 before t_1 and rate r_2 following the speciation event, we obtain

$$\mathbb{P}(T \geq s + t | T \geq s) = e^{-(r_1 s + r_2 t)} / e^{-r_1 s} = e^{-r_2 t}$$

Hence, if the simulated waiting time reaches beyond the speciation event, the clock is advanced to the speciation event and a new waiting time is drawn with the rates after the speciation event. In practice, a new random number to obtain the time step $\Delta\tau'$ with the updated rates after the speciation event. In the new interval $[\tau_s, \tau_s - \Delta\tau']$ we again have to check for speciation events. Since the speciation events are known *a priori* from the dated species tree S , they are held in a priority queue in temporal

order. The final result is a dated gene tree T , i.e., each event is unambiguously associated with a time stamp. The simulation also completely determines the reconciliation map μ .

We simulated 2000 pairs of species and gene trees, where $|L(S)|$ was drawn uniformly from the interval $[3, 50]$. The duplication and loss rates were (independently) drawn from $[0.5, 1.0]$.

5.3 Modeling rate imbalances

In order to produce realistic (sequence) data, an evolution rate ω_e has to be associated with each edge e of T . To this end we use a hierarchical model that first determines a baseline gene substitution rate ω_e^0 for each edge e of the species tree S in order to simulate effects such as variations of population size and generation time. This introduces a correlation between the rates of all genes in the same lineage of S . These base rates are then modified by gene-specific contributions that capture effects such as differences in selection pressures that depend on gene function and rate differences in the wake of duplications such as neofunctionalization and subfunctionalization [Force et al. \[1999\]](#). In detail, we use the following parametrization:

- mean substitution rate of the conserved members of a gene family (default 1.0).
- variance σ_0^2 for the baseline substitution rate in S (default 0.2).
- a gamma distribution for the substitution rates > 1 of divergent genes. The parameters are estimated from data for the whole genome duplication in saccharomycete yeasts [Byrne and Wolfe \[2007\]](#). Alternatively, a uniform distribution on $(1; r_{max}]$ can be selected.
- weights for the relative frequency of the possible fates of duplicates (functional conservation, subfunctionalization, neofunctionalization; default equal weights 1/3).

We determine the baseline substitution rates ω_{uv}^0 for the edge $uv \in E(S)$ as follows: We simply assign the mean substitution rate to the planted edge $0_S \rho_S$ (i.e. 1.0 by default). We traverse S in pre-order and draw for each edge $uv \in E(S) \setminus \{0_S \rho_S\}$ the logarithm $\ln \omega_{uv}^0$ of the rate of evolution from a normal distribution with variance $\sigma^2 = \sigma_0^2(\tau(u) - \tau(v))$. To avoid bias towards higher or lower rates, we normalize the mean of the normal distribution such that $E(\omega_{uv}^0) = \omega_{\text{par}(u)u}^0$.

For the gene specific rates we first sort all vertices $u \in V(T)$ by $\tau(u)$ in descending temporal order. We keep track of the current number of genes in each branch of the species tree. During the simulation, the edges of T will be marked as either **conserved** or **divergent** depending on the fate of the branch after a duplication event. For each edge $e = uv \in E(T)$ in the gene tree, we initialize an empty list \mathfrak{L}_e of ordered pairs of the form (τ, ω) to record the gene-specific evolution rates ω and the corresponding time points τ at which they become valid during the existence of e . This allows us to reset the **divergent** status of a gene in case it is the last survivor in a given species. At present, we do not consider other events that change the rate of evolution of a gene within the edge e . The framework, however, can easily accommodate such rules in future refinements of the model. We denote by $\mathfrak{L}_{e,i}$ the i^{th} ordered pair (τ_i, ω_i) in \mathfrak{L}_e and define $\tau(\mathfrak{L}_{e,i}) := \tau_i$ and $\omega(\mathfrak{L}_{e,i}) := \omega_i$.

Recall that $0_T \rho_T$ is the first (planted) edge in T . To initialize the simulation, we mark $0_T \rho_T$ as **conserved** and append $(\tau(0_T), 1.0)$ to $\mathfrak{L}_{0_T \rho_T}$. Then for each vertex u in the sorted list we proceed as follows:

(1) *u is a speciation event*

Mark all edges uv with $v \in \text{child}(u)$ the same as $\text{par}(u)u$. To \mathfrak{L}_{uv} we append the pair $(\tau(u), \omega)$ with $\omega = 1.0$ (uv is **conserved**) or ω Gamma-distributed (uv is **divergent**), respectively.

(2) *u is a duplication event*

If the edge $\text{par}(u)u$ is marked as **divergent**, then all edges uv with $v \in \text{child}(u)$ are also marked as **divergent** and corresponding pairs $(\tau(u), \omega)$ are appended to \mathfrak{L}_{uv} , where the values of ω are drawn i.i.d. from the Gamma distribution.

If $\text{par}(u)u$ is marked as **conserved**, we choose between (a) conservation, (b) subfunctionalization and (c) neofunctionalization with the specified weights. For (a) mark both incident edges below u as **conserved**, for (b) as **divergent** and for (c) one edge is **conserved** and the other is **divergent**. To \mathfrak{L}_{uv} we append the pair $(\tau(u), \omega)$ with $\omega = 1.0$ (uv is **conserved**) or ω Gamma-distributed (uv is **divergent**), respectively.

(3) *u is a loss event*

If a single copy is left in the respective species after the loss: Let e^* be the corresponding edge of the remaining copy at $\tau(u)$. Mark e^* as **conserved** and append the pair $(\tau(u), 1.0)$ to \mathfrak{L}_{e^*} .

(4) u is an HGT event

Let v_1 be the copy that remains in the species and v_2 the transferred copy. Mark uv_1 the same as $\text{par}(u)u$ and append $(\tau(u), \omega)$ to \mathfrak{L}_{uv_1} where ω is the last rate that was appended to $\mathfrak{L}_{\text{par}(u)u}$. Mark uv_2 as **divergent** and append $(\tau(u), \omega)$ to \mathfrak{L}_{uv_2} with ω Gamma-distributed.

For each edge $e = uv$ in T we finalize \mathfrak{L}_e by appending $(\tau(v), \omega)$ where ω is the last rate that was appended to \mathfrak{L}_e so far. We then define the edge length $\ell(e)$ for each edge e in T as

$$\ell(e) = \omega_f^0 \sum_{i=1}^{|\mathfrak{L}_e|-1} \omega(\mathfrak{L}_{e,i}) \cdot (\tau(\mathfrak{L}_{e,i}) - \tau(\mathfrak{L}_{e,i+1})) \quad (5)$$

where f is the edge in the species tree S into which e is embedded. The resulting distance function $\ell : E(T) \rightarrow \mathbb{R}^+$ defines an additive metric on the set of vertices $V(T)$. We denote by \mathbf{D} a distance matrix on the set of non-loss leaves in T , i.e., the extant genes at time $\tau = 0$.

5.4 Simulation of measurement noise

In order to simulate measurement noise we consider three strategies:

(1) It is easy to see that a convex combination

$(1 - \epsilon)\mathbf{D} + \epsilon\mathbf{D}'$, $0 \leq \epsilon \leq 1$ of two metrics \mathbf{D} and \mathbf{D}' is again a metric (i.e., in particular, satisfies the triangle inequality). Even if both \mathbf{D} and \mathbf{D}' are additive, however, their convex combination is not additive in general. This yields a distance that is affected by a systematic bias corresponding to the noise contribution $\epsilon\mathbf{D}'$.

(2) Adding i.i.d. random noise ε_{xy} with mean 1 and standard deviation s to the additive distance, i.e., substituting $D'(x, y) = D'(y, x) = \varepsilon_{xy}D(x, y)$, in general violates the triangle inequality. We therefore use the following simple algorithm: choose $x \neq y \in L$ at random. If D' satisfies the triangle inequality, then accept the addition of ε_{xy} ; otherwise, reject the change and choose a new random perturbation. We repeat this until $\binom{|L|}{2}$ perturbations have been accepted. An alternative approach is to first introduce perturbations to all distances in \mathbf{D} and then to extract a corrected distance matrix $\hat{\mathbf{D}}$ using one of several algorithms for the ‘‘metric repair problem’’, see e.g. [Brickell et al. \[2008\]](#), [Gilbert and Jain \[2017\]](#). A cursory test showed that the trees reconstructed from distance matrices processed with these methods tend to be more different from the reference than with our approach of enforcing the triangle inequality immediately. We therefore did not pursue them further in this contribution.

(3) The distance $\ell(e)$ can also be interpreted as the number of evolutionary events per site. These are then simulated directly using the generation tool `pyvolve` [Spielman and Wilke \[2015\]](#). We generated nucleic acid sequences of length 200 with equal transition and transversion rates and of length 2000 with a transition:transversion ratio of 2 : 1. To assess saturation effects, we scaled the rates μ_e by 1/4, 1/2, 1, and 2, respectively. To better connect this work with protein-based orthology assessment pipelines, we simulated aminoacid sequences of length 500 using the WAG model [Whelan and Goldman \[2001\]](#). Distances were then estimated in `biopython` with the BLOSUM62 matrix [Eddy \[2004\]](#).

5.5 Estimation of quartets from sequence data

In order to estimate quartets directly from aligned sequence data, we use the approach of statistical geometry [Eigen et al. \[1988\]](#), [Nieselt-Struwe \[1997\]](#). We start from a multiple alignment of the sequences x, y', y'' , and z , which we assume to appear in this order. The alignment itself is produced by the sequence simulator and thus does not need to be recomputed. Each alignment column belongs to one of the 15 categories determined by which of the four sequence x, y', y'' , and z feature the same

character:

	\mathcal{C}_1	\mathcal{C}_2	\mathcal{C}_3	\mathcal{C}_4	\mathcal{C}_5	\mathcal{C}_6	\mathcal{C}_7	\mathcal{C}_8	\mathcal{C}_9	\mathcal{C}_{10}	\mathcal{C}_{11}	\mathcal{C}_{12}	\mathcal{C}_{13}	\mathcal{C}_{14}	\mathcal{C}_{15}
x	a	a	a	a	b	a	a	a	a	a	a	b	b	b	a
y'	a	a	a	b	a	a	b	b	a	b	b	a	a	c	b
y''	a	a	b	a	a	b	a	b	b	a	c	a	c	a	c
z	a	b	a	a	a	b	b	a	c	c	a	c	a	a	d

The categories \mathcal{C}_1 through \mathcal{C}_5 and \mathcal{C}_{15} do not convey phylogenetic information. Of the remaining ones, \mathcal{C}_6 , \mathcal{C}_9 , and \mathcal{C}_{14} support $(xy'|y''z)$, \mathcal{C}_7 , \mathcal{C}_{10} , and \mathcal{C}_{13} support $(xy''|y'z)$, and \mathcal{C}_8 , \mathcal{C}_{11} , and \mathcal{C}_{12} support $(xz|y'y'')$ Nieselt-Struwe and von Haeseler [2001]. Denoting by d_{aaaa} , etc., the number of alignment columns belonging to a given category, the support scores for *geometry mapping* Nieselt-Struwe and von Haeseler [2001] are

$$\begin{aligned}
S(xy'|y''z) &= d_{aabb} + \frac{1}{2}(d_{aabc} + d_{bcaa}) \\
S(xy''|y'z) &= d_{abab} + \frac{1}{2}(d_{abac} + d_{baca}) \\
S(xz|y'y'') &= d_{abba} + \frac{1}{2}(d_{abca} + d_{baac})
\end{aligned} \tag{6}$$

Using $S := S(xy'|y''z) + S(xy''|y'z) + S(xz|y'y'')$, normalized scores are defined as $s(xy'|y''z) := S(xy'|y''z)/S$. This unweighted version can be extended to a weighted version when a non-trivial distance measure D on the underlying alphabet is given. As derived in Nieselt-Struwe and von Haeseler [2001], a support value for the three possible quartets can be computed separately for each alignment column i as the isolation index for the distances on the four characters:

$$\begin{aligned}
2\beta_i(xy'|y''z) &= D_i^* - (D(x_i, y'_i) + D(y''_i, z_i)) \\
2\beta_i(xy''|y'z) &= D_i^* - (D(x_i, y''_i) + D(y'_i, z_i)) \\
2\beta_i(xz|y'y'') &= D_i^* - (D(x_i, z_i) + D(y'_i, y''_i))
\end{aligned} \tag{7}$$

Here D_i^* is the largest of the three distance sums appearing in the equation above. Summing up the $\beta_i(\cdot)$ values over all alignment columns i yields aggregated support scores $\beta(\cdot)$. These are conveniently normalized to relative values as in the unweighted case. The relative support scores for the weighted model reduce to the unweighted ones if $D(a, b) = 1 - \delta - a, b$ is the trivial metric Nieselt-Struwe and von Haeseler [2001]. If no quartet can be inferred unambiguously, then we default to the assumption $\text{lca}(x, y') = \text{lca}(x, y'')$.

Competing interests

The authors declare that they have no competing interests.

Author's contributions

PFS designed the study, PFS, MG, MH, DS, and MHR derived the mathematical results, DS implemented the generation of weighted scenarios, AL, MGL, and DIV performed the sequence-based simulations. All authors contributed to the interpretation of results and the writing of the manuscript.

Funding

This work was supported in part by the German Federal Ministry of Education and Research (BMBF, project no. 031A538A, de.NBI-RBC) and the Mexican Consejo Nacional de Ciencia y Tecnología (CONACyT, 278966 FONCICYT 2). Publication costs are covered by the DFG through the Open Access Fund at Universität Leipzig.

Additional Files

Additional file 1 — Alternative construction of Γ

In order to further investigate the inaccuracies introduced by unresolved quartets, we considered alternative constructions of the auxiliary graph Γ . In addition to the default method, we omitted all edges defined for quartets classified as unresolved (\times), and we ignored the contribution of outgroups that lead to unresolved and used a majority vote only for the remaining choices of the outgroup. All non-trivial sinks in Γ were then interpreted as best matches, i.e., isolated vertices in Γ were ignored. Both variants perform worse than the default method. Data are compared for short nucleic acid sequences with rates of sequence divergence scaled by 1 (OR), 1/2 (D2), 1/4 (D4), and 2 (M2).

[AdditionalFile_1.pdf](#)

References

- A.V. Aho, Y. Sagiv, T.G. Szymanski, and J.D. Ullman. Inferring a tree from lowest common ancestors with an application to the optimization of relational expressions. *SIAM J Comput*, 10:405–421, 1981. doi: 10.1137/0210030.
- A. M. Altenhoff and C. Dessimoz. Phylogenetic and functional assessment of orthologs inference projects and methods. *PLoS Comput Biol*, 5:e1000262, 2009. doi: 10.1371/journal.pcbi.1000262.
- Adrian M. Altenhoff, Brigitte Boeckmann, Salvador Capella-Gutierrez, Daniel A. Dalquen, Todd DeLuca, Kristoffer Forslund, Huerta-Cepas Jaime, Benjamin Linard, Cécile Pereira, Leszek P. Prysycz, Fabian Schreiber, Alan Sousa da Silva, Damian Szklarczyk, Clément-Marie Train, Peer Bork, Odile Lecompte, Christian von Mering, Ioannis Xenarios, Kimmen Sjölander, Lars Juhl Jensen, Maria J. Martin, Matthieu Muffato, Toni Gabaldón, Suzanna E. Lewis, Paul D. Thomas, Erik Sonnhammer, and Christophe Dessimoz. Standardized benchmarking in the quest for orthologs. *Nature Methods*, 13:425–430, 2016. doi: 10.1038/nmeth.3830.
- Kevin Atteson. The performance of Neighbor-Joining methods of phylogenetic reconstruction. *Algorithmica*, 25:251–278, 1999. doi: 10.1007/PL00008277.
- Sebastian Böcker and Andreas W. M. Dress. Recovering symbolically dated, rooted trees from symbolic ultrametrics. *Adv. Math.*, 138:105–125, 1998. doi: 10.1006/aima.1998.1743.
- P Bork, T Dandekar, Y Diaz-Lazcoz, F Eisenhaber, M Huynen, and Y Yuan. Predicting function: from genes to genomes and back. *J Mol Biol*, 283:707–725, 1998. doi: 10.1006/jmbi.1998.2144.
- Justin Brickell, Inderjit S. Dhillon, S. Sra, and Joel A. Tropp. The metric nearness problem. *SIAM J. Matrix Analysis Appl.*, 30:375–396, 2008. doi: 10.1137/060653391.
- Peter Buneman. Note on the metric properties of trees. *J. Comb. Th. B*, 17:48–50, 1974. doi: 10.1016/0095-8956(74)90047-1.
- Kevin P. Byrne and Kenneth H. Wolfe. Consistent patterns of rate asymmetry and gene loss indicate widespread neofunctionalization of yeast genes after whole-genome duplication. *Genetics*, 175:1341–1350, 2007. doi: 10.1534/genetics.106.066951.
- S Cherlin, T M W Nye, R J Boys, S E Heaps, T A Williams, and T M Embley. The effect of non-reversibility on inferring rooted phylogenies. *Mol Biol Evol*, 35:984–1002, 2018. doi: 10.1093/molbev/msx294.
- J-P Doyon, V Ranwez, V Daubin, and V Berry. Models, algorithms and programs for phylogeny reconciliation. *Brief Bioinform.*, 12:392–400, 2011. doi: 10.1093/bib/bbr045.

- A J Drummond, S Y W Ho, M J Phillips, and A Rambaut. Relaxed phylogenetics and dating with confidence. *PLoS Biol*, 4:699–710, 2006. doi: 10.1371/journal.pbio.0040088.
- Sean R Eddy. Where did the BLOSUM62 alignment score matrix come from? *Nature Biotech.*, 22: 1035–1036, 2004. doi: 10.1038/nbt0804-1035.
- Manfred Eigen, Ruth Winkler-Oswatitsch, and Andreas W. M. Dress. Statistical geometry in sequence space: a method of quantitative comparative sequence analysis. *Proc Natl Acad Sci USA*, 85:5913–5917, 1988. doi: 10.1073/pnas.85.16.5913.
- W M Fitch. Distinguishing homologous from analogous proteins. *Syst Zool*, 19:99–113, 1970. doi: 10.2307/2412448.
- W. M. Fitch. A non-sequential method for constructing trees and hierarchical classifications. *J. Mol. Evol.*, 18:30–37, 1981. doi: 10.1007/BF01733209.
- Walter M. Fitch. Homology: a personal view on some of the problems. *Trends Genet.*, 16:227–231, 2000. doi: 10.1016/S0168-9525(00)02005-9.
- Allan Force, Michael Lynch, F. Bryan Pickett, Angel Amores, Yi-lin Yan, and John Postlethwait. Preservation of duplicate genes by complementary, degenerative mutations. *Genetics*, 151:1531–1545, 1999.
- Toni Gabaldón and Eugene V. Koonin. Functional and evolutionary implications of gene orthology. *Nat Rev Genet.*, 14:360–366, 2013. doi: 10.1038/nrg3456.
- M. Geiß, P. F. Stadler, and M Hellmuth. Reciprocal best match graphs. Technical Report arXiv:1903.07920v4, arXiv, 2019a.
- Manuela Geiß, Edgar Chávez, Marcos González, Alitzel López, Bärbel M R Stadler, Dulce Valdivia, Marc Hellmuth, Maribel Hernández Rosales, and Peter F Stadler. Best match graphs. *J. Math. Biol.*, 78:2015–2057, 2019b. doi: 10.1007/s00285-019-01332-9.
- Manuela Geiß, Marcos E. González Laffitte, Alitzel López Sánchez, Dulce I. Valdivia, Marc Hellmuth, Maribel Hernández Rosales, and Peter F. Stadler. Best match graphs and reconciliation of gene trees with species trees. Technical Report arXiv:1904.12021v2, arXiv, 2019c.
- Anna C. Gilbert and Lalit Jain. If it ain’t broke, don’t fix it: Sparse metric repair. In *55th Annual Allerton Conference on Communication, Control, and Computing*, pages 612–619, 2017. doi: 10.1109/ALLERTON.2017.8262793.
- Daniel T. Gillespie. Exact stochastic simulation of coupled chemical reactions. *J. Phys. Chem.*, 81: 2340–2361, 1977. doi: 10.1021/j100540a008.
- Pawel Górecki and Jerzy Tiuryn. DLS-trees: A model of evolutionary scenarios. *Theor. Comp. Sci.*, 359:378–399, 2006. doi: 10.1016/j.tcs.2006.05.019.
- M Hasegawa, H Kishino, and T Yano. Dating of human-ape splitting by a molecular clock of mitochondrial DNA. *J. Mol. Evol.*, 22:160–174, 1985. doi: 10.1007/BF02101694.
- M. Hellmuth. Biologically feasible gene trees, reconciliation maps and informative triples. *Alg. Mol. Biol.*, 12:23, 2017a. doi: 10.1186/s13015-017-0114-z.
- Marc Hellmuth. Biologically feasible gene trees, reconciliation maps and informative triples. *Alg. Mol. Biol.*, 12:23, 2017b. doi: 10.1186/s13015-017-0114-z.
- Maribel Hernandez-Rosales, Marc Hellmuth, Nicolas Wieseke, Katharina T. Huber, Vincent Moulton, and Peter F. Stadler. From event-labeled gene trees to species trees. *BMC Bioinformatics*, 13 (Suppl. 19):S6, 2012. doi: 10.1186/1471-2105-13-S19-S6.

- Pablo N Hess and Claudia A de Moraes Russo. An empirical test of the midpoint rooting method. *Biol. J. Linnean Soc.*, 92:669–674, 2007. doi: 10.1111/j.1095-8312.2007.00864.x.
- C T Hittinger and S B Carroll. Gene duplication and the adaptive evolution of a classic genetic switch. *Nature*, 449:677–681, 2007. doi: 10.1038/nature06151.
- B R Holland, D Penny, and M D Hendy. Outgroup misplacement and phylogenetic inaccuracy under a molecular clock — a simulation study. *Syst. Biol.*, 52:229–238, 2003. doi: 10.1080/10635150390192771.
- J P Huelsenbeck, B Larget, R E Miller, and F Ronquist. Potential applications and pitfalls of Bayesian inference of phylogeny. *Syst Biol.*, 51:673–688, 2002. doi: 10.1080/10635150290102366.
- T H Jukes and C R Cantor. Evolution of protein molecules. In H N Munro, editor, *Mammalian Protein Metabolism*, pages 21–132. Academic Press, New York, 1969.
- L A Katz, J R Grant, L W Parfrey, and J G Burleigh. Turning the crown upside down: gene tree parsimony roots the eukaryotic tree of life. *Syst. Biol.*, 61:653–660, 2012. doi: 10.1093/sysbio/sys026.
- Y Kawahara and T. Imanishi. A genome-wide survey of changes in protein evolutionary rates across four closely related species of *Saccharomyces sensu stricto* group. *BMC Evol Biol.*, 7:9, 2007. doi: 10.1186/1471-2148-7-9.
- Stephanie Keller-Schmidt and Konstantin Klemm. A model of macroevolution as a branching process based on innovations. *Adv. Complex Syst.*, 15:1250043, 2012. doi: 10.1142/S0219525912500439.
- M Kimura. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.*, 16:111–120, 1980. doi: 10.1007/BF01731581.
- T Kinene, J Wainaina, S Maina, and LM Boykin. Rooting trees, methods for. In Richard M. Kliman, editor, *Encyclopedia of Evolutionary Biology*, volume 3, page 489. Elsevier, Amsterdam, NL, 2016. doi: 10.1016/B978-0-12-800049-6.00215-8.
- Eugene Koonin. Orthologs, paralogs, and evolutionary genomics. *Ann. Rev. Genetics*, 39:309–338, 2005. doi: 10.1146/annurev.genet.39.073003.114725.
- Sudhir Kumar. Molecular clocks: four decades of evolution. *Nat Rev Genet*, 6:654–662, 2005. doi: 10.1038/nrg1659.PMID16136655.
- Marcus Lechner, Sven Findeiß, Lydia Steiner, Manja Marz, Peter F. Stadler, and Sonja J. Prohaska. **Proteinortho**: detection of (co-)orthologs in large-scale analysis. *BMC Bioinformatics*, 12:124, 2011. doi: 10.1186/1471-2105-12-124.
- Uyen Mai, Erfan Sayyari, and Siavash Mirarab. Minimum variance rooting of phylogenetic trees and implications for species tree reconstruction. *PLoS ONE*, 12:e0182238. doi: 10.1371/journal.pone.0182238.
- K. Nieselt-Struwe. Graphs in sequence spaces: a review of statistical geometry. *Biophys Chem.*, 66: 111–131, 1997. doi: 10.1016/S0301-4622(97)00064-1.
- Kay Nieselt-Struwe and Arndt von Haeseler. Quartet-mapping, a generalization of the likelihood-mapping procedure. *Mol Biol Evol*, 18:1204–1219, 2001. doi: 10.1093/oxfordjournals.molbev.a003907.
- R Overbeek, M Fonstein, M D’Souza, G D Pusch, and N Maltsev. The use of gene clusters to infer functional coupling. *Proc Natl Acad Sci USA*, 96:2896–2901, 1999. doi: 10.1073/pnas.96.6.2896.
- David Penny. Criteria for optimising phylogenetic trees and the problem of determining the root of a tree. *J. Mol. Evol.*, 8:95–116, 1976. doi: 10.1007/BF01739097.

- Nancy Retzlaff and Peter F. Stadler. Phylogenetics beyond biology. *Th. Biosci.*, 137:133–143, 2018. doi: 10.1007/s12064-018-0264-7.
- L. Y. Rusin, E. Lyubetskaya, K. Y. Gorbunov, and V. Lyubetsky. Reconciliation of gene and species trees. *BioMed Res Int.*, 2014:642089, 2014. doi: 10.1155/2014/642089.
- N. Saitou and M. Nei. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol*, 4:406–425, 1987. doi: 10.1093/oxfordjournals.molbev.a040454.
- S. Sattah and A. Tversky. Additive similarity trees. *Psychometrika*, 42:319–345, 1977. doi: 10.1007/BF02293654.
- Charles Semple and Mike Steel. *Phylogenetics*. Oxford University Press, Oxford UK, 2003.
- L Shavit, D Penny, M D Hendy, and B R Holland. The problem of rooting rapid radiations. *Mol Biol Evol*, 24:2400–2411, 2007. doi: 10.1093/molbev/msm178.
- J. M. S. Simões-Pereira. A note on the tree realizability of a distance matrix. *J. Combin. Theory*, 6: 303–310, 1969. doi: 10.1016/S0021-9800(69)80092-X.
- P S Soria, K L McGary, and A. Rokas. Functional divergence for every paralog. *Mol Biol Evol.*, 31: 984–992, 2014. doi: 10.1093/molbev/msu050.
- S J Spielman and C O Wilke. Pyvolve: A flexible python module for simulating sequences along phylogenies. *PLoS One*, 10:e0139047, 2015. doi: 10.1371/journal.pone.0139047.
- Mike Steel. The complexity of reconstructing trees from qualitative characters and subtreess. *J. Classification*, 9:91–116, 1992.
- D L Swofford, G J Olsen, P J Waddell, and D M Hillis. Phylogenetic inference. In D M Hillis, C Moritz, and B K Mable, editors, *Molecular systematics*, pages 407–514. Sinauer Associates, Sunderland, MA, 1996.
- K Tamura. Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G+C content biases. *Mol. Biol. Evol.*, 9:678–687, 1992. doi: 10.1093/oxfordjournals.molbev.a040752.
- R. L. Tatusov, E. V. Koonin, and D. J. Lipman. A genomic perspective on protein families. *Science*, 278:631–637, 1997. doi: 10.1126/science.278.5338.631.
- Clément-Marie Train, Natasha M Glover, Gaston H Gonnet, Adrian M Altenhoff, and Christophe Dessimoz. Orthologous matrix (OMA) algorithm 2.0: more robust to asymmetric evolutionary rates and more scalable hierarchical orthologous group inference. *Bioinformatics*, 33:i75–i82. doi: 10.1093/bioinformatics/btx229.
- Günter P. Wagner, Kazuhiko Takahashi, Vincent Lynch, Sonja J. Prohaska, Claudia Fried, Peter F. Stadler, and Chris T. Amemiya. Molecular evolution of duplicated ray finned fish *hoxa* clusters: Increased synonymous substitution rate and asymmetrical co-divergence of coding and non-coding sequences. *J. Mol. Evol.*, pages 665–676, 2005.
- D P Wall, H B Fraser, and A E Hirsh. Detecting putative orthologs. *Bioinformatics*, 19:1710–1711, 2003. doi: 10.1093/bioinformatics/btg213.
- S Whelan and N Goldman. A general empirical model of protein evolution derived from multiple protein families using a maximum-likelihood approach. *Mol Biol Evol*, 18:691–699, 2001. doi: 10.1093/oxfordjournals.molbev.a003851.

- Tom A. Williams, Sarah E. Heaps, Svetlana Cherlin, Tom M. W. Nye, Richard J. Boys, and T. Martin Embley. New substitution models for rooting phylogenetic trees. *Philos Trans R Soc Lond B Biol Sci*, 370:20140336, 2015. doi: 10.1098/rstb.2014.0336.
- Chenggang Yu, Nela Zavaljevski, Valmik Desai, and Jaques Reifman. QuartetS: a fast and accurate algorithm for large-scale orthology detection. *Nucleic Acids Res.*, 39:e88, 2011. doi: 10.1093/nar/gkr308.
- E. Zuckerkandl and L. B. Pauling. Molecular disease, evolution, and genic heterogeneity. In M. Kasha and B Pullman, editors, *Horizons in Biochemistry*, pages 189–225. Academic Press, New York, 1962.