

On the probability of Boolean functions of events in the n -dimensional Euclidean space

Jinwook Lee^{a,*}

^aDecision Sciences and MIS, LeBow College of Business, Drexel University, Philadelphia, PA 19104, United States

Abstract

It is shown in [15] and [16] that for the union of N orthants in R^n ($N \gg n$) there exists an efficient and systematic way to find the exact value, using a suitable partial order relation construction. In this paper our events are hyperrectangles (or n -orthotopes), the Cartesian product of intervals – another important sets (or events) in both theory and practice. We have discovered a new efficient algorithm for the union of such events. With other important Boolean functions we present optimization problem formulations for both hyperrectangles and orthants.

Keywords: Boolean functions of events; hyperrectangle; orthotope; binomial moment problem

1. Introduction

Let A_1, \dots, A_N be events in an arbitrary probability space. In many applications we are interested to find probabilities of Boolean functions of them, among which prominent are the following:

- at least one of them occurs;
- at least r of them occur;
- exactly r of them occur.

In reliability theory, where consecutive events play important role, further Boolean functions of events can be used. Our results can be applied for probabilities of these other Boolean functions, as well, but here we restrict ourselves to the above-mentioned three.

Given such Boolean functions of events, related probabilities and conditional expectations are useful in decision making. For example, If there are N divisions of a business organization, then a massive failure of just a few of the divisions will possibly collapse the entire organization. For this reason, a problem of “at least r ($1 \leq r \ll N$)” events comes into play for suitable decision making, using the probabilities of such events and their related conditional expectations (e.g., probability weighted sum of losses (or profits)), etc.

Our events in this paper are hyperrectangles (or n -orthotopes) in the n -space, designated by

$$A_1 = A(z^{(1)}), \dots, A_N = A(z^{(N)}), \quad (1)$$

where $z^{(1)}, \dots, z^{(N)}$ are the pairs of lower and upper vertices of hyperrectangles R^n . $z^{(i)} = [l^{(i)}, u^{(i)}]$ where $l^{(i)}, u^{(i)}$ for $i \in R^n, i = 1, \dots, N$. Each of the $z^{(i)}$'s is a pair of vertices.

For simplicity, let us consider a cube in R^3 :

$$z^{(1)} = [(1, 2, 3), (4, 5, 6)].$$

Then our event is:

$$\begin{aligned} A(z^{(1)}) &= A([(1, 2, 3), (4, 5, 6)]) \\ &= \{(z_1, z_2, z_3) \mid (1, 2, 3) \leq (z_1, z_2, z_3) \leq (4, 5, 6)\}, \end{aligned} \quad (2)$$

Note that there are eight vertices for the cube $z^{(1)}$, but we only need a pair. It is easy to see that no matter what dimension we have, say n some large positive integer, we only need a pair (i.e., lower and upper corners) out of 2^n vertices.

(2) can also be represented by the Cartesian product:

$$A(z^{(1)}) = Z^{(1)} = Z_1^{(1)} \times Z_2^{(1)} \times Z_3^{(1)}, \quad (3)$$

where $Z_1 = \{z_1 \mid 1 \leq z_1 \leq 4\}$, $Z_2 = \{z_2 \mid 2 \leq z_2 \leq 5\}$ and $Z_3 = \{z_3 \mid 3 \leq z_3 \leq 6\}$. Equations (2) and (3) are equivalent to

$$\{(z_1, z_2, z_3) \mid z_1 \in [1, 4] \wedge z_2 \in [2, 5] \wedge z_3 \in [3, 6]\}. \quad (4)$$

Let f denote the p.d.f. of a random vector $Z \in R^3$. Then we can write

$$P(A(z^{(1)})) = \int_1^4 \int_2^5 \int_3^6 f(z_1, z_2, z_3) dz_3 dz_2 dz_1. \quad (5)$$

2. Vertex comparison for empty intersections of hyperrectangles

Theorem 1. A pairwise intersection of N hyperrectangles A_1, \dots, A_N in R^n is not empty if the condition (i) or (ii) is met.

(i) For every $k = 1, \dots, n$, we have:

$$\max\{l_k^{(i)}, l_k^{(j)}\} \leq \min\{u_k^{(i)}, u_k^{(j)}\},$$

where integers i, j such that $1 \leq i < j \leq N$.

*Corresponding author, Phone: 1-215-571-4672, Fax: 1-215-895-2907
Email address: j13539@drexel.edu (Jinwook Lee)

(ii) In case of a continuous distribution, for every $k = 1, \dots, n$, we have:

$$\max\{l_k^{(i)}, l_k^{(j)}\} < \min\{u_k^{(i)}, u_k^{(j)}\},$$

where integers i, j such that $1 \leq i < j \leq N$.

Proof. Suppose not. Then there exists k such that

$$\max\{l_k^{(i)}, l_k^{(j)}\} > \min\{u_k^{(i)}, u_k^{(j)}\},$$

which can't make a nonempty intersection. \square

Remark 1 (Screening test). *Theorem 1 can be used as a screening test. We can remove empty intersections from the inclusion-exclusion formula.*

Example 1. For simplicity let us consider a continuous uniform random vector $(X_1, X_2)^T$ with support set $\{(x_1, x_2) \mid 0 \leq x_i \leq 10, i = 1, 2\}$. Suppose $A_1 = [(5, 6), (9, 9)]$, $A_2 = [(2, 4), (6, 7)]$, $A_3 = [(1, 3), (4, 8)]$, $A_4 = [(3, 2), (7, 10)]$, $A_5 = [(0, 1), (10, 5)]$. Note that they are depicted in Figure 1.

In Example 1 there are 10 pairs $A_i A_j$, $i < j$, for $i, j = 1, \dots, 5$ and their emptiness testing is as follows.

Pairs	Are conditions of Theorem 1 met?
$A_1 A_2 = [(5, 6), (6, 7)]$	yes
$A_1 A_3 = [(5, 6), (4, 8)]$	no good
$A_1 A_4 = [(5, 6), (7, 9)]$	yes
$A_1 A_5 = [(5, 6), (9, 5)]$	no good
$A_2 A_3 = [(2, 4), (4, 7)]$	yes
$A_2 A_4 = [(3, 4), (6, 7)]$	yes
$A_2 A_5 = [(2, 4), (6, 5)]$	yes
$A_3 A_4 = [(3, 3), (4, 8)]$	yes
$A_3 A_5 = [(1, 3), (4, 5)]$	yes
$A_4 A_5 = [(3, 2), (7, 5)]$	yes

(6)

We can draw Figure 2 based on (6) which shows $A_1 A_3$ and $A_1 A_5$ are empty. In Figure 2 there are eight 2-vertex clique, five 3-vertex clique and one 4-vertex clique. In Example 1 there are 10 triples $A_i A_j A_k$, $i < j < k$, for $i, j, k = 1, \dots, 5$ where there might be some empty ones. Using the information of (6) we can remove empty triples including the empty pairs of A_1, A_3 and A_1, A_5 . Therefore, we remove the following triples: $A_1 A_2 A_3, A_1 A_3 A_4, A_1 A_3 A_5$ and $A_1 A_2 A_5, A_1 A_3 A_5, A_1 A_4 A_5$ which are empty.

Triples	Are conditions of Theorem 1 met?
$A_1 A_2 A_4 = [(5, 6), (6, 7)]$	yes
$A_2 A_3 A_4 = [(3, 4), (4, 7)]$	yes
$A_2 A_3 A_5 = [(2, 4), (4, 5)]$	yes
$A_2 A_4 A_5 = [(3, 4), (6, 5)]$	yes
$A_3 A_4 A_5 = [(3, 3), (4, 5)]$	yes

(7)

Using the information of (7) we find out that there is only one 4 tuple $A_2 A_3 A_4 A_5$ and no 5 tuple.

4-tuple	Are conditions of Theorem 1 met?
$A_2 A_3 A_4 A_5 = [(3, 4), (4, 5)]$	yes

(8)

With (6), (7), (8), we can write up an inclusion exclusion formula: $S_1 - S'_2 + S'_3 - S'_4$, where we have only 19 terms compared to 31 terms in the original inclusion-exclusion formula.




Figure 1: Description of Example 1.




Figure 2: A graph with nonempty intersections. The clique number of the graph is 4.

Example 2. For simplicity let us consider a continuous uniform random vector $(X_1, X_2, X_3)^T$ with support set $\{(x_1, x_2, x_3) \mid 0 \leq x_i \leq 5, i = 1, 2, 3\}$. Suppose $A_1 = [(0, 0, 0), (2, 2, 2)]$, $A_2 = [(3, 1, 3), (5, 3, 5)]$, $A_3 = [(1, 3, 3), (3, 5, 5)]$, $A_4 = [(4, 4, 4), (5, 5, 5)]$, $A_5 = [(2, 2, 2), (3, 3, 4)]$, $A_6 = [(1, 4, 1), (2, 5, 2)]$, $A_7 = [(4, 1, 4), (5, 2, 5)]$.

In Example 1 there are 21 pairs $A_i A_j$, $i < j$, for $i, j = 1, \dots, 7$

and their emptiness testing is as follows.

Pairs	Are conditions of Theorem 1 met?
$A_1A_2 = [(3, 1, 3), (2, 2, 2)]$	no good
$A_1A_3 = [(1, 3, 3), (2, 2, 2)]$	no good
$A_1A_4 = [(4, 4, 4), (2, 2, 2)]$	no good
$A_1A_5 = [(2, 2, 2), (2, 2, 2)]$	no good
$A_1A_6 = [(1, 4, 1), (2, 2, 2)]$	no good
$A_1A_7 = [(4, 1, 4), (2, 2, 2)]$	no good
$A_2A_3 = [(3, 3, 3), (3, 3, 5)]$	no good
$A_2A_4 = [(4, 4, 4), (5, 3, 5)]$	no good
$A_2A_5 = [(3, 2, 3), (3, 3, 4)]$	no good
$A_2A_6 = [(3, 4, 3), (2, 3, 2)]$	no good
$A_2A_7 = [(4, 1, 4), (5, 2, 5)]$	yes
$A_3A_4 = [(4, 4, 4), (3, 5, 5)]$	no good
$A_3A_5 = [(2, 3, 3), (3, 3, 4)]$	no good
$A_3A_6 = [(1, 4, 3), (2, 5, 2)]$	no good
$A_3A_7 = [(4, 3, 4), (3, 2, 5)]$	no good
$A_4A_5 = [(4, 4, 4), (3, 3, 4)]$	no good
$A_4A_6 = [(4, 4, 4), (2, 5, 2)]$	no good
$A_4A_7 = [(4, 4, 4), (5, 2, 5)]$	no good
$A_5A_6 = [(2, 4, 2), (2, 3, 2)]$	no good
$A_5A_7 = [(4, 2, 4), (3, 2, 4)]$	no good
$A_6A_7 = [(4, 4, 4), (2, 2, 2)]$	no good

(9)

It is shown in (9) that there is only one nonempty pair A_2A_7 which can be depicted as in Figure 3. And the probability of union can be calculated only by $S_1 - S_2 = \sum_{i=1}^7 P(A_i) - P(A_2A_7)$ which has only 8 terms instead of 127 terms.




Figure 3: A graph with nonempty intersections. The clique number of the graph is 2.

3. Probability bounding using linear program

[18], [20] and [21] observed that the sharp probability bounds, using S_1, \dots, S_m , are optimum values of LP's that he called binomial moment problems. In doing so, he opened a new research area: the discrete moment problems. The term binomial moment comes from the fact that if ξ is the number of events in A_1, \dots, A_N , which occur, then

$$S_k = E \left[\binom{\xi}{k} \right], k = 1, \dots, N. \quad (10)$$

By convention we write $S_0 = 1$ with which (10) holds also for $k = 0$. Equation (10) is a classical theorem and it is not known who proved it first.

Starting from (10) and introducing the notations

$$p_i = P(\xi = i), i = 0, 1, \dots, N, \quad (11)$$

The binomial problems for:

- at least one of them occurs;
- at least r of them occur;
- exactly r of them occur.

can be formulated as the following.

$$\begin{aligned} \min(\max) \quad & \sum_{i=1}^N p_i \\ \text{subject to} \quad & \sum_{i=0}^N \binom{i}{k} p_i = S_k, k = 0, 1, \dots, m \\ & p_i \geq 0, i = 0, 1, \dots, N, \end{aligned} \quad (12)$$

$$\begin{aligned} \min(\max) \quad & \sum_{i=r}^N p_i \\ \text{subject to} \quad & \sum_{i=0}^N \binom{i}{k} p_i = S_k, k = 0, 1, \dots, m \\ & p_i \geq 0, i = 0, 1, \dots, N, \end{aligned} \quad (13)$$

$$\begin{aligned} \min(\max) \quad & p_r \\ \text{subject to} \quad & \sum_{i=0}^N \binom{i}{k} p_i = S_k, k = 0, 1, \dots, m \\ & p_i \geq 0, i = 0, 1, \dots, N. \end{aligned} \quad (14)$$

Instead of problem (12) [18] used a simpler problem, removing p_0 and the first constraint:

$$\begin{aligned} \min(\max) \quad & \sum_{i=1}^N p_i \\ \text{subject to} \quad & \sum_{i=1}^N \binom{i}{k} p_i = S_k, k = 1, \dots, m \\ & p_i \geq 0, i = 1, \dots, N, \end{aligned} \quad (15)$$

[18] characterized the dual feasible bases and showed that there can be used closed form formulas for $m = 2, 3$. For the case of a general m he gave elegant dual algorithms to obtain bounds for the probabilities in question. The results have been used by [4] to obtain closed form formulas for the case $m = 4$ and also for the case of a general m .

Another direction in probability bounding problems is to use the individual probabilities $p_{i_1 \dots i_k} = P(A_{i_1} \dots A_{i_k}), k \leq m$, rather than the aggregated values S_1, \dots, S_m . [2] formulated probability bounding problems using individual probabilities that we call today LP's and further developed it in (1868). It was [9]

who formulated the general LP that we call today Boolean probability bounding problem. In order to present it in a simple form we introduce the notations:

$$\begin{aligned} p_I &= P\left(\bigcap_{j \in I} A_j\right), I, J \subset \{1, \dots, N\}, \\ a_{IJ} &= \begin{cases} 1, & \text{if } I \subset J \\ 0, & \text{if } I \not\subset J, I, J \subset \{1, \dots, N\} \end{cases}, \\ x_J &= P\left(\left(\bigcap_{j \in J} A_j\right) \cap \left(\bigcap_{j \notin J} \bar{A}_j\right)\right), \end{aligned}$$

The probability p_I means that all events A_j , $j \in K$ occur and the probability x_J means that all events A_j , $j \in I$ occur but the other do not occur. The Boolean lower and upper bounding problems for the union are the following:

$$\begin{aligned} &\min(\max) \sum_{\emptyset \neq J \subset \{1, \dots, N\}} x_J \\ &\text{subject to} \\ &\sum_{J \subset \{1, \dots, N\}} a_{IJ} x_J = p_I, I \subset \{1, \dots, N\}, |I| \leq m \\ &x_J \geq 0, J \subset \{1, \dots, N\}. \end{aligned}$$

A counterpart of it also exist, where we use $I, J \neq \emptyset$. The first closed form bounding formula was given by [10] and [23]. In this connection we also mention [22], [8], [12], [6], [5]. [22] has shown that the Hunter-Worsley bounds in the objective function value corresponding to a dual feasible basis in the modified Boolean minimization problem. [6] have shown the same about their upper bound.

4. The binomial moment problem formulations for the probabilities that at least k and exactly k events occur

In case of orthants, the bounds of the probability that “at least k events occur” can be found by the following

$$\begin{aligned} &\min(\max) \sum_{i=k}^N p_i \\ &\sum_{i=1}^N p_i = Q \\ &\sum_{i=1}^N i p_i = S_1 \\ &\sum_{i=1}^N \binom{i}{2} p_i = S_2 \\ &\sum_{i=1}^N \binom{i}{3} p_i = S_3 \\ &p_i \geq 0, i = 1, \dots, N, \end{aligned} \quad (16)$$

where Q denotes the probability of union that can be calculated by by Theorem 9 of [15]. The worst case time complexity of calculating Q is $O(n^2 N^2 + n \cdot N \log N)$ for R^n , $n \geq 3$ ($O(n^2 N^2)$ for sorted sets). Note that in practice we typically have $N \gg n$. If we assume that $N \geq n^2$, then it can be written $O(N^3)$ for sorted sets. In this case, the above formulation would be

more efficient and give better bounds than the original binomial moment problem formulation. In the bivariate case Q can be calculated in a constant time for sorted sets since we have $Q = S_1 - S'_2$, where S'_2 denotes the sum of incomparable pairwise intersections (see [16] for more details).

Similarly, the bounds of the probability that “exactly k events occur” can be found by the following

$$\begin{aligned} &\min(\max) p_k \\ &\text{subject to} \sum_{i=1}^N p_i = Q \\ &\sum_{i=1}^N i p_i = S_1 \\ &\sum_{i=1}^N \binom{i}{2} p_i = S_2 \\ &\sum_{i=1}^N \binom{i}{3} p_i = S_3 \\ &p_i \geq 0, i = 1, \dots, N. \end{aligned} \quad (17)$$

In case of hyperrectangles, we can just use (13), (14) and (15) with a lot less terms.

References

- [1] Bonferroni, C. E., 1937. Teoria statistics delle classi e calcolo delle probabilita. Volume in onore di Riccardo Dalla Volta, Universita di Firenze, 1–62.
- [2] Boole, G., 1854. Laws of Thought. American reprint of 854 edition. Dover, New York.
- [3] Boole, G., 1868. Of Propositions Numerically Definite. In Transactions of the CambridgePhilosophical Society, Part II, XI, reprinted as StudyIV in the next reference.
- [4] Boros, E., Prékopa, A., 1989. Closed form two-sided bounds for probabilities that exactly r and at least r out of n events occur. Math. Oper. Res. 14, 317–342.
- [5] Boros, E., Scozzari, A., Tardella, F., Veneziani, P., 2014. Polynomially computable bounds for the probability of the union of events. Math. Oper. Res. 39 (4), 1311–1329.
- [6] Bukszár, J., Prékopa, A., 2001. Probability bounds with cherry trees. Math. Oper. Res. 26 (1), 174–192.
- [7] Dawson, D., Sankoff, A., 1967. An inequality for probabilities. Proceedings of the American Mathematical Society 18, 504–507.
- [8] de Caen, D., 1997. A lower bound on the probability of union. Discrete Math. 169, 217–220.
- [9] Hailperin, T., 1965. Best possible inequalities for the probability of a logical function of events. The American Mathematical Monthly 72, 343–359.
- [10] Hunter, D., 1976. Bounds for the probability of a union. J. of Appl. Prob. 13, 597–603.
- [11] Jordan, C., 1927. The foundations of the theory of probability. Math. Phys. Lapok (in Hungarian) 34, 109–136.
- [12] Kuai, H., Alajaji, F., Takahara, G., 2000. A lower bound on the probability of a finite union of events. Discrete Math. 215, 147–158.
- [13] Kwerel, S. M., 1975a. Most stringent bounds on aggregated probabilities of partially specified dependent probability systems. Journal of the American Statistical Association 70, 472–479.
- [14] Kwerel, S. M., 1975b. Bounds on the probability of a union and intersection of m events. Advances of Applied Probability 7, 431–448.
- [15] Lee, J., 2017. Computing the probability of union in the n -dimensional Euclidean space for application of the multivariate quantile: p -level efficient points. Oper. Res. Lett. 45 (3), 242–247.
- [16] Lee, J., Prékopa, A., 2017. On the probability of union in the n -space. Oper. Res. Lett 45 (1), 19–24.

- [17] Poincaré, H., 1896. Calcul des probabilités. Gauthier-Villars, Paris.
- [18] Prékopa, A., 1988. Boole-Bonferroni inequalities and linear programming. *Oper. Res.* 36 (1), 145–162.
- [19] Prékopa, A., 1990. Dual method for a one-stage stochastic programming problem with random rhs obeying a discrete probability distribution. *Oper. Res.* 34, 441–461.
- [20] Prékopa, A., 1990a. Sharp bounds on probabilities using linear programming. *Oper. Res.* 38 (2), 227–239.
- [21] Prékopa, A., 1990b. The discrete moment problem and linear programming. *Discrete Appl. Math.* 27, 235–254.
- [22] Prékopa, A., 1992. Inequalities on expectations based on the knowledge of multivariate moments. In: *Stochastic Inequalities* (M. Shaked, Y.L. Tong, eds.), Institute of Mathematical Statistics, Lecture Notes - Monograph Series 22, 309–331.
- [23] Worsley, K., 1982. An improved Bonferroni inequality and applications. *Biometrika* 69, 297–302.