
Optimal quadrature formulas for non-periodic functions in Sobolev space and its application to CT image reconstruction

Abdullo R. Hayotov*, Soomin Jeon,
Chang-Ock Lee, Kholmat M. Shadimetov

Received: date / Accepted: date

Abstract In the present paper, optimal quadrature formulas in the sense of Sard are constructed for numerical integration of the integral $\int_a^b e^{2\pi i\omega x} \varphi(x) dx$ with $\omega \in \mathbb{R}$ in the Sobolev space $L_2^{(m)}[a, b]$ of complex-valued functions which are square integrable with m -th order derivative. Here, using the discrete analogue of the differential operator $\frac{d^{2m}}{dx^{2m}}$, the explicit formulas for optimal coefficients are obtained. The order of convergence of the obtained optimal quadrature formula is $O(h^m)$. As an application, we implement the filtered back-projection (FBP) algorithm, which is a well-known image reconstruction algorithm for computed tomography (CT). By approximating Fourier transforms and its inversion using the proposed optimal quadrature formula of the second and third orders, we observe that the accuracy of the reconstruction algorithm is improved. In numerical experiments, we compare the quality of the reconstructed image obtained by using the proposed optimal quadrature formulas with the conventional FBP, in which fast Fourier transform is used for the calculation of Fourier transform and its inversion. In the noise test,

Abdullo R. Hayotov

*Corresponding author:

Department of Mathematical Sciences, KAIST, 291 Daehak-ro, Yuseong-gu, Daejeon 34141, Republic of Korea,

V.I.Romanovskiy Institute of Mathematics, Uzbekistan Academy of Sciences, 81 M.Ulugbek str., Tashkent 100170, Uzbekistan,

E-mail: hayotov@mail.ru.

Soomin Jeon

Department of Mathematical Sciences, KAIST, 291 Daehak-ro, Yuseong-gu, Daejeon 34141, Republic of Korea

E-mail: soominjeon@kaist.ac.kr.

Chang-Ock Lee

Department of Mathematical Sciences, KAIST, 291 Daehak-ro, Yuseong-gu, Daejeon 34141, Republic of Korea

E-mail: colee@kaist.edu.

Kholmat M.Shadimetov

V.I.Romanovskiy Institute of Mathematics, Uzbekistan Academy of Sciences, 81 M.Ulugbek str., Tashkent 100170, Uzbekistan,

Tashkent Railway Engineering Institute, 1 Odilxojaev str., Tashkent 100167, Uzbekistan

E-mail: kholmatshadimetov@mail.ru.

the proposed algorithm provides more reliable results against the noise than the conventional FBP.

MSC: 41A05, 41A15, 65D30, 65D32.

Keywords: Optimal quadrature formula, square integrable function, error functional, Fourier transform, Radon transform, the filtered back-projection, CT image reconstruction.

1 Introduction

It is known that when complete continuous X-ray data are available Computed Tomography (CT) image can be reconstructed exactly using the filtered back-projection formula (see, for instance, [9, 15, 22, 28]). This formula gives interactions between the Radon transform, the Fourier transform and the back-projection transform. A description of the filtered back-projection formula along [22, Chapter 3] is provided below.

In the Cartesian system with x, y -axes consider a unit vector $(\cos \theta, \sin \theta)$. Then the line perpendicular to this vector with the distance t to the origin can be expressed as $\ell_{t,\theta}$: $x \cos \theta + y \sin \theta = t$. Assume the object is represented by a two variable function $\mu(x, y)$, which denotes the attenuation coefficient in X-ray CT applications. Then, the θ -view projection along the line $\ell_{t,\theta}$ can be expressed as

$$P(t, \theta) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \mu(x, y) \delta(x \cos \theta + y \sin \theta - t) dx dy,$$

where δ denotes the Dirac delta-function. The function $P(t, \theta)$ is known as the Radon transform of $\mu(x, y)$. A projection is formed by combining a set of line integrals. The simplest projection is a collection of parallel ray integrals as is given by $P(t, \theta)$ for a constant θ . This is known as a parallel beam projection. It should be noted that there are fan-beam in 2D and cone-beam in 3D projections [9, 15, 22].

The problem of CT is to reconstruct the function $\mu(x, y)$ from its projections $P(t, \theta)$. There are analytic and iterative methods for CT reconstruction. One of the widely used analytic methods of CT reconstruction is the filtered back-projection method. It can be modeled by

$$\mu(x, y) = \int_0^{\pi} \int_{-\infty}^{\infty} S(\omega, \theta) |\omega| e^{2\pi i \omega (x \cos \theta + y \sin \theta)} d\omega d\theta, \quad (1.1)$$

where

$$S(\omega, \theta) = \int_{-\infty}^{\infty} P(t, \theta) e^{-2\pi i \omega t} dt \quad (1.2)$$

is the 1D Fourier transform of $P(t, \theta)$. The inner integral of (1.1),

$$Q(t, \theta) = \int_{-\infty}^{\infty} S(\omega, \theta) |\omega| e^{2\pi i \omega t} d\omega, \quad (1.3)$$

is a 1D inverse Fourier transform of the product $S(\omega, \theta) |\omega|$, which represents a projection filtered by a 1D filter whose frequency representation is $|\omega|$. The outer integral performs back-projection. Therefore, the filtered back-projection consists of two steps: filtration and then back-projection.

Thus, the Fourier transforms play the main role in (1.1)-(1.3). But in practice, due to the fact that we have discrete values of the Radon transform, we have to approximately calculate the Fourier transforms in the filtered back-projection. For this purpose, it is necessary to consider the problem of approximate calculation of the integral

$$I(\varphi) = \int_a^b e^{2\pi i \omega x} \varphi(x) dx \quad (1.4)$$

with $\omega \in \mathbb{R}$. This type of integrals are called *highly oscillating integrals*. In most cases it is impossible to get the exact values of such integrals. Thus, they can be approximately calculated using the formulas of numerical integration. However, standard methods of numerical integration cannot be successfully applied for that. Therefore special effective methods should be developed for approximation of highly oscillating integrals. One of the first numerical integration formula for the integral (1.4) was obtained by Filon [16] in 1928 using a quadratic spline. Since then, for integrals of different types of highly oscillating functions many special effective methods have been developed, such as Filon-type method, Clenshaw-Curtis-Filon type method, Levin type methods, modified Clenshaw-Curtis method, generalized quadrature rule, and Gauss-Laguerre quadrature (see, for example, [1, 3-5, 21, 26, 29, 46, 47], for more review see, for instance, [14, 27, 30] and references therein). Recently, in [6-8], based on Sobolev's method, the problem of construction of optimal quadrature formulas in the sense of Sard for numerical calculation of integrals (1.4) with integer ω was studied in Hilbert spaces $L_2^{(m)}$ and $W_2^{(m, m-1)}$. Here, we consider the Sobolev space $L_2^{(m)}[a, b]$ of non-periodic, complex-valued functions defined on the interval $[a, b]$, which possess an absolute continuous $(m - 1)$ -th derivative on $[a, b]$, and whose m -th order derivative is square integrable [43, 45]. The space $L_2^{(m)}[a, b]$ is a Hilbert space with the inner product

$$\langle \varphi, \psi \rangle = \int_a^b \varphi^{(m)}(x) \bar{\psi}^{(m)}(x) dx, \quad (1.5)$$

where $\varphi^{(m)}$ is the m -th order derivative of the function φ with respect to x , $\bar{\psi}$ is the complex conjugate function to the function ψ and the norm of the function φ is correspondingly defined by the formula

$$\|\varphi\|_{L_2^{(m)}[a, b]} = \langle \varphi, \varphi \rangle^{1/2}.$$

The aim of the present work is to construct optimal quadrature formulas in the sense of Sard in the Sobolev space $L_2^{(m)}$ for numerical integration of the integral (1.4) with real ω using the results of the work [8]. Then to approximate reconstruction of CT image, we apply the obtained optimal quadrature formulas to the Fourier transform and its inversion in (1.1). Recently we got the results when $m = 1$ [20]. There is a work by Reider and Faridani [31] for an optimal reconstruction algorithm, where the optimal L^2 -convergence rates of filtered back-projection

(FBP) algorithm was provided with assumption that the Radon transform was analytically computable.

The rest of the paper is organized as follows. Section 2 is devoted to construction of optimal quadrature formulas in the sense of Sard in the space $L_2^{(m)}$ for numerical calculation of Fourier integrals. There are obtained analytic formulas for optimal coefficients using the discrete analogue of the differential operator $\frac{d^{2m}}{dx^{2m}}$. In section 3, the obtained optimal quadrature formulas for the cases $m = 2$ and $m = 3$ are applied for CT image reconstruction by approximating Fourier transforms in the filtered back-projection formula.

2 Optimal quadrature formulas for Fourier integrals in the space $L_2^{(m)}[a, b]$

In $L_2^{(m)}[a, b]$ space for approximation of the integral (1.4), we consider the following quadrature formula

$$\int_a^b e^{2\pi i \omega x} \varphi(x) dx \cong \sum_{\beta=0}^N C_\beta \varphi(x_\beta) \quad (2.1)$$

with the error

$$(\ell, \varphi) = \int_a^b e^{2\pi i \omega x} \varphi(x) dx - \sum_{\beta=0}^N C_\beta \varphi(x_\beta), \quad (2.2)$$

where $(\ell, \varphi) = \int_{-\infty}^{\infty} \ell(x) \varphi(x) dx$ is the value of the error functional ℓ at the function φ . Here the error functional ℓ has the form

$$\ell(x) = e^{2\pi i \omega x} \varepsilon_{[a,b]}(x) - \sum_{\beta=0}^N C_\beta \delta(x - x_\beta), \quad (2.3)$$

C_β are coefficients, $x_\beta = h\beta + a$ ($\in [a, b]$) are nodes of the formula (2.1), $h = \frac{b-a}{N}$, $N \in \mathbb{N}$, $i^2 = -1$, $\omega \in \mathbb{R}$, $\varepsilon_{[a,b]}(x)$ is the characteristic function of the interval $[a, b]$, and δ is the Dirac delta-function. We mention that the coefficients C_β depend on ω , h and m .

The error of the formula (2.1) defines a linear functional in $L_2^{(m)*}[a, b]$, where $L_2^{(m)*}[a, b]$ is the conjugate space to the space $L_2^{(m)}[a, b]$. Since the functional (2.3) is defined on the space $L_2^{(m)}$, the conditions

$$(\ell, x^\alpha) = 0, \quad \alpha = 0, 1, \dots, m-1 \quad (2.4)$$

should be fulfilled. The conditions (2.4) mean the exactness of the quadrature formula (2.1) for all algebraic polynomials of degree less than or equal to $m-1$. Hence we get that for a function from the space $C^m[a, b]$ the order of convergence of the optimal quadrature formula (2.1) is $O(h^m)$.

It should be mentioned that from equalities (2.4) one can get the condition $N+1 \geq m$ for existence of the optimal quadrature formula of the form (2.1) in the space $L_2^{(m)}$.

Sard's optimization problem of numerical integration formulas of the form (2.1) in the space $L_2^{(m)}[a, b]$ is the problem of finding the minimum of the norm of the error functional ℓ by coefficients C_β , i.e., to find coefficients C_β satisfying the equality

$$\|\mathring{\ell}\|_{L_2^{(m)*}[a, b]} = \inf_{C_\beta} \|\ell\|_{L_2^{(m)*}[a, b]}. \quad (2.5)$$

The coefficients satisfying the last equality are called *optimal coefficients* and they are denoted as \mathring{C}_β . The quadrature formula with coefficients \mathring{C}_β is called *the optimal quadrature formula in the sense of Sard*, and $\mathring{\ell}$ is the error functional corresponding to the optimal quadrature formula.

The solution of Sard's problem gives the sharp upper bound for the error (2.2) of functions φ from the space $L_2^{(m)}[a, b]$ as follows

$$|(\ell, \varphi)| \leq \|\varphi\|_{L_2^{(m)}[a, b]} \|\mathring{\ell}\|_{L_2^{(m)*}[a, b]}.$$

This problem, for the quadrature formulas of the form (2.1) with $\omega = 0$, was first studied by Sard [32] in the space $L_2^{(m)}$ for some m . Since then, it was investigated by many authors (see, for instance, [11, 13, 12, 18, 24, 25, 37]) using spline method, ϕ -function method, and Sobolev's method. Finally, in the works [23, 34, 36] this problem was solved for any $m \in \mathbb{N}$ with equally spaced nodes and the explicit expressions for the optimal coefficients have been obtained (see Theorem 4).

It should be noted that the problem of construction of lattice optimal cubature formulas in the space $L_2^{(m)}$ of multi-variable functions was first stated and investigated by Sobolev [43, 45]. Further, in this section, based on the results of the work [8], we solve Sard's problem on construction of optimal quadrature formulas of the form (2.1) for $\omega \in \mathbb{R}$ with $\omega \neq 0$, first for the interval $[0, 1]$ and then using a linear transformation for the interval $[a, b]$. For this we use the following auxiliary results.

2.1 Preliminaries

We need the concept of discrete argument functions and operations on them. For this we give the definition for functions of discrete argument by following [43, 45].

Assume that the nodes x_β are equally spaced, i.e., $x_\beta = h\beta$, h is a small positive parameter, and φ and ψ are complex-valued functions defined on the real line \mathbb{R} or on an interval of \mathbb{R} . The function $\varphi(h\beta)$ given on some set of integer values of β is called *a function of discrete argument*. The *inner product* of two discrete argument functions $\varphi(h\beta)$ and $\psi(h\beta)$ is defined by

$$[\varphi(h\beta), \psi(h\beta)] = \sum_{\beta=-\infty}^{\infty} \varphi(h\beta) \cdot \bar{\psi}(h\beta)$$

if the series on the right hand side of the last equality converges absolutely. The *convolution* of two functions $\varphi(h\beta)$ and $\psi(h\beta)$ is the following inner product

$$\varphi(h\beta) * \psi(h\beta) = [\varphi(h\gamma), \psi(h\beta - h\gamma)] = \sum_{\gamma=-\infty}^{\infty} \varphi(h\gamma) \cdot \bar{\psi}(h\beta - h\gamma).$$

We note that coefficients of optimal quadrature formulas and interpolation splines in the spaces $L_2^{(m)}$ and $W_2^{(m, m-1)}$ depend on the roots of the Euler-Frobenius type polynomials (see, for instance, [2, 8, 10, 23, 33, 37–42, 44, 45]). The Euler-Frobenius polynomials $E_k(x)$, $k = 1, 2, \dots$, are defined as follows (see, for instance, [17, 45]):

$$E_k(x) = \frac{(1-x)^{k+2}}{x} \left(x \frac{d}{dx} \right)^k \frac{x}{(1-x)^2}, \quad k = 0, 1, 2, \dots \quad (2.6)$$

The coefficients of the Euler-Frobenius polynomial $E_k(x) = \sum_{s=0}^k a_s x^s$ of degree k are expressed by the following formula which was obtained by Euler:

$$a_s = \sum_{j=0}^s (-1)^j \binom{k+2}{j} (s+1-j)^{k+1}.$$

In [17] it was shown that all roots q_j , $j = 1, 2, \dots, k$, of the polynomial $E_k(x)$ are real, negative and distinct, that is:

$$q_1 < q_2 < \dots < q_k < 0.$$

Furthermore, these roots satisfy the relation

$$q_j \cdot q_{k+1-j} = 1, \quad j = 1, 2, \dots, k.$$

For the Euler-Frobenius polynomials $E_k(x)$ the following identity holds

$$E_k(x) = x^k E_k\left(\frac{1}{x}\right), \quad (2.7)$$

and also the following is true.

Theorem 1 (Lemma 3 of [33]). *Polynomial $Q_k(x)$ which is defined by the formula*

$$Q_k(x) = (x-1)^{k+1} \sum_{i=0}^{k+1} \frac{\Delta^i 0^{k+1}}{(x-1)^i} \quad (2.8)$$

is the Euler-Frobenius polynomial (2.6) of degree k , i.e., $Q_k(x) = E_k(x)$, where $\Delta^i 0^k = \sum_{l=1}^i (-1)^{i-l} \binom{i}{l} l^k$.

We also use the formula

$$\sum_{\gamma=0}^{n-1} q^\gamma \gamma^k = \frac{1}{1-q} \sum_{i=0}^k \left(\frac{q}{1-q} \right)^i \Delta^i 0^k - \frac{q^n}{1-q} \sum_{i=0}^k \left(\frac{q}{1-q} \right)^i \Delta^i \gamma^k |_{\gamma=n}, \quad (2.9)$$

which is given in [19], where $\Delta^i \gamma^k$ is the finite difference of order i of γ^k and q is the ratio of a geometric progression.

In finding the analytic formulas for coefficients of optimal formulas in the space $L_2^{(m)}$ by Sobolev method the discrete analogue $D_m(h\beta)$ of the operator $\frac{d^{2m}}{dx^{2m}}$ plays the main role. This discrete analogue satisfies the equality

$$hD_m(h\beta) * G_m(h\beta) = \delta_d(h\beta), \quad (2.10)$$

where $G_m(h\beta)$ is the discrete argument function for the function

$$G_m(x) = \frac{|x|^{2m-1}}{2(2m-1)!}, \quad (2.11)$$

and $\delta_d(h\beta)$ is equal to 0 when $\beta \neq 0$ and 1 when $\beta = 0$.

We note that the operator $D_m(h\beta)$ was introduced and studied by Sobolev [43]. In [35] the discrete function $D_m(h\beta)$ was constructed and the following was proved.

Theorem 2 *The discrete analogue of the differential operator $\frac{d^{2m}}{dx^{2m}}$ has the form*

$$D_m(h\beta) = p \begin{cases} \sum_{k=1}^{m-1} A_k q_k^{|\beta|-1} & \text{for } |\beta| \geq 2, \\ 1 + \sum_{k=1}^{m-1} A_k & \text{for } |\beta| = 1, \\ C + \sum_{k=1}^{m-1} \frac{A_k}{q_k} & \text{for } \beta = 0, \end{cases} \quad (2.12)$$

where

$$p = \frac{(2m-1)!}{h^{2m}}, \quad A_k = \frac{(1-q_k)^{2m+1}}{E_{2m-1}(q_k)}, \quad C = -2^{2m-1}, \quad (2.13)$$

$E_{2m-1}(x)$ is the Euler-Frobenius polynomial of degree $2m-1$, q_k are the roots of the Euler-Frobenius polynomial $E_{2m-2}(x)$, $|q_k| < 1$, and h is a small positive parameter.

In addition, some properties of $D_m(h\beta)$ were studied in the works [35, 43]. Here we give the following.

Theorem 3 *The discrete argument function $D_m(h\beta)$ and the monomials $(h\beta)^k$ are related to each other as follows:*

$$\sum_{\beta=-\infty}^{\infty} D_m(h\beta)(h\beta)^k = \begin{cases} 0 & \text{for } 0 \leq k \leq 2m-1, \\ (2m)! & \text{for } k = 2m. \end{cases} \quad (2.14)$$

Now we give the results of the works [34, 33] on the optimal quadrature formulas of the form (2.1) in the sense of Sard and on the norm of the optimal error functional corresponding to the case $\omega = 0$.

Theorem 4 *Coefficients of the optimal quadrature formulas of the form*

$$\int_0^1 \varphi(x) dx \cong \sum_{\beta=0}^N C_\beta \varphi(h\beta) \quad (2.15)$$

in the space $L_2^{(m)}[0, 1]$ have the form

$$\begin{aligned} \mathring{C}_0 &= h \left(\frac{1}{2} - \sum_{k=1}^{m-1} d_k \frac{q_k - q_k^N}{1 - q_k} \right), \\ \mathring{C}_\beta &= h \left(1 + \sum_{k=1}^{m-1} d_k \left(q_k^\beta + q_k^{N-\beta} \right) \right), \quad \beta = 1, 2, \dots, N-1, \\ \mathring{C}_N &= h \left(\frac{1}{2} - \sum_{k=1}^{m-1} d_k \frac{q_k - q_k^N}{1 - q_k} \right), \end{aligned} \quad (2.16)$$

where d_k satisfy the system

$$\sum_{k=1}^{m-1} d_k \sum_{i=1}^j \frac{q_k + (-1)^{i+1} q_k^{N+i}}{(q_k - 1)^{i+1}} \Delta^i 0^j = \frac{B_{j+1}}{j+1}, \quad j = 1, 2, \dots, m-1,$$

q_k are roots of the Euler-Frobenius polynomial $E_{2m-2}(x)$ of degree $2m-2$ with $|q_k| < 1$, and $\Delta^i 0^j = \sum_{l=1}^i (-1)^{i-l} \binom{i}{l} l^j$.

Theorem 5 The norm of the error functional of the optimal quadrature formula (2.15) on the space $L_2^{(m)}[0, 1]$ has the form

$$\|\ell\|_{L_2^{(m)*}[0,1]}^2 = (-1)^{m+1} \left(\frac{h^{2m} B_{2m}}{(2m)!} + \frac{2h^{2m+1}}{(2m)!} \sum_{k=1}^{m-1} d_k \sum_{i=1}^{2m} \frac{-q_k^{N+i} + (-1)^i q_k}{(1-q_k)^{i+1}} \Delta^i 0^{2m} \right),$$

where B_{2m} is the Bernoulli number, and q_k, d_k and $\Delta^i 0^{2m}$ are given in Theorem 4.

2.2 Construction of optimal quadrature formulas for the interval $[0, 1]$

Here we obtain optimal quadrature formulas of the form (2.1) for the interval $[0, 1]$ when $\omega \in \mathbb{R}$ and $\omega \neq 0$. In the space $L_2^{(m)}[0, 1]$, using the results of Sections 2, 3 and 5 of [8], for the coefficients of the optimal quadrature formulas in the sense of Sard of the form

$$\int_0^1 e^{2\pi i \omega x} \varphi(x) dx \cong \sum_{\beta=0}^N C_\beta \varphi(h\beta) \quad (2.17)$$

for $\omega \in \mathbb{R}$ with $\omega \neq 0$, we get the following system of linear equations

$$\sum_{\gamma=0}^N C_\gamma G_m(h\beta - h\gamma) + P_{m-1}(h\beta) = f_m(h\beta), \quad \beta = 0, 1, \dots, N, \quad (2.18)$$

$$\sum_{\gamma=0}^N C_\gamma (h\gamma)^\alpha = g_\alpha, \quad \alpha = 0, 1, \dots, m-1, \quad (2.19)$$

where $P_{m-1}(h\beta) = \sum_{\alpha=0}^{m-1} p_\alpha (h\beta)^\alpha$ is a polynomial of degree $m-1$ with complex coefficients,

$$f_m(h\beta) = \int_0^1 e^{2\pi i \omega x} G_m(x - h\beta) dx \quad (2.20)$$

$$= - \sum_{\alpha=0}^{2m-1} \frac{(h\beta)^{2m-1-\alpha} (-1)^\alpha g_\alpha}{2\alpha! (2m-1-\alpha)!} + \frac{e^{2\pi i \omega h\beta}}{(2\pi i \omega)^{2m}} - \sum_{k=0}^{2m-1} \frac{(h\beta)^{2m-1-k}}{(2m-1-k)! (2\pi i \omega)^{k+1}},$$

$$g_\alpha = \int_0^1 e^{2\pi i \omega x} x^\alpha dx \quad (2.21)$$

$$= \sum_{k=0}^{\alpha-1} \frac{(-1)^k \alpha! e^{2\pi i \omega}}{(\alpha-k)! (2\pi i \omega)^{k+1}} + \frac{(-1)^\alpha \alpha!}{(2\pi i \omega)^{\alpha+1}} (e^{2\pi i \omega} - 1), \quad \alpha = 0, 1, \dots,$$

$G_m(x)$ is defined by (2.11), and $h = \frac{1}{N}$ for $N \in \mathbb{N}$.

In the system (2.18)-(2.19) unknowns are the optimal coefficients \mathring{C}_β , $\beta = 0, 1, \dots, N$ and p_α , $\alpha = 0, 1, \dots, m-1$. We point out that when $N+1 \geq m$ the system (2.18)-(2.19) has a unique solution. This solution satisfies conditions (2.4) and the equality (2.5). It should be noted that the existence and uniqueness of the solution for such type of systems were studied, for example, in [40, 43, 45].

We are interested in finding explicit formulas for the optimal coefficients \mathring{C}_β , $\beta = 0, 1, \dots, N$ and unknown polynomial $P_{m-1}(h\beta)$ satisfying the system (2.18)-(2.19). The system (2.18)-(2.19) is solved similarly as the system (34)-(35) of [8] by Sobolev's method, using the discrete analogue $D_m(h\beta)$ of the differential operator $\frac{d^{2m}}{dx^{2m}}$.

We formulate the results of this section as the following two theorems.

Theorem 6 For real ω with $\omega h \notin \mathbb{Z}$, the coefficients of optimal quadrature formulas of the form (2.1) in the space $L_2^{(m)}[0, 1]$ when $N+1 \geq m$ are expressed by formulas

$$\mathring{C}_0 = h \left(\frac{e^{2\pi i \omega h} K_{\omega, m}}{e^{2\pi i \omega h} - 1} - \frac{1}{2\pi i \omega h} + \sum_{k=1}^{m-1} \left(a_k \frac{q_k}{q_k - 1} + b_k \frac{q_k^N}{1 - q_k} \right) \right), \quad (2.22)$$

$$\mathring{C}_\beta = h \left(e^{2\pi i \omega h \beta} K_{\omega, m} + \sum_{k=1}^{m-1} \left(a_k q_k^\beta + b_k q_k^{N-\beta} \right) \right), \quad \beta = 1, 2, \dots, N-1, \quad (2.23)$$

$$\mathring{C}_N = h \left(\frac{e^{2\pi i \omega} K_{\omega, m}}{1 - e^{2\pi i \omega h}} + \frac{e^{2\pi i \omega}}{2\pi i \omega h} + \sum_{k=1}^{m-1} \left(a_k \frac{q_k^N}{1 - q_k} + b_k \frac{q_k}{q_k - 1} \right) \right), \quad (2.24)$$

where a_k and b_k are defined by the following system of $(2m-2)$ linear equations

$$\begin{aligned} & \sum_{k=1}^{m-1} a_k \left[\sum_{t=1}^j \frac{q_k \Delta^t 0^j}{(q_k - 1)^{t+1}} \right] + \sum_{k=1}^{m-1} b_k \left[\sum_{t=1}^j \frac{q_k^{N+t} \Delta^t 0^j}{(1 - q_k)^{t+1}} \right] \\ &= \frac{j!}{(2\pi i \omega h)^{j+1}} - \sum_{t=1}^j \frac{e^{2\pi i \omega h} K_{\omega, m} \Delta^t 0^j}{(e^{2\pi i \omega h} - 1)^{t+1}}, \quad j = 1, 2, \dots, m-1, \\ & \sum_{k=1}^{m-1} a_k \left[\sum_{t=1}^j \frac{q_k^t \Delta^t 0^j}{(1 - q_k)^{t+1}} - \sum_{\alpha=1}^j h^{\alpha-j} (j)_\alpha \sum_{t=1}^\alpha \frac{q_k^{N+t} \Delta^t 0^\alpha}{(1 - q_k)^{t+1}} \right] \\ &+ \sum_{k=1}^{m-1} b_k \left[\sum_{t=1}^j \frac{q_k^{N+1} \Delta^t 0^j}{(q_k - 1)^{t+1}} - \sum_{\alpha=1}^j h^{\alpha-j} (j)_\alpha \sum_{t=1}^\alpha \frac{q_k \Delta^t 0^\alpha}{(q_k - 1)^{t+1}} \right] \\ &= \frac{(-1)^{j+1} j!}{(2\pi i \omega h)^{j+1}} + \sum_{\alpha=1}^j h^{\alpha-j} \frac{(-1)^\alpha j! e^{2\pi i \omega}}{(j-\alpha)! (2\pi i \omega h)^{\alpha+1}} - \frac{K_{\omega, m}}{1 - e^{2\pi i \omega h}} \sum_{t=1}^j \left(\frac{e^{2\pi i \omega h}}{1 - e^{2\pi i \omega h}} \right)^t \Delta^t 0^j \\ &+ \frac{K_{\omega, m} e^{2\pi i \omega}}{1 - e^{2\pi i \omega h}} \sum_{\alpha=1}^j h^{\alpha-j} (j)_\alpha \sum_{t=1}^\alpha \left(\frac{e^{2\pi i \omega h}}{1 - e^{2\pi i \omega h}} \right)^t \Delta^t 0^\alpha, \quad j = 1, 2, \dots, m-1, \end{aligned} \quad (2.25)$$

q_k are roots of the Euler-Frobenius polynomial $E_{2m-2}(x)$ with $|q_k| < 1$, and

$$K_{\omega, m} = \left(\frac{\sin \pi \omega h}{\pi \omega h} \right)^{2m} \frac{(2m-1)!}{2 \sum_{\alpha=0}^{m-2} a_\alpha \cos[2\pi \omega h(m-1-\alpha)] + a_{m-1}}. \quad (2.26)$$

Here $a_\alpha = \sum_{j=0}^\alpha (-1)^j \binom{2m}{j} (s+1-j)^{2m-1}$ are the coefficients of the Euler-Frobenius polynomial $E_{2m-2}(x)$ of degree $2m-2$.

Theorem 7 For $\omega h \in \mathbb{Z}$ with $\omega \neq 0$, the coefficients of optimal quadrature formulas of the form (2.1) in the space $L_2^{(m)}[0, 1]$ when $N + 1 \geq m$ are expressed by formulas

$$\begin{aligned}\check{C}_0 &= h \left(-\frac{1}{2\pi i \omega h} + \sum_{k=1}^{m-1} \left(a_k \frac{q_k}{q_k - 1} + b_k \frac{q_k^N}{1 - q_k} \right) \right), \\ \check{C}_\beta &= h \sum_{k=1}^{m-1} \left(a_k q_k^\beta + b_k q_k^{N-\beta} \right), \quad \beta = 1, 2, \dots, N-1, \\ \check{C}_N &= h \left(\frac{e^{2\pi i \omega}}{2\pi i \omega h} + \sum_{k=1}^{m-1} \left(a_k \frac{q_k^N}{1 - q_k} + b_k \frac{q_k}{q_k - 1} \right) \right),\end{aligned}$$

where a_k and b_k , $k = 1, 2, \dots, m-1$, are defined by the following system of $(2m-2)$ linear equations:

$$\begin{aligned}& \sum_{k=1}^{m-1} a_k \left[\sum_{t=1}^j \frac{q_k \Delta^t 0^j}{(q_k - 1)^{t+1}} \right] + \sum_{k=1}^{m-1} b_k \left[\sum_{t=1}^j \frac{q_k^{N+t} \Delta^t 0^j}{(1 - q_k)^{t+1}} \right] = \frac{j!}{(2\pi i \omega h)^{j+1}}, \quad j = 1, 2, \dots, m-1, \\ & \sum_{k=1}^{m-1} a_k \left[\sum_{t=1}^j \frac{q_k^t \Delta^t 0^j}{(1 - q_k)^{t+1}} - \sum_{\alpha=1}^j h^{\alpha-j} \binom{j}{\alpha} \sum_{t=1}^{\alpha} \frac{q_k^{N+t} \Delta^t 0^\alpha}{(1 - q_k)^{t+1}} \right] \\ & + \sum_{k=1}^{m-1} b_k \left[\sum_{t=1}^j \frac{q_k^{N+1} \Delta^t 0^j}{(q_k - 1)^{t+1}} - \sum_{\alpha=1}^j h^{\alpha-j} \binom{j}{\alpha} \sum_{t=1}^{\alpha} \frac{q_k \Delta^t 0^\alpha}{(q_k - 1)^{t+1}} \right] \\ & = \frac{(-1)^{j+1} j!}{(2\pi i \omega h)^{j+1}} + \sum_{\alpha=1}^j h^{\alpha-j} \frac{(-1)^\alpha j! e^{2\pi i \omega}}{(j-\alpha)! (2\pi i \omega h)^{\alpha+1}}, \quad j = 1, 2, \dots, m-1.\end{aligned}$$

Here q_k are the roots of the Euler-Frobenius polynomial $E_{2m-2}(x)$ and $|q_k| < 1$.

We note that Theorem 6 is generalization of Theorem 6 in [6] for real ω with $\omega h \notin \mathbb{Z}$ while Theorem 7 for $\omega h \in \mathbb{Z}$ with $\omega \neq 0$ is the same with Theorem 7 of the work [6]. Theorem 6 is proved similarly as Theorem 6 of [6]. Therefore, it is sufficient to give a brief proof of Theorem 6.

The brief proof of Theorem 6. First, such as in Theorem 5 of the work [8], using the discrete function $D_m(h\beta)$, for optimal coefficients \check{C}_β , $\beta = 1, 2, \dots, N-1$, when ω is real and $\omega h \notin \mathbb{Z}$ we get the following formula

$$\check{C}_\beta = h \left(K_{\omega, m} e^{2\pi i \omega h \beta} + \sum_{k=1}^{m-1} \left(a_k q_k^\beta + b_k q_k^{N-\beta} \right) \right), \quad \beta = 1, 2, \dots, N-1, \quad (2.27)$$

where a_k , b_k , $k = 1, 2, \dots, m-1$, $K_{\omega, m}$ are unknowns, and q_k are roots of the Euler-Frobenius polynomial $E_{2m-2}(x)$ of degree $2m-2$ with $|q_k| < 1$.

Next, it is sufficient to find a_k , b_k , $k = 1, 2, \dots, m-1$, $K_{\omega, m}$, optimal coefficients \check{C}_0 , \check{C}_N and unknown polynomial $P_{m-1}(h\beta)$ of degree $m-1$. Now putting the form (2.27) of optimal coefficients \check{C}_β , $\beta = 1, 2, \dots, N-1$, into (2.18), using (2.7)-(2.9),

after some simplifications, we get the following identity with respect to $(h\beta)$

$$\begin{aligned}
& e^{2\pi i\omega h\beta} \left(\frac{h^{2m} K_{\omega,m}}{(2m-1)!} \sum_{i=0}^{2m-1} \frac{e^{2\pi i\omega h} \Delta^i 0^{2m-1}}{(e^{2\pi i\omega h} - 1)^{i+1}} \right) \\
& + \frac{(h\beta)^{2m-1}}{(2m-1)!} \left(\mathring{C}_0 - \frac{K_{\omega,m} h e^{2\pi i\omega h}}{e^{2\pi i\omega h} - 1} - h \sum_{k=1}^{m-1} \frac{a_k q_k - b_k q_k^N}{q_k - 1} \right) \\
& - \sum_{j=1}^{2m-1} \frac{(h\beta)^{2m-1-j}}{j!(2m-1-j)!} h^{j+1} \left(\sum_{i=0}^j \frac{K_{\omega,m} e^{2\pi i\omega h} \Delta^i 0^j}{(e^{2\pi i\omega h} - 1)^{i+1}} + \sum_{k=1}^{m-1} \sum_{i=0}^j \frac{a_k q_k + (-1)^{i+1} b_k q_k^{N+i}}{(q_k - 1)^{i+1}} \Delta^i 0^j \right) \\
& - \sum_{j=m}^{2m-1} \frac{(h\beta)^{2m-1-j}}{j!(2m-1-j)!} \frac{(-1)^j}{2} \sum_{\gamma=0}^N \mathring{C}_\gamma (h\gamma)^j + P_{m-1}(h\beta) \\
& = \frac{e^{2\pi i\omega h\beta}}{(2\pi i\omega)^{2m}} - \frac{(h\beta)^{2m-1}}{(2m-1)!(2\pi i\omega)} \\
& \quad - \sum_{j=1}^{2m-1} \frac{(h\beta)^{2m-1-j}}{j!(2m-1-j)!} \frac{j!}{(2\pi i\omega)^{j+1}} - \sum_{j=m}^{2m-1} \frac{(h\beta)^{2m-1-j}}{j!(2m-1-j)!} \frac{(-1)^j}{2} g_j.
\end{aligned}$$

Hence, the formulas (2.22) and (2.26) for \mathring{C}_0 and $K_{\omega,m}$ are obtained by equating the coefficients of the terms $(h\beta)^{2m-1}$ and $e^{2\pi i\omega h\beta}$, respectively. Then (2.24) is found from (2.19) when $\alpha = 0$, using (2.22) and (2.27). From the last identity equating the coefficients of the terms $(h\beta)^{2m-1-j}$ for $j = 1, 2, \dots, m-1$ and using (2.19) for $\alpha = 1, 2, \dots, m-1$, taking into account (2.22), (2.24) and (2.27), we obtain the system (2.25). Finally, the polynomial $P_{m-1}(h\beta)$ is obtained from the identity by equating the coefficients of the terms $(h\beta)^{2m-1-j}$ for $j = m, m+1, \dots, 2m-1$. Theorem is proved. \square

2.3 Optimal quadrature formulas for the interval $[a, b]$

In this section we obtain the optimal quadrature formulas for the interval $[a, b]$ in the space $L_2^{(m)}$ by a linear transformation from the results of the previous section.

We consider construction of the optimal quadrature formulas of the form

$$\int_a^b e^{2\pi i\omega x} \varphi(x) \, dx \cong \sum_{\beta=0}^N C_{\beta,\omega}[a, b] \varphi(x_\beta) \quad (2.28)$$

in the Sobolev space $L_2^{(m)}[a, b]$. Here $C_{\beta,\omega}[a, b]$ are coefficients, $x_\beta = h\beta + a$ ($\in [a, b]$) are nodes of the formula (2.28), $\omega \in \mathbb{R}$, $i^2 = -1$, and $h = \frac{b-a}{N}$ for $N+1 \geq m$.

Now, by a linear transformation $x = (b-a)y + a$, where $0 \leq y \leq 1$, we obtain

$$\int_a^b e^{2\pi i\omega x} \varphi(x) \, dx = (b-a) e^{2\pi i\omega a} \int_0^1 e^{2\pi i\omega(b-a)y} \varphi((b-a)y + a) \, dy. \quad (2.29)$$

Then, applying Theorems 4, 6 and 7 to the integral on the right-hand side of the last equality, we have the following results which are optimal quadrature formulas of the form (2.28) in the sense of Sard in the space $L_2^{(m)}[a, b]$ for all real ω .

For the case $\omega = 0$, using Theorem 4 in (2.29), we get

Theorem 8 *Coefficients of the optimal quadrature formulas of the form*

$$\int_a^b \varphi(x) dx \cong \sum_{\beta=0}^N C_{\beta,0}[a,b] \varphi(h\beta + a)$$

in the space $L_2^{(m)}[a,b]$ have the form

$$\begin{aligned} \mathring{C}_{0,0}[a,b] &= h \left(\frac{1}{2} - \sum_{k=1}^{m-1} d_k \frac{q_k - q_k^N}{1 - q_k} \right), \\ \mathring{C}_{\beta,0}[a,b] &= h \left(1 + \sum_{k=1}^{m-1} d_k \left(q_k^\beta + q_k^{N-\beta} \right) \right), \quad \beta = 1, 2, \dots, N-1, \\ \mathring{C}_{N,0}[a,b] &= h \left(\frac{1}{2} - \sum_{k=1}^{m-1} d_k \frac{q_k - q_k^N}{1 - q_k} \right), \end{aligned}$$

where $h = \frac{b-a}{N}$ and d_k satisfy the system

$$\sum_{k=1}^{m-1} d_k \sum_{i=1}^j \frac{q_k + (-1)^{i+1} q_k^{N+i}}{(q_k - 1)^{i+1}} \Delta^i 0^j = \frac{B_{j+1}}{j+1}, \quad j = 1, 2, \dots, m-1.$$

Here B_{j+1} is the Bernoulli number, q_k are roots of the Euler-Frobenius polynomial $E_{2m-2}(x)$ of degree $(2m-2)$ with $|q_k| < 1$, and $\Delta^i 0^j = \sum_{l=1}^i (-1)^{i-l} \binom{i}{l} l^j$.

For the case $\omega \in \mathbb{R}$ and $\omega h \notin \mathbb{Z}$, applying Theorem 6 to the integral in (2.29), we obtain

Theorem 9 *For real ω with $\omega h \notin \mathbb{Z}$, the coefficients of optimal quadrature formulas of the form (2.28) in the space $L_2^{(m)}[a,b]$ when $N+1 \geq m$ are expressed by formulas*

$$\begin{aligned} \mathring{C}_{0,\omega}[a,b] &= h \left(\frac{e^{2\pi i \omega(a+h)} K_{\omega,m}}{e^{2\pi i \omega h} - 1} - \frac{e^{2\pi i \omega a}}{2\pi i \omega h} + \sum_{k=1}^{m-1} \left(a_k \frac{q_k}{q_k - 1} + b_k \frac{q_k^N}{1 - q_k} \right) \right), \\ \mathring{C}_{\beta,\omega}[a,b] &= h \left(e^{2\pi i \omega(h\beta+a)} K_{\omega,m} + \sum_{k=1}^{m-1} \left(a_k q_k^\beta + b_k q_k^{N-\beta} \right) \right), \quad \beta = 1, 2, \dots, N-1, \\ \mathring{C}_{N,\omega}[a,b] &= h \left(\frac{e^{2\pi i \omega b} K_{\omega,m}}{1 - e^{2\pi i \omega h}} + \frac{e^{2\pi i \omega a}}{2\pi i \omega h} + \sum_{k=1}^{m-1} \left(a_k \frac{q_k^N}{1 - q_k} + b_k \frac{q_k}{q_k - 1} \right) \right), \end{aligned}$$

where a_k and b_k are defined by the following system of $(2m - 2)$ linear equations:

$$\begin{aligned} & \sum_{k=1}^{m-1} a_k \left[\sum_{t=1}^j \frac{q_k \Delta^t 0^j}{(q_k - 1)^{t+1}} \right] + \sum_{k=1}^{m-1} b_k \left[\sum_{t=1}^j \frac{q_k^{N+t} \Delta^t 0^j}{(1 - q_k)^{t+1}} \right] \\ &= \frac{j! e^{2\pi i \omega a}}{(2\pi i \omega h)^{j+1}} - \sum_{t=1}^j \frac{e^{2\pi i \omega (a+h)} K_{\omega, m} \Delta^t 0^j}{(e^{2\pi i \omega h} - 1)^{t+1}}, \quad j = 1, 2, \dots, m-1, \\ & \sum_{k=1}^{m-1} a_k \left[\sum_{t=1}^j \frac{q_k^t \Delta^t 0^j}{(1 - q_k)^{t+1}} - \sum_{\alpha=1}^j \left(\frac{h}{b-a} \right)^{\alpha-j} (j)_{\alpha} \sum_{t=1}^{\alpha} \frac{q_k^{N+t} \Delta^t 0^{\alpha}}{(1 - q_k)^{t+1}} \right] \\ &+ \sum_{k=1}^{m-1} b_k \left[\sum_{t=1}^j \frac{q_k^{N+1} \Delta^t 0^j}{(q_k - 1)^{t+1}} - \sum_{\alpha=1}^j \left(\frac{h}{b-a} \right)^{\alpha-j} (j)_{\alpha} \sum_{t=1}^{\alpha} \frac{q_k \Delta^t 0^{\alpha}}{(q_k - 1)^{t+1}} \right] \\ &= \frac{(-1)^{j+1} j! e^{2\pi i \omega a}}{(2\pi i \omega h)^{j+1}} + \sum_{\alpha=1}^j \left(\frac{h}{b-a} \right)^{\alpha-j} \frac{(-1)^{\alpha} j! e^{2\pi i \omega b}}{(j-\alpha)! (2\pi i \omega h)^{\alpha+1}} - \frac{K_{\omega, m} e^{2\pi i \omega a}}{1 - e^{2\pi i \omega h}} \sum_{t=1}^j \left(\frac{e^{2\pi i \omega h}}{1 - e^{2\pi i \omega h}} \right)^t \Delta^t 0^j \\ &+ \frac{K_{\omega, m} e^{2\pi i \omega b}}{1 - e^{2\pi i \omega h}} \sum_{\alpha=1}^j \left(\frac{h}{b-a} \right)^{\alpha-j} (j)_{\alpha} \sum_{t=1}^{\alpha} \left(\frac{e^{2\pi i \omega h}}{1 - e^{2\pi i \omega h}} \right)^t \Delta^t 0^{\alpha}, \quad j = 1, 2, \dots, m-1, \end{aligned}$$

q_k are roots of the Euler-Frobenius polynomial $E_{2m-2}(x)$ with $|q_k| < 1$, and $K_{\omega, m}$ is defined by (2.26).

Lastly, for the case $\omega h \in \mathbb{Z}$ with $\omega \neq 0$ application Theorem 7 in (2.29) gives the following.

Theorem 10 For $\omega h \in \mathbb{Z}$ with $\omega \neq 0$, the coefficients of optimal quadrature formulas of the form (2.28) in the space $L_2^{(m)}[a, b]$ when $N + 1 \geq m$ are expressed by formulas

$$\begin{aligned} \mathring{C}_{0, \omega}[a, b] &= h \left(-\frac{e^{2\pi i \omega a}}{2\pi i \omega h} + \sum_{k=1}^{m-1} \left(a_k \frac{q_k}{q_k - 1} + b_k \frac{q_k^N}{1 - q_k} \right) \right), \\ \mathring{C}_{\beta, \omega}[a, b] &= h \sum_{k=1}^{m-1} \left(a_k q_k^{\beta} + b_k q_k^{N-\beta} \right), \quad \beta = 1, 2, \dots, N-1, \\ \mathring{C}_{N, \omega}[a, b] &= h \left(\frac{e^{2\pi i \omega b}}{2\pi i \omega h} + \sum_{k=1}^{m-1} \left(a_k \frac{q_k^N}{1 - q_k} + b_k \frac{q_k}{q_k - 1} \right) \right), \end{aligned}$$

where a_k and b_k , $k = 1, 2, \dots, m-1$, are defined by the following system of $(2m - 2)$ linear equations

$$\begin{aligned} & \sum_{k=1}^{m-1} a_k \left[\sum_{t=1}^j \frac{q_k \Delta^t 0^j}{(q_k - 1)^{t+1}} \right] + \sum_{k=1}^{m-1} b_k \left[\sum_{t=1}^j \frac{q_k^{N+t} \Delta^t 0^j}{(1 - q_k)^{t+1}} \right] = \frac{j! e^{2\pi i \omega a}}{(2\pi i \omega h)^{j+1}}, \quad j = 1, 2, \dots, m-1, \\ & \sum_{k=1}^{m-1} a_k \left[\sum_{t=1}^j \frac{q_k^t \Delta^t 0^j}{(1 - q_k)^{t+1}} - \sum_{\alpha=1}^j \left(\frac{h}{b-a} \right)^{\alpha-j} (j)_{\alpha} \sum_{t=1}^{\alpha} \frac{q_k^{N+t} \Delta^t 0^{\alpha}}{(1 - q_k)^{t+1}} \right] \\ &+ \sum_{k=1}^{m-1} b_k \left[\sum_{t=1}^j \frac{q_k^{N+1} \Delta^t 0^j}{(q_k - 1)^{t+1}} - \sum_{\alpha=1}^j \left(\frac{h}{b-a} \right)^{\alpha-j} (j)_{\alpha} \sum_{t=1}^{\alpha} \frac{q_k \Delta^t 0^{\alpha}}{(q_k - 1)^{t+1}} \right] \\ &= \frac{(-1)^{j+1} j! e^{2\pi i \omega a}}{(2\pi i \omega h)^{j+1}} + \sum_{\alpha=1}^j \left(\frac{h}{b-a} \right)^{\alpha-j} \frac{(-1)^{\alpha} j! e^{2\pi i \omega b}}{(j-\alpha)! (2\pi i \omega h)^{\alpha+1}}, \quad j = 1, 2, \dots, m-1. \end{aligned}$$

Here q_k are the roots of the Euler-Frobenius polynomial $E_{2m-2}(x)$ and $|q_k| < 1$.

From a practical point of view, to have the formulas for coefficients of the optimal quadrature formulas (2.28) for the first several values of m is very useful. Below we present explicit formulas of the optimal coefficients for $m = 2$ and 3, using Theorems 8, 9 and 10.

Remark 1. In the space $L_2^{(1)}[a, b]$ the coefficients of the optimal quadrature formulas (2.28) have the same forms as derived in [20], where it was shown that for functions with a continuous second derivative the convergence order of this optimal quadrature formula is $O(h^2)$.

Remark 2. The optimal quadrature formulas (2.28) for $m = 1$ and 2 have the same convergence order.

Corollary 1 *In the space $L_2^{(2)}[a, b]$ the coefficients of the optimal quadrature formulas (2.28) for $\omega = 0$ are written as*

$$\begin{aligned}\mathring{C}_{0,0}[a, b] &= h \left(\frac{1}{2} + \frac{q_1 - q_1^N}{2(1 - q_1)(1 + q_1^N)} \right), \\ \mathring{C}_{\beta,0}[a, b] &= h \left(1 - \frac{q_1^\beta + q_1^{N-\beta}}{2(1 + q_1^N)} \right), \quad \beta = 1, 2, \dots, N - 1, \\ \mathring{C}_{N,0}[a, b] &= h \left(\frac{1}{2} + \frac{q_1 - q_1^N}{2(1 - q_1)(1 + q_1^N)} \right),\end{aligned}$$

for $\omega \in \mathbb{R}$ and $\omega h \notin \mathbb{Z}$ are expressed as

$$\begin{aligned}\mathring{C}_{0,\omega}[a, b] &= h \left(\frac{e^{2\pi i \omega(a+h)} K_{\omega,2}}{e^{2\pi i \omega h} - 1} - \frac{e^{2\pi i \omega a}}{2\pi i \omega h} + a_1 \frac{q_1}{q_1 - 1} + b_1 \frac{q_1^N}{1 - q_1} \right), \\ \mathring{C}_{\beta,\omega}[a, b] &= h \left(e^{2\pi i \omega(h\beta+a)} K_{\omega,2} + a_1 q_1^\beta + b_1 q_1^{N-\beta} \right), \quad \beta = 1, 2, \dots, N - 1, \\ \mathring{C}_{N,\omega}[a, b] &= h \left(\frac{e^{2\pi i \omega b} K_{\omega,2}}{1 - e^{2\pi i \omega h}} + \frac{e^{2\pi i \omega b}}{2\pi i \omega h} + a_1 \frac{q_1^N}{1 - q_1} + b_1 \frac{q_1}{q_1 - 1} \right),\end{aligned}$$

and for $\omega h \in \mathbb{Z}$ with $\omega \neq 0$ have the form

$$\begin{aligned}\mathring{C}_{0,\omega}[a, b] &= h \left(-\frac{e^{2\pi i \omega a}}{2\pi i \omega h} + a_1 \frac{q_1}{q_1 - 1} + b_1 \frac{q_1^N}{1 - q_1} \right), \\ \mathring{C}_{\beta,\omega}[a, b] &= h \left(a_1 q_1^\beta + b_1 q_1^{N-\beta} \right), \quad \beta = 1, 2, \dots, N - 1, \\ \mathring{C}_{N,\omega}[a, b] &= h \left(\frac{e^{2\pi i \omega b}}{2\pi i \omega h} + a_1 \frac{q_1^N}{1 - q_1} + b_1 \frac{q_1}{q_1 - 1} \right),\end{aligned}$$

where $q_1 = \sqrt{3} - 2$,

$$\begin{aligned}a_1 &= \frac{B_\omega(e^{2\pi i \omega a} - e^{2\pi i \omega b} q_1^N)}{1 - q_1^{2N}}, & b_1 &= \frac{B_\omega(e^{2\pi i \omega b} - e^{2\pi i \omega a} q_1^N)}{1 - q_1^{2N}}, \\ B_\omega &= 6 \left(\frac{1}{(2\pi \omega h)^2} - \frac{K_{\omega,2}}{2 - 2 \cos 2\pi \omega h} \right), & K_{\omega,2} &= \left(\frac{\sin \pi \omega h}{\pi \omega h} \right)^4 \frac{3}{2 + \cos 2\pi \omega h}.\end{aligned}$$

Corollary 2 In the space $L_2^{(3)}[a, b]$ the coefficients of the optimal quadrature formulas (2.28) for $\omega = 0$ are written as

$$\begin{aligned}\mathring{C}_{0,0}[a, b] &= h \left(\frac{1}{2} - \sum_{k=1}^2 d_k \frac{q_k - q_k^N}{1 - q_k} \right), \\ \mathring{C}_{\beta,0}[a, b] &= h \left(1 + \sum_{k=1}^2 d_k \left(q_k^\beta + q_k^{N-\beta} \right) \right), \quad \beta = 1, 2, \dots, N-1, \\ \mathring{C}_{N,0}[a, b] &= h \left(\frac{1}{2} - \sum_{k=1}^2 d_k \frac{q_k - q_k^N}{1 - q_k} \right),\end{aligned}$$

where d_k are defined from the system

$$\begin{aligned}\sum_{k=1}^2 d_k \frac{q_k + q_k^{N+1}}{(q_k - 1)^2} &= \frac{1}{12}, \\ \sum_{k=1}^2 d_k \frac{q_k - q_k^{N+2}}{(q_k - 1)^3} &= \frac{-1}{24},\end{aligned}\tag{2.30}$$

for $\omega \in \mathbb{R}$ and $\omega h \notin \mathbb{Z}$ are expressed as

$$\begin{aligned}\mathring{C}_{0,\omega}[a, b] &= h \left(\frac{e^{2\pi i \omega (a+h)} K_{\omega,3}}{e^{2\pi i \omega h} - 1} - \frac{e^{2\pi i \omega a}}{2\pi i \omega h} + \sum_{k=1}^2 \left(a_k \frac{q_k}{q_k - 1} + b_k \frac{q_k^N}{1 - q_k} \right) \right), \\ \mathring{C}_{\beta,\omega}[a, b] &= h \left(e^{2\pi i \omega (h\beta+a)} K_{\omega,3} + \sum_{k=1}^2 \left(a_k q_k^\beta + b_k q_k^{N-\beta} \right) \right), \quad \beta = 1, 2, \dots, N-1, \\ \mathring{C}_{N,\omega}[a, b] &= h \left(\frac{e^{2\pi i \omega b} K_{\omega,3}}{1 - e^{2\pi i \omega h}} + \frac{e^{2\pi i \omega b}}{2\pi i \omega h} + \sum_{k=1}^2 \left(a_k \frac{q_k^N}{1 - q_k} + b_k \frac{q_k}{q_k - 1} \right) \right),\end{aligned}$$

where a_k and b_k are determined from the system

$$\begin{aligned}\sum_{k=1}^2 a_k \left[\frac{q_k}{(q_k - 1)^2} \right] + \sum_{k=1}^2 b_k \left[\frac{q_k^{N+1}}{(1 - q_k)^2} \right] &= \frac{e^{2\pi i \omega a}}{(2\pi i \omega h)^2} - \frac{e^{2\pi i \omega (h+a)} K_{\omega,3}}{(e^{2\pi i \omega h} - 1)^2}, \\ \sum_{k=1}^2 a_k \left[\frac{q_k}{(q_k - 1)^3} \right] + \sum_{k=1}^2 b_k \left[\frac{q_k^{N+2}}{(1 - q_k)^3} \right] &= \frac{e^{2\pi i \omega a}}{(2\pi i \omega h)^3} - \frac{e^{2\pi i \omega a}}{2(2\pi i \omega h)^2} - \frac{e^{2\pi i \omega (h+a)} K_{\omega,3}}{(e^{2\pi i \omega h} - 1)^3}, \\ \sum_{k=1}^2 a_k \left[\frac{q_k^{N+1}}{(1 - q_k)^2} \right] + \sum_{k=1}^2 b_k \left[\frac{q_k}{(q_k - 1)^2} \right] &= \frac{e^{2\pi i \omega b}}{(2\pi i \omega h)^2} - \frac{e^{2\pi i \omega (h+b)} K_{\omega,3}}{(1 - e^{2\pi i \omega h})^2}, \\ \sum_{k=1}^2 a_k \left[\frac{q_k^2 - q_k^{N+2}}{(1 - q_k)^3} \right] + \sum_{k=1}^2 b_k \left[\frac{q_k^{N+1} - q_k}{(q_k - 1)^3} \right] &= (e^{2\pi i \omega b} - e^{2\pi i \omega a}) \left(\frac{1}{(2\pi i \omega h)^3} + \frac{1}{2(2\pi i \omega h)^2} + \frac{e^{4\pi i \omega h} K_{\omega,3}}{(1 - e^{2\pi i \omega h})^3} \right),\end{aligned}\tag{2.31}$$

and for $\omega h \in \mathbb{Z}$ with $\omega \neq 0$ have the form

$$\begin{aligned}\mathring{C}_{0,\omega}[a, b] &= h \left(-\frac{e^{2\pi i \omega a}}{2\pi i \omega h} + \sum_{k=1}^2 \left(a_k \frac{q_k}{q_k - 1} + b_k \frac{q_k^N}{1 - q_k} \right) \right), \\ \mathring{C}_{\beta,\omega}[a, b] &= h \left(\sum_{k=1}^2 \left(a_k q_k^\beta + b_k q_k^{N-\beta} \right) \right), \quad \beta = 1, 2, \dots, N-1, \\ \mathring{C}_{N,\omega}[a, b] &= h \left(\frac{e^{2\pi i \omega b}}{2\pi i \omega h} + \sum_{k=1}^2 \left(a_k \frac{q_k^N}{1 - q_k} + b_k \frac{q_k}{q_k - 1} \right) \right),\end{aligned}$$

where a_k and b_k are defined as follows

$$\begin{aligned}\sum_{k=1}^2 a_k \left[\frac{q_k}{(q_k - 1)^2} \right] + \sum_{k=1}^2 b_k \left[\frac{q_k^{N+1}}{(1 - q_k)^2} \right] &= \frac{e^{2\pi i \omega a}}{(2\pi i \omega h)^2}, \\ \sum_{k=1}^2 a_k \left[\frac{q_k}{(q_k - 1)^3} \right] + \sum_{k=1}^2 b_k \left[\frac{q_k^{N+2}}{(1 - q_k)^3} \right] &= \frac{e^{2\pi i \omega a}}{(2\pi i \omega h)^3} - \frac{e^{2\pi i \omega a}}{2(2\pi i \omega h)^2}, \\ \sum_{k=1}^2 a_k \left[\frac{q_k^{N+1}}{(1 - q_k)^2} \right] + \sum_{k=1}^2 b_k \left[\frac{q_k}{(q_k - 1)^2} \right] &= \frac{e^{2\pi i \omega b}}{(2\pi i \omega h)^2}, \\ \sum_{k=1}^2 a_k \left[\frac{q_k^2 - q_k^{N+2}}{(1 - q_k)^3} \right] + \sum_{k=1}^2 b_k \left[\frac{q_k^{N+1} - q_k}{(q_k - 1)^3} \right] &= (e^{2\pi i \omega b} - e^{2\pi i \omega a}) \left(\frac{1}{(2\pi i \omega h)^3} + \frac{1}{2(2\pi i \omega h)^2} \right),\end{aligned}\tag{2.32}$$

$q_k, k = 1, 2$, are roots of the Euler-Frobenius polynomial

$$E_4(x) = x^4 + 26x^3 + 66x^2 + 26x + 1\tag{2.33}$$

with $|q_k| < 1$, and

$$K_{\omega,3} = \left(\frac{\sin \pi \omega h}{\pi \omega h} \right)^6 \frac{5!}{2[\cos 4\pi \omega h + 26 \cos 2\pi \omega h] + 66}.\tag{2.34}$$

3 Application: CT image reconstruction

In this section, we give numerical results of CT image reconstruction by applying the second and third-order optimal quadrature formulas (Corollaries 1 and 2) to calculate the Fourier transform and its inversion. One of the most commonly used CT reconstruction algorithms is FBP [9, 15, 22, 28] and as in [22], the implementation of the FBP consists of four steps: (1) sinogram acquisition, (2) Fourier transform of the sinogram, (3) application of Ram-Lak filter and the Fourier inversion, and (4) back-projection. To show the effect of the proposed optimal quadrature formula, we implement the FBP in two different ways: one uses *fft* and *ifft* for the Fourier transform and its inversion, respectively, and the other uses the proposed optimal quadrature formulas. Then, we compare the quality of the resulting

Algorithm 1 Reconstruction algorithm with optimal quadrature formula

- 1: A sinogram $P(t_m, \theta_k)$ for $t_m \in [a, b]$, $\theta \in [0, \pi]$ is given as a discrete form.
- 2: Compute the Fourier transform using the proposed optimal quadrature:

$$S(\omega, \theta) \cong S(\omega, \theta_k) = \sum_{m=0}^M \hat{C}_{m, -\omega} P(t_m, \theta_k), \quad \omega \in \mathbb{R}.$$

- 3: Compute the inverse Fourier transform using the proposed optimal quadrature:

$$Q(t, \theta) \cong Q(t, \theta_k) = \sum_{n=0}^N \hat{C}_{n, t} S(\omega_n, \theta_k) |\omega_n|.$$

- 4: Reconstruct the CT image using back-projection:

$$f(x, y) = \int_0^\pi Q(t, \theta) d\theta \cong \frac{\pi}{K} \sum_{k=0}^{K-1} Q(t, \theta_k).$$




Fig. 1 Shepp-Logan phantom images of size 512×512 : without noise (left) and with noise (right)

reconstructed images. Also, we test the performance of the proposed algorithm by adding the 10% of Poisson noise to the given sinogram. In this section, we deal with the second and third order optimal quadrature formulas only. For the numerical results with the first order optimal quadrature formula, see [20].

Algorithm 1 is the pseudo code of the algorithm for CT image reconstruction using optimal quadrature formulas. For *Step 2* and *Step 3*, the optimal quadrature formulas are used for approximating Fourier integrals. For the numerical experiment, we use the Shepp-Logan phantom (SL) in [22] and we also consider the case with noise (see Fig. 1). Both are of size 512×512 and the sinograms are generated using half rotation sampling with sampling angle 0.5° . For the implementation, we use $q_k = \frac{t_k + \sqrt{t_k^2 - 4}}{2}$ ($k = 1, 2$), where $t_k = -13 \pm \sqrt{105}$, the roots of (2.33) with $|q_k| < 1$. For q_1 and q_2 , d_k are obtained from the 2×2 linear system (2.30), and a_k and b_k are obtained from the 4×4 linear systems (2.31) and (2.32). For the numerical experiments, MATLAB 2019b is used. For the image quality analysis, we compare maximum error (E_{\max}), mean squared error (MSE), and the peak




Fig. 2 Reconstruction results of the conventional FBP using *fft* and *ifft* (left), using the second order optimal quadrature formula (middle), and using the third order optimal quadrature formula (right).




Fig. 3 Profiles of the reconstructed image of the Shepp-Logan: horizontal (left) and vertical (right).

signal-to-noise ratio (PSNR):

$$E_{\max}(I) = \max_{i,j} |I(i,j) - I_{ref}(i,j)|,$$

$$\text{MSE}(I) = \frac{1}{mn} \sum_{i=1}^m \sum_{j=1}^n |I(i,j) - I_{ref}(i,j)|^2,$$

$$\text{PSNR}(I) = 10 \log_{10} \left(\frac{I_{\max}^2}{\text{MSE}(I)} \right),$$

where I_{\max} is the maximum pixel value of the image I . Images of original simulated phantoms are denoted by I_{ref} (Fig. 1).




Fig. 4 Noise test: reconstruction results of the conventional FBP using *fft* and *ifft* (left), using the second order optimal quadrature formula (middle), and using the third order optimal quadrature formula (right).




Fig. 5 Noise test: Profiles of the reconstructed image of the Shepp-Logan: horizontal (left) and vertical (right).

	Shepp-Logan without noise			Shepp-Logan with noise		
	conv-FBP <i>fft</i> and <i>ifft</i>	OQF $m = 2$	OQF $m = 3$	conv-FBP <i>fft</i> and <i>ifft</i>	OQF $m = 2$	OQF $m = 3$
E_{\max}	0.3458	0.3526	0.3307	0.3722	0.3634	0.3472
MSE	7.9648e-04	7.2111e-04	6.5084e-04	7.9088e-04	7.4509e-04	6.4990e-04
PSNR	30.9883	31.4200	31.8652	31.0189	31.2779	31.8715

Table 1 Quantitative analysis for the reconstructed CT image from FBP using conventional *fft-ifft* and the second and third order optimal quadrature formulas in Corollaries 1 and 2.

Figs. 2 and 3 show the results of CT image reconstruction using optimal quadrature formulas of the second and third orders and profiles of the results, respectively. As shown in Fig. 3, the profile lines of conv-FBP and optimal quadrature formula almost coincide. Figs. 4 and 5 show the results of noise test and we observe that the resulting reconstructed images are very similar, visually. Table 1 shows E_{\max} , MSE, and PSNR for the reconstruction results using the second and third order optimal quadrature formulas. The second order optimal quadrature formula performs similar to or slightly better than the conventional FBP. The proposed third order optimal quadrature formula produces more improved quality than the conventional FBP particularly in terms of MSE and PSNR. The third order optimal quadrature formula also shows the best performance when noise was added.

Remark 3. The computational complexity of optimal quadrature formulas for Fourier integration is $\mathcal{O}(n^2)$ for the $n \times n$ matrix multiplication and addition and it is larger than $\mathcal{O}(n \log n)$ for the fast Fourier transformation. However, once the coefficients of optimal quadrature formulas for Fourier integral are computed, they can be used several times for the same size of input signals. Moreover, they can be well optimized using parallelization.

Remark 4. It should be noted that since the coefficients of the optimal quadrature formulas of the form (2.28) are continuous functions of the parameter ω , it is not needed any interpolation in Algorithm 1.

Acknowledgements

The work has been done while A.R. Hayotov was visiting Department of Mathematical Sciences at KAIST, Daejeon, Republic of Korea. A.R. Hayotov's work was supported by the 'Korea Foundation for Advanced Studies'/'Chey Institute for Advanced Studies' International Scholar Exchange Fellowship for academic year of 2018-2019. C.-O. Lee's work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. NRF-2017R1A2B4011627).

References

1. V. A. Avdeyenko and A. A. Malyukov. A quadrature formula for the Fourier integral based on the use of a cubic spline. *USSR Computational Mathematics and Mathematical Physics*, 29:101–103, 1989.
2. S. S. Babaev and A. R. Hayotov. Optimal interpolation formulas in the space $W_2^{(m,m-1)}$. *Calcolo*, 56:23, 2019.
3. I. Babuška. Optimal quadrature formulae. In *Doklady A. N.*, volume 149, pages 227–229. Russian Academy of Sciences, 1963.
4. I. Babuška, M. Práger, and E. Vitásek. *Numerical processes in differential equations*. Sntl Prague, 1966.
5. N. S. Bakhvalov and L. G. Vasil'eva. Evaluation of the integrals of oscillating functions by interpolation at nodes of Gaussian quadratures. *USSR Computational Mathematics and Mathematical Physics*, 8:241–249, 1968.
6. N. D. Boltaev, A. R. Hayotov, G. Milovanović, and K. M. Shadimetov. Optimal quadrature formulas for fourier coefficients in $W_2^{(m,m-1)}$ space. *Journal of Applied Analysis and Computation*, 7:1233–1266, 2017.
7. N. D. Boltaev, A. R. Hayotov, and K. M. Shadimetov. Construction of optimal quadrature formula for numerical calculation of Fourier coefficients in Sobolev space $L_2^{(1)}$. *American Journal of Numerical Analysis*, 4:1–7, 2016.

8. N. D. Boltaev, A. R. Hayotov, and K. M. Shadimetov. Construction of optimal quadrature formulas for Fourier coefficients in Sobolev space $L_2^{(m)}(0, 1)$. *Numerical Algorithms*, 74:307–336, 2017.
9. T. M. Buzug and D. Mihailidis. Computed tomography from photon statistics to modern cone beam CT. *Medical Physics*, 36:3858–3858, 2009.
10. A. Cabada, A. R. Hayotov, and K. M. Shadimetov. Construction of D^m -splines in $L_2^{(m)}(0, 1)$ space by Sobolev method. *Applied Mathematics and Computation*, 244:542–551, 2014.
11. T. Catinaş and G. Coman. Optimal quadrature formulas based on the ϕ -function method. *Studia. Universitatis Babeş-Bolyai. Mathematica*, 51:49–64, 2006.
12. G. Coman. Monosplines and optimal quadrature formulae in L_p . *Rendiconti di Matematica e delle sue Applicazioni*, 5:567–577, 1972.
13. G. Coman. Quadrature formulas of Sard type (Romanian). *Studia. Universitatis Babeş-Bolyai. Mathematica*, 17:73–77, 1972.
14. A. Deaño, D. Huybrechs, and A. Iserles. *Computing Highly Oscillatory Integrals*. Other Titles in Applied Mathematics. Society for Industrial and Applied Mathematics, Philadelphia, 2018.
15. T. G. Feeman. *The mathematics of medical imaging*. Springer Undergraduate Texts in Mathematics and Technology. Springer, Switzerland, 2015.
16. L. N. G. Filon. Iii.on a quadrature formula for trigonometric integrals. *Proceedings of the Royal Society of Edinburgh*, 49:38–47, 1930.
17. G. Frobenius. *Über die Bernoullischen Zahlen und die Eulerschen polynome*. In: *Gesammelte Abhandlungen*, Vol. III. J-P. Serre, editor. Berlin-Heidelberg-New York: Springer, 1968.
18. A. Ghizzetti and A. Ossicini. *Quadrature formulae*. International Series of Numerical Mathematics. Birkhäuser, Basel, 1970.
19. R. Hamming. *Numerical Methods for Scientists and Engineers, Second Edition*. Dover Books on Mathematics. Dover Publications, New York: McGraw Bill Book Company, Inc., USA, 1973.
20. A. R. Hayotov, S. Jeon, and C.-O. Lee. On an optimal quadrature formula for approximation of fourier integrals in the space $L_2^{(1)}$. *Journal of Computational and Applied Mathematics*, 372:112713, 2020.
21. A. Iserles and S. P. Nørsett. Efficient quadrature of highly oscillatory integrals using derivatives. *Proceedings of the Royal Society A*, 461:1383–1399, 2005.
22. A. C. Kak and M. Slaney. *Principles of Computerized Tomographic Imaging*. IEEE, New York, 188.
23. P. Köhler. On the weights of Sard’s quadrature formulas. *Calcolo*, 25:169–186, 1988.
24. F. Lanzara. On optimal quadrature formulae. *Journal of Inequalities and Applications*, 5:201–225, 2000.
25. A. A. Maljukov and I. I. Orlov. Construction of coefficients of the best quadrature formula for the class $W_{L_2}^{(2)}(M; ON)$ with equally spaced nodes. *Optimization Methods and Operations Research, Applied Mathematics*, 191:174–177, 1976.
26. G. V. Milovanović. Numerical calculation of integrals involving oscillatory and singular kernels and some applications of quadratures. *Computers & Mathematics with Applications*, 36:19–39, 1998.
27. G. V. Milovanović and M. P. Stanić. Numerical integration of highly oscillating functions. In *Analytic Number Theory, Approximation Theory, and Special Functions*, pages 613–649. Springer, New York, 2014.
28. F. Natterer. *The Mathematics of Computerized Tomography*. Society for Industrial and Applied Mathematics, Philadelphia, 2001
29. E. Novak, M. Ullrich, and H. Woźniakowski. Complexity of oscillatory integration for univariate Sobolev spaces. *Journal of Complexity*, 31:15–41, 2015.
30. S. Olver. *Numerical approximation of highly oscillatory integrals*. PhD thesis, University of Cambridge, 2008.
31. A. Rieder and A. Faridani. The semidiscrete filtered backprojection algorithm is optimal for tomographic inversion. *SIAM Journal on Numerical Analysis*, 41:869–892, 2003.
32. A. Sard. Best approximate integration formulas; best approximation formulas. *American Journal of Mathematics*, 71:80–91, 1949.
33. K. M. Shadimetov. Optimal formulas of approximate integration for differentiable functions. *Candidate dissertation, Novosibirsk*, 140, 1983.

34. K. M. Shadimetov. Optimal quadrature formulas in $L_2^m(\omega)$ and $L_2^m(r^1)$. In *Dokl. Akad. Nauk UzSSR*, volume 3, pages 5–8, 1983.
35. K. M. Shadimetov. The discrete analogue of the differential operator d^{2m}/dx^{2m} and its construction. *Questions of Computations and Applied Mathematics, Tashkent*, pages 22–35, 1985.
36. K. M. Shadimetov. Weight optimal cubature formulas in Sobolev's periodic space. *Sibirskii Zhurnal Vychislitel'noi Matematiki*, 2:185–196, 1999.
37. K. M. Shadimetov and A. R. Hayotov. Optimal quadrature formulas with positive coefficients in $L_2^{(m)}(0, 1)$ space. *Journal of Computational and Applied Mathematics*, 235(5):1114–1128, 2011.
38. K. M. Shadimetov and A. R. Hayotov. Construction of interpolation splines minimizing semi-norm in $W_2^{(m, m-1)}(0, 1)$ space. *BIT Numerical Mathematics*, 53(2):545–563, 2013.
39. K. M. Shadimetov and A. R. Hayotov. Optimal quadrature formulas in the sense of Sard in $W_2^{(m, m-1)}(0, 1)$ space. *Calcolo*, 51(2):211–243, 2014.
40. K. M. Shadimetov, A. R. Hayotov, and D. M. Akhmedov. Optimal quadrature formulas for Cauchy type singular integrals in Sobolev space. *Applied Mathematics and Computation*, 263:302–314, 2015.
41. K. M. Shadimetov, A. R. Hayotov, and F. A. Nuraliev. On an optimal quadrature formula in Sobolev space $L_2^{(m)}(0, 1)$. *Journal of Computational and Applied Mathematics*, 243:91–112, 2013.
42. K. M. Shadimetov, A. R. Hayotov, and F. A. Nuraliev. Optimal quadrature formulas of Euler–Maclaurin type. *Applied Mathematics and Computation*, 276:340–355, 2016.
43. S. L. Sobolev. Introduction to the theory of cubature formulas, 1974.
44. S. L. Sobolev. The coefficients of optimal quadrature formulas. In *Selected works of S. L. Sobolev*, pages 561–565. Springer, 2006.
45. S. L. Sobolev and V. L. Vaskevich. The theory of cubature formulas. *Kluwer Academic Publishers Group, Dordrecht*, 1997.
46. Z. Xu, G. V. Milovanović, and S. Xiang. Efficient computation of highly oscillatory integrals with Hankel kernel. *Applied Mathematics and Computation*, 261:312–322, 2015.
47. S. Zhang and E. Novak. Optimal quadrature formulas for the Sobolev space H^1 . *Journal of Scientific Computing*, 78:274–289, 2019.