

CONVERGENCE OF LARGE ATOMIC CONGESTION GAMES

ROBERTO COMINETTI[‡], MARCO SCARSINI[¶], MARC SCHRÖDER^{*}, AND NICOLÁS STIER-MOSES[§]

ABSTRACT. We study the convergence of sequences of atomic unsplittable congestion games with an increasing number of players. We consider two situations. In the first setting, each player has a weight that tends to zero, in which case the mixed equilibria of the finite games converge to the set of Wardrop equilibria of the corresponding nonatomic limit game. In the second case, players have unit weights, but participate in the game with a probability that tends to zero. In this case, the mixed equilibria converge to the set of Wardrop equilibria of another nonatomic game with suitably defined costs, which can be seen as a Poisson game in the sense of Myerson (1998b). In both settings we show that the price of anarchy of the sequence of games converges to the price of anarchy of the nonatomic limit. Beyond the case of congestion games, we establish a general result on the convergence of large games with random players towards a Poisson game.

1. INTRODUCTION

Nonatomic congestion games were introduced by Wardrop (1952) as a model for traffic networks with many drivers, where each single player has a negligible impact on congestion. The model is stated directly in terms of continuous flows and is motivated as the limit case when the number of players grows to infinity while their weights tend to zero. Although such convergence is intuitive and expected, only some special cases have been formally addressed in the literature, mainly for atomic splittable games. Following the motivation of many of the real-world applications such as road traffic and telecommunications, it is important to consider the convergence of *unsplittable* routing games with finitely many players who must route a given load over a single path chosen deterministically or at random using a mixed strategy. We consider the more general class of *atomic unsplittable congestion games* (not necessarily routing games) and allow for heterogeneous players. The main question is whether Nash equilibria for these games converge towards a Wardrop equilibrium of a limiting nonatomic congestion game.

In a *weighted congestion game*, there is a finite number of players who are characterized by a type and a weight. The type determines the set of feasible strategies of a player and the weight determines the player's impact on the costs. Moreover, there is a finite set of resources and each strategy corresponds to a subset of these resources. Players of a given type have the same set of available strategies. A strategy profile for all players induces a flow on each strategy equal to the total weight of players choosing it, as well as a load on each resource equal to the total weight of players using that resource as part of their strategy. The cost of using a resource is a weakly increasing function of its load. This defines a finite cost-minimization game. As shown by Rosenthal (1973), every congestion game in which all the players have the same weight admits a potential and has equilibria in pure strategies. In general, only mixed equilibria are guaranteed to exist (Nash, 1950).

As a special case of congestion games, a routing game features a finite directed network whose edges represent the resources. The origin-destination pairs encode the types, and the corresponding origin-destination paths provide the strategies. To illustrate, consider a routing game over a

simple network composed of two parallel edges with the same strictly-increasing cost function $c(\cdot)$, as shown in Fig. 1, and suppose that there are n players who need to choose an edge to route an identical weight of $w \equiv d/n$. Here, d denotes the total weight or demand. In a symmetric




Figure 1. Parallel edge network.

equilibrium every player randomizes by choosing each route with probability $1/2$. Consequently, the number of players on each route is distributed as a $\text{Binomial}(n, 1/2)$ random variable. The total load on each edge is therefore d/n times a $\text{Binomial}(n, 1/2)$, and converges in distribution to $d/2$, which is precisely the Wardrop equilibrium for a total demand of d units of flow. Notice that already in this simple example there is a multitude of other equilibria where n_1 and n_2 players, respectively, choose the upper and lower edges for sure (with $n_1, n_2 \leq n/2$), and the remaining players (if any) randomize appropriately so as to equalize the expected cost of both routes. As the number of players n tends to infinity, all these different equilibria converge to the unique Wardrop equilibrium with a $(d/2, d/2)$ -split of the flow. This includes the special case of a pure equilibrium in which half of the players (up to one unit if n is odd) take each route.

Angelidakis et al. (2013) and Cominetti et al. (2019) studied *Bernoulli congestion games* in which each player participates in the game with an independent probability. Motivated by this model, an alternative way of thinking about the limit of large games with an increasing number of players, is to consider the stochastic case in which each player has a unit weight but is present in the game with a small probability. Arguably, this might be a more natural way of modeling traffic. In fact, the contribution of an additional car to congestion is small but nonnegligible, while the congestion experienced by an agent depends on how many drivers are on the road at the same time. If we focus on a small interval around the time at which a player joins the game, the probability for the concurrent presence of other players emerges as a naturally small parameter. Taking the limit in this setting yields a different limit game in which the random loads on the edges converge to a family of Poisson random variables, and whose expected values can be again characterized as a Wardrop equilibrium of a suitably defined nonatomic game. Moreover, these limit loads can be interpreted as an equilibrium for a Poisson game in the sense of Myerson (1998b).

To provide some insight, consider again the parallel-edge example of Fig. 1, except that now each of the n players has a unit weight, but is present in the game with a small probability d/n . In this case the effective demand is a random variable $D^n \sim \text{Binomial}(n, d/n)$, which converges as $n \rightarrow \infty$ to a random variable $D \sim \text{Poisson}(d)$. Also, in a symmetric equilibrium where each player chooses an edge uniformly at random, the load on each route is distributed $\text{Binomial}(n, d/(2n))$ and converges to a $\text{Poisson}(d/2)$, whose expected value is again $d/2$. We will show that this convergence holds for asymmetric equilibria and general congestion games with an increasing number of heterogeneous players who are active with different vanishing probabilities.

Note that in both cases—vanishing weights and vanishing probabilities—the equilibria of the nonatomic limit games characterize the *expected values* of the random loads. However, in the Poisson regime the resource loads remain random in the limit, whereas in the Wardrop limit these random loads converge to a constant. Thus, the Poisson limit seems to provide a more accurate

description of the traffic flows observed in real networks, which exhibit stochastic variability. This is in agreement with some theoretical models—as well as empirical evidence based on traffic counts— which exhibit a Poissonian behavior at least under moderate congestion conditions.

1.1. Our contribution. After introducing the relevant classes of congestion games in [Section 2](#), [Section 3](#) shows that under very weak conditions any Nash-equilibrium loads in a sequence of weighted congestion games converge to a Wardrop equilibrium of a limiting nonatomic game. More precisely, if the number of players grows to infinity and their weights tend to zero in a way that the aggregate demands converge, then, for any sequence of mixed equilibria in the finite games, the random variables that represent the resource loads converge in distribution to a (deterministic) Wardrop equilibrium for a nonatomic limit game. We stress that players are not assumed to be symmetric and that they may have different weights and strategy sets. As long as their weights converge to zero, the random resource loads converge to some constants which are precisely a Wardrop equilibrium for the limit game. This provides stronger support to Wardrop’s model as a sensible approximation for large games with many small players.

[Section 4](#) deals with sequences of Bayesian Nash equilibria for Bernoulli congestion games when the participation probability of players tends to zero, and establishes their convergence to the set of Wardrop equilibria of another nonatomic game with suitably defined costs. This nonatomic game has a demand that is random and is distributed according to a Poisson random variable. In [Section 5](#) we show that the nonatomic limit games in the case of Bernoulli demands are related to Poisson games as introduced by [Myerson \(1998b\)](#). In fact, the Wardrop equilibria of the limit game are in one-to-one correspondence with the equilibria of an associated Poisson game. Poisson games were introduced axiomatically and not as a limit of a sequence of finite games. We close our paper by filling this gap, establishing a general result on convergence of sequences of games with Bernoulli demands (not necessarily congestion games) towards Poisson games. This constitutes a novel alternative justification for Wardrop’s model.

In both [Sections 3](#) and [4](#) we provide nonasymptotic estimates for the distance between the sequence of equilibria and their limits, and we establish the convergence of the price of anarchy, as a measure of the inefficiency of equilibria.

1.2. Related literature. Although the convergence of congestion games to nonatomic games has been considered in the past, most of the previous work assumed that players can split their weight over the available strategies. [Haurie and Marcotte \(1985\)](#) were the first to deal with issues of convergence for atomic splittable routing games, [Milchtaich \(2000\)](#) studied the convergence of replicas of crowding games, [Jacquot and Wan \(2018\)](#) considered splittable routing on parallel networks with heterogeneous players, and [Jacquot and Wan \(2019\)](#) studied the approximation of nonatomic aggregative games by a sequence of finite splittable games. Recently, [Chen et al. \(2014\)](#) defined altruism for nonatomic games by considering nonatomic games as the limit of sequences of atomic games. [Feldman et al. \(2016\)](#) used ideas of (λ, μ) -smoothness to study the convergence of the price of anarchy of atomic congestion games to the one of nonatomic games, when the number n of players increases and their weight is $1/n$, and they show a multiplicative error of $1+o(1/n)$. The relation between Nash equilibria and a Wardrop-like notion of equilibrium in aggregative games with finitely many players, was also studied recently by [Paccagnan et al. \(2019\)](#).

Several old papers considered the stochastic aspects of traffic, both theoretically and empirically, and the role of the Poisson distribution for modeling it (see, e.g., [Adams, 1936](#), [Mayne, 1954](#),

Oliver, 1961, Buckley, 1967, Miller, 1970). More recently, various authors have studied congestion games with stochastic features, focusing on the efficiency of equilibria under incomplete information. For instance, Gairing et al. (2008) studied the inefficiency of equilibria for congestion games in which the weight of a player is private information. Looking at the cost uncertainty, Nikolova and Stier-Moses (2014) and Piliouras et al. (2016) considered players' risk attitudes in the nonatomic and atomic cases, respectively. Roughgarden (2015) showed that whenever player types are independent, the inefficiency bounds for complete information games extend to Bayesian Nash equilibria of the incomplete information game. Wang et al. (2014) and Correa et al. (2019) looked at similar questions for nonatomic routing games. This trend does not only include congestion games: Stidham (2014) studied the efficiency of some classical queueing models on various networks, whereas Hassin et al. (2018) examined a queueing model with heterogeneous agents and studied how the inefficiency of equilibria varies with the intensity function.

Related to our results about convergence of Bernoulli games, Angelidakis et al. (2013) considered congestion games with stochastic players who are risk-averse, restricting their attention to the case of parallel edges. In the same spirit, Cominetti et al. (2019) studied Bernoulli congestion games, where each player i takes part in the game independently with probability p_i , and found sharp bounds for the price of anarchy as a function of the maximum p_i .

Games with random number of players were introduced by Myerson (1998a,b, 2000, 2002), with the main goal of analyzing elections with a large number of voters. In his seminal paper, Myerson (1998b) showed that the case where the number of players has a Poisson distribution is of particular relevance, as it is the only case where an environmental equivalence holds, i.e., the belief of a player of any type about the type profile of the other active players coincides with the belief of an external game theorist. Myerson (2000) dealt with large Poisson games in which the parameter of the Poisson distribution diverges. His approach differs from ours in the sense that he starts with a Poisson distribution, axiomatically justified, and lets the expectation of this distribution go to infinity. In our case we start with a finite number of players and let their number diverge in such a way that in the limit we get a Poisson distribution, but not necessarily with a large parameter. The asymptotic approach we take to Poisson games is based on results from Poisson approximation theory. In probability, articles about this topic abound. For an overview of this literature, we refer the reader to the books and surveys by Barbour et al. (1992), Barbour and Chen (2005), Novak (2019).

Several other papers have studied the properties of games with random number of players and Poisson games in particular. Among them, Milchtaich (2004) provides a sophisticated analysis of general games with a random number of players. De Sinopoli and Pimienta (2009), De Sinopoli et al. (2014), Meroni and Pimienta (2017) deal with various properties of equilibria in Poisson games, such as stability and existence of equilibria in undominated strategies. Other papers apply Poisson games in different settings, not necessarily related to elections. For instance, Lim and Matros (2009) study contests with finitely many players, where each player takes part in the game independently with the same probability, while Du and Gong (2016) use a Poisson game to model parking congestion and propose a decentralized and coordinated online parking mechanism to reduce congestion. Let us finally mention Kordonis and Papavassilopoulos (2015) and Bernhard and Deschamps (2017) who study dynamic games where players arrive at random over time.

2. CONGESTION GAMES

Informally, a *congestion game* is played over a set of resources whose costs depend on the mass of players using the resource. Each player chooses a subset of resources among the allowed subsets, seeking to achieve the minimum possible cost. Throughout this paper we consider a fixed finite set of resources $\mathcal{E}$, where each $e \in \mathcal{E}$ is associated with a weakly increasing continuous cost function $c_e : \mathbb{R}_+ \rightarrow \mathbb{R}_+$. We also fix a finite set of types $\mathcal{T}$ where each type $t \in \mathcal{T}$ is associated with a set of feasible actions $\mathcal{S}_t \subseteq 2^{\mathcal{E}}$, which describes the pure strategies.

To name a standard example of a congestion game, routing games capture the topology of a network structure. Games of this family are defined over the set of edges of a finite graph, encoded by $\mathcal{E}$, and the set of origin-destination (OD) pairs, encoded by $\mathcal{T}$. The set of actions $\mathcal{S}_t$ contains the feasible paths for the OD pair representing type t .

The structural objects

$$\mathcal{G} = (\mathcal{E}, (c_e)_{e \in \mathcal{E}}, \mathcal{T}, (\mathcal{S}_t)_{t \in \mathcal{T}}) \quad (2.1)$$

will be the same in all the congestion games considered hereafter, and the only differences will be in how we describe the set of players and their behavior. In the nonatomic framework, players are considered to be infinitesimally small and the model is stated in terms of the aggregate mass of players that use each strategy and resource. In contrast, for weighted congestion games, as well as for Bernoulli congestion games, we have a finite set of players who behave strategically. The rest of this section describes the three different models precisely.

As a guide for the notation used in the sequel, we write $\Delta(\mathcal{S}_t)$ for the simplex of all probability distributions over the strategy set $\mathcal{S}_t$. We use capital letters for random variables and lower case for their expected values. For instance, X_e will represent a random load on a resource $e \in \mathcal{E}$ with expected value $x_e = \mathbb{E}[X_e]$, and we will add a hat $\hat{x}_e$ when referring to an equilibrium.

2.1. Nonatomic congestion games. A *nonatomic congestion game* (NCG) is given by a pair $\Gamma^\infty = (\mathcal{G}, \mathbf{d})$, where $\mathcal{G}$ stands for the structural objects of the game as in Eq. (2.1), and $\mathbf{d} = (d_t)_{t \in \mathcal{T}}$ is a vector of demands with $d_t \geq 0$ representing the aggregate demand of type t . The *total demand* is given by the sum over all types $d_{\text{tot}} = \sum_{t \in \mathcal{T}} d_t$.

Each demand d_t is split over the corresponding strategies $\mathcal{S}_t$ and induces loads on the resources. Specifically, a *strategy flow* vector $\mathbf{y} := (y_{t,s})_{t \in \mathcal{T}, s \in \mathcal{S}_t}$ and a *resource load* vector $\mathbf{x} := (x_e)_{e \in \mathcal{E}}$ are called *feasible* if they satisfy the following constraints:

$$\forall t \in \mathcal{T}, \quad d_t = \sum_{s \in \mathcal{S}_t} y_{t,s}, \quad \text{and} \quad \forall e \in \mathcal{E}, \quad x_e = \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} y_{t,s} \mathbb{1}_{\{e \in s\}}. \quad (2.2)$$

The set of such *feasible flow-load pairs* $(\mathbf{y}, \mathbf{x})$ is denoted by $\mathcal{F}(\mathbf{d})$. Note that the resource loads are uniquely defined by the strategy flows, but not vice versa. Nevertheless, instead of only considering strategy flows, we will refer to flow-load pairs since some concepts are easier to express in terms of flows while others are defined in terms of loads. The notation and nomenclature is inspired by network games, and most of our examples will be of this type since they are intuitive and well-studied. However, we will use these terms in the more general setting of congestion games, even when there is no network structure of resources and strategies.

A *Wardrop equilibrium* (WE) is defined as a feasible flow-load pair $(\hat{\mathbf{y}}, \hat{\mathbf{x}}) \in \mathcal{F}(\mathbf{d})$ for which the prevailing cost of all used strategies is minimal, or, mathematically,

$$\forall t \in \mathcal{T}, \forall s, s' \in \mathcal{S}_t \quad \hat{y}_{t,s} > 0 \Rightarrow \sum_{e \in s} c_e(\hat{x}_e) \leq \sum_{e \in s'} c_e(\hat{x}_e). \quad (2.3)$$

Since, for each type, only strategies with minimum cost are used, all the used strategies have the same cost. As a consequence, any strategy flow decomposition of $\widehat{\mathbf{x}}$ yields an equilibrium. The set of Wardrop equilibria of the game Γ^∞ will be denoted by $\text{WE}(\Gamma^\infty)$.

2.2. Weighted congestion games. A *weighted congestion game* (WCG) is defined as a tuple $\Gamma_w = (\mathcal{G}, (w_i, t_i)_{i \in \mathcal{N}})$, where $\mathcal{N}$ is a finite set of players and each player $i \in \mathcal{N}$ has a weight $w_i \in \mathbb{R}_+$ and a type $t_i \in \mathcal{T}$ that determines her strategy set $\mathcal{S}_{t_i}$. Our use of the term “type” slightly differs from what is common in game theory, where a type is a random variable associated to each player, whose distribution is common knowledge but whose realization is private information. Here, the players’ types are deterministic and the type of each player is common knowledge.

The aggregate demand for each type t and the total demand are given by

$$d_t = \sum_{i: t_i=t} w_i, \quad \text{and} \quad d_{\text{tot}} = \sum_{t \in \mathcal{T}} d_t = \sum_{i \in \mathcal{N}} w_i. \quad (2.4)$$

Let $\sigma = (\sigma_i)_{i \in \mathcal{N}}$ be a *mixed strategy profile*, where $\sigma_i \in \Delta(\mathcal{S}_{t_i})$ represents the mixed strategy used by player $i \in \mathcal{N}$, and let $\Sigma := \times_{i \in \mathcal{N}} \Delta(\mathcal{S}_{t_i})$ be the set of mixed strategy profiles. Call E_σ the expectation with respect to the product probability measure $P_\sigma := \otimes_{i \in \mathcal{N}} \sigma_i$ over the set of pure strategy profiles $\mathcal{S} = \times_{i \in \mathcal{N}} \mathcal{S}_{t_i}$. If S_i is the random strategy of player i , whose distribution is σ_i , then the probability that player i uses a given resource e is

$$\sigma_{i,e} := P_\sigma(e \in S_i) = \sum_{s \in \mathcal{S}_{t_i}} \sigma_i(s) \mathbb{1}_{\{e \in s\}}. \quad (2.5)$$

Accordingly, the strategy flows $Y_{t,s}$ and the resource loads X_e become random variables

$$Y_{t,s} = \sum_{i: t_i=t} w_i \mathbb{1}_{\{S_i=s\}}, \quad \text{and} \quad X_e = \sum_{i \in \mathcal{N}} w_i \mathbb{1}_{\{e \in S_i\}}, \quad (2.6)$$

given by the random realizations of S_i . The expected values are

$$y_{t,s} := E_\sigma[Y_{t,s}] = \sum_{i: t_i=t} w_i \sigma_i(s), \quad \text{and} \quad x_e := E_\sigma[X_e] = \sum_{i \in \mathcal{N}} w_i \sigma_{i,e}. \quad (2.7)$$

A straightforward calculation shows that the pair $(\mathbf{y}, \mathbf{x})$ satisfies Eq. (2.2), so that $(\mathbf{y}, \mathbf{x}) \in \mathcal{F}(\mathbf{d})$.

To take the perspective of a fixed player i , we assume the player already selected resource e and define the conditional load

$$X_{i,e} := w_i + \sum_{j \neq i} w_j \mathbb{1}_{\{e \in S_j\}}. \quad (2.8)$$

Using this, the expected cost of player i conditional on the player using the resource e is

$$E_\sigma[c_e(X_e) \mid e \in S_i] = E_\sigma[c_e(X_{i,e})]. \quad (2.9)$$

A strategy profile $\widehat{\sigma} \in \Sigma$ is a *mixed Nash equilibrium* (MNE) if

$$\forall i \in \mathcal{N}, \forall s, s' \in \mathcal{S}_{t_i} \quad \widehat{\sigma}_i(s) > 0 \Rightarrow \sum_{e \in s} E_{\widehat{\sigma}}[c_e(X_{i,e})] \leq \sum_{e \in s'} E_{\widehat{\sigma}}[c_e(X_{i,e})]. \quad (2.10)$$

The set of mixed Nash equilibria of Γ_w is denoted by $\text{MNE}(\Gamma_w)$.

When all the players have the same weight $w_i \equiv w$, Rosenthal (1973) showed that Γ_w is a potential game and, as a consequence, pure Nash equilibria (PNE) are guaranteed to exist (see also Monderer and Shapley, 1996). Moreover, Fotakis et al. (2005) showed that every weighted congestion game with affine costs admits an exact potential. Conversely, Harks et al. (2011) proved

that if $\mathcal{C}$ is a class of cost functions such that every weighted congestion game with costs in $\mathcal{C}$ admits a potential, then $\mathcal{C}$ only contains affine functions. Existence of pure equilibria in weighted congestion games was further studied by [Harks and Klimm \(2012\)](#). Beyond these cases, one can only establish existence of equilibria in mixed strategies ([Nash \(1950\)](#)).

2.3. Bernoulli congestion games. In a weighted congestion game the randomness arises only from the players' mixed strategies. In this section, we add another stochastic element: players may not be present in the game. A *Bernoulli congestion game* (BCG) is a congestion game in which each player $i \in \mathcal{N}$ has a unit weight $w_i \equiv 1$, but takes part in the game only with some probability $r_i \in (0, 1)$ and otherwise remains inactive and incurs no cost. The participation events, assumed to be independent, are encoded in random variables $U_i \sim \text{Bernoulli}(r_i)$, which indicate whether player $i \in \mathcal{N}$ is active or not. A Bernoulli congestion game is denoted by $\Gamma_b = (\mathcal{G}, (r_i, t_i)_{i \in \mathcal{N}})$.

The framework is similar to a weighted congestion game in which the w_i 's are replaced by random weights $U_i \in \{0, 1\}$ with expected value r_i , so that the per-type demands become the random variables $D_t = \sum_{i:t_i=t} U_i$ with expected values $d_t = \mathbb{E}[D_t] = \sum_{i:t_i=t} r_i$. The formulas are therefore very similar, with w_i replaced by U_i , or by r_i when taking expectations. Nevertheless, we will see later on that the two classes of games behave differently in some respects.

Let $\sigma = (\sigma_i)_{i \in \mathcal{N}} \in \Sigma$ describe the profile of mixed strategies used by each player. We assume that each player chooses her mixed strategy before knowing whether she will participate. Now randomness is induced both by the random participation and by the mixed strategies. This is described by a discrete probability space $(\Omega, 2^\Omega, P_\sigma)$, where $\Omega = \{0, 1\}^\mathcal{N} \times \mathcal{S}$ with $\mathcal{S} = \times_{i \in \mathcal{N}} \mathcal{S}_i$ as before, and P_σ is now the probability measure induced by σ and by the random participation of players; that is, for $\mathbf{u} \in \{0, 1\}^\mathcal{N}$ and $\mathbf{s} \in \mathcal{S}$ we have

$$P_\sigma(\mathbf{u}, \mathbf{s}) = \prod_{i \in \mathcal{N}} P_i(u_i) \sigma_i(s_i), \quad (2.11)$$

with $P_i(1) = r_i$ and $P_i(0) = 1 - r_i$. The corresponding expectation operator will be denoted $\mathbb{E}_\sigma$.

As before, $\mathbb{1}_{\{e \in S_i\}}$ is a Bernoulli random variable indicating whether the random strategy S_i includes resource e , with the notation in [Eq. \(2.5\)](#) still in place. Additionally, let $U_{i,e} = U_i \mathbb{1}_{\{e \in S_i\}}$ indicate whether player i is active and chooses resource e , for which we have $\mathbb{E}_\sigma[U_{i,e}] = r_i \sigma_{i,e}$. Then, the total number of active players of type $t \in \mathcal{T}$ using strategy $\mathbf{s} \in \mathcal{S}_t$, and the total load on resource $e \in \mathcal{E}$, are now the random variables

$$Y_{t,s} = \sum_{i:t_i=t} U_i \mathbb{1}_{\{S_i=s\}}, \quad \text{and} \quad X_e = \sum_{i \in \mathcal{N}} U_{i,e}. \quad (2.12)$$

The expected strategy flows and resource loads are

$$y_{t,s} := \mathbb{E}_\sigma[Y_{t,s}] = \sum_{i:t_i=t} r_i \sigma_i(s), \quad \text{and} \quad x_e := \mathbb{E}_\sigma[X_e] = \sum_{i \in \mathcal{N}} r_i \sigma_{i,e}. \quad (2.13)$$

Once again the pair $(\mathbf{y}, \mathbf{x})$ satisfies [Eq. \(2.2\)](#), so that $(\mathbf{y}, \mathbf{x}) \in \mathcal{F}(\mathbf{d})$.

When $U_{i,e} = 1$, conditional on player i selecting resource $e \in \mathcal{E}$, its load is $X_{i,e} = 1 + Z_{i,e}$, where $Z_{i,e}$ represents the number of other players using that resource, that is

$$Z_{i,e} = \sum_{j \neq i} U_{j,e}. \quad (2.14)$$

Then the conditional expected cost for player $i \in \mathcal{N}$ when using this resource is

$$\mathbb{E}_\sigma[c_e(X_e) \mid U_{i,e} = 1] = \mathbb{E}_\sigma[c_e(X_{i,e})] = \mathbb{E}_\sigma[c_e(1 + Z_{i,e})]. \quad (2.15)$$

Notice that in this setting all players have unit weight so that the resource loads are integer-valued and therefore the costs $c_e : \mathbb{N} \rightarrow \mathbb{R}_+$ need only be defined over the integers.

A strategy profile $\hat{\sigma} \in \Sigma$ is a *Bayesian Nash equilibrium* (BNE) if

$$\forall i \in \mathcal{N}, \forall s, s' \in \mathcal{S}_{t_i} \quad \hat{\sigma}_i(s) > 0 \Rightarrow \sum_{e \in s} \mathbb{E}_{\hat{\sigma}}[c_e(1 + Z_{i,e})] \leq \sum_{e \in s'} \mathbb{E}_{\hat{\sigma}}[c_e(1 + Z_{i,e})]. \quad (2.16)$$

The set of all Bayesian Nash equilibria of Γ_b is denoted by $\text{BNE}(\Gamma_b)$.

Remark 2.1. [Cominetti et al. \(2019, Proposition 3.3\)](#) showed that Γ_b is a potential game which drives the class of BCGs apart from the class of WCGs, which admits a potential only in special cases. In particular, a BCG always has equilibria in pure strategies. However, here we will consider both pure and mixed equilibria.

2.4. Price of anarchy and price of stability. To understand how the additional variability induced by player participation affects congestion games, we would like to understand if the ensuing equilibria are closer to optimality than in other games. Previous research looked at deterministic atomic and nonatomic congestion games as defined in earlier sections. [Cominetti et al. \(2019\)](#) looked at the case of finite games with stochastic player participation. Here, we extend the analysis to look at the limits of games when the number of players increases.

For this analysis we need to be able to measure the social cost in the congestion games, which is defined as the sum of all costs. This provides us with a yardstick with which we can quantify the efficiency of equilibria as first proposed by [Koutsoupias and Papadimitriou \(1999\)](#). The *price of anarchy* (PoA) is the worst-case ratio of the social cost of the equilibrium to the optimum. Here, worst case is with respect to all possible equilibria. Instead, the *price of stability* (PoS) is defined accordingly with respect to the best equilibria.

Starting with NCGs, their *social cost* (SC) is given by

$$\forall (\mathbf{y}, \mathbf{x}) \in \mathcal{F}(\mathbf{d}) \quad \text{SC}(\mathbf{y}, \mathbf{x}) := \sum_{e \in \mathcal{E}} x_e c_e(x_e), \quad (2.17)$$

from where the social optimum is $\text{Opt}(\Gamma^\infty) := \min_{(\mathbf{y}, \mathbf{x}) \in \mathcal{F}(\mathbf{d})} \text{SC}(\mathbf{y}, \mathbf{x})$. It is well known (see [Beckmann et al., 1956](#)) that, whenever the cost functions c_e are weakly increasing, all Wardrop equilibria have the same social cost, so that defining $\text{Eq}(\Gamma^\infty) := \text{SC}(\hat{\mathbf{y}}, \hat{\mathbf{x}})$ for any $(\hat{\mathbf{y}}, \hat{\mathbf{x}}) \in \text{WE}(\Gamma^\infty)$, it follows that the PoA and PoS for a nonatomic game coincide and are given by

$$\text{PoA}(\Gamma^\infty) = \text{PoS}(\Gamma^\infty) := \frac{\text{Eq}(\Gamma^\infty)}{\text{Opt}(\Gamma^\infty)}. \quad (2.18)$$

The corresponding definitions for WCGs and BCGs are similar, adjusted for the fact that now these games include stochastic realizations. The *expected social cost* (ESC) is

$$\forall \sigma \in \Sigma \quad \text{ESC}(\sigma) := \mathbb{E}_\sigma \left[\sum_{e \in \mathcal{E}} X_e c_e(X_e) \right], \quad (2.19)$$

where $(X_e)_{e \in \mathcal{E}}$ are the random resource loads induced by the mixed strategy profile σ . The optimum cost is $\text{Opt}(\Gamma) := \min_{\sigma \in \Sigma} \text{ESC}(\sigma)$, and the social optimum is $\tilde{\sigma} \in \arg \min_{\sigma \in \Sigma} \text{ESC}(\sigma)$. Considering the best and worst social cost at equilibrium, the price of anarchy and stability are:

$$\text{PoA}(\Gamma) := \min_{\hat{\sigma} \in \text{MNE}(\Gamma)} \frac{\text{ESC}(\hat{\sigma})}{\text{Opt}(\Gamma)}, \quad \text{and} \quad \text{PoS}(\Gamma) := \max_{\hat{\sigma} \in \text{MNE}(\Gamma)} \frac{\text{ESC}(\hat{\sigma})}{\text{Opt}(\Gamma)}. \quad (2.20)$$

3. CONVERGENCE OF WEIGHTED CONGESTION GAMES

Now that we laid out the games we are considering, we study the convergence of WCGs to NCGs when the number of players increases and their weights decrease. The only stochastic element appearing in this section is the fact that players randomize by considering mixed strategies. Under mild and natural conditions, we prove that equilibria converge as there are more players in the game, and that the optimal social cost, as well as the PoA, converge to the corresponding values for the limit nonatomic game. We also compute the speed of convergence for these limits.

3.1. Sequences of weighted congestion games. Consider a sequence of WCGs

$$\Gamma_{\mathcal{W}}^n = (\mathcal{G}, (w_i^n, t_i^n)_{i \in \mathcal{N}^n}). \quad (3.1)$$

The elements that vary over the sequence are the set $\mathcal{N}^n$ of players, their weights w_i^n , and their types t_i^n . We want to study if and how the equilibria for this sequence converge. All the notations in Section 2.2 will remain in place, by simply adding the superscript n .

We assume that the number of players goes to infinity, while the sequence of weights goes to zero in such a way that the aggregate demand for each type converges. In other words, as there are more players, no player becomes dominant and the demands remain bounded. This is captured by the following asymptotic behavior as $n \rightarrow \infty$:

$$|\mathcal{N}^n| \rightarrow \infty, \quad (3.2a)$$

$$w^n := \max_{i \in \mathcal{N}^n} w_i^n \rightarrow 0, \quad (3.2b)$$

$$d_t^n := \sum_{i: t_i^n = t} w_i^n \rightarrow d_t. \quad (3.2c)$$

Under these conditions, the random loads at equilibrium for the sequence of games $\Gamma_{\mathcal{W}}^n$ converge to the loads of Wardrop equilibria of a corresponding nonatomic limit game.

Theorem 3.1. *Let $\Gamma_{\mathcal{W}}^n$ be a sequence of WCGs satisfying the assumptions in Eq. (3.2), and let $\hat{\sigma}^n \in \text{MNE}(\Gamma_{\mathcal{W}}^n)$ be an arbitrary sequence of mixed Nash equilibria. Then the corresponding sequence of expected flow-load pairs $(\hat{\mathbf{y}}^n, \hat{\mathbf{x}}^n)$ is bounded and every accumulation point $(\hat{\mathbf{y}}, \hat{\mathbf{x}})$ is a Wardrop equilibrium of the NCG $\Gamma^\infty = (\mathcal{G}, (d_t)_{t \in \mathcal{T}})$. Furthermore, along every convergent subsequence, the random flow-load pairs (Y^n, X^n) converge in distribution to $(\hat{\mathbf{y}}, \hat{\mathbf{x}})$.*

Proof. As for every mixed strategy profile, the expected flow-load pair $(\hat{\mathbf{y}}^n, \hat{\mathbf{x}}^n)$ belongs to $\mathcal{F}(\mathbf{d}^n)$ so that we have Eq. (2.2), adding the superindex n to all the terms involved. Since $d_t^n \rightarrow d_t$, it follows from Eq. (2.2) that the sequence $(\hat{\mathbf{y}}^n, \hat{\mathbf{x}}^n)$ is bounded, and then passing to the limit in these equations we conclude that every accumulation point $(\hat{\mathbf{y}}, \hat{\mathbf{x}})$ belongs to $\mathcal{F}(\mathbf{d})$.

For simplicity of notation, take a convergent subsequence and rename it so that the full sequence converges $(\hat{\mathbf{y}}^n, \hat{\mathbf{x}}^n) \rightarrow (\hat{\mathbf{y}}, \hat{\mathbf{x}})$. From Eq. (2.6) we get

$$\text{Var}_{\hat{\sigma}^n}[X_e^n] = \text{Var}_{\hat{\sigma}^n} \left[\sum_{i \in \mathcal{N}^n} w_i^n \mathbb{1}_{\{e \in S_i^n\}} \right] = \sum_{i \in \mathcal{N}^n} (w_i^n)^2 \hat{\sigma}_{i,e}^n (1 - \hat{\sigma}_{i,e}^n) \leq \sum_{i \in \mathcal{N}^n} (w_i^n)^2. \quad (3.3)$$

It can be shown similarly that

$$\text{Var}_{\hat{\sigma}^n}[Y_{t,s}^n] \leq \sum_{i \in \mathcal{N}^n} (w_i^n)^2. \quad (3.4)$$

Eq. (3.2) in turn implies

$$\sum_{i \in \mathcal{N}^n} (w_i^n)^2 \leq w^n \sum_{i \in \mathcal{N}^n} w_i^n = w^n \sum_{t \in \mathcal{T}} d_t^n \rightarrow 0, \quad (3.5)$$

so that $\text{Var}_{\hat{\sigma}^n}[Y_{t,s}^n] \rightarrow 0$ and $\text{Var}_{\hat{\sigma}^n}[X_e^n] \rightarrow 0$, from which convergence in distribution follows.

It remains to show that $(\hat{\mathbf{y}}, \hat{\mathbf{x}})$ is a WE, that is, we need to establish Eq. (2.3). If $\hat{y}_{t,s} > 0$, then, for n large enough, we have $\hat{y}_{t,s}^n > 0$ and there exists some player $i \in \mathcal{N}^n$ of type $t_i^n = t$ with $\hat{\sigma}_i^n(s) > 0$. Note that i actually depends on n , so we should write i^n . We omit the superscript for the sake of simplicity. The equilibrium condition in $\Gamma_{\mathbf{w}}^n$ implies that, for each alternative strategy $s' \in \mathcal{S}_t$ and for this player i ,

$$\sum_{e \in s} \mathbb{E}_{\hat{\sigma}^n}[c_e(X_{i,e}^n)] \leq \sum_{e \in s'} \mathbb{E}_{\hat{\sigma}^n}[c_e(X_{i,e}^n)]. \quad (3.6)$$

Since $|X_{i,e}^n - X_e^n| \leq w_i^n \leq w^n \rightarrow 0$, it follows that $X_{i,e}^n \rightarrow \hat{x}_e$ in distribution. Moreover, the loads $X_e^n \geq 0$ are bounded above by the total demands $d_{\text{tot}}^n = \sum_{t \in \mathcal{T}} d_t^n$, which converge to $d_{\text{tot}} = \sum_{t \in \mathcal{T}} d_t$, so that both X_e^n and $X_{i,e}^n$ are uniformly bounded. It follows that $\mathbb{E}_{\hat{\sigma}^n}[c_e(X_{i,e}^n)] \rightarrow c_e(\hat{x}_e)$ and, letting $n \rightarrow \infty$ in Eq. (3.6), we obtain Eq. (2.3) as required. $\square$

For games with strictly-increasing cost functions $c_e(\cdot)$ the equilibrium loads $\hat{x}_e$ are unique, which yields the following direct consequence.

Corollary 3.2. *Suppose that the resource costs $c_e(\cdot)$ are strictly increasing. Then, for every sequence $\Gamma_{\mathbf{w}}^n$ of WCGs satisfying Eq. (2.2) and each $\hat{\sigma}^n \in \text{MNE}(\Gamma_{\mathbf{w}}^n)$, the random loads X_e^n converge in distribution to the unique Wardrop equilibrium loads $\hat{x}_e$ of the nonatomic limit game $\Gamma^\infty = (\mathcal{G}, (d_t)_{t \in \mathcal{T}})$.*

Example 3.1. We illustrate the previous result on the Wheatstone network shown in Fig. 2. There is a single OD pair and $n \geq 2$ identical players, each one with weight $w_i \equiv 1/n$ so that the total demand is $d_{\text{tot}} = 1$. For each n , there is a unique symmetric pure Nash equilibrium in which




Figure 2. Wheatstone network.

all players take the zig-zag path e_1, e_3, e_5 and pay a cost equal to 2. For $n = 2$, we also have a pure equilibrium in which one player takes the upper path while the other takes the lower path, and they both pay 3/2. For $n \geq 3$, the only mixed equilibrium (modulo permutation of the players) is when $n - 1$ players take the zig-zag path and the last player mixes in any possible way over the three paths. In the limit when $n \rightarrow \infty$, all these equilibria converge to the WE of the corresponding nonatomic game with a unit flow over the zig-zag path.

3.2. Rate of convergence. The previous results showed that the random loads X_e^n in the finite games converge in distribution to the WE loads $\widehat{x}_e$. Under mild additional assumptions, which call for a lower and upper bounds on the derivative of the cost functions, we can find nonasymptotic estimates for the speed of convergence.

We start by establishing some estimates for the distance between Wardrop equilibria in nonatomic games with different demands, as well as between exact and approximate equilibria. We recall that an ε -approximate Wardrop equilibrium $(\mathbf{y}, \mathbf{x}) \in \mathcal{F}(\mathbf{d})$, or ε -WE, is defined exactly like a WE but with an additive constant:

$$\forall t \in \mathcal{T}, \forall s, s' \in \mathcal{S}_t \quad y_{t,s} > 0 \Rightarrow \sum_{e \in s} c_e(x_e) \leq \sum_{e \in s'} c_e(x_e) + \varepsilon. \quad (3.7)$$

Proposition 3.3. *Let $(\widehat{\mathbf{y}}, \widehat{\mathbf{x}})$ be a WE for a nonatomic congestion game Γ^∞ with $d_{\text{tot}} \leq \alpha$. Suppose that there exists $\beta > 0$ such that $c'_e(x) \geq \beta$ for all $e \in \mathcal{E}$ and $x \in [0, \alpha]$, and let $\Xi = \sqrt{2C/\beta}$ with $C = \max_s \sum_{e \in s} c_e(\alpha)$, where the maximum is taken over all the feasible strategies $s \in \cup_{t \in \mathcal{T}} \mathcal{S}_t$.*

- (a) *If $(\mathbf{y}, \mathbf{x}) \in \mathcal{F}(\mathbf{d})$ is an ε -approximate Wardrop equilibrium, then $\|\mathbf{x} - \widehat{\mathbf{x}}\|_2 \leq \sqrt{\varepsilon\alpha/\beta}$.*
- (b) *If $(\widehat{\mathbf{y}}', \widehat{\mathbf{x}}')$ is a WE for perturbed demands $\mathbf{d}'$ with $d'_{\text{tot}} \leq \alpha$, then $\|\widehat{\mathbf{x}}' - \widehat{\mathbf{x}}\|_2 \leq \Xi \cdot \sqrt{\|\mathbf{d}' - \mathbf{d}\|_1}$.*

Proof. (a) Defining $C_s := \sum_{e \in s} c_e(x_e)$ and $\eta_t := \min_{s \in \mathcal{S}_t} C_s$, we bound the squared-distance by

$$\begin{aligned} \sum_{e \in \mathcal{E}} \beta (x_e - \widehat{x}_e)^2 &\leq \sum_{e \in \mathcal{E}} (c_e(x_e) - c_e(\widehat{x}_e))(x_e - \widehat{x}_e) \\ &\leq \sum_{e \in \mathcal{E}} c_e(x_e)(x_e - \widehat{x}_e) \\ &= \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} \left(\sum_{e \in s} c_e(x_e) \right) (y_{t,s} - \widehat{y}_{t,s}) \\ &= \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} (C_s - \eta_t)(y_{t,s} - \widehat{y}_{t,s}) \\ &\leq \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} (C_s - \eta_t) y_{t,s}, \end{aligned}$$

where in the second inequality we dropped the sum $\sum_{e \in \mathcal{E}} c_e(\widehat{x}_e)(x_e - \widehat{x}_e)$, which is nonnegative since $(\widehat{\mathbf{y}}, \widehat{\mathbf{x}})$ is a WE for Γ^∞ and $(\mathbf{y}, \mathbf{x}) \in \mathcal{F}(\mathbf{d})$, while the subsequent equalities follow by expressing the resource loads in terms of the strategy flows and exchanging the order of summation, and then using the fact that both $(\widehat{\mathbf{y}}, \widehat{\mathbf{x}})$ and $(\mathbf{y}, \mathbf{x})$ are feasible. Now, using Eq. (3.7), we conclude

$$\beta \|\mathbf{x} - \widehat{\mathbf{x}}\|_2^2 \leq \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} \varepsilon y_{t,s} = \varepsilon \sum_{t \in \mathcal{T}} d_t = \varepsilon d_{\text{tot}} \leq \varepsilon \alpha.$$

(b) Proceeding as in the previous part, we have

$$\begin{aligned} \beta \|\widehat{\mathbf{x}}' - \widehat{\mathbf{x}}\|_2^2 &\leq \sum_{e \in \mathcal{E}} (c_e(\widehat{x}'_e) - c_e(\widehat{x}_e))(\widehat{x}'_e - \widehat{x}_e) \\ &= \sum_{e \in \mathcal{E}} c_e(\widehat{x}'_e)(\widehat{x}'_e - \widehat{x}_e) + \sum_{e \in \mathcal{E}} c_e(\widehat{x}_e)(\widehat{x}_e - \widehat{x}'_e). \end{aligned}$$

Let the two sums in the RHS be denoted by $\Psi(\widehat{\mathbf{x}}', \widehat{\mathbf{x}})$ and $\Psi(\widehat{\mathbf{x}}, \widehat{\mathbf{x}}')$, respectively. To bound $\Psi(\widehat{\mathbf{x}}', \widehat{\mathbf{x}})$, we exploit the fact that $\widehat{\mathbf{x}}'$ is a WE for the demand $\mathbf{d}'$. Since $(\widehat{\mathbf{y}}, \widehat{\mathbf{x}})$ is feasible for $\mathbf{d}$, but not for $\mathbf{d}'$,

for each type t with $d_t > 0$ we consider the rescaled flows $y_{t,s} = \widehat{y}_{t,s} d'_t / d_t$, whereas when $d_t = 0$ we simply take $y_{t,s} = \widehat{y}_{t,s}$. Letting $x_e = \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} y_{t,s} \mathbb{1}_{\{e \in s\}}$ denote the corresponding resource loads, we have that $(\mathbf{y}, \mathbf{x}) \in \mathcal{F}(\mathbf{d}')$, and therefore

$$\begin{aligned} \Psi(\widehat{\mathbf{x}}', \widehat{\mathbf{x}}) &= \sum_{e \in \mathcal{E}} c_e(\widehat{x}'_e)(\widehat{x}'_e - x_e) + \sum_{e \in \mathcal{E}} c_e(\widehat{x}'_e)(x_e - \widehat{x}_e) \\ &\leq \sum_{e \in \mathcal{E}} c_e(\widehat{x}'_e)(x_e - \widehat{x}_e) \\ &= \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} \left(\sum_{e \in s} c_e(\widehat{x}'_e) \right) (y_{t,s} - \widehat{y}_{t,s}) \end{aligned}$$

where in the inequality we dropped the first sum which is nonpositive since $(\widehat{\mathbf{y}}', \widehat{\mathbf{x}}')$ is a WE and $(\mathbf{y}, \mathbf{x})$ is feasible for $\mathbf{d}'$, whereas the last equality follows by expressing the resource loads in terms of the strategy flows and exchanging the order of the sums.

We now analyze each term in the outer sum over $t \in \mathcal{T}$. When $d_t > 0$ the inner double sum can be bounded as

$$\begin{aligned} \sum_{s \in \mathcal{S}_t} \left(\sum_{e \in s} c_e(\widehat{x}'_e) \right) (y_{t,s} - \widehat{y}_{t,s}) &= \sum_{s \in \mathcal{S}_t} \left(\sum_{e \in s} c_e(\widehat{x}'_e) \right) \widehat{y}_{t,s} (d'_t / d_t - 1) \\ &\leq \sum_{s \in \mathcal{S}_t} C \widehat{y}_{t,s} |d'_t / d_t - 1| \\ &= C |d'_t - d_t|, \end{aligned}$$

whereas, when $d_t = 0$, we have $y_{t,s} = \widehat{y}_{t,s}$ and $\widehat{y}_{t,s} = 0$ so that

$$\sum_{s \in \mathcal{S}_t} \left(\sum_{e \in s} c_e(\widehat{x}'_e) \right) (y_{t,s} - \widehat{y}_{t,s}) = \sum_{s \in \mathcal{S}_t} \left(\sum_{e \in s} c_e(\widehat{x}'_e) \right) \widehat{y}_{t,s} \leq C d'_t = C |d'_t - d_t|. \quad (3.8)$$

Summing these estimates over all $t \in \mathcal{T}$ we get $\Psi(\widehat{\mathbf{x}}', \widehat{\mathbf{x}}) \leq C \|\mathbf{d}' - \mathbf{d}\|_1$. Symmetrically, we have $\Psi(\widehat{\mathbf{x}}, \widehat{\mathbf{x}}') \leq C \|\mathbf{d} - \mathbf{d}'\|_1$, from which the result follows. $\square$

We now prove that any mixed Nash equilibrium for a weighted congestion game Γ_w is close to a Wardrop equilibrium in the nonatomic game Γ^∞ with the same aggregate demands. The distance depends—in a way that the theorem makes precise—on the topology of the instance, on the cost functions, and on the magnitude of the weights.

Theorem 3.4. *Let Γ_w be a WCG with aggregate demands $(d_t)_{t \in \mathcal{T}}$ given by Eq. (2.4) and $d_{\text{tot}} \leq \alpha$. Suppose that $\beta \leq c'_e(x) \leq \zeta$ and $|c''_e(x)| \leq \gamma$ for all $x \in [0, \alpha]$, with suitable constants $\beta > 0$, ζ , and γ . Define $\Theta = \sqrt{\alpha/4} + \sqrt{2\alpha\kappa(\zeta + \gamma\alpha/4)/\beta}$ with κ the cardinality of the largest feasible strategy $s \in \cup_{t \in \mathcal{T}} \mathcal{S}_t$. Let X_e be the random loads in a mixed Nash equilibrium $\widehat{\sigma} \in \text{MNE}(\Gamma_w)$, and let $\widehat{x}_e$ be the unique resource loads in the Wardrop equilibrium for the nonatomic game $\Gamma^\infty = (\mathcal{G}, (d_t)_{t \in \mathcal{T}})$ with the same aggregate demands as Γ_w . Then, letting $w = \max_{i \in \mathcal{N}} w_i$, we have*

$$\|X_e - \widehat{x}_e\|_{L^2} \leq \Theta \cdot \sqrt{w}. \quad (3.9)$$

Proof. Let $x_e = \mathbb{E}_{\widehat{\sigma}}[X_e]$ so that

$$\|X_e - \widehat{x}_e\|_{L^2} \leq \|X_e - x_e\|_{L^2} + |x_e - \widehat{x}_e|, \quad (3.10)$$

with

$$\|X_e - x_e\|_{L^2}^2 = \text{Var}_{\widehat{\sigma}}(X_e) = \sum_{i \in \mathcal{N}} w_i^2 \sigma_{i,e} (1 - \sigma_{i,e}) \leq \frac{1}{4} \sum_{i \in \mathcal{N}} w_i^2 \leq \frac{w d_{\text{tot}}}{4} \leq \frac{\alpha w}{4}. \quad (3.11)$$

With this bound in place, we proceed to estimate $|x_e - \widehat{x}_e|$ by showing that $(\mathbf{y}, \mathbf{x})$ is an ε -WE with

$$\varepsilon = 2\kappa(\zeta + \gamma\alpha/4)w. \quad (3.12)$$

To this end, we first observe that a second order expansion of the costs $c_e(\cdot)$ allows us to get the maximum bias of the cost:

$$-\gamma(X_e - x_e)^2 \leq c_e(X_e) - c_e(x_e) - c'_e(x_e)(X_e - x_e) \leq \gamma(X_e - x_e)^2, \quad (3.13)$$

so that taking expectations, and noting that the expected value of the linear part vanishes, we get

$$|\mathbb{E}_{\widehat{\sigma}}[c_e(X_e)] - c_e(x_e)| \leq \gamma \text{Var}_{\widehat{\sigma}}(X_e) \leq \gamma\alpha w/4. \quad (3.14)$$

Now, if $y_{t,s} > 0$, then there exists some player $i \in \mathcal{N}$ with $t_i = t$ and $\widehat{\sigma}_i(s) > 0$, so that the equilibrium condition in Γ_w implies that for each alternative strategy $s' \in \mathcal{S}_t$ we have

$$\sum_{e \in s} \mathbb{E}_{\widehat{\sigma}}[c_e(X_{i,e})] \leq \sum_{e \in s'} \mathbb{E}_{\widehat{\sigma}}[c_e(X_{i,e})]. \quad (3.15)$$

Now, since $|X_{i,e} - X_e| \leq w$ and $c_e(\cdot)$ is ζ -Lipschitz, by using (3.14) we get

$$|\mathbb{E}_{\widehat{\sigma}}[c_e(X_{i,e})] - c_e(x_e)| \leq \zeta w + |\mathbb{E}_{\widehat{\sigma}}[c_e(X_e)] - c_e(x_e)| \leq (\zeta + \gamma\alpha/4)w \quad (3.16)$$

and then we can approximate Eq. (3.15) with respect to costs:

$$\sum_{e \in s} c_e(x_e) \leq \sum_{e \in s'} c_e(x_e) + 2\kappa(\zeta + \gamma\alpha/4)w. \quad (3.17)$$

This shows that $(\mathbf{y}, \mathbf{x})$ is an ε -WE for the nonatomic game with demands $\mathbf{d}$, and then invoking Proposition 3.3(a) we get

$$|x_e - \widehat{x}_e| \leq \|\mathbf{x} - \widehat{\mathbf{x}}\|_2 \leq \sqrt{2\alpha\kappa(\zeta + \gamma\alpha/4)w/\beta}. \quad (3.18)$$

Plugging Eq. (3.18) and Eq. (3.11) into Eq. (3.10) we obtain the final estimate Eq. (3.9). $\square$

Using the previous result we obtain an explicit estimate for the L^2 distance between the random resource loads X_e^n in a sequence of mixed Nash equilibria for Γ_w^n , and the loads in the Wardrop equilibrium of the nonatomic limit game Γ^∞ . The distance depends on what is given by the previous result plus another term that captures the distance between the actual demand and the limiting one. In other words, one needs the weights to be small and close to the limit for the resulting equilibrium to approach the limit.

Corollary 3.5. *Let Γ_w^n be a sequence of WCGs satisfying Eq. (3.2) with $d_{\text{tot}}^n \leq \alpha$ for all n . Let the costs $c_e(\cdot)$ and Θ be as in Theorem 3.4, and take $\Xi = \sqrt{2C/\beta}$ with $C = \max_s \sum_{e \in s} c_e(\alpha)$, where the maximum is taken over all the feasible strategies $s \in \cup_{t \in \mathcal{T}} \mathcal{S}_t$. Let X_e^n be the random loads in a sequence of mixed Nash equilibria $\widehat{\sigma}^n \in \text{MNE}(\Gamma_w^n)$, and $\widehat{x}_e$ the unique resource loads in the WE for the nonatomic limit game $\Gamma^\infty = (\mathcal{G}, (d_t)_{t \in \mathcal{T}})$. Then, with w^n defined as in Eq. (3.2b), we have*

$$\|X_e^n - \widehat{x}_e\|_{L^2} \leq \Theta \cdot \sqrt{w^n} + \Xi \cdot \sqrt{\|\mathbf{d}^n - \mathbf{d}\|_1}. \quad (3.19)$$

Proof. Take $(\hat{\mathbf{y}}^n, \hat{\mathbf{x}}^n)$ a Wardrop equilibrium for the nonatomic game Γ^∞ with demands $\mathbf{d}^n$. A triangle inequality gives $\|X_e^n - \hat{x}_e\|_{L^2} \leq \|X_e^n - \hat{x}_e^n\|_{L^2} + |\hat{x}_e^n - \hat{x}_e|$, so the result follows from the estimate $\|X_e^n - \hat{x}_e^n\|_{L^2} \leq \Theta\sqrt{w^n}$ in [Theorem 3.4](#), and the bound $|\hat{x}_e^n - \hat{x}_e| \leq \Xi\sqrt{\|\mathbf{d}^n - \mathbf{d}\|_1}$ from [Proposition 3.3\(b\)](#). $\square$

3.3. Convergence of the price of anarchy. In this section, we investigate the convergence of the PoA for the sequence $\Gamma_{\mathbf{w}}^n$. Although we already know from the previous sections that the sequence of equilibria converges, we need to prove that the associated social cost converges, besides also proving that the sequence of optimal social costs also converges. To this end, we start by proving that for any sequence of converging expected flow-load pairs, the social cost of the sequence converges to that of the limiting flow-load pair.

Lemma 3.6. *Let $\Gamma_{\mathbf{w}}^n$ be a sequence of WCGs satisfying the conditions in [Eq. \(3.2\)](#), and σ^n an arbitrary sequence of mixed strategy profiles (not necessarily equilibria). Let $(Y_{t,s}^n, X_e^n)$ be the corresponding random flow-load pairs with expected values $(y_{t,s}^n, x_e^n)$. Then, along any subsequence of $(\mathbf{y}^n, \mathbf{x}^n)$ converging to some $(\mathbf{y}, \mathbf{x})$, the expected social cost $\text{ESC}(\sigma^n)$ converges to $\text{SC}(\mathbf{y}, \mathbf{x}) = \sum_{e \in \mathcal{E}} x_e c_e(x_e)$.*

Proof. Take a convergent subsequence and rename it so that $(\mathbf{y}^n, \mathbf{x}^n) \rightarrow (\mathbf{y}, \mathbf{x})$. By conditioning on $\mathbb{1}_{i,e}^n := \mathbb{1}_{\{e \in S_i^n\}}$, which indicates whether player i selects a strategy including e , we get

$$\text{ESC}(\sigma^n) = \sum_{e \in \mathcal{E}} \sum_{i \in \mathcal{N}^n} \mathbb{E}_{\sigma^n} [w_i^n \mathbb{1}_{i,e}^n c_e(X_e^n)] = \sum_{e \in \mathcal{E}} \sum_{i \in \mathcal{N}^n} w_i^n \sigma_{i,e}^n \mathbb{E}_{\sigma^n} [c_e(X_{i,e}^n)]. \quad (3.20)$$

Note that $|X_{i,e}^n - X_e^n| \leq w^n \rightarrow 0$ and $0 \leq X_e^n \leq d_{\text{tot}}^n$ with $d_{\text{tot}}^n \rightarrow d_{\text{tot}}$. Since $c_e(\cdot)$ is continuous, and hence uniformly continuous on compact intervals, it follows that $\mathbb{E}_{\sigma^n} [c_e(X_{i,e}^n)] - \mathbb{E}_{\sigma^n} [c_e(X_e^n)]$ converges to 0 uniformly in i , that is,

$$\delta_e^n := \max_{i \in \mathcal{N}^n} |\mathbb{E}_{\sigma^n} [c_e(X_{i,e}^n)] - \mathbb{E}_{\sigma^n} [c_e(X_e^n)]| \rightarrow 0. \quad (3.21)$$

Hence, using the identity

$$x_e^n = \sum_{i \in \mathcal{N}^n} w_i^n \sigma_{i,e}^n, \quad (3.22)$$

we obtain

$$\begin{aligned} \left| \text{ESC}(\sigma^n) - \sum_{e \in \mathcal{E}} x_e^n c_e(x_e^n) \right| &\leq \sum_{e \in \mathcal{E}} \sum_{i \in \mathcal{N}^n} w_i^n \sigma_{i,e}^n |\mathbb{E}_{\sigma^n} [c_e(X_{i,e}^n)] - c_e(x_e^n)| \\ &\leq \sum_{e \in \mathcal{E}} \sum_{i \in \mathcal{N}^n} w_i^n \sigma_{i,e}^n (\delta_e^n + |\mathbb{E}_{\sigma^n} [c_e(X_e^n)] - c_e(x_e^n)|) \\ &= \sum_{e \in \mathcal{E}} x_e^n (\delta_e^n + |\mathbb{E}_{\sigma^n} [c_e(X_e^n)] - c_e(x_e^n)|). \end{aligned}$$

The conclusion follows since $x_e^n \rightarrow x_e$ and $X_e^n \xrightarrow{\mathcal{D}} x_e$, so that $\mathbb{E}_{\sigma^n} [c_e(X_e^n)] \rightarrow c_e(x_e)$. $\square$

With this, we are ready to prove the convergence of optima.

Proposition 3.7. *Let $\Gamma_{\mathbf{w}}^n$ be a sequence of WCGs satisfying [Eq. \(3.2\)](#). Then $\text{Opt}(\Gamma_{\mathbf{w}}^n) \rightarrow \text{Opt}(\Gamma^\infty)$.*

Proof. Let $(\tilde{\mathbf{y}}, \tilde{\mathbf{x}})$ be a social optimum flow-load pair in the limiting game Γ^∞ . We convert the strategy flow $\tilde{\mathbf{y}}$ into mixed strategies $\tilde{\sigma}_t \in \Delta(\mathcal{S}_t)$ by setting

$$\forall s \in \mathcal{S}_t \quad \tilde{\sigma}_t(s) = \tilde{y}_{t,s} / d_t \quad (3.23)$$

when $d_t > 0$, and otherwise taking an arbitrary $\tilde{\sigma}_t \in \Delta(\mathcal{S}_t)$ for each type with $d_t = 0$.

Let $\tilde{\sigma}^n$ be the strategy profile for $\Gamma_{\mathcal{W}}^n$ in which player i plays $\tilde{\sigma}_i^n = \tilde{\sigma}_{t_i^n}$. For each t such that $d_t = 0$ we have $\tilde{y}_{t,s}^n \rightarrow 0 = \tilde{y}_{t,s}$ for all $s \in \mathcal{S}_t$, whereas when $d_t > 0$ we have

$$\tilde{y}_{t,s}^n = \frac{\tilde{y}_{t,s} d_t^n}{d_t} \rightarrow \tilde{y}_{t,s}. \quad (3.24)$$

Hence, $(\tilde{y}^n, \tilde{x}^n)$ converges to $(\tilde{y}, \tilde{x})$, and [Lemma 3.6](#) implies that $\text{ESC}(\tilde{\sigma}^n) \rightarrow \text{SC}(\tilde{y}, \tilde{x})$.

Now, take a sequence $\tilde{\sigma}^n$ of optimal mixed strategies in $\Gamma_{\mathcal{W}}^n$ and let $(\tilde{y}^n, \tilde{x}^n)$ be the corresponding expected loads. From the optimality of $\tilde{\sigma}^n$ we have $\text{ESC}(\tilde{\sigma}^n) \leq \text{ESC}(\tilde{\sigma}^n)$, so that

$$\limsup_{n \rightarrow \infty} \text{ESC}(\tilde{\sigma}^n) \leq \limsup_{n \rightarrow \infty} \text{ESC}(\tilde{\sigma}^n) = \text{SC}(\tilde{y}, \tilde{x}) = \text{Opt}(\Gamma^\infty). \quad (3.25)$$

On the other hand, taking a subsequence along which we attain the $\liminf_{n \rightarrow \infty} \text{ESC}(\tilde{\sigma}^n)$ and extracting a further subsequence so that $(\tilde{y}^n, \tilde{x}^n)$ converges to a certain limit $(\mathbf{y}, \mathbf{x})$, it follows that

$$\liminf_{n \rightarrow \infty} \text{ESC}(\tilde{\sigma}^n) = \text{SC}(\mathbf{y}, \mathbf{x}) \geq \text{Opt}(\Gamma^\infty), \quad (3.26)$$

which, combined with [Eq. \(3.25\)](#), yields the result. $\square$

We finish our analysis of WCGs with the convergence of the cost of equilibria and therefore that of the price of anarchy.

Theorem 3.8. *Let $\Gamma_{\mathcal{W}}^n$ be a sequence of WCG satisfying the conditions in [Eq. \(3.2\)](#) and let $\hat{\sigma}^n \in \text{MNE}(\Gamma_{\mathcal{W}}^n)$. Then, $\text{ESC}(\hat{\sigma}^n) \rightarrow \text{Eq}(\Gamma^\infty)$, and therefore $\text{PoA}(\Gamma_{\mathcal{W}}^n) \rightarrow \text{PoA}(\Gamma^\infty)$ and $\text{PoS}(\Gamma_{\mathcal{W}}^n) \rightarrow \text{PoA}(\Gamma^\infty)$.*

Proof. Every accumulation point of $(\hat{y}^n, \hat{x}^n)$ is a WE for Γ^∞ ; hence, by [Lemma 3.6](#), the full sequence $\text{ESC}(\hat{\sigma}^n)$ converges to $\text{Eq}(\Gamma^\infty)$ as $n \rightarrow \infty$. $\square$

Example 3.2. Consider the sequence of games in [Example 3.1](#) and the different equilibria described there. The social cost is minimized by splitting half of the players between the upper and lower paths (up to 1 player when n is odd). So for $n = 2$, we have $\text{PoA}(\Gamma_n) = 4/3$ and $\text{PoS}(\Gamma_n) = 1$, whereas for $n \geq 3$, by setting $\delta_n = 1$ if n is odd and $\delta_n = 0$ otherwise, we get $\text{PoA}(\Gamma^n) = 4n^2/(3n^2 + \delta_n)$ and $\text{PoS}(\Gamma^n) = (4n^2 - 2n + 2)/(3n^2 + \delta_n)$, and both converge to $\text{PoA}(\Gamma^\infty) = \text{PoS}(\Gamma^\infty) = 4/3$ as $n \rightarrow \infty$.

4. CONVERGENCE OF BERNOULLI CONGESTION GAMES

4.1. Sequences of Bernoulli congestion games. In this section we analyze the convergence of a sequence of BCGs

$$\Gamma_{\mathcal{B}}^n = (\mathcal{G}, (r_i^n, t_i^n)_{i \in \mathcal{N}^n}). \quad (4.1)$$

The elements that vary over the sequence are the set $\mathcal{N}^n$ of players, their participation probabilities r_i^n , and their types t_i^n , as well as the underlying probability spaces encoding the random participation of players and their mixed strategies. We proceed to study the convergence of a corresponding sequence of Bayes-Nash equilibria $\hat{\sigma}^n \in \text{BNE}(\Gamma_{\mathcal{B}}^n)$.

We assume that the number of players goes to infinity while the sequence of participation probabilities goes to zero in such a way that the aggregate expected demand for each type converges.

This is captured by the following asymptotic behavior as $n \rightarrow \infty$:

$$|\mathcal{N}^n| \rightarrow \infty, \quad (4.2a)$$

$$r^n := \max_{i \in \mathcal{N}^n} r_i^n \rightarrow 0, \quad (4.2b)$$

$$d_t^n := \sum_{i: t_i^n = t} r_i^n \rightarrow d_t. \quad (4.2c)$$

Let $Y_{t,s}^n$ and X_e^n be given by Eq. (2.12), with $U_i = U_i^n$, $t_i = t_i^n$, and $S_i = S_i^n$ drawn independently at random according to σ_i^n . As before, let $(\mathbf{y}^n, \mathbf{x}^n)$ be the vector of expected loads and observe that from Eq. (2.13) it follows that

$$d_t^n = \sum_{s \in \mathcal{S}_t} y_{t,s}^n, \quad \text{and} \quad x_e^n = \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} y_{t,s}^n \mathbb{1}_{\{e \in s\}}. \quad (4.3)$$

Our analysis is based on results on Poisson approximations for sums of Bernoulli random variables. We recall that a sequence of probability measures Q^n on $\mathbb{N}$ converges in total variation to the probability measure Q , written as $Q^n \xrightarrow{\text{TV}} Q$, whenever $\rho_{\text{TV}}(Q^n, Q) \rightarrow 0$ where

$$\rho_{\text{TV}}(Q^n, Q) := \sup_{A \subset \mathbb{N}} |Q^n(A) - Q(A)| = \frac{1}{2} \sum_{k \in \mathbb{N}} |Q^n(k) - Q(k)|. \quad (4.4)$$

Similarly, a sequence of random variables T^n converges in total variation to T if $\mathcal{L}(T^n) \xrightarrow{\text{TV}} \mathcal{L}(T)$. This concept of convergence is stronger than convergence in distribution—concept that we used in Section 3—and is suitable for situations where the limit distribution is discrete. It is known that, under the conditions in Eq. (4.2), the per-type demands $D_t^n = \sum_{i: t_i^n = t} U_i^n$ converge in total variation to a Poisson variable $D_t \sim \text{Poisson}(d_t)$ (see, e.g., Adell and Lekuona, 2005, Corollary 3.1). For instance, in the simple case where the probabilities of taking part in the game are the same for all players i of type t , say $r_i^n = d_t/N_t^n$, with $N_t^n = |\{i \in \mathcal{N}^n : t_i^n = t\}|$, we have $D_t^n \sim \text{Binomial}(N_t^n, d_t/N_t^n)$, which is well known to converge to $D_t \sim \text{Poisson}(d_t)$ as $N_t^n \rightarrow \infty$. Appendix A collects some further results that are used in our analysis.

Proposition 4.1. *Let Γ_B^n be a sequence of BCGs satisfying the conditions in Eq. (4.2), and let σ^n be an arbitrary sequence of mixed strategy profiles. Then*

- (a) *The sequence $(\mathbf{y}^n, \mathbf{x}^n)_{n \in \mathbb{N}}$ is bounded and each accumulation point $(\mathbf{y}, \mathbf{x})$ belongs to $\mathcal{F}(\mathbf{d})$.*
- (b) *Along any convergent subsequence of $(\mathbf{y}^n, \mathbf{x}^n)$, the random flows $Y_{t,s}^n$ and loads X_e^n converge in total variation to $Y_{t,s} \sim \text{Poisson}(y_{t,s})$ and $X_e \sim \text{Poisson}(x_e)$, respectively.*
- (c) *The limit variables $(Y_{t,s})_{t \in \mathcal{T}, s \in \mathcal{S}_t}$ are independent.*

Proof. (a) Since the expected demands d_t^n are convergent, this follows directly from Eq. (4.3).

(b) Take a convergent subsequence and for simplicity rename it to be the full sequence $(\mathbf{y}^n, \mathbf{x}^n) \rightarrow (\mathbf{y}, \mathbf{x})$. Note that Theorem A.4(a) gives

$$\rho_{\text{TV}}(\mathcal{L}(X_e^n), \mathcal{L}(V_e^n)) \leq r^n, \quad (4.5)$$

where $V_e^n \sim \text{Poisson}(x_e^n)$, and then, using the triangle inequality and Eq. (A.2), we conclude

$$\rho_{\text{TV}}(\mathcal{L}(X_e^n), \mathcal{L}(X_e)) \leq r^n + |x_e^n - x_e| \rightarrow 0. \quad (4.6)$$

A similar argument shows that $Y_{t,s}^n$ converges in total variation to $Y_{t,s}$.

(c) Consider the joint moment generating function M^n of the random variables $Y_{t,s}^n$

$$M^n(\lambda) = \mathbb{E}_{\sigma^n} \left[\exp \left(\sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} \lambda_{t,s} Y_{t,s}^n \right) \right], \quad (4.7)$$

where $\lambda = (\lambda_{t,s})_{t \in \mathcal{T}, s \in \mathcal{S}_t}$. Since

$$Y_{t,s}^n = \sum_{i: t_i^n = t} U_i^n \mathbb{1}_{\{S_i^n = s\}} \quad (4.8)$$

and $U_i^n \mathbb{1}_{\{S_i^n = s\}}$ are independent across players and types (although not across strategies), we have

$$\begin{aligned} M^n(\lambda) &= \mathbb{E}_{\sigma^n} \left[\exp \left(\sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} \sum_{i: t_i^n = t} \lambda_{t,s} U_i^n \mathbb{1}_{\{S_i^n = s\}} \right) \right] \\ &= \prod_{t \in \mathcal{T}} \prod_{i: t_i^n = t} \mathbb{E}_{\sigma^n} \left[\exp \left(U_i^n \sum_{s \in \mathcal{S}_t} \lambda_{t,s} \mathbb{1}_{\{S_i^n = s\}} \right) \right] \\ &= \prod_{t \in \mathcal{T}} \prod_{i: t_i^n = t} \left(1 - r_i^n + r_i^n \mathbb{E}_{\sigma^n} \left[\exp \left(\sum_{s \in \mathcal{S}_t} \lambda_{t,s} \mathbb{1}_{\{S_i^n = s\}} \right) \right] \right) \\ &= \prod_{t \in \mathcal{T}} \prod_{i: t_i^n = t} \left(1 - r_i^n + r_i^n \sum_{s \in \mathcal{S}_t} \sigma_i^n(s) \exp(\lambda_{t,s}) \right). \end{aligned}$$

Taking logarithms and using the fact that $\ln(1 + y) = y + O(y^2)$, it follows that

$$\begin{aligned} \ln M^n(\lambda) &= \sum_{t \in \mathcal{T}} \sum_{i: t_i^n = t} \ln \left(1 + r_i^n \sum_{s \in \mathcal{S}_t} \sigma_i^n(s) [\exp(\lambda_{t,s}) - 1] \right) \\ &= \sum_{t \in \mathcal{T}} \sum_{i: t_i^n = t} \left(\sum_{s \in \mathcal{S}_t} r_i^n \sigma_i^n(s) [\exp(\lambda_{t,s}) - 1] + O((r_i^n)^2) \right) \\ &= \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} y_{t,s}^n [\exp(\lambda_{t,s}) - 1] + \sum_{i \in \mathcal{N}^n} O((r_i^n)^2). \end{aligned}$$

In view of the conditions in [Eq. \(4.2\)](#) the sum $\sum_{i \in \mathcal{N}^n} O((r_i^n)^2)$ converges to zero, and, as a consequence,

$$\lim_{n \rightarrow \infty} M^n(\lambda) = \prod_{t \in \mathcal{T}} \prod_{s \in \mathcal{S}_t} \exp(y_{t,s} [\exp(\lambda_{t,s}) - 1]), \quad (4.9)$$

which is the moment generating function of a family of independent Poisson random variables with parameters $y_{t,s}$. $\square$

Remark 4.1. Even if the strategy flows $Y_{t,s}$ are independent, in general the resource loads X_e are not. For instance in a trivial network with a single OD pair connected by a unique path, all the edges will carry exactly the same load.

4.2. The Poisson limit game. Let us recall that in BCGs the resource costs $c_e : \mathbb{N} \rightarrow \mathbb{R}_+$ are only defined over the integers. We introduce the auxiliary costs $\check{c}_e : \mathbb{R}_+ \rightarrow \mathbb{R}_+ \cup \{+\infty\}$ where $\check{c}_e(x)$ is defined for $x \geq 0$ by taking a random variable $X \sim \text{Poisson}(x)$ and setting

$$\check{c}_e(x) := \mathbb{E}[c_e(1 + X)] = \sum_{k=0}^{\infty} c_e(1 + k) e^{-x} \frac{x^k}{k!}. \quad (4.10)$$

For these costs to be finite and smooth, we impose the following mild assumption:

$$\exists v \in \mathbb{R} \text{ and } \alpha > \sum_{t \in \mathcal{T}} d_t \text{ with } \mathbb{E}[|\Delta^2 c_e(1 + V)|] \leq v \text{ for all } e \in \mathcal{E} \text{ and } V \sim \text{Poisson}(\alpha), \quad (4.11)$$

where $\Delta^2 c_e(k) = c_e(k + 2) - 2c_e(k + 1) + c_e(k)$. This condition holds in particular for costs with subexponential growth $c_e(k) \leq b \exp(ak)$ for some $a, b \in \mathbb{R}_+$, which includes all polynomials. The condition fails for rapidly growing functions such as $c_e(k) = k!$ or $c_e(k) = \exp(\exp(k))$. The following is the main consequence of Eq. (4.11) used in the sequel.

Lemma 4.2. *Assume Eq. (4.11) and let $\zeta = (e^\alpha - 1)v + \max_{e \in \mathcal{E}}[c_e(2) - c_e(1)]$. Then, the auxiliary costs $\check{c}_e(\cdot)$ are finite and of class C^2 on $[0, \alpha]$ with $0 \leq \check{c}'_e(x) \leq \zeta$ for all $x \in [0, \alpha]$. Moreover, let Γ_B be a BCG with $d_{\text{tot}} \leq \alpha$ and let $r = \max_{i \in \mathcal{N}} r_i$. Let X_e be the random loads in a mixed strategy profile σ , and $Z_{i,e} = X_e - U_{i,e}$ the loads excluding player i . Then, the expected values $x_e = \mathbb{E}_\sigma[X_e]$ and $z_{i,e} = \mathbb{E}_\sigma[Z_{i,e}]$ satisfy $|z_{i,e} - x_e| = \mathbb{E}_\sigma[U_{i,e}] \leq r$, and we have*

$$|\mathbb{E}_\sigma[c_e(1 + Z_{i,e})] - \check{c}_e(x_e)| \leq \Lambda(r) := \frac{\alpha v}{2} \frac{r e^r}{(1 - r)^2} + \zeta r. \quad (4.12)$$

Proof. The smoothness and the bound $0 \leq \check{c}'_e(x) \leq \zeta$ for the auxiliary costs follow directly from Proposition A.2(b) and Corollary A.3 in the Appendix. In particular, $\check{c}_e(\cdot)$ is ζ -Lipschitz, and then Eq. (4.12) follows by using a triangle inequality and Theorem A.4(b). $\square$

We now proceed to prove that the random loads at the equilibrium of games with Bernoulli demands converge to a Wardrop equilibrium for the nonatomic congestion game $\check{\Gamma}^\infty = (\check{\mathcal{G}}, (d_t)_{t \in \mathcal{T}})$ defined by the costs $\check{c}_e(\cdot)$ and the limiting demands d_t .

Theorem 4.3. *Let Γ_B^n be a sequence of BCGs satisfying Eqs. (4.2) and (4.11), and let $\hat{\sigma}^n \in \text{BNE}(\Gamma_B^n)$ be an arbitrary sequence of Bayesian Nash equilibria. Then the corresponding sequence of expected flow-load pairs $(\hat{\mathbf{y}}^n, \hat{\mathbf{x}}^n)$ is bounded and every accumulation point $(\hat{\mathbf{y}}, \hat{\mathbf{x}})$ is a Wardrop equilibrium for the nonatomic congestion game $\check{\Gamma}^\infty$. Furthermore, along every such convergent subsequence the random flows $Y_{t,s}^n$ and loads X_e^n converge in total variation to Poisson limits $Y_{t,s} \sim \text{Poisson}(\hat{y}_{t,s})$ and $X_e \sim \text{Poisson}(\hat{x}_e)$, where the variables $Y_{t,s}$ are independent.*

Proof. By Proposition 4.1, it suffices to show that every accumulation point is a WE for $\check{\Gamma}^\infty$. Take a convergent subsequence and rename it to be the full sequence $(\hat{\mathbf{y}}^n, \hat{\mathbf{x}}^n) \rightarrow (\hat{\mathbf{y}}, \hat{\mathbf{x}})$. We must prove that, for each type t and each pair of strategies s, s' in $\mathcal{S}_t$, we have

$$\hat{y}_{t,s} > 0 \Rightarrow \sum_{e \in s} \check{c}_e(\hat{x}_e) \leq \sum_{e \in s'} \check{c}_e(\hat{x}_e). \quad (4.13)$$

A strict inequality $\hat{y}_{t,s} > 0$ implies that for all n large enough we have $\hat{y}_{t,s}^n > 0$ and there must be a player $i = i^n$ of type $t_i^n = t$ with $\hat{\sigma}_i^n(s) > 0$. The equilibrium condition in Γ_B^n implies that for

each alternative strategy $s' \in \mathcal{S}_i$ for player i we have

$$\sum_{e \in s} \mathbb{E}_{\hat{\sigma}^n} [c_e(1 + Z_{i,e}^n)] \leq \sum_{e \in s'} \mathbb{E}_{\hat{\sigma}^n} [c_e(1 + Z_{i,e}^n)]. \quad (4.14)$$

Using Lemma 4.2, we get

$$|\mathbb{E}_{\hat{\sigma}^n} [c_e(1 + Z_{i,e}^n)] - \check{c}_e(x_e^n)| \leq \Lambda(r^n) \rightarrow 0, \quad (4.15)$$

and, since $x_e^n \rightarrow \hat{x}_e$, it follows that $\mathbb{E}_{\hat{\sigma}^n} [c_e(1 + Z_{i,e}^n)] \rightarrow \check{c}_e(\hat{x}_e)$. Thus, letting $n \rightarrow \infty$ in Eq. (4.14), we obtain Eq. (4.13). $\square$

Moreover, under mild additional conditions, the equilibrium loads of the corresponding nonatomic game are unique and every sequence of equilibria converges. The following corollary is a direct consequence of Theorem 4.3 and a property of $\check{c}_e(\cdot)$ (see Corollary A.3(b)).

Corollary 4.4. *Let $c_e : \mathbb{N} \rightarrow \mathbb{R}$ be weakly increasing and nonconstant for $k \geq 1$ and assume that the conditions Eq. (4.11) hold. Then, the extended functions $\check{c}_e(\cdot)$ are strictly increasing and the loads $\hat{x}_e$ are the same in every WE for $\check{\Gamma}^\infty$. Moreover, for every sequence Γ_B^n of BCGs satisfying the conditions in Eq. (4.2) and every sequence $\hat{\sigma}^n \in \text{BNE}(\Gamma_B^n)$, the random loads X_e^n converge in total variation to a random variable $X_e \sim \text{Poisson}(\hat{x}_e)$.*

Example 4.1. Consider a sequence $(\Gamma_B^n)_{n \in \mathbb{N}}$ of games played on the Wheatstone network in Fig. 2, with n identical players and $r_i \equiv r = 1/n$ so that the total expected demand is $d_{\text{tot}} = 1$. Notice that the conditional expected cost of a player $i \in \mathcal{N}$ when using e_1 or e_5 equals $1 + r(x - 1)$, where x is the number of players using the edge. It follows that the zig-zag path (e_1, e_3, e_5) is strictly dominated by a linear combination of the upper and lower paths, and no player will use the zig-zag path. This is in sharp contrast with the weighted congestion games described in Example 3.1, where all but one players were choosing the zig-zag path. This difference is explained by the fact that each player has a unit weight and has a significant impact on the costs.

If the number of players $n = 2k$ is even, there is a unique PNE (modulo permutations of players), where half of the players choose the upper path and the other half take the lower path. In this equilibrium, the expected cost for each player is $(2.5n - 1)/n^2$. For an odd number of players $n = 2k + 1$, there is a Nash equilibrium (NE) where each edge gets k players for sure, and the last player randomizes arbitrarily between these two paths. For all n , there is also a symmetric mixed equilibrium where every player mixes with $q = 1/2$ between the upper and lower paths, and the expected cost of each player is $(5n - 1)/(2n^2)$. In general, one can prove that every MNE is of the form where k_1 players take the upper path, k_2 players take the lower path, and the remaining $k_3 = n - k_1 - k_2$ players, with $k_3 - 1 > |k_2 - k_1|$, randomize between these two paths with exactly the same probabilities $(q, 1 - q)$ with $q = \frac{1}{2} \left(1 + \frac{k_2 - k_1}{k_3 - 1} \right)$, so as to equalize their costs.

As $n \rightarrow \infty$, all these equilibria converge to the WE of a nonatomic game where the costs of e_1 and e_5 become $\check{c}(x) = 1 + x$. In this WE, half of the demand goes on the upper path and the remaining half on the lower path.

4.3. Rate of convergence. Analogously to Section 3.2, we now find nonasymptotic estimates for the speed of convergence under a mild additional assumption. Eq. (4.6) provides a simpler—but more difficult to quantify—bound on the distance between the resource-loads in the finite games Γ_B^n and the Poisson limits. Since r^n is a primitive parameter of the model, its proximity to zero is readily available. In contrast, the second term $|\hat{x}_e^n - \hat{x}_e|$ is only known to converge to zero asymptotically.

We first show that the distance between a Bayesian Nash equilibrium for a Bernoulli congestion game Γ_B and a Wardrop equilibrium for the nonatomic game $\check{\Gamma}^\infty$ with the same aggregate demands is bounded by an expression that depends on the topology and a function $\Lambda(r)$ defined in Eq. (4.12) evaluated at the maximum participation probability $r = \max_{i \in \mathcal{N}} r_i$.

Theorem 4.5. *Let Γ_B be a BCG with expected demands $d_t = \sum_{i:t_i=t} r_i$ and $d_{\text{tot}} \leq \alpha$. Suppose that Eq. (4.11) holds and that $\check{c}'_e(x) \geq \beta > 0$ for all $x \in [0, \alpha]$. Let $\check{\Theta} = \sqrt{2\alpha\kappa/\beta}$ with κ the cardinality of the largest feasible strategy $s \in \cup_{t \in \mathcal{T}} \mathcal{S}_t$. Let X_e be the random loads in a Bayesian Nash equilibrium $\hat{\sigma} \in \text{BNE}(\Gamma_B)$ with expected flow-loads $(\mathbf{y}, \mathbf{x})$, and $\hat{X}_e \sim \text{Poisson}(\hat{x}_e)$ with $\hat{\mathbf{x}}$ the unique Wardrop equilibrium loads in the nonatomic game $\check{\Gamma}^\infty$ with costs $\check{c}_e(\cdot)$ and with the same aggregate demands d_t as Γ_B . Then, $\|\mathbf{x} - \hat{\mathbf{x}}\|_2 \leq \check{\Theta} \cdot \sqrt{\Lambda(r)}$ and*

$$\rho_{\text{TV}}(\mathcal{L}(X_e), \mathcal{L}(\hat{X}_e)) \leq r + \check{\Theta} \cdot \sqrt{\Lambda(r)}. \quad (4.16)$$

Proof. The bound $\|\mathbf{x} - \hat{\mathbf{x}}\|_2 \leq \check{\Theta} \sqrt{\Lambda(r)}$ follows from Proposition 3.3(a) since $\mathbf{x}$ is an ε -WE for $\check{\Gamma}^\infty$ with $\varepsilon = 2\kappa\Lambda(r)$. Indeed, if $y_{t,s} > 0$ there exists some player $i \in \mathcal{N}$ with $t_i = t$ and $\hat{\sigma}_i(s) > 0$, so that the equilibrium condition yields for all $s' \in \mathcal{S}_t$

$$\sum_{e \in s} \mathbb{E}_{\hat{\sigma}}[c_e(1 + Z_{i,e})] \leq \sum_{e \in s'} \mathbb{E}_{\hat{\sigma}}[c_e(1 + Z_{i,e})], \quad (4.17)$$

and then Eq. (4.12) implies the conditions for ε -WE

$$\sum_{e \in s} \check{c}_e(x_e) \leq \sum_{e \in s'} \check{c}_e(x_e) + 2\kappa\Lambda(r).$$

Now, taking $V_e \sim \text{Poisson}(x_e)$, Theorem A.4(a) gives $\rho_{\text{TV}}(\mathcal{L}(X_e), \mathcal{L}(V_e)) \leq r$, while Eq. (A.2) implies $\rho_{\text{TV}}(\mathcal{L}(V_e), \mathcal{L}(\hat{X}_e)) \leq |x_e - \hat{x}_e|$, so that Eq. (4.16) follows from a triangle inequality. $\square$

Remark 4.2. The assumption $\check{c}'_e(x) \geq \beta$ for $x \in [0, \alpha]$ holds when $c_e(k+1) \geq c_e(k) + \beta$ for all $e \in \mathcal{E}$ and $k \geq 1$. A milder condition is $0 < \delta := \min_{e \in \mathcal{E}} c_e(2) - c_e(1)$ since then one can take $\beta = \delta e^{-\alpha}$.

Using the previous result we may bound the distance between the random loads in a sequence Γ_B^n of BCGs and the corresponding Wardrop equilibrium in the limit game $\check{\Gamma}^\infty$. Besides depending on the previous bound, there is an additional term that captures the distance between the participation probabilities in the instance and those in the limit. Having an equilibrium close to the limit thus requires small participation probabilities together with them being close to the limit.

Corollary 4.6. *Let Γ_B^n be a sequence of BCG satisfying the conditions in Eqs. (4.2) and (4.11), with total expected demands bounded as $d_{\text{tot}}^n \leq \alpha$ for all n . Suppose that $\check{c}'_e(x) \geq \beta > 0$ for all $e \in \mathcal{E}$ and $x \in [0, \alpha]$, and let $\check{\Theta} = \sqrt{2\alpha\kappa/\beta}$ and $\check{\Xi} = \sqrt{2C/\beta}$ with κ the cardinality of the largest feasible strategy $s \in \cup_{t \in \mathcal{T}} \mathcal{S}_t$ and $C = \max_s \sum_{e \in s} \check{c}_e(\alpha)$. Let X_e^n be the random loads in a sequence of Bayesian Nash equilibria $\hat{\sigma}^n \in \text{BNE}(\Gamma_B^n)$, and $X_e \sim \text{Poisson}(\hat{x}_e)$ with $\hat{\mathbf{x}}$ the unique resource loads in the WE for the nonatomic limit game $\check{\Gamma}^\infty$. Then,*

$$\rho_{\text{TV}}(\mathcal{L}(X_e^n), \mathcal{L}(X_e)) \leq r^n + \check{\Theta} \cdot \sqrt{\Lambda(r^n)} + \check{\Xi} \cdot \sqrt{\|\mathbf{d}^n - \mathbf{d}\|_1}. \quad (4.18)$$

Proof. This follows by considering $\hat{X}_e^n \sim \text{Poisson}(\hat{x}_e^n)$ with $(\hat{\mathbf{y}}^n, \hat{\mathbf{x}}^n)$ a Wardrop equilibrium for the game $\check{\Gamma}^\infty$ with demands $\mathbf{d}^n$, and then using a triangle inequality and applying Theorem 4.5 and Proposition 3.3(b). $\square$

4.4. Convergence of the price of anarchy. We now investigate the convergence of the price of anarchy for the sequence of BCGs Γ_B^n . To this end, we exploit the following fact.

Lemma 4.7. *Let Γ_B^n be a sequence of BCGs satisfying the conditions in Eqs. (4.2) and (4.11), and let σ^n be an arbitrary sequence of mixed strategies. Let $Y_{t,s}^n$ and X_e^n be the corresponding random loads with expected values $y_{t,s}^n$ and x_e^n . Then, along any subsequence of $(\mathbf{y}^n, \mathbf{x}^n)$ converging to some $(\mathbf{y}, \mathbf{x})$, the expected social cost $\text{ESC}(\sigma^n)$ converges to $\check{\text{SC}}(\mathbf{y}, \mathbf{x}) := \sum_{e \in \mathcal{E}} x_e \check{c}_e(x_e)$.*

Proof. Take a convergent subsequence and rename it so that $(\mathbf{y}^n, \mathbf{x}^n) \rightarrow (\mathbf{y}, \mathbf{x})$. By conditioning on the event $U_{i,e}^n = 1$, we have

$$\text{ESC}(\sigma^n) = \sum_{e \in \mathcal{E}} \sum_{i \in \mathcal{N}^n} \mathbb{E}_{\sigma^n} [U_{i,e}^n c_e(X_e^n)] = \sum_{e \in \mathcal{E}} \sum_{i \in \mathcal{N}^n} r_i^n \sigma_{i,e}^n \mathbb{E}_{\sigma^n} [c_e(1 + Z_{i,e}^n)]. \quad (4.19)$$

Using the identity $x_e^n = \sum_{i \in \mathcal{N}^n} r_i^n \sigma_{i,e}^n$ and invoking Lemma 4.2, we obtain

$$\begin{aligned} \left| \text{ESC}(\sigma^n) - \sum_{e \in \mathcal{E}} x_e^n \check{c}_e(x_e^n) \right| &\leq \sum_{e \in \mathcal{E}} \sum_{i \in \mathcal{N}^n} r_i^n \sigma_{i,e}^n |\mathbb{E}_{\sigma^n} [c_e(1 + Z_{i,e}^n)] - \check{c}_e(x_e^n)| \\ &\leq \sum_{e \in \mathcal{E}} \sum_{i \in \mathcal{N}^n} r_i^n \sigma_{i,e}^n \Lambda(r^n) = \sum_{e \in \mathcal{E}} x_e^n \Lambda(r^n) \rightarrow 0, \end{aligned}$$

and then the conclusion follows from $x_e^n \rightarrow x_e$ and the continuity of $\check{c}_e(\cdot)$. $\square$

Proposition 4.8. *Let Γ_B^n be a sequence of BCGs satisfying Eqs. (4.2) and (4.11). Then $\text{Opt}(\Gamma_B^n) \rightarrow \text{Opt}(\check{\Gamma}^\infty)$.*

Proof. It suffices to repeat the proof of Proposition 3.7 step-by-step, replacing $\text{SC}(\cdot)$ with $\check{\text{SC}}(\cdot)$ and invoking Lemma 4.7 instead of Lemma 3.6. $\square$

From Lemma 4.2 we have that $\check{c}'_e(x) \geq 0$ for $x \in [0, \alpha]$ so that the extended costs $\check{c}'_e(\cdot)$ are weakly increasing, and therefore the social cost $\text{SC}(\hat{\mathbf{y}}, \hat{\mathbf{x}}) \equiv \text{Eq}(\check{\Gamma}^\infty)$ is the same in every WE. As in the case of WCGs we obtain the following theorem.

Theorem 4.9. *Let Γ_B^n be a sequence of BCGs satisfying the conditions in Eqs. (4.2) and (4.11). Then, for every sequence $\hat{\sigma}^n \in \text{BNE}(\Gamma_B^n)$, the expected social cost $\text{ESC}(\hat{\sigma}^n)$ converges to $\text{Eq}(\check{\Gamma}^\infty)$. As a consequence, both $\text{PoA}(\Gamma_B^n)$ and $\text{PoS}(\Gamma_B^n)$ converge to $\text{PoA}(\check{\Gamma}^\infty) = \text{PoS}(\check{\Gamma}^\infty)$.*

Remark 4.3. For polynomial costs $c_e(\cdot)$ of degree at most d we have that $\check{c}_e(x)$ are again polynomials of the same degrees (though with different coefficients), so that the results in Roughgarden (2002, 2003) imply

$$\lim_{n \rightarrow \infty} \text{PoA}(\Gamma_B^n) = \text{PoA}(\check{\Gamma}^\infty) \leq B(d) := \frac{(d+1)\sqrt[d]{d+1}}{(d+1)\sqrt[d]{d+1} - d}. \quad (4.20)$$

In fact, for $d = 1$, the bound $\text{PoA}(\Gamma_B^n) \leq 4/3$ is valid as soon as $r^n \leq 1/4$ (see Cominetti et al. (2019)). For higher degrees we conjecture the existence of a threshold for r^n under which $\text{PoA}(\Gamma_B^n)$ already falls below the nonatomic bound $B(d)$. The current result implies that we only have this as an asymptotic bound when $r^n \rightarrow 0$.

Example 4.2. Consider the sequence of games Γ_B^n on the Wheatstone network in Example 4.1. The social cost is minimized by splitting half of the players between the upper and lower routes (up to 1 player when n is odd). Since this strategy profile is also a pure Nash equilibrium, we have

$\text{PoS}(\Gamma_B^n) = 1$ for all $n \in \mathbb{N}$. As far as the PoA is concerned, the worst equilibrium occurs when each player chooses the upper and lower routes with equal probability $1/2$. Setting $\delta_n = 1/n$ when n is odd and $\delta_n = 0$ otherwise, we obtain $\text{PoA}(\Gamma_B^n) = \frac{5n-1}{5n-2+\delta_n}$, which converges to $\text{PoA}(\tilde{\Gamma}^\infty) = 1$ as $n \rightarrow \infty$.

Remark 4.4. The [Examples 3.2](#) and [4.2](#) may suggest that the price of anarchy in the nonatomic game obtained as a limit of WCGs would be larger than the price of anarchy in the nonatomic limit game for the BCGs. This is not true in general. Consider for instance the Pigou network in [Fig. 3](#) with a demand of 1. The WE of the standard nonatomic game is optimal, so $\text{PoA}(\Gamma^\infty) = 1$.




Figure 3. Pigou network.

The WE of the nonatomic limit game of the Bernoulli game, in which the auxiliary cost function on the upper edge is now $\check{c}_1(x) = 1 + x$, sends all demand on the upper path, whereas in the social optimum (SO) the demand is split over the upper and lower path. So, we have that $\text{PoA}(\tilde{\Gamma}^\infty) = 8/7$.

5. GAMES WITH POPULATION UNCERTAINTY AND POISSON GAMES

In this final section we establish a connection between the nonatomic game $\tilde{\Gamma}^\infty$ obtained as a limit of Bernoulli congestion games, and Poisson games in the sense of [Myerson \(1998b\)](#).

Let us first recall the concept of a *game with population uncertainty*, as first introduced by [Myerson \(1998b\)](#), which is given by a tuple $\Gamma_p = (\mathcal{T}, (\mathcal{S}_t)_{t \in \mathcal{T}}, \mu, C)$, where $\mathcal{T}$ is a finite set of types and each $t \in \mathcal{T}$ has an associated strategy set $\mathcal{S}_t$ and a random number of players N_t that take part in the game.¹ Players are not aware of the specific identities of the other players and they only know the joint probability distribution μ over $\mathbb{N}^{\mathcal{T}}$ according to which the number $\mathbf{N} = (N_t)_{t \in \mathcal{T}}$ of players of each type t are drawn. This is why in these games players are treated symmetrically by assuming that all players of type t adopt the same mixed strategy $\sigma_t \in \Delta(\mathcal{S}_t)$, which only depends on their type. The cost $C(Y^{-t} \mid t, s)$ for an active player of type t choosing an action $s \in \mathcal{S}_t$, depends on the strategy flows Y^{-t} induced by the *other* players present in the game, as described below.

The measure μ , together with $\sigma = (\sigma_t)_{t \in \mathcal{T}}$, determines the distribution of the strategy flows $Y = (Y_{t,s})_{t \in \mathcal{T}, s \in \mathcal{S}_t}$ over types and strategies. Conditional on $N_t = \bar{n}_t$ the flows $Y_t = (Y_{t,s})_{s \in \mathcal{S}_t}$ for type $t \in \mathcal{T}$ are distributed on strategies as independent multinomials $Y_t \sim \text{Multinomial}(\bar{n}_t, \sigma_t)$. Thus, if for each $\mathbf{n} = (n_{t,s})_{t \in \mathcal{T}, s \in \mathcal{S}_t}$, we define $\bar{\mathbf{n}} := (\bar{n}_t)_{t \in \mathcal{T}}$ with $\bar{n}_t := \sum_{s \in \mathcal{S}_t} n_{t,s}$, then we have

$$P_{\mu, \sigma}(Y = \mathbf{n}) = \mu(\bar{\mathbf{n}}) \prod_{t \in \mathcal{T}} \left(\bar{n}_t! \prod_{s \in \mathcal{S}_t} \frac{\sigma_t(s)^{n_{t,s}}}{n_{t,s}!} \right), \quad (5.1)$$

where $P_{\mu, \sigma}$ is the probability measure induced by μ and σ .

¹To avoid trivialities we suppose that N_t is not identically zero.

Now, consider a generic player of a given type t who happens to be active in the game, and let $\mathbf{N}^{-t} = (\mathbf{N}_{t'}^{-t})_{t' \in \mathcal{T}}$ denote the random vector giving the number of *other players* for each type, excluding this active player. As shown by Myerson (1998b) (see also Milchtaich, 2004), the posterior distribution of $\mathbf{N}^{-t}$ assessed by such a generic player is given by

$$\mu(\bar{\mathbf{n}} \mid t) := P_{\mu, \sigma}(\mathbf{N}^{-t} = \bar{\mathbf{n}}) = \frac{(\bar{n}_t + 1) \mu(\bar{\mathbf{n}} + \boldsymbol{\delta}_t)}{E_{\mu}[N_t]}, \quad (5.2)$$

where $\bar{\mathbf{n}} + \boldsymbol{\delta}_t$ denotes the vector $\bar{\mathbf{n}}$ with $\bar{n}_t$ replaced by $\bar{n}_t + 1$. Accordingly, the posterior distribution of the strategy flows $\mathbf{Y}^{-t} = (\mathbf{Y}_{t', s}^{-t})_{t' \in \mathcal{T}, s \in \mathcal{S}_{t'}}$ induced by the remaining active players is given by Eq. (5.1), with $\mu(\bar{\mathbf{n}})$ replaced by $\mu(\bar{\mathbf{n}} \mid t)$.

The expected cost for a player of type t is computed according to this posterior distribution, and an equilibrium is then defined as a family of type-dependent mixed strategies $\hat{\sigma} = (\hat{\sigma}_t)_{t \in \mathcal{T}}$ with $\hat{\sigma}_t \in \Delta(\mathcal{S}_t)$ such that

$$\forall t \in \mathcal{T}, \forall s, s' \in \mathcal{S}_t \quad \hat{\sigma}_t(s) > 0 \Rightarrow E_{\mu, \hat{\sigma}}[C(\mathbf{Y}^{-t} \mid t, s)] \leq E_{\mu, \hat{\sigma}}[C(\mathbf{Y}^{-t} \mid t, s')]. \quad (5.3)$$

Remark 5.1. The Bernoulli congestion games in Section 2.3 fall in the framework of games with population uncertainty where $N_t = \sum_{i: t_i=t} U_i$ is the sum of independent nonhomogeneous Bernoulli random variables. However, we considered not only strategies defined by the player's type, but also asymmetric equilibria in which players choose their strategies individually.

Example 5.1. Consider the game $\Gamma_{\mathbf{b}}^n$ on the Wheatstone network of Fig. 2 in Example 4.1. Recall that there are n players—all of the same type t —and each one is present with probability $1/n$. Assume that all players play a mixed strategy in which with probability $1/2$ they choose the upper path, and with probability $1/2$ they choose the lower path. Let $\mathbf{Y} = (Y_1, Y_2, Y_3)$ denote the random vector that gives the number of players on the paths $s_1 = (e_1, e_4)$, $s_2 = (e_1, e_3, e_5)$ and $s_3 = (e_2, e_5)$, respectively. Then $Y_1, Y_3 \sim \text{Binomial}(n, (2n)^{-1})$ and $Y_2 \equiv 0$, while the corresponding posterior distributions are $Y_1^{-t}, Y_3^{-t} \sim \text{Binomial}(n-1, (2n-2)^{-1})$ and $Y_2^{-t} \equiv 0$.

An important subclass of games with population uncertainty are the *Poisson games* which are characterized by independent variables $N_t \sim \text{Poisson}(d_t)$, with $d_t > 0$, that is,

$$\mu(\bar{\mathbf{n}}) = P(N_t = \bar{n}_t, \forall t \in \mathcal{T}) = \prod_{t \in \mathcal{T}} e^{-d_t} \frac{(d_t)^{\bar{n}_t}}{\bar{n}_t!}. \quad (5.4)$$

It is not difficult to see that Poisson games are characterized by the fact that the posteriors $\mu(\cdot \mid t)$ coincide with μ for every $t \in \mathcal{T}$. Moreover, in this case the loads $Y_{t,s}$ are also independent with $Y_{t,s} \sim \text{Poisson}(d_t \sigma_t(s))$. In fact, as shown in Myerson (1998b, Theorem 1), in a game with population uncertainty the variables $Y_{t,s}$ are independent if and only if the game is Poisson.

The nonatomic game $\check{\Gamma}^\infty$ in Section 4.2, obtained as a limit of a sequence of Bernoulli congestion games, can be interpreted as a Poisson game defined by the costs

$$C(\mathbf{Y}^{-t} \mid t, s) = \sum_{e \in s} c_e(1 + X_e^{-t}), \quad X_e^{-t} = \sum_{t' \in \mathcal{T}} \sum_{s \in \mathcal{S}_{t'}} Y_{t', s}^{-t} \mathbb{1}_{\{e \in s\}}. \quad (5.5)$$

We state this observation in the following result.

Theorem 5.1. *Let σ be a strategy profile in the Poisson game defined by the costs in Eq. (5.5) and the demands given in Eq. (5.4) with $d_t > 0$. Define $(\mathbf{y}, \mathbf{x})$ as $y_{t,s} = d_t \sigma_t(s)$ and $x_e = \sum_{t \in \mathcal{T}} \sum_{s \in \mathcal{S}_t} y_{t,s} \mathbb{1}_{\{e \in s\}}$. Then σ is an equilibrium in the Poisson game if and only if $(\mathbf{y}, \mathbf{x})$ is a Wardrop equilibrium for the nonatomic game $\check{\Gamma}^\infty$ in Section 4.2.*

Proof. It suffices to note that the posterior distribution of Y^{-t} is Poisson with independent components and expected values $E_{\mu,\sigma}[Y_{t',s}^{-t}] = y_{t',s}$, so that $X_e^{-t} \sim \text{Poisson}(x_e)$. Then, taking expectation, we get precisely

$$E_{\mu,\sigma}[C(Y^{-t} \mid t, s)] = \sum_{e \in S} E[c_e(1 + X_e^{-t})] = \sum_{e \in S} \check{c}_e(x_e), \quad (5.6)$$

where $\check{c}_e(\cdot)$ is defined as in Eq. (4.10). $\square$

Notice that the costs in Eq. (5.5) depend on the strategy flows Y^{-t} only through the aggregate resource loads X_e^{-t} , whereas Myerson (1998b) considers more general cost functions $C(Y^{-t} \mid t, s)$. However, to ensure that the expected costs are well defined, these costs were assumed bounded. In our case this boundedness assumption is relaxed and replaced by the conditions in Eq. (4.11). On the other hand, as mentioned in the Introduction, Myerson (1998b) introduced Poisson games axiomatically and not as limits of a sequence of finite games. The following extension of Theorem 4.3, to the general setting of games with population uncertainty, fills this gap by establishing the convergence of sequences of games with Bernoulli players towards Poisson games.

Let $\mathcal{T}$ be a finite set of types with strategy sets $\mathcal{S}_t$ for $t \in \mathcal{T}$. Consider a sequence of games Γ^n with finitely many players $i \in \mathcal{N}^n$, with types t_i^n , and probabilities r_i^n of being active. Let U_i^n be Bernoulli random variables with $P(U_i^n = 1) = r_i^n$, and let player i choose $S_i^n \in \mathcal{S}_{t_i^n}$ at random using a mixed strategy σ_i^n . Let Y^n be the random vector of strategy flows $Y_{t,s}^n = \sum_{j: t_j^n=t} U_j^n \mathbb{1}_{\{S_j^n=s\}}$, and define $Y^{-i,n}$ similarly by excluding player i . The expected cost of an action $s \in \mathcal{S}_{t_i^n}$ for player i is given by $E_{\sigma^n}[C(Y^{-i,n} \mid t_i^n, s)]$ with $C(\cdot \mid t, s)$ a family of *bounded* functions. A Nash equilibrium $\hat{\sigma}^n$ is defined as usual by the condition

$$\forall i \in \mathcal{N}^n, \forall s, s' \in \mathcal{S}_{t_i^n} \quad \hat{\sigma}_i^n(s) > 0 \Rightarrow E_{\hat{\sigma}^n}[C(Y^{-i,n} \mid t_i^n, s)] \leq E_{\hat{\sigma}^n}[C(Y^{-i,n} \mid t_i^n, s')]. \quad (5.7)$$

Theorem 5.2. *Consider a sequence Γ^n as above, and assume the conditions in Eq. (4.2) with $d_t > 0$. Then, for every sequence $\hat{\sigma}^n$ of Nash equilibria, the expected loads $y_{t,s}^n = E_{\hat{\sigma}^n}[Y_{t,s}^n]$ are bounded and each accumulation point $\hat{\mathbf{y}} = (\hat{y}_{t,s})_{t \in \mathcal{T}, s \in \mathcal{S}_t}$ corresponds to an equilibrium $\hat{\sigma}$ in the Poisson game by setting $\hat{\sigma}_t(s) = \hat{y}_{t,s}/d_t$ for all $s \in \mathcal{S}_t$.*

Proof. The boundedness of the expected strategy loads $y_{t,s}^n$ follows from $\sum_{s \in \mathcal{S}_t} y_{t,s}^n = d_t^n \rightarrow d_t$. Take a convergent subsequence and rename it so that $\mathbf{y}^n \rightarrow \hat{\mathbf{y}}$, and define $\hat{\sigma}$ as in the statement. If $\hat{\sigma}_t(s) > 0$ for some $s \in \mathcal{S}_t$, then $\hat{y}_{t,s} > 0$, so that, for all n large we have $y_{t,s}^n > 0$ and there must be a player $i^n \in \mathcal{N}^n$ with type t and $\hat{\sigma}_{i^n}^n(s) > 0$. The equilibrium condition for i^n implies that

$$\forall s' \in \mathcal{S}_t \quad E_{\hat{\sigma}^n}[C(Y^{-i^n,n} \mid t, s)] \leq E_{\hat{\sigma}^n}[C(Y^{-i^n,n} \mid t, s')]. \quad (5.8)$$

As in Proposition 4.1, it follows that the variables $Y^{-i^n,n}$ converge in total variation, and hence in distribution, to a random vector Y with independent Poisson components $Y_{t,s} \sim \text{Poisson}(\hat{y}_{t,s})$. Letting $n \rightarrow \infty$ in Eq. (5.8) we obtain Eq. (5.3), from which the result follows. $\square$

Remark 5.2. Theorem 5.2 remains valid when some of the demands converge to zero $d_t^n \rightarrow d_t = 0$, by considering the Poisson game restricted to the nontrivial types with $d_t > 0$.

6. SUMMARY AND PERSPECTIVES

In this paper we studied the convergence of equilibria of atomic-unsplittable congestion games with an increasing number of players, towards a WE for a limiting nonatomic game. For the case where players have vanishing weights, the random loads (Y^n, X^n) at a mixed equilibrium in the

finite games were shown to converge in distribution to constant loads $(\hat{\mathbf{y}}, \hat{\mathbf{x}})$, which are a Wardrop equilibrium of the nonatomic congestion game. In contrast, if players have a fixed unit weight but are present in the game with vanishing probabilities, then (Y^n, X^n) converge in total variation to Poisson variables (Y, X) , whose expected values $(\hat{\mathbf{y}}, \hat{\mathbf{x}})$ are again characterized as a WE for a nonatomic congestion game with auxiliary cost functions. In the latter case, the limit variables (Y, X) can also be interpreted in terms of an equilibrium for a Poisson game in the sense of Myerson (1998b).

These convergence results were completed by showing that in both frameworks the price of anarchy and the price of stability converge to the price of anarchy of the limit game. Under additional conditions we also established explicit estimates for the distance between the random loads (Y^n, X^n) and their limits $(\hat{\mathbf{y}}, \hat{\mathbf{x}})$ and (Y, X) , respectively.

We did not address the combined case in which both the weights and the presence probabilities vary across players. Such situations may be relevant for routing games where cars and trucks have a different impact on traffic, or in the presence of heterogeneous drivers that may be slower or faster inducing more or less congestion. Other settings in which players are naturally heterogeneous arise in telecommunications, where packets come in different sizes, and in processor sharing, where tasks arriving to a server have different workloads. In such cases, one might still expect to obtain a limit game which is likely to yield a *weighted nonatomic game*, possibly involving weighted sums of Poisson distributions.

Another direction not explored in this paper concerns the case of oligopolistic competition in which some players, e.g., TomTom, Waze, FedEx, UPS, etc. may control a nonnegligible fraction of the demand, while simultaneously another fraction of the demand behaves as individual selfish players. A natural conjecture is that in the vanishing weight limit one may converge to a composite game as those studied by Cominetti et al. (2009) and Sorin and Wan (2016), with co-existence of atomic-splittable and nonatomic players. Similarly, in the case of fixed weights and vanishing probabilities one may expect to converge to some form of composite game involving Poisson random variables, which remains to be discovered.

Acknowledgments. This collaboration started in Dagstuhl at the Seminar on Dynamic Traffic Models in Transportation Science in 2018. Roberto Cominetti gratefully acknowledges the support of FONDECYT 1171501 and Proyecto Anillo ACT192094. Marco Scarsini is a member of GNAMPA-INdAM. His work was partially supported by the the INdAM-GNAMPA Project 2019 “Markov chains and games on networks,” and the Italian MIUR PRIN 2017 Project ALGADIMAR “Algorithms, Games, and Digital Markets.” This research also received partial support from the COST action GAMENET.

REFERENCES

- D. J. Adams. 1936. Road traffic considered as a random series. *J. Inst. Civ. Engrs.* 4, 1 (1936), 121–130.
- José Antonio Adell and Alberto Lekuona. 2005. Sharp estimates in signed Poisson approximation of Poisson mixtures. *Bernoulli* 11, 1 (2005), 47–65. <https://doi.org/10.3150/bj/1110228242>
- Haris Angelidakis, Dimitris Fotakis, and Thanasis Lianeas. 2013. Stochastic congestion games with risk-averse players. In *Algorithmic Game Theory*. Lecture Notes in Comput. Sci., Vol. 8146. Springer, Heidelberg, 86–97. https://doi.org/10.1007/978-3-642-41392-6_8
- A. D. Barbour and Louis H. Y. Chen (Eds.). 2005. *An introduction to Stein’s method*. Singapore University Press, Singapore. xii+225 pages. <https://doi.org/10.1142/9789812567680>
- A. D. Barbour and Peter Hall. 1984. On the rate of Poisson convergence. *Math. Proc. Cambridge Philos. Soc.* 95, 3 (1984), 473–480. <https://doi.org/10.1017/S0305004100061806>

- A. D. Barbour, Lars Holst, and Svante Janson. 1992. *Poisson Approximation*. The Clarendon Press, Oxford University Press, New York. x+277 pages.
- Martin J. Beckmann, C.B McGuire, and Christopher B Winsten. 1956. *Studies in the Economics of Transportation*. Yale University Press, New Haven, CT.
- Pierre Bernhard and Marc Deschamps. 2017. On dynamic games with randomly arriving players. *Dyn. Games Appl.* 7, 3 (2017), 360–385. <https://doi.org/10.1007/s13235-016-0197-z>
- I. S. Borisov and P. S. Ruzankin. 2002. Poisson approximation for expectations of unbounded functions of independent random variables. *Ann. Probab.* 30, 4 (2002), 1657–1680. <https://doi.org/10.1214/aop/1039548369>
- D.J. Buckley. 1967. Road traffic counting distributions. *Transportation Research* 1, 2 (1967), 105 – 116. [https://doi.org/10.1016/0041-1647\(67\)90166-9](https://doi.org/10.1016/0041-1647(67)90166-9)
- Po-An Chen, Bart De Keijzer, David Kempe, and Guido Schäfer. 2014. Altruism and its impact on the price of anarchy. *ACM Transactions on Economics and Computation (TEAC)* 2, 4 (2014), 1–45.
- R. Cominetti, J.R. Correa, and N.E. Stier-Moses. 2009. The Impact of Oligopolistic Competition in Networks. *Operations Research* 57, 6 (2009), 1421–1437. <https://doi.org/10.1287/opre.1080.0653>
- R. Cominetti, M. Scarsini, M. Schröder, and N. Stier-Moses. 2019. Price of anarchy in stochastic atomic congestion games with affine costs. (2019). <http://arxiv.org/abs/1903.03309> arXiv 1903.03309.
- José Correa, Ruben Hoeksma, and Marc Schröder. 2019. Network congestion games are robust to variable demand. *Transportation Res. Part B* 119 (2019), 69–78.
- Francesco De Sinopoli, Claudia Meroni, and Carlos Pimienta. 2014. Strategic stability in Poisson games. *J. Econom. Theory* 153 (2014), 46–63. <https://doi.org/10.1016/j.jet.2014.05.005>
- Francesco De Sinopoli and Carlos Pimienta. 2009. Undominated (and) perfect equilibria in Poisson games. *Games Econom. Behav.* 66, 2 (2009), 775–784. <https://doi.org/10.1016/j.geb.2008.09.029>
- Lili Du and Siyuan Gong. 2016. Stochastic Poisson game for an online decentralized and coordinated parking mechanism. *Transportation Research Part B: Methodological* 87 (2016), 44–63. <https://doi.org/10.1016/j.trb.2016.02.006>
- Michal Feldman, Nicole Immorlica, Brendan Lucier, Tim Roughgarden, and Vasilis Syrgkanis. 2016. The price of anarchy in large games. In *STOC’16—Proceedings of the 48th Annual ACM SIGACT Symposium on Theory of Computing*. ACM, New York, 963–976.
- Dimitris Fotakis, Spyros Kontogiannis, and Paul Spirakis. 2005. Selfish unsplittable flows. *Theoret. Comput. Sci.* 348, 2-3 (2005), 226–239. <https://doi.org/10.1016/j.tcs.2005.09.024>
- Martin Gairing, Burkhard Monien, and Karsten Tiemann. 2008. Selfish routing with incomplete information. *Theory Comput. Syst.* 42, 1 (2008), 91–130. <https://doi.org/10.1007/s00224-007-9015-8>
- Tobias Harks and Max Klimm. 2012. On the existence of pure Nash equilibria in weighted congestion games. *Math. Oper. Res.* 37, 3 (2012), 419–436. <https://doi.org/10.1287/moor.1120.0543>
- Tobias Harks, Max Klimm, and Rolf H. Möhring. 2011. Characterizing the existence of potential functions in weighted congestion games. *Theory Comput. Syst.* 49, 1 (2011), 46–70. <https://doi.org/10.1007/s00224-011-9315-x>
- Refael Hassin, Irit Nowik, and Yair Y. Shaki. 2018. On the price of anarchy in a single-server queue with heterogeneous service valuations induced by travel costs. *European J. Oper. Res.* 265, 2 (2018), 580–588. <https://doi.org/10.1016/j.ejor.2017.08.020>
- A. Haurie and P. Marcotte. 1985. On the relationship between Nash-Cournot and Wardrop equilibria. *Networks* 15, 3 (1985), 295–308. <https://doi.org/10.1002/net.3230150303>
- Paulin Jacquot and Cheng Wan. 2018. Routing game on parallel networks: the convergence of atomic to nonatomic. (2018). <http://arxiv.org/abs/1804.03081>
- Paulin Jacquot and Cheng Wan. 2019. Nonatomic aggregative games with infinitely many types. (2019). <https://arxiv.org/abs/1712.08622>
- Ioannis Kordonis and George P. Papavassilopoulos. 2015. LQ Nash games with random entrance: an infinite horizon major player and minor players of finite horizons. *IEEE Trans. Automat. Control* 60, 6 (2015), 1486–1500.

- <https://doi.org/10.1109/TAC.2015.2396642>
- Elias Koutsoupias and Christos Papadimitriou. 1999. Worst-case equilibria. In *STACS 99 (Trier)*. Lecture Notes in Comput. Sci., Vol. 1563. Springer, Berlin, 404–413. https://doi.org/10.1007/3-540-49116-3_38
- Wooyoung Lim and Alexander Matros. 2009. Contests with a stochastic number of players. *Games Econom. Behav.* 67, 2 (2009), 584–597. <https://doi.org/10.1016/j.geb.2009.03.012>
- Alan J. Mayne. 1954. Some further results in the theory of pedestrians and road traffic. *Biometrika* 41 (1954), 375–389. <https://doi.org/10.1093/biomet/41.3-4.375>
- Claudia Meroni and Carlos Pimienta. 2017. The structure of Nash equilibria in Poisson games. *J. Econom. Theory* 169 (2017), 128–144. <https://doi.org/10.1016/j.jet.2017.02.003>
- Igal Milchtaich. 2000. Generic uniqueness of equilibrium in large crowding games. *Math. Oper. Res.* 25, 3 (2000), 349–364. <https://doi.org/10.1287/moor.25.3.349.12220>
- Igal Milchtaich. 2004. Random-player games. *Games Econom. Behav.* 47, 2 (2004), 353–388. <https://doi.org/10.1016/j.geb.2003.05.002>
- Alan J. Miller. 1970. An empirical model for multilane road traffic. *Transportation Sci.* 4, 2 (1970), 164–186. <https://doi.org/10.1287/trsc.4.2.164> arXiv:<https://doi.org/10.1287/trsc.4.2.164>
- D. Monderer and L. S. Shapley. 1996. Potential games. *Games Econom. Behav.* 14, 1 (1996), 124–143. <https://doi.org/10.1006/game.1996.0044>
- R. B. Myerson. 1998a. Extended Poisson games and the Condorcet jury theorem. *Games Econom. Behav.* 25, 1 (1998), 111–131. <https://doi.org/10.1006/game.1997.0610>
- R. B. Myerson. 1998b. Population uncertainty and Poisson games. *Internat. J. Game Theory* 27, 3 (1998), 375–392. <https://doi.org/10.1007/s001820050079>
- R. B. Myerson. 2000. Large Poisson games. *J. Econom. Theory* 94, 1 (2000), 7–45. <https://doi.org/10.1006/jeth.1998.2453>
- R. B. Myerson. 2002. Comparison of scoring rules in Poisson voting games. *J. Econom. Theory* 103, 1 (2002), 219–251. <https://doi.org/10.1006/jeth.2001.2830> Political science.
- John F. Nash, Jr. 1950. Equilibrium points in n -person games. *Proc. Nat. Acad. Sci. U.S.A.* 36 (1950), 48–49. <https://doi.org/10.1073/pnas.36.1.48>
- Evdokia Nikolova and Nicolás E. Stier-Moses. 2014. A mean-risk model for the traffic assignment problem with stochastic travel times. *Oper. Res.* 62, 2 (2014), 366–382. <https://doi.org/10.1287/opre.2013.1246>
- S. Y. Novak. 2019. Poisson approximation. *Probab. Surv.* 16 (2019), 228–276. <https://doi.org/10.1214/18-PS318>
- Robert M. Oliver. 1961. A traffic counting distribution. *Operations Res.* 9 (1961), 802–810. <https://doi.org/10.1287/opre.9.6.802>
- Dario Paccagnan, Basilio Gentile, Francesca Parise, Maryam Kamgarpour, and John Lygeros. 2019. Nash and Wardrop equilibria in aggregative games with coupling constraints. *IEEE Trans. Automat. Control* 64, 4 (2019), 1373–1388.
- Georgios Piliouras, Evdokia Nikolova, and Jeff S. Shamma. 2016. Risk Sensitivity of Price of Anarchy Under Uncertainty. *ACM Trans. Econ. Comput.* 5, 1, Article 5 (Oct. 2016), 27 pages. <https://doi.org/10.1145/2930956>
- R. W. Rosenthal. 1973. A class of games possessing pure-strategy Nash equilibria. *Internat. J. Game Theory* 2 (1973), 65–67. <http://dx.doi.org/10.1007/BF01737559>
- Tim Roughgarden. 2002. The price of anarchy is independent of the network topology. In *Proceedings of the Thirty-Fourth Annual ACM Symposium on Theory of Computing*. ACM, New York, 428–437. <https://doi.org/10.1145/509907.509971>
- Tim Roughgarden. 2003. The price of anarchy is independent of the network topology. *J. Comput. System Sci.* 67, 2 (2003), 341–364.
- Tim Roughgarden. 2015. The price of anarchy in games of incomplete information. *ACM Trans. Econ. Comput.* 3, 1 (2015), Art. 6, 1–20. <https://doi.org/10.1145/2737816>

- Sylvain Sorin and Cheng Wan. 2016. Finite composite games: equilibria and dynamics. *J. Dyn. Games* 3, 1 (2016), 101–120. <https://doi.org/10.3934/jdg.2016005>
- Shaler Stidham. 2014. The price of anarchy for a network of queues in heavy traffic. In *Essays in Production, Project Planning and Scheduling: A Festschrift in Honor of Salah Elmaghraby*, P. Simin Pulat, Subhash C. Sarin, and Reha Uzsoy (Eds.). Springer, Boston, MA, 91–121.
- Chenlan Wang, Xuan Vinh Doan, and Bo Chen. 2014. Price of anarchy for non-atomic congestion games with stochastic demands. *Transportation Res. Part B* 70 (2014), 90–111. <https://doi.org/10.1016/j.trb.2014.08.009>
- J. G. Wardrop. 1952. Some theoretical aspects of road traffic research. In *Proceedings of the Institute of Civil Engineers, Part II*, Vol. 1. 325–378.

APPENDIX A. POISSON APPROXIMATION

This appendix collects some known facts on the Poisson approximation for sums of Bernoulli random variables. The main results are taken from [Adell and Lekuona \(2005\)](#), [Barbour and Hall \(1984\)](#), and [Borisov and Ruzankin \(2002\)](#), suitably adapted to our goals. We recall that $X \sim \text{Poisson}(x)$ with parameter $x \geq 0$ iff

$$P(X = k) = e^{-x} \frac{x^k}{k!} \quad \forall k \in \mathbb{N}. \quad (\text{A.1})$$

As usual we denote $\mathcal{L}(X)$ the law of X .

Two Poisson variables $X \sim \text{Poisson}(x)$ and $Y \sim \text{Poisson}(y)$ are close when $x \approx y$. More precisely, their total variation distance can be estimated as (see [Adell and Lekuona, 2005](#))

$$\rho_{\text{TV}}(\mathcal{L}(X), \mathcal{L}(Y)) \leq 1 - e^{-|x-y|} \leq |x-y|, \quad (\text{A.2})$$

where the total variation distance ρ_{TV} is defined as in [Eq. \(4.4\)](#).

A.1. Poisson distributions and expected values. Given a function $h : \mathbb{N} \rightarrow \mathbb{R}$, for each $X \sim \text{Poisson}(x)$ with $E[|h(X)|] < \infty$, we define

$$\check{h}(x) = E[h(X)] = \sum_{k=0}^{\infty} h(k) e^{-x} \frac{x^k}{k!}. \quad (\text{A.3})$$

We also define $h \mapsto \Delta h$ the operator that takes the function h into $\Delta h(k) = h(k+1) - h(k)$. Below we state two useful facts that are used in the sequel.

Proposition A.1. *Let $X \sim \text{Poisson}(x)$. Then*

- (a) $E[|h(Y)|] \leq e^{x-y} E[|h(X)|]$ for each $Y \sim \text{Poisson}(y)$ with $y \leq x$.
- (b) For each $j = 1, 2, \dots$ we have²

$$E[|\Delta^j h(X)|] < \infty \Leftrightarrow E[|h(X+j)|] < \infty \Leftrightarrow E[X^j |h(X)|] < \infty. \quad (\text{A.4})$$

Proof. Property (a) is just the monotonicity of $x \mapsto e^x \cdot E[|h(X)|] = \sum_{k=0}^{\infty} |h(k)| \frac{x^k}{k!}$, while (b) can be found in [Borisov and Ruzankin \(2002, Proposition 1\)](#). $\square$

Proposition A.2. *Let $h : \mathbb{N} \rightarrow \mathbb{R}$ and $V \sim \text{Poisson}(\alpha)$.*

- (a) *If $E[|h(V)|] < \infty$ then $\check{h}(x)$ is well defined and continuous for $x \in [0, \alpha]$.*
- (b) *If $E[|\Delta^j h(V)|] < \infty$ for some $j \in \mathbb{N}$ then $\check{h}$ is of class C^j on $[0, \alpha]$ and $\check{h}^{(j)}(x) = E[\Delta^j h(X)]$.*

Proof. (a) It suffices to note that the series

$$f(x) = e^x \check{h}(x) = \sum_{k=0}^{\infty} h(k) \frac{x^k}{k!} \quad (\text{A.5})$$

is well defined and continuous. This follows since the partial sums $f_n(x) = \sum_{k=0}^n h(k) x^k / k!$ converge uniformly to $f(x)$. Indeed,

$$\sup_{x \in [0, \alpha]} |f(x) - f_n(x)| \leq \sup_{x \in [0, \alpha]} \sum_{k=n+1}^{\infty} |h(k)| \frac{x^k}{k!} \leq \sum_{k=n+1}^{\infty} |h(k)| \frac{\alpha^k}{k!}, \quad (\text{A.6})$$

where the latter tail of the series tends to 0 as $n \rightarrow \infty$ since $E[|h(V)|] < \infty$.

²Note that if $E[X^j |h(X)|] < \infty$ holds for a certain j , it also holds for $j' = 1, \dots, j$.

(b) Consider first the case $j = 1$. We note that the derivatives

$$f'_n(x) = \sum_{k=0}^{n-1} h(k+1) \frac{x^k}{k!} \quad (\text{A.7})$$

converge uniformly towards $g(x) = \sum_{k=0}^{\infty} h(k+1) x^k / k!$. This follows from part (a) since by [Proposition A.1\(b\)](#) with $j = 1$ we have $E[|h(V+1)|] < \infty$. Hence f is C^1 with $f'(x) = g(x)$ and then $\check{h}(x) = e^{-x} f(x)$ is also C^1 with

$$\check{h}'(x) = e^{-x} f'(x) - e^{-x} f(x) = \sum_{k=0}^{\infty} (h(k+1) - h(k)) e^{-x} \frac{x^k}{k!} = E[\Delta h(X)]. \quad (\text{A.8})$$

This establishes the case $j = 1$. Applying this property to Δh we obtain the result for $j = 2$, and then the cases $j = 3, 4, \dots$ follow by induction. $\square$

Corollary A.3. *Let $V \sim \text{Poisson}(\alpha)$ and suppose that $E[|\Delta^2 h(V)|] \leq \nu < \infty$.*

- (a) *For all $x \in [0, \alpha]$ we have $|\check{h}'(x)| \leq (e^\alpha - 1)\nu + |h(1) - h(0)|$.*
- (b) *If $h(\cdot)$ is weakly increasing then for all $x \in [0, \alpha]$ we have $\check{h}'(x) \geq 0$, with strict inequality except when $h(\cdot)$ is constant.*

Proof.

(a) Combining [Propositions A.2\(b\)](#) and [A.1\(a\)](#) we get $|\check{h}''(x)| \leq e^{\alpha-x}\nu$, so that by integration it follows that $|\check{h}'(x) - \check{h}'(0)| \leq (e^\alpha - 1)\nu$, and we conclude since $\check{h}'(0) = h(1) - h(0)$.

(b) This follows from [Proposition A.2\(b\)](#). $\square$

A.2. Poisson approximation for sums of Bernoulli random variables. Let $S = X_1 + \dots + X_n$ be a sum of independent Bernoulli random variables with $P(X_i = 1) = p_i \in (0, 1)$, and denote $x = E[S] = p_1 + \dots + p_n$. Consider a Poisson variable $X \sim \text{Poisson}(x)$ with the same expectation. The following result shows that S and X are close when the p_i 's are small.

Theorem A.4. *Let $p = \max\{p_1, \dots, p_n\}$. Then*

- (a) *The following double inequality holds:*

$$\rho_{\text{TV}}(\mathcal{L}(S), \mathcal{L}(X)) \leq (1 - e^{-x}) x^{-1} \sum_{i=1}^n p_i^2 \leq p.$$

- (b) *If $h : \mathbb{N} \rightarrow \mathbb{R}$ is such that $E[|\Delta^2 h(X)|] \leq \nu < \infty$, then*

$$|E[h(S)] - E[h(X)]| \leq \frac{x\nu}{2} \frac{p e^p}{(1-p)^2}. \quad (\text{A.9})$$

Proof. These properties follow from [Barbour and Hall \(1984, Theorem 1\)](#) and [Borisov and Ruzankin \(2002, Corollary 4\)](#), respectively. $\square$

APPENDIX B. LIST OF SYMBOLS

$\text{BNE}(\Gamma_{\text{B}}^n)$	set of Bayesian Nash equilibria of the game Γ_{B}^n
c_e	cost function of edge e
$\check{c}_e$	cost function of edge e in the Poisson limit game, defined in Eq. (4.10)
$\mathbf{d}$	demand vector
d_t	demand of type t
d_t^n	aggregate demand of type t in the games Γ_{W}^n and Γ_{B}^n
d_{tot}	total demand
D	random demand
$\mathcal{E}$	set of resources
$\mathbb{E}_{\sigma}$	expectation induced by σ
$\text{Eq}(\Gamma^{\infty})$	equilibrium cost of Γ^{∞}
$\text{ESC}(\sigma)$	expected social cost of σ , defined in Eq. (2.19)
$\mathcal{F}(\mathbf{d})$	feasible pairs for demand $\mathbf{d}$
$\mathcal{L}(X)$	law of the random variable X
$\text{MNE}(\Gamma)$	set of mixed Nash equilibria of Γ
N_e	$\sum_{i:e \in S_i} U_i$, i.e., random number of players who use resource e
N_t	random number of players of type t in a Poisson game
$\mathbf{N}$	$(N_t)_{t \in \mathcal{T}}$
$\mathcal{N}$	set of players
$\text{Opt}(\Gamma^{\infty})$	optimum social cost in Γ^{∞}
$\text{Opt}(\Gamma)$	optimum expected social cost in Γ
$\mathbb{P}_{\sigma}$	probability measure induced by σ
$\text{PNE}(\Gamma)$	set of pure Nash equilibria of Γ
PoA	price of anarchy
PoS	price of stability
r_i	probability of player i being active
r^n	$\max_{i \in \mathcal{N}^n} r_i^n$, defined in Eq. (4.2b)
U_i	indicator of player i being active
$U_{i,e}$	$U_i \mathbb{1}_{\{e \in S_i\}}$
$\mathcal{S}_t$	set of strategies for type t
$\mathbf{s}$	strategy profile
s_i	strategy of player i
S_i	random strategy of player i
$\text{SC}(\mathbf{y}, \mathbf{x})$	social cost of $(\mathbf{y}, \mathbf{x})$, defined in Eq. (2.17)
$\mathcal{T}$	set of types
t_i	type of player i
w_i	weight of player i
w^n	$\max_{i \in \mathcal{N}^n} w_i^n$, defined in Eq. (3.2b)
$\mathbf{x}$	load vector
x_e	load of edge e
X_e	random load of edge e
$X_{i,e}$	$w_i + \sum_{j \neq i} w_j \mathbb{1}_{\{e \in S_j\}}$, defined in Eq. (2.8)
$\tilde{\mathbf{x}}$	optimum load in Γ^{∞}

$(\widehat{\mathbf{y}}, \widehat{\mathbf{x}})$	equilibrium flow-load pair
$\mathbf{y}$	flow vector
$y_{t,s}$	flow of type t on strategy s
$Y_{t,s}$	random flow of type t on strategy s
$Z_{i,e}$	$\sum_{j \neq i} U_{j,e}$, defined in Eq. (2.14)
α	upper bound on the demand
β	lower bound on c'_e
γ	upper bound on $ c''_e $
Γ	game
Γ_B	Bernoulli congestion game
Γ_P	game with population uncertainty
Γ_W	weighted congestion game
Γ^∞	nonatomic congestion game
Δ	difference operator
$\Delta(\mathcal{S}_t)$	simplex of probability measures on $\mathcal{S}_t$
ζ	Lipschitz constant
Θ	$\sqrt{\alpha/4} + \sqrt{2\alpha\kappa(\zeta + \gamma\alpha/4)/\beta}$, defined in Theorem 3.4
$\check{\Theta}$	$\sqrt{2\alpha\kappa/\beta}$, defined in Theorem 4.5
κ	cardinality of the largest feasible strategy $s \in \cup_{t \in \mathcal{T}} \mathcal{S}_t$, defined in Theorem 3.4
$\Lambda(r)$	$\frac{\alpha v}{2} \frac{r e^r}{(1-r)^2} + \zeta r$, defined in Eq. (4.12)
μ	joint probability distribution of the number of players in a Poisson game
Ξ	$\sqrt{2C/\beta}$, defined in Proposition 3.3 with $C = \max_s \sum_{e \in s} c_e(\alpha)$
$\check{\Xi}$	$\sqrt{2C/\beta}$, defined in Corollary 4.6 with $C = \max_s \sum_{e \in s} \check{c}_e(\alpha)$
σ	mixed strategy profile
σ_i	mixed strategy of player i
$\sigma_{i,e}$	$P_\sigma(e \in S_i)$, defined in Eq. (2.5)
$\widehat{\sigma}$	mixed Nash equilibrium
$\widetilde{\sigma}$	optimum mixed strategy in Γ
Σ	set of mixed strategy profiles

[‡] FACULTAD DE INGENIERÍA Y CIENCIAS, UNIVERSIDAD ADOLFO IBÁÑEZ, SANTIAGO, CHILE.
E-mail address: roberto.cominetti@uai.cl

[¶] DIPARTIMENTO DI ECONOMIA E FINANZA, LUISS, VIALE ROMANIA 32, 00197 ROMA, ITALY.
E-mail address: marco.scarsini@luiss.it

* RWTH AACHEN UNIVERSITY, AACHEN, GERMANY.
E-mail address: marc.schroeder@oms.rwth-aachen.de

[§] FACEBOOK CORE DATA SCIENCE, MENLO PARK, USA
E-mail address: nicostier@yahoo.com