

Psychological effect can lead to bistability in epidemics [☆]

Shaoli Wang^{*a}, Xiyan Bai^a

^a*School of Mathematics and Statistics, Henan University, Kaifeng 475001, Henan, PR China*

Abstract

In this paper, we study the psychological effect in a SIS epidemic model. The basic reproduction number is obtained. However, the disease free equilibrium is always asymptotically stable, which doesn't depend on the basic reproduction number. The system has a saddle-node bifurcation appear and displays bistable behavior, which is a new phenomenon in epidemic dynamics and different from the backward bifurcation behavior.

Key words: SIS model, Psychological effect; Saddle-node bifurcation; Bistability behavior

1. Introduction

In classic disease transmission model, the incidence rate is bilinear in the infectious fraction I and the susceptible fraction S . Recently nonlinear incidence functions in epidemic models attracted much attention [1–9, 11–14].

Capasso and Serio [1], Ruan and Wang [5] show the incidence function $g(I)$ can interpret the “psychological” effect: for a very large number of infective individuals the infection force may decrease as the number of infective individuals increases, because in the presence of large number of infective the population may tend to reduce the number of contacts per unit time. Xiao and Ruan [8] studied an epidemic model with nonmonotonic incidence rate, which describes the psychological effect of certain serious diseases on the community when the number of infectives is getting larger. Lu et al. [9] provided a more reasonable incidence function, which first increases to a maximum when a new infectious disease emerges or an old infectious disease reemerges, then decreases due to psychological effect, and eventually tends to a saturation level due to crowding effect.

In this paper we will discuss the psychological effect in epidemics in a different way. The general SIS epidemic model takes the following form

$$\begin{cases} \frac{dS}{dt} = b - dS - k_1 S g(I) + \gamma I, \\ \frac{dI}{dt} = k_1 S g(I) - (d + \mu + \gamma)I, \end{cases} \quad (1.1)$$

where b is natural birth rate, d is natural decay rate, k_1 is transmission rate for naive susceptible, μ is disease related death rate, γ is the rate of infective individuals lose immunity and move into susceptible compartment. For the incidence rate $g(I)$, we have following cases, some are based on the work of Andrews [10].

- (I) If $g(I) = I$, then system (1.1) is the classic SIS model;
- (II) If we choose $g(I)S$ as following

$$g(I)S = \frac{S}{1 + \frac{k_s}{I}} = \frac{SI}{k_s + I},$$

which is the saturated incidence rate in epidemic models [1, 11, 12]. Here k_s is the saturation constant of infected population concentration;

[☆]This work is supported by NSFC (No.U1604180), Key Scientific and Technological Research Projects in Henan Province (No.192102310089), Foundation of Henan Educational Committee (No.19A110009) and Grant of Bioinformatics Center of Henan University (No.2019YLXKJC02).

*Corresponding author.

Email address: wslheda@163.com (Shaoli Wang)

(III) If

$$g(I)S = \frac{S}{1 + \frac{k_s}{I} + \frac{I}{k_i}} = \frac{k_i S I}{k_i k_s + k_i I + I^2},$$

which is the nonmonotone incidence rate, where k_i is the inhibition constant of infected population concentration. The special case is

$$g(I)S = \frac{S}{\frac{k_s}{I} + \frac{I}{k_i}} = \frac{k_i S I}{k_i k_s + I^2} = \frac{\frac{1}{k_s} S I}{1 + \frac{1}{k_i k_s} I^2},$$

which was studied by Xiao and Ruan [8].

(IV) If we choose $g(I)S$ as following

$$g(I)S = \frac{S I}{1 + \frac{k_s}{I} + \frac{I}{k_i}} = \frac{k_i S I^2}{k_i k_s + k_i I + I^2},$$

which is the generalized nonmonotone and saturated incidence rate [9]. Especially, if

$$g(I)S = \frac{S I}{\frac{k_s}{I} + \frac{I}{k_i}} = \frac{k_i S I^2}{k_i k_s + I^2} = \frac{\frac{1}{k_s} S I^2}{1 + \frac{1}{k_i k_s} I^2},$$

which was studied by Ruan and Wang [5], and Tang et al.[6].

(V) If we choose $g(I)S$ as following

$$g(I)S = \frac{S I^r}{1 + \frac{k_s}{I^p} + \frac{I^{p'}}{k_i}} = \frac{k_i S I^{p+r}}{k_i k_s + k_i I^p + I^q},$$

which is the general incidence rate. Here r, p, p' and $q = p + p'$ are nonnegative. The special case is

$$g(I)S = \frac{S I^r}{\frac{k_s}{I^p} + \frac{I^{p'}}{k_i}} = \frac{k_i S I^{p+r}}{k_i k_s + I^q} = \frac{\frac{1}{k_s} S I^{p+r}}{1 + \frac{1}{k_i k_s} I^q},$$

which was studied by a number of authors [3, 4, 13, 14].

In this paper, we only consider the case $p = 1, q = 2, r = 1$. Denote $k_1 k_i = k, k_i k_s = \alpha, k_i = \beta$, then system (1.1) can be written by following model

$$\begin{cases} \frac{dS}{dt} = b - dS - \frac{kSI^2}{\alpha + \beta I + I^2} + \gamma I, \\ \frac{dI}{dt} = \frac{kSI^2}{\alpha + \beta I + I^2} - (\mu + \gamma + d)I. \end{cases} \quad (1.2)$$

Here, β is the inhibition psychological effect constant of infected population and α, β are positive.

2. Equilibria and thresholds

It can be verified that the nonnegative orthant $\mathbb{R}_2^+ = \{(S, I) : S \geq 0, I > 0\}$ is positively invariant with respect to system (1.2) and the model is well posed.

Denote

$$R_0 = \frac{bk}{\beta d(\mu + \gamma + d)} = \frac{b}{d} \cdot k_1 \cdot \frac{1}{\mu + \gamma + d}$$

be the basic reproduction number, which determining whether or not the disease dies out in classical SIS epidemic models. We also denote

$$R_c = R_0 - \frac{2}{\beta} \sqrt{\alpha \left(1 + \frac{k(\mu + d)}{d(\mu + \gamma + d)}\right)},$$

and

$$R_{cc} = R_0 + \frac{2}{\beta} \sqrt{\alpha \left(1 + \frac{k(\mu + d)}{d(\mu + \gamma + d)}\right)}.$$

It is easy to see that $R_c < R_0 < R_{cc}$.

(i) System (1.2) always has a disease-free equilibrium $E_0 = (\frac{b}{d}, 0)$.

(ii) To obtain the positive equilibria of system (1.2), we solve the following equations:

$$\begin{aligned} b - dS - \frac{kSI^2}{\alpha + \beta I + I^2} + \gamma I &= 0, \\ \frac{kSI}{\alpha + \beta I + I^2} - (\mu + \gamma + d) &= 0. \end{aligned} \quad (2.1)$$

Solving the first equation of (2.1), we have

$$S = \frac{b - (\mu + d)I}{d},$$

substituting which into the second equation of (2.1) yields

$$AI^2 + BI + \alpha = 0, \quad (2.2)$$

where

$$\begin{aligned} A &= 1 + \frac{k(\mu + d)}{d(\mu + \gamma + d)}, \\ B &= \beta(1 - R_0). \end{aligned}$$

Denote $\Delta = B^2 - 4A\alpha$. If $\Delta > 0$, then $R_{cc} < 1$ or $R_c > 1$. If $B < 0$, then $R_0 > 1$. When $R_c > 1$, equation (2.2) has two positive roots:

$$I_{\pm}^* = \frac{-B \pm \sqrt{\Delta}}{2A}.$$

Theorem 2.1 (i) System (1.2) always has a disease-free equilibrium E_0 ;

(ii) If $R_c > 1$, system (1.2) also has two positive equilibria $E_+(S_+, I_+^*)$, $E_-(S_-, I_-^*)$, where

$$\begin{aligned} S_+^* &= \frac{b - (\mu + d)I_+^*}{d}, I_+^* = \frac{-B + \sqrt{\Delta}}{2A}, \\ S_-^* &= \frac{b - (\mu + d)I_-^*}{d}, I_-^* = \frac{-B - \sqrt{\Delta}}{2A}. \end{aligned}$$

The existence of positive equilibria are summarized in Table 1.

Table 1: The existence of the positive equilibria of system (1.2)

	$R_c < 1$	$R_c > 1$
E_0	exist	exist
E_+^*	—	exist
E_-^*	—	exist

3. Stability analysis

Let \tilde{E} be any arbitrary equilibrium of system (1.2). The Jacobian matrix associated with system (1.2) is

$$\mathcal{J}_{\tilde{E}} = \begin{bmatrix} -d - \frac{k\tilde{I}^2}{\alpha + \beta\tilde{I} + \tilde{I}^2} & \gamma - \frac{k\tilde{S}\tilde{I}(2\alpha + \beta\tilde{I})}{(\alpha + \beta\tilde{I} + \tilde{I}^2)^2} \\ \frac{k\tilde{I}^2}{\alpha + \beta\tilde{I} + \tilde{I}^2} & \frac{k\tilde{S}\tilde{I}(2\alpha + \beta\tilde{I})}{(\alpha + \beta\tilde{I} + \tilde{I}^2)^2} - (\mu + \gamma + d) \end{bmatrix}.$$

The characteristic equation of system (1.2) at \tilde{E} is $|\lambda I - \mathcal{J}_{\tilde{E}}| = 0$.

3.1. Stability analysis of the disease-free equilibrium

Theorem 3.1 The disease-free equilibrium E_0 of system (1.2) is always locally asymptotically stable.

Proof. The characteristic equation of system of (1.2) at the disease-free equilibrium E_0 is obtained as

$$(\lambda + d)(\lambda + \mu + \gamma + d) = 0.$$

The characteristic polynomial has two roots $-d$, $-(\mu + \gamma + d)$. Since the two roots are all negative, the disease-free equilibrium E_0 of system (1.2) is locally asymptotically stable. \square

3.2. Stability analysis of positive equilibria

Theorem 3.2 If $R_c > 1$, $a_1 > 0$, system (1.2) has two positive equilibria E_+^* and E_-^* , where E_+^* is a locally asymptotically stable and E_-^* is unstable.

Proof. Denote an arbitrary positive equilibrium of system (1.2) as E^* . The characteristic equation of the system (1.2) at the arbitrary positive equilibrium E^* is obtained as

$$\lambda^2 + a_1\lambda + a_2 = 0,$$

where

$$\begin{aligned} a_1 &= \mu + \gamma + 2d + \frac{k(I^*)^2}{\alpha + \beta I^* + (I^*)^2} - \frac{(\mu + \gamma + d)(2\alpha + \beta I^*)}{\alpha + \beta I^* + (I^*)^2}, \\ a_2 &= d(\mu + \gamma + d) + (\mu + \gamma + d) \frac{k(I^*)^2}{\alpha + \beta I^* + (I^*)^2} - d \frac{(\mu + \gamma + d)(2\alpha + \beta I^*)}{\alpha + \beta I^* + (I^*)^2} - \gamma \frac{k(I^*)^2}{\alpha + \beta I^* + (I^*)^2}. \end{aligned}$$

(i) For equilibrium E_+^* , we have

$$\begin{aligned} & d(I_+^*)^2 + k(I_+^*)^2 - d\alpha - \gamma \frac{k(I_+^*)^2}{(d+\delta)(d+\mu)}, \\ &= d\left(1 + \frac{k}{d}\left(1 - \frac{\gamma}{\mu + \gamma + d}\right)\right)(I_+^*)^2 - d\alpha, \\ &= d\left(1 + \frac{k(\mu + d)}{d(\mu + \gamma + d)}\right) \frac{(\beta(R_0 - 1) + \sqrt{\Delta})^2}{4A^2} - d\alpha, \\ &= \frac{d(\beta(R_0 - 1) + \sqrt{\Delta})^2}{4A} - d\alpha. \end{aligned}$$

It follows from

$$\frac{d(\beta(R_0 - 1) + \sqrt{\Delta})^2}{4A} - d\alpha = \frac{d\Delta + d\beta(R_0 - 1)\sqrt{\Delta}}{2A}$$

that $\frac{d(\beta(R_0 - 1) + \sqrt{\Delta})^2}{4A} - d\alpha > 0$. Then,

$$\begin{aligned} & \frac{d(\beta(R_0 - 1) + \sqrt{\Delta})^2}{4A} - d\alpha > 0, \\ \Leftrightarrow & d + \frac{k(I_+^*)^2}{\alpha + \beta I_+^* + (I_+^*)^2} - \frac{d(2\alpha + \beta I_+^*)}{\alpha + \beta I_+^* + (I_+^*)^2} - \frac{\gamma}{\mu + \gamma + d} \cdot \frac{k(I_+^*)^2}{\alpha + \beta I_+^* + (I_+^*)^2} > 0, \\ \Leftrightarrow & a_2 > 0. \end{aligned}$$

Clearly, $a_2 > 0$, and we also have $a_1 > 0$. By the Routh-Hurwitz Criterion, we know that the positive equilibrium E_+^* is a locally asymptotically stable node.

(ii) For equilibrium E_-^* , we have

$$\begin{aligned} & d(I_-^*)^2 + k(I_-^*)^2 - d\alpha - \gamma \frac{k(I_-^*)^2}{(d+\delta)(d+\mu)}, \\ &= d\left(1 + \frac{k}{d}\left(1 - \frac{\gamma}{\mu + \gamma + d}\right)\right)(I_-^*)^2 - d\alpha, \\ &= d\left(1 + \frac{k(\mu + d)}{d(\mu + \gamma + d)}\right) \frac{(\beta(R_0 - 1) - \sqrt{\Delta})^2}{4A^2} - d\alpha, \\ &= \frac{d(\beta(R_0 - 1) - \sqrt{\Delta})^2}{4A} - d\alpha, \\ &< \frac{d[2\beta^2(R_0 - 1)^2 - 8A\alpha - 2(\beta^2(R_0 - 1)^2 - 4A\alpha)]}{4A} = 0. \end{aligned}$$

Thus, $a_2 < 0$. By the Routh-Hurwitz Criterion, we know in this case the positive equilibrium E_-^* is an unstable saddle. \square

Table 2: The stabilities of the equilibria and the behaviors of system (1.2) .

	E_0	E_+^*	E_-^*	System (1.2)
$R_c < 1$	LAS	—	—	Converges to E_0
$R_c > 1$	LAS	LAS	US	Bistable

4. Saddle-node bifurcation

In this section, we discuss the bifurcation behavior of system (1.2). The conditions for saddle-node bifurcation are derived. If $R_c = 1$, system (1.2) undergoes a saddle-node bifurcation. The positive equilibrium E_+^* and E_-^* collide to each other and system (1.2) has a unique instantaneous positive equilibrium \bar{E} . Also one of the eigenvalues of the Jacobian evaluated at the instantaneous positive equilibrium $\bar{E} = (\bar{S}, \bar{I})$ is zero. Here $\bar{S} = \frac{b-(\mu+d)\bar{I}}{d}$, $\bar{I} = \frac{\beta(R_0-1)}{2A}$.

Theorem 4.1 If $R_c = 1$ or $R_0 = 1 + \frac{2}{\beta} \sqrt{\alpha(1 + \frac{k(\mu+d)}{d(\mu+\gamma+d)})} \triangleq R_0^{[sn]}$, system (1.2) undergoes a saddle-node bifurcation around instantaneous positive equilibrium $\bar{E} = (\bar{S}, \bar{I})$.

Proof. Let R_0 be the bifurcation parameter. We use the Sotomayor's theorem to prove that system (1.2) undergoes a saddle-node bifurcation. The Jacobian matrix at the saddle-node must have a zero eigenvalue and two eigenvalues with negative real parts. Let $F = (f_1, f_2)^T$ with

$$\begin{aligned} f_1 &= b - dS - \frac{kSI^2}{\alpha+\beta I+I^2} + \gamma I \\ &= b - dS - R_0 \frac{\beta d(\mu+\gamma+d)SI^2}{b(\alpha+\beta I+I^2)} + \delta I, \\ f_2 &= \frac{kSI^2}{\alpha+\beta I+I^2} - (\mu + \gamma + d)I \\ &= R_0 \frac{\beta d(\mu+\gamma+d)SI^2}{b(\alpha+\beta I+I^2)} - (\mu + \gamma + d)I. \end{aligned}$$

The Jacobian matrix of system (1.2) at \bar{E} is given by

$$\mathcal{J}_{\bar{E}} = \begin{bmatrix} -d - \frac{k\bar{I}^2}{\alpha+\beta\bar{I}+\bar{I}^2} & \gamma - \frac{(\mu+\gamma+d)(2\alpha+\beta\bar{I})}{\alpha+\beta\bar{I}+\bar{I}^2} \\ \frac{k\bar{I}^2}{\alpha+\beta\bar{I}+\bar{I}^2} & \frac{(\mu+\gamma+d)(2\alpha+\beta\bar{I})}{\alpha+\beta\bar{I}+\bar{I}^2} - (\mu + \gamma + d) \end{bmatrix}.$$

The matrix has a simple zero eigenvalue, which requires that $\det(\mathcal{J}_{\bar{E}}) = 0$ at $R_0 = R_0^{[sn]}$. If V and W represent eigenvectors corresponding to the eigenvectors of $\mathcal{J}_{\bar{E}}$ and $\mathcal{J}_{\bar{E}}^T$ corresponding to the zero eigenvalue, respectively, then they are given by

$$\begin{aligned} V &= \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -\frac{\mu+d}{d} \\ 1 \end{bmatrix}, \\ W &= \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 + \frac{(\mu+2d)(\alpha+\beta\bar{I}+\bar{I}^2)}{k\bar{I}^2 + (\mu+\gamma+d)(\alpha-\bar{I}^2)} \end{bmatrix}. \end{aligned}$$

Thus we get

$$\begin{aligned} F_{R_0}(\bar{E}, R_0^{[sn]}) &= \begin{bmatrix} -\frac{\beta d(\mu+\gamma+d)\bar{S}\bar{I}^2}{b(\alpha+\beta\bar{I}+\bar{I}^2)} \\ \frac{\beta d(\mu+\gamma+d)\bar{S}\bar{I}^2}{b(\alpha+\beta\bar{I}+\bar{I}^2)} \end{bmatrix}, \\ D^2 F(\bar{E}, R_0^{[sn]})(V, V) &= \begin{bmatrix} \frac{(\mu+\gamma+d)(\beta\bar{I}^2+4\alpha\bar{I}+\beta\alpha)}{(\alpha+\beta\bar{I}+\bar{I}^2)^2} + 2\frac{k\bar{I}(2\alpha+\beta\bar{I})}{(\alpha+\beta\bar{I}+\bar{I}^2)^2} \cdot \frac{\mu+d}{d} \\ -\frac{(\mu+\gamma+d)(\beta\bar{I}^2+4\alpha\bar{I}+\beta\alpha)}{(\alpha+\beta\bar{I}+\bar{I}^2)^2} - 2\frac{k\bar{I}(2\alpha+\beta\bar{I})}{(\alpha+\beta\bar{I}+\bar{I}^2)^2} \cdot \frac{\mu+d}{d} \end{bmatrix}. \end{aligned}$$

Clearly,

$$W^T F_{R_0}(\bar{E}, R_0^{[sn]}) = \frac{\mu + 2d}{k\bar{I}^2 + (\mu + \gamma + d)(\alpha - \bar{I}^2)} \cdot \frac{\beta d(\mu + \gamma + d)\bar{S}\bar{I}^2}{b} \neq 0,$$

$$W^T D^2 F(\bar{E}, R_0^{[sn]})(V, V) = -\left(\frac{(\mu+\gamma+d)(\beta\bar{I}^2+4\alpha\bar{I}+\beta\alpha)}{\alpha+\beta\bar{I}+\bar{I}^2} + 2\frac{k\bar{I}(2\alpha+\beta\bar{I})}{\alpha+\beta\bar{I}+\bar{I}^2} \cdot \frac{\mu+d}{d}\right) \cdot \frac{\mu+2d}{k\bar{I}^2+(\mu+\gamma+d)(\alpha-\bar{I}^2)} \neq 0.$$

Therefore, from the Sotomayor's theorem, system (1.2) undergoes a saddle-node bifurcation around instantaneous positive equilibrium $\bar{E} = (\bar{S}, \bar{I})$ at $R_0 = R_0^{[sn]}$. Hence, we can conclude that when the parameter a passes from one side of $R_0 = R_0^{[sn]}$ to the other side, the number of positive equilibria of system (1.2) changes from zero to two.

5. Numerical simulations and Discussion

To verify our analytical results, we carry out some numerical simulations. In the following, we fix the parameter values as follows[8, 15]:

$$b = 1, d = 0.12, k = 0.2, \gamma = 0.05, \mu = 0.15, \alpha = 0.5. \quad (5.1)$$

If we choose $\beta = 3$, the thresholds $R_0^{[sn]} \approx 1.73$ and $R_c \approx 1.005$. In this case, we have a saddle-node bifurcation (Figure 1). When $\beta = 2.8$, $R_0 = 1.86$, two equilibria of the model E_+^* and E_0 are stable (Figure 2). If we choose $\beta = 3.5$, such that $R_0 = 1.49$, then we have only one equilibrium E_0 which is stable (Figure 3);




Figure 1: Bistability and saddle-node bifurcation diagram of system (1.2). In this case, $R_0^{[sn]} \approx 1.73$. The system displays two stable equilibria E_0 (the blue solid line at the bottom) and E_+^* (the above blue curve), indicating bistable behaviour. Here, \bar{E} is the saddle point, where the two equilibria converge and display saddle-node bifurcation. The point E_-^* (dashed lines) on the bottom half of the curve is unstable, and the point E_+^* (solid line) on the top half of the curve is stable. Here, $\beta = 3$ and other parameter values are listed in (5.1).

In this paper, we consider a SIS model with psychological effect and performed mathematical studies. We found that the system displays bistable behaviors. System (1.2) admits an disease-free equilibrium E_0 , and two positive equilibria E_+^* and E_-^* . We obtain two thresholds, the basic reproduction number $R_0 = \frac{bk_1}{d(\mu+\gamma+d)}$ and $R_c = \frac{bk}{\beta d(\mu+\gamma+d)} - \frac{2}{\beta} \sqrt{\alpha(1 + \frac{k(\mu+d)}{d(\mu+\gamma+d)})}$. We find that the system always admits a disease free equilibrium E_0 which is always asymptotically stable, indicating that there is no infective in the system and all individuals are susceptible. Thus, if there is no disease, then the uninfected state will remain stable for a long time. When $R_c > 1$, both E_+^* and E_-^* exist, where E_+^* is locally asymptotically stable and E_-^* is unstable, which implies the coexistence of susceptible, infective individuals. Choosing R_0 as the branching parameter, our investigation implies that if $R_c = 1$ or $R_0 = R_0^{[sn]}$ system (1.2) undergoes a saddle-node bifurcation. The positive equilibria E_+^* and E_-^* collide to each other and system (1.2) has the unique instantaneous endemic equilibrium \bar{E} . From the branch diagram in figure 1, we find that when $R_0 > R_0^{[sn]}$, the system has two stable equilibria E_+^* and E_0 appear. The system displays bistable behavior. When $R_0 < R_0^{[sn]}$, the system has only one equilibrium point E_0 , suggesting that infectious diseases will die out eventually.

Castillo-Chavez and Song [16] proposed the backward bifurcation to illustrate that even if the basic reproduction number $R_0 < 1$, disease outbreaks are still possible. The backward bifurcation indicates that the system displays bistable behavior when the bifurcation point $R_c < R_0 < 1$. However, when $R_0 > 1$, the system has only one positive equilibrium point, which is stable, and the disease-free equilibrium point is unstable.

In this paper, we investigated a SIS model with psychological effect. We find that (i) the disease-free equilibrium is always stable. (ii) When $1 < R_0 < R_0^{[sn]}$, the model does not have positive equilibrium point. (iii)




Figure 2: For $\beta = 2.8, R_0 = 1.86$ and other parameter values listed in (5.1), we can see that in the case of different initial values, S, I converge to either E_0 or E_+^* . At this interval, the system display two stable equilibria E_0 and E_+^* , indicating bistable behavior.




Figure 3: For $\beta = 3.5, R_0 = 1.49$ and other parameter values listed in (5.1), we can see that S, I converge to E_0 . Here E_0 is a locally asymptotically stable point.

When $R_0 > R_0^{[sm]}$, the system always display bistability behavior. Our investigation implies that psychological effect is a kind of self-protection behavior of human during the outbreak of a disease. Such self-protection behavior may lead to bistable behavior, i.e., there may or may not be a disease outbreak.

References

- [1] V. Capasso, G. Serio, A generalization of the Kermack–Mckendrick deterministic epidemic model, *Math. Biosci.* 42 (1978) 43–61.
- [2] W. Liu, H.W. Hethcote, S.A. Levin, Dynamical behavior of epidemiological models with nonlinear incidence rates, *J. Math. Biol.* 25 (1987) 359–380.
- [3] W. Liu, H.W. Hethcote, S.A. Levin, Influence of nonlinear incidence rates upon the behavior of SIRS epidemiological models, *J. Math. Biol.* 23 (1986) 187–204.

- [4] H.W. Hethcote, P. van den Driessche, Some epidemiological models with nonlinear incidence, *J. Math. Biol.* 29 (1991) 271–287.
- [5] S.G. Ruan, W.D. Wang, Dynamical behavior of an epidemic model with a nonlinear incidence rate, *J. Differential Equations* 188 (2003) 135–163.
- [6] Y.L. Tang, D.Q. Huang, S.G. Ruan, W.N. Zhang, Coexistence of limit cycles and homoclinic loops in a SIRS model with a nonlinear incidence rate, *SIAM J. Appl. Math.* 69 (2008) 621–639.
- [7] J.H. Li, Z.D. Teng, Bifurcations of an SIRS model with generalized non-monotone incidence rate, *Adv. Differ. Equ-Ny.* (2018) 2018:217.
- [8] D.M. Xiao, S.G. Ruan, Global analysis of an epidemic model with nonmonotone incidence rate, *Math. Biosci.* 208 (2007) 419–429.
- [9] M. Lu, J.C. Huang, S.G. Ruan, P. Yu, Bifurcation analysis of an SIRS epidemic model with a generalized nonmonotone and saturated incidence rate, *J. Differ. Equations* 267 (2019) 1859–1898.
- [10] J. Andrews, A mathematical model for the continuous culture of microorganisms utilizing inhibitory substrates, *Biotechnol. Bioeng.* 10(1968):707–723.
- [11] H.W. Hethcote, H.W. Stech, P. van den Driessche, Stability analysis for models of diseases without immunity, *J. Math. Biol.* 13 (1981) 185–198.
- [12] S. Busenberg, K.L. Cooke, The population dynamics of two vertically transmitted infections, *Theor. Popul. Biol.* 33 (1988) 181–198.
- [13] W.R. Derrick, P. van den Driessche, A disease transmission model in a nonconstant population, *J. Math. Biol.* 31 (1993) 495–512.
- [14] H.W. Hethcote, The mathematics of infectious disease, *SIAM Rev.* 42 (2000) 599–653.
- [15] D.Z. Gao, Travel frequency and infectious diseases, *SIAM J. Appl. Math.* (2019) 1581–1606.
- [16] C. Castillo-Chavez, B.J. Song, Dynamical models of tuberculosis and their applications, *Math. Biosci. Eng.* 1 (2004)