

SHARP RATE OF CONVERGENCE FOR DEEP NEURAL NETWORK CLASSIFIERS UNDER THE TEACHER-STUDENT SETTING

BY TIANYANG HU^{*}, ZUOFENG SHANG[†] AND GUANG CHENG^{*}

Purdue University^{} and New Jersey Institute of Technology[†]*

Classifiers built with neural networks handle large-scale high dimensional data, such as facial images from computer vision, extremely well while traditional statistical methods often fail miserably. In this paper, we attempt to understand this empirical success in high dimensional classification by deriving the convergence rates of excess risk. In particular, a teacher-student framework is proposed that assumes the Bayes classifier to be expressed as ReLU neural networks. In this setup, we obtain a sharp rate of convergence, i.e., $\tilde{O}_d(n^{-2/3})^*$, for classifiers trained using either 0-1 loss or hinge loss. This rate can be further improved to $\tilde{O}_d(n^{-1})$ when the data distribution is separable. Here, n denotes the sample size. An interesting observation is that the data dimension only contributes to the $\log(n)$ term in the above rates. This may provide one theoretical explanation for the empirical successes of deep neural networks in high dimensional classification, particularly for structured data.

1. Introduction. Deep learning has gained tremendous success in classification problems such as image classifications [Deng et al., 2009b]. With the introduction of convolutional neural network [Krizhevsky et al., 2012] and residual neural network [He et al., 2016], various benchmarks in computer vision have been revolutionized and neural network based methods have achieved better-than-human performance [Nguyen et al., 2017]. For instance, AlexNet [Krizhevsky et al., 2012] and its variants [Zeiler and Fergus, 2014, Simonyan and Zisserman, 2014] have demonstrated superior performance in ImageNet data [Deng et al., 2009a, Russakovsky et al., 2015], where the data dimension is huge, i.e., each image has pixel size 256×256 and hence is an 65536-dimensional vector. Traditional statistical thinking sounds an alarm when facing such high-dimension data as the “curse of dimensionality” usually prevents nonparametric classification achieving fast convergence rates. This work attempts to provide a theoretical explanation for the empirical success of deep neural networks (DNN) in (especially high dimensional) classification, beyond the existing statistical theories.

^{*}The notation \tilde{O}_d means “up to a logarithmic factor depending on the data dimension d .”

In the context of nonparametric regression, similar investigations have been recently carried out. Among others [Farrell et al., 2018, Suzuki, 2018, Nakada and Imaizumi, 2019, Oono and Suzuki, 2019, Chen et al., 2019, Liu et al., 2019], Schmidt-Hieber [2019] shows that deep ReLU neural networks can achieve minimax rate of convergence when the underlying regression function possesses a certain compositional smooth structure; Bauer et al. [2019], Kohler and Langer [2019] show a similar result by instead considering hierarchical interaction models; Imaizumi and Fukumizu [2018] demonstrate the superiority of neural networks in estimating a class of non-smooth functions, in which case no linear methods, e.g. kernel smoothing, Gaussian process, can achieve the optimal convergence rate. The aforementioned works all build on the traditional smoothness assumption and the convergence rates derived therein are still subject to curse of dimensionality.

Classification and regression are fundamentally different due to the discrete nature of class labels. Specifically, in nonparametric regression, we are interested in recovering the whole underlying function while in classification, the focus is on the nonparametric estimation of sets corresponding to different classes. As a result, it is well known that many established results on regression cannot be directly translated to classification. The goal of this paper is hence to fill this gap by investigating how well neural network based classifiers can perform in theory and further provide a theoretical explanation for the “break-the-curse-of-dimensionality” phenomenon. To this end, we propose to study neural network based classifiers in a teacher-student setting where the traditional smoothness assumption is no longer present.

The teacher-student framework has originated from statistical mechanics [Saad and Solla, 1996, Mace and Coolen, 1998, Engel and Broeck, 2002] and recently gained increasing interest [Hinton et al., 2015, Ba and Caruana, 2014, Goldt et al., 2019, Aubin et al., 2018]. In this setup, one neural network, called student net, is trained on data generated by another neural network, called teacher net. While worst-case analysis for arbitrary data distributions may not be suitable for real structured dataset, adopting this framework can facilitate the understanding of how deep neural networks work as it provides an explicit target function with bounded complexity. Furthermore, assuming the target classifier to be a teacher network of an explicit architecture may provide insights on what specific architecture of the student classifier is needed to achieve an optimal excess risk. At the same time, by comparing the two networks, both optimization and generalization can be handled more elegantly. Existing works on how well student network can learn from the teacher mostly focus on regression problems and study how the student network evolves during training from computational aspects, e.g., [Tian, 2018,

2019, Goldt et al., 2019, Zhang et al., 2019, Cao and Gu, 2019]. Still, there is a lack of statistical understanding in this important direction, particularly on classification aspects.

In this paper, we consider binary classification, and focus on the teacher-student framework where the optimal decision region is defined by ReLU neural networks. This setting is closely related to the classical smooth boundary assumption where the neural networks are substituted by smooth functions. Specifically, a well-adopted assumption called as “boundary fragment” [Mammen et al., 1999, Tsybakov et al., 2004, Imaizumi and Fukumizu, 2018] assumes the smooth function to be linear in one of the dimensions (see Appendix 6.1). Our teacher-student network setting is more general as it does not impose any special structures on the decision boundary. Moreover, by the universal approximation property [Cybenko, 1989, Arora et al., 2016, Lu et al., 2017], the teacher network can sufficiently approximate any continuous function given large enough size.

In the above setting, an un-improvable rate of convergence is derived as $\tilde{O}_d(n^{-2/3})$ for the excess risk of the empirical 0-1 loss minimizer, given that the student network is deeper and larger than the teacher network (unless the teacher network has a limited capacity in some sense to be specified later). When data are separable, the rate improves to $\tilde{O}_d(n^{-1})$. In contrast, under the smooth boundary assumption, Mammen et al. [1999] establish the optimal risk bound to be $O(n^{-\beta(\kappa+1)/[\beta(\kappa+2)+(d-1)\kappa]})$ where $\beta > 0$ represents the smoothness of the “boundary fragments” and $\kappa > 0$ is the so-called Tsybakov noise exponent. Clearly, this rate suffers from the “curse of dimensionality” but interestingly, coincides with our rate when $\kappa = 1$ and $\beta \rightarrow \infty$ (up to a logarithmic factor). If we further allow $\kappa \rightarrow \infty$ (corresponding to separable data), the classical rate above recovers $\tilde{O}(n^{-1})$ (up to a logarithmic factor). Please see the Appendix 6.1 for detail.

Furthermore, we extend our analysis to a specific surrogate loss, i.e., hinge loss, and show that the convergence rate remains the same (up to higher order logarithmic terms) while allowing *deeper* student and teacher nets. The obtained sharp risk bounds may explain the empirical success of deep neural networks in high-dimensional classification as the data dimension d only appears in the $\log(n)$ terms. Our main technical novelty is the nontrivial entropy calculation for nonparametric set estimation based on combinatorial analysis of ReLU neural networks.

Existing Works. We review some related works on classification using neural networks. The first class of literature demonstrate that deep neural network (DNN) classifiers can be efficiently optimized in different senses. For example, Liang et al. [2018] study the loss surface of neural networks in classification

and provide conditions that guarantee zero training error at all local minima of appropriately chosen surrogate loss functions. Additionally, [Lyu and Li \[2019\]](#) show that under exponential loss [[Soudry et al., 2018](#), [Gunasekar et al., 2018](#)], gradient descent on homogeneous neural network has implicit bias towards the maximum L_2 margin solution. In all these optimization works, sharp bounds are not derived for either generalization error or convergence rate of the excess risk. From a nonparametric perspective, [Kim et al. \[2018\]](#) derive the excess risk bound for DNN classifiers as $\tilde{O}(n^{-\beta(\kappa+1)/[\beta(\kappa+2)+(d-1)(\kappa+1)]})$, which is suboptimal in the sense of [Tsybakov et al. \[2004\]](#). Clearly, such a dimension-dependent bound, indicating exponential dependence on the sample size, may not support the empirical success of deep learning in high dimension classification.

Notations. For any function $f : \mathcal{X} \rightarrow \mathbb{R}$, denote let $\|f\|_\infty = \sup_{\mathbf{x} \in \mathcal{X}} |f(\mathbf{x})|$ and $\|f\|_p = (\int |f|^p)^{1/p}$ for $p \in \mathbb{N}$. For two given sequences $\{a_n\}_{n \in \mathbb{N}}$ and $\{b_n\}_{n \in \mathbb{N}}$ of real numbers, we write $a_n \lesssim b_n$ if there exists a constant $C > 0$ such that $a_n \leq Cb_n$ for all sufficiently large n , which is also denoted as $a_n = O(b_n)$. Let $\Omega(\cdot)$ be the counterpart of $O(\cdot)$ that $a_n = \Omega(b_n)$ means $a_n \gtrsim b_n$. In addition, we write $a_b \asymp b_n$ if $a_n \lesssim b_n$ and $a_n \gtrsim b_n$. For $a, b \in \mathbb{R}$, denote $a \vee b = \max\{a, b\}$ and $a \wedge b = \min\{a, b\}$. Let $\lfloor a \rfloor$ represent $\max\{n \in \mathbb{N} : n \leq a\}$. \mathbb{I} denotes the indicator function. Independently, identically distributed is abbreviated as i.i.d..

2. Preliminary.

2.1. *Neural Network Setup.* We consider deep neural networks with Rectified Linear Unit (ReLU) activation that $\sigma(x) = \max\{x, 0\}$. For an L hidden layer ReLU neural network $f(\cdot)$, let the width of each layer be n_0, n_1, \dots, n_L , where $n_0 = d$ is the input dimension, and denote the weight matrices and bias vectors in each layer to be $\mathbf{W}^{(l)}$ and $\mathbf{b}^{(l)}$, respectively. Let $\sigma_{(\mathbf{W}, \mathbf{b})}(\mathbf{x}) = \sigma(\mathbf{W} \cdot \mathbf{x} + \mathbf{b})$ and \circ represent function composition. Then, the ReLU DNN can be written as

$$f(\mathbf{x}|\Theta) = \mathbf{W}^{(L+1)} \sigma_{(\mathbf{W}^{(L)}, \mathbf{b}^{(L)})} \circ \dots \circ \sigma_{(\mathbf{W}^{(1)}, \mathbf{b}^{(1)})}(\mathbf{x}) + \mathbf{b}^{(L+1)}, \quad \mathbf{x} \in \mathbb{R}^d,$$

where $\Theta = \{(\mathbf{W}^{(l)}, \mathbf{b}^{(l)})\}_{l=1, \dots, L+1}$ denotes the parameter set.

For any given Θ , let $|\Theta|$ be the number of hidden layers in Θ , and $N_{\max}(\Theta)$ be the maximum width. We define $\|\Theta\|_0$ as the number of nonzero parameters:

$$\|\Theta\|_0 = \sum_{l=1}^{L+1} \left(\|\text{vec}(\mathbf{W}^{(l)})\|_0 + \|\mathbf{b}^{(l)}\|_0 \right),$$

where $\text{vec}(\mathbf{W}^{(l)})$ transforms the matrix $\mathbf{W}^{(l)}$ into the corresponding vector by concatenating the column vectors. Similarly, we define $\|\Theta\|_\infty$ as the largest absolute value of the parameters in Θ ,

$$\|\Theta\|_\infty = \max \left\{ \max_{1 \leq l \leq L+1} \|\text{vec}(\mathbf{W}^{(l)})\|_\infty, \max_{1 \leq l \leq L+1} \|\mathbf{b}^{(l)}\|_\infty \right\}.$$

For any given n , let \mathcal{F}_n be

$$\begin{aligned} \mathcal{F}_n &= \mathcal{F}^{\text{DNN}}(L_n, N_n, S_n, B_n, F_n) \\ &= \{f(\mathbf{x}|\Theta) : |\Theta| \leq L_n, N_{\max}(\Theta) \leq N_n, \|\Theta\|_0 \leq S_n, \\ &\quad \|\Theta\|_\infty \leq B_n, \|f(\cdot|\Theta)\|_\infty \leq F_n\}. \end{aligned}$$

2.2. Binary Classification. Consider binary classification with a feature vector $\mathbf{x} \in \mathcal{X} \subset \mathbb{R}^d$ and a label $y \in \{-1, 1\}$. Assume $\mathbf{x}|y = 1 \sim p(\mathbf{x})$, $\mathbf{x}|y = -1 \sim q(\mathbf{x})$ where p and q are two bounded densities on \mathcal{X} w.r.t. base measure \mathbb{Q} . If p, q have disjoint support, we say the data distribution or the classification problem is separable. For simplicity, assume that \mathbb{Q} is Lebesgue measure, and positive and negative labels are equally likely to appear, i.e., balanced labels.

The objective of classification is to find an optimal classifier (called the Bayes classifier) C^* within some classifier family \mathcal{C} , that minimizes the 0-1 loss defined as

$$C^* = \underset{C \in \mathcal{C}}{\text{argmin}} R(C) := \underset{C \in \mathcal{C}}{\text{argmin}} \mathbb{E} [\mathbb{I}\{C(\mathbf{x}) \neq y\}].$$

We can estimate C^* based on the training data by minimizing the empirical 0-1 risk as follows

$$\hat{C}_n = \underset{C \in \mathcal{C}_n}{\text{argmin}} R_n(C) := \underset{C \in \mathcal{C}_n}{\text{argmin}} \sum_{i=1}^n \mathbb{I}\{C(\mathbf{x}_i) \neq y_i\}/n,$$

where \mathcal{C}_n is a given class of classifiers possibly depending on the sample size n . In practice, the above 0-1 loss is often replaced by its (computationally feasible) surrogate counterparts [Bartlett et al., 2006], such as hinge loss $\phi(z) = (1 - z)_+ = \max\{1 - z, 0\}$ or logistic loss $\phi(z) = \log(1 + \exp(-z))$.

Given a surrogate loss ϕ , we first obtain \hat{f}_ϕ by minimizing the empirical risk

$$R_{\phi,n}(f) = \sum_{i=1}^n \phi(y_i f(\mathbf{x}_i))/n$$

over \mathcal{F} , and then construct a classifier by $\widehat{C}_\phi(\mathbf{x}) = \text{sign}(\widehat{f}_\phi(\mathbf{x}))$. Accordingly, define an optimal f_ϕ^* as $f_\phi^* = \text{argmin}_{f \in \mathcal{F}} R_\phi(f)$, where $R_\phi(f) := \mathbb{E}R_{\phi,n}(f)$ is the population risk. Given that $C(\mathbf{x}) = \text{sign}(f(\mathbf{x}))$, with a slight abuse of notation, we write $R(C)$ and $R(f)$ interchangeably. A classifier C is evaluated by its excess risk defined as the difference of the population risk between C and the Bayes optimal classifier C^* that $\mathcal{E}(C, C^*) = R(C) - R(C^*)$. Our goal is to derive sharp convergence rates of $\mathcal{E}(C, C^*)$ under different losses.

3. Teacher-Student Framework for Classification. In this section, we set up the teacher-student framework for classification under which sharp rates for the excess risk are developed. Such a teacher-student bound sets an algorithmic independent benchmark for various deep neural network classifiers and also helps understand the role of input data dimension in the classification performance. The Bayes classifier C^* is defined via the optimal decision region $G^* := \{\mathbf{x} \in \mathcal{X}, p(\mathbf{x}) - q(\mathbf{x}) \geq 0\}$. The set estimate $\widehat{G} = \{\mathbf{x} \in \mathcal{X}, \widehat{f}(\mathbf{x}) \geq 0\}$ can be constructed through deep neural network classifiers $\widehat{f} : \mathbb{R}^d \rightarrow \mathbb{R}$ trained using either 0-1 loss or surrogate loss. Accordingly, a natural teacher network assumption is that $p(\mathbf{x}) - q(\mathbf{x})$ can be expressed by some neural network $f_n^* \in \mathcal{F}_n^*$. Here, the underlying densities are indexed by n , but such an assumption is not uncommon in high-dimensional statistics, where population quantities may depend on the sample size n , e.g., [Zhao and Yu \[2006\]](#).

3.1. *Training with 0-1 Loss.* In this section, we focus on, for the theoretical purpose, DNN classifiers trained with the empirical 0-1 loss. Denote

$$\widehat{f}_n = \text{argmin}_{f \in \mathcal{F}_n} \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{y_i f(\mathbf{x}_i) < 0\},$$

given a certain DNN family \mathcal{F}_n .

It is important to control the complexity of the underlying classification problem. Otherwise, the student network would not be able to recover the Bayes classifier [[Telgarsky, 2015](#)] with sufficient accuracy. To this end, we impose the following teacher network assumptions on $(p(\mathbf{x}) - q(\mathbf{x}))$:

(A1) p, q have compact supports.

(A2) $p(\mathbf{x}) - q(\mathbf{x})$ is representable by some teacher ReLU DNN $f_n^* \in \mathcal{F}_n^*$ with

$$N_n^* = O(\log n)^{m_*}, \quad L_n^* = O(1) \quad \text{for some } m_* \geq 1.$$

(A3) For any n , there exists $c_n, 1/T_n = O(\log n)^{m_* d^2 L_n^*}$ such that for all $0 \leq t \leq T_n$.

$$\mathbb{Q}\{\mathbf{x} \in \mathcal{X} : |f_n^*(\mathbf{x})| \leq t\} \leq c_n t$$

Assumption (A3) characterizes how concentrated the data are around the decision boundary, which can be seen as an extension to the classical Tsybakov noise condition [Mammen et al., 1999]. The difference is that in our case, the underlying densities are indexed by sample size and thus c_n and T_n are allowed to vary with n . Assumption (A3) is not unrealistic as we will show that it holds with high probability if the teacher network is random as stated in the following lemma (see Appendix 6.3 for detail).

Lemma 3.1. Let f_n^* be the teacher network with structures specified in assumption (A2). Suppose that all weights of f_n^* are i.i.d. with any continuous distribution, e.g. Gaussian, truncated Gaussian, etc.. Then, with probability at least $1 - \delta$, assumption (A3) holds with $c_n, 1/T_n \leq A(\delta)(\log n)^{m^*d^2L_n^*}$ where $A(\delta)$ is some constant depending on δ .

The following theorem characterizes how well the student network of proper size can learn from the teacher in terms of the excess risk.

Theorem 3.2. Under the teacher assumptions (A1) through (A3), denote all such (p, q) pairs to be \mathcal{P}_n^* and let the corresponding Bayes classifier be C_n^* . Let \mathcal{F}_n be a student ReLU DNN family with $N_n = O(\log n)^m$ and $L_n = O(1)$ for some $m \geq m_*$ and assume the student network is larger than the teacher network in the sense that $L_n \geq L_n^*, S_n \geq S_n^*, N_n \geq N_n^*, B_n \geq B_n^*$. Then the excess risk for $\hat{f}_n \in \mathcal{F}_n$ satisfies

$$\sup_{(p,q) \in \mathcal{P}_n^*} \mathbb{E}[\mathcal{E}(\hat{f}_n, C_n^*)] = \tilde{O}_d \left(\frac{1}{n} \right)^{\frac{2}{3}},$$

where \tilde{O}_d hides the $\log n$ terms, which depend on d .

The dependence on the dimension d is in the order of $O(\log n)^{d^2}$. We further argue that under the present setting, the rate $n^{-2/3}$ in Theorem 3.2 cannot be further improved.

Theorem 3.3. Under the same assumptions of p, q as in Theorem 3.2 that $(p, q) \in \tilde{\mathcal{F}}_n^*$. Let $\tilde{\mathcal{F}}_n$ be an arbitrary function space, then

$$\inf_{\tilde{f}_n \in \tilde{\mathcal{F}}_n} \sup_{(p,q) \in \tilde{\mathcal{F}}_n^*} \mathbb{E}[\mathcal{E}(\tilde{f}_n, f_n^*)] = \tilde{\Omega}_d \left(\frac{1}{n} \right)^{\frac{2}{3}},$$

where $\tilde{\Omega}_d$ hides the $\log n$ terms, which depend on d .

Theorem 3.3 shows that the convergence rate achieved by the empirical 0-1 loss minimizer cannot be further improved (up to a logarithmic term). If p and q have disjoint supports, i.e. separable, which could be true in some

image data, the rate improves to n^{-1} , as stated in the following corollary. This rate improvement is not surprising since the classification task becomes much easier for separable data.

Corollary 3.4. Under the same setting as in Theorem 3.2, if we further assume p, q have disjoint supports, then the rate of convergence of the empirical 0-1 loss minimizer improves to

$$\inf_{f_n \in \mathcal{F}_n} \sup_{(p,q) \in \tilde{\mathcal{F}}_n^*} \mathbb{E}[\mathcal{E}(f_n, f_n^*)] \asymp \tilde{O}\left(\frac{1}{n}\right).$$

Remark 1 (Disjoint Support). Given that data are separable, Srebro et al. [2010] derived the excess risk bound as $O(D \log n/n)$ (under a smooth loss) where D is the VC-subgraph-dimension of the estimation family. Additionally, separability implies that the noise exponent κ in Tsybakov noise condition [Mammen et al., 1999, Tsybakov et al., 2004] can be arbitrarily large, which also gives $O(1/n)$ rate under the “boundary fragments” assumption.

The imposed relation between teacher and student nets in Theorem 3.2 is referred to as “over-realization” in Goldt et al. [2019], Tian [2019], Bai et al. [2019]: at each layer, the number of student nodes is larger than that of teacher nodes given the same depth. In other words, the student network is larger than the teacher in order to obtain zero approximation error.

On the other hand, such a requirement is not necessary as long as the corresponding Bayes classifier is not too complicated. A ReLU neural network is a continuous piecewise linear function, i.e. its domain can be divided into connected regions (pieces) within where the function is linear. If the ReLU neural network crosses 0 on one piece, we call that piece as being *active* (see Figure 1 for an illustration). One way to measure the complexity of the teacher network is the number of active pieces. The following Corollary says that the teacher network can be much larger and deeper as long as the number of active pieces are in a logarithmic order with respect to n .

Corollary 3.5. The same result in Theorem 3.2 holds when the teacher network is larger than the student network, i.e. $L_n \leq L_n^*, S_n \leq S_n^*, N_n \leq N_n^*, B_n \leq B_n^*$, given that the total number of active pieces in the teacher network is of the following order

$$(3.1) \quad o\left(\left(\prod_{l=1}^{L_n-1} \lfloor \frac{n_l}{d} \rfloor^d\right) \sum_{j=0}^d \binom{n_{L_n}}{j}\right),$$

where n_1, \dots, n_{L_n} are the width of each hidden layer of the student network.




FIG 1. Example of a ReLU DNN function in $[0, 1]$. There are 5 pieces p_1, p_2, \dots, p_5 and among them, only p_1, p_4, p_5 cross value 0 (horizontal line). There are 3 active pieces in this example and they are colored red.

The number of active pieces is the key quantity in controlling the complexity of the optimal set G^* . The expression in (3.1) comes from the lower bound developed by Montufar et al. [2014] on the maximum number of linear pieces for a ReLU neural network (Lemma 3.8). This lower bound is determined by the structure of the student network. If the number of active pieces of the teacher network is on this order, i.e. within the capacity of the student, then the corresponding optimal set can still be recovered by an even smaller student network, which ensures zero approximation error. Since the student network in consideration satisfies $N_n = O(\log n)^m$, the required order for the number of active pieces is in the order of $o(\log n)^{mdL_n}$.

3.2. Proof of Theorem 3.2. We first present some preliminary lemmas. As we mentioned before, classification can be thought of as nonparametric estimation of sets. For this, we define two distances over sets. The first one is the usual symmetric difference of sets: for any $G_1, G_2 \subset \mathbb{R}^d$,

$$d_{\Delta}(G_1, G_2) = \mathbb{Q}(G_1 \Delta G_2) = \mathbb{Q}((G_1 \setminus G_2) \cup (G_2 \setminus G_1)),$$

where \mathbb{Q} denotes the Lebesgue measure. The second one is induced by densities p, q : for any $G_1, G_2 \subset \mathbb{R}^d$,

$$d_{p,q}(G_1, G_2) = \int_{G_1 \Delta G_2} |p(\mathbf{x}) - q(\mathbf{x})| \mathbb{Q}(d\mathbf{x}).$$

There are two key factors governing the rate of convergence in classification:

- How concentrated the data are around the decision boundary;
- The complexity of the set \mathcal{G}^* where the optimal G^* resides.

For the first factor, the following Tsybakov noise condition [Mammen et al., 1999] quantifies how close p and q are:

(N) There exists constant $c > 0$ and $\kappa \in [0, \infty]$ such that for any $0 \leq t \leq T$

$$\mathbb{P}(\{\mathbf{x} : |p(\mathbf{x}) - q(\mathbf{x})| \leq t\}) \leq ct^\kappa.$$

The parameter $\kappa > 0$ is referred to as the *noise exponent*. The bigger the κ , the less concentrated the data are around the decision boundary and hence the easier the classification. In the extreme case that p, q have different supports, κ can be arbitrarily large and the classification is easy. To another extreme where $\mathbb{Q}\{\mathbf{x} \in \mathcal{X} : p(\mathbf{x}) = q(\mathbf{x})\} > 0$, there exists a region where different classes are indistinguishable. In this case, $\kappa = 0$ and the classification is hard in that region.

For the second factor, we use bracketing entropy to measure the complexity of a collection of subsets \mathcal{G} in \mathbb{R}^d . For any $\delta > 0$, the bracketing number $\mathcal{N}_B(\delta, \mathcal{G}, d_\Delta)$ is the minimal number of set pairs (U_j, V_j) such that, (a) for each j , $U_j \subset V_j$ and $d_\Delta(U_j, V_j) \leq \delta$; (b) for any $G \in \mathcal{G}$, there exists a pair (U_j, V_j) such that $U_j \subset G \subset V_j$. Simply denote $\mathcal{N}_B(\delta) = \mathcal{N}_B(\delta, \mathcal{G}, d_\Delta)$ if no confusion arises. The bracketing entropy is defined as $H_B(\delta) = \log \mathcal{N}_B(\delta, \mathcal{G}, d_\Delta)$.

Lemma 3.6 characterizes the complexity of a special collection of sets.

Lemma 3.6. Let $\mathcal{X} = [0, 1]^d$ and \mathcal{G} be a collection of polyhedrons with at most S vertices in \mathbb{R}^d . Then the bracketing entropy of $\bar{\mathcal{G}} = \mathcal{G} \cap \mathcal{X}$ satisfies

$$H_B(\delta, \bar{\mathcal{G}}, d_\Delta) = \log \mathcal{N}_B(\delta, \bar{\mathcal{G}}, d_\Delta) \lesssim d^2 S \log(d^{3/2} S / \delta)$$

PROOF OF LEMMA 3.6. Let's first introduce some notations and terminologies. For any $\delta > 0$, let M_δ denote the smallest integer such that $M_\delta > 1/\delta$. Consider the set of lattice points $\mathbf{X}_\delta^d = \{(i_1/M_\delta, \dots, i_d/M_\delta) : i_1, \dots, i_d = 0, 1, \dots, M_\delta\}$ which has cardinality $(M_\delta + 1)^d$. Let $G(\mathbf{x}_1, \dots, \mathbf{x}_s)$ denote a polyhedron with vertices $\mathbf{x}_1, \dots, \mathbf{x}_s \in [0, 1]^d$ where $s \leq S$. (the \mathbf{x}_i 's are not necessarily distinct). Any convex polyhedron G in \mathbb{R}^d is the intersection of multiple $(d-1)$ -dimensional hyperplanes. If we move all such hyperplanes inwards (to the direction perpendicular to the hyperplanes) by a small distance δ , they produce another polyhedron, denoted $G_{-\delta}$, called as the δ -contraction of G . Note that $G_{-\delta}$ can be empty if δ is not small enough.

We prove the result for $d = 1$, in which $\bar{\mathcal{G}}$ is a collection of subintervals in $[0, 1]$. For any subinterval $[a, b] \subset [0, 1]$, there exist $x_i, x_j \in \mathbf{X}_\delta^1$ such that

$$x_i \leq a \leq x_{i+1}, \quad x_j \leq b \leq x_{j+1}.$$

(By convention, $[x_i, x_j]$ is empty if $x_i > x_j$.) Then $([x_i, x_{j+1}], [x_{i+1}, x_j])$ is a 2δ -bracket of $[a, b]$ since obviously

$$(3.2) \quad [x_{i+1}, x_j] \subset [a, b] \subset [x_i, x_{j+1}], \quad d_\Delta([x_i, x_{j+1}], [x_{i+1}, x_j]) \leq 2\delta.$$




FIG 2. Grid in 2D and the outer cover (green) constructed for with grid points for a polygon (blue).

There are $\binom{M_\delta+1}{2}$ different choices of $[x_i, x_j]$, hence, $\binom{M_\delta+1}{2}$ different choices of the pairs $([x_i, x_{j+1}], [x_{i+1}, x_j])$. Any $[a, b] \subset [0, 1]$ can be 2δ bracketed by one of such pairs in the sense of (3.2). This shows that $H_B(2\delta) \leq \log \binom{M_\delta+1}{2} \leq 2 \log(1/\delta)$.

When $d \geq 2$, any $G \in \bar{\mathcal{G}}$ has at most S vertices, so $\bar{G} := G \cap [0, 1]^d$ has at most dS vertices where the factor d is due to the fact that each edge of G intersects at most d edges of $[0, 1]^d$ therefore creates at most dS vertices for \bar{G} . For any polygon $G(\mathbf{x}_1, \dots, \mathbf{x}_s)$ where $s \leq dS$, denote $G_{-\sqrt{d}\delta}(\mathbf{x}_1, \dots, \mathbf{x}_s) = G(\mathbf{x}_1^-, \dots, \mathbf{x}_s^-)$. Each vertex must be in one of the grids in \mathbf{X}_δ^d . It is easy to see that there exist $\mathbf{v}_1^1, \dots, \mathbf{v}_1^d, \mathbf{v}_2^1, \dots, \mathbf{v}_2^d, \dots, \mathbf{v}_s^1, \dots, \mathbf{v}_s^d \in \mathbf{X}_\delta^d$, where $\mathbf{v}_i^1, \dots, \mathbf{v}_i^d$ are in the same grid, such that

- $G(\mathbf{x}_1, \dots, \mathbf{x}_s) \subset G(\mathbf{v}_1^1, \dots, \mathbf{v}_1^d, \mathbf{v}_2^1, \dots, \mathbf{v}_2^d, \dots, \mathbf{v}_s^1, \dots, \mathbf{v}_s^d)$;
- $\|\mathbf{v}_i^j - \mathbf{x}_i\|_2 \leq \sqrt{d}\delta$ for $i = 1, 2, \dots, s$ and $j = 1, 2, \dots, d$.

See Figure 2 for an illustration when $d = 2$. Similarly for $G(\mathbf{x}_1^-, \dots, \mathbf{x}_s^-)$, there exist $\mathbf{u}_1^1, \dots, \mathbf{u}_1^d, \mathbf{u}_2^1, \dots, \mathbf{u}_2^d, \dots, \mathbf{u}_s^1, \dots, \mathbf{u}_s^d \in \mathbf{X}_\delta^d$ such that

- $G(\mathbf{x}_1^-, \dots, \mathbf{x}_s^-) \subset G(\mathbf{u}_1^1, \dots, \mathbf{u}_1^d, \mathbf{u}_2^1, \dots, \mathbf{u}_2^d, \dots, \mathbf{u}_s^1, \dots, \mathbf{u}_s^d)$;
- $\|\mathbf{u}_i^j - \mathbf{x}_i^-\|_2 \leq \sqrt{d}\delta$ for $i = 1, 2, \dots, s$ and $j = 1, 2, \dots, d$.

By the definition of $G_{-\sqrt{d}\delta}$, we have $\|\mathbf{x}_i - \mathbf{x}_i^-\|_2 \geq \sqrt{d}\delta$. Thus $\|\mathbf{u}_i^j - \mathbf{x}_i^-\|_2 \leq \sqrt{d}\delta$ implies $G(\mathbf{u}_1^1, \dots, \mathbf{u}_1^d, \dots, \mathbf{u}_s^1, \dots, \mathbf{u}_s^d) \subset G(\mathbf{x}_1, \dots, \mathbf{x}_s)$. On the other

hand,

$$\begin{aligned} & d_\Delta(G(\mathbf{u}_1^1, \dots, \mathbf{u}_1^d, \dots, \mathbf{u}_s^1, \dots, \mathbf{u}_s^d), G(\mathbf{v}_1^1, \dots, \mathbf{v}_1^d, \dots, \mathbf{v}_s^1, \dots, \mathbf{v}_s^d)) \\ & \leq d_\Delta(G_{+\sqrt{d}\delta}(\mathbf{x}_1, \dots, \mathbf{x}_s), G_{-\sqrt{d}\delta}(\mathbf{x}_1, \dots, \mathbf{x}_s)) \\ & \leq s \cdot 2\sqrt{d}\delta, \end{aligned}$$

where the term s is due to the fact that $G(\mathbf{x}_1, \dots, \mathbf{x}_s)$ has at most $O(s)$ faces. Notice that

$$G(\mathbf{u}_1^1, \dots, \mathbf{u}_1^d, \dots, \mathbf{u}_s^1, \dots, \mathbf{u}_s^d), G(\mathbf{v}_1^1, \dots, \mathbf{v}_1^d, \dots, \mathbf{v}_s^1, \dots, \mathbf{v}_s^d) \in \bar{\mathcal{G}},$$

and $s \leq dS$. Thus, with at most $(M_\delta + 1)^{d^2 S}$ pairs of subsets in $\bar{\mathcal{G}}$, we can $2d^{3/2}S\delta$ -bracket any $\bar{G} \in \bar{\mathcal{G}}$. Therefore,

$$\log \mathcal{N}_B((2d^{3/2}S\delta), \bar{\mathcal{G}}, d_\Delta) \lesssim \log \left((M_\delta + 1)^{d^2 S} \right),$$

which implies

$$\log \mathcal{N}_B(\delta, \bar{\mathcal{G}}, d_\Delta) \lesssim d^2 S \log(d^{3/2}S/\delta).$$

□

Lemma 3.7 (Theorem 1 in [Serra et al., 2017]). Consider a deep ReLU network with L layers, n_l ReLU nodes at each layer l , and an input of dimension n_0 . The maximal number of linear pieces of this neural network is at most

$$\sum_{(j_1, \dots, j_L) \in J} \prod_{l=1}^L \binom{n_l}{j_l},$$

where $J = \{(j_1, \dots, j_L) \in \mathbb{Z}^L : 0 \leq j_l \leq \min\{n_0, n_1 - j_1, \dots, n_{l-1} - j_{l-1}, n_l\} \forall l = 1, \dots, L\}$. This bound is tight when $L = 1$. When $n_0 = O(1)$ and all layers have the same width N , we have the same best known asymptotic bound $O(N^{L n_0})$ first presented in [Raghu et al., 2017].

Consider a deep ReLU network with $n_0 = d$ inputs and L hidden layers of widths $n_i \geq n_0$ for all $i \in [L]$. The following lemma establishes a lower bound for the maximal number of linear pieces of deep ReLU networks:

Lemma 3.8 (Theorem 4 in [Montufar et al., 2014]). The maximal number of linear pieces of a ReLU network with n_0 input units, L hidden layers, and $n_i \geq n_0$ rectifiers on the i -th layer, is lower bounded by

$$\left(\prod_{i=1}^{L-1} \left\lfloor \frac{n_i}{n_0} \right\rfloor^{n_0} \right) \sum_{j=0}^{n_0} \binom{n_L}{j}.$$




FIG 3. *Demonstration of how a polygon in $d = 2$ case can be divided into basic triangles. The union of the two brackets form a bracket of the original polygon. The blue shade is the symmetric difference.*

Lemma 3.9. Let \mathcal{F} be a class of ReLU neural networks, defined on $\mathcal{X} = [0, 1]^d$, with at most L layers and N neurons per layer. Let $G^f = \{\mathbf{x} \in \mathcal{X} : f(\mathbf{x}) \geq 0\}$ and $\mathcal{G}^{\mathcal{F}} = \{G^f : f \in \mathcal{F}\}$. Then the bracketing number of $\mathcal{G}^{\mathcal{F}}$ satisfies

$$\log \mathcal{N}_B(\delta, \mathcal{G}^{\mathcal{F}}, d_{\Delta}) \lesssim N^{Ld^2} d^3 (Ld^2 \log(N) \vee \log(1/\delta)).$$

PROOF. The proof relies on Lemma 3.6 for which we need to control the number of vertexes of G^f based on the number of pieces (linear regions) of the ReLU neural network. Since ReLU neural networks are piecewise linear, G^f is a collection of sets of polyhedrons. Define the subgraph of a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ to be the set of points in \mathbb{R}^{d+1} :

$$\text{sub}(f) = \{(\mathbf{x}, t) : f(\mathbf{x}) \geq t\}.$$

In this sense, $\text{sub}(f) \cap \{(\mathbf{x}, 0) : \mathbf{x} \in \mathcal{X}\} = \{(\mathbf{x}, 0) : \mathbf{x} \in G^f\}$, a slice of the subgraph. Denote all the pieces to be p_1, p_2, \dots, p_s . Each piece is a d -dimensional polyhedron on which $f(x)$ is linear. To control the complexity of G^f , consider the most extreme case that the function crosses zero on each piece, i.e. for any $i = 1, \dots, s$, $\{(\mathbf{x}, f(\mathbf{x})) : \mathbf{x} \in p_i\} \cap \{(\mathbf{x}, 0) : \mathbf{x} \in \mathcal{X}\} \neq \emptyset$. Each intersection resides in a $(d - 1)$ -dimensional hyperplane, e.g. dot for $d = 1$, line segment for $d = 2$ and so on. So the number of such $(d - 1)$ -dimensional hyperplanes in G^f is at most s .

A vertex of a polyhedron in $[0, 1]^d$ can be thought of as the intersection of at least d hyperplanes of dimension $d - 1$. Thus, with at most s hyperplanes there are at most $\binom{s}{d} < s^d$ vertices in G^f . In order to apply Lemma 3.6, we break the collection of polyhedrons into the so-called *basic polyhedrons* each with $d + 1$ vertices. For instance, the basic polyhedrons are intervals when $d = 2$, are triangles when $d = 3$, and so on.

A polyhedron G with at most s vertices can be divided into at most s disjoint basic polyhedrons B_1, \dots, B_s . For instance, Figure 3 demonstrates

the $d = 2$ case. Therefore, the bracketing number of the polyhedrons can be derived by bracketing the basic polyhedrons. For a basic polyhedron B , denote its δ -bracketing pair to be $(U_{B,\delta}, V_{B,\delta})$, i.e., $U_{B,\delta} \subset B \subset V_{B,\delta}$. Then $(U_{G,\delta}, V_{G,\delta})$, defined as below

$$\begin{aligned} U_{G,\delta} &= U_{B_1,\delta} \cup U_{B_2,\delta} \cup \cdots \cup U_{B_s,\delta} \\ V_{G,\delta} &= V_{B_1,\delta} \cup V_{B_2,\delta} \cup \cdots \cup V_{B_s,\delta}, \end{aligned}$$

form a $(s\delta)$ -bracket of G . Hence, the bracketing number of all polyhedrons is controlled by the s -th power of the bracketing number of all basic polyhedrons. Applying Lemma 3.7 we know $s = O(N^{Ld})$ and the number of vertices is at most $S = O(N^{Ld^2})$. Together with Lemma 3.6, we therefore get that

$$\log \mathcal{N}_B(S\delta, \mathcal{G}^{\mathcal{F}}, d_{\Delta}) \lesssim S(d+1)d^2 \log((d+1)d^{3/2}/\delta),$$

which implies

$$\begin{aligned} \log \mathcal{N}_B(\delta, \mathcal{G}^{\mathcal{F}}, d_{\Delta}) &\lesssim N^{Ld^2} d^3 \log(N^{Ld^2} d^3 / \delta) \\ &\lesssim N^{Ld^2} d^3 (Ld^2 \log(N) \vee \log(1/\delta)). \end{aligned}$$

□

More discussions about Lemma 3.9 can be found in Appendix 6.2. Next, we present some lemmas that can take advantage of the obtained entropy bound and eventually take us to the proof of the excess risk convergence rate.

Lemma 3.10 (Theorem 5.11 in Van De Geer [2000]). For some function space \mathcal{H} with $\sup_{h \in \mathcal{H}} \|h(\mathbf{x})\|_{\infty} \leq K$ and $\sup_{h \in \mathcal{H}} \|h(\mathbf{x})\|_{L_2(P)} \leq R$ where P is the distribution of \mathbf{x} . Take $a > 0$ satisfying (1) $a \leq C_1 \sqrt{n} R^2 / K$; (2) $a \leq 8\sqrt{n} R$;

$$(3) \quad a \geq C_0 \left(\int_{a/64\sqrt{n}}^R H_B^{1/2}(u, \mathcal{F}, L_2(P)) du \vee R \right);$$

and (4) $C_0^2 \geq C^2(C_1 + 1)$. Then

$$\mathbb{P} \left(\sup_{h \in \mathcal{H}} \left| \sqrt{n} \int h d(P_n - P) \right| \geq a \right) \leq C \exp \left(-\frac{a^2}{C^2(C_1 + 1)R^2} \right),$$

where P_n is the empirical counterpart of P .

So far, the presented lemmas are only concerned with the general case, i.e. set G^* , p , q , etc. that does not depend on n . However, in our teacher-student framework, the optimal set G_n^* is indexed by n as it's determined by the teacher network \mathcal{F}_n^* . In the remaining part of the proof, we will consider specifically for our teacher network case. The next lemma investigates the modulus of continuity of the empirical process. It's similar to Lemma 5.13 in Van De Geer [2000] but with a key difference in the entropy assumption (3.3), where the entropy bound contains n .

Lemma 3.11. For a probability measure P , let \mathcal{H}_n be a class of uniformly bounded (by 1) functions h in $L_2(P)$ depending on n . Suppose that the δ -entropy with bracketing $H_B(\delta, \mathcal{H}_n, L_2(P))$ satisfies, for some $A_n > 0$, the inequality

$$(3.3) \quad H_B(\delta, \mathcal{H}_n, L_2(P)) \leq A_n \log(1/\delta)$$

for all $\delta > 0$ small enough. Let h_{n0} be a fixed element in \mathcal{H}_n . Let $\mathcal{H}_n(\delta) = \{h_n \in \mathcal{H}_n : \|h_n - h_{n0}\|_{L_2(P)} \leq \delta\}$. Then there exist constants $D_1 > 0, D_2 > 0$ such that for a sequence of i.i.d. random variables $\mathbf{x}_1, \dots, \mathbf{x}_n$ with probability distribution P , it holds that

$$\mathbb{P} \left(\sup_{\substack{h_n \in \mathcal{H}_n; \\ \|h_n - h_{n0}\| > \sqrt{\frac{A_n}{n}}}} \frac{\sum_{i=1}^n [(h_n - h_{n0})(\mathbf{x}_i) - \mathbb{E}(h_n - h_{n0})(\mathbf{x}_i)]}{\sqrt{\frac{A_n}{n}} \|h_n - h_{n0}\|_2 \log(\|h_n - h_{n0}\|_2^{-1})} > D_1 x \right) \leq D_2 e^{-A_n x}$$

for all $x \geq 1$.

PROOF. The main tool for the proof is Lemma 3.10. Replace \mathcal{H} with $\mathcal{H}_n(\delta)$ in Lemma 3.10 and take $K = 4$, $R = \sqrt{2}\delta$ and $a = \frac{1}{2}C_1\sqrt{A_n}\delta \log(1/\delta)$, with $C_1 = 2\sqrt{2}C_0$. Then (1) is satisfied if

$$(3.4) \quad \frac{\sqrt{A_n}}{\delta} \log\left(\frac{1}{\delta}\right) \leq \sqrt{n}.$$

Under (3.4), condition (2) and (3) are satisfied automatically. Choosing C_0 sufficiently large will ensure (4). Thus, for all δ satisfying (3.4), we have

$$\mathbb{P} \left(\sup_{h_n \in \mathcal{H}_n(\delta)} \left| \sqrt{n} \int (h_n - h_{n0}) d(P_n - P) \right| \geq \frac{C_1}{2} \sqrt{A_n} \delta \log(1/\delta) \right) \leq C \exp \left(-\frac{C_1 A_n \log^2(1/\delta)}{16C^2} \right)$$

Notice that (3.4) holds if $\delta \geq \sqrt{A_n/n}$. Let $B = \min\{b > 1 : 2^{-b} \leq \sqrt{A_n/n}\}$ and apply the peeling device. Then,

$$\begin{aligned} & \mathbb{P} \left(\sup_{\substack{h_n \in \mathcal{H}_n; \\ \|h_n - h_{n0}\| > \sqrt{A_n/n}}} \frac{|\sqrt{n} \int (h_n - h_{n0}) d(P_n - P)|}{\sqrt{A_n} \|h_n - h_{n0}\| \log(1/\|h_n - h_{n0}\|)} \geq \frac{C_2}{2} \right) \\ & \leq \sum_{b=1}^B \mathbb{P} \left(\sup_{h_n \in \mathcal{H}_n(2^{-b})} \left| \sqrt{n} \int (h_n - h_{n0}) d(P_n - P) \right| \geq \frac{C_1}{2} \sqrt{A_n} 2^{-b} \log(2^b) \right) \\ & \leq \sum_{b=1}^B C \exp \left(-\frac{C_1 A_n b^2 \log^2(2)}{16C^2} \right) \leq 2C \exp \left(-\frac{C_1 A_n}{16C^2} \right), \end{aligned}$$

if $C_1 A_n$ is sufficiently large. \square

We then present a lemma that establishes the connection between d_Δ and $d_{p,q}$, which is adapted from Lemma 2 in [Mammen et al. \[1999\]](#) to our teacher network setting. Corresponding to assumption (A3), we define (N_n) as an extension to the classical Tsybakov noise condition (N).

(N_n) There exists $c_n > 0$ depending on n and $\kappa \in [0, \infty]$ such that for any $0 \leq t \leq T_n$

$$\mathbb{P}(\{\mathbf{x} : |p_n(\mathbf{x}) - q_n(\mathbf{x})| \leq t\}) \leq c_n t^\kappa.$$

Note that the (N) is a special case of (N_n) with T_n and c_n being absolute constant.

Lemma 3.12. Assume (N_n) and p_n, q_n are bounded by $b_2 > 0$. Then, there exists absolute constants $b_1(\kappa) > 0$ depending on κ such that for any Lebesgue measurable subsets G_1 and G_2 of \mathcal{X} ,

$$b_1(\kappa) \left(T_n \wedge c_n^{-1/\kappa} \right) d_\Delta^{(\kappa+1)/\kappa}(G_1, G_2) \leq d_{p_n, q_n}(G_1, G_2) \leq 2b_2 d_\Delta(G_1, G_2).$$

PROOF. The second inequality is trivial given that p, q are bounded by b_2 . For the first inequality, since $\mathbb{Q}(|p_n - q_n| \leq t) \leq c_n t^\kappa$ for all $0 \leq t \leq T_n$, the boundedness of $\mathbb{Q}(\mathcal{X})$ implies that

$$\mathbb{Q}(|p_n - q_n| \leq t) \leq A_n t^\kappa, \quad \forall t > 0,$$

where $A_n = \left(\frac{\mathbb{Q}(\mathcal{X})}{T_n^\kappa} \vee c_n\right)$. Then,

$$\begin{aligned}
& d_{p_n, q_n}(G_1, G_2) \\
& \geq \int_{G_1 \Delta G_2} |p_n - q_n| \mathbb{I}\{|p_n - q_n| \geq \left(\frac{d_\Delta(G_1, G_2)}{2A_n}\right)^{1/\kappa}\} d\mathbb{Q} \\
& \geq \left(\frac{d_\Delta(G_1, G_2)}{2A_n}\right)^{1/\kappa} \left[\mathbb{Q}(G_1 \Delta G_2) - \mathbb{Q}(|p_n - q_n| < \left(\frac{d_\Delta(G_1, G_2)}{2A_n}\right)^{1/\kappa}) \right] \\
& \geq \frac{d_\Delta(G_1, G_2)^{1+1/\kappa}}{(2A_n)^{1/\kappa}} - 1/2 \frac{d_\Delta(G_1, G_2)^{(\kappa+1)/\kappa}}{(2A_n)^{1/\kappa}} \\
& \geq \frac{2^{-(\kappa+1)/\kappa}}{A_n^{1/\kappa}} d_\Delta(G_1, G_2)^{(\kappa+1)/\kappa}.
\end{aligned}$$

□

Our goal in classification is to estimate G_n^* by $\widehat{G}_n = \operatorname{argmin}_{G \in \mathcal{G}_n} R_n(G)$, where \mathcal{G}_n is some collection of sets associated with the student network \mathcal{F}_n and

$$R_n(G) = \frac{1}{2n} \sum_{i=1}^n (\mathbb{I}\{\mathbf{x}_i \in G | y_i = 1\}(\mathbf{x}) + \mathbb{I}\{\mathbf{x}_i \notin G | y_i = -1\}(\mathbf{x})).$$

Similar to Theorem 1 in [Mammen et al. \[1999\]](#), we have the following lemma regarding the upper bound on the rate of convergence.

Lemma 3.13. Suppose $0 < \mathbb{Q}(\mathcal{X}) < \infty$ and let \mathcal{G}_n^* be a collection of subsets of $\mathcal{X} \subset \mathbb{R}^d$. Define

$$\begin{aligned}
(3.5) \quad \mathcal{D}_n^{\mathcal{G}_n^*} &= \{(p_n, q_n) : \mathbb{Q}\{\mathbf{x} \in \mathcal{X} : |p_n(\mathbf{x}) - q_n(\mathbf{x})| \leq t\} \leq c_n t^\kappa \text{ for } 0 \leq t \leq T_n, \\
&\quad \{\mathbf{x} \in \mathcal{X} : p_n(\mathbf{x}) \geq q_n(\mathbf{x})\} \in \mathcal{G}_n^*, p_n(\mathbf{x}), q_n(\mathbf{x}) \leq b_2 \text{ for } \mathbf{x} \in \mathcal{X}\},
\end{aligned}$$

where b_2 is an absolute constant. Let \mathcal{G}_n be another class of subsets satisfying $\mathcal{G}_n^* \subset \mathcal{G}_n$. Suppose there exist positive constants $A_n > 0$ depending on n such that for any $\delta > 0$ small enough,

$$(3.6) \quad H_B(\delta, \mathcal{G}_n, d_\Delta) \leq A_n \log(1/\delta).$$

Then we have

$$(3.7) \quad \lim_{n \rightarrow \infty} \sup_{(p_n, q_n) \in \mathcal{D}_n^{\mathcal{G}_n^*}} \left(\frac{A_n \log^2 n}{n}\right)^{-\frac{\kappa+1}{\kappa+2}} \left(T_n \wedge c_n^{-1/\kappa}\right)^{\frac{\kappa}{\kappa+2}} \mathbb{E}[d_{p_n, q_n}(\widehat{G}_n, G_n^*)] < \infty.$$

PROOF. For $(p_n, q_n) \in \mathcal{F}_n^{G_n^*}$, let $G_n^* = \{\mathbf{x} \in \mathcal{X} : p_n(\mathbf{x}) \geq q_n(\mathbf{x})\}$. For a given set $G \in \mathcal{X}$, let $h_G(\mathbf{x}) = \mathbb{I}\{\mathbf{x} \in G\}$. In particular, let $h_n^* = h_{G_n^*}$. Let $\|h\|_p^2 = \int h^2(\mathbf{x})p(\mathbf{x})\mathbb{Q}(d\mathbf{x})$. Since both p_n and q_n are bounded,

$$(3.8) \quad \begin{aligned} \|h_{G_n} - h_n^*\|_p^2 &= \int_{G_n \Delta G_n^*} p_n(\mathbf{x})\mathbb{Q}(d\mathbf{x}) \leq b_2 d_\Delta(G_n, G_n^*), \\ \|h_{G_n} - h_n^*\|_q^2 &= \int_{G_n \Delta G_n^*} q_n(\mathbf{x})\mathbb{Q}(d\mathbf{x}) \leq b_2 d_\Delta(G_n, G_n^*). \end{aligned}$$

Consider the random variable

$$V_n = -\sqrt{n} \frac{R_n(\widehat{G}_n) - R_n(G_n^*) - \mathbb{E}(R_n(\widehat{G}_n) - R_n(G_n^*))}{\sqrt{A_n d_\Delta(G_n^*, \widehat{G}_n) \log(1/d_\Delta(G_n^*, \widehat{G}_n))}}.$$

Since $\mathcal{G}_n^* \subset \mathcal{G}_n$, we have $R_n(\widehat{G}_n) \leq R_n(G_n^*)$. Thus

$$(3.9) \quad \frac{\sqrt{n} \mathbb{E}(R_n(\widehat{G}_n) - R_n(G_n^*))}{\sqrt{A_n d_\Delta(G_n^*, \widehat{G}_n) \log(1/d_\Delta(G_n^*, \widehat{G}_n))}} \leq V_n.$$

Note that

$$\begin{aligned} R_n(G_n) - R_n(G_n^*) &= \frac{1}{2n} \sum_{i=1}^n \mathbb{I}_{\{y_i=1\}} (h_n^* - h_{G_n})(\mathbf{x}_i) \\ &\quad + \frac{1}{2n} \sum_{i=1}^n \mathbb{I}_{\{y_i=-1\}} (h_{G_n} - h_n^*)(\mathbf{x}_i). \end{aligned}$$

Then V_n can be written as

$$\begin{aligned} V_n &= \frac{(1/2n) \sum_{i=1}^n \mathbb{I}_{\{y_i=1\}} (h_{\widehat{G}_n} - h_n^*)(\mathbf{x}_i) - \mathbb{E}(\mathbb{I}_{\{y=1\}} (h_{\widehat{G}_n} - h_n^*)(\mathbf{x}))}{\sqrt{A_n d_\Delta(G_n^*, \widehat{G}_n)/n \log(1/d_\Delta(G_n^*, \widehat{G}_n))}} + \\ &\quad \frac{(1/2n) \sum_{i=1}^n \mathbb{I}_{\{y_i=-1\}} (h_n^* - h_{\widehat{G}_n})(\mathbf{x}_i) - \mathbb{E}(\mathbb{I}_{\{y=-1\}} (h_n^* - h_{\widehat{G}_n})(\mathbf{x}))}{\sqrt{A_n d_\Delta(G_n^*, \widehat{G}_n)/n \log(1/d_\Delta(G_n^*, \widehat{G}_n))}}. \end{aligned}$$

Consider the event $E_n = \{d_\Delta(G_n^*, \widehat{G}_n) > \sqrt{A_n/n}\}$ and let $\widetilde{\mathcal{G}}_n = \{G \in \mathcal{G}_n :$

$d_\Delta(G, G_n^*) > \sqrt{A_n/n}$. If E_n holds, then

$$\begin{aligned}
V_n &= -\sqrt{n} \frac{R_n(\widehat{G}_n) - R_n(G_n^*) - \mathbb{E}(R_n(\widehat{G}_n) - R_n(G_n^*))}{\sqrt{A_n d_\Delta(G_n^*, \widehat{G}_n) \log(1/d_\Delta(G_n^*, \widehat{G}_n))}} \\
&\leq \sup_{G_n \in \widetilde{\mathcal{G}}_n} \sqrt{n} \frac{R_n(G_n^*) - R_n(G_n) - \mathbb{E}(R_n(G_n) - R_n(G_n^*))}{\sqrt{A_n d_\Delta(G_n^*, \widehat{G}_n) \log(1/d_\Delta(G_n^*, \widehat{G}_n))}} \\
&\leq \sup_{G_n \in \widetilde{\mathcal{G}}_n} \frac{|(1/2n) \sum_{i=1}^n \mathbb{I}_{\{y_i=1\}}(h_{G_n} - h_n^*)(\mathbf{x}_i) - \mathbb{E}(\mathbb{I}_{\{y=1\}}(h_{G_n} - h_n^*)(\mathbf{x}))|}{\sqrt{A_n d_\Delta(G_n^*, \widehat{G}_n)/n \log(1/d_\Delta(G_n^*, \widehat{G}_n))}} + \\
&\quad \sup_{G_n \in \widetilde{\mathcal{G}}_n} \frac{|(1/2n) \sum_{i=1}^n \mathbb{I}_{\{y_i=-1\}}(h_{G_n} - h_n^*)(\mathbf{x}_i) - \mathbb{E}(\mathbb{I}_{\{y=-1\}}(h_{G_n} - h_n^*)(\mathbf{x}))|}{\sqrt{A_n d_\Delta(G_n^*, \widehat{G}_n)/n \log(1/d_\Delta(G_n^*, \widehat{G}_n))}} \\
&\leq \sup_{h_n \in \mathcal{H}_n} \frac{|(1/2n) \sum_{i=1}^n \mathbb{I}_{\{y_i=1\}}(h_n - h_n^*)(\mathbf{x}_i) - \mathbb{E}(\mathbb{I}_{\{y=1\}}(h_n - h_n^*)(\mathbf{x}))|}{2b_2^{-1/2} \sqrt{A_n/n} \|h_n - h_n^*\|_p \log(\sqrt{b_2}/\|h_n - h_n^*\|_p)} + \\
&\quad \sup_{h_n \in \mathcal{H}_n} \frac{|(1/2n) \sum_{i=1}^n \mathbb{I}_{\{y_i=1\}}(h_n - h_n^*)(\mathbf{x}_i) - \mathbb{E}(\mathbb{I}_{\{y=1\}}(h_n - h_n^*)(\mathbf{x}))|}{2b_2^{-1/2} \sqrt{A_n/n} \|h_n - h_n^*\|_q \log(\sqrt{b_2}/\|h_n - h_n^*\|_q)},
\end{aligned}$$

where $\mathcal{H}_n = \{h_n(\mathbf{x}) = \mathbb{I}\{\mathbf{x} \in G_n\} : G_n \in \mathcal{G}_n\}$. The last inequality follow from the fact that $\sqrt{x} \log(1/x)$ is strictly increasing when $x < 1$. Notice that h_n 's are uniformly bounded by 1 and the L_2 norm squared of $h_{G_1} - h_{G_2}$ is $d_\Delta(G_1, G_2)$. Applying Lemma 3.11, we have

$$(3.10) \quad \mathbb{E}[V_n \mathbb{I}(E_n)] \leq C$$

for some finite constant C . Now we use this inequality to prove the main result. From (3.9), we know that

$$d_{p_n, q_n}(\widehat{G}_n, G_n^*) \leq V_n (A_n/n)^{1/2} d_\Delta(G_n^*, \widehat{G}_n)^{1/2} \log(1/d_\Delta(G_n^*, \widehat{G}_n)),$$

which, together with Lemma 3.12, yields that

$$\begin{aligned}
d_{p_n, q_n}(\widehat{G}_n, G_n^*) &\lesssim V_n (A_n/n)^{1/2} \left(T_n \wedge c_n^{-1/\kappa}\right)^{-\frac{\kappa}{2(\kappa+1)}} d_{p_n, q_n}(\widehat{G}_n, G_n^*)^{\frac{\kappa}{2(\kappa+1)}} \\
&\quad \cdot [\log(1/d_{p_n, q_n}(\widehat{G}_n, G_n^*)) + \log(b_1(\kappa)(T_n \wedge c_n^{-1/\kappa}))],
\end{aligned}$$

which simplifies to be

$$d_{p_n, q_n}(\widehat{G}_n, G_n^*) \lesssim V_n^{\frac{2\kappa+2}{\kappa+2}} \left(\frac{A_n \log^2 n}{n}\right)^{\frac{\kappa+1}{\kappa+2}} \left(T_n \wedge c_n^{-1/\kappa}\right)^{-\frac{\kappa}{\kappa+2}}.$$

where we used the fact that $d_{p_n, q_n}(\widehat{G}_n, G_n^*) \gtrsim 1/n$. Therefore, under E_n , (3.10) implies that

$$\mathbb{E}[d_{p_n, q_n}(\widehat{G}_n, G_n^*)] \lesssim \left(\frac{A_n \log^2 n}{n} \right)^{\frac{\kappa+1}{\kappa+2}} \left(T_n \wedge c_n^{-1/\kappa} \right)^{-\frac{\kappa}{\kappa+2}}.$$

On the other hand, under E_n^c , we have

$$d_\Delta(\widehat{G}_n, G_n^*) \leq \sqrt{A_n/n}.$$

By Lemma 3.12 we know $d_{p, q}(\widehat{G}_n, G_n^*)$ is also bounded by $\sqrt{A_n/n}$. Since $(\kappa + 1)/(\kappa + 2) \leq 1$, the rate under E_n dominates and the proof is complete. \square

PROOF OF THEOREM 3.2. First, we verify that the Tsybakov noise condition holds for $\kappa = 1$ in our setting. The proof is based on the fact that a ReLU network is piecewise linear and the number of linear pieces is quantifiable. Assumption (A3) implies (N_n) with $c_n, 1/T_n = O(\log n)^{m^* d^2 L_n^*}$ and $\kappa = 1$. In the case where p, q have disjoint support, obviously κ can be arbitrarily large.

Next, we consider the bracketing number of \mathcal{G}_n defined via \mathcal{F}_n that $\mathcal{G}_n = \{\mathbf{x} \in \mathcal{X} : f(\mathbf{x}) \geq 0, f \in \mathcal{F}_n\}$. From Lemma 3.9 we have

$$\log \mathcal{N}_B(\delta, \mathcal{G}_n, d_\Delta) \lesssim N^{L d^2} d^2 (L d^2 \log(N) \vee \log(1/\delta)).$$

Thus, $A_n = O(N_n)^{d^2 L_n}$ as in (3.6) if $\delta \ll 1/N$. Recall from assumption (A2) and (A3) that $L_n = O(1)$, $N_n = O(\log n)^m$ and $1/T_n, c_n = O(\log n)^{m^* d^2 L_n^*}$. Applying Lemma 3.13 with $\kappa = 1$ we have that the excess risk has upper bound

$$\begin{aligned} & \sup_{(p, q) \in \widetilde{\mathcal{F}}_n^*} \mathbb{E}[\mathcal{E}(\widehat{f}_n, C_n^*)] \\ & \lesssim \left(\frac{A_n \log^2 n}{n} \right)^{\frac{2}{3}} (T_n^{-1} \wedge c_n)^{\frac{1}{3}} \\ & \lesssim \left(\frac{1}{n} \right)^{\frac{2}{3}} (\log n)^{\frac{2}{3}(m d^2 L_n + 2) + \frac{1}{3} m^* d^2 L_n^*}. \end{aligned}$$

\square

PROOF OF COROLLARY 3.4. Corollary 3.4 easily follows from the fact that p, q having disjoint support implies $\kappa = \infty$ in (N_n) . \square

3.3. *Proof of Theorem 3.3.* We will show that the lower bound holds in special case that (1) assumption (A3) satisfies $c_n, 1/T_n$ being absolute constants that doesn't depend on n ; (2) instead of general ReLU neural network $f_n^* \in \mathcal{F}^*$, we consider a special structure where f_n^* is linear in one of the dimensions, reminiscent of the ‘‘boundary fragment’’ assumption. In this special case, we are able to show the best possible convergence rate already matches that in Theorem 3.2. For ease of notation, we omit the subscript n and write p_n, q_n as p, q if no confusion arises.

PROOF. Without loss of generality, let $\mathcal{X} = [0, 1]^d$. Consider the ‘‘boundary fragment’’ setting and let $\tilde{\mathcal{G}}_n$ be a set defined by a ReLU network family $\tilde{\mathcal{F}}_n$ containing functions from \mathbb{R}^{d-1} to \mathbb{R} :

$$\tilde{\mathcal{G}}_n = \{(x_1, \dots, x_d) \in [0, 1]^d : 0 \leq x_j \leq h(\mathbf{x}_{-j}), h \in \tilde{\mathcal{F}}_n, j = 1, \dots, d\},$$

where $\mathbf{x}_{-j} = (x_1, \dots, x_{j-1}, x_j, \dots, x_d)$. Notice that if $h(\mathbf{x}_{-j})$ is a ReLU network on \mathbb{R}^{d-1} , then $\tilde{h}(\mathbf{x}) = h(\mathbf{x}_{-j}) - x_j$ is a ReLU network on \mathbb{R}^d . Thus $\tilde{\mathcal{G}}_n$ is a subset of \mathcal{G}_n which corresponds to the student network that

$$(3.11) \quad \mathcal{G}_n = \{\mathbf{x} \in \mathcal{X} : f(\mathbf{x}) > 0, f \in \mathcal{F}_n\}$$

Let \tilde{G}_n denote the empirical 0-1 loss minimizer over $\tilde{\mathcal{G}}_n$. To show the lower bound, consider the subset of $\mathcal{D}^{\tilde{\mathcal{G}}_n}$ (3.5) that contains all pairs like (p, q_0) , where $p \in \mathcal{F}_1, q_0$ will be specified later. Then,

$$\begin{aligned} \sup_{(p,q) \in \mathcal{D}^{\tilde{\mathcal{G}}_n}} \mathbb{E}d_{\Delta}(\tilde{G}_n, G^*) &\geq \sup_{(p,q_0): p \in \mathcal{F}_1} \mathbb{E}d_{\Delta}(\tilde{G}_n, G^*) \\ &\geq \mathbb{E}_{q_0} \left[\frac{1}{|\mathcal{F}_1|} \sum_{p \in \mathcal{F}_1} \mathbb{E}_p[d_{\Delta}(\tilde{G}_n, G^*) | \mathcal{D}_{q_0}] \right], \end{aligned}$$

where \mathcal{F}_1 is a finite set to be specified later, p, q_0 are the underlying densities for the two labels and \mathcal{D}_{q_0} denotes all the data generated from q_0 .

For ease of presentation, we first give the proof for the case $d = 2$ and then extend to general d . Let $\phi(t)$ be a piecewise linear function supported on $[-1, 1]$ defined as

$$\phi(t) = \begin{cases} t+1 & -1 < t \leq 0, \\ -t+1 & 0 < t < 1, \\ 0 & |t| \geq 1. \end{cases}$$

Rewrite ϕ as $\phi(t) = \sigma(t+1) - \sigma(t) + \sigma(-t+1) - \sigma(-t) - 2$, which is a one hidden layer ReLU neural network with 11 non-zero weights that are either 1 or -1 . For $\mathbf{x} = (x_1, x_2) \in [0, 1]^2$, define

$$q_0(\mathbf{x}) = (1 - \eta_0 - b_1)\mathbb{I}\{0 \leq x_2 < 1/2\} + \mathbb{I}\{1/2 \leq x_2 < 1/2 + e^{-M}\} \\ + (1 + \eta_0 + b_2)\mathbb{I}\{1/2 + e^{-M} \leq x_2 \leq 1\},$$

where $M \geq 2$ is an integer to be specified later. Let $b_1 = c_2^{-1/\kappa} e^{-M/\kappa}$ and $b_2 > 0$ be chosen such that q_0 integrates to 1 (so q_0 is a valid probability density).

For $j = 1, 2, \dots, M$ and $t \in [0, 1]$, let

$$\psi_j(t) = e^{-M} \phi\left(M \left[t - \frac{j-1}{M}\right]\right).$$

Note that ψ_j is only supported on $[\frac{j-1}{M}, \frac{j}{M}]$. For any vector $\omega = (\omega_1, \dots, \omega_M) \in \Omega := \{0, 1\}^M$, define

$$b_\omega(t) = \sum_{j=1}^M \omega_j \psi_j(t),$$

and

$$p_\omega(\mathbf{x}) = 1 + \left[\frac{1/2 + e^{-M} - x_2}{c_2}\right]^{1/\kappa} \mathbb{I}\{1/2 \leq x_2 \leq 1/2 + b_\omega(x_1)\} \\ - b_3(\omega) \mathbb{I}\{1/2 + b_\omega(x_1) < x_2 \leq 1\},$$

where $b_3(\omega) > 0$ is a constant depending on ω chosen such that $p_\omega(x)$ integrates to 1. Let $\mathcal{F}_1 = \{p_\omega : \omega \in \Omega\}$ and we will show that $(p_\omega, q_0) \in \mathcal{D}^{\tilde{\mathcal{G}}_n}$ for all $\omega \in \Omega$.

To this end, we need to verify that

- (a) $p_\omega(\mathbf{x}) \leq c_1$ for $\mathbf{x} \in [0, 1]^2$;
- (b) $\{\mathbf{x} \in \mathcal{X} : p_\omega(\mathbf{x}) \geq q_0(\mathbf{x})\} \in \mathcal{G}_n$;
- (c) $\mathbb{Q}\{\mathbf{x} \in \mathcal{X} : |p_\omega(\mathbf{x}) - q_0(\mathbf{x})| \leq \eta\} \leq c_2 \eta^\kappa$.

For (a), since p_ω integrates to 1,

$$b_3(\omega) \leq \max_{\{1/2 \leq x_2 \leq 1/2 + b_\omega(x_1)\}} \left[\frac{1/2 + e^{-M} - x_2}{c_2}\right]^{1/\kappa} = O(e^{-M/\kappa}).$$

Thus, $p_\omega(\mathbf{x}) \leq c_1$ for a large enough M and some absolute constant c_1 . For (b), notice that

$$\{\mathbf{x} : p_\omega(\mathbf{x}) \geq q_0(\mathbf{x})\} = \{\mathbf{x} : 0 \leq x_2 \leq 1/2 + b_\omega(x_1)\} \\ = \{\mathbf{x} \in [0, 1]^2 : b_\omega(x_1) - \sigma(x_2) + 1/2 \geq 0\} \in \mathcal{G}_n,$$

where the last inclusion follows from the definition of \mathcal{G}_n (3.11) and the fact that $b_\omega(x_1) - \sigma(x_2) + 1/2$ is a ReLU neural network with one hidden layer, whose width and number of non-zero weights are both $O(M)$. Later we will see that $M = O(\log n)$, and thus the constructed neural network satisfies all the size constraints in Theorem 3.2. For (c), it follows that

$$\begin{aligned} & Q\{\mathbf{x} \in \mathcal{X} : |p_\omega(\mathbf{x}) - q_0(\mathbf{x})| \leq \eta\} \\ & \leq Q\{\mathbf{x} \in \mathcal{X} : 1/2 \leq x_2 \leq 1/2 + e^{-M}, \left[\frac{1/2 + e^{-M} - x_2}{c_2} \right]^{1/\kappa} \leq \eta\} \\ & \leq Q\{\mathbf{x} \in \mathcal{X} : 1/2 + e^{-M} - c_2\eta^\kappa \leq x_2 \leq 1/2 + e^{-M}\} \\ & \leq c_2\eta^\kappa. \end{aligned}$$

Since the above (a)-(c) hold and by the definition of $\mathcal{D}^{\tilde{\mathcal{G}}_n}$ (3.5), we conclude that $(p_\omega, q_0) \in \mathcal{D}^{\tilde{\mathcal{G}}_n}$ for all $\omega \in \Omega$. We next establish how fast $S := |\mathcal{F}_1|^{-1} \sum_{p \in \mathcal{F}_1} \mathbb{E}_p[d_\Delta(\tilde{G}_n, G^*) | \mathcal{D}_{q_0}]$ can converge to zero. To this end, we use the Assouad's lemma stated in [Korostelev and Tsybakov, 2012] which is adapted to the estimation of sets.

For $j = 1, \dots, M$ and $\omega = (\omega_1, \dots, \omega_M) \in \Omega$, let

$$\begin{aligned} \omega_{j0} &= (\omega_1, \dots, \omega_{j-1}, 0, \omega_{j+1}, \dots, \omega_M) \\ \omega_{j1} &= (\omega_1, \dots, \omega_{j-1}, 1, \omega_{j+1}, \dots, \omega_M) \end{aligned}$$

For $i = 0$ and $i = 1$, let P_{ji} be the probability measure corresponding to the distribution of x_1, \dots, x_n when the underlying density is $f_{\omega_{ji}}$. Denote the expectation w.r.t. P_{ji} as \mathbb{E}_{ji} . Let

$$\begin{aligned} \mathcal{D}_j &= \{\mathbf{x} \in \mathcal{X} : 1/2 + b_{\omega_{j0}}(x_1) < x_2 \leq 1/2 + b_{\omega_{j1}}(x_1)\} \\ &= \{\mathbf{x} \in \mathcal{X} : b_{\omega_{j0}}(x_1) < x_2 - 1/2 \leq b_{\omega_{j0}}(x_1) + \psi_j(x_1)\}. \end{aligned}$$

Then

$$\begin{aligned} S &\geq 1/2 \sum_{j=1}^M \mathbb{Q}(\mathcal{D}_j) \int \min\{dP_{j1}, dP_{j0}\} \\ &\geq 1/2 \sum_{j=1}^M \int_0^1 \psi_j(x_1) dx_1 \int \min\{dP_{j1}, dP_{j0}\} \\ &\geq 1/2 \sum_{j=1}^M e^{-M} \int \phi(Mt) dt \int \min\{dP_{j1}, dP_{j0}\} \\ &\geq \frac{1}{4} \sum_{j=1}^M e^{-M} \int \phi(Mt) dt [1 - H^2(P_{10}, P_{11})/2]^n, \end{aligned}$$

where $H(\cdot, \cdot)$ denotes the Hellinger distance. Then it holds that

$$\begin{aligned}
H^2(P_{10}, P_{11}) &= \int \left[\sqrt{f_{\omega_{10}}(\mathbf{x})} - \sqrt{f_{\omega_{11}}(\mathbf{x})} \right]^2 d\mathbf{x} \\
&\leq \int_0^1 \left\{ \int_{1/2}^{1/2+\psi_1(x_1)} \left[1 - \sqrt{1 + \left(\frac{1/2 + e^{-M} - x_2}{c_2} \right)^{1/\kappa}} \right]^2 dx_2 \right. \\
&\quad \left. + \int_{1/2}^1 \left[\sqrt{1 - b_3(\omega_{10})} - \sqrt{1 - b_3(\omega_{11})} \right]^2 dx_2 \right\} dx_1 \\
&\leq \int_0^1 \int_{e^{-M}-\psi_1(x_1)}^{e^{-M}} \left[1 - \sqrt{1 + \left(\frac{v}{c_2} \right)^{1/\kappa}} \right]^2 dv dx_1 \\
&\quad + |b_3(\omega_{10}) - b_3(\omega_{11})|^2.
\end{aligned}$$

We will analyze the last two terms. For the first term,

$$\begin{aligned}
&\int_0^1 \int_{e^{-M}-\psi_1(x_1)}^{e^{-M}} \left[1 - \sqrt{1 + \left(\frac{v}{c_2} \right)^{1/\kappa}} \right]^2 dv dx_1 \\
&\leq \int_0^1 \int_{e^{-M}-\psi_1(x_1)}^{e^{-M}} \left(\frac{v}{c_2} \right)^{2/\kappa} dv dx_1 \\
&\leq \frac{\kappa c_2^{-2/\kappa}}{\kappa + 2} \int_0^1 (e^{-M})^{1+2/\kappa} - (e^{-M} - \psi_1(x_1))^{1+2/\kappa} dx_1 \\
&\leq \frac{\kappa c_2^{-2/\kappa}}{\kappa + 2} (e^{-M})^{1+2/\kappa} \int (1 - (1 - \phi(Mt))^{1+2/\kappa}) dt \\
&= O\left(\frac{1}{M} e^{-M(1+2/\kappa)} \right).
\end{aligned}$$

For the second term, notice that

$$\int_0^1 \int_{1/2}^{1/2+b_\omega(x_1)} \left[\frac{1/2 + e^{-M} - x_2}{c_2} \right]^{1/\kappa} dx_2 dx_1 = b_3(\omega) [1/2 - b_\omega(x_1)]$$

which yields

$$\begin{aligned}
b_3(\omega_{11}) &= \frac{1}{1/2 - b_{\omega_{11}}(x_1)} \int_0^1 \int_{1/2}^{1/2+b_{\omega_{11}}(x_1)} \left[\frac{1/2 + e^{-M} - x_2}{c_2} \right]^{1/\kappa} dx_2 dx_1 \\
&\leq \frac{M c_2^{-1/\kappa}}{1/2 - e^{-M}} \int_0^1 \int_{e^{-M}(1-\phi(Mx_1))}^{e^{-M}} u^{1/\kappa} du dx_1 \\
&= \frac{M c_2^{-1/\kappa}}{(1/2 - e^{-M})(1 + 1/\kappa)} e^{-M(1+1/\kappa)} \int (1 - (1 - \phi(Mt))^{1+1/\kappa}) dt \\
&\leq \frac{c_2^{-1/\kappa}}{(1/2 - e^{-M})(1 + 1/\kappa)} e^{-M(1+1/\kappa)} \\
&= O\left(e^{-M(1+1/\kappa)}\right).
\end{aligned}$$

Hence, $|b_3(\omega_{11}) - b_3(\omega_{10})| = O\left(e^{-M(1+1/\kappa)}\right)$. Unifying the above, we have

$$\begin{aligned}
H^2(P_{10}, P_{11}) &= O\left(\frac{1}{M} e^{-M(1+2/\kappa)} \vee e^{-M(2+2/\kappa)}\right) \\
&= O\left(\frac{1}{M} e^{-M(1+2/\kappa)}\right).
\end{aligned}$$

Now choose M as the smallest integer such that

$$M \geq \frac{\kappa}{\kappa + 2} \log n.$$

Then we have $H^2(P_{10}, P_{11}) \leq C^* n^{-1} (1 + o(1))$ for some constant C^* depending only on κ, c_2, ϕ , and

$$\int \min\{dP_{j1}, dP_{j0}\} \geq 1/2 \left[1 - \frac{C^*}{2} n^{-1} (1 + o(1)) \right]^n \geq C_1^*$$

for n large enough and C_1^* is another absolute constant depending only on C^* . Thus for n large enough,

$$S \geq \frac{1}{4} C_1^* e^{-M} \int \phi(t) dt \geq C_2^* n^{-\frac{\kappa}{\kappa+2}},$$

in which the constant C_2^* only depends on κ, c_2 and ϕ .

Combining all the results so far we get that

$$\liminf_{n \rightarrow \infty} \inf_{\tilde{G}_n} \sup_{(p,q) \in \mathcal{D}_{\tilde{G}_n}} n^{\frac{\kappa}{\kappa+2}} \mathbb{E}[d_{\Delta}(\tilde{G}_n, G^*)] > 0,$$

which holds when $d = 2$. Using Lemma 3.12, we have

$$\liminf_{n \rightarrow \infty} \inf_{\tilde{G}_n} \sup_{(p,q) \in \mathcal{D}_{\tilde{G}_n}} n^{\frac{\kappa+1}{\kappa+2}} \mathbb{E}[d_{p,q}(\tilde{G}_n, G^*)] > 0.$$

Using the same argument as in the proof of Theorem 3.2, we get $\kappa = 1$, which will give us the rate $2/3$.

The proof for general d can be derived similarly. We treat the last dimension x_d as x_2 in the $d = 2$ case and treat $\mathbf{x}_{-d} := (x_1, \dots, x_{d-1})$ as x_1 in the $d = 2$ case. Define

$$\begin{aligned} q_0(\mathbf{x}) &= (1 - \eta_0 - b_1) \mathbb{I}\{0 \leq x_d < 1/2\} + \mathbb{I}\{1/2 \leq x_d < 1/2 + e^{-M}\} \\ &\quad + (1 + \eta_0 + b_2) \mathbb{I}\{1/2 + e^{-M} \leq x_d \leq 1\}, \end{aligned}$$

and

$$\begin{aligned} p_\omega(\mathbf{x}) &= 1 + \left[\frac{1/2 + e^{-M} - x_2}{c_2} \right]^{1/\kappa} \mathbb{I}\{1/2 \leq x_d \leq 1/2 + \mathbf{b}_\omega(\mathbf{x}_{-d})\} \\ &\quad - b_3(\omega) \mathbb{I}\{1/2 + \mathbf{b}_\omega(\mathbf{x}_{-d}) < x_d \leq 1\}, \end{aligned}$$

where $\mathbf{b}_\omega(\mathbf{x}_{-d})$ is constructed similarly as a shallow ReLU neural network that

$$\mathbf{b}_\omega(\mathbf{x}_{-d}) = \sum_{j_1, \dots, j_{d-1}=1}^M \omega_{j_1, \dots, j_{d-1}} \psi_{j_1, \dots, j_{d-1}}(\mathbf{x}_{-d}),$$

where $\omega_{j_1, \dots, j_{d-1}}$ are binary 0, 1 variables and

$$\psi_{j_1, \dots, j_{d-1}}(\mathbf{x}_{-d}) = e^{-M} \phi \left(M \left[\mathbf{x}_{-d} - \left(\frac{j_1 - 1}{M}, \dots, \frac{j_{d-1} - 1}{M} \right) \right] \right),$$

where $\phi(\cdot)$ is a shallow ReLU neural network with input dimension $d - 1$ satisfying the following conditions:

- $\phi = 0$ outside $[-1, 1]^d$ and $\phi \leq 1$ on $[-1, 1]^d$;
- $\max_{\mathbf{x}_{-d} \in [-1, 1]^d} \phi(\mathbf{x}_{-d}) \leq 1$ and $\phi(\mathbf{0}) = 1$.

Such a construction is similar to the ‘‘spike’’ function in Yarotsky and Zhevnerchuk [2019] and it requires $O(d^2)$ non-zero weights. The rest of the proof follows the $d = 2$ case. \square

4. Training with Surrogate Loss. In this section, we consider deep classifiers trained under the hinge loss $\phi(z) = (1 - z)_+ = \max\{1 - z, 0\}$. This kind of surrogate loss has been widely used for “maximum-margin” classification, most notably for support vector machines [Cortes and Vapnik, 1995]. An desirable property of hinge loss is that its optimal classifier coincides with that under 0-1 loss [Lin, 2002], i.e. $f_\phi^*(\mathbf{x}) = C^*(\mathbf{x})$. Hence, a lot of arguments for 0-1 loss can be easily carried over. Additionally, minimizing the sample average of an appropriately behaved loss function has a regularizing effect [Bartlett et al., 2006]. It is thus possible to obtain uniform upper bounds on the risk of a function that minimizes the empirical average of the loss ϕ , even for rich classes that no such upper bounds are possible for the minimizer of the empirical average of the 01 loss.

Under the surrogate loss, our requirement on the size of the teacher network is relaxed from (A2) as follows:

(A2 $_\phi$) $p(\mathbf{x}) - q(\mathbf{x})$ is representable by some teacher ReLU DNN $f_n^* \in \mathcal{F}_n^*$ with

$$N_n^* = O(\log n)^{m_*}, \quad L_n^* = O(\log n), \quad B_n^*, F_n^* = O(\sqrt{n})$$

for some $m_* \geq 1$.

The following theorem says that the same un-improvable rate can be obtained for the empirical hinge loss minimizer $\hat{f}_{\phi,n} \in \mathcal{F}_n$.

Theorem 4.1. Suppose the underlying densities p and q satisfy assumptions (A1), (A2 $_\phi$), (A3) and denote all such (p, q) pairs as $\tilde{\mathcal{F}}_n^*$. Let \mathcal{F}_n be a student ReLU DNN family with $L_n = O(\log n)$, $N_n = O(\log n)^m$ and $B_n, F_n = O(\log n)$ for some $m \geq m_*$. Assume the student network is larger than the teacher network, i.e., $L_n \geq L_n^*$, $S_n \geq S_n^*$, $N_n \geq N_n^*$, $B_n \geq B_n^*$, $F_n \geq F_n^*$. Then the excess risk for $\hat{f}_{\phi,n} \in \mathcal{F}_n$ satisfies

$$\sup_{(p,q) \in \tilde{\mathcal{F}}_n^*} \mathbb{E}[\mathcal{E}(\hat{f}_{\phi,n}, C_n^*)] \asymp \tilde{O}_d \left(\frac{1}{n} \right)^{\frac{2}{3}}$$

Similarly, results in Corollary 3.4 and 3.5 hold for the empirical hinge loss minimizer. Specifically, when p, q are disjoint, the convergence rate of excess risk improves to n^{-1} , and all conclusions hold when the teacher network is larger but with bounded active pieces.

Remark 2 (Network Depth). Training with surrogate loss such as hinge loss, unlike 0-1 loss, doesn’t involve any hard thresholding, i.e. $\mathbb{I}\{yf(\mathbf{x}) < 0\}$. As a result, to control the complexity of the student network, Lemma 4.4 is used instead of Lemma 3.9, which allows us to use deeper neural networks ($L_n = O(\log n)$) for both the student and teacher network.

4.1. *Proof of Theorem 4.1.* One important observation to be used in the proof is that the Bayes classifier under hinge loss is the same as that under 0-1 loss, i.e. $f_\phi^*(\mathbf{x}) = C^*(\mathbf{x})$. To show the upper bound on excess risk convergence rate, we utilize the following lemma from Kim et al. [2018]. Let $\eta(\mathbf{x})$ denote the conditional probability of label 1 that $\eta(\mathbf{x}) = \mathbb{P}(y = 1|\mathbf{x})$.

Lemma 4.2. [Theorem 6 of [Kim et al., 2018]] Let ϕ be the hinge loss. Assume (N) with the noise exponent $\kappa \in [0, \infty]$, and that following conditions (C1) through (C4) hold.

- (C1) For a positive sequence $a_n = O(n^{-a_0})$ as $n \rightarrow \infty$ for some $a_0 > 0$, there exists a sequence of function classes $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ such that $\mathcal{E}_\phi(f_n, f_\phi^*) \leq a_n$ for some $f_n \in \mathcal{F}_n$.
- (C2) There exists a real valued sequence $\{F_n\}_{n \in \mathbb{N}}$ with $F_n \gtrsim 1$ such that $\sup_{f \in \mathcal{F}_n} \|f\|_\infty \leq F_n$.
- (C3) There exists a constant $\nu \in (0, 1]$ such that for any $f \in \mathcal{F}_n$ and any $n \in \mathbb{N}$,

$$\mathbb{E} \left[\left\{ \phi(Yf(\mathbf{X})) - \phi(Yf_\phi^*(\mathbf{X})) \right\}^2 \right] \leq C_2 F_n^{2-\nu} \{\mathcal{E}_\phi(f, f_\phi^*)\}^\nu$$

for a constant $C_2 > 0$ depending only on ϕ and $\eta(\cdot)$.

- (C4) For a positive constant $C_3 > 0$, there exists a sequence $\{\delta_n\}_{n \in \mathbb{N}}$ such that

$$H_B(\delta_n, \mathcal{F}_n, \|\cdot\|_2) \leq C_3 n \left(\frac{\delta_n}{F_n} \right)^{2-\nu},$$

for $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ in (C1), $\{F_n\}_{n \in \mathbb{N}}$ in (C2), and ν in (C3).

Let $\epsilon_n^2 \asymp \max\{a_n, \delta_n\}$. Assume that $n^{1-\iota} (\epsilon_n^2 / F_n)^{(\kappa+2)/(\kappa+1)} \gtrsim 1$ for an arbitrarily small constant $\iota > 0$. Then, the empirical ϕ -risk minimizer $\widehat{f}_{\phi,n}$ over \mathcal{F}_n satisfies

$$\mathbb{E} \left[\mathcal{E}(\widehat{f}_{\phi,n}, C^*) \right] \lesssim \epsilon_n^2.$$

In Lemma 4.2, condition (C1) guarantees the approximation error of f_n to f_ϕ^* to be sufficiently small. For condition (C3), we introduce the following lemma, which is reminiscent of Lemma 3.12 in the sense that it characterizes the relationship between the $\mathcal{E}_\phi(f, f_\phi^*)$ and the some other distance measure between f and f_ϕ^* .

Lemma 4.3 (Lemma 6.1 of Steinwart et al. [2007]). Assume (N) with the Tsybakov noise exponent $\kappa \in [0, \infty]$. Assume $\|f\|_\infty \leq F$ for any $f \in \mathcal{F}$.

Under the hinge loss ϕ , for any $f \in \mathcal{F}$,

$$\begin{aligned} & \mathbb{E} \left[(\phi(Yf(\mathbf{x})) - \phi(Yf_\phi^*(\mathbf{x})))^2 \right] \\ & \leq C_{\eta, \kappa} (F + 1)^{(\kappa+2)/(\kappa+1)} \left(\mathbb{E} [\phi(Yf(\mathbf{x})) - \phi(Yf_\phi^*(\mathbf{x}))] \right)^{\kappa/\kappa+1}, \end{aligned}$$

where $C_{\eta, \kappa} = (\|(2\eta - 1)^{-1}\|_{\kappa, \infty}^\kappa + 1) \mathbb{I}(\kappa > 0) + 1$ and $\|(2\eta - 1)^{-1}\|_{\kappa, \infty}^\kappa$ is defined by

$$\|(2\eta - 1)^{-1}\|_{\kappa, \infty}^\kappa = \sup_{t > 0} (t^\kappa \Pr(\{\mathbf{x} : |(2\eta(\mathbf{x}) - 1)^{-1}| > t\})).$$

For condition (C4) in Lemma 4.2, we present the following lemma.

Lemma 4.4. [Lemma 3 in Suzuki [2018]] For any $\delta > 0$, the covering number of $\mathcal{F}^{DNN}(L, N, S, B)$ (in sup-norm) satisfies

$$\begin{aligned} & \log \mathcal{N}(\delta, \mathcal{F}^{DNN}(L, N, S, B), \|\cdot\|_\infty) \\ & \leq 2L(S + 1) \log(\delta^{-1}(L + 1)(N + 1)(B \vee 1)). \end{aligned}$$

PROOF OF THEOREM 4.1. The lower bound directly follows from Theorem 3.3, as the constructed ReLU neural network in the proof also satisfy assumption (A2 $_\phi$).

For the upper bound on the convergence rate, we utilize Lemma 4.2 and check the conditions (C1) through (C4). Since the student network is larger than the teacher, (C1) and (C2) trivially hold with arbitrarily small a_n and $F_n = O(\log n)$ as assumed. To apply Lemma 4.3, notice that $C_{\eta, \kappa} = O(c_n) = O(\log n)^{m^* d^2 L_n^*}$ by assumption (A3) and $F = O(\log n)$, we have (C3) holds for $\nu = \kappa/(\kappa + 1) + \epsilon_n$, where $\epsilon_n = (2 + m^* d^2 L_n^*) \log \log n / \log n$. The term ϵ_n is to deal with the fact that $C_{\eta, \kappa}$ can also diverge at an $O(\log n)^{m^* d^2 L_n^*}$ rate.

For (C4), by Lemma 4.4,

$$\begin{aligned} & \log \mathcal{N}(\delta_n, \mathcal{F}^{DNN}(L_n, N_n, S_n, B_n, F_n), \|\cdot\|_\infty) \\ & \leq 2L_n(S_n + 1) \log(\delta_n^{-1}(L_n + 1)(N_n + 1)(B_n \vee 1)) \\ & \lesssim (\log n)^{2m+2} \log(\delta_n^{-1} \vee \log^m(n)). \end{aligned}$$

Therefore, (4.2) implies that (C3) is satisfied if we choose δ_n with

$$\delta_n^{\frac{\kappa+2}{\kappa+1}} \gtrsim \frac{(\log n)^{2m+2+(\kappa+2)/(\kappa+1)+2+m^* d^2 L_n^*+1}}{n},$$

which can be satisfied by choosing

$$\delta_n = \left(\frac{(\log n)^{2m+m^* d^2 L_n^*+7}}{n} \right)^{\frac{\kappa+1}{\kappa+2}}.$$

Similar to the proof of Theorem 3.2, the Tsybakov exponent $\kappa = 1$. Thus, by Lemma 4.2 with $\epsilon_n^2 = \delta_n$, the proof of Theorem 4.1 is completed. \square

5. Discussion and Future Directions. In this work, we obtain a sharp rate of convergence for the excess risk under both empirical 0-1 loss and hinge loss minimizer in the teacher-student setting. Our current results for training under 0-1 loss only hold for student networks with $O(1)$ layers and the assumption that $f_n^* \in \mathcal{F}_n$, i.e. zero approximation, is required. In the future, we aim to relax these two constraints and provide more comprehensive analysis of the teacher-student network. Additionally, we would like to

- explore other type of neural networks such as convolutional neural network and residual neural network, which are both very successful at image classification;
- consider the implicit bias of training algorithms, e.g. stochastic gradient descent, to regularize the complexity of larger and deeper neural networks in the teacher-student setting;
- consider the more general improper learning scenario where the Bayes classifier is not necessarily in the student neural network;
- consider other popular surrogate losses such as exponential loss or cross entropy loss.

Further investigation under the teacher-student network setting may facilitate a better understanding of how deep neural network works and shed light on its empirical success especially in high-dimensional image classification.

References.

- Raman Arora, Amitabh Basu, Poorya Mianjy, and Anirbit Mukherjee. Understanding deep neural networks with rectified linear units. *arXiv preprint arXiv:1611.01491*, 2016.
- Benjamin Aubin, Antoine Maillard, Jean Barbier, Florent Krzakala, Nicolas Macris, and Lenka Zdeborová. The committee machine: Computational to statistical gaps in learning a two-layers neural network. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems 31*, pages 3223–3234. Curran Associates, Inc., 2018.
- Jean-Yves Audibert, Alexandre B Tsybakov, et al. Fast learning rates for plug-in classifiers. *The Annals of statistics*, 35(2):608–633, 2007.
- Jimmy Ba and Rich Caruana. Do deep nets really need to be deep? In Z. Ghahramani, M. Welling, C. Cortes, N. D. Lawrence, and K. Q. Weinberger, editors, *Advances in Neural Information Processing Systems 27*, pages 2654–2662. Curran Associates, Inc., 2014.
- Jincheng Bai, Qifan Song, and Guang Cheng. Rate optimal variational bayesian inference for sparse dnn. *arXiv preprint arXiv:1910.04355*, 2019.
- Peter L Bartlett, Michael I Jordan, and Jon D McAuliffe. Convexity, classification, and risk bounds. *Journal of the American Statistical Association*, 101(473):138–156, 2006.

- Benedikt Bauer, Michael Kohler, et al. On deep learning as a remedy for the curse of dimensionality in nonparametric regression. *The Annals of Statistics*, 47(4):2261–2285, 2019.
- Yuan Cao and Quanquan Gu. Tight sample complexity of learning one-hidden-layer convolutional neural networks. In *Advances in Neural Information Processing Systems*, pages 10611–10621, 2019.
- Minshuo Chen, Haoming Jiang, Wenjing Liao, and Tuo Zhao. Efficient approximation of deep relu networks for functions on low dimensional manifolds. In *Advances in Neural Information Processing Systems*, pages 8172–8182, 2019.
- Corinna Cortes and Vladimir Vapnik. Support-vector networks. *Machine learning*, 20(3): 273–297, 1995.
- George Cybenko. Approximations by superpositions of a sigmoidal function. *Mathematics of Control, Signals and Systems*, 2:183–192, 1989.
- J. Deng, W. Dong, R. Socher, L.-J. Li, K. Li, and L. Fei-Fei. ImageNet: A Large-Scale Hierarchical Image Database. In *CVPR09*, 2009a.
- Jia Deng, Wei Dong, Richard Socher, Li-Jia Li, Kai Li, and Li Fei-Fei. Imagenet: A large-scale hierarchical image database. In *2009 IEEE conference on computer vision and pattern recognition*, pages 248–255. Ieee, 2009b.
- A Engel and C.V. Broeck. *Statistical Mechanics of Learning*. Cambridge University Press, 2002.
- Max H Farrell, Tengyuan Liang, and Sanjog Misra. Deep neural networks for estimation and inference: Application to causal effects and other semiparametric estimands. *arXiv preprint arXiv:1809.09953*, 2018.
- Sebastian Goldt, Madhu Advani, Andrew M Saxe, Florent Krzakala, and Lenka Zdeborová. Dynamics of stochastic gradient descent for two-layer neural networks in the teacher-student setup. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d’Alché-Buc, E. Fox, and R. Garnett, editors, *Advances in Neural Information Processing Systems 32*, pages 6979–6989. Curran Associates, Inc., 2019.
- Suriya Gunasekar, Jason D Lee, Daniel Soudry, and Nati Srebro. Implicit bias of gradient descent on linear convolutional networks. In *Advances in Neural Information Processing Systems*, pages 9461–9471, 2018.
- Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep residual learning for image recognition. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 770–778, 2016.
- Geoffrey Hinton, Oriol Vinyals, and Jeff Dean. Distilling the knowledge in a neural network. *arXiv preprint arXiv:1503.02531*, 2015.
- Masaaki Imaizumi and Kenji Fukumizu. Deep neural networks learn non-smooth functions effectively. *arXiv preprint arXiv:1802.04474*, 2018.
- Yongdai Kim, Ilsang Ohn, and Dongha Kim. Fast convergence rates of deep neural networks for classification. *arXiv preprint arXiv:1812.03599*, 2018.
- Michael Kohler and Sophie Langer. On the rate of convergence of fully connected very deep neural network regression estimates. *arXiv preprint arXiv:1908.11133*, 2019.
- Aleksandr Petrovich Korostelev and Alexandre B Tsybakov. *Minimax theory of image reconstruction*, volume 82. Springer Science & Business Media, 2012.
- Alex Krizhevsky, Ilya Sutskever, and Geoffrey E Hinton. Imagenet classification with deep convolutional neural networks. In *Advances in neural information processing systems*, pages 1097–1105, 2012.
- Shiyu Liang, Ruoyu Sun, Yixuan Li, and Rayadurgam Srikant. Understanding the loss surface of neural networks for binary classification. *arXiv preprint arXiv:1803.00909*, 2018.

- Yi Lin. Support vector machines and the bayes rule in classification. *Data Mining and Knowledge Discovery*, 6(3):259–275, 2002.
- Ruiqi Liu, Ben Boukai, and Zuofeng Shang. Optimal nonparametric inference via deep neural network. *arXiv preprint arXiv:1902.01687*, 2019.
- Zhou Lu, Hongming Pu, Feicheng Wang, Zhiqiang Hu, and Liwei Wang. The expressive power of neural networks: A view from the width. In *Advances in neural information processing systems*, pages 6231–6239, 2017.
- Kaifeng Lyu and Jian Li. Gradient descent maximizes the margin of homogeneous neural networks. *arXiv preprint arXiv:1906.05890*, 2019.
- C. W. H. Mace and A. C. C. Coolen. Statistical mechanical analysis of the dynamics of learning in perceptrons. *Statistics and Computing*, 8(1):55–88, 1998. . URL <https://doi.org/10.1023/A:1008896910704>.
- Enno Mammen, Alexandre B Tsybakov, et al. Smooth discrimination analysis. *The Annals of Statistics*, 27(6):1808–1829, 1999.
- Guido F Montufar, Razvan Pascanu, Kyunghyun Cho, and Yoshua Bengio. On the number of linear regions of deep neural networks. In *Advances in neural information processing systems*, pages 2924–2932, 2014.
- Ryumei Nakada and Masaaki Imaizumi. Adaptive approximation and estimation of deep neural network to intrinsic dimensionality. *arXiv preprint arXiv:1907.02177*, 2019.
- Kien Nguyen, Clinton Fookes, Arun Ross, and Sridha Sridharan. Iris recognition with off-the-shelf cnn features: A deep learning perspective. *IEEE Access*, 6:18848–18855, 2017.
- Kenta Oono and Taiji Suzuki. Approximation and non-parametric estimation of resnet-type convolutional neural networks. *arXiv preprint arXiv:1903.10047*, 2019.
- Maithra Raghu, Ben Poole, Jon Kleinberg, Surya Ganguli, and Jascha Sohl Dickstein. On the expressive power of deep neural networks. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pages 2847–2854. JMLR. org, 2017.
- Olga Russakovsky, Jia Deng, Hao Su, Jonathan Krause, Sanjeev Satheesh, Sean Ma, Zhiheng Huang, Andrej Karpathy, Aditya Khosla, Michael Bernstein, Alexander C. Berg, and Li Fei-Fei. ImageNet Large Scale Visual Recognition Challenge. *International Journal of Computer Vision (IJCV)*, 115(3):211–252, 2015. .
- David Saad and Sara A. Solla. Dynamics of on-line gradient descent learning for multilayer neural networks, 1996.
- Johannes Schmidt-Hieber. Nonparametric regression using deep neural networks with relu activation function. *The Annals of Statistics*, 2019. Forthcoming.
- Thiago Serra, Christian Tjandraatmadja, and Srikumar Ramalingam. Bounding and counting linear regions of deep neural networks. *arXiv preprint arXiv:1711.02114*, 2017.
- Karen Simonyan and Andrew Zisserman. Very deep convolutional networks for large-scale image recognition. *arXiv preprint arXiv:1409.1556*, 2014.
- Daniel Soudry, Elad Hoffer, Mor Shpigel Nacson, Suriya Gunasekar, and Nathan Srebro. The implicit bias of gradient descent on separable data. *The Journal of Machine Learning Research*, 19(1):2822–2878, 2018.
- Nathan Srebro, Karthik Sridharan, and Ambuj Tewari. Optimistic rates for learning with a smooth loss. *arXiv preprint arXiv:1009.3896*, 2010.
- Ingo Steinwart, Clint Scovel, et al. Fast rates for support vector machines using gaussian kernels. *The Annals of Statistics*, 35(2):575–607, 2007.
- Taiji Suzuki. Adaptivity of deep relu network for learning in besov and mixed smooth besov spaces: optimal rate and curse of dimensionality. *arXiv preprint arXiv:1810.08033*, 2018.
- Matus Telgarsky. Representation benefits of deep feedforward networks. *arXiv preprint*

- arXiv:1509.08101*, 2015.
- Yuandong Tian. A theoretical framework for deep locally connected relu network. *arXiv preprint arXiv:1809.10829*, 2018.
- Yuandong Tian. Over-parameterization as a catalyst for better generalization of deep relu network. *arXiv preprint arXiv:1909.13458*, 2019.
- Alexander B Tsybakov et al. Optimal aggregation of classifiers in statistical learning. *The Annals of Statistics*, 32(1):135–166, 2004.
- Alexandre B Tsybakov, Sara A van de Geer, et al. Square root penalty: adaptation to the margin in classification and in edge estimation. *The Annals of Statistics*, 33(3): 1203–1224, 2005.
- Sara Van De Geer. *Empirical Processes in M-estimation*. Cambridge University Press, 2000.
- Dmitry Yarotsky and Anton Zhevnerchuk. The phase diagram of approximation rates for deep neural networks. *arXiv preprint arXiv:1906.09477*, 2019.
- Matthew D Zeiler and Rob Fergus. Visualizing and understanding convolutional networks. In *European conference on computer vision*, pages 818–833. Springer, 2014.
- Xiao Zhang, Yaodong Yu, Lingxiao Wang, and Quanquan Gu. Learning one-hidden-layer relu networks via gradient descent. *Proceedings of the 22nd International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2019.
- Peng Zhao and Bin Yu. On model selection consistency of lasso. *Journal of Machine learning research*, 7(Nov):2541–2563, 2006.

6. Appendix.

6.1. *Smooth Boundary Condition.* In this section we review some existing work under smooth boundary condition in details and point out its connection to our proposed teacher-student neural network.

Smooth Functions. A function has Hölder smoothness index β if all partial derivatives up to order $\lfloor \beta \rfloor$ exist and are bounded, and the partial derivatives of order $\lfloor \beta \rfloor$ are $\beta - \lfloor \beta \rfloor$ Lipschitz. The ball of β -Hölder functions with radius R is then defined as

$$\mathcal{H}_r^\beta(R) = \left\{ f : \mathbb{R}^r \rightarrow \mathbb{R} : \sum_{\alpha: |\alpha| < \beta} \|\partial^\alpha f\|_\infty + \sum_{\alpha: |\alpha| = \lfloor \beta \rfloor} \sup_{\substack{\mathbf{x}, \mathbf{y} \in D \\ \mathbf{x} \neq \mathbf{y}}} \frac{|\partial^\alpha f(\mathbf{x}) - \partial^\alpha f(\mathbf{y})|}{|\mathbf{x} - \mathbf{y}|_\infty^{\beta - \lfloor \beta \rfloor}} \leq R \right\},$$

where $\partial^\alpha = \partial^{\alpha_1} \dots \partial^{\alpha_r}$ with $\alpha = (\alpha_1, \dots, \alpha_r) \in \mathbb{N}^r$ and $|\alpha| := |\alpha|_1$.

Boundary Assumption. It is known that estimating the classifier directly instead of the conditional class probability helps achieve fast convergence rates [Mammen et al., 1999, Tsybakov et al., 2004, 2005, Audibert et al., 2007]. Classification in this case can be thought of as nonparametric estimation of sets where we directly estimate the decision regions for different labels, e.g., G for label 1. Then, the classifier is determined by attributing \mathbf{x} to label 1 if $\mathbf{x} \in G$ and to label -1 otherwise, i.e.,

$$C(\mathbf{x}) = 2 \cdot \mathbb{I}_G(\mathbf{x}) - 1.$$

In this case, the Bayes risk can be written as

$$R(G) = 1/2 \left(\int_{G^c} p(\mathbf{x}) \mathbb{Q}(d\mathbf{x}) + \int_G q(\mathbf{x}) \mathbb{Q}(d\mathbf{x}) \right).$$

Denote $G^* = \{\mathbf{x} : p(\mathbf{x}) \geq q(\mathbf{x})\}$ to be the Bayes risk minimizer, and the classification problem is equivalent to estimation of the optimal set G^* . Given p, q , let

$$\mathcal{X}_+ = \{\mathbf{x} \in \mathcal{X} : p(\mathbf{x}) \geq q(\mathbf{x})\}, \quad \mathcal{X}_- = \{\mathbf{x} \in \mathcal{X} : p(\mathbf{x}) < q(\mathbf{x})\}$$

The optimal decision rule is to assign label 1 to $\mathbf{x} \in \mathcal{X}_+$ and -1 to $\mathbf{x} \in \mathcal{X}_-$. The decision boundary in this case is $\{\mathbf{x} \in \mathcal{X} : p(\mathbf{x}) = q(\mathbf{x})\}$. To characterize the smoothness of the boundary, it is usually assumed that \mathcal{X}_+ consists of union and intersection of smooth hyper-surfaces [Kim et al., 2018, Tsybakov et al., 2004]. Specifically, the following assumption is widely

adopted [Mammen et al., 1999, Tsybakov et al., 2004, Kim et al., 2018, Imaizumi and Fukumizu, 2018] and referred to as "boundary fragment". Let \mathcal{H} be some smooth function space from $\mathbb{R}^{d-1} \rightarrow \mathbb{R}$. Define sets $\mathcal{G}^{\mathcal{H}}$ as

$$(6.1) \quad \mathcal{G}^{\mathcal{H}} = \{\mathbf{x} \in \mathcal{X} : x_j > h(\mathbf{x}_{-j}), h \in \mathcal{H}, j \in \{1, 2, \dots, d\}\}$$

where $\mathbf{x}_{-j} = (x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_d)$. It is assumed that \mathcal{X}_+ is composed of finite union and intersection of sets in $\mathcal{G}^{\mathcal{H}}$. The seemingly odd form (6.1) enforces special structures on the indicator function and reduces the complexity of the corresponding sets.

A more general assumption on the decision boundary is that the set, denoted as \mathcal{G} , containing all possible \mathcal{X}_+ cannot be too large. This is measured by the bracketing entropy H_B of the metric space $(\mathcal{G}, d_{\Delta})$. The more general assumption of the decision boundary is stated as

(B) There exists $A > 0$ and $\rho \in [0, \infty]$ such that

$$H_B(\delta, \mathcal{G}, d_{\Delta}) \leq A\delta^{-\rho}.$$

Our proposed teacher-student network setting follows this more general assumption and makes more sense in high-dimensional classification.

Remark 3. If \mathcal{H} are β -smooth functions in (6.1), then (B) holds with $\rho = (d-1)/\beta$. On the other hand, Lemma 3.9 gives $H_B \lesssim A_n \vee (B_n \log(1/\delta))$ where in terms of δ , the order is $\log(1/\delta)$, which corresponds to $\rho \rightarrow 0$ and $\beta \rightarrow \infty$. However, this doesn't necessarily imply the boundary fragment set (6.1) is larger, since A_n, B_n depend on n and are not absolute constants as in (B).

6.2. *Comments on Lemma 3.9.* Lemma 3.9 is the main result for controlling the bracketing entropy of the estimation sets. Below we point out some key property of this result and compare to other entropy bounds of neural networks.

Exponential Dependence on Depth. The bracketing entropy of $\mathcal{G}^{\mathcal{F}}$ developed in Lemma 3.9 is much larger than that of \mathcal{F} itself with respect to $\|\cdot\|_{\infty}$, as described in Lemma 4.4. The main difference is the dependence on the number of layers L : the dependence is linear in Lemma 4.4 while exponential in Lemma 3.9. Thus, even though $\mathcal{G}^{\mathcal{F}}$ is a slice of the subgraph of \mathcal{F} , $\mathcal{G}^{\mathcal{F}}$ is much more complicated than \mathcal{F} in term of entropy. We argue that this gap cannot be closed even in the special case $d = 1$.

Montufar et al. [2014] establish a lower bound on the maximum number of linear pieces for a ReLU neural network (Lemma 3.8). Consider a 1-dimensional ReLU DNN function with L layers and 2 nodes on each layer.




FIG 4. Example of a ReLU function in 1D. The induced set where $f > 0$ is colored red and it's a union of two intervals $(a_1, b_1), (a_2, b_2)$. All pieces cross 0 so there are all active.

Corollary 5 of Montufar et al. [2014] show that there exists some f with $s = \Omega(2^{L-1})$ pieces on $[0, 1]$. With scaling and shifting, assume that on each piece the linear function crosses 0. Then, \mathcal{G}^f will be at least $\lfloor s/2 \rfloor = \Omega(2^{L-2})$ intervals. Denote these disjoint intervals to be $\{(a_i, b_i)\}_{i=1}^{\lfloor s/2 \rfloor}$. Since they are disjoint, to construct a δ -bracket of all the intervals, we need to δ -cover all the a_i 's and b_i 's. Similar to the grid argument from the proof of Lemma 3.6, we need at least

$$\binom{1/\delta}{s} = \Omega((1/\delta - s)^s)$$

different combinations of the s grid points. Hence the bracketing entropy must be in the order of

$$\log((1/\delta - s)^s) = 2^{L-2} \log(1/\delta).$$

The exponential dependence of depth L in the entropy stems from the fact that the number of linear regions of ReLU DNNs scales exponentially with L .

Independent of Weights Magnitude. We also want to point out that the entropy of $\mathcal{G}^{\mathcal{F}}$ is not concerned with the magnitude of the neural network weights, in contrast to the bound in Lemma 4.4. This is because any scaling of the function doesn't change how it intercepts with zero. Hence, unlike \mathcal{F} , the entropy of $\mathcal{G}^{\mathcal{F}}$ doesn't depend on the weight maximum B .

The Use of ReLU Activation. The reason why we can even bound the entropy of $\mathcal{G}^{\mathcal{F}}$ critically relies on the fact that we are considering the ReLU activation function. If we consider smooth nonlinear activation functions, e.g. hyperbolic tangent, sigmoid, instead of the order $\log(1/\delta)$, we can only get the entropy of a much larger order

$$H_B(\delta, \mathcal{G}^{\mathcal{F}}, d_{\Delta}) \leq A\delta^{-\alpha}$$

for some constant $A > 0$ and $\alpha > 0$. To see this, consider the case $d = 2$. Instead of polygons, which can be controlled by the vertices, the regions have smooth boundary and will require $O(1/\delta)$ many grid points to cover. Thus the covering number is of order

$$\binom{1/\delta^2}{1/\delta} = O\left(\left(\frac{1}{\delta}\right)^{2/\delta}\right).$$

Thus, the entropy is in a polynomial order of $1/\delta$.

6.3. Illustration and Proof of Lemma 3.1. In this section, Assumption (A3) will be examined in the setting that the teacher network f_n^* has random weights. We will argue that with probability at least $1 - \delta$, f_n^* will satisfy assumption (A3) with $T_n = A(\delta)/(\log n)^{m^*d^2L_n^*}$ and $c_n = B(\delta)(\log n)^{m^*d^2L_n^*}$, where $A(\delta), B(\delta)$ are constants depending only on δ and the distribution of the random weights, e.g. normal, truncated normal, etc. Hence, the results which assume Assumption (A3) will hold with high probability.

A Toy Case. To illustrate the intuition, consider the case where $d = 1$ and f_n^* is the following one hidden-layer ReLU neural network

$$(6.2) \quad f_n^*(x) = \sum_{j=1}^{N_n^*} w_{2j} \sigma(w_{1j}x + b_j) + b, \quad x \in [0, 1],$$

with $L_n^* = 1$, $N_n^* = O(\log n)$ and w_{1j}, w_{2j}, b_j, b are i.i.d. standard Gaussian. Since all the weights are almost surely nonzero, we omit the zero weight cases for the analysis. Let $p_i = (u_i, v_i)$, $i = 1, 2, \dots, s$, denote the active pieces of (6.2). By Lemma 3.7, we know that $s = O(\log n)$. For each p_i , define the following quantities:

1. $k_i =$ the slope of $f_n^*(x)$ on $x \in p_i$;
2. $t_i = \max_{x \in p_i} f_n^*(x) \wedge \max_{x \in p_i} -f_n^*(x)$.

See Figure 5 for an illustration. Then, assumption (A3) is satisfied if

$$(6.3) \quad \min_i \{|k_i|\} = \Omega(1/\log n) \quad \text{and} \quad \min_i \{t_i\} = \Omega(1/\log n).$$

Next we will rigorously examine (6.3).

From (6.2), each k_i can be expressed as $w_{1j}w_{2j}$ for some $j \in \{1, 2, \dots, N_n^*\}$. Therefore, $\min_{1 \leq i \leq N_n^*} \{|k_i|\} = \min_{1 \leq j \leq N_n^*} \{|w_{1j}w_{2j}|\}$. Since w_{1j}, w_{2j} are i.i.d.




FIG 5. Example of a ReLU function in $[0, 1]$. There are two active pieces p_1, p_2 . On each active piece, t_i, k_i are illustrated in color red.

standard Gaussian, we have

$$\begin{aligned} \mathbb{P}(\min_{1 \leq i \leq N_n^*} \{|k_i|\} < k) &= \mathbb{P}(\min_{1 \leq j \leq N_n^*} \{|w_{1j}w_{2j}|\} < k) \\ &\leq \sum_{j=1}^{N_n^*} \mathbb{P}(|w_{1j}w_{2j}| < k) \leq 2N_n^* \mathbb{P}(|w_{11}| < \sqrt{k}) \leq 2N_n^* \sqrt{k}. \end{aligned}$$

By choosing $k = \left(\frac{\delta}{2N_n^*}\right)^2$, we have $\min_{1 \leq i \leq N_n^*} \{|k_i|\} = \Omega(1/\log n)$ with probability at least $1 - \delta$.

On the other hand, for any $i = 1, \dots, s$, $t_i = |f_n^*(x_{h_i})|$ for some $h_i \in \{1, \dots, N_n^*\}$, where $x_{h_i} = -b_{h_i}/w_{1h_i}$. Hence

$$\min_{1 \leq i \leq s} \{t_i\} \geq \min_{1 \leq j \leq N_n^*} \{|f_n^*(x_j)|\}.$$

Let $W_1 = \{w_{1j}, b_j\}_{j=1}^{N_n^*}$. Then, $f_n^*(x_i) | W_1 \sim N(0, \sigma_{x_i}^2)$, where $\sigma_{x_i}^2$ has an expression of $\sum_{j=1}^{N_n^*} \sigma(w_{1j}x_i + b_j)^2 + 1$. Hence, for any $t > 0$,

$$\begin{aligned} \mathbb{P}(\min_{i \leq N_n^*} \{|f_n^*(x_i)|\} < t | W_1) &\leq \sum_{i=1}^{N_n^*} \mathbb{P}(|f_n^*(x_i)| < t | W_1) \\ &= N_n^* \mathbb{P}(|f_n^*(x_i)| < t | W_1) \leq N_n^* \left(\frac{t}{\sigma_{x_i}}\right). \end{aligned}$$

Since $\sigma_{x_i} \geq 1$, by taking $t = \delta/N_n^*$, we have that with probability at least $1 - \delta$, $\min_i \{t_i\} \geq t$ and $t = \Omega(1/\log n)$. Therefore, (6.3) holds with high probability, so that assumption (A3) holds by setting $1/c_n = \min_i \{|k_i|\}$ and $T_n = \min_i \{t_i\}$, which are both in the order of $\Omega(1/\log n)$.

General Case. Now we consider the general case $d > 1$ and $L_n^* > 1$. The teacher network has an expression

$$f_n^*(\mathbf{x}) = \mathbf{W}^{(L_n^*+1)} \sigma_{(\mathbf{W}^{(L_n^*)}, \mathbf{b}^{(L_n^*)})} \circ \cdots \circ \sigma_{(\mathbf{W}^{(1)}, \mathbf{b}^{(1)})}(\mathbf{x}) + \mathbf{b}^{(L_n^*+1)}, \mathbf{x} \in [0, 1]^d.$$

Let $N_n^* = O(\log n)^{m^*}$. By Lemma 3.7, f_n^* has linear pieces p_1, \dots, p_s for $s = O(\log n)^{m^* L_n^* d}$. Let $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{v_s}\}$ be the collection of vertices of $\{p_1, \dots, p_s\}$. We call such $\mathbf{x}_i \in \mathbb{R}^d$ a *piece vertex* and it's not the same as the vertex of $\{\mathbf{x} \in \mathcal{X} : f_n(\mathbf{x}) \geq 0\}$, which is closely examined in the proof of Lemma 3.9. The following lemma states that $v_s = O(\log n)^{m^* L_n^* d^2}$ in our setting.

Lemma 6.1. Let f be a ReLU neural network with d -dimensional input, L hidden layers and width N for every layer. Then, $v_s = O(N)^{Ld^2}$.

PROOF. Recall that $\mathbf{w}_i^{(l)}$ and $b_i^{(l)}$ for $i = 1, \dots, N$, $1 \leq l \leq L$ are the weight vectors and biases on the l -th hidden layer. For $i = 1, \dots, N$, define

$$f_i^{(l-1)}(\mathbf{x}) = \mathbf{w}_i^{(l)} \sigma_{(\mathbf{W}^{(l-1)}, \mathbf{b}^{(l-1)})} \circ \cdots \circ \sigma_{(\mathbf{W}^{(1)}, \mathbf{b}^{(1)})}(\mathbf{x}) + b_i^{(l)},$$

which maps $\mathbb{R}^d \rightarrow \mathbb{R}$. We can rewrite f as

$$(6.4) \quad f(\mathbf{x}) = \sum_{i=1}^N w_i^{(L+1)} \sigma(f_i^{(L-1)}(\mathbf{x})) + b^{(L+1)},$$

In other words, $f_i^{(L-1)}(\mathbf{x})$ represents the inputs to the i -th ReLU unit in the last hidden layer of f and itself is an $(L-1)$ -hidden-layer ReLU neural network.

The key idea of the proof is by induction. Notice that the piece vertices of f can only come from the following two ways: Type I: The piece vertices of $f_1^{(L-1)}, f_2^{(L-1)}, \dots, f_N^{(L-1)}$, in whose local neighbourhoods, the ReLU units in the last layer doesn't change sign; Type II: By activations of the ReLU unit in the last layer. i.e. $f_i^{(L-1)}(\mathbf{x}) = 0$ for some $i = 1, \dots, N$. Let $V(l)$ be the maximum number of piece vertices of an l -hidden-layer ReLU neural network with width N and let $U(l)$ be the maximum number of Type II piece vertices created at layer l . Then for $1 < l \leq L$ we have

$$(6.5) \quad V(l) \leq NV(l-1) + U(l).$$

For $U(l)$, the key is to connect the Type II piece vertices of f to the vertices of $\{\mathbf{x} \in \mathcal{X} : f_i^{(L-1)}(\mathbf{x}) \geq 0\}$, which has been extensively studied in

Lemma 3.9. To this end, we define another quantity. On the i -th ReLU unit in the l -th hidden layer, let $R_i^{(l)} := \{\mathbf{x} \in \mathcal{X} : f_i^{(l)}(\mathbf{x}) = 0\}$, which consists of $(d-1)$ -dimensional hyperplane segments. To be specific, denote all the active pieces of $f_i^{(l)}(\mathbf{x})$ to be $\{p_{ij}^{(l)} : j = 1, \dots, s_i^{(l)}\}$, where $s_i^{(l)} = O(N)^{(l-1)d}$ according to Lemma 3.7 for any $1 \leq i \leq N$. On each active piece $p_{ij}^{(l)}$, denote

$$h_{ij}^{(l)} = \{(\mathbf{x}, f_i^{(l)}(\mathbf{x})) : \mathbf{x} \in p_{ij}^{(l)}\} \cap \{(\mathbf{x}, 0) : \mathbf{x} \in p_{ij}^{(l)}\},$$

which is part of a $(d-1)$ -dimensional hyperplane. Then we have $R_i^{(l)} = \{h_{ij}^{(l)} : j = 1, \dots, s_i^{(l)}\}$, a collection of $(d-1)$ -dimensional hyperplane segments. Let $R^{(l)} = \cup_{i=1}^N R_i^{(l)}$, which corresponds to the piece boundaries of f^{l+1} .

By definition, all Type II pieces vertices must reside in at least one of the the activation sets ($z = 0$ in $\sigma(z)$) of the ReLU units in the last layer. $R^{(L)}$ contains all such activation sets for the last hidden layer, i.e. for any $h \in R$, there exists $1 \leq i \leq N$ such that $f_i(\mathbf{x}) = 0, \forall \mathbf{x} \in h$. The Type II pieces vertices are jointly determined by such activation sets and the piece boundary of f_i 's (dimension $d-1$), i.e. $R_i^{(L-2)}$. Therefore, the total number of such piece vertices can be bounded by

$$U(l) \leq \binom{|R^{(l-1)}| + |R^{(l-2)}|}{d} = O(N)^{(l-1)d^2+d},$$

where $|R^{(l)}|$ denotes the number of elements in $R^{(l)}$, which is bounded by $O(N)^{(l-1)d+1}$.

For $V(L)$, we first conclude that $V(1) = O(N^d)$. For a 1-hidden layer ReLU network, the decision boundary of every ReLU unit is a $(d-1)$ -dimension hyperplane ($\mathbf{w}_1 \mathbf{x} + b_1 = 0$). The maximum number of piece vertices is bounded by $\binom{N}{d} = O(N^d)$. Then, (6.5) can be repeatedly broken down to

$$\begin{aligned} V(L) &\leq NV(L-1) + U(L) \\ &\leq N^2V(L-2) + NU(L-1) + U(L) \leq \dots \\ &\leq N^{L-1}V(1) + \sum_{l=0}^{L-1} N^l U(L-l) \\ &= O\left(N^{L-1+d}\right) + O\left(\sum_{l=0}^{L-1} N^{(L-l-1)d^2+d+l}\right) \\ &= O\left(N^{(L-1)d^2+d}\right) = O\left(N^{Ld^2}\right) \end{aligned}$$

□

As an extension to the toy case, for any $1 \leq i \leq N_n^*$, define

1. $k_i = \min_{j=1, \dots, d} \left\{ \left| \frac{\partial f_n^*(\mathbf{x})}{\partial x_j} \right| : \mathbf{x} \in p_i \right\}$;
2. $t_0 = \min_{1 \leq i \leq v_s} \{|f_n^*(\mathbf{x}_i)|\}$.

That is, k_i is the minimal absolute values of the directional derivatives of f_n^* on piece p_i . Assumption (A3) is satisfied if the following holds:

$$(6.6) \quad \min_{1 \leq i \leq s} \{k_i\}, t_0 = \Omega(\log n)^{m^* d^2 L_n^*}.$$

We will check (6.6). Since the partial derivative of $f_n^*(\mathbf{x})$ for $\mathbf{x} \in p_i$ can be expressed as the product of the random weights, we have

$$\min_{1 \leq i \leq s} \{k_i\} \geq \min_{1 \leq j_l \leq N_n^*} \left| \prod_{l=1}^{L_n^*+1} w_{i_l j_l}^{(l)} \right|,$$

where $w_{i_l j_l}^{(l)}$ is an element from $\mathbf{W}^{(l)}$. Since

$$\min_{1 \leq j_l \leq N_n^*} \prod_{l=1}^{L_n^*+1} |w_{i_l j_l}^{(l)}| \geq \min_{l \leq L_n^*+1; i_l, j_l \leq N_n^*} |w_{i_l j_l}^{(l)}|^{L_n^*+1},$$

we get that

$$\begin{aligned} \mathbb{P}(\min_{1 \leq i \leq s} \{k_i\} < k) &\leq \mathbb{P}\left(\min_{l \leq L_n^*+1; j_l \leq N_n^*} |w_{l j_l}| \leq k^{1/(L_n^*+1)}\right) \\ &\leq (L_n^* + 1)(N_n^*)^2 \mathbb{P}(|w_{l j_l}| < k^{1/(L_n^*+1)}) \\ &\leq (L_n^* + 1)(N_n^*)^2 k^{1/(L_n^*+1)}. \end{aligned}$$

By taking

$$k_0 = \left(\frac{\delta}{(N_n^*)^2 (L_n^* + 1)} \right)^{L_n^*+1},$$

we have that with probability at least $1 - \delta$, $\min_{1 \leq i \leq s} \{k_i\} \geq k_0$ and $k_0 = \Omega(1/\log n)^{2m^*(L_n^*+1)}$.

On the other hand, for any t_i , there exist $j = 1, \dots, v_s$ such that $t_i = f_n^*(\mathbf{x}_j)$. Hence

$$\min_{i=1, \dots, v_s} \{t_i\} \geq \min_{j=1, \dots, v_s} \{|f_n^*(\mathbf{x}_j)|\}.$$

Let $\mathbf{W}_{-L_n^*} := \{\mathbf{W}^{(l)}, \mathbf{b}^{(l)}\}_{l=1}^{L_n^*}$. Then we have $f_n^*(\mathbf{x}_j) | \mathbf{W}_{-L_n^*} \sim N(0, \sigma_{\mathbf{x}_j}^2)$, where $\sigma_{\mathbf{x}_j}^2$ depends on $\mathbf{W}_{-L_n^*}$ and $\sigma_{\mathbf{x}_j}^2 \geq 1$ that

$$\sigma_{\mathbf{x}_j}^2 | \mathbf{W}_{-L_n^*} := \sum_{i=1}^{N_{L_n^*}} \sigma_i^2(\mathbf{x}_j) + 1,$$

which is reminiscent of (6.4) and $N_{L_n^*}$ is the width of the last layer and $\sigma_j(\cdot)$'s are outputs (post-activations) from the last layer given $\mathbf{W}_{-L_n^*}$. Therefore, for any $t > 0$, we have

$$\begin{aligned} \mathbb{P}(\min_{1 \leq j \leq v_s} \{|f_n^*(\mathbf{x}_j)|\} < t \mid \mathbf{W}_{-L_n^*}) &\leq \sum_{j=1}^{v_s} \mathbb{P}(|f_n^*(\mathbf{x}_j)| < t \mid \mathbf{W}_{-L_n^*}) \\ &= v_s \mathbb{P}(|f_n^*(\mathbf{x}_1)| < t \mid \mathbf{W}_{-L_n^*}) \\ &\leq v_s \left(\frac{t}{\sigma_{x_i}} \right) \leq t(N_n^*)^{d^2 L_n^*}. \end{aligned}$$

Thus by taking $t = \delta/(N_n^*)^{d^2 L_n^*}$, we have that with probability at least $1 - \delta$, $\min_i \{t_i\} \geq t$ and $t = \Omega(1/\log n)^{m^* d^2 L_n^*}$. Therefore, (6.6) holds. That is to say, when $d \geq 2$, with high probability, Assumption (A3) holds in which $c_n, 1/T_n = O(\log n)^{m^* d^2 L_n^*}$.

Notice that the probability arguments used in this section don't rely on Gaussian distribution. As long as all weights are i.i.d. with distribution that doesn't have a point mass at 0, our claim holds.