

FOUR NEW FORMS OF THE TAYLOR–ITO AND TAYLOR–STRATONOVICH EXPANSIONS AND ITS APPLICATION TO THE HIGH-ORDER STRONG NUMERICAL METHODS FOR ITO STOCHASTIC DIFFERENTIAL EQUATIONS

DMITRIY F. KUZNETSOV

ABSTRACT. The problem of the Taylor–Ito and Taylor–Stratonovich expansions of the Ito stochastic processes in a neighborhood of a fixed moment of time is considered. The classical forms of the Taylor–Ito and Taylor–Stratonovich expansions are transformed to the four new representations, which includes the minimal sets of different types of iterated Ito and Stratonovich stochastic integrals. Therefore, these representations (the so-called unified Taylor–Ito and Taylor–Stratonovich expansions) are more convenient for constructing of high-order strong numerical methods for Ito stochastic differential equations. Explicit one-step strong numerical schemes with the orders of convergence 1.0, 1.5, 2.0, 2.5, and 3.0 based on the unified Taylor–Ito and Taylor–Stratonovich expansions are derived. Effective mean-square approximations of iterated Ito and Stratonovich stochastic integrals from these numerical schemes are constructed on the base of the multiple Fourier–Legendre series with multiplicities 1 to 6.

CONTENTS

1.	Introduction	2
2.	Integration Order Replacement Technique for Iterated Ito Stochastic Integrals	6
3.	The Taylor–Ito Expansion	13
4.	The First Form of the Unified Taylor–Ito Expansion	15
5.	The Second Form of the Unified Taylor–Ito Expansion	18
6.	The Taylor–Stratonovich Expansion	20

MATHEMATICS SUBJECT CLASSIFICATION: 60H05, 60H10, 42B05.

KEYWORDS: TAYLOR–ITO EXPANSION, TAYLOR–STRATONOVICH EXPANSION, UNIFIED TAYLOR–ITO EXPANSION, UNIFIED TAYLOR–STRATONOVICH EXPANSION, ITO STOCHASTIC DIFFERENTIAL EQUATION, HIGH-ORDER STRONG NUMERICAL METHOD FOR ITO SDEs, ITERATED ITO STOCHASTIC INTEGRAL, ITERATED STRATONOVICH STOCHASTIC INTEGRAL, GENERALIZED MULTIPLE FOURIER SERIES, MULTIPLE FOURIER–LEGENDRE SERIES, MEAN-SQUARE APPROXIMATION, EXPANSION.

7.	The First Form of the Unified Taylor–Stratonovich Expansion	23
8.	The Second Form of the Unified Taylor–Stratonovich Expansion	26
9.	Comparison of the Unified Taylor–Ito and Taylor–Stratonovich Expansions With the Classical Taylor–Ito and Taylor–Stratonovich Expansions	27
10.	Application of First Form of the Unified Taylor–Ito Expansion to the High-Order Strong Numerical Methods for Ito SDEs	31
11.	Application of First Form of the Unified Taylor–Stratonovich Expansion to the High-Order Strong Numerical Methods for Ito SDEs	36
12.	Method of the Mean-Square Approximation of Iterated Ito and Stratonovich Stochastic Integrals Based on Generalized Multiple Fourier Series	41
13.	Theorems 6–8, 10–13 from Point of View of the Wong–Zakai Approximation	82
	References	86

1. INTRODUCTION

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space, let $\{\mathcal{F}_t, t \in [0, T]\}$ be a nondecreasing right-continuous family of σ -subfields of \mathcal{F} , and let \mathbf{f}_t be a standard m -dimensional Wiener stochastic process, which is \mathcal{F}_t -measurable for any $t \in [0, T]$. We assume that the components $\mathbf{f}_t^{(i)}$ ($i = 1, \dots, m$) of this process are independent. Consider an Ito stochastic differential equation (SDE) in the integral form

$$(1) \quad \mathbf{x}_t = \mathbf{x}_0 + \int_0^t \mathbf{a}(\mathbf{x}_\tau, \tau) d\tau + \int_0^t B(\mathbf{x}_\tau, \tau) d\mathbf{f}_\tau, \quad \mathbf{x}_0 = \mathbf{x}(0, \omega).$$

Here \mathbf{x}_t is some n -dimensional stochastic process satisfying to Ito SDE (1). The nonrandom functions $\mathbf{a} : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^n$, $B : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^{n \times m}$ guarantee the existence and uniqueness up to stochastic equivalence of a solution of (1) [1]. The second integral on the right-hand side of (1) is interpreted as an Ito stochastic integral. Let \mathbf{x}_0 be an n -dimensional random variable, which is \mathcal{F}_0 -measurable and $\mathbb{M}\{|\mathbf{x}_0|^2\} < \infty$ (\mathbb{M} denotes a mathematical expectation). Also we assume that \mathbf{x}_0 and $\mathbf{f}_t - \mathbf{f}_0$ are independent when $t > 0$.

It is well known [2]–[5] that Ito SDEs are adequate mathematical models of dynamic systems of different physical origin that are affected by random perturbations. For example, Ito SDEs are used as mathematical models in stochastic mathematical finance, hydrology, seismology, geophysics,

chemical kinetics, population dynamics, electrodynamics, medicine and other fields [2]-[5]. Numerical integration of Ito SDEs based on the strong convergence criterion of approximations [2] is widely used for the numerical simulation of sample trajectories of solutions to Ito SDEs (which is required for constructing new mathematical models on the basis of such equations and for the numerical solution of different mathematical problems connected with Ito SDEs). Among these problems, we note the following: filtering of signals under influence of random noises in various statements (linear Kalman–Bucy filtering, nonlinear optimal filtering, filtering of continuous time Markov chains with a finite space of states, etc.), optimal stochastic control (including incomplete data control), testing estimation procedures of parameters of stochastic systems, stochastic stability and bifurcations analysis [2], [3].

Exact solutions of Ito SDEs are known in rather rare cases. It is for this reason that it becomes necessary to construct numerical procedures for solving these equations.

In this paper, a promising approach [2]-[5] to the numerical integration of Ito SDEs based on the stochastic analogues of the Taylor formula (Taylor–Ito and Taylor–Stratonovich expansions) [6]-[9] is used. This approach uses a finite discretization of the time variable and the numerical simulation of the solution to the Ito SDE at discrete instants of time using the stochastic analogues of the Taylor formula mentioned above. A number of works (e.g., [2]-[5]) describe numerical schemes with the strong orders of convergence 1.5, 2.0, 2.5, and 3.0 for the Ito SDEs; however, they do not contain efficient procedures of the mean-square approximation of iterated stochastic integrals involved in these schemes for the case of non-commutative noise.

In this paper we consider the unified Taylor–Ito and Taylor–Stratonovich expansions [8], [9] which makes it possible (in contrast with its classical analogues [6], [7]) to use the minimal sets of iterated Ito and Stratonovich stochastic integrals; this is a simplifying factor for the numerical methods implementation. We prove the unified Taylor–Ito expansion [8] with using the slightly different approach (which is taken from [9]) in comparison with the approach from [8]. Moreover, we obtain another (second) version of the unified Taylor–Ito expansion [10], [11]. In addition, we construct two new forms of the Taylor–Stratonovich expansion (the so-called unified Taylor–Stratonovich expansions [9]). Furthermore, in this paper we study methods [12]-[63] of numerical simulation for iterated Ito and Stratonovich stochastic integrals of multiplicities 1, 2, 3, 4, 5, 6, ... used in the strong numerical methods for Ito SDEs [2]-[5], [46]-[50], [56]-[61]. To approximate the iterated Ito and Stratonovich stochastic integrals appearing in the numerical schemes with the strong orders of convergence 1.0, 1.5, 2.0, 2.5, 3.0 etc., the method of generalized multiple Fourier series and especially method of multiple Fourier–Legendre series are studied in [12]-[63]. It is important to note that the method of generalized multiple Fourier series [12]-[63] does not lead to the partitioning of the integration interval of the iterated Ito and Stratonovich stochastic integrals under consideration; this interval is the integration step of the numerical methods used to solve Ito SDEs; therefore, it is already fairly small and does not need to be partitioned. Computational experiments [46] show that the application of numerical simulation for iterated stochastic integrals (in which the interval of integration is partitioned) leads to unacceptably high computational cost and accumulation of computation errors. Also note that the Legendre polynomials have essential advantage over the trigonometric functions (see [23], [39]) in the framework of the method of generalized multiple Fourier series [12]-[63] for the mean-square approximation of iterated Ito and Stratonovich stochastic integrals.

The rest of the article is organized as follows. In the introduction (below) we consider a brief review of the literature on the problem of construction of the Taylor–Ito and Taylor–Stratonovich expansions for the solutions of Ito SDEs. Sect. 2 is devoted to the integration order replacement technique for iterated Ito stochastic integrals. In Sect. 3, we consider the classical Taylor–Ito expansion while Sect. 4 and Sect. 5 are devoted to the first and second forms of the so-called unified Taylor–Ito expansion correspondingly. The classical Taylor–Stratonovich expansion is considered in Sect. 6. The first and second forms of the unified Taylor–Stratonovich expansion are derived in Sect. 7 and Sect. 8. In Sect. 9, we give a comparative analysis of the unified Taylor–Ito and Taylor–Stratonovich expansions with the classical Taylor–Ito and Taylor–Stratonovich expansions. Application of the first form of the unified Taylor–Ito expansion to the high-order strong numerical methods for Ito SDEs is considered

in Sect. 10. In Sect. 11, we construct the high-order strong numerical methods for Ito SDEs on the base of the first form of the unified Taylor–Stratonovich expansion. Sect. 12 is devoted to the effective method of the mean-square approximation of iterated Ito and Stratonovich stochastic integrals based on generalized multiple Fourier series. In Sect. 13, we discuss the connection between Theorems 8, 10–13 (see Sect. 12) and Wong–Zakai approximation.

Let us consider the following iterated Ito and Stratonovich stochastic integrals

$$(2) \quad J[\psi^{(k)}]_{s,t} = \int_t^s \psi_k(t_k) \dots \int_t^{t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_k}^{(i_k)},$$

$$(3) \quad J^*[\psi^{(k)}]_{s,t} = \int_t^{*s} \psi_k(t_k) \dots \int_t^{*t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_k}^{(i_k)},$$

where $0 \leq t < s \leq T$, every $\psi_l(\tau)$ ($l = 1, \dots, k$) is a nonrandom function at the interval $[t, T]$, $\mathbf{w}_\tau^{(i)} = \mathbf{f}_\tau^{(i)}$ for $i = 1, \dots, m$ and $\mathbf{w}_\tau^{(0)} = \tau$, $i_1, \dots, i_k = 0, 1, \dots, m$,

$$\int \quad \text{and} \quad \int^*$$

denote Ito and Stratonovich stochastic integrals, respectively. In this paper we use the definition of the Stratonovich stochastic integral from [2].

It could be noted that one of the main problems when constructing the high-order strong numerical methods for Ito SDEs on the base of the Taylor–Ito and Taylor–Stratonovich expansions is the mean-square approximation of iterated Ito and Stratonovich stochastic integrals. Obviously, in the absence of procedures for the numerical simulation of stochastic integrals, the mentioned numerical methods are unrealizable in practice. For this reason, in Sect. 12 we give a brief overview to the effective method of the mean-square approximation of iterated Ito and Stratonovich stochastic integrals (2) and (3) of arbitrary multiplicity k ($k \in \mathbb{N}$), which is proposed and developed by the author of this article in a number of publications [12]–[63]. This method is based on the generalized multiple Fourier series converging in the mean-square sense. The extensive practical material on expansions and mean-square approximations of iterated Ito and Stratonovich stochastic integrals of multiplicities 1 to 6 from the Taylor–Ito and Taylor–Stratonovich expansions is given in Sect. 12. In the mentioned section, the main focus is on approximations based on multiple Fourier–Legendre series. Such approximations is more effective in comparison with the trigonometric approximations [23], [39] at least for the numerical methods with the strong order 1.5 of convergence and higher [23], [39].

Let us give a brief review of the literature on the problem of construction of the Taylor–Ito and Taylor–Stratonovich expansions for the solutions of Ito SDEs. A few variants of a stochastic analog of the Taylor formula have been obtained in [2]–[7] for the stochastic processes in the form $R(\mathbf{x}_s, s)$, where \mathbf{x}_s is a solution of the Ito SDE (1) and $R: \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^1$ is a nonrandom sufficiently smooth function.

The first result in this direction called the Ito–Taylor expansion has been obtained in [6], [7]. This result gives an expansion of the process $R(\mathbf{x}_s, s)$ into a series such that every term (if $k > 0$) contains an iterated Ito stochastic integral

$$(4) \quad \int_t^s \dots \int_t^{t_2} d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_k}^{(i_k)}$$

as a factor, where $0 \leq t < s \leq T$, $i_1, \dots, i_k = 0, 1, \dots, m$. Obviously that the iterated Ito stochastic integral (4) is a particular case of (2) for $\psi_1(\tau), \dots, \psi_k(\tau) \equiv 1$.

In [7], another expansion of the stochastic process $R(\mathbf{x}_s, s)$ in a series has been derived. Instead of the Ito integrals, the iterated Stratonovich stochastic integrals

$$(5) \quad \int_t^{*s} \dots \int_t^{*t_2} d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_k}^{(i_k)},$$

were used; the corresponding expansion was called the Stratonovich–Taylor expansion. In the formula (5), the indices i_1, \dots, i_k take values $0, 1, \dots, m$.

In [8] the Ito–Taylor expansion from [6], [7] is reduced to the interesting and unexpected form (called the unified Taylor–Ito expansion) with the help of special transformations [66] (also see [33], [67], [68]). Every term of this expansion (if $k > 0$) contains an iterated Ito stochastic integral of the form

$$(6) \quad \int_t^s (s - t_k)^{l_k} \dots \int_t^{t_2} (s - t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)},$$

where $0 \leq t < s \leq T$, $l_1, \dots, l_k = 0, 1, 2, \dots$ and $i_1, \dots, i_k = 1, \dots, m$.

It is worth to mention another form of the unified Taylor–Ito expansion [10], [11], [67] (also see [46]–[50], [54]–[61]). Terms of the latter expansion contain iterated Ito stochastic integrals of the form

$$(7) \quad \int_t^s (t - t_k)^{l_k} \dots \int_t^{t_2} (t - t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)},$$

where $l_1, \dots, l_k = 0, 1, 2, \dots$ and $i_1, \dots, i_k = 1, \dots, m$.

In this paper, we derive two new forms of the Taylor–Ito expansions (the so-called unified Taylor–Ito expansions [10], [11], [67] (also see [46]–[50], [54]–[61])) using an approach which is taken from [9]. Obviously that some of iterated Ito stochastic integrals of the form (4) or (5) are connected by linear relations, while this is not the case for integrals of the form (6), (7). In this sense, the total quantity of stochastic integrals of the form (6) or (7) is minimal. Futhermore, in this article we construct two new forms of the Taylor–Stratonovich expansion (the so-called unified Taylor–Stratonovich expansions [9]) such that every term (if $k > 0$) contains as a multiplier an iterated Stratonovich stochastic integral of one of two types

$$(8) \quad \int_t^{*s} (t - t_k)^{l_k} \dots \int_t^{*t_2} (t - t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)},$$

$$(9) \quad \int_t^{*s} (s - t_k)^{l_k} \dots \int_t^{*t_2} (s - t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)},$$

where $l_1, \dots, l_k = 0, 1, 2, \dots$, $i_1, \dots, i_k = 1, \dots, m$, and $k = 1, 2, \dots$.

It is not difficult to see that for the sets of iterated Stratonovich stochastic integrals (8) and (9) the property of minimality (see above) also holds as for the sets of iterated Ito stochastic integrals (6), (7).

As we noted above, the main problem in implementation of high-order strong numerical methods for Ito SDEs is the mean-square approximation of iterated stochastic integrals (4)–(9). Obviously, these stochastic integrals are particular cases of the stochastic integrals (2), (3).

Taking into account the results of [12]–[65] and the minimality of the sets of stochastic integrals of the forms (6)–(9), we conclude that the unified Taylor–Ito and Taylor–Stratonovich expansions based on the iterated stochastic integrals (6)–(9) may be useful for constructing of high-order strong numerical methods with the orders of convergence 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, ... for Ito SDEs.

2. INTEGRATION ORDER REPLACEMENT TECHNIQUE FOR ITERATED ITO STOCHASTIC INTEGRALS

Let $f_\tau, \tau \in [0, T]$ be a scalar standard Wiener process that is F_τ -measurable for every $\tau \in [0, T]$. We introduce a class $\mathfrak{M}_2([t, T])$ ($t \geq 0$) of random functions $\xi(\tau, \omega) \stackrel{\text{def}}{=} \xi_\tau : [t, T] \times \Omega \rightarrow \mathbb{R}^1$ having the following properties: these functions are measurable with respect to the pair (τ, ω) of variables, F_τ -measurable for every $\tau \in [t, T]$, and satisfy the conditions

$$\int_t^T \mathbf{M} \{ \xi_\tau^2 \} d\tau < \infty$$

and $\mathbf{M} \{ \xi_\tau^2 \} < \infty$ for any $\tau \in [t, T]$.

On the class $\mathfrak{M}_2([t, T])$ ($t \geq 0$), we introduce the Hilbert norm

$$\|\xi\|_{2,T,t} = \left(\int_t^T \mathbf{M} \{ \xi_\tau^2 \} d\tau \right)^{1/2}.$$

Let $\{ \tau_j^{(N)} \}_{j=1}^N$ be a partition of the interval $[t, T]$ such that

$$(10) \quad t = \tau_0^{(N)} < \tau_1^{(N)} < \dots < \tau_N^{(N)} = T, \quad \Delta_N = \max_{0 \leq j \leq N-1} \left| \tau_{j+1}^{(N)} - \tau_j^{(N)} \right| \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

Let $\xi^{(N)}(\tau, \omega)$ be a sequence of step functions from the space $\mathfrak{M}_2([t, T])$ defined as follows

$$\xi^{(N)}(\tau, \omega) = \xi_j(\omega) \quad \text{w. p. 1 for } \tau \in \left[\tau_j^{(N)}, \tau_{j+1}^{(N)} \right), \quad j = 0, 1, \dots, N-1,$$

where here and further w. p. 1 means with probability 1.

It is known [1] that for any function $\xi_\tau \in \mathfrak{M}_2([t, T])$ there exists a sequence $\xi^{(N)}(\tau, \omega) \in \mathfrak{M}_2([t, T])$, which converges to the function ξ_τ in the sense of norm $\|\cdot\|_{2,T,t}$.

The mean-square limit

$$(11) \quad \text{l.i.m.}_{N \rightarrow \infty} \sum_{j=0}^{N-1} \xi^{(N)}(\tau_j^{(N)}, \omega) \left(f(\tau_{j+1}^{(N)}, \omega) - f(\tau_j^{(N)}, \omega) \right) \stackrel{\text{def}}{=} \int_t^T \xi_\tau df_\tau$$

is called [1] the Ito stochastic integral of a function $\xi_\tau \in \mathfrak{M}_2([t, T])$. Here $\xi^{(N)}(\tau, \omega)$ is an arbitrary sequence of step functions from the class $\mathfrak{M}_2([t, T])$ converging to the function $\xi(\tau, \omega)$ in the sense of norm $\|\cdot\|_{2,T,t}$, i.e.

$$(12) \quad \lim_{N \rightarrow \infty} \int_t^T \mathbf{M} \left\{ \left| \xi^{(N)}(\tau, \omega) - \xi(\tau, \omega) \right|^2 \right\} d\tau = 0.$$

We introduce the class $\mathfrak{Q}_m([t, T])$ ($t \geq 0$) of Ito processes $\eta_\tau, \tau \in [t, T]$ of the form

$$(13) \quad \eta_\tau = \eta_t + \int_t^\tau a_s ds + \int_t^\tau b_s df_s,$$

where $(a_\tau)^m, (b_\tau)^m \in \mathfrak{M}_2([t, T])$ and

$$\lim_{s \rightarrow \tau} \mathbf{M} \left\{ |b_s - b_\tau|^4 \right\} = 0$$

for all $\tau \in [t, T]$.

Let $C^{2,1}(\mathbb{R}^1 \times [t, T])$ ($t \geq 0$) be the space of functions $F(x, \tau) : \mathbb{R}^1 \times [t, T] \rightarrow \mathbb{R}^1$ with the following property: these functions are twice differentiable in x and have one derivative in τ . Moreover, all these derivatives are bounded.

The mean-square limit

$$(14) \quad \text{l.i.m}_{N \rightarrow \infty} \sum_{j=0}^{N-1} F \left(\frac{1}{2} \left(\eta_{\tau_j^{(N)}} + \eta_{\tau_{j+1}^{(N)}} \right), \tau_j^{(N)} \right) \left(f_{\tau_{j+1}^{(N)}} - f_{\tau_j^{(N)}} \right) \stackrel{\text{def}}{=} \int_t^{*T} F(\eta_\tau, \tau) df_\tau$$

is called [69] the Stratonovich stochastic integral of the process $F(\eta_\tau, \tau)$, $\tau \in [t, T]$ ($t \geq 0$), where $F(x, \tau) \in C^{2,1}(\mathbb{R}^1 \times [t, T])$. We apply in the formula (14) the same notations as in the formula (11).

It is known [69] (also see [2]) that under proper conditions, the following relation holds

$$(15) \quad \int_t^{*T} F(\eta_\tau, \tau) df_\tau = \int_t^T F(\eta_\tau, \tau) df_\tau + \frac{1}{2} \int_t^T \frac{\partial F}{\partial x}(\eta_\tau, \tau) b_\tau d\tau \quad \text{w. p. 1.}$$

If the Wiener processes in the formulas (13) and (14) are independent, then

$$(16) \quad \int_t^{*T} F(\eta_\tau, \tau) df_\tau = \int_t^T F(\eta_\tau, \tau) df_\tau \quad \text{w. p. 1.}$$

Note that a possible variant of conditions providing the correctness of the formulas (15) and (16) consists of the following conditions

$$\eta_\tau \in \mathfrak{Q}_4([t, T]), \quad F(\eta_\tau, \tau) \in \mathfrak{M}_2([t, T]), \quad \text{and} \quad F(x, \tau) \in C^{2,1}(\mathbb{R}^1 \times [t, T]).$$

Note that if $F(x, \tau) = F_1(x)F_2(\tau)$, then the smoothness condition $F(x, \tau) \in C^{2,1}(\mathbb{R}^1 \times [t, T])$ can be weakened. Namely, it suffices to replace the condition with respect to τ by continuity with respect to this variable.

A theorem allowing the change of order of integration in the iterated Ito stochastic integrals has been proved in [66]-[68] (also see [33]). In what follows, we apply this theorem; let us cite its exact formulation and the notations.

It is well known that the Ito stochastic integral exists in the mean-square sense (see (11)), if the stochastic process $\xi(\tau, \omega) \in \mathfrak{M}_2([0, T])$, that is, perhaps this process does not satisfy the property of the mean-square continuity on the interval $[0, T]$. Let us formulate the theorem on integration order replacement for the special class of iterated Ito stochastic integrals. At the same time, the condition of the mean-square continuity of integrand in the innermost stochastic integral will be significant.

Let $\mathfrak{S}_2([t, T])$ ($t \geq 0$) be the class of functions $\xi : [t, T] \times \Omega \rightarrow \mathbb{R}^1$, which satisfy the conditions:

1. $\xi_\tau \in \mathfrak{M}_2([t, T])$.
2. ξ_τ is the mean-square continuous stochastic process at the interval $[t, T]$.

Let us introduce the following class of iterated Ito stochastic integrals

$$J[\phi, \psi^{(k)}]_{T,t} = \int_t^T \psi_1(t_1) \dots \int_t^{t_{k-1}} \psi_k(t_k) \int_t^{t_k} \phi_\tau dw_\tau^{(k+1)} dw_{t_k}^{(k)} \dots dw_{t_1}^{(1)},$$

where $\phi_\tau \in \mathfrak{S}_2([t, T])$, every $\psi_l(\tau)$ ($l = 1, \dots, k$) is a continuous nonrandom function at the interval $[t, T]$, here and further $w_\tau^{(l)} = f_\tau$ or $w_\tau^{(l)} = \tau$ for $\tau \in [t, T]$ ($l = 1, \dots, k+1$), $(\psi_1, \dots, \psi_k) \stackrel{\text{def}}{=} \psi^{(k)}$, $\psi^{(1)} \stackrel{\text{def}}{=} \psi_1$.

In [69] Stratonovich introduced the definition of the so-called combined stochastic integral for the specific class of integrated processes. Taking this definition as a foundation, let us consider the following construction of stochastic integral

$$(17) \quad \text{l.i.m.}_{N \rightarrow \infty} \sum_{j=0}^{N-1} \phi_{\tau_j} (f_{\tau_{j+1}} - f_{\tau_j}) \theta_{\tau_{j+1}} \stackrel{\text{def}}{=} \int_t^T \phi_\tau df_\tau \theta_\tau,$$

where $\phi_\tau, \theta_\tau \in \mathfrak{S}_2([t, T])$, $\{\tau_j\}_{j=0}^N$ is the partition of the interval $[t, T]$, which satisfies the condition (10) (for simplicity we write here and sometimes further τ_j instead of $\tau_j^{(N)}$).

Further, we will use integrals of the type (17) ($\phi_\tau \in \mathfrak{S}_2([t, T])$ and θ_τ from a little bit narrower class of stochastic processes than $\mathfrak{S}_2([t, T])$) for formulation the theorem on integration order replacement for iterated Ito stochastic integrals $J[\phi, \psi^{(k)}]_{T,t}$, $k \geq 1$.

Note that under the appropriate conditions the following properties of stochastic integrals defined by the formula (17) can be proved

$$\int_t^T \phi_\tau df_\tau g(\tau) = \int_t^T \phi_\tau g(\tau) df_\tau \quad \text{w. p. 1,}$$

where $g(\tau)$ is a continuous nonrandom function at the interval $[t, T]$,

$$\int_t^T (\alpha \phi_\tau + \beta \psi_\tau) df_\tau \theta_\tau = \alpha \int_t^T \phi_\tau df_\tau \theta_\tau + \beta \int_t^T \psi_\tau df_\tau \theta_\tau \quad \text{w. p. 1,}$$

$$\int_t^T \phi_\tau df_\tau (\alpha \theta_\tau + \beta \psi_\tau) = \alpha \int_t^T \phi_\tau df_\tau \theta_\tau + \beta \int_t^T \phi_\tau df_\tau \psi_\tau \quad \text{w. p. 1,}$$

where $\alpha, \beta \in \mathbb{R}^1$. At that, we suppose that the stochastic processes ϕ_τ, θ_τ , and ψ_τ are such that all integrals (included in the mentioned properties) exist.

Let us define the stochastic integrals $\hat{I}[\psi^{(k)}]_{T,s}$, $k \geq 1$ of the form

$$\hat{I}[\psi^{(k)}]_{T,s} = \int_s^T \psi_k(t_k) dw_{t_k}^{(k)} \int_{t_k}^T \psi_{k-1}(t_{k-1}) dw_{t_{k-1}}^{(k-1)} \dots \int_{t_2}^T \psi_1(t_1) dw_{t_1}^{(1)}$$

in accordance with the definition (17) by the following recurrence relation

$$(18) \quad \hat{I}[\psi^{(k)}]_{T,t} \stackrel{\text{def}}{=} \text{l.i.m.}_{N \rightarrow \infty} \sum_{l=0}^{N-1} \psi_k(\tau_l) \left(w_{\tau_{l+1}}^{(k)} - w_{\tau_l}^{(k)} \right) \hat{I}[\psi^{(k-1)}]_{T,\tau_{l+1}},$$

where $k \geq 1$, $\hat{I}[\psi^{(0)}]_{T,s} \stackrel{\text{def}}{=} 1$, and $[s, T] \subseteq [t, T]$.

Then, we will define the iterated stochastic integral $\hat{J}[\phi, \psi^{(k)}]_{T,t}$, $k \geq 1$

$$\hat{J}[\phi, \psi^{(k)}]_{T,t} = \int_t^T \phi_s dw_s^{(k+1)} \hat{I}[\psi^{(k)}]_{T,s}$$

similarly in accordance with the definition (17)

$$\hat{J}[\phi, \psi^{(k)}]_{T,t} \stackrel{\text{def}}{=} \text{l.i.m.}_{N \rightarrow \infty} \sum_{l=0}^{N-1} \phi_{\tau_l} \left(w_{\tau_{l+1}}^{(k+1)} - w_{\tau_l}^{(k+1)} \right) \hat{I}[\psi^{(k)}]_{T,\tau_{l+1}}.$$

Let us formulate the theorem on integration order replacement for iterated Ito stochastic integrals.

Theorem 1 [66]-[68] (also see [33]). *Suppose that $\phi_\tau \in \mathfrak{S}_2([t, T])$ and every $\psi_l(\tau)$ ($l = 1, \dots, k$) is a continuous nonrandom function at the interval $[t, T]$. Then, the stochastic integral $\hat{J}[\phi, \psi^{(k)}]_{T,t}$ ($k \geq 1$) exists and*

$$J[\phi, \psi^{(k)}]_{T,t} = \hat{J}[\phi, \psi^{(k)}]_{T,t} \quad w. p. 1.$$

Let us consider some propositions related to Theorem 1.

Proposition 1 [66]-[68] (also see [33]). *Let the conditions of Theorem 1 are fulfilled and $h(\tau)$ is a continuous nonrandom function at the interval $[t, T]$. Then*

$$(19) \quad \int_t^T \phi_\tau dw_\tau^{(k+1)} h(\tau) \hat{I}[\psi^{(k)}]_{T,\tau} = \int_t^T \phi_\tau h(\tau) dw_\tau^{(k+1)} \hat{I}[\psi^{(k)}]_{T,\tau} \quad w. p. 1,$$

and the integrals on the left-hand side of (19) as well as on the right-hand side of (19) exist.

Proposition 2 [66]-[68] (also see [33]). *Under the conditions of Theorem 1 the following equality is satisfied*

$$(20) \quad \begin{aligned} & \int_t^T h(t_1) \int_t^{t_1} \phi_\tau dw_\tau^{(k+2)} dw_{t_1}^{(k+1)} \hat{I}[\psi^{(k)}]_{T,t_1} = \\ & = \int_t^T \phi_\tau dw_\tau^{(k+2)} \int_\tau^T h(t_1) dw_{t_1}^{(k+1)} \hat{I}[\psi^{(k)}]_{T,t_1} \quad w. p. 1. \end{aligned}$$

Moreover, the stochastic integrals in (20) exist.

Using the integration order replacement technique for iterated Ito stochastic integrals (Theorem 1), we can obtain different equalities for iterated Ito stochastic integrals. At that, the mentioned

technique is essentially simpler (for the specific class of Ito processes which are the iterated Ito stochastic integrals) in application than the Ito formula. Let us consider two examples on application of the integration order replacement technique for iterated Ito stochastic integrals.

Example 1. *Using Theorem 1 and Proposition 1, we obtain*

$$\begin{aligned}
& \int_t^T \int_t^{t_3} \int_t^{t_2} df_{t_1} df_{t_2} dt_3 = \int_t^T df_{t_1} \int_{t_1}^T df_{t_2} \int_{t_2}^T dt_3 = \\
& = \int_t^T df_{t_1} \int_{t_1}^T df_{t_2} (T - t_2) = \int_t^T df_{t_1} \int_{t_1}^T (T - t_2) df_{t_2} = \\
& = \int_t^T (T - t_2) \int_t^{t_2} df_{t_1} df_{t_2} \quad \text{w. p. 1.}
\end{aligned}$$

Example 2. *Using Theorem 1 and Proposition 1, we obtain*

$$\begin{aligned}
& \int_t^T \int_t^{t_4} \int_t^{t_3} \int_t^{t_2} df_{t_1} dt_2 df_{t_3} dt_4 = \int_t^T df_{t_1} \int_{t_1}^T dt_2 \int_{t_2}^T df_{t_3} \int_{t_3}^T dt_4 = \\
& = \int_t^T df_{t_1} \int_{t_1}^T dt_2 \int_{t_2}^T df_{t_3} (T - t_3) = \int_t^T df_{t_1} \int_{t_1}^T dt_2 \int_{t_2}^T (T - t_3) df_{t_3} = \\
& = \int_t^T (T - t_3) \int_t^{t_3} \int_t^{t_2} df_{t_1} dt_2 df_{t_3} = \int_t^T (T - t_3) \left(\int_t^{t_3} \int_t^{t_2} df_{t_1} dt_2 \right) df_{t_3} = \\
& = \int_t^T (T - t_3) \left(\int_t^{t_3} df_{t_1} \int_{t_1}^{t_3} dt_2 \right) df_{t_3} = \int_t^T (T - t_3) \left(\int_t^{t_3} df_{t_1} (t_3 - t_1) \right) df_{t_3} = \\
& = \int_t^T (T - t_3) \left(\int_t^{t_3} (t_3 - t_1) df_{t_1} \right) df_{t_3} = \int_t^T (T - t_2) \int_t^{t_2} (t_2 - t_1) df_{t_1} df_{t_2} \quad \text{w. p. 1.}
\end{aligned}$$

Let us apply Theorem 1 to deriving of one property for Ito stochastic integrals.

Lemma 1. *Let $h(\tau), g(\tau), G(\tau) : [t, s] \rightarrow \mathbb{R}^1$ be continuous nonrandom functions at the interval $[t, s]$ and let $G(\tau)$ be a antiderivative of the function $g(\tau)$. Furthermore, let $\xi_\tau \in \mathfrak{S}_2([t, s])$. Then*

$$(21) \quad \int_t^s g(\tau) \int_t^\tau h(\theta) \int_t^\theta \xi_u d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} d\tau = \int_t^s (G(s) - G(\theta)) h(\theta) \int_t^\theta \xi_u d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)}$$

w. p. 1, where $i, j = 1, 2$ and $\mathbf{f}_\tau^{(1)}, \mathbf{f}_\tau^{(2)}$ are independent standard Wiener processes that are F_τ -measurable for all $\tau \in [t, s]$.

Proof. Applying Theorem 1 twice and Proposition 1, we get the following relations

$$\begin{aligned}
& \int_t^s g(\tau) \int_t^\tau h(\theta) \int_t^\theta \xi_u d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} d\tau = \int_t^s \xi_u d\mathbf{f}_u^{(i)} \int_u^s h(\theta) d\mathbf{f}_\theta^{(j)} \int_\theta^s g(\tau) d\tau = \\
& = G(s) \int_t^s \xi_u d\mathbf{f}_u^{(i)} \int_u^s h(\theta) d\mathbf{f}_\theta^{(j)} - \int_t^s \xi_u d\mathbf{f}_u^{(i)} \int_u^s G(\theta) h(\theta) d\mathbf{f}_\theta^{(j)} = \\
& = G(s) \int_t^s h(\theta) \int_t^\theta \xi_u d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} - \int_t^s G(\theta) h(\theta) \int_t^\theta \xi_u d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} = \\
(22) \quad & = \int_t^s (G(s) - G(\theta)) h(\theta) \int_t^\theta \xi_u d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} \quad \text{w. p. 1.}
\end{aligned}$$

The proof of Lemma 1 is completed. Let us consider an analogue of Lemma 1 for Stratonovich stochastic integrals.

Lemma 2 [9]. *Let $h(\tau), g(\tau), G(\tau) : [t, s] \rightarrow \mathbb{R}^1$ be continuous nonrandom functions at the interval $[t, s]$ and let $G(\tau)$ be an antiderivative of the function $g(\tau)$. Let $\xi_\tau^{(l)} \in \mathfrak{Q}_4([t, s])$ and*

$$\xi_\tau^{(l)} = \int_t^\tau a_u du + \int_t^\tau b_u d\mathbf{f}_u^{(l)}, \quad l = 1, 2.$$

Then

$$(23) \quad \int_t^s g(\tau) \int_t^{*\tau} h(\theta) \int_t^{*\theta} \xi_u^{(l)} d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} d\tau = \int_t^s (G(s) - G(\theta)) h(\theta) \int_t^{*\theta} \xi_u^{(l)} d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)}$$

w. p. 1, where $i, j, l = 1, 2$ and $\mathbf{f}_\tau^{(1)}, \mathbf{f}_\tau^{(2)}$ are independent standard Wiener processes that are F_τ -measurable for all $\tau \in [t, s]$.

Proof. Under the conditions of Lemma 2, we can apply equalities (15) and (16) with $F(x, \theta) \equiv xh(\theta)$,

$$\eta_\theta = \int_t^{*\theta} \xi_u^{(l)} d\mathbf{f}_u^{(i)},$$

since the function $xh(\theta)$ is sufficiently smooth and the following obvious inclusions hold: $\eta_\theta \in \mathfrak{Q}_4([t, s])$ and $\eta_\theta h(\theta) \in \mathfrak{M}_2([t, s])$. Thus, we have the equalities

$$(24) \quad \int_t^{*\tau} h(\theta) \int_t^{*\theta} \xi_u^{(l)} d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} = \int_t^\tau h(\theta) \int_t^\theta \xi_u^{(l)} d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} + \frac{1}{2} \mathbf{1}_{\{i=j\}} \int_t^\tau h(\theta) \xi_\theta^{(l)} d\theta,$$

$$(25) \quad \int_t^{*\theta} \xi_u^{(l)} d\mathbf{f}_u^{(i)} = \int_t^\theta \xi_u^{(l)} d\mathbf{f}_u^{(i)} + \frac{1}{2} \mathbf{1}_{\{l=i\}} \int_t^\theta b_u du$$

w. p. 1, where $\mathbf{1}_A$ is the indicator of a set A . Substituting formulas (24) and (25) into the left-hand side of equality (23) and applying Theorem 1 twice and Proposition 1, we get the following relations

$$\begin{aligned}
& \int_t^s g(\tau) \int_t^{*\tau} h(\theta) \int_t^{*\theta} \xi_u^{(l)} d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} d\tau = \\
& = \int_t^s \xi_u^{(l)} d\mathbf{f}_u^{(i)} \int_u^s h(\theta) d\mathbf{f}_\theta^{(j)} \int_\theta^s g(\tau) d\tau + \\
& + \frac{1}{2} \mathbf{1}_{\{l=i\}} \int_t^s b_u du \int_u^s h(\theta) d\mathbf{f}_\theta^{(j)} \int_\theta^s g(\tau) d\tau + \frac{1}{2} \mathbf{1}_{\{i=j\}} \int_t^s h(\theta) \xi_\theta^{(l)} d\theta \int_\theta^s g(\tau) d\tau = \\
& = G(s) \left(\int_t^s \xi_u^{(l)} d\mathbf{f}_u^{(i)} \int_u^s h(\theta) d\mathbf{f}_\theta^{(j)} + \frac{1}{2} \mathbf{1}_{\{i=j\}} \int_t^s h(\theta) \xi_\theta^{(l)} d\theta + \right. \\
& \quad \left. + \frac{1}{2} \mathbf{1}_{\{l=i\}} \int_t^s b_u du \int_u^s h(\theta) d\mathbf{f}_\theta^{(j)} \right) - \\
& - \left(\int_t^s \xi_u^{(l)} d\mathbf{f}_u^{(i)} \int_u^s G(\theta) h(\theta) d\mathbf{f}_\theta^{(j)} + \frac{1}{2} \mathbf{1}_{\{i=j\}} \int_t^s G(\theta) h(\theta) \xi_\theta^{(l)} d\theta + \right. \\
& \quad \left. + \frac{1}{2} \mathbf{1}_{\{l=i\}} \int_t^s b_u du \int_u^s h(\theta) G(\theta) d\mathbf{f}_\theta^{(j)} \right) = \\
& = G(s) \left(\int_t^s h(\theta) \int_t^\theta \xi_u^{(l)} d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} + \frac{1}{2} \mathbf{1}_{\{i=j\}} \int_t^s h(\theta) \xi_\theta^{(l)} d\theta + \right. \\
& \quad \left. + \frac{1}{2} \mathbf{1}_{\{l=i\}} \int_t^s h(\theta) \int_t^\theta b_u du d\mathbf{f}_\theta^{(j)} \right) - \\
& - \left(\int_t^s G(\theta) h(\theta) \int_t^\theta \xi_u^{(l)} d\mathbf{f}_u^{(i)} d\mathbf{f}_\theta^{(j)} + \frac{1}{2} \mathbf{1}_{\{i=j\}} \int_t^s G(\theta) h(\theta) \xi_\theta^{(l)} d\theta + \right. \\
& \quad \left. + \frac{1}{2} \mathbf{1}_{\{l=i\}} \int_t^s h(\theta) G(\theta) \int_t^\theta b_u du d\mathbf{f}_\theta^{(j)} \right)
\end{aligned} \tag{26}$$

w. p. 1. Applying successively the formulas (24), (25) together with the formula (24) in which $h(\theta)$ replaced by $G(\theta)h(\theta)$ as well as the relation (26), we obtain the equality (23).

3. THE TAYLOR–ITO EXPANSION

In this section, we cite the Taylor-Ito expansion [7] and introduce some necessary notations. At that, we will use the original notation introduced by the author of this paper.

Let \mathfrak{L} be the class of functions $R(\mathbf{x}, t) : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^1$ with the following property: these functions are twice continuously differentiable in \mathbf{x} and have one continuous derivative in t . We consider the following operators on the space \mathfrak{L}

$$(27) \quad \begin{aligned} LR(\mathbf{x}, t) &= \frac{\partial R}{\partial t}(\mathbf{x}, t) + \sum_{i=1}^n a^{(i)}(\mathbf{x}, t) \frac{\partial R}{\partial \mathbf{x}^{(i)}}(\mathbf{x}, t) + \\ &+ \frac{1}{2} \sum_{j=1}^m \sum_{l, i=1}^n B^{(lj)}(\mathbf{x}, t) B^{(ij)}(\mathbf{x}, t) \frac{\partial^2 R}{\partial \mathbf{x}^{(l)} \partial \mathbf{x}^{(i)}}(\mathbf{x}, t), \end{aligned}$$

$$(28) \quad G_0^{(i)} R(\mathbf{x}, t) = \sum_{j=1}^n B^{(ji)}(\mathbf{x}, t) \frac{\partial R}{\partial \mathbf{x}^{(j)}}(\mathbf{x}, t), \quad i = 1, \dots, m.$$

By the Ito formula, we have the equality

$$(29) \quad R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \int_t^s LR(\mathbf{x}_\tau, \tau) d\tau + \sum_{i=1}^m \int_t^s G_0^{(i)} R(\mathbf{x}_\tau, \tau) d\mathbf{f}_\tau^{(i)}$$

w. p. 1, where $0 \leq t < s \leq T$. In the formula (29) it is assumed that the functions $\mathbf{a}(\mathbf{x}, t)$, $B(\mathbf{x}, t)$, and $R(\mathbf{x}, t)$ satisfy the following condition: $LR(\mathbf{x}_\tau, \tau)$, $G_0^{(i)} R(\mathbf{x}_\tau, \tau) \in \mathfrak{M}_2([0, T])$ for $i = 1, \dots, m$.

Introduce the following notation

$$(30) \quad {}^{(k)}A = \left\| A^{(i_1 \dots i_k)} \right\|_{i_1=1, \dots, i_k=1}^{m_1 \dots m_k}, \quad m_1, \dots, m_k \geq 1,$$

$${}^{(k+l)}A \cdot {}^{(l)}B^{(k)} = \begin{cases} \left\| \sum_{i_1=1}^{m_1} \dots \sum_{i_l=1}^{m_l} A^{(i_1 \dots i_{k+l})} B^{(i_1 \dots i_l)} \right\|_{i_{l+1}=1, \dots, i_{l+k}=1}^{m_{l+1} \dots m_{l+k}} & \text{for } k \geq 1 \\ \sum_{i_1=1}^{m_1} \dots \sum_{i_l=1}^{m_l} A^{(i_1 \dots i_l)} B^{(i_1 \dots i_l)} & \text{for } k = 0 \end{cases},$$

$$(31) \quad \left\| A_{k+1} D_k^{(i_k)} A_k \dots A_2 D_1^{(i_1)} A_1 R(\mathbf{x}, t) \right\|_{i_1=1, \dots, i_k=1}^{m_1 \dots m_k} = {}^{(k)}A_{k+1} D_k A_k \dots A_2 D_1 A_1 R(\mathbf{x}, t),$$

where A_p and $D_q^{(i_q)}$ are operators defined on the space \mathfrak{L} for $p = 1, \dots, k+1$, $q = 1, \dots, k$, and $i_q = 1, \dots, m_q$. It is assumed that the left-hand side of (31) exists. The symbol \cdot is treated as the

usual multiplication. If $m_l = 0$ in (30) for some $l \in \{1, \dots, k\}$, then the right-hand side of (30) is treated as

$$\left\| A^{(i_1 \dots i_{l-1} i_{l+1} \dots i_k)} \right\|_{i_1=1, \dots, i_{l-1}=1, i_{l+1}=1, \dots, i_k=1}^{m_1 \dots m_{l-1} m_{l+1} \dots m_k},$$

(shortly, $(k-1)A$).

We also introduce the following notation

$$\left\| Q_{\lambda_l}^{(i_l)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}, t) \right\|_{i_1=\lambda_1, \dots, i_l=\lambda_l}^{m\lambda_1 \dots m\lambda_l} \stackrel{\text{def}}{=} {}^{(p_l)}Q_{\lambda_l} \dots Q_{\lambda_1} R(\mathbf{x}, t),$$

$${}^{(p_k)}J_{(\lambda_k \dots \lambda_1)s,t} = \left\| J_{(\lambda_k \dots \lambda_1)s,t}^{(i_k \dots i_1)} \right\|_{i_1=\lambda_1, \dots, i_k=\lambda_k}^{m\lambda_1 \dots m\lambda_k},$$

$$M_k = \left\{ (\lambda_k, \dots, \lambda_1) : \lambda_l = 1 \text{ or } \lambda_l = 0; l = 1, \dots, k \right\}, \quad k \geq 1,$$

$$J_{(\lambda_k \dots \lambda_1)s,t}^{(i_k \dots i_1)} = \int_t^s \dots \int_t^{t_2} d\mathbf{w}_{t_1}^{(i_k)} \dots d\mathbf{w}_{t_k}^{(i_1)}, \quad k \geq 1,$$

where $\lambda_l = 1$ or $\lambda_l = 0$, $Q_{\lambda_l}^{(i_l)} = L$ and $i_l = 0$ for $\lambda_l = 0$, $Q_{\lambda_l}^{(i_l)} = G_0^{(i_l)}$ and $i_l = 1, \dots, m$ for $\lambda_l = 1$,

$$p_l = \sum_{j=1}^l \lambda_j \quad \text{for } l = 1, \dots, r+1, \quad r \in \mathbb{N},$$

$\mathbf{w}_\tau^{(i)}$ ($i = 1, \dots, m$) are F_τ -measurable for all $\tau \in [0, T]$ independent standard Wiener processes and $\mathbf{w}_\tau^{(0)} = \tau$.

Applying the formula (29) to the process $R(\mathbf{x}_s, s)$ repeatedly, we obtain the following Taylor–Ito expansion [7]

$$(32) \quad R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{k=1}^r \sum_{(\lambda_k, \dots, \lambda_1) \in M_k} {}^{(p_k)}Q_{\lambda_k} \dots Q_{\lambda_1} R(\mathbf{x}_t, t) {}^{p_k} {}^{(p_k)}J_{(\lambda_k \dots \lambda_1)s,t} + (D_{r+1})_{s,t}$$

w. p. 1, where

$$(33) \quad (D_{r+1})_{s,t} = \sum_{(\lambda_{r+1}, \dots, \lambda_1) \in M_{r+1}} \int_t^s \dots \left(\int_t^{t_2} {}^{(p_{r+1})}Q_{\lambda_{r+1}} \dots Q_{\lambda_1} R(\mathbf{x}_{t_1}, t_1) {}^{\lambda_{r+1}} \dots {}^{\lambda_1} d\mathbf{w}_{t_1} \right) \dots {}^{\lambda_1} d\mathbf{w}_{t_{r+1}}.$$

It is assumed that the right-hand sides of (32), (33) exist.

A possible variant of the conditions under which the right-hand sides of (32), (33) exist is as follows

- (i) $Q_{\lambda_l}^{(i_l)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}, t) \in \mathfrak{L}$ for all $(\lambda_l, \dots, \lambda_1) \in \bigcup_{g=1}^r M_g$;

(ii) $Q_{\lambda_l}^{(i_l)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}_\tau, \tau) \in \mathfrak{M}_2([0, T])$ for all $(\lambda_l, \dots, \lambda_1) \in \bigcup_{g=1}^{r+1} M_g$.

Let us write the expansion (32) in the another form

$$R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{k=1}^r \sum_{(\lambda_k, \dots, \lambda_1) \in M_k} \sum_{i_1=\lambda_1}^{m\lambda_1} \dots \sum_{i_k=\lambda_k}^{m\lambda_k} Q_{\lambda_k}^{(i_k)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}_t, t) J_{(\lambda_k \dots \lambda_1)_{s,t}}^{(i_k \dots i_1)} + (D_{r+1})_{s,t} \quad \text{w. p. 1.}$$

Denote

$$G_{rk} = \left\{ (\lambda_k, \dots, \lambda_1) : r+1 \leq 2k - \lambda_1 - \dots - \lambda_k \leq 2r \right\},$$

$$E_{qk} = \left\{ (\lambda_k, \dots, \lambda_1) : 2k - \lambda_1 - \dots - \lambda_k = q \right\},$$

where $\lambda_l = 1$ or $\lambda_l = 0$ ($l = 1, \dots, k$).

The Taylor–Ito expansion ordered according to the order of smallness (in the mean-square sense when $s \downarrow t$) of its terms has the form

$$R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{q,k=1}^r \sum_{(\lambda_k, \dots, \lambda_1) \in E_{qk}} \sum_{i_1=\lambda_1}^{m\lambda_1} \dots \sum_{i_k=\lambda_k}^{m\lambda_k} Q_{\lambda_k}^{(i_k)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}_t, t) J_{(\lambda_k \dots \lambda_1)_{s,t}}^{(i_k \dots i_1)} + (H_{r+1})_{s,t} \quad \text{w. p. 1,}$$

where

$$(H_{r+1})_{s,t} = \sum_{k=1}^r \sum_{(\lambda_k, \dots, \lambda_1) \in G_{rk}} \sum_{i_1=\lambda_1}^{m\lambda_1} \dots \sum_{i_k=\lambda_k}^{m\lambda_k} Q_{\lambda_k}^{(i_k)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}_t, t) J_{(\lambda_k \dots \lambda_1)_{s,t}}^{(i_k \dots i_1)} + (D_{r+1})_{s,t}.$$

4. THE FIRST FORM OF THE UNIFIED TAYLOR–ITO EXPANSION

In this section, we transform the right-hand side of (32) with the help of Theorem 1 and Lemma 1 to a representation including iterated Ito stochastic integrals of the form (7).

Denote

$$(35) \quad I_{l_1 \dots l_{k s,t}}^{(i_1 \dots i_k)} = \int_t^s (t - t_k)^{l_k} \dots \int_t^{t_2} (t - t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)} \quad \text{for } k \geq 1$$

and

$$I_{l_1 \dots l_{k s,t}}^{(i_1 \dots i_k)} = 1 \quad \text{for } k = 0,$$

where $i_1, \dots, i_k = 1, \dots, m$. Moreover, let

$$(36) \quad \begin{aligned} {}^{(k)}I_{l_1 \dots l_{k,s,t}} &= \left\| I_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)} \right\|_{i_1, \dots, i_k=1}^m, \\ G_p^{(i)} &\stackrel{\text{def}}{=} \frac{1}{p} \left(G_{p-1}^{(i)} L - L G_{p-1}^{(i)} \right), \quad p = 1, 2, \dots, \quad i = 1, \dots, m, \end{aligned}$$

where L and $G_0^{(i)}$, $i = 1, \dots, m$, are determined by the equalities (27), (28). Denote

$$\begin{aligned} A_q &\stackrel{\text{def}}{=} \left\{ (k, j, l_1, \dots, l_k) : k + j + \sum_{p=1}^k l_p = q; k, j, l_1, \dots, l_k = 0, 1, \dots \right\}, \\ \left\| G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} L^j R(\mathbf{x}, t) \right\|_{i_1, \dots, i_k=1}^m &\stackrel{\text{def}}{=} {}^{(k)}G_{l_1} \dots G_{l_k} L^j R(\mathbf{x}, t), \\ L^j R(\mathbf{x}, t) &\stackrel{\text{def}}{=} \begin{cases} \underbrace{L \dots L}_j R(\mathbf{x}, t) & \text{for } j \geq 1 \\ R(\mathbf{x}, t) & \text{for } j = 0 \end{cases}. \end{aligned}$$

Theorem 2. *Let conditions (i), (ii) be satisfied. Then for any $s, t \in [0, T]$ such that $s > t$ and for any positive integer r , the following expansion takes place w. p. 1*

$$(37) \quad \begin{aligned} R(\mathbf{x}_s, s) &= R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k, j, l_1, \dots, l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} L^j R(\mathbf{x}_t, t) I_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)} + \\ &+ (D_{r+1})_{s,t}, \end{aligned}$$

where $(D_{r+1})_{s,t}$ has the form (33).

Proof. We claim that

$$(38) \quad \begin{aligned} &\sum_{(\lambda_q, \dots, \lambda_1) \in M_q} {}^{(p_q)}Q_{\lambda_q} \dots Q_{\lambda_1} R(\mathbf{x}_t, t) {}^{p_q} J_{(\lambda_q \dots \lambda_1)_{s,t}} = \\ &= \sum_{(k, j, l_1, \dots, l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} L^j R(\mathbf{x}_t, t) I_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)} \end{aligned}$$

w. p. 1. The equality (38) is valid for $q = 1$. Assume that (38) is valid for some $q > 1$. In this case, using the induction hypothesis, we obtain

$$\begin{aligned}
& \sum_{(\lambda_{q+1}, \dots, \lambda_1) \in M_{q+1}} {}^{(p_{q+1})} Q_{\lambda_1} \dots Q_{\lambda_{q+1}} R(\mathbf{x}_t, t) {}^{p_{q+1}} J_{(\lambda_1 \dots \lambda_{q+1})s, t} = \\
& = \sum_{\lambda_{q+1} \in \{1, 0\}} \int_t^s \sum_{(\lambda_q, \dots, \lambda_1) \in M_q} \left({}^{(p_{q+1})} Q_{\lambda_1} \dots Q_{\lambda_{q+1}} R(\mathbf{x}_t, t) {}^{p_q} J_{(\lambda_1 \dots \lambda_q)\theta, t} \right)^{\lambda_{q+1}} d\mathbf{w}_\theta = \\
& = \sum_{\lambda_{q+1} \in \{1, 0\}} \int_t^s \sum_{(k, j, l_1, \dots, l_k) \in A_q} \frac{(\theta - t)^j}{j!} \times \\
& \quad \times \left({}^{(k+\lambda_{q+1})} G_{l_1} \dots G_{l_k} L^j Q_{\lambda_{q+1}} R(\mathbf{x}_t, t) {}^{(k)} I_{l_1 \dots l_{k, s, t}} \right)^{\lambda_{q+1}} d\mathbf{w}_\theta = \\
& = \sum_{(k, j, l_1, \dots, l_k) \in A_q} \left({}^{(k)} G_{l_1} \dots G_{l_k} L^{j+1} R(\mathbf{x}_t, t) {}^{(k)} \int_t^s \frac{(\theta - t)^j}{j!} {}^{(k)} I_{l_1 \dots l_{k, \theta, t}} d\theta + \right. \\
(39) \quad & \left. + \left({}^{(k+1)} G_{l_1} \dots G_{l_k} L^j G_0 R(\mathbf{x}_t, t) {}^{(k)} \int_t^s \frac{(\theta - t)^j}{j!} {}^{(k)} I_{l_1 \dots l_{k, \theta, t}} \right) {}^1 d\mathbf{f}_\theta \right)
\end{aligned}$$

w. p. 1.

Using Lemma 1, we obtain

$$\begin{aligned}
& \int_t^s \frac{(\theta - t)^j}{j!} {}^{(k)} I_{l_1 \dots l_{k, \theta, t}} d\theta = \\
(40) \quad & = \frac{1}{(j+1)!} \begin{cases} (s-t)^{j+1} & \text{for } k=0 \\ (s-t)^{j+1} \cdot {}^{(k)} I_{l_1 \dots l_{k, s, t}} - (-1)^{j+1} \cdot {}^{(k)} I_{l_1 \dots l_{k-1} \ l_{k+j+1}, s, t} & \text{for } k>0 \end{cases}
\end{aligned}$$

w. p. 1. In addition (see (35)), we get

$$(41) \quad \int_t^s \frac{(\theta - t)^j}{j!} I_{l_1 \dots l_{k, \theta, t}}^{(i_1 \dots i_k)} d\mathbf{f}_\theta^{(i_{k+1})} = \frac{(-1)^j}{j!} I_{l_1 \dots l_{k, s, t}}^{(i_1 \dots i_k i_{k+1})}$$

in the notations just introduced. Substitute the relations (40) and (41) into the formula (39). Grouping summands of the obtained expression with equal lower indices at iterated Ito stochastic integrals and using (36) and the equality

$$(42) \quad G_p^{(i)} R(\mathbf{x}, t) = \frac{1}{p!} \sum_{q=0}^p (-1)^q C_p^q L^q G_0^{(i)} L^{p-q} R(\mathbf{x}, t), \quad \text{where } C_p^q = \frac{p!}{q!(p-q)!},$$

(this equality follows from (36)), we note that the obtained expression is equal to

$$\sum_{(k,j,l_1,\dots,l_k) \in A_{q+1}} \frac{(s-t)^j}{j!} {}^{(k)}G_{l_1} \dots G_{l_k} L^j \{\eta_t\} \cdot {}^{(k)}I_{l_1 \dots l_{k_{s,t}}}$$

w. p. 1. Summing the equalities (38) for $q = 1, 2, \dots, r$ and applying the formula (32), we obtain the expression (37). The proof is completed.

Let us order terms of the expansion (37) according to their smallness orders as $s \downarrow t$ in the mean-square sense

$$(43) \quad R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k,j,l_1,\dots,l_k) \in D_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} L^j R(\mathbf{x}_t, t) I_{l_1 \dots l_{k_{s,t}}}^{(i_1 \dots i_k)} + (H_{r+1})_{s,t} \quad \text{w. p. 1,}$$

where

$$(44) \quad (H_{r+1})_{s,t} = \sum_{(k,j,l_1,\dots,l_k) \in U_r} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} L^j R(\mathbf{x}_t, t) I_{l_1 \dots l_{k_{s,t}}}^{(i_1 \dots i_k)} + (D_{r+1})_{s,t},$$

$$D_q = \left\{ (k, j, l_1, \dots, l_k) : k + 2 \left(j + \sum_{p=1}^k l_p \right) = q; k, j, l_1, \dots, l_k = 0, 1, \dots \right\},$$

$$(45) \quad U_r = \left\{ (k, j, l_1, \dots, l_k) : k + j + \sum_{p=1}^k l_p \leq r, k + 2 \left(j + \sum_{p=1}^k l_p \right) \geq r + 1; k, j, l_1, \dots, l_k = 0, 1, \dots \right\},$$

and $(D_{r+1})_{s,t}$ has the form (33). Note that the remainder term $(H_{r+1})_{s,t}$ in (43) has a higher order of smallness in the mean-square sense as $s \downarrow t$ than the terms of the main part of expansion (43).

5. THE SECOND FORM OF THE UNIFIED TAYLOR–ITO EXPANSION

Consider iterated Ito stochastic integrals of the form

$$J_{l_1 \dots l_{k_{s,t}}}^{(i_1 \dots i_k)} = \int_t^s (s-t_k)^{l_k} \dots \int_t^{t_2} (s-t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)} \quad \text{for } k \geq 1$$

and

$$J_{l_1 \dots l_{k_{s,t}}}^{(i_1 \dots i_k)} = 1 \quad \text{for } k = 0,$$

where $i_1, \dots, i_k = 1, \dots, m$.

The additive property of stochastic integrals and the Newton binomial formula imply the following equality

$$(46) \quad I_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)} = \sum_{j_1=0}^{l_1} \dots \sum_{j_k=0}^{l_k} \prod_{g=1}^k C_{l_g}^{j_g} (t-s)^{l_1+\dots+l_k-j_1-\dots-j_k} J_{j_1 \dots j_{k,s,t}}^{(i_1 \dots i_k)} \quad \text{w. p. 1,}$$

where

$$C_l^k = \frac{l!}{k!(l-k)!}$$

is the binomial coefficient. Thus, the Taylor–Ito expansion of the process $\eta_s = R(\mathbf{x}_s, s)$ can be constructed either using the iterated stochastic integrals $I_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)}$ similarly to the previous section or using the iterated stochastic integrals $J_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)}$. This is the main subject of this section.

Denote

$$\left\| J_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)} \right\|_{i_1, \dots, i_k=1}^m \stackrel{\text{def}}{=} {}^{(k)} J_{l_1 \dots l_{k,s,t}},$$

$$\left\| L^j G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} R(\mathbf{x}, t) \right\|_{i_1, \dots, i_k=1}^m \stackrel{\text{def}}{=} {}^{(k)} L^j G_{l_1} \dots G_{l_k} R(\mathbf{x}, t).$$

Theorem 3. *Let conditions (i), (ii) be satisfied. Then for any $s, t \in [0, T]$ such that $s > t$ and for any positive integer r , the following expansion is valid w. p. 1*

$$(47) \quad R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k,j,l_1, \dots, l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m L^j G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} R(\mathbf{x}_t, t) J_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)} + (D_{r+1})_{s,t},$$

where $(D_{r+1})_{s,t}$ has the form (33).

Proof. To prove the theorem, we check the equalities

$$(48) \quad \sum_{(k,j,l_1, \dots, l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m L^j G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} R(\mathbf{x}_t, t) J_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)} = \sum_{(k,j,l_1, \dots, l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} L^j R(\mathbf{x}_t, t) I_{l_1 \dots l_{k,s,t}}^{(i_1 \dots i_k)} \quad \text{w. p. 1}$$

for $q = 1, 2, \dots, r$. To check (48), substitute the expression (46) into the right-hand side of (48) and then use the formulas (36), (42).

Let us rank terms of the expansion (47) according to their orders of smallness in the mean-square sense as $s \downarrow t$

$$\begin{aligned}
R(\mathbf{x}_s, s) &= R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k,j,l_1,\dots,l_k) \in D_q} \frac{(s-t)^j}{j!} \sum_{i_1,\dots,i_k=1}^m L^j G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} R(\mathbf{x}_t, t) J_{l_1 \dots l_k s, t}^{(i_1 \dots i_k)} + \\
&\quad + (H_{r+1})_{s,t} \quad \text{w. p. 1,}
\end{aligned}$$

where

$$(H_{r+1})_{s,t} = \sum_{(k,j,l_1,\dots,l_k) \in U_r} \frac{(s-t)^j}{j!} \sum_{i_1,\dots,i_k=1}^m L^j G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} R(\mathbf{x}_t, t) J_{l_1 \dots l_k s, t}^{(i_1 \dots i_k)} + (D_{r+1})_{s,t}.$$

The term $(D_{r+1})_{s,t}$ has the form (33); the terms D_q and U_r have the forms (44) and (45), respectively. Finally, we note that the convergence w. p. 1 of the truncated Taylor–Ito expansion (32) (without the remainder term $(D_{r+1})_{s,t}$) to the process $R(\mathbf{x}_s, s)$ as $r \rightarrow \infty$ for all $s, t \in [0, T]$ such that $s > t$ and $T < \infty$ has been proved in [2] (Proposition 5.9.2). Since expansions (37) and (47) are obtained from the Taylor–Ito expansion (32) without any additional conditions, the truncated expansions (37) and (47) (without the remainder term $(D_{r+1})_{s,t}$) under the conditions of [2] (Proposition 5.9.2) converge to the process $R(\mathbf{x}_s, s)$ w. p. 1 as $r \rightarrow \infty$ for all $s, t \in [0, T]$ such that $s > t$ and $T < \infty$.

6. THE TAYLOR–STRATONOVICH EXPANSION

In this section, we cite the Taylor–Stratonovich expansion [7] and introduce some necessary notations. At that, we will use the original notations introduced by the author of this paper.

Assume that $LR(\mathbf{x}_\tau, \tau)$, $G_0^{(i)} R(\mathbf{x}_\tau, \tau) \in \mathfrak{M}_2([0, T])$ for $i = 1, \dots, m$, and consider the Ito formula in the form (29).

In addition, suppose that the function $G_0^{(i)} R(\mathbf{x}, t)$ ($i = 1, \dots, m$) is such that the formulas (15) and (16) can be applied. In this case, the relations (15) and (16) imply that

$$(49) \quad \int_t^s G_0^{(i)} R(\mathbf{x}_\tau, \tau) d\mathbf{f}_\tau^{(i)} = \int_t^{*s} G_0^{(i)} R(\mathbf{x}_\tau, \tau) d\mathbf{f}_\tau^{(i)} - \frac{1}{2} \int_t^s G_0^{(i)} G_0^{(i)} R(\mathbf{x}_\tau, \tau) d\tau$$

w. p. 1 for $i = 1, \dots, m$.

Using the relation (49), let us write the formula (29) in the following form

$$(50) \quad R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \int_t^s \bar{L}R(\mathbf{x}_\tau, \tau) d\tau + \sum_{i=1}^m \int_t^{*s} G_0^{(i)} R(\mathbf{x}_\tau, \tau) d\mathbf{f}_\tau^{(i)} \quad \text{w. p. 1,}$$

where

$$(51) \quad \bar{L}R(\mathbf{x}, t) = LR(\mathbf{x}, t) - \frac{1}{2} \sum_{i=1}^m G_0^{(i)} G_0^{(i)} R(\mathbf{x}, t).$$

Introduce the following notation

$$\left\| D_{\lambda_l}^{(i_l)} \dots D_{\lambda_1}^{(i_1)} R(\mathbf{x}, t) \right\|_{i_1=\lambda_1, \dots, i_l=\lambda_l}^{m\lambda_1 \dots m\lambda_l} \stackrel{\text{def}}{=} {}^{(p_l)} D_{\lambda_l} \dots D_{\lambda_1} R(\mathbf{x}, t),$$

$${}^{(p_k)} J_{(\lambda_k \dots \lambda_1) s, t}^* = \left\| J_{(\lambda_k \dots \lambda_1) s, t}^{*(i_k \dots i_1)} \right\|_{i_1=\lambda_1, \dots, i_k=\lambda_k}^{m\lambda_1 \dots m\lambda_k},$$

$$M_k = \left\{ (\lambda_k, \dots, \lambda_1) : \lambda_l = 1 \text{ or } \lambda_l = 0; l = 1, \dots, k \right\}, \quad k \geq 1,$$

$$J_{(\lambda_k \dots \lambda_1) s, t}^{*(i_k \dots i_1)} = \int_t^{*s} \dots \int_t^{*t_2} d\mathbf{w}_{t_1}^{(i_k)} \dots d\mathbf{w}_{t_k}^{(i_1)}, \quad k \geq 1,$$

where $\lambda_l = 1$ or $\lambda_l = 0$, $D_{\lambda_l}^{(i_l)} = \bar{L}$ and $i_l = 0$ for $\lambda_l = 0$, $D_{\lambda_l}^{(i_l)} = G_0^{(i_l)}$ and $i_l = 1, \dots, m$ for $\lambda_l = 1$,

$$p_l = \sum_{j=1}^l \lambda_j \quad \text{for } l = 1, \dots, r+1, \quad r \in \mathbb{N},$$

$\mathbf{w}_\tau^{(i)}$ ($i = 1, \dots, m$) are \mathbb{F}_τ -measurable for all $\tau \in [0, T]$ independent standard Wiener processes and $\mathbf{w}_\tau^{(0)} = \tau$.

Applying the formula (50) to the process $R(\mathbf{x}_s, s)$ repeatedly, we obtain the following Taylor–Stratonovich expansion [7]

$$(52) \quad R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{k=1}^r \sum_{(\lambda_k, \dots, \lambda_1) \in M_k} {}^{(p_k)} D_{\lambda_k} \dots D_{\lambda_1} R(\mathbf{x}_t, t) \cdot {}^{(p_k)} J_{(\lambda_k \dots \lambda_1) s, t}^* + (D_{r+1})_{s, t}$$

w. p. 1, where

$$(53) \quad (D_{r+1})_{s, t} = \sum_{(\lambda_{r+1}, \dots, \lambda_1) \in M_{r+1}} \int_t^{*s} \dots \left(\int_t^{*t_2} {}^{(p_{r+1})} D_{\lambda_{r+1}} \dots D_{\lambda_1} R(\mathbf{x}_{t_1}, t_1) \cdot {}^{\lambda_{r+1}} dw_{t_1} \right) \dots \cdot {}^{\lambda_1} dw_{t_{r+1}}.$$

It is assumed that the right-hand sides of (52), (53) exist.

A possible variant of the conditions under which the right-hand sides of (52), (53) exist is as follows

$$(i^*) \quad Q_{\lambda_l}^{(i_l)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}, t) \in \mathcal{L} \text{ for all } (\lambda_l, \dots, \lambda_1) \in \bigcup_{g=1}^r M_g;$$

(ii*)

$$(54) \quad \left| Q_{\lambda_l}^{(i_l)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}, t) - Q_{\lambda_l}^{(i_l)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{y}, t) \right| \leq K |\mathbf{x} - \mathbf{y}|,$$

$$(55) \quad \left| Q_{\lambda_l}^{(i_l)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}, t) \right| \leq K(1 + |\mathbf{x}|),$$

and

$$\left| Q_{\lambda_l}^{(i_l)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}, t) - Q_{\lambda_l}^{(i_l)} \dots Q_{\lambda_1}^{(i_1)} R(\mathbf{x}, s) \right| \leq K |t - s|^\nu (1 + |\mathbf{x}|)$$

for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, $t, s \in [0, T]$, $(\lambda_l, \dots, \lambda_1) \in \bigcup_{g=1}^{r+1} M_g$ and for some $\nu > 0$, where $K < \infty$ is a constant, $Q_{\lambda_l}^{(i_l)} = L$ and $i_l = 0$ for $\lambda_l = 0$, $Q_{\lambda_l}^{(i_l)} = G_0^{(i_l)}$ and $i_l = 1, \dots, m$ for $\lambda_l = 1$;

(iii*) the functions $\mathbf{a}(\mathbf{x}, t)$ and $B(\mathbf{x}, t)$ are measurable with respect to all of the variables and satisfy the conditions (54) and (55);

(iv*) \mathbf{x}_0 is F_0 -measurable and $M \{|\mathbf{x}_0|^8\} < \infty$.

Let us write the expansion (52) in the another form

$$\begin{aligned} R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{k=1}^r \sum_{(\lambda_k, \dots, \lambda_1) \in M_k} \sum_{i_1=\lambda_1}^{m\lambda_1} \dots \sum_{i_k=\lambda_k}^{m\lambda_k} D_{\lambda_k}^{(i_k)} \dots D_{\lambda_1}^{(i_1)} R(\mathbf{x}_t, t) J_{(\lambda_k \dots \lambda_1) s, t}^{*(i_k \dots i_1)} + \\ + (D_{r+1})_{s, t} \quad \text{w. p. 1.} \end{aligned}$$

Denote

$$G_{rk} = \left\{ (\lambda_k, \dots, \lambda_1) : r+1 \leq 2k - \lambda_1 - \dots - \lambda_k \leq 2r \right\},$$

$$E_{qk} = \left\{ (\lambda_k, \dots, \lambda_1) : 2k - \lambda_1 - \dots - \lambda_k = q \right\},$$

where $\lambda_l = 1$ or $\lambda_l = 0$ ($l = 1, \dots, k$).

The Taylor–Stratonovich expansion ordered according to the order of smallness (in the mean-square sense when $s \downarrow t$) of its terms has the form

$$\begin{aligned} R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{q,k=1}^r \sum_{(\lambda_k, \dots, \lambda_1) \in E_{qk}} \sum_{i_1=\lambda_1}^{m\lambda_1} \dots \sum_{i_k=\lambda_k}^{m\lambda_k} D_{\lambda_k}^{(i_k)} \dots D_{\lambda_1}^{(i_1)} R(\mathbf{x}_t, t) J_{(\lambda_k \dots \lambda_1) s, t}^{*(i_k \dots i_1)} + \\ (56) \quad + (H_{r+1})_{s, t} \quad \text{w. p. 1,} \end{aligned}$$

where

$$\begin{aligned} (H_{r+1})_{s, t} = \sum_{k=1}^r \sum_{(\lambda_k, \dots, \lambda_1) \in G_{rk}} \sum_{i_1=\lambda_1}^{m\lambda_1} \dots \sum_{i_k=\lambda_k}^{m\lambda_k} D_{\lambda_k}^{(i_k)} \dots D_{\lambda_1}^{(i_1)} R(\mathbf{x}_t, t) J_{(\lambda_k \dots \lambda_1) s, t}^{*(i_k \dots i_1)} + \\ + (D_{r+1})_{s, t}. \end{aligned}$$

7. THE FIRST FORM OF THE UNIFIED TAYLOR–STRATONOVICH EXPANSION

In this section, we transform the right-hand side of (52) with the help of Theorem 1 and Lemma 2 to a representation including iterated Stratonovich stochastic integrals of the form (8).

Denote

$$(57) \quad I_{l_1 \dots l_{k s, t}}^{*(i_1 \dots i_k)} = \int_t^{*s} (t - t_k)^{l_k} \dots \int_t^{*t_2} (t - t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)} \quad \text{for } k \geq 1$$

and

$$I_{l_1 \dots l_{k s, t}}^{*(i_1 \dots i_k)} = 1 \quad \text{for } k = 0,$$

where $i_1, \dots, i_k = 1, \dots, m$. Moreover, let

$$(58) \quad \begin{aligned} {}^{(k)}I_{l_1 \dots l_{k s, t}}^* &= \left\| I_{l_1 \dots l_{k s, t}}^{*(i_1 \dots i_k)} \right\|_{i_1, \dots, i_k=1}^m, \\ \bar{G}_p^{(i)} &\stackrel{\text{def}}{=} \frac{1}{p} \left(\bar{G}_{p-1}^{(i)} \bar{L} - \bar{L} \bar{G}_{p-1}^{(i)} \right), \quad p = 1, 2, \dots, \quad i = 1, \dots, m, \end{aligned}$$

where $\bar{G}_0^{(i)} \stackrel{\text{def}}{=} G_0^{(i)}$, $i = 1, \dots, m$. The operators \bar{L} and $G_0^{(i)}$, $i = 1, \dots, m$, are determined by the equalities (27), (28), and (51). Denote

$$\begin{aligned} A_q &\stackrel{\text{def}}{=} \left\{ (k, j, l_1, \dots, l_k) : k + j + \sum_{p=1}^k l_p = q; k, j, l_1, \dots, l_k = 0, 1, \dots \right\}, \\ \left\| \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} \bar{L}^j R(\mathbf{x}, t) \right\|_{i_1, \dots, i_k=1}^m &\stackrel{\text{def}}{=} {}^{(k)}\bar{G}_{l_1} \dots \bar{G}_{l_k} \bar{L}^j R(\mathbf{x}, t), \\ \bar{L}^j R(\mathbf{x}, t) &\stackrel{\text{def}}{=} \begin{cases} \underbrace{\bar{L} \dots \bar{L}}_j R(\mathbf{x}, t) & \text{for } j \geq 1 \\ R(\mathbf{x}, t) & \text{for } j = 0 \end{cases}. \end{aligned}$$

Theorem 4. *Let conditions (i*)–(iv*) be satisfied. Then for any $s, t \in [0, T]$ such that $s > t$ and for any positive integer r , the following expansion takes place w. p. 1*

$$(59) \quad \begin{aligned} R(\mathbf{x}_s, s) &= R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k, j, l_1, \dots, l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} \bar{L}^j R(\mathbf{x}_t, t) I_{l_1 \dots l_{k s, t}}^{*(i_1 \dots i_k)} + \\ &\quad + (D_{r+1})_{s, t}, \end{aligned}$$

where $(D_{r+1})_{s,t}$ has the form (53).

Proof. We claim that

$$\begin{aligned}
& \sum_{(\lambda_q, \dots, \lambda_1) \in M_q} {}^{(p_q)}D_{\lambda_q} \dots D_{\lambda_1} R(\mathbf{x}_t, t) \cdot {}^{p_q} J_{(\lambda_q \dots \lambda_1) s, t}^* = \\
(60) \quad & = \sum_{(k, j, l_1, \dots, l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} \bar{L}^j R(\mathbf{x}_t, t) I_{l_1 \dots l_k s, t}^{*(i_1 \dots i_k)}
\end{aligned}$$

w. p. 1. The equality (60) is valid for $q = 1$. Assume that (60) is valid for some $q > 1$. In this case, using the induction hypothesis, we obtain

$$\begin{aligned}
& \sum_{(\lambda_{q+1}, \dots, \lambda_1) \in M_{q+1}} {}^{(p_{q+1})}D_{\lambda_1} \dots D_{\lambda_{q+1}} R(\mathbf{x}_t, t) \cdot {}^{p_{q+1}} J_{(\lambda_1 \dots \lambda_{q+1}) s, t}^* = \\
& = \sum_{\lambda_{q+1} \in \{1, 0\}} \int_t^{*s} \sum_{(\lambda_q, \dots, \lambda_1) \in M_q} \left({}^{(p_{q+1})}D_{\lambda_1} \dots D_{\lambda_{q+1}} R(\mathbf{x}_t, t) \cdot {}^{p_q} J_{(\lambda_1 \dots \lambda_q) \theta, t}^* \right)^{\lambda_{q+1}} d\mathbf{w}_\theta = \\
& = \sum_{\lambda_{q+1} \in \{1, 0\}} \int_t^{*s} \sum_{(k, j, l_1, \dots, l_k) \in A_q} \frac{(\theta-t)^j}{j!} \times \\
& \quad \times \left({}^{(k+\lambda_{q+1})} \bar{G}_{l_1} \dots \bar{G}_{l_k} \bar{L}^j D_{\lambda_{q+1}} R(\mathbf{x}_t, t) \cdot {}^{(k)} I_{l_1 \dots l_k s, t}^* \right)^{\lambda_{q+1}} d\mathbf{w}_\theta = \\
& = \sum_{(k, j, l_1, \dots, l_k) \in A_q} \left({}^{(k)} \bar{G}_{l_1} \dots \bar{G}_{l_k} \bar{L}^{j+1} R(\mathbf{x}_t, t) \cdot \int_t^s \frac{(\theta-t)^j}{j!} {}^{(k)} I_{l_1 \dots l_k \theta, t}^* d\theta + \right. \\
(61) \quad & \left. + \left({}^{(k+1)} \bar{G}_{l_1} \dots \bar{G}_{l_k} \bar{L}^j \bar{G}_0 R(\mathbf{x}_t, t) \cdot \int_t^{*s} \frac{(\theta-t)^j}{j!} {}^{(k)} I_{l_1 \dots l_k \theta, t}^* \right)^1 d\mathbf{f}_\theta \right)
\end{aligned}$$

w. p. 1.

Using Lemma 1, we obtain

$$\begin{aligned}
& \int_t^s \frac{(\theta-t)^j}{j!} {}^{(k)} I_{l_1 \dots l_k \theta, t}^* d\theta = \\
(62) \quad & = \frac{1}{(j+1)!} \begin{cases} (s-t)^{j+1} & \text{for } k = 0 \\ (s-t)^{j+1} \cdot {}^{(k)} I_{l_1 \dots l_k s, t}^* - (-1)^{j+1} \cdot {}^{(k)} I_{l_1 \dots l_{k-1} l_{k+j+1} s, t}^* & \text{for } k > 0 \end{cases}
\end{aligned}$$

w. p. 1. In addition (see (57)), we get

$$(63) \quad \int_t^{*s} \frac{(\theta-t)^j}{j!} I_{l_1 \dots l_k \theta, t}^{*(i_1 \dots i_k)} d\mathbf{f}_\theta^{(i_{k+1})} = \frac{(-1)^j}{j!} I_{l_1 \dots l_k j, t}^{*(i_1 \dots i_k i_{k+1})}$$

in the notations just introduced. Substitute the relations (62) and (63) into the formula (61). Grouping summands of the obtained expression with equal lower indices at iterated Stratonovich stochastic integrals and using (58) and the equality

$$(64) \quad \bar{G}_p^{(i)} R(\mathbf{x}, t) = \frac{1}{p!} \sum_{q=0}^p (-1)^q C_p^q \bar{L}^q \bar{G}_0^{(i)} \bar{L}^{p-q} R(\mathbf{x}, t), \quad \text{where} \quad C_p^q = \frac{p!}{q!(p-q)!},$$

(this equality follows from (58)), we note that the obtained expression is equal to

$$\sum_{(k, j, l_1, \dots, l_k) \in A_{q+1}} \frac{(s-t)^j}{j!} {}^{(k)}\bar{G}_{l_1} \dots \bar{G}_{l_k} \bar{L}^j \{\eta_t\} \cdot {}^{(k)} I_{l_1 \dots l_k, t}^*$$

w. p. 1. Summing the equalities (60) for $q = 1, 2, \dots, r$ and applying the formula (52), we obtain the expression (59). The proof is completed.

Let us order terms of the expansion (59) according to their smallness orders as $s \downarrow t$ in the mean-square sense

$$(65) \quad R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k, j, l_1, \dots, l_k) \in D_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} \bar{L}^j R(\mathbf{x}_t, t) I_{l_1 \dots l_k, t}^{*(i_1 \dots i_k)} + (H_{r+1})_{s,t} \quad \text{w. p. 1,}$$

where

$$(66) \quad (H_{r+1})_{s,t} = \sum_{(k, j, l_1, \dots, l_k) \in U_r} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} \bar{L}^j R(\mathbf{x}_t, t) I_{l_1 \dots l_k, t}^{*(i_1 \dots i_k)} + (D_{r+1})_{s,t},$$

$$D_q = \left\{ (k, j, l_1, \dots, l_k) : k + 2 \left(j + \sum_{p=1}^k l_p \right) = q; k, j, l_1, \dots, l_k = 0, 1, \dots \right\},$$

$$(67) \quad U_r = \left\{ (k, j, l_1, \dots, l_k) : k + j + \sum_{p=1}^k l_p \leq r, k + 2 \left(j + \sum_{p=1}^k l_p \right) \geq r + 1; k, j, l_1, \dots, l_k = 0, 1, \dots \right\},$$

and $(D_{r+1})_{s,t}$ has the form (53). Note that the remainder term $(H_{r+1})_{s,t}$ in (65) has a higher order of smallness in the mean-square sense as $s \downarrow t$ than the terms of the main part of expansion (65).

8. THE SECOND FORM OF THE UNIFIED TAYLOR–STRATONOVICH EXPANSION

Consider iterated Stratonovich stochastic integrals of the form

$$J_{l_1 \dots l_{k,s,t}}^{*(i_1 \dots i_k)} = \int_t^{*s} (s - t_k)^{l_k} \dots \int_t^{*t_2} (s - t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)} \quad \text{for } k \geq 1$$

and

$$J_{l_1 \dots l_{k,s,t}}^{*(i_1 \dots i_k)} = 1 \quad \text{for } k = 0,$$

where $i_1, \dots, i_k = 1, \dots, m$.

The additive property of stochastic integrals and the Newton binomial formula imply the following equality

$$(68) \quad I_{l_1 \dots l_{k,s,t}}^{*(i_1 \dots i_k)} = \sum_{j_1=0}^{l_1} \dots \sum_{j_k=0}^{l_k} \prod_{g=1}^k C_{l_g}^{j_g} (t - s)^{l_1 + \dots + l_k - j_1 - \dots - j_k} J_{j_1 \dots j_{k,s,t}}^{*(i_1 \dots i_k)} \quad \text{w. p. 1,}$$

where

$$C_l^k = \frac{l!}{k!(l-k)!}$$

is the binomial coefficient. Thus, the Taylor–Stratonovich expansion of the process $\eta_s = R(\mathbf{x}_s, s)$, $s \in [0, T]$ can be constructed either using the iterated stochastic integrals $I_{l_1 \dots l_{k,s,t}}^{*(i_1 \dots i_k)}$ similarly to the previous section or using the iterated stochastic integrals $J_{l_1 \dots l_{k,s,t}}^{*(i_1 \dots i_k)}$. This is the main subject of this section.

Denote

$$\left\| J_{l_1 \dots l_{k,s,t}}^{*(i_1 \dots i_k)} \right\|_{i_1, \dots, i_k=1}^m \stackrel{\text{def}}{=} {}^{(k)} J_{l_1 \dots l_{k,s,t}}^*,$$

$$\left\| \bar{L}^j \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} R(\mathbf{x}, t) \right\|_{i_1, \dots, i_k=1}^m \stackrel{\text{def}}{=} {}^{(k)} \bar{L}^j \bar{G}_{l_1} \dots \bar{G}_{l_k} R(\mathbf{x}, t).$$

Theorem 5. *Let conditions (i*)–(iv*) be satisfied. Then for any $s, t \in [0, T]$ such that $s > t$ and for any positive integer r , the following expansion is valid w. p. 1*

$$(69) \quad R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k,j,l_1, \dots, l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m \bar{L}^j \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} R(\mathbf{x}_t, t) J_{l_1 \dots l_{k,s,t}}^{*(i_1 \dots i_k)} +$$

$$+ (D_{r+1})_{s,t},$$

where $(D_{r+1})_{s,t}$ has the form (53).

Proof. To prove the theorem, we check the equalities

$$\begin{aligned}
& \sum_{(k,j,l_1,\dots,l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1,\dots,i_k=1}^m \bar{L}^j \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} R(\mathbf{x}_t, t) J_{l_1 \dots l_{k_s,t}}^{*(i_1 \dots i_k)} = \\
(70) \quad & \sum_{(k,j,l_1,\dots,l_k) \in A_q} \frac{(s-t)^j}{j!} \sum_{i_1,\dots,i_k=1}^m \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} \bar{L}^j R(\mathbf{x}_t, t) I_{l_1 \dots l_{k_s,t}}^{*(i_1 \dots i_k)} \quad \text{w. p. 1}
\end{aligned}$$

for $q = 1, 2, \dots, r$. To check (70), substitute the expression (68) into the right-hand side of (70) and then use the formulas (58), (64).

Let us rank terms of the expansion (69) according to their orders of smallness in the mean-square sense as $s \downarrow t$

$$\begin{aligned}
R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k,j,l_1,\dots,l_k) \in D_q} \frac{(s-t)^j}{j!} \sum_{i_1,\dots,i_k=1}^m \bar{L}^j \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} R(\mathbf{x}_t, t) J_{l_1 \dots l_{k_s,t}}^{*(i_1 \dots i_k)} + \\
+ (H_{r+1})_{s,t} \quad \text{w. p. 1,}
\end{aligned}$$

where

$$(H_{r+1})_{s,t} = \sum_{(k,j,l_1,\dots,l_k) \in U_r} \frac{(s-t)^j}{j!} \sum_{i_1,\dots,i_k=1}^m \bar{L}^j \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} R(\mathbf{x}_t, t) J_{l_1 \dots l_{k_s,t}}^{*(i_1 \dots i_k)} + (D_{r+1})_{s,t}.$$

The term $(D_{r+1})_{s,t}$ has the form (53); the terms D_q and U_r have the forms (66) and (67), respectively. Finally, we note that the convergence w. p. 1 of the truncated Taylor–Stratonovich expansion (52) (without the remainder term $(D_{r+1})_{s,t}$) to the process $R(\mathbf{x}_s, s)$ as $r \rightarrow \infty$ for all $s, t \in [0, T]$ such that $s > t$ and $T < \infty$ has been proved in [2] (Proposition 5.10.2). Since the expansions (59) and (69) are obtained from the Taylor–Stratonovich expansion (52) without any additional conditions, the truncated expansions (59) and (69) (without the remainder term $(D_{r+1})_{s,t}$) under the conditions of [2] (Proposition 5.10.2) converge to the process $R(\mathbf{x}_s, s)$ w. p. 1 as $r \rightarrow \infty$ for all $s, t \in [0, T]$ such that $s > t$ and $T < \infty$.

9. COMPARISON OF THE UNIFIED TAYLOR–ITO AND TAYLOR–STRATONOVICH EXPANSIONS WITH THE CLASSICAL TAYLOR–ITO AND TAYLOR–STRATONOVICH EXPANSIONS

Note that the truncated unified Taylor–Ito and Taylor–Stratonovich expansions contain the less number of various iterated Ito and Stratonovich stochastic integrals (moreover, their major part will have less multiplicity) in comparison with the classical Taylor–Ito and Taylor–Stratonovich expansions [7].

It is easy to notice that the stochastic integrals from the families (4), (5) are connected by linear relations. However, the stochastic integrals from the families (6), (7) cannot be connected by linear relations. This holds for the stochastic integrals from the families (8), (9). Therefore, we will call the families (6)–(9) as the *stochastic bases*.

Let us call the numbers $\text{rank}_A(r)$ and $\text{rank}_D(r)$ of various iterated Ito and Stratonovich stochastic integrals which are included in the families (6)–(9) as the *ranks of stochastic bases* when summation

in the stochastic expansions is performed using the sets A_q ($q = 1, \dots, r$) and D_q ($q = 1, \dots, r$) correspondently. Here r is a fixed natural number.

At the beginning, let us analyze several examples related to the Taylor–Ito expansions (obviously, the same conclusions will hold for the Taylor–Stratonovich expansions).

Assume that summation in the unified Taylor–Ito expansions is performed using the sets D_q ($q = 1, \dots, r$). It is easy to see that the truncated unified Taylor–Ito expansion (43), where summation is performed with respect to the sets D_q when $r = 3$ includes 4 ($\text{rank}_D(3) = 4$) various iterated Ito stochastic integrals

$$I_{0s,t}^{(i_1)}, \quad I_{00s,t}^{(i_1 i_2)}, \quad I_{1s,t}^{(i_1)}, \quad I_{000s,t}^{(i_1 i_2 i_3)}.$$

The same truncated classical Taylor–Ito expansion (34) [2] contains 5 various iterated Ito stochastic integrals

$$J_{(1)s,t}^{(i_1)}, \quad J_{(11)s,t}^{(i_1 i_2)}, \quad J_{(10)s,t}^{(i_1 0)}, \quad J_{(01)s,t}^{(0 i_1)}, \quad J_{(111)s,t}^{(i_1 i_2 i_3)}.$$

For $r = 4$ we have 7 ($\text{rank}_D(4) = 7$) integrals

$$I_{0s,t}^{(i_1)}, \quad I_{00s,t}^{(i_1 i_2)}, \quad I_{1s,t}^{(i_1)}, \quad I_{000s,t}^{(i_1 i_2 i_3)}, \quad I_{01s,t}^{(i_1 i_2)}, \quad I_{10s,t}^{(i_1 i_2)}, \quad I_{0000s,t}^{(i_1 i_2 i_3 i_4)}$$

against 9 stochastic integrals

$$J_{(1)s,t}^{(i_1)}, \quad J_{(11)s,t}^{(i_1 i_2)}, \quad J_{(10)s,t}^{(i_1 0)}, \quad J_{(01)s,t}^{(0 i_1)}, \quad J_{(111)s,t}^{(i_1 i_2 i_3)}, \quad J_{(101)s,t}^{(i_1 0 i_3)}, \quad J_{(110)s,t}^{(i_1 i_2 0)}, \quad J_{(011)s,t}^{(0 i_1 i_2)}, \quad J_{(1111)s,t}^{(i_1 i_2 i_3 i_4)}.$$

For $r = 5$ ($\text{rank}_D(5) = 12$) we get 12 integrals against 17 integrals and for $r = 6$ and $r = 7$ we have 20 against 29 and 33 against 50 correspondently.

We will obtain the same results when compare the unified Taylor–Stratonovich expansions [9], [46]–[50], [54]–[61] with their classical analogues [2], [7] (see previous sections).

Note that summation according to the sets D_q is usually used while constructing strong numerical methods (built according to the mean-square criterion of convergence) for Ito SDEs [2], [4], [46]–[50], [56]–[61]. Summation according to the sets A_q is usually used when building weak numerical methods (built in accordance with the weak criterion of convergence) for Ito SDEs [2], [4]. For example, $\text{rank}_A(4) = 15$, while the total number of various iterated Ito stochastic integrals (included in the classical Taylor–Ito expansion [2] when $r = 4$) equals to 26.

Let us show that [48]–[50], [56]–[61]

$$\text{rank}_A(r) = 2^r - 1.$$

Let (l_1, \dots, l_k) be an ordered set such that $l_1, \dots, l_k = 0, 1, \dots$ and $k = 1, 2, \dots$. Consider $S(k) \stackrel{\text{def}}{=} l_1 + \dots + l_k = p$ (p is a fixed natural number or zero). Let $N(k, p)$ be a number of all ordered combinations (l_1, \dots, l_k) such that $l_1, \dots, l_k = 0, 1, \dots$, $k = 1, 2, \dots$, and $S(k) = p$. First let us show that

$$N(k, p) = C_{p+k-1}^{k-1},$$

where

$$C_n^m = \frac{n!}{m!(n-m)!}$$

is a binomial coefficient.

It is not difficult to see that

$$N(1, p) = 1 = C_{p+1-1}^{1-1},$$

$$N(2, p) = p + 1 = C_{p+2-1}^{2-1},$$

$$N(3, p) = \frac{(p+1)(p+2)}{2} = C_{p+3-1}^{3-1}.$$

Moreover,

$$N(k+1, p) = \sum_{l=0}^p N(k, l) = \sum_{l=0}^p C_{l+k-1}^{k-1} = C_{p+k}^k,$$

where we used the induction assumption and the well known property of binomial coefficients.

Then

$$\begin{aligned} \text{rank}_A(r) &= \\ &= N(1, 0) + (N(1, 1) + N(2, 0)) + (N(1, 2) + N(2, 1) + N(3, 0)) + \dots \\ &\quad \dots + (N(1, r-1) + N(2, r-2) + \dots + N(r, 0)) = \\ &= C_0^0 + (C_1^0 + C_1^1) + (C_2^0 + C_2^1 + C_2^2) + \dots \\ &\quad \dots + (C_{r-1}^0 + C_{r-1}^1 + C_{r-1}^2 + \dots + C_{r-1}^{r-1}) = \\ &= 2^0 + 2^1 + 2^2 + \dots + 2^{r-1} = 2^r - 1. \end{aligned}$$

Let $n_M(r)$ be the total number of various iterated stochastic integrals included in the classical Taylor–Ito expansion (32) [2], where summation is performed with respect to the set

$$\bigcup_{k=1}^r M_k.$$

If we exclude from the consideration the integrals which are equal to $(s-t)^j/j!$, then

$$\begin{aligned} n_M(r) &= \\ &= (2^1 - 1) + (2^2 - 1) + (2^3 - 1) + \dots + (2^r - 1) = \\ &= 2(1 + 2 + 2^2 + \dots + 2^{r-1}) - r = 2(2^r - 1) - r. \end{aligned}$$

It means that

$$\lim_{r \rightarrow \infty} \frac{n_M(r)}{\text{rank}_A(r)} = 2.$$

In Table 1 we can see the numbers

$$\text{rank}_A(r), \quad n_M(r), \quad f(r) = n_M(r)/\text{rank}_A(r)$$

for various values r .

Let us show that [48]-[50], [56]-[61]

ТАБЛИЦА 1. Numbers $\text{rank}_A(r)$, $n_M(r)$, $f(r) = n_M(r)/\text{rank}_A(r)$

r	1	2	3	4	5	6	7	8	9	10
$\text{rank}_A(r)$	1	3	7	15	31	63	127	255	511	1023
$n_M(r)$	1	4	11	26	57	120	247	502	1013	2036
$f(r)$	1	1.3333	1.5714	1.7333	1.8387	1.9048	1.9449	1.9686	1.9824	1.9902

$$(71) \quad \text{rank}_D(r) = \begin{cases} \sum_{s=0}^{r-1} \sum_{l=s}^{(r-1)/2+[s/2]} C_l^s & \text{for } r = 1, 3, 5, \dots \\ \sum_{s=0}^{r-1} \sum_{l=s}^{r/2-1+[(s+1)/2]} C_l^s & \text{for } r = 2, 4, 6, \dots \end{cases},$$

where $[x]$ is an integer part of a number x , and C_n^m is a binomial coefficient.

For proving (71) we write the condition

$$k + 2(j + S(k)) \leq r,$$

where $S(k) \stackrel{\text{def}}{=} l_1 + \dots + l_k$ ($k, j, l_1, \dots, l_k = 0, 1, \dots$) in the form $j + S(k) \leq (r - k)/2$, and perform the consideration of all possible combinations with respect to $k = 1, \dots, r$. Moreover, we take into account the above reasoning.

Let us calculate the number $n_E(r)$ of all different iterated Ito stochastic integrals from the classical Taylor–Ito expansion (34) [2] if the summation in this expansion is performed with respect to the set

$$\bigcup_{q,k=1}^r E_{qk}.$$

The summation condition can be written in this case in the form: $0 \leq p + 2q \leq r$, where q is a total number of integrations with respect to time while p is a total number of integrations with respect to the Wiener processes in the selected iterated stochastic integral from the Taylor–Ito expansion (34) [2]. At that, the multiplicity of the mentioned stochastic integral equals to $p + q$ and it is not more than r . Let us write the above condition ($0 \leq p + 2q \leq r$) in the form: $0 \leq q \leq (r - p)/2 \Leftrightarrow 0 \leq q \leq [(r - p)/2]$, where $[x]$ means an integer part of a real number x . Then, performing the consideration of all possible combinations with respect to $p = 1, \dots, r$ and using the combinatorial reasoning, we obtain the formula

$$(72) \quad n_E(r) = \sum_{s=1}^r \sum_{l=0}^{[(r-s)/2]} C_{[(r-s)/2]+s-l}^s,$$

where $[x]$ means an integer part of a real number x .

In Table 2 we can see the numbers

$$\text{rank}_D(r), \quad n_E(r), \quad g(r) = n_E(r)/\text{rank}_D(r)$$

for various values r .

ТАБЛИЦА 2. Numbers $\text{rank}_{\mathbb{D}}(r)$, $n_{\mathbb{E}}(r)$, $g(r) = n_{\mathbb{E}}(r)/\text{rank}_{\mathbb{D}}(r)$

r	1	2	3	4	5	6	7	8	9	10
$\text{rank}_{\mathbb{D}}(r)$	1	2	4	7	12	20	33	54	88	143
$n_{\mathbb{E}}(r)$	1	2	5	9	17	29	50	83	138	261
$g(r)$	1	1	1.2500	1.2857	1.4167	1.4500	1.5152	1.5370	1.5682	1.8252

10. APPLICATION OF FIRST FORM OF THE UNIFIED TAYLOR–ITO EXPANSION TO THE HIGH-ORDER STRONG NUMERICAL METHODS FOR ITO SDES

Let us write (43) for all $s, t \in [0, T]$ such that $s > t$ in the following form

$$(73) \quad R(\mathbf{x}_s, s) = R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k,j,l_1,\dots,l_k) \in \mathbb{D}_q} \frac{(s-t)^j}{j!} \sum_{i_1,\dots,i_k=1}^m G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} L^j R(\mathbf{x}_t, t) I_{l_1 \dots l_k s, t}^{(i_1 \dots i_k)} + \\ + \mathbf{1}_{\{r=2d-1, d \in \mathbb{N}\}} \frac{(s-t)^{(r+1)/2}}{((r+1)/2)!} L^{(r+1)/2} R(\mathbf{x}_t, t) + (\bar{H}_{r+1})_{s,t} \quad \text{w. p. 1,}$$

where

$$(\bar{H}_{r+1})_{s,t} = (H_{r+1})_{s,t} - \mathbf{1}_{\{r=2d-1, d \in \mathbb{N}\}} \frac{(s-t)^{(r+1)/2}}{((r+1)/2)!} L^{(r+1)/2} R(\mathbf{x}_t, t).$$

Consider the partition $\{\tau_p\}_{p=0}^N$ of the interval $[0, T]$ such that

$$0 = \tau_0 < \tau_1 < \dots < \tau_N = T, \quad \Delta_N = \max_{0 \leq j \leq N-1} |\tau_{j+1} - \tau_j|.$$

From (73) for $s = \tau_{p+1}$, $t = \tau_p$ we obtain the following representation of explicit one-step strong numerical scheme for Ito SDE (1), which is based on the first form of the unified Taylor–Ito expansion

$$(74) \quad \mathbf{y}_{p+1} = \mathbf{y}_p + \sum_{q=1}^r \sum_{(k,j,l_1,\dots,l_k) \in \mathbb{D}_q} \frac{(\tau_{p+1} - \tau_p)^j}{j!} \sum_{i_1,\dots,i_k=1}^m G_{l_1}^{(i_1)} \dots G_{l_k}^{(i_k)} L^j \mathbf{y}_p \hat{I}_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{(i_1 \dots i_k)} + \\ + \mathbf{1}_{\{r=2d-1, d \in \mathbb{N}\}} \frac{(\tau_{p+1} - \tau_p)^{(r+1)/2}}{((r+1)/2)!} L^{(r+1)/2} \mathbf{y}_p,$$

where $\hat{I}_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{(i_1 \dots i_k)}$ is an approximation of the iterated Ito stochastic integral $I_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{(i_1 \dots i_k)}$ defined as

$$I_{l_1 \dots l_k s, t}^{(i_1 \dots i_k)} = \int_t^s (t - t_k)^{l_k} \dots \int_t^{t_2} (t - t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)}.$$

Note that we understand the equality (74) componentwise with respect to the components $\mathbf{y}_p^{(i)}$ of the column \mathbf{y}_p . Also for simplicity we put $\tau_p = p\Delta$, $\Delta = T/N$, $T = \tau_N$, $p = 0, 1, \dots, N$.

It is known [2] that under the appropriate conditions the numerical scheme (74) has strong order of convergence $r/2$ ($r \in \mathbb{N}$).

Let $B_j(\mathbf{x}, t)$ is the j -th column of the matrix function $B(\mathbf{x}, t)$.

Below we consider particular cases of the numerical scheme (74) for $r = 2, 3, 4, 5$, and 6 , i.e. explicit one-step strong numerical schemes for Ito SDE (1) with orders $1.0, 1.5, 2.0, 2.5$, and 3.0 of convergence. At that, for simplicity we will write \mathbf{a} , $L\mathbf{a}$, B_i , $G_0^{(i)}B_j$ etc. instead of $\mathbf{a}(\mathbf{y}_p, \tau_p)$, $L\mathbf{a}(\mathbf{y}_p, \tau_p)$, $B_i(\mathbf{y}_p, \tau_p)$, $G_0^{(i)}B_j(\mathbf{y}_p, \tau_p)$ etc. correspondingly. Moreover, the operators L and $G_0^{(i)}$, $i = 1, \dots, m$, are determined by the equalities (27), (28) as before.

Scheme with strong order 1.0

$$(75) \quad \mathbf{y}_{p+1} = \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1} \hat{I}_{0\tau_{p+1}, \tau_p}^{(i_1)} + \Delta \mathbf{a} + \sum_{i_1, i_2=1}^m G_0^{(i_2)} B_{i_1} \hat{I}_{00\tau_{p+1}, \tau_p}^{(i_2 i_1)}.$$

Scheme with strong order 1.5

$$(76) \quad \begin{aligned} \mathbf{y}_{p+1} = & \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1} \hat{I}_{0\tau_{p+1}, \tau_p}^{(i_1)} + \Delta \mathbf{a} + \sum_{i_1, i_2=1}^m G_0^{(i_2)} B_{i_1} \hat{I}_{00\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \\ & + \sum_{i_1=1}^m \left[G_0^{(i_1)} \mathbf{a} \left(\Delta \hat{I}_{0\tau_{p+1}, \tau_p}^{(i_1)} + \hat{I}_{1\tau_{p+1}, \tau_p}^{(i_1)} \right) - LB_{i_1} \hat{I}_{1\tau_{p+1}, \tau_p}^{(i_1)} \right] + \\ & + \sum_{i_1, i_2, i_3=1}^m G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{000\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} + \\ & + \frac{\Delta^2}{2} L\mathbf{a}. \end{aligned}$$

Scheme with strong order 2.0

$$\begin{aligned} \mathbf{y}_{p+1} = & \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1} \hat{I}_{0\tau_{p+1}, \tau_p}^{(i_1)} + \Delta \mathbf{a} + \sum_{i_1, i_2=1}^m G_0^{(i_2)} B_{i_1} \hat{I}_{00\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \\ & + \sum_{i_1=1}^m \left[G_0^{(i_1)} \mathbf{a} \left(\Delta \hat{I}_{0\tau_{p+1}, \tau_p}^{(i_1)} + \hat{I}_{1\tau_{p+1}, \tau_p}^{(i_1)} \right) - LB_{i_1} \hat{I}_{1\tau_{p+1}, \tau_p}^{(i_1)} \right] + \\ & + \sum_{i_1, i_2, i_3=1}^m G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{000\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} + \frac{\Delta^2}{2} L\mathbf{a} + \end{aligned}$$

$$\begin{aligned}
& + \sum_{i_1, i_2=1}^m \left[G_0^{(i_2)} L B_{i_1} \left(\hat{I}_{10\tau_{p+1}, \tau_p}^{(i_2 i_1)} - \hat{I}_{01\tau_{p+1}, \tau_p}^{(i_2 i_1)} \right) - L G_0^{(i_2)} B_{i_1} \hat{I}_{10\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \right. \\
& \quad \left. + G_0^{(i_2)} G_0^{(i_1)} \mathbf{a} \left(\hat{I}_{01\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \Delta \hat{I}_{00\tau_{p+1}, \tau_p}^{(i_2 i_1)} \right) \right] + \\
(77) \quad & + \sum_{i_1, i_2, i_3, i_4=1}^m G_0^{(i_4)} G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{0000\tau_{p+1}, \tau_p}^{(i_4 i_3 i_2 i_1)}.
\end{aligned}$$

Scheme with strong order 2.5

$$\begin{aligned}
\mathbf{y}_{p+1} = \mathbf{y}_p & + \sum_{i_1=1}^m B_{i_1} \hat{I}_{0\tau_{p+1}, \tau_p}^{(i_1)} + \Delta \mathbf{a} + \sum_{i_1, i_2=1}^m G_0^{(i_2)} B_{i_1} \hat{I}_{00\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \\
& + \sum_{i_1=1}^m \left[G_0^{(i_1)} \mathbf{a} \left(\Delta \hat{I}_{0\tau_{p+1}, \tau_p}^{(i_1)} + \hat{I}_{1\tau_{p+1}, \tau_p}^{(i_1)} \right) - L B_{i_1} \hat{I}_{1\tau_{p+1}, \tau_p}^{(i_1)} \right] + \\
& + \sum_{i_1, i_2, i_3=1}^m G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{000\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} + \frac{\Delta^2}{2} L \mathbf{a} + \\
& + \sum_{i_1, i_2=1}^m \left[G_0^{(i_2)} L B_{i_1} \left(\hat{I}_{10\tau_{p+1}, \tau_p}^{(i_2 i_1)} - \hat{I}_{01\tau_{p+1}, \tau_p}^{(i_2 i_1)} \right) - L G_0^{(i_2)} B_{i_1} \hat{I}_{10\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \right. \\
& \quad \left. + G_0^{(i_2)} G_0^{(i_1)} \mathbf{a} \left(\hat{I}_{01\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \Delta \hat{I}_{00\tau_{p+1}, \tau_p}^{(i_2 i_1)} \right) \right] + \\
& + \sum_{i_1, i_2, i_3, i_4=1}^m G_0^{(i_4)} G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{0000\tau_{p+1}, \tau_p}^{(i_4 i_3 i_2 i_1)} + \\
& + \sum_{i_1=1}^m \left[G_0^{(i_1)} L \mathbf{a} \left(\frac{1}{2} \hat{I}_{2\tau_{p+1}, \tau_p}^{(i_1)} + \Delta \hat{I}_{1\tau_{p+1}, \tau_p}^{(i_1)} + \frac{\Delta^2}{2} \hat{I}_{0\tau_{p+1}, \tau_p}^{(i_1)} \right) + \right. \\
& \quad \left. + \frac{1}{2} L L B_{i_1} \hat{I}_{2\tau_{p+1}, \tau_p}^{(i_1)} - L G_0^{(i_1)} \mathbf{a} \left(\hat{I}_{2\tau_{p+1}, \tau_p}^{(i_1)} + \Delta \hat{I}_{1\tau_{p+1}, \tau_p}^{(i_1)} \right) \right] + \\
& + \sum_{i_1, i_2, i_3=1}^m \left[G_0^{(i_3)} L G_0^{(i_2)} B_{i_1} \left(\hat{I}_{100\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} - \hat{I}_{010\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} \right) + \right. \\
& \quad \left. + G_0^{(i_3)} G_0^{(i_2)} L B_{i_1} \left(\hat{I}_{010\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} - \hat{I}_{001\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} \right) + \right. \\
& \quad \left. + G_0^{(i_3)} G_0^{(i_2)} G_0^{(i_1)} \mathbf{a} \left(\Delta \hat{I}_{000\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} + \hat{I}_{001\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} \right) - \right.
\end{aligned}$$

$$\begin{aligned}
& -LG_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{100\tau_{p+1},\tau_p}^{(i_3i_2i_1)}] + \\
& + \sum_{i_1,i_2,i_3,i_4,i_5=1}^m G_0^{(i_5)}G_0^{(i_4)}G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{00000\tau_{p+1},\tau_p}^{(i_5i_4i_3i_2i_1)} + \\
(78) \quad & + \frac{\Delta^3}{6}LLa.
\end{aligned}$$

Scheme with strong order 3.0

$$\begin{aligned}
\mathbf{y}_{p+1} &= \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1}\hat{I}_{0\tau_{p+1},\tau_p}^{(i_1)} + \Delta\mathbf{a} + \sum_{i_1,i_2=1}^m G_0^{(i_2)}B_{i_1}\hat{I}_{00\tau_{p+1},\tau_p}^{(i_2i_1)} + \\
& + \sum_{i_1=1}^m \left[G_0^{(i_1)}\mathbf{a} \left(\Delta\hat{I}_{0\tau_{p+1},\tau_p}^{(i_1)} + \hat{I}_{1\tau_{p+1},\tau_p}^{(i_1)} \right) - LB_{i_1}\hat{I}_{1\tau_{p+1},\tau_p}^{(i_1)} \right] + \\
& + \sum_{i_1,i_2,i_3=1}^m G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{000\tau_{p+1},\tau_p}^{(i_3i_2i_1)} + \frac{\Delta^2}{2}La + \\
& + \sum_{i_1,i_2=1}^m \left[G_0^{(i_2)}LB_{i_1} \left(\hat{I}_{10\tau_{p+1},\tau_p}^{(i_2i_1)} - \hat{I}_{01\tau_{p+1},\tau_p}^{(i_2i_1)} \right) - LG_0^{(i_2)}B_{i_1}\hat{I}_{10\tau_{p+1},\tau_p}^{(i_2i_1)} + \right. \\
& \quad \left. + G_0^{(i_2)}G_0^{(i_1)}\mathbf{a} \left(\hat{I}_{01\tau_{p+1},\tau_p}^{(i_2i_1)} + \Delta\hat{I}_{00\tau_{p+1},\tau_p}^{(i_2i_1)} \right) \right] + \\
(79) \quad & + \sum_{i_1,i_2,i_3,i_4=1}^m G_0^{(i_4)}G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{0000\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)} + \mathbf{q}_{p+1,p} + \mathbf{r}_{p+1,p},
\end{aligned}$$

where

$$\begin{aligned}
\mathbf{q}_{p+1,p} &= \sum_{i_1=1}^m \left[G_0^{(i_1)}La \left(\frac{1}{2}\hat{I}_{2\tau_{p+1},\tau_p}^{(i_1)} + \Delta\hat{I}_{1\tau_{p+1},\tau_p}^{(i_1)} + \frac{\Delta^2}{2}\hat{I}_{0\tau_{p+1},\tau_p}^{(i_1)} \right) + \right. \\
& \quad \left. + \frac{1}{2}LLB_{i_1}\hat{I}_{2\tau_{p+1},\tau_p}^{(i_1)} - LG_0^{(i_1)}\mathbf{a} \left(\hat{I}_{2\tau_{p+1},\tau_p}^{(i_1)} + \Delta\hat{I}_{1\tau_{p+1},\tau_p}^{(i_1)} \right) \right] + \\
& + \sum_{i_1,i_2,i_3=1}^m \left[G_0^{(i_3)}LG_0^{(i_2)}B_{i_1} \left(\hat{I}_{100\tau_{p+1},\tau_p}^{(i_3i_2i_1)} - \hat{I}_{010\tau_{p+1},\tau_p}^{(i_3i_2i_1)} \right) + \right. \\
& \quad \left. + G_0^{(i_3)}G_0^{(i_2)}LB_{i_1} \left(\hat{I}_{010\tau_{p+1},\tau_p}^{(i_3i_2i_1)} - \hat{I}_{001\tau_{p+1},\tau_p}^{(i_3i_2i_1)} \right) + \right.
\end{aligned}$$

$$\begin{aligned}
& +G_0^{(i_3)}G_0^{(i_2)}G_0^{(i_1)}\mathbf{a}\left(\Delta\hat{I}_{000\tau_{p+1},\tau_p}^{(i_3i_2i_1)} + \hat{I}_{001\tau_{p+1},\tau_p}^{(i_3i_2i_1)}\right) - \\
& \quad -LG_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{100\tau_{p+1},\tau_p}^{(i_3i_2i_1)}\Big] + \\
& + \sum_{i_1,i_2,i_3,i_4,i_5=1}^m G_0^{(i_5)}G_0^{(i_4)}G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{00000\tau_{p+1},\tau_p}^{(i_5i_4i_3i_2i_1)} + \\
& \quad + \frac{\Delta^3}{6}LL\mathbf{a},
\end{aligned}$$

and

$$\begin{aligned}
\mathbf{r}_{p+1,p} &= \sum_{i_1,i_2=1}^m \left[G_0^{(i_2)}G_0^{(i_1)}L\mathbf{a}\left(\frac{1}{2}\hat{I}_{02\tau_{p+1},\tau_p}^{(i_2i_1)} + \Delta\hat{I}_{01\tau_{p+1},\tau_p}^{(i_2i_1)} + \frac{\Delta^2}{2}\hat{I}_{00\tau_{p+1},\tau_p}^{(i_2i_1)}\right) + \right. \\
& \quad \left. + \frac{1}{2}LLG_0^{(i_2)}B_{i_1}\hat{I}_{20\tau_{p+1},\tau_p}^{(i_2i_1)} \right. \\
& \quad \left. + G_0^{(i_2)}LG_0^{(i_1)}\mathbf{a}\left(\hat{I}_{11\tau_{p+1},\tau_p}^{(i_2i_1)} - \hat{I}_{02\tau_{p+1},\tau_p}^{(i_2i_1)} + \Delta\left(\hat{I}_{10\tau_{p+1},\tau_p}^{(i_2i_1)} - \hat{I}_{01\tau_{p+1},\tau_p}^{(i_2i_1)}\right)\right) + \right. \\
& \quad \left. + LG_0^{(i_2)}LB_{i_1}\left(\hat{I}_{11\tau_{p+1},\tau_p}^{(i_2i_1)} - \hat{I}_{20\tau_{p+1},\tau_p}^{(i_2i_1)}\right) + \right. \\
& \quad \left. + G_0^{(i_2)}LLB_{i_1}\left(\frac{1}{2}\hat{I}_{02\tau_{p+1},\tau_p}^{(i_2i_1)} + \frac{1}{2}\hat{I}_{20\tau_{p+1},\tau_p}^{(i_2i_1)} - \hat{I}_{11\tau_{p+1},\tau_p}^{(i_2i_1)}\right) - \right. \\
& \quad \left. - LG_0^{(i_2)}G_0^{(i_1)}\mathbf{a}\left(\Delta\hat{I}_{10\tau_{p+1},\tau_p}^{(i_2i_1)} + \hat{I}_{11\tau_{p+1},\tau_p}^{(i_2i_1)}\right)\right] + \\
& + \sum_{i_1,i_2,i_3,i_4=1}^m \left[G_0^{(i_4)}G_0^{(i_3)}G_0^{(i_2)}G_0^{(i_1)}\mathbf{a}\left(\Delta\hat{I}_{0000\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)} + \hat{I}_{0001\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)}\right) + \right. \\
& \quad \left. + G_0^{(i_4)}G_0^{(i_3)}LG_0^{(i_2)}B_{i_1}\left(\hat{I}_{0100\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)} - \hat{I}_{0010\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)}\right) - \right. \\
& \quad \left. - LG_0^{(i_4)}G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{1000\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)} + \right. \\
& \quad \left. + G_0^{(i_4)}LG_0^{(i_3)}G_0^{(i_2)}B_{i_1}\left(\hat{I}_{1000\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)} - \hat{I}_{0100\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)}\right) + \right. \\
& \quad \left. + G_0^{(i_4)}G_0^{(i_3)}G_0^{(i_2)}LB_{i_1}\left(\hat{I}_{0010\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)} - \hat{I}_{0001\tau_{p+1},\tau_p}^{(i_4i_3i_2i_1)}\right)\right] + \\
& + \sum_{i_1,i_2,i_3,i_4,i_5,i_6=1}^m G_0^{(i_6)}G_0^{(i_5)}G_0^{(i_4)}G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{000000\tau_{p+1},\tau_p}^{(i_6i_5i_4i_3i_2i_1)}.
\end{aligned}$$

It is well known [2] that under the standard conditions the numerical schemes (75)–(79) have strong orders of convergence 1.0, 1.5, 2.0, 2.5, and 3.0 correspondingly. Among these conditions we consider only the condition for approximations of iterated Ito stochastic integrals from the numerical schemes (75)–(79) [2], [57]–[61]

$$\mathbb{M} \left\{ \left(I_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{(i_1 \dots i_k)} - \hat{I}_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{(i_1 \dots i_k)} \right)^2 \right\} \leq C \Delta^{r+1},$$

where $r/2$ are strong orders of convergence for the numerical schemes (75)–(79), i.e. $r/2 = 1.0, 1.5, 2.0, 2.5,$ and 3.0 . Moreover, constant C does not depends on Δ .

As we mentioned above, the numerical schemes (75)–(79) are unrealizable in practice without procedures for the numerical simulation of iterated Ito stochastic integrals from (73). In Sect. 12, we give a brief overview to the effective method of the mean-square approximation of iterated Ito and Stratonovich stochastic integrals of arbitrary multiplicity k ($k \in \mathbb{N}$).

11. APPLICATION OF FIRST FORM OF THE UNIFIED TAYLOR–STRATONOVICH EXPANSION TO THE HIGH-ORDER STRONG NUMERICAL METHODS FOR ITO SDES

Let us write (65) for all $s, t \in [0, T]$ such that $s > t$ in the following from

$$\begin{aligned} R(\mathbf{x}_s, s) &= R(\mathbf{x}_t, t) + \sum_{q=1}^r \sum_{(k, j, l_1, \dots, l_k) \in \mathbb{D}_q} \frac{(s-t)^j}{j!} \sum_{i_1, \dots, i_k=1}^m \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} \bar{L}^j R(\mathbf{x}_t, t) I_{l_1 \dots l_k s, t}^{*(i_1 \dots i_k)} + \\ (80) \quad &+ \mathbf{1}_{\{r=2d-1, d \in \mathbb{N}\}} \frac{(s-t)^{(r+1)/2}}{((r+1)/2)!} L^{(r+1)/2} R(\mathbf{x}_t, t) + (\bar{H}_{r+1})_{s, t} \quad \text{w. p. 1,} \end{aligned}$$

where

$$(\bar{H}_{r+1})_{s, t} = (H_{r+1})_{s, t} - \mathbf{1}_{\{r=2d-1, d \in \mathbb{N}\}} \frac{(s-t)^{(r+1)/2}}{((r+1)/2)!} L^{(r+1)/2} R(\mathbf{x}_t, t).$$

Consider the partition $\{\tau_p\}_{p=0}^N$ of the interval $[0, T]$ such that

$$0 = \tau_0 < \tau_1 < \dots < \tau_N = T, \quad \Delta_N = \max_{0 \leq j \leq N-1} |\tau_{j+1} - \tau_j|.$$

From (80) for $s = \tau_{p+1}$, $t = \tau_p$ we obtain the following representation of explicit one-step strong numerical scheme for Ito SDE (1), which is based on the first form of the unified Taylor–Stratonovich expansion

$$\begin{aligned} \mathbf{y}_{p+1} &= \mathbf{y}_p + \sum_{q=1}^r \sum_{(k, j, l_1, \dots, l_k) \in \mathbb{D}_q} \frac{(\tau_{p+1} - \tau_p)^j}{j!} \sum_{i_1, \dots, i_k=1}^m \bar{G}_{l_1}^{(i_1)} \dots \bar{G}_{l_k}^{(i_k)} \bar{L}^j \mathbf{y}_p \hat{I}_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{*(i_1 \dots i_k)} + \\ (81) \quad &+ \mathbf{1}_{\{r=2d-1, d \in \mathbb{N}\}} \frac{(\tau_{p+1} - \tau_p)^{(r+1)/2}}{((r+1)/2)!} L^{(r+1)/2} \mathbf{y}_p, \end{aligned}$$

where $\hat{I}_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{*(i_1 \dots i_k)}$ is an approximation of the iterated Stratonovich stochastic integral $I_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{*(i_1 \dots i_k)}$ defined as

$$I_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{*(i_1 \dots i_k)} = \int_t^{*s} (t - t_k)^{l_k} \dots \int_t^{*t_2} (t - t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)}.$$

Note that we understand the equality (81) componentwise with respect to the components $\mathbf{y}_p^{(i)}$ of the column \mathbf{y}_p . Also for simplicity we put $\tau_p = p\Delta$, $\Delta = T/N$, $T = \tau_N$, $p = 0, 1, \dots, N$.

It is known [2] that under the appropriate conditions the numerical scheme (81) has strong order of convergence $r/2$ ($r \in \mathbb{N}$).

Denote

$$\bar{\mathbf{a}}(\mathbf{x}, t) = \mathbf{a}(\mathbf{x}, t) - \frac{1}{2} \sum_{j=1}^m G_0^{(j)} B_j(\mathbf{x}, t),$$

where $B_j(\mathbf{x}, t)$ is the j -th column of the matrix function $B(\mathbf{x}, t)$.

Below we consider particular cases of the numerical scheme (81) for $r = 2, 3, 4, 5$, and 6, i.e. explicit one-step strong numerical schemes for Ito SDE (1) with orders 1.0, 1.5, 2.0, 2.5, and 3.0 of convergence. At that for simplicity we will write $\bar{\mathbf{a}}$, $\bar{L}\bar{\mathbf{a}}$, $L\mathbf{a}$, B_i , $G_0^{(i)} B_j$ etc. instead of $\bar{\mathbf{a}}(\mathbf{y}_p, \tau_p)$, $\bar{L}\bar{\mathbf{a}}(\mathbf{y}_p, \tau_p)$, $L\mathbf{a}(\mathbf{y}_p, \tau_p)$, $B_i(\mathbf{y}_p, \tau_p)$, $G_0^{(i)} B_j(\mathbf{y}_p, \tau_p)$ etc. correspondingly. Moreover, the operators \bar{L} and $G_0^{(i)}$, $i = 1, \dots, m$, are determined by the equalities (27), (28), and (51) as before.

Scheme with strong order 1.0

$$(82) \quad \mathbf{y}_{p+1} = \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1} \hat{I}_{0\tau_{p+1}, \tau_p}^{*(i_1)} + \Delta \bar{\mathbf{a}} + \sum_{i_1, i_2=1}^m G_0^{(i_2)} B_{i_1} \hat{I}_{00\tau_{p+1}, \tau_p}^{*(i_2 i_1)}.$$

Scheme with strong order 1.5

$$(83) \quad \begin{aligned} \mathbf{y}_{p+1} = & \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1} \hat{I}_{0\tau_{p+1}, \tau_p}^{*(i_1)} + \Delta \bar{\mathbf{a}} + \sum_{i_1, i_2=1}^m G_0^{(i_2)} B_{i_1} \hat{I}_{00\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \\ & + \sum_{i_1=1}^m \left[G_0^{(i_1)} \bar{\mathbf{a}} \left(\Delta \hat{I}_{0\tau_{p+1}, \tau_p}^{*(i_1)} + \hat{I}_{1\tau_{p+1}, \tau_p}^{*(i_1)} \right) - \bar{L} B_{i_1} \hat{I}_{1\tau_{p+1}, \tau_p}^{*(i_1)} \right] + \\ & + \sum_{i_1, i_2, i_3=1}^m G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{000\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} + \frac{\Delta^2}{2} L\mathbf{a}. \end{aligned}$$

Scheme with strong order 2.0

$$\mathbf{y}_{p+1} = \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1} \hat{I}_{0\tau_{p+1}, \tau_p}^{*(i_1)} + \Delta \bar{\mathbf{a}} + \sum_{i_1, i_2=1}^m G_0^{(i_2)} B_{i_1} \hat{I}_{00\tau_{p+1}, \tau_p}^{*(i_2 i_1)} +$$

$$\begin{aligned}
& + \sum_{i_1=1}^m \left[G_0^{(i_1)} \bar{\mathbf{a}} \left(\Delta \hat{I}_{0\tau_{p+1}, \tau_p}^{*(i_1)} + \hat{I}_{1\tau_{p+1}, \tau_p}^{*(i_1)} \right) - \bar{L} B_{i_1} \hat{I}_{1\tau_{p+1}, \tau_p}^{*(i_1)} \right] + \\
& \quad + \sum_{i_1, i_2, i_3=1}^m G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{000\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} + \frac{\Delta^2}{2} \bar{L} \bar{\mathbf{a}} + \\
& + \sum_{i_1, i_2=1}^m \left[G_0^{(i_2)} \bar{L} B_{i_1} \left(\hat{I}_{10\tau_{p+1}, \tau_p}^{*(i_2 i_1)} - \hat{I}_{01\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \right) - \bar{L} G_0^{(i_2)} B_{i_1} \hat{I}_{10\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \right. \\
& \quad \left. + G_0^{(i_2)} G_0^{(i_1)} \bar{\mathbf{a}} \left(\hat{I}_{01\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \Delta \hat{I}_{00\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \right) \right] + \\
(84) \quad & \quad + \sum_{i_1, i_2, i_3, i_4=1}^m G_0^{(i_4)} G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{0000\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)}.
\end{aligned}$$

Scheme with strong order 2.5

$$\begin{aligned}
\mathbf{y}_{p+1} & = \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1} \hat{I}_{0\tau_{p+1}, \tau_p}^{*(i_1)} + \Delta \bar{\mathbf{a}} + \sum_{i_1, i_2=1}^m G_0^{(i_2)} B_{i_1} \hat{I}_{00\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \\
& + \sum_{i_1=1}^m \left[G_0^{(i_1)} \bar{\mathbf{a}} \left(\Delta \hat{I}_{0\tau_{p+1}, \tau_p}^{*(i_1)} + \hat{I}_{1\tau_{p+1}, \tau_p}^{*(i_1)} \right) - \bar{L} B_{i_1} \hat{I}_{1\tau_{p+1}, \tau_p}^{*(i_1)} \right] + \\
& \quad + \sum_{i_1, i_2, i_3=1}^m G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{000\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} + \frac{\Delta^2}{2} \bar{L} \bar{\mathbf{a}} + \\
& + \sum_{i_1, i_2=1}^m \left[G_0^{(i_2)} \bar{L} B_{i_1} \left(\hat{I}_{10\tau_{p+1}, \tau_p}^{*(i_2 i_1)} - \hat{I}_{01\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \right) - \bar{L} G_0^{(i_2)} B_{i_1} \hat{I}_{10\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \right. \\
& \quad \left. + G_0^{(i_2)} G_0^{(i_1)} \bar{\mathbf{a}} \left(\hat{I}_{01\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \Delta \hat{I}_{00\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \right) \right] + \\
& \quad + \sum_{i_1, i_2, i_3, i_4=1}^m G_0^{(i_4)} G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{0000\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} + \\
& + \sum_{i_1=1}^m \left[G_0^{(i_1)} \bar{L} \bar{\mathbf{a}} \left(\frac{1}{2} \hat{I}_{2\tau_{p+1}, \tau_p}^{*(i_1)} + \Delta \hat{I}_{1\tau_{p+1}, \tau_p}^{*(i_1)} + \frac{\Delta^2}{2} \hat{I}_{0\tau_{p+1}, \tau_p}^{*(i_1)} \right) + \right. \\
& \quad \left. + \frac{1}{2} \bar{L} \bar{L} B_{i_1} \hat{I}_{2\tau_{p+1}, \tau_p}^{*(i_1)} - \bar{L} G_0^{(i_1)} \bar{\mathbf{a}} \left(\hat{I}_{2\tau_{p+1}, \tau_p}^{*(i_1)} + \Delta \hat{I}_{1\tau_{p+1}, \tau_p}^{*(i_1)} \right) \right] + \\
& \quad + \sum_{i_1, i_2, i_3=1}^m \left[G_0^{(i_3)} \bar{L} G_0^{(i_2)} B_{i_1} \left(\hat{I}_{100\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} - \hat{I}_{010\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} \right) + \right.
\end{aligned}$$

$$\begin{aligned}
& +G_0^{(i_3)}G_0^{(i_2)}\bar{L}B_{i_1}\left(\hat{I}_{010\tau_{p+1},\tau_p}^{*(i_3i_2i_1)} - \hat{I}_{001\tau_{p+1},\tau_p}^{*(i_3i_2i_1)}\right) + \\
& +G_0^{(i_3)}G_0^{(i_2)}G_0^{(i_1)}\bar{\mathbf{a}}\left(\Delta\hat{I}_{000\tau_{p+1},\tau_p}^{*(i_3i_2i_1)} + \hat{I}_{001\tau_{p+1},\tau_p}^{*(i_3i_2i_1)}\right) - \\
& \quad -\bar{L}G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{100\tau_{p+1},\tau_p}^{*(i_3i_2i_1)}\Big] + \\
& + \sum_{i_1,i_2,i_3,i_4,i_5=1}^m G_0^{(i_5)}G_0^{(i_4)}G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{00000\tau_{p+1},\tau_p}^{*(i_5i_4i_3i_2i_1)} + \\
& \quad + \frac{\Delta^3}{6}LL\mathbf{a}.
\end{aligned} \tag{85}$$

Scheme with strong order 3.0

$$\begin{aligned}
\mathbf{y}_{p+1} & = \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1}\hat{I}_{0\tau_{p+1},\tau_p}^{*(i_1)} + \Delta\bar{\mathbf{a}} + \sum_{i_1,i_2=1}^m G_0^{(i_2)}B_{i_1}\hat{I}_{00\tau_{p+1},\tau_p}^{*(i_2i_1)} + \\
& + \sum_{i_1=1}^m \left[G_0^{(i_1)}\bar{\mathbf{a}}\left(\Delta\hat{I}_{0\tau_{p+1},\tau_p}^{*(i_1)} + \hat{I}_{1\tau_{p+1},\tau_p}^{*(i_1)}\right) - \bar{L}B_{i_1}\hat{I}_{1\tau_{p+1},\tau_p}^{*(i_1)} \right] + \\
& \quad + \sum_{i_1,i_2,i_3=1}^m G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{000\tau_{p+1},\tau_p}^{*(i_3i_2i_1)} + \frac{\Delta^2}{2}\bar{L}\bar{\mathbf{a}} + \\
& + \sum_{i_1,i_2=1}^m \left[G_0^{(i_2)}\bar{L}B_{i_1}\left(\hat{I}_{10\tau_{p+1},\tau_p}^{*(i_2i_1)} - \hat{I}_{01\tau_{p+1},\tau_p}^{*(i_2i_1)}\right) - \bar{L}G_0^{(i_2)}B_{i_1}\hat{I}_{10\tau_{p+1},\tau_p}^{*(i_2i_1)} + \right. \\
& \quad \left. + G_0^{(i_2)}G_0^{(i_1)}\bar{\mathbf{a}}\left(\hat{I}_{01\tau_{p+1},\tau_p}^{*(i_2i_1)} + \Delta\hat{I}_{00\tau_{p+1},\tau_p}^{*(i_2i_1)}\right) \right] + \\
& + \sum_{i_1,i_2,i_3,i_4=1}^m G_0^{(i_4)}G_0^{(i_3)}G_0^{(i_2)}B_{i_1}\hat{I}_{0000\tau_{p+1},\tau_p}^{*(i_4i_3i_2i_1)} + \mathbf{q}_{p+1,p} + \mathbf{r}_{p+1,p},
\end{aligned} \tag{86}$$

where

$$\begin{aligned}
\mathbf{q}_{p+1,p} & = \sum_{i_1=1}^m \left[G_0^{(i_1)}\bar{L}\bar{\mathbf{a}}\left(\frac{1}{2}\hat{I}_{2\tau_{p+1},\tau_p}^{*(i_1)} + \Delta\hat{I}_{1\tau_{p+1},\tau_p}^{*(i_1)} + \frac{\Delta^2}{2}\hat{I}_{0\tau_{p+1},\tau_p}^{*(i_1)}\right) + \right. \\
& \quad \left. + \frac{1}{2}\bar{L}\bar{L}B_{i_1}\hat{I}_{2\tau_{p+1},\tau_p}^{*(i_1)} - \bar{L}G_0^{(i_1)}\bar{\mathbf{a}}\left(\hat{I}_{2\tau_{p+1},\tau_p}^{*(i_1)} + \Delta\hat{I}_{1\tau_{p+1},\tau_p}^{*(i_1)}\right) \right] +
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i_1, i_2, i_3=1}^m \left[G_0^{(i_3)} \bar{L} G_0^{(i_2)} B_{i_1} \left(\hat{I}_{100\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} - \hat{I}_{010\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} \right) + \right. \\
& \quad + G_0^{(i_3)} G_0^{(i_2)} \bar{L} B_{i_1} \left(\hat{I}_{010\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} - \hat{I}_{001\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} \right) + \\
& \quad + G_0^{(i_3)} G_0^{(i_2)} G_0^{(i_1)} \bar{\mathbf{a}} \left(\Delta \hat{I}_{000\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} + \hat{I}_{001\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} \right) - \\
& \quad \left. - \bar{L} G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{100\tau_{p+1}, \tau_p}^{*(i_3 i_2 i_1)} \right] + \\
& + \sum_{i_1, i_2, i_3, i_4, i_5=1}^m G_0^{(i_5)} G_0^{(i_4)} G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{00000\tau_{p+1}, \tau_p}^{*(i_5 i_4 i_3 i_2 i_1)} + \\
& \quad + \frac{\Delta^3}{6} \bar{L} \bar{L} \bar{\mathbf{a}},
\end{aligned}$$

and

$$\begin{aligned}
\mathbf{r}_{p+1, p} & = \sum_{i_1, i_2=1}^m \left[G_0^{(i_2)} G_0^{(i_1)} \bar{L} \bar{\mathbf{a}} \left(\frac{1}{2} \hat{I}_{02\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \Delta \hat{I}_{01\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \frac{\Delta^2}{2} \hat{I}_{00\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \right) + \right. \\
& \quad + \frac{1}{2} \bar{L} \bar{L} G_0^{(i_2)} B_{i_1} \hat{I}_{20\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \\
& \quad + G_0^{(i_2)} \bar{L} G_0^{(i_1)} \bar{\mathbf{a}} \left(\hat{I}_{11\tau_{p+1}, \tau_p}^{*(i_2 i_1)} - \hat{I}_{02\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \Delta \left(\hat{I}_{10\tau_{p+1}, \tau_p}^{*(i_2 i_1)} - \hat{I}_{01\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \right) \right) + \\
& \quad + \bar{L} G_0^{(i_2)} \bar{L} B_{i_1} \left(\hat{I}_{11\tau_{p+1}, \tau_p}^{*(i_2 i_1)} - \hat{I}_{20\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \right) + \\
& \quad + G_0^{(i_2)} \bar{L} \bar{L} B_{i_1} \left(\frac{1}{2} \hat{I}_{02\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \frac{1}{2} \hat{I}_{20\tau_{p+1}, \tau_p}^{*(i_2 i_1)} - \hat{I}_{11\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \right) - \\
& \quad \left. - \bar{L} G_0^{(i_2)} G_0^{(i_1)} \bar{\mathbf{a}} \left(\Delta \hat{I}_{10\tau_{p+1}, \tau_p}^{*(i_2 i_1)} + \hat{I}_{11\tau_{p+1}, \tau_p}^{*(i_2 i_1)} \right) \right] + \\
& + \sum_{i_1, i_2, i_3, i_4=1}^m \left[G_0^{(i_4)} G_0^{(i_3)} G_0^{(i_2)} G_0^{(i_1)} \bar{\mathbf{a}} \left(\Delta \hat{I}_{0000\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} + \hat{I}_{0001\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} \right) + \right. \\
& \quad + G_0^{(i_4)} G_0^{(i_3)} \bar{L} G_0^{(i_2)} B_{i_1} \left(\hat{I}_{0100\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} - \hat{I}_{0010\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} \right) - \\
& \quad - \bar{L} G_0^{(i_4)} G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{1000\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} + \\
& \quad \left. + G_0^{(i_4)} \bar{L} G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \left(\hat{I}_{1000\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} - \hat{I}_{0100\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} \right) + \right.
\end{aligned}$$

$$\begin{aligned}
& + G_0^{(i_4)} G_0^{(i_3)} G_0^{(i_2)} \bar{L} B_{i_1} \left(\hat{I}_{0010\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} - \hat{I}_{0001\tau_{p+1}, \tau_p}^{*(i_4 i_3 i_2 i_1)} \right) \Big] + \\
& + \sum_{i_1, i_2, i_3, i_4, i_5, i_6=1}^m G_0^{(i_6)} G_0^{(i_5)} G_0^{(i_4)} G_0^{(i_3)} G_0^{(i_2)} B_{i_1} \hat{I}_{000000\tau_{p+1}, \tau_p}^{*(i_6 i_5 i_4 i_3 i_2 i_1)}.
\end{aligned}$$

It is well known [2] that under the standard conditions the numerical schemes (82)–(86) have strong orders of convergence 1.0, 1.5, 2.0, 2.5, and 3.0 correspondingly. Among these conditions we consider only the condition for approximations of iterated Stratonovich stochastic integrals from the numerical schemes (82)–(86) [2], [57]–[61]

$$\mathbb{M} \left\{ \left(\hat{I}_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{*(i_1 \dots i_k)} - \hat{I}_{l_1 \dots l_k \tau_{p+1}, \tau_p}^{*(i_1 \dots i_k)} \right)^2 \right\} \leq C \Delta^{r+1},$$

where $r/2$ are strong orders of convergence for the numerical schemes (82)–(86), i.e. $r/2 = 1.0, 1.5, 2.0, 2.5,$ and 3.0 . Moreover, constant C does not depends on Δ .

As we mentioned above, the numerical schemes (82)–(86) are unrealizable in practice without procedures for the numerical simulation of iterated Stratonovich stochastic integrals from (80). In the next section, we give a brief overview to the effective method of the mean-square approximation of iterated Ito and Stratonovich stochastic integrals of arbitrary multiplicity k ($k \in \mathbb{N}$).

12. METHOD OF THE MEAN-SQUARE APPROXIMATION OF ITERATED ITO AND STRATONOVICH STOCHASTIC INTEGRALS BASED ON GENERALIZED MULTIPLE FOURIER SERIES

It should be noted that there is an approach to the mean-square approximation of iterated stochastic integrals based on multiple integral sums (see, for example, [4]). This method implies the partitioning of the integration interval of the iterated stochastic integral under consideration; this interval is the integration step of the numerical methods used to solve Ito SDEs; therefore, it is already fairly small and does not need to be partitioned. Computational experiments [46] show that the application of the method [4] to stochastic integrals with multiplicities $k \geq 2$ leads to unacceptably high computational cost and accumulation of computation errors. Another well-known method is based on the Karhunen–Loeve expansion of the Brownian bridge process [4]. This method has no the mentioned drawback (also see [2], [3]) but leads to iterated application of the operation of limit transition. So, the mentioned method may not converge in the mean-square sense to appropriate iterated stochastic integrals for some methods of series summation (see discussion in Sect. 13 for details).

The difficulties noted above can be overcome with a different and more effective method proposed and developed by the author in [46] (also see [12]–[43], [47]–[63]). The idea of this method is as follows: the iterated Ito stochastic integral $J[\psi^{(k)}]_{T,t}$ of the form (2) with multiplicity k is represented as a multiple stochastic integral from the nonrandom discontinuous function $K(t_1, \dots, t_k)$ of k variables (see (87) below) defined on the hypercube $[t, T]^k$, where here and further $[t, T]$ is an interval of integration of the iterated Ito stochastic integral. Then, the function $K(t_1, \dots, t_k)$ is expanded in the hypercube $[t, T]^k$ into the generalized multiple Fourier series converging in the mean-square sense in the space $L_2([t, T]^k)$. After a number of nontrivial transformations we come (see Theorems 6, 7 below) to the mean-square converging expansion of the iterated Ito stochastic integral into the multiple series of products of standard Gaussian random variables. The coefficients of this series

are the coefficients of generalized multiple Fourier series for the function $K(t_1, \dots, t_k)$, which can be calculated using the explicit formula regardless of multiplicity k of the iterated Ito stochastic integral. Hereinafter, this method is referred to as the method of generalized multiple Fourier series.

Suppose that $\{\phi_j(x)\}_{j=0}^{\infty}$ is a complete orthonormal system of functions in the space $L_2([t, T])$. Define the following function on the hypercube $[t, T]^k$

$$(87) \quad K(t_1, \dots, t_k) = \begin{cases} \psi_1(t_1) \dots \psi_k(t_k), & t_1 < \dots < t_k \\ 0, & \text{otherwise} \end{cases} = \prod_{l=1}^k \psi_l(t_l) \prod_{l=1}^{k-1} \mathbf{1}_{\{t_l < t_{l+1}\}},$$

where $t_1, \dots, t_k \in [t, T]$ ($k \geq 2$), $K(t_1) \equiv \psi_1(t_1)$ for $t_1 \in [t, T]$, $\psi_1(\tau), \dots, \psi_k(\tau) : [t, T] \rightarrow \mathbb{R}$ are continuous nonrandom functions (the case $\psi_1(\tau), \dots, \psi_k(\tau) \in L_2([t, T])$ will be considered below in this section). Here $\mathbf{1}_A$ denotes the indicator of the set A .

The function $K(t_1, \dots, t_k)$ of the form (87) is piecewise continuous in the hypercube $[t, T]^k$. At this situation it is well known that the generalized multiple Fourier series of $K(t_1, \dots, t_k) \in L_2([t, T]^k)$ is converging to $K(t_1, \dots, t_k)$ in the hypercube $[t, T]^k$ in the mean-square sense, i.e.

$$(88) \quad \lim_{p_1, \dots, p_k \rightarrow \infty} \left\| K(t_1, \dots, t_k) - \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1} \prod_{l=1}^k \phi_{j_l}(t_l) \right\|_{L_2([t, T]^k)} = 0,$$

where

$$(89) \quad C_{j_k \dots j_1} = \int_{[t, T]^k} K(t_1, \dots, t_k) \prod_{l=1}^k \phi_{j_l}(t_l) dt_1 \dots dt_k$$

is the Fourier coefficient, and

$$\|f\|_{L_2([t, T]^k)} = \left(\int_{[t, T]^k} f^2(t_1, \dots, t_k) dt_1 \dots dt_k \right)^{1/2}.$$

Consider the partition $\{\tau_j\}_{j=0}^N$ of $[t, T]$ such that

$$(90) \quad t = \tau_0 < \dots < \tau_N = T, \quad \Delta_N = \max_{0 \leq j \leq N-1} \Delta\tau_j \rightarrow 0 \text{ if } N \rightarrow \infty, \quad \Delta\tau_j = \tau_{j+1} - \tau_j.$$

Theorem 6 [46] (2006) (also see [12]-[43], [47]-[62]). *Suppose that every $\psi_l(\tau)$ ($l = 1, \dots, k$) is a continuous nonrandom function on $[t, T]$ and $\{\phi_j(x)\}_{j=0}^{\infty}$ is a complete orthonormal system of continuous functions in the space $L_2([t, T])$. Then*

$$(91) \quad \begin{aligned} J[\psi^{(k)}]_{T,t} &= \text{l.i.m.}_{p_1, \dots, p_k \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1} \left(\prod_{l=1}^k \zeta_{j_l}^{(i_l)} - \right. \\ &\left. - \text{l.i.m.}_{N \rightarrow \infty} \sum_{(l_1, \dots, l_k) \in G_k} \phi_{j_{l_1}}(\tau_{l_1}) \Delta \mathbf{w}_{\tau_{l_1}}^{(i_1)} \dots \phi_{j_{l_k}}(\tau_{l_k}) \Delta \mathbf{w}_{\tau_{l_k}}^{(i_k)} \right), \end{aligned}$$

where

$$\begin{aligned} \mathbf{G}_k &= \mathbf{H}_k \setminus \mathbf{L}_k, \quad \mathbf{H}_k = \left\{ (l_1, \dots, l_k) : l_1, \dots, l_k = 0, 1, \dots, N-1 \right\}, \\ \mathbf{L}_k &= \left\{ (l_1, \dots, l_k) : l_1, \dots, l_k = 0, 1, \dots, N-1; l_g \neq l_r \ (g \neq r); g, r = 1, \dots, k \right\}, \end{aligned}$$

l.i.m. is a limit in the mean-square sense, $i_1, \dots, i_k = 0, 1, \dots, m$,

$$(92) \quad \zeta_j^{(i)} = \int_t^T \phi_j(s) d\mathbf{w}_s^{(i)}$$

are independent standard Gaussian random variables for various i or j (if $i \neq 0$), $C_{j_k \dots j_1}$ is the Fourier coefficient (89), $\Delta \mathbf{w}_{\tau_j}^{(i)} = \mathbf{w}_{\tau_{j+1}}^{(i)} - \mathbf{w}_{\tau_j}^{(i)}$ ($i = 0, 1, \dots, m$), $\{\tau_j\}_{j=0}^N$ is a partition of $[t, T]$, which satisfies the condition (90).

In order to evaluate the significance of Theorem 6 for practice we will demonstrate its transformed particular cases for $k = 1, \dots, 6$ [46] (2006) (also see [12]-[43], [47]-[62])

$$(93) \quad J[\psi^{(1)}]_{T,t} = \text{l.i.m.}_{p_1 \rightarrow \infty} \sum_{j_1=0}^{p_1} C_{j_1} \zeta_{j_1}^{(i_1)},$$

$$(94) \quad J[\psi^{(2)}]_{T,t} = \text{l.i.m.}_{p_1, p_2 \rightarrow \infty} \sum_{j_1=0}^{p_1} \sum_{j_2=0}^{p_2} C_{j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \right),$$

$$(95) \quad \begin{aligned} J[\psi^{(3)}]_{T,t} &= \text{l.i.m.}_{p_1, \dots, p_3 \rightarrow \infty} \sum_{j_1=0}^{p_1} \sum_{j_2=0}^{p_2} \sum_{j_3=0}^{p_3} C_{j_3 j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \right. \\ &\quad \left. - \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right), \end{aligned}$$

$$(96) \quad \begin{aligned} J[\psi^{(4)}]_{T,t} &= \text{l.i.m.}_{p_1, \dots, p_4 \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_4=0}^{p_4} C_{j_4 \dots j_1} \left(\prod_{l=1}^4 \zeta_{j_l}^{(i_l)} - \right. \\ &\quad - \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \\ &\quad - \mathbf{1}_{\{i_1=i_4 \neq 0\}} \mathbf{1}_{\{j_1=j_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} - \\ &\quad - \mathbf{1}_{\{i_2=i_4 \neq 0\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_3=i_4 \neq 0\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} + \\ &\quad + \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_4 \neq 0\}} \mathbf{1}_{\{j_3=j_4\}} + \\ &\quad + \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_4 \neq 0\}} \mathbf{1}_{\{j_2=j_4\}} + \\ &\quad \left. + \mathbf{1}_{\{i_1=i_4 \neq 0\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \right), \end{aligned}$$

applied in Theorem 6 [12], [48]-[61]. The generalization of Theorem 6 for complete orthonormal with weighth $r(t_1) \dots r(t_k) \geq 0$ systems of functions in the space $L_2([t, T]^k)$ can be found in [24], [57]-[61]. Another modification of Theorem 6 is connected with the mean-square approximation of iterated stochastic integrals with respect to the infinite-dimensional Q -Wiener process [31], [32], [43]. The latters play the key role for implemetation of the high-order strong numerical methods for non-commutative semilinear stochastic partial differential equations with multiplicative trace class noise [70]-[73].

For further consideration, let us consider the generalization of formulas (93)–(98) for the case of an arbitrary multiplicity k ($k \in \mathbb{N}$) of the iterated Ito stochastic integral $J[\psi^{(k)}]_{T,t}$ defined by (2). In order to do this, let us introduce some notations. Let us consider the unordered set $\{1, 2, \dots, k\}$ and separate it into two parts: the first part consists of r unordered pairs (sequence order of these pairs is also unimportant) and the second one consists of the remaining $k - 2r$ numbers. So, we have

$$(99) \quad (\underbrace{\{\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}\}}_{\text{part 1}}, \underbrace{\{q_1, \dots, q_{k-2r}\}}_{\text{part 2}}),$$

where

$$\{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\},$$

braces mean an unordered set, and parentheses mean an ordered set.

We will say that (99) is a partition and consider the sum with respect to all possible partitions

$$(100) \quad \sum_{\substack{(\{\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}\}, \{q_1, \dots, q_{k-2r}\}) \\ \{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\}}} a_{g_1 g_2, \dots, g_{2r-1} g_{2r}, q_1 \dots q_{k-2r}},$$

where $a_{g_1 g_2, \dots, g_{2r-1} g_{2r}, q_1 \dots q_{k-2r}} \in \mathbb{R}^1$.

Below there are several examples of sums in the form (100)

$$\begin{aligned} & \sum_{\substack{(\{g_1, g_2\}) \\ \{g_1, g_2\} = \{1, 2\}}} a_{g_1 g_2} = a_{12}, \\ & \sum_{\substack{(\{\{g_1, g_2\}, \{g_3, g_4\}\}) \\ \{g_1, g_2, g_3, g_4\} = \{1, 2, 3, 4\}}} a_{g_1 g_2, g_3 g_4} = a_{12, 34} + a_{13, 24} + a_{23, 14}, \\ & \sum_{\substack{(\{g_1, g_2\}, \{q_1, q_2\}) \\ \{g_1, g_2, q_1, q_2\} = \{1, 2, 3, 4\}}} a_{g_1 g_2, q_1 q_2} = \\ & = a_{12, 34} + a_{13, 24} + a_{14, 23} + a_{23, 14} + a_{24, 13} + a_{34, 12}, \\ & \sum_{\substack{(\{g_1, g_2\}, \{q_1, q_2, q_3\}) \\ \{g_1, g_2, q_1, q_2, q_3\} = \{1, 2, 3, 4, 5\}}} a_{g_1 g_2, q_1 q_2 q_3} = \\ & = a_{12, 345} + a_{13, 245} + a_{14, 235} + a_{15, 234} + a_{23, 145} + a_{24, 135} + \\ & + a_{25, 134} + a_{34, 125} + a_{35, 124} + a_{45, 123}, \end{aligned}$$

$$\begin{aligned}
& \sum_{\substack{(\{g_1, g_2\}, \{g_3, g_4\}), \{q_1\} \\ \{g_1, g_2, g_3, g_4, q_1\} = \{1, 2, 3, 4, 5\}}} a_{g_1 g_2, g_3 g_4, q_1} = \\
& = a_{12, 34, 5} + a_{13, 24, 5} + a_{14, 23, 5} + a_{12, 35, 4} + a_{13, 25, 4} + a_{15, 23, 4} + \\
& + a_{12, 54, 3} + a_{15, 24, 3} + a_{14, 25, 3} + a_{15, 34, 2} + a_{13, 54, 2} + a_{14, 53, 2} + \\
& + a_{52, 34, 1} + a_{53, 24, 1} + a_{54, 23, 1}.
\end{aligned}$$

Now we can write (91) as

$$\begin{aligned}
(101) \quad & J[\psi^{(k)}]_{T,t} = \text{l.i.m.}_{p_1, \dots, p_k \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1} \left(\prod_{l=1}^k \zeta_{j_l}^{(i_l)} + \sum_{r=1}^{[k/2]} (-1)^r \times \right. \\
& \times \sum_{\substack{(\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}), \{q_1, \dots, q_{k-2r}\} \\ \{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\}}} \prod_{s=1}^r \mathbf{1}_{\{i_{g_{2s-1}} = i_{g_{2s}} \neq 0\}} \mathbf{1}_{\{j_{g_{2s-1}} = j_{g_{2s}}\}} \prod_{l=1}^{k-2r} \zeta_{j_{q_l}}^{(i_{q_l})} \Big),
\end{aligned}$$

where $\prod_{\emptyset}^{\text{def}} \equiv 1$, $\sum_{\emptyset}^{\text{def}} \equiv 0$, $[x]$ is an integer part of a real number x ; another notations are the same as in Theorem 6.

In particular, from (101) for $k = 5$ we obtain

$$\begin{aligned}
& J[\psi^{(5)}]_{T,t} = \text{l.i.m.}_{p_1, \dots, p_5 \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_5=0}^{p_5} C_{j_5 \dots j_1} \left(\prod_{l=1}^5 \zeta_{j_l}^{(i_l)} - \right. \\
& - \sum_{\substack{(\{g_1, g_2\}, \{q_1, q_2, q_3\}) \\ \{g_1, g_2, q_1, q_2, q_3\} = \{1, 2, 3, 4, 5\}}} \mathbf{1}_{\{i_{g_1} = i_{g_2} \neq 0\}} \mathbf{1}_{\{j_{g_1} = j_{g_2}\}} \prod_{l=1}^3 \zeta_{j_{q_l}}^{(i_{q_l})} + \\
& + \sum_{\substack{(\{g_1, g_2\}, \{g_3, g_4\}), \{q_1\} \\ \{g_1, g_2, g_3, g_4, q_1\} = \{1, 2, 3, 4, 5\}}} \mathbf{1}_{\{i_{g_1} = i_{g_2} \neq 0\}} \mathbf{1}_{\{j_{g_1} = j_{g_2}\}} \mathbf{1}_{\{i_{g_3} = i_{g_4} \neq 0\}} \mathbf{1}_{\{j_{g_3} = j_{g_4}\}} \zeta_{j_{q_1}}^{(i_{q_1})} \Big).
\end{aligned}$$

The last equality obviously agrees with (97).

Let us consider the generalization of Theorem 6 for the case of an arbitrary complete orthonormal systems of functions in the space $L_2([t, T])$ and $\psi_1(\tau), \dots, \psi_k(\tau) \in L_2([t, T])$.

Theorem 7 [12] (Sect. 15), [58] (Sect. 1.11), [62]. *Suppose that $\psi_1(\tau), \dots, \psi_k(\tau) \in L_2([t, T])$ and $\{\phi_j(x)\}_{j=0}^{\infty}$ is an arbitrary complete orthonormal system of functions in the space $L_2([t, T])$. Then the following expansion*

$$\begin{aligned}
(102) \quad & J[\psi^{(k)}]_{T,t} = \text{l.i.m.}_{p_1, \dots, p_k \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1} \left(\prod_{l=1}^k \zeta_{j_l}^{(i_l)} + \sum_{r=1}^{[k/2]} (-1)^r \times \right. \\
& \times \sum_{\substack{(\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}), \{q_1, \dots, q_{k-2r}\} \\ \{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\}}} \prod_{s=1}^r \mathbf{1}_{\{i_{g_{2s-1}} = i_{g_{2s}} \neq 0\}} \mathbf{1}_{\{j_{g_{2s-1}} = j_{g_{2s}}\}} \prod_{l=1}^{k-2r} \zeta_{j_{q_l}}^{(i_{q_l})} \Big)
\end{aligned}$$

converging in the mean-square sense is valid, where $\prod_{\emptyset}^{\text{def}} \equiv 1$, $\sum_{\emptyset}^{\text{def}} \equiv 0$, $[x]$ is an integer part of a real number x ; another notations are the same as in Theorem 6.

It should be noted that an analogue of Theorem 7 was considered in [74]. Note that we use another notations in comparison with [74]. Moreover, the proof of an analogue of Theorem 7 from [74] is different from the proof given in [12] (Sect. 15), [58] (Sect. 1.11), [62].

In a number of works of the author [16]-[22], [30], [34], [40], [42], [51]-[61], [63] Theorems 6, 7 have been adapted for the iterated Stratonovich stochastic integrals (3). Let us collect some of these results (old results) in the following statement.

Theorem 8 [16]-[22], [30], [34], [40], [42], [51]-[61]. *Suppose that $\{\phi_j(x)\}_{j=0}^{\infty}$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$. At the same time $\psi_2(\tau)$ is a continuously differentiable function on $[t, T]$ and $\psi_1(\tau), \psi_3(\tau)$ are two times continuously differentiable functions on $[t, T]$. Then*

$$(103) \quad J^*[\psi^{(2)}]_{T,t} = \text{l.i.m.}_{p_1, p_2 \rightarrow \infty} \sum_{j_1=0}^{p_1} \sum_{j_2=0}^{p_2} C_{j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \quad (i_1, i_2 = 1, \dots, m),$$

$$(104) \quad J^*[\psi^{(3)}]_{T,t} = \text{l.i.m.}_{p_1, p_2, p_3 \rightarrow \infty} \sum_{j_1=0}^{p_1} \sum_{j_2=0}^{p_2} \sum_{j_3=0}^{p_3} C_{j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \quad (i_1, i_2, i_3 = 0, 1, \dots, m),$$

$$(105) \quad J^*[\psi^{(3)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \quad (i_1, i_2, i_3 = 1, \dots, m),$$

$$(106) \quad J^*[\psi^{(4)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, \dots, j_4=0}^p C_{j_4 j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \quad (i_1, \dots, i_4 = 0, 1, \dots, m),$$

where $J^*[\psi^{(k)}]_{T,t}$ is defined by (3) and $\psi_l(\tau) \equiv 1$ ($l = 1, \dots, 4$) in (104), (106); another notations are the same as in Theorem 6.

Note that the formula (103) is generalized to the case of continuous functions $\psi_1(\tau), \psi_2(\tau)$ in [58] (Sect. 2.1.4).

Recently, a new approach to the expansion and mean-square approximation of iterated Stratonovich stochastic integrals has been obtained [16] (Sect. 13–19), [21] (Sect. 5–11), [22] (Sect. 7–13), [58] (Sect. 2.10–2.16), [61] (Sect. 2.10–2.16), [63], [64]. Let us introduce some notations and formulate the main theorem of the approach noted above.

Consider the Fourier coefficient

$$C_{j_k \dots j_1} = \int_t^T \psi_k(t_k) \phi_{j_k}(t_k) \dots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \dots dt_k$$

corresponding to the function (87), where $\{\phi_j(x)\}_{j=0}^{\infty}$ is a complete orthonormal system of functions in the space $L_2([t, T])$. At that we suppose $\phi_0(x) = 1/\sqrt{T-t}$.

Denote

$$C_{j_k \dots j_{l+1} j_l j_{l-2} \dots j_1} \Big|_{(j_l j_l) \sim (\cdot)} \stackrel{\text{def}}{=}$$

$$\begin{aligned}
(107) \quad & \stackrel{\text{def}}{=} \int_t^T \psi_k(t_k) \phi_{j_k}(t_k) \cdots \int_t^{t_{l+2}} \psi_{l+1}(t_{l+1}) \phi_{j_{l+1}}(t_{l+1}) \int_t^{t_{l+1}} \psi_l(t_l) \psi_{l-1}(t_l) \times \\
& \times \int_t^{t_l} \psi_{l-2}(t_{l-2}) \phi_{j_{l-2}}(t_{l-2}) \cdots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \cdots dt_{l-2} dt_l t_{l+1} \cdots dt_k = \\
& = \sqrt{T-t} \int_t^T \psi_k(t_k) \phi_{j_k}(t_k) \cdots \int_t^{t_{l+2}} \psi_{l+1}(t_{l+1}) \phi_{j_{l+1}}(t_{l+1}) \int_t^{t_{l+1}} \psi_l(t_l) \psi_{l-1}(t_l) \phi_0(t_l) \times \\
& \times \int_t^{t_l} \psi_{l-2}(t_{l-2}) \phi_{j_{l-2}}(t_{l-2}) \cdots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \cdots dt_{l-2} dt_l t_{l+1} \cdots dt_k = \\
& = \sqrt{T-t} \hat{C}_{j_k \dots j_{l+1} 0 j_{l-2} \dots j_1},
\end{aligned}$$

i.e. $\sqrt{T-t} \hat{C}_{j_k \dots j_{l+1} 0 j_{l-2} \dots j_1}$ is again the Fourier coefficient of type $C_{j_k \dots j_1}$ but with a new shorter multi-index $j_k \dots j_{l+1} 0 j_{l-2} \dots j_1$ and new weight functions $\psi_1(\tau), \dots, \psi_{l-2}(\tau), \sqrt{T-t} \psi_{l-1}(\tau) \psi_l(\tau), \psi_{l+1}(\tau), \dots, \psi_k(\tau)$ (also we suppose that $\{l, l-1\}$ is one of the pairs $\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}$ (see the relation (99)).

Denote

$$\begin{aligned}
& \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}} \stackrel{\text{def}}{=} \\
& \stackrel{\text{def}}{=} \sum_{j_{g_{2r-1}}=p+1}^{\infty} \sum_{j_{g_{2r-3}}=p+1}^{\infty} \cdots \sum_{j_{g_3}=p+1}^{\infty} \sum_{j_{g_1}=p+1}^{\infty} C_{j_k \dots j_1} \Big|_{j_{g_1}=j_{g_2}, \dots, j_{g_{2r-1}}=j_{g_{2r}}}.
\end{aligned}$$

Introduce the following notation

$$\begin{aligned}
& S_l \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}} \right\} \stackrel{\text{def}}{=} \frac{1}{2} \mathbf{1}_{\{g_{2l}=g_{2l-1}+1\}} \sum_{j_{g_{2r-1}}=p+1}^{\infty} \sum_{j_{g_{2r-3}}=p+1}^{\infty} \cdots \\
& \cdots \sum_{j_{g_{2l+1}}=p+1}^{\infty} \sum_{j_{g_{2l-3}}=p+1}^{\infty} \cdots \sum_{j_{g_3}=p+1}^{\infty} \sum_{j_{g_1}=p+1}^{\infty} C_{j_k \dots j_1} \Big|_{(j_{g_{2l}} j_{g_{2l-1}}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, \dots, j_{g_{2r-1}}=j_{g_{2r}}}.
\end{aligned}$$

Note that the operation S_l ($l = 1, 2, \dots, r$) acts on the value

$$(108) \quad \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}}$$

as follows: S_l multiplies (108) by $\mathbf{1}_{\{g_{2l}=g_{2l-1}+1\}}/2$, removes the summation

$$\sum_{j_{g_{2l-1}}=p+1}^{\infty},$$

and replaces

$$C_{j_k \dots j_1} \Big|_{j_{g_1}=j_{g_2}, \dots, j_{g_{2r-1}}=j_{g_{2r}}}$$

with

$$(109) \quad C_{j_k \dots j_1} \Big|_{(j_{g_{2l}} j_{g_{2l-1}}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, \dots, j_{g_{2r-1}}=j_{g_{2r}}}.$$

Note that we write

$$C_{j_k \dots j_1} \Big|_{(j_{g_1} j_{g_2}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}} = C_{j_k \dots j_1} \Big|_{(j_{g_1} j_{g_1}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}},$$

$$C_{j_k \dots j_1} \Big|_{(j_{g_1} j_{g_2}) \curvearrowright (\cdot), (j_{g_3} j_{g_4}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, j_{g_3}=j_{g_4}} = C_{j_k \dots j_1} \Big|_{(j_{g_1} j_{g_1}) \curvearrowright (\cdot), (j_{g_3} j_{g_3}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, j_{g_3}=j_{g_4}}, \dots$$

Since (109) is again the Fourier coefficient, then the action of superposition $S_l S_m$ on (109) is obvious. For example, for $r = 3$

$$S_3 S_2 S_1 \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_5, g_6} \right\} =$$

$$= \frac{1}{2^3} \prod_{s=1}^3 \mathbf{1}_{\{g_{2s}=g_{2s-1}+1\}} C_{j_k \dots j_1} \Big|_{(j_{g_2} j_{g_1}) \curvearrowright (\cdot), (j_{g_4} j_{g_3}) \curvearrowright (\cdot), (j_{g_6} j_{g_5}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, j_{g_3}=j_{g_4}, j_{g_5}=j_{g_6}},$$

$$S_3 S_1 \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_5, g_6} \right\} =$$

$$= \frac{1}{2^2} \mathbf{1}_{\{g_6=g_5+1\}} \mathbf{1}_{\{g_2=g_1+1\}} \sum_{j_{g_3}=p+1}^{\infty} C_{j_k \dots j_1} \Big|_{(j_{g_2} j_{g_1}) \curvearrowright (\cdot), (j_{g_6} j_{g_5}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, j_{g_3}=j_{g_4}, j_{g_5}=j_{g_6}},$$

$$S_2 \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_5, g_6} \right\} =$$

$$= \frac{1}{2} \mathbf{1}_{\{g_4=g_3+1\}} \sum_{j_{g_1}=p+1}^{\infty} \sum_{j_{g_5}=p+1}^{\infty} C_{j_k \dots j_1} \Big|_{(j_{g_4} j_{g_3}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, j_{g_3}=j_{g_4}, j_{g_5}=j_{g_6}}.$$

Theorem 9 [16] (Sect. 13), [21] (Sect. 5), [22] (Sect. 7), [58] (Sect. 2.10), [63] (Sect. 4). *Assume that the continuously differentiable functions $\psi_1(\tau), \dots, \psi_k(\tau)$ and the complete orthonormal system*

$\{\phi_j(x)\}_{j=0}^\infty$ of continuous functions ($\phi_0(x) = 1/\sqrt{T-t}$) in the space $L_2([t, T])$ are such that the following conditions are satisfied:

1. The equality

$$(110) \quad \frac{1}{2} \int_t^s \Phi_1(t_1) \Phi_2(t_1) dt_1 = \sum_{j_1=0}^\infty \int_t^s \Phi_2(t_2) \phi_{j_1}(t_2) \int_t^{t_2} \Phi_1(t_1) \phi_{j_1}(t_1) dt_1 dt_2$$

holds for all $s \in (t, T]$, where the nonrandom functions $\Phi_1(\tau)$, $\Phi_2(\tau)$ are continuously differentiable on $[t, T]$ and the series on the right-hand side of (110) converges absolutely.

2. The estimates

$$\left| \int_t^s \phi_j(\tau) \Phi_1(\tau) d\tau \right| \leq \frac{\Psi_1(s)}{j^{1/2+\alpha}}, \quad \left| \int_s^T \phi_j(\tau) \Phi_2(\tau) d\tau \right| \leq \frac{\Psi_1(s)}{j^{1/2+\alpha}},$$

$$\left| \sum_{j=p+1}^\infty \int_t^s \Phi_2(\tau) \phi_j(\tau) \int_t^\tau \Phi_1(\theta) \phi_j(\theta) d\theta d\tau \right| \leq \frac{\Psi_2(s)}{p^\beta}$$

hold for all $s \in (t, T)$ and for some $\alpha, \beta > 0$, where $\Phi_1(\tau)$, $\Phi_2(\tau)$ are continuously differentiable nonrandom functions on $[t, T]$, $j, p \in \mathbb{N}$, and

$$\int_t^T \Psi_1^2(\tau) d\tau < \infty, \quad \int_t^T |\Psi_2(\tau)| d\tau < \infty.$$

3. The condition

$$\lim_{p \rightarrow \infty} \sum_{\substack{j_1, \dots, j_q, \dots, j_k=0 \\ q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}}}^p \left(S_{l_1} S_{l_2} \dots S_{l_d} \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}} \right\} \right)^2 = 0$$

holds for all possible $g_1, g_2, \dots, g_{2r-1}, g_{2r}$ (see (99)) and l_1, l_2, \dots, l_d such that $l_1, l_2, \dots, l_d \in \{1, 2, \dots, r\}$, $l_1 > l_2 > \dots > l_d$, $d = 0, 1, 2, \dots, r-1$, where $r = 1, 2, \dots, [k/2]$ and

$$S_{l_1} S_{l_2} \dots S_{l_d} \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}} \right\} \stackrel{\text{def}}{=} \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}}$$

for $d = 0$.

Then, for the iterated Stratonovich stochastic integral of arbitrary multiplicity k

$$(111) \quad J^*[\psi^{(k)}]_{T,t} = \int_t^*{}^T \psi_k(t_k) \dots \int_t^*{}^{t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_k}^{(i_k)}$$

the following expansion

$$(112) \quad J^*[\psi^{(k)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, \dots, j_k=0}^p C_{j_k \dots j_1} \prod_{l=1}^k \zeta_{j_l}^{(i_l)}$$

that converges in the mean-square sense is valid, where

$$C_{j_k \dots j_1} = \int_t^T \psi_k(t_k) \phi_{j_k}(t_k) \dots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \dots dt_k$$

is the Fourier coefficient, l.i.m. is a limit in the mean-square sense, $i_1, \dots, i_k = 0, 1, \dots, m$,

$$\zeta_j^{(i)} = \int_t^T \phi_j(\tau) d\mathbf{w}_\tau^{(i)}$$

are independent standard Gaussian random variables for various i or j (in the case when $i \neq 0$), $\mathbf{w}_\tau^{(i)} = \mathbf{f}_\tau^{(i)}$ for $i = 1, \dots, m$ and $\mathbf{w}_\tau^{(0)} = \tau$.

Note that (110) is fulfilled for the case of an arbitrary complete orthonormal system of functions in $L_2([t, T])$ and $\Phi_1(\tau), \Phi_2(\tau) \in L_2([t, T])$ (see [58], Sect. 2.1.4 or [77]).

Consider the following four theorems, which were proved in [16], [21], [22], [58], [61], [63], [64].

Theorem 10 [16], [21], [22], [58], [63]. *Suppose that $\{\phi_j(x)\}_{j=0}^\infty$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$. Furthermore, let $\psi_1(\tau), \psi_2(\tau), \psi_3(\tau)$ are continuously differentiable nonrandom functions on $[t, T]$. Then, for the iterated Stratonovich stochastic integral of third multiplicity*

$$J^*[\psi^{(3)}]_{T,t} = \int_t^{*T} \psi_3(t_3) \int_t^{*t_3} \psi_2(t_2) \int_t^{*t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} d\mathbf{w}_{t_2}^{(i_2)} d\mathbf{w}_{t_3}^{(i_3)} \quad (i_1, i_2, i_3 = 0, 1, \dots, m)$$

the following relations

$$(113) \quad J^*[\psi^{(3)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)},$$

$$(114) \quad \mathbb{M} \left\{ \left(J^*[\psi^{(3)}]_{T,t} - \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \right)^2 \right\} \leq \frac{C}{p}$$

are fulfilled, where $i_1, i_2, i_3 = 0, 1, \dots, m$ in (113) and $i_1, i_2, i_3 = 1, \dots, m$ in (114), constant C is independent of p ,

$$C_{j_3 j_2 j_1} = \int_t^T \psi_3(t_3) \phi_{j_3}(t_3) \int_t^{t_3} \psi_2(t_2) \phi_{j_2}(t_2) \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 dt_2 dt_3$$

and

$$\zeta_j^{(i)} = \int_t^T \phi_j(\tau) d\mathbf{f}_\tau^{(i)}$$

are independent standard Gaussian random variables for various i or j (in the case when $i \neq 0$); another notations are the same as in Theorem 6.

Theorem 11 [16], [21], [22], [58], [63]. Let $\{\phi_j(x)\}_{j=0}^\infty$ be a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$. Furthermore, let $\psi_1(\tau), \dots, \psi_4(\tau)$ be continuously differentiable nonrandom functions on $[t, T]$. Then, for the iterated Stratonovich stochastic integral of fourth multiplicity

$$(115) \quad J^*[\psi^{(4)}]_{T,t} = \int_t^{*T} \psi_4(t_4) \int_t^{*t_4} \psi_3(t_3) \int_t^{*t_3} \psi_2(t_2) \int_t^{*t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} d\mathbf{w}_{t_2}^{(i_2)} d\mathbf{w}_{t_3}^{(i_3)} d\mathbf{w}_{t_4}^{(i_4)}$$

the following relations

$$(116) \quad J^*[\psi^{(4)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)},$$

$$(117) \quad \mathbb{M} \left\{ \left(J^*[\psi^{(4)}]_{T,t} - \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \right)^2 \right\} \leq \frac{C}{p^{1-\varepsilon}}$$

are fulfilled, where $i_1, \dots, i_4 = 0, 1, \dots, m$ in (115), (116) and $i_1, \dots, i_4 = 1, \dots, m$ in (117), constant C does not depend on p , ε is an arbitrary small positive real number for the case of complete orthonormal system of Legendre polynomials in the space $L_2([t, T])$ and $\varepsilon = 0$ for the case of complete orthonormal system of trigonometric functions in the space $L_2([t, T])$,

$$C_{j_4 j_3 j_2 j_1} = \int_t^T \psi_4(t_4) \phi_{j_4}(t_4) \int_t^{t_4} \psi_3(t_3) \phi_{j_3}(t_3) \int_t^{t_3} \psi_2(t_2) \phi_{j_2}(t_2) \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 dt_2 dt_3 dt_4;$$

another notations are the same as in Theorem 10.

Theorem 12 [16], [21], [22], [58], [63]. Assume that $\{\phi_j(x)\}_{j=0}^\infty$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$ and $\psi_1(\tau), \dots, \psi_5(\tau)$ are continuously differentiable nonrandom functions on $[t, T]$. Then, for the iterated Stratonovich stochastic integral of fifth multiplicity

$$(118) \quad J^*[\psi^{(5)}]_{T,t} = \int_t^{*T} \psi_5(t_5) \dots \int_t^{*t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_5}^{(i_5)}$$

the following relations

$$(119) \quad J^*[\psi^{(5)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, \dots, j_5=0}^p C_{j_5 \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_5}^{(i_5)},$$

$$(120) \quad \mathbb{M} \left\{ \left(J^*[\psi^{(5)}]_{T,t} - \sum_{j_1, \dots, j_5=0}^p C_{j_5 \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_5}^{(i_5)} \right)^2 \right\} \leq \frac{C}{p^{1-\varepsilon}}$$

are fulfilled, where $i_1, \dots, i_5 = 0, 1, \dots, m$ in (118), (119) and $i_1, \dots, i_5 = 1, \dots, m$ in (120), constant C is independent of p , ε is an arbitrary small positive real number for the case of complete orthonormal system of Legendre polynomials in the space $L_2([t, T])$ and $\varepsilon = 0$ for the case of complete orthonormal system of trigonometric functions in the space $L_2([t, T])$,

$$C_{j_5 \dots j_1} = \int_t^T \psi_5(t_5) \phi_{j_5}(t_5) \dots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \dots dt_5;$$

another notations are the same as in Theorem 10, 11.

Theorem 13 [16], [21], [22], [58], [64]. Suppose that $\{\phi_j(x)\}_{j=0}^\infty$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$. Then, for the iterated Stratonovich stochastic integral of sixth multiplicity

$$(121) \quad J_{T,t}^{*(i_1 \dots i_6)} = \int_t^{*T} \dots \int_t^{*t_2} d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_6}^{(i_6)}$$

the following expansion

$$J_{T,t}^{*(i_1 \dots i_6)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, \dots, j_6=0}^p C_{j_6 \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_6}^{(i_6)}$$

that converges in the mean-square sense is valid, where $i_1, \dots, i_6 = 0, 1, \dots, m$,

$$C_{j_6 \dots j_1} = \int_t^T \phi_{j_6}(t_6) \dots \int_t^{t_2} \phi_{j_1}(t_1) dt_1 \dots dt_6;$$

another notations are the same as in Theorems 10–12.

On the base of Theorems 8–13 in [22], [55]–[61] the following hypothesis was formulated.

Hypothesis 1 [22], [55]–[61]. Assume that $\{\phi_j(x)\}_{j=0}^\infty$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$. Then, for the iterated Stratonovich stochastic integral of k th multiplicity

$$(122) \quad J^*[\psi^{(k)}]_{T,t} = \int_t^{*T} \psi_k(t_k) \dots \int_t^{*t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_k}^{(i_k)} \quad (i_1, \dots, i_k = 0, 1, \dots, m)$$

the following converging in the mean-square sense expansion

$$(123) \quad J^*[\psi^{(k)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, \dots, j_k=0}^p C_{j_k \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_k}^{(i_k)}$$

holds, where the Fourier coefficient $C_{j_k \dots j_1}$ has the form

$$C_{j_k \dots j_1} = \int_t^T \psi_k(t_k) \phi_{j_k}(t_k) \dots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \dots dt_k,$$

l.i.m. is a limit in the mean-square sense,

$$\zeta_j^{(i)} = \int_t^T \phi_j(\tau) d\mathbf{w}_\tau^{(i)}$$

are independent standard Gaussian random variables for various i or j (if $i \neq 0$), $\mathbf{w}_\tau^{(i)} = \mathbf{f}_\tau^{(i)}$ are independent standard Wiener processes ($i = 1, \dots, m$) and $\mathbf{w}_\tau^{(0)} = \tau$.

The hypothesis 1 allows to approximate the iterated Stratonovich stochastic integral $J^*[\psi^{(k)}]_{T,t}$ by the sum

$$(124) \quad J^*[\psi^{(k)}]_{T,t}^p = \sum_{j_1, \dots, j_k=0}^p C_{j_k \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_k}^{(i_k)},$$

where

$$\lim_{p \rightarrow \infty} \mathbb{M} \left\{ \left(J^*[\psi^{(k)}]_{T,t} - J^*[\psi^{(k)}]_{T,t}^p \right)^2 \right\} = 0.$$

The following theorem shows how to calculate exactly the mean-square approximation error for iterated Ito stochastic integrals in Theorems 6, 7.

Theorem 14 [14] (Sect. 6). *Suppose that $\psi_1(\tau), \dots, \psi_k(\tau) \in L_2([t, T])$ and $\{\phi_j(x)\}_{j=0}^\infty$ is an arbitrary complete orthonormal system of functions in the space $L_2([t, T])$. Then*

$$(125) \quad \mathbb{M} \left\{ \left(J[\psi^{(k)}]_{T,t} - J[\psi^{(k)}]_{T,t}^p \right)^2 \right\} = \int_{[t, T]^k} K^2(t_1, \dots, t_k) dt_1 \dots dt_k -$$

$$- \sum_{j_1=0}^p \dots \sum_{j_k=0}^p C_{j_k \dots j_1} \mathbb{M} \left\{ J[\psi^{(k)}]_{T,t} \sum_{(j_1, \dots, j_k)} \int_t^T \phi_{j_k}(t_k) \dots \int_t^{t_2} \phi_{j_1}(t_1) d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)} \right\},$$

where $J[\psi^{(k)}]_{T,t}$ is the iterated Ito stochastic integral (2),

$$(126) \quad J[\psi^{(k)}]_{T,t}^p = \sum_{j_1=0}^p \dots \sum_{j_k=0}^p C_{j_k \dots j_1} \left(\prod_{l=1}^k \zeta_{j_l}^{(i_l)} + \sum_{r=1}^{[k/2]} (-1)^r \times \right.$$

$$\left. \sum_{\substack{(\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}), \{q_1, \dots, q_{k-2r}\} \\ \{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\}}} \prod_{s=1}^r \mathbf{1}_{\{i_{g_{2s-1}} = i_{g_{2s}} \neq 0\}} \mathbf{1}_{\{j_{g_{2s-1}} = j_{g_{2s}}\}} \prod_{l=1}^{k-2r} \zeta_{j_{q_l}}^{(i_{q_l})} \right),$$

$i_1, \dots, i_k = 1, \dots, m$, the expression

$$\sum_{(j_1, \dots, j_k)}$$

means the sum with respect to all possible permutations (j_1, \dots, j_k) . At the same time if j_r swapped with j_q in the permutation (j_1, \dots, j_k) , then i_r swapped with i_q in the permutation (i_1, \dots, i_k) ; another notations are the same as in Theorems 6, 7.

Denote

$$E_k^p \stackrel{\text{def}}{=} \mathbb{M} \left\{ \left(J[\psi^{(k)}]_{T,t} - J[\psi^{(k)}]_{T,t}^p \right)^2 \right\}.$$

Note that

$$\begin{aligned} & \mathbb{M} \left\{ J[\psi^{(k)}]_{T,t} \int_t^T \phi_{j_k}(t_k) \cdots \int_t^{t_2} \phi_{j_1}(t_1) d\mathbf{f}_{t_1}^{(i_1)} \cdots d\mathbf{f}_{t_k}^{(i_k)} \right\} = \\ & = \int_t^T \psi_k(t_k) \phi_{j_k}(t_k) \cdots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \cdots dt_k = C_{j_k \dots j_1}. \end{aligned}$$

Then from Theorem 14 for pairwise different i_1, \dots, i_k and for $i_1 = \dots = i_k$ we obtain

$$(127) \quad \begin{aligned} E_k^p &= I_k - \sum_{j_1, \dots, j_k=0}^p C_{j_k \dots j_1}^2, \\ E_k^p &= I_k - \sum_{j_1, \dots, j_k=0}^p C_{j_k \dots j_1} \left(\sum_{(j_1, \dots, j_k)} C_{j_k \dots j_1} \right) \end{aligned}$$

correspondingly.

Consider some examples of application of Theorem 14 ($i_1, \dots, i_5 = 1, \dots, m$)

$$(128) \quad E_2^p = I_2 - \sum_{j_1, j_2=0}^p C_{j_2 j_1}^2 - \sum_{j_1, j_2=0}^p C_{j_2 j_1} C_{j_1 j_2} \quad (i_1 = i_2),$$

$$(129) \quad E_3^p = I_3 - \sum_{j_3, j_2, j_1=0}^p C_{j_3 j_2 j_1}^2 - \sum_{j_3, j_2, j_1=0}^p C_{j_3 j_1 j_2} C_{j_3 j_2 j_1} \quad (i_1 = i_2 \neq i_3),$$

$$(130) \quad E_3^p = I_3 - \sum_{j_3, j_2, j_1=0}^p C_{j_3 j_2 j_1}^2 - \sum_{j_3, j_2, j_1=0}^p C_{j_2 j_3 j_1} C_{j_3 j_2 j_1} \quad (i_1 \neq i_2 = i_3),$$

$$E_3^p = I_3 - \sum_{j_3, j_2, j_1=0}^p C_{j_3 j_2 j_1}^2 - \sum_{j_3, j_2, j_1=0}^p C_{j_3 j_2 j_1} C_{j_1 j_2 j_3} \quad (i_1 = i_3 \neq i_2),$$

$$E_4^p = I_4 - \sum_{j_1, \dots, j_4=0}^p C_{j_4 \dots j_1} \left(\sum_{(j_1, j_2)} C_{j_4 \dots j_1} \right) \quad (i_1 = i_2 \neq i_3, i_4; i_3 \neq i_4),$$

$$E_4^p = I_4 - \sum_{j_1, \dots, j_4=0}^p C_{j_4 \dots j_1} \left(\sum_{(j_1, j_3)} C_{j_4 \dots j_1} \right) \quad (i_1 = i_3 \neq i_2, i_4; i_2 \neq i_4),$$

$$E_4^p = I_4 - \sum_{j_1, \dots, j_4=0}^p C_{j_4 \dots j_1} \left(\sum_{(j_1, j_4)} C_{j_4 \dots j_1} \right) \quad (i_1 = i_4 \neq i_2, i_3; i_2 \neq i_3),$$

$$E_4^p = I_4 - \sum_{j_1, \dots, j_4=0}^p C_{j_4 \dots j_1} \left(\sum_{(j_2, j_3)} C_{j_4 \dots j_1} \right) \quad (i_2 = i_3 \neq i_1, i_4; i_1 \neq i_4),$$

$$E_4^p = I_4 - \sum_{j_1, \dots, j_4=0}^p C_{j_4 \dots j_1} \left(\sum_{(j_2, j_4)} C_{j_4 \dots j_1} \right) \quad (i_2 = i_4 \neq i_1, i_3; i_1 \neq i_3),$$

$$E_4^p = I_4 - \sum_{j_1, \dots, j_4=0}^p C_{j_4 \dots j_1} \left(\sum_{(j_3, j_4)} C_{j_4 \dots j_1} \right) \quad (i_3 = i_4 \neq i_1, i_2; i_1 \neq i_2),$$

$$E_4^p = I_4 - \sum_{j_1, \dots, j_4=0}^p C_{j_4 \dots j_1} \left(\sum_{(j_1, j_3, j_4)} C_{j_4 \dots j_1} \right) \quad (i_1 = i_3 = i_4 \neq i_2),$$

$$E_5^p = I_5 - \sum_{j_1, \dots, j_5=0}^p C_{j_5 \dots j_1} \left(\sum_{(j_2, j_4)} \left(\sum_{(j_3, j_5)} C_{j_5 \dots j_1} \right) \right) \quad (i_1 \neq i_2 = i_4 \neq i_3 = i_5 \neq i_1),$$

$$E_5^p = I_5 - \sum_{j_1, \dots, j_5=0}^p C_{j_5 \dots j_1} \left(\sum_{(j_4, j_5)} \left(\sum_{(j_1, j_2, j_3)} C_{j_5 \dots j_1} \right) \right) \quad (i_1 = i_2 = i_3 \neq i_4 = i_5),$$

$$E_5^p = I_5 - \sum_{j_1, \dots, j_5=0}^p C_{j_5 \dots j_1} \left(\sum_{(j_1, j_3, j_4, j_5)} C_{j_5 \dots j_1} \right) \quad (i_1 = i_3 = i_4 = i_5 \neq i_2).$$

Let $J[\psi^{(k)}]_{T,t}^{p_1, \dots, p_k}$ be the expression on the right-hand side of (102) before passing to the limit. Denote

$$I_k \stackrel{\text{def}}{=} \int_{[t, T]^k} K^2(t_1, \dots, t_k) dt_1 \dots dt_k.$$

In [12], [14], [57]–[61] it was shown that

$$(131) \quad \mathbb{M} \left\{ \left(J[\psi^{(k)}]_{T,t} - J[\psi^{(k)}]_{T,t}^{p_1, \dots, p_k} \right)^2 \right\} \leq k! \left(I_k - \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1}^2 \right),$$

where $i_1, \dots, i_k = 1, \dots, m$ and $0 < T - t < \infty$ or $i_1, \dots, i_k = 0, 1, \dots, m$ and $0 < T - t < 1$.

Moreover [12], [48]–[61],

$$\mathbb{M} \left\{ \left(J[\psi^{(k)}]_{T,t} - J[\psi^{(k)}]_{T,t}^{p_1, \dots, p_k} \right)^{2n} \right\} \leq C_{n,k} \left(I_k - \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1}^2 \right)^n,$$

where $C_{n,k} = (k!)^{2n} (n(2n-1))^{n(k-1)} (2n-1)!!$, $n \in \mathbb{N}$.

Below we provide practical material (based on Theorems 6–8, 10–13) concerning expansions and approximations of iterated Ito and Stratonovich stochastic integrals of multiplicities 1 to 6 (the case of Legendre polynomial system). The question about what kind of functions (polynomial or trigonometric) is more convenient for the mean-square approximation of iterated stochastic integrals is also considered.

Let us introduce more convenient (for further) notations for the iterated Ito and Stratonovich stochastic integrals

$$(132) \quad I_{(l_1 \dots l_k)T,t}^{(i_1 \dots i_k)} = \int_t^T (t-t_k)^{l_k} \dots \int_t^{t_2} (t-t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)},$$

$$(133) \quad I_{(l_1 \dots l_k)T,t}^* = \int_t^*{}^T (t-t_k)^{l_k} \dots \int_t^*{}^{t_2} (t-t_1)^{l_1} d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)},$$

where $i_1, \dots, i_k = 1, \dots, m$, $l_1, \dots, l_k = 0, 1, \dots$

The complete orthonormal system of Legendre polynomials in the space $L_2([t, T])$ looks as follows

$$(134) \quad \phi_j(x) = \sqrt{\frac{2j+1}{T-t}} P_j \left(\left(x - \frac{T+t}{2} \right) \frac{2}{T-t} \right), \quad j = 0, 1, 2, \dots,$$

where $P_j(x)$ is the Legendre polynomial.

Using the system of functions (134) and Theorems 6–8, 10–13, we obtain the following expansions of iterated Ito and Stratonovich stochastic integrals (132), (133) [13], [15], [36]–[39], [46]–[63] (also see early publications [44] (2000), [45] (2001), [65] (1997), [67] (1998))

$$I_{(0)T,t}^{(i_1)} = \sqrt{T-t} \zeta_0^{(i_1)},$$

$$(135) \quad I_{(1)T,t}^{(i_1)} = -\frac{(T-t)^{3/2}}{2} \left(\zeta_0^{(i_1)} + \frac{1}{\sqrt{3}} \zeta_1^{(i_1)} \right),$$

$$(136) \quad I_{(2)T,t}^{(i_1)} = \frac{(T-t)^{5/2}}{3} \left(\zeta_0^{(i_1)} + \frac{\sqrt{3}}{2} \zeta_1^{(i_1)} + \frac{1}{2\sqrt{5}} \zeta_2^{(i_1)} \right),$$

$$(137) \quad I_{(00)T,t}^*(i_1 i_2) = \frac{T-t}{2} \left(\zeta_0^{(i_1)} \zeta_0^{(i_2)} + \sum_{i=1}^{\infty} \frac{1}{\sqrt{4i^2-1}} \left(\zeta_{i-1}^{(i_1)} \zeta_i^{(i_2)} - \zeta_i^{(i_1)} \zeta_{i-1}^{(i_2)} \right) \right),$$

$$I_{(00)T,t}^{(i_1 i_2)} = \frac{T-t}{2} \left(\zeta_0^{(i_1)} \zeta_0^{(i_2)} + \sum_{i=1}^{\infty} \frac{1}{\sqrt{4i^2-1}} \left(\zeta_{i-1}^{(i_1)} \zeta_i^{(i_2)} - \zeta_i^{(i_1)} \zeta_{i-1}^{(i_2)} \right) - \mathbf{1}_{\{i_1=i_2\}} \right),$$

$$I_{(01)T,t}^*(i_1 i_2) = -\frac{T-t}{2} I_{(00)T,t}^*(i_1 i_2) - \frac{(T-t)^2}{4} \left(\frac{1}{\sqrt{3}} \zeta_0^{(i_1)} \zeta_1^{(i_2)} + \sum_{i=0}^{\infty} \left(\frac{(i+2) \zeta_i^{(i_1)} \zeta_{i+2}^{(i_2)} - (i+1) \zeta_{i+2}^{(i_1)} \zeta_i^{(i_2)}}{\sqrt{(2i+1)(2i+5)(2i+3)}} - \frac{\zeta_i^{(i_1)} \zeta_i^{(i_2)}}{(2i-1)(2i+3)} \right) \right),$$

$$(138) \quad I_{(10)T,t}^{*(i_1 i_2)} = -\frac{T-t}{2} I_{(00)T,t}^{*(i_1 i_2)} - \frac{(T-t)^2}{4} \left(\frac{1}{\sqrt{3}} \zeta_0^{(i_2)} \zeta_1^{(i_1)} + \sum_{i=0}^{\infty} \left(\frac{(i+1)\zeta_{i+2}^{(i_2)} \zeta_i^{(i_1)} - (i+2)\zeta_i^{(i_2)} \zeta_{i+2}^{(i_1)}}{\sqrt{(2i+1)(2i+5)(2i+3)}} + \frac{\zeta_i^{(i_1)} \zeta_i^{(i_2)}}{(2i-1)(2i+3)} \right) \right),$$

or

$$I_{(01)T,t}^{*(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{01} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)},$$

$$I_{(10)T,t}^{*(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{10} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)},$$

where

$$C_{j_2 j_1}^{01} = \frac{\sqrt{(2j_1+1)(2j_2+1)}}{8} (T-t)^2 \bar{C}_{j_2 j_1}^{01},$$

$$C_{j_2 j_1}^{10} = \frac{\sqrt{(2j_1+1)(2j_2+1)}}{8} (T-t)^2 \bar{C}_{j_2 j_1}^{10},$$

$$\bar{C}_{j_2 j_1}^{01} = -\int_{-1}^1 (1+y) P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy,$$

$$\bar{C}_{j_2 j_1}^{10} = -\int_{-1}^1 P_{j_2}(y) \int_{-1}^y (1+x) P_{j_1}(x) dx dy;$$

$$I_{(10)T,t}^{(i_1 i_2)} = I_{(10)T,t}^{*(i_1 i_2)} + \frac{1}{4} \mathbf{1}_{\{i_1=i_2\}} (T-t)^2 \quad \text{w. p. 1,}$$

$$I_{(01)T,t}^{(i_1 i_2)} = I_{(01)T,t}^{*(i_1 i_2)} + \frac{1}{4} \mathbf{1}_{\{i_1=i_2\}} (T-t)^2 \quad \text{w. p. 1,}$$

$$I_{(01)T,t}^{(i_1 i_2)} = -\frac{T-t}{2} I_{(00)T,t}^{(i_1 i_2)} - \frac{(T-t)^2}{4} \left(\frac{1}{\sqrt{3}} \zeta_0^{(i_1)} \zeta_1^{(i_2)} + \sum_{i=0}^{\infty} \left(\frac{(i+2)\zeta_i^{(i_1)} \zeta_{i+2}^{(i_2)} - (i+1)\zeta_{i+2}^{(i_1)} \zeta_i^{(i_2)}}{\sqrt{(2i+1)(2i+5)(2i+3)}} - \frac{\zeta_i^{(i_1)} \zeta_i^{(i_2)}}{(2i-1)(2i+3)} \right) \right),$$

$$I_{(10)T,t}^{(i_1 i_2)} = -\frac{T-t}{2} I_{(00)T,t}^{(i_1 i_2)} - \frac{(T-t)^2}{4} \left(\frac{1}{\sqrt{3}} \zeta_0^{(i_2)} \zeta_1^{(i_1)} + \sum_{i=0}^{\infty} \left(\frac{(i+1)\zeta_{i+2}^{(i_2)} \zeta_i^{(i_1)} - (i+2)\zeta_i^{(i_2)} \zeta_{i+2}^{(i_1)}}{\sqrt{(2i+1)(2i+5)(2i+3)}} + \frac{\zeta_i^{(i_1)} \zeta_i^{(i_2)}}{(2i-1)(2i+3)} \right) \right),$$

or

$$I_{(01)T,t}^{(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{01} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \right),$$

$$I_{(10)T,t}^{(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{10} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \right),$$

$$I_{(000)T,t}^{*(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)},$$

$$I_{(000)T,t}^{(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right),$$

(139)

$$I_{(000)T,t}^{(i_1 i_1 i_1)} = \frac{1}{6} (T-t)^{3/2} \left(\left(\zeta_0^{(i_1)} \right)^3 - 3\zeta_0^{(i_1)} \right) \quad \text{w. p. 1,}$$

$$I_{(000)T,t}^{*(i_1 i_1 i_1)} = \frac{1}{6} (T-t)^{3/2} \left(\zeta_0^{(i_1)} \right)^3 \quad \text{w. p. 1,}$$

where

$$C_{j_3 j_2 j_1} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)}}{8} (T-t)^{3/2} \bar{C}_{j_3 j_2 j_1},$$

$$\bar{C}_{j_3 j_2 j_1} = \int_{-1}^1 P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz;$$

(141)

$$I_{(000)T,t}^{(i_1 i_2 i_3)} = I_{(000)T,t}^{*(i_1 i_2 i_3)} + \mathbf{1}_{\{i_1=i_2 \neq 0\}} \frac{1}{2} I_{(1)T,t}^{(i_3)}$$

$$-\mathbf{1}_{\{i_2=i_3 \neq 0\}} \frac{1}{2} \left((T-t)I_{(0)T,t}^{(i_1)} + I_{(1)T,t}^{(i_1)} \right) \quad \text{w. p. 1,}$$

$$\begin{aligned} I_{(02)T,t}^{*(i_1 i_2)} &= -\frac{(T-t)^2}{4} I_{(00)T,t}^{*(i_1 i_2)} - (T-t)I_{(01)T,t}^{*(i_1 i_2)} + \frac{(T-t)^3}{8} \left[\frac{2}{3\sqrt{5}} \zeta_2^{(i_2)} \zeta_0^{(i_1)} + \right. \\ &+ \frac{1}{3} \zeta_0^{(i_1)} \zeta_0^{(i_2)} + \sum_{i=0}^{\infty} \left(\frac{(i+2)(i+3) \zeta_{i+3}^{(i_2)} \zeta_i^{(i_1)} - (i+1)(i+2) \zeta_i^{(i_2)} \zeta_{i+3}^{(i_1)}}{\sqrt{(2i+1)(2i+7)(2i+3)(2i+5)}} + \right. \\ &\left. \left. + \frac{(i^2+i-3) \zeta_{i+1}^{(i_2)} \zeta_i^{(i_1)} - (i^2+3i-1) \zeta_i^{(i_2)} \zeta_{i+1}^{(i_1)}}{\sqrt{(2i+1)(2i+3)(2i-1)(2i+5)}} \right) \right], \end{aligned} \quad (142)$$

$$\begin{aligned} I_{(20)T,t}^{*(i_1 i_2)} &= -\frac{(T-t)^2}{4} I_{(00)T,t}^{*(i_1 i_2)} - (T-t)I_{(10)T,t}^{*(i_1 i_2)} + \frac{(T-t)^3}{8} \left[\frac{2}{3\sqrt{5}} \zeta_0^{(i_2)} \zeta_2^{(i_1)} + \right. \\ &+ \frac{1}{3} \zeta_0^{(i_1)} \zeta_0^{(i_2)} + \sum_{i=0}^{\infty} \left(\frac{(i+1)(i+2) \zeta_{i+3}^{(i_2)} \zeta_i^{(i_1)} - (i+2)(i+3) \zeta_i^{(i_2)} \zeta_{i+3}^{(i_1)}}{\sqrt{(2i+1)(2i+7)(2i+3)(2i+5)}} + \right. \\ &\left. \left. + \frac{(i^2+3i-1) \zeta_{i+1}^{(i_2)} \zeta_i^{(i_1)} - (i^2+i-3) \zeta_i^{(i_2)} \zeta_{i+1}^{(i_1)}}{\sqrt{(2i+1)(2i+3)(2i-1)(2i+5)}} \right) \right], \end{aligned} \quad (143)$$

$$\begin{aligned} I_{(11)T,t}^{*(i_1 i_2)} &= -\frac{(T-t)^2}{4} I_{(00)T,t}^{*(i_1 i_2)} - \frac{(T-t)}{2} \left(I_{(10)T,t}^{*(i_1 i_2)} + I_{(01)T,t}^{*(i_1 i_2)} \right) + \\ &+ \frac{(T-t)^3}{8} \left[\frac{1}{3} \zeta_1^{(i_1)} \zeta_1^{(i_2)} + \sum_{i=0}^{\infty} \left(\frac{(i+1)(i+3) \left(\zeta_{i+3}^{(i_2)} \zeta_i^{(i_1)} - \zeta_i^{(i_2)} \zeta_{i+3}^{(i_1)} \right)}{\sqrt{(2i+1)(2i+7)(2i+3)(2i+5)}} + \right. \right. \\ &\left. \left. + \frac{(i+1)^2 \left(\zeta_{i+1}^{(i_2)} \zeta_i^{(i_1)} - \zeta_i^{(i_2)} \zeta_{i+1}^{(i_1)} \right)}{\sqrt{(2i+1)(2i+3)(2i-1)(2i+5)}} \right) \right], \end{aligned} \quad (144)$$

or

$$I_{(02)T,t}^{*(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{02} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)},$$

$$I_{(20)T,t}^{*(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{20} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)},$$

$$I_{(11)T,t}^{*(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{11} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)},$$

where

$$C_{j_2 j_1}^{02} = \frac{\sqrt{(2j_1+1)(2j_2+1)}}{16} (T-t)^3 \bar{C}_{j_2 j_1}^{02},$$

$$C_{j_2 j_1}^{20} = \frac{\sqrt{(2j_1+1)(2j_2+1)}}{16} (T-t)^3 \bar{C}_{j_2 j_1}^{20},$$

$$C_{j_2 j_1}^{11} = \frac{\sqrt{(2j_1+1)(2j_2+1)}}{16} (T-t)^3 \bar{C}_{j_2 j_1}^{11},$$

$$\bar{C}_{j_2 j_1}^{02} = \int_{-1}^1 P_{j_2}(y)(y+1)^2 \int_{-1}^y P_{j_1}(x) dx dy,$$

$$\bar{C}_{j_2 j_1}^{20} = \int_{-1}^1 P_{j_2}(y) \int_{-1}^y P_{j_1}(x)(x+1)^2 dx dy,$$

$$\bar{C}_{j_2 j_1}^{11} = \int_{-1}^1 P_{j_2}(y)(y+1) \int_{-1}^y P_{j_1}(x)(x+1) dx dy,$$

$$I_{(11)T,t}^{*(i_1 i_1)} = \frac{1}{2} \left(I_{(1)T,t}^{(i_1)} \right)^2 \quad \text{w. p. 1,}$$

$$I_{(02)T,t}^{(i_1 i_2)} = I_{(02)T,t}^{*(i_1 i_2)} - \frac{1}{6} \mathbf{1}_{\{i_1=i_2\}} (T-t)^3 \quad \text{w. p. 1,}$$

$$I_{(20)T,t}^{(i_1 i_2)} = I_{(20)T,t}^{*(i_1 i_2)} - \frac{1}{6} \mathbf{1}_{\{i_1=i_2\}} (T-t)^3 \quad \text{w. p. 1,}$$

$$I_{(11)T,t}^{(i_1 i_2)} = I_{(11)T,t}^{*(i_1 i_2)} - \frac{1}{6} \mathbf{1}_{\{i_1=i_2\}} (T-t)^3 \quad \text{w. p. 1,}$$

$$\begin{aligned}
I_{(02)T,t}^{(i_1 i_2)} &= -\frac{(T-t)^2}{4} I_{(00)T,t}^{(i_1 i_2)} - (T-t) I_{01T,t}^{(i_1 i_2)} + \frac{(T-t)^3}{8} \left[\frac{2}{3\sqrt{5}} \zeta_2^{(i_2)} \zeta_0^{(i_1)} + \right. \\
&+ \frac{1}{3} \zeta_0^{(i_1)} \zeta_0^{(i_2)} + \sum_{i=0}^{\infty} \left(\frac{(i+2)(i+3) \zeta_{i+3}^{(i_2)} \zeta_i^{(i_1)} - (i+1)(i+2) \zeta_i^{(i_2)} \zeta_{i+3}^{(i_1)}}{\sqrt{(2i+1)(2i+7)(2i+3)(2i+5)}} + \right. \\
&\left. \left. + \frac{(i^2+i-3) \zeta_{i+1}^{(i_2)} \zeta_i^{(i_1)} - (i^2+3i-1) \zeta_i^{(i_2)} \zeta_{i+1}^{(i_1)}}{\sqrt{(2i+1)(2i+3)(2i-1)(2i+5)}} \right) \right] - \\
&\quad - \frac{1}{24} \mathbf{1}_{\{i_1=i_2\}} (T-t)^3,
\end{aligned}$$

$$\begin{aligned}
I_{(20)T,t}^{(i_1 i_2)} &= -\frac{(T-t)^2}{4} I_{(00)T,t}^{(i_1 i_2)} - (T-t) I_{(10)T,t}^{(i_1 i_2)} + \frac{(T-t)^3}{8} \left[\frac{2}{3\sqrt{5}} \zeta_0^{(i_2)} \zeta_2^{(i_1)} + \right. \\
&+ \frac{1}{3} \zeta_0^{(i_1)} \zeta_0^{(i_2)} + \sum_{i=0}^{\infty} \left(\frac{(i+1)(i+2) \zeta_{i+3}^{(i_2)} \zeta_i^{(i_1)} - (i+2)(i+3) \zeta_i^{(i_2)} \zeta_{i+3}^{(i_1)}}{\sqrt{(2i+1)(2i+7)(2i+3)(2i+5)}} + \right. \\
&\left. \left. + \frac{(i^2+3i-1) \zeta_{i+1}^{(i_2)} \zeta_i^{(i_1)} - (i^2+i-3) \zeta_i^{(i_2)} \zeta_{i+1}^{(i_1)}}{\sqrt{(2i+1)(2i+3)(2i-1)(2i+5)}} \right) \right] - \\
&\quad - \frac{1}{24} \mathbf{1}_{\{i_1=i_2\}} (T-t)^3,
\end{aligned}$$

$$\begin{aligned}
I_{(11)T,t}^{(i_1 i_2)} &= -\frac{(T-t)^2}{4} I_{(00)T,t}^{(i_1 i_2)} - \frac{T-t}{2} \left(I_{(10)T,t}^{(i_1 i_2)} + I_{(01)T,t}^{(i_1 i_2)} \right) + \frac{(T-t)^3}{8} \left[\frac{1}{3} \zeta_1^{(i_1)} \zeta_1^{(i_2)} + \right. \\
&+ \sum_{i=0}^{\infty} \left(\frac{(i+1)(i+3) \left(\zeta_{i+3}^{(i_2)} \zeta_i^{(i_1)} - \zeta_i^{(i_2)} \zeta_{i+3}^{(i_1)} \right)}{\sqrt{(2i+1)(2i+7)(2i+3)(2i+5)}} + \right. \\
&\left. \left. + \frac{(i+1)^2 \left(\zeta_{i+1}^{(i_2)} \zeta_i^{(i_1)} - \zeta_i^{(i_2)} \zeta_{i+1}^{(i_1)} \right)}{\sqrt{(2i+1)(2i+3)(2i-1)(2i+5)}} \right) \right] - \\
&\quad - \frac{1}{24} \mathbf{1}_{\{i_1=i_2\}} (T-t)^3,
\end{aligned}$$

or

$$I_{(02)T,t}^{(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{02} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \right),$$

$$I_{(20)T,t}^{(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{20} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \right),$$

$$I_{(11)T,t}^{(i_1 i_2)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1}^{11} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \right),$$

$$I_{(3)T,t}^{(i_1)} = -\frac{(T-t)^{7/2}}{4} \left(\zeta_0^{(i_1)} + \frac{3\sqrt{3}}{5} \zeta_1^{(i_1)} + \frac{1}{\sqrt{5}} \zeta_2^{(i_1)} + \frac{1}{5\sqrt{7}} \zeta_3^{(i_1)} \right),$$

$$I_{(0000)T,t}^{*(i_1 i_2 i_3 i_4)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)},$$

$$(145) \quad I_{(0000)T,t}^{(i_1 i_2 i_3 i_4)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} + \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} + \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} + \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \right),$$

$$I_{(0000)T,t}^{(i_1 i_1 i_1 i_1)} = \frac{1}{24} (T-t)^2 \left(\left(\zeta_0^{(i_1)} \right)^4 - 6 \left(\zeta_0^{(i_1)} \right)^2 + 3 \right) \quad \text{w. p. 1,}$$

$$I_{(0000)T,t}^{*(i_1 i_1 i_1 i_1)} = \frac{1}{24} (T-t)^2 \left(\zeta_0^{(i_1)} \right)^4 \quad \text{w. p. 1,}$$

where

$$(146) \quad C_{j_4 j_3 j_2 j_1} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)(2j_4+1)}}{16} (T-t)^2 \bar{C}_{j_4 j_3 j_2 j_1},$$

$$(147) \quad \bar{C}_{j_4 j_3 j_2 j_1} = \int_{-1}^1 P_{j_4}(u) \int_{-1}^u P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz du;$$

$$I_{(001)T,t}^{*(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1}^{001} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)},$$

$$I_{(010)T,t}^{*(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1}^{010} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)},$$

$$I_{(100)T,t}^{*(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1}^{100} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)},$$

$$(148) \quad I_{(001)T,t}^{(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1}^{001} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \right. \\ \left. - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right),$$

$$(149) \quad I_{(010)T,t}^{(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1}^{010} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \right. \\ \left. - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right),$$

$$(150) \quad I_{(100)T,t}^{(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1}^{100} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \right. \\ \left. - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right),$$

where

$$C_{j_3 j_2 j_1}^{001} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)}}{16} (T-t)^{5/2} \bar{C}_{j_3 j_2 j_1}^{001},$$

$$C_{j_3 j_2 j_1}^{010} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)}}{16} (T-t)^{5/2} \bar{C}_{j_3 j_2 j_1}^{010},$$

$$C_{j_3 j_2 j_1}^{100} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)}}{16} (T-t)^{5/2} \bar{C}_{j_3 j_2 j_1}^{100},$$

$$\bar{C}_{j_3 j_2 j_1}^{100} = - \int_{-1}^1 P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x)(x+1) dx dy dz,$$

$$\bar{C}_{j_3 j_2 j_1}^{010} = - \int_{-1}^1 P_{j_3}(z) \int_{-1}^z P_{j_2}(y)(y+1) \int_{-1}^y P_{j_1}(x) dx dy dz,$$

$$\bar{C}_{j_3 j_2 j_1}^{001} = - \int_{-1}^1 P_{j_3}(z)(z+1) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz;$$

$$I_{(ll)T,t}^{(i_1 i_1 i_1)} = \frac{1}{6} \left(\left(I_{(l)T,t}^{(i_1)} \right)^3 - 3 I_{(l)T,t}^{(i_1)} \Delta_{l(T,t)} \right) \quad \text{w. p. 1,}$$

$$I_{(ll)T,t}^{*(i_1 i_1 i_1)} = \frac{1}{6} \left(I_{(l)T,t}^{(i_1)} \right)^3 \quad \text{w. p. 1,}$$

$$I_{(ull)T,t}^{(i_1 i_1 i_1 i_1)} = \frac{1}{24} \left(\left(I_{(l)T,t}^{(i_1)} \right)^4 - 6 \left(I_{(l)T,t}^{(i_1)} \right)^2 \Delta_{l(T,t)} + 3 \left(\Delta_{l(T,t)} \right)^2 \right) \quad \text{w. p. 1,}$$

$$I_{(ull)T,t}^{*(i_1 i_1 i_1 i_1)} = \frac{1}{24} \left(I_{(l)T,t}^{(i_1)} \right)^4 \quad \text{w. p. 1,}$$

where

$$I_{(l)T,t}^{(i_1)} = \sum_{j=0}^l C_j^l \zeta_j^{(i_1)} \quad \text{w. p. 1,}$$

$$\Delta_{l(T,t)} = \int_t^T (t-s)^{2l} ds, \quad C_j^l = \int_t^T (t-s)^l \phi_j(s) ds;$$

$$I_{(00000)T,t}^{*(i_1 i_2 i_3 i_4 i_5)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4, j_5=0}^p C_{j_5 j_4 j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)},$$

$$\begin{aligned}
I_{(00000)T,t}^{(i_1 i_2 i_3 i_4 i_5)} &= \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4, j_5=0}^p C_{j_5 j_4 j_3 j_2 j_1} \left(\prod_{l=1}^5 \zeta_{j_l}^{(i_l)} - \right. \\
&- \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \\
&- \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_1=j_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \\
&- \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} - \\
&- \mathbf{1}_{\{i_2=i_5\}} \mathbf{1}_{\{j_2=j_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_5}^{(i_5)} - \\
&- \mathbf{1}_{\{i_3=i_5\}} \mathbf{1}_{\{j_3=j_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_4=i_5\}} \mathbf{1}_{\{j_4=j_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} + \\
&+ \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_5\}} \mathbf{1}_{\{j_3=j_5\}} \zeta_{j_4}^{(i_4)} + \\
&+ \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_4=i_5\}} \mathbf{1}_{\{j_4=j_5\}} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_5}^{(i_5)} + \\
&+ \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_5\}} \mathbf{1}_{\{j_2=j_5\}} \zeta_{j_4}^{(i_4)} + \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_4=i_5\}} \mathbf{1}_{\{j_4=j_5\}} \zeta_{j_2}^{(i_2)} + \\
&+ \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_5\}} \mathbf{1}_{\{j_2=j_5\}} \zeta_{j_3}^{(i_3)} + \\
&+ \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_3=i_5\}} \mathbf{1}_{\{j_3=j_5\}} \zeta_{j_2}^{(i_2)} + \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_4}^{(i_4)} + \\
&+ \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_2}^{(i_2)} + \\
&+ \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_4=i_5\}} \mathbf{1}_{\{j_4=j_5\}} \zeta_{j_1}^{(i_1)} + \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_3=i_5\}} \mathbf{1}_{\{j_3=j_5\}} \zeta_{j_1}^{(i_1)} + \\
&\left. + \mathbf{1}_{\{i_2=i_5\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \right),
\end{aligned} \tag{151}$$

$$I_{(00000)T,t}^{(i_1 i_1 i_1 i_1 i_1)} = \frac{1}{120} (T-t)^{5/2} \left(\left(\zeta_0^{(i_1)} \right)^5 - 10 \left(\zeta_0^{(i_1)} \right)^3 + 15 \zeta_0^{(i_1)} \right) \quad \text{w. p. 1,}$$

$$I_{(00000)T,t}^{*(i_1 i_1 i_1 i_1 i_1)} = \frac{1}{120} (T-t)^{5/2} \left(\zeta_0^{(i_1)} \right)^5 \quad \text{w. p. 1,}$$

where

$$C_{j_5 j_4 j_3 j_2 j_1} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)(2j_4+1)(2j_5+1)}}{32} (T-t)^{5/2} \bar{C}_{j_5 j_4 j_3 j_2 j_1},$$

$$\bar{C}_{j_5 j_4 j_3 j_2 j_1} = \int_{-1}^1 P_{j_5}(v) \int_{-1}^v P_{j_4}(u) \int_{-1}^u P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz dudv;$$

$$I_{(0001)T,t}^{*(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1}^{0001} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)},$$

$$I_{(0010)T,t}^{*(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1}^{0010} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)},$$

$$I_{(0100)T,t}^{*(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1}^{0100} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)},$$

$$I_{(1000)T,t}^{*(i_1 i_2 i_3)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1}^{1000} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)},$$

$$\begin{aligned} I_{(0001)T,t}^{(i_1 i_2 i_3 i_4)} &= \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1}^{0001} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \right. \\ &- \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \\ &- \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} - \\ &- \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} + \\ &+ \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} + \\ &+ \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} + \\ &\left. + \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \right), \end{aligned}$$

$$\begin{aligned} I_{(0010)T,t}^{(i_1 i_2 i_3 i_4)} &= \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1}^{0010} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \right. \\ &- \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \\ &- \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} - \\ &- \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} + \\ &+ \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} + \\ &+ \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} + \\ &\left. + \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \right), \end{aligned}$$

$$\begin{aligned} I_{(0100)T,t}^{(i_1 i_2 i_3 i_4)} &= \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1}^{0100} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \right. \\ &- \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \end{aligned}$$

$$\begin{aligned}
& -\mathbf{1}_{\{i_1=i_4\}}\mathbf{1}_{\{j_1=j_4\}}\zeta_{j_2}^{(i_2)}\zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3\}}\mathbf{1}_{\{j_2=j_3\}}\zeta_{j_1}^{(i_1)}\zeta_{j_4}^{(i_4)} - \\
& -\mathbf{1}_{\{i_2=i_4\}}\mathbf{1}_{\{j_2=j_4\}}\zeta_{j_1}^{(i_1)}\zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_3=i_4\}}\mathbf{1}_{\{j_3=j_4\}}\zeta_{j_1}^{(i_1)}\zeta_{j_2}^{(i_2)} + \\
& \quad +\mathbf{1}_{\{i_1=i_2\}}\mathbf{1}_{\{j_1=j_2\}}\mathbf{1}_{\{i_3=i_4\}}\mathbf{1}_{\{j_3=j_4\}} + \\
& \quad +\mathbf{1}_{\{i_1=i_3\}}\mathbf{1}_{\{j_1=j_3\}}\mathbf{1}_{\{i_2=i_4\}}\mathbf{1}_{\{j_2=j_4\}} + \\
& \quad +\mathbf{1}_{\{i_1=i_4\}}\mathbf{1}_{\{j_1=j_4\}}\mathbf{1}_{\{i_2=i_3\}}\mathbf{1}_{\{j_2=j_3\}} \Big),
\end{aligned}$$

$$\begin{aligned}
I_{(1000)T,t}^{(i_1 i_2 i_3 i_4)} &= \text{li.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1}^{1000} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \right. \\
& -\mathbf{1}_{\{i_1=i_2\}}\mathbf{1}_{\{j_1=j_2\}}\zeta_{j_3}^{(i_3)}\zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_3\}}\mathbf{1}_{\{j_1=j_3\}}\zeta_{j_2}^{(i_2)}\zeta_{j_4}^{(i_4)} - \\
& -\mathbf{1}_{\{i_1=i_4\}}\mathbf{1}_{\{j_1=j_4\}}\zeta_{j_2}^{(i_2)}\zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3\}}\mathbf{1}_{\{j_2=j_3\}}\zeta_{j_1}^{(i_1)}\zeta_{j_4}^{(i_4)} - \\
& -\mathbf{1}_{\{i_2=i_4\}}\mathbf{1}_{\{j_2=j_4\}}\zeta_{j_1}^{(i_1)}\zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_3=i_4\}}\mathbf{1}_{\{j_3=j_4\}}\zeta_{j_1}^{(i_1)}\zeta_{j_2}^{(i_2)} + \\
& \quad +\mathbf{1}_{\{i_1=i_2\}}\mathbf{1}_{\{j_1=j_2\}}\mathbf{1}_{\{i_3=i_4\}}\mathbf{1}_{\{j_3=j_4\}} + \\
& \quad +\mathbf{1}_{\{i_1=i_3\}}\mathbf{1}_{\{j_1=j_3\}}\mathbf{1}_{\{i_2=i_4\}}\mathbf{1}_{\{j_2=j_4\}} + \\
& \quad \left. +\mathbf{1}_{\{i_1=i_4\}}\mathbf{1}_{\{j_1=j_4\}}\mathbf{1}_{\{i_2=i_3\}}\mathbf{1}_{\{j_2=j_3\}} \right),
\end{aligned}$$

where

$$C_{j_4 j_3 j_2 j_1}^{0001} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)(2j_4+1)}}{32} (T-t)^3 \bar{C}_{j_4 j_3 j_2 j_1}^{0001},$$

$$C_{j_3 j_2 j_1}^{0010} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)(2j_4+1)}}{32} (T-t)^3 \bar{C}_{j_4 j_3 j_2 j_1}^{0010},$$

$$C_{j_4 j_3 j_2 j_1}^{0100} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)(2j_4+1)}}{32} (T-t)^3 \bar{C}_{j_3 j_2 j_1}^{0100},$$

$$C_{j_4 j_3 j_2 j_1}^{1000} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)(2j_4+1)}}{32} (T-t)^3 \bar{C}_{j_4 j_3 j_2 j_1}^{1000},$$

$$\bar{C}_{j_4 j_3 j_2 j_1}^{1000} = - \int_{-1}^1 P_{j_4}(u) \int_{-1}^u P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x)(x+1) dx dy dz du,$$

$$\bar{C}_{j_4 j_3 j_2 j_1}^{0100} = - \int_{-1}^1 P_{j_4}(u) \int_{-1}^u P_{j_3}(z) \int_{-1}^z P_{j_2}(y)(y+1) \int_{-1}^y P_{j_1}(x) dx dy dz du,$$

$$\bar{C}_{j_4 j_3 j_2 j_1}^{0010} = - \int_{-1}^1 P_{j_4}(u) \int_{-1}^u P_{j_3}(z)(z+1) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz du,$$

$$\bar{C}_{j_4 j_3 j_2 j_1}^{0001} = - \int_{-1}^1 P_{j_4}(u)(u+1) \int_{-1}^u P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz du;$$

$$I_{(000000)T,t}^{*(i_1 i_2 i_3 i_4 i_5 i_6)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4, j_5, j_6=0}^p C_{j_6 j_5 j_4 j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)},$$

$$\begin{aligned} I_{(000000)T,t}^{(i_1 i_2 i_3 i_4 i_5 i_6)} &= \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4, j_5, j_6=0}^p C_{j_6 j_5 j_4 j_3 j_2 j_1} \left(\prod_{l=1}^6 \zeta_{j_l}^{(i_l)} - \right. \\ &- \mathbf{1}_{\{j_1=j_6\}} \mathbf{1}_{\{i_1=i_6\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{j_2=j_6\}} \mathbf{1}_{\{i_2=i_6\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \\ &- \mathbf{1}_{\{j_3=j_6\}} \mathbf{1}_{\{i_3=i_6\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{j_4=j_6\}} \mathbf{1}_{\{i_4=i_6\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} - \\ &- \mathbf{1}_{\{j_5=j_6\}} \mathbf{1}_{\{i_5=i_6\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)} - \\ &- \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)} - \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)} - \\ &- \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_6}^{(i_6)} - \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)} - \\ &- \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)} - \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_6}^{(i_6)} - \\ &- \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)} - \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} \zeta_{j_6}^{(i_6)} - \\ &- \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_6}^{(i_6)} + \\ &+ \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)} + \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_4}^{(i_4)} \zeta_{j_6}^{(i_6)} + \\ &+ \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_3}^{(i_3)} \zeta_{j_6}^{(i_6)} + \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)} + \\ &+ \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \zeta_{j_4}^{(i_4)} \zeta_{j_6}^{(i_6)} + \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_6}^{(i_6)} + \\ &+ \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \zeta_{j_5}^{(i_5)} \zeta_{j_6}^{(i_6)} + \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \zeta_{j_3}^{(i_3)} \zeta_{j_6}^{(i_6)} + \\ &+ \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_6}^{(i_6)} + \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \zeta_{j_4}^{(i_4)} \zeta_{j_6}^{(i_6)} + \\ &+ \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \zeta_{j_3}^{(i_3)} \zeta_{j_6}^{(i_6)} + \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_6}^{(i_6)} + \\ &+ \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_6}^{(i_6)} + \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_6}^{(i_6)} + \\ &+ \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_6}^{(i_6)} + \mathbf{1}_{\{j_6=j_1\}} \mathbf{1}_{\{i_6=i_1\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_5}^{(i_5)} + \\ &+ \mathbf{1}_{\{j_6=j_1\}} \mathbf{1}_{\{i_6=i_1\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} + \mathbf{1}_{\{j_6=j_1\}} \mathbf{1}_{\{i_6=i_1\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} + \\ &+ \mathbf{1}_{\{j_6=j_1\}} \mathbf{1}_{\{i_6=i_1\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{j_6=j_1\}} \mathbf{1}_{\{i_6=i_1\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} + \\ &+ \mathbf{1}_{\{j_6=j_1\}} \mathbf{1}_{\{i_6=i_1\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{j_6=j_2\}} \mathbf{1}_{\{i_6=i_2\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} + \\ &+ \mathbf{1}_{\{j_6=j_2\}} \mathbf{1}_{\{i_6=i_2\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{j_6=j_2\}} \mathbf{1}_{\{i_6=i_2\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_5}^{(i_5)} + \\ &+ \mathbf{1}_{\{j_6=j_2\}} \mathbf{1}_{\{i_6=i_2\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} + \mathbf{1}_{\{j_6=j_2\}} \mathbf{1}_{\{i_6=i_2\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} + \\ &+ \mathbf{1}_{\{j_6=j_2\}} \mathbf{1}_{\{i_6=i_2\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{j_6=j_3\}} \mathbf{1}_{\{i_6=i_3\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} + \end{aligned}$$

$$\begin{aligned}
& + \mathbf{1}_{\{j_6=j_3\}} \mathbf{1}_{\{i_6=i_3\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} + \mathbf{1}_{\{j_6=j_3\}} \mathbf{1}_{\{i_6=i_3\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_5}^{(i_5)} + \\
& + \mathbf{1}_{\{j_6=j_3\}} \mathbf{1}_{\{i_6=i_3\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} + \mathbf{1}_{\{j_6=j_3\}} \mathbf{1}_{\{i_6=i_3\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_5}^{(i_5)} + \\
& + \mathbf{1}_{\{j_6=j_3\}} \mathbf{1}_{\{i_6=i_3\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{j_6=j_4\}} \mathbf{1}_{\{i_6=i_4\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} + \\
& + \mathbf{1}_{\{j_6=j_4\}} \mathbf{1}_{\{i_6=i_4\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{j_6=j_4\}} \mathbf{1}_{\{i_6=i_4\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_5}^{(i_5)} + \\
& + \mathbf{1}_{\{j_6=j_4\}} \mathbf{1}_{\{i_6=i_4\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{j_6=j_4\}} \mathbf{1}_{\{i_6=i_4\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_5}^{(i_5)} + \\
& + \mathbf{1}_{\{j_6=j_4\}} \mathbf{1}_{\{i_6=i_4\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{j_6=j_5\}} \mathbf{1}_{\{i_6=i_5\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} + \\
& + \mathbf{1}_{\{j_6=j_5\}} \mathbf{1}_{\{i_6=i_5\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{j_6=j_5\}} \mathbf{1}_{\{i_6=i_5\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} + \\
& + \mathbf{1}_{\{j_6=j_5\}} \mathbf{1}_{\{i_6=i_5\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{j_6=j_5\}} \mathbf{1}_{\{i_6=i_5\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} + \\
& \quad + \mathbf{1}_{\{j_6=j_5\}} \mathbf{1}_{\{i_6=i_5\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \\
& \quad - \mathbf{1}_{\{j_6=j_1\}} \mathbf{1}_{\{i_6=i_1\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_1\}} \mathbf{1}_{\{i_6=i_1\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_1\}} \mathbf{1}_{\{i_6=i_1\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_2\}} \mathbf{1}_{\{i_6=i_2\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_2\}} \mathbf{1}_{\{i_6=i_2\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_2\}} \mathbf{1}_{\{i_6=i_2\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_3\}} \mathbf{1}_{\{i_6=i_3\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_3\}} \mathbf{1}_{\{i_6=i_3\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} - \\
& \quad - \mathbf{1}_{\{j_3=j_6\}} \mathbf{1}_{\{i_3=i_6\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_4\}} \mathbf{1}_{\{i_6=i_4\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_4\}} \mathbf{1}_{\{i_6=i_4\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_4\}} \mathbf{1}_{\{i_6=i_4\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_5\}} \mathbf{1}_{\{i_6=i_5\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_5\}} \mathbf{1}_{\{i_6=i_5\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} - \\
& \quad - \mathbf{1}_{\{j_6=j_5\}} \mathbf{1}_{\{i_6=i_5\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \Big),
\end{aligned}$$

$$I_{(000000)T,t}^{(i_1 i_1 i_1 i_1 i_1 i_1)} = \frac{1}{720} (T-t)^3 \left(\left(\zeta_0^{(i_1)} \right)^6 - 15 \left(\zeta_0^{(i_1)} \right)^4 + 45 \left(\zeta_0^{(i_1)} \right)^2 - 15 \right) \quad \text{w. p. 1,}$$

$$I_{(000000)T,t}^{*(i_1 i_1 i_1 i_1 i_1 i_1)} = \frac{1}{720} (T-t)^3 \left(\zeta_0^{(i_1)} \right)^6 \quad \text{w. p. 1,}$$

where

$$C_{j_6 j_5 j_4 j_3 j_2 j_1} = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)(2j_4+1)(2j_5+1)(2j_6+1)}}{64} (T-t)^3 \bar{C}_{j_6 j_5 j_4 j_3 j_2 j_1},$$

$$\bar{C}_{j_6 j_5 j_4 j_3 j_2 j_1} = \int_{-1}^1 P_{j_6}(w) \int_{-1}^w P_{j_5}(v) \int_{-1}^v P_{j_4}(u) \int_{-1}^u P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz du dv dw.$$

Let us analyze the approximation $I_{(00)T,t}^{*(i_1 i_2)q}$ of the iterated stochastic integral $I_{(00)T,t}^{*(i_1 i_2)}$ obtained from (137) by replacing ∞ on q .

It is easy to prove that

$$(152) \quad \mathbb{M} \left\{ \left(I_{(00)T,t}^{*(i_1 i_2)} - I_{(00)T,t}^{*(i_1 i_2)q} \right)^2 \right\} = \frac{(T-t)^2}{2} \left(\frac{1}{2} - \sum_{i=1}^q \frac{1}{4i^2 - 1} \right) \quad (i_1 \neq i_2).$$

Moreover, using Theorem 14, we obtain for $i_1 \neq i_2$

$$\begin{aligned} & \mathbb{M} \left\{ \left(I_{(10)T,t}^{*(i_1 i_2)} - I_{(10)T,t}^{*(i_1 i_2)q} \right)^2 \right\} = \mathbb{M} \left\{ \left(I_{(01)T,t}^{*(i_1 i_2)} - I_{(01)T,t}^{*(i_1 i_2)q} \right)^2 \right\} = \\ & = \frac{(T-t)^4}{16} \left(\frac{5}{9} - 2 \sum_{i=2}^q \frac{1}{4i^2 - 1} - \sum_{i=1}^q \frac{1}{(2i-1)^2(2i+3)^2} - \sum_{i=0}^q \frac{(i+2)^2 + (i+1)^2}{(2i+1)(2i+5)(2i+3)^2} \right). \end{aligned}$$

For the case $i_1 = i_2$, using Theorem 14, we have

$$(153) \quad \begin{aligned} & \mathbb{M} \left\{ \left(I_{(10)T,t}^{(i_1 i_1)} - I_{(10)T,t}^{(i_1 i_1)q} \right)^2 \right\} = \mathbb{M} \left\{ \left(I_{(01)T,t}^{(i_1 i_1)} - I_{(01)T,t}^{(i_1 i_1)q} \right)^2 \right\} = \\ & = \frac{(T-t)^4}{16} \left(\frac{1}{9} - \sum_{i=0}^q \frac{1}{(2i+1)(2i+5)(2i+3)^2} - 2 \sum_{i=1}^q \frac{1}{(2i-1)^2(2i+3)^2} \right). \end{aligned}$$

On the basis of the presented expansions of iterated stochastic integrals we can see that increasing of multiplicities of these integrals or degree indexes of their weight functions leads to noticeable complication of formulas for the mentioned expansions.

However, increasing of the mentioned parameters leads to increasing of orders of smallness with respect to $T - t$ in the mean-square sense for iterated stochastic integrals. This leads to a sharp decrease of member quantities in expansions of iterated stochastic integrals, which are required for achieving the acceptable accuracy of approximation. In the context of it let us consider the approach to approximation of iterated stochastic integrals, which provides a possibility to obtain the mean-square approximations of the required accuracy without using the complex expansions.

Let us analyze the following approximation of triple stochastic integral using (139)

$$(154) \quad \begin{aligned} I_{(000)T,t}^{(i_1 i_2 i_3)q_1} = & \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_3 j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \right. \\ & \left. - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right), \end{aligned}$$

where $C_{j_3 j_2 j_1}$ is defined by (140), (141).

In particular, from (154) when $i_1 \neq i_2, i_2 \neq i_3, i_1 \neq i_3$ we obtain

$$(155) \quad I_{(000)T,t}^{(i_1 i_2 i_3)q_1} = \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)}.$$

Using (127), (129)–(131), we get

$$(156) \quad \begin{aligned} & \mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q_1} \right)^2 \right\} = \\ & = \frac{(T-t)^3}{6} - \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_3 j_2 j_1}^2 \quad (i_1 \neq i_2, i_1 \neq i_3, i_2 \neq i_3), \end{aligned}$$

$$(157) \quad \begin{aligned} & \mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q_1} \right)^2 \right\} = \\ & = \frac{(T-t)^3}{6} - \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_3 j_2 j_1}^2 - \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_2 j_3 j_1} C_{j_3 j_2 j_1} \quad (i_1 \neq i_2 = i_3), \end{aligned}$$

$$(158) \quad \begin{aligned} & \mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q_1} \right)^2 \right\} = \\ & = \frac{(T-t)^3}{6} - \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_3 j_2 j_1}^2 - \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_3 j_2 j_1} C_{j_1 j_2 j_3} \quad (i_1 = i_3 \neq i_2), \end{aligned}$$

$$(159) \quad \begin{aligned} & \mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q_1} \right)^2 \right\} = \\ & = \frac{(T-t)^3}{6} - \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_3 j_2 j_1}^2 - \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_3 j_1 j_2} C_{j_3 j_2 j_1} \quad (i_1 = i_2 \neq i_3), \end{aligned}$$

$$(160) \quad \mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q_1} \right)^2 \right\} \leq 6 \left(\frac{(T-t)^3}{6} - \sum_{j_1, j_2, j_3=0}^{q_1} C_{j_3 j_2 j_1}^2 \right) \quad (i_1, i_2, i_3 = 1, \dots, m).$$

We may act similarly with more complicated iterated stochastic integrals. For example, for the approximation of stochastic integral $I_{(0000)T,t}^{(i_1 i_2 i_3 i_4)}$ we may write (see (145))

$$\begin{aligned}
I_{(0000)T,t}^{(i_1 i_2 i_3 i_4)q_2} = & \sum_{j_1, j_2, j_3, j_4=0}^{q_2} C_{j_4 j_3 j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \right. \\
& - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \\
& - \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} - \\
& - \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} + \\
& + \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_4\}} \mathbf{1}_{\{j_3=j_4\}} + \\
& + \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_2=j_4\}} + \\
& \left. + \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \right),
\end{aligned}$$

where $C_{j_4 j_3 j_2 j_1}$ is defined by (146), (147). Moreover, according to (131)

$$\mathbb{M} \left\{ \left(I_{(0000)T,t}^{(i_1 i_2 i_3 i_4)} - I_{(0000)T,t}^{(i_1 i_2 i_3 i_4)q_2} \right)^2 \right\} \leq 24 \left(\frac{(T-t)^4}{24} - \sum_{j_1, j_2, j_3, j_4=0}^{q_2} C_{j_4 j_3 j_2 j_1}^2 \right) \quad (i_1, i_2, i_3, i_4 = 1, \dots, m).$$

For pairwise different $i_1, i_2, i_3, i_4 = 1, \dots, m$ from (127) we obtain

$$(161) \quad \mathbb{M} \left\{ \left(I_{(0000)T,t}^{(i_1 i_2 i_3 i_4)} - I_{(0000)T,t}^{(i_1 i_2 i_3 i_4)q_2} \right)^2 \right\} = \frac{(T-t)^4}{24} - \sum_{j_1, j_2, j_3, j_4=0}^{q_2} C_{j_4 j_3 j_2 j_1}^2.$$

Using Theorem 14, we can calculate exactly the left-hand side of (161) for any possible combinations of i_1, i_2, i_3, i_4 . These relations were obtained in [14], [57]-[61].

In Tables 3–5, we have some examples of exact values of the Fourier–Legendre coefficients (here and further the Fourier–Legendre coefficients have been calculated exactly using DERIVE (computer algebra system)). Note that in [75], [76] the database with 270,000 exactly calculated Fourier–Legendre coefficients was described. This database was used in the software package, which is written in the Python programming language for the implementation of the numerical schemes (75)–(79), (82)–(86).

Assume that $q_1 = 6$. Calculating the value of expression (156) for $q_1 = 6$, $i_1 \neq i_2$, $i_1 \neq i_3$, $i_3 \neq i_2$, we obtain

$$\mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q_1} \right)^2 \right\} \approx 0.01956(T-t)^3.$$

Let us choose, for example, $q_2 = 2$. In the case of pairwise different i_1, i_2, i_3, i_4 we have from (161) the following approximate equality

$$(162) \quad \mathbb{M} \left\{ \left(I_{(0000)T,t}^{(i_1 i_2 i_3 i_4)} - I_{(0000)T,t}^{(i_1 i_2 i_3 i_4)q_2} \right)^2 \right\} \approx 0.0236084(T-t)^4.$$

Let us analyze the approximations

Table 3. Coefficients $\bar{C}_{3j_2j_1}$.

$j_2^{j_1}$	0	1	2	3	4	5	6
0	0	$\frac{2}{105}$	0	$-\frac{4}{315}$	0	$\frac{2}{693}$	0
1	$\frac{4}{105}$	0	$-\frac{2}{315}$	0	$-\frac{8}{3465}$	0	$\frac{10}{9009}$
2	$\frac{2}{35}$	$-\frac{2}{105}$	0	$\frac{4}{3465}$	0	$-\frac{74}{45045}$	0
3	$\frac{2}{315}$	0	$-\frac{2}{3465}$	0	$\frac{16}{45045}$	0	$-\frac{10}{9009}$
4	$-\frac{2}{63}$	$\frac{46}{3465}$	0	$-\frac{32}{45045}$	0	$\frac{2}{9009}$	0
5	$-\frac{10}{693}$	0	$\frac{38}{9009}$	0	$-\frac{4}{9009}$	0	$\frac{122}{765765}$
6	0	$-\frac{10}{3003}$	0	$\frac{20}{9009}$	0	$-\frac{226}{765765}$	0

Table 4. Coefficients $\bar{C}_{21j_2j_1}$.

$j_2^{j_1}$	0	1	2
0	$\frac{2}{21}$	$-\frac{2}{45}$	$\frac{2}{315}$
1	$\frac{2}{315}$	$\frac{2}{315}$	$-\frac{225}{2}$
2	$-\frac{10}{105}$	$\frac{2}{225}$	$\frac{2}{1155}$

Table 5. Coefficients $\bar{C}_{101j_2j_1}$.

$j_2^{j_1}$	0	1
0	$\frac{4}{315}$	0
1	$\frac{4}{315}$	$-\frac{8}{945}$

$$I_{(001)T,t}^{(i_1 i_2 i_3)q_3}, \quad I_{(010)T,t}^{(i_1 i_2 i_3)q_3}, \quad I_{(100)T,t}^{(i_1 i_2 i_3)q_3}, \quad I_{(00000)T,t}^{(i_1 i_2 i_3 i_4 i_5)q_4}$$

based on the expansions (148)–(151).

Assume that $q_3 = 2, q_4 = 1$. In the case of pairwise different i_1, \dots, i_5 we obtain

$$M \left\{ \left(I_{(100)T,t}^{(i_1 i_2 i_3)} - I_{(100)T,t}^{(i_1 i_2 i_3)q_3} \right)^2 \right\} = \frac{(T-t)^5}{60} - \sum_{j_1, j_2, j_3=0}^2 (C_{j_3 j_2 j_1}^{100})^2 \approx 0.00815429(T-t)^5,$$

$$M \left\{ \left(I_{(010)T,t}^{(i_1 i_2 i_3)} - I_{(010)T,t}^{(i_1 i_2 i_3)q_3} \right)^2 \right\} = \frac{(T-t)^5}{20} - \sum_{j_1, j_2, j_3=0}^2 (C_{j_3 j_2 j_1}^{010})^2 \approx 0.0173903(T-t)^5,$$

$$M \left\{ \left(I_{(001)T,t}^{(i_1 i_2 i_3)} - I_{(001)T,t}^{(i_1 i_2 i_3)q_3} \right)^2 \right\} = \frac{(T-t)^5}{10} - \sum_{j_1, j_2, j_3=0}^2 (C_{j_3 j_2 j_1}^{001})^2 \approx 0.0252801(T-t)^5,$$

$$M \left\{ \left(I_{(00000)T,t}^{(i_1 i_2 i_3 i_4 i_5)} - I_{(00000)T,t}^{(i_1 i_2 i_3 i_4 i_5)q_4} \right)^2 \right\} =$$

$$= \frac{(T-t)^5}{120} - \sum_{j_1, j_2, j_3, j_4, j_5=0}^1 C_{j_5 j_4 j_3 j_2 j_1}^2 \approx 0.00759105(T-t)^5.$$

Note that using (131), we can write

$$\mathbb{M} \left\{ \left(I_{(00000)T,t}^{(i_1 i_2 i_3 i_4 i_5)} - I_{(00000)T,t}^{(i_1 i_2 i_3 i_4 i_5)q_4} \right)^2 \right\} \leq 120 \left(\frac{(T-t)^5}{120} - \sum_{j_1, j_2, j_3, j_4, j_5=0}^{q_4} C_{j_5 j_4 j_3 j_2 j_1}^2 \right),$$

where $i_1, \dots, i_5 = 1, \dots, m$.

Moreover, from (131) we obtain the following useful estimates

$$\mathbb{M} \left\{ \left(I_{(01)T,t}^{(i_1 i_2)} - I_{(01)T,t}^{(i_1 i_2)q} \right)^2 \right\} \leq 2 \left(\frac{(T-t)^4}{4} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{01})^2 \right),$$

$$\mathbb{M} \left\{ \left(I_{(10)T,t}^{(i_1 i_2)} - I_{(10)T,t}^{(i_1 i_2)q} \right)^2 \right\} \leq 2 \left(\frac{(T-t)^4}{12} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{10})^2 \right),$$

$$\mathbb{M} \left\{ \left(I_{(100)T,t}^{(i_1 i_2 i_3)} - I_{(100)T,t}^{(i_1 i_2 i_3)q} \right)^2 \right\} \leq 6 \left(\frac{(T-t)^5}{60} - \sum_{j_1, j_2, j_3=0}^q (C_{j_3 j_2 j_1}^{100})^2 \right),$$

$$\mathbb{M} \left\{ \left(I_{(010)T,t}^{(i_1 i_2 i_3)} - I_{(010)T,t}^{(i_1 i_2 i_3)q} \right)^2 \right\} \leq 6 \left(\frac{(T-t)^5}{20} - \sum_{j_1, j_2, j_3=0}^q (C_{j_3 j_2 j_1}^{010})^2 \right),$$

$$\mathbb{M} \left\{ \left(I_{(001)T,t}^{(i_1 i_2 i_3)} - I_{(001)T,t}^{(i_1 i_2 i_3)q} \right)^2 \right\} \leq 6 \left(\frac{(T-t)^5}{10} - \sum_{j_1, j_2, j_3=0}^q (C_{j_3 j_2 j_1}^{001})^2 \right),$$

$$\mathbb{M} \left\{ \left(I_{(20)T,t}^{(i_1 i_2)} - I_{(20)T,t}^{(i_1 i_2)q} \right)^2 \right\} \leq 2 \left(\frac{(T-t)^6}{30} - \sum_{j_2, j_1=0}^q (C_{j_2 j_1}^{20})^2 \right),$$

$$\mathbb{M} \left\{ \left(I_{(11)T,t}^{(i_1 i_2)} - I_{(11)T,t}^{(i_1 i_2)q} \right)^2 \right\} \leq 2 \left(\frac{(T-t)^6}{18} - \sum_{j_2, j_1=0}^q (C_{j_2 j_1}^{11})^2 \right),$$

$$\mathbb{M} \left\{ \left(I_{(02)T,t}^{(i_1 i_2)} - I_{(02)T,t}^{(i_1 i_2)q} \right)^2 \right\} \leq 2 \left(\frac{(T-t)^6}{6} - \sum_{j_2, j_1=0}^q (C_{j_2 j_1}^{02})^2 \right),$$

$$\mathbb{M} \left\{ \left(I_{(1000)T,t}^{(i_1 i_2 i_3 i_4)} - I_{(1000)T,t}^{(i_1 i_2 i_3 i_4)q} \right)^2 \right\} \leq 24 \left(\frac{(T-t)^6}{360} - \sum_{j_1, j_2, j_3, j_4=0}^q (C_{j_4 j_3 j_2 j_1}^{1000})^2 \right),$$

$$\begin{aligned} \mathbb{M} \left\{ \left(I_{(0100)T,t}^{(i_1 i_2 i_3 i_4)} - I_{(0100)T,t}^{(i_1 i_2 i_3 i_4)q} \right)^2 \right\} &\leq 24 \left(\frac{(T-t)^6}{120} - \sum_{j_1, j_2, j_3, j_4=0}^q (C_{j_4 j_3 j_2 j_1}^{0100})^2 \right), \\ \mathbb{M} \left\{ \left(I_{(0010)T,t}^{(i_1 i_2 i_3 i_4)} - I_{(0010)T,t}^{(i_1 i_2 i_3 i_4)q} \right)^2 \right\} &\leq 24 \left(\frac{(T-t)^6}{60} - \sum_{j_1, j_2, j_3, j_4=0}^q (C_{j_4 j_3 j_2 j_1}^{0010})^2 \right), \\ \mathbb{M} \left\{ \left(I_{(0001)T,t}^{(i_1 i_2 i_3 i_4)} - I_{(0001)T,t}^{(i_1 i_2 i_3 i_4)q} \right)^2 \right\} &\leq 24 \left(\frac{(T-t)^6}{36} - \sum_{j_1, j_2, j_3, j_4=0}^q (C_{j_4 j_3 j_2 j_1}^{0001})^2 \right), \\ \mathbb{M} \left\{ \left(I_{(000000)T,t}^{(i_1 i_2 i_3 i_4 i_5 i_6)} - I_{(000000)T,t}^{(i_1 i_2 i_3 i_4 i_5 i_6)q} \right)^2 \right\} &\leq 720 \left(\frac{(T-t)^6}{720} - \sum_{j_1, j_2, j_3, j_4, j_5, j_6=0}^q C_{j_6 j_5 j_4 j_3 j_2 j_1}^2 \right). \end{aligned}$$

In addition, from (128) we get

$$\begin{aligned} \mathbb{M} \left\{ \left(I_{(10)T,t}^{(i_1 i_2)} - I_{(10)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^4}{12} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{10})^2 - \sum_{j_1, j_2=0}^q C_{j_2 j_1}^{10} C_{j_1 j_2}^{10} \quad (i_1 = i_2), \\ \mathbb{M} \left\{ \left(I_{(10)T,t}^{(i_1 i_2)} - I_{(10)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^4}{12} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{10})^2 \quad (i_1 \neq i_2), \\ \mathbb{M} \left\{ \left(I_{(01)T,t}^{(i_1 i_2)} - I_{(01)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^4}{4} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{01})^2 - \sum_{j_1, j_2=0}^q C_{j_2 j_1}^{01} C_{j_1 j_2}^{01} \quad (i_1 = i_2), \\ \mathbb{M} \left\{ \left(I_{(01)T,t}^{(i_1 i_2)} - I_{(01)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^4}{4} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{01})^2 \quad (i_1 \neq i_2), \\ \mathbb{M} \left\{ \left(I_{(20)T,t}^{(i_1 i_2)} - I_{(20)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^6}{30} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{20})^2 - \sum_{j_1, j_2=0}^q C_{j_2 j_1}^{20} C_{j_1 j_2}^{20} \quad (i_1 = i_2), \\ \mathbb{M} \left\{ \left(I_{(20)T,t}^{(i_1 i_2)} - I_{(20)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^6}{30} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{20})^2 \quad (i_1 \neq i_2), \\ \mathbb{M} \left\{ \left(I_{(11)T,t}^{(i_1 i_2)} - I_{(11)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^6}{18} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{11})^2 - \sum_{j_1, j_2=0}^q C_{j_2 j_1}^{11} C_{j_1 j_2}^{11} \quad (i_1 = i_2), \end{aligned}$$

$$\begin{aligned} \mathbb{M} \left\{ \left(I_{(11)T,t}^{(i_1 i_2)} - I_{(11)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^6}{18} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{11})^2 \quad (i_1 \neq i_2), \\ \mathbb{M} \left\{ \left(I_{(02)T,t}^{(i_1 i_2)} - I_{(02)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^6}{6} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{02})^2 - \sum_{j_1, j_2=0}^q C_{j_2 j_1}^{02} C_{j_1 j_2}^{02} \quad (i_1 = i_2), \\ \mathbb{M} \left\{ \left(I_{(02)T,t}^{(i_1 i_2)} - I_{(02)T,t}^{(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^6}{6} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{02})^2 \quad (i_1 \neq i_2), \end{aligned}$$

Clearly, expansions for iterated Stratonovich stochastic integrals (see Theorems 8, 10-12) are simpler than expansions for iterated Ito stochastic integrals (see Theorems 6, 7, and (93)–(98)). However, the calculation of the mean-square approximation error for iterated Stratonovich stochastic integrals turns out to be much more difficult than for iterated Ito stochastic integrals (see Theorem 14 and (131)). Below we consider how we can estimate or calculate exactly (for some particular cases) the mean-square approximation error for iterated Stratonovich stochastic integrals (the development of these results is contained in Chapter 5 [58] (also see Theorems 10–13 from this paper)).

As we mentioned above, on the basis of the presented approximations of iterated Stratonovich stochastic integrals we can see that increasing of multiplicities of these integrals leads to increasing of orders of smallness with respect to $T-t$ in the mean-square sense for iterated Stratonovich stochastic integrals ($T-t \ll 1$ because the length of integration interval $[t, T]$ of the iterated Stratonovich stochastic integrals plays the role of integration step for the numerical methods for Ito SDEs, so $T-t$ is already fairly small). This leads to a sharp decrease of member quantities in the approximations of iterated Stratonovich stochastic integrals, which are required for achieving the acceptable accuracy of approximation.

From (152) ($i_1 \neq i_2$) we obtain

$$\begin{aligned} \mathbb{M} \left\{ \left(I_{(00)T,t}^{*(i_1 i_2)} - I_{(00)T,t}^{*(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^2}{2} \sum_{i=q+1}^{\infty} \frac{1}{4i^2 - 1} \leq \\ (163) \quad &\leq \frac{(T-t)^2}{2} \int_q^{\infty} \frac{1}{4x^2 - 1} dx = -\frac{(T-t)^2}{8} \ln \left| 1 - \frac{2}{2q+1} \right| \leq C_1 \frac{(T-t)^2}{q}, \end{aligned}$$

where C_1 is a constant.

Since $T-t \ll 1$, then it is easy to notice that there exists such a constant C_2 that

$$(164) \quad \mathbb{M} \left\{ \left(I_{(l_1 \dots l_k)T,t}^{*(i_1 \dots i_k)} - I_{(l_1 \dots l_k)T,t}^{*(i_1 \dots i_k)q} \right)^2 \right\} \leq C_2 \mathbb{M} \left\{ \left(I_{(00)T,t}^{*(i_1 i_2)} - I_{(00)T,t}^{*(i_1 i_2)q} \right)^2 \right\},$$

where $I_{(l_1 \dots l_k)T,t}^{*(i_1 \dots i_k)q}$ is an approximation of the iterated Stratonovich stochastic integral $I_{(l_1 \dots l_k)T,t}^{*(i_1 \dots i_k)}$.

From (163) and (164) we finally obtain

$$(165) \quad \mathbb{M} \left\{ \left(I_{(l_1 \dots l_k)T,t}^{*(i_1 \dots i_k)} - I_{(l_1 \dots l_k)T,t}^{*(i_1 \dots i_k)q} \right)^2 \right\} \leq C \frac{(T-t)^2}{q},$$

where constant C does not depend on $T-t$. Note that, in contrast to the estimate (165), the constant C in Theorems 10–12 does not depend on q .

The same idea can be found in [2] in the framework of the method based on the trigonometric expansion of the Brownian bridge process.

We can get more information about the numbers q (these numbers are different for different iterated Stratonovich stochastic integrals) using the another approach. Since for pairwise different $i_1, \dots, i_k = 1, \dots, m$

$$J^*[\psi^{(k)}]_{T,t} = J[\psi^{(k)}]_{T,t} \quad \text{w. p. 1,}$$

where $J[\psi^{(k)}]_{T,t}$, $J^*[\psi^{(k)}]_{T,t}$ are defined by (2) and (3) correspondingly, then for pairwise different $i_1, \dots, i_k = 1, \dots, m$ we can write (see (127))

$$\begin{aligned} \mathbb{M} \left\{ \left(I_{(01)T,t}^{*(i_1 i_2)} - I_{(01)T,t}^{*(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^4}{4} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{01})^2, \\ \mathbb{M} \left\{ \left(I_{(10)T,t}^{*(i_1 i_2)} - I_{(10)T,t}^{*(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^4}{12} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{10})^2, \\ \mathbb{M} \left\{ \left(I_{(000)T,t}^{*(i_1 i_2 i_3)} - I_{(000)T,t}^{*(i_1 i_2 i_3)q} \right)^2 \right\} &= \frac{(T-t)^3}{6} - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1}^2, \\ \mathbb{M} \left\{ \left(I_{(0000)T,t}^{*(i_1 i_2 i_3 i_4)} - I_{(0000)T,t}^{*(i_1 i_2 i_3 i_4)q} \right)^2 \right\} &= \frac{(T-t)^4}{24} - \sum_{j_1, j_2, j_3, j_4=0}^q C_{j_4 j_3 j_2 j_1}^2, \\ \mathbb{M} \left\{ \left(I_{(100)T,t}^{*(i_1 i_2 i_3)} - I_{(100)T,t}^{*(i_1 i_2 i_3)q} \right)^2 \right\} &= \frac{(T-t)^5}{60} - \sum_{j_1, j_2, j_3=0}^q (C_{j_3 j_2 j_1}^{100})^2, \\ \mathbb{M} \left\{ \left(I_{(010)T,t}^{*(i_1 i_2 i_3)} - I_{(010)T,t}^{*(i_1 i_2 i_3)q} \right)^2 \right\} &= \frac{(T-t)^5}{20} - \sum_{j_1, j_2, j_3=0}^q (C_{j_3 j_2 j_1}^{010})^2, \\ \mathbb{M} \left\{ \left(I_{(001)T,t}^{*(i_1 i_2 i_3)} - I_{(001)T,t}^{*(i_1 i_2 i_3)q} \right)^2 \right\} &= \frac{(T-t)^5}{10} - \sum_{j_1, j_2, j_3=0}^q (C_{j_3 j_2 j_1}^{001})^2, \\ \mathbb{M} \left\{ \left(I_{(00000)T,t}^{*(i_1 i_2 i_3 i_4 i_5)} - I_{(00000)T,t}^{*(i_1 i_2 i_3 i_4 i_5)q} \right)^2 \right\} &= \frac{(T-t)^5}{120} - \sum_{j_1, j_2, j_3, j_4, j_5=0}^q C_{j_5 i_4 i_3 i_2 j_1}^2, \\ \mathbb{M} \left\{ \left(I_{(20)T,t}^{*(i_1 i_2)} - I_{(20)T,t}^{*(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^6}{30} - \sum_{j_2, j_1=0}^q (C_{j_2 j_1}^{20})^2, \\ \mathbb{M} \left\{ \left(I_{(11)T,t}^{*(i_1 i_2)} - I_{(11)T,t}^{*(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^6}{18} - \sum_{j_2, j_1=0}^q (C_{j_2 j_1}^{11})^2, \end{aligned}$$

$$\begin{aligned}
\mathbb{M} \left\{ \left(I_{(02)T,t}^{*(i_1 i_2)} - I_{(02)T,t}^{*(i_1 i_2)q} \right)^2 \right\} &= \frac{(T-t)^6}{6} - \sum_{j_2, j_1=0}^q (C_{j_2 j_1}^{02})^2, \\
\mathbb{M} \left\{ \left(I_{(1000)T,t}^{*(i_1 i_2 i_3 i_4)} - I_{(1000)T,t}^{*(i_1 i_2 i_3 i_4)q} \right)^2 \right\} &= \frac{(T-t)^6}{360} - \sum_{j_1, j_2, j_3, j_4=0}^q (C_{j_4 j_3 j_2 j_1}^{1000})^2, \\
\mathbb{M} \left\{ \left(I_{(0100)T,t}^{*(i_1 i_2 i_3 i_4)} - I_{(0100)T,t}^{*(i_1 i_2 i_3 i_4)q} \right)^2 \right\} &= \frac{(T-t)^6}{120} - \sum_{j_1, j_2, j_3, j_4=0}^q (C_{j_4 j_3 j_2 j_1}^{0100})^2, \\
\mathbb{M} \left\{ \left(I_{(0010)T,t}^{*(i_1 i_2 i_3 i_4)} - I_{(0010)T,t}^{*(i_1 i_2 i_3 i_4)q} \right)^2 \right\} &= \frac{(T-t)^6}{60} - \sum_{j_1, j_2, j_3, j_4=0}^q (C_{j_4 j_3 j_2 j_1}^{0010})^2, \\
\mathbb{M} \left\{ \left(I_{(0001)T,t}^{*(i_1 i_2 i_3 i_4)} - I_{(0001)T,t}^{*(i_1 i_2 i_3 i_4)q} \right)^2 \right\} &= \frac{(T-t)^6}{36} - \sum_{j_1, j_2, j_3, j_4=0}^q (C_{j_4 j_3 j_2 j_1}^{0001})^2, \\
\mathbb{M} \left\{ \left(I_{(000000)T,t}^{*(i_1 i_2 i_3 i_4 i_5 i_6)} - I_{(000000)T,t}^{*(i_1 i_2 i_3 i_4 i_5 i_6)q} \right)^2 \right\} &= \frac{(T-t)^6}{720} - \sum_{j_1, j_2, j_3, j_4, j_5, j_6=0}^q C_{j_6 j_5 j_4 j_3 j_2 j_1}^2.
\end{aligned}$$

For example [46] (also see [26]-[29], [38], [47]-[61]),

$$\begin{aligned}
\mathbb{M} \left\{ \left(I_{(000)T,t}^{*(i_1 i_2 i_3)} - I_{(000)T,t}^{*(i_1 i_2 i_3)6} \right)^2 \right\} &= \frac{(T-t)^3}{6} - \sum_{j_3, j_2, j_1=0}^6 C_{j_3 j_2 j_1}^2 \approx 0.01956000(T-t)^3, \\
\mathbb{M} \left\{ \left(I_{(0000)T,t}^{*(i_1 i_2 i_3 i_4)} - I_{(0000)T,t}^{*(i_1 i_2 i_3 i_4)2} \right)^2 \right\} &= \frac{(T-t)^4}{24} - \sum_{j_1, j_2, j_3, j_4=0}^2 C_{j_4 j_3 j_2 j_1}^2 \approx 0.02360840(T-t)^4, \\
\mathbb{M} \left\{ \left(I_{(100)T,t}^{*(i_1 i_2 i_3)} - I_{(100)T,t}^{*(i_1 i_2 i_3)2} \right)^2 \right\} &= \frac{(T-t)^5}{60} - \sum_{j_1, j_2, j_3=0}^2 (C_{j_3 j_2 j_1}^{100})^2 \approx 0.00815429(T-t)^5, \\
\mathbb{M} \left\{ \left(I_{(010)T,t}^{*(i_1 i_2 i_3)} - I_{(010)T,t}^{*(i_1 i_2 i_3)2} \right)^2 \right\} &= \frac{(T-t)^5}{20} - \sum_{j_1, j_2, j_3=0}^2 (C_{j_3 j_2 j_1}^{010})^2 \approx 0.0173903(T-t)^5, \\
\mathbb{M} \left\{ \left(I_{(001)T,t}^{*(i_1 i_2 i_3)} - I_{(001)T,t}^{*(i_1 i_2 i_3)2} \right)^2 \right\} &= \frac{(T-t)^5}{10} - \sum_{j_1, j_2, j_3=0}^2 (C_{j_3 j_2 j_1}^{001})^2 \approx 0.0252801(T-t)^5, \\
\mathbb{M} \left\{ \left(I_{(00000)T,t}^{*(i_1 i_2 i_3 i_4 i_5)} - I_{(00000)T,t}^{*(i_1 i_2 i_3 i_4 i_5)1} \right)^2 \right\} &=
\end{aligned}$$

$$= \frac{(T-t)^5}{120} - \sum_{j_1, j_2, j_3, j_4, j_5=0}^1 C_{j_5 j_4 j_3 j_2 j_1}^2 \approx 0.00759105(T-t)^5.$$

Let us consider expansions of the Ito stochastic integrals $I_{(1)T,t}^{(i_1)}$, $I_{(2)T,t}^{(i_1)}$ based on the approach from [4] (also see [2])

$$(166) \quad I_{(1)T,t}^{(i_1)q} = -\frac{(T-t)^{3/2}}{2} \left(\zeta_0^{(i_1)} - \frac{\sqrt{2}}{\pi} \left(\sum_{r=1}^q \frac{1}{r} \zeta_{2r-1}^{(i_1)} + \sqrt{\alpha_q} \xi_q^{(i_1)} \right) \right),$$

$$(167) \quad I_{(2)T,t}^{(i_1)q} = (T-t)^{5/2} \left(\frac{1}{3} \zeta_0^{(i_1)} + \frac{1}{\sqrt{2}\pi^2} \left(\sum_{r=1}^q \frac{1}{r^2} \zeta_{2r}^{(i_1)} + \sqrt{\beta_q} \mu_q^{(i_1)} \right) - \frac{1}{\sqrt{2}\pi} \left(\sum_{r=1}^q \frac{1}{r} \zeta_{2r-1}^{(i_1)} + \sqrt{\alpha_q} \xi_q^{(i_1)} \right) \right),$$

where $\zeta_j^{(i)}$ is defined by the formula (92), $\phi_j(\tau)$ is a complete orthonormal system of trigonometric functions in the space $L_2([t, T])$, and $\zeta_0^{(i)}$, $\zeta_{2r}^{(i)}$, $\zeta_{2r-1}^{(i)}$, $\xi_q^{(i)}$, $\mu_q^{(i)}$ ($r = 1, \dots, q$, $i = 1, \dots, m$) are independent standard Gaussian random variables, $i_1 = 1, \dots, m$,

$$\xi_q^{(i)} = \frac{1}{\sqrt{\alpha_q}} \sum_{r=q+1}^{\infty} \frac{1}{r} \zeta_{2r-1}^{(i)}, \quad \alpha_q = \frac{\pi^2}{6} - \sum_{r=1}^q \frac{1}{r^2},$$

$$\mu_q^{(i)} = \frac{1}{\sqrt{\beta_q}} \sum_{r=q+1}^{\infty} \frac{1}{r^2} \zeta_{2r}^{(i)}, \quad \beta_q = \frac{\pi^4}{90} - \sum_{r=1}^q \frac{1}{r^4}.$$

It is obvious that (166), (167) significantly more complicated compared to (135), (136).

Another example of obvious advantage of the Legendre polynomials over the trigonometric functions (in the framework of the considered problem) is the truncated expansion of the iterated Stratonovich stochastic integral $I_{(10)T,t}^{*(i_1 i_2)}$ obtained by Theorem 8, in which instead of the double Fourier–Legendre series (see (137), (138)) is taken the double trigonometric Fourier series

$$I_{(10)T,t}^{*(i_1 i_2)q} = -(T-t)^2 \left(\frac{1}{6} \zeta_0^{(i_1)} \zeta_0^{(i_2)} - \frac{1}{2\sqrt{2}\pi} \sqrt{\alpha_q} \xi_q^{(i_2)} \zeta_0^{(i_1)} + \frac{1}{2\sqrt{2}\pi^2} \sqrt{\beta_q} \left(\mu_q^{(i_2)} \zeta_0^{(i_1)} - 2\mu_q^{(i_1)} \zeta_0^{(i_2)} \right) + \frac{1}{2\sqrt{2}} \sum_{r=1}^q \left(-\frac{1}{\pi r} \zeta_{2r-1}^{(i_2)} \zeta_0^{(i_1)} + \frac{1}{\pi^2 r^2} \left(\zeta_{2r}^{(i_2)} \zeta_0^{(i_1)} - 2\zeta_{2r}^{(i_1)} \zeta_0^{(i_2)} \right) \right) \right) -$$

$$\begin{aligned}
& -\frac{1}{2\pi^2} \sum_{\substack{r,l=1 \\ r \neq l}}^q \frac{1}{r^2 - l^2} \left(\zeta_{2r}^{(i_1)} \zeta_{2l}^{(i_2)} + \frac{l}{r} \zeta_{2r-1}^{(i_1)} \zeta_{2l-1}^{(i_2)} \right) + \\
& + \sum_{r=1}^q \left(\frac{1}{4\pi r} \left(\zeta_{2r}^{(i_1)} \zeta_{2r-1}^{(i_2)} - \zeta_{2r-1}^{(i_1)} \zeta_{2r}^{(i_2)} \right) + \right. \\
(168) \quad & \left. + \frac{1}{8\pi^2 r^2} \left(3\zeta_{2r-1}^{(i_1)} \zeta_{2r-1}^{(i_2)} + \zeta_{2r}^{(i_2)} \zeta_{2r}^{(i_1)} \right) \right),
\end{aligned}$$

where the meaning of the notations included in (166), (12) is preserved.

A deep comparative analysis of the efficiency of application of Legendre polynomials and trigonometric functions to the numerical integration of Ito SDEs is given in [23], [39].

13. THEOREMS 6–8, 10–13 FROM POINT OF VIEW OF THE WONG–ZAKAI APPROXIMATION

The iterated Ito stochastic integrals and solutions of Ito SDEs are complex and important functionals from the independent components $\mathbf{f}_s^{(i)}$, $i = 1, \dots, m$ of the multidimensional Wiener process \mathbf{f}_s , $s \in [0, T]$. Let $\mathbf{f}_s^{(i)p}$, $p \in \mathbb{N}$ be some approximation of $\mathbf{f}_s^{(i)}$, $i = 1, \dots, m$. Suppose that $\mathbf{f}_s^{(i)p}$ converges to $\mathbf{f}_s^{(i)}$, $i = 1, \dots, m$ if $p \rightarrow \infty$ in some sense and has differentiable sample trajectories.

A natural question arises: if we replace $\mathbf{f}_s^{(i)}$ by $\mathbf{f}_s^{(i)p}$, $i = 1, \dots, m$ in the functionals mentioned above, will the resulting functionals converge to the original functionals from the components $\mathbf{f}_s^{(i)}$, $i = 1, \dots, m$ of the multidimensional Wiener process \mathbf{f}_s ? The answer to this question is negative in the general case. However, in the pioneering works of Wong E. and Zakai M. [78], [79], it was shown that under the special conditions and for some types of approximations of the Wiener process the answer is affirmative with one peculiarity: the convergence takes place to the iterated Stratonovich stochastic integrals and solutions of Stratonovich SDEs and not to iterated Ito stochastic integrals and solutions of Ito SDEs. The piecewise linear approximation as well as the regularization by convolution [78]–[80] relate the mentioned types of approximations of the Wiener process. The above approximation of stochastic integrals and solutions of SDEs is often called the Wong–Zakai approximation.

Let \mathbf{f}_s , $s \in [0, T]$ be an m -dimensional standard Wiener process with independent components $\mathbf{f}_s^{(i)}$, $i = 1, \dots, m$. It is well known that the following representation takes place [81], [82]

$$(169) \quad \mathbf{f}_\tau^{(i)} - \mathbf{f}_t^{(i)} = \sum_{j=0}^{\infty} \int_t^\tau \phi_j(s) ds \zeta_j^{(i)}, \quad \zeta_j^{(i)} = \int_t^T \phi_j(s) d\mathbf{f}_s^{(i)},$$

where $\tau \in [t, T]$, $t \geq 0$, $\{\phi_j(x)\}_{j=0}^{\infty}$ is an arbitrary complete orthonormal system of functions in the space $L_2([t, T])$, and $\zeta_j^{(i)}$ are independent standard Gaussian random variables for various i or j . Moreover, the series (169) converges for any $\tau \in [t, T]$ in the mean-square sense.

Let $\mathbf{f}_\tau^{(i)p} - \mathbf{f}_t^{(i)p}$ be the mean-square approximation of the process $\mathbf{f}_\tau^{(i)} - \mathbf{f}_t^{(i)}$, which has the following form

$$(170) \quad \mathbf{f}_\tau^{(i)p} - \mathbf{f}_t^{(i)p} = \sum_{j=0}^p \int_t^\tau \phi_j(s) ds \zeta_j^{(i)}.$$

From (170) we obtain

$$(171) \quad d\mathbf{f}_\tau^{(i)p} = \sum_{j=0}^p \phi_j(\tau) \zeta_j^{(i)} d\tau.$$

Consider the following iterated Riemann–Stieltjes integral

$$(172) \quad \int_t^T \psi_k(t_k) \dots \int_t^{t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)p_1} \dots d\mathbf{w}_{t_k}^{(i_k)p_k},$$

where $p_1, \dots, p_k \in \mathbb{N}$, $i_1, \dots, i_k = 0, 1, \dots, m$,

$$(173) \quad d\mathbf{w}_\tau^{(i)p} = \begin{cases} d\mathbf{f}_\tau^{(i)p} & \text{for } i = 1, \dots, m \\ d\tau & \text{for } i = 0 \end{cases},$$

and $d\mathbf{f}_\tau^{(i)p}$ is defined by the relation (171).

Let us substitute (171) into (172)

$$(174) \quad \int_t^T \psi_k(t_k) \dots \int_t^{t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)p_1} \dots d\mathbf{w}_{t_k}^{(i_k)p_k} = \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1} \prod_{l=1}^k \zeta_{j_l}^{(i_l)},$$

where

$$\zeta_j^{(i)} = \int_t^T \phi_j(s) d\mathbf{w}_s^{(i)}$$

are independent standard Gaussian random variables for various i or j (in the case when $i \neq 0$), $\mathbf{w}_s^{(i)} = \mathbf{f}_s^{(i)}$ for $i = 1, \dots, m$ and $\mathbf{w}_s^{(0)} = s$,

$$C_{j_k \dots j_1} = \int_t^T \psi_k(t_k) \phi_{j_k}(t_k) \dots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \dots dt_k$$

is the Fourier coefficient.

To best of our knowledge [78]–[80] the approximations of the Wiener process in the Wong–Zakai approximation must satisfy fairly strong restrictions [80] (see Definition 7.1, pp. 480–481). Moreover, approximations of the Wiener process that are similar to (170) were not considered in [78], [79] (also see [80], Theorems 7.1, 7.2). Therefore, the proof of analogs of Theorems 7.1 and 7.2 [80] for approximations of the Wiener process based on its series expansion (169) should be carried out separately. Thus, the mean-square convergence of the right-hand side of (174) to the iterated Stratonovich stochastic integral (3) does not follow from the results of the papers [78], [79] (also see [80], Theorems 7.1, 7.2).

From the other hand, Theorems 8, 10–13 from this paper can be considered as the proof of the Wong–Zakai approximation for the iterated Stratonovich stochastic integrals (3) of multiplicities 2 to 6 based on the approximation (170) of the Wiener process. At that, the Riemann–Stieltjes integrals

(172) converge (according to Theorems 8, 10–13) to the appropriate Stratonovich stochastic integrals (3). Recall that $\{\phi_j(x)\}_{j=0}^{\infty}$ (see (169), (170), and Theorems 8, 10–13) is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$.

To illustrate the above reasoning, consider two examples for the case $k = 2$, $\psi_1(s), \psi_2(s) \equiv 1$; $i_1, i_2 = 1, \dots, m$.

The first example relates to the piecewise linear approximation of the multidimensional Wiener process (these approximations were considered in [78]–[80]).

Let $\mathbf{b}_{\Delta}^{(i)}(t)$, $t \in [0, T]$ be the piecewise linear approximation of the i th component $\mathbf{f}_t^{(i)}$ of the multidimensional standard Wiener process \mathbf{f}_t , $t \in [0, T]$ with independent components $\mathbf{f}_t^{(i)}$, $i = 1, \dots, m$, i.e.

$$\mathbf{b}_{\Delta}^{(i)}(t) = \mathbf{f}_{k\Delta}^{(i)} + \frac{t - k\Delta}{\Delta} \Delta \mathbf{f}_{k\Delta}^{(i)},$$

where

$$\Delta \mathbf{f}_{k\Delta}^{(i)} = \mathbf{f}_{(k+1)\Delta}^{(i)} - \mathbf{f}_{k\Delta}^{(i)}, \quad t \in [k\Delta, (k+1)\Delta], \quad k = 0, 1, \dots, N-1.$$

Note that w. p. 1

$$(175) \quad \frac{d\mathbf{b}_{\Delta}^{(i)}}{dt}(t) = \frac{\Delta \mathbf{f}_{k\Delta}^{(i)}}{\Delta}, \quad t \in [k\Delta, (k+1)\Delta], \quad k = 0, 1, \dots, N-1.$$

Consider the following iterated Riemann–Stieltjes integral

$$\int_0^T \int_0^s d\mathbf{b}_{\Delta}^{(i_1)}(\tau) d\mathbf{b}_{\Delta}^{(i_2)}(s), \quad i_1, i_2 = 1, \dots, m.$$

Using (175) and the additive property of Riemann–Stieltjes integrals, we can write w. p. 1

$$\begin{aligned} & \int_0^T \int_0^s d\mathbf{b}_{\Delta}^{(i_1)}(\tau) d\mathbf{b}_{\Delta}^{(i_2)}(s) = \int_0^T \int_0^s \frac{d\mathbf{b}_{\Delta}^{(i_1)}}{d\tau}(\tau) d\tau \frac{d\mathbf{b}_{\Delta}^{(i_2)}}{ds}(s) ds = \\ & = \sum_{l=0}^{N-1} \int_{l\Delta}^{(l+1)\Delta} \left(\sum_{q=0}^{l-1} \int_{q\Delta}^{(q+1)\Delta} \frac{\Delta \mathbf{f}_{q\Delta}^{(i_1)}}{\Delta} d\tau + \int_{l\Delta}^s \frac{\Delta \mathbf{f}_{l\Delta}^{(i_1)}}{\Delta} d\tau \right) \frac{\Delta \mathbf{f}_{l\Delta}^{(i_2)}}{\Delta} ds = \\ & = \sum_{l=0}^{N-1} \sum_{q=0}^{l-1} \Delta \mathbf{f}_{q\Delta}^{(i_1)} \Delta \mathbf{f}_{l\Delta}^{(i_2)} + \frac{1}{\Delta^2} \sum_{l=0}^{N-1} \Delta \mathbf{f}_{l\Delta}^{(i_1)} \Delta \mathbf{f}_{l\Delta}^{(i_2)} \int_{l\Delta}^{(l+1)\Delta} \int_{l\Delta}^s d\tau ds = \\ (176) \quad & = \sum_{l=0}^{N-1} \sum_{q=0}^{l-1} \Delta \mathbf{f}_{q\Delta}^{(i_1)} \Delta \mathbf{f}_{l\Delta}^{(i_2)} + \frac{1}{2} \sum_{l=0}^{N-1} \Delta \mathbf{f}_{l\Delta}^{(i_1)} \Delta \mathbf{f}_{l\Delta}^{(i_2)}. \end{aligned}$$

Using (176) and the standard relation between Stratonovich and Ito stochastic integrals, it is not difficult to show that

$$\begin{aligned}
 \text{l.i.m.}_{N \rightarrow \infty} \int_0^T \int_0^s d\mathbf{b}_\Delta^{(i_1)}(\tau) d\mathbf{b}_\Delta^{(i_2)}(s) &= \int_0^T \int_0^s d\mathbf{f}_\tau^{(i_1)} d\mathbf{f}_s^{(i_2)} + \frac{1}{2} \mathbf{1}_{\{i_1=i_2\}} \int_0^T ds = \\
 (177) \qquad \qquad \qquad &= \int_0^{*T} \int_0^{*s} d\mathbf{f}_\tau^{(i_1)} d\mathbf{f}_s^{(i_2)},
 \end{aligned}$$

where $\Delta \rightarrow 0$ if $N \rightarrow \infty$ ($N\Delta = T$).

Obviously, (177) agrees with Theorem 7.1 (see [80], p. 486).

The next example relates to the approximation of the Wiener process based on its series expansion (169) for $t = 0$, where $\{\phi_j(x)\}_{j=0}^\infty$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([0, T])$.

Consider the following iterated Riemann–Stieltjes integral

$$(178) \qquad \int_0^T \int_0^s d\mathbf{f}_\tau^{(i_1)p} d\mathbf{f}_s^{(i_2)p}, \quad i_1, i_2 = 1, \dots, m,$$

where $d\mathbf{f}_\tau^{(i)p}$ is defined by the relation (171).

Let us substitute (171) into (178)

$$(179) \qquad \int_0^T \int_0^s d\mathbf{f}_\tau^{(i_1)p} d\mathbf{f}_s^{(i_2)p} = \sum_{j_1, j_2=0}^p C_{j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)},$$

where

$$C_{j_2 j_1} = \int_0^T \phi_{j_2}(s) \int_0^s \phi_{j_1}(\tau) d\tau ds$$

is the Fourier coefficient; another notations are the same as in (174).

As we noted above, approximations of the Wiener process that are similar to (170) were not considered in [78], [79] (also see Theorems 7.1, 7.2 in [80]). Furthermore, the extension of the results of Theorems 7.1 and 7.2 [80] to the case under consideration is not obvious.

However, the authors of the works [2] (Sect. 5.8, pp. 202–204), [3] (pp. 82–84), [83] (pp. 438–439), [84] (pp. 263–264) use the Wong–Zakai approximation [78]–[80] (without rigorous proof) within the frames of the approach [4] based on the series expansion of the Brownian bridge process.

On the other hand, we can apply the theory built in Chapters 1 and 2 of the monographs [58]–[61]. More precisely, using Theorem 8 from this paper we obtain from (179) the desired result

$$\text{l.i.m.}_{p \rightarrow \infty} \int_0^T \int_0^s d\mathbf{f}_\tau^{(i_1)p} d\mathbf{f}_s^{(i_2)p} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} =$$

$$(180) \quad = \int_0^{*T} \int_0^{*s} d\mathbf{f}_\tau^{(i_1)} d\mathbf{f}_s^{(i_2)}.$$

From the other hand, by Theorem 6 (see (94)) for the case $k = 2$ we obtain from (179) the following relation

$$(181) \quad \begin{aligned} & \text{l.i.m.}_{p \rightarrow \infty} \int_0^T \int_0^s d\mathbf{f}_\tau^{(i_1)p} d\mathbf{f}_s^{(i_2)p} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} = \\ & = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2=0}^p C_{j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \right) + \mathbf{1}_{\{i_1=i_2\}} \sum_{j_1=0}^{\infty} C_{j_1 j_1} = \\ & = \int_0^T \int_0^s d\mathbf{f}_\tau^{(i_1)} d\mathbf{f}_s^{(i_2)} + \mathbf{1}_{\{i_1=i_2\}} \sum_{j_1=0}^{\infty} C_{j_1 j_1}. \end{aligned}$$

Since

$$\sum_{j_1=0}^{\infty} C_{j_1 j_1} = \frac{1}{2} \sum_{j_1=0}^{\infty} \left(\int_0^T \phi_j(\tau) d\tau \right)^2 = \frac{1}{2} \left(\int_0^T \phi_0(\tau) d\tau \right)^2 = \frac{1}{2} \int_0^T ds,$$

then from (181) and the standard relation between Stratonovich and Ito stochastic integrals we obtain (180).

REFERENCES

- [1] Gihman I.I., Skorohod A.V. Stochastic Differential Equations and its Applications. Kiev, Naukova Dumka, 1982, 612 pp.
- [2] Kloeden P.E., Platen E. Numerical Solution of Stochastic Differential Equations. Berlin, Springer, 1992, 632 pp.
- [3] Kloeden P.E., Platen E., Schurz H. Numerical Solution of SDE Through Computer Experiments. Springer, Berlin, 1994, 292 pp.
- [4] Milstein G.N. Numerical Integration of Stochastic Differential Equations. Sverdlovsk, Ural University Press, 1988, 225 pp.
- [5] Milstein G.N., Tretyakov M.V. Stochastic Numerics for Mathematical Physics. Berlin, Springer, 2004, 616 pp.
- [6] Platen E., Wagner W. On a Taylor formula for a class of Ito processes. Probab. Math. Statist. 3 (1982), 37-51.
- [7] Kloeden P.E., Platen E. The Stratonovich and Ito-Taylor expansions. Math. Nachr. 151 (1991), 33-50.
- [8] Kulchitskiy O.Yu., Kuznetsov D.F. The unified Taylor-Ito expansion. Journal of Mathematical Sciences (N.Y.). 99, 2 (2000), 1130-1140. DOI: <http://doi.org/10.1007/BF02673635>
- [9] Kuznetsov D.F. New representations of the Taylor-Stratonovich expansions. Journal of Mathematical Sciences (N.Y.). 118, 6 (2003), 5586-5596. DOI: <http://doi.org/10.1023/A:1026138522239>
- [10] Kulchitski O.Yu., Kuznetsov D.F. Expansion of Ito processes into a Taylor-Ito series in a neighborhood of a fixed time moment, Dep. VINITI, No. 2637-93, 1993, 25 pp.
- [11] Kuznetsov D.F. Methods of numerical simulation of stochastic differential Ito equations solutions in problems of mechanics. Ph. D., Saint-Petersburg, 1996. 260 pp.
- [12] Kuznetsov D.F. Expansion of iterated Ito stochastic integrals of arbitrary multiplicity based on generalized multiple Fourier series converging in the mean. arXiv:1712.09746 [math.PR]. 2023, 144 pp. [in English].

- [13] Kuznetsov D.F. Development and application of the Fourier method to the mean-square approximation of iterated Ito and Stratonovich stochastic integrals. arXiv:1712.08991 [math.PR]. 2023, 58 pp. [in English].
- [14] Kuznetsov D.F. Exact calculation of the mean-square error in the method of approximation of iterated Ito stochastic integrals based on generalized multiple Fourier series. arXiv:1801.01079 [math.PR]. 2023, 71 pp. [in English].
- [15] Kuznetsov D.F. Mean-square approximation of iterated Ito and Stratonovich stochastic integrals of multiplicities 1 to 6 from the Taylor-Ito and Taylor-Stratonovich expansions using Legendre polynomials. arXiv:1801.00231 [math.PR]. 2017, 106 pp. [in English].
- [16] Kuznetsov D.F. Expansions of iterated Stratonovich stochastic integrals based on generalized multiple Fourier series: multiplicities 1 to 6 and beyond. arXiv:1712.09516 [math.PR]. 2023, 222 pp. [in English].
- [17] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals of arbitrary multiplicity based on generalized iterated Fourier series converging pointwise. arXiv:1801.00784 [math.PR]. 2023, 80 pp. [in English].
- [18] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals of multiplicities 1 to 3 based on generalized multiple Fourier series converging in the mean: general case of series summation. arXiv:1801.01564 [math.PR]. 2018, 66 pp. [in English].
- [19] Kuznetsov D.F. Expansions of iterated Stratonovich stochastic integrals of multiplicities 1 to 4. Combined approach based on generalized multiple and iterated Fourier series. arXiv:1801.05654 [math.PR]. 2018, 46 pp. [In English].
- [20] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals of multiplicity 2. Combined approach based on generalized multiple and iterated Fourier series. arXiv:1801.07248 [math.PR]. 2018, 20 pp. [In English].
- [21] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals of fifth and sixth multiplicity based on generalized multiple Fourier series. arXiv:1802.00643 [math.PR]. 2023, 148 pp. [in English].
- [22] Kuznetsov D.F. The hypotheses on expansions of iterated Stratonovich stochastic integrals of arbitrary multiplicity and their partial proof. arXiv:1801.03195 [math.PR]. 2023, 158 pp. [in English].
- [23] Kuznetsov D.F. Comparative analysis of the efficiency of application of Legendre polynomials and trigonometric functions to the numerical integration of Ito stochastic differential equations. arXiv:1901.02345 [math.GM], 2019, 40 pp. [In English].
- [24] Kuznetsov D.F. Expansion of iterated stochastic integrals with respect to martingale Poisson measures and with respect to martingales based on generalized multiple Fourier series. arXiv:1801.06501 [math.PR]. 2018, 40 pp. [In English].
- [25] Kuznetsov D.F. To numerical modeling with strong orders 1.0, 1.5, and 2.0 of convergence for multidimensional dynamical systems with random disturbances. arXiv:1802.00888 [math.PR]. 2018, 29 pp. [In English].
- [26] Kuznetsov D.F. Numerical simulation of 2.5-set of iterated Stratonovich stochastic integrals of multiplicities 1 to 5 from the Taylor-Stratonovich expansion. arXiv:1806.10705 [math.PR]. 2018, 29 pp. [In English].
- [27] Kuznetsov D.F. Numerical simulation of 2.5-set of iterated Ito stochastic integrals of multiplicities 1 to 5 from the Taylor-Ito expansion. arXiv:1805.12527 [math.PR]. 2018, 29 pp. [In English].
- [28] Kuznetsov D.F. Explicit one-step strong numerical methods of orders 2.0 and 2.5 for Ito stochastic differential equations based on the unified Taylor-Ito and Taylor-Stratonovich expansions. arXiv: 1802.04844 [math.PR]. 2018, 37 pp. [in English].
- [29] Kuznetsov D.F. Strong numerical methods of orders 2.0, 2.5, and 3.0 for Ito stochastic differential equations based on the unified stochastic Taylor expansions and multiple Fourier-Legendre series. arXiv:1807.02190 [math.PR], 2018, 44 pp. [in English].
- [30] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals of multiplicity 2 based on double Fourier-Legendre series summarized by Pringsheim method arXiv:1801.01962 [math.PR]. 2023, 49 pp. [in English].
- [31] Kuznetsov D.F. Application of the method of approximation of iterated Ito stochastic integrals based on generalized multiple Fourier series to the high-order strong numerical methods for non-commutative semilinear stochastic partial differential equations. arXiv:1905.03724 [math.GM], 2019, 41 pp. [In English].
- [32] Kuznetsov D.F. Application of multiple Fourier-Legendre series to implementation of strong exponential Milstein and Wagner-Platen methods for non-commutative semilinear stochastic partial differential equations. arXiv:1912.02612 [math.PR], 2019, 32 pp. [In English].
- [33] Kuznetsov D.F. Integration order replacement technique for iterated Ito stochastic integrals and iterated stochastic integrals with respect to martingales. arXiv:1801.04634 [math.PR], 2018, 27 pp. [In English].
- [34] Kuznetsov D.F. Expansions of iterated Stratonovich stochastic integrals from the Taylor-Stratonovich expansion based on multiple trigonometric Fourier series. Comparison with the Milstein expansion. arXiv:1801.08862 [math.PR], 2018, 36 p. [In English].
- [35] Kuznetsov D.F. New simple method of expansion of iterated Ito stochastic integrals of multiplicity 2 based on expansion of the Brownian motion using Legendre polynomials and trigonometric functions. arXiv:1807.00409 [math.PR], 2019, 23 pp. [In English].
- [36] Kuznetsov D.F. Development and application of the Fourier method for the numerical solution of Ito stochastic differential equations. [In English]. Computational Mathematics and Mathematical Physics, 58, 7 (2018), 1058-1070. DOI: <http://doi.org/10.1134/S0965542518070096>

- [37] Kuznetsov D.F. On numerical modeling of the multidimensional dynamic systems under random perturbations with the 1.5 and 2.0 orders of strong convergence [In English]. *Automation and Remote Control*, 79, 7 (2018), 1240-1254. DOI: <http://doi.org/10.1134/S0005117918070056>
- [38] Kuznetsov D.F. On numerical modeling of the multidimensional dynamic systems under random perturbations with the 2.5 order of strong convergence. [In English]. *Automation and Remote Control*, 80, 5 (2019), 867-881. DOI: <http://doi.org/10.1134/S0005117919050060>
- [39] Kuznetsov D.F. Comparative analysis of the efficiency of application of Legendre polynomials and trigonometric functions to the numerical integration of Itô stochastic differential equations. [In English]. *Computational Mathematics and Mathematical Physics*, 59, 8 (2019), 1236-1250. DOI: <http://doi.org/10.1134/S0965542519080116>
- [40] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals based on generalized multiple Fourier series. [In English]. *Ufa Mathematical Journal*, 11, 4 (2019), 49-77. DOI: <http://doi.org/10.13108/2019-11-4-49> Available at: http://matem.anrb.ru/en/article?art_id=604
- [41] Kuznetsov D.F. Explicit one-step numerical method with the order of strong convergence 2.5 for Ito stochastic differential equations with multidimensional nonadditive noise, based on the Taylor-Stratonovich expansion. *Computational Mathematics and Mathematical Physics*, 60, 3 (2020), 379-389. DOI: <http://doi.org/10.1134/S0965542520030100>
- [42] Kuznetsov D.F. Expansion of multiple Stratonovich stochastic integrals of second multiplicity, based on double Fourier-Legendre series summarized by Prinsheim method [In Russian]. *Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online)*, 1 (2018), 1-34. Available at: <http://diffjournal.spbu.ru/EN/numbers/2018.1/article.1.1.html>
- [43] Kuznetsov D.F. Application of the method of approximation of iterated stochastic Ito integrals based on generalized multiple Fourier series to the high-order strong numerical methods for non-commutative semilinear stochastic partial differential equations [In English]. *Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online)*, 3 (2019), 18-62. Available at: <http://diffjournal.spbu.ru/EN/numbers/2019.3/article.1.2.html>
- [44] Kuznetsov D.F. Mean square approximation of solutions of stochastic differential equations using Legendres polynomials. [In English]. *Journal of Automation and Information Sciences (Begell House)*, 2000, 32 (Issue 12), 69-86. DOI: <http://doi.org/10.1615/JAutomatInfScien.v32.i12.80>
- [45] Kuznetsov D.F. New representations of explicit one-step numerical methods for jump-diffusion stochastic differential equations. [In English]. *Computational Mathematics and Mathematical Physics*, 41, 6 (2001), 874-888. Available at: <http://www.sde-kuznetsov.spb.ru/01b.pdf>
- [46] Kuznetsov D.F. *Numerical Integration of Stochastic Differential Equations. 2.* [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2006, 764 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-227> Available at: <http://www.sde-kuznetsov.spb.ru/06.pdf> (ISBN 5-7422-1191-0)
- [47] Kuznetsov D.F. *Stochastic Differential Equations: Theory and Practice of Numerical Solution.* With MatLab programs, 1st Edition. [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2007, 778 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-228> Available at: <http://www.sde-kuznetsov.spb.ru/07b.pdf> (ISBN 5-7422-1394-8)
- [48] Kuznetsov D.F. *Stochastic Differential Equations: Theory and Practice of Numerical Solution.* With MatLab programs, 2nd Edition. [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2007, XXXII+770 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-229> Available at: <http://www.sde-kuznetsov.spb.ru/07a.pdf> (ISBN 5-7422-1439-1)
- [49] Kuznetsov D.F. *Stochastic Differential Equations: Theory and Practice of Numerical Solution.* With MatLab programs, 3rd Edition. [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2009, XXXIV+768 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-230> Available at: <http://www.sde-kuznetsov.spb.ru/09.pdf> (ISBN 978-5-7422-2132-6)
- [50] Kuznetsov D.F. *Stochastic Differential Equations: Theory and Practice of Numerical Solution.* With MatLab programs. 4th Edition. [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2010, XXX+786 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-231> Available at: <http://www.sde-kuznetsov.spb.ru/10.pdf> (ISBN 978-5-7422-2448-8)
- [51] Kuznetsov D.F. Multiple Stochastic Ito and Stratonovich Integrals and Multiple Fourier Series. [In Russian]. *Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online)*, 3 (2010), A.1-A.257. DOI: <http://doi.org/10.18720/SPBPU/2/z17-7> Available at: <http://diffjournal.spbu.ru/EN/numbers/2010.3/article.2.1.html>
- [52] Kuznetsov D.F. *Strong Approximation of Multiple Ito and Stratonovich Stochastic Integrals: Multiple Fourier Series Approach.* 1st Edition. [In English]. Polytechnical University Publishing House, Saint-Petersburg, 2011, 250 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-232> Available at: <http://www.sde-kuznetsov.spb.ru/11b.pdf> (ISBN 978-5-7422-2988-9)
- [53] Kuznetsov D.F. *Strong Approximation of Multiple Ito and Stratonovich Stochastic Integrals: Multiple Fourier Series Approach.* 2nd Edition. [In English]. Polytechnical University Publishing House, Saint-Petersburg, 2011,

- 284 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-233> Available at: <http://www.sde-kuznetsov.spb.ru/11a.pdf> (ISBN 978-5-7422-3162-2)
- [54] Kuznetsov D.F. Multiple Ito and Stratonovich stochastic integrals: approximations, properties, formulas. [In English]. Polytechnical University Publishing House, Saint-Petersburg, 2013, 382 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-234> Available at: <http://www.sde-kuznetsov.spb.ru/13.pdf> (ISBN 978-5-7422-3973-4)
- [55] Kuznetsov D.F. Multiple Ito and Stratonovich Stochastic Integrals: Fourier-Legendre and Trigonometric Expansions, Approximations, Formulas. [In English]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 1 (2017), A.1-A.385. DOI: <http://doi.org/10.18720/SPBPU/2/z17-3> Available at: <http://diffjournal.spbu.ru/EN/numbers/2017.1/article.2.1.html>
- [56] Kuznetsov D.F. Stochastic Differential Equations: Theory and Practice of Numerical Solution. With Programs on MATLAB, 5th Edition. [In Russian]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 2 (2017), A.1-A.1000. DOI: <http://doi.org/10.18720/SPBPU/2/z17-4> Available at: <http://diffjournal.spbu.ru/EN/numbers/2017.2/article.2.1.html>
- [57] Kuznetsov D.F. Stochastic Differential Equations: Theory and Practice of Numerical Solution. With MATLAB Programs, 6th Edition. [In Russian]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 4 (2018), A.1-A.1073. Available at: <http://diffjournal.spbu.ru/EN/numbers/2018.4/article.2.1.html>
- [58] Kuznetsov D.F. Strong Approximation of Iterated Itô and Stratonovich Stochastic Integrals Based on Generalized Multiple Fourier Series. Application to Numerical Solution of Itô SDEs and Semilinear SPDEs. arXiv:2003.14184v45 [math.PR], 2023, 996 pp. [In English].
- [59] Kuznetsov D.F. Strong Approximation of Iterated Itô and Stratonovich Stochastic Integrals Based on Generalized Multiple Fourier Series. Application to Numerical Solution of Itô SDEs and Semilinear SPDEs. [In English]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 4 (2020), A.1-A.606. Available at: <http://diffjournal.spbu.ru/EN/numbers/2020.4/article.1.8.html>
- [60] Kuznetsov D.F. Mean-Square Approximation of Iterated Itô and Stratonovich Stochastic Integrals Based on Generalized Multiple Fourier Series. Application to Numerical Integration of Itô SDEs and Semilinear SPDEs. [In English]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 4 (2021), A.1-A.788. Available at: <http://diffjournal.spbu.ru/EN/numbers/2021.4/article.1.9.html>
- [61] Kuznetsov D.F. Mean-Square Approximation of Iterated Itô and Stratonovich Stochastic Integrals Based on Generalized Multiple Fourier Series. Application to Numerical Integration of Itô SDEs and Semilinear SPDEs (Third Edition). [In English]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 1 (2023), A.1-A.947. Available at: <http://diffjournal.spbu.ru/EN/numbers/2023.1/article.1.10.html>
- [62] Kuznetsov D.F. A new proof of the expansion of iterated Ito stochastic integrals with respect to the components of a multidimensional Wiener process based on generalized multiple Fourier series and Hermite polynomials. arXiv:2307.11006 [math.PR], 2023, 58 pp. [In English].
- [63] Kuznetsov D.F. A new approach to the series expansion of iterated Stratonovich stochastic integrals of arbitrary multiplicity with respect to components of the multidimensional Wiener process. [In English]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 2 (2022), 83-186. Available at: <http://diffjournal.spbu.ru/EN/numbers/2022.2/article.1.6.html>
- [64] Kuznetsov, D.F. A new approach to the series expansion of iterated Stratonovich stochastic integrals of arbitrary multiplicity with respect to components of the multidimensional Wiener process. II. [In English]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 4 (2022), 135-194. Available at: <http://diffjournal.spbu.ru/EN/numbers/2022.4/article.1.9.html>
- [65] Kuznetsov D. F. A method of expansion and approximation of repeated stochastic Stratonovich integrals based on multiple Fourier series on full orthonormal systems. [In Russian]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 1 (1997), 18-77. Available at: <http://diffjournal.spbu.ru/EN/numbers/1997.1/article.1.2.html>
- [66] Kuznetsov D.F. Theorems about integration order replacement in iterated stochastic integrals. Dep. VINITI. 3607-V97, 1997, 31 pp.
- [67] Kuznetsov D.F. Problems of the numerical analysis of Ito stochastic differential equations. [In Russian]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 1 (1998), 66-367. Available at: <http://diffjournal.spbu.ru/EN/numbers/1998.1/article.1.3.html> Hard Cover Edition: 1998, SPbGTU Publishing House, 204 pp. (ISBN 5-7422-0045-5)
- [68] Kuznetsov D.F. Integration order replacement in iterated stochastic integrals with respect to martingale. Preprint. Saint-Petersburg: SPbGTU Publ., 1999, 11 pp. Available at: <http://www.sde-kuznetsov.spb.ru/99c.pdf>

- [69] Stratonovich R.L. Conditional Markov Processes and Their Application to the Theory of Optimal Control [in Russian], Moscow University Press, Moscow, 1966, 320 pp.
- [70] Jentzen A. and Röckner M. A Milstein scheme for SPDEs. *Foundations Comp. Math.* 15, 2 (2015), 313-362.
- [71] Becker S., Jentzen A. and Kloeden P.E. An exponential Wagner-Platen type scheme for SPDEs. *SIAM J. Numer. Anal.* 54, 4 (2016), 2389-2426.
- [72] Jentzen A. and Röckner M. Regularity analysis of stochastic partial differential equations with nonlinear multiplicative trace class noise. *J. Differ. Eq.* 252, 1 (2012), 114-136.
- [73] Da Prato G., Jentzen A. and Röckner M. A mild Itô formula for SPDEs. *arXiv:1009.3526 [math.PR]* (2012), 39 pp.
- [74] Rybakov K.A. Orthogonal expansion of multiple Itô stochastic integrals. *Differential Equations and Control Processes*, 3 (2021), 109-140. Available at: <http://diffjournal.spbu.ru/EN/numbers/2021.3/article.1.8.html>
- [75] Kuznetsov M.D., Kuznetsov D.F. SDE-MATH: A software package for the implementation of strong high-order numerical methods for Ito SDEs with multidimensional non-commutative noise based on multiple Fourier-Legendre series. [In English]. *Electronic Journal "Differential Equations and Control Processes"* ISSN 1817-2172 (online), 1 (2021), 93-422. Available at: <http://diffjournal.spbu.ru/EN/numbers/2021.1/article.1.5.html>
- [76] Kuznetsov M.D., Kuznetsov D.F. Implementation of strong numerical methods of orders 0.5, 1.0, 1.5, 2.0, 2.5, and 3.0 for Ito SDEs with non-commutative noise based on the unified Taylor-Ito and Taylor-Stratonovich expansions and multiple Fourier-Legendre series. *arXiv:2009.14011 [math.PR]*, 2020, 342 pp. [In English].
- [77] Rybakov, K.A. On traces of linear operators with symmetrized Volterra-type kernels. *Symmetry*, 15, 1821 (2023), 1-18. DOI: <http://doi.org/10.3390/sym15101821>
- [78] Wong E., Zakai M. On the convergence of ordinary integrals to stochastic integrals. *Ann. Math. Stat.*, 5, 36 (1965), 1560-1564.
- [79] Wong E., Zakai M. On the relation between ordinary and stochastic differential equations. *Int. J. Eng. Sci.*, 3 (1965), 213-229.
- [80] Ikeda N., Watanabe S. *Stochastic Differential Equations and Diffusion Processes*. 2nd Edition. North-Holland Publishing Company, Amsterdam, Oxford, New-York, 1989. 555 pp.
- [81] Liptser R.Sh., Shirjaev A.N. *Statistics of Stochastic Processes: Nonlinear Filtering and Related Problems*. [In Russian]. Moscow, Nauka, 1974. 696 pp.
- [82] Luo W. Wiener chaos expansion and numerical solutions of stochastic partial differential equations. PhD thesis, California Inst. of Technology, 2006, 225 pp.
- [83] Kloeden P.E., Platen E., Wright I.W. The approximation of multiple stochastic integrals. *Stoch. Anal. Appl.* 10, 4 (1992), 431-441.
- [84] Platen E., Bruti-Liberati N. *Numerical Solution of Stochastic Differential Equations with Jumps in Finance*. Springer, Berlin-Heidelberg, 2010. 868 pp.

DMITRIY FELIKSOVICH KUZNETSOV
PETER THE GREAT SAINT-PETERSBURG POLYTECHNIC UNIVERSITY,
POLYTECHNICHESKAYA UL., 29,
195251, SAINT-PETERSBURG, RUSSIA
Email address: sde_kuznetsov@inbox.ru