## LIMIT THEOREMS FOR A STABLE SAUSAGE

WOJCIECH CYGAN, NIKOLA SANDRIĆ, AND STJEPAN ŠEBEK

Abstract. In this article, we study fluctuations of the volume of a stable sausage defined via a d-dimensional rotationally invariant  $\alpha$ -stable process. As the main results, we establish a functional central limit theorem (in the case when  $d/\alpha > 3/2$ ) with a standard one-dimensional Brownian motion in the limit, and Khintchine's and Chung's laws of the iterated logarithm (in the case when  $d/\alpha > 9/5$ ).

### 1. INTRODUCTION

Let  $X = \{X_t\}_{t\geq 0}$  be a Lévy process in  $\mathbb{R}^d$  defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . A Lévy sausage associated with the process X and a given compact set  $K \subset \mathbb{R}^d$ , on the time interval [s, t],  $0 \leq s \leq t$ , is the random set defined as

$$
\mathcal{S}^K[s,t] = \bigcup_{s \le u \le t} \{X_u + K\}.
$$

If  $s = 0$  we use the notation  $S_t^K = S^K[0, t]$ . Let  $\lambda(dx)$  be the Lebesgue measure on  $\mathbb{R}^d$ and let us denote by

<span id="page-0-1"></span>
$$
\mathcal{V}^K[s,t] \,=\, \lambda(\mathcal{S}^K[s,t])
$$

the volume of the Lévy sausage  $S^K[s, t]$  (we write  $\mathcal{V}_t^K = \lambda(S_t^K)$ ). Already Spitzer [\[28\]](#page-33-0) linked  $\mathcal{V}_t^K$  with the first hitting time  $\tau_K = \inf\{s \geq 0 : X_s \in K\}$  via the identity

(1.1) 
$$
\mathbb{E}[\mathcal{V}_t^K] = \int_{\mathbb{R}^d} \mathbb{P}_x(\tau_K \le t) \, \mathrm{d}x, \qquad t \ge 0,
$$

where  $\mathbb{P}_x$  is the probability measure related to the process X started at  $x \in \mathbb{R}^d$ . Port and Stone  $[22,$  Theorem 11.1] proved that if X is transient then

(1.2) 
$$
\lim_{t \nearrow \infty} \frac{\mathbb{E}[\mathcal{V}_t^K]}{t} = \text{Cap}(K),
$$

where  $\text{Cap}(K)$  is the capacity of K associated with the process X. Hawkes [\[11\]](#page-32-0) observed that in view of the subadditivity of the process  $\{\mathcal{V}_t^K\}_{t \geq 0}$ , that is,

<span id="page-0-2"></span><span id="page-0-0"></span>
$$
\mathcal{V}_{s+t}^K \leq \mathcal{V}_s^K + \mathcal{V}^K[s, s+t], \qquad s, t \geq 0,
$$

eq. [\(1.2\)](#page-0-0) combined with Kingman's ergodic theorem (cf. [\[16,](#page-33-2) Theorem Ch. I, 5.6]) and [\[15,](#page-33-3) Proposition 3.12] implies the following strong law of large numbers

(1.3) 
$$
\lim_{t \nearrow \infty} \frac{\mathcal{V}_t^K}{t} = \text{Cap}(K) \qquad \mathbb{P}\text{-a.s.}
$$

More satisfactory limit theorems for the volume of a Lévy sausage are known if  $X$  is a standard Brownian motion. In this case  $S_t^K$  is called a Wiener sausage, and there is a vast amount of literature concerning its asymptotic behavior. The pioneering work [\[6\]](#page-32-1) was due to Donsker and Varadhan were they established a large deviation principle for the volume of a Wiener sausage. Their result was extended by Eisele and Lang [\[8\]](#page-32-2) to the

<sup>2010</sup> *Mathematics Subject Classification.* 60F05, 60G52, 60F17.

*Key words and phrases.* functional central limit theorem, law of the iterated logarithm, stable process, stable sausage.

case when the driving process is a standard Brownian motion with drift, and to a class of elliptic diffusions by Sznitman  $[29]$ , while Okura investigated similar questions for a certain class of symmetric Lévy processes. Le Gall  $[17]$  obtained a central limit theorem for the volume of a Wiener sausage in dimensions  $d \geq 2$ , with different normalizing sequences and distributions in the limit for  $d = 2$ ,  $d = 3$  and  $d \geq 4$ , respectively. More recently, van den Berg, Bolthausen and den Hollander [\[34\]](#page-33-6) studied the problem of intersections of two Wiener sausages, see also [\[31\]](#page-33-7), [\[32\]](#page-33-8) and [\[33\]](#page-33-9). For further limit theorems for the volume of a Wiener sausage see  $\lbrack 3\rbrack$ ,  $\lbrack 12\rbrack$  and  $\lbrack 35\rbrack$ . We remark that first studies on a Wiener sausage were motivated by its applications in physics  $[14]$ . We refer the reader to the book by Simon [\[27\]](#page-33-12) for a comprehensive discussion on this topic.

In the present article, we focus on the limit behavior of the volume of a stable sausage, that is, a Lévy sausage corresponding to a stable Lévy process. Asymptotic behavior of stable sausages has not been extensively studied yet. In the seminal paper [\[6\]](#page-32-1) Donsker and Varadhan obtained a large deviation principle for the volume of a stable sausage. Some other works were concerned with the expansion of the expected volume of a stable sausage. More precisely, Getoor [\[9\]](#page-32-5) proved eq. [\(1.2\)](#page-0-0) for rotationally invariant  $\alpha$ -stable processes with  $d > \alpha$  and for any compact set K. He also investigated the first order asymptotics of the difference  $\mathbb{E}[\mathcal{V}_t^K] - t \operatorname{Cap}(K)$ , whose form depends on the value of the ratio  $d/\alpha$ , see [\[9,](#page-32-5) Theorem 2]. The second order terms in this expansion were found by Port [\[21\]](#page-33-13) for all strictly stable processes satisfying some extra assumptions. In [\[24\]](#page-33-14) Rosen established asymptotic expansions for the volume of a stable sausage in the plane with the coefficients represented by n-fold self-intersections of the stable process. In this article, we obtain a central limit theorem for the volume of a stable sausage. We then apply this result to study convergence of the volume process in the Skorohod space, and establish the corresponding functional central limit theorem. Finally, we also obtain Khintchine's and Chung's laws of the iterated logarithm for this process.

Before we formulate our results, we briefly recall some basic notation from the potential theory of stable processes. Let  $X$  be a rotationally invariant stable Lévy process of index  $\alpha \in (0, 2]$ , that is, a Lévy process whose bounded continuous transition density  $p(t, x)$  is uniquely determined by the Fourier transform

$$
e^{-t|\xi|^{\alpha}} = \int_{\mathbb{R}^d} e^{i(x,\xi)} p(t,x) dx,
$$

where  $(x,\xi)$  stands for the inner product in  $\mathbb{R}^d$ ,  $|x| = (x,x)^{1/2}$  is the Euclidean norm, and  $dx = \lambda(dx)$ . We assume that X is transient, which holds if (and only if)  $d > \alpha$ . Its Green function is then given by  $G(x) = \int_0^\infty p(t, x) dt$ . Let  $\mathfrak{B}(\mathbb{R}^d)$  denote the family of all Borel subsets of  $\mathbb{R}^d$ . For each  $B \in \mathfrak{B}(\mathbb{R}^d)$  there exists a unique Borel measure  $\mu_B(\mathrm{d}x)$ supported on  $B \in \mathfrak{B}(\mathbb{R}^d)$  such that

(1.4) 
$$
\mathbb{P}_x(\tau_B < \infty) = \int_{\mathbb{R}^d} G(x - y) \,\mu_B(\mathrm{d}y).
$$

The measure  $\mu_B(dx)$  is called the equilibrium measure of B, and its capacity Cap(B) is defined as the total mass of  $\mu_B(dx)$ , that is,  $Cap(B) = \mu_B(B)$ . We denote by  $\mathcal{B}(x, r)$  the closed Euclidean ball centered at  $x \in \mathbb{R}^d$  of radius  $r > 0$ . In the case when  $r = 1$  and  $x = 0$ , we write  $B = B(0, 1)$ . If  $B = B(0, r)$  then the measure  $\mu_B(dy)$  has a density which is proportional to  $(r - |y|^2)^{-\alpha/2}$ . In particular, we have (see for instance [\[30\]](#page-33-15))

<span id="page-1-0"></span>
$$
Cap(\mathcal{B}) = \frac{\Gamma(d/2)}{\Gamma(\alpha/2)\Gamma(1 + (d - \alpha)/2)}.
$$

In the case when  $K = \mathcal{B}$ , we simply write  $\mathcal{V}_t$  instead of  $\mathcal{V}_t^{\mathcal{B}}$  $t^{\mathcal{B}}$  (and similarly  $\mathcal{S}_t$  for  $\mathcal{S}_t^{\mathcal{B}}$  $\binom{\mathcal{B}}{t}$ . Let  $\mathcal{N}(0, 1)$  denote the Gaussian random variable with mean zero and variance one. Our

central limit theorem (see Theorem [2.1\)](#page-3-0) for the volume of a stable sausage asserts that if  $d/\alpha > 3/2$  then there exists a constant  $\sigma = \sigma(d, \alpha) > 0$  such that

<span id="page-2-0"></span>(1.5) 
$$
\frac{\mathcal{V}_t - t \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{t}} \xrightarrow[t \to \infty]{(\mathrm{d})} \mathcal{N}(0,1),
$$

where convergence holds in distribution. The cornerstone of the proof of eq.  $(1.5)$  is to represent  $V_t$  as a sum of independent random variables plus an error term. For this we use inclusion-exclusion formula together with the Markov property and rotational invariance of the process X. More precisely, for  $t, s \geq 0$ , we have

<span id="page-2-2"></span>(1.6) 
$$
\mathcal{V}_{t+s} = \lambda \big( \mathcal{S}_t \cup \mathcal{S}[t, t+s] \big) = \lambda \big( (\mathcal{S}_t - X_t) \cup (\mathcal{S}[t, t+s] - X_t) \big) = \mathcal{V}_t^{(1)} + \mathcal{V}_s^{(2)} - \lambda \big( \mathcal{S}_t^{(1)} \cap \mathcal{S}_s^{(2)} \big),
$$

where  $V_t^{(1)}$  and  $V_s^{(2)}$   $(S_t^{(1)}$  and  $S_s^{(2)})$  are independent and have the same law as  $V_t$  and  $\mathcal{V}_s$  ( $\mathcal{S}_t$  and  $\mathcal{S}_s$ ), respectively. This decomposition allows us to apply the Lindeberg-Feller central limit theorem in the present context. The first key step is to find estimates for the error term  $\lambda(\mathcal{S}_t^{(1)} \cap \mathcal{S}_s^{(2)})$ , which we give in Section [2.1.](#page-3-1) The second step is to control the variance of the volume of a stable sausage which is achieved in Section [2.2.](#page-5-0)

Let us emphasize that the present article has been mainly inspired by Le Gall's work [\[17\]](#page-33-5) where he studied fluctuations of the volume of a Wiener sausage (the case  $\alpha = 2$ ). Among other results, he established the central limit theorem in eq. [\(1.5\)](#page-2-0) for dimensions  $d \geq 4$ . Still another source of motivation was the article [\[18\]](#page-33-16) by Le Gall and Rosen where they proved a corresponding central limit theorem for the range of stable random walks and mentioned that it is plausible that similar result holds for stable sausages, see  $[18,$ Page 654]. Both of these articles were also concerned with the lower-dimensional case  $d < 4$  and  $d/\alpha \leq 3/2$ , respectively. In the present article we are only interested in the case when  $d/\alpha > 3/2$ , and we postpone the study of the remaining values of the ratio  $d/\alpha$ to follow-up articles.

As an application of eq. [\(1.5\)](#page-2-0) we obtain a functional central limit theorem (see Theorem [3.1\)](#page-14-0) which states that under the same assumptions, and with the same constant  $\sigma > 0$ ,

<span id="page-2-1"></span>(1.7) 
$$
\left\{ \frac{\mathcal{V}_{nt} - nt \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{n}} \right\}_{t \geq 0} \xrightarrow[n \nearrow \infty]{(J_1)} \{W_t\}_{t \geq 0}.
$$

Here, convergence holds in the Skorohod space  $\mathcal{D}([0,\infty),\mathbb{R})$  endowed with the J<sub>1</sub> topology, and  $\{W_t\}_{t\geq0}$  is a standard Brownian motion in R. The proof of eq. [\(1.7\)](#page-2-1) is performed according to a general two-step scheme: (i) convergence of finite-dimensional distributions, which follows from eq.  $(1.5)$ ; (ii) tightness, which we investigate by employing the wellknown Aldous criterion, see Section [3](#page-14-1) for details.

It is remarkable that results in eqs. [\(1.5\)](#page-2-0) and [\(1.7\)](#page-2-1) correspond to analogous results for the range (and its capacity) of stable random walks on the integer lattice  $\mathbb{Z}^d$  which we discussed in [\[4\]](#page-32-6) and [\[5\]](#page-32-7), respectively.

We finally use eq.  $(1.5)$  to study growth of the paths of the volume of the stable sausage. In the case when  $d/\alpha > 9/5$ , we prove Khintchine's law of the iterated logarithm

<span id="page-2-3"></span>(1.8) 
$$
\liminf_{t \nearrow \infty} \frac{\mathcal{V}_t - t \operatorname{Cap}(\mathcal{B})}{\sqrt{2\sigma^2 t \log \log t}} = -1 \quad \text{and} \quad \limsup_{t \nearrow \infty} \frac{\mathcal{V}_t - t \operatorname{Cap}(\mathcal{B})}{\sqrt{2\sigma^2 t \log \log t}} = 1 \quad \mathbb{P}\text{-a.s.},
$$

as well as Chung's law of the iterated logarithm

<span id="page-2-4"></span>(1.9) 
$$
\liminf_{t \nearrow \infty} \frac{\sup_{0 \le s \le t} |\mathcal{V}_s - s \operatorname{Cap}(\mathcal{B})|}{\sqrt{\sigma^2 t / \log \log t}} = \frac{\pi}{\sqrt{8}} \qquad \mathbb{P}\text{-a.s.}
$$

Our proof is based on the approach which was developed by Jain and Pruitt [\[13\]](#page-32-8) (see also [\[1\]](#page-32-9)) in the context of the range of random walks and then successfully applied in [\[35\]](#page-33-10) to obtain Khintchine's and Chung's laws of the iterated logarithm for a Wiener sausage in dimensions  $d \geq 4$ .

The rest of the paper is organized as follows. In Section [2](#page-3-2) we prove the central limit theorem in eq.  $(1.5)$ . For this we first deal with the error terms derived from eq.  $(1.6)$ , and in the second part we show that the variance of the volume of a stable sausage behaves linearly at infinity. Section [3](#page-14-1) is devoted to the proof of eq.  $(1.7)$ , and in Section [4](#page-16-0) we prove eqs.  $(1.8)$  and  $(1.9)$ . In Section [5](#page-20-0) we present the proofs of some technical results which we need in the course of the study.

#### 2. Central limit theorem

<span id="page-3-2"></span>The goal of this section is to prove the following central limit theorem. We assume that X is a rotationally invariant stable Lévy process in  $\mathbb{R}^d$  of index  $\alpha \in (0, 2]$  satisfying  $d/\alpha > 3/2$ .

<span id="page-3-0"></span>**Theorem 2.1.** Under the above assumptions, there exists a constant  $\sigma = \sigma(d, \alpha) > 0$ such that

$$
\frac{\mathcal{V}_t - t \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{t}} \xrightarrow[t \to \infty]{(\mathrm{d})} \mathcal{N}(0,1).
$$

We remark that Theorem [2.1](#page-3-0) holds for any closed ball  $\mathcal{B}(x, r)$ , with a possibly different constant  $\sigma > 0$ . Moreover, as indicated by eq. [\(1.2\)](#page-0-0), the statement of the theorem remains valid if we replace the term  $t \operatorname{Cap}(\mathcal{B})$  with  $\mathbb{E}[\mathcal{V}_t].$ 

Before we embark on the proof of the theorem, which is given at the end of the section, we first need to find satisfactory estimates for the error term in decomposition eq. [\(1.6\)](#page-2-2). Next step is to investigate the variance  $\text{Var}(\mathcal{V}_t)$  of the volume of a stable sausage and to show that it behaves as  $\sigma t$  at infinity.

<span id="page-3-1"></span>2.1. Error term estimates. We assume that X is defined on the canonical space  $\Omega =$  $\mathcal{D}([0,\infty),\mathbb{R}^d)$  of all càdlàg functions  $\omega : [0,\infty) \to \mathbb{R}^d$ . It is endowed with the Borel  $\sigma$ -algebra  $\mathcal F$  generated by the Skorokhod J<sub>1</sub> topology. Then X is understood as the coordinate process, that is,  $X_t(\omega) = \omega(t)$ , and the shift operator  $\theta_t$  acting on  $\Omega$  is defined by

$$
\theta_t \,\omega(s) \,=\, \omega(t+s), \qquad t,s \geq 0.
$$

In what follows we use notation

$$
\mathcal{S}^K[t,\infty) = \bigcup_{s \ge t} \{X_s + K\}, \qquad t \ge 0.
$$

We also write  $S_{\infty}^{K} = S^{K}[0, \infty)$ ,  $\mathcal{V}_{\infty}^{K} = \lambda(\mathcal{S}_{\infty}^{K})$  and  $\mathcal{V}^{K}[t, \infty) = \lambda(\mathcal{S}^{K}[t, \infty))$ ,  $t \geq 0$ . We start with a lemma which enables us to represent the expected volume of the intersection of two sausages in terms of the difference  $\mathbb{E}[\mathcal{V}_t^K] - t \operatorname{Cap}(K)$ .

<span id="page-3-3"></span>**Lemma 2.2.** For any compact set K and all  $t \geq 0$  it holds

$$
\mathbb{E}[\mathcal{V}_t^K] - t \operatorname{Cap}(K) = \mathbb{E}[\lambda(\mathcal{S}_t^K \cap \mathcal{S}^K[t,\infty))].
$$

Proof. We clearly have

$$
\mathcal{V}_{\infty}^K = \mathcal{V}_{t}^K + \mathcal{V}^K[t, \infty) - \lambda(\mathcal{S}_{t}^K \cap \mathcal{S}^K[t, \infty))
$$

which implies

$$
\lambda(\mathcal{S}_t^K \cap \mathcal{S}^K[t,\infty)) = \mathcal{V}_t^K - \lambda(\mathcal{S}_t^K \setminus \mathcal{S}^K[t,\infty)).
$$

Hence, it suffices to show that

<span id="page-4-5"></span>(2.1) 
$$
\mathbb{E}[\lambda(\mathcal{S}_t^K \setminus \mathcal{S}^K[t,\infty)] = t \operatorname{Cap}(K), \qquad t \ge 0.
$$

By rotational invariance of the process X we have

$$
\mathbb{E}[\lambda(\mathcal{S}_t^K \setminus \mathcal{S}^K[t,\infty)] = \int_{\mathbb{R}^d} \mathbb{P}(x \in \mathcal{S}_t^K \setminus \mathcal{S}^K[t,\infty)) dx
$$
  

$$
= \int_{\mathbb{R}^d} \mathbb{P}_x \Big( \bigcup_{0 \le s \le t} \{X_s \in K\}, \bigcap_{s \ge t} \{X_s \notin K\} \Big) dx
$$
  
(2.2)  

$$
= \int_{\mathbb{R}^d} \mathbb{P}_x(0 < \eta_K \le t) dx,
$$

<span id="page-4-0"></span>where  $\eta_K$  is the last exit time of the process X from the set K, that is,

$$
\eta_K = \begin{cases} \sup\{t > 0 : X_t \in K\}, & \tau_K < \infty, \\ 0, & \tau_K = \infty. \end{cases}
$$

We observe that  $\{\eta_K > t\} = {\{\tau_K \circ \theta_t < \infty\}}$  which together with eq. [\(1.4\)](#page-1-0) yields

$$
\mathbb{P}_x(\eta_K > t) = \int_{\mathbb{R}^d} p(t, x, y) \, \mathbb{P}_y(\tau_K < \infty) \, \mathrm{d}y = \int_{\mathbb{R}^d} \int_t^{\infty} p(s, x, z) \, \mathrm{d}s \, \mu_K(\mathrm{d}z),
$$

where we used notation  $p(t, x, y) = p(t, y - x)$ . We obtain

$$
\mathbb{P}_x(0<\eta_K\leq t) = \int_{\mathbb{R}^d} \int_0^t p(s,x,y) \,ds \,\mu_K(\mathrm{d}y).
$$

This and eq. [\(2.2\)](#page-4-0) imply

$$
\mathbb{E}[\lambda(\mathcal{S}_t^K \setminus \mathcal{S}^K[t,\infty))] = \int_{\mathbb{R}^d} \int_0^t \int_{\mathbb{R}^d} p(s,y,x) \,dx \,ds \,\mu_K(\mathrm{d}y) = t \operatorname{Cap}(K),
$$
  
and the proof is finished.

<span id="page-4-4"></span>In the following lemma we show how one can easily estimate the higher moments of the expected volume of the intersection of two sausages through the first moment estimate.

**Lemma 2.3.** Let X' be an independent copy of the process X such that  $X_0 = X'_0$ , and let  $S'_t$ ,  $t \geq 0$ , denote the sausage associated with X'. Then for all  $k \in \mathbb{N}$  and  $t \geq 0$  it holds

$$
\mathbb{E}[\lambda(\mathcal{S}_t \cap \mathcal{S}'_\infty)^k] \leq 2^{k-1} (k!)^2 \left( \mathbb{E}[\lambda(\mathcal{S}_t \cap \mathcal{S}'_\infty)] \right)^k.
$$

Proof. We observe that

<span id="page-4-3"></span>
$$
(2.3) \quad \mathbb{E}[\lambda(\mathcal{S}_t \cap \mathcal{S}'_{\infty})] = \int_{\mathbb{R}^d} \mathbb{P}(x \in \mathcal{S}_t) \, \mathbb{P}(x \in \mathcal{S}'_{\infty}) \, dx = \int_{\mathbb{R}^d} \mathbb{P}_x(\tau_{\mathcal{B}} \le t) \, \mathbb{P}_x(\tau_{\mathcal{B}} < \infty) \, dx,
$$

where we used rotational invariance of X. Similarly, for  $k \geq 1$  we have

<span id="page-4-2"></span>
$$
(2.4) \quad \mathbb{E}[\lambda(\mathcal{S}_t \cap \mathcal{S}_{\infty}')^k] = \int_{\mathbb{R}^d} \cdots \int_{\mathbb{R}^d} \mathbb{P}(x_1,\ldots,x_k \in \mathcal{S}_t) \, \mathbb{P}(x_1,\ldots,x_k \in \mathcal{S}_{\infty}) \, dx_1 \cdots dx_k.
$$

By the strong Markov property, we obtain

<span id="page-4-1"></span>
$$
\mathbb{P}(x_1, ..., x_k \in \mathcal{S}_t) = \mathbb{P}(\tau_{\mathcal{B}-x_1} \le t, ..., \tau_{\mathcal{B}-x_k} \le t)
$$
\n
$$
\le k! \mathbb{P}(\tau_{\mathcal{B}-x_1} \le \cdots \le \tau_{\mathcal{B}-x_k} \le t)
$$
\n
$$
\le k! \mathbb{P}(\tau_{\mathcal{B}-x_1} \le \cdots \le \tau_{\mathcal{B}-x_{k-1}} \le t) \sup_{z \in \mathcal{B}-x_{k-1}} \mathbb{P}_z(\tau_{\mathcal{B}-x_k} \le t).
$$

For any  $w \in \mathcal{B}$  we have  $\mathcal{B} - w \subseteq \mathcal{B}(0, 2)$  and whence

$$
\mathbb{P}_{w-x_{k-1}}(\tau_{\mathcal{B}-x_k} \leq t) = \mathbb{P}_{x_k-x_{k-1}}(\tau_{\mathcal{B}-w} \leq t) \leq \mathbb{P}_{x_k-x_{k-1}}(\tau_{\mathcal{B}(0,2)} \leq t).
$$

For  $x \in \mathbb{R}^d$  and  $B \in \mathfrak{B}(\mathbb{R}^d)$  we set  $\tau_B^x = \inf\{t \geq 0 : x + X_t \in B\}$  and show that  $\tau_{\mathcal{B}(0,2)}^{x} \stackrel{\text{(d)}}{=} 2^{\alpha} \tau_{\mathcal{B}}^{x/2}$ , that is, the random variables  $\tau_{\mathcal{B}(0,2)}^{x}$  and  $2^{\alpha} \tau_{\mathcal{B}}^{x/2}$  are equal in distribution. Indeed, the easy calculation yields

$$
\tau_{\mathcal{B}(0,2)}^{x} = \inf\{t \ge 0 : |x + X_t| \le 2\} = \inf\left\{t \ge 0 : \left|\frac{x}{2} + \frac{X_t}{2}\right| \le 1\right\}
$$

$$
\stackrel{\text{(d)}}{=} \inf\left\{t \ge 0 : \left|\frac{x}{2} + X_{t/2^{\alpha}}\right| \le 1\right\} = 2^{\alpha} \inf\left\{s \ge 0 : \left|\frac{x}{2} + X_s\right| \le 1\right\} = 2^{\alpha} \tau_{\mathcal{B}}^{x/2}.
$$

This implies that for arbitrary  $w \in \mathcal{B}$ ,

$$
\mathbb{P}_{w-x_{k-1}}(\tau_{\mathcal{B}-x_k} \leq t) \leq \mathbb{P}(\tau_{\mathcal{B}(0,2)}^{x_k-x_{k-1}} \leq t) = \mathbb{P}(2^{\alpha} \tau_{\mathcal{B}}^{(x_k-x_{k-1})/2} \leq t) \leq \mathbb{P}_{\frac{x_k-x_{k-1}}{2}}(\tau_{\mathcal{B}} \leq t).
$$

In particular,

$$
\sup_{z \in \mathcal{B} - x_{k-1}} \mathbb{P}_z(\tau_{\mathcal{B} - x_k} \le t) \le \mathbb{P}_{\frac{x_k - x_{k-1}}{2}}(\tau_{\mathcal{B}} \le t).
$$

By combining this with eq.  $(2.5)$  and iterating the whole procedure, we obtain

$$
\mathbb{P}(x_1,\ldots,x_k\in\mathcal{S}_t) \leq k!\,\mathbb{P}_{x_1}(\tau_{\mathcal{B}}\leq t)\,\mathbb{P}_{\frac{x_2-x_1}{2}}(\tau_{\mathcal{B}}\leq t)\cdots\mathbb{P}_{\frac{x_k-x_{k-1}}{2}}(\tau_{\mathcal{B}}\leq t).
$$

Similarly, it follows that

<span id="page-5-4"></span>
$$
\mathbb{P}(x_1,\ldots,x_k\in\mathcal{S}_{\infty}) \leq k!\,\mathbb{P}_{x_1}(\tau_{\mathcal{B}}<\infty)\,\mathbb{P}_{\frac{x_2-x_1}{2}}(\tau_{\mathcal{B}}<\infty)\cdots\mathbb{P}_{\frac{x_k-x_{k-1}}{2}}(\tau_{\mathcal{B}}<\infty).
$$

<span id="page-5-5"></span>Applying the last two inequalities to eq.  $(2.4)$  and using eq.  $(2.3)$  finishes the proof.  $\square$ 

Corollary 2.4. In the notation of Lemma [2.3,](#page-4-4) for all  $k \in \mathbb{N}$  and  $t > 0$  large enough there is a constant  $c = c(d, \alpha) > 0$  such that

(2.6) 
$$
\mathbb{E}[\lambda(\mathcal{S}_t \cap \mathcal{S}'_\infty)^k] \leq 2^{k-1}(k!)^2 c^k h(t)^k,
$$

where the function  $h : (0, \infty) \to (0, \infty)$  is defined as

<span id="page-5-1"></span>(2.7) 
$$
h(t) = \begin{cases} 1, & d/\alpha > 2, \\ \log(t + e), & d/\alpha = 2, \\ t^{2 - d/\alpha}, & d/\alpha \in (1, 2). \end{cases}
$$

*Proof.* It follows from [\[9,](#page-32-5) Theorem 2] that there is a constant  $c(d, \alpha) > 0$  such that for  $t > 0$  large enough

(2.8) 
$$
\int_{\mathbb{R}^d} \mathbb{P}_x(\tau_{\mathcal{B}} \leq t) \, dx - t \operatorname{Cap}(\mathcal{B}) \leq c(d, \alpha) h(t),
$$

where the function  $h(t)$  is given in eq. [\(2.7\)](#page-5-1). We observe that by the Markov property and translation invariance of  $\lambda(dx)$  we have

<span id="page-5-2"></span>
$$
\mathbb{E}[\lambda(\mathcal{S}_t \cap \mathcal{S}'_\infty)] = \mathbb{E}[\lambda((\mathcal{S}_t - X_t) \cap (\mathcal{S}[t,\infty) - X_t))] = \mathbb{E}[\lambda(\mathcal{S}_t \cap \mathcal{S}[t,\infty))].
$$

Thus, eq. [\(1.1\)](#page-0-1) combined with Lemma [2.2](#page-3-3) and eq. [\(2.8\)](#page-5-2) implies the assertion for  $k = 1$ . For  $k > 1$  the result follows by Lemma [2.3.](#page-4-4)

<span id="page-5-0"></span>2.2. Variance of the volume of a stable sausage. Our aim in this section is to determine the constant  $\sigma$  in Theorem [2.1.](#page-3-0) We can easily adapt the approach of [\[4,](#page-32-6) Lemma 4.3] to the present setting and combine it with [\[10,](#page-32-10) Theorem 2] to conclude that the limit below exists

<span id="page-5-3"></span>(2.9) 
$$
\lim_{t \nearrow \infty} \frac{\text{Var}(\mathcal{V}_t)}{t} = \sigma^2.
$$

The main difficulty is to show that  $\sigma$  is strictly positive, and this is obtained in the following crucial lemma. We adapt the proof of  $[17, \text{Lemma } 4.2]$  but let us emphasize that it is a laborious task to adjust it to the case of stable processes.

<span id="page-6-4"></span>**Lemma 2.5.** The constant  $\sigma$  in eq. [\(2.9\)](#page-5-3) is strictly positive.

Proof. We split the proof into several steps and we notice that it clearly suffices to restrict our attention to integer values of the parameter  $t$ .

Step 1. We start by finding a handy decomposition of the variance  $\text{Var}(\mathcal{V}_n)$  expressed as a sum of specific random variables, see eq.  $(2.18)$ . We assume that  $X_0 = 0$ , and we set

(2.10) 
$$
\mathcal{S}[s,t] = \mathcal{S}[s,t] \setminus \mathcal{S}_s, \qquad 0 \le s \le t < \infty.
$$

For  $n, N \in \mathbb{N}$  such that  $1 \leq n \leq N$  we have

<span id="page-6-2"></span>
$$
\mathcal{V}_n + \lambda(\widehat{\mathcal{S}}[n,N]) = \mathcal{V}_N.
$$

Let  $\mathcal{F}_t = \sigma(X_s : s \le t)$ . Since  $\mathcal{V}_n$  is  $\mathcal{F}_n$ -measurable, we obtain

$$
\mathcal{V}_n + \mathbb{E}[\lambda(\widehat{\mathcal{S}}[n,N]) \mid \mathcal{F}_n] = \mathbb{E}[\mathcal{V}_N \mid \mathcal{F}_n]
$$

and by taking expectations and subtracting[1](#page-6-0)

$$
\langle \mathcal{V}_n \rangle + \langle \mathbb{E}[\lambda(\widehat{\mathcal{S}}[n,N]) | \mathcal{F}_n] \rangle = \mathbb{E}[\mathcal{V}_N | \mathcal{F}_n] - \mathbb{E}[\mathcal{V}_N].
$$

Hence

(2.11) 
$$
\langle \mathcal{V}_n \rangle + \langle \mathbb{E}[\lambda(\widehat{\mathcal{S}}[n,N]) | \mathcal{F}_n] \rangle = \sum_{k=1}^n U_k^N,
$$

where

<span id="page-6-1"></span>
$$
U_k^N = \mathbb{E}[\mathcal{V}_N | \mathcal{F}_k] - \mathbb{E}[\mathcal{V}_N | \mathcal{F}_{k-1}].
$$

We first discuss the second term on the left-hand side of eq.  $(2.11)$ . We claim that

$$
(2.12) \qquad \langle \mathbb{E}[\lambda(\widehat{\mathcal{S}}[n,N]) \mid \mathcal{F}_n] \rangle = -\langle \mathbb{E}[\lambda(\mathcal{S}[n,N] \cap \mathcal{S}_n) \mid \mathcal{F}_n] \rangle.
$$

Indeed, by eq.  $(2.10)$  we have

<span id="page-6-3"></span>
$$
\lambda(\widehat{\mathcal{S}}[n,N]) = \lambda(\mathcal{S}[n,N]) - \lambda(\mathcal{S}[n,N] \cap \mathcal{S}_n)
$$

and the independence of the increments of the process X implies that

$$
\mathbb{E}[\lambda(\widehat{\mathcal{S}}[n,N]) \mid \mathcal{F}_n] = \mathbb{E}[\lambda(\mathcal{S}[n,N]) | \mathcal{F}_n] - \mathbb{E}[\lambda(\mathcal{S}[n,N] \cap \mathcal{S}_n) \mid \mathcal{F}_n]
$$
  
\n
$$
= \mathbb{E}[\lambda(\mathcal{S}[n,N] - X_n) | \mathcal{F}_n] - \mathbb{E}[\lambda(\mathcal{S}[n,N] \cap \mathcal{S}_n) \mid \mathcal{F}_n]
$$
  
\n
$$
= \mathbb{E}[\lambda(\mathcal{S}[n,N] - X_n)] - \mathbb{E}[\lambda(\mathcal{S}[n,N] \cap \mathcal{S}_n) \mid \mathcal{F}_n]
$$
  
\n
$$
= \mathbb{E}[\lambda(\mathcal{S}[n,N])] - \mathbb{E}[\lambda(\mathcal{S}[n,N] \cap \mathcal{S}_n) \mid \mathcal{F}_n].
$$

Taking expectation and then subtracting the two relations yields eq. [\(2.12\)](#page-6-3).

Next we deal with the random variables  $U_k^N$  for  $k = 1, ..., N$ . By the independence of the increments of the process X, we obtain

$$
U_k^N = \mathbb{E}[\lambda(\mathcal{S}_k) + \lambda(\mathcal{S}[k, N]) - \lambda(\mathcal{S}_k \cap \mathcal{S}[k, N]) | \mathcal{F}_k]
$$
  
\n
$$
- \mathbb{E}[\lambda(\mathcal{S}_{k-1}) + \lambda(\mathcal{S}[k-1, N-1]) - \lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1, N-1]) | \mathcal{F}_{k-1}]
$$
  
\n
$$
- \mathbb{E}[\lambda(\widehat{\mathcal{S}}[N-1, N]) | \mathcal{F}_{k-1}]
$$
  
\n
$$
= \lambda(\mathcal{S}_k) - \lambda(\mathcal{S}_{k-1}) + \mathbb{E}[\lambda(\mathcal{S}[k, N])] - \mathbb{E}[\lambda(\mathcal{S}[k-1, N-1])]
$$
  
\n
$$
- \mathbb{E}[\lambda(\widehat{\mathcal{S}}[N-1, N]) | \mathcal{F}_{k-1}] + \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1, N-1]) | \mathcal{F}_{k-1}]
$$

<span id="page-6-0"></span><sup>1</sup>For any random variable  $Y \in L^1$  we write  $\langle Y \rangle = Y - \mathbb{E}[Y]$ .

$$
- \mathbb{E}[\lambda(\mathcal{S}_k \cap \mathcal{S}[k,N]) | \mathcal{F}_k]
$$
  
=  $\lambda(\widehat{\mathcal{S}}[k-1,k]) - \mathbb{E}[\lambda(\widehat{\mathcal{S}}[N-1,N]) | \mathcal{F}_{k-1}] + \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,N-1]) | \mathcal{F}_{k-1}]$   
-  $\mathbb{E}[\lambda(\mathcal{S}_k \cap \mathcal{S}[k,N]) | \mathcal{F}_k].$ 

Let  $\mathcal{F}_{s,t}$  denote the  $\sigma$ -algebra generated by the increments of X on [s, t],  $0 \le s \le t$ . Then, by a reversibility argument,

<span id="page-7-0"></span>
$$
\mathbb{E}[\lambda(\widehat{\mathcal{S}}[N-1,N]) \mid \mathcal{F}_{k-1}] \stackrel{\text{(d)}}{=} \mathbb{E}[\lambda(\mathcal{S}_1 \setminus \mathcal{S}[1,N]) \mid \mathcal{F}_{N-k+1,N}].
$$

Moreover, the following convergence in  $\mathbf{L}^1$  holds

(2.13) 
$$
\mathbb{E}[\lambda(\mathcal{S}_1 \setminus \mathcal{S}[1,N]) | \mathcal{F}_{N-k+1,N}] \xrightarrow[N \to \infty]{\mathbb{L}^1} \mathbb{E}[\lambda(\mathcal{S}_1 \setminus \mathcal{S}[1,\infty))].
$$

The proof of eq.  $(2.13)$  is postponed to Section [5,](#page-20-0) Lemma [5.1.](#page-20-1) In view of eq.  $(2.1)$  it follows that

$$
\mathbb{E}[\lambda(\widehat{\mathcal{S}}[N-1,N]) \mid \mathcal{F}_{k-1}] \xrightarrow[N \nearrow \infty]{\mathbb{L}^1} \text{Cap}(\mathcal{B}).
$$

We thus obtain

<span id="page-7-1"></span>(2.14) 
$$
U_k^N \xrightarrow[N \to \infty]{\mathbb{L}^1} \lambda(\widehat{\mathcal{S}}[k-1,k]) - \text{Cap}(\mathcal{B}) + \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) | \mathcal{F}_{k-1}] - \mathbb{E}[\lambda(\mathcal{S}_k \cap \mathcal{S}[k,\infty)) | \mathcal{F}_k].
$$

Further, we observe that

$$
\lambda(\mathcal{S}_k \cap \mathcal{S}[k,\infty)) = \lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) - \lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,k]) \n+ \lambda(\mathcal{S}_k \cap \mathcal{S}[k,\infty) \cap \mathcal{S}_{k-1} \cap \mathcal{S}[k-1,k]) + \lambda(\mathcal{S}[k-1,k] \cap \mathcal{S}[k,\infty)) \n- \lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty) \cap \mathcal{S}[k-1,k] \cap \mathcal{S}[k,\infty)) \n= \lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) - \lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,k]) \n+ \lambda(\mathcal{S}[k-1,k] \cap \mathcal{S}[k,\infty)).
$$

This and eq. [\(2.14\)](#page-7-1) imply

$$
U_k^N \xrightarrow[N \to \infty]{\text{L}^1} \lambda(\mathcal{S}[k-1,k] \cap \mathcal{S}_{k-1}^c) - \text{Cap}(\mathcal{B}) + \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) \mid \mathcal{F}_{k-1}]
$$
  
\n
$$
- \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) \mid \mathcal{F}_k] + \lambda(\mathcal{S}[k-1,k] \cap \mathcal{S}_{k-1})
$$
  
\n
$$
- \mathbb{E}[\lambda(\mathcal{S}[k-1,k] \cap \mathcal{S}[k,\infty)) \mid \mathcal{F}_k]
$$
  
\n
$$
= \lambda(\mathcal{S}[k-1,k]) - \mathbb{E}[\lambda(\mathcal{S}[k-1,k] \cap \mathcal{S}[k,\infty)) \mid \mathcal{F}_k] - \text{Cap}(\mathcal{B})
$$
  
\n
$$
+ \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) \mid \mathcal{F}_{k-1}] - \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) \mid \mathcal{F}_k]
$$
  
\n
$$
= \langle \mathbb{E}[\lambda(\mathcal{S}[k-1,k] \setminus \mathcal{S}[k,\infty)) \mid \mathcal{F}_k] \rangle + \mathbb{E}[\lambda(\mathcal{S}[k-1,k] \setminus \mathcal{S}[k,\infty))] - \text{Cap}(\mathcal{B})
$$
  
\n
$$
+ \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) \mid \mathcal{F}_{k-1}] - \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) \mid \mathcal{F}_k]
$$
  
\n
$$
= \langle \mathbb{E}[\lambda(\mathcal{S}[k-1,k] \setminus \mathcal{S}[k,\infty)) \mid \mathcal{F}_k] \rangle
$$
  
\n
$$
+ \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) \mid \mathcal{F}_{k-1}] - \mathbb{E}
$$

where in the last line we used eq.  $(2.1)$ . Hence, by eqs.  $(2.11)$  and  $(2.12)$ ,

<span id="page-7-2"></span>(2.15) 
$$
\langle \mathcal{V}_n \rangle = \langle \mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty)) | \mathcal{F}_n] \rangle + \sum_{k=1}^n Y_k,
$$

where

<span id="page-7-3"></span>(2.16) 
$$
Y_k = \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) | \mathcal{F}_{k-1}] - \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) | \mathcal{F}_k] + \langle \mathbb{E}[\lambda(\mathcal{S}[k-1,k] \setminus \mathcal{S}[k,\infty)) | \mathcal{F}_k] \rangle.
$$

From eq. [\(2.15\)](#page-7-2) it follows that the variance of  $\mathcal{V}_n$  is equal to

<span id="page-8-5"></span>
$$
\operatorname{Var}(\mathcal{V}_n) = \operatorname{Var}(\mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty)) | \mathcal{F}_n]) + \mathbb{E}\Big[\Big(\sum_{k=1}^n Y_k\Big)^2\Big] + 2 \mathbb{E}\Big[\langle \mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty)) | \mathcal{F}_n] \rangle \sum_{k=1}^n Y_k\Big].
$$

Clearly,  $Y_k$  is  $\mathcal{F}_k$ -measurable and  $\mathbb{E}[Y_l | \mathcal{F}_k] = 0$  for  $k < l$ . It follows that

$$
\mathbb{E}\Big[\Big(\sum_{k=1}^n Y_k\Big)^2\Big] = \sum_{k=1}^n \mathbb{E}[Y_k^2].
$$

Jensen's inequality and eq. [\(2.6\)](#page-5-4) with  $d > 3\alpha/2$  yield

$$
\lim_{n \nearrow \infty} \frac{1}{n} \text{Var}(\mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty)) | \mathcal{F}_n]) \leq \lim_{n \nearrow \infty} \frac{1}{n} \mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty))^{2}] = 0.
$$

The sequence  $\left\{\frac{1}{n}\right\}$  $\frac{1}{n}\sum_{k=1}^n \mathbb{E}[Y_k^2]\}_{n\geq 1}$  is bounded (the proof is given in Section [5,](#page-20-0) Lemma [5.2\)](#page-20-2), and thus by the Cauchy-Schwarz inequality we conclude that

<span id="page-8-0"></span>
$$
\lim_{n \nearrow \infty} \frac{1}{n} \mathbb{E} \Big[ \langle \mathbb{E} [\lambda(S_n \cap \mathcal{S}[n,\infty)) \mid \mathcal{F}_n] \rangle \sum_{k=1}^n Y_k \Big] = 0.
$$

We have shown that

(2.18) 
$$
\lim_{n \nearrow \infty} \frac{\text{Var}(\mathcal{V}_n)}{n} = \lim_{n \nearrow \infty} \frac{1}{n} \sum_{k=1}^n \mathbb{E}[Y_k^2].
$$

Step 2. In this step we prove that the limit on the right-hand side of eq.  $(2.18)$  is strictly positive. Let X' be an independent copy of the original process X such that  $X'_0 = 0$  and it has càglàd paths. We consider a process  $\bar{X} = {\{\bar{X}_t\}}_{t \in \mathbb{R}}$  by setting  $\bar{X}_t = X'_{-t}$  for  $t \leq 0$ , and  $\bar{X}_t = X_t$  for  $t \geq 0$ . Clearly, the process  $\bar{X}$  has càdlàg paths, and stationary and independent increments. The sausages  $S[s, t]$ ,  $S(-\infty, s]$  and  $S[s, \infty)$  corresponding to  $\bar{X}$ are defined for all  $s, t \in \mathbb{R}$ ,  $s \leq t$ . Recall that  $\mathcal{S}_{\infty} = \mathcal{S}[0, \infty)$ . We assume that the process  $\bar{X}$  is defined on the canonical space  $\Omega = \mathcal{D}(\mathbb{R}, \mathbb{R}^d)$  of all càdlàg functions  $\omega \colon \mathbb{R} \to \mathbb{R}^d$  as the coordinate process  $\bar{X}_t(\omega) = \omega(t)$ . We define a shift operator  $\vartheta$  acting on  $\Omega$  by

(2.19) 
$$
\vartheta \omega(t) = \omega(1+t) - \omega(1), \qquad t \in \mathbb{R},
$$

and we notice that it is a P-preserving mapping. For  $t \in \mathbb{R}$ , we define

<span id="page-8-3"></span><span id="page-8-1"></span>
$$
\mathcal{G}_t = \sigma(\bar{X}_s : -\infty < s \leq t),
$$

and for  $k \in \mathbb{N}$ ,

<span id="page-8-4"></span>(2.20) 
$$
Z_k = \mathbb{E}[\lambda(\mathcal{S}[-k,0]\cap\mathcal{S}_{\infty}) | \mathcal{G}_0] - \mathbb{E}[\lambda(\mathcal{S}[-k,0]\cap\mathcal{S}_{\infty}) | \mathcal{G}_1] + \langle \mathbb{E}[\lambda(\mathcal{S}_1\setminus\mathcal{S}[1,\infty)) | \mathcal{G}_1] \rangle
$$
.  
By eqs. (2.16) and (2.19) it follows that

(2.21) 
$$
Y_k = Z_{k-1} \circ \vartheta^{k-1}.
$$

In the sequel, we prove that there exists a random variable Z such that  $\mathbb{E}[|Z|] > 0$  and

$$
(2.22) \t\t Z_k \xrightarrow[k \nearrow \infty]{\mathbb{L}^1} Z.
$$

Step 2a. We start by proving the existence of Z. This is evident if  $d > 2\alpha$  as in this case  $h(t) = 1$  and whence using eq. [\(2.6\)](#page-5-4) we obtain

<span id="page-8-2"></span>
$$
\mathbb{E}[\lambda(\mathcal{S}(-\infty,0]\cap\mathcal{S}_{\infty}]<\infty.
$$

This implies [\(2.22\)](#page-8-2) with

(2.23) 
$$
Z = \mathbb{E}[\lambda(\mathcal{S}(-\infty,0] \cap \mathcal{S}_{\infty}) | \mathcal{G}_0] - \mathbb{E}[\lambda(\mathcal{S}(-\infty,0] \cap \mathcal{S}_{\infty}) | \mathcal{G}_1] + \langle \mathbb{E}[\lambda(\mathcal{S}_1 \setminus \mathcal{S}[1,\infty)) | \mathcal{G}_1] \rangle.
$$

We next consider the case  $3\alpha/2 < d \leq 2\alpha$ . Then  $\mathbb{E}[\lambda(\mathcal{S}(-\infty,0] \cap \mathcal{S}_{\infty})]$  is not finite and we cannot define  $Z$  as in the previous case. We notice that

<span id="page-9-1"></span>
$$
\lambda(\mathcal{S}[-k,0]\cap\mathcal{S}_{\infty})\,=\,\lambda(\mathcal{S}[-k,0]\cap\mathcal{S}[1,\infty))+\lambda(\mathcal{S}[-k,0]\cap(\mathcal{S}_1\setminus\mathcal{S}[1,\infty))
$$

and thus we can rewrite  $Z_k$  as follows

$$
Z_k = \langle \mathbb{E}[\lambda(\mathcal{S}_1 \setminus \mathcal{S}[1,\infty)) | \mathcal{G}_1] \rangle - \mathbb{E}[\lambda((\mathcal{S}_1 \setminus \mathcal{S}[1,\infty]) \cap \mathcal{S}[-k,0]) | \mathcal{G}_1] + \mathbb{E}[\lambda(\mathcal{S}[-k,0] \cap \mathcal{S}_{\infty}) | \mathcal{G}_0] - \mathbb{E}[\lambda(\mathcal{S}[-k,0] \cap \mathcal{S}[1,\infty)) | \mathcal{G}_1].
$$

Before we let  $k$  tend to infinity in the above expression, we rewrite the expression from the second line. We observe that

$$
\lambda(\mathcal{S}[-k,0]\cap\mathcal{S}_{\infty})\,=\,\int_{\mathbb{R}^d}\mathbb{1}_{\mathcal{S}[-k,0]}(y)\,\mathbb{1}_{\mathcal{S}_{\infty}}(y)\,\mathrm{d}y.
$$

By taking conditional expectation with respect to  $\mathcal{G}_0$ , we obtain

$$
\mathbb{E}[\lambda(\mathcal{S}[-k,0]\cap\mathcal{S}_{\infty})\mid\mathcal{G}_0]=\int_{\mathbb{R}^d}\mathbb{1}_{\mathcal{S}[-k,0]}(y)\,\mathbb{E}[\mathbb{1}_{\mathcal{S}_{\infty}}(y)]\,\mathrm{d}y\,=\,\int_{\mathbb{R}^d}\mathbb{1}_{\mathcal{S}[-k,0]}(y)\,\phi(y)\,\mathrm{d}y,
$$

where we set

$$
\phi(y) = \mathbb{P}(y \in \mathcal{S}_{\infty}).
$$

Similarly, we write

$$
\lambda(\mathcal{S}[-k,0]\cap \mathcal{S}[1,\infty))\,=\, \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}[-k,0]-X_1}(y) \, \mathbb{1}_{\mathcal{S}[1,\infty)-X_1}(y) \, \mathrm{d}y
$$

and we take conditional expectation with respect to  $\mathcal{G}_1$  which yields

<span id="page-9-2"></span>
$$
\mathbb{E}[\lambda(\mathcal{S}[-k,0]\cap\mathcal{S}[1,\infty)) \mid \mathcal{G}_1] = \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}[-k,0]-X_1}(y) \mathbb{E}[\mathbb{1}_{\mathcal{S}[1,\infty)-X_1}(y)] dy
$$
  
= 
$$
\int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}[-k,0]}(y) \phi(y-X_1) dy.
$$

It follows that

<span id="page-9-0"></span>
$$
\mathbb{E}[\lambda(\mathcal{S}[-k,0] \cap \mathcal{S}_{\infty}) \mid \mathcal{G}_0] - \mathbb{E}[\lambda(\mathcal{S}[-k,0] \cap \mathcal{S}[1,\infty)) \mid \mathcal{G}_1]
$$
  
= 
$$
\int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}[-k,0]}(y) \left(\phi(y) - \phi(y - X_1)\right) dy.
$$

We prove in Lemma [5.4](#page-24-0) that the right-hand side integral in eq.  $(2.25)$  is a well-defined random variable in  $L^1$ . Thus, the dominated convergence theorem implies eq.  $(2.22)$  with

$$
Z = \langle \mathbb{E}[\lambda(\mathcal{S}_1 \setminus \mathcal{S}[1,\infty)) | \mathcal{G}_1] \rangle - \mathbb{E}[\lambda((\mathcal{S}_1 \setminus \mathcal{S}[1,\infty]) \cap \mathcal{S}(-\infty,0]) | \mathcal{G}_1] + \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0]}(y) (\phi(y) - \phi(y - X_1)) dy.
$$

Step 2b. We next show that  $\mathbb{E}(|Z| > 0$  and we remark that the following arguments apply to all  $d > 3\alpha/2$ . From eqs.  $(2.15)$  and  $(2.21)$  we have

$$
\langle \mathcal{V}_n \rangle = \sum_{k=0}^{n-1} Z \circ \vartheta^k + H_n,
$$

where

(2.26) 
$$
H_n = \langle \mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty)) | \mathcal{G}_n] \rangle + \sum_{k=0}^{n-1} (Z_k - Z) \circ \vartheta^k.
$$

Equation  $(2.1)$  yields

<span id="page-10-2"></span>
$$
\langle \mathcal{V}_n \rangle = \mathcal{V}_n - \mathbb{E}[\mathcal{V}_n] \leq \mathcal{V}_n - \mathbb{E}[\lambda(\mathcal{S}_n \setminus \mathcal{S}[n,\infty))] = \mathcal{V}_n - n \operatorname{Cap}(\mathcal{B}).
$$

This implies

<span id="page-10-0"></span>
$$
\mathcal{V}_n \ge n \operatorname{Cap}(\mathcal{B}) + \sum_{k=0}^{n-1} Z \circ \vartheta^k + H_n.
$$

We aim to prove that there is  $\bar{c} > 0$  (which does not depend on n) such that for all  $n \in \mathbb{N}$ 

(2.27) 
$$
\mathbb{P}(\lbrace \mathcal{V}_n \leq \bar{c} \rbrace \cap \lbrace |H_n| \leq \bar{c} \rbrace) > 0.
$$

Notice that if  $\mathbb{E}[|Z|] = 0$  then the inequality  $\mathcal{V}_n \ge n \operatorname{Cap}(\mathcal{B}) + H_n$  and eq. [\(2.27\)](#page-10-0) would imply that  $Cap(B) = 0$  which is a contradiction. Therefore, it is enough to show eq. [\(2.27\)](#page-10-0). From eqs.  $(2.20)$  and  $(2.23)$  we obtain

<span id="page-10-1"></span>
$$
\sum_{k=0}^{n-1} (Z - Z_k) \circ \vartheta^k = \int_{\mathbb{R}^d} \mathbb{1}_{S(-\infty,0]}(y) \mathbb{E}[\mathbb{1}_{S_{\infty}}(y) | \mathcal{G}_0] dy
$$

$$
- \sum_{k=0}^{n-2} \int_{\mathbb{R}^d} \left( \mathbb{1}_{S(-\infty,0] \setminus S_k}(y) \mathbb{E}[\mathbb{1}_{S[k,\infty)}(y) | \mathcal{G}_{k+1}] - \mathbb{1}_{S(-\infty,0] \setminus S_{k+1}}(y) \mathbb{E}[\mathbb{1}_{S[k+1,\infty)}(y) | \mathcal{G}_{k+1}] \right) dy
$$

$$
- \int_{\mathbb{R}^d} \mathbb{1}_{S(-\infty,0] \setminus S_{n-1}}(y) \mathbb{E}[\mathbb{1}_{S[n-1,\infty)}(y) | \mathcal{G}_n] dy.
$$

We next observe that

$$
\int_{\mathbb{R}^d} \left( \mathbb{1}_{\mathcal{S}(-\infty,0] \setminus \mathcal{S}_k}(y) \, \mathbb{E}[\mathbb{1}_{\mathcal{S}[k,\infty)}(y) \mid \mathcal{G}_{k+1}] \right. \\
\left. - \mathbb{1}_{\mathcal{S}(-\infty,0] \setminus \mathcal{S}_{k+1}}(y) \, \mathbb{E}[\mathbb{1}_{\mathcal{S}[k+1,\infty)}(y) \mid \mathcal{G}_{k+1}] \right) \mathrm{d}y \\
= \int_{\mathbb{R}^d} \left( \mathbb{1}_{\mathcal{S}(-\infty,0]}(y) \, \mathbb{1}_{\mathcal{S}_k^c}(y) \, \mathbb{E}[\mathbb{1}_{\mathcal{S}[k,k+1]}(y) + \mathbb{1}_{\mathcal{S}[k+1,\infty)}(y) \, \mathbb{1}_{\mathcal{S}[k,k+1]^c}(y) \mid \mathcal{G}_{k+1}] \right. \\
\left. - \mathbb{1}_{\mathcal{S}(-\infty,0]}(y) \, \mathbb{1}_{\mathcal{S}_{k+1}^c}(y) \, \mathbb{E}[\mathbb{1}_{\mathcal{S}[k+1,\infty)}(y) \mid \mathcal{G}_{k+1}] \right) \mathrm{d}y \\
= \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0]}(y) \left( \mathbb{1}_{\mathcal{S}_k^c}(y) \, \mathbb{1}_{\mathcal{S}[k,k+1]}(y) \right. \\
\left. + \mathbb{1}_{\mathcal{S}_k^c}(y) \, \mathbb{1}_{\mathcal{S}[k,k+1]^c}(y) \, \mathbb{E}[\mathbb{1}_{\mathcal{S}[k+1,\infty)}(y) \mid \mathcal{G}_{k+1}] \right. \\
\left. - \mathbb{1}_{\mathcal{S}_{k+1}^c}(y) \, \mathbb{E}[\mathbb{1}_{\mathcal{S}[k+1,\infty)}(y) \mid \mathcal{G}_{k+1}] \right) \mathrm{d}y \\
= \lambda(\mathcal{S}(-\infty,0] \cap (\mathcal{S}[k,k+1] \setminus \mathcal{S}_k)).
$$

We clearly have  $\mathbb{1}_{\mathcal{S}[n-1,\infty)}(y) = \mathbb{1}_{\mathcal{S}[n,\infty)}(y) + \mathbb{1}_{\mathcal{S}[n-1,n]\setminus\mathcal{S}[n,\infty)}(y)$ . This identity and a similar argument as in eq. [\(2.24\)](#page-9-2) yield

<span id="page-11-0"></span>
$$
\int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0] \setminus \mathcal{S}_{n-1}}(y) \mathbb{E}[\mathbb{1}_{\mathcal{S}[n-1,\infty)}(y) | \mathcal{G}_n] dy
$$
\n
$$
= \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0] \setminus \mathcal{S}_{n-1}}(y) \mathbb{E}[\mathbb{1}_{\mathcal{S}[n,\infty)}(y) | \mathcal{G}_n] dy
$$
\n(2.30)\n
$$
+ \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0] \setminus \mathcal{S}_{n-1}}(y) \mathbb{E}[\mathbb{1}_{\mathcal{S}[n-1,n] \setminus \mathcal{S}[n,\infty)}(y) | \mathcal{G}_n] dy
$$
\n
$$
= \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0]}(y) \phi(y - X_n) dy - \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0] \cap \mathcal{S}_{n-1}}(y) \phi(y - X_n) dy
$$
\n
$$
+ \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0] \setminus \mathcal{S}_{n-1}}(y) \mathbb{E}[\mathbb{1}_{\mathcal{S}[n-1,n] \setminus \mathcal{S}[n,\infty)}(y) | \mathcal{G}_n] dy.
$$

By combining eqs.  $(2.28)$  to  $(2.30)$ , we arrive at

<span id="page-11-1"></span>
$$
\sum_{k=0}^{n-1} (Z - Z_k) \circ \vartheta^k
$$
\n
$$
(2.31) \qquad = \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0]}(y) \left(\phi(y) - \phi(y - X_n)\right) \mathrm{d}y + \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0] \cap \mathcal{S}_{n-1}}(y) \phi(y - X_n) \mathrm{d}y
$$
\n
$$
- \int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0] \setminus \mathcal{S}_{n-1}}(y) \, \mathbb{E}[\mathbb{1}_{\mathcal{S}[n-1,n] \setminus \mathcal{S}[n,\infty)}(y) \mid \mathcal{G}_n] \mathrm{d}y - \lambda(\mathcal{S}(-\infty,0] \cap \mathcal{S}_{n-1}).
$$

We claim that there is a constant  $\tilde{c} > 0$  such that for all  $n \in \mathbb{N}$ 

<span id="page-11-2"></span>
$$
(2.32) \quad \mathbb{P}\left(\left\{\sup_{0\leq s\leq n}|X_s|\leq 1\right\}\cap\left\{\left|\int_{\mathbb{R}^d}\mathbb{J}_{\mathcal{S}(-\infty,0]}(y)\left(\phi(y)-\phi(y-X_n)\right)\mathrm{d}y\right|\leq \tilde{c}+1\right\}\right)>0.
$$

If  $\sup_{0\leq s\leq n}|X_n|\leq 1$ , then clearly  $\lambda(\mathcal{S}_n)\leq \lambda(\mathcal{B}(0, 2))$  and this allows us to estimate the first term on the right-hand side of eq. [\(2.26\)](#page-10-2) and, similarly, the three last terms on the right-hand side of eq. [\(2.31\)](#page-11-1) by a constant. We infer that there exists a constant  $\bar{c} > 0$ such that

$$
\left\{\sup_{0\leq s\leq n}|X_s|\leq 1\right\}\cap\left\{\left|\int_{\mathbb{R}^d}\mathbb{1}_{\mathcal{S}(-\infty,0]}(y)\left(\phi(y)-\phi(y-X_n)\right)\mathrm{d}y\right|\leq \tilde{c}+1\right\}
$$

$$
\subseteq \{\mathcal{V}_n\leq \bar{c}\}\cap\{H_n\leq \bar{c}\}.
$$

To finish the proof of eq. [\(2.27\)](#page-10-0), we are only left to show eq. [\(2.32\)](#page-11-2). In view of the Markov inequality it is enough to prove that under  $\sup_{0 \le s \le n} |X_n| \le 1$  we have

$$
\mathbb{E}\Big[\Big|\int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0]}(y) \left(\phi(y) - \phi(y - X_n)\right) \mathrm{d}y\Big|\Big] < \infty.
$$

This holds as, under  $\sup_{0 \le s \le n} |X_n| \le 1$ ,

$$
\mathbb{E}\Big[\Big|\int_{\mathbb{R}^d} \mathbb{1}_{\mathcal{S}(-\infty,0]}(y)\,\big(\phi(y)-\phi(y-X_n)\big)\,\mathrm{d} y\,\Big|\Big]\leq \sup_{x\in\mathcal{B}}\int_{\mathbb{R}^d}\phi(y)|\phi(y)-\phi(y-x)|\,\mathrm{d} y\,<\,\infty,
$$

where convergence of the last integral is established in Lemma [5.5.](#page-26-0)

*Step 2c.* We finally show that the limit in eq.  $(2.18)$  is positive. Equation  $(2.22)$  implies

$$
\lim_{n \nearrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathbb{E}[|Z_k|] = \mathbb{E}[|Z|].
$$

Hence, by Jensen's inequality,

$$
\lim_{n \nearrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathbb{E}[Z_k^2] \ge \lim_{n \nearrow \infty} \mathbb{E}\Big[\Big(\frac{1}{n} \sum_{k=0}^{n-1} |Z_k|\Big)^2\Big] \ge \lim_{n \nearrow \infty} \Big(\mathbb{E}\Big[\frac{1}{n} \sum_{k=0}^{n-1} |Z_k|\Big]\Big)^2
$$

$$
= \lim_{n \nearrow \infty} \Big(\frac{1}{n} \sum_{k=0}^{n-1} \mathbb{E}[|Z_k|]\Big)^2 = (\mathbb{E}[|Z|])^2.
$$

By eq.  $(2.21)$ , we have

$$
\lim_{n \nearrow \infty} \frac{1}{n} \sum_{k=1}^{n} \mathbb{E}[Y_k^2] = \lim_{n \nearrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathbb{E}[Z_k^2] \ge (\mathbb{E}[|Z|])^2 > 0,
$$

and this finishes the proof of the lemma.

2.3. Proof of the central limit theorem. In this paragraph, we prove Theorem [2.1.](#page-3-0) In the proof we apply the Lindeberg-Feller central limit theorem which we include for completeness.

<span id="page-12-1"></span>**Lemma 2.6** ([\[7,](#page-32-11) Theorem 3.4.5]). For each  $n \in \mathbb{N}$  let  $\{X_{n,i}\}_{1\leq i\leq n}$  be a sequence of independent random variables with zero mean. If the following conditions are satisfied

(i) 
$$
\lim_{n \to \infty} \sum_{i=1}^{n} \mathbb{E}[X_{n,i}^2] = \sigma^2 > 0, \text{ and}
$$
  
\n(ii) for every  $\varepsilon > 0$ , 
$$
\lim_{n \to \infty} \sum_{i=1}^{n} \mathbb{E}[X_{n,i}^2 \mathbb{1}_{\{|X_{n,i}| > \varepsilon\}}] = 0,
$$
  
\nthen  $X_{n,1} + \cdots + X_{n,n} \xrightarrow[n \to \infty]{(d)} \sigma \mathcal{N}(0, 1).$ 

*Proof of Theorem [2.1.](#page-3-0)* For  $t > 0$  large enough we choose  $n = n(t) = |\log(t)|$ . We have

$$
\mathcal{V}_t = \lambda(\mathcal{S}_t) = \lambda(\mathcal{S}_{t/n} \cup \mathcal{S}[t/n, t]) = \lambda((\mathcal{S}_{t/n} - X_{t/n}) \cup (\mathcal{S}[t/n, t] - X_{t/n})).
$$

By the Markov property,

$$
S_{t/n}^{(1)} = S_{t/n} - X_{t/n}
$$
 and  $S_{(n-1)t/n}^{(2)} = S[t/n, t] - X_{t/n}$ 

are independent, and  $\mathcal{S}_{(n-1)t/n}^{(2)}$  has the same law as  $\mathcal{S}_{(n-1)t/n}$ . Rotational invariance of X implies that  $\mathcal{S}_{t/n}^{(1)}$  is equal in law to  $\mathcal{S}_{t/n}$ . Hence,

<span id="page-12-0"></span>
$$
\mathcal{V}_t = \lambda(\mathcal{S}_{t/n}^{(1)}) + \lambda(\mathcal{S}_{(n-1)t/n}^{(2)}) - \lambda(\mathcal{S}_{t/n}^{(1)} \cap \mathcal{S}_{(n-1)t/n}^{(2)}).
$$

By iterating this procedure, we obtain

(2.33) 
$$
\mathcal{V}_t = \sum_{i=1}^n \lambda(\mathcal{S}_{t/n}^{(i)}) - \sum_{i=1}^{n-1} \lambda(\mathcal{S}_{t/n}^{(i)} \cap \mathcal{S}_{(n-i)t/n}^{(i+1)}).
$$

We denote

$$
\mathcal{V}_{t/n}^{(i)} = \lambda(\mathcal{S}_{t/n}^{(i)}) \quad \text{and} \quad \mathcal{R}(t) = \sum_{i=1}^{n-1} \lambda(\mathcal{S}_{t/n}^{(i)} \cap \mathcal{S}_{(n-i)t/n}^{(i+1)}),
$$

and we notice that  $\{V_{t/n}^{(i)}\}_{1 \leq t \leq n}$  are i.i.d. random variables. By taking expectation in eq. [\(2.33\)](#page-12-0) and then subtracting, we obtain

(2.34) 
$$
\langle \mathcal{V}_t \rangle = \sum_{i=1}^n \langle \mathcal{V}_{t/n}^{(i)} \rangle - \langle \mathcal{R}(t) \rangle.
$$

We first show that

(2.35) 
$$
\frac{\langle \mathcal{R}(t) \rangle}{\sqrt{t}} \xrightarrow[t \to \infty]{} 0.
$$

Since  $\mathcal{R}(t) \geq 0$ , we clearly have  $\mathbb{E}[|\langle \mathcal{R}(t)\rangle|] \leq 2 \mathbb{E}[\mathcal{R}(t)]$ . By Corollary [2.4,](#page-5-5)

<span id="page-13-0"></span>
$$
\mathbb{E}[\mathcal{R}(t)] \leq \sum_{i=1}^{n-1} \mathbb{E}\left[\lambda \big(\mathcal{S}_{t/n}^{(i)} \cap \mathcal{S}_{\infty}^{(i+1)}\big)\right] \leq cn \, h(t/n),
$$

for all  $t > 0$  large enough. Hence, eq.  $(2.35)$  follows by eq.  $(2.7)$ , and the fact that  $n = |\log(t)|$  and  $d/\alpha > 3/2$ . Next we prove that

(2.36) 
$$
\frac{1}{\sqrt{t}} \sum_{i=1}^{n} \langle \mathcal{V}_{t/n}^{(i)} \rangle \xrightarrow[t \to \infty]{(\text{d})} \sigma \mathcal{N}(0,1).
$$

For this we introduce the random variables

<span id="page-13-1"></span>
$$
X_{n,i} = \frac{\langle \mathcal{V}_{t/n}^{(i)} \rangle}{\sqrt{t}}, \qquad i = 1, \dots, n,
$$

and we check the validity of conditions (i) and (ii) from Lemma [2.6.](#page-12-1) Condition (i) follows by Lemma [2.5,](#page-6-4)

(2.37) 
$$
\lim_{n \nearrow \infty} \sum_{i=1}^{n} \mathbb{E}[X_{n,i}^2] = \lim_{n \nearrow \infty} \frac{n}{t} \text{Var}(\mathcal{V}_{t/n}) = \sigma^2.
$$

To establish condition (ii) we apply the Cauchy-Schwartz inequality and obtain that for every  $\varepsilon > 0$ ,

$$
\mathbb{E}\left[X_{n,i}^2\,1_{\{|X_{n,i}|>\varepsilon\}}\right] \leq \frac{1}{t}\left(\mathbb{E}\left[\langle \mathcal{V}_{t/n}\rangle^4\right]\mathbb{P}\left(\left|\langle \mathcal{V}_{t/n}\rangle\right|>\varepsilon\sqrt{t}\right)\right)^{1/2}
$$

.

By Chebyshev's inequality combined with Lemma [2.5](#page-6-4) and the fact that  $n = |\log(t)|$ , there is a constant  $c_1 > 0$  such that

$$
\mathbb{P}\left(|\langle \mathcal{V}_{t/n}\rangle| > \varepsilon\sqrt{t}\right) \leq \frac{\text{Var}(\mathcal{V}_{t/n})}{\varepsilon^2 t} \leq \frac{c_1 t/n}{\varepsilon^2 t} = \frac{c_1}{\varepsilon^2 n}.
$$

This together with Lemma [5.6](#page-27-0) imply that there are constants  $c_2, c_3 > 0$  such that

$$
(2.38) \qquad \lim_{n \nearrow \infty} \sum_{i=1}^{n} \mathbb{E}\left[X_{n,i}^{2} \mathbb{1}_{\{|X_{n,i}| > \varepsilon\}}\right] \leq \lim_{n \nearrow \infty} \sum_{i=1}^{n} \frac{1}{t} \Big(c_{2} \Big(\frac{t}{n}\Big)^{2} \frac{c_{1}}{\varepsilon^{2} n}\Big)^{1/2} \leq \lim_{n \nearrow \infty} \frac{c_{3}}{\sqrt{n}} = 0.
$$

Thus, eq. [\(2.36\)](#page-13-1) follows and we conclude that

$$
\frac{\langle \mathcal{V}_t \rangle}{\sqrt{t}} \xrightarrow[t \to \infty]{} \sigma \mathcal{N}(0,1).
$$

We finally observe that

$$
\frac{\mathcal{V}_t - t \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{t}} = \frac{\langle \mathcal{V}_t \rangle}{\sigma \sqrt{t}} + \frac{\mathbb{E}[\mathcal{V}_t] - t \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{t}},
$$

which allows us to finish the proof in view of Lemma [2.2](#page-3-3) and Corollary [2.4.](#page-5-5)  $\Box$ 

### 3. Functional central limit theorem

<span id="page-14-1"></span>The goal of this section is to prove the functional central limit theorem in eq.  $(1.7)$ . To prove this statement we adapt the proof of  $[5,$  Theorem 1.1, which is concerned with the functional central limit theorem for the capacity of the range of a stable random walk.

We again assume that X is a stable rotationally invariant Lévy process in  $\mathbb{R}^d$  of index  $\alpha \in (0, 2]$  satisfying  $d/\alpha > 3/2$ . We follow the classical two-step scheme (see [\[15,](#page-33-3) Theorem 16.10 and Theorem 16.11]). Let  $\{Y^n\}_{n\geq 1}$  be a sequence of random elements in the Skorohod space  $\mathcal{D}([0,\infty),\mathbb{R})$  endowed with the Skorohod J<sub>1</sub> topology. The sequence  ${Y^n}_{n \geq 1}$  converges weakly to a random element Y (in  $\mathcal{D}([0,\infty),\mathbb{R})$ ) if the following two conditions are satisfied:

- (i) The finite dimensional distributions of  ${Y<sup>n</sup>}_{n\geq 1}$  converge weakly to the finite dimensional distributions of Y.
- (ii) For any bounded sequence  $\{T_n\}_{n\geq 1}$  of  $\{Y^n\}_{n\geq 1}$ -stopping times and any non-negative sequence  ${b_n}_{n>1}$  converging to zero,

$$
\lim_{n \nearrow \infty} \mathbb{P}\left( |Y_{T_n + b_n}^n - Y_{T_n}^n| \ge \varepsilon \right) = 0, \qquad \varepsilon > 0.
$$

<span id="page-14-0"></span>Theorem 3.1. Under the above assumptions, the following convergence holds

$$
\left\{\frac{\mathcal{V}_{nt}-nt\operatorname{Cap}(\mathcal{B})}{\sigma\sqrt{n}}\right\}_{t\geq 0} \xrightarrow[n\nearrow\infty]{(J_1)} \{W_t\}_{t\geq 0},
$$

where  $\sigma$  is the constant from Theorem [2.1.](#page-3-0)

Proof. We consider the following sequence of random elements which are defined in the space  $\mathcal{D}([0,\infty),\mathbb{R}),$ 

(3.1) 
$$
Y_t^n = \frac{\mathcal{V}_{nt} - nt \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{n}}, \qquad n \in \mathbb{N},
$$

where  $\sigma$  is the constant from Theorem [2.1.](#page-3-0) Let us start by showing condition (i). *Condition (i)*. By Theorem [2.1,](#page-3-0) we have

<span id="page-14-2"></span>
$$
Y_t^n = \frac{\mathcal{V}_{nt} - nt \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{n}} = \sqrt{t} \frac{\mathcal{V}_{nt} - nt \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{nt}} \xrightarrow[n \to \infty]{\text{(d)}} \mathcal{N}(0, t).
$$

Let  $k \in \mathbb{N}$  be arbitrary and choose  $0 = t_0 < t_1 < t_2 < \cdots < t_k$ . We need to prove that

$$
(Y_{t_1}^n, Y_{t_2}^n, \ldots, Y_{t_k}^n) \xrightarrow[n \nearrow \infty]{(\mathrm{d})} (W_{t_1}, W_{t_2}, \ldots, W_{t_k}).
$$

In view of the Cramér-Wold theorem  $[15,$  Corollary 5.5] it suffices to show that

$$
\sum_{j=1}^k \xi_j Y_{t_j}^n \xrightarrow[n \nearrow \infty]{(\mathrm{d})} \sum_{j=1}^k \xi_j W_{t_j}, \qquad (\xi_1, \xi_2, \dots, \xi_k) \in \mathbb{R}^k.
$$

Using a similar reasoning as in the beginning of the proof of Theorem [2.1,](#page-3-0) we obtain for  $j \in \{1, 2, \ldots, k\},\$ 

$$
\mathcal{V}_{nt_j} \,=\, \sum_{i=1}^{j} \mathcal{V}_{n(t_i - t_{i-1})}^{(i)} - \sum_{i=1}^{j-1} \mathcal{R}_{nt_j}^{(i)},
$$

where

$$
\mathcal{V}_{n(t_i-t_{i-1})}^{(i)} = \lambda(\mathcal{S}_{n(t_i-t_{i-1})}^{(i)}) \quad \text{and} \quad \mathcal{R}_{nt_j}^{(i)} = \lambda(\mathcal{S}_{n(t_i-t_{i-1})}^{(i)} \cap \mathcal{S}_{n(t_j-t_i)}^{(i+1)}).
$$

The random variables  $\mathcal{V}_{n(t)}^{(i)}$  $n(t_{i-t_{i-1}}),$  for  $i \in \{1, 2, ..., k\}$ , are independent,  $\mathcal{S}_{n(t)}^{(i)}$  $_{n(t_i-t_{i-1})}^{(t)}$  has the same law as  $\mathcal{S}_{n(t_i-t_{i-1})}$ , and  $\mathcal{R}_{nt_i}^{(i)}$  $\mathcal{S}_{nt_j}^{(i)}$  has the same law as  $\lambda(\mathcal{S}_{n(t_i-t_{i-1})} \cap \mathcal{S}_{n(t_j-t_i)}')$ , with  $\mathcal{S}_{n(t_j-t_i)}'$ being an independent copy of  $\mathcal{S}_{n(t_j-t_i)}$ . For arbitrary  $(\xi_1, \xi_2, \ldots, \xi_k) \in \mathbb{R}^k$  we have

$$
\sum_{j=1}^{k} \xi_{j} Y_{t_{j}}^{n} = \sum_{j=1}^{k} \xi_{j} \left( \frac{\mathcal{V}_{nt_{j}} - nt_{j} \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{n}} \right)
$$
\n
$$
= \frac{1}{\sigma \sqrt{n}} \sum_{j=1}^{k} \xi_{j} \left( \sum_{i=1}^{j} \mathcal{V}_{n(t_{i}-t_{i-1})}^{(i)} - \sum_{i=1}^{j-1} \mathcal{R}_{nt_{j}}^{(i)} - \sum_{i=1}^{j} n(t_{i}-t_{i-1}) \operatorname{Cap}(\mathcal{B}) \right)
$$
\n
$$
= \sum_{j=1}^{k} \xi_{j} \sum_{i=1}^{j} \frac{\mathcal{V}_{n(t_{i}-t_{i-1})}^{(i)} - n(t_{i}-t_{i-1}) \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{n}} - \frac{1}{\sigma} \sum_{j=1}^{k} \xi_{j} \sum_{i=1}^{j-1} \frac{\mathcal{R}_{nt_{j}}^{(i)}}{\sqrt{n}}
$$
\n
$$
= \sum_{i=1}^{k} \left( \sum_{j=i}^{k} \xi_{j} \right) \frac{\mathcal{V}_{n(t_{i}-t_{i-1})}^{(i)} - n(t_{i}-t_{i-1}) \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{n}} - \frac{1}{\sigma} \sum_{j=1}^{k} \xi_{j} \sum_{i=1}^{j-1} \frac{\mathcal{R}_{nt_{j}}^{(i)}}{\sqrt{n}}.
$$

Theorem [2.1](#page-3-0) provides that

$$
\frac{\mathcal{V}_{n(t_i-t_{i-1})}^{(i)} - n(t_i - t_{i-1})\operatorname{Cap}(\mathcal{B})}{\sigma\sqrt{n}} \xrightarrow[n \to \infty]{\text{(d)}} \mathcal{N}(0, t_i - t_{i-1}).
$$

Markov's inequality combined with Corollary [2.4](#page-5-5) implies that for every  $\varepsilon > 0$ ,

$$
\mathbb{P}\left(\frac{\mathcal{R}_{nt_j}^{(i)}}{\sqrt{n}} > \varepsilon\right) \leq \frac{\mathbb{E}[\lambda(\mathcal{S}_{n(t_i-t_{i-1})} \cap \mathcal{S}_{n(t_j-t_i)})]}{\varepsilon \sqrt{n}} \leq \frac{\mathbb{E}[\lambda(\mathcal{S}_{nt_j} \cap \mathcal{S}_{\infty}')] }{\varepsilon \sqrt{n}} \leq \frac{c \, h(nt_j)}{\varepsilon \sqrt{n}},
$$

which converges to zero, as n tends to infinity. Since for  $i \in \{1, 2, ..., k\}$  the random variables  $\mathcal{V}_{n(t)}^{(i)}$  $n(t_i-t_{i-1})$  are independent, we obtain

$$
\sum_{j=1}^k \xi_j Y_{t_j}^n \xrightarrow[n \to \infty]{\text{(d)}} \mathcal{N}\left(0, \sum_{i=1}^k \Big(\sum_{j=i}^k \xi_j\Big)^2 (t_i - t_{i-1})\right).
$$

It follows that the finite dimensional distributions of  ${Y^n}_{n \geq 1}$  converge weakly to the finite dimensional distributions of a one-dimensional standard Brownian motion.

Condition (ii). Let  ${T_n}_{n\geq 1}$  be a bounded sequence of  ${Y^n}_{n\geq 1}$ -stopping times, and let  ${b_n}_{n\geq 1} \subset [0,\infty)$  be an arbitrary sequence which converges to zero. We aim to prove that

$$
Y^n_{T_n+b_n} - Y^n_{T_n} \xrightarrow[n \nearrow \infty]{\mathbb{P}} 0,
$$

where the convergence holds in probability. By eq.  $(3.1)$ , we have

(3.2) 
$$
Y_{T_n+b_n}^n - Y_{T_n}^n = \frac{\mathcal{V}_{n(T_n+b_n)} - n(T_n+b_n)\operatorname{Cap}(\mathcal{B})}{\sigma\sqrt{n}} - \frac{\mathcal{V}_{nT_n} - nT_n\operatorname{Cap}(\mathcal{B})}{\sigma\sqrt{n}}.
$$

The Markov property and rotational invariance of X yield

<span id="page-15-0"></span>
$$
\mathcal{V}_{n(T_n+b_n)} - \mathcal{V}_{nT_n} = \lambda ((\mathcal{S}_{nT_n} \cup \mathcal{S}[nT_n, n(T_n+b_n)]) - X_{nT_n}) - \lambda (\mathcal{S}_{nT_n} - X_{nT_n})
$$
  
=  $\lambda(\mathcal{S}_{nT_n}^{(1)}) + \lambda(\mathcal{S}_{n b_n}^{(2)}) - \lambda(\mathcal{S}_{nT_n}^{(1)} \cap \mathcal{S}_{n b_n}^{(2)}) - \lambda(\mathcal{S}_{nT_n}^{(1)})$   
=  $\lambda(\mathcal{S}_{n b_n}^{(2)}) - \lambda(\mathcal{S}_{nT_n}^{(1)} \cap \mathcal{S}_{n b_n}^{(2)}),$ 

where  $\mathcal{S}_{nT}^{(1)}$  $\begin{smallmatrix} \mathcal{S}^{(1)}_{nT_{n}} \text{ and } \mathcal{S}^{(2)}_{nb_{n}} \end{smallmatrix}$  $n_{nb_n}^{(2)}$  are independent and have the same distribution as  $S_{nT_n}$  and  $S_{nb_n}$ , respectively. Equation [\(3.2\)](#page-15-0) implies

$$
Y_{T_n+b_n}^n - Y_{T_n}^n = \frac{\lambda(\mathcal{S}_{nb_n}^{(2)}) - \lambda(\mathcal{S}_{nT_n}^{(1)} \cap \mathcal{S}_{nb_n}^{(2)}) - nb_n \operatorname{Cap}(\mathcal{B})}{\sigma \sqrt{n}}.
$$

With a slight abuse of notation we write  $\mathcal{V}_{nb_n} = \lambda(\mathcal{S}_{nb_n}^{(2)})$  $\binom{1}{n}$ . By Lemma [2.2,](#page-3-3) we obtain

<span id="page-16-1"></span>
$$
(3.3) \tY_{T_n+b_n}^n-Y_{T_n}^n=\frac{\mathcal{V}_{nb_b}-\mathbb{E}[\mathcal{V}_{nb_n}]}{\sigma\sqrt{n}}+\frac{\mathbb{E}[\lambda(\mathcal{S}_{nb_n}\cap\mathcal{S}[nb_b,\infty))]}{\sigma\sqrt{n}}-\frac{\lambda(\mathcal{S}_{nT_n}^{(1)}\cap\mathcal{S}_{nb_n}^{(2)})}{\sigma\sqrt{n}}.
$$

We prove that the three terms on the right-hand side of eq.  $(3.3)$  converge to zero in probability. For the first term, Chebyshev's inequality yields that for every  $\varepsilon > 0$ ,

<span id="page-16-2"></span>
$$
\mathbb{P}\left(\left|\frac{\mathcal{V}_{nb_n} - \mathbb{E}[\mathcal{V}_{nb_n}]}{\sqrt{n}}\right| > \varepsilon\right) \le \frac{\text{Var}(\mathcal{V}_{nb_n})}{\varepsilon^2 n},
$$

and we are left to show that

(3.4) 
$$
\lim_{n \nearrow \infty} \frac{\text{Var}(\mathcal{V}_{nb_n})}{n} = 0.
$$

This follows by Lemma [2.5.](#page-6-4) Indeed, there exist  $t_1, c_1 > 0$ , such that for every  $t \geq t_1$ , we have  $Var(V_t) \leq c_1 t$ , and whence for  $nb_n \geq t_1$ ,  $Var(V_{nb_n}) \leq c_1 nb_n$ . For  $nb_n < t_1$  we observe that  $Var(\mathcal{V}_{nb_n}) \leq \mathbb{E}[\mathcal{V}_{nb_n}^2] \leq \mathbb{E}[\mathcal{V}_{t_1}^2]$ . This trivially implies eq. [\(3.4\)](#page-16-2).

By Corollary [2.4,](#page-5-5) similarly as above, we show that there is  $t_2 > 0$  such that for all  $n \in \mathbb{N}$ 

$$
\mathbb{E}[\lambda(\mathcal{S}_{nb_n} \cap \mathcal{S}[nb_n,\infty))] \le ch(nb_n) + \mathbb{E}[\mathcal{V}_{t_2}].
$$

We then easily conclude that the second term on the right-hand side of eq.  $(3.3)$  converges to zero in probability.

There exists  $c_2 > 0$  such that  $\sup_{n\geq 1} T_n \leq c_2$ . By the Markov inequality and Corol-lary [2.4,](#page-5-5) we obtain that for every  $\varepsilon > 0$ 

$$
\mathbb{P}\left(\frac{\lambda(\mathcal{S}_{nT_n}^{(1)}\cap\mathcal{S}_{nb_n}^{(2)})}{\sigma\sqrt{n}} > \varepsilon\right) \leq \frac{\mathbb{E}[\lambda(\mathcal{S}_{nT_n}^{(1)}\cap\mathcal{S}_{nb_n}^{(2)})]}{\varepsilon\sigma\sqrt{n}} \leq \frac{\mathbb{E}[\lambda(\mathcal{S}_{c_2n}^{(1)}\cap\mathcal{S}_{\infty}^{(2)})]}{\varepsilon\sigma\sqrt{n}} \leq \frac{c\,h(c_2n)}{\varepsilon\sigma\sqrt{n}},
$$

<span id="page-16-0"></span>which converges to zero, as  $n$  tends to infinity. This shows that the last term on the right-hand side of eq.  $(3.3)$  goes to zero in probability and the proof is finished.

### 4. Laws of the iterated logarithm

<span id="page-16-3"></span>This section is devoted to the proof of the following result.

**Theorem 4.1.** If  $d/\alpha > 9/5$ , then the process  $\{\mathcal{V}_t\}_{t>0}$  satisfies Khintchine's and Chung's law of the iterated logarithm, that is, eqs.  $(1.8)$  and  $(1.9)$ , respectively.

<span id="page-16-4"></span>We start with the proof of eq.  $(1.8)$ , which is based on the following lemma.

**Lemma 4.2** ([\[20,](#page-33-17) Chapter X, Theorem 2]). Let  $\{Y_n\}_{n\geq 1}$  be a sequence of independent random variables with mean 0 and finite variance. Set

<span id="page-16-5"></span>
$$
S_n = \sum_{i=1}^n Y_i
$$
, and  $s_n^2 = \sum_{i=1}^n \mathbb{E}[Y_i^2]$ .

Suppose  $\lim_{n \to \infty} s_n = \infty$ , and that for any  $\varepsilon > 0$ ,

(4.1) 
$$
\lim_{n \nearrow \infty} \frac{1}{s_n^2} \sum_{i=i_0}^n \mathbb{E} \left[ Y_i^2 1\!\!1_{\{|Y_i| \ge \varepsilon \sqrt{s_i^2/\log \log s_i^2}\}} \right] = 0
$$

and

(4.2) 
$$
\sum_{i=i_0}^{\infty} \frac{1}{s_i^2 \log \log s_i^2} \mathbb{E}\left[Y_i^2 1_{\{|Y_i| \ge \varepsilon \sqrt{s_i^2/\log \log s_i^2}\}}\right] < \infty,
$$

where  $i_0 = \min\{i \ge 1 : \log \log s_j^2 > 0, \ j \ge i\}$ . Then

<span id="page-17-1"></span>
$$
\limsup_{n \nearrow \infty} \frac{S_n}{\sqrt{2s_n^2 \log \log s_n^2}} = 1 \qquad \mathbb{P}\text{-}a.s.
$$

*Proof of Theorem [4.1](#page-16-3) - Khintchine's law of the iterated logarithm.* By  $|a|$  we denote the greatest integer less than or equal to  $a \in \mathbb{R}$ . We consider a sequence  $\{n_i\}_{i\geq 0}$  of nonnegative integers such that if  $2^k \leq n_i < 2^{k+1}$ , then  $n_i$  runs over all consecutive integers of the form  $2^k + \lfloor j2^{k/2}/k \rfloor$ , for  $k \in \mathbb{N}$  and  $j = 0, 1, \ldots, \lfloor k2^{k/2} \rfloor$ . We set  $n_0 = 0$ . Clearly, if  $2^k \leq n_i < 2^{k+1}$  then  $0 \leq n_{i+1} - n_i \leq 2^{k/2}/k + 1$ . Hence,

(4.3) 
$$
\lim_{i \nearrow \infty} \frac{n_{i+1}}{n_i} = 1 \quad \text{and} \quad n_{i+1} - n_i = \mathcal{O}(n_i^{1/2}/\log n_i).
$$

Since

<span id="page-17-2"></span><span id="page-17-0"></span>
$$
\sup_{n_i \leq t \leq n_{i+1}} |\langle \mathcal{V}_t \rangle - \langle \mathcal{V}_{n_i} \rangle| \leq \mathcal{V}[n_i, n_{i+1}] + \mathbb{E}[\mathcal{V}[n_i, n_{i+1}]],
$$

we see that in the case when  $d/\alpha > 1$  (transience) eqs. [\(1.2\)](#page-0-0), [\(1.3\)](#page-0-2) and [\(4.3\)](#page-17-0) imply that

(4.4) 
$$
\lim_{i \nearrow \infty} \frac{\sup_{n_i \le t \le n_{i+1}} |\langle \mathcal{V}_t \rangle - \langle \mathcal{V}_{n_i} \rangle|}{\sqrt{n_i/\log \log n_i}} = 0 \quad \mathbb{P}\text{-a.s.}
$$

Thus, it suffices to prove that P-a.s.

$$
\liminf_{i \nearrow \infty} \frac{\langle \mathcal{V}_{n_i} \rangle}{\sqrt{2\sigma^2 n_i \log \log n_i}} = -1 \quad \text{and} \quad \limsup_{i \nearrow \infty} \frac{\langle \mathcal{V}_{n_i} \rangle}{\sqrt{2\sigma^2 n_i \log \log n_i}} = 1.
$$

We only discuss the second relation as the first one can be handled in an analogous way. For  $i \geq 0$  we set

<span id="page-17-4"></span>(4.5) 
$$
\mathcal{V}(n_i, n_{i+1}] = \lambda(\mathcal{S}(n_i, n_{i+1}]) \quad \text{and} \quad \mathcal{J}_{n_i} = \lambda(\mathcal{S}(n_i, n_{i+1}] \cap \mathcal{S}_{n_i}),
$$

where  $S(s,t) = \bigcup_{s. Observe that  $\{\mathcal{V}(n_i, n_{i+1}]\}_{i\geq 0}$  forms a sequence of$ independent random variables, and for  $i \geq 1$  we have

(4.6) 
$$
\mathcal{V}_{n_i} = \sum_{j=0}^{i-1} \mathcal{V}(n_j, n_{j+1}] - \sum_{j=0}^{i-1} \mathcal{J}_{n_j},
$$

which yields

<span id="page-17-3"></span>
$$
\frac{\langle \mathcal{V}_{n_i} \rangle}{\sqrt{2\sigma^2 n_i \log \log n_i}} = \frac{\sum_{j=0}^{i-1} \langle \mathcal{V}(n_j, n_{j+1}] \rangle}{\sqrt{2\sigma^2 n_i \log \log n_i}} - \frac{\sum_{j=0}^{i-1} \langle \mathcal{J}_{n_j} \rangle}{\sqrt{2\sigma^2 n_i \log \log n_i}}
$$

.

In Lemma [5.8](#page-29-0) we show that if  $d/\alpha > 9/5$ , then the last term on the right-hand side of the above identity converges to zero P-a.s. Thus we are left to prove that

$$
\limsup_{i \nearrow \infty} \frac{\sum_{j=0}^{i-1} \langle \mathcal{V}(n_j, n_{j+1}] \rangle}{\sqrt{2\sigma^2 n_i \log \log n_i}} = 1 \quad \mathbb{P}\text{-a.s.}
$$

When  $d/\alpha \in (1, 2)$  we set  $\Lambda = 2 - d/\alpha$  which satisfies  $\Lambda \in (0, 1)$ . We apply Lemma [5.7](#page-28-0) and obtain that for  $n_i \in [2^k, 2^{k+1}]$  there are constants  $c_1, c_2, c_3, c_4 > 0$  such that

$$
\operatorname{Var}\left(\sum_{j=0}^{i-1} \mathcal{V}(n_j, n_{j+1}]\right) \le \sigma^2 n_i + c_1 \left(\sum_{j=0}^{i-1} (n_{j+1} - n_j)^{1/2} h(n_{j+1} - n_j)\right)
$$

$$
\leq \sigma^2 n_i + c_2 \left( \sum_{l=1}^k 2^{3l/4} h(2^{l/2}/l) l^{1/2} \right)
$$
  
\n
$$
\leq \begin{cases} \sigma^2 n_i + c_3 \left( \sum_{l=1}^k 2^{3l/4} l^{1/2} \right), & d/\alpha > 2, \\ \sigma^2 n_i + c_3 \left( \sum_{l=1}^k 2^{3l/4} l^{3/2} \right), & d/\alpha = 2, \\ \sigma^2 n_i + c_3 \left( \sum_{l=1}^k 2^{(2\Lambda+3)l/4} l^{1/2-\Lambda} \right), & d/\alpha \in (1,2). \end{cases}
$$
  
\n
$$
\leq \begin{cases} \sigma^2 n_i + c_4 2^{3k/4} k^{3/2}, & d/\alpha \geq 2, \\ \sigma^2 n_i + c_4 2^{(2\Lambda+3)k/4} k^{1/2}, & d/\alpha \in (1,2). \end{cases}
$$
  
\n
$$
= \begin{cases} \sigma^2 n_i + \mathcal{O}(n_i^{3/4} (\log n_i)^{3/2}), & d/\alpha \geq 2, \\ \sigma^2 n_i + \mathcal{O}(n_i^{(2\Lambda+3)/4} (\log n_i)^{1/2}), & d/\alpha \in (1,2). \end{cases}
$$

Hence, for  $d/\alpha > 3/2$ ,

(4.7) 
$$
\operatorname{Var}\left(\sum_{j=0}^{i-1} \mathcal{V}(n_j, n_{j+1}]\right) = \sigma^2 n_i + o(n_i).
$$

This enables us to apply Lemma [4.2.](#page-16-4) We only need to show that

$$
\sum_{i=2}^{\infty} \frac{1}{n_i} \mathbb{E} \left[ \langle \mathcal{V}(n_i, n_{i+1}] \rangle^2 \mathbb{1}_{\{|\langle \mathcal{V}(n_i, n_{i+1}] \rangle| \ge \varepsilon \sqrt{n_i/\log \log n_i} \}} \right] < \infty
$$

as then Kronecker's lemma implies both eqs.  $(4.1)$  and  $(4.2)$ . By Lemma [5.6](#page-27-0) we obtain

$$
\mathbb{E}[\langle \mathcal{V}(n_i, n_{i+1}] \rangle^4] \leq c_5(n_{i+1} - n_i)^2 \leq c_5 n_i
$$

for some  $c_5 > 0$ . Finally, we have

$$
\sum_{i=2}^{\infty} \frac{1}{n_i} \mathbb{E} \left[ \langle \mathcal{V}(n_i, n_{i+1}] \rangle^2 \mathbb{1}_{\{|\langle \mathcal{V}(n_i, n_{i+1}] \rangle| \ge \varepsilon \sqrt{n_i/\log \log n_i} \}} \right]
$$
\n
$$
\le \sum_{i=2}^{\infty} \frac{\log \log n_i}{\varepsilon^2 n_i^2} \mathbb{E} \left[ \langle \mathcal{V}(n_i, n_{i+1}] \rangle^4 \right]
$$
\n
$$
\le \frac{c_5}{\varepsilon^2} \sum_{i=2}^{\infty} \frac{\log \log n_i}{n_i}
$$
\n
$$
\le \frac{c_5}{\varepsilon^2} \sum_{k=1}^{\infty} \sum_{l=0}^{k2^{k/2}} \frac{\log \log(2^k + l2^{k/2}/k)}{2^k + l2^{k/2}/k - 1}
$$
\n
$$
\le \frac{c_5}{\varepsilon^2} \sum_{k=1}^{\infty} \frac{k \log \log 2^{k+1}}{2^{k/2} - 2^{-k/2}} < \infty,
$$

which completes the proof.  $\Box$ 

<span id="page-18-0"></span>The proof of Chung's law of the iterated logarithm is based on the following result.

**Lemma 4.3** ([\[26,](#page-33-18) Theorem A]). Let  $\{Y_n\}_{n\geq 1}$  be a sequence of independent random variables with mean 0 and finite variance. Set

$$
S_n = \sum_{i=1}^n Y_i
$$
, and  $s_n^2 = \sum_{i=1}^n \mathbb{E}[Y_i^2]$ .

Suppose that  $\lim_{n \to \infty} s_n = \infty$ ,  $\mathbb{E}[Y_n^2] = o(s_n^2/\log \log s_n^2)$ , and that  $\{Y_n^2/\mathbb{E}[Y_n^2]\}_{n \ge 1}$  is uniformly integrable. Then

$$
\liminf_{n \nearrow \infty} \frac{\max_{1 \le i \le n} |S_i|}{\sqrt{s_n^2 / \log \log s_n^2}} = \frac{\pi}{\sqrt{8}} \qquad \mathbb{P}\text{-}a.s.
$$

Proof of Theorem [4.1](#page-16-3) - Chung's law of the iterated logarithm. We observe that for  $n_i \leq$  $t < n_{i+1}$  it holds

$$
\sup_{0\leq s\leq t}|\langle \mathcal{V}_s\rangle| \leq \max_{0\leq j\leq i}|\langle \mathcal{V}_{n_j}\rangle| + \max_{0\leq j\leq i}\sup_{n_j\leq s\leq n_{j+1}}|\langle \mathcal{V}_s\rangle - \langle \mathcal{V}_{n_j}\rangle|.
$$

We first claim that

<span id="page-19-0"></span>(4.8) 
$$
\lim_{i \nearrow \infty} \frac{\max_{0 \le j \le i} \sup_{n_j \le s \le n_{j+1}} |\langle \mathcal{V}_s \rangle - \langle \mathcal{V}_{n_j} \rangle|}{\sqrt{n_i/\log \log n_i}} = 0 \quad \mathbb{P}\text{-a.s.}
$$

Indeed, according to eq. [\(4.4\)](#page-17-2) (if  $d/\alpha > 1$ ), there is  $\overline{\Omega} \subseteq \Omega$  such that  $\mathbb{P}(\overline{\Omega}) = 1$ , and for any  $\omega \in \overline{\Omega}$  and any  $\varepsilon > 0$  there exists  $j_0 = j_0(\omega, \varepsilon) \in \mathbb{N}$  for which

$$
\max_{j\geq j_0}\frac{\sup_{n_j\leq s\leq n_{j+1}}|\langle\mathcal{V}_s\rangle(\omega)-\langle\mathcal{V}_{n_j}\rangle(\omega)|}{\sqrt{n_j/\log\log n_j}} \ \leq \ \frac{\varepsilon}{2}.
$$

We then write

$$
\frac{\max_{0 \le j \le i} \sup_{n_j \le s \le n_{j+1}} |\langle \mathcal{V}_s \rangle(\omega) - \langle \mathcal{V}_{n_j} \rangle(\omega)|}{\sqrt{n_i/\log \log n_i}} \le \frac{\max_{0 \le j \le j_0} \sup_{n_j \le s \le n_{j+1}} |\langle \mathcal{V}_s \rangle(\omega) - \langle \mathcal{V}_{n_j} \rangle(\omega)|}{\sqrt{n_i/\log \log n_i}} + \frac{\max_{j_0 \le j \le i} \sup_{n_j \le s \le n_{j+1}} |\langle \mathcal{V}_s \rangle(\omega) - \langle \mathcal{V}_{n_j} \rangle(\omega)|}{\sqrt{n_i/\log \log n_i}}.
$$

We next choose  $i_0 = i_0(\omega, \varepsilon) \in \mathbb{N}$  such that

$$
\frac{\max_{0 \le j \le j_0} \sup_{n_j \le s \le n_{j+1}} |\langle \mathcal{V}_s \rangle(\omega) - \langle \mathcal{V}_{n_j} \rangle(\omega)|}{\sqrt{n_i/\log \log n_i}} \le \frac{\varepsilon}{2}, \qquad i \ge i_0,
$$

and we infer eq. [\(4.8\)](#page-19-0). We are thus left to show that

$$
\liminf_{i \nearrow \infty} \frac{\max_{0 \le j \le i} |\langle \mathcal{V}_{n_j} \rangle|}{\sqrt{\sigma^2 n_i / \log \log n_i}} = \frac{\pi}{\sqrt{8}} \qquad \mathbb{P}\text{-a.s.}
$$

From eq. [\(4.6\)](#page-17-3) we have

$$
\max_{0 \leq j \leq i} \left| \sum_{k=0}^{j-1} \langle \mathcal{V}(n_k, n_{k+1}] \rangle \right| - \max_{0 \leq j \leq i} \left| \sum_{k=0}^{j-1} \langle \mathcal{J}_{n_k} \rangle \right| \leq \max_{0 \leq j \leq i} |\langle \mathcal{V}_{n_j} \rangle|
$$
  

$$
\leq \max_{0 \leq j \leq i} \left| \sum_{k=0}^{j-1} \langle \mathcal{V}(n_k, n_{k+1}] \rangle \right| + \max_{0 \leq j \leq i} \left| \sum_{k=0}^{j-1} \langle \mathcal{J}_{n_k} \rangle \right|.
$$

If we proceed similarly as in the proof of eq. [\(4.8\)](#page-19-0) and apply Lemma [5.8](#page-29-0) instead of eq. [\(4.4\)](#page-17-2), we arrive at  $\frac{1}{2}$  + 1

$$
\lim_{i \nearrow \infty} \frac{\max_{0 \le j \le i} |\sum_{k=0}^{j-1} \langle \mathcal{J}_{n_k} \rangle|}{\sqrt{n_i/\log \log n_i}} = 0 \quad \mathbb{P}\text{-a.s.}
$$

Thus it suffices to show that

$$
\liminf_{i \nearrow \infty} \frac{\max_{0 \le j \le i} |\sum_{k=0}^{j-1} \langle \mathcal{V}(n_k, n_{k+1}] \rangle|}{\sqrt{\sigma^2 n_i / \log \log n_i}} = \frac{\pi}{\sqrt{8}} \qquad \mathbb{P}\text{-a.s.}
$$

To this end, we apply Lemma [4.3.](#page-18-0) Recall that according to Lemma [5.6,](#page-27-0) there is  $c > 0$ such that

$$
\mathbb{E}[\langle \mathcal{V}(n_i, n_{i+1}] \rangle^4] \le c (n_{i+1} - n_i)^2.
$$

This combined with Lemma [5.7](#page-28-0) gives that

$$
\sup_{i\geq 1}\frac{\mathbb{E}[\langle \mathcal{V}(n_i,n_{i+1}]\rangle^4]}{\mathbb{E}[\langle \mathcal{V}(n_i,n_{i+1}]\rangle^2]^2} < \infty
$$

which implies that the sequence  $\{\langle \mathcal{V}(n_i, n_{i+1}]\rangle^2/\mathbb{E}[\langle \mathcal{V}(n_i, n_{i+1}]\rangle^2]\}_{i\geq 1}$  is uniformly integrable. In view of Lemma [5.7](#page-28-0) and eq. [\(4.3\)](#page-17-0),

$$
\mathbb{E}[\langle \mathcal{V}(n_i, n_{i+1}]\rangle^2] = \mathsf{o}(\sqrt{n_i/\log\log n_i}),
$$

<span id="page-20-0"></span>and the proof is finished.  $\square$ 

# 5. Technical results

<span id="page-20-1"></span>**Lemma 5.1.** In the notation of the proof of Lemma [2.5,](#page-6-4) for any  $k \in \mathbb{N}$  it holds that

$$
\mathbb{E}[\lambda(\mathcal{S}_1\setminus\mathcal{S}[1,N]) | \mathcal{F}_{N-k+1,N}] \xrightarrow[N \nearrow \infty]{\mathbb{L}^1} \mathbb{E}[\lambda(\mathcal{S}_1\setminus\mathcal{S}[1,\infty))].
$$

*Proof.* Set  $m_N = \lambda(S_1 \setminus S[1, N])$  and  $m_\infty = \lambda(S_1 \setminus S[1, \infty))$ . We have

$$
\lim_{N \nearrow \infty} \mathbb{E}[|\mathbb{E}[m_N | \mathcal{F}_{N-k+1,N}] - \mathbb{E}[m_{\infty}]|]
$$
\n
$$
\leq \lim_{N \nearrow \infty} \mathbb{E}[|\mathbb{E}[m_N - m_{\infty} | \mathcal{F}_{N-k+1,N}]| + |\mathbb{E}[m_{\infty} | \mathcal{F}_{N-k+1,N}] - \mathbb{E}[m_{\infty}]]]
$$
\n
$$
\leq \lim_{N \nearrow \infty} (\mathbb{E}[\mathbb{E}[m_N - m_{\infty} | \mathcal{F}_{N-k+1,N}]] + \mathbb{E}[|\mathbb{E}[m_{\infty} | \mathcal{F}_{N-k+1,N}] - \mathbb{E}[m_{\infty}]]])
$$
\n
$$
= \lim_{N \nearrow \infty} \mathbb{E}[|m_N - m_{\infty}|] + \lim_{N \nearrow \infty} \mathbb{E}[|\mathbb{E}[m_{\infty} | \mathcal{F}_{N-k+1,N}] - \mathbb{E}[m_{\infty}]]].
$$

Since  $m_N$  clearly converges to  $m_\infty$  in  $L^1$ , we are left to prove that the second limit in the expression above is zero. For a fixed  $k \in \mathbb{N}$  we define

<span id="page-20-3"></span> $\mathcal{H}_N = \sigma(X_t : t \geq N - k + 1), \qquad N \geq k.$ 

We observe that  $\mathcal{F}_{N-k+1,N} \subseteq \mathcal{H}_N$  and  $\mathcal{H}_N$  is a decreasing family of  $\sigma$ -algebras. Moreover, according to Kolmogorov's  $0-1$  law, for every  $H \in \mathcal{H}_{\infty} = \bigcap_{N \geq 1} \mathcal{H}_N$ , we have  $\mathbb{P}(H) \in$  $\{0, 1\}$ . From Levi's theorem (see [\[23,](#page-33-19) Ch. II, Corollary 2.4]) we infer that  $\mathbb{P}\text{-a.s.}$ 

(5.1) 
$$
\lim_{N \nearrow \infty} \mathbb{E}[m_{\infty} | \mathcal{H}_N] = \mathbb{E}[m_{\infty} | \mathcal{H}_{\infty}] = \mathbb{E}[m_{\infty}].
$$

Notice that by eq. [\(2.1\)](#page-4-5),  $\mathbb{E}[m_{\infty}] = \text{Cap}(\mathcal{B})$ . Since the family  $\{|\mathbb{E}[m_{\infty} | \mathcal{H}_N]| \}_{N>1}$  is uniformly integrable, we infer that the convergence in eq.  $(5.1)$  holds also in  $L^1$ , see [\[7,](#page-32-11) Theorem 5.5.1]. We finally obtain

$$
\lim_{N \nearrow \infty} \mathbb{E} \big[ \big| \mathbb{E} [m_{\infty} | \mathcal{F}_{N-k+1,N}] - \mathbb{E} [m_{\infty}] \big| \big]
$$
\n
$$
= \lim_{N \nearrow \infty} \mathbb{E} \big[ \big| \mathbb{E} [\mathbb{E} [m_{\infty} | \mathcal{H}_N] | \mathcal{F}_{N-k+1,N}] - \mathbb{E} [m_{\infty}] \big| \big]
$$
\n
$$
\leq \lim_{N \nearrow \infty} \mathbb{E} \big[ \mathbb{E} [ \mathbb{E} [m_{\infty} | \mathcal{H}_N] - \mathbb{E} [m_{\infty}] | | \mathcal{F}_{N-k+1,N}] \big]
$$
\n
$$
= \lim_{N \nearrow \infty} \mathbb{E} \big[ \big| \mathbb{E} [m_{\infty} | \mathcal{H}_N] - \mathbb{E} [m_{\infty}] \big| \big] = 0,
$$

<span id="page-20-2"></span>and the proof is finished.  $\square$ 

**Lemma 5.2.** In the notation of the proof of Lemma [2.5,](#page-6-4) the sequence  $\{\frac{1}{n}\}$  $\frac{1}{n} \sum_{k=1}^{n} \mathbb{E}[Y_k^2] \}_{n \geq 1}$ is bounded.

*Proof.* We set  $\Delta = d/\alpha - 3/2$  and recall that it is a positive number. We present the proof in the case  $\Delta \in (0, 1/2)$  as the proof for  $\Delta \geq 1/2$  is similar. For  $\Delta \in (0, 1/2)$ , the function h(t) defined in eq. [\(2.7\)](#page-5-1) is given by  $h(t) = t^{1/2-\Delta}$ . By the Cauchy-Schwarz inequality,

$$
\begin{aligned}\n&\left|\mathbb{E}\Big[\langle \mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty)) \mid \mathcal{F}_n] \rangle \sum_{k=1}^n Y_k\Big]\right| \\
&\leq \left(\text{Var}(\mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty)) \mid \mathcal{F}_n])\right)^{1/2} \left(\sum_{k=1}^n \mathbb{E}[Y_k^2]\right)^{1/2} \\
&\leq 2\sqrt{2} \, c \, n^{1/2-\Delta} \left(\sum_{k=1}^n \mathbb{E}[Y_k^2]\right)^{1/2}.\n\end{aligned}
$$

This combined with eq. [\(2.17\)](#page-8-5) yields

$$
\frac{\text{Var}(\mathcal{V}_n)}{n} \geq \frac{1}{n} \text{Var}\big(\mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty)) \mid \mathcal{F}_n]\big) + \frac{1}{n} \sum_{k=1}^n \mathbb{E}[Y_k^2] - \frac{4\sqrt{2}c}{n^{\Delta}} \left(\frac{1}{n} \sum_{k=1}^n \mathbb{E}[Y_k^2]\right)^{1/2}.
$$

We suppose, for the sake of contradiction, that there exists a subsequence  ${n_m}_{m\geq 1} \subseteq \mathbb{N}$ such that

$$
\lim_{m \nearrow \infty} \frac{1}{n_m} \sum_{k=1}^{n_m} \mathbb{E}[Y_k^2] = \infty.
$$

Since

$$
\lim_{n \nearrow \infty} \frac{\text{Var}(\mathcal{V}_n)}{n} = \sigma^2 \quad \text{and} \quad \lim_{n \nearrow \infty} \frac{1}{n} \text{Var}\big(\mathbb{E}[\lambda(\mathcal{S}_n \cap \mathcal{S}[n,\infty)) \mid \mathcal{F}_n]\big) = 0,
$$

it follows that

$$
\lim_{m \nearrow \infty} \frac{1}{n_m^{\Delta}} \left( \frac{1}{n_m} \sum_{k=1}^{n_m} \mathbb{E}[Y_k^2] \right)^{1/2} = \infty.
$$

We deduce that  $\{\frac{1}{n_n}\}$  $\frac{1}{n_m}\sum_{k=1}^{n_m} \mathbb{E}[Y_k^2]\}_{m\geq 1}$  diverges faster to infinity than  $\{n_m^{2\Delta}\}_{m\geq 1}$ . Since

$$
\lim_{n \nearrow \infty} \frac{\text{Var}(\mathcal{V}_n)}{n^{1+2\Delta}} = 0,
$$

we can again use eq. [\(2.17\)](#page-8-5) to obtain

$$
\frac{\text{Var}(\mathcal{V}_{n_m})}{n_m^{1+2\Delta}} \ge \frac{1}{n_m^{1+2\Delta}} \text{Var}\big(\mathbb{E}[\lambda(\mathcal{S}_{n_m} \cap \mathcal{S}[n_m,\infty)) \mid \mathcal{F}_{n_m}]\big) + \frac{1}{n_m^{1+2\Delta}} \sum_{k=1}^{n_m} \mathbb{E}[Y_k^2] - \frac{4\sqrt{2}c}{n_m^{3\Delta}} \left(\frac{1}{n_m} \sum_{k=1}^{n_m} \mathbb{E}[Y_k^2]\right)^{1/2}.
$$

We infer that  $\{\frac{1}{n_n}\}$  $\frac{1}{n_m}\sum_{k=1}^{n_m} \mathbb{E}[Y_k^2] \}_{m\geq 1}$  grows faster to infinity than  $\{n_m^{6\Delta}\}_{m\geq 1}$ . By iterating this procedure, we conclude that

(5.2) 
$$
\lim_{m \nearrow \infty} \frac{1}{n_m^2} \sum_{k=1}^{n_m} \mathbb{E}[Y_k^2] = \infty.
$$

On the other hand, from eq.  $(2.16)$  we have

$$
|Y_k| \leq \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) | \mathcal{F}_{k-1}] + \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) | \mathcal{F}_k]
$$
  
+  $\mathbb{E}[\lambda(\mathcal{S}[k-1,k] \setminus \mathcal{S}[k,\infty)) | \mathcal{F}_k] + \mathbb{E}[\lambda(\mathcal{S}[k-1,k] \setminus \mathcal{S}[k,\infty))]$   
 $\leq \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) | \mathcal{F}_{k-1}] + \mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty)) | \mathcal{F}_k]$ 

$$
+\lambda(\mathcal{S}[k-1,k]) + \operatorname{Cap}(\mathcal{B}),
$$

where in the last line we used monotonicity and eq.  $(2.1)$ . By Jensen's inequality,

$$
\mathbb{E}[|Y_k|^2] \le 4\big(2\mathbb{E}[\lambda(\mathcal{S}_{k-1} \cap \mathcal{S}[k-1,\infty))^2] + \mathbb{E}[\lambda(\mathcal{S}[k-1,k])^2] + \mathrm{Cap}^2(\mathcal{B})\big) \le 64 c^2 h(k-1)^2 + 4\mathbb{E}[\mathcal{V}_1^2] + 4(\mathrm{Cap}(\mathcal{B}))^2 \le c_1 k,
$$

<span id="page-22-1"></span>for a constant  $c_1 > 0$ . This yields  $\sum_{k=1}^{n_m} \mathbb{E}[Y_k^2] \le c_1 n_m^2$ , which gives a contradiction.  $\Box$ 

**Lemma 5.3.** In the notation of the proof of Lemma [2.5,](#page-6-4) for any  $\beta \in (0,1]$  there exists a constant  $c(d, \alpha, \beta) > 0$  such that

$$
|\phi(y) - \phi(y - x)| \le c(d, \alpha, \beta) (\phi(y) + \phi(y - x)) \left( \frac{1 + |x|^\beta}{|y|^\beta} \wedge 1 \right), \qquad x, y \in \mathbb{R}^d.
$$

*Proof.* Recall that  $\phi(y) = \mathbb{P}(y \in S_{\infty})$ . This yields

(5.3) 
$$
|\phi(y) - \phi(y - x)| \le \phi(y) + \phi(y - x), \quad x, y \in \mathbb{R}^d.
$$

To establish the second non-trivial part of the claimed inequality, that is, for  $|y|^{\beta} > 1+|x|^{\beta}$ , we first observe that by rotational invariance of X it holds  $\phi(y) = \mathbb{P}_y(\tau_{\mathcal{B}} < \infty)$ . Moreover, by eq.  $(1.4)$ ,

<span id="page-22-0"></span>
$$
\phi(y) = a_{d,\alpha} \int_{\mathcal{B}} |y - w|^{\alpha - d} (1 - |w|^2)^{-\alpha/2} dw, \qquad y \in \mathbb{R}^d,
$$

where

$$
a_{d,\alpha} = \frac{\sin \frac{\pi \alpha}{2} \Gamma((d-\alpha)/2) \Gamma(d/2)}{2^{\alpha} \pi^{d+1} \Gamma(\alpha/2)},
$$

see e.g. [\[30\]](#page-33-15). We fix  $\beta \in (0,1]$  and  $x \in \mathbb{R}^d$ . For any  $y \in \mathcal{B}^c(0,1+|x|)$  we have

$$
|y - w| \ge |y| - |w| \ge |y| - 1 > |x|, \quad w \in \mathcal{B}.
$$

There exists  $x_0 \in \mathcal{B}(0, |y-w|)$  lying on the line going through the origin, determined by the vector  $y - w$ , and such that

$$
\frac{\left||y-w|^{\alpha-d}-|y-w-x|^{\alpha-d}\right||y-w|^{\beta}}{|y-w|^{\alpha-d}+|y-w-x|^{\alpha-d}} = \frac{\left||y-w|^{d-\alpha}-|y-w-x|^{d-\alpha}\right||y-w|^{\beta}}{|y-w|^{d-\alpha}+|y-w-x|^{d-\alpha}} \frac{1+|x|^{\beta}}{1+|x|^{\beta}} \le \frac{\left||y-w|^{d-\alpha}-|y-w-x|^{d-\alpha}}{|y-w|^{d-\alpha}-|y-w-x_0|^{d-\alpha}} \frac{|y-w|^{\beta}}{1+|x_0|^{\beta}}.
$$

Since  $x_0$  is necessarily of the form  $x_0 = \frac{y-w}{|y-w|}$  $\frac{y-w}{|y-w|}$ *o*, for some  $\rho \in [-|y-w|, |y-w|]$ , we have

$$
\frac{\left| |y-w|^{\alpha-d} - |y-w-x|^{\alpha-d} \right|}{|y-w|^{\alpha-d} + |y-w-x|^{\alpha-d}} \frac{|y-w|^\beta}{1+|x|^\beta} \le \frac{\left| |y-w|^{d-\alpha} - \left( |y-w| - \varrho \right)^{d-\alpha} \right|}{|y-w|^{d-\alpha} + \left( |y-w| - \varrho \right)^{d-\alpha}} \frac{|y-w|^\beta}{1+|\varrho|^\beta}.
$$

We investigate the two following cases.

Case 1. We first assume that  $d - \alpha \leq 1$ . If  $\varrho \in [0, |y - w|/2]$  then, by the concavity of the function  $r \mapsto r^{d-\alpha}$ , we obtain

$$
\frac{|y - w|^{\alpha - d} - |y - w - x|^{\alpha - d}|}{|y - w|^{\alpha - d} + |y - w - x|^{\alpha - d}} \frac{|y - w|^{\beta}}{1 + |x|^{\beta}} \le \frac{(d - \alpha)\varrho(|y - w| - \varrho)^{d - \alpha - 1}}{|y - w|^{d - \alpha} + (|y - w| - \varrho)^{d - \alpha}} \frac{|y - w|^{\beta}}{1 + \varrho^{\beta}}
$$
  

$$
\le (d - \alpha) \frac{\varrho}{|y - w| - \varrho} \frac{|y - w|^{\beta}}{1 + \varrho^{\beta}}
$$
  

$$
\le 2(d - \alpha) \frac{\varrho}{|y - w|} \frac{|y - w|^{\beta}}{1 + \varrho^{\beta}}
$$

$$
\leq 2(d-\alpha)\frac{\varrho^{\beta}}{|y-w|^{\beta}}\frac{|y-w|^{\beta}}{1+\varrho^{\beta}}
$$
  

$$
\leq 2(d-\alpha).
$$

If  $\varrho \in [|y - w|/2, |y - w|]$ , then

$$
\frac{\big||y-w|^{\alpha-d}-|y-w-x|^{\alpha-d}\big|}{|y-w|^{\alpha-d}+|y-w-x|^{\alpha-d}}\frac{|y-w|^\beta}{1+|x|^\beta}\,\le\,\frac{|y-w|^\beta}{1+2^{-\beta}|y-w|^\beta}\,\le\,2^\beta
$$

If  $\varrho \in [-|y-w|,0]$  then we again use the concavity argument which yields

$$
\frac{||y - w|^{\alpha - d} - |y - w - x|^{\alpha - d}|}{|y - w|^{\alpha - d} + |y - w - x|^{\alpha - d}} \frac{||y - w|^{\beta}}{1 + |x|^{\beta}} \le \frac{(d - \alpha) |\varrho| |y - w|^{d - \alpha - 1}}{|y - w|^{d - \alpha} + (|y - w| - \varrho)^{d - \alpha}} \frac{||y - w|^{\beta}}{1 + |\varrho|^{\beta}} \le (d - \alpha) \frac{|\varrho|}{|y - w|} \frac{||y - w|^{\beta}}{1 + |\varrho|^{\beta}} \le (d - \alpha) \frac{|\varrho|^{\beta}}{|y - w|^{\beta}} \frac{||y - w|^{\beta}}{1 + |\varrho|^{\beta}} \le d - \alpha.
$$

Case 2. Assume that  $d - \alpha > 1$ . If  $\varrho \in [0, |y - w|]$  then the function  $r \mapsto r^{d-\alpha}$  is convex and we obtain

$$
\frac{\left| |y - w|^{\alpha - d} - |y - w - x|^{\alpha - d} \right| |y - w|^{\beta}}{|y - w|^{\alpha - d} + |y - w - x|^{\alpha - d}} \frac{|y - w|^{\beta}}{1 + |x|^{\beta}} \le \frac{(d - \alpha) \varrho |y - w|^{d - \alpha - 1}}{|y - w|^{d - \alpha} + (|y - w| - \varrho)^{d - \alpha}} \frac{|y - w|^{\beta}}{1 + \varrho^{\beta}}
$$
  

$$
\le (d - \alpha) \frac{\varrho}{|y - w|} \frac{|y - w|^{\beta}}{1 + \varrho^{\beta}}
$$
  

$$
\le (d - \alpha) \frac{\varrho^{\beta}}{|y - w|^{\beta}} \frac{|y - w|^{\beta}}{1 + \varrho^{\beta}}
$$
  

$$
\le d - \alpha.
$$

If  $\varrho \in [-|y-w|, 0]$  then again in view of the convexity we have

$$
\frac{|y - w|^{\alpha - d} - |y - w - x|^{\alpha - d}|}{|y - w|^{\alpha - d} + |y - w - x|^{\alpha - d}} \frac{|y - w|^{\beta}}{1 + |x|^{\beta}} \le \frac{(d - \alpha) |\varrho| (|y - w| + |\varrho|)^{d - \alpha - 1}}{|y - w|^{d - \alpha} + (|y - w| + |\varrho|)^{d - \alpha}} \frac{|y - w|^{\beta}}{1 + |\varrho|^{\beta}}
$$
  
\n
$$
\le (d - \alpha) \frac{|\varrho|}{|y - w| + |\varrho|} \frac{|y - w|^{\beta}}{1 + |\varrho|^{\beta}}
$$
  
\n
$$
\le (d - \alpha) \frac{|\varrho|}{|y - w|} \frac{|y - w|^{\beta}}{1 + |\varrho|^{\beta}}
$$
  
\n
$$
\le (d - \alpha) \frac{|\varrho|^{\beta}}{|y - w|^{\beta}} \frac{|y - w|^{\beta}}{1 + |\varrho|^{\beta}}
$$
  
\n
$$
\le d - \alpha.
$$

Finally, for  $y \in \mathcal{B}^c(0, 1+|x|) \cap \mathcal{B}^c(0, 2)$  we obtain

$$
\frac{||y - w|^{\alpha - d} - |y - w - x|^{\alpha - d}|}{|y - w|^{\alpha - d} + |y - w - x|^{\alpha - d}} \frac{||y|^{\beta}}{1 + |x|^{\beta}}\n= \frac{||y - w|^{\alpha - d} - |y - w - x|^{\alpha - d}|}{|y - w|^{\alpha - d} + |y - w - x|^{\alpha - d}} \frac{||y - w|^{\beta}}{1 + |x|^{\beta}} \frac{||y|^{\beta}}{||y - w|^{\beta}}\n\leq 2^{1 + \beta} (d - \alpha) \vee 2^{2\beta}.
$$

On the other hand, if  $y \in \mathcal{B}^c(0, 1+|x|) \cap \mathcal{B}(0, 2)$  then  $x \in \mathcal{B}$  and

(5.5) 
$$
\frac{1+|x|^{\beta}}{|y|^{\beta}} \geq 2^{-\beta}.
$$

Equations [\(5.3\)](#page-22-0) to [\(5.5\)](#page-24-1) imply the result.  $\square$ 

<span id="page-24-0"></span>Lemma 5.4. In the notation of the proof of Lemma [2.5,](#page-6-4) it holds that

<span id="page-24-1"></span>
$$
\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} p(1,x) \, \phi(y) \, \big| \phi(y) - \phi(y-x) \big| \, \mathrm{d}y \, \mathrm{d}x \, < \, \infty.
$$

Proof. We split the integral into three parts

<span id="page-24-2"></span>(5.6)  
\n
$$
\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} p(1, x) \phi(y) |\phi(y) - \phi(y - x)| dy dx
$$
\n
$$
= \int_{\mathbb{R}^d} \int_{\mathbb{B}^c(0, 1+|x|)} p(1, x) \phi(y) |\phi(y) - \phi(y - x)| dy dx
$$
\n
$$
+ \int_{\mathbb{R}^d} \int_{\mathbb{B}(0, 1+|x|) \cap \mathbb{B}} p(1, x) \phi(y) |\phi(y) - \phi(y - x)| dy dx
$$
\n
$$
+ \int_{\mathbb{R}^d} \int_{\mathbb{B}(0, 1+|x|) \cap \mathbb{B}^c} p(1, x) \phi(y) |\phi(y) - \phi(y - x)| dy dx.
$$

According to Lemma [5.3,](#page-22-1) by setting  $\beta = \alpha/2$  we obtain

$$
|\phi(y) - \phi(y - x)| \le c_1(\phi(y) + \phi(y - x)) \frac{1 + |x|^{\alpha/2}}{|y|^{\alpha/2}}, \qquad y \in \mathcal{B}^c(0, 1 + |x|),
$$

where  $c_1 = c(d, \alpha, \beta)$ . By [\[19,](#page-33-20) Lemma 2.5], there exists a constant  $c_2 = c_2(d, \alpha) > 0$  such that  $\phi(w) \leq c_2 |w|^{\alpha - d}$ , for any  $w \in \mathcal{B}^c$ . Thus,

$$
\int_{\mathbb{R}^d} \int_{\mathbb{B}^c(0,1+|x|)} p(1,x) \phi(y) \left| \phi(y) - \phi(y-x) \right| dy dx
$$
\n
$$
\leq c_1 c_2^2 \int_{\mathbb{R}^d} \int_{\mathbb{B}^c(0,1+|x|)} p(1,x) \frac{1}{|y|^{d-\alpha}} \left( \frac{1}{|y|^{d-\alpha}} + \frac{1}{|y-x|^{d-\alpha}} \right) \frac{1+|x|^{\alpha/2}}{|y|^{\alpha/2}} dy dx
$$
\n
$$
\leq 2^{d-\alpha/2} (1+2^{d-\alpha}) c_1 c_2^2 \int_{\mathbb{R}^d} \int_{\mathbb{B}^c(0,1+|x|)} p(1,x) (1+|x|^{\alpha/2}) \frac{1}{|y-x|^{2d-3\alpha/2}} dy dx
$$
\n
$$
\leq 2^{d-\alpha/2} (1+2^{d-\alpha}) c_1 c_2^2 \int_{\mathbb{R}^d} p(1,x) (1+|x|^{\alpha/2}) dx \int_{\mathbb{B}^c} \frac{1}{|z|^{2d-3\alpha/2}} dz
$$
\n
$$
= 2^{d-\alpha/2} (1+2^{d-\alpha}) c_1 c_2^2 d\lambda(\mathbb{B}) \int_{\mathbb{R}^d} p(1,x) (1+|x|^{\alpha/2}) dx \int_{1}^{\infty} \frac{1}{r^{d-3\alpha/2+1}} dr,
$$

where in the second step we used the fact that  $|y - x| \le |y| + |x| \le |y| + |y| - 1 \le 2|y|$ . The last integral is finite as  $d/\alpha > 3/2$  and X has finite  $\beta$ -moment for any  $\beta < \alpha$  (see [\[25,](#page-33-21) Example 25.10]).

For the second integral on the right-hand side of eq.  $(5.6)$  we observe that

$$
\int_{\mathbb{R}^d} \int_{\mathcal{B}(0,1+|x|)\cap \mathcal{B}} p(1,x) \, \phi(y) \, \big| \phi(y) - \phi(y-x) \big| \, \mathrm{d}y \, \mathrm{d}x \, \leq \, 2\,\lambda(\mathcal{B}).
$$

The third integral on the right-hand side of eq. [\(5.6\)](#page-24-2) is most demanding. We start by splitting this integral into two parts

<span id="page-25-0"></span>(5.7) 
$$
\int_{\mathbb{R}^d} \int_{\mathbb{B}(0,1+|x|)\cap \mathbb{B}^c} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx
$$

$$
= \int_{\mathbb{B}(0,\Lambda)} \int_{\mathbb{B}(0,1+|x|)\cap \mathbb{B}^c} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx
$$

$$
+ \int_{\mathbb{B}^c(0,\Lambda)} \int_{\mathbb{B}(0,1+|x|)\cap \mathbb{B}^c} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx.
$$

For the first integral in this decomposition we have

$$
\int_{\mathcal{B}(0,\Lambda)} \int_{\mathcal{B}(0,1+|x|)\cap\mathcal{B}^c} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx
$$
  
\n
$$
\leq 2 c_2 \int_{\mathcal{B}(0,\Lambda)} \int_{\mathcal{B}(0,1+|x|)\cap\mathcal{B}^c} p(1,x) \frac{1}{|y|^{d-\alpha}} dy dx
$$
  
\n
$$
\leq 2 c_2 \int_{\mathcal{B}(0,1+\Lambda)\cap\mathcal{B}^c} \frac{1}{|y|^{d-\alpha}} dy
$$
  
\n
$$
\leq 2 c_2 \lambda(\mathcal{B}(0,1+\Lambda)).
$$

The second integral on the right-hand side of eq. [\(5.7\)](#page-25-0) we decompose further as follows

<span id="page-25-1"></span>
$$
\int_{\mathbb{B}^{c}(0,\Lambda)} \int_{\mathbb{B}(0,1+|x|)\cap\mathbb{B}^{c}} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx
$$
\n
$$
= \int_{\mathbb{B}^{c}(0,\Lambda)} \int_{\mathbb{B}(0,1+|x|)\cap\mathbb{B}^{c}\cap\{z:\|z-x|>\|z\| \}} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx
$$
\n
$$
+ \int_{\mathbb{B}^{c}(0,\Lambda)} \int_{\mathbb{B}(0,1+|x|)\cap\mathbb{B}^{c}\cap\{z:\|z-x|\le|z|\}} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx
$$
\n
$$
+ \int_{\mathbb{B}^{c}(0,\Lambda)} \int_{\mathbb{B}(0,1+|x|)\cap\mathbb{B}^{c}\cap\{z:\|z-x|<1\}} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx.
$$

It is well-known that for any  $\Lambda > 1$  large enough there is  $c_3 = c_3(d, \alpha, \Lambda) > 0$  such that

$$
p(1,x) \le \frac{c_3}{|x|^{d+\alpha}}, \qquad x \in \mathcal{B}^c(0,\Lambda).
$$

We set  $A_1 = \mathcal{B}(0, 1 + |x|) \cap \mathcal{B}^c \cap \{z : |z - x| > |z|\}$  and estimate the first integral on the right-hand side of eq. [\(5.8\)](#page-25-1) as follows

$$
\int_{\mathcal{B}^{c}(0,\Lambda)} \int_{A_1} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx
$$
\n
$$
\leq c_2^2 c_3 \int_{\mathcal{B}^{c}(0,\Lambda)} \int_{A_1} \frac{1}{|x|^{d+\alpha}} \frac{1}{|y|^{d-\alpha}} \left( \frac{1}{|y|^{d-\alpha}} + \frac{1}{|y-x|^{d-\alpha}} \right) dy dx
$$
\n
$$
\leq 2 c_2^2 c_3 \int_{\mathcal{B}^{c}(0,\Lambda)} \int_{A_1} \frac{1}{|x|^{d+\alpha}} \frac{1}{|y|^{2d-2\alpha}} dy dx
$$
\n
$$
\leq 2 c_2^2 c_3 \int_{\mathcal{B}^{c}(0,\Lambda)} \int_{\mathcal{B}(0,1+|x|)\cap\mathcal{B}^{c}} \frac{1}{|x|^{d+\alpha}} \frac{1}{|y|^{2d-2\alpha}} dy dx
$$
\n
$$
\leq 2 c_2^2 c_3 d\lambda(\mathcal{B}) \int_{\mathcal{B}^{c}(0,\Lambda)} \frac{1}{|x|^{d+\alpha}} \int_{1}^{1+|x|} \frac{1}{r^{d-2\alpha+1}} dr dx.
$$

The last integral is finite as  $d/\alpha > 3/2$ .

We set  $A_2 = \mathcal{B}(0, 1 + |x|) \cap \mathcal{B}^c \cap \{z : 1 \leq |z - x| \leq |z|\}$  and for the second integral on the right-hand side of eq. [\(5.8\)](#page-25-1) we have

$$
\int_{\mathcal{B}^{c}(0,\Lambda)} \int_{A_2} p(1,x) \phi(y) |\phi(y) - \phi(y-x)| dy dx
$$
\n
$$
\leq c_2^2 c_3 \int_{\mathcal{B}^{c}(0,\Lambda)} \int_{A_2} \frac{1}{|x|^{d+\alpha}} \frac{1}{|y|^{d-\alpha}} \left( \frac{1}{|y|^{d-\alpha}} + \frac{1}{|y-x|^{d-\alpha}} \right) dy dx
$$
\n
$$
\leq 2^{d+\alpha+1} c_2^2 c_3 \int_{\mathcal{B}^{c}(0,\Lambda)} \int_{A_2} \frac{1}{|y|^{2d}} \frac{1}{|y-x|^{d-\alpha}} dy dx
$$
\n
$$
\leq 2^{d+\alpha+1} c_2^2 c_3 \int_{\mathcal{B}^c} \int_{\{z:1 \leq |z-x| \leq |z|\}} \frac{1}{|y|^{2d}} \frac{1}{|y-x|^{d-\alpha}} dx dy
$$
\n
$$
\leq 2^{d+\alpha+1} c_2^2 c_3 \int_{\mathcal{B}^c} \frac{1}{|y|^{2d}} \int_{\mathcal{B}(0,|y|) \cap \mathcal{B}^c} \frac{1}{|z|^{d-\alpha}} dz dy
$$
\n
$$
\leq 2^{d+\alpha+1} c_2^2 c_3 d \alpha^{-1} \lambda(\mathcal{B}) \int_{\mathcal{B}^c} \frac{1}{|y|^{2d-\alpha}} dy
$$
\n
$$
= 2^{d+\alpha+1} c_2^2 c_3 d^2 \alpha^{-1} \lambda(\mathcal{B})^2 \int_1^\infty \frac{1}{r^{d-\alpha+1}} dr,
$$

where in the second step we used the fact that  $|y| \leq 2|x|$ .

Finally, we set  $A_3 = \mathcal{B}(0, 1 + |x|) \cap \mathcal{B}^c \cap \{z : |z - x| < 1\}$  and for the third integral on the right-hand side of eq. [\(5.8\)](#page-25-1) we proceed as follows

$$
\int_{\mathcal{B}^c(0,\Lambda)} \int_{A_3} p(1,x) \phi(y) |\phi(y) - \phi(y - x)| dy dx
$$
  
\n
$$
\leq 2 c_2 c_3 \int_{\mathcal{B}^c(0,\Lambda)} \int_{A_3} \frac{1}{|x|^{d+\alpha}} \frac{1}{|y|^{d-\alpha}} dy dx
$$
  
\n
$$
\leq 2^{1+d-\alpha} c_2 c_3 \int_{\mathcal{B}^c(0,\Lambda)} \int_{\{z:|z-x|<1\}} \frac{1}{|x|^{2d}} dy dx
$$
  
\n
$$
\leq 2^{1+d-\alpha} c_2 c_3 \lambda(\mathcal{B}) \int_{\Lambda}^{\infty} \frac{1}{r^{d+1}} dr,
$$

<span id="page-26-0"></span>where in the second step we used the fact that  $|x| \le |y - x| + |y| \le 1 + |y| \le 2|y|$ .  $\Box$ **Lemma 5.5.** In the notation of the proof of Lemma  $2.5$ , it holds that

$$
\sup_{x\in\mathcal{B}}\int_{\mathbb{R}^d}\phi(y)|\phi(y)-\phi(y-x)|\,\mathrm{d}y\,<\,\infty.
$$

Proof. We split the integral as follows

$$
\int_{\mathbb{R}^d} \phi(y) |\phi(y) - \phi(y - x)| \, dy = \int_{\mathcal{B}(0, 1+|x|)} \phi(y) |\phi(y) - \phi(y - x)| \, dy + \int_{\mathcal{B}^c(0, 1+|x|)} \phi(y) |\phi(y) - \phi(y - x)| \, dy.
$$

For any  $x \in \mathcal{B}$  one has  $\mathcal{B}(0, 1+|x|) \subseteq \mathcal{B}(0, 2)$ , and whence

$$
\sup_{x\in\mathcal{B}}\int_{\mathcal{B}(0,1+|x|)}\phi(y)|\phi(y)-\phi(y-x)|\,\mathrm{d}y\,\leq\,2\,\lambda(\mathcal{B}(0,2)).
$$

By Lemma [5.3](#page-22-1) with  $\beta = \alpha/2$ , for a constant  $c_1 = c(d, \alpha, \beta)$ ,

$$
|\phi(y) - \phi(y - x)| \le c_1(\phi(y) + \phi(y - x)) \frac{1 + |x|^{\alpha/2}}{|y|^{\alpha/2}}, \qquad y \in \mathcal{B}^c(0, 1 + |x|).
$$

By [\[19,](#page-33-20) Lemma 2.5], there exists a constant  $c_2 = c_2(d, \alpha) > 0$  such that  $\phi(w) \le c_2 |w|^{\alpha-d}$ , for any  $w \in \mathcal{B}^c$ . Thus, for  $x \in \mathcal{B}$  we have

$$
\int_{\mathcal{B}^c(0,1+|x|)} \phi(y)|\phi(y) - \phi(y-x)| \,dy
$$
\n
$$
\leq 2^{d-\alpha/2+1} (1+2^{d-\alpha}) c_1 c_2^2 \int_{\mathcal{B}^c(0,1+|x|)} \frac{1}{|y-x|^{2d-3\alpha/2}} \,dy
$$
\n
$$
\leq 2^{d-\alpha/2+1} (1+2^{d-\alpha}) c_1 c_2^2 d\lambda(\mathcal{B}) \int_1^\infty \frac{1}{r^{d-3\alpha/2+1}} \,dr,
$$

<span id="page-27-0"></span>where we used the fact that  $|y - x| \le 2|y|$ . The assertion follows as  $d/\alpha > 3/2$ .

**Lemma 5.6.** There exists a constant  $\tilde{c} > 0$  such that for all  $t > 0$  large enough,

$$
\mathbb{E}[\langle \mathcal{V}_t \rangle^4] \leq \tilde{c} t^2.
$$

*Proof.* By setting  $n = 2$  in eq.  $(2.33)$ , we have

$$
\mathcal{V}_t = \lambda(\mathcal{S}_{t/2}^{(1)}) + \lambda(\mathcal{S}_{t/2}^{(2)}) - \lambda(\mathcal{S}_{t/2}^{(1)} \cap \mathcal{S}_{t/2}^{(2)}),
$$

where  $\mathcal{S}_{t/2}^{(1)}$  $t_{t/2}^{(1)}$  and  $\mathcal{S}_{t/2}^{(2)}$ (2) are independent, and have the same law as  $\mathcal{S}_{t/2}$ . Let  $\mathcal{V}_{t/2}^{(i)} = \lambda(\mathcal{S}_{t/2}^{(i)})$  $t_{t/2}^{(i)}$ ), for  $i = 1, 2$ . Taking expectation in the last equation and then subtracting the two relations yields

<span id="page-27-2"></span>
$$
\langle \mathcal{V}_t \rangle \,=\, \langle \mathcal{V}^{(1)}_{t/2} \rangle + \langle \mathcal{V}^{(2)}_{t/2} \rangle - \langle \lambda(\mathcal{S}^{(1)}_{t/2} \cap \mathcal{S}^{(2)}_{t/2}) \rangle.
$$

By the triangle inequality,<sup>[2](#page-27-1)</sup>

(5.9) 
$$
\|\langle \mathcal{V}_t \rangle\|_4 \leq \|\langle \mathcal{V}_{t/2}^{(1)} \rangle + \langle \mathcal{V}_{t/2}^{(2)} \rangle\|_4 + \|\langle \lambda(\mathcal{S}_{t/2}^{(1)} \cap \mathcal{S}_{t/2}^{(2)}) \rangle\|_4,
$$

Jensen's inequality and Lemma [2.3](#page-4-4) imply that there is a constant  $c_1 > 0$  such that

$$
(5.10) \quad \|\langle \lambda(\mathcal{S}_{t/2}^{(1)} \cap \mathcal{S}_{t/2}^{(2)}) \rangle \|_{4} \leq \|\lambda(\mathcal{S}_{t/2}^{(1)} \cap \mathcal{S}_{t/2}^{(2)}) \|_{4} + \mathbb{E}[\lambda(\mathcal{S}_{t/2}^{(1)} \cap \mathcal{S}_{t/2}^{(2)})] \leq 2 \|\lambda(\mathcal{S}_{t/2}^{(1)} \cap \mathcal{S}_{t/2}^{(2)}) \|_{4}
$$
  

$$
\leq 2 \|\lambda(\mathcal{S}_{t/2}^{(1)} \cap \mathcal{S}_{\infty}^{(2)}) \|_{4} \leq c_{1} h(t/2) \leq c_{1} \sqrt{t}.
$$

By the independence of the variables  $\langle V_{t/2}^{(1)} \rangle$  and  $\langle V_{t/2}^{(2)} \rangle$ , we have

$$
\mathbb{E}\left[\left(\langle \mathcal{V}^{(1)}_{t/2}\rangle+\langle \mathcal{V}^{(2)}_{t/2}\rangle\right)^4\right]=\mathbb{E}\left[\langle \mathcal{V}^{(1)}_{t/2}\rangle^4\right]+\mathbb{E}\left[\langle \mathcal{V}^{(2)}_{t/2}\rangle^4\right]+6\,\mathbb{E}\left[\langle \mathcal{V}^{(1)}_{t/2}\rangle^2\right]\mathbb{E}\left[\langle \mathcal{V}^{(2)}_{t/2}\rangle^2\right].
$$

By Lemma [2.5,](#page-6-4) there exists  $N \in \mathbb{N}$  large enough such that  $Var(\mathcal{V}_t) \le c_2 t$  for all  $t \ge 2^N$ and some  $c_2 > 0$ . Hence, for  $t \geq 2^{N+1}$ ,

$$
\mathbb{E}\left[\left(\langle \mathcal{V}^{(1)}_{t/2} \rangle + \langle \mathcal{V}^{(2)}_{t/2} \rangle \right)^4\right] \leq \mathbb{E}\left[\langle \mathcal{V}^{(1)}_{t/2} \rangle^4\right] + \mathbb{E}\left[\langle \mathcal{V}^{(2)}_{t/2} \rangle^4\right] + 6\left(\frac{c_2 t}{2}\right)^2
$$

.

By combining this with the elementary inequality  $(a + b)^{1/4} \le a^{1/4} + b^{1/4}$ , we arrive at

(5.11) 
$$
\|\langle \mathcal{V}_{t/2}^{(1)} \rangle + \langle \mathcal{V}_{t/2}^{(2)} \rangle\|_4 \le \left( \mathbb{E}\left[\langle \mathcal{V}_{t/2}^{(1)} \rangle^4\right] + \mathbb{E}\left[\langle \mathcal{V}_{t/2}^{(2)} \rangle^4\right] \right)^{1/4} + c_3 \sqrt{t}
$$

with  $c_3 = (3c_2^2/2)^{1/4}$ . From eqs. [\(5.9\)](#page-27-2) to [\(5.11\)](#page-27-3) it follows that there is  $c_4 > 0$  such that

<span id="page-27-3"></span>
$$
\|\langle \mathcal{V}_t \rangle\|_4 \leq \left(\mathbb{E}\left[\langle \mathcal{V}_{t/2}^{(1)} \rangle^4\right] + \mathbb{E}\left[\langle \mathcal{V}_{t/2}^{(2)} \rangle^4\right]\right)^{1/4} + c_4\sqrt{t}
$$

For  $k > N$  we set

$$
\gamma_k = \sup \{ ||\langle \mathcal{V}_t \rangle ||_4 : 2^k \le t < 2^{k+1} \}.
$$

<span id="page-27-1"></span><sup>2</sup>For a random variable Y we write  $||Y||_p = (E[|Y|^p])^{1/p}$ , for any  $p \ge 1$ .

Thus, for  $k \geq N+1$  and for every  $2^k \leq t < 2^{k+1}$  we have

$$
\|\langle \mathcal{V}_t \rangle\|_4 \le (\gamma_{k-1}^4 + \gamma_{k-1}^4)^{1/4} + c_5 2^{k/2}
$$

with  $c_5 = \sqrt{2} c_4$ . Taking supremum over  $2^k \le t < 2^{k+1}$  yields  $\gamma_k \leq 2^{1/4} \gamma_{k-1} + c_5 2^{k/2}.$ 

We set  $\delta_k = \gamma_k/2^{k/2}$  and we divide the last inequality by  $2^{k/2}$ . We thus have

$$
\delta_k \le \frac{2^{1/4} \gamma_{k-1}}{2^{1/2} 2^{(k-1)/2}} + c_5 = 2^{-1/4} \delta_{k-1} + c_5.
$$

<span id="page-28-0"></span>By iterating this inequality we finally conclude the result.  $\Box$ 

Lemma 5.7. The following expansion is valid

$$
Var(\mathcal{V}_t) = \sigma^2 t + \mathcal{O}(t^{1/2}h(t)), \qquad t \ge 1,
$$

where the function  $h(t)$  is defined in eq.  $(2.7)$ .

*Proof.* For every  $s, t \geq 0$  we have

$$
\mathcal{V}_{s+t} = \lambda \big( \mathcal{S}_s \cup \mathcal{S}[s, s+t] \big) = \lambda(\mathcal{S}_s^{(1)}) + \lambda(\mathcal{S}_t^{(2)}) - \lambda(\mathcal{S}_s^{(1)} \cap \mathcal{S}_t^{(2)}).
$$

This implies

$$
\mathcal{V}^{(1)}_s + \mathcal{V}^{(2)}_t - \lambda \big( \mathcal{S}^{(1)}_{s+t} \cap \mathcal{S}^{(2)}_{s+t} \big) \, \leq \, \mathcal{V}_{s+t} \, \leq \, \mathcal{V}^{(1)}_s + \mathcal{V}^{(2)}_t,
$$

and whence

$$
\langle \mathcal{V}_s^{(1)} \rangle + \langle \mathcal{V}_t^{(2)} \rangle - \lambda \big( \mathcal{S}_{s+t}^{(1)} \cap \mathcal{S}_{s+t}^{(2)} \big) \leq \langle \mathcal{V}_{s+t} \rangle \leq \langle \mathcal{V}_s^{(1)} \rangle + \langle \mathcal{V}_t^{(2)} \rangle + \mathbb{E}[\lambda(\mathcal{S}_{s+t}^{(1)} \cap \mathcal{S}_{s+t}^{(2)})].
$$

Here,  $S_s^{(1)}$  and  $S_t^{(2)}$  are independent and have the same law as  $S_s$  and  $S_t$ , respectively. We set  $\mathcal{I}_t = \lambda(\mathcal{S}_t \cap \mathcal{S}'_t)$ , where  $\mathcal{S}'_t$  is an independent copy of  $\mathcal{S}_t$ . From the previous relation we obtain

<span id="page-28-2"></span>
$$
|\langle \mathcal{V}_{s+t} \rangle - (\langle \mathcal{V}_s^{(1)} \rangle + \langle \mathcal{V}_t^{(2)} \rangle)| \leq \mathcal{I}_{s+t} + \mathbb{E}[\mathcal{I}_{s+t}] \leq \mathcal{I}_{s+t} + ||\mathcal{I}_{s+t}||_2.
$$

Hence

(5.12) 
$$
\|\langle \mathcal{V}_{s+t} \rangle - (\langle \mathcal{V}_s^{(1)} \rangle + \langle \mathcal{V}_t^{(2)} \rangle)\|_2 \leq 2\|\mathcal{I}_{s+t}\|_2,
$$

and

<span id="page-28-1"></span>
$$
(5.13) \quad \|\langle \mathcal{V}_{s+t} \rangle\|_2^2 \le \|\langle \mathcal{V}_s \rangle\|_2^2 + \|\langle \mathcal{V}_t \rangle\|_2^2 + 4 \left( \|\langle \mathcal{V}_s \rangle\|_2^2 + \|\langle \mathcal{V}_t \rangle\|_2^2 \right)^{1/2} \|\mathcal{I}_{s+t}\|_2 + 4 \|\mathcal{I}_{s+t}\|_2^2.
$$

By eq. [\(2.6\)](#page-5-4), there are  $c_1 > 0$  and  $t_1 > 1$  such that  $\|\mathcal{I}_t\|_2 \leq c_1 h(t)$  for  $t \geq t_1$ . For  $t \in [1, t_1]$ we clearly have  $\mathcal{I}_t \leq \mathcal{I}_{t_1}$ . Thus, there is a constant  $c_2 > 0$  such that,

 $1/2$ 

$$
\|\mathcal{I}_t\|_2 \le c_2 h(t), \qquad t \ge 1.
$$

Moreover, from eq. [\(2.9\)](#page-5-3) we have that there exist  $c_3 > 0$  and  $t_2 > 1$  such that  $Var(\mathcal{V}_t) \le c_3 t$ for  $t \ge t_2$ , and for  $t \in [1, t_2]$  we have  $\text{Var}(\mathcal{V}_t) \le \mathbb{E}[\mathcal{V}_t^2] \le \mathbb{E}[\mathcal{V}_{t_2}^2] t$ . Hence

$$
Var(\mathcal{V}_t) \le (c_3 + \mathbb{E}[\mathcal{V}_{t_2}^2])t, \qquad t \ge 1.
$$

We conclude that there is  $c_4 > 0$  such that

(5.14)  $\|\langle \mathcal{V}_t \rangle\|_2 \leq c_4 \sqrt{2}$ and  $||\mathcal{I}_t||_2 \le c_4 h(t), \quad t \ge 1.$ 

By eq.  $(5.13)$ , we obtain

<span id="page-28-3"></span>
$$
\begin{aligned} \|\langle \mathcal{V}_{s+t} \rangle\|_2^2 &\leq \|\langle \mathcal{V}_s \rangle\|_2^2 + \|\langle \mathcal{V}_t \rangle\|_2^2 + 4c_4^2\sqrt{s+t} \, h(s+t) + 4c_4^2(h(s+t))^2 \\ &\leq \|\langle \mathcal{V}_s \rangle\|_2^2 + \|\langle \mathcal{V}_t \rangle\|_2^2 + c_5\sqrt{s+t} \, h(s+t), \end{aligned}
$$

for some constant  $c_5 > 0$ . Similarly as above, in view of eq. [\(5.12\)](#page-28-2) we have

$$
\|\langle \mathcal{V}_s^{(1)} \rangle + \langle \mathcal{V}_t^{(2)} \rangle\|_2 \leq \|\langle \mathcal{V}_{s+t} \rangle\|_2 + \|\langle \mathcal{V}_{s+t} \rangle - (\langle \mathcal{V}_s^{(1)} \rangle + \langle \mathcal{V}_t^{(2)} \rangle)\|_2
$$

 $\leq \|\langle \mathcal{V}_{s+t}\rangle\|_2 + 2\|\mathcal{I}_{s+t}\|_2,$ 

which implies

$$
\|\langle \mathcal{V}_s \rangle\|_2^2 + \|\langle \mathcal{V}_t \rangle\|_2^2 \le \|\langle \mathcal{V}_{s+t} \rangle\|_2^2 + 4 \|\langle \mathcal{V}_{s+t} \rangle\|_2 \|\mathcal{I}_{s+t}\|_2 + 4 \|\mathcal{I}_{s+t}\|_2^2.
$$
  
By eq. (5.14),

$$
\begin{aligned} \|\langle \mathcal{V}_s \rangle\|_2^2 + \|\langle \mathcal{V}_t \rangle\|_2^2 &\leq \|\langle \mathcal{V}_{s+t} \rangle\|_2^2 + 4c_4^2\sqrt{s+t} \, h(s+t) + 4c_4^2(h(s+t))^2 \\ &\leq \|\langle \mathcal{V}_{s+t} \rangle\|_2^2 + +c_5\sqrt{s+t} \, h(s+t). \end{aligned}
$$

We set

$$
x_t = \text{Var}(\mathcal{V}_t) = ||\langle \mathcal{V}_t \rangle||_2^2 \quad \text{and} \quad b_t = c_5 \sqrt{t} \, h(t), \qquad t > 0,
$$
\nand we have shown that

$$
x_s + x_t - b_{s+t} \le x_{s+t} \le x_s + x_t + b_{s+t}, \qquad s, t \ge 1.
$$

By Lemma [2.5](#page-6-4) we know that

$$
\lim_{t \nearrow \infty} \frac{x_t}{t} = \sigma^2 > 0.
$$

Take  $s = t = 2^{k-1}r$  for  $k \in \mathbb{N}$  and  $r \in \mathbb{R}$ ,  $r \ge 1$ . We easily verify that

$$
\left|\frac{x_{2^kr}}{2^kr} - \frac{x_{2^{k-1}r}}{2^{k-1}r}\right| \le \frac{b_{2^kr}}{2^kr}, \qquad k \in \mathbb{N}, \ r \ge 1.
$$

Next, we observe that

$$
\sum_{k=1}^{\infty} \left( \frac{x_{2^k r}}{2^k r} - \frac{x_{2^{k-1} r}}{2^{k-1} r} \right) = \lim_{N \nearrow \infty} \sum_{k=1}^N \left( \frac{x_{2^k r}}{2^k r} - \frac{x_{2^{k-1} r}}{2^{k-1} r} \right) = \sigma^2 - \frac{x_r}{r}, \qquad r \ge 1,
$$

and whence

$$
\left|\frac{x_t}{t} - \sigma^2\right| \ = \ \left|\sum_{k=1}^{\infty} \left(\frac{x_{2^kt}}{2^kt} - \frac{x_{2^{k-1}t}}{2^{k-1}t}\right)\right| \ \leq \ \sum_{k=1}^{\infty} \frac{b_{2^kt}}{2^kt}, \qquad t \geq 1.
$$

This yields

$$
\left|\frac{x_t}{t} - \sigma^2\right| \le \sum_{k=1}^{\infty} \frac{c_5 \sqrt{2^k t} \, h(2^k t)}{2^k t} \le \frac{c_5}{\sqrt{t}} \sum_{k=1}^{\infty} \frac{h(2^k t)}{2^{k/2}}, \qquad t \ge 1.
$$

*Case (i)*. For  $\Delta \in (0, 1/2)$  we have

$$
\left|\frac{x_t}{t} - \sigma^2\right| \le \frac{c_5}{\sqrt{t}} \sum_{k=1}^{\infty} \frac{(2^k t)^{1/2 - \Delta}}{2^{k/2}} = \frac{c_5}{t^{\Delta}} \sum_{k=1}^{\infty} (2^{-\Delta})^k = c_6 t^{-\Delta}, \qquad t \ge 1,
$$

where  $c_6 = c_5 \sum_{k=1}^{\infty} 2^{-\Delta k}$ . It follows that

$$
|x_t - \sigma^2 t| \le c_6 t^{1-\Delta} = c_6 t^{1/2} h(t), \qquad t \ge 1.
$$

Case (ii). If  $\Delta \geq 1/2$ , then  $h(t)$  is slowly varying. According to [\[2,](#page-32-12) Theorem 1.5.6] there is a constant  $c_7 > 0$  such that  $h(2^k t) \le c_7 2^{k/4} h(t)$  for all  $k \in \mathbb{N}$  and  $t \ge 1$ . We obtain

$$
\left|\frac{x_t}{t} - \sigma^2\right| \le \frac{c_5}{\sqrt{t}} \sum_{k=1}^{\infty} \frac{c_7 2^{k/4} h(t)}{2^{k/2}} = \frac{c_5 c_7 h(t)}{\sqrt{t}} \sum_{k=1}^{\infty} 2^{-k/4} = c_8 t^{-1/2} h(t), \qquad t \ge 1,
$$

<span id="page-29-0"></span>with  $c_8 = c_5 c_7 \sum_{k=1}^{\infty} 2^{-k/4}$ , and the proof is finished.

**Lemma 5.8.** Assume that  $d/\alpha > 9/5$ . Then, for the process  $\{\mathcal{J}_{n_i}\}_{i\geq 0}$  defined in eq. [\(4.5\)](#page-17-4), it holds that

$$
\lim_{i \nearrow \infty} \frac{\left| \sum_{j=0}^{i-1} \langle \mathcal{J}_{n_j} \rangle \right|}{\sqrt{n_i/\log \log n_i}} = 0 \qquad \mathbb{P}\text{-}a.s.
$$

*Proof.* For  $i \geq 2$  we clearly have

$$
\text{Var}\left(\sum_{j=1}^{i-1}\mathcal{J}_{n_j}\right) \leq \sum_{j=1}^{i-1}\mathbb{E}[\mathcal{J}_{n_j}^2]+2\sum_{j=1}^{i-1}\sum_{k=1}^{j-1}\mathbb{E}[\mathcal{J}_{n_j}\mathcal{J}_{n_k}]-2\sum_{j=1}^{i-1}\sum_{k=1}^{j-1}\mathbb{E}[\mathcal{J}_{n_j}]\,\mathbb{E}[\mathcal{J}_{n_k}].
$$

Let  $S_t$  be an independent copy of  $S_t$ . Then

$$
\mathcal{J}_{n_j} \stackrel{\text{(d)}}{=} \lambda(\mathcal{S}'[0, n_{j+1} - n_j] \cap \mathcal{S}[0, n_j])
$$

for  $j = 1, \ldots, i - 1$ . Jensen's inequality and Corollary [2.4](#page-5-5) imply that, for some  $c_1 > 0$ ,

$$
\mathbb{E}[\mathcal{J}_{n_j}]^2 \leq \mathbb{E}[\mathcal{J}_{n_j}^2] \leq c_1 h(n_i)^2.
$$

Further, for any  $j = 1, \ldots, i - 1$  and  $k = 1, \ldots, j - 1$ , it holds that

$$
\mathcal{J}_{n_j} = \lambda(\mathcal{S}(n_j, n_{j+1}] \cap \mathcal{S}[0, n_j])
$$
  
=  $\lambda(\mathcal{S}(n_j, n_{j+1}] \cap \mathcal{S}[n_{k+1}, n_j]) + \lambda(\mathcal{S}(n_j, n_{j+1}] \cap (\mathcal{S}[0, n_{k+1}] \setminus \mathcal{S}[n_{k+1}, n_j])).$ 

We set

$$
\mathcal{J}_{j,k}^{(1)} = \lambda(\mathcal{S}(n_j, n_{j+1}] \cap \mathcal{S}[n_{k+1}, n_j]),
$$
  

$$
\mathcal{J}_{j,k}^{(2)} = \lambda(\mathcal{S}(n_j, n_{j+1}] \cap (\mathcal{S}[0, n_{k+1}] \setminus \mathcal{S}[n_{k+1}, n_j])).
$$

Due to independence,

$$
\mathbb{E}[\mathcal{J}_{n_k}\mathcal{J}_{j,k}^{(1)}] = \mathbb{E}[\mathcal{J}_{n_k}]\,\mathbb{E}[\mathcal{J}_{j,k}^{(1)}].
$$

Thus,

$$
\operatorname{Var}\left(\sum_{j=1}^{i-1} \mathcal{J}_{n_j}\right) \leq \sum_{j=1}^{i-1} \mathbb{E}[\mathcal{J}_{n_j}^2] + 2 \sum_{j=1}^{i-1} \sum_{k=1}^{j-1} \mathbb{E}[\mathcal{J}_{n_k} \mathcal{J}_{j,k}^{(2)}] - 2 \sum_{j=1}^{i-1} \sum_{k=1}^{j-1} \mathbb{E}[\mathcal{J}_{n_k}] \mathbb{E}[\mathcal{J}_{j,k}^{(2)}].
$$

Further,

$$
\sum_{j=1}^{i-1} \sum_{k=0}^{j-1} \mathcal{J}_{n_k} \mathcal{J}_{j,k}^{(2)}
$$
\n
$$
= \sum_{j=1}^{i-1} \sum_{k=0}^{j-1} \lambda(\mathcal{S}(n_k, n_{k+1}] \cap \mathcal{S}[0, n_k]) \lambda(\mathcal{S}(n_j, n_{j+1}] \cap (\mathcal{S}[0, n_{k+1}] \setminus \mathcal{S}[n_{k+1}, n_j]))
$$
\n
$$
= \sum_{k=0}^{i-2} \lambda(\mathcal{S}(n_k, n_{k+1}] \cap \mathcal{S}[0, n_k]) \sum_{j=k+1}^{i-1} \lambda(\mathcal{S}(n_j, n_{j+1}] \cap (\mathcal{S}[0, n_{k+1}] \setminus \mathcal{S}[n_{k+1}, n_j]))
$$
\n
$$
\leq \sum_{k=0}^{i-2} \lambda(\mathcal{S}(n_k, n_{k+1}] \cap \mathcal{S}[0, n_k]) \lambda(\mathcal{S}(n_{k+1}, n_i] \cap \mathcal{S}[0, n_{k+1}]).
$$

Similarly as before, by Corollary [2.4,](#page-5-5) we obtain

$$
\mathbb{E}[\lambda(\mathcal{S}(n_k, n_{k+1}] \cap \mathcal{S}[0, n_k])^2] \le c_1 h(n_i)^2,
$$
  

$$
\mathbb{E}[\lambda(\mathcal{S}(n_{k+1}, n_i] \cap \mathcal{S}[0, n_{k+1}])^2] \le c_1 h(n_i)^2.
$$

This implies

$$
\sum_{j=1}^{i-1} \sum_{k=0}^{j-1} \mathbb{E}[\mathcal{J}_{n_k} \mathcal{J}_{j,k}^{(2)}] \leq c_1 i h(n_i)^2.
$$

Analogously we can show that

$$
\sum_{j=1}^{i-1} \sum_{k=0}^{j-1} \mathbb{E}[\mathcal{J}_{n_k}] \, \mathbb{E}[\mathcal{J}_{j,k}^{(2)}] \, \leq \, c_1 i h(n_i)^2.
$$

Thus,

$$
\operatorname{Var}\left(\sum_{j=0}^{i-1} \mathcal{J}_{n_j}\right) \le c_2 ih(n_i)^2
$$

for some  $c_2 > 0$ .

Let  $2^k \leq n_i < 2^{k+1}$ , and  $\Lambda = 2 - d/\alpha$  when  $d/\alpha \in (1, 2)$ . Then

$$
ih(n_i)^2 \le h(2^{k+1})^2 \sum_{j=1}^k j2^{j/2} = \begin{cases} \mathcal{O}(n_i^{1/2} \log n_i), & d/\alpha > 2, \\ \mathcal{O}(n_i^{1/2} (\log n_i)^3), & d/\alpha = 2, \\ \mathcal{O}(n_i^{2\Lambda + 1/2} \log n_i), & d/\alpha \in (1, 2), \end{cases}
$$

and, for any  $\varepsilon > 0$ ,

$$
\mathbb{P}\left(\left|\sum_{j=0}^{i-1}\langle\mathcal{J}_{n_j}\rangle\right| \geq \varepsilon\sqrt{n_i/\log\log n_i}\right) = \begin{cases}\n\mathcal{O}(n_i^{-1/2}(\log n_i)\log\log n_i), & d/\alpha > 2, \\
\mathcal{O}(n_i^{-1/2}(\log n_i)^3\log\log n_i), & d/\alpha = 2, \\
\mathcal{O}(n_i^{2\Lambda - 1/2}(\log n_i)\log\log n_i), & d/\alpha \in (1, 2) \\
\mathcal{O}(2^{-k/2}k\log k), & d/\alpha > 2, \\
\mathcal{O}(2^{-k/2}k^3\log k), & d/\alpha = 2, \\
\mathcal{O}(2^{(2\Lambda - 1/2)k}k\log k), & d/\alpha \in (1, 2).\n\end{cases}
$$

Let  $\epsilon \in (0,1)$  be arbitrary and let us consider a subsequence  $\{n_{i_l}\}_{l\geq 1}$  which consists of every  $\lfloor 2^{(1-\epsilon)k/2} \rfloor$ -th member of  $\{n_i\}_{i\geq 0}$  in  $\lfloor 2^k, 2^{k+1} \rfloor$ . Clearly, there are at most  $k2^{\epsilon k/2}$ members of this subsequence in  $[2^k, 2^{k+1}]$ . We have

$$
\sum_{l=1}^{\infty} \mathbb{P}\left(\left|\sum_{j=0}^{i_l-1} \langle \mathcal{J}_{n_j} \rangle\right| \geq \varepsilon \sqrt{n_{i_l}/\log \log n_{i_l}}\right)
$$
  

$$
\leq c_3 \begin{cases} \sum_{k=1}^{\infty} 2^{(\epsilon-1)k/2} k^2 \log k, & d/\alpha > 2, \\ \sum_{k=1}^{\infty} 2^{(\epsilon-1)k/2} k^4 \log k, & d/\alpha = 2, \\ \sum_{k=1}^{\infty} 2^{(4\Lambda + \epsilon - 1)k/2} k^2 \log k, & d/\alpha \in (1, 2) \end{cases}
$$

for some  $c_3 > 0$ . When  $d/\alpha \geq 2$  we take an arbitrary  $\epsilon \in (0,1)$ , and when  $d/\alpha \in (1,2)$ we take  $\epsilon \in (0,1)$  such that  $\epsilon < 1-4\Lambda$ . Observe that in the former case it is necessary that  $\Lambda < 1/4$  (that is,  $d/\alpha \in (7/4, 2)$ ). In this case,

<span id="page-31-0"></span>
$$
\sum_{l=1}^{\infty} \mathbb{P}\left(\left|\sum_{j=0}^{i_l-1} \langle \mathcal{J}_{n_j}\rangle\right| \geq \varepsilon \sqrt{n_{i_l}/\log\log n_{i_l}}\right) < \infty.
$$

Borel-Cantelli lemma implies that

(5.15) 
$$
\lim_{l \nearrow \infty} \frac{|\sum_{j=0}^{i_l-1} \langle \mathcal{J}_{n_j} \rangle|}{\sqrt{n_{i_l}/\log \log n_{i_l}}} = 0 \quad \mathbb{P}\text{-a.s.}
$$

We finally prove that eq. [\(5.15\)](#page-31-0) holds for the sequence  $\{n_i\}_{i>0}$ . If  $2^k \leq n_{i} \leq n_i \leq n_{i_{l+1}} \leq$  $2^{k+1}$  then

$$
\sum_{j=0}^{i_l-1} \langle \mathcal{J}_{n_j} \rangle - \sum_{j=i_l}^{i_{l+1}-1} \mathbb{E}[\mathcal{J}_{n_j}] \leq \sum_{j=0}^{i-1} \langle \mathcal{J}_{n_j} \rangle \leq \sum_{j=0}^{i_{l+1}-1} \langle \mathcal{J}_{n_j} \rangle + \sum_{j=i_l}^{i_{l+1}-1} \mathbb{E}[\mathcal{J}_{n_j}].
$$

From Corollary [2.4](#page-5-5) and eq. [\(4.3\)](#page-17-0) it follows that

$$
\mathbb{E}[\mathcal{J}_{n_j}] = \mathcal{O}(h(n_{j+1} - n_j)) = \begin{cases} \mathcal{O}(1), & d/\alpha > 2, \\ \mathcal{O}(\log(n_j^{1/2}/\log n_j)), & d/\alpha = 2, \\ \mathcal{O}((n_j^{1/2}/\log n_j)^{\Delta}), & d/\alpha \in (1, 2). \end{cases}
$$

Hence

$$
\sum_{j=i_l}^{i_{l+1}-1} \mathbb{E}[\mathcal{J}_{n_j}] = \begin{cases} \mathcal{O}(n_i^{(1-\epsilon)/2}), & d/\alpha > 2, \\ \mathcal{O}(n_i^{(1-\epsilon)/2} \log n_i), & d/\alpha = 2, \\ \mathcal{O}(n_i^{(\Lambda+1-\epsilon)/2}), & d/\alpha \in (1,2). \end{cases}
$$

By choosing an arbitrary  $\epsilon \in (0,1)$  in the case when  $d/\alpha \geq 2$ , and  $\epsilon \in (0,1) \cap (\Lambda, 1-4\Lambda)$ in the case when  $d/\alpha \in (1, 2)$ , we obtain

$$
\sum_{j=i_l}^{i_{l+1}-1} \mathbb{E}[\mathcal{J}_{n_j}] = \mathsf{o}(\sqrt{n_i/\log\log n_i}),
$$

which concludes the proof. We observe that  $\Lambda < 1 - 4\Lambda$  if, and only if,  $\Lambda < 1/5$ , that is,  $d/\alpha \in (9/5, 2)$  $d/\alpha \in (9/5, 2).$ 

Acknowledgement. This work has been supported by Deutscher Akademischer Austauschdienst (DAAD) and Ministry of Science and Education of the Republic of Croatia (MSE) via project Random Time-Change and Jump Processes.

Financial support through the Alexander von Humboldt Foundation and Croatian Science Foundation under projects 8958 and 4197 (for N. Sandrić), and the Austrian Science Fund (FWF) under project P31889-N35 and Croatian Science Foundation under project 4197 (for S. Sebek) is gratefully acknowledged.

#### **REFERENCES**

- <span id="page-32-9"></span>[1] R. F. Bass and T. Kumagai. Laws of the iterated logarithm for the range of random walks in two and three dimensions. *Ann. Probab.*, 30(3):1369–1396, 2002.
- <span id="page-32-12"></span>[2] N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular variation*. Cambridge University Press, Cambridge, 1989.
- <span id="page-32-3"></span>[3] E. Csáki and Y. Hu. Strong approximations of three-dimensional Wiener sausages. *Acta Math. Hungar.*, 114(3):205–226, 2007.
- <span id="page-32-6"></span>[4] W. Cygan, N. Sandrić, and S. Šebek. CLT for the capacity of the range of stable random walks. Preprint (2019), arXiv:1904.05695.
- <span id="page-32-7"></span>[5] W. Cygan, N. Sandrić, and S. Šebek. Functional CLT for the range of stable random walks. To appear in Bull. Malays. Math. Sci. Soc.
- <span id="page-32-1"></span>[6] M. D. Donsker and S. R. S. Varadhan. Asymptotics for the Wiener sausage. *Comm. Pure Appl. Math.*, 28(4):525–565, 1975.
- <span id="page-32-11"></span>[7] R. Durrett. *Probability: theory and examples*. Cambridge Series in Statistical and Probabilistic Mathematics, fourth edition, 2010.
- <span id="page-32-2"></span>[8] T. Eisele and R. Lang. Asymptotics for the Wiener sausage with drift. *Probab. Theory Related Fields*, 74(1):125–140, 1987.
- <span id="page-32-5"></span>[9] R. K. Getoor. Some asymptotic formulas involving capacity. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 4:248–252 (1965), 1965.
- <span id="page-32-10"></span>[10] J. M. Hammersley. Generalization of the fundamental theorem on sub-additive functions. *Proc. Cambridge Philos. Soc.*, 58:235–238, 1962.
- <span id="page-32-0"></span>[11] J. Hawkes. Some geometric aspects of potential theory. In *Stochastic analysis and applications (Swansea, 1983)*, volume 1095 of *Lecture Notes in Math.*, pages 130–154. Springer, Berlin, 1984.
- <span id="page-32-4"></span>[12] J. B. Hough and Y. Peres. An LIL for cover times of disks by planar random walk and Wiener sausage. *Trans. Amer. Math. Soc.*, 359(10):4653–4668, 2007.
- <span id="page-32-8"></span>[13] N. C. Jain and W. E. Pruitt. The law of the iterated logarithm for the range of random walk. *Ann. Math. Statist.*, 43:1692–1697, 1972.

.

- <span id="page-33-11"></span><span id="page-33-3"></span>[14] M. Kac and J. M. Luttinger. Bose-Einstein condensation in the presence of impurities. II. *J. Mathematical Phys.*, 15:183–186, 1974.
- <span id="page-33-2"></span>[15] O. Kallenberg. *Foundations of modern probability*. Springer-Verlag, New York, 1997.
- <span id="page-33-5"></span>[16] U. Krengel. *Ergodic theorems*. Walter de Gruyter & Co., Berlin, 1985.
- <span id="page-33-16"></span>[17] J.-F. Le Gall. Fluctuation results for the Wiener sausage. *Ann. Probab.*, 16(3):991–1018, 1988.
- <span id="page-33-20"></span>[18] J.-F. Le Gall and J. Rosen. The range of stable random walks. *Ann. Probab.*, 19(2):650–705, 1991.
- [19] A. Mimica and Z. Vondraˇcek. Unavoidable collections of balls for isotropic L´evy processes. *Stochastic Process. Appl.*, 124(3):1303–1334, 2014.
- <span id="page-33-17"></span>[20] V. V. Petrov. *Sums of independent random variables*. Springer-Verlag, New York-Heidelberg, 1975. Translated from the Russian by A. A. Brown, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 82.
- <span id="page-33-13"></span><span id="page-33-1"></span>[21] S. C. Port. Asymptotic expansions for the expected volume of a stable sausage. *Ann. Probab.*, 18(2):492–523, 1990.
- <span id="page-33-19"></span>[22] S. C. Port and C. J. Stone. Infinitely divisible processes and their potential theory. *Ann. Inst. Fourier (Grenoble)*, 21(2):157–275; ibid. 21 (1971), no. 4, 179–265, 1971.
- [23] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*. Springer-Verlag, Berlin, third edition, 1999.
- <span id="page-33-21"></span><span id="page-33-14"></span>[24] J. Rosen. The asymptotics of stable sausages in the plane. *Ann. Probab.*, 20(1):29–60, 1992.
- [25] K. Sato. *L´evy processes and infinitely divisible distributions*, volume 68. Cambridge University Press, Cambridge, 1999.
- <span id="page-33-18"></span>[26] Q. M. Shao. A Chung type law of the iterated logarithm for subsequences of a Wiener process. *Stochastic Process. Appl.*, 59(1):125–142, 1995.
- <span id="page-33-12"></span>[27] B. Simon. *Functional integration and quantum physics*. AMS Chelsea Publishing, Providence, RI, second edition, 2005.
- <span id="page-33-0"></span>[28] F. Spitzer. Electrostatic capacity, heat flow, and Brownian motion. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 3:110–121, 1964.
- <span id="page-33-4"></span>[29] A.-S. Sznitman. Some bounds and limiting results for the measure of Wiener sausage of small radius associated with elliptic diffusions. *Stochastic Process. Appl.*, 25(1):1–25, 1987.
- <span id="page-33-15"></span>[30] J. Takeuchi. On the sample paths of the symmetric stable processes in spaces. *J. Math. Soc. Japan*, 16:109–127, 1964.
- <span id="page-33-7"></span>[31] M. van den Berg. On the expected volume of intersection of independent Wiener sausages and the asymptotic behaviour of some related integrals. *J. Funct. Anal.*, 222(1):114–128, 2005.
- <span id="page-33-8"></span>[32] M. van den Berg. On the volume of intersection of three independent Wiener sausages. *Ann. Inst. Henri Poincar´e Probab. Stat.*, 46(2):313–337, 2010.
- <span id="page-33-9"></span>[33] M. van den Berg. On the volume of the intersection of two independent Wiener sausages. *Potential Anal.*, 34(1):57–79, 2011.
- <span id="page-33-6"></span>[34] M. van den Berg, E. Bolthausen, and F. den Hollander. On the volume of the intersection of two Wiener sausages. *Ann. of Math. (2)*, 159(2):741–782, 2004.
- <span id="page-33-10"></span>[35] Y. Q. Wang and F. Q. Gao. Laws of the iterated logarithm for high-dimensional Wiener sausage. *Acta Math. Sin. (Engl. Ser.)*, 27(8):1599–1610, 2011.

(Wojciech Cygan) Institut für Mathematische Stochastik, Technische Universität Dresden, Dresden, Germany & Instytut Matematyczny, Uniwersytet Wrocławski, Wrocław, **POLAND** 

*Email address*: wojciech.cygan@uwr.edu.pl

(Nikola Sandrić) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZAGREB, ZAGREB, CROATIA *Email address*: nsandric@math.hr

(Stjepan Šebek) INSTITUTE OF DISCRETE MATHEMATICS, GRAZ UNIVERSITY OF TECHNOLOGY, Graz, Austria & Department of Applied Mathematics, Faculty of Electrical Engineering and Computing, University of Zagreb, Zagreb, Croatia

*Email address*: stjepan.sebek@fer.hr