Second order subexponentiality and infinite divisibility

Toshiro Watanabe

Abstract

We characterize the second order subexponentiality of an infinitely divisible distribution on the real line under an exponential moment assumption. We investigate the asymptotic behaviour of the difference between the tails of an infinitely divisible distribution and its Lévy measure. Moreover, we study the second order asymptotic behaviour of the tail of the t-th convolution power of an infinitely divisible distribution. The density version for a self-decomposable distribution on the real line without an exponential moment assumption is also given. Finally, the regularly varying case for a self-decomposable distribution on the half line is discussed.

2010 Mathematics Subject Classification : 60E07, 60G50

Keywords and phrases : second order subexponentiality, local subexponentiality, infinite divisibility

1 Introduction and results

The subexponentiality of infinitely divisible distributions on the half line was characterized by Embrechts et al. [6] and on the real line by Pakes [16]. The subexponentiality of an infinitely divisible distribution implies the asymptotic equivalence between the tails of the distribution and its Lévy measure. In this paper, we characterize the second order subexponentiality of an infinitely divisible distribution on the real line in terms of its Lévy measure under an exponential moment assumption. The second order subexponentiality yields a higher asymptotic relation than the usual subexponentiality between the tails of an infinitely divisible distribution and its Lévy measure.

In what follows, we denote by \mathbb{R} the real line and by \mathbb{R}_+ the half line $[0, \infty)$. Denote by \mathbb{N} the totality of positive integers. The symbol $\delta_a(dx)$ stands for the delta measure at $a \in \mathbb{R}$. Let η and ρ be probability distributions on \mathbb{R} . We denote by $\eta * \rho$ the convolution of η and ρ and by ρ^{n*} the *n*-th convolution power of ρ with the understanding that $\rho^{0*}(dx) = \delta_0(dx)$. Denote by $m(\rho)$ the mean of ρ . The characteristic function of ρ is denoted by $\hat{\rho}(z)$, namely, for $z \in \mathbb{R}$,

$$\widehat{\rho}(z) := \int_{-\infty}^{\infty} e^{izx} \rho(dx)$$

For a measure ξ on \mathbb{R} , we denote by $\overline{\xi}(x)$ the tail $\xi((x,\infty))$ for x > 0. For positive functions f(x) and g(x) on $[a,\infty)$ for some $a \in \mathbb{R}$, we define the relation $f(x) \sim g(x)$ by $\lim_{x\to\infty} f(x)/g(x) = 1$. We say that $f(x,A) \sim cf(x)$ as $x \to \infty$ and then $A \to \infty$, if

$$\lim_{A \to \infty} \lim_{x \to \infty} f(x, A) / f(x) = c > 0.$$

We say that f(x, A) = o(f(x)) as $x \to \infty$ and then $A \to \infty$, if

$$\limsup_{A \to \infty} \limsup_{x \to \infty} |f(x, A)| / f(x) = 0$$

Definition 1.1. (i) A nonnegative measurable function g(x) on \mathbb{R} belongs to the class **L** if $g(x + a) \sim g(x)$ for every $a \in \mathbb{R}$.

(ii) Let $\Delta := (0, c]$ with c > 0. A distribution ρ on \mathbb{R} belongs to the class \mathcal{L}_{Δ} if $\rho((x, x + c]) \in \mathbf{L}$. A distribution ρ on \mathbb{R} belongs to the class \mathcal{L}_{loc} if $\rho \in \mathcal{L}_{\Delta}$ for each $\Delta := (0, c]$ with c > 0.

(iii) Let $\Delta := (0, c]$ with c > 0. A distribution ρ on \mathbb{R} belongs to the class \mathcal{S}_{Δ} if $\rho \in \mathcal{L}_{\Delta}$ and $\rho^{2*}((x, x + c]) \sim 2\rho((x, x + c])$. A distribution ρ on \mathbb{R} belongs to the class \mathcal{S}_{loc} if $\rho \in \mathcal{S}_{\Delta}$ for each $\Delta := (0, c]$ with c > 0.

If a distribution ρ on \mathbb{R} belongs to the class \mathcal{L}_{loc} , then, for c > 0,

$$\rho((x, x+c]) \sim c\rho((x, x+1])$$

and, for every $\delta > 0$, $e^{\delta x}\rho((x, x + 1]) \to \infty$ as $x \to \infty$. See (2.6) in the proof of Theorem 2.1 of Watanabe and Yamamuro [23] and Lemma 2.17 of Foss et al. [7]. A distribution ρ on \mathbb{R} belongs to the class S if $\bar{\rho}(x) \in \mathbf{L}$ and $\bar{\rho}^{2*}(x) \sim 2\bar{\rho}(x)$. Distributions in the classes S and S_{loc} are called *subexponential* and *locally* subexponential, respectively.

Definition 1.2. A distribution ρ on \mathbb{R} belongs to the class S_{loc}^2 if the following three conditions hold :

(1)
$$\rho \in S_{loc}$$
.
(2) $\int_{-\infty}^{\infty} |x|\rho(dx) < \infty$.
(3) We have
 $\overline{\rho^{2*}}(x) = 2\overline{\rho}(x) + 2m(\rho)\rho((x, x+1]) + o(\rho((x, x+1]))$ (1.1)

as $x \to \infty$.

The subclasses S_{Δ} , S_{loc} , and S_{loc}^2 of the class S were respectively introduced by Asmussen et al. [1], Watanabe and Yamamuro [23], and Lin [13]. Lin [13] treated the one-sided case and used the symbol S_2 for the class S_{loc}^2 . Distributions in the class S_{loc}^2 are called *second order subexponential*. Infinitely divisible distributions on \mathbb{R} in the classes S_{Δ} and S_{loc} are found in Watanabe and Yamamuro [22, 23] and Shimura and Watanabe [18]. Lin [13] gave some sufficient conditions in order that a distribution on \mathbb{R}_+ belongs to the class S_{loc}^2 . See Proposition 2.4 and Corollary 2.1 of [13]. He showed that the lognormal distribution, Weibull distribution with parameter $\beta \in (0, 1)$, and Pareto distribution with parameter $\alpha > 1$ belong to the class S_{loc}^2 . Geluk and Pakes [9] and Geluk [8] treated another second order subexponentiality. Let μ be an infinitely divisible distribution on \mathbb{R} . Then, its characteristic function $\hat{\mu}(z)$ is represented as

$$\widehat{\mu}(z) = \exp\left(\int_{-\infty}^{\infty} (e^{izx} - 1 - \frac{izx}{1 + x^2})\nu(dx) + i\gamma z - \frac{1}{2}az^2\right),$$

where $\gamma \in \mathbb{R}$, $a \ge 0$, and ν is a measure on \mathbb{R} satisfying $\nu(\{0\}) = 0$ and

$$\int_{-\infty}^{\infty} \frac{x^2}{1+x^2} \nu(dx) < \infty.$$

The measure ν is called Lévy measure of μ . See Sato [17]. Throughout the paper, we assume that the tail $\bar{\nu}(c)$ is positive for all c > 0. For c > 0, define a normalized distribution $\nu_{(c)}$ as

$$\nu_{(c)}(dx) := 1_{(c,\infty)}(x) \frac{\nu(dx)}{\bar{\nu}(c)}.$$

Here the symbol $1_{(c,\infty)}(x)$ stands for the indicator function of the set (c,∞) . Denote by μ^{t*} the *t*-th convolution power of μ for t > 0. Note that μ^{t*} is the distribution of X_t for a certain Lévy process $\{X_t\}$.

Theorem 1.1. Let μ be an infinitely divisible distribution on \mathbb{R} with Lévy measure ν . Assume that there exists $\epsilon > 0$ such that $\int_{-\infty}^{\infty} \exp(-\epsilon x)\mu(dx) < \infty$. Then, we have the following :

 $\begin{array}{l} (i) \ \mu \in \mathcal{S}^2_{loc} \ if \ and \ only \ if \ \nu_{(1)} \in \mathcal{S}^2_{loc}. \\ (ii) \ If \ \mu \in \mathcal{S}^2_{loc}, \ then \end{array}$

$$\bar{\nu}(x) = \bar{\mu}(x) - m(\mu)\mu((x, x+1]) + o(\mu((x, x+1]))$$
(1.2)

as $x \to \infty$, equivalently,

$$\bar{\mu}(x) = \bar{\nu}(x) + m(\mu)\nu((x, x+1]) + o(\nu((x, x+1]))$$
(1.3)

as $x \to \infty$.

(iii) Conversely, if (1.2) with finite $m(\mu)$, $\mu \in S_{loc}$, and

$$(\bar{\mu}(x))^2 = o(\mu((x, x+1]))$$

as $x \to \infty$ hold, then $\mu \in \mathcal{S}^2_{loc}$.

Remark 1.1. An exponential moment assumption in the above theorem is necessary for the restriction of the class S_{loc} in the two sided case. See Jian et al. [10] for the detailed account.

Corollary 1.1. Let μ be an infinitely divisible distribution on \mathbb{R} with Lévy measure ν . Assume that there exists $\epsilon > 0$ such that $\int_{-\infty}^{\infty} \exp(-\epsilon x)\mu(dx) < \infty$. Then, we have the following :

Then, we have the following : (i) $\mu \in S^2_{loc}$ if and only if $\mu^{t*} \in S^2_{loc}$ for some t > 0, equivalently, for all t > 0.

(ii) If $\mu \in \mathcal{S}_{loc}^2$, then, for all t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x) + (t^2 - t)m(\mu)\mu((x, x+1]) + o(\mu((x, x+1]))$$
(1.4)

as $x \to \infty$.

Remark 1.2. Let μ be an infinitely divisible distribution on \mathbb{R}_+ with Lévy measure ν . If $\mu \in S_{loc}$, $m(\mu) < \infty$, and μ satisfies (1.4) for $t = t_0, t_0 + 1$ with some $t_0 > 0$, then $\mu \in S_{loc}^2$.

The organization of this paper is as follows. In Sect. 2, we give preliminaries for the proof of the results. In Sect. 3, we prove Theorem 1.1 and its corollary. In Sect. 4, we treat the self-decomposable case. In Sect. 5, three examples of the results are given. In Sect. 6, we give some remarks on the regularly varying case.

2 Preliminaries

Watanabe and Yamamuro [23] used the main results of Watanabe [20] on the convolution equivalence of infinitely divisible distributions on \mathbb{R} to prove the following two lemmas.

Lemma 2.1. (Corollary 2.1 of [23]) Let μ be an infinitely divisible distribution on \mathbb{R} with Lévy measure ν . Assume that there exists $\epsilon > 0$ such that $\int_{-\infty}^{\infty} \exp(-\epsilon x)\mu(dx) < \infty$. Then, the following are equivalent :

(1)
$$\mu \in S_{loc}$$
.
(2) $\nu_{(1)} \in S_{loc}$.
(3) $\nu_{(1)} \in \mathcal{L}_{loc}$ and $\mu((x, x + c]) \sim \nu((x, x + c])$ for all $c > 0$.

Remark 2.1. Since $S_{loc} \subset S$, we see that if condition (1) holds in the above lemma, then

$$\bar{\mu}(x) \sim \bar{\nu}(x) \in \mathbf{L}.$$

Lemma 2.2. (Corollary 3.1 of [23]) Let μ be an infinitely divisible distribution on \mathbb{R} with Lévy measure ν . Assume that there exists $\epsilon > 0$ such that $\int_{-\infty}^{\infty} \exp(-\epsilon x)\mu(dx) < \infty$. If $\mu^{t*} \in S_{loc}$ for some t > 0, then $\mu^{t*} \in S_{loc}$ for all t > 0 and

$$\mu^{t*}((x,x+c]) \sim t\mu((x,x+c])$$

for all t > 0 and for all c > 0.

Lin [13] proved the following three lemmas.

Lemma 2.3. (Theorem 2.1 of [13]) Let ρ be a distribution on \mathbb{R}_+ . Let $\{p_n\}_{n=0}^{\infty}$ be a nonnegative sequence with $p_n > 0$ for some $n \ge 2$ and $\sum_{n=0}^{\infty} p_n = 1$ satisfying $\sum_{n=0}^{\infty} p_n (1+\epsilon_1)^n < \infty$ for some $\epsilon_1 > 0$. Define a distribution η on \mathbb{R}_+ as

$$\eta(dx) := \sum_{n=0}^{\infty} p_n \rho^{n*}(dx).$$

Then we have the following:

(i) If $\rho \in S^2_{loc}$, then we have $\eta \in S^2_{loc}$ and

$$\overline{\eta}(x) = \left(\sum_{n=1}^{\infty} np_n\right)\overline{\rho}(x) + \left(\sum_{n=2}^{\infty} n(n-1)p_n\right)m(\rho)\rho((x,x+1]) + o(\rho((x,x+1])) \quad (2.1)$$

as $x \to \infty$.

(ii) Conversely, if (2.1) with finite $m(\rho)$, $\rho \in S_{loc}$, and

$$(\bar{\rho}(x))^2 = o(\rho((x, x+1]))$$

as $x \to \infty$ hold, then $\rho \in \mathcal{S}^2_{loc}$.

Remark 2.2. We can see from the proof of Theorem 2.1 of [13] that even in the case of $p_n < 0$ for some $n \ge 0$, assertion (i) of the above lemma is still true if $\sum_{n=0}^{\infty} |p_n|(1+\epsilon_1)^n < \infty$ for some $\epsilon_1 > 0$.

Lemma 2.4. (Proposition 2.3 of [13]) Let ρ and η be distributions on \mathbb{R}_+ . If $\rho \in S_{loc}^2$, and there are K > 0 and $c \in \mathbb{R}$ such that

$$\lim_{x \to \infty} \frac{\bar{\eta}(x) - K\bar{\rho}(x)}{\rho((x, x+1])} = c,$$

then $\eta \in \mathcal{S}_{loc}^2$.

Lemma 2.5. (Lemma 3.4 of [13]) Let ρ be a distribution on \mathbb{R}_+ . Assume that $m(\rho) < \infty, \ \rho \in \mathcal{L}_{loc}$ and $(\bar{\rho}(x))^2 = o(\rho(x, x + 1])$ as $x \to \infty$. Then the relation (1.1) implies $\rho \in \mathcal{S}_{loc}$.

Let $\delta := \bar{\nu}(c)$ for c > 0. Define a compound Poisson distribution μ_1 and a distribution σ on \mathbb{R}_+ as

$$\mu_1 := e^{-\delta} \sum_{n=0}^{\infty} \frac{\delta^n}{n!} (\nu_{(c)})^{n*}$$
(2.2)

and

$$\sigma := \frac{e^{-\delta}}{1 - e^{-\delta}} \sum_{n=1}^{\infty} \frac{\delta^n}{n!} (\nu_{(c)})^{n*}.$$
(2.3)

Lemma 2.6. We can choose sufficiently large c > 0 such that $0 < e^{\delta} - 1 < 1$ and we have

$$\nu_{(c)} = -\frac{1}{\delta} \sum_{n=1}^{\infty} \frac{(1-e^{\delta})^n}{n} \sigma^{n*}.$$
(2.4)

Proof. We define a signed measure η as

$$\eta := -\frac{1}{\delta} \sum_{n=1}^{\infty} \frac{(1-e^{\delta})^n}{n} \sigma^{n*}.$$

Let ρ be a signed measure on \mathbb{R}_+ . Denote by $L_{\rho}(t)$ for $t \geq 0$ the Laplace transform of ρ , that is, $L_{\rho}(t) := \int_{0-}^{\infty} e^{-tx} \rho(dx)$. We have

$$L_{\eta}(t) = -\frac{1}{\delta} \sum_{n=1}^{\infty} \frac{(1-e^{\delta})^n}{n} (L_{\sigma}(t))^n$$
$$= \frac{1}{\delta} \log(1-(1-e^{\delta})L_{\sigma}(t)).$$

We see from (2.3) that

$$L_{\sigma}(t) = (e^{\delta} - 1)^{-1} (\exp(\delta L_{\nu_{(c)}}(t)) - 1).$$

Thus we have

$$L_{\eta}(t) = \frac{1}{\delta} \log(\exp(\delta L_{\nu_{(c)}}(t)))$$
$$= L_{\nu_{(c)}}(t)$$

and hence we have $\eta = \nu_{(c)}$, that is, (2.4).

3 Proof of Theorem 1.1 and its corollary

Proof of Theorem 1.1. Let μ be an infinitely divisible distribution on \mathbb{R} with Lévy measure ν . As in Lemma 2.6, we choose sufficiently large c > 0 such that $0 < e^{\delta} - 1 < 1$. We define an infinitely divisible distribution μ_2 by $\mu = \mu_1 * \mu_2$. Assume that there exists $\epsilon \in (0, 1)$ such that $\int_{-\infty}^{\infty} \exp(-\epsilon x)\mu(dx) < \infty$. Then we see from Theorem 25.17 of Sato [17] that $\int_{-\infty}^{\infty} \exp(-\epsilon x)\mu_2(dx) < \infty$ and,

for every b > 0, $\int_{-\infty}^{\infty} \exp(bx)\mu_2(dx) < \infty$. Hence, for every b > 0, $\overline{\mu_2}(x) = o(e^{-bx})$ as $x \to \infty$. We find from Lemma 2.1 that $\mu \in \mathcal{S}_{loc}$ if and only if $\mu_1 \in \mathcal{S}_{loc}$. Since $\int_{-\infty}^{\infty} \exp(-\epsilon x)\mu(dx) < \infty$, we have $\int_{-\infty}^{\infty} |x|\mu(dx) < \infty$ if and only if $\int_{-\infty}^{\infty} |x|\mu_1(dx) < \infty$. Suppose that $\mu \in \mathcal{S}_{loc}$ and $\int_{-\infty}^{\infty} |x|\mu(dx) < \infty$, that is, $\mu_1 \in \mathcal{S}_{loc}$ and $\int_{-\infty}^{\infty} |x|\mu_1(dx) < \infty$. We have

$$\begin{split} \bar{\mu}(x) &- \bar{\mu}_1(x) \\ &= \overline{\mu_1 * \mu_2}(x) - \bar{\mu}_1(x) \\ &= \int_{0-}^{\infty} \mu_1((x-y,x]) \mu_2(dy) - \int_{-\infty}^{0-} \mu_1((x,x-y]) \mu_2(dy) \\ &= I_1 - I_2. \end{split}$$
(3.1)

If $\int_{0-}^{\infty} y\mu_2(dy) = 0$, then $I_1 = 0$ and if $\int_{-\infty}^{0-} |y|\mu_2(dy) = 0$, then $I_2 = 0$. Thus we can assume that $\int_{0-}^{\infty} y\mu_2(dy) > 0$ and $\int_{-\infty}^{0-} |y|\mu_2(dy) > 0$. We find that

$$I_1 := I_{11} + I_{12} + I_{13}$$

4 1

where, for A > 0,

$$I_{11} := \int_{0-}^{A+} \mu_1((x-y,x])\mu_2(dy),$$
$$I_{12} := \int_{A+}^{x/2+} \mu_1((x-y,x])\mu_2(dy),$$

and

$$I_{13} := \int_{x/2+}^{\infty} \mu_1((x-y,x])\mu_2(dy).$$

We have by $\mu_1 \in \mathcal{S}_{loc} \subset \mathcal{L}_{loc}$

$$I_{11} \sim \mu_1((x, x+1]) \int_{0-}^{A+} y\mu_2(dy)$$
$$\sim \mu_1((x, x+1]) \int_{0-}^{\infty} y\mu_2(dy)$$

as $x \to \infty$ and then $A \to \infty$. For any $\epsilon_1 \in (0, 1)$, there is $C_1 > 0$ such that, for $0 \le y \le x/2$ and for sufficiently large x > 0,

$$\frac{\mu_1((x-y,x])}{\mu_1((x,x+1])} \le C_1 e^{\epsilon_1 y}.$$

Thus we see that

$$I_{12} \le \mu_1((x, x+1])C_1 \int_{A+}^{x/2+} e^{\epsilon_1 y} \mu_2(dy)$$

= $o(\mu_1((x, x+1])),$

as $x \to \infty$ and then $A \to \infty$. We have

$$I_{13} \le \bar{\mu}_2(x/2) = o(e^{-x}) = o(\mu_1((x, x+1]))$$

as $x \to \infty$. Thus we have

$$I_1 \sim \mu_1((x, x+1]) \int_{0-}^{\infty} y \mu_2(dy).$$
(3.2)

For any $\epsilon_2 \in (0, \epsilon)$, there is $C_2 > 0$ such that, for y < 0 and for sufficiently large x > 0,

$$\frac{\mu_1(x, x-y])}{\mu_1((x, x+1])} \le C_2 e^{\epsilon_2|y|}.$$

Thus, by dominated convergence theorem, we see that

$$I_2 \sim \mu_1((x, x+1]) \int_{-\infty}^{0-} |y| \mu_2(dy).$$
(3.3)

Hence, we find from (3.1), (3.2), and (3.3) that

$$\bar{\mu}(x) - \bar{\mu}_1(x) = m(\mu_2)\mu_1((x, x+1]) + o(\mu_1((x, x+1]))$$
(3.4)

as $x \to \infty$. By argument analogous to the above equation, we have

$$\overline{\mu^{2*}}(x) - \overline{\mu_1^{2*}}(x) = m(\mu_2^{2*})\mu_1^{2*}((x,x+1]) + o(\mu_1^{2*}((x,x+1]))$$

as $x \to \infty$. Since $\mu_1^{2*}((x, x+1]) \sim 2\mu_1((x, x+1])$ and $m(\mu_2^{2*}) = 2m(\mu_2)$,

$$\overline{\mu^{2*}}(x) - \overline{\mu_1^{2*}}(x) = 4m(\mu_2)\mu_1((x,x+1]) + o(\mu_1((x,x+1]))$$

as $x \to \infty$. Thus we see from (3.4) that

$$\overline{\mu^{2*}}(x) - 2\overline{\mu}(x) = \overline{\mu_1^{2*}}(x) - 2\overline{\mu_1}(x) + 2m(\mu_2)\mu_1((x,x+1]) + o(\mu_1((x,x+1]))$$
(3.5)

as $x \to \infty$. Since $m(\mu) = m(\mu_1) + m(\mu_2)$ and we find from Lemma 2.1 that

$$\mu((x, x+1])) \sim \nu((x, x+1])) \sim \mu_1((x, x+1]), \tag{3.6}$$

we have by (3.5)

$$\overline{\mu^{2*}}(x) = 2\overline{\mu}(x) + 2m(\mu)\mu((x,x+1]) + o(\mu((x,x+1]))$$

as $x \to \infty$ if and only if

$$\overline{\mu_1^{2*}}(x) = 2\overline{\mu}_1(x) + 2m(\mu_1)\mu_1((x,x+1]) + o(\mu_1((x,x+1]))$$

as $x \to \infty$. Hence $\mu \in S_{loc}^2$ if and only if $\mu_1 \in S_{loc}^2$. Since, for x > 0,

$$\frac{\overline{\mu_1}(x)}{1 - e^{-\delta}} = \bar{\sigma}(x)$$

we see from Lemma 2.4 that $\sigma \in S_{loc}^2$ if and only if $\mu_1 \in S_{loc}^2$. We find from (2.3), Lemma 2.6, and Remark 2.2 that if $\sigma \in S_{loc}^2$, then $\nu_{(c)} \in S_{loc}^2$ for sufficiently large c > 0. We see from (2.2) and Lemma 2.3 that if $\nu_{(c)} \in S_{loc}^2$, then $\mu_1 \in S_{loc}^2$. Thus, for sufficiently large c > 0, $\mu \in S_{loc}^2$ if and only if $\nu_{(c)} \in S_{loc}^2$. Since, for sufficiently large x > 0,

$$\overline{\nu_{(c)}}(x) = \frac{\overline{\nu}(1)}{\overline{\nu}(c)}\overline{\nu_{(1)}}(x),$$

we obtain from Lemma 2.4 that $\nu_{(1)} \in S_{loc}^2$ if and only if $\nu_{(c)} \in S_{loc}^2$ for sufficiently large c > 0. Thus we have $\mu \in S_{loc}^2$ if and only if $\nu_{(1)} \in S_{loc}^2$. We have proved assertion (i). Next, we prove assertion (ii). Assume that $\mu \in S_{loc}^2$, equivalently, $\nu_{(c)} \in S_{loc}^2$ for c > 0. Note that $m(\mu_1) = \delta m(\nu_{(c)})$. We see from Lemma 2.3 that

$$\overline{\mu_{1}}(x) = e^{-\delta} \sum_{n=1}^{\infty} \frac{\delta^{n}}{(n-1)!} \overline{\nu_{(c)}}(x) + e^{-\delta} \sum_{n=2}^{\infty} \frac{\delta^{n}}{(n-2)!} m(\nu_{(c)}) \nu_{(c)}((x,x+1]) + o(\nu_{(c)}((x,x+1])) = \overline{\nu}(x) + m(\mu_{1})\nu((x,x+1]) + o(\nu((x,x+1]))$$
(3.7)

as $x \to \infty$. Thus we obtain (1.2) and (1.3) from (3.4) and (3.6). Next we prove assertion (iii). We see from (3.4) that the assumption that (1.2) with finite $m(\mu)$, $\mu \in S_{loc}$, and $(\bar{\mu}(x))^2 = o(\mu((x, x + 1]))$ as $x \to \infty$ is equivalent to that (3.7) with finite $m(\nu_{(c)})$, $\nu_{(c)} \in S_{loc}$, and $(\overline{\nu_{(c)}}(x))^2 = o(\nu_{(c)}((x, x + 1]))$ as $x \to \infty$. This implies from Lemma 2.3 that $\nu_{(c)} \in S_{loc}^2$, equivalently, $\mu \in S_{loc}^2$.

Proof of Corollary 1.1. We see from Theorem 1.1 that $\mu^{t*} \in S^2_{loc}$ for some t > 0, equivalently, for all t > 0 if and only if $\nu_{(1)} \in S^2_{loc}$. Hence assertion (i) is true. Next we prove assertion (ii). Suppose that $\mu \in S^2_{loc}$. Then we find from (i) that $\mu^{t*} \in S^2_{loc}$ for all t > 0. We see from (1.2) that

$$\overline{\mu^{t*}}(x) = t\bar{\nu}(x) + m(\mu^{t*})\mu^{t*}((x,x+1]) + o(\mu^{t*}((x,x+1]))$$

as $x \to \infty$. Note that $m(\mu^{t*}) = tm(\mu)$ and from Lemma 2.2 that

$$\mu^{t*}((x, x+1]) \sim t\mu((x, x+1]).$$

Thus we have by (1.2)

$$\overline{u^{t*}}(x) = t\overline{\nu}(x) + t^2 m(\mu)\mu((x, x+1]) + o(\mu((x, x+1]))$$
$$= t\overline{\mu}(x) + (t^2 - t)m(\mu)\mu((x, x+1]) + o(\mu((x, x+1]))$$

as $x \to \infty$. We have proved (1.4).

Proof of Remark 1.2. Assume that $\mu \in S_{loc}$, $m(\mu) < \infty$, and μ satisfies (1.4) for $t = t_0, t_0 + 1$ with some $t_0 > 0$. Then we have

$$\overline{\mu^{(t_0+1)*}}(x) - (t_0+1)\overline{\mu}(x) - t_0(\overline{\mu^{2*}}(x) - 2\overline{\mu}(x))$$

$$= \int_{0-}^{x+} \overline{\mu^{t_0*}}(x-y)\mu(dy) + \overline{\mu}(x) - (t_0+1)\overline{\mu}(x)$$

$$- t_0 \int_{0-}^{x+} \overline{\mu}(x-y)\mu(dy) - t_0\overline{\mu}(x) + 2t_0\overline{\mu}(x)$$

$$= \int_{0-}^{x+} (\overline{\mu^{t_0*}}(x-y) - t_0\overline{\mu}(x-y))\mu(dy)$$

$$= I_1 + I_2 + I_3,$$
(3.8)

 \square

where, for 0 < 2A < x,

$$I_1 := \int_{0-}^{A+} (\overline{\mu^{t_0*}}(x-y) - t_0 \overline{\mu}(x-y)) \mu(dy),$$

$$I_2 := \int_{A+}^{(x-A)+} (\overline{\mu^{t_0*}}(x-y) - t_0 \overline{\mu}(x-y)) \mu(dy),$$

and

$$I_3 := \int_{(x-A)+}^{x+} (\overline{\mu^{t_0*}}(x-y) - t_0 \overline{\mu}(x-y)) \mu(dy)$$

We divide the proof into three cases: $t_0 > 1$; $t_0 = 1$; and $0 < t_0 < 1$. Let $t_0 > 1$. By the assumption, we see that

$$I_{1} \sim t_{0}(t_{0} - 1)m(\mu) \int_{0-}^{A+} \mu((x - y, x - y + 1])\mu(dy)$$

 $\sim t_{0}(t_{0} - 1)m(\mu)\mu((x, x + 1])$ (3.9)

as $x \to \infty$ and then $A \to \infty$. We find from $\mu \in S_{loc}$ that there is $\epsilon > 0$ such that

$$|I_2| \le (1+\epsilon)t_0(t_0-1)m(\mu) \int_{A+}^{(x-A)+} \mu((x-y,x-y+1])\mu(dy)$$

= $o(\mu((x,x+1]))$ (3.10)

as $x \to \infty$ and then $A \to \infty$. By using integration by parts, we have

$$I_{3} = \int_{0-}^{A+} (\bar{\mu}(x-y) - \bar{\mu}(x))\mu^{t_{0}*}(dy)$$

- $t_{0} \int_{0-}^{A+} (\bar{\mu}(x-y) - \bar{\mu}(x))\mu(dy)$
+ $(\overline{\mu^{t_{0}*}}(A) - t_{0}\bar{\mu}(A))(\bar{\mu}(x-A) - \bar{\mu}(x))$
= $K_{1} - K_{2} + K_{3}.$

As $x \to \infty$ and then $A \to \infty$, we have

$$K_1 \sim m(\mu^{t_0*})\mu((x,x+1]) = t_0 m(\mu)\mu((x,x+1]),$$

and

$$K_2 \sim t_0 m(\mu) \mu((x, x+1]).$$

Note from $m(\mu) < \infty$ that $\overline{\mu^{t_0*}}(A)A \to 0$ and $\overline{\mu}(A)A \to 0$ as $A \to \infty$. Thus we see that

$$\limsup_{A \to \infty} \limsup_{x \to \infty} \frac{|K_3|}{\mu((x, x+1])} \le \limsup_{A \to \infty} (\overline{\mu^{t_0*}}(A) + t_0\overline{\mu}(A))A = 0.$$

Thus we have

$$I_3 = o(\mu((x, x+1])) \tag{3.11}$$

as $x \to \infty$ and then $A \to \infty$. Thus we obtain from (3.8), (3.9), (3.10), and (3.11) and the assumption that

$$(t_0 + 1)t_0 m(\mu)\mu((x, x+1]) - t_0(\overline{\mu^{2*}}(x) - 2\overline{\mu}(x)) = t_0(t_0 - 1)m(\mu)\mu((x, x+1]) + o(\mu((x, x+1]))$$

$$(3.12)$$

as $x \to \infty$. Hence we have

$$\overline{\mu^{2*}}(x) = 2\overline{\mu}(x) + 2m(\mu)\mu((x,x+1]) + o(\mu((x,x+1]))$$
(3.13)

as $x \to \infty$. That is, $\mu \in S_{loc}^2$. Next, let $t_0 = 1$. Then we have (3.13) and hence $\mu \in S_{loc}^2$. Finally, let $0 < t_0 < 1$. In the same way, we see that, as $x \to \infty$ and then $A \to \infty$,

$$-I_1 \sim t_0(1 - t_0)m(\mu)\mu((x, x + 1]),$$
$$I_2 = o(\mu((x, x + 1])),$$

and

$$I_3 = o(\mu((x, x+1]))$$

Thus we have (3.12) and (3.13) and hence $\mu \in S_{loc}^2$.

4 Self-decomposable case

Let f(x) and g(x) be probability density functions on \mathbb{R} . We denote by $f \otimes g(x)$ the convolution of f(x) and g(x) and by $f^{n\otimes}(x)$ the *n*-th convolution power of f(x) for $n \in \mathbb{N}$.

Definition 4.1. (i) A probability density function g(x) on \mathbb{R} belongs to the class \mathcal{L}_d if $g(x) \in \mathbf{L}$.

(ii) A probability density function g(x) on \mathbb{R} belongs to the class \mathcal{S}_d if $g(x) \in \mathcal{L}_d$ and $g^{2\otimes}(x) \sim 2g(x)$.

Definition 4.2. A probability density function g(x) on \mathbb{R} belongs to the class S_d^2 if the following three conditions hold :

(1)
$$g(x) \in S_d$$
.
(2) $\int_{-\infty}^{\infty} |x|g(x)dx < \infty$.
(3) For $\rho(dx) := g(x)dx$,
 $\overline{\rho^{2*}}(x) = 2\overline{\rho}(x) + 2m(\rho)g(x) + o(g(x))$,

as $x \to \infty$.

The classes S_d and S_d^2 were introduced by Chover et al. [5] and Omey and Willekens [15], respectively. Densities in the classes S_d and S_d^2 are called *subexponential* and *second order subexponential*, respectively. See also Foss et al. [7] and Klüppelberg [12] for the class S_d . An infinitely divisible distribution on \mathbb{R}_+ with its density in the class S_d is found in Watanabe [21]. Omey and Willekens [15] studied an infinitely divisible distribution on \mathbb{R}_+ with the density of the normalized Lévy measure in the class S_d^2 . However, they could not characterize the density of an infinitely divisible distribution on \mathbb{R}_+ with its density in the class S_d^2 because they did not know Lemma 2.1 and Lemma 4.1 below. An infinitely divisible distribution μ on \mathbb{R} is called *self-decomposable* if, for every $b \in (0, 1)$, there is a distribution ρ_b on \mathbb{R} such that

$$\widehat{\mu}(z) = \widehat{\mu}(bz)\widehat{\rho}_b(z).$$

An infinitely divisible distribution μ on \mathbb{R} is self-decomposable if and only if $\nu(dx) = k(x)/|x|dx$ with k(x) being nonnegative and increasing on $(-\infty, 0)$

and nonnegative and decreasing on $(0, \infty)$. An infinitely divisible distribution μ on \mathbb{R} is non-degenerate if it is not a delta measure. Every non-degenerate self-decomposable distribution μ on \mathbb{R} is absolutely continuous and unimodal. Many important statistical distributions are known to be self-decomposable. However their Lévy measures and the *t*-th convolution powers are often not explicitly known. See Sato [17]. Let $\mu(dx) = p(x)dx$ be a non-degenerate self-decomposable distribution on \mathbb{R} . We assume that k(x) is positive for all x > 0. We define self-decomposable distributions $\xi_1(dx) = p_1(x)dx$ and $\xi_2(dx) = p_2(x)dx$ as $\mu = \xi_1 * \xi_2$ and

$$\widehat{\xi}_1(z) := \exp\left(\int_0^\infty (e^{izx} - 1) \frac{k(x \lor d)}{x} dx\right).$$

for sufficiently large d > 0. Watanabe and Yamamuro [23] proved the following two lemmas.

Lemma 4.1. (Theorem 1.3 of [23] and its proof) Let $\mu(dx) = p(x)dx$ be a self-decomposable distribution on \mathbb{R} with $\nu(dx) = k(x)/|x|dx$. The following are equivalent :

(1) $\mu \in S_{loc}$. (2) $p(x) \in S_d$. (3) $p_1(x) \in S_d$. (4) $\frac{1}{\bar{\nu}(1)} 1_{(1,\infty)}(x) k(x) / x \in S_d$. (5) $k(x) \in \mathbf{L}$ and $p(x) \sim p_1(x) \sim k(x) / x$.

Remark 4.1. Let $\mu(dx) = p(x)dx$ be a self-decomposable distribution on \mathbb{R} with Lévy measure $\nu(dx) = k(x)/|x|dx$. We see from Lemma 4.1 that $\mu \in S_{loc}^2$ if and only if $p(x) \in S_d^2$ and that $\nu_{(1)} \in S_{loc}^2$ if and only if $\frac{1}{\nu(1)} \mathbb{1}_{(1,\infty)}(x)k(x)/x \in S_d^2$.

Lemma 4.2. (Theorem 1.4 of [23]) Let $\mu(dx) = p(x)dx$ be a self-decomposable distribution on \mathbb{R} . Let $p^t(x)$ be the density of μ^{t*} for t > 0. If $p^t(x) \in S_d$ for some t > 0, then $p^t(x) \in S_d$ for all t > 0 and

$$p^t(x) \sim tp(x)$$

for all t > 0.

Proposition 4.1. Let $\mu(dx) = p(x)dx$ be a self-decomposable distribution on \mathbb{R}_+ . If $(\bar{\mu}(x))^2 = o(\mu((x, x + 1]))$ as $x \to \infty$, $m(\mu) < \infty$, and

$$\overline{\mu^{2*}}(x) = 2\bar{\mu}(x) + 2m(\mu)\mu((x,x+1]) + o(\mu((x,x+1]))$$

as $x \to \infty$, then $p(x) \in \mathcal{S}^2_d$.

Proof. Assume that $(\bar{\mu}(x))^2 = o(\mu((x, x+1]))$ as $x \to \infty$, $m(\mu) < \infty$, and

$$\overline{\mu^{2*}}(x) = 2\overline{\mu}(x) + 2m(\mu)\mu((x, x+1]) + o(\mu((x, x+1]))$$

as $x \to \infty$. Note that

$$\overline{\mu^{2*}}(x) - 2\overline{\mu}(x) + (\overline{\mu}(x))^2 = \int_{0-}^{x+} (\overline{\mu}(x-y) - \overline{\mu}(x))\mu(dy)$$

Thus, by the assumption, we have

$$\int_{0-}^{x+} (\bar{\mu}(x-y) - \bar{\mu}(x))\mu(dy) \sim 2m(\mu)\mu((x,x+1])$$

We shall prove that, for every $m \in \mathbb{N}$,

$$\lim_{x \to \infty} \frac{\mu((x - m, x - m + 1])}{\mu((x, x + 1])} = 1.$$
(4.1)

Since μ is unimodal, we see that, for every $m \in \mathbb{N}$,

$$\liminf_{x \to \infty} \frac{\mu((x - m, x - m + 1])}{\mu((x, x + 1])} \ge 1.$$

Suppose that there are some c > 1, $m_0 \in \mathbb{N}$, and a increasing sequence $\{x_n\}_{n=1}^{\infty}$ with $\lim_{n\to\infty} x_n = \infty$ such that

$$\lim_{n \to \infty} \frac{\mu((x_n - m_0, x_n - m_0 + 1])}{\mu((x_n, x_n + 1])} = c.$$

we have

$$\int_{0-}^{x_n+} (\bar{\mu}(x_n-y) - \bar{\mu}(x_n))\mu(dy) = I_1 + I_2 + I_3,$$

where

$$\begin{split} I_1 &:= \int_{0-}^{m+} (\bar{\mu}(x_n - y) - \bar{\mu}(x_n)) \mu(dy), \\ I_2 &:= \int_{m+}^{(x_n - m)+} (\bar{\mu}(x_n - y) - \bar{\mu}(x_n)) \mu(dy), \end{split}$$

and

$$I_3 := \int_{(x_n - m)+}^{x_n +} (\bar{\mu}(x_n - y) - \bar{\mu}(x_n))\mu(dy).$$

By the unimodality, we have for $0 \leq y \leq m_0$

$$\liminf_{n \to \infty} \frac{\mu((x_n - y, x_n])}{\mu((x_n, x_n + 1])} \ge y$$

and for $y \ge m_0$

$$\liminf_{n \to \infty} \frac{\mu((x_n - y, x_n])}{\mu((x_n, x_n + 1])} \ge c(y - m_0) + m_0$$

Thus we have by Fatou's lemma

$$\liminf_{m \to \infty} \liminf_{n \to \infty} \frac{I_1}{\mu((x_n, x_n + 1])} \\
\geq \int_{0^-}^{m_0^+} \liminf_{n \to \infty} \frac{\mu((x_n - y, x_n])}{\mu((x_n, x_n + 1])} \mu(dy) \\
+ \liminf_{m \to \infty} \int_{m_0^+}^{m^+} \liminf_{n \to \infty} \frac{\mu((x_n - y, x_n])}{\mu((x_n, x_n + 1])} \mu(dy) \\
\geq \int_{0^-}^{m_0^+} y\mu(dy) + \liminf_{m \to \infty} \int_{m_0^+}^{m^+} (c(y - m_0) + m_0)\mu(dy) \\
= \int_{0^-}^{m_0^+} y\mu(dy) + \int_{m_0^+}^{\infty} (c(y - m_0) + m_0)\mu(dy) > m(\mu).$$
(4.2)

Clearly, we have

$$\liminf_{n \to \infty} \frac{I_2}{\mu((x_n, x_n + 1])} \ge 0.$$
(4.3)

By using integration by parts, we see that, for sufficiently large n,

$$I_3 = I_1 + (\bar{\mu}(x_n - m) - \bar{\mu}(x_n))(\bar{\mu}(m) - \bar{\mu}(x_n)) \ge I_1.$$

Thus we obtain from (4.2) that

$$\liminf_{m \to \infty} \liminf_{n \to \infty} \frac{I_3}{\mu((x_n, x_n + 1])} > m(\mu).$$
(4.4)

Hence we have by (4.2), (4.3), and (4.4)

$$\liminf_{n \to \infty} \frac{1}{\mu((x_n, x_n + 1])} \int_{0-}^{x_n+} (\bar{\mu}(x_n - y) - \bar{\mu}(x_n))\mu(dy) > 2m(\mu).$$

This is a contradiction. Thus we have proved (4.1). By the unimodality, it implies that $p(x) \in \mathcal{L}_d$. Thus, by Lemma 2.5, we have proved $\mu \in \mathcal{S}_{loc}^2$ and hence $p(x) \in \mathcal{S}_d^2$.

Theorem 4.1. Let $\mu(dx) = p(x)dx$ be a self-decomposable distribution on \mathbb{R} with Lévy measure $\nu(dx) = k(x)/|x|dx$. Assume that $\int_{-\infty}^{0-} |x|\mu(dx) < \infty$. Then, the following hold :

the following hold : (i) $p(x) \in S_d^2$ if and only if $\nu_{(1)} \in S_{loc}^2$, equivalently, $\frac{1}{\bar{\nu}(1)} \mathbb{1}_{(1,\infty)}(x)k(x)/x \in S_d^2$.

 $\mathcal{S}_{\overline{d}}^{\cdot}$. (ii) If $p(x) \in \mathcal{S}_{d}^{2}$, then

$$\bar{\nu}(x) = \bar{\mu}(x) - m(\mu)p(x) + o(p(x))$$
(4.5)

as $x \to \infty$, equivalently,

$$\bar{\mu}(x) = \bar{\nu}(x) + m(\mu)k(x)/x + o(k(x)/x)$$
(4.6)

as $x \to \infty$.

(iii) Conversely, if (4.5) with finite $m(\mu)$, $p(x) \in S_d$, and $(\bar{\mu}(x))^2 = o(p(x))$ as $x \to \infty$ hold, then $p(x) \in S_d^2$.

Proof. Since the support of the Lévy measure of ξ_2 has an upper bound, we find from Theorem 25.17 of Sato [17] that, for every b > 0, $\int_0^\infty e^{bx}\xi_2(dx) < \infty$ and hence $\overline{\xi_2}(x) = o(e^{-bx})$ as $x \to \infty$. We have

$$\begin{split} \bar{\mu}(x) &- \bar{\xi}_1(x) \\ &= \overline{\xi_1 * \xi_2}(x) - \bar{\xi}_1(x) \\ &= \int_0^\infty \xi_1((x - y, x]) \xi_2(dy) - \int_{-\infty}^0 \xi_1((x, x - y]) \xi_2(dy) \\ &= I_1 - I_2. \end{split}$$

If $\int_0^\infty y\xi_2(dy) = 0$, then $I_1 = 0$ and if $\int_{-\infty}^0 |y|\xi_2(dy) = 0$, then $I_2 = 0$. Thus we can assume that $\int_0^\infty y\xi_2(dy) > 0$ and $\int_{-\infty}^0 |y|\xi_2(dy) > 0$. Suppose that $p(x) \in \mathcal{S}_d$, equivalently by Lemma 4.1, $p_1(x) \in \mathcal{S}_d$. Note that $p_1(x) \in \mathbf{L}$ and ξ_1 is unimodal. Thus there are C > 0 and $\epsilon > 0$ such that, for 0 < y < x/2 and for sufficiently large x > 0,

$$\xi_1((x-y,x]) \le Ce^{\epsilon y} p_1(x).$$

Note that $\int_0^\infty e^{\epsilon y} \xi_2(dy) < \infty$ and

$$\int_{x/2}^{\infty} \xi_1((x-y,x])\xi_2(dy) \le \bar{\xi}_2(x/2) = o(e^{-x}) = o(p_1(x))$$

as $x \to \infty$. Thus, by dominated convergence theorem,

$$I_1 \sim p_1(x) \int_0^\infty y \xi_2(dy)$$

Since ξ_1 is unimodal, we have, for y < 0 and for sufficiently large x > 0,

$$\xi_1((x, x - y]) \le p_1(x)|y|.$$

Since $\int_{-\infty}^{0} |y| \mu(dy) < \infty$, we see from Theorem 25.3 of Sato [17] that $\int_{-\infty}^{\infty} |y| \xi_2(dy) < \infty$. Thus, by dominated convergence theorem,

$$I_2 \sim p_1(x) \int_{-\infty}^0 |y| \xi_2(dy).$$

Note from Lemma 4.1 that $p_1(x) \sim p(x)$. Hence we see that

$$\bar{\mu}(x) = \xi_1(x) + m(\xi_2)p_1(x) + o(p_1(x))$$

= $\bar{\xi}_1(x) + m(\xi_2)p(x) + o(p(x))$ (4.7)

as $x \to \infty$. Note that $p^{2\otimes}(x) \sim 2p(x)$ and $m(\xi_2^{2*}) = 2m(\xi_2)$. In the same way, we have

$$\overline{\mu^{2*}}(x) = \overline{\xi_1^{2*}}(x) + m(\xi_2^{2*})p^{2\otimes}(x) + o(p^{2\otimes}(x))$$
$$= \overline{\xi_1^{2*}}(x) + 4m(\xi_2)p(x) + o(p(x))$$

as $x \to \infty$. Hence we obtain from (4.7) that

$$\overline{\mu^{2*}(x)} - 2\overline{\mu}(x) = \overline{\xi_1^{2*}}(x) - 2\overline{\xi_1}(x) + 2m(\xi_2)p(x) + o(p(x)).$$

as $x \to \infty$. Since $\int_{-\infty}^{\infty} |y|\xi_2(dy) < \infty$, we see from Theorem 25.3 of Sato [17] that $\int_{-\infty}^{\infty} |x| \mu(dx) < \infty$ if and only if $0 < m(\xi_1) < \infty$. Thus we have

$$\overline{\mu^{2*}}(x) = 2\overline{\mu}(x) + 2m(\mu)p(x) + o(p(x))$$

as $x \to \infty$ if and only if

$$\overline{\xi_1^{2*}}(x) = 2\overline{\xi_1}(x) + 2m(\xi_1)p_1(x) + o(p_1(x))$$

as $x \to \infty$. Thus under the assumption of $\int_{-\infty}^{0} |y| \mu(dy) < \infty$, we have $p(x) \in \mathcal{S}_{d}^{2}$ if and only if $p_{1}(x) \in \mathcal{S}_{d}^{2}$, equivalently $\xi_{1} \in \mathcal{S}_{loc}^{2}$. We find from Theorem 1.1

that $\xi_1 \in S_{loc}^2$ if and only if $\nu_{(1)} \in S_{loc}^2$. That is, $p(x) \in S_d^2$ if and only if $\nu_{(1)} \in S_{loc}^2$, equivalently, $\frac{1}{\bar{\nu}(1)} \mathbb{1}_{(1,\infty)}(x) k(x) / x \in S_d^2$. Next we prove assertion (ii). If $p(x) \in S_d^2$, then $\nu_{(1)} \in S_{loc}^2$ and hence by Theorem 1.1 we have

$$\bar{\xi}_{1}(x) = \bar{\nu}(x) + m(\xi_{1})\nu((x, x+1]) + o(\nu((x, x+1]))$$

$$= \bar{\nu}(x) + m(\xi_{1})p(x) + o(p(x))$$

$$= \bar{\nu}(x) + m(\xi_{1})p_{1}(x) + o(p_{1}(x))$$
(4.8)

as $x \to \infty$. Thus it follows from (4.7) that (4.5) and (4.6) hold. Next we prove assertion (iii). The assumption that (4.5) with finite $m(\mu)$, $p(x) \in S_d^2$, and $(\bar{\mu}(x))^2 = o(p(x))$ as $x \to \infty$ implies that (4.8) with finite $m(\xi_1)$, $\xi_1 \in S_{loc}$, and $(\bar{\xi}_1(x))^2 = o(\xi_1((x, x + 1]))$ as $x \to \infty$. Thus we see from (iii) of Theorem 1.1 that $\xi_1 \in S_{loc}^2$, that is, $p_1(x) \in S_d^2$. It follows from the proof of (i) that $p(x) \in S_d^2$.

Corollary 4.1. Let $\mu(dx) = p(x)dx$ be a self-decomposable distribution on \mathbb{R} with Lévy measure ν . Then, the following hold :

(i) $p(x) \in S_d^2$ if and only if $p^t(x) \in S_d^2$ for some t > 0, equivalently, for all t > 0.

(ii) If $p(x) \in S_d^2$, then, for all t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x) + (t^2 - t)m(\mu)p(x) + o(p(x))$$

as $x \to \infty$.

Proof. By argument analogous to the proof of Corollary 1.1, we can easily prove the corollary from Theorem 4.1 and Lemma 4.2. $\hfill \Box$

5 Examples

By using a method of Klüppelberg [11] and Baltrunas [2], Lin [13] proved that the standard lognormal distribution, Weibull distribution with parameter $\beta \in (0, 1)$, and Pareto distribution with parameter $\alpha > 1$ belong to the class S_{loc}^2 . Those distributions are all self-decomposable, so their densities also belong to the class S_d^2 . See Sato [17] and Steutel and van Harn [19] for their selfdecomposability. The following examples are direct consequence of Theorem 1.1 and Corollary 1.1 and hence their proofs are omitted.

Example 5.1. Let μ be the standard lognormal distribution with Lévy measure $\nu(dx) = k(x)/xdx$. Then we have the density

$$p(x) := \frac{1}{\sqrt{2\pi}x} \exp\left(-\frac{(\log x)^2}{2}\right)$$

for x > 0. Embrechts et al. [6] showed that μ is subexponential and that

$$\bar{\nu}(x) \sim \bar{\mu}(x) \sim \frac{x}{\log x} p(x)$$

and

$$\overline{\mu^{t*}}(x) \sim t\bar{\mu}(x)$$

Watanabe and Yamamuro [23] proved a conjecture of Bondesson [4]. That is,

$$k(x) \sim xp(x).$$

We have

$$\bar{\nu}(x) = \bar{\mu}(x) \left(1 - \sqrt{e} \frac{\log x}{x} + o\left(\frac{\log x}{x}\right) \right)$$

as $x \to \infty$ and, for t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x) \left(1 + (t-1)\sqrt{e\frac{\log x}{x}} + o\left(\frac{\log x}{x}\right) \right)$$

as $x \to \infty$.

Example 5.2. Let μ be Weibull distribution with Lévy measure ν and parameter $\beta \in (0, 1)$. Then we have

$$\bar{\mu}(x) := \exp(-x^{\beta})$$

for $x \in \mathbb{R}_+$,

$$\bar{\nu}(x) = \bar{\mu}(x)(1 - \Gamma(\beta^{-1})x^{\beta - 1} + o(x^{\beta - 1}))$$

as $x \to \infty$, and, for t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x)(1 + (t-1)\Gamma(\beta^{-1})x^{\beta-1} + o(x^{\beta-1}))$$

as $x \to \infty$.

Example 5.3. Let μ be Pareto distribution with Lévy measure ν and parameter $\alpha > 1$. Then we have

$$\bar{\mu}(x) := (1+x)^{-\alpha}$$

for $x \in \mathbb{R}_+$,

$$\bar{\nu}(x) = \bar{\mu}(x) \left(1 - \frac{\alpha}{\alpha - 1} x^{-1} + o(x^{-1}) \right)$$

as $x \to \infty$, and, for t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x) \left(1 + (t-1)\frac{\alpha}{\alpha-1}x^{-1} + o(x^{-1}) \right)$$

as $x \to \infty$.

6 Remarks on the regularly varying case

We cannot find from our results the relations of Example 5.3 for Pareto distribution with parameter $0 < \alpha \leq 1$ because it does not belong to the class S_{loc}^2 . However, we can get the analogous relations by using the following lemma of Omey and Willekens [14]. Theorem 4.3 of [14] is a direct consequence from Theorem 2.3 of [14] for a compound Poisson distribution on \mathbb{R}_+ , but there is a mistake in the case of finite mean for an infinitely divisible distribution on \mathbb{R}_+ . **Lemma 6.1.** (Theorem 4.3 of [14]) Let μ be an infinitely divisible distribution on \mathbb{R}_+ with Lévy measure ν . Assume that $\nu(dx)$ has a density q(x) on $(1,\infty)$ such that $q(x) \sim x^{-\alpha-1}l(x)$ for $0 \leq \alpha \leq 1$ with l(x) being slowly varying as $x \to \infty$. Define a constant $C(\alpha)$ for $0 < \alpha < 1$ as

$$C(\alpha) := \frac{(1-\alpha)(2\alpha-1)(\Gamma(1-\alpha))^2}{2\alpha\Gamma(2-2\alpha)}.$$

(i) We have for $0 < \alpha < 1$

$$\lim_{x \to \infty} \frac{\bar{\mu}(x) - \bar{\nu}(x)}{q(x) \int_1^x \bar{\nu}(u) du} = C(\alpha).$$
(6.1)

(ii) For $\alpha = 1$, if $\int_1^{\infty} \bar{\nu}(u) du = \infty$, then we have

$$\lim_{x \to \infty} \frac{\bar{\mu}(x) - \bar{\nu}(x)}{q(x) \int_{1}^{x} \bar{\nu}(u) du} = 1.$$
(6.2)

(iii) For $\alpha = 1$, if $\int_{1}^{\infty} \bar{\nu}(u) du < \infty$, then we have

$$\lim_{x \to \infty} \frac{\bar{\mu}(x) - \bar{\nu}(x)}{q(x)m(\mu)} = 1.$$
(6.3)

(iv) For $\alpha = 0$, then we have

$$\lim_{x \to \infty} \frac{\bar{\mu}(x) - \bar{\nu}(x)}{(\bar{\nu}(x))^2} = -\frac{1}{2}.$$
(6.4)

Proof. Let μ be an infinitely divisible distribution on \mathbb{R}_+ with Lévy measure ν . Assume that $\nu(dx)$ has a density q(x) on $(1,\infty)$ such that $q(x) \sim x^{-\alpha-1}l(x)$ for $0 \leq \alpha \leq 1$ with l(x) being slowly varying as $x \to \infty$. Define a compound Poisson distribution μ_1 on \mathbb{R}_+ as (2.2) for c = 1. Define an infinitely divisible distributions μ_2 on \mathbb{R}_+ as $\mu = \mu_1 * \mu_2$. Then we have by Theorem 2.3 of [14], for $0 \leq \alpha \leq 1$, the lemma is true by substituting μ_1 for μ . Thus we can assume that $\mu_2(dx) \neq \delta_0(dx)$. We see from Theorem 25.17 of Sato [17] that, for every b > 0, $\int_{0^-}^{\infty} \exp(bx)\mu_2(dx) < \infty$ and hence $\overline{\mu_2}(x) = o(e^{-bx})$ as $x \to \infty$. We have

$$\begin{split} \bar{\mu}(x) &- \bar{\mu}_1(x) \\ &= \overline{\mu_1 * \mu_2}(x) - \bar{\mu}_1(x) \\ &= \int_{0-}^{\infty} \mu_1((x-y,x]) \mu_2(dy) \\ &= I_1 + I_2 + I_3, \end{split}$$

where

$$I_{1} := \int_{0-}^{A+} \mu_{1}((x-y,x])\mu_{2}(dy),$$
$$I_{2} := \int_{A+}^{x/2+} \mu_{1}((x-y,x])\mu_{2}(dy),$$

and

$$I_3 := \int_{x/2+}^{\infty} \mu_1((x-y,x])\mu_2(dy).$$

Since $q(x) \sim x^{-\alpha-1}l(x), \nu_{(1)} \in \mathcal{S}_{loc}$ and hence, by Lemma 2.1, $\mu_1 \in \mathcal{S}_{loc}$. Thus,

$$I_1 \sim \mu_1((x, x+1]) \int_{0-}^{A+} y\mu_2(dy)$$
$$\sim \mu_1((x, x+1]) \int_{0-}^{\infty} y\mu_2(dy)$$

as $x \to \infty$ and then $A \to \infty$. Since $\mu_1 \in \mathcal{S}_{loc}$, there are C > 0 and $\epsilon > 0$ such that, for $0 \le y \le x/2$ and for sufficiently large x > 0,

$$\mu_1((x - y, x]) \le Ce^{\epsilon y} \mu_1((x, x + 1]).$$

Thus we have

$$I_2 \le \mu_1((x, x+1]) \int_{A+}^{x/2+} C e^{\epsilon y} \mu_2(dy)$$

= $o(\mu_1((x, x+1]))$

as $x \to \infty$ and then $A \to \infty$.

$$I_3 \le \bar{\mu}_2(x/2) = o(e^{-x}) = o(\mu_1((x, x+1]))$$

as $x \to \infty$. Thus we see that

$$\bar{\mu}(x) - \bar{\mu}_1(x) \sim m(\mu_2)\mu_1((x, x+1]).$$
 (6.5)

Note from Lemma 4.1 that, for $0 < \alpha < 1$ or $\alpha = 1$ with $\int_1^{\infty} \bar{\nu}(u) du = \infty$,

$$\mu_1((x, x+1]) \sim q(x) = o(q(x) \int_1^x \bar{\nu}(u) du)$$

as $x \to \infty$. For $\alpha = 0$, we have by Lemma 4.1

$$\mu_1((x, x+1]) \sim q(x) = o((\bar{\nu}(x))^2)$$

as $x \to \infty$. Thus except the case of $\alpha = 1$ with finite $m(\mu_1)$, the lemma is true. In the case of $\alpha = 1$ with finite $m(\mu_1)$, we see from (6.3) with substituting μ_1 for μ and (6.5) that the lemma is true.

Proposition 6.1. Let $\mu(dx) = p(x)dx$ be a self-decomposable distribution on \mathbb{R}_+ . Assume that $p(x) \sim x^{-\alpha-1}l(x)$ for $0 \leq \alpha \leq 1$ with l(x) being slowly varying as $x \to \infty$. Define slowly varying functions $l^*(x)$ and $l_*(x) = \int_1^x l(u)/udu$ and $l_*(x) = \int_x^\infty l(u)/udu$ for x > 1. Then we have the following : (i) Let $0 < \alpha < 1$ and define $K(\alpha)$ as

$$K(\alpha) := \frac{(2\alpha - 1)(\Gamma(1 - \alpha))^2}{2\alpha\Gamma(2 - 2\alpha)}$$

Then we have

$$\bar{\nu}(x) = \bar{\mu}(x) \left(1 - K(\alpha) x^{-\alpha} l(x) + o(x^{-\alpha} l(x)) \right)$$
(6.6)

as $x \to \infty$, and, for t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x) \left(1 + (t-1)K(\alpha)x^{-\alpha}l(x) + o(x^{-\alpha}l(x)) \right)$$
(6.7)

as $x \to \infty$.

(ii) Let $\alpha = 1$. Assume that $l^*(\infty) = \infty$. Then we have

$$\bar{\nu}(x) = \bar{\mu}(x) \left(1 - \frac{l^*(x)}{x} + o\left(\frac{l^*(x)}{x}\right) \right)$$
(6.8)

as $x \to \infty$, and, for t > 0,

$$\overline{\mu^{t*}}(x) = t\bar{\mu}(x) \left(1 + (t-1)\frac{l^*(x)}{x} + o\left(\frac{l^*(x)}{x}\right) \right)$$
(6.9)

as $x \to \infty$.

(iii) Let $\alpha = 1$. Assume that $l^*(\infty) < \infty$. Then we have

$$\bar{\nu}(x) = \bar{\mu}(x) \left(1 - \frac{m(\mu)}{x} + o\left(\frac{1}{x}\right) \right)$$
(6.10)

as $x \to \infty$, and, for t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x)\left(1 + (t-1)\frac{m(\mu)}{x} + o\left(\frac{1}{x}\right)\right) \tag{6.11}$$

as $x \to \infty$.

(iv) Let $\alpha = 0$. Then we have

$$\bar{\nu}(x) = \bar{\mu}(x) \left(1 + \frac{l_*(x)}{2} + o(l_*(x)) \right)$$
(6.12)

as $x \to \infty$, and, for t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x) \left(1 - (t-1)\frac{l_*(x)}{2} + o(l_*(x)) \right)$$
(6.13)

Proof. Assume that $p(x) \sim x^{-\alpha-1}l(x)$ for $0 \leq \alpha \leq 1$ with l(x) being slowly varying as $x \to \infty$. First we prove (i). Let $0 < \alpha < 1$. Since $p(x) \in S_d$, we have by Lemma 4.1

$$q(x) \sim x^{-\alpha - 1} l(x).$$

By Karamata's theorem (Theorem 1.5.11 of [3]), we have

$$\bar{\nu}(x) \sim \bar{\mu}(x) \sim \frac{x^{-\alpha} l(x)}{\alpha}$$

and

$$\int_1^x \bar{\nu}(u) du \sim \frac{x^{1-\alpha} l(x)}{\alpha(1-\alpha)}.$$

Thus we see from (6.1) of Lemma 6.1 that

$$\lim_{x \to \infty} \frac{\bar{\mu}(x) - \bar{\nu}(x)}{x^{-2\alpha}(l(x))^2} = \frac{K(\alpha)}{\alpha}.$$

Thus we have (6.6). In the same way, we have

$$\lim_{x \to \infty} \frac{\overline{\mu^{t*}(x) - t\bar{\nu}(x)}}{x^{-2\alpha}(l(x))^2} = t^2 \frac{K(\alpha)}{\alpha}.$$

Hence we get (6.7) by (6.6). Next we prove (ii). Assume that $p(x) \sim x^{-2}l(x)$. Then, by Karamata's theorem, we have $\bar{\mu}(x) \sim x^{-1}l(x)$. We have by Lemma 4.1

$$q(x) \sim x^{-2}l(x).$$

We see from Karamata's theorem that $\bar{\nu}(x) \sim x^{-1}l(x)$ and

$$\int_1^x \bar{\nu}(u) du \sim l^*(x)$$

and that $\int_{1}^{\infty} \bar{\nu}(u) du = \infty$ from $l^{*}(\infty) = \infty$. Thus we see from (6.2) of Lemma 6.1 that

$$\lim_{x \to \infty} \frac{\bar{\mu}(x) - \bar{\nu}(x)}{x^{-2} l(x) l^*(x)} = 1.$$

Thus we have (6.8). In the same way, we have

$$\lim_{x \to \infty} \frac{\overline{\mu^{t*}}(x) - t\overline{\nu}(x)}{x^{-2}l(x)l^*(x)} = t^2.$$

Hence we get (6.9) by (6.8). Next we prove (iii). As in (ii), we have $q(x) \sim p(x) \sim x^{-2}l(x)$, $\bar{\nu}(x) \sim \bar{\mu}(x) \sim x^{-1}l(x)$, and

$$\int_1^x \bar{\nu}(u) du \sim l^*(x).$$

We see that $\int_1^{\infty} \bar{\nu}(u) du < \infty$ from $l^*(\infty) < \infty$. Thus we find from (6.3) of Lemma 6.1 that

$$\lim_{x \to \infty} \frac{\bar{\mu}(x) - \bar{\nu}(x)}{x^{-2}l(x)m(\mu)} = 1.$$

Thus we have (6.10). In the same way, we have

$$\lim_{x \to \infty} \frac{\overline{\mu^{t*}}(x) - t\overline{\nu}(x)}{x^{-2}l(x)m(\mu)} = t^2.$$

Hence we get (6.11) by (6.10). Next we prove (iv). Assume that $p(x) \sim x^{-1}l(x)$. Then, we see from Lemma 4.1 that $q(x) \sim x^{-1}l(x)$. Thus we have

$$\bar{\mu}(x) \sim \bar{\nu}(x) \sim l_*(x).$$

We find from (6.4) of Lemma 6.1 that

$$\lim_{x \to \infty} \frac{\bar{\mu}(x) - \bar{\nu}(x)}{(l_*(x))^2} = -\frac{1}{2}.$$

Thus we have (6.12). In the same way, we have

$$\lim_{x \to \infty} \frac{\overline{\mu^{t*}}(x) - t\bar{\nu}(x)}{(l_*(x))^2} = -\frac{t^2}{2}.$$

Hence we get (6.13) by (6.12).

Finally, we give the relations for Pareto distribution with parameter $0 < \alpha \leq 1$ as an example of Proposition 6.1. They are different from the relations of Example 5.3.

Example 6.1. Let μ be Pareto distribution with Lévy measure ν and parameter $0 < \alpha \leq 1$. Then we have

$$\bar{\mu}(x) := (1+x)^{-c}$$

for $x \in \mathbb{R}_+$.

(i) Let $0 < \alpha < 1$. Then we have

$$\bar{\nu}(x) = \bar{\mu}(x) \left(1 - \alpha K(\alpha) x^{-\alpha} + o(x^{-\alpha}) \right)$$

as $x \to \infty$, and, for t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x) \left(1 + (t-1)\alpha K(\alpha)x^{-\alpha} + o(x^{-\alpha}) \right)$$

as $x \to \infty$.

(ii) Let $\alpha = 1$. Then we have

$$\bar{\nu}(x) = \bar{\mu}(x) \left(1 - \frac{\log x}{x} + o\left(\frac{\log x}{x}\right) \right)$$

as $x \to \infty$, and, for t > 0,

$$\overline{\mu^{t*}}(x) = t\overline{\mu}(x)\left(1 + (t-1)\frac{\log x}{x} + o\left(\frac{\log x}{x}\right)\right)$$

as $x \to \infty$.

References

- S. Asmussen, S. Foss and D. Korshunov, 'Asymptotics for sums of random variables with local subexponential behaviour', J. Theoret. Probab. 16 (2003), 489-518.
- [2] A. Baltrunas, 'Second order behaviour of ruin probabilities', Scand. Actuarial J. 2 (1999), 120133.
- [3] N.H. Bingham, C.M. Goldie and J.L. Teugels, *Regular variation*, Cambridge University Press 1984.
- [4] L. Bondesson, 'On the Lvy measure of the lognormal and the logcauchy distributions', Methodology And Computing In Applied Probability 4 (2002), 243256.
- [5] J. Chover, P. Ney and S. Wainger, 'Functions of probability measures', J. Analyse Math. 26 (1973), 255-302.
- [6] P. Embrechts, C.M. Goldie and N. Veraverbeke, 'Subexponentiality and infinite divisibility', Z. Wahrscheinlichkeitheorie Verw. Gebiet. 49 (1979), 335-347.
- [7] S. Foss, D. Korshunov and S. Zachary, An introduction to heavy-tailed and subexponential distributions, Second edition. Springer Series in Operations Research and Financial Engineering. Springer, New York, 2013.
- [8] J.L. Geluk, 'Second order tail behaviour of a subordinated probability distribution', Stoch. Process. Appl. 40 (1992), 325-337.

- [9] J.L. Geluk and A.G. Pakes, 'Second order subexponential distributions', J. Austral. Math. Soc., 51 (1991), 73-87.
- [10] T. Jiang, Y. Wang Z. Cui and Y. Chen 'On the almost decrease of a subexponential density', *Statist. Probab. Letters* **153** (2019), 71-79.
- [11] C. Klüppelberg, 'Subexponential distributions and integrated tails', J. Appl. Probab. 25 (1988), 132-141.
- [12] C. Klüppelberg, 'Subexponential distributions and characterizations of related classes', Probab. Theory Related Fields 82 (1989), 259-269.
- [13] J. Lin, 'Second order subexponential distributions with finite mean and their applications to subordinated distributions', J. Theoret. Probab. 25 (2012), 834853.
- [14] E. Omey and E. Willekens, 'Second order behaviour of the tail of a subordinated probability distribution', Stoch. Process. Appl. 21 (1986), 339-353.
- [15] E. Omey and E. Willekens, 'Second-order behaviour of distributions subordinate to distribution with finite mean', *Commun. Statist. Stochastic Models* 3 (1987), 311342.
- [16] A.G. Pakes, 'Convolution equivalence and infinite divisibility'. J. Appl. Probab. 41 (2004), 407-424.
- [17] K. Sato, Lévy processes and infinitely divisible distributions, Cambridge Studies in Advanced Mathematics, 68 Cambridge Univ. Press. 2013.
- [18] T. Shimura and T. Watanabe, 'Subexponential densities of compound Poisson sums and the supremum of a random walk', submitted.
- [19] F.W. Steutel and K. van Harn, Infinite divisibility of probability distributions on the real line, Pure and applied mathematics : a series of monographs and textbooks, 259. Marcel Dekker Inc. 2004.
- [20] T. Watanabe, 'Convolution equivalence and distributions of random sums', Probab. Theory Related Fields 142 (2008), 367-397.
- [21] T. Watanabe, 'Subexponential densities of infinitely divisible distributions on the half line', arXiv preprint arXiv:1912.08366, 2019.
- [22] T. Watanabe and K. Yamamuro, 'Local subexponentiality of infinitely divisible distributions', J. Math-for-Ind. 1 (2009), 81-90.
- [23] T. Watanabe and K. Yamamuro, 'Local subexponentiality and self-decomposability', J. Theoret. Probab. 23 (2010), 1039-1067.

Toshiro Watanabe

Center for Mathematical Sciences, The University of Aizu, Aizu-Wakamatsu Fukushima 965-8580, Japan e-mail: t-watanb@u-aizu.ac.jp

22