

Generative Modeling with Denoising Auto-Encoders and Langevin Sampling

Adam Block¹, Youssef Mroueh², and Alexander Rakhlin¹

¹MIT

²IBM Research & MIT-IBM Watson AI Lab

Abstract

We study convergence of a generative modeling method that first estimates the score function of the distribution using Denoising Auto-Encoders (DAE) or Denoising Score Matching (DSM) and then employs Langevin diffusion for sampling. We show that both DAE and DSM provide estimates of the score of the Gaussian smoothed population density, allowing us to apply the machinery of Empirical Processes. We overcome the challenge of relying only on L^2 bounds on the score estimation error and provide finite-sample bounds in the Wasserstein distance between the law of the population distribution and the law of this sampling scheme. We then apply our results to the homotopy method of [SE19] and provide theoretical justification for its empirical success.

1 Introduction

Recent empirical successes of generative modeling range from high-fidelity image generation with Generative Adversarial Networks (GANs) [GPAM⁺20] to protein folding with differentiable simulators [SEJ⁺20, IRSM19]. GANs are implicit likelihood models, in the sense that they do not directly model the likelihood. On the other hand, explicit generative models directly estimate the likelihood or the score function (the gradient of the log likelihood). Recent works, including [SE19, NCB⁺17, GWJ⁺19], show that successful image generation can be achieved by estimating the score function from data (using Denoising Score Matching, Denoising Auto-Encoders, and pre-trained classifiers respectively) and by using Langevin dynamics for sampling. Conditional text generation, as well as protein folding can be accomplished using a similar approach, see e.g [DML⁺19] for text generation and [IRSM19] for learning 3D protein structures from sequences. Motivated by the recent resurgence of those explicit methods and their empirical success in wide range of applications we focus on this family of explicit generative models and address their theoretical properties.

Formally, we consider the problem of data generation from an unknown distribution. The algorithm we consider is in two parts. In the first part, we estimate the score of the data using a Denoising Auto-Encoder (DAE), while in the second part we plug our estimate of the score of the data into a discretized approximation to the Langevin dynamics stochastic differential equation, generating a sample. In this paper we bound the Wasserstein 2-distance between the law of our sampling algorithm and the population law. The algorithm is a variation on the method of [SE19], which produced state-of-the-art results on standard vision datasets.

If we consider a distribution with density p with respect to the Lebesgue measure, then Langevin Dynamics provide a well-known and much studied way to sample from this distribution. Supposing that $\nabla \log p$ is $\frac{M}{2}$ -Lipschitz, we note that under mild conditions, the Langevin diffusion given by

$$dW(t) = \nabla \log p(W(t))dt + \sqrt{2}dB_t \quad W(0) \sim \mu_0 \quad (1)$$

where B_t is a standard Brownian motion in \mathbb{R}^d , converges in law as $t \rightarrow \infty$ to the population distribution p . We consider the setting where p is unknown, but we have access to n i.i.d. samples $X_1, \dots, X_n \sim p$. In this case, $\nabla \log p$, often called the score of the distribution, must be estimated. We build on the observations of [Vin11, AB14] and show that DAEs trained on the sample provide estimators of the score of a smoothed distribution that are close in the sense of L^2 . We then show that this estimate suffices to bring the Langevin process associated to our estimate

close to the Langevin process that is actually associated with the population distribution. Finally we note that our error decomposition lends itself naturally to the homotopy method used in [SE19] and we show that this approach significantly helps the sampling scheme with respect to Wasserstein-2 distance. The new contributions are as follows:

- We show that an estimator of score that is close in the L^2 sense still furnishes us with a Langevin diffusion whose law at a fixed time $t > 0$ remains close in the sense of Wasserstein to the law of the Langevin diffusion driven by the score of the population distribution. In particular, many statistical estimators are only guaranteed to have small L^2 error, rather than small uniform error, and so the ability to provide estimates of Wasserstein distance for a Langevin sample using such estimates has the potential for significant general application. While the theory is more straightforward for an estimator that is uniformly close, a case that is studied in [RRT17], to our knowledge this is the first theoretical justification for the L^2 -close regime.
- We exhibit a decomposition of the error between the law of our sampling algorithm and the law of the population distribution that makes the benefits of a homotopy method explicit, thereby providing the first rigorous, theoretical justification that we know of for the success realized by the algorithm in [SE19].
- We shed new light on what the Denoising Auto-Encoder (DAE) learns from a distribution, showing that optimizing DAE loss is equivalent to optimizing Denoising score-matching (DSM) loss, which then implies that the population level DAE provides an unbiased estimator of the score of the population distribution convolved with a Gaussian. Moreover we provide a condition to guarantee that the convolved distribution satisfies a log-Sobolev inequality, the first such result of which we know.
- We use our connection between DAE and DSM losses to provide finite sample, high probability estimates of the error of a DAE. To our knowledge, these are the first such finite sample bounds.

We consider a sampling scheme where we fix in advance a sequence (η_i, σ_i^2) with both η_i and σ_i^2 non-increasing, $1 \leq i \leq N$. We then consider a sequence of DAEs $\tilde{f}_1, \dots, \tilde{f}_N$ trained on the data with the variance parameter of \tilde{f}_i equal to σ_i^2 . Let $\hat{f}_i(x) = \frac{1}{\sigma_i^2}(\tilde{f}_i(x) - x)$. Then we apply a homotopy method of discrete Langevin sampling with warm restarts where on the i^{th} leg of the homotopy, we use \hat{f}_i as an estimate of score and a step length of η_i . This is the identical algorithm proposed and empirically justified by [SE19], up to the fact that we consider the DAE criterion, while they consider the DSM criterion.

2 Notation and Preliminaries

There are two steps to our sampling scheme: the estimation of the score and the Langevin sampling. For the first, we have

Definition 1. *If p is a density on \mathbb{R}^d , we call $\nabla \log p$ the score of the distribution. We let g_{σ^2} denote the density of a centred Gaussian with variance σ^2 and let $p_{\sigma^2} = p * g_{\sigma^2}$ denote the convolved distribution. If $r : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a function, we denote the DAE error by*

$$L_{DAE}(r) = \mathbb{E}_{\substack{X \sim p \\ \epsilon \sim g_{\sigma^2}}} [||r(X + \epsilon) - X||^2] \quad (2)$$

We add a hat to indicate that we are considering the empirical distribution:

$$\hat{L}(r) = \frac{1}{n} \sum_{i=1}^n ||r(X_i + \epsilon_i) - X_i||^2 \quad (3)$$

where ϵ_i are i.i.d. centred Gaussians of variance σ^2 . We define the Denoising Score-Matching (DSM) loss as

$$L_{DSM}(s) = \mathbb{E}_{X \sim p_{\sigma^2}} [||s(X) - \nabla \log p_{\sigma^2}(X)||^2] \quad (4)$$

For the entirety of the paper, we assume that $\nabla \log p$ is $\frac{M}{2}$ -Lipschitz. Regarding the Langevin process, we have

Definition 2. We define the Langevin diffusion started at some distribution μ_0 as the (guaranteed unique by say [KS91]) solution $W_{\sigma^2}(t)$ to

$$dW_{\sigma^2}(t) = \nabla \log p_{\sigma^2}(W_{\sigma^2}(t))dt + \sqrt{2}dB_t \quad W_{\sigma^2}(0) \sim \mu_0 \quad (5)$$

where we drop the σ^2 when context allows. If \hat{f} is an $\frac{M}{2}$ -Lipschitz estimate of $\nabla \log p_{\sigma^2}$, then we let $\widehat{W}(t)$ be the unique solution to

$$d\widehat{W}(t) = \hat{f}(\widehat{W}(t))dt + \sqrt{2}dB_t \quad \widehat{W}(0) \sim \mu_0 \quad (6)$$

For a fixed small step length $\eta > 0$, we define

$$W_{k+1} = \eta \hat{f}(W_k) + \sqrt{2\eta} \xi_k \quad (7)$$

where ξ_k are i.i.d. standard Gaussians in \mathbb{R}^d .

We make use of the following definition

Definition 3. For constants $m, b > 0$, we say that a vector field $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is (m, b) -dissipative if for all $x \in \mathbb{R}^d$, we have

$$\langle f(x), x \rangle \geq m\|x\|^2 - b \quad (8)$$

We also introduce some of the notation related to the theory of empirical processes.

Definition 4. Let \mathcal{G} be a class of real valued functions on \mathbb{R}^d and let $S = (X_1, \dots, X_n)$ be n samples from \mathbb{R}^d . We define the Rademacher average with respect to the sample as

$$\widehat{\mathfrak{R}}_n(\mathcal{G}, S) = \mathbb{E}_\varepsilon \left[\sup_{g \in \mathcal{G}} \frac{1}{n} \sum_{i=1}^n \varepsilon_i g(X_i) \right] \quad (9)$$

where ε_i are i.i.d random variables independent of the X_i , taking values $\{\pm 1\}$ with probability $\frac{1}{2}$ each and the expectation is conditional on the X_i . We define the Rademacher complexity of the function class \mathcal{G} as

$$\mathfrak{R}_n(\mathcal{G}) = \sup_{S \subset (\mathbb{R}^d)^n} \widehat{\mathfrak{R}}_n(\mathcal{G}, S). \quad (10)$$

For a class of \mathbb{R}^k -valued functions \mathcal{G} , we denote by $\mathfrak{R}_n(\mathcal{G}) = \sum_{i=1}^k \mathfrak{R}_n(\mathcal{G}_i)$ where \mathcal{G}_i is the restriction of \mathcal{G} to the i -th coordinate.

We make use of the following assumptions on the population distribution:

Assumption 1. The density p is positive everywhere on \mathbb{R}^d .

Assumption 2. The vector field $\nabla \log p$ is $\frac{M}{2}$ Lipschitz for some $M > 0$ and there exist $\tilde{\sigma}_{max}^2, M > 0$ such that for all $0 \leq \sigma^2 \leq \tilde{\sigma}_{max}^2$, the vector field $\nabla \log \frac{p_{\sigma^2}}{p}$ is $\sigma^2 \frac{M}{2}$ -Lipschitz and similarly for \hat{f} .

Assumption 3. The vector fields $-\nabla \log p$ and \hat{f} are (m, b) -dissipative for some positive constants m, b .

The first two assumptions are standard and are used to ensure that there exists a unique strong solution to the Langevin diffusion. The last assumption has seen increased use in recent work to bound the log-Sobolev constant of p when $-\log p$ is not strongly convex, as in, for example, [RRT17]. Note that the third condition coupled with the fundamental theorem of calculus implies that p is $\frac{2}{m}$ -sub-Gaussian, as per Lemma 45.

Regarding the initialization of the Langevin algorithm, i.e. the distribution of $W(0)$ in Equation (1), we make the assumption

Assumption 4. The initialization of $W_0 = W(0)$ satisfies the condition that for some $\alpha \geq 2M^2$ with M the Lipschitz constant above that there is some k_α such that $\mathbb{E} \left[e^{\alpha \|W(0)\|^2} \right] = k_\alpha < \infty$ as well as $KL(\mu_0, p_{\sigma^2}) < \infty$.

Note that in practice $W(0)$ is often initialized by a Gaussian with sufficiently small variance, certainly satisfies the above assumption. This technical presupposition is required in order to control Wasserstein distance by relative entropy as will be seen below. Below, we always take $\alpha < m$ when doing computations; the $\alpha = 2M^2$ case is only to allow us to apply Girsanov's theorem and thus the fact that $k_\alpha < \infty$ is sufficient.

3 Estimating the Score

Our first result connects the DAE to Denoising Score-Matching (DSM), showing that the objectives are equivalent up to an affine transformation, a variant on a result in [Vin11]; while [Vin11] proves the following for a particular DAE parameterization, we show that the below is true in general:

Proposition 5. *Let p be a differentiable density. Then the DAE loss*

$$L_{DAE}(r) = \mathbb{E}_{x \sim p} \mathbb{E}_{\epsilon \sim g_{\sigma^2}} [\|r(x + \epsilon) - x\|^2] \quad (11)$$

and the DSM loss

$$L_{DSM}(s) = \mathbb{E}_{p_{\sigma^2}} [\|s(x) - \nabla \log p_{\sigma^2}(x)\|^2] \quad (12)$$

with

$$s(x) = \frac{r(x) - x}{\sigma^2} \quad (13)$$

are equivalent up to a term that does not depend on r or s .

The proof of Proposition 5 is a simple application of the divergence theorem and Stein's lemma (Lemma 46) and can be found in Appendix D.

An easy corollary of Proposition 5 is similar to a result in [AB14], which relates the population DAE to the score of the population distribution. Instead, we consider the score of the population distribution smoothed by the addition of Gaussian noise; obviously, without knowledge of p , the DSM loss cannot be explicitly evaluated, so this equivalence allows for a loss that can be evaluated in practice. Below, we establish Corollary 6 using Proposition 5; an alternate, direct proof, is included in Appendix D for those interested.

Corollary 6. *Let p be a population density with respect to the Lebesgue measure. Let $r_{\sigma^2}(x)$ denote the optimal DAE with Gaussian noise of variance σ^2 . Then*

$$r_{\sigma^2}(x) = x + \sigma^2 \nabla \log p_{\sigma^2}(x) \quad (14)$$

Proof. Clearly $s(x) = \nabla \log p_{\sigma^2}(x)$ minimizes the L_{DSM} loss. Note that, by Proposition 5, we have that $r(x) = x + \sigma^2 s(x)$ then minimizes L_{DAE} , the DAE loss. The result follows. ■

Later, in our analysis of the Langevin dynamics, we will need to assume dissipative conditions not just on the score of the population distribution p , but also on the score of the smoothed distribution p_{σ^2} ; in particular, we will need to establish a log-Sobolev inequality for the smoothed distribution. For reasons to become clear below, we define

$$\sigma_{max}^2 = \frac{m}{2M} \wedge \tilde{\sigma}_{max}^2 \quad (15)$$

where $\tilde{\sigma}_{max}^2$ is as appears in Assumption 2. Then we have

Proposition 7. *Suppose that $-\nabla \log p$ is $\frac{M}{2}$ -Lipschitz and (m, b) -dissipative. Then for $\sigma^2 \leq \sigma_{max}^2$, there is a constant B such that $-\nabla \log p_{\sigma^2}$ is $(m_{\sigma^2}, b_{\sigma^2})$ -dissipative where*

$$m_{\sigma^2} = \frac{m - \sigma^2 M}{2} \quad b_{\sigma^2} = b + \frac{B^2}{2(m - \sigma^2 M)} \quad (16)$$

Remark 8. *Note that as $\sigma_{max}^2 \leq \frac{m}{2M}$, we can bound the dissipativity constants to say that $-\nabla \log p_{\sigma^2}$ is $\left(\frac{m}{4}, b + \frac{mB^2}{4M^2}\right)$ -dissipative uniformly in $\sigma^2 \leq \sigma_{max}^2$.*

The proposition follows from applying Cauchy-Schwarz and the definition of dissipativity, along with Assumption 2; details are provided in Appendix D. Both Assumption 2 and Proposition 7 are necessary for the analysis in Section 4, the first to show the existence of the Langevin diffusion and the second to show exponential convergence to the stationary distribution.

The above analysis deals with the population risk, but in reality we are only given n i.i.d samples from p . We have the following result:

Proposition 9. Let \mathcal{F} be a class of \mathbb{R}^d -valued functions, all of which are $\frac{M}{2}$ -Lipschitz, bounded coordinate wise by $R > 0$, containing arbitrarily good uniform approximations of $\nabla \log p_{\sigma^2}$ on the ball of radius R . Let $\sigma^2 < \sigma_{max}^2$ and suppose we have n i.i.d samples from p_{σ^2} , X_1, \dots, X_n . Let

$$\hat{s} \in \operatorname{argmin}_{s \in \mathcal{F}} \frac{1}{n} \sum_{i=1}^n \|s(X_i) - \nabla \log p_{\sigma^2}(X_i)\|^2 \quad (17)$$

Then with probability at least $1 - 4\delta - Cne^{-\frac{R^2}{m_{\sigma^2}}}$ on the randomness due to the sample,

$$\mathbb{E}_{X \sim p_{\sigma^2}} [\|\hat{s}(X) - \nabla \log p_{\sigma^2}(X)\|^2] \leq C(MR + B)^2(\log^3 n \cdot \mathfrak{R}_n^2(\mathcal{F}) + \beta_n d) \quad (18)$$

where C is a universal constant, m_{σ^2} can be found in Proposition 7, B is from Lemma 38, and

$$\beta_n = \frac{\log \frac{1}{\delta} + \log \log n}{n} \quad (19)$$

Remark 10. Note while B can be taken to be a bound on the norm of $\nabla \log p$ at the origin, in reality, by replacing R with $2R$ we can take B to be the infimum of $\|\nabla \log p$ on the ball of radius R about the origin. By dissipativity, if R is sufficiently large, then this constant is small.

Remark 11. In order to have a high probability estimate, we need $Cne^{-\frac{R^2}{m_{\sigma^2}}}$ to be small; as such we see that $R^2 = \Omega(\log n)$. Thus, up to factors polynomial in $\log n$, we see that the L^2 error of the estimate is $\tilde{O}(\mathfrak{R}_n^2(\mathcal{F}))$.

From the equivalence between DSM and DAE loss, we have as an immediate corollary

Proposition 12. Suppose we are in the setting of Proposition 9. Let

$$\hat{r} \in \operatorname{argmin}_{r \in \mathcal{F}} \frac{1}{n} \sum_{i=1}^n \|r(X_i + \sigma \xi_i) - X_i\|^2 \quad (20)$$

where ξ_i are iid standard Gaussians. Then with probability at least $1 - 4\delta - Cne^{-\frac{R^2}{m_{\sigma^2}}}$

$$\mathbb{E}_{X \sim p_{\sigma^2}} \left[\left\| \frac{(\hat{r}(X) - X)}{\sigma^2} - \nabla \log p_{\sigma^2}(X) \right\|^2 \right] \leq \frac{1}{\sigma^4} C(MR + B)^2(\log^3 n \cdot \mathfrak{R}_n^2(\mathcal{F}) + \beta_n d) \quad (21)$$

where the notation is as in Proposition 9.

Proof. Let $r^*(x) = x + \sigma^2 \nabla \log p_{\sigma^2}(x)$ be the population optimal DAE. Using the identical analysis as in Proposition 9, we get that

$$\mathbb{E}_{X \sim p_{\sigma^2}} \left[\left\| \hat{r}(X) - X - \sigma^2 \nabla \log p_{\sigma^2}(X) \right\|^2 \right] \leq C(MR + B)^2(\log^3 n \cdot \mathfrak{R}_n^2(\mathcal{F}) + \beta_n d) \quad (22)$$

Dividing by $\sigma^4 > 0$ on both sides of the above inequality concludes the proof. \blacksquare

The proof of Proposition 9 is an application of a result on lower isometry in [RST17]. We provide a sketch below, with full details appearing in Appendix A.

Proof. (Sketch of Proposition 9) In order to apply the desired result from empirical processes theory, we require our function class to be bounded; while this does not hold, we can provide bounds in high probability. By the fundamental theorem of calculus and the dissipativity assumption, we have that p has Gaussian tails, as proved in Lemma 45. Thus with probability at least $1 - Cne^{-\frac{2R^2}{m_{\sigma^2}}}$ all the samples fall in a ball of radius R , on which \mathcal{F} is certainly bounded. Then the minimizer of the population DSM loss is given by $\nabla \log p_{\sigma^2}$ and we are in the well-specified case. Breaking the error up coordinate-wise, we bound the squared error by the product of the dimension and the largest coordinate-wise error. Applying [RST17, Lemmas 8, 9] to this coordinate concludes the proof. \blacksquare

The generality of Proposition 9 with regard to the function class is potentially helpful in fine-grained analysis of how DAEs are used in practice: usually \mathcal{F} is a class of neural networks; combining known results on the complexity of such classes with Corollary 24 gives high-probability bounds on the DAE error. If we consider the special case where \mathcal{F} is the class of Lipschitz functions bounded by R in each coordinate, then we can apply known results on the complexity of this class [Tik93, Theorem XIII] to get a rate of $\tilde{O}\left(n^{-\frac{2}{d}}\right)$, ignoring factors polynomial in $\log n$. On the other hand, norm-based bounds for Rademacher averages of neural networks in [BFT17, GRS17, NTS15] can imply the faster n^{-1} rate, as long as the empirical error in Equation (20) is small.

4 The Langevin Process: Approximation and Convergence

In this section, we analyze one section of the homotopy method described above. As such, we fix an η and a σ^2 and bound $\mathcal{W}_2(\mu_k, p)$, where we recall that μ_k is the law of W_k , the k^{th} iterate in the discrete Langevin sampling scheme described in the introduction. We have the following theorem:

Theorem 13. *Let $d \geq 3$ and suppose that $-\nabla \log p$ is $\frac{M}{2}$ -Lipschitz and (m, b) -dissipative. Suppose that \hat{f} is an estimate of $\nabla \log p_{\sigma^2}$ that is also $\frac{M}{2}$ -Lipschitz and (m, b) -dissipative and whose squared error is bounded by ε^2 :*

$$\mathbb{E}_{X \sim p_{\sigma^2}} \left[\left\| \hat{f}(X) - \nabla \log p_{\sigma^2}(X) \right\|^2 \right] \leq \varepsilon^2 \quad (23)$$

If we initialize W_0 according to a distribution μ_0 such that $\mathcal{W}_2(\mu_0, p_{\sigma^2}) < \infty$ and μ_0 is concentrated in high probability on a ball of radius R and satisfying Assumption 4 for some α, k_α , and we let μ_k denote the law of a discrete Langevin sampling scheme with constant step size η run for k iterations with score estimate \hat{f} , then, for sufficiently small $\varepsilon > 0$ with $\tau = k\eta$, we have

$$\mathcal{W}_2(\mu_k, p) \leq \sigma\sqrt{d} + A_{M,d,B}(\eta, \tau) + \sqrt{c_{LS}(\sigma^2) \cdot KL(\mu_0 \| p_{\sigma^2})} \cdot e^{-\frac{\tau}{c_{LS}(\sigma^2)}} + C_{\sigma^2, M, R, B, d}(\tau, \varepsilon) \quad (24)$$

where $c_{LS}(\sigma^2)$ can be found in Proposition 19,

$$A_{M,d,B}(\eta, \tau) = C\sqrt{d\eta\tau}e^{\frac{M^2}{2}\tau} \quad (25)$$

and

$$C_{\sigma^2, M, R, B, d, \alpha}(\tau, \varepsilon) = C\sqrt{(b+d)\tau} \left(\varepsilon\tau + C\|p_{\sigma^2}\|_\infty^{\frac{1}{2}-\frac{1}{d}} e^{\frac{M\sqrt{d}}{4}\tau} \sqrt{\tau\varepsilon^{\frac{1}{d}}} \right)^{\frac{1}{4}} \quad (26)$$

where B is a constant from Lemma 38 and C depends on $M, B, m, b, \alpha, k_\alpha$ and $\mathbb{E}[\|W(0)\|^4]$ in both expressions above, with explicit dependence found in Appendix D and Appendix C respectively.

Remark 14. *The ε in Equation (23) above is controlled in high probability by Proposition 9 or Proposition 12 as rates depending on the number of samples and the complexity of the function class over which we are optimizing. Thus, combining Proposition 9 instantiated on a given function class and Theorem 13 gives explicit high probability bounds on how well the Langevin sampling algorithm with an estimate trained on n samples approximates the population distribution.*

Remark 15. *Note that the condition that ε is sufficiently small is included only to give a nice functional form to $C_{\sigma^2, M, R, B, d, \alpha}$ and can be relaxed if we a term with a square root dependence in addition to the fourth root, as seen in Appendix C.*

Remark 16. *While the exponential dependence on τ in the second and last terms of Equation (24) may seem bad, note that the exponential convergence in the third term tells us that if we want to get δ -close in Wasserstein distance, then τ needs only be poly $(\log \frac{1}{\delta})$ and so the exponential dependence on τ is only a polynomial dependence on error.*

The exponential growth with respect to dimension is a touch more serious. If we make more than a Lipschitz assumption on the score function and assume moreover that $|\Delta \log p| \leq C$ then $M\sqrt{d}$ can be replaced by C in the factor under the square root above. While we leave to future work the job of determining if the $\varepsilon^{\frac{1}{2d}}$ factor is tight, we suspect that, without further assumptions, an exponential dependence on dimension cannot be avoided. In similar work that makes no assumption of convexity, such as [RRT17], polynomial dependence on dimension is not achieved

and there is reason to believe that it cannot be true in general. Regarding sample complexity, minimax results on Wasserstein estimation (see e.g. [NWR19, GGPW19]) suggest that exponential dependence in dimension cannot be improved without further assumptions.

The remainder of this section is devoted to a sketch of the proof of Theorem 13; the rigorous proof is relegated to the appendices. With respect to the approximation of the continuous Langevin diffusion by a discrete process, we apply the standard technique to produce an explicit coupling. Note that our result, in contradistinction to earlier work, requires only that we have access to an estimate of the score that is $L^2(p)$ -close, rather than uniformly close, significantly increasing the difficulty of proving the bound.

In order to prove Theorem 13, we consider several intermediate measures. Let ν_t be the law of $W(t)$ and $\widehat{\nu}_t$ the law of $\widehat{W}(t)$ at a fixed time t . Then by the triangle inequality we can decompose

$$\mathcal{W}_2(\mu_k, p) \leq \mathcal{W}_2(\mu_k, \widehat{\nu}_\tau) + \mathcal{W}_2(\widehat{\nu}_\tau, \nu_\tau) + \mathcal{W}_2(\nu_\tau, \nu_\infty) + \mathcal{W}_2(\nu_\infty, p) \quad (27)$$

Note that $\nu_\infty = p_{\sigma^2}$ and so the last error term is controlled by

Lemma 17. *Let p be a measure on \mathbb{R}^d . Then*

$$\mathcal{W}_2(p, p_{\sigma^2}) \leq \sigma\sqrt{d} \quad (28)$$

Proof. This follows immediately from considering the coupling $(X, X + \xi)$ where $X \sim p$ and $\xi \sim g_{\sigma^2}$ is independent of X . The variance of ξ is $\sigma\sqrt{d}$, concluding the proof. ■

We now bound the other three terms.

The first term in Equation (27) comes from the error introduced by the fact that our sampling algorithm is only an approximation of the continuous Langevin process. We have as a standard result, with proof in Appendix D:

Proposition 18. *With the notation as above and assuming that \widehat{f} is $\frac{M}{2}$ -Lipschitz, we have that there is a constant C depending on M, m, b and linearly on $\mathbb{E}[|W(0)|^2]$ such that*

$$\mathcal{W}_2(\mu_k, \widehat{\nu}_\tau)^2 \leq Cd\eta\tau e^{M^2\tau}$$

Having dispensed with the first and last terms in Equation (27), we are now ready to tackle the middle terms, the error due to the lack of convergence to the stationary distribution and the error due to the difference between the estimated and population Langevin diffusions.

4.1 Log-Sobolev Inequalities and Exponential Convergence in Wasserstein Distance

In order to deal with convergence in Wasserstein distance to the stationary distribution, we show that for sufficiently small $\sigma^2 \geq 0$, p_{σ^2} satisfies a log-Sobolev inequality. This will also be helpful in bounding the second term in Equation (27), as we shall see in the following section. We will use the Lyapunov function criterion as proved in [CGW09]. The key result is the proof that if $-\nabla \log p$ is (m, b) -dissipative, then p satisfies a log-Sobolev inequality with a constant bounded in terms of m and b , a result proved in [RRT17]. We have as a translation of Proposition 3.2 from [RRT17]:

Proposition 19. *Let $-\nabla \log p$ be $\frac{M}{2}$ -Lipschitz and (m, b) -dissipative. Then for $0 \leq \sigma^2 \leq \sigma_{max}^2$, the smoothed distribution p_{σ^2} satisfies a log-Sobolev inequality with constant*

$$c_{LS}(\sigma^2) \leq \frac{8M}{m_{\sigma^2}} + \frac{2}{M} + c_P \left(2 + \frac{6M}{m_{\sigma^2}} (b + d) \right) \quad (29)$$

Where m_{σ^2} is the constant appearing in Proposition 7 and

$$c_P(\sigma^2) \leq \frac{2}{m_{\sigma^2}(d+b)} \left(1 + C(d+b)^2 e^{\frac{8(M+B)(d+b)}{m_{\sigma^2}}} \right) \quad (30)$$

where the constant B appears in Lemma 38.

Remark 20. Note that without further assumptions, the exponential dependence on dimension is unavoidable. This is because were this not the case, we would mix in polynomial time, which would allow for general nonconvex optimization in polynomial time, even though the problem is NP-hard. Thus at the very least, the above bound has the tightest functional dependence on dimension.

We replicate the proof used in [RRT17] in detail, in Appendix B. With the log-Sobolev constant established, the convergence in Wasserstein distance is immediate:

Proposition 21. Let $-\nabla \log p$ be $\frac{M}{2}$ -Lipschitz and (m, b) -dissipative. Then for $0 \leq \sigma^2 \leq \sigma_{\max}^2$, if ν_t is the law of $W(t)$, then

$$\mathcal{W}_2(\nu_t, p_{\sigma^2}) \leq \sqrt{c_{LS}(\sigma^2) \cdot KL(\nu_t, p_{\sigma^2})} \leq \sqrt{c_{LS}(\sigma^2) \cdot KL(\mu_0, p_{\sigma^2})} \cdot e^{-\frac{t}{c_{LS}(\sigma^2)}} \quad (31)$$

The proof of this result is well known given the Otto-Villani theorem [OV00, Theorem 1] and the exponential convergence of Langevin in $KL(\cdot, p_{\sigma^2})$ under a log-sobolev inequality; see [BGL14] for details.

4.2 Running a Langevin Diffusion with a Score Estimator

The second term in Equation (27) is the error due to the difference between running a continuous Langevin diffusion with drift \hat{f} and that with drift $\nabla \log p_{\sigma^2}$. The details are in Appendix C, but a sketch of the argument is below. Let $\hat{\nu}_t$ be the law of $\hat{W}(t)$ and let ν_t be the law of $W(t)$ at a fixed time t . We recall that the classic theorem of Bolley and Villani (found in [BV05]) tells us that for all $t > 0$,

$$\mathcal{W}_2(\hat{\nu}_t, \nu_t) \leq C \left(\sqrt{KL(\nu_t, \hat{\nu}_t)} + \left(\frac{KL(\nu_t, \hat{\nu}_t)}{2} \right)^{\frac{1}{4}} \right) \quad (32)$$

where

$$C = 2 \inf_{\alpha > 0} \left(\frac{1}{\alpha} \left(\frac{3}{2} + \log \mathbb{E} \left[e^{\alpha \|\hat{W}(t)\|^2} \right] \right) \right)^{\frac{1}{2}} \quad (33)$$

and

$$KL(\nu_t, \hat{\nu}_t) = \mathbb{E}_{\nu_t} \left[\log \frac{d\nu_t}{d\hat{\nu}_t} \right] \quad (34)$$

is the relative entropy. The constant C is controlled in Proposition 30 in Appendix C using the identical technique as that applied in [RRT17]. Thus it suffices to bound the relative entropy.

In order to compute the relative entropy between the laws of two diffusions with the same noise but different drifts, we can apply Girsanov's theorem and take expectations, as in [Dal14, DT12, RRT17]. In particular, this tells us that

Proposition 22. Let $W(t), \hat{W}(t)$ be as above and assume that $\hat{f}, \nabla \log p_{\sigma^2}$ is $\frac{M}{2}$ -Lipschitz. Then

$$KL(\nu_t, \hat{\nu}_t) = \mathbb{E} \left[\frac{1}{2} \int_0^t \left\| \nabla \log p_{\sigma^2}(W(s)) - \hat{f}(W(s)) \right\|^2 ds \right] \quad (35)$$

Thus it suffices to control this last quantity. If we had that \hat{f} were uniformly close to $\nabla \log p$ or even that we were close in the $L^2(W(t))$ sense, we would be done; unfortunately, we only have that the estimate and the score are close in the sense of $L^2(p_{\sigma^2})$ and so it is not a priori obvious that the above integral can be controlled. In order to get around this, we use the concept of local time and expected occupation density, as seen in [KS91, GH80], as well as bounds on the transition density of a diffusion.

Proposition 23. Let \hat{f} be an estimator of $\nabla \log p_{\sigma^2}$ such that both the estimator and the score are $\frac{M}{2}$ -Lipschitz and (m, b) -dissipative and the error is bounded:

$$\mathbb{E}_{X \sim p_{\sigma^2}} \left[\left\| \hat{f}(X) - \nabla \log p_{\sigma^2}(X) \right\|^2 \right] \leq \varepsilon^2 \quad (36)$$

Suppose we initialize $W(0) \sim \mu_0$ which is concentrated in high probability on a ball of radius R . Then with high probability under the randomness due to initialization at x ,

$$\mathbb{E}_x \left[\int_0^t \left\| \nabla \log p(W(s)) - \hat{f}(W(s)) \right\|^2 ds \right] \leq \varepsilon t + C \|p_{\sigma^2}\|_{L^\infty}^{\frac{1}{2} - \frac{1}{d}} e^{\frac{M\sqrt{d}}{4}t} \sqrt{t} \varepsilon^{\frac{1}{d}} \quad (37)$$

for a constant C depending on M, B, m, b, R and $\mathbb{E}[\|W(0)\|^4]$, whose explicit dependence is given in Appendix C.

A full proof can be found in Appendix C, but we provide a sketch without details.

Proof. (Sketch) In order to bound the desired expected value, we consider the set $U \subset \mathbb{R}^d$ where $\left\| \widehat{f}(x) - \nabla \log p_{\sigma^2}(x) \right\| > \varepsilon$. We can break the integral into the times when $W(s) \in U$ and the times when $W(s) \notin U$. In the latter case we can apply a uniform bound of εt . In the former case, we can apply Cauchy-Schwarz to bound this term by the square root of the product of the fourth moment of $\left\| \widehat{f}(W(s)) - \nabla \log p_{\sigma^2}(W(s)) \right\|$ and the expected amount of time that $W(s)$ spends in U . The former follows easily from the Lipschitz bounds of Lemma 38 and the moment bound of Lemma 39. The second comes from a bound on expected occupation times, which are time integrals of the transition density of the Langevin diffusion. We bound the Radon-Nikodym derivative of these transition densities with respect to the population distribution with the Girsanov theorem, and then integrate with respect to time. Finally, we apply a generalization of a famous inequality of Hardy and Littlewood, giving the result. ■

Combining Proposition 23 with the theorem of Bolley and Villani and Lemma 42 yields

Corollary 24. *Let \widehat{f} be an estimator of $\nabla \log p_{\sigma^2}$ such that both the estimator and the score are $\frac{M}{2}$ -Lipschitz and (m, b) -dissipative. Suppose further that the error is bounded:*

$$\mathbb{E}_{X \sim p_{\sigma^2}} \left[\left\| \widehat{f}(X) - \nabla \log p_{\sigma^2}(X) \right\|^2 \right] \leq \varepsilon^2. \quad (38)$$

Suppose we initialize $W(0) \sim \mu_0$ which is concentrated in high probability on a ball of radius R and satisfies Assumption 4. Then, with high probability under the randomness due to initialization at x , we have

$$\mathcal{W}_2(\widehat{\nu}_t, \nu_t) \leq C \sqrt{(b+d)t} \left(\left(\varepsilon t + C \|p\|_{\infty}^{\frac{1}{2}-\frac{1}{d}} e^{\frac{M\sqrt{d}}{4}t} \sqrt{t\varepsilon^{\frac{1}{d}}} \right)^{\frac{1}{2}} + \left(\varepsilon t + C \|p\|_{\infty}^{\frac{1}{2}-\frac{1}{d}} e^{\frac{M\sqrt{d}}{4}t} \sqrt{t\varepsilon^{\frac{1}{d}}} \right)^{\frac{1}{4}} \right) \quad (39)$$

where C depends on α, k_{α} in Assumption 4 with the explicit dependence given in Appendix C.

We are finally ready to prove Theorem 13:

Proof. (Proof of Theorem 13) Using the triangle inequality, we need to bound each of the four terms in Equation (27). These terms are bounded in Proposition 18, Corollary 24, Proposition 21, and Lemma 17. If $\varepsilon > 0$ is sufficiently small then

$$\left(\varepsilon t + C \|p\|_{\infty}^{\frac{1}{2}-\frac{1}{d}} e^{\frac{M\sqrt{d}}{4}t} \sqrt{t\varepsilon^{\frac{1}{d}}} \right)^{\frac{1}{2}} \leq \left(\varepsilon t + C \|p\|_{\infty}^{\frac{1}{2}-\frac{1}{d}} e^{\frac{M\sqrt{d}}{4}t} \sqrt{t\varepsilon^{\frac{1}{d}}} \right)^{\frac{1}{4}} \quad (40)$$

and thus up to a factor of 2 we can keep only the larger term. This concludes the proof. ■

5 Homotopy and Annealing

The above section and the discussion surrounding Theorem 13 focuses on one leg of the homotopy; this section uses Theorem 13 to analyze the effect that the homotopy method and annealing the DAE has on the Wasserstein distance between the sample and the population distribution.

We consider the following sampling scheme. For fixed k , we fix a sequence $\{(\eta_i, \sigma_i^2) \mid 1 \leq i \leq N\}$ where η_i, σ_i^2 are both decreasing in i . We train a DAE with variance parameter σ_i^2, r_i , and set $s_i(x) = \frac{1}{\sigma_i^2}(r_i(x) - x)$. Then we initialize $W^{(1)} \sim \mu_0$ and evolve with $\widehat{f} = s_1$ with step size η_1 for k iterations. Then, we use warm restarts and for $2 \leq i \leq N$, we evolve a Langevin sampling scheme $W^{(i)}$ with $\widehat{f} = s_i$, step size η_i , and $W_0^{(i)} = W_k^{(i-1)}$. Our final sample is $W_k^{(N)}$.

The decomposition in Equation (27) provides clues as to why the homotopy method described above speeds up the Langevin sampling. Consider, first, the effect that η has on the decomposition. According to Theorem 13, we have that the error is bounded by

$$\mathcal{W}_2(\mu_k, p_{\sigma^2}) \leq \sigma \sqrt{d} + A_{M,d,B}(\eta, \tau) + \sqrt{c_{LS}(\sigma^2) \cdot KL(\mu_0, p_{\sigma^2})} e^{-\frac{\tau}{c_{LS}(\sigma^2)}} + C_{\sigma^2, M, R, B, d, \alpha}(\tau, \varepsilon) \quad (41)$$

As η increases, since $\tau = k\eta$, we see that τ increases and we note that both $A(\eta, \tau)$ and $C(\tau, \varepsilon)$ increase, but the third term decreases. Thus with all other constants fixed, the optimal η can be determined as $\eta_{opt} > 0$.

With regard to score estimation, the greater value of σ^2 makes it easier to estimate the score. Consider that, in our regime, we are training a DAE and then using the transformation in Corollary 24 to plug the score estimate into Theorem 13. With a fixed number of samples, doing the proof of Proposition 12 in reverse, we note that if the DAE has squared error $\tilde{\varepsilon}^2$, then the score estimate has squared error $\frac{1}{\sigma^4}\tilde{\varepsilon}^2$. Thus, if we fix $\tilde{\varepsilon}^2$ as the achievable error of a DAE trained on n samples, then as σ^2 increases, ε^2 decreases and thus so, too, does $C_{\sigma^2, M, R, B, d, \alpha}(\tau, \varepsilon)$.

The effect of σ^2 on the log-Sobolev constant remains a bit more mysterious from a rigorous point of view. While Proposition 7 provides a bound on $c_{LS}(\sigma^2)$, it is likely not tight, as it amounts to a ‘worst-case’ analysis of the effect that the Gaussian smoothing has on the population distribution, using a crude argument involving Cauchy-Schwarz. If we make further assumptions, these results can be tightened. For example, if we suppose that p has compact support, then [BGMZ15] gives a bound on the log-Sobolev constant of the smoothed distribution that decreases with larger variance, thereby accelerating the convergence of the Langevin diffusion to its stationary distribution; while their bound is dimension-dependent, they suggest that future work may lose this handicap. Thus, in this case, just as there exists an η_{opt} that minimizes the right hand side of Equation (41), there is too such a σ_{opt}^2 . We leave to future work the task of better controlling $c_{LS}(\sigma^2)$ in the general case. Thus the annealing and the homotopy method combine to provide a form of dynamic optimization of the upper bound of Theorem 13, significantly decreasing the error and simultaneously speeding up the naive Langevin sampling that does not involve homotopy or annealing. The above is empirically indicated by the success of the annealed score matching in [SE19]; as we saw in Proposition 5, though, the annealed score matching is equivalent to DAE loss, so the empirical success in one area transfers to the other *mutatis mutandis*.

6 Conclusion and Further Directions

We have provided rigorous justification above for two empirical approaches that have recently generated excitement: the use of score estimators to run Langevin sampling and the homotopy method of [SE19]. While the bounds in Theorem 13 and proposition 9 allow for high probability guarantees with finite samples, there is a question of tightness. First, the dependence on the dimension of the bound in eq. (24) is potentially suboptimal: it is possible that a more detailed analysis of the argument proving Proposition 23 would substantially improve this dependence on the dimension. Second, while we have proved that the smoothed distribution p_{σ^2} satisfies a log-Sobolev inequality, we almost certainly do not have the optimal log-Sobolev constant. In fact, as discussed in the above section, while our bound on this constant gets worse with more noise, it is likely that the log-Sobolev constant actually improves with increased variance, which would explain fully the benefits of the annealing of DSM estimators that is so successful in [SE19]. Third, we could likely improve the algorithm by using lower variance estimators of the score function and a higher-order method for approximating the continuous Langevin diffusion. There has been some recent theoretical work in this direction (see, for example, [LWME19]) and we suspect that, especially in practical application, this would considerably accelerate the algorithm.

While the unconditional generative modeling studied in this work is certainly interesting in its own right, practitioners tend to focus on the benefits of conditional generative modeling, i.e., where there are two variables x, y and we wish to input y and generate samples from the conditional distribution $p(x|y)$. It is highly likely that, given the right conditions on the joint distribution of x and y to ensure a uniformity in y to the dissipativity and Lipschitz nature of the conditional distribution, many of the results above could be extended to this regime.

Acknowledgements

We would like to acknowledge the support from the MIT-IBM Watson AI Lab. In addition, this material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. 1122374. We would also like to thank Thibaut Le Gouic and Sinho Chewi for pointing out an error in an earlier draft.

References

- [AB14] Guillaume Alain and Yoshua Bengio. What regularized auto-encoders learn from the data-generating distribution. *J. Mach. Learn. Res.*, 15(1):3563–3593, January 2014.

- [BBCG08] Dominique Bakry, Franck Barthe, Patrick Cattiaux, and Arnaud Guillin. A simple proof of the poincaré inequality for a large class of probability measures. *Electronic Communications in Probability*, 13(0):60–66, 2008.
- [BEL18] Sébastien Bubeck, Ronen Eldan, and Joseph Lehec. Sampling from a log-concave distribution with projected langevin monte carlo. *Discrete & Computational Geometry*, 59(4):757–783, April 2018.
- [BFT17] Peter L Bartlett, Dylan J Foster, and Matus J Telgarsky. Spectrally-normalized margin bounds for neural networks. In *Advances in Neural Information Processing Systems*, pages 6240–6249, 2017.
- [BGL14] Dominique Bakry, Ivan Gentil, and Michel Ledoux. *Analysis and geometry of Markov diffusion operators*. Springer, Cham, 2014.
- [BGMZ15] Jean-Baptiste Bardet, Nathaël Gozlan, Florent Malrieu, and Pierre-André Zitt. Functional inequalities for gaussian convolutions of compactly supported measures: explicit bounds and dimension dependence, 2015.
- [BS88] Colin Bennett and Robert Sharpley. *Interpolation of Operators*. Academic Press, Boston, 1988.
- [BV05] Francois Bolley and Cedric Villani. Weighted csizar-kullback-pinsker inequalities and applications to transportation inequalities. *Annales de la Faculte des sciences de Toulouse : Mathematiques*, Ser. 6, 14(3):331–352, 2005.
- [CGW09] Patrick Cattiaux, Arnaud Guillin, and Li-Ming Wu. A note on talagrand’s transportation inequality and logarithmic sobolev inequality. *Probability Theory and Related Fields*, 148(1-2):285–304, June 2009.
- [Dal14] Arnak Dalalyan. Theoretical guarantees for approximate sampling from smooth and log-concave densities. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79, 12 2014.
- [DML⁺19] Sumanth Dathathri, Andrea Madotto, Janice Lan, Jane Hung, Eric Frank, Piero Molino, Jason Yosinski, and Rosanne Liu. Plug and play language models: A simple approach to controlled text generation, 2019.
- [Dow08] Andrew N. Downes. Bounds for the transition density of time-homogeneous diffusion processes, 2008.
- [DT12] A.S. Dalalyan and A.B. Tsybakov. Sparse regression learning by aggregation and langevin monte-carlo. *Journal of Computer and System Sciences*, 78(5):1423–1443, September 2012.
- [GGPW19] Ziv Goldfeld, Kristjan Greenewald, Yury Polyanskiy, and Jonathan Weed. Convergence of smoothed empirical measures with applications to entropy estimation, 2019.
- [GH80] Donald Geman and Joseph Horowitz. Occupation densities. *Ann. Probab.*, 8(1):1–67, 02 1980.
- [GPAM⁺20] Ian Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial networks. *Communications of the ACM*, 63(11):139–144, 2020.
- [GRS17] Noah Golowich, Alexander Rakhlin, and Ohad Shamir. Size-independent sample complexity of neural networks. *arXiv preprint arXiv:1712.06541*, 2017.
- [GWJ⁺19] Will Grathwohl, Kuan-Chieh Wang, Jörn-Henrik Jacobsen, David Duvenaud, Mohammad Norouzi, and Kevin Swersky. Your classifier is secretly an energy based model and you should treat it like one. *arXiv preprint arXiv:1912.03263*, 2019.
- [IRSM19] John Ingraham, Adam Riesselman, Chris Sander, and Debora Marks. Learning protein structure with a differentiable simulator. In *International Conference on Learning Representations*, 2019.
- [KS91] Ioannis Karatzas and Steven Shreve. *Brownian motion and stochastic calculus*. Springer-Verlag, New York, 1991.
- [LS77] R.S Lipster and A.N. Shiriyayev. *Statistics of random processes*. Springer-Verlag, New York, 1977.

- [LWME19] Xuechen Li, Denny Wu, Lester Mackey, and Murat A. Erdogdu. Stochastic runge-kutta accelerates langevin monte carlo and beyond, 2019.
- [MP10] Peter Mörters and Yuval Peres. *Brownian motion*. Cambridge University Press, Cambridge New York, 2010.
- [NCB⁺17] Anh Nguyen, Jeff Clune, Yoshua Bengio, Alexey Dosovitskiy, and Jason Yosinski. Plug & play generative networks: Conditional iterative generation of images in latent space. *2017 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, Jul 2017.
- [NTS15] Behnam Neyshabur, Ryota Tomioka, and Nathan Srebro. Norm-based capacity control in neural networks. In *Conference on Learning Theory*, pages 1376–1401, 2015.
- [NWR19] Jonathan Niles-Weed and Philippe Rigollet. Estimation of wasserstein distances in the spiked transport model, 2019.
- [OV00] F. Otto and C. Villani. Generalization of an inequality by talagrand and links with the logarithmic sobolev inequality. *Journal of Functional Analysis*, 173(2):361–400, June 2000.
- [RRT17] Maxim Raginsky, Alexander Rakhlin, and Matus Telgarsky. Non-convex learning via stochastic gradient langevin dynamics: a nonasymptotic analysis. In Satyen Kale and Ohad Shamir, editors, *Proceedings of the 2017 Conference on Learning Theory*, volume 65 of *Proceedings of Machine Learning Research*, pages 1674–1703, Amsterdam, Netherlands, 07–10 Jul 2017. PMLR.
- [RST17] Alexander Rakhlin, Karthik Sridharan, and Alexandre B. Tsybakov. Empirical entropy, minimax regret and minimax risk. *Bernoulli*, 23(2):789–824, May 2017.
- [SE19] Yang Song and Stefano Ermon. Generative modeling by estimating gradients of the data distribution. *ArXiv*, abs/1907.05600, 2019.
- [SEJ⁺20] Andrew W Senior, Richard Evans, John Jumper, James Kirkpatrick, Laurent Sifre, Tim Green, Chongli Qin, Augustin Židek, Alexander WR Nelson, Alex Bridgland, et al. Improved protein structure prediction using potentials from deep learning. *Nature*, pages 1–5, 2020.
- [Ste81] Charles M. Stein. Estimation of the mean of a multivariate normal distribution. *The Annals of Statistics*, 9(6):1135–1151, November 1981.
- [Tik93] V. M. Tikhomirov. *ϵ -Entropy and ϵ -Capacity of Sets In Functional Spaces*, pages 86–170. Springer Netherlands, Dordrecht, 1993.
- [Vin11] Pascal Vincent. A connection between score matching and denoising autoencoders. *Neural Comput.*, 23(7):1661–1674, July 2011.
- [WGD04] L. Wu, A. Guillin, and H. Djellout. Transportation cost-information inequalities and applications to random dynamical systems and diffusions. *The Annals of Probability*, 32(3B):2702–2732, July 2004.

A Empirical Processes and Proving Proposition 9

We briefly sketch a few definitions from the theory of empirical processes. Denote by \mathbb{E} as expectation with respect to p_{σ^2} and by $\widehat{\mathbb{E}}$ as expectation with respect to the empirical measure. We have already defined the Rademacher complexity above. Given an $r > 0$, a function class \mathcal{F} , and a sample of n points $X_1, \dots, X_n \in \mathbb{R}^d$, we let

$$\mathcal{F}[r, S] = \left\{ f \in \mathcal{F} : \frac{1}{n} \sum_{i=1}^n f(X_i) \leq r \right\} \quad (42)$$

We call a function $\phi_n : [0, \infty) \rightarrow \mathbb{R}_{\geq 0}$ an upper function for \mathcal{F} if for all $r > 0$,

$$\sup_{S \subset (\mathbb{R}^d)^n} \widehat{\mathfrak{R}}_n(\mathcal{F}[r, S], S) \leq \phi_n(r) \quad (43)$$

We define the (nonunique) localization radius r^* as an upper bound on the maximal solution to the equation $\phi_n(r) = r$. We recall two results:

Lemma 25. [RST17, Lemma 8.i] For any class \mathcal{F} of real valued functions with image in the interval $[0, 1]$, with $n \geq 2$, we may take as localization radius of $\mathcal{G} = \{(f - f')^2 | f, f' \in \mathcal{F}\}$

$$r^* = C \log^3(n) \mathfrak{R}_n(\mathcal{F})^2 \quad (44)$$

where C is a universal constant.

Lemma 26. [RST17, Lemma 9] For any class \mathcal{F} of real valued functions with image contained in the unit interval, and $\delta > 0$, with probability at least $1 - 4\delta$, if $X_1, \dots, X_n \sim p$ i.i.d., for all $f, f' \in \mathcal{F}$

$$\mathbb{E}[(f - f')^2] \leq 2\widehat{\mathbb{E}}(f - f')^2 + C(r^* + \beta) \quad (45)$$

where r^* is the localization radius of $\mathcal{G} = \{(f - f')^2 | f, f' \in \mathcal{F}\}$ and

$$\beta = \frac{\log \frac{1}{\delta} + \log \log n}{n} \quad (46)$$

and C is a universal constant.

With these results, we can prove Proposition 9:

Proof. (Proof of Proposition 9) We prove for the case $\sigma^2 = 0$, as we rely only on the dissipativity and Lipschitz assumptions; thus Assumption 2 and Proposition 7 allow us to apply the same argument with slightly different constants.

By the fundamental theorem of calculus and the dissipativity assumption, we may invoke Lemma 45 to get that for sufficiently large $R > 0$ that there is a constant C such that the probability that $X \sim p$ has norm bigger than R is bounded above by $Ce^{-\frac{mR^2}{4}}$. By a union bound, with probability at least $1 - Cne^{-\frac{R^2}{m}}$ all of the samples lie in the ball bounded by R . On this event then, the norm of $\nabla \log p$ evaluated on the data is bounded by $MR + B$ by Lemma 38. Thus up to a factor of $MR + B$ we are in the situation of Lemma 25 and Lemma 26 with $\nabla \log p \mathbf{1}_{\|X\| \leq R} \in \mathcal{F}$. Let $s^* = \nabla \log p$ and

$$\widehat{s} \in \operatorname{argmin}_{s \in \mathcal{F}} \widehat{\mathbb{E}}[\|s(X) - s^*(X)\|^2]. \quad (47)$$

Clearly,

$$\widehat{\mathbb{E}}[\|\widehat{s}(X) - s^*(X)\|^2] = 0 \quad (48)$$

since $s^* \in \mathcal{F}$ gives that

$$0 \leq \widehat{\mathbb{E}}[\|\widehat{s}(X) - s^*(X)\|^2] \leq \widehat{\mathbb{E}}[\|s^*(X) - s^*(X)\|^2] = 0 \quad (49)$$

We have

$$\mathbb{E}[\|\widehat{s}(X) - s^*(X)\|^2] = \sum_{i=1}^d \mathbb{E}[(\widehat{s}(X)_i - s^*(X)_i)^2] \quad (50)$$

where the subscript i denotes coordinates. Applying Lemma 26 yields

$$\mathbb{E}[(\widehat{s}(X)_i - s^*(X)_i)^2] \leq 2\widehat{\mathbb{E}}[(\widehat{s}(X)_i - s^*(X)_i)^2] + C(MR + B)(r_i^* + \beta) \quad (51)$$

where r_i^* is the localization radius for the coordinate restriction \mathcal{F}_i . Applying Lemma 25 to bound $\sum r_i^*$ and noting that we are off by a factor of $(MR + B)$ from the result in [RST17], gives that

$$\sum_{i=1}^d r_i^* \leq (MR + B)C \log^3(n) \cdot \sum_{i=1}^d \mathfrak{R}_n(\mathcal{F}_i)^2 \leq (MR + B)C \log^3(n) \cdot \mathfrak{R}_n(\mathcal{F})^2 \quad (52)$$

which concludes the proof. ■

B The log-Sobolev Constant and Convergence to the Stationary Distribution

Background on log-Sobolev inequalities can be found in [BGL14]. Recall that the generator of the process $W(t)$ is given by a second order differential operator \mathcal{L} acting on a test function u by

$$\mathcal{L}u = \Delta u + \langle \nabla \log p_{\sigma^2}, \nabla u \rangle \quad (53)$$

We call the Dirichlet form evaluated on a function f :

$$\mathcal{E}(f) = \int \|\nabla f\|^2 p_{\sigma^2} dx \quad (54)$$

Note that $W(t)$, if $\nabla \log p_{\sigma^2}$ is Lipschitz, has a unique invariant distribution of p_{σ^2} . We say that p_{σ^2} satisfies a Poincaré inequality with constant c_P if for all measures $\mu \ll p_{\sigma^2}$, we have

$$\int \left| \frac{d\mu}{dp_{\sigma^2}} - 1 \right|^2 p_{\sigma^2} \leq c_P \mathcal{E} \left(\sqrt{\frac{d\mu}{dp_{\sigma^2}}} \right) \quad (55)$$

We say that p_{σ^2} satisfies a log-Sobolev inequality with constant c_{LS} if for all $\mu \ll p_{\sigma^2}$, we have

$$KL(\mu, p_{\sigma^2}) \leq c_{LS} \mathcal{E} \left(\sqrt{\frac{d\mu}{dp_{\sigma^2}}} \right) \quad (56)$$

where $KL(\mu, \nu) = \mathbb{E}_\mu \left[\log \frac{d\mu}{d\nu} \right]$ is the relative entropy. If $\nabla \log p$ were strongly concave, then there are well-known bounds on the log-Sobolev constant; as we assume no such convexity, we, as in [RRT17], use a dissipativity condition and the Lyapunov function criteria found in [BBCG08, CGW09], presented in the following two theorems:

Theorem 27. ([BBCG08]) *Let $V : \mathbb{R}^d \rightarrow [1, \infty)$ be a real valued function, and let \mathcal{L} be the generator of the diffusion W with stationary distribution p . If there are constants $\lambda_1, \lambda_2, R > 0$ such that*

$$\frac{\mathcal{L}V(x)}{V(x)} \leq -\lambda_1 + \lambda_2 \mathbf{1}_{B_R}(x) \quad (57)$$

Then p satisfies a Poincaré inequality with constant

$$c_P \leq \frac{1}{\lambda_1} \left(1 + C \lambda_2 R^2 e^{\text{osc}_R(\log p)} \right) \quad (58)$$

where $C > 0$ is a universal constant and for a continuous, real-valued function f , we let $\text{osc}_R(f) = \max_{B_R} f - \min_{B_R} f$.

Theorem 28. ([CGW09]) *Suppose that p is a measure such that $\nabla^2 \log p \succeq -KI_d$ for $K \geq 0$ in the sense of matrices and that p satisfies a Poincaré inequality with constant c_P . Further suppose that there is a Lyapunov function $V : \mathbb{R}^d \rightarrow [1, \infty)$ such that*

$$\frac{\mathcal{L}V(x)}{V(x)} \leq \kappa - \gamma \|x\|^2 \quad (59)$$

Then p satisfies a log-Sobolev inequality with constant

$$c_{LS} \leq \frac{2K}{\gamma} + \frac{2}{K} + c_P \left(2 + \frac{2K}{\gamma} (\kappa + \gamma \mathbb{E}_{X \sim p} [\|X\|^2]) \right) \quad (60)$$

Further discussion regarding both theorems can be found in the appendix of [RRT17]. With this in hand, we are ready to prove Proposition 19:

Proof. (Proof of Proposition 19) Given that we rely on the (m, b) -dissipativity and the $\frac{M}{2}$ -Lipschitz of p_{σ^2} , it suffices to prove in the case of $\sigma^2 = 0$ and apply Proposition 7. We apply in sequence the theorems from [BBCG08, CGW09]. Consider the following Lyapunov function: $V(x) = e^{\frac{m}{4}\|x\|^2}$. We compute:

$$\frac{\mathcal{L}V}{V} = \Delta \left(\frac{m}{4}\|x\|^2 \right) + \left\| \nabla \left(\frac{m}{4}\|x\|^2 \right) \right\|^2 + \left\langle \frac{m}{4}x, \nabla \log p(x) \right\rangle \quad (61)$$

$$= \frac{md}{2} + \frac{m^2}{4}\|x\|^2 - \frac{m}{2}\langle x, -\nabla \log p(x) \rangle \quad (62)$$

$$\leq \frac{m}{2}(d+b) - \frac{m^2}{4}\|x\|^2 \quad (63)$$

by the dissipativity assumption. Let

$$R^2 = \frac{4(d + \frac{b}{2})}{m} \quad \lambda_1 = m(d+b) \quad \lambda_2 = \frac{m}{2}(d+b) \quad (64)$$

in the theorem of [BBCG08]. Then we see that by Lemma 38 and the fundamental theorem of calculus, we may choose B such that

$$\text{osc}_R(\log p) \leq (M+B)R^2 + B \quad (65)$$

Thus we have

$$c_P \leq \frac{2}{m(d+b)} \left(1 + C(d+b)^2 e^{\frac{8(M+B)(d+b)}{m}} \right) \quad (66)$$

Now, we may apply the theorem of [CGW09] with the same Lyapunov function. Note that as $\nabla \log p$ is $\frac{M}{2}$ -Lipschitz, we have $\nabla^2 \log p \geq -MI_d$. Then if we set

$$\kappa = \frac{m}{2}(d+b) \quad \gamma = \frac{m^2}{4} \quad (67)$$

then the above computation with the Lyapunov function shows that the assumptions of the theorem of [CGW09] hold. In order to conclude, we need to bound $\mathbb{E}_p[\|X\|^2]$. Let $W(t)$ be the Langevin diffusion initialized on a measure with finite second moment. Then by Lemma 41, we have

$$\mathbb{E}_p[\|X\|^2] = \lim_{t \rightarrow \infty} \mathbb{E}[\|W(t)\|^2] \leq \lim_{t \rightarrow \infty} \mathbb{E}[\|W(0)\|^2] e^{-2mt} + \frac{b+d}{m}(1 - e^{-2mt}) = \frac{b+d}{m} \quad (68)$$

The result follows. ■

C Bounding the Distance between $\widehat{W}(t)$ and $W(t)$

In this appendix, we provide a detailed account of the results appearing in Section 4.2, in particular a proof of the key proposition, Proposition 23. We assume that \widehat{f} is Lipschitz and, for the sake of convenience, we assume that $\widehat{f}, \nabla \log p_{\sigma^2}$ have the same Lipschitz constant $\frac{M}{2}$. Again we let $W(t)$ evolve according to the continuous Langevin process, and we consider the process $\widehat{W}(t)$, the unique solution

$$d\widehat{W}(t) = \widehat{f}(\widehat{W}(t)) dt + \sqrt{2}dB(t) \quad \widehat{W}(0) = W(0)$$

First, we need to control the Wasserstein distance in terms of the relative entropy. In order to do this, we apply the following result of Bolley and Villani:

Theorem 29 (Corollary 2.3 from [BV05]). *Let $X \sim \mu$ be a random variable in \mathbb{R}^d and suppose that*

$$C = 2 \inf_{\alpha > 0} \left(\frac{1}{\alpha} \left(\frac{3}{2} + \log \mathbb{E} \left[e^{\alpha \|X\|^2} \right] \right) \right)^{\frac{1}{2}} < \infty \quad (69)$$

Then for all $\nu \ll \mu$

$$W_2(\nu, \mu) \leq C \left(KL(\nu, \mu)^{\frac{1}{2}} + \left(\frac{KL(\nu, \mu)}{2} \right)^{\frac{1}{4}} \right) \quad (70)$$

As a corollary, we have

Proposition 30. *Suppose that \widehat{f} is (m, b) -dissipative and $\frac{M}{2}$ -Lipschitz, and that we are in the setting of Assumption 4 for some $\alpha < m$. Then if ν_t is the law of $W(t)$ and $\mu \ll \nu_t$ then*

$$W_2(\mu, \widehat{\nu}_t) \leq 2 \left(\frac{3}{2\alpha} + \frac{k_\alpha}{\alpha} + 2(b+d)t \right)^{\frac{1}{2}} \left(\sqrt{KL(\mu, \widehat{\nu}_t)} + \left(\frac{KL(\mu, \widehat{\nu}_t)}{2} \right)^{\frac{1}{4}} \right) \quad (71)$$

In particular, if $KL(\mu, \widehat{\nu}_t) \leq 1$ then

$$W_2(\mu, \widehat{\nu}_t) \leq 32 \left(\frac{3}{2\alpha} + \frac{k_\alpha}{\alpha} + 2(b+d)t \right)^{\frac{1}{2}} (KL(\mu, \widehat{\nu}_t))^{\frac{1}{4}} \leq C_\alpha \sqrt{(b+d)t} KL(\mu, \widehat{\nu}_t)^{\frac{1}{4}} \quad (72)$$

Proof. The second statement follows immediately from the first and the fact that for $c \leq 1$, $c^{\frac{1}{2}} \leq c^{\frac{1}{4}}$. The first statement follows from Theorem 29 if we can bound the constant C . This bound follows immediately from Lemma 42, concluding the proof. \blacksquare

Thus we are left with bounding the relative entropy between ν_t , the law of $W(t)$ and $\widehat{\nu}_t$ the law of $\widehat{W}(t)$. We follow the method used in [RRT17, BEL18] of the application of the Girsanov theorem. In particular, we restate and prove Proposition 22:

Proposition 31. *Let $W(t), \widehat{W}(t)$ be as above and assume that $\nabla \log p_{\sigma^2}$ is $\frac{M}{2}$ -Lipschitz. Suppose further that $W(0) = \widehat{W}(0) \sim \mu_0$ such that*

$$\mathbb{E}_{X \sim \mu_0} \left[e^{2M^2 \|X\|^2} \right] < \infty \quad (73)$$

Then

$$KL(\nu_t, \widehat{\nu}_t) = \mathbb{E} \left[\frac{1}{4} \int_0^t \left\| \nabla \log p_{\sigma^2}(W(s)) - \widehat{f}(W(s)) \right\|^2 ds \right] \quad (74)$$

Proof. We apply the version of Girsanov's theorem found in Theorem 7.20 in [LS77]. Note that

$$\mathbb{E} \left[\exp \left(\frac{1}{4} \int_0^t \|\widehat{f}(\widehat{W}(s)) - \nabla \log p_{\sigma^2}(\widehat{W}(s))\|^2 ds \right) \right] \leq \mathbb{E} \left[\exp \left(2M^2 \int_0^t \|\widehat{W}(t)\|^2 ds \right) \right] < \infty \quad (75)$$

by Lemma 42. Thus by Girsanov's theorem, we have that

$$\frac{d\nu_t}{d\widehat{\nu}_t} = \exp \left(\frac{1}{\sqrt{2}} \int_0^t (\widehat{f}(\widehat{W}(s)) - \nabla \log p_{\sigma^2}(\widehat{W}(s))) dB_s - \frac{1}{4} \int_0^t \|\widehat{f}(\widehat{W}(s)) - \nabla \log p_{\sigma^2}(\widehat{W}(s))\|^2 ds \right) \quad (76)$$

and so

$$KL(\nu_t, \widehat{\nu}_t) = \mathbb{E}_{\nu_t} \left[\log \frac{\nu_t}{d\widehat{\nu}_t} \right] \quad (77)$$

$$= \mathbb{E} \left[\int_0^t (\widehat{f}(W(s)) - \nabla \log p_{\sigma^2}(W(s))) dB_s + \frac{1}{4} \int_0^t \|\widehat{f}(W(s)) - \nabla \log p_{\sigma^2}(W(s))\|^2 ds \right] \quad (78)$$

But the first term is a real martingale by Novikov's condition so has expectation zero, yielding the result. \blacksquare

Thus it suffices to bound

$$\mathbb{E} \left[\int_0^t \|\widehat{f}(W(s)) - \nabla \log p_{\sigma^2}(W(s))\|^2 ds \right]$$

We first need the following identity:

Lemma 32. *Let $U \subset \mathbb{R}^d$ be measurable and let $W(s)$ be as above. Let $\pi_s(x, y)$ be the transition density of $W(s)$. Then if $\mathbb{E}_x[\cdot]$ denotes expectation with respect to the measure associated with $W(s)$ started at $W(0) = x$, then*

$$\mathbb{E}_x \left[\int_0^t \mathbf{1}_U(W(s)) ds \right] = \int_U \left(\int_0^t \pi_s(x, y) ds \right) dy \quad (79)$$

Proof. Following the proof of [MP10, Theorem 3.32] and appealing to [GH80] for justification in the case that $W(t)$ is not just Brownian motion, we have

$$\mathbb{E}_x \left[\int_0^t \mathbf{1}_U(W(s)) ds \right] = \int_0^t \mathbb{E}_x [\mathbf{1}_U(W(s))] ds = \int_0^t \int_U \pi_s(x, y) dy ds = \int_A \left(\int_0^t \pi_s(x, y) ds \right) dy \quad (80)$$

by Fubini's theorem and the definition of the transition density. \blacksquare

In order to construct a bound on the relevant integral, we need a bound on $\pi_t(x, y)$. In order to construct a Gaussian bound, we adapt an argument of [Dow08]:

Lemma 33. *For fixed x , let $W(t)$ evolve as above and let $W(0) = x$. If $\nabla \log p_{\sigma^2}$ is $\frac{M}{2}$ -Lipschitz, $\pi_t(x, y)$ is the transition density of $W(t)$ for $y \in \mathbb{R}^d$, then we have*

$$\pi_t(x, y) \leq \frac{p_{\sigma^2}(y)}{p_{\sigma^2}(x)} e^{\frac{M\sqrt{d}}{2}t} g_t(x, y) \quad (81)$$

where $g_t(x, y)$ is the standard Gaussian heat kernel.

Proof. We adapt a proof in [Dow08] to the case of higher dimensions. By the same argument as in the proof of Proposition 22, we may apply Girsanov's theorem. Let \mathbb{Q}_x be a measure under which $W(t) = \tilde{B}_t$ is a \mathbb{Q}_x Brownian motion started at x . Let \mathbb{P}_x be the original measure pertaining to the Brownian motion B_t that drives $W(t)$. Then by Girsanov's theorem, we have

$$\left(\frac{d\mathbb{P}_x}{d\mathbb{Q}_x} \right)_t = \widehat{\mathbb{E}}_x \left[\exp \left(\int_0^t \nabla \log p_{\sigma^2}(W(s)) dW(s) - \frac{1}{2} \int_0^t \|\nabla \log p_{\sigma^2}(W(s))\|^2 ds \right) \right] \quad (82)$$

$$= \widehat{\mathbb{E}}_x \left[\exp \left(\int_0^t \nabla \log p_{\sigma^2}(W(s)) d\tilde{B}_s - \frac{1}{2} \int_0^t \|\nabla \log p_{\sigma^2}(W(s))\|^2 ds \right) \right] \quad (83)$$

where $\widehat{\mathbb{E}}_x$ denotes expectation with respect to \mathbb{Q}_x . Now, by Rademacher's theorem we may apply Ito's lemma to $\log p_{\sigma^2}(\cdot)$, thus we get

$$\log p_{\sigma^2}(W(t)) - \log p_{\sigma^2}(W(0)) = \int_0^t \nabla \log p_{\sigma^2}(W(s)) dW(s) + \frac{1}{2} \int_0^t \Delta \log p_{\sigma^2}(W(s)) ds \quad (84)$$

Rearranging, we get that

$$\int_0^t \nabla \log p_{\sigma^2}(W(s)) d\tilde{B}_s = \log \left(\frac{p_{\sigma^2}(W(t))}{p_{\sigma^2}(x)} \right) - \frac{1}{2} \int_0^t \Delta \log p_{\sigma^2}(W(s)) ds \quad (85)$$

By the fact that $\nabla \log p_{\sigma^2}$ is $\frac{M}{2}$ -Lipschitz, we have that $|\Delta \log p_{\sigma^2}(y)| \leq \sqrt{d}M$ for all $y \in \mathbb{R}^d$. Because $\|\nabla \log p_{\sigma^2}(\cdot)\|^2 \geq 0$, we have by the above work and the fact that the transition density of Brownian motion is $g_t(x, y)$, that

$$\pi_t(x, y) \leq \frac{p_{\sigma^2}(y)}{p_{\sigma^2}(x)} e^{\frac{M\sqrt{d}}{2}t} g_t(x, y) \quad (86)$$

as desired. \blacksquare

In order to bound the right hand side in Lemma 32, we need to introduce the notion of asymmetric decreasing rearrangements. A full exposition on the topic can be found in [BS88]. We have the following definition

Definition 34. *Let μ be a probability measure on \mathbb{R}^d and let f be a nonnegative, measurable function $f : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$. We define for all $s \geq 0$,*

$$\mu^f(s) = \mu(\{f(x) > s\}) \quad (87)$$

We define the decreasing rearrangement of f to be

$$f^*(t) = \inf \{s > 0 \mid \mu^f(s) \leq t\} \quad (88)$$

Note that the specific case when μ is the Lebesgue measure is known as a symmetric decreasing rearrangement and is well known to geometric analysts. In this special case, there is a well known inequality, the Hardy-Littlewood inequality that governs integrals of products of rearrangements. A generalization to the arbitrary μ case, whose proof can be found in [BS88], is

Theorem 35. [BS88, Theorem 2.2] *Let $f, g : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$ be measurable functions. Then*

$$\int_{\mathbb{R}^d} f(x)g(x)d\mu(x) \leq \int_0^\infty f^*(t)g^*(t)dt \quad (89)$$

With this result in hand, we are able to prove the following proposition:

Proposition 36. *Considering the setting of Lemma 32, let $U \subset \mathbb{R}^d$ be a measurable set. If $d \geq 3$, then*

$$\mathbb{E}_x \left[\int_0^t \mathbf{1}_U(W(s))ds \right] \leq \frac{3e^{\frac{1}{e}}}{2\pi} \|p_{\sigma^2}\|_\infty^{1-\frac{2}{d}} \frac{e^{\frac{M\sqrt{d}}{2}t}}{p_{\sigma^2}(x)} p_{\sigma^2}(U)^{\frac{2}{d}} \quad (90)$$

Proof. By Lemmas 32 and 33, we have

$$\mathbb{E}_x \left[\int_0^t \mathbf{1}_U(W(s))ds \right] = \int_U \int_0^t \pi_s(x, y)dsdy \leq \int_U \int_0^t \frac{p_{\sigma^2}(y)}{p_{\sigma^2}(x)} e^{\frac{M\sqrt{d}}{2}s} g_s(x, y)dsdy \quad (91)$$

$$\leq \frac{e^{\frac{M\sqrt{d}}{2}t}}{p_{\sigma^2}(x)} \int_U \int_0^t g_s(x, y)p_{\sigma^2}(y)dsdy \quad (92)$$

Note now, that for $d \geq 3$,

$$\int_0^t g_s(x, y)ds \leq \int_0^\infty g_s(x, y)ds = \frac{\Gamma(\frac{d}{2} - 1)}{2\pi^{\frac{d}{2}}} \|x - y\|^{2-d} \quad (93)$$

by, for instance, [MP10, Theorem 3.33]. Let $f(y) = \mathbf{1}_U(y)$ and let $g(y) = \|x - y\|^{2-d}$. Then we note that

$$p_{\sigma^2}^f(s) = \begin{cases} p_{\sigma^2}(U) & s < 1 \\ 0 & \text{otherwise} \end{cases} \quad p_{\sigma^2}^g(s) = p_{\sigma^2} \left(B_d \left(x, s^{-\frac{1}{d-2}} \right) \right) \quad (94)$$

where $B_d(x, r)$ denotes the d -dimensional ball centred at x with radius r . Thus we have that

$$f^*(t) = \mathbf{1}_{\{s < p_{\sigma^2}(U)\}}(t) \quad g^*(t) = \inf \left\{ s | p_{\sigma^2} \left(B_d(x, s^{-\frac{1}{d-2}}) \right) \leq t \right\} \quad (95)$$

Let $\omega_d = \frac{\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2}+1)}$ be the Lebesgue volume of the unit ball. Then we note that if $s = \left(\frac{t}{\omega_d \|p_{\sigma^2}\|_\infty} \right)^{\frac{2}{d}-1}$, then we have

$$p_{\sigma^2} \left(B_d(x, s^{-\frac{1}{d-2}}) \right) \leq \|p_{\sigma^2}\|_\infty \omega_d s^{-\frac{d}{d-2}} = t \quad (96)$$

Thus

$$g^*(t) \leq \left(\frac{t}{\omega_d \|p_{\sigma^2}\|_\infty} \right)^{\frac{2}{d}-1} \quad (97)$$

Now, applying Theorem 35, with $d\mu(y) = p_{\sigma^2}(y)dy$, we have

$$\int_U \int_0^t g_s(x, y)ds p_{\sigma^2}(y)dy \leq \int_U \frac{\Gamma(\frac{d}{2} - 1)}{2\pi^{\frac{d}{2}}} \|x - y\|^{2-d} p_{\sigma^2}(y)dy \quad (98)$$

$$\leq \frac{\Gamma(\frac{d}{2} - 1)}{2\pi^{\frac{d}{2}}} \int_0^\infty \mathbf{1}_{\{s < p_{\sigma^2}(U)\}}(t) \left(\frac{t}{\omega_d \|p_{\sigma^2}\|_\infty} \right)^{\frac{2}{d}-1} dt \quad (99)$$

$$= \frac{\Gamma(\frac{d}{2} - 1)}{2\pi^{\frac{d}{2}}} \int_0^{p_{\sigma^2}(U)} \left(\frac{t}{\omega_d \|p_{\sigma^2}\|_\infty} \right)^{\frac{2}{d}-1} dt \quad (100)$$

$$= \frac{\Gamma(\frac{d}{2} - 1)}{2\pi^{\frac{d}{2}}} \omega_d^{1-\frac{2}{d}} \|p_{\sigma^2}\|_\infty^{1-\frac{2}{d}} \frac{d}{2} p_{\sigma^2}(U)^{\frac{2}{d}} \quad (101)$$

Now, note that

$$\frac{\Gamma(\frac{d}{2}-1)}{2\pi^{\frac{d}{2}}}\omega_d = \frac{2}{d(d-2)} \quad (102)$$

using the fact that $\Gamma(x+1) = x\Gamma(x)$. Moreover, we have

$$\omega_d^{-\frac{2}{d}} = \left(\frac{\Gamma(\frac{d}{2}+1)}{\pi^{\frac{d}{2}}}\right)^{\frac{2}{d}} = \frac{(\frac{d}{2})^{\frac{2}{d}}\Gamma(\frac{d}{2})^{\frac{2}{d}}}{\pi} \leq \frac{e^{\frac{1}{2}}\frac{d}{2}}{\pi} \quad (103)$$

by the fact that $x^{\frac{1}{x}} \leq e^{\frac{1}{e}}$ and the fact that $\Gamma(x) \leq x^x$. Thus,

$$\int_U \int_0^t g_s(x, y) ds dy \leq \|p_{\sigma^2}\|_{\infty}^{1-\frac{2}{d}} \frac{e^{\frac{1}{e}}}{\pi} \frac{d}{2} \frac{2}{d(d-2)} \frac{d}{2} p_{\sigma^2}(U)^{\frac{2}{d}} \leq \frac{3e^{\frac{1}{e}}}{2\pi} \|p_{\sigma^2}\|_{\infty}^{1-\frac{2}{d}} p_{\sigma^2}(U)^{\frac{2}{d}} \quad (104)$$

Plugging this into the first set of inequalities above concludes the proof. \blacksquare

With this in mind, we are able to prove the key lemma:

Lemma 37. *Let $\phi : \mathbb{R}^d \rightarrow \mathbb{R}_{\geq 0}$ be measurable and $W(t)$ as above and suppose that $\mathbb{E}[\phi(X)] \leq \varepsilon^2$, where X is distributed according to p_{σ^2} . Let $x \in \mathbb{R}^d$. Then*

$$\mathbb{E}_x \left[\int_0^t \phi(W(s)) ds \right] \leq \varepsilon t + \sqrt{\left(\int_0^t \mathbb{E}_x [\phi(W(s))^2] ds \right) \frac{3e^{\frac{1}{e}}}{2\pi} \|p_{\sigma^2}\|_{\infty}^{1-\frac{2}{d}} \frac{e^{\frac{M\sqrt{d}}{2}} t}{p_{\sigma^2}(x)} \varepsilon^{\frac{2}{d}}} \quad (105)$$

where \mathbb{E}_x denotes expectation with respect to $W(t)$ started at $W(0) = x$.

Proof. Let $U = \{y \in \mathbb{R}^d | \phi(y) > \varepsilon\}$. Then we have

$$\int_0^t \phi(W(s)) ds = \int_0^t \phi(W(s)) \mathbf{1}_{U^c} ds + \int_0^t \phi(W(s)) \mathbf{1}_U ds \leq \varepsilon t + \int_0^t \phi(W(s)) \mathbf{1}_U ds \quad (106)$$

By Cauchy-Schwarz,

$$\mathbb{E}_x \left[\int_0^t \phi(W(s)) \mathbf{1}_U ds \right] \leq \sqrt{\int_0^t \mathbb{E}_x [\phi(W(s))^2] ds \int_0^t \mathbb{E}_x [\mathbf{1}_U] ds} \quad (107)$$

By assumption, we have

$$\varepsilon^2 \geq \mathbb{E}[\phi(X)] \geq \mathbb{E}[\phi(X) \mathbf{1}_U] \geq \varepsilon \int_U p_{\sigma^2}(y) dy \quad (108)$$

Thus we have that $p_{\sigma^2}(U) \leq \varepsilon$. Applying Proposition 36 to bound the second factor under the square root concludes the proof. \blacksquare

Finally, we are able to prove Proposition 23.

Proof. (Proof of Proposition 23) We apply Lemma 37 to $\phi(x) = \|\nabla \log p(x) - \hat{f}(x)\|^2$. By the Lipschitz condition and Lemma 38, we note that

$$\phi(x)^2 \leq 16M^4 \|x\|^4 + 16B^4 \quad (109)$$

and by Lemma 41, we have then

$$\int_0^t \mathbb{E}_x [\phi(W(s))^2] ds \leq \left(16M^4 \left(\mathbb{E} [\|W(0)\|^4] + \frac{(b+d+2)(\mathbb{E} [\|W(0)\|^2] + \frac{b+d}{m})}{m} \right) + 16B^4 \right) t \quad (110)$$

Thus there exists a constant C depending on the initialization and the smoothness parameters m, b, M, B such that

$$\int_0^t \mathbb{E}_x [\phi(W(s))^2] ds \leq Cd^2t \quad (111)$$

Now, note that since $W(0)$ is concentrated with high probability in a ball of radius R , we have by Lemma 38 that $p_{\sigma^2}(W(0)) \geq Ce^{-MR^2 - BR}$. Thus by Lemma 37, we have that

$$\mathbb{E} \left[\int_0^t \phi(W(s)) ds \right] \leq \varepsilon t + \sqrt{Cd^2te^{MR^2 + BR} \frac{3e^{\frac{1}{\varepsilon}}}{2\pi} \|p_{\sigma^2}\|_{\infty}^{1 - \frac{2}{d}} e^{\frac{M\sqrt{d}}{2}t} \varepsilon^{\frac{2}{d}}} \leq \varepsilon t + C \|p\|_{\infty}^{\frac{1}{2} - \frac{1}{d}} e^{\frac{M\sqrt{d}}{4}t} \sqrt{t} \varepsilon^{\frac{1}{d}} \quad (112)$$

as desired. \blacksquare

D Miscellaneous Proofs

Proof. (Proof of Proposition 5) Let $\varepsilon \sim g_{\sigma^2}$ and $X \sim p$. Then we have, letting $y = x + \varepsilon$,

$$L_{DAE}(r) = \mathbb{E}_X \mathbb{E}_{\varepsilon} [\|r(x + \varepsilon) - x\|^2] = \int \int \|r(y) - y + \varepsilon\|^2 p(y - \varepsilon) g(\varepsilon) dy d\varepsilon \quad (113)$$

$$= \int \int \|r(y) - y\|^2 p(y - \varepsilon) g(\varepsilon) dy d\varepsilon + \int \int 2\langle \varepsilon, r(y) - y \rangle p(y - \varepsilon) g(\varepsilon) dy d\varepsilon \quad (114)$$

$$+ \int \int \|\varepsilon\|^2 p(y - \varepsilon) g(\varepsilon) dy d\varepsilon \quad (115)$$

The last term above does not depend on r and so we may ignore it. We focus now on the second term. Let $\xi \sim g_1$ be a standard Gaussian and let $s'(x) = r(x) - x$. By Lemma 46, we have that

$$\int \langle \varepsilon, s'(x + \varepsilon) \rangle g(\varepsilon) d\varepsilon = \sigma \int \langle \xi, s'(x + \sigma\xi) \rangle g_1(\xi) d\xi \quad (116)$$

$$= \sigma^2 \frac{1}{\sigma} \int \langle \xi, s'(x + \sigma\xi) \rangle g_1(\xi) d\xi \quad (117)$$

$$= \sigma^2 \operatorname{div}(\int s'(x + \varepsilon) g(\varepsilon) d\varepsilon) \quad (118)$$

where we used Gaussian Stein identity above. Now, note that as we know that p_{σ^2} is a density, it must tend to zero as $\|x\| \rightarrow \infty$. Thus we may apply the divergence theorem to get

$$\int \int 2\langle \varepsilon, r(y) - y \rangle p(y - \varepsilon) g(\varepsilon) dy d\varepsilon = 2\sigma^2 \int p(x) \operatorname{div}(\mathbb{E}_{\varepsilon}[s'(x + \varepsilon)]) dx \quad (119)$$

$$= 2\sigma^2 \int \operatorname{div}(s'(y)) p_{\sigma^2}(y) dy \quad (120)$$

$$= -2\sigma^2 \int \left\langle s'(x), \frac{\nabla p_{\sigma^2}(x)}{p_{\sigma^2}(x)} \right\rangle p_{\sigma^2}(x) dx \quad (121)$$

Thus we have that

$$L_{DAE}(r) = \mathbb{E}_{X \sim p_{\sigma^2}} [\|r(X) - X\|^2 - 2\sigma^2 \langle s'(X), \nabla \log p_{\sigma^2}(X) \rangle] + C(p, \sigma^2) \quad (122)$$

$$= \mathbb{E}_{X \sim p_{\sigma^2}} [\|s'(X) - \sigma^2 \nabla \log p_{\sigma^2}(X)\|^2] + C(p, \sigma^2) - \sigma^4 \mathbb{E}_{X \sim p_{\sigma^2}} [\|\nabla \log p_{\sigma^2}\|^2] \quad (123)$$

$$= \mathbb{E}_{X \sim p_{\sigma^2}} [\|s'(X) - \sigma^2 \nabla \log p_{\sigma^2}(X)\|^2] + C'(p, \sigma^2) \quad (124)$$

where $C(p, \sigma^2)$, $C'(p, \sigma^2)$ do not depend on r . Dividing by σ^2 and setting $s(x) = \frac{s'(x)}{\sigma^2}$ shows that

$$L_{DSM}(s) = \mathbb{E}_{p_{\sigma^2}} [\|s(x) - \nabla \log p_{\sigma^2}\|^2] = \frac{1}{\sigma^2} L_{DAE}(r) + C(p, \sigma^2) \quad (125)$$

Thus, the two losses are equivalent to minimize with respect to r or s . \blacksquare

Proof. (Alternate proof of Corollary 6) For $y \in \mathbb{R}^d$, the loss of the DAE is given by

$$\int_{\mathbb{R}^d} \mathbb{E}_\epsilon [p(y) \|r(y + \epsilon) - y\|^2] dy = \int_{\mathbb{R}^d} \mathbb{E}_\epsilon [p(x - \epsilon) \|r(x) - x + \epsilon\|^2] dx \quad (126)$$

where we substituted $x = y + \epsilon$. Now by the calculus of variations, it suffices to minimize the integrand with respect to $r(x)$ for each $x \in \mathbb{R}^d$. Taking the derivative and setting it equal to zero gives

$$r_{\sigma^2}(x) = \frac{\mathbb{E}_\epsilon [p(x - \epsilon)(x - \epsilon)]}{\mathbb{E}_\epsilon [p(x - \epsilon)]} \quad (127)$$

the result given by [AB14, Theorem 1]. By linearity, then we have

$$r_{\sigma^2}(x) = \frac{\mathbb{E}_\epsilon [xp(x - \epsilon)]}{\mathbb{E}_\epsilon [p(x - \epsilon)]} - \frac{\mathbb{E}_\epsilon [\epsilon p(x - \epsilon)]}{\mathbb{E}_\epsilon [p(x - \epsilon)]} = x - \frac{\mathbb{E}_\epsilon [\epsilon p(x - \epsilon)]}{\mathbb{E}_\epsilon [p(x - \epsilon)]} \quad (128)$$

But we have

$$\mathbb{E}_\epsilon [\epsilon p(x - \epsilon)] = \int \epsilon p(x - \epsilon) g_{\sigma^2}(\epsilon) d\epsilon = \sigma^2 \int \nabla g_{\sigma^2}(\epsilon) p(x - \epsilon) d\epsilon = -\sigma^2 \int g_{\sigma^2}(\epsilon) \nabla p(x - \epsilon) d\epsilon \quad (129)$$

by Lemma 46. But then we have

$$\frac{\mathbb{E}_\epsilon [\epsilon p(x - \epsilon)]}{\mathbb{E}_\epsilon [p(x - \epsilon)]} = -\sigma^2 \frac{\mathbb{E}_\epsilon [-\nabla_x p(x - \epsilon)]}{\mathbb{E}_\epsilon [p(x - \epsilon)]} = \sigma^2 \frac{\nabla p * g_{\sigma^2}(x)}{p * g_{\sigma^2}(x)} = \sigma^2 \nabla \log p_{\sigma^2} \quad (130)$$

Putting this together yields the result. \blacksquare

Proof. (Proof of Proposition 7) Let $\sigma^2 \leq \sigma_{max}^2$. Let $\eta_{\sigma^2}(X) = \nabla \log p_{\sigma^2}(X) - \nabla \log p(X)$. By Assumption 2, η_{σ^2} is Lipschitz with constant $\sigma^2 \frac{M}{2}$. Thus, we have:

$$\langle -\nabla \log p_{\sigma^2}(x), x \rangle = \langle -\nabla \log p(x), x \rangle - \langle \eta_{\sigma^2}(X), x \rangle \quad (131)$$

By Lemma 38 we know that there is some constant, which, by raising B if necessary, we may take to be equal to B , such that

$$\|\eta_{\sigma^2}(x)\| \leq \sigma^2 M \|x\| + B \quad (132)$$

By Cauchy-Schwarz,

$$|\langle \eta_{\sigma^2}(X), x \rangle| \leq \|\eta_{\sigma^2}(x)\| \cdot \|x\| \leq \sigma^2 M \|x\|^2 + B \|x\| \quad (133)$$

and thus

$$\langle \eta_{\sigma^2}(X), x \rangle \geq -\sigma^2 M \|x\|^2 - B \|x\| \quad (134)$$

By the dissipativity assumption, we have

$$\langle -\nabla \log p(x), x \rangle \geq m \|x\|^2 - b \quad (135)$$

Thus we have

$$\langle -\nabla \log p_{\sigma^2}(x), x \rangle \geq (m - \sigma^2 M) \|x\|^2 - b - B \|x\| \quad (136)$$

$$\geq \frac{m - \sigma^2 M}{2} \|x\|^2 - b + \frac{m - \sigma^2 M}{2} \|x\|^2 - B \|x\| \quad (137)$$

$$\geq \frac{m - \sigma^2 M}{2} \|x\|^2 - b - \frac{B^2}{2(m - \sigma^2 M)} \quad (138)$$

where the last inequality follows by the fact that

$$\frac{m - \sigma^2 M}{2} \|x\|^2 - B \|x\| = \frac{m - \sigma^2 M}{2} \left(\|x\| - \frac{B}{m - \sigma^2 M} \right)^2 - \frac{B^2}{2(m - \sigma^2 M)} \quad (139)$$

$$\geq -\frac{B^2}{2(m - \sigma^2 M)} \quad (140)$$

\blacksquare

Proof. (Proof of Proposition 18) Consider the coupling where the Brownian motion driving $W(t)$ also generates the Gaussians in W_k . Let $\widehat{W}(s)$ be the continuous time process such that $\widehat{W}(s) = W_{\lfloor \frac{s}{\eta} \rfloor \eta}$ and $\widehat{W}(0) = W(0)$. Recall $\tau = k\eta$. Then we can compute

$$\begin{aligned} \mathbb{E} \left[\|W(\tau) - W_k\|^2 \right] &= \mathbb{E} \left[\|W(\tau) - \widehat{W}(\tau)\|^2 \right] = \mathbb{E} \left[\left\| \int_0^\tau \nabla \log p(W(s)) - \nabla \log p(\widehat{W}(s)) ds \right\|^2 \right] \\ &\leq \int_0^\tau \mathbb{E} \left[\left\| \nabla \log p(W(s)) - \nabla \log p(\widehat{W}(s)) \right\|^2 \right] ds \\ &\leq M^2 \sum_{j=0}^{k-1} \int_{j\eta}^{(j+1)\eta} \mathbb{E} \left[\|W(s) - W_j\|^2 \right] ds \end{aligned}$$

Now note that for $j\eta \leq s < (j+1)\eta$, we have

$$\mathbb{E} \left[\|W(s) - W_j\|^2 \right] \leq 2\mathbb{E} \left[\|W(s) - W(j\eta)\|^2 \right] + 2\mathbb{E} \left[\|W(j\eta) - W_j\|^2 \right]$$

For the first term, we have

$$\mathbb{E} \left[\|W(s) - W_j\|^2 \right] = \mathbb{E} \left[\left\| \int_{j\eta}^s \nabla \log p(W(u)) du + \sqrt{2} \int_{j\eta}^s dB_u \right\|^2 \right] \quad (141)$$

$$\leq 2\eta M^2 \mathbb{E} \left[\|W(s)\|^2 \right] + 4d\eta \leq 2\eta M^2 \mathbb{E} \left[\|W(\tau)\|^2 \right] + 4d\eta \quad (142)$$

where we recall that $\tau = k\eta$. Plugging this last bound into our above inequality, we get

$$\mathbb{E} \left[\|W(\tau) - W_k\|^2 \right] \leq M^2 \tau (2\eta M^2 \mathbb{E} \left[\|W(\tau)\|^2 \right] + 4d\eta) + M^2 \eta \sum_{j=0}^{k-1} \mathbb{E} \left[\|W(j\eta) - W_{j\eta}\|^2 \right]$$

Applying the discrete Gronwall lemma, we have

$$\mathbb{E} \left[\|W(\tau) - W_k\|^2 \right] \leq M^2 \tau (2\eta M^2 \mathbb{E} \left[\|W(\tau)\|^2 \right] + 4d\eta) e^{M^2 \tau} \quad (143)$$

To conclude, we apply Lemma 41 to bound $\mathbb{E} \left[\|W(\tau)\|^2 \right]$ and the result follows from the fact that the Wasserstein distance is bounded by the above explicit coupling of the two laws μ_k and ν_τ to get

$$\mathcal{W}_2(\mu_k, \nu_\tau)^2 \leq M^2 \tau \eta \left(4d + M^2 \left(\mathbb{E} \left[\|W(0)\|^2 \right] + \frac{b+d}{m} \right) \right) e^{M^2 \tau} \leq Cd\eta\tau e^{M^2 \tau} \quad (144)$$

with C depending on M, b, m , and $\mathbb{E} \left[\|W(0)\|^2 \right]$ as desired. \blacksquare

E Auxiliary Lemmata

Lemma 38. *If $\nabla \log p$ is $\frac{M}{2}$ -Lipschitz, then there is some constant B such that*

$$\|\nabla \log p(x)\|^p \leq M^p \|x\|^p + B^p \quad (145)$$

for all $p \geq 1$.

Proof. By the definition of Lipschitz, we have that

$$\|\nabla \log p(x)\| \leq \|\nabla \log p(x) - \nabla \log p(0)\| + \|\nabla \log p(0)\| \leq \frac{M}{2} \|x\| + \|\nabla \log p(0)\|$$

Applying Minkowski's inequality concludes the proof. \blacksquare

Lemma 39. For all $p > 1$ and all $t > 0$, we have that if $\nabla \log p$ is $\frac{M}{2}$ Lipschitz, then

$$\mathbb{E} [||W(t)||^p] \leq \left((2dt)^{\frac{p}{2}} + B^p t \right) e^{M^p t} \quad (146)$$

where B is as appears in Lemma 38.

Proof. By the triangle inequality and Lemma 38, we have

$$||W(t)||^p = \left\| \sqrt{2}B_t + \int_0^t \nabla \log p(W(s)) ds \right\|^p \leq 2^{\frac{p}{2}} ||B_t||^p + \int_0^t ||\nabla \log p(W(s))||^p ds \quad (147)$$

$$\leq 2^{\frac{p}{2}} ||B_t||^p + \int_0^t (M^p ||W(s)||^p + B^p) ds \quad (148)$$

Taking expected values and applying Fubini, we get

$$\mathbb{E} [||W(t)||^p] \leq 2^{\frac{p}{2}} \mathbb{E} [||B_t||^p] + \int_0^t (M^p \mathbb{E} [||W(s)||^p] + B^p) ds \quad (149)$$

$$\leq (2dt)^{\frac{p}{2}} + B^p t + M^p \int_0^t \mathbb{E} [||W(s)||^p] ds \quad (150)$$

Applying Gronwall's inequality finishes the proof. ■

Remark 40. The following lemma will demonstrate that the bound in Lemma 39 is not tight as an inductive argument applied to the result below would show that each moment of $W(t)$ is actually bounded uniformly in time. The utility of the above lemma is that we have finiteness of all moments without the extra difficulties of iterating the proof below.

Lemma 41. There exists a constant κ depending only on the initialization of $W_{\sigma^2}(0)$ such that for all t ,

$$\mathbb{E} [||W_{\sigma^2}(t)||^2] \leq \mathbb{E} [||W(0)||^2] e^{-2mt} + \frac{b+d}{m} \quad (151)$$

$$\mathbb{E} [||W_{\sigma^2}(t)||^4] \leq \mathbb{E} [||W(0)||^4] + \frac{(b+d+2) (\mathbb{E} [||W(0)||^2] + \frac{b+d}{m})}{m} \quad (152)$$

Proof. We adapt the proof of [RRT17, Lemma 3.2]. Without loss of generality, we take $\sigma^2 = 0$ as the proof relies only on the Lipschitz and dissipative constants.

Let $Y(t) = ||W(t)||^4$. By Ito's lemma, then, we have

$$dY(t) = 4||W(t)||^2 \langle W(t), \nabla \log p(W(t)) \rangle dt + 4(d+2)||W(t)||^2 + 4||W(t)||^2 W(t) \sqrt{2} dB_t \quad (153)$$

Thus

$$d(e^{4mt} Y(t)) = 4me^{4mt} ||W(t)||^4 dt + 4e^{4mt} ||W(t)||^2 \langle W(t), \nabla \log p(W(t)) \rangle dt \quad (154)$$

$$+ 4e^{4mt} (d+2)||W(t)||^2 + 4||W(t)||^2 W(t) \sqrt{2} e^{4mt} dB_t \quad (155)$$

Thus we have

$$Y(t) = e^{-4mt} Y(0) + \int_0^t e^{4m(s-t)} 4||W(s)||^2 (\langle W(s), \nabla \log p(W(s)) \rangle + m||W(s)||^2) ds \quad (156)$$

$$+ \int_0^t e^{4m(s-t)} 4(d+2)||W(s)||^2 ds + \int_0^t e^{4m(s-t)} \sqrt{2} 4||W(s)||^2 W(s) dB_s \quad (157)$$

By Lemma 39, the last term is an actual martingale and so has expectation zero. By the dissipativity assumption, we have

$$\langle W(s), \nabla \log p(W(s)) \rangle + m||W(s)||^2 = -\langle -\nabla \log p(W(s)), W(s) \rangle + m||W(s)||^2 \quad (158)$$

$$\leq -m||W(s)||^2 + b + m||W(s)||^2 = b \quad (159)$$

Thus we have

$$\mathbb{E}[Y(t)] \leq e^{-4mt} \mathbb{E}[Y(0)] + \int_0^t e^{4m(s-t)} 4(b+d+2) \mathbb{E}[\|W(s)\|^2] ds \quad (160)$$

Now, we need to bound $\mathbb{E}[\|W(s)\|^2]$ independently of s . We repeat the same trick from [RRT17]. We define $Y'(t) = \|W(t)\|^2$. Then

$$dY'(t) = 2\langle W(t), \nabla \log p(W(t)) \rangle dt + 2ddt + \sqrt{2}W(t)dB_t \quad (161)$$

Thus

$$d(e^{2mt}Y'(t)) = 2me^{2mt}\|W(t)\|^2 dt + 2\langle W(t), \nabla \log p(W(t)) \rangle dt + 2ddt + \sqrt{2}W(t)dB_t \quad (162)$$

Again, we have

$$Y'(t) = e^{-2mt}Y'(0) + \int_0^t e^{2m(s-t)} 2(\langle W(s), \nabla \log p(W(s)) \rangle + m\|W(s)\|^2) ds \quad (163)$$

$$+ \int_0^t e^{2m(s-t)} 2ddt + \int_0^t e^{2m(s-t)} \sqrt{2}W(t)dB_t \quad (164)$$

Again, we note that the first integrand is bounded by $e^{2m(s-t)}2b$ by dissipativity and the last integral drops out in expectation. Thus

$$\mathbb{E}[\|W(t)\|^2] \leq e^{-2mt} \mathbb{E}[\|W(0)\|^2] + \int_0^t e^{2m(s-t)} 2(b+d) dt = e^{-2mt} \mathbb{E}[\|W(0)\|^2] + \frac{b+d}{m} (1 - e^{-2mt}) \quad (165)$$

which establishes the first inequality. For the second inequality, we see that

$$\mathbb{E}[\|W(t)\|^4] \leq e^{-4mt} \mathbb{E}[\|W(0)\|^4] + \int_0^t e^{4m(s-t)} 4(b+d+2) \left(\mathbb{E}[\|W(0)\|^2] + \frac{b+d}{m} \right) ds \quad (166)$$

$$= e^{-4mt} \mathbb{E}[\|W(0)\|^4] + \frac{(b+d+2) \left(\mathbb{E}[\|W(0)\|^2] + \frac{b+d}{m} \right)}{m} (1 - e^{-4mt}) \quad (167)$$

$$\leq \mathbb{E}[\|W(0)\|^4] + \frac{(b+d+2) \left(\mathbb{E}[\|W(0)\|^2] + \frac{b+d}{m} \right)}{m} \quad (168)$$

as desired. ■

Lemma 42. *Suppose that $-\nabla \log p_{\sigma^2}$ is $\frac{M}{2}$ -Lipschitz and (m, b) -dissipative. Suppose that $W(t)$ is a solution to Equation (1) initialized at $W(0)$ such that there is some $0 < \alpha \leq m$ such that $\log \mathbb{E} \left[e^{\alpha \|W(0)\|^2} \right] = k_\alpha < \infty$. Then for all $t > 0$,*

$$\log \mathbb{E} \left[e^{\alpha \|W(t)\|^2} \right] \leq k_\alpha + 2\alpha(b+d)t. \quad (169)$$

If \hat{f} is also $\frac{M}{2}$ -Lipschitz and (m, b) -dissipative, then the same result holds for $\widehat{W}(t)$, i.e.,

$$\log \mathbb{E} \left[e^{\alpha \|\widehat{W}(t)\|^2} \right] \leq k_\alpha + 2\alpha(b+d)t. \quad (170)$$

Remark 43. *Note that this is essentially [RRT17, Lemma 3.3] in our setting and is proved in the same way. The proof is included for the sake of completeness.*

Proof. As above, without loss of generality, we take $\sigma^2 = 0$. Let $Y(t) = e^{\alpha \|W(t)\|^2}$. By the Ito calculus, we have

$$dY(t) = 2\alpha W(t)Y(t)dW(t) + 2d\alpha Y(t)dt + 2\alpha^2 \|W(t)\|^2 Y(t)dt \quad (171)$$

$$= (2\langle W(t), \nabla \log p(W(t)) \rangle + 2d + 2\alpha \|W(t)\|^2) \alpha Y(t)dt + 2\sqrt{2}\alpha Y(t)W(t)dB_t \quad (172)$$

Thus

$$Y(t) = Y(0) + \int_0^t (2\langle W(s), \nabla \log p(W(s)) \rangle + 2d + 2\alpha \|W(s)\|^2) \alpha Y(s) ds + \int_0^t 2\sqrt{2}\alpha Y(s) W(s) dB_s \quad (173)$$

By [WGD04, Corollary 4.1], we know $\int_0^t \mathbb{E}[|Y(s)W(s)|^2] ds < \infty$ and thus the last term above is a real martingale. Taking expectations, we have by assumption that $\mathbb{E}[Y(0)] = e^{k\alpha}$ and thus we have

$$\mathbb{E}[Y(t)] = e^{k\alpha} + \mathbb{E} \left[\int_0^t (2\langle W(s), \nabla \log p(W(s)) \rangle + 2d + 2\alpha \|W(s)\|^2) \alpha Y(s) ds \right] \quad (174)$$

Now, we note that by dissipativity, we have

$$2\langle W(s), \nabla \log p(W(s)) \rangle + 2d + 2\alpha \|W(s)\|^2 \leq 2b - 2m \|W(s)\|^2 + 2d + 2\alpha \|W(s)\|^2 \leq 2b + 2d \quad (175)$$

by the assumption that $\alpha \leq m$. Thus we have

$$\mathbb{E}[Y(t)] \leq e^{k\alpha} + \int_0^t Y(s) 2\alpha(b+d) ds \quad (176)$$

By Gronwall's inequality, the first result follows. Applying the identical argument to $\widehat{W}(t)$ concludes the proof. ■

Lemma 44. *Let $-\nabla \log p$ be (m, b) -dissipative. Then*

$$\log p(x) \leq -\frac{m}{4} \|x\|^2 + \frac{2b^2}{m} + \log p(0) \quad (177)$$

Proof. By the fundamental theorem of calculus,

$$\log p(x) = \log p(0) + \int_0^1 \frac{d}{dt} (\log p(tx)) dt = \log p(0) + \int_0^1 \langle \nabla \log p(tx), x \rangle dt \quad (178)$$

$$\leq \log p(0) - \int_0^1 mt^2 \|x\|^2 + b dt = \log p(0) + b - \frac{m}{2} \|x\|^2 \quad (179)$$

$$= -\frac{m}{4} \|x\|^2 + \log p(0) - \frac{m}{4} \|x\|^2 + b \|x\| \quad (180)$$

by the dissipativity assumption. Maximizing the last two terms with respect to $\|x\|$ yields the result. ■

Lemma 45. *Let $-\nabla \log p$ be (m, b) -dissipative. Then there is a constant C such that for all $\|x\|$,*

$$p(x) \leq C e^{-\frac{m\|x\|^2}{4}} \quad (181)$$

In particular, p is $\frac{2}{m}$ -sub-Gaussian.

Proof. The first statement follows immediately from Lemma 44. The second follows from a Gaussian tail-bound. ■

Lemma 46. (*Gaussian Stein Identity, [Ste81]*) *Let $\xi \sim N(0, I_d)$ and let $g : \mathbb{R}^d \rightarrow \mathbb{R}$ be an almost everywhere differentiable function with $\mathbb{E}_\xi[|\nabla g(\xi)|] < \infty$. Then*

$$\mathbb{E}_\xi [g(\xi)\xi] = \mathbb{E}_\xi [\nabla g(\xi)] \quad (182)$$