

Bayesian Reliability Analysis Of The Power Law Process With Respect To The Higgins-Tsokos Loss Function For Modeling Software Failure Times

Freeh Alenezi¹ and Chris. Tsokos²

¹PhD Candidtae, Department of Mathematics and Statistics, *College of Arts and Sciences , University of South Florida*, Tampa, 33647, USA

²Distinguished University Professor, Department of Mathematics and Statistics, *College of Arts and Sciences , University of South Florida*, Tampa, 33647, USA

1 abstract

The Power Law Process, also known as Non-Homogeneous Poisson Process, has been used in various aspects, one of which is the software reliability assessment. Specifically, by using its intensity function to compute the rate of change of a software reliability as time-varying function. Justification of Bayesian analysis applicability to the Power Law Process was shown using real data. The probability distribution that best characterizes the behavior of the key parameter of the intensity function was first identified, then the likelihood-based Bayesian reliability estimate of the Power Law Process under the Higgins-Tsokos loss function was obtained. As a result of a simulation study and using real data, the Bayesian estimate shows an outstanding performance compared to the maximum likelihood estimate using different sample sizes. In addition, a sensitivity analysis was performed, resulting in the Bayesian estimate being sensitive to the prior selection; whether parametric or non-parametric.

Index terms— Reliability growth; intensity function; non-homogeneous Poisson process; kernel density; loss function; robustness

2 Introduction

Software reliability growth is often tested during the software development process to insure a good quality product. Repairable software is tested until a failure is detected, then fixed, and tested again until a new failure is detected. This reliability improvement of software has been studied for decades. Duane (1964) [1] introduced the "learning curve approach", which is a plot of the failure rate (or

the intensity function) of a system as a function of time. It is used to assess software reliability improvement over time. For example, software reliability has improved when we observe a negative curve, whereas a positive curve means that reliability is deteriorating. Stability in software reliability is achieved when there is no curve, i.e. the graph is a horizontal line. The number of failures in the interval $(0, t]$, $N(t)$, is considered a Poisson counting process after satisfying the following conditions [9, 6]:

1. $N(t = 0) = 0$.
2. Independent increment (counts of disjoint time intervals are independent).
3. It has an intensity function $V(t) = \lim_{\Delta t \rightarrow 0} \frac{P(N(t, t+\Delta t)=1)}{\Delta t}$.
4. Simultaneous failures do not exist ($\lim_{\Delta t \rightarrow 0} \frac{P(N(t, t+\Delta t)=2)}{\Delta t} = 0$).

The probability of a random value $N(t) = n$ is given by:

$$P(N(t) = n) = \frac{\exp\left\{-\int_0^t V(t)dt\right\} \left\{\int_0^t V(t)dt\right\}^n}{n!}, \quad t > 0. \quad (2.1)$$

Crow (1974) proposed a non-homogeneous Poisson process (NHPP) [3], which is a Poisson process with a time-varying intensity function, given by:

$$V(t) = V(t; \beta, \theta) = \frac{\beta}{\theta} \left(\frac{t}{\theta}\right)^{\beta-1}, \quad t > 0, \beta > 0, \theta > 0, \quad (2.2)$$

with β and θ as the shape and scale parameters, respectively. This NHPP is also known as the power law process (PLP).

The joint probability density function (PDF) of the ordered failure times T_1, T_2, \dots, T_n from an NHPP with intensity function $V(t; \beta, \theta)$ is given by:

$$f(t_1, \dots, t_n) = \prod_{i=1}^n V(t_i; \beta, \theta) \exp\left\{-\int_0^w V(t; \beta, \theta) dt\right\}, \quad (2.3)$$

where w is the so-called stopping time. Considering the failure truncation case ($w = t_n$), the conditional reliability function of the failure time T_n given $T_1 = t_1, T_2 = t_2, T_3 = t_3, \dots, T_{n-2} = t_{n-2}, T_{n-1} = t_{n-1}$ is a function of $V(t; \beta, \theta)$.

To monitor software reliability growth over time, an engineer can use the estimate of the β value, the key parameter in the intensity function, since it plays a significant role during the testing process. For $\beta > 1$, the number of failures would increase because the intensity function is increasing. On the other hand, if the intensity function is decreasing, $\beta < 1$ means that the number of failures would decrease, indicating improved software reliability. Note that in the case of a homogeneous Poisson process pertains when $\beta = 1$, in which case the intensity function will be $\frac{1}{\theta}$ and whatever changes have been made have had no effect on the outcome.

The NHPP has been used for analyzing software failure times, and for predicting the next failure event. Several publications show the effectiveness and usefulness of this model in assessing reliability growth [2, 10, 11, 6]. In addition, NHPP has been used to study drug effectiveness in breast cancer treatment [21] and in the formulation of a software cost model [13].

Since the intensity function is driving the NHPP, improving the existing methods to estimate the key parameter β will certainly improve the accuracy of reliability growth assessment and help the structuring of maintenance strategies. Molinares and Tsokos [5], obtained a Bayesian estimate of the parameter β and compared it with its approximate maximum likelihood estimate (MLE). The authors derived the Bayesian estimates with respect to squared-error loss function, using Burr, Jeffreys, and inverted gamma probability distributions as the prior PDFs for β .

In performing Bayesian analysis on a real world problem, we need some sort of justification for pursuing this particular type of analysis. Once we have identified the probability distribution that characterized the probabilistic behavior of the failure time, we need to identify the prior PDF of β and a loss function. The squared-error loss function is the most popular loss function used in Bayesian analysis because of its analytical tractability. It places a small weight on the estimates around the true value, but proportionally more weight on estimates far from the true value. Higgins and Tsokos [7] proposed a new loss function that places exponential weight on extreme deviations from the true value, while remaining mathematically tractable.

In the present study, we investigate the effectiveness of Bayesian analysis in using the Higgins-Tsokos (H-T) loss function (that puts the loss at the end of the process) for modeling software failure times. To accomplish this, we use the NHPP as the underlying failure distribution subject to using the Burr PDF as a prior of β . In addition, we utilize the H-T loss function to perform sensitive analysis of prior selections. We employ parametric and non-parametric priors, namely Burr, inverted gamma, Jeffery, and two kernel PDFs. Therefore, the primary objective of the study is to answer the following questions within a Bayesian framework:

1. What is the performance of the Bayesian estimate of β under the H-T loss function compared to its MLE when modeling software failure times using PLP?
2. Is the Bayesian estimate of β , using the H-T loss function in the PLP, sensitive to the selection of the prior PDF, both parametric and non-parametric?

The paper is organized as follows: Section 2 describes the theory and development of the Bayesian reliability model; Section 3 presents the results and discussion; Section 4 are the conclusions.

3 Theory and Bayesian Estimates

3.1 Review of the Analytical Power Law Process

The probability of achieving n failures of a given system in the time interval $(0, t]$ can be written as, [5, 6]:

$$P(x = n; t) = \frac{\exp\left\{-\int_0^t V(t; \beta, \theta) dt\right\} \left\{\int_0^t V(t; \beta, \theta) dt\right\}^n}{n!}, \quad t > 0, \quad (3.1.1)$$

where $V(t; \beta, \theta)$ is the intensity function given by (2.2). The reduced expression

$$P(x = n; t) = \frac{1}{n!} \exp\left\{-\frac{t^\beta}{\theta}\right\} \frac{t^{n\beta}}{\theta}, \quad (3.1.2)$$

is the PLP that is commonly known as the Weibull or NHPP.

If the PLP is the underlying failure model of the failure times $t_1, t_2, t_3, \dots, t_{n-1}$, and t_n , the conditional reliability function of t_n given $t_1, t_2, t_3, \dots, t_{n-1}$ can be written as, [5, 6]:

$$R(t_n | t_1, t_2, \dots, t_{n-1}) = \exp\left\{\int_{t_{n-1}}^{t_n} V(t; \beta, \theta) dt\right\}, \quad t_n > t_{n-1} > 0, \quad (3.1.3)$$

since it is independent of $t_1, t_2, t_3, \dots, t_{n-2}$.

Since the reliability function, equation (3.1.3), is written mathematically as a function of the intensity function, estimating the parameter β in the $V(t; \beta, \theta)$ leads to estimation of the reliability function.

The (MLE) of β is a function of the largest failure time and the MLE of θ is also a function of the MLE of β . Let T_1, T_2, \dots, T_n denote the first n failure times of the PLP, where $T_1 < T_2 < \dots < T_n$ are total times since the initial startup of the system. Thus, the truncated conditional PDF, $f_i(t | t_1, \dots, t_{i-1})$, in

the Weibull process and is given by, [5, 6]:

$$f_i(t|t_1, \dots, t_{i-1}) = \frac{\beta}{\theta} \left(\frac{t}{\theta}\right)^{\beta-1} \exp\left\{-\frac{t^\beta}{\theta} + \frac{t_{i-1}^\beta}{\theta}\right\}, \quad t_{i-1} < t. \quad (3.1.4)$$

With $t = (t_1, t_2, \dots, t_n)$, the likelihood function for the first n failure times of the PLP $T_1 = t_1, T_2 = t_2, \dots, T_n = t_n$ can be written as:

$$L(t; \beta) = \exp\left(-\left(\frac{t_n}{\theta}\right)^\beta\right) \left(\frac{\beta}{\theta}\right)^n \prod_{i=1}^n \left(\frac{t_i}{\theta}\right)^{\beta-1}. \quad (3.1.5)$$

The MLE for the shape parameter is given by, [2, 3, 5, 6]:

$$\hat{\beta}_n = \frac{n}{\sum_{i=1}^n \log\left(\frac{t_n}{t_i}\right)}, \quad (3.1.6)$$

and for the scale parameter is:

$$\hat{\theta}_n = \frac{t_n}{n^{1/\hat{\beta}_n}}. \quad (3.1.7)$$

Note that the MLE of θ depends on the MLE of β using the largest (last) observed failure time.

3.2 Development of the Bayesian Estimates

Crow [2, 3] failure data from a system undergoing developmental testing was used, by Molinares & Tsokos [5], to show how β varied depending on the last failure time (largest time), thus they proposed a Bayesian approach to the PLP. The authors also found that the MLE of β follows a four-parameter Burr probability distribution, $g(\beta; \alpha, \gamma, \delta, \kappa)$, known as the four-parameter Burr type XII probability distribution, with a PDF given by:

$$g_B(\beta) = g(\beta; \alpha, \gamma, \delta, \kappa) = \begin{cases} \frac{\alpha\kappa\left(\frac{\beta-\gamma}{\delta}\right)^{\alpha-1}}{\delta\left(1+\left(\frac{\beta-\gamma}{\delta}\right)^\alpha\right)^{\kappa+1}}, & \gamma \leq \beta < \infty \\ 0, & \text{otherwise} \end{cases}, \quad (3.2.1)$$

where the hyperparameters α, γ, δ and κ are being estimated using MLE in the goodness of fit (GOF) test applied to the β estimates. The Crow successive failure data for his system is given in Table 1.

Table 1: Crow's failure times of a system under development.

Failure times						
0.7	3.7	13.2	17.6	54.5	99.2	112.2
120.9	151	163	174.5	191.6	282.8	355.2
486.3	490.5	513.3	558.4	678.1	688	785.9
887	1010.7	1029.1	1034.4	1136.1	1178.9	1259.7
1297.9	1419.7	1571.7	1629.8	1702.4	1928.9	2072.3
2525.2	2928.5	3016.4	3181	3256.3	–	–

According to the reliability growth failure data, the system failed for the first time at 0.7 units of time, $t_1 = 0.7$, and it failed the 40th time at 3256.3 units of time, $t_{40} = 3256.3$. The MLE of the parameter β for $n = 40$ is, [5, 6]:

$$\hat{\beta}_{40} = \frac{40}{\sum_{i=1}^{40} \log\left(\frac{3256.3}{t_i}\right)} \simeq 0.49. \quad (3.2.2)$$

If β were treated in a non-Bayesian setting, its MLE would be given by Eq. (3.2.2).

In an experimental process, the largest time to failure could occur at any point in the series of failures for a given system. Therefore, consider the case where the largest failure is $t_{39} = 3181$. In such a case, the estimate of β_{39} is 0.48.

The largest failure time always affects the MLE of β . Thus, it is recommended that β not to be thought of as an unknown constant [5], but rather as an unknown random variable. This recommendation provides the opportunity to study Bayesian analysis in the PLP with respect to various selections of loss functions and priors.

The Bayesian estimates of β will be derived using H-T loss functions.

3.2.1 Bayesian Estimates Using the Higgins-Tsokos Loss Function

The H-T loss function (1976) is given by, [5]:

$$L(\hat{\xi}, \xi) = \frac{f_1 \exp\{f_2(\hat{\xi} - \xi)\} + f_2 \exp\{-f_1(\hat{\xi} - \xi)\}}{f_1 + f_2} - 1, \quad f_1, f_2 > 0. \quad (3.2.1)$$

Higgins and Tsokos [7] showed that it places more weight on the extreme underestimation and overestimation of the true value when $f_1 > f_2$ and $f_1 < f_2$, respectively. The risk using the H-T loss function, where $\xi = \beta$ represents the estimate of $\hat{\xi} = \hat{\beta}$, is given by:

$$E[L(\hat{\beta}, \beta)] = \int_{-\infty}^{\infty} \left[\frac{f_1 \exp\{f_2(\hat{\beta} - \beta)\} + f_2 \exp\{-f_1(\hat{\beta} - \beta)\}}{f_1 + f_2} - 1 \right] h(\beta|t) d\beta. \quad (3.2.2)$$

By differentiating $E[L(\hat{\beta}, \beta)]$ with respect to β and setting it equal to zero we solve for $\hat{\beta}$, the Bayesian estimate of β with respect to the H-T loss function, is given by:

$$\hat{\beta}_{B.TH} = \frac{1}{f_1 + f_2} \ln \left[\frac{\int_{-\infty}^{\infty} \exp \{f_1 \beta\} h(\beta|t) d\beta}{\int_{-\infty}^{\infty} \exp \{-f_2 \beta\} h(\beta|t) d\beta} \right]. \quad (3.2.3)$$

The Bayesian estimate of β with respect to the H-T loss function and Burr probability distribution, as the prior, has $h(\beta|t)$ given by:

$$h(\beta|t) = \frac{\int_{\gamma}^{\infty} \left(\frac{\beta}{\theta}\right)^n \exp \left\{ -\left(\frac{t_n}{\theta}\right)^{\beta} \right\} \prod_{i=1}^n \left(\frac{t_i}{\theta}\right)^{\beta-1} \frac{\left(\frac{\beta-\gamma}{\delta}\right)^{\alpha-1}}{\left(1+\left(\frac{\beta-\gamma}{\delta}\right)^{\alpha}\right)^{\kappa+1}} d\beta}{\int_{\gamma}^{\infty} \left(\frac{\beta}{\theta}\right)^n \exp \left\{ -\left(\frac{t_n}{\theta}\right)^{\beta} \right\} \prod_{i=1}^n \left(\frac{t_i}{\theta}\right)^{\beta-1} \frac{\left(\frac{\beta-\gamma}{\delta}\right)^{\alpha-1}}{\left(1+\left(\frac{\beta-\gamma}{\delta}\right)^{\alpha}\right)^{\kappa+1}} d\beta}. \quad (3.2.4)$$

With the use of Eq. (3.1.3), the conditional reliability of t_i , the analytical structure of the conditional Bayesian reliability estimate for the PLP that is subject to the above information, is given by:

$$\hat{R}_B(t_i|t_1, t_2, \dots, t_{i-1}) = \exp \left\{ - \int_{t_{i-1}}^{t_i} \hat{V}'_B(t; \beta, \theta) dt \right\}, \quad t_i > t_{i-1} > 0, \quad (3.2.5)$$

where

$$\hat{V}'_B(t; \beta_{B.TH}, \theta) = \frac{\hat{\beta}_{B.TH}}{\theta} \left(\frac{t}{\theta}\right)^{\hat{\beta}_{B.TH}-1}, \quad \theta > 0, t > 0, \quad (3.2.6)$$

where $\hat{\beta}_{B.TH}$ is the Bayesian estimate of β using the H-T loss function. We are also interested in comparing the Bayesian estimate, using the H-T loss function, with MLE of the subject parameter for different parametric and non-parametric priors, assuming β has a random behavior and θ is known; and also comparing Eq. (3.1.7) with an adjusted MLE considered as a function of β .

3.3 Sensitivity Analysis: Prior Selection

In this section, we seek the answer to the following question: Is the Bayesian estimate of β , using the H-T loss function in the PLP, sensitive to the selection of the prior, with parametric or non-parametric priors? Assuming β is a random variable, using simulated data, sensitivity analysis was done for the following parametric and non-parametric priors:

1. Jeffreys' prior [[8]]:

Jeffreys' prior is proportional to the square root of the determinant of the Fisher information matrix ($I(\beta)$). It is a non-informative prior, where the Jeffreys' prior for the PLP, considering that β , $I(\beta)$ is scalar in this case, is given by:

$$g_J(\beta) \propto \sqrt{I(\beta)} = \sqrt{-E\left(\frac{\partial^2 \text{Log}L(t; \beta)}{\partial \beta^2}\right)} \propto \frac{1}{\beta}, \beta > 0. \quad (3.3.1)$$

2. The inverted gamma:

The PLP and inverted gamma probability distributions belong to the exponential family of probability distributions, which makes the latter a logical choice for an informative parametric prior for β . The inverted gamma probability distribution is given by:

$$g_{IG}(\beta) \propto \left(\frac{\mu}{\beta}\right)^{v+1} \frac{1}{\mu \Gamma(v)} \exp\left\{-\frac{\mu}{\beta}\right\}, \beta > 0, \mu > 0, v > 0, \quad (3.3.2)$$

where v and μ are the shape and scale parameters.

3. Kernel' prior:

The kernel probability density estimation is a non-parametric method to approximately estimate the PDF of β using a finite data set. It is given by:

$$g_K(\beta) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{\beta - \beta_i}{h}\right), \quad (3.3.3)$$

where K is the kernel function and h is a positive number called the bandwidth.

3.3.1 The Jeffreys' Prior:

Assuming Jeffreys' PDF, Eq. (3.3.1), as the prior of β and using the likelihood function (3.1.5), the posterior density of β is given by:

$$h_J(\bar{t}|\beta) = \frac{\exp\left\{\left(\frac{t_n}{\theta}\right)^\beta\right\} \frac{\beta^{n-1}}{\theta^{n\beta}} \prod_{i=1}^n (t_i)^{\beta-1}}{\int_0^\infty \exp\left\{\left(\frac{t_n}{\theta}\right)^\beta\right\} \frac{\beta^{n-1}}{\theta^{n\beta}} \prod_{i=1}^n (t_i)^{\beta-1} d\beta}. \quad (3.3.1.1)$$

Thus, the Jeffreys' Bayesian estimate of β in $V(t; \beta, \theta)$ under the H-T loss function, using Eq. (3.2.3), is given by:

$$\hat{\beta}_{J.HT} = \frac{1}{f_1 + f_2} \ln\left[\frac{\int_\gamma^\infty \exp\{f_1\beta\} h_J(\bar{t}|\beta) d\beta}{\int_\gamma^\infty \exp\{-f_2\beta\} h_J(\bar{t}|\beta) d\beta}\right]. \quad (3.3.1.2)$$

We cannot obtain a closed analytical form of the Bayesian estimate, $\hat{\beta}_{J.HT}$, thus we must utilize numerical method to obtain the subject estimate. Also note that the estimate depends on knowing or being able to estimate the scale parameter θ .

3.3.2 The Inverted Gamma Prior:

We proceed with our study with the prior probability density of β given by the inverted gamma distribution Eq. (3.3.2). Using the likelihood Eq. (3.1.5), the posterior density of β is given by:

$$h_{IG}(t|\beta) = \frac{\frac{\beta^{n-v-1}}{\theta^{n\beta}} \exp\left\{-\left(\frac{t_n}{\theta}\right)^\beta - \frac{\mu}{\beta}\right\} \prod_{i=1}^n (t_i)^{\beta-1}}{\int_0^\infty \frac{\beta^{n-v-1}}{\theta^{n\beta}} \exp\left\{-\left(\frac{t_n}{\theta}\right)^\beta - \frac{\mu}{\beta}\right\} \prod_{i=1}^n (t_i)^{\beta-1} d\beta}. \quad (3.3.2.1)$$

Thus, the Bayesian estimate of β under the inverted gamma distribution with respect to the H-T loss function, using Eq. (3.2.3) and Eq. (3.3.2.1), is given by:

$$\hat{\beta}_{IG.HT} = \frac{1}{f_1 + f_2} \ln\left[\frac{\int_\gamma^\infty \exp\{f_1\beta\} h_{IG}(t|\beta) d\beta}{\int_\gamma^\infty \exp\{-f_2\beta\} h_{IG}(t|\beta) d\beta}\right]. \quad (3.3.2.2)$$

Here as well, we must rely on a numerical estimation of $\hat{\beta}_{IG.HT}$ because we cannot obtain a closed form of the above equation. Also note that the estimate depends on knowing or being able to estimate the scale parameter θ .

3.3.3 The Kernel' Prior:

Here, we shall assume the non-parametric kernel probability density Eq. (3.3.3) as the prior PDF of β ; using the likelihood Eq. (3.1.5), the posterior density of β is given by:

$$h_k(\bar{t}|\beta) = \frac{\exp\left\{\left(\frac{t_n}{\theta}\right)^\beta\right\} \frac{\beta^n}{\theta^{n\beta}} \prod_{i=1}^n (t_i)^{\beta-1} \frac{1}{nh} \sum_{i=1}^n K\left(\frac{\beta-\beta_i}{h}\right)}{\int_0^\infty \exp\left\{\left(\frac{t_n}{\theta}\right)^\beta\right\} \frac{\beta^n}{\theta^{n\beta}} \prod_{i=1}^n (t_i)^{\beta-1} \frac{1}{nh} \sum_{i=1}^n K\left(\frac{\beta-\beta_i}{h}\right) d\beta}. \quad (3.3.3.1)$$

Thus, the kernel Bayesian estimate of the key parameter β in $V(t; \beta, \theta)$ under the H-T loss function, using Eq. (3.2.3) and Eq. (3.3.1.1), is given by:

$$\hat{\beta}_{K,HT} = \frac{1}{f_1 + f_2} \ln \left[\frac{\int_\gamma^\infty \exp\{f_1\beta\} h_k(\bar{t}|\beta) d\beta}{\int_\gamma^\infty \exp\{-f_2\beta\} h_k(\bar{t}|\beta) d\beta} \right]. \quad (3.3.3.2)$$

We must rely on a numerical estimation because we cannot obtain a closed form solution for $\hat{\beta}_{K,HT}$. In addition, the kernel function, $K(u)$, and bandwidth, h , will be chosen to minimize the asymptotic mean integrated squared error (AMISE) given by:

$$AMISE(\hat{f}(\beta)) = \int E \left[\left(\hat{f}(\beta) - f(\beta) \right)^2 \right] d\beta, \quad (3.3.3.3)$$

where $\hat{f}(\beta)$ and $f(\beta)$ are the estimated probability density of β and the true probability density of β respectively. Below are details of the analysis we conducted using Monte Carlo simulation to generate data governed by a PLP, followed by using actual data.


4 Results and Discussion

4.1 Numerical Simulation

A Monte Carlo simulation was used to compare the Bayesian (under H-T loss functions) and the MLE approaches. The parameter β of the intensity function for the PLP was calculated using numerical integration techniques in conjunction with a Monte Carlo simulation to obtain its Bayesian estimate. Substituting these estimates in the intensity function, we obtained the Bayesian intensity function estimates, from which the reliability function can be estimated.

For a given value of the parameter θ , a stochastic value for the parameter β was generated from the Burr PDF. For each pair of values of θ and β , 400 samples of 40 failure times that followed a PLP

were generated. This procedure was repeated 200 times for three distinct values of θ . The procedure is summarized in the algorithm (Algorithm 1) given below.



Algorithm 1. Simulation to analyze Bayesian estimates of β for a given θ .

For each sample of size 40, the Bayesian estimates and MLEs of the parameter were calculated when $\theta \in \{0.5, 1.7441, 4\}$. The comparison is based on the mean squared error (MSE) averaged over the 100,000 repetitions. The results are given in Table 2.

Table 2: MSE for Bayesian estimates under the H-T loss function and MLE of β , for each assumed θ value.

θ	MSE of $\hat{\beta}$	MSE of $\hat{\beta}_{B,HT}$
0.5	0.0112436	0.000507356
1.7441	0.0110573	0.000516057
4	0.010961	0.000518632

It is observed that $\hat{\beta}_{B,HT}$ maintains a good accuracy, and is superior to $\hat{\beta}$ in estimating β for the different values of θ . For various sample sizes, the Bayesian estimate under the H-T loss function and the MLE of the parameter β were calculated and averaged over 10,000 repetitions. Table 3 displays the simulated result of comparing a true value of β with respect to its MLE and Bayesian estimates for $n = 20, 30, \dots, 160$.

Again, the Bayesian estimate is uniformly closer to the true value of β than its MLE, even for a very small sample size of $n = 20$. A graphical comparison of the true value of β along with the

Table 3: Bayesian estimates, under H-T loss function, and MLEs for the parameter $\beta = 0.7054$ averaged over 10,000 repetitions

n	β_{Fixed}	$\hat{\beta}$	$\hat{\beta}_{B,HT}$
20	0.7054	0.784026	0.675263
30	0.7054	0.756617	0.690189
40	0.7054	0.743982	0.696467
50	0.7054	0.73531	0.699158
60	0.7054	0.729563	0.700642
70	0.7054	0.725977	0.70169
80	0.7054	0.723338	0.702382
100	0.7054	0.719117	0.703165
120	0.7054	0.716315	0.703585
140	0.7054	0.714821	0.70398
160	0.7054	0.713641	0.704244

Bayesian and MLE estimates as functions of sample size is given by Figure 2.




Figure 2: β estimates versus sample size.

Figure 2 shows the MLE of β tends to overestimate whereas the Bayesian estimate tends to underestimate the true value of β , particularly when considering small sample sizes. The MSEs of the MLE and Bayesian estimates of β is given below by Figure 3.

Regardless of sample size, the MSE of the Bayesian estimate of the key parameter β is significantly smaller than the MSE of the MLE of β (Figure 3).

Since the Bayesian estimate under the H-T loss function for β is better than its MLE, Molinares and Tsokos proposed to adjust the MLE of the parameter θ using (3.1.6) with a Bayesian estimate of




Figure 3: β estimates versus sample size.

β instead of its MLE, both of which are needed to estimate the $V(t; \beta, \theta)$, as given below:

$$\hat{\theta}_{B.HT} = \frac{t_n}{n^{1/\hat{\beta}_{B.HT}}}. \tag{4.1.1}$$

For various sample sizes and the same β ($\beta = 0.7054$), the Bayesian MLE and MLE of the parameter θ and their corresponding MSEs were computed, averaging over the 10,000 repetitions, using the MLE of θ ($\theta = 1.7441$) of the Crow data.

Table 4: MLE and Bayesian estimates under the H-T loss function for the parameter $\theta = 1.7441$ averaged over 10,000 repetitions.

n	θ	$\hat{\theta}_{MLE}$	$\hat{\theta}_{B.HT}$
20	1.7441	3.17139	1.36422
30	1.7441	2.908	1.5097
40	1.7441	2.73107	1.58115
50	1.7441	2.59245	1.61985
60	1.7441	2.48865	1.64406
70	1.7441	2.41782	1.66084
80	1.7441	2.36522	1.67294
100	1.7441	2.26774	1.68902
120	1.7441	2.20117	1.69923
140	1.7441	2.15539	1.70659
160	1.7441	2.11872	1.71193

Table 4 shows the inferior performance for the MLE of θ and the slow convergence of its average values to $\theta = 1.7441$, whereas the adjusted estimate of θ ($\hat{\theta}_{B.HT}$) using the Bayesian estimate of β under the H-T loss function performed better in estimating the true value of θ . The MLE of and the Bayesian MLE estimate of θ had a tendency to overestimate and underestimate the parameter θ ,

respectively.

As expected, based on the Bayesian influence on β , $\hat{\theta}_{B.HT}$ is a better estimate than the MLE of θ ($\hat{\theta}$). This can be seen in Figure 4 where the MSEs of θ estimates were plotted against various sample sizes, which demonstrates the excellent performance of $\hat{\theta}_{B.HT}$.




Figure 4: MSE of θ : Bayesian and MLE estimates versus sample size.

We also computed the proposed estimate for the parameter θ ($\hat{\theta}_{B.HT}$) and its MSE over 100,000 repetitions for different values of θ (0.5, 1.7441, 4) and sample size $n = 40$. The results are given by Table 5. The θ values (including 1.7441) were selected for this simulation are smaller and larger than the MLE of θ of the Crow data.

Table 5 below shows that the $\hat{\theta}_{B.HT}$ performed well for the selected θ values. This is particularly true for the small and medium value of θ values.

Table 5: MSE of θ estimates: Bayesian under the H-T loss function, and MLE of β .

θ	$\hat{\theta}_{B.HT}$	MSE of $\hat{\theta}_{B.HT}$
0.5	0.503314	0.00691164
1.7441	1.7509	0.0827802
4	4.01025	0.439035

For a fixed value of $\theta = 1.7441$ and a sample size similar to the size of the collected data, $n = 40$, the estimates of the intensity function $\hat{V}_{MLE}(t)$ and $\hat{V}_{B.HT}(t)$ were obtained using $\hat{\beta}$ and $\hat{\beta}_{B.HT}$, respectively, in Eq. (2.2). That is,

$$\hat{V}'_{MLE}(t) = \frac{\hat{\beta}}{\theta} \left(\frac{t}{\theta} \right)^{\hat{\beta}-1}, \theta > 0, t > 0. \tag{4.1.2}$$




Figure 5: Graph for $\theta = 1.7441$ and the corresponding β Bayesian estimate and MLE's used in \hat{V}'_{MLE} and $\hat{V}'_{B,HT}$, estimates of $V(t; \beta, \theta)$ with $n = 40$.

and

$$\hat{V}'_{B,HT}(t) = \frac{\hat{\beta}_{B,HT}}{\theta} \left(\frac{t}{\theta} \right)^{\hat{\beta}_{B,HT}-1}, \quad \theta > 0, t > 0. \quad (4.1.3)$$

Their graphs (Figure 5) reveal the superior performance of $\hat{V}'_{B,HT}(t)$.

In order to obtain Bayesian estimates of the intensity function, $\hat{V}^*_{B,SE}$ and $\hat{V}^*_{B,HT}$, we substituted the Bayesian estimates of β and its corresponding θ MLE in Eq. (2.2). That is,

$$\hat{V}^*_{B,HT}(t) = \frac{\hat{\beta}_{B,HT}}{\hat{\theta}} \left(\frac{t}{\hat{\theta}} \right)^{\hat{\beta}_{B,HT}-1}, \quad t > 0. \quad (4.1.4)$$

The MLE of the intensity function, \hat{V}_{MLE} , is obtained using the MLEs of β and θ . That is,

$$\hat{V}_{MLE}(t) = \frac{\hat{\beta}}{\hat{\theta}} \left(\frac{t}{\hat{\theta}} \right)^{\hat{\beta}-1}, \quad t > 0. \quad (4.1.5)$$

The Bayesian MLE of the intensity function under the influence of the Bayesian estimate of β , denoted by $\hat{V}_{B,HT}$, is obtained by substituting $\hat{\beta}_{B,HT}$ and $\hat{\theta}_{B,HT}$ in Eq. (2.2):

$$\hat{V}_{B,HT}(t) = \frac{\hat{\beta}_{B,HT}}{\hat{\theta}_{B,HT}} \left(\frac{t}{\hat{\theta}_{B,HT}} \right)^{\hat{\beta}_{B,HT}-1}, \quad t > 0. \quad (4.1.6)$$

To measure the robustness of $\hat{V}_{B,HT}$ with respect to \hat{V}_{MLE} , we calculated the relative efficiency

(RE) of the estimate $\hat{V}_{B.HT}$ compared to the estimate \hat{V}_{MLE} , which is defined as

$$RE(\hat{V}_{B.HT}, \hat{V}_{MLE}) = \frac{IMSE(\hat{V}_{B.HT})}{IMSE(\hat{V}_{MLE})} = \frac{\int_{-\infty}^{\infty} [\hat{V}_{B.HT}(t) - V(t)]^2 dt}{\int_{-\infty}^{\infty} [\hat{V}_{MLE}(t) - V(t)]^2 dt}. \quad (4.1.7)$$

If $RE = 1$, $\hat{V}_{B.HT}$ and \hat{V}_{MLE} will be interpreted as equally effective. If $RE < 1$, $\hat{V}_{B.HT}$ is more efficient than \hat{V}_{MLE} , contrary to $RE > 1$, in which case $\hat{V}_{B.HT}$ is less efficient than \hat{V}_{MLE} .

Bayesian estimates and MLEs for the parameters $\beta = 0.7054$ and $\theta = 1.7441$ (Table 6), averaged over 10,000 repetitions, were used, for $n = 40$, to compare $\hat{V}_{B.TH}$ and \hat{V}_{MLE} using Eq. (4.1.7).

Table 6: Averages of the Bayesian (under the H-T loss function) and MLE estimates of β and θ

β	$\hat{\beta}$	$\hat{\beta}_{B.HT}$	θ	$\hat{\theta}$	$\hat{\theta}_{B.HT}$
0.7054	0.743982	0.696467	1.7441	2.73107	1.58115

Table 6 above reveals that the averages of the Bayesian and Bayesian MLE estimates of the parameters β and θ under the H-T loss function, respectively, are closer to the true values than their corresponding MLE estimates. The results of the comparison among the $\hat{V}_{B.TH}$ and \hat{V}_{MLE} using Eq. (4.1.7) are given in Tables 7 and 8.

The analytical forms of the $V(t)$, \hat{V}_{MLE} , $\hat{V}^*_{B.TH}$, and $\hat{V}_{B.TH}$ (Table 7) were derived by substituting the initialized values, MLE estimates of both parameters, Bayesian estimate β and MLE of θ , and Bayesian estimates, respectively.

Table 8 shows the comparison result of $\hat{V}_{B.HT}$ and \hat{V}_{MLE} , where the $RE(\hat{V}_{B.HT}, \hat{V}_{MLE})$ is less than 1, which implies that the intensity function using $\hat{\beta}_{B.HT}$ is more efficient than the intensity function under $\hat{\beta}_{MLE}$, establishing the superior relative efficiency of Bayesian estimates under the H-T loss function over MLE estimates. The corresponding graph for the intensity functions is given by Figure 6. In addition, $\hat{V}^*_{B.HT}$ computed using a Bayesian estimate for β and MLE estimate for θ , is less efficient compared to \hat{V}_{MLE} , and $\hat{V}_{B.HT}$.

The $\hat{V}_{B.HT}$ is a better estimate of $V(t)$, compared to the \hat{V}_{MSE} and $\hat{V}^*_{B.HT}$. Based on the results of this section, the Bayesian estimates under the H-T loss function will be used to analyze real data in the following section.

Table 7: Intensity functions with Bayesian and MLE estimates for β and θ

$V(t)$	\hat{V}_{MLE}	$\hat{V}^*_{B.HT}$	$\hat{V}_{B.HT}$
$0.476465 \cdot t^{-0.2946}$	$0.352321 \cdot t^{-0.256018}$	$0.345946 \cdot t^{-0.303533}$	$0.5062 \cdot t^{-0.303533}$

Table 8: Relative efficiency of $\hat{V}_{B.HT}$ compared to \hat{V}_{MLE} .

$RE(\hat{V}_{B.HT}, \hat{V}_{MLE})$	$RE(\hat{V}_{B.HT}, \hat{V}^*_{B.HT})$
0.0761919	0.00550275




Figure 6: Estimates of the intensity function using values in Table 6, $n = 40$.

4.2 Using Real Data

Using the software reliability growth data from Table 1, we computed $\hat{\beta}_{B.HT}$ and the adjusted estimate of θ ($\hat{\theta}_{B.HT}$) in order to obtain a Bayesian estimate of the intensity function under the H-T loss function. We followed the algorithm given below (Algorithm 2) to obtain the Bayesian intensity function for the given real data.


For the failure data of Crow, provided in Table 1, $\hat{\beta}_{B.HT}$ is 0.501199 and $\hat{\theta}_{B.HT}$ is 2.07144. Therefore, with the use of $\hat{\theta}_{B.HT}$, the Bayesian MLE of the intensity function for the data is given by:

$$\hat{V}_{B.HT}(t) = 0.347933 \cdot t^{-0.498801}, t > 0. \tag{4.2.1}$$

A graphical display of $\hat{V}_{B.HT}(t)$ is given below.

Figure 8 shows the Bayesian MLE estimate of the intensity function ($V(t; \beta, \theta)$) under the H-T loss function ($\hat{V}^*_{B.HT}$), which indicates the improvement of the software reliability over time.

To obtain a Bayesian MLE for the reliability function under the H-T loss function, we use this Bayesian estimate for the intensity function. The analytical form for the corresponding Bayesian



Algorithm 2. Estimate of the intensity function using Crow data in Table 1

reliability estimate, based on the real data, is given by:

$$\hat{R}_{B,HT}(t_i | t_1, \dots, t_{i-1}) = \exp \left\{ -0.347933 \int_{t_{i-1}}^{t_i} x^{-0.498801} dx \right\}, \quad t_i > t_{i-1} > 0. \quad (4.2.2)$$

Thus far, we demonstrated not only the applicability of the Bayesian analysis to the PLP, but also, using real data, the superiority of its performance and influence compared to the MLE of the parameters β and θ , respectively, assuming the Burr PDF is the prior knowledge of the key parameter β . Next section, we study the sensitivity of the prior selections, in which an engineer might lack a prior knowledge of parameter β .

4.3 Sensitivity of Prior Selection

In the implementation of the simulation procedure we followed Algorithm 1. Random failure times (time to failures) distributed according to the PLP are simulated for a realization of the stochastic scale parameter β , which follows a Burr type XII probability distribution. Informative parametric priors were considered, such as inverted gamma and Burr probability distributions, whereas the Jefferys prior was




Figure 8: Estimate of the intensity function for the real data in Table 1, using $\hat{\beta}_{B,HT}$ and $\hat{\theta}_{B,HT}$.

chosen as a non-informative prior. In addition, non-parametric priors like kernel density were applied during the sensitivity analysis study. Kernel density estimation depends on several variables, including sample size, bandwidth, and kernel function. In this study, the optimal bandwidth (h^*) and kernel function were chosen such that the asymptotic mean integrated squared error (AMISE) is minimized. The simplified analytical form of AMISE, Eq. (3.3.3.3), is given by:

$$AMISE\left(\hat{f}(\beta)\right) = \frac{C(K)}{n \cdot h} + \left(\frac{1}{4} \cdot h^4 \cdot k_2^2 \cdot R\left(f^{(2)}(\beta)\right)\right) \quad (4.3.1)$$

Where:

- $C(K) = \int (K(u))^2 du$.
- n : sample size.
- h : bandwidth.
- $k_2 = \int_{-\infty}^{+\infty} u^2 \cdot K(u) du$.
- $f^{(2)}(\beta)$ is the second derivative of Burr PDF.
- $R(f^{(2)}(\beta)) = \int (f^{(2)}(\beta))^2 d\beta$.

$$\bullet h^* = \left[\frac{C(K)}{k_2^2 \cdot R(f^{(2)}(\beta))} \right]^{1/5} \cdot n^{-1/5}.$$

The minimum AMISE corresponds to the Epanechnikov kernel function ($K(u) = \frac{3}{4}(1-u^2)I_{|u| \leq 1}$), [6]. In addition to the Epanechnikov kernel function, the Gaussian kernel function ($K(u) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{u^2}{2}\right) I_{\mathbb{R}}$) was also used in the calculation since it is commonly used for its analytical tractability.

Numerical integration techniques were used to compute the Bayesian estimates of β under the H-T loss function, according to the equations in Section 3.3, for each of the five densities and three distinct values of θ . Samples of sizes of 20, 30, 40, 50, 60, 70, 80, 100, 120, 140, and 160 were generated, where the parameter θ was assumed to be the MLE of θ (1.7441) using Crow’s data. The results, for 10,000 repetitions, are presented by Table 9 and Figure 9.

Table 9: Averages of the Bayesian estimates (using the subject prior PDFs), under H-T loss functions, and MLEs of the parameter β over 10,000 repetitions

n	β	$\hat{\beta}_{MLE}$	$\hat{\beta}_{B.HT}$	$\hat{\beta}_{IG.HT}$	$\hat{\beta}_{J.HT}$	$\hat{\beta}_{KG.HT}$	$\hat{\beta}_{KE.HT}$
20	0.7054	0.781303	0.674095	0.799535	0.710047	0.675698	0.675693
30	0.7054	0.753465	0.688951	0.762536	0.707341	0.693849	0.693881
40	0.7054	0.740665	0.695087	0.745074	0.70651	0.698597	0.698631
50	0.7054	0.732738	0.697886	0.734888	0.705885	0.699751	0.69976
60	0.7054	0.728669	0.699823	0.728731	0.705847	0.700727	0.700699
70	0.7054	0.725499	0.70101	0.724471	0.705776	0.70148	0.701405
80	0.7054	0.723539	0.701979	0.721552	0.705882	0.702323	0.7022
100	0.7054	0.719621	0.70285	0.717324	0.705664	0.70338	0.703159
120	0.7054	0.717465	0.703468	0.71478	0.705632	0.704258	0.703959
140	0.7054	0.716224	0.703954	0.713141	0.705692	0.704928	0.70457
160	0.7054	0.714739	0.704188	0.711865	0.705629	0.7053	0.704893

It can be observed that the Bayesian estimate of β for the Jeffreys prior PDF under the H-T loss function converges to the true value faster than other prior PDFs; followed very closely by the Bayesian estimates using the kernel prior PDFs, which tend to converge faster than the inverted gamma prior PDF and are superior to the maximum likelihood approach.

Figure 9 presents the graphical behavior of the MLE and Bayesian estimates represented by their average values, where the average values of the Jeffery Bayesian estimates were the closest to the true value of β for all sample sizes. Followed by the average values of the kernel Bayesian and the Burr Bayesian estimates, which they tend to have similar behavior in estimating for sample sizes of 50 and larger. Bayesian estimates seems to have insignificant difference for sample sizes of 100 and larger except the inverted gamma Bayesian estimate which performed as poor as the MLE in estimating the key parameter β .

Table 10 shows the MSE of the Bayesian estimates of β under the H-T loss function and the subject




Figure 9: Averages of β estimates for different sample sizes.

prior PDFs with respect to various sample sizes.

Table 10: MSE of the Bayesian estimates, under the H-T loss functions, and MLEs of the parameter β averaged over 1000 repetitions for different priors.

n	MSE of $\hat{\beta}_{MLE}$	MSE of $\hat{\beta}_{B,HT}$	MSE of $\hat{\beta}_{IG,HT}$	MSE of $\hat{\beta}_{J,HT}$	MSE of $\hat{\beta}_{KG,HT}$	MSE of $\hat{\beta}_{KE,HT}$
20	0.0417095	0.00362378	0.0115968	0.00285493	0.0037072	0.00371019
30	0.0238951	0.00167959	0.00460464	0.00139859	0.00166486	0.00166402
40	0.016305	0.00100242	0.00242657	0.000885863	0.000976117	0.000974269
50	0.0123248	0.00067513	0.00146266	0.000612733	0.000653313	0.000653343
60	0.00997557	0.000502602	0.000998874	0.000467943	0.000476869	0.000478856
70	0.00831476	0.000394076	0.000726943	0.000372566	0.00036205	0.000365206
80	0.00701888	0.000320076	0.000561229	0.000307048	0.000284152	0.000288374
100	0.00546558	0.000227799	0.000358957	0.000220557	0.000194561	0.000199899
120	0.00452469	0.000177734	0.000259043	0.000173553	0.000149927	0.000156087
140	0.00386396	0.000145074	0.00020085	0.000142729	0.000121572	0.000128407
160	0.00329065	0.000120261	0.00015916	0.000118617	0.00010159	0.000108463

For a sample size of 20, the Jeffery Bayesian estimate of β is the best estimate based on the MSE, followed by the Burr and kernel Bayesian estimates (Table 10). The MSE of the Gaussian Kernel Bayesian estimate was the lowest for moderate to large sample sizes ($n = 70, \dots, 160$). The inverted gamma Bayesian estimate had the lowest performance compared to other Bayesian estimates. The MSE of the MLE of the key parameter β indicated its poor performance compared to the Bayesian estimates.

Figure 10 shows the MLE and Bayesian estimates of β . It can be observed the Bayesian estimates under the H-T loss function are superior to the MLE of the key parameter β for all sample sizes considered in this study. Moreover, the Bayesian estimates are presented without MLE in Figure 11 to look closely at the performance of Bayesian estimates, under the H-T loss function and the subject




Figure 10: MSEs of β estimates for different sample sizes.

prior PDFs, based on their MSEs.




Figure 11: MSEs of β Bayesian estimates for different sample sizes.

The MSEs of the Jeffreys prior, for various sample sizes, were the smallest, followed by MSEs of the kernels and Burr prior PDFs. Bayesian estimates using Gaussian and Epanechnikov kernel probability densities as priors performed similarly, whereas the Bayesian estimate using the inverted gamma PDF had the lowest performance compared to other Bayesian estimates. It also shows that for a sample size of $n = 30$ and larger, the Bayesian estimates using Burr, Jeffery, and kernel PDFs are similar in their convergence to the true value. For a sample size of $n = 70$ and larger, all Bayesian estimates tend to converge to the true value of the key parameter β . The Bayesian estimates of β under Jeffreys

and inverted gamma prior PDFs tend to overestimate β , whereas Burr and kernel prior PDFs tend to underestimate β .

For each sample of size 40 based on Monte Carlo simulation, the Bayesian estimates and MLEs of β were calculated when $\theta \in \{0.5, 1.7441, 4\}$. The comparison is based on the MSE averaged over the 2,000 simulated samples. The results are given by Table 11.

Table 11: MSE of β Bayesian estimates with Burr, Jeffreys, inverted gamma, and kernel PDFs as priors under the H-T loss function. MSE of MLE estimate of the parameter β in an NHPP with samples of $n = 40$ and different values of the parameter θ .

θ	MSE of $\hat{\beta}_{MLE}$	MSE of $\hat{\beta}_{B.HT}$	MSE of $\hat{\beta}_{IG.HT}$	MSE of $\hat{\beta}_{J.HT}$	MSE of $\hat{\beta}_{KG.HT}$	MSE of $\hat{\beta}_{KE.HT}$
0.5	0.0106298	0.000520244	0.00229073	0.000577579	0.000535304	0.000534979
1.7441	0.00996687	0.000524716	0.00225675	0.000585273	0.000516497	0.000516599
4	0.0112937	0.000558022	0.00246984	0.000632173	0.000556695	0.000556804

It can be observed that all Bayesian estimates are superior to MLE ($\hat{\beta}$) in estimating β , with sample size $n = 40$, while maintaining a consistent behavior for the different values of θ (Table 11). For small value of θ , Jeffrey Bayesian estimate had the lowest MSE value, whereas the MSE of the Gaussian kernel Bayesian estimate was the lowest for moderate and large values of θ .

For the case in which we misleadingly assumed the true probability distribution of the key parameter β , we found that the Jeffreys and kernel Bayesian estimates of β had the best performance compared to the inverted gamma Bayesian estimate of β , indicating that the Bayesian estimate of β is sensitive to the choice of its prior PDF.

As expected, the adjusted MLE of θ produced a better estimate under the mentioned Bayesian influence with respect to the H-T loss function. The average values of the MLE and Bayesian estimates of θ using the MLE and Bayesian estimates of β for various sample sizes, respectively, are presented by Figure 12. where the average values of the Jeffery Bayesian estimates were the closest to the true value of θ for all sample sizes. Followed by the average values of the kernel Bayesian and the Burr Bayesian estimates, which tend to have similar behavior in estimating for sample sizes of 40 and larger. When considering sample sizes of 100 and larger, the Bayesian estimates seems to have insignificant difference except for the inverted gamma Bayesian estimate which which still performs better than the MLE in estimating the parameter θ .

The MSE of θ estimates using MLE and Bayesian estimates of β , namely $\hat{\theta}_{MLE}$, $\hat{\theta}_{B.HT}$, $\hat{\theta}_{IG.HT}$, $\hat{\theta}_{J.HT}$, $\hat{\theta}_{KG.HT}$, and $\hat{\theta}_{KE.HT}$, are shown in Table 12 with various sample sizes. For a small sample, moderate, and large sample sizes, $n = 20, 70, 160$ respectively, the Jeffrey Bayesian estimate of θ ($\hat{\theta}_{J.HT}$) performed better than the other estimates, followed by the kernel Bayesian and Burr Bayesian estimates.




Figure 12: Averages of MLEs of θ using the Bayesian estimates of the parameter β with respect to different priors.

Table 12: MSE of the Bayesian estimates, under the H-T loss functions, and MLEs of the parameter θ averaged over 1000 repetitions for different priors.

n	MSE of $\hat{\theta}_{MLE}$	MSE of $\hat{\theta}_{B,HT}$	MSE of $\hat{\theta}_{IG,HT}$	MSE of $\hat{\theta}_{J,HT}$	MSE of $\hat{\theta}_{KG,HT}$	MSE of $\hat{\theta}_{KE,HT}$
20	11.5735	0.175991	1.0789	0.0310651	0.174974	0.175081
30	8.6721	0.0726953	0.50118	0.0166961	0.0538952	0.0537907
40	6.67926	0.0378416	0.27733	0.0107884	0.0264298	0.026306
50	5.45282	0.0233418	0.171955	0.00777231	0.0182648	0.018213
60	4.55577	0.015811	0.115456	0.00593216	0.0139622	0.0139218
70	3.92543	0.0114843	0.0820851	0.0047144	0.0110305	0.0109659
80	3.38362	0.00867134	0.0610179	0.00382098	0.00880575	0.00869148
100	2.73005	0.00566115	0.0369537	0.0027899	0.00591856	0.00572456
120	2.33339	0.00402454	0.0245551	0.00215943	0.004114	0.00387668
140	2.03741	0.00302931	0.0174259	0.00174116	0.003041	0.00277349
160	1.75235	0.00238185	0.0129141	0.0014346	0.0023649	0.00206666

While the inverted gamma Bayesian estimate of θ outperformed that of the MLE, it did not perform as well as the other Bayesian estimates based on their MSE values across all sample size. This indicates the MLE of θ had the poorest performance compared to the Bayesian estimates.

The MSEs of θ estimates using the MLE and Bayesian estimates of the parameter β with respect to different priors is displayed in Figure 13, from which it can be seen that the MLE of θ was extremely weak estimator since it has the largest MSEs across the sample sizes. The adjusted θ estimates were displayed without the MLE estimates by Figure 14.

It can be noted that the θ estimate using the inverted gamma Bayesian estimate of β had the lowest performance compared to other estimates that used Bayesian estimates of β . In addition, above the sample size $n = 40$, the estimate of θ using Burr kernels Bayesian estimates of β performed similarly in estimating the true value of θ . All θ estimates using Bayesian estimates of β tend to converge to the true value in similar trajectories, whereas the θ estimate using the Jeffrey Bayesian estimate of β




Figure 13: MSE of the MLEs of θ using MLE and Bayesian estimates of β with respect to different prior β




Figure 14: MSE of the MLEs of θ using Bayesian estimates of β with respect to different prior β .

converges slightly faster.

Therefore, the Bayesian analysis to the PLP under the H-T loss function is sensitive to the prior selection. Based on this finding, an engineer, for example, is recommended to use the Jeffreys or kernel PDFs if he/she lacks any prior knowledge of β .

5 Conclusion

In the present study, we developed the analytical Bayesian form of the key parameter β , under the H-T loss function, in the intensity function, where the underlying failure distribution is the PLP that

is used for software reliability assessment, among others. The reliability function of the subject model is written analytically as a function of the intensity function.

The behavior of β is characterized by the Burr type XII probability distribution. Real data and numerical simulation were used to illustrate the efficiency improvement in the estimation of the intensity function of PLP under the H-T loss function ($\hat{V}_{B,HT}(t)$). Based on the 100,000 samples of software failure times, using Monte Carlo simulations and sample size of 40, we found that the Bayesian estimate of β under the H-T loss function ($\hat{\beta}_{B,HT}$) performed better than the MLE of β with respect to three different values of θ (0.5, 1.7441, 4). Even for different sample sizes (20, 30, 40, 50, 60, 70, 80, 100, 120, 140, and 160), similar results were achieved using $\beta=0.7054$, $\theta=1.7441$, and averaged over 10,000 samples of software failure times.

Because the MLE of the second parameter (θ) in the intensity function depends on the estimate of β , the adjusted estimate of θ $\hat{\beta}_{B,HT}$, as expected, performed better than MLE of θ . Moreover, by comparing the relative efficiency metric, mainly using MLEs for both β and θ ($\hat{V}_{MLE}(t)$), using Bayesian estimates of β under the H-T loss function, and Bayesian MLE of θ ($\hat{V}_{B,HT}(t)$), we found that $\hat{V}_{B,HT}(t)$ is more efficient in estimating the intensity function $V(t; \beta, \theta)$.

In section 4.3, we answered the second research question: Is the Bayesian estimate of the key parameter, β , using the H-T loss function in the PLP, sensitive to the selection of the prior, whether parametric or non-parametric? The parametric priors were Burr, Jefferys, and inverted gamma probability distributions. The non-parametric priors were the Gaussian and Epanechnikov kernel densities. The priors' parameters were estimated using Crow failure times. Additionally, the optimal bandwidth and kernel functions were selected to minimize the asymptotic mean integrated squared error.

Using the proposed algorithm, the Bayesian estimate of β under the H-T loss function and Burr PDF as a prior, $\hat{\beta}_{B,HT}$, performed slightly better than the other Bayesian estimates using different prior PDFs for small value of θ . The Bayesian estimate of β under H-T loss function and Gaussian kernel PDF as a prior, $\hat{\beta}_{KG,HT}$, had the smallest MSE comparing to other prior PDFs for moderate and large values of θ . MLE of β continued the poor performance comparing to Bayesian estimates using the subject prior PDFs. The Bayesian estimate of β under H-T loss function and inverted gamma PDF as a prior performed the lowest among other Bayesian estimates using Burr, Jeffrey, Gaussian, and Epanechnikov PDFs as priors whereas the latter have slightly different MSEs values. Even for different sample sizes, the MLE of β has the poorest performance comparing to Bayesian estimates using different prior PDFs. For small to moderate sample sizes ($n = 20, 30, 40, 60$), the Bayesian estimate of β under H-T loss function and Jeffrey PDF as a prior, $\hat{\beta}_{J,HT}$, has the smallest MSE value,

followed closely by the Bayesian estimates using the Gaussian and Epanechnikov kernels, and Burr PDFs as priors. For moderate to large sample sizes ($n = 70, 80, \dots, 160$), the Bayesian estimate of β under H-T loss function and Gaussian kernel PDF as a prior, $\hat{\beta}_{KG.HT}$, has the smallest MSE value, followed by the Bayesian estimates using the Epanechnikov kernel, Jeffrey, and Burr PDFs as priors.

The Bayesian estimates under H-T loss function and the parametric and non-parametric priors were used to compute the adjusted estimate of θ , where the adjusted θ estimate using the Jeffrey Bayesian estimate of β performed the superior estimate comparing to MLE and other Bayesian estimates for various sample sizes. Followed by Bayesian estimates using Epanechnikov kernel, Burr, and Gaussian kernel PDFs as priors for moderate to large sample sizes whereas using the Gaussian kernel Bayesian estimate of β to compute the adjusted estimate of θ performed slightly better comparing to the usage of Burr Bayesian estimate of β .

Thus, based on this aspect of our analysis, we can conclude that the Bayesian analysis approach under Higgins-Tsokos loss function is superior to the maximum likelihood approach in estimating the reliability function of the Power Law Process. Therefore, the results of this study have the potential to contribute not only to the reliability analysis field but also to other fields that employ the Power Law Process.

References

- [1] J. T. Duane, Learning curve approach to reliability monitoring, *IEEE Trans. Aerospace* **2** (1964) 563–566.
- [2] L. H. Crow, Tracking reliability growth, in *Proc. Twentieth Conf. Design of Experiments*, Report 75-2 (US Army Research Office, Research Triangle Park, North Carolina, 1975), pp. 741–754.
- [3] L. H. Crow, Reliability analysis for complex, repairable systems, in *Reliability and Biometry*, eds. F. Proschan and R. J. Serfling (SIAM, Philadelphia, Pennsylvania, 1974).
- [4] H. E. Ascher and H. Feingold, Repairable Systems Reliability: Modeling, Inference, *Misconceptions and Their Causes* (Marcel-Dekker, New York, 1984).
- [5] C. A. Moltnares and C. P. Tsokos, Bayesian Reliability Approach To The Power Law Process With Sensitive Analysis To Prior Selection, *International Journal of Reliability, Quality and Safety Engineering* (World Scientific, 2013).

- [6] F. N. Alenezi and C. P. Tsokos, The Effectiveness of The Squared Error And Higgins-Tsokos Loss Functions On The Bayesian Reliability Analysis Of Software Failure Times Under The Power Law Process, *Engineering* **11**, 272- 299 (Scientific Research Publishing, 2019).
- [7] J. J. Higgins and C. P. Tsokos, A study of the effect of the loss function on Bayes estimates of failure intensity, MTBF, and Reliability, *Applied Mathematics and Computation* **6** (1980) 145–166.
- [8] Harold Jeffreys, F.R.S., An invariant form for the prior probability in estimation problems, *Royal Society* (1946).
- [9] D. J. Daley and D. Vere-Jones, An introduction to the theory of point processes, *Vol. I. Probability and its Applications* (Springer, New York, second edition, 2003).
- [10] S. Yamada, M. u Ohba and S. Osaki, S-Shaped Reliability Growth Modeling for Software Error Detection, *IEEE Trans. Reliab.* **R-32** (1983) 475–484.
- [11] A. L. Goel K. Okumoto, Time-Dependent Error-Detection Rate Model for Software Reliability and Other Performance Measures, *IEEE Trans. Reliab.* **R-28** (1979) 206–211.
- [12] M. Engelhardt and J. L. Bain, Statistical analysis of a compound power law model for repairable systems, *IEEE Trans. Reliab.* **R-36** (1987) 392–396.
- [13] M. Kimura, T. Toyota and S. Yamada, Economic analysis of software release problems with warranty cost and reliability requirement, *Reliability Engineering and System Safety* **66** (1999) 49–55.
- [14] M. Engelhardt and J. L. Bain, Prediction intervals for the Weibull process, *Technometrics* **20** (1978) 167–169.
- [15] S. E. Rigdon and A. P. Basu, The effect of assuming a homogeneous Poisson process when the true process is a power law process, *J. Qual. Tech.* **22** (1990) 111–117.
- [16] S. E. Rigdon and A. P. Basu, The power law process: A model for the reliability of repairable systems, *J. Qual. Tech.* **21** (1990) 251–260.
- [17] S. E. Rigdon and A. P. Basu, Estimating the intensity function of a power law process at the current time: Time truncated case, *Commun. Stat.-Simulat. Comput.* **19** (1990) 1079–1104.
- [18] H. E. Ascher, T. T. Y. Lin and D. P. Siewiorek, Modification of error log analysis: Statistical modeling and heuristic trend analysis, *IEEE Trans. Reliab.* **41** (1992) 599–601.

- [19] J. Kyparisis and N. D. Singpurwalla, Bayesian inference for the Weibull process with applications to assessing software reliability growth and predicting software failures, in *Computer Science and Statistics: Proc. Sixteenth Symp. Interface* (1985).
- [20] W. M. Bassin, Increasing hazard functions and overhaul policy, in *Proc. 1969 Annual Symp. Reliability*, Chicago, Vol. 8 (IEEE, 1969), pp. 173–178.
- [21] Y. Xu and C. P. Tsokos, Non-homogeneous Poisson process for evaluating stage I and II ductal breast cancer treatment, *Journal of Modern Applied Statistical Methods*, 10 (2), pp. 646–655, (2011)
- [22] M. Evans, N. Hastings and B. Peacock, *Statistical Distributions* (John Wiley and Sons, Inc., New York, 2000).
- [23] C. P. Tsokos, Recent Advances in Life-Testing and Reliability: *A Volume in Honor of Alonso Clifford Cohenm Jr.*, ed. N. Balakrishnan (CRC Press, Boca Ratón, 1995).
- [24] L. J. Bain and M. Engelhardt, *Statistical Analysis of Reliability and Life Testing Models* (Marcel-Dekker, New York, 1991).
- [25] S. K. Bar-Lev, I. Lavi and B. Reiser, Bayesian inference for the power law process, *Ann. Inst. Stat. Math.* **44** (1992) 623–639.
- [26] V. A. Camara and C. P. Tsokos, Bayesian reliability modeling with a new loss function and numerical simulation, *Statistica* **61** (2001) 619–630.
- [27] J. M. Finkelstein, Confidence bounds on the parameters of the Weibull process, *Technometrics* **18** (1978) 115–117.
- [28] M. S. Hamada, A. G. Wilson, C. Shane Reese and H. F. Martz, *Bayesian Reliability* (Springer Science, New York, 2008).
- [29] H. Kim, S. Choi and S. Kim, Bayesian inference for the power law process with the power prior, *J. Korean Statist. Soc.* **34** (2005) 331–344.
- [30] K. Kim, S. W. Kim and H. Kim, Intrinsic Bayes factors for model selection in nonhomogeneous Poisson process, *Far East J. Theor. Stat.* **11(1)** (2003) 15–30.
- [31] S. W. Kim and D. Sun, Intrinsic priors for model selection using an encompassing model with applications to censored failure time data, *Lifetime Data Anal.* **6** (2000) 251–269.

- [32] J. J. Higgins and C. P. Tsokos, A quasi-Bayes estimate of the failure intensity of a reliability-growth model, *IEEE Trans. Reliab.* **R-30** (1981) 471–475.
- [33] L. Lee and S. K. Lee, Some results on inference for the Weibull process, *Technometrics* **20** (1978) 41–45.
- [34] R. T. Lingham and S. Sivaganesan, Testing hypotheses about the power law process under failure truncation using intrinsic Bayes factors, *Ann. Inst. Stat. Math.* **49** (1997) 693–710.
- [35] H. Raiffa and R. Schlaifer, *Applied Statistical Decision Theory* (Harvard Graduate School of Business Administration, Boston, 1961).
- [36] S. K. Bhattacharya, Bayesian approach to life testing and reliability estimation, *J. Amer. Stat. Assoc.* **62** (1967) 48–62.
- [37] W. A. Thompson Jr., On the foundations of reliability, *Technometrics* **23** (1981) 1–23.
- [38] C. P. Tsokos and A. N. V. Rao, Estimation of failure intensity for the Weibull process, *Reliab. Eng. Syst. Safety* **45** (1994) 271–275.