

## GREEDY QUASI-NEWTON METHODS WITH EXPLICIT SUPERLINEAR CONVERGENCE\*

ANTON RODOMANOV<sup>†</sup> AND YURII NESTEROV<sup>‡</sup>

**Abstract.** In this paper, we study greedy variants of quasi-Newton methods. They are based on the updating formulas from a certain subclass of the Broyden family. In particular, this subclass includes the well-known DFP, BFGS and SR1 updates. However, in contrast to the classical quasi-Newton methods, which use the difference of successive iterates for updating the Hessian approximations, our methods apply basis vectors, greedily selected so as to maximize a certain measure of progress. For greedy quasi-Newton methods, we establish an explicit non-asymptotic bound on their rate of local superlinear convergence, as applied to minimizing strongly convex and strongly self-concordant functions (and, in particular, to strongly convex functions with Lipschitz continuous Hessian). The established superlinear convergence rate contains a contraction factor, which depends on the square of the iteration counter. We also show that greedy quasi-Newton methods produce Hessian approximations whose deviation from the exact Hessians linearly converges to zero.

**Key words.** quasi-Newton methods, Broyden family, SR1, DFP, BFGS, superlinear convergence, local convergence, rate of convergence

**AMS subject classifications.** 90C53, 90C30, 68Q25

**DOI.** 10.1137/20M1320651

### 1. Introduction.

**1.1. Motivation.** Quasi-Newton methods have a reputation of the most efficient numerical schemes for smooth unconstrained optimization. The main idea of these algorithms is to approximate the standard Newton method by replacing the exact Hessian with some approximation, which is updated between iterations according to special formulas. There exist numerous variants of quasi-Newton algorithms that differ mainly in the rules of updating Hessian approximations. The three most popular are the *Davidon–Fletcher–Powell (DFP)* method [1, 2], the *Broyden–Fletcher–Goldfarb–Shanno (BFGS)* method [6–10], and the *Symmetric Rank 1 (SR1)* method [1, 3]. For a general overview of the topic, see [14] and [25, Ch. 6]; also see [28] for the application of quasi-Newton methods for non-smooth optimization.

The most attractive feature of quasi-Newton methods is their *superlinear convergence*, which was first established in the 1970s [11–13]. Namely, for several standard quasi-Newton methods (such as DFP and BFGS), it was proved that the ratio of successive residuals tends to zero as the number of iterations goes to infinity. However, the authors did not obtain any explicit bounds on the corresponding *rate* of this superlinear convergence. For example, it is unknown whether the residuals convergence like  $O(c^{k^2})$ , where  $c \in (0, 1)$  is some constant and  $k$  is the iteration counter, or  $O(c^{k^3})$ , or  $O(k^{-k})$ , or somehow else. Thus, despite the qualitative usefulness of the mentioned

---

\*Received by the editors February 20, 2020; accepted for publication (in revised form) November 18, 2020; published electronically March 1, 2021.

<https://doi.org/10.1137/20M1320651>

**Funding:** The research results of this paper were obtained with support of ERC Advanced Grant 788368.

<sup>†</sup>Institute of Information and Communication Technologies, Electronics and Applied Mathematics (ICTEAM), Catholic University of Louvain (UCL), 1348 Louvain-la-Neuve, Belgium ([anton.rodomanov@uclouvain.be](mailto:anton.rodomanov@uclouvain.be)).

<sup>‡</sup>Center for Operations Research and Econometrics (CORE), Catholic University of Louvain (UCL), 1348 Louvain-la-Neuve, Belgium ([yurii.nesterov@uclouvain.be](mailto:yurii.nesterov@uclouvain.be)).

result, it still lacks quantitative estimates of the rate of convergence. Although many other works on quasi-Newton methods have appeared since then, to our knowledge, all of them still contain only asymptotic results (see e.g. [16–19, 21, 23, 24, 26, 31, 34, 35]). Thus, up to now, there are still no *explicit* and *non-asymptotic* estimates of the rate of superlinear convergence of quasi-Newton methods.

In this work, we make a first step towards obtaining such estimates. We propose new quasi-Newton methods, which are based on the updating formulas from a certain subclass of the Broyden family [3]. In particular, this subclass contains the DFP, BFGS and SR1 updates. However, in contrast to the classical quasi-Newton methods, which use the difference of successive iterates for updating the Hessian approximations, our methods apply *basis vectors*, greedily selected to maximize a certain measure of progress. For greedy quasi-Newton methods, we establish an explicit non-asymptotic bound on their rate of local superlinear convergence, which contains a contraction factor, depending on the *square* of the iteration counter. We also show that these methods produce Hessian approximations whose deviation from the exact Hessians converges to zero at a *linear* rate. In contrast, it is known that the standard quasi-Newton methods, in general, cannot ensure the convergence of the Hessian approximations to the true Hessian (see e.g. [13])<sup>1</sup>.

The idea of using basis vectors in quasi-Newton methods goes back at least to so-called *methods of dual directions* [15], for which it is also possible to prove both local superlinear convergence of the iterates and convergence of the Hessian approximations. However, similarly to the standard quasi-Newton methods, all corresponding results are only asymptotic. In any case, despite to the fact that the greedy quasi-Newton methods, presented in this paper, are based on the same idea, their construction and analysis are significantly different.

Finally, let us mention that recently there have been proposed some randomized variants of quasi-Newton algorithms, which also use nonstandard directions for updating Hessian approximations [29, 30, 36].

**1.2. Contents.** In Section 2, we discuss a class of quasi-Newton updating rules for approximating a self-adjoint positive definite linear operator. We present a special greedy strategy for selecting an update direction, which ensures a linear convergence rate in approximating the target operator. In Section 3, we analyze greedy quasi-Newton methods, applied to the problem of minimizing a quadratic function. We show that these methods have a global linear convergence rate, comparable to that of the standard gradient descent, and also a superlinear convergence rate, which contains a contraction factor, depending on the square of the iteration counter. In Section 4, we show that similar results also hold in a more general setting of minimizing a strongly convex and strongly self-concordant function (and, in particular, a strongly convex function with Lipschitz continuous Hessian), provided that the starting point is chosen sufficiently close to the solution. The main difficulty here, compared to the quadratic case, is that the Hessian of the objective function is no longer constant, resulting in the necessity to apply a special *correction strategy* to keep the Hessian approximations under control. Finally, in Section 5, we present some preliminary computational results.

**1.3. Notation.** In what follows,  $\mathbb{E}$  denotes an arbitrary  $n$ -dimensional real vector space. Its dual space, composed of all linear functionals on  $\mathbb{E}$ , is denoted by  $\mathbb{E}^*$ .

---

<sup>1</sup>However, it is worth mentioning that there are some settings, in which the standard SR1 method indeed yields convergence to the true Hessian (for more details, see [20]).

The value of a linear function  $s \in \mathbb{E}^*$ , evaluated at a point  $x \in \mathbb{E}$ , is denoted by  $\langle s, x \rangle$ .

For a smooth function  $f : \mathbb{E} \rightarrow \mathbb{R}$ , we denote by  $\nabla f(x)$  and  $\nabla^2 f(x)$  its gradient and Hessian respectively, evaluated at a point  $x \in \mathbb{E}$ . Note that  $\nabla f(x) \in \mathbb{E}^*$ , and  $\nabla^2 f(x)$  is a self-adjoint linear operator from  $\mathbb{E}$  to  $\mathbb{E}^*$ .

The partial ordering of self-adjoint linear operators is defined in the standard way. We write  $A \preceq A_1$  for  $A, A_1 : \mathbb{E} \rightarrow \mathbb{E}^*$  if  $\langle (A_1 - A)x, x \rangle \geq 0$  for all  $x \in \mathbb{E}$ , and  $W \preceq W_1$  for  $W, W_1 : \mathbb{E}^* \rightarrow \mathbb{E}$  if  $\langle s, (W_1 - W)s \rangle \geq 0$  for all  $s \in \mathbb{E}^*$ .

Any self-adjoint positive definite linear operator  $A : \mathbb{E} \rightarrow \mathbb{E}^*$  induces in the spaces  $\mathbb{E}$  and  $\mathbb{E}^*$  the following pair of conjugate Euclidean norms:

$$(1.1) \quad \begin{aligned} \|h\|_A &\stackrel{\text{def}}{=} \langle Ah, h \rangle^{1/2}, & h \in \mathbb{E}, \\ \|s\|_A^* &\stackrel{\text{def}}{=} \langle s, A^{-1}s \rangle^{1/2}, & s \in \mathbb{E}^*. \end{aligned}$$

When  $A = \nabla^2 f(x)$ , where  $f : \mathbb{E} \rightarrow \mathbb{R}$  is a smooth function with positive definite Hessian, and  $x \in \mathbb{E}$ , we prefer to use notation  $\|\cdot\|_x$  and  $\|\cdot\|_x^*$ , provided that there is no ambiguity with the reference function  $f$ .

Sometimes, in the formulas, involving products of linear operators, it is convenient to treat  $x \in \mathbb{E}$  as a linear operator from  $\mathbb{R}$  to  $\mathbb{E}$ , defined by  $x\alpha = \alpha x$ , and  $x^*$  as a linear operator from  $\mathbb{E}^*$  to  $\mathbb{R}$ , defined by  $x^*s = \langle s, x \rangle$ . In this case,  $xx^*$  is a rank-one self-adjoint linear operator from  $\mathbb{E}^*$  to  $\mathbb{E}$ , acting as follows:

$$(xx^*)s = \langle s, x \rangle x, \quad s \in \mathbb{E}^*.$$

Likewise, any  $s \in \mathbb{E}^*$  can be treated as a linear operator from  $\mathbb{R}$  to  $\mathbb{E}^*$ , defined by  $s\alpha = \alpha s$ , and  $s^*$  as a linear operator from  $\mathbb{E}$  to  $\mathbb{R}$ , defined by  $s^*x = \langle s, x \rangle$ . Then,  $ss^*$  is a rank-one self-adjoint linear operator from  $\mathbb{E}$  to  $\mathbb{E}^*$ .

For two self-adjoint linear operators  $A : \mathbb{E} \rightarrow \mathbb{E}^*$  and  $W : \mathbb{E}^* \rightarrow \mathbb{E}$ , define

$$\langle W, A \rangle \stackrel{\text{def}}{=} \text{Trace}(WA).$$

Note that  $WA$  is a linear operator from  $\mathbb{E}$  to itself, and hence its trace is well-defined (it coincides with the trace of the matrix representation of  $WA$  with respect to an arbitrary chosen basis in the space  $\mathbb{E}$ , and the result is independent of the particular choice of the basis). Observe that  $\langle \cdot, \cdot \rangle$  is a bilinear form, and for any  $x \in \mathbb{E}$ , we have

$$(1.2) \quad \langle Ax, x \rangle = \langle xx^*, A \rangle.$$

When  $A$  is invertible, we also have

$$(1.3) \quad \langle A^{-1}, A \rangle = n.$$

If the operator  $W$  is positive semidefinite, and  $A \preceq A_1$  for some self-adjoint linear operator  $A_1 : \mathbb{E} \rightarrow \mathbb{E}^*$ , then  $\langle W, A \rangle \leq \langle W, A_1 \rangle$ . Similarly, if  $A$  is positive semidefinite and  $W \preceq W_1$  for some self-adjoint linear operator  $W_1 : \mathbb{E}^* \rightarrow \mathbb{E}$ , then  $\langle W, A \rangle \leq \langle W_1, A \rangle$ . When  $A$  is positive definite, and  $R : \mathbb{E} \rightarrow \mathbb{E}^*$  is a self-adjoint linear operator,  $\langle A^{-1}, R \rangle$  equals the sum of the eigenvalues of  $R$  with respect to the operator  $A$ . In particular, if  $R$  is positive semidefinite, then all its eigenvalues with respect to  $A$  are non-negative, and the maximal one can be bounded by the trace:

$$(1.4) \quad R \preceq \langle A^{-1}, R \rangle A.$$

**2. Greedy Quasi-Newton Updates.** Let  $A : \mathbb{E} \rightarrow \mathbb{E}^*$  be a self-adjoint positive definite linear operator. In this section, we consider a class of quasi-Newton updating rules for approximating  $A$ .

Let  $G : \mathbb{E} \rightarrow \mathbb{E}^*$  be a self-adjoint linear operator, such that

$$(2.1) \quad A \preceq G,$$

and let  $u \in \mathbb{E}$  be a direction. Consider the following family of updates, parameterized by a scalar  $\tau \in \mathbb{R}$ . If  $Gu \neq Au$ , define

$$(2.2) \quad \text{Broyd}_\tau(G, A, u) \stackrel{\text{def}}{=} \tau \left[ G - \frac{Auu^*G + Guu^*A}{\langle Au, u \rangle} + \left( \frac{\langle Gu, u \rangle}{\langle Au, u \rangle} + 1 \right) \frac{Auu^*A}{\langle Au, u \rangle} \right] \\ + (1 - \tau) \left[ G - \frac{(G-A)uu^*(G-A)}{\langle (G-A)u, u \rangle} \right].$$

Otherwise, if  $Gu = Au$ , define  $\text{Broyd}_\tau(G, A, u) \stackrel{\text{def}}{=} G$ .

Note that, for  $\tau = 0$ , formula (2.2) corresponds to the well-known *SR1 update*,

$$(2.3) \quad \text{SR1}(G, A, u) \stackrel{\text{def}}{=} G - \frac{(G-A)uu^*(G-A)}{\langle (G-A)u, u \rangle},$$

and, for  $\tau = 1$ , it corresponds to the well-known *DFP update*:

$$(2.4) \quad \text{DFP}(G, A, u) \stackrel{\text{def}}{=} G - \frac{Auu^*G + Guu^*A}{\langle Au, u \rangle} + \left( \frac{\langle Gu, u \rangle}{\langle Au, u \rangle} + 1 \right) \frac{Auu^*A}{\langle Au, u \rangle}.$$

Thus, (2.2) describes the *Broyden family* of quasi-Newton updates (see [25, Section 6.3]), and can be written as the linear combination of DFP and SR1 updates:<sup>2</sup>

$$\text{Broyd}_\tau(G, A, u) = \tau \text{DFP}(G, A, u) + (1 - \tau) \text{SR1}(G, A, u).$$

Our main interest will be in the class, described by the values  $\tau \in [0, 1]$ , i.e. in the convex combination of the DFP and SR1 updates. Note in particular, that this subclass includes another well-known update—*BFGS*. Indeed, for

$$(2.5) \quad \tau_{\text{BFGS}} \stackrel{\text{def}}{=} \frac{\langle Au, u \rangle}{\langle Gu, u \rangle} \stackrel{(2.1)}{\in} (0, 1),$$

we have  $1 - \tau_{\text{BFGS}} = \frac{\langle (G-A)u, u \rangle}{\langle Gu, u \rangle}$ , and thus

$$(2.6) \quad \text{Broyd}_{\tau_{\text{BFGS}}}(G, A, u) = G - \frac{\langle (G-A)u, u \rangle}{\langle Gu, u \rangle} \frac{(G-A)uu^*(G-A)}{\langle (G-A)u, u \rangle} \\ + \frac{\langle Au, u \rangle}{\langle Gu, u \rangle} \left[ -\frac{Auu^*G + Guu^*A}{\langle Au, u \rangle} + \left( \frac{\langle Gu, u \rangle}{\langle Au, u \rangle} + 1 \right) \frac{Auu^*A}{\langle Au, u \rangle} \right] \\ = G - \frac{(G-A)uu^*(G-A)}{\langle Gu, u \rangle} - \frac{Auu^*G + Guu^*A}{\langle Gu, u \rangle} \\ + \left( \frac{\langle Gu, u \rangle}{\langle Au, u \rangle} + 1 \right) \frac{Auu^*A}{\langle Gu, u \rangle} \\ = G - \frac{Guu^*G}{\langle Gu, u \rangle} + \frac{Auu^*A}{\langle Au, u \rangle} \stackrel{\text{def}}{=} \text{BFGS}(G, A, u).$$

This is the classic BFGS formula for direction  $u$ .

Let us show that the Broyden family is monotonic in the parameter  $\tau$ .

<sup>2</sup>Usually, the Broyden family is defined as the linear combination of the DFP and BFGS updates. Here we use alternative (but equivalent) parametrization of this family, which is more convenient for our purposes.

LEMMA 2.1. *If (2.1) holds, then, for any  $u \in \mathbb{E}$ ,  $\tau_1, \tau_2 \in \mathbb{R}$ , such that  $\tau_1 \leq \tau_2$ ,*

$$\text{Broyd}_{\tau_1}(G, A, u) \preceq \text{Broyd}_{\tau_2}(G, A, u).$$

*Proof.* Suppose that  $Gu \neq Au$  since otherwise the claim is trivial. Then,

$$\begin{aligned} \text{Broyd}_{\tau}(G, A, u) &\stackrel{(2.2)}{=} G - \frac{(G-A)uu^*(G-A)}{\langle(G-A)u, u\rangle} \\ &\quad + \tau \left[ \frac{(G-A)uu^*(G-A)}{\langle(G-A)u, u\rangle} - \frac{Auu^*G+Guu^*A}{\langle Au, u\rangle} + \left( \frac{\langle Gu, u\rangle}{\langle Au, u\rangle} + 1 \right) \frac{Auu^*A}{\langle Au, u\rangle} \right]. \end{aligned}$$

Denote  $s \stackrel{\text{def}}{=} \frac{(G-A)u}{\langle(G-A)u, u\rangle} - \frac{Au}{\langle Au, u\rangle}$ . Then,

$$\begin{aligned} \langle(G-A)u, u\rangle ss^* &= \frac{(G-A)uu^*(G-A)}{\langle(G-A)u, u\rangle} + \frac{\langle(G-A)u, u\rangle}{\langle Au, u\rangle} \frac{Auu^*A}{\langle Au, u\rangle} - \frac{(G-A)uu^*A + Auu^*(G-A)}{\langle Au, u\rangle} \\ &= \frac{(G-A)uu^*(G-A)}{\langle(G-A)u, u\rangle} - \frac{Auu^*G+Guu^*A}{\langle Au, u\rangle} + \left( \frac{\langle Gu, u\rangle}{\langle Au, u\rangle} + 1 \right) \frac{Auu^*A}{\langle Au, u\rangle}. \end{aligned}$$

Therefore,

$$\text{Broyd}_{\tau}(G, A, u) = G - \frac{(G-A)uu^*(G-A)}{\langle(G-A)u, u\rangle} + \tau \langle(G-A)u, u\rangle ss^*.$$

The claim now follows from the fact that  $\langle(G-A)u, u\rangle ss^* \succeq 0$  in view of (2.1).  $\square$

Next, let us show that the relation (2.1) can be preserved after applying to  $G$  any update from the class of our interest. Moreover, each update from this class does not increase the deviation from the target operator  $A$ .

LEMMA 2.2. *If, for some  $\eta \geq 1$ , we have*

$$(2.7) \quad A \preceq G \preceq \eta A,$$

*then, for any  $u \in \mathbb{E}$  and any  $\tau \in [0, 1]$ , we also have*

$$(2.8) \quad A \preceq \text{Broyd}_{\tau}(G, A, u) \preceq \eta A.$$

*Proof.* Denote  $G_+ \stackrel{\text{def}}{=} \text{Broyd}_{\tau}(G, A, u)$  and assume that  $Gu \neq Au$  since otherwise the claim is trivial. Using that  $\tau \geq 0$  and applying Lemma 2.1, we obtain

$$G_+ \succeq \text{SR1}(G, A, u) \stackrel{(2.3)}{=} G - \frac{(G-A)uu^*(G-A)}{\langle(G-A)u, u\rangle}.$$

Let  $R \stackrel{\text{def}}{=} G - A \succeq 0$ , and let  $I_{\mathbb{E}}, I_{\mathbb{E}^*}$  be the identity operators in the spaces  $\mathbb{E}, \mathbb{E}^*$  respectively. Then,

$$G_+ - A \succeq R - \frac{Ruu^*R}{\langle Ru, u\rangle} = \left( I_{\mathbb{E}^*} - \frac{Ruu^*}{\langle Ru, u\rangle} \right) R \left( I_{\mathbb{E}} - \frac{uu^*R}{\langle Ru, u\rangle} \right) \succeq 0,$$

Thus, the first relation in (2.8) is proved. To prove the second relation, we apply Lemma 2.1, using that  $\tau \leq 1$ , to obtain

$$\begin{aligned} G_+ &\preceq \text{DFP}(G, A, u) \stackrel{(2.4)}{=} G + \left( \frac{\langle Gu, u\rangle}{\langle Au, u\rangle} + 1 \right) \frac{Auu^*A}{\langle Au, u\rangle} - \frac{Auu^*G+Guu^*A}{\langle Au, u\rangle} \\ &= \frac{Auu^*A}{\langle Au, u\rangle} + \left( I_{\mathbb{E}^*} - \frac{Auu^*}{\langle Au, u\rangle} \right) G \left( I_{\mathbb{E}} - \frac{uu^*A}{\langle Au, u\rangle} \right) \\ &\stackrel{(2.7)}{\preceq} \frac{Auu^*A}{\langle Au, u\rangle} + \eta \left( I_{\mathbb{E}^*} - \frac{Auu^*}{\langle Au, u\rangle} \right) A \left( I_{\mathbb{E}} - \frac{uu^*A}{\langle Au, u\rangle} \right) \\ &= \frac{Auu^*A}{\langle Au, u\rangle} + \eta \left( A - \frac{Auu^*A}{\langle Au, u\rangle} \right) = \eta A - (\eta - 1) \frac{Auu^*A}{\langle Au, u\rangle} \preceq \eta A. \end{aligned}$$

The proof is now finished.  $\square$

*Remark 2.3.* Similar results to the one from Lemma 2.2 have been known for some time in the literature for different quasi-Newton updating formulas. For example, in [4] and [5], it was proved for the SR1 update that if  $A \preceq G$  (respectively,  $G \preceq A$ ), then  $A \preceq G_+$  (respectively,  $G_+ \preceq A$ ), where  $G_+$  is the result of the corresponding update. An even stronger property was established in [8] for the *convex Broyden class* (composed of all convex combinations of the BFGS and DFP updates); in particular, it was shown that if  $\eta_1 A \preceq G \preceq \eta_2 A$  for some  $0 < \eta_1 \leq 1 \leq \eta_2$ , then  $\eta_1 A \preceq G_+ \preceq \eta_2 A$ .

Interestingly, from Lemma 2.1 and Lemma 2.2, it follows that, if (2.1) holds, then

$$A \preceq \text{SR1}(G, A, u) \preceq \text{BFGS}(G, A, u) \preceq \text{DFP}(G, A, u).$$

In other words, the approximation, produced by SR1, is better than the one, produced by BFGS, which is in turn better than the one, produced by DFP.

Let us now justify the efficiency of update (2.2) with  $\tau \in [0, 1]$  in ensuring convergence  $G \rightarrow A$ . For this, we introduce the following measure of progress:

$$(2.9) \quad \sigma_A(G) \stackrel{\text{def}}{=} \langle A^{-1}, G - A \rangle \stackrel{(1.3)}{=} \langle A^{-1}, G \rangle - n.$$

Thus,  $\sigma_A(G)$  is the sum of the eigenvalues of the difference  $G - A$ , measured with respect to the operator  $A$ . Clearly, for  $G$ , satisfying (2.1), we have  $\sigma_A(G) \geq 0$  with  $\sigma_A(G) = 0$  if and only if  $G = A$ . Therefore, we need to ensure that  $\sigma_A(G) \rightarrow 0$  by choosing an appropriate sequence of update directions  $u$ .

First, let us estimate the decrease in the measure  $\sigma_A$  for an arbitrary direction.

LEMMA 2.4. *Let (2.1) hold. Then, for any  $u \in \mathbb{E}$  and any  $\tau \in [0, 1]$ , we have*

$$(2.10) \quad \sigma_A(G) - \sigma_A(\text{Broyd}_\tau(G, A, u)) \geq \frac{\langle (G-A)u, u \rangle}{\langle Au, u \rangle}.$$

*Proof.* Denote  $G_+ \stackrel{\text{def}}{=} \text{Broyd}_\tau(G, A, u)$  and assume that  $Gu \neq Au$  since otherwise the claim is trivial. By Lemma 2.1, we have

$$G - G_+ \succeq G - \text{DFP}(G, A, u) \stackrel{(2.4)}{=} \frac{Auu^*G + Guu^*A}{\langle Au, u \rangle} - \left( \frac{\langle Gu, u \rangle}{\langle Au, u \rangle} + 1 \right) \frac{Auu^*A}{\langle Au, u \rangle}.$$

Therefore,

$$\begin{aligned} \sigma_A(G) - \sigma_A(G_+) &\stackrel{(2.9)}{=} \langle A^{-1}, G - G_+ \rangle \geq 2 \frac{\langle Gu, u \rangle}{\langle Au, u \rangle} - \left( \frac{\langle Gu, u \rangle}{\langle Au, u \rangle} + 1 \right) \\ &= \frac{\langle Gu, u \rangle}{\langle Au, u \rangle} - 1 = \frac{\langle (G-A)u, u \rangle}{\langle Au, u \rangle}. \end{aligned}$$

The proof is now finished.  $\square$

According to Lemma 2.4, the choice of the updating direction  $u$  directly influences the bound on the decrease in the measure  $\sigma_A$ . Ideally, we would like to select a direction  $u$ , which maximizes the right-hand side in (2.10). However, this requires finding an eigenvector, corresponding to the maximal eigenvalue of  $G$  with respect to  $A$ , which might be computationally a difficult problem. Therefore, let us consider another approach.

Let us fix in the space  $\mathbb{E}$  some basis:

$$e_1, \dots, e_n \in \mathbb{E}.$$

With respect to this basis, we can define the following greedily selected direction:

$$(2.11) \quad \bar{u}_A(G) \stackrel{\text{def}}{=} \operatorname{argmax}_{u \in \{e_1, \dots, e_n\}} \frac{\langle (G-A)u, u \rangle}{\langle Au, u \rangle} = \operatorname{argmax}_{u \in \{e_1, \dots, e_n\}} \frac{\langle Gu, u \rangle}{\langle Au, u \rangle}.$$

Thus,  $\bar{u}_A(G)$  is a basis vector, which maximizes the right-hand side in (2.10). Note that for certain choices of the basis, the computation of  $\bar{u}_A(G)$  might be relatively simple. For example, if  $\mathbb{E} = \mathbb{R}^n$ , and  $e_1, \dots, e_n$  are coordinate directions, then the calculation of  $\bar{u}_A(G)$  requires computing only the *diagonals* of the matrix representations of the operators  $G$  and  $A$ . The update (2.2), applying the rule (2.11), is called the *greedy quasi-Newton update*.

Let us show that the greedy quasi-Newton update decreases the measure  $\sigma_A$  with a *linear* rate. For this, define

$$(2.12) \quad B \stackrel{\text{def}}{=} \left( \sum_{i=1}^n e_i e_i^* \right)^{-1}.$$

Note that  $B$  is a self-adjoint positive definite linear operator from  $\mathbb{E}$  to  $\mathbb{E}^*$ .

**THEOREM 2.5.** *Let (2.1) hold, and let  $\mu, L > 0$  be such that*

$$(2.13) \quad \mu B \preceq A \preceq LB.$$

*Then, for any  $\tau \in [0, 1]$ , we have*

$$(2.14) \quad \sigma_A(\text{Broyd}_\tau(G, A, \bar{u}_A(G))) \leq \left(1 - \frac{\mu}{nL}\right) \sigma_A(G).$$

*Proof.* Denote  $G_+ \stackrel{\text{def}}{=} \text{Broyd}_\tau(G, A, \bar{u}_A(G))$ , and  $R \stackrel{\text{def}}{=} G - A$ . By Lemma 2.4,

$$\begin{aligned} \sigma_A(G) - \sigma_A(G_+) &\geq \frac{\langle R\bar{u}_A(G), \bar{u}_A(G) \rangle}{\langle A\bar{u}_A(G), \bar{u}_A(G) \rangle} \stackrel{(2.11)}{=} \max_{1 \leq i \leq n} \frac{\langle Re_i, e_i \rangle}{\langle Ae_i, e_i \rangle} \stackrel{(2.13)}{\geq} \frac{1}{L} \max_{1 \leq i \leq n} \langle Re_i, e_i \rangle \\ &\geq \frac{1}{nL} \sum_{i=1}^n \langle Re_i, e_i \rangle \stackrel{(1.2)}{=} \frac{1}{nL} \sum_{i=1}^n \langle e_i e_i^*, R \rangle \stackrel{(2.12)}{=} \frac{1}{nL} \langle B^{-1}, R \rangle \\ &\stackrel{(2.13)}{\geq} \frac{\mu}{nL} \langle A^{-1}, R \rangle \stackrel{(2.9)}{=} \frac{\mu}{nL} \sigma_A(G). \end{aligned}$$

The proof is now finished.  $\square$

*Remark 2.6.* A simple modification of the above proof shows that the factor  $nL$  in (2.14) can be improved up to  $\langle B^{-1}, A \rangle$ . However, to simplify the future analysis, we prefer to work directly with constant  $L$ .

**3. Unconstrained Quadratic Minimization.** Let us demonstrate how we can apply the quasi-Newton updates, described in the previous section, for minimizing the quadratic function

$$(3.1) \quad f(x) \stackrel{\text{def}}{=} \frac{1}{2} \langle Ax, x \rangle - \langle b, x \rangle, \quad x \in \mathbb{E},$$

where  $A : \mathbb{E} \rightarrow \mathbb{E}^*$  is a self-adjoint positive definite linear operator, and  $b \in \mathbb{E}^*$ .

Let  $B$  be the operator, defined in (2.12), and let  $\mu, L > 0$  be such that

$$(3.2) \quad \mu B \preceq A \preceq LB.$$

Thus,  $\mu$  is the *constant of strong convexity* of  $f$ , and  $L$  is the *Lipschitz constant* of the gradient of  $f$ , both measured with respect to the operator  $B$ .

Consider the following quasi-Newton scheme:

<b>Initialization:</b> Choose $x_0 \in \mathbb{E}$ . Set $G_0 = LB$ .
<b>For <math>k \geq 0</math> iterate:</b>
1. Update $x_{k+1} = x_k - G_k^{-1} \nabla f(x_k)$ .
2. Choose $u_k \in \mathbb{E}$ and $\tau_k \in [0, 1]$ .
3. Compute $G_{k+1} = \text{Broyd}_{\tau_k}(G_k, A, u_k)$ .

(3.3)

Note that scheme (3.3) starts with  $G_0 = LB$ . Therefore, its first iteration is identical to that one of the standard *gradient method*:

$$x_1 = x_0 - \frac{1}{L} B^{-1} \nabla f(x_0).$$

Also, from (3.2), it follows that  $A \preceq G_0$ . Hence, in view of Lemma 2.2, we have

$$(3.4) \quad A \preceq G_k$$

for all  $k \geq 0$ . In particular, all  $G_k$  are positive definite, and scheme (3.3) is well-defined.

*Remark 3.1.* For avoiding the  $O(n^3)$  complexity for computing  $G_k^{-1} \nabla f(x_k)$ , it is typical for practical implementation of scheme (3.3) to work directly with the inverse operators  $G_k^{-1}$  (or, alternatively, with the Cholesky decomposition of  $G_k$ ). Due to a low-rank structure of the updates (2.2), it is possible to compute efficiently  $G_{k+1}^{-1}$  via  $G_k^{-1}$  at the cost  $O(n^2)$ .

To estimate the convergence rate of scheme (3.3), let us look at the norm of the gradient of  $f$ , measured with respect to  $A$ :

$$(3.5) \quad \lambda_f(x) \stackrel{\text{def}}{=} \|\nabla f(x)\|_A^* \stackrel{(1.1)}{=} \langle \nabla f(x), A^{-1} \nabla f(x) \rangle^{1/2}, \quad x \in \mathbb{E}.$$

Note that this measure of optimality is directly related to the functional residual. Indeed, let  $x^* = A^{-1}b$  be the minimizer of (3.1). Then, using Taylor's formula, we obtain

$$\begin{aligned} f(x) - f^* &= \frac{1}{2} \langle A(x - x^*), x - x^* \rangle = \frac{1}{2} \langle Ax - b, A^{-1}(Ax - b) \rangle \\ &\stackrel{(3.1)}{=} \frac{1}{2} \langle \nabla f(x), A^{-1} \nabla f(x) \rangle \stackrel{(3.5)}{=} \frac{1}{2} \lambda_f^2(x). \end{aligned}$$

The following lemma shows how  $\lambda_f$  changes after one iteration of process (3.3).

LEMMA 3.2. *Let  $k \geq 0$ , and let  $\eta_k \geq 1$  be such that*

$$(3.6) \quad G_k \preceq \eta_k A.$$

Then,

$$\lambda_f(x_{k+1}) \leq \left(1 - \frac{1}{\eta_k}\right) \lambda_f(x_k) = \frac{\eta_k - 1}{\eta_k} \lambda_f(x_k).$$



*Proof.* Indeed,

$$\nabla f(x_{k+1}) = \nabla f(x_k) + A(x_{k+1} - x_k) \stackrel{(3.3)}{=} A(A^{-1} - G_k^{-1})\nabla f(x_k).$$

Therefore,

$$\lambda_f(x_{k+1}) \stackrel{(3.5)}{=} \langle \nabla f(x_k), (A^{-1} - G_k^{-1})A(A^{-1} - G_k^{-1})\nabla f(x_k) \rangle^{1/2}.$$

Note that

$$\frac{1}{\eta_k}A^{-1} \stackrel{(3.6)}{\preceq} G_k^{-1} \stackrel{(3.4)}{\preceq} A^{-1}.$$

Therefore,

$$(3.7) \quad 0 \preceq A^{-1} - G_k^{-1} \preceq \left(1 - \frac{1}{\eta_k}\right)A^{-1}.$$

Consequently,

$$(A^{-1} - G_k^{-1})A(A^{-1} - G_k^{-1}) \preceq \left(1 - \frac{1}{\eta_k}\right)^2 A^{-1},$$

and

$$\lambda_f(x_{k+1}) \stackrel{(3.7)}{\leq} \left(1 - \frac{1}{\eta_k}\right) \langle \nabla f(x_k), A^{-1}\nabla f(x_k) \rangle^{1/2} \stackrel{(3.5)}{=} \left(1 - \frac{1}{\eta_k}\right) \lambda_f(x_k).$$

The proof is now finished.  $\square$

Thus, to estimate how fast  $\lambda_f(x_k)$  converges to zero, we need to upper bound  $\eta_k$ . There are two ways to proceed, depending on the choice of directions  $u_k$  in (3.3).

First, consider the general situation, when we do not impose any restrictions on  $u_k$ . In this case, we can guarantee that  $\eta_k$  stays uniformly bounded, and  $\lambda_f(x_k) \rightarrow 0$  at a *linear* rate.

**THEOREM 3.3.** *For all  $k \geq 0$ , in scheme (3.3), we have*

$$(3.8) \quad A \preceq G_k \preceq \frac{L}{\mu}A,$$

and

$$(3.9) \quad \lambda_f(x_k) \leq \left(1 - \frac{\mu}{L}\right)^k \lambda_f(x_0).$$

*Proof.* Since  $G_0 = LB$ , in view of (3.2), we have

$$A \preceq G_0 \preceq \frac{L}{\mu}A.$$

By Lemma 2.2, this implies (3.8). Applying now Lemma 3.2, we obtain

$$\lambda_f(x_{k+1}) \leq \left(1 - \frac{\mu}{L}\right) \lambda_f(x_k)$$

for all  $k \geq 0$ , and (3.9) follows.  $\square$

Note that (3.9) is exactly the convergence rate of the standard gradient method. Thus, according to Theorem 3.3, the convergence rate of scheme (3.3) is at least as good as that of the gradient method.

Now assume that the directions  $u_k$  in scheme (3.3) are chosen in accordance to the greedy strategy (2.11). Recall that, in this case, we can guarantee that  $G_k \rightarrow A$  (Theorem 2.5). Therefore, we can expect faster convergence from scheme (3.3).

THEOREM 3.4. *Suppose that, for each  $k \geq 0$ , we choose  $u_k = \bar{u}_A(G_k)$  in scheme (3.3). Then, for all  $k \geq 0$ , we have*

$$(3.10) \quad A \preceq G_k \preceq \left(1 + \left(1 - \frac{\mu}{nL}\right)^k \frac{nL}{\mu}\right) A,$$

and

$$(3.11) \quad \lambda_f(x_{k+1}) \leq \left(1 - \frac{\mu}{nL}\right)^k \frac{nL}{\mu} \cdot \lambda_f(x_k).$$

*Proof.* We already know that  $A \preceq G_k$ . Hence,

$$G_k - A \stackrel{(1.4)}{\preceq} \langle A^{-1}, G_k - A \rangle A \stackrel{(2.9)}{=} \sigma_A(G_k)A,$$

or, equivalently,

$$G_k \preceq (1 + \sigma_A(G_k))A.$$

At the same time, by Theorem 2.5, we have

$$\sigma_A(G_k) \leq \left(1 - \frac{\mu}{nL}\right)^k \sigma_A(G_0).$$

Note that

$$\sigma_A(G_0) \stackrel{(2.9)}{=} \langle A^{-1}, G_0 \rangle - n \stackrel{(3.8)}{\leq} \langle A^{-1}, \frac{L}{\mu}A \rangle - n \stackrel{(1.3)}{=} n \left(\frac{L}{\mu} - 1\right) \leq \frac{nL}{\mu}.$$

Thus, (3.10) is proved. Applying now Lemma 3.2 and using the fact that  $\frac{\eta-1}{\eta} \leq \eta - 1$  for any  $\eta \geq 1$ , we obtain (3.11).  $\square$

Theorem 3.4 shows that the convergence rate of  $\lambda_f(x_k)$  is *superlinear*. Let us now combine this result with Theorem 3.3 and write down the final efficiency estimate. Denote by  $k_0 \geq 0$  the number of the first iteration, for which

$$(3.12) \quad \left(1 - \frac{\mu}{nL}\right)^{k_0} \frac{nL}{\mu} \leq \frac{1}{2}.$$

Clearly,  $k_0 \leq \frac{nL}{\mu} \ln \frac{2nL}{\mu}$ . According to Theorem 2.5, during the first  $k_0$  iterations,

$$(3.13) \quad \lambda_f(x_k) \leq \left(1 - \frac{\mu}{nL}\right)^k \lambda_f(x_0).$$

After that, by Theorem 3.4, for all  $k \geq 0$ , we have

$$\lambda_f(x_{k_0+k+1}) \stackrel{(3.11)}{\leq} \left(1 - \frac{\mu}{nL}\right)^{k_0+k} \frac{nL}{\mu} \lambda_f(x_{k_0+k}) \stackrel{(3.12)}{\leq} \left(1 - \frac{\mu}{nL}\right)^k \frac{1}{2} \lambda_f(x_{k_0+k}),$$

or, more explicitly,

$$\begin{aligned} \lambda_f(x_{k_0+k}) &\leq \lambda_f(x_{k_0}) \prod_{i=0}^{k-1} \left[ \left(1 - \frac{\mu}{nL}\right)^i \frac{1}{2} \right] = \left(1 - \frac{\mu}{nL}\right)^{\sum_{i=0}^{k-1} i} \left(\frac{1}{2}\right)^k \lambda_f(x_{k_0}) \\ &= \left(1 - \frac{\mu}{nL}\right)^{\frac{k(k-1)}{2}} \left(\frac{1}{2}\right)^k \lambda_f(x_{k_0}) \\ &\stackrel{(3.13)}{\leq} \left(1 - \frac{\mu}{nL}\right)^{\frac{k(k-1)}{2}} \left(\frac{1}{2}\right)^k \left(1 - \frac{\mu}{nL}\right)^{k_0} \lambda_f(x_0). \end{aligned}$$

Note that the first factor in this estimate depends on the *square* of the iteration counter.

To conclude, let us mention one important property of scheme (3.3) with greedily selected  $u_k$ . It turns out that, in the particular case, when  $\tau_k = 0$  for all  $k \geq 0$ , i.e. when scheme (3.3) corresponds to the greedy *SR1* method, it will identify the operator  $A$ , and consequently, the minimizer  $x^*$  of (3.1), in a *finite* number of steps.

**THEOREM 3.5.** *Suppose that, in scheme (3.3), for each  $k \geq 0$ , we choose  $u_k = \bar{u}_A(G_k)$  and  $\tau_k = 0$ . Then  $G_k = A$  for some  $0 \leq k \leq n$ .*

*Proof.* Suppose that  $R_k \stackrel{\text{def}}{=} G_k - A \neq 0$  for all  $0 \leq k \leq n$ . Since  $R_k \succeq 0$  (see (3.4)), we must have  $u_k \notin \text{Ker}(R_k)$  in view of (2.11), and

$$R_{k+1} \stackrel{(2.3)}{=} R_k - \frac{R_k u_k u_k^* R_k}{\langle R_k u_k, u_k \rangle}$$

for all  $0 \leq k \leq n$ . From this formula, it is easily seen that

- (1)  $\text{Ker}(R_k) \subseteq \text{Ker}(R_{k+1})$ ,
- (2)  $u_k \in \text{Ker}(R_{k+1})$ .

Thus, the dimension of  $\text{Ker}(R_k)$  grows at least by 1 at every iteration. In particular, the dimension of  $\text{Ker}(R_{n+1})$  must be at least  $n+1$ , which is impossible, since the operator  $R_{n+1}$  acts in an  $n$ -dimensional vector space.  $\square$

It is worth noting that for other updates, such as (2.4) and (2.6), the inclusion  $\text{Ker}(R_k) \subseteq \text{Ker}(R_{k+1})$  is, in general, no longer valid.

**4. Minimization of General Functions.** Now consider a general problem of unconstrained minimization:

$$(4.1) \quad \min_{x \in \mathbb{E}} f(x),$$

where  $f : \mathbb{E} \rightarrow \mathbb{R}$  is a twice differentiable function with positive definite Hessian. Our goal is to extend the results, obtained in the previous section, onto the problem (4.1), assuming that the methods can start from a sufficiently good initial point  $x_0$ .

Our main assumption is that the Hessians of  $f$  are close to each other in the sense that there exists a constant  $M \geq 0$ , such that

$$(4.2) \quad \nabla^2 f(y) - \nabla^2 f(x) \preceq M \|y - x\|_z \nabla^2 f(w)$$

for all  $x, y, z, w \in \mathbb{E}$ . We call such a function  $f$  *strongly self-concordant*. Note that strongly self-concordant functions form a subclass of self-concordant functions. Indeed, let us choose a point  $x \in \mathbb{E}$  and a direction  $h \in \mathbb{E}$ . Then, for all  $t > 0$ , we have

$$\langle [\nabla^2 f(x + th) - \nabla^2 f(x)]h, h \rangle \leq Mt \|h\|_x^3.$$

Dividing this inequality by  $t$  and computing the limit as  $t \downarrow 0$ , we obtain

$$D^3 f(x)[h, h, h] \leq M \|h\|_x^3.$$

for all  $h \in \mathbb{E}$ . Thus, function  $f$  is self-concordant with constant  $\frac{1}{2}M$  (see [22, 32]).

The main example of a strongly self-concordant function is a strongly convex function with Lipschitz continuous Hessian. Note however that strong self-concordancy is an *affine-invariant* property.

*Example 4.1.* Let  $C : \mathbb{E} \rightarrow \mathbb{E}^*$  be a self-adjoint positive definite operator. Suppose there exist  $\beta > 0$  and  $L_2 \geq 0$ , such that the function  $f$  is  $\beta$ -strongly convex and its Hessian is  $L_2$ -Lipschitz continuous with respect to the norm  $\|\cdot\|_C$ . Then  $f$  is strongly self-concordant with constant  $M = \frac{L_2}{\beta^{3/2}}$ .

*Proof.* By strong convexity of  $f$ , we have

$$(4.3) \quad \beta C \preceq \nabla^2 f(x)$$

for all  $x \in \mathbb{E}$ . Therefore, using the Lipschitz continuity of the Hessian, we obtain

$$\begin{aligned} \nabla^2 f(y) - \nabla^2 f(x) &\preceq L_2 \|y - x\|_C C \stackrel{(1.1)}{=} L_2 \langle C(y - x), y - x \rangle^{1/2} C \\ &\stackrel{(4.3)}{\preceq} \frac{L_2}{\mu^{1/2}} \langle \nabla^2 f(z)(y - x), y - x \rangle^{1/2} C \\ &\stackrel{(1.1)}{=} \frac{L_2}{\mu^{1/2}} \|y - x\|_z C \stackrel{(4.3)}{\preceq} \frac{L_2}{\mu^{3/2}} \|y - x\|_z \nabla^2 f(w) \end{aligned}$$

for all  $x, y, z, w \in \mathbb{E}$ .  $\square$

Let us establish some useful relations for strongly self-concordant functions.

LEMMA 4.2. *Let  $x, y \in \mathbb{E}$ , and let  $r \stackrel{\text{def}}{=} \|y - x\|_x$ . Then,*

$$(4.4) \quad \frac{\nabla^2 f(x)}{1+Mr} \preceq \nabla^2 f(y) \preceq (1+Mr) \nabla^2 f(x).$$

Also, for  $J \stackrel{\text{def}}{=} \int_0^1 \nabla^2 f(x + t(y - x)) dt$  and any  $v \in \{x, y\}$ , we have

$$(4.5) \quad \frac{\nabla^2 f(v)}{1+\frac{Mr}{2}} \preceq J \preceq \left(1 + \frac{Mr}{2}\right) \nabla^2 f(v).$$

*Proof.* Denote  $h \stackrel{\text{def}}{=} y - x$ . Taking  $z = w = x$  in (4.2), we obtain

$$\nabla^2 f(y) - \nabla^2 f(x) \preceq Mr \nabla^2 f(x),$$

which gives us the second relation in (4.4) after moving  $\nabla^2 f(x)$  into the right-hand side. Interchanging now  $x$  and  $y$  in (4.2) and taking  $z = x, w = y$ , we get

$$\nabla^2 f(x) - \nabla^2 f(y) \preceq Mr \nabla^2 f(y),$$

which gives us the first relation in (4.4) after moving  $\nabla^2 f(x)$  into the right-hand side and then dividing by  $1 + Mr$ .

Let us now prove (4.5) for  $v = x$  (the proof for  $v = y$  is similar). Choosing  $y = x + th$  in (4.2) for  $t > 0$ , and  $w = z = x$ , we obtain

$$\nabla^2 f(x + th) - \nabla^2 f(x) \preceq M \|th\|_x \nabla^2 f(x) = Mrt \nabla^2 f(x).$$

This gives us the second relation in (4.5) after integrating for  $t$  from 0 to 1 and moving  $\nabla^2 f(x)$  into the right-hand side. Interchanging  $x$  and  $y$  in (4.2) and taking  $y = x + th$  for  $t > 0, z = x$ , while leaving  $w$  arbitrary, we get

$$\nabla^2 f(x) - \nabla^2 f(x + th) \preceq M \| -th \|_x \nabla^2 f(w) = Mrt \nabla^2 f(w).$$

Hence, by integrating for  $t$  from 0 to 1, we see that

$$\nabla^2 f(x) - J \preceq \frac{Mr}{2} \nabla^2 f(w).$$

Taking now  $w = x + th$  and integrating again, we obtain

$$\nabla^2 f(x) - J \preceq \frac{Mr}{2} \int_0^1 \nabla^2 f(x + th) dt = \frac{Mr}{2} J,$$

and the first inequality in (4.5) follows after moving  $J$  to the right-hand side and dividing by  $1 + \frac{Mr}{2}$ .  $\square$

Let us now estimate the progress of a general quasi-Newton step. As before, for measuring the progress, we use the *local norm of the gradient*:

$$(4.6) \quad \lambda_f(x) \stackrel{\text{def}}{=} \|\nabla f(x)\|_x^* \stackrel{(1.1)}{=} \langle \nabla f(x), \nabla^2 f(x)^{-1} \nabla f(x) \rangle^{1/2}, \quad x \in \mathbb{E}.$$

LEMMA 4.3. *Let  $x \in \mathbb{E}$ , and let  $G : \mathbb{E} \rightarrow \mathbb{E}^*$  be a self-adjoint linear operator, such that*

$$(4.7) \quad \nabla^2 f(x) \preceq G \preceq \eta \nabla^2 f(x)$$

for some  $\eta \geq 1$ . Let

$$(4.8) \quad x_+ \stackrel{\text{def}}{=} x - G^{-1} \nabla f(x),$$

and let  $\lambda \stackrel{\text{def}}{=} \lambda_f(x)$  be such that  $M\lambda \leq 2$ . Then,  $r \stackrel{\text{def}}{=} \|x_+ - x\|_x \leq \lambda$ , and

$$\lambda_f(x_+) \leq \left(1 + \frac{M\lambda}{2}\right) \frac{\eta^{-1} + \frac{M\lambda}{2}}{\eta} \lambda.$$

*Proof.* Denote  $J \stackrel{\text{def}}{=} \int_0^1 \nabla^2 f(x + t(x_+ - x)) dt$ . Applying Taylor's formula, we obtain

$$(4.9) \quad \nabla f(x_+) = \nabla f(x) + J(x_+ - x) \stackrel{(4.8)}{=} J(J^{-1} - G^{-1})\nabla f(x).$$

Note that

$$\begin{aligned} r &= \|x_+ - x\|_x \stackrel{(4.8)}{=} \|G^{-1} \nabla f(x)\|_x \stackrel{(1.1)}{=} \langle \nabla f(x), G^{-1} \nabla^2 f(x) G^{-1} \nabla f(x) \rangle^{1/2} \\ &\stackrel{(4.7)}{\leq} \langle \nabla f(x), G^{-1} \nabla f(x) \rangle^{1/2} \stackrel{(4.7)}{\leq} \langle \nabla f(x), \nabla^2 f(x)^{-1} \nabla f(x) \rangle^{1/2} \stackrel{(4.6)}{=} \lambda. \end{aligned}$$

Hence, in view of Lemma 4.2, we have

$$(4.10) \quad \frac{\nabla^2 f(x)}{1 + \frac{M\lambda}{2}} \preceq J \preceq \left(1 + \frac{M\lambda}{2}\right) \nabla^2 f(x), \quad J \preceq \left(1 + \frac{M\lambda}{2}\right) \nabla^2 f(x_+).$$

Therefore,

$$(4.11) \quad \begin{aligned} \lambda_f(x_+) &\stackrel{(4.6)}{=} \langle \nabla f(x_+), \nabla^2 f(x_+)^{-1} \nabla f(x_+) \rangle^{1/2} \\ &\stackrel{(4.10)}{\leq} \sqrt{1 + \frac{M\lambda}{2}} \langle \nabla f(x_+), J^{-1} \nabla f(x_+) \rangle^{1/2} \\ &\stackrel{(4.9)}{=} \sqrt{1 + \frac{M\lambda}{2}} \langle \nabla f(x), (J^{-1} - G^{-1}) J (J^{-1} - G^{-1}) \nabla f(x) \rangle^{1/2}. \end{aligned}$$

Further,

$$\frac{1}{1 + \frac{M\lambda}{2}} J \stackrel{(4.10)}{\preceq} \nabla^2 f(x) \stackrel{(4.7)}{\preceq} G \stackrel{(4.7)}{\preceq} \eta \nabla^2 f(x) \stackrel{(4.10)}{\preceq} \eta \left(1 + \frac{M\lambda}{2}\right) J.$$

Hence,

$$\frac{1}{(1 + \frac{M\lambda}{2})\eta} J^{-1} \preceq G^{-1} \preceq \left(1 + \frac{M\lambda}{2}\right) J^{-1},$$

and

$$-\left(1 - \frac{1}{(1 + \frac{M\lambda}{2})\eta}\right) J^{-1} \preceq G^{-1} - J^{-1} \preceq \frac{M\lambda}{2} J^{-1}.$$

Note that

$$1 - \frac{1}{(1 + \frac{M\lambda}{2})\eta} \leq 1 - \frac{1 - \frac{M\lambda}{2}}{\eta} = \frac{\eta - 1 + \frac{M\lambda}{2}}{\eta},$$

and, since  $M\lambda \leq 2$ ,

$$\frac{M\lambda}{2} = 1 - \left(1 - \frac{M\lambda}{2}\right) \leq 1 - \frac{1 - \frac{M\lambda}{2}}{\eta} = \frac{\eta - 1 + \frac{M\lambda}{2}}{\eta}.$$

Therefore,

$$-\frac{\eta - 1 + \frac{M\lambda}{2}}{\eta} J^{-1} \preceq G^{-1} - J^{-1} \preceq \frac{\eta - 1 + \frac{M\lambda}{2}}{\eta} J^{-1}.$$

Consequently,

$$(G^{-1} - J^{-1})J(G^{-1} - J^{-1}) \preceq \left(\frac{\eta - 1 + \frac{M\lambda}{2}}{\eta}\right)^2 J^{-1}.$$

and thus,

$$\begin{aligned} \lambda_f(x_+) &\stackrel{(4.11)}{\leq} \sqrt{1 + \frac{M\lambda}{2} \frac{\eta - 1 + \frac{M\lambda}{2}}{\eta}} \langle \nabla f(x), J^{-1} \nabla f(x) \rangle^{1/2} \\ &\stackrel{(4.10)}{\leq} \left(1 + \frac{M\lambda}{2}\right) \frac{\eta - 1 + \frac{M\lambda}{2}}{\eta} \langle \nabla f(x), \nabla^2 f(x)^{-1} \nabla f(x) \rangle^{1/2} \\ &\stackrel{(4.6)}{=} \left(1 + \frac{M\lambda}{2}\right) \frac{\eta - 1 + \frac{M\lambda}{2}}{\eta} \lambda. \end{aligned}$$

The proof is now finished.  $\square$

Now we need to analyze what happens with the Hessian approximation after a quasi-Newton update. Let  $G$  be the current approximation of  $\nabla^2 f(x)$ , satisfying, as usual, the condition

$$(4.12) \quad \nabla^2 f(x) \preceq G.$$

Using this approximation, we can compute the new test point

$$x_+ = x - G^{-1} \nabla f(x).$$

After that, we would like to update  $G$  into a new operator  $G_+$ , approximating the Hessian  $\nabla^2 f(x_+)$  at the new point and satisfying the condition

$$\nabla^2 f(x_+) \preceq G_+.$$

A natural idea is, of course, to set

$$(4.13) \quad G_+ = \text{Broyd}_\tau(G, \nabla^2 f(x_+), u)$$

for some  $u \in \mathbb{E}$  and  $\tau \in [0, 1]$ . However, we cannot do this, since update (4.13) is well-defined only when

$$\nabla^2 f(x_+) \preceq G$$

(see Section 2), which may not be true, even though (4.12) holds. To avoid this problem, let us apply the following *correction strategy*:

1. Choose some  $\delta \geq 0$ , and set  $\tilde{G} = (1 + \delta)G$ .
2. Compute  $G_+$ , using (4.13) with  $G$  replaced by  $\tilde{G}$ .

Clearly, for some value of  $\delta$ , the condition  $\nabla^2 f(x_+) \preceq \tilde{G}$  will be valid. If, at the same time, this  $\delta$  is sufficiently small, then the above correction strategy should not introduce too big error.

LEMMA 4.4. *Let  $x \in \mathbb{E}$ , and let  $G : \mathbb{E} \rightarrow \mathbb{E}^*$  be a self-adjoint linear operator, such that*

$$(4.14) \quad \nabla^2 f(x) \preceq G \preceq \eta \nabla^2 f(x)$$

for some  $\eta \geq 1$ . Let  $x_+ \in \mathbb{E}$ , let  $r \stackrel{\text{def}}{=} \|x_+ - x\|_x$ . Then

$$(4.15) \quad \tilde{G} \stackrel{\text{def}}{=} (1 + Mr)G \succeq \nabla^2 f(x_+),$$

and, for all  $u \in \mathbb{E}$  and  $\tau \in [0, 1]$ , we have

$$\nabla^2 f(x_+) \preceq \text{Broyd}_\tau(\tilde{G}, \nabla^2 f(x_+), u) \preceq [(1 + Mr)^2 \eta] \nabla^2 f(x_+),$$

*Proof.* Note that

$$\nabla^2 f(x_+) \stackrel{(4.4)}{\preceq} (1 + Mr) \nabla^2 f(x) \stackrel{(4.14)}{\preceq} (1 + Mr)G = \tilde{G},$$

and,

$$\tilde{G} = (1 + Mr)G \stackrel{(4.14)}{\preceq} (1 + Mr)\eta \nabla^2 f(x) \stackrel{(4.4)}{\preceq} (1 + Mr)^2 \eta \nabla^2 f(x_+).$$

Thus,

$$\nabla^2 f(x_+) \preceq \tilde{G} \preceq (1 + Mr)^2 \eta \nabla^2 f(x_+),$$

and the claim now follows from Lemma 2.2.  $\square$

Let us now make one more assumption about the function  $f$ . We assume that, with respect to the operator  $B$ , defined by (2.12), the function  $f$  is *strongly convex*, and its gradient is *Lipschitz continuous*, i.e. there exist  $\mu, L > 0$ , such that, for all  $x, y \in \mathbb{E}$ , we have

$$(4.16) \quad \mu B \preceq \nabla^2 f(x) \preceq LB.$$

*Remark 4.5.* In fact, for our purposes, it is enough to require that conditions (4.2), (4.16) hold only in a neighborhood of a solution, but, for the sake of simplicity, we do not do this.

We are ready to write down the scheme of our quasi-Newton methods. For simplicity, we assume that the constants  $M$  and  $L$  are available.

**Initialization:** Choose  $x_0 \in \mathbb{E}$ . Set  $G_0 = LB$ .

**For  $k \geq 0$  iterate:**

- (4.17)
1. Update  $x_{k+1} = x_k - G_k^{-1} \nabla f(x_k)$ .
  2. Compute  $r_k = \|x_{k+1} - x_k\|_{x_k}$  and set  $\tilde{G}_k = (1 + Mr_k)G_k$ .
  2. Choose  $u_k \in \mathbb{E}$  and  $\tau_k \in [0, 1]$ .
  4. Compute  $G_{k+1} = \text{Broyd}_{\tau_k}(\tilde{G}_k, \nabla^2 f(x_{k+1}), u_k)$ .

*Remark 4.6.* Similarly to Remark 3.1, in a practical implementation of scheme (4.17), one should work directly with the inverse operators  $G_k^{-1}$ , or with the Cholesky decomposition of  $G_k$ . Note that the correction step  $\tilde{G}_k = (1 + Mr_k)G_k$  does not affect the complexity of the iteration.

As before, we present two convergence results for scheme (4.17). The first one establishes linear convergence and can be seen as a generalization of Theorem 3.3. Note that for this result the directions  $u_k$  in the method (4.17) can be chosen arbitrarily.

**THEOREM 4.7.** *Suppose the initial point  $x_0$  is sufficiently close to the solution:*

$$(4.18) \quad M\lambda_f(x_0) \leq \frac{\ln \frac{3}{2} \frac{\mu}{L}}{4}.$$

Then, for all  $k \geq 0$ , we have

$$(4.19) \quad \nabla^2 f(x_k) \preceq G_k \preceq e^{2M \sum_{i=0}^{k-1} \lambda_f(x_i)} \frac{L}{\mu} \nabla^2 f(x_k) \preceq \frac{3L}{2\mu} \nabla^2 f(x_k),$$

and

$$(4.20) \quad \lambda_f(x_k) \leq \left(1 - \frac{\mu}{2L}\right)^k \lambda_f(x_0).$$

*Proof.* In view of (4.16), we have

$$\nabla^2 f(x_0) \preceq G_0 \preceq \frac{L}{\mu} \nabla^2 f(x_0).$$

Therefore, for  $k = 0$ , both (4.19) and (4.20) are satisfied.

Now let  $k \geq 0$ , and suppose (4.19), (4.20) have already been proved for all  $0 \leq k' \leq k$ . Denote  $\lambda_k \stackrel{\text{def}}{=} \lambda_f(x_k)$ ,  $r_k \stackrel{\text{def}}{=} \|x_{k+1} - x_k\|_{x_k}$ , and

$$(4.21) \quad \eta_k \stackrel{\text{def}}{=} e^{2M \sum_{i=0}^{k-1} \lambda_i} \frac{L}{\mu}.$$

Note that

$$(4.22) \quad M \sum_{i=0}^k \lambda_i \stackrel{(4.20)}{\leq} M\lambda_0 \sum_{i=0}^k \left(1 - \frac{\mu}{2L}\right)^i \leq \frac{2L}{\mu} M\lambda_0 \stackrel{(4.18)}{\leq} \frac{\ln \frac{3}{2}}{2}.$$

Applying Lemma 4.3, we obtain that

$$(4.23) \quad r_k \leq \lambda_k$$



and

$$(4.24) \quad \lambda_{k+1} \leq \left(1 + \frac{M\lambda_k}{2}\right) \frac{\eta_k^{-1} + \frac{M\lambda_k}{2}}{\eta_k} \lambda_k = \left(1 + \frac{M\lambda_k}{2}\right) \left(1 - \frac{1 - \frac{M\lambda_k}{2}}{\eta_k}\right) \lambda_k.$$

Note (using the inequality  $1 - t \geq e^{-2t}$ , valid at least for  $0 \leq t \leq \frac{1}{2}$ ) that

$$\frac{1 - \frac{M\lambda_k}{2}}{\eta_k} \geq e^{-M\lambda_k} \eta_k^{-1} \stackrel{(4.21)}{=} e^{-M\lambda_k - 2M \sum_{i=0}^{k-1} \lambda_i \frac{\mu}{L}} \geq e^{-2M \sum_{i=0}^k \lambda_i \frac{\mu}{L}} \stackrel{(4.22)}{\geq} \frac{2\mu}{3L},$$

and also (using  $\ln(1+t) \leq t$ , valid for  $t \geq 0$ ) that

$$\frac{M\lambda_k}{2} \stackrel{(4.18)}{\leq} \frac{\ln \frac{3}{2} \frac{\mu}{L}}{\frac{3}{8} \frac{\mu}{L}} \leq \frac{\mu}{16L}.$$

Hence,

$$\left(1 + \frac{M\lambda_k}{2}\right) \left(1 - \frac{1 - \frac{M\lambda_k}{2}}{\eta_k}\right) \leq \left(1 + \frac{\mu}{16L}\right) \left(1 - \frac{2\mu}{3L}\right) \leq 1 - \left(\frac{2}{3} - \frac{1}{16}\right) \frac{\mu}{L} \leq 1 - \frac{\mu}{2L}.$$

Consequently,

$$\lambda_{k+1} \stackrel{(4.24)}{\leq} \left(1 - \frac{\mu}{2L}\right) \lambda_k \stackrel{(4.20)}{\leq} \left(1 - \frac{\mu}{2L}\right)^{k+1} \lambda_0.$$

Finally, from Lemma 4.4, it follows that

$$(4.25) \quad \begin{aligned} \nabla^2 f(x_{k+1}) &\preceq G_{k+1} \preceq (1 + Mr_k)^2 \eta_k \nabla^2 f(x_{k+1}) \\ &\stackrel{(4.23)}{\preceq} (1 + M\lambda_k)^2 \eta_k \nabla^2 f(x_{k+1}) \preceq e^{2M\lambda_k} \eta_k \nabla^2 f(x_{k+1}) \\ &\stackrel{(4.21)}{=} e^{2M \sum_{i=0}^k \lambda_i \frac{L}{\mu}} \nabla^2 f(x_{k+1}) \stackrel{(4.22)}{\preceq} \frac{3L}{2\mu} \nabla^2 f(x_{k+1}). \end{aligned}$$

Thus, (4.19), (4.20) are valid for  $k' = k + 1$ , and we can continue by induction.  $\square$

Now let us analyze the greedy strategy. First, we analyze how the Hessian approximation measure (2.9) changes after one iteration.

LEMMA 4.8. *Let  $x \in \mathbb{E}$ , and let  $G : \mathbb{E} \rightarrow \mathbb{E}^*$  be a self-adjoint linear operator, such that be such that  $\nabla^2 f(x) \preceq G$ . Let  $x_+ \in \mathbb{E}$ , let  $r \stackrel{\text{def}}{=} \|x_+ - x\|_x$ , and let*

$$(4.26) \quad \tilde{G} \stackrel{\text{def}}{=} (1 + Mr)G.$$

Then, for any  $\tau \in [0, 1]$ , we have

$$\sigma_{x_+}(\text{Broyd}_\tau(\tilde{G}, \nabla^2 f(x_+), \bar{u}_{x_+}(G))) \leq \left(1 - \frac{\mu}{nL}\right) (1 + Mr)^2 \left(\sigma_x(G) + \frac{2nMr}{1 + Mr}\right).$$

*Proof.* We already know from Lemma 4.4 that  $\nabla^2 f(x_+) \preceq \tilde{G}$ . Also note that  $\bar{u}_{x_+}(\tilde{G}) = \bar{u}_{x_+}(G)$  (see (2.11)). Hence, by Theorem 2.5, we have

$$\sigma_{x_+}(\text{Broyd}_\tau(\tilde{G}, \nabla^2 f(x_+), \bar{u}_{x_+}(G))) \leq \left(1 - \frac{\mu}{nL}\right) \sigma_{x_+}(\tilde{G}).$$

Further,

$$\begin{aligned}
\sigma_{x_+}(\tilde{G}) &\stackrel{(2.9)}{=} \langle \nabla^2 f(x_+)^{-1}, \tilde{G} \rangle - n \stackrel{(4.26)}{=} (1+Mr) \langle \nabla^2 f(x_+)^{-1}, G \rangle - n \\
&\stackrel{(4.4)}{\leq} (1+Mr)^2 \langle \nabla^2 f(x)^{-1}, G \rangle - n \stackrel{(2.9)}{=} (1+Mr)^2 (\sigma_x(G) + n) - n \\
&= (1+Mr)^2 \sigma_x(G) + n((1+Mr)^2 - 1) \\
&= (1+Mr)^2 \sigma_x(G) + 2nMr \left(1 + \frac{Mr}{2}\right) \\
&\leq (1+Mr)^2 \left( \sigma_x(G) + \frac{2nMr}{1+Mr} \right).
\end{aligned}$$

The proof is now finished.  $\square$

Now we can prove superlinear convergence. In what follows, we assume that  $n \geq 2$ .

**THEOREM 4.9.** *Suppose that, in scheme (4.17), for each  $k \geq 0$  we take  $u_k = \bar{u}_{x_{k+1}}(G_k)$ . And suppose that the initial point  $x_0$  is sufficiently close to the solution:*

$$(4.27) \quad M\lambda_f(x_0) \leq \frac{\ln 2}{4(2n+1)} \frac{\mu}{L} \quad \left( \leq \frac{\ln \frac{3}{4}}{4} \frac{\mu}{L} \right).$$

Then, for all  $k \geq 0$ , we have

$$(4.28) \quad \nabla^2 f(x_k) \preceq G_k \preceq \left(1 + \left(1 - \frac{\mu}{nL}\right)^k \frac{2nL}{\mu}\right) \nabla^2 f(x_k),$$

and

$$(4.29) \quad \lambda_f(x_{k+1}) \leq \left(1 - \frac{\mu}{nL}\right)^k \frac{2nL}{\mu} \cdot \lambda_f(x_k).$$

*Proof.* Denote  $\lambda_k \stackrel{\text{def}}{=} \lambda_f(x_k)$  and  $\sigma_k \stackrel{\text{def}}{=} \sigma_{x_k}(G_k)$  for  $k \geq 0$ . In view of Theorem 4.7, the first relation in (4.28) is indeed true, and also

$$(4.30) \quad M \sum_{i=0}^k \lambda_i \leq M\lambda_0 \sum_{i=0}^k \left(1 - \frac{\mu}{2L}\right)^i \leq \frac{2L}{\mu} \lambda_0 \stackrel{(4.27)}{\leq} \frac{\ln 2}{2(2n+1)}.$$

for all  $k \geq 0$ .

Let us show by induction that, for all  $k \geq 0$ , we have

$$(4.31) \quad \sigma_k + 2nM\lambda_k \leq \theta_k.$$

where

$$(4.32) \quad \theta_k \stackrel{\text{def}}{=} \left(1 - \frac{\mu}{nL}\right)^k e^{2(2n+1)M \sum_{i=0}^{k-1} \lambda_i} \frac{nL}{\mu} \stackrel{(4.30)}{\leq} \left(1 - \frac{\mu}{nL}\right)^k \frac{2nL}{\mu}.$$

Indeed, since  $\nabla^2 f(x_0) \preceq G_0 \preceq \frac{L}{\mu} \nabla^2 f(x_0)$  (see (4.16)), we have

$$\begin{aligned}
\sigma_0 + 2nM\lambda_0 &\stackrel{(2.9)}{=} \langle \nabla^2 f(x_0)^{-1}, G_0 \rangle - n + 2nM\lambda_0 \\
&\leq \langle \nabla^2 f(x_0)^{-1}, \frac{L}{\mu} \nabla^2 f(x_0) \rangle - n + 2nM\lambda_0 \\
&\stackrel{(1.3)}{=} n \left( \frac{L}{\mu} - 1 \right) + 2nM\lambda_0 \stackrel{(4.27)}{\leq} n \left( \frac{L}{\mu} - 1 \right) + \frac{n \ln 2}{2(2n+1)} \leq \frac{nL}{\mu}.
\end{aligned}$$

Therefore, for  $k = 0$ , inequality (4.31) is satisfied. Now suppose that it is also satisfied for some  $k \geq 0$ . Since  $\nabla^2 f(x_k) \preceq G_k$ , we know that

$$G_k - \nabla^2 f(x_k) \stackrel{(1.4)}{\preceq} \sigma_k \nabla^2 f(x_k),$$

or, equivalently,

$$(4.33) \quad G_k \preceq (1 + \sigma_k) \nabla^2 f(x_k).$$

Therefore, applying Lemma 4.3, we obtain that

$$(4.34) \quad r_k \stackrel{\text{def}}{=} \|x_{k+1} - x_k\|_{x_k} \leq \lambda_k,$$

and

$$(4.35) \quad \begin{aligned} \lambda_{k+1} &\leq \left(1 + \frac{M\lambda_k}{2}\right) \frac{\sigma_k + \frac{M\lambda_k}{2}}{1 + \sigma_k} \lambda_k \leq \left(1 + \frac{M\lambda_k}{2}\right) (\sigma_k + 2nM\lambda_k) \lambda_k \\ &\stackrel{(4.31)}{\leq} \left(1 + \frac{M\lambda_k}{2}\right) \theta_k \lambda_k \leq e^{\frac{M\lambda_k}{2}} \theta_k \lambda_k \leq e^{2M\lambda_k} \theta_k \lambda_k. \end{aligned}$$

Further, by Lemma 4.8, we have

$$\begin{aligned} \sigma_{k+1} &\leq \left(1 - \frac{\mu}{nL}\right) (1 + Mr_k)^2 \left(\sigma_k + \frac{2nMr_k}{1 + Mr_k}\right) \\ &\stackrel{(4.34)}{\leq} \left(1 - \frac{\mu}{nL}\right) (1 + M\lambda_k)^2 \left(\sigma_k + \frac{2nM\lambda_k}{1 + M\lambda_k}\right) \\ &\leq \left(1 - \frac{\mu}{nL}\right) (1 + M\lambda_k)^2 (\sigma_k + 2nM\lambda_k) \\ &\stackrel{(4.31)}{\leq} \left(1 - \frac{\mu}{nL}\right) (1 + M\lambda_k)^2 \theta_k \leq \left(1 - \frac{\mu}{nL}\right) e^{2M\lambda_k} \theta_k. \end{aligned}$$

Note that  $\frac{1}{2} \leq 1 - \frac{\mu}{nL}$  since  $n \geq 2$ . Therefore,

$$\begin{aligned} \sigma_{k+1} + 2nM\lambda_{k+1} &\leq \left(1 - \frac{\mu}{nL}\right) e^{2M\lambda_k} \theta_k + e^{2M\lambda_k} \theta_k 2nM\lambda_k \\ &\leq \left(1 - \frac{\mu}{nL}\right) e^{2M\lambda_k} \theta_k + \left(1 - \frac{\mu}{nL}\right) e^{2M\lambda_k} \theta_k 4nM\lambda_k \\ &= \left(1 - \frac{\mu}{nL}\right) e^{2M\lambda_k} (1 + 4nM\lambda_k) \theta_k \\ &\leq \left(1 - \frac{\mu}{nL}\right) e^{2(2n+1)M\lambda_k} \theta_k \stackrel{(4.32)}{=} \theta_{k+1}. \end{aligned}$$

Thus, (4.31) is proved.

Let us fix now some  $k \geq 0$ . Since  $\lambda_k \geq 0$ , we have

$$\sigma_k \leq \sigma_k + 2M\lambda_k \stackrel{(4.31)}{\leq} \theta_k \stackrel{(4.32)}{\leq} \left(1 - \frac{\mu}{nL}\right)^k \frac{2nL}{\mu}.$$

This proves the second relation in (4.28) in view of (4.33). Finally,

$$\lambda_{k+1} \stackrel{(4.35)}{\leq} e^{2M\lambda_k} \theta_k \lambda_k \leq e^{2(2n+1)M\lambda_k} \theta_k \lambda_k \stackrel{(4.32)}{=} \frac{\theta_{k+1}}{1 - \frac{\mu}{nL}} \lambda_k \stackrel{(4.32)}{\leq} \left(1 - \frac{\mu}{nL}\right)^k \frac{2nL}{\mu} \lambda_k,$$

and we obtain (4.29).  $\square$

Similarly to the quadratic case, combining Theorem 4.7 with Theorem 4.9, we obtain the following final efficiency estimate:

$$\lambda_f(x_{k_0+k}) \leq \left(1 - \frac{\mu}{nL}\right)^{\frac{k(k-1)}{2}} \left(\frac{1}{2}\right)^k \left(1 - \frac{\mu}{2L}\right)^{k_0} \lambda_f(x_0), \quad k \geq 0,$$

where  $k_0 \leq \frac{nL}{\mu} \ln \frac{2nL}{\mu}$ .

## 5. Numerical Experiments.

**5.1. Regularized Log-Sum-Exp.** In this section, we present preliminary computational results for greedy quasi-Newton methods, applied to the following test function<sup>3</sup>:

$$(5.1) \quad f(x) \stackrel{\text{def}}{=} \ln \left( \sum_{j=1}^m e^{\langle c_j, x \rangle - b_j} \right) + \frac{1}{2} \sum_{j=1}^m \langle c_j, x \rangle^2 + \frac{\gamma}{2} \|x\|^2, \quad x \in \mathbb{R}^n,$$

where  $c_1, \dots, c_m \in \mathbb{R}^n$ ,  $b_1, \dots, b_m \in \mathbb{R}$ , and  $\gamma > 0$ .

We compare scheme (4.17) (which realizes GrDFP, GrBFGS and GrSR1, depending on the choice of  $\tau_k$ ) with the usual gradient method (GM)<sup>4</sup> and standard quasi-Newton methods DFP, BFGS and SR1.

All the standard methods need access only to the gradient of function  $f$ :

$$(5.2) \quad \nabla f(x) = g(x) + \sum_{j=1}^m \langle c_j, x \rangle c_j + \gamma x, \quad g(x) \stackrel{\text{def}}{=} \sum_{j=1}^m \pi_j(x) c_j,$$

where

$$\pi_j(x) \stackrel{\text{def}}{=} \frac{e^{\langle c_j, x \rangle - b_j}}{\sum_{j'=1}^m e^{\langle c_{j'}, x \rangle - b_{j'}}} \in [0, 1], \quad j = 1, \dots, m.$$

Note that, for a given point  $x \in \mathbb{R}^n$ ,  $\nabla f(x)$  can be computed in  $O(mn)$  operations.

For greedy methods, to implement the Hessian approximation update, at every iteration, we need to carry out some additional operations with the Hessian

$$(5.3) \quad \begin{aligned} \nabla^2 f(x) &= \sum_{j=1}^m \pi_j(x) c_j c_j^T - g(x) g(x)^T + \sum_{j=1}^m c_j c_j^T + \gamma I \\ &= \sum_{j=1}^m (\pi_j(x) + 1) c_j c_j^T - g(x) g(x)^T + \gamma I. \end{aligned}$$

Namely, given a point  $x \in \mathbb{R}^n$ , we need to be able to perform the following two actions:

- For all  $1 \leq i \leq n$ , compute the values

$$\langle \nabla^2 f(x) e_i, e_i \rangle \stackrel{(5.3)}{=} \sum_{j=1}^m (\pi_j(x) + 1) \langle c_j, e_i \rangle^2 - \langle g(x), e_i \rangle^2 + \gamma,$$

where  $e_1, \dots, e_n$  are the basis vectors.

- For a given direction  $h \in \mathbb{R}^n$ , compute the Hessian-vector product

$$\nabla^2 f(x) h \stackrel{(5.3)}{=} \sum_{j=1}^m (\pi_j(x) + 1) \langle c_j, h \rangle c_j - \langle g(x), h \rangle g(x) + \gamma h.$$

Let us take the basis  $e_1, \dots, e_n$ , comprised of the standard coordinate directions:

$$(5.4) \quad e_i \stackrel{\text{def}}{=} (0, \dots, 0, 1, 0, \dots, 0)^T, \quad 1 \leq i \leq n.$$

Then, both the above operations have a cost of  $O(mn)$ . Thus, the cost of one iteration for all the methods under our consideration is comparable.

<sup>3</sup>Note that we work in the space  $\mathbb{E} = \mathbb{R}^n$  and identify  $\mathbb{E}^*$  with  $\mathbb{E}$  in such a way that  $\langle \cdot, \cdot \rangle$  is the standard dot product, and  $\| \cdot \|$  is the standard Euclidean norm. Linear operators from  $\mathbb{E}$  to  $\mathbb{E}^*$  are identified with  $n \times n$  matrices.

<sup>4</sup>For GM, we use the constant step size  $\frac{1}{L}$ , where  $L$  is the corresponding estimate of the Lipschitz constant of the gradient, given by (5.5).

Note that for basis (5.4), the matrix  $B$ , defined by (2.12), is the identity matrix:

$$B = I.$$

Hence, the Lipschitz constant of the gradient of  $f$  with respect to  $B$  can be taken as follows (see (5.3)):

$$(5.5) \quad L = 2 \sum_{j=1}^m \|c_j\|^2 + \gamma.$$

All quasi-Newton methods in our comparison start from the same initial Hessian approximation  $G_0 = LB$ , and use unit step sizes.

Finally, for greedy quasi-Newton methods, we also need to provide an estimate of the strong self-concordancy parameter. Note that, with respect to the operator  $\sum_{j=1}^m c_j c_j^T$ , the function  $f$  is 1-strongly convex and its Hessian is 2-Lipschitz continuous (see e.g. [33, Ex. 1]). Hence, in view of Example 4.1, the strong self-concordancy parameter can be chosen as follows:

$$M = 2.$$

The data, defining the test function (5.1), is randomly generated in the following way. First, we generate a collection of random vectors

$$\hat{c}_1, \dots, \hat{c}_m$$

with entries, uniformly distributed in the interval  $[-1, 1]$ . Then we generate  $b_1, \dots, b_m$  from the same distribution. Using this data, we form a preliminary function

$$\hat{f}(x) \stackrel{\text{def}}{=} \ln \left( \sum_{j=1}^m e^{\langle \hat{c}_j, x \rangle - b_j} \right),$$

and finally define

$$c_j \stackrel{\text{def}}{=} \hat{c}_j - \nabla \hat{f}(0), \quad j = 1, \dots, m.$$

Note that by construction

$$\nabla f(0) \stackrel{(5.2)}{=} \frac{1}{\sum_{j=1}^m e^{-b_j}} \sum_{j=1}^m e^{-b_j} (\hat{c}_j - \nabla \hat{f}(0)) = 0,$$

so the unique minimizer of our test function (5.1) is  $x^* = 0$ . The starting point  $x_0$  for all methods is the same and generated randomly from the uniform distribution on the standard Euclidean sphere of radius  $1/n$  (this choice is motivated by (4.27)) centered at the minimizer.

Thus, our test function (5.1) has three parameters: the dimension  $n$ , the number  $m$  of linear functions, and the regularization coefficient  $\gamma$ . Let us present computational results for different values of these parameters. The termination criterion for all methods is  $f(x_k) - f(x^*) \leq \epsilon(f(x_0) - f(x^*))$ .

In the tables below, for each method, we display the number of iterations until its termination. The minus sign ( $-$ ) means that the method has not been able to achieve the required accuracy after  $1000n$  iterations.

**Table 1:**  $n = m = 50, \gamma = 1$ 

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	79	4	4	3	45	35	34
$10^{-3}$	1812	777	57	18	342	57	52
$10^{-5}$	5263	1866	107	29	738	72	58
$10^{-7}$	8873	2836	158	39	917	83	63
$10^{-9}$	12532	3911	203	48	1028	93	67

**Table 2:**  $n = m = 50, \gamma = 0.1$ 

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	76	4	4	3	44	33	33
$10^{-3}$	2732	1278	78	23	512	70	56
$10^{-5}$	29785	12923	254	57	3850	126	72
$10^{-7}$	—	23245	346	74	6794	169	81
$10^{-9}$	—	32441	381	79	8216	204	87

**Table 3:**  $n = m = 250, \gamma = 1$ 

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	444	4	4	3	214	158	157
$10^{-3}$	10351	4743	98	21	3321	264	251
$10^{-5}$	73685	31468	288	55	15637	350	274
$10^{-7}$	159391	58138	450	82	21953	413	296
$10^{-9}$	249492	85218	627	110	25500	464	314

**Table 4:**  $n = m = 250, \gamma = 0.1$ 

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	442	4	4	3	209	155	155
$10^{-3}$	9312	4175	91	21	2686	258	251
$10^{-5}$	207978	102972	488	87	60461	556	346
$10^{-7}$	—	—	1003	170	147076	792	391
$10^{-9}$	—	—	1407	233	212100	976	419

We see that all quasi-Newton methods outperform the gradient method and demonstrate superlinear convergence (from some moment, the difference in the number of iterations between successive rows in the table becomes smaller and smaller). Among quasi-Newton methods (both the standard and the greedy ones), SR1 is always better than BFGS, while DFP is significantly worse than the other two. At the first few iterations, the greedy methods loose to the standard ones, but later they catch up. However, the classical SR1 method always remains the best. Nevertheless, the greedy methods are quite competitive.

Now let us look at the quality of Hessian approximations, produced by the quasi-Newton methods. In the tables below, we display the desired accuracy  $\epsilon$  vs the final Hessian approximation error (defined as the operator norm of  $G_k - \nabla^2 f(x_k)$ , measured with respect to  $\nabla^2 f(x_k)$ ). We look at the same problems as in Table 1 and Table 3.

**Table 5:**  $n = m = 50, \gamma = 1$ 

$\epsilon$	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-0}$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$
$10^{-1}$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$2.7 \cdot 10^3$	$1.5 \cdot 10^3$	$1.5 \cdot 10^3$
$10^{-3}$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.2 \cdot 10^3$	$1.2 \cdot 10^1$	$3.8 \cdot 10^0$
$10^{-5}$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$2.1 \cdot 10^2$	$7.2 \cdot 10^0$	$2.6 \cdot 10^0$
$10^{-7}$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$9.1 \cdot 10^1$	$5.6 \cdot 10^0$	$2.2 \cdot 10^0$
$10^{-9}$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$1.6 \cdot 10^3$	$5.2 \cdot 10^1$	$4.1 \cdot 10^0$	$1.8 \cdot 10^0$

**Table 6:**  $n = m = 250, \gamma = 1$ 

$\epsilon$	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-0}$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$
$10^{-1}$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$7.1 \cdot 10^4$	$3.8 \cdot 10^4$	$3.9 \cdot 10^4$
$10^{-3}$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$6.8 \cdot 10^4$	$6.6 \cdot 10^1$	$1.7 \cdot 10^1$
$10^{-5}$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$9.4 \cdot 10^3$	$3.7 \cdot 10^1$	$1.2 \cdot 10^1$
$10^{-7}$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$3.1 \cdot 10^3$	$2.8 \cdot 10^1$	$9.7 \cdot 10^0$
$10^{-9}$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$4.1 \cdot 10^4$	$1.7 \cdot 10^3$	$2.2 \cdot 10^1$	$7.3 \cdot 10^0$

As we can see from these tables, for standard quasi-Newton methods the Hessian approximation error always stays at the initial level. In contrast, for the greedy ones, it decreases relatively fast (especially for GrBFGS and GrSR1). Note also that sometimes the initial residual slightly increases at the first several iterations (which is noticeable only for GrDFP). This happens due to the fact that the objective function is non-quadratic, and we apply the correction strategy.

Note that in all the above tests we have used the same values for the parameters  $n$  and  $m$ . Let us briefly illustrate what happens when, for example,  $m > n$ .

**Table 7:**  $n = 50, m = 100, \gamma = 0.1$ 

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	84	4	4	3	46	37	37
$10^{-3}$	897	316	32	11	183	53	52
$10^{-5}$	2421	833	67	19	334	63	58
$10^{-7}$	4087	1304	98	25	423	71	62
$10^{-9}$	5810	1859	132	32	473	78	66

**Table 8:**  $n = 50, m = 200, \gamma = 0.1$ 

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	108	4	4	3	45	46	46
$10^{-3}$	479	101	17	7	97	53	52
$10^{-5}$	1059	338	39	12	154	62	59
$10^{-7}$	1817	615	62	18	206	67	64
$10^{-9}$	2659	807	81	21	234	73	68

Comparing these tables with Table 2, we see that, with the increase of  $m$ , all the methods generally terminate faster. However, the overall picture is still the same as before. The results for  $m < n$  are similar, so we do not include them.

Finally, let us present the results for the *randomized* version of scheme (4.17), in which, at every step, we select the update direction uniformly at random from the

standard Euclidean sphere:

$$(5.6) \quad u_k \sim \text{Unif}(\mathcal{S}^{n-1}),$$

where  $\mathcal{S}^{n-1} \stackrel{\text{def}}{=} \{x \in \mathbb{R}^n : \|x\| = 1\}$ . We call the corresponding methods RaDFP, RaBFGS and RaSR1.

**Table 9:**  $n = m = 50, \gamma = 1$

$\epsilon$	RaDFP	RaBFGS	RaSR1
$10^{-1}$	35	29	34
$10^{-3}$	566	102	64
$10^{-5}$	1156	125	77
$10^{-7}$	1481	142	85
$10^{-9}$	1698	156	91

**Table 10:**  $n = m = 250, \gamma = 1$

$\epsilon$	RaDFP	RaBFGS	RaSR1
$10^{-1}$	261	144	158
$10^{-3}$	4276	366	287
$10^{-5}$	19594	517	346
$10^{-7}$	33293	619	376
$10^{-9}$	41177	698	396

It is instructive to compare these tables with Table 1 and Table 3, which contain the results for the greedy methods on the same problems. We see that the randomized methods are slightly slower than the greedy ones. However, the difference is not really significant, and, what is especially interesting, the randomized methods do not loose superlinear convergence.

**5.2. Logistic Regression.** Now let us consider another test function, namely  $l_2$ -regularized logistic regression, which is popular in the field of machine learning:

$$(5.7) \quad f(x) \stackrel{\text{def}}{=} \sum_{j=1}^m \ln(1 + e^{-b_j \langle c_j, x \rangle}) + \frac{\gamma}{2} \|x\|^2, \quad x \in \mathbb{R}^n,$$

where  $c_1, \dots, c_m \in \mathbb{R}^n$ ,  $b_1, \dots, b_m \in \{-1, 1\}$ , and  $\gamma > 0$ .

Note that the structure of the function (5.7) is similar to the one of (5.1). In particular, both the diagonal of the Hessian and the Hessian-vector product for this function can be computed with the similar complexity of that for computing the gradient. Also it can be shown that the Lipschitz constant of the gradient of  $f$  can be chosen in accordance with (5.5) but with the coefficient  $\frac{1}{4}$  instead of 2.

We follow the same experiment design as before with only a couple of differences. First, instead of generating the data, defining the function (5.7), artificially, now we take it from the LIBSVM collection of real-world data sets for binary classification problems<sup>5</sup> [27]. Second, we have found it better in practice not to apply the correction strategy in the greedy methods (i.e. simply set  $\tilde{G}_k = G_k$  in scheme (4.17)). This is the only heuristic that we use. For the regularization coefficient, we always use the value  $\gamma = 1$ , which is a standard choice.

Let us look at the results.

**Table 11:** Data set *ijcnn1* ( $n = 22, m = 49990$ )

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	246	43	8	6	25	19	18
$10^{-3}$	1925	672	45	16	71	25	23
$10^{-5}$	5123	2007	85	25	145	32	23
$10^{-7}$	8966	2738	102	29	192	38	23
$10^{-9}$	12815	3269	118	33	215	43	24

<sup>5</sup>The original labels  $b_i$  in the *mushrooms* data set are “1” and “2” instead of “1” and “-1”. Therefore, we renamed in advance the class label “2” into “-1”.



**Table 12:** Data set *mushrooms* ( $n = 112, m = 8124$ )

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	4644	936	15	6	230	83	82
$10^{-3}$	77103	30594	105	24	1185	149	113
$10^{-5}$	—	58221	166	34	1700	170	113
$10^{-7}$	—	83740	217	42	1945	182	113
$10^{-9}$	—	107471	257	48	2088	194	114

**Table 13:** Data set *a9a* ( $n = 123, m = 32561$ )

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	160	32	10	6	110	81	81
$10^{-3}$	18690	9229	145	38	2203	127	117
$10^{-5}$	—	79014	411	88	23715	316	123
$10^{-7}$	—	—	553	113	35700	441	124
$10^{-9}$	—	—	581	118	38285	475	124

**Table 14:** Data set *w8a* ( $n = 300, m = 49749$ )

$\epsilon$	GM	DFP	BFGS	SR1	GrDFP	GrBFGS	GrSR1
$10^{-1}$	10148	3531	35	10	694	300	300
$10^{-3}$	194813	86315	178	34	1426	307	301
$10^{-5}$	—	188561	300	54	1849	327	301
$10^{-7}$	—	255224	387	68	2036	339	301
$10^{-9}$	—	264346	399	69	2057	340	301

As we can see, the general picture is the same as for the previous test function. In particular, the DFP update is always much worse than BFGS and SR1. The greedy methods are competitive with the standard ones and often outperform them for high values of accuracy.

**6. Discussion.** We have presented the greedy quasi-Newton methods, that are based on the updating formulas from the Broyden family and use greedily selected basis vectors for updating Hessian approximations. For these methods, we have established explicit non-asymptotic rate of local superlinear convergence for the iterates and also a linear convergence for the deviations of Hessian approximations from the correct Hessians.

Clearly, there is a number of open questions. First, at every iteration, our methods need to compute the greedily selected basis vector. This requires additional information beyond just the gradient of the objective function (such as the diagonal of the Hessian). However, many problems, that arise in applications, possess certain structure (separable, sparse, etc.), for which the corresponding computations have a cost similar to that of the gradient evaluation (such as the test function in our experiments). Nevertheless, ideally it is desirable to get rid of the necessity in this auxiliary information at all. A natural idea might be to replace the greedy strategy with a *randomized* one. Indeed, as can be seen from our experiments, the corresponding scheme (4.17), (5.6) demonstrate almost the same performance as the greedy one. Therefore, one can expect that it should be possible to establish similar theoretical results about its superlinear convergence. Nevertheless, at the moment, we do not know how to do this. Although it is not difficult to show that, in terms of *expectations*, the randomized strategy still preserves the linear convergence of Hessian approximations

(see [30]), it is not clear how to proceed after this in proving the superlinear convergence of the iterates, even in the quadratic case. The main difficulty, arising in the analysis, is that, at some moment, one needs to take the expectation of the product of random variables with known expectations, but the random variables themselves are *non-independent*.

Second, we have analyzed together a whole class of Hessian approximation updates by essentially upper bounding all its members via the worst one—DFP. Thus, all the efficiency guarantees, that we have established, might be too pessimistic for other members of this class such as BFGS, and especially SR1. Indeed, in our experiments, we have seen that the convergence properties of these three methods might differ quite significantly. It is therefore desirable to refine our current analysis and obtain separate estimates for different updates.

Third, note that our current results do not prove anything about the rate of superlinear convergence of the standard quasi-Newton methods. Of course, it would be interesting to obtain the corresponding estimates and compare them to the ones, that we have established in this work.

Finally, apart from the quadratic case, we have not addressed at all the question of global convergence.

In any case, we believe that the ideas and the theoretical analysis, presented in this paper, will be useful for future advances in the theory of quasi-Newton methods.

**Acknowledgments.** The authors would like to thank two anonymous referees for their useful comments and suggestions.

#### REFERENCES

- [1] W. Davidon. Variable metric method for minimization. Argonne National Laboratory Research and Development Report 5990 (1959).
- [2] R. Fletcher and M. Powell. A rapidly convergent descent method for minimization. *Computer Journal*, **6**(2), 163-168 (1963).
- [3] C. Broyden. Quasi-Newton methods and their application to function minimization. *Mathematics of Computation*, **21**(99), 368-381 (1967).
- [4] W. Davidon. Variance algorithm for minimization. *Computer Journal*, **10**(4), 406-410 (1968).
- [5] D. Goldfarb. Sufficient conditions for the convergence of a variable metric algorithm. *Optimization*, ed. R. Fletcher, 273-281, Academic Press, London (1969).
- [6] C. Broyden. The convergence of a class of double-rank minimization algorithms: 1. General considerations. *IMA Journal of Applied Mathematics*, **6**(1), 76-90 (1970).
- [7] C. Broyden. The convergence of a class of double-rank minimization algorithms: 2. The new algorithm. *IMA Journal of Applied Mathematics*, **6**(3), 222-231 (1970).
- [8] R. Fletcher. A new approach to variable metric algorithms. *Computer Journal*, **13**(3), 317-322 (1970).
- [9] D. Goldfarb. A family of variable-metric methods derived by variational means. *Mathematics of Computation*, **24**(109), 23-26 (1970).
- [10] D. Shanno. Conditioning of quasi-Newton methods for function minimization. *Mathematics of Computation*, **24**(111), 647-656 (1970).
- [11] M. Powell. On the convergence of the variable metric algorithm. *IMA Journal of Applied Mathematics*, **7**(1), 21-36 (1971).
- [12] C. Broyden, J. Dennis, and J. Moré. On the local and superlinear convergence of quasi-Newton methods. *IMA Journal of Applied Mathematics*, **12**(3), 223-245 (1973).
- [13] J. Dennis and J. Moré. A characterization of superlinear convergence and its application to quasi-Newton methods. *Mathematics of Computation*, **28**(126), 549-560 (1974).
- [14] J. Dennis and J. Moré. Quasi-Newton methods, motivation and theory. *SIAM Review*, **19**(1), 46-89 (1977).
- [15] B. Pshenichnyi and I. Danilin. Numerical methods in extremal problems. *Mir Publishers* (1978).
- [16] A. Stachurski. Superlinear convergence of Broyden's bounded  $\theta$ -class of methods. *Mathematical Programming*, **20**(1), 196-212 (1981).

- [17] A. Griewank and P. Toint. Local convergence analysis for partitioned quasi-Newton updates. *Numerische Mathematik*, **39**(3), 429-448 (1982).
- [18] R. Byrd, J. Nocedal, and Y. Yuan. Global convergence of a class of quasi-Newton methods on convex problems. *SIAM Journal on Numerical Analysis*, **24**(5), 1171-1190 (1987).
- [19] R. Byrd and J. Nocedal. A tool for the analysis of quasi-Newton methods with application to unconstrained minimization. *SIAM Journal on Numerical Analysis*, **26**(3), 727-739 (1989).
- [20] A. Conn, N. Gould, and P. Toint. Convergence of quasi-Newton matrices generated by the symmetric rank one update. *Mathematical Programming*, **50**(1-3), 177-195 (1991).
- [21] J. Engels and H. Martínez. Local and superlinear convergence for partially known quasi-Newton methods. *SIAM Journal on Optimization*, **1**(1), 42-56 (1991).
- [22] Y. Nesterov and A. Nemirovskii. Interior-point polynomial algorithms in convex programming. *SIAM*, **13** (1994).
- [23] H. Yabe and N. Yamaki. Local and superlinear convergence of structured quasi-Newton methods for nonlinear optimization. *Journal of the Operations Research Society of Japan*, **39**(4), 541-557 (1996).
- [24] Z. Wei, G. Yu, G. Yuan, and Z. Lian. The superlinear convergence of a modified BFGS-type method for unconstrained optimization. *Computational Optimization and Applications*, **29**(3), 315-332 (2004).
- [25] J. Nocedal and S. Wright. Numerical optimization. *Springer Science & Business Media* (2006).
- [26] H. Yabe, H. Ogasawara, and M. Yoshino. Local and superlinear convergence of quasi-Newton methods based on modified secant conditions. *Journal of Computational and Applied Mathematics*, **205**(1), 617-632 (2007).
- [27] C. Chang and C. Lin. LIBSVM: A library for support vector machines. *ACM transactions on intelligent systems and technology (TIST)*, **2**(3), 1-27 (2011).
- [28] A. Lewis and M. Overton. Nonsmooth optimization via quasi-Newton methods. *Mathematical Programming*, **141**(1-2), 135-163 (2013).
- [29] R. Gower, D. Goldfarb, and P. Richtárik. Stochastic block BFGS: squeezing more curvature out of data. *International Conference on Machine Learning*, 1869-1878 (2016).
- [30] R. Gower and P. Richtárik. Randomized quasi-Newton updates are linearly convergent matrix inversion algorithms. *SIAM Journal on Matrix Analysis and Applications*, **38**(4), 1380-1409 (2017).
- [31] A. Mokhtari, M. Eisen, and A. Ribeiro. IQN: An incremental quasi-Newton method with local superlinear convergence rate. *SIAM Journal on Optimization*, **28**(2), 1670-1698 (2018).
- [32] Y. Nesterov. Lectures on convex optimization. *Springer*, **137** (2018).
- [33] N. Doikov and Y. Nesterov. Minimizing uniformly convex functions by cubic regularization of Newton method. *arXiv*, 1905.02671 (2019)
- [34] T. Sun and Q. Tran-Dinh. Generalized self-concordant functions: a recipe for Newton-type methods. *Mathematical Programming*, **178**, 145-213 (2019).
- [35] W. Gao and D. Goldfarb. Quasi-Newton methods: superlinear convergence without line searches for self-concordant functions. *Optimization Methods and Software*, **34**(1), 194-217 (2019).
- [36] D. Kovalev, R. Gower, P. Richtárik, and A. Rogozin. Fast linear convergence of randomized BFGS. *arXiv preprint arXiv:2002.11337* (2020).