

The strong topology of ω -plurisubharmonic functions

Antonio Trusiani*

Abstract

On (X, ω) compact Kähler manifold, given a model type envelope $\psi \in PSH(X, \omega)$ (i.e. a singularity type) we prove that the Monge-Ampère operator is a homeomorphism between the set of ψ -relative finite energy potentials and the set of ψ -relative finite energy measures endowed with their strong topologies given as the coarsest refinements of the weak topologies such that the relative energies become continuous. Moreover, given a totally ordered family \mathcal{A} of model type envelopes with positive total mass representing different singularities types, the sets $X_{\mathcal{A}}, Y_{\mathcal{A}}$ given respectively as the union of all ψ -relative finite energy potentials and of all ψ -relative finite energy measures varying $\psi \in \overline{\mathcal{A}}$ have two natural strong topologies which extends the strong topologies on each component of the unions. We show that the Monge-Ampère operator produces a homeomorphism between $X_{\mathcal{A}}$ and $Y_{\mathcal{A}}$. As an application we also prove the strong stability of a sequence of solutions of complex Monge-Ampère equations when the measures have uniformly L^p -bounded densities for $p > 1$ and the prescribed singularities are totally ordered.

Keywords: Complex Monge-Ampère equations, compact Kähler manifolds, quasi-psh functions.

2020 Mathematics subject classification: 32W20 (primary); 32U05, 32Q15 (secondary).

1 Introduction

Let (X, ω) be a compact Kähler manifold where ω is a fixed Kähler form, and let \mathcal{H}_{ω} denote the set of all Kähler potentials, i.e. all $\varphi \in C^{\infty}$ such that $\omega + dd^c\varphi$ is a Kähler form, the pioneering work of Yau ([26]) shows that the Monge-Ampère operator

$$MA_{\omega} : \mathcal{H}_{\omega, norm} \longrightarrow \left\{ dV \text{ volume form} : \int_X dV = \int_X \omega^n \right\}, \quad (1)$$

$MA_{\omega}(\varphi) := (\omega + dd^c\varphi)^n$ is a bijection, where for any subset $A \subset PSH(X, \omega)$ of all ω -plurisubharmonic functions we use the notation $A_{norm} := \{u \in A : \sup_X u = 0\}$. Note that the assumption on the total mass of the volume forms in (1) is necessary since $\mathcal{H}_{\omega, norm}$ represents all Kähler forms in the cohomology class $\{\omega\}$ and the quantity $\int_X \omega^n$ is cohomological.

In [18] the authors extended the Monge-Ampère operator using the *non-pluripolar product* (as denominated successively in [7]) and the bijection (1) to

$$MA_{\omega} : \mathcal{E}_{norm}(X, \omega) \longrightarrow \left\{ \mu \text{ non-pluripolar positive measure} : \mu(X) = \int_X \omega^n \right\} \quad (2)$$

where $\mathcal{E}(X, \omega) := \{u \in PSH(X, \omega) : \int_X MA_{\omega}(u) = \int_X MA_{\omega}(0)\}$ is the set of all ω -psh functions with full Monge-Ampère mass.

The set $PSH(X, \omega)$ is naturally endowed with the L^1 -topology which we will call *weak*, but the Monge-Ampère operator in (2) is not continuous even if the set of measures is endowed with the weak topology. Thus in [3], setting $V_0 := \int_X MA_{\omega}(0)$, two strong topologies were respectively introduced for

$$\begin{aligned} \mathcal{E}^1(X, \omega) &:= \{u \in \mathcal{E}(X, \omega) : E(u) > -\infty\} \\ \mathcal{M}^1(X, \omega) &:= \left\{ V_0 \mu : \mu \text{ is a probability measure satisfying } E^*(\mu) < +\infty \right\} \end{aligned}$$

as the coarsest refinements of the weak topologies such that respectively the Monge-Ampère energy $E(u)$ ([1], [2], [7]) and the energy for probability measures E^* ([4], [3]) becomes continuous. The map

$$MA_{\omega} : (\mathcal{E}_{norm}^1(X, \omega), strong) \longrightarrow (\mathcal{M}^1(X, \omega), strong) \quad (3)$$

*antonio.trusiani91@gmail.com;

is then a homeomorphism. Later Darvas ([12]) showed that actually $(\mathcal{E}^1(X, \omega), \text{strong})$ coincides with the metric closure of \mathcal{H}_ω endowed with the Finsler metric $|f|_{1, \varphi} := \int_X |f| MA_\omega(\varphi)$, $\varphi \in \mathcal{H}_\omega$, $f \in T_\varphi \mathcal{H}_\omega \simeq C^\infty(X)$ and associated distance

$$d(u, v) := E(u) + E(v) - 2E(P_\omega(u, v))$$

where $P_\omega(u, v)$ is the rooftop envelope given basically as the largest ω -psh function bounded above by $\min(u, v)$ ([22]). This metric topology has played an important role in the last decade to characterize the existence of special metrics ([16], [5], [9], [10], [11]).

It is also important and natural to solve complex Monge-Ampère equations requiring that the solutions have some prescribed behavior, for instance along a divisor.

We first need to recall that on $PSH(X, \omega)$ there is a natural partial order \preceq given as $u \preceq v$ if $u \leq v + O(1)$, and the total mass through the Monge-Ampère operator respects such partial order, i.e. $V_u := \int_X MA_\omega(u) \leq V_v$ if $u \preceq v$ ([7], [24]). Thus in [13] the authors introduced the ψ -relative analogs of the sets $\mathcal{E}(X, \omega)$, $\mathcal{E}^1(X, \omega)$ for $\psi \in PSH(X, \omega)$ fixed as

$$\begin{aligned} \mathcal{E}(X, \omega, \psi) &:= \{u \in PSH(X, \omega) : u \preceq \psi \text{ and } V_u = V_\psi\} \\ \mathcal{E}^1(X, \omega, \psi) &:= \{u \in \mathcal{E}(X, \omega, \psi) : E_\psi(u) > -\infty\} \end{aligned}$$

where E_ψ is the ψ -relative energy, and they proved that

$$MA_\omega : \mathcal{E}_{norm}(X, \omega, \psi) \longrightarrow \left\{ \mu \text{ non-pluripolar positive measure} : \mu(X) = V_\psi \right\} \quad (4)$$

is a bijection if and only if ψ , up to a bounded function, is a *model type envelope*, i.e. $\psi = (\lim_{C \rightarrow +\infty} P(\psi + C, 0))^*$, satisfying $V_\psi > 0$ (the star is for the upper semicontinuous regularization). There are plenty of these functions, for instance to any ω -psh function ψ with analytic singularities is associated a unique model type envelope. We denote by \mathcal{M} the set of all model type envelopes and with \mathcal{M}^+ those elements ψ such that $V_\psi > 0$.

Letting $\psi \in \mathcal{M}^+$, in [23], we proved that $\mathcal{E}^1(X, \omega, \psi)$ can be endowed with a natural metric topology given by the complete distance $d(u, v) := E_\psi(u) + E_\psi(v) - 2E_\psi(P_\omega(u, v))$.

Analogously to E^* , we introduce in section §5 a natural ψ -relative energy for probability measures E_ψ^* , thus the set

$$\mathcal{M}^1(X, \omega, \psi) := \{V_\psi \mu : \mu \text{ is a probability measure satisfying } E_\psi^*(\mu) < +\infty\}$$

can be endowed with its strong topology given as the coarsest refinement of the weak topology such that E_ψ^* becomes continuous.

Theorem A. *Let $\psi \in \mathcal{M}^+$. Then*

$$MA_\omega : (\mathcal{E}_{norm}^1(X, \omega, \psi), d) \rightarrow (\mathcal{M}^1(X, \omega, \psi), \text{strong}) \quad (5)$$

is a homeomorphism.

Then it is natural to wonder if one can extend the bijections (2), (4) to bigger subsets of $PSH(X, \omega)$. Given $\psi_1, \psi_2 \in \mathcal{M}^+$ such that $\psi_1 \neq \psi_2$ the sets $\mathcal{E}(X, \omega, \psi_1)$, $\mathcal{E}(X, \omega, \psi_2)$ are disjoint (Theorem 1.3 [13] quoted below as Theorem 2.1) but it may happen that $V_{\psi_1} = V_{\psi_2}$. So in these situations, at least one of $\mathcal{E}_{norm}^1(X, \omega, \psi_1)$, $\mathcal{E}_{norm}^1(X, \omega, \psi_2)$ must be ruled out to extend (4). However, given a totally ordered family $\mathcal{A} \subset \mathcal{M}^+$ of model type envelopes, the map $\mathcal{A} \ni \psi \rightarrow V_\psi$ is injective (again by Theorem 1.3 [13]), i.e.

$$MA_\omega : \bigsqcup_{\psi \in \mathcal{A}} \mathcal{E}_{norm}(X, \omega, \psi) \longrightarrow \left\{ \mu \text{ non-pluripolar positive measure} : \mu(X) = V_\psi \text{ for } \psi \in \mathcal{A} \right\}$$

is a bijection.

In [23] we introduced a complete distance $d_{\mathcal{A}}$ on

$$X_{\mathcal{A}} := \bigsqcup_{\psi \in \overline{\mathcal{A}}} \mathcal{E}^1(X, \omega, \psi)$$

where $\overline{\mathcal{A}} \subset \mathcal{M}$ is the weak closure of \mathcal{A} and where we identify $\mathcal{E}^1(X, \omega, \psi_{\min})$ with a point $P_{\psi_{\min}}$ if $\psi_{\min} \in \mathcal{M} \setminus \mathcal{M}^+$ (since in this case $E_\psi \equiv 0$, see Remark 2.7). Here ψ_{\min} is given as the smallest element in

$\overline{\mathcal{A}}$, observing that the Monge-Ampère operator $MA_\omega : \overline{\mathcal{A}} \rightarrow MA_\omega(\overline{\mathcal{A}})$ is a homeomorphism when the range is endowed with the weak topology (Lemma 3.12). We call strong topology on $X_{\mathcal{A}}$ the metric topology given by $d_{\mathcal{A}}$ since $d_{\mathcal{A}}|_{\mathcal{E}^1(X, \omega, \psi) \times \mathcal{E}^1(X, \omega, \psi)} = d$. The precise definition of $d_{\mathcal{A}}$ is quite technical (in section §2 we will recall many of its properties) but the strong topology is natural since it is the coarsest refinement of the weak topology such that $E(\cdot)$ becomes continuous as Theorem 6.2 shows. In particular the strong topology is independent on the set \mathcal{A} chosen.

Also the set

$$Y_{\mathcal{A}} := \bigsqcup_{\psi \in \overline{\mathcal{A}}} \mathcal{M}^1(X, \omega, \psi)$$

has a natural strong topology given as the coarsest refinement of the weak topology such that $E^*(\cdot)$ becomes continuous.

Theorem B. *The Monge-Ampère map*

$$MA_\omega : (X_{\mathcal{A}, norm}, d_{\mathcal{A}}) \rightarrow (Y_{\mathcal{A}}, strong)$$

is a homeomorphism.

Obviously in Theorem B we define $MA_\omega(P_{\psi_{\min}}) := 0$ if $V_{\psi_{\min}} = 0$.

Note that by Hartogs' Lemma and Theorem 6.2 the metric subspace $X_{\mathcal{A}, norm}$ is complete and it represents the set of all closed and positive $(1, 1)$ -currents $T = \omega + dd^c u$ such that $u \in X_{\mathcal{A}}$, where $P_{\psi_{\min}}$ encases all currents whose potentials u are more singular than ψ_{\min} if $V_{\psi_{\min}} = 0$.

Finally, as an application of Theorem B we study an example of the stability of solutions of complex Monge-Ampère equations. Other important situations will be dealt in a future work.

Theorem C. *Let $\mathcal{A} := \{\psi_k\}_{k \in \mathbb{N}} \subset \mathcal{M}^+$ be totally ordered, and let $\{f_k\}_{k \in \mathbb{N}} \subset L^1 \setminus \{0\}$ a sequence of non-negative functions such that $f_k \rightarrow f \in L^1 \setminus \{0\}$ and such that $\int_X f_k \omega^n = V_{\psi_k}$ for any $k \in \mathbb{N}$. Assume also that there exists $p > 1$ such that $\|f_k\|_{L^p}, \|f\|_{L^p}$ are uniformly bounded. Then $\psi_k \rightarrow \psi \in \mathcal{M}^+$ weakly, the sequence $\{u_k\}_{k \in \mathbb{N}}$ of solutions of*

$$\begin{cases} MA_\omega(u_k) = f_k \omega^n \\ u_k \in \mathcal{E}_{norm}^1(X, \omega, \psi_k) \end{cases} \quad (6)$$

converges strongly to $u \in X_{\mathcal{A}}$ (i.e. $d_{\mathcal{A}}(u_k, u) \rightarrow 0$), which is the unique solution of

$$\begin{cases} MA_\omega(u) = f \omega^n \\ u \in \mathcal{E}_{norm}^1(X, \omega, \psi). \end{cases}$$

In particular $u_k \rightarrow u$ in capacity.

The existence of the solutions of (6) follows by Theorem A in [14], while the fact that the strong convergence implies the convergence in capacity is our Theorem 6.3. Note also that the convergence in capacity of Theorem C was already obtained in [15] (see Remark 7.1).

1.1 Structure of the paper

Section §2 is dedicated to introduce some preliminaries, and in particular all necessary results presented in [23]. In section §3 we extend some known uniform estimates for $\mathcal{E}^1(X, \omega)$ to the relative setting, and we prove the key upper-semicontinuity of the relative energy functional $E(\cdot)$ in $X_{\mathcal{A}}$. Section §4 regards the properties of the action of measures on $PSH(X, \omega)$ and in particular their continuity. Then Section §5 is dedicated to prove Theorem A. We use a variational approach to show the bijection, then we need some further important properties of the strong topology on $\mathcal{E}^1(X, \omega, \psi)$ to conclude the proof. Section §6 is the heart of the article where we extends the results proved in the previous section to $X_{\mathcal{A}}$ and we present our main Theorem B. Finally in the last Section §7 we show Theorem C.

1.2 Future developments

As said above, in a future work we will present some strong stability results of more general solutions of complex Monge-Ampère equations with prescribed singularities than Theorem C, starting the study of a kind of *continuity method* when also the singularities will vary. As an application we will study the existence of (log) Kähler-Einstein metrics with prescribed singularities with a particular focus on the relationships among them varying the singularities.

1.3 Acknowledgments

I want to thank David Witt Nyström and Stefano Trapani for their suggestions and comments. I am also grateful to Hoang-Chinh Lu to have pointed me out a minor mistake in the previous version.

2 Preliminaries

We recall that given (X, ω) a Kähler complex compact manifold, the set $PSH(X, \omega)$ is the set of all ω -plurisubharmonic functions (ω -psh), i.e. all $u \in L^1$ given locally as sum of a smooth function and of a plurisubharmonic function such that $\omega + dd^c u \geq 0$ as $(1, 1)$ -current. Here $d^c := \frac{i}{2\pi}(\bar{\partial} - \partial)$ so that $dd^c = \frac{i}{\pi}\partial\bar{\partial}$. For any couple of ω -psh functions u, v the function

$$P_\omega[u](v) := \left(\lim_{C \rightarrow \infty} P_\omega(u + C, v) \right)^* = \left(\sup\{w \in PSH(X, \omega) : w \preceq u, w \leq v\} \right)^*$$

is ω -psh where the star is for the upper semicontinuous regularization and $P_\omega(u, v) := \left(\sup\{w \in PSH(X, \omega) : w \leq \min(u, v)\} \right)^*$. Then the set of all model type envelopes is defined as

$$\mathcal{M} := \{\psi \in PSH(X, \omega) : \psi = P_\omega[\psi](0)\}.$$

We also recall that \mathcal{M}^+ denotes the elements $\psi \in \mathcal{M}$ such that $V_\psi > 0$ where, as said in the Introduction, $V_\psi := \int_X MA_\omega(\psi)$.

The class of ψ -relative full mass functions $\mathcal{E}(X, \omega, \psi)$ complies the following characterization.

Theorem 2.1 (Theorem 1.3, [13]). *Suppose $v \in PSH(X, \omega)$ such that $V_v > 0$ and $u \in PSH(X, \omega)$ more singular than v . The followings are equivalent:*

- (i) $u \in \mathcal{E}(X, \omega, v)$;
- (ii) $P_\omega[u](v) = v$;
- (iii) $P_\omega[u](0) = P_\omega[v](0)$.

The clear inclusion $\mathcal{E}(X, \omega, v) \subset \mathcal{E}(X, \omega, P_\omega[v](0))$ may be strict, and it seems more natural in many cases to consider only functions $\psi \in \mathcal{M}$. For instance as showed in [13] ψ being a model type envelope is a necessary assumption to make the equation

$$\begin{cases} MA_\omega(u) = \mu \\ u \in \mathcal{E}(X, \omega, \psi) \end{cases}$$

always solvable where μ is a non-pluripolar measure such that $\mu(X) = V_\psi$. It is also worth to recall that there are plenty of elements in \mathcal{M} since $P_\omega[P_\omega[\psi]] = P_\omega[\psi]$ for any $\psi \in PSH(X, \omega)$ with $\int_X MA_\omega(\psi) > 0$ (Theorem 3.12, [13]). Indeed $v \rightarrow P_\omega[v]$ may be thought as a projection from the set of negative ω -psh functions with positive Monge-Ampère mass to \mathcal{M}^+ .

We also retrieve the following useful result.

Theorem 2.2 (Theorem 3.8, [13]). *Let $u, \psi \in PSH(X, \omega)$ such that $u \succcurlyeq \psi$. Then*

$$MA_\omega(P_\omega[\psi](u)) \leq \mathbb{1}_{\{P_\omega[\psi](u)=u\}} MA_\omega(u).$$

In particular if $\psi \in \mathcal{M}$ then $MA_\omega(\psi) \leq \mathbb{1}_{\{\psi=0\}} MA_\omega(0)$.

Note also that in Theorem 2.2 the equality holds if u is continuous with bounded distributional laplacian with respect to ω as a consequence of [17]. In particular $MA_\omega(\psi) = \mathbb{1}_{\{\psi=0\}} MA_\omega(0)$ for any $\psi \in \mathcal{M}$.

2.1 The metric space $(\mathcal{E}^1(X, \omega, \psi), d)$.

In this subsection we assume $\psi \in \mathcal{M}^+$ where $\mathcal{M}^+ := \{\psi \in \mathcal{M} : V_\psi > 0\}$.

As in [13] we also denote by $PSH(X, \omega, \psi)$ the set of all ω -psh functions which are more singular than ψ , and we recall that a function $u \in PSH(X, \omega, \psi)$ has ψ -relative minimal singularities if $|u - \psi|$ is globally bounded on X . We also use the notation

$$MA_\omega(u_1^{j_1}, \dots, u_l^{j_l}) := (\omega + dd^c u_1)^{j_1} \wedge \dots \wedge (\omega + dd^c u_l)^{j_l}$$

for $u_1, \dots, u_l \in PSH(X, \omega)$ where $j_1, \dots, j_l \in \mathbb{N}$ such that $j_1 + \dots + j_l = n$.

Definition 2.3 (Section §4.2, [13]). *The ψ -relative energy functional $E_\psi : PSH(X, \omega, \psi) \rightarrow \mathbb{R} \cup \{-\infty\}$ is defined as*

$$E_\psi(u) := \frac{1}{n+1} \sum_{j=0}^n \int_X (u - \psi) MA_\omega(u^j, \psi^{n-j})$$

if u has ψ -relative minimal singularities, and as

$$E_\psi(u) := \inf\{E_\psi(v) : v \in \mathcal{E}(X, \omega, \psi) \text{ with } \psi\text{-relative minimal singularities, } v \geq u\}$$

otherwise. The subset $\mathcal{E}^1(X, \omega, \psi) \subset \mathcal{E}(X, \omega, \psi)$ is defined as

$$\mathcal{E}^1(X, \omega, \psi) := \{u \in \mathcal{E}(X, \omega, \psi) : E_\psi(u) > -\infty\}.$$

When $\psi = 0$ the ψ -relative energy functional is the *Aubin-Mabuchi energy functional*, also called *Monge-Ampère energy* (see [1],[21]).

Proposition 2.4. *The following properties hold:*

- (i) E_ψ is non decreasing (Theorem 4.10, [13]);
- (ii) $E_\psi(u) = \lim_{j \rightarrow \infty} E_\psi(\max(u, \psi - j))$ (Lemma 4.12, [13]);
- (iii) E_ψ is continuous along decreasing sequences (Lemma 4.14, [13]);
- (iv) E_ψ is concave along affine curves (Theorem 4.10, Corollary 4.16, [13]);
- (v) $u \in \mathcal{E}^1(X, \omega, \psi)$ if and only if $u \in \mathcal{E}(X, \omega, \psi)$ and $\int_X (u - \psi) MA_\omega(u) > -\infty$ (Lemma 4.13, [13]);
- (vi) $E_\psi(u) \geq \limsup_{k \rightarrow \infty} E_\psi(u_k)$ if $u_k, u \in \mathcal{E}^1(X, \omega, \psi)$ and $u_k \rightarrow u$ with respect to the weak topology (Proposition 4.19, [13]);
- (vii) letting $u \in \mathcal{E}^1(X, \omega, \psi)$, $\chi \in C^0(X)$ and $u_t := \sup\{v \in PSH(X, \omega) : v \leq u + t\chi\}^*$ for any $t > 0$, then $t \rightarrow E_\psi(u_t)$ is differentiable and its derivative is given by

$$\frac{d}{dt} E_\psi(u_t) = \int_X \chi MA_\omega(u_t)$$

(Proposition 4.20, [13]);

(viii) if $u, v \in \mathcal{E}^1(X, \omega, \psi)$ then

$$E_\psi(u) - E_\psi(v) = \frac{1}{n+1} \sum_{j=0}^n \int_X (u - v) MA_\omega(u^j, v^{n-j})$$

and the function $\mathbb{N} \ni j \rightarrow \int_X (u - v) MA_\omega(u^j, v^{n-j})$ is decreasing. In particular

$$\int_X (u - v) MA_\omega(u) \leq E_\psi(u) - E_\psi(v) \leq \int_X (u - v) MA_\omega(v)$$

(Theorem 4.10, [13]);

(ix) if $u \leq v$ then $E_\psi(u) - E_\psi(v) \leq \frac{1}{n+1} \int_X (u - v) MA_\omega(u)$ (Theorem 4.10, [13]).

Remark 2.5. All the properties of Proposition 2.4 are showed in [13] assuming ψ having *small unbounded locus*, but the general integration by parts formula proved in [25] and Proposition 2.7 in [23] allow to extend these properties to the general case as described in Remark 2.10 in [23].

Recalling that for any $u, v \in \mathcal{E}^1(X, \omega, \psi)$ the function $P_\omega(u, v) = \sup\{w \in PSH(X, \omega) : w \leq \min(u, v)\}^*$ belongs to $\mathcal{E}^1(X, \omega, \psi)$ (see Proposition 2.13. in [23]), the function $d : \mathcal{E}^1(X, \omega, \psi) \times \mathcal{E}^1(X, \omega, \psi) \rightarrow \mathbb{R}_{\geq 0}$ defined as

$$d(u, v) = E_\psi(u) + E_\psi(v) - 2E_\psi(P_\omega(u, v))$$

assumes finite values. Moreover it is a complete distance as the next result shows.

Theorem 2.6 (Theorem A, [23]). *$(\mathcal{E}^1(X, \omega, \psi), d)$ is a complete metric space.*

We call *strong topology* on $\mathcal{E}^1(X, \omega, \psi)$ the metric topology given by the distance d . Note that by construction $d(u_k, u) \rightarrow 0$ as $k \rightarrow \infty$ if $u_k \searrow u$, and that $d(u, v) = d(u, w) + d(w, v)$ if $u \leq w \leq v$ (see Lemma 3.1 in [23]).

Moreover as a consequence of Proposition 2.4 it follows that for any $C \in \mathbb{R}_{>0}$ the set

$$\mathcal{E}_C^1(X, \omega, \psi) := \{u \in \mathcal{E}^1(X, \omega, \psi) : \sup_X u \leq C \text{ and } E_\psi(u) \geq -C\}$$

is a weakly compact convex set.

Remark 2.7. As described in Remark 3.10 in [23], if $\psi \in \mathcal{M} \setminus \mathcal{M}^+$ then $\mathcal{E}^1(X, \omega, \psi) = PSH(X, \omega, \psi)$ since $E_\psi \equiv 0$ by definition. In particular $d \equiv 0$ and it is natural to identify $(\mathcal{E}^1(X, \omega, \psi), d)$ with a point P_ψ . Moreover we recall that $\mathcal{E}^1(X, \omega, \psi_1) \cap \mathcal{E}^1(X, \omega, \psi_2) = \emptyset$ if $\psi_1, \psi_2 \in \mathcal{M}$, $\psi_1 \neq \psi_2$ and $V_{\psi_2} > 0$.

2.2 The space $(X_{\mathcal{A}}, d_{\mathcal{A}})$.

From now on we assume $\mathcal{A} \subset \mathcal{M}^+$ to be a totally ordered set of model type envelopes, and we denote by $\overline{\mathcal{A}}$ its closure as subset of $PSH(X, \omega)$ endowed with the weak topology. Note that $\overline{\mathcal{A}} \subset PSH(X, \omega)$ is compact by Lemma 2.6 in [23]. Indeed we will prove in Lemma 3.12 that actually $\overline{\mathcal{A}}$ is homeomorphic to its image through the Monge-Ampère operator MA_ω when the set of measure is endowed with the weak topology, which yields that $\overline{\mathcal{A}}$ is also homeomorphic to a closed set contained in $[0, \int_X \omega^n]$ through the map $\psi \rightarrow V_\psi$.

Definition 2.8. We define the set

$$X_{\mathcal{A}} := \bigsqcup_{\psi \in \overline{\mathcal{A}}} \mathcal{E}^1(X, \omega, \psi)$$

if $\psi_{\min} := \inf \mathcal{A}$ satisfies $V_{\psi_{\min}} > 0$, and

$$X_{\mathcal{A}} := P_{\psi_{\min}} \sqcup \bigsqcup_{\psi' \in \overline{\mathcal{A}}, \psi' \neq \psi_{\min}} \mathcal{E}^1(X, \omega, \psi')$$

if $V_{\psi_{\min}} = 0$, where $P_{\psi_{\min}}$ is a singleton.

$X_{\mathcal{A}}$ can be endowed with a natural metric structure as section 4 in [23] shows.

Theorem 2.9 (Theorem B, [23]). $(X_{\mathcal{A}}, d_{\mathcal{A}})$ is a complete metric space such that $d_{\mathcal{A}|_{\mathcal{E}^1(X, \omega, \psi) \times \mathcal{E}^1(X, \omega, \psi)}} = d$ for any $\psi \in \overline{\mathcal{A}} \cap \mathcal{M}^+$.

We call *strong topology* on $X_{\mathcal{A}}$ the metric topology given by the distance $d_{\mathcal{A}}$. Note that the denomination is coherent with that of subsection 2.1 since the induced topology on $\mathcal{E}^1(X, \omega, \psi) \subset X_{\mathcal{A}}$ coincides with the strong topology given by d .

We will also need the following contraction property which is the starting point to construct $d_{\mathcal{A}}$.

Proposition 2.10 (Lemma 4.2., Proposition 4.3., [23]). Let $\psi_1, \psi_2, \psi_3 \in \mathcal{M}$ such that $\psi_1 \preceq \psi_2 \preceq \psi_3$. Then $P_\omega[\psi_1](P_\omega[\psi_2](u)) = P_\omega[\psi_1](u)$ for any $u \in \mathcal{E}^1(X, \omega, \psi_3)$ and $|P_\omega[\psi_1](u) - \psi_1| \leq C$ if $|u - \psi_3| \leq C$. Moreover the map

$$P_\omega[\psi_1](\cdot) : \mathcal{E}^1(X, \omega, \psi_2) \rightarrow PSH(X, \omega, \psi_1)$$

has image in $\mathcal{E}^1(X, \omega, \psi_1)$ and it is a Lipschitz map of constant 1 when the sets $\mathcal{E}^1(X, \omega, \psi_i)$, $i = 1, 2$, are endowed with the d distances, i.e.

$$d(P_\omega[\psi_1](u), P_\omega[\psi_1](v)) \leq d(u, v)$$

for any $u, v \in \mathcal{E}^1(X, \omega, \psi_2)$.

Here we report some properties of the distance $d_{\mathcal{A}}$ and some consequences which will be useful in the sequel.

Proposition 2.11. The following properties hold:

i) if $u \in \mathcal{E}^1(X, \omega, \psi_1), v \in \mathcal{E}^1(X, \omega, \psi_2)$ for $\psi_1, \psi_2 \in \overline{\mathcal{A}}$, $\psi_1 \succcurlyeq \psi_2$ then

$$d_{\mathcal{A}}(u, v) \geq d(P_\omega[\psi_2](u), v)$$

(Proposition 4.14, [23]);

- ii) if $\{\psi_k\}_{k \in \mathbb{N}} \subset \mathcal{M}^+$, $\psi \in \mathcal{M}$ with $\psi_k \searrow \psi$ (resp. $\psi_k \nearrow \psi$ a.e.), $u_k \searrow u$, $v_k \searrow v$ (resp. $u_k \nearrow u$ a.e., $v_k \nearrow v$ a.e.) for $u_k, v_k \in \mathcal{E}^1(X, \omega, \psi_k)$, $u, v \in \mathcal{E}^1(X, \omega, \psi)$ and $|u_k - v_k|$ is uniformly bounded, then

$$d(u_k, v_k) \rightarrow d(u, v)$$

(Lemma 4.6, [23]);

- iii) if $\{\psi_k\}_{k \in \mathbb{N}} \subset \mathcal{M}^+$, $\psi \in \mathcal{M}$ such that $\psi_k \rightarrow \psi$ monotonically a.e., then for any $\psi' \in \mathcal{M}$ such that $\psi' \succcurlyeq \psi_k$ for any $k \gg 1$ big enough, and for any strongly compact set $K \subset (\mathcal{E}^1(X, \omega, \psi'), d)$,

$$d(P_\omega[\psi_k](\varphi_1), P_\omega[\psi_k](\varphi_2)) \rightarrow d(P_\omega[\psi](\varphi_1), P_\omega[\psi](\varphi_2))$$

uniformly on $K \times K$, i.e. varying $(\varphi_1, \varphi_2) \in K \times K$. In particular if $\psi_k, \psi \in \overline{\mathcal{A}}$ then

$$d_{\mathcal{A}}(P_\omega[\psi](u), P_\omega[\psi_k](u)) \rightarrow 0$$

$$d(P_\omega[\psi_k](u), P_\omega[\psi_k](v)) \rightarrow d(P_\omega[\psi](u), P_\omega[\psi](v))$$

monotonically for any $(u, v) \in \mathcal{E}^1(X, \omega, \psi') \times \mathcal{E}^1(X, \omega, \psi')$ (Proposition 4.5, [23]);

- iv) $d_{\mathcal{A}}(u_1, u_2) \geq |V_{\psi_1} - V_{\psi_2}|$ if $u_1 \in \mathcal{E}^1(X, \omega, \psi_1)$, $u_2 \in \mathcal{E}^1(X, \omega, \psi_2)$ and the equality holds if $u_1 = \psi_1$, $u_2 = \psi_2$ (by definition of $d_{\mathcal{A}}$, see section §4.2 in [23]).

The following Lemma is a special case of Theorem 2.2 in [25] (see also Lemma 4.1. in [13]).

Lemma 2.12 (Proposition 2.7, [23]). *Let $\{\psi_k\}_{k \in \mathbb{N}} \subset \mathcal{M}^+$, $\psi \in \mathcal{M}$ such that $\psi_k \rightarrow \psi$ monotonically almost everywhere. Let also $u_k, v_k \in \mathcal{E}^1(X, \omega, \psi_k)$ converging in capacity respectively to $u, v \in \mathcal{E}^1(X, \omega, \psi)$. Then for any $j = 0, \dots, n$*

$$MA_\omega(u_k^j, v_k^{n-j}) \rightarrow MA_\omega(u^j, v^{n-j})$$

weakly. Moreover if $|u_k - v_k|$ is uniformly bounded, then for any $j = 0, \dots, n$

$$(u_k - v_k)MA_\omega(u_k^j, v_k^{n-j}) \rightarrow (u - v)MA_\omega(u^j, v^{n-j})$$

weakly.

It is well-known that the set of Kähler potentials $\mathcal{H}_\omega := \{\varphi \in PSH(X, \omega) \cap C^\infty(X) : \omega + dd^c \varphi > 0\}$ is dense into $(\mathcal{E}^1(X, \omega), d)$. The same holds for $P_\omega[\psi](\mathcal{H}_\omega)$ into $(\mathcal{E}^1(X, \omega, \psi), d)$.

Lemma 2.13 (Lemma 4.8, [23]). *The set $\mathcal{P}_{\mathcal{H}_\omega}(X, \omega, \psi) := P_\omega[\psi](\mathcal{H}) \subset \mathcal{P}(X, \omega, \psi)$ is dense in $(\mathcal{E}^1(X, \omega, \psi), d)$.*

The following Lemma shows that, for $u \in PSH(X, \omega)$ fixed, the map $\mathcal{M}^+ \ni \psi \rightarrow P_\omega[\psi](u)$ is weakly continuous over any totally ordered set of model type envelopes that are more singular than u .

Lemma 2.14. *Let $u \in PSH(X, \omega)$, and let $\{\psi_k\}_{k \in \mathbb{N}} \subset \mathcal{M}^+$ be a totally ordered sequence of model type envelopes converging to $\psi \in \mathcal{M}$. Assume also that $\psi_k \preccurlyeq u$ for any $k \gg 1$ big enough. Then $P_\omega[\psi_k](u) \rightarrow P_\omega[\psi](u)$ weakly.*

Proof. As $\{\psi_k\}_{k \in \mathbb{N}}$ is totally ordered, without loss of generality we may assume that $\psi_k \rightarrow \psi$ monotonically almost everywhere. Set $\tilde{u} := \lim_{k \rightarrow \infty} P_\omega[\psi_k](u)$, and we want to prove that $\tilde{u} = P_\omega[\psi](u)$.

Suppose $\psi_k \searrow \psi$. It is immediate to check that $P_\omega[\psi_k](u) \leq P_\omega[\psi_k](\sup_X u) = \psi_k + \sup_X u$, which implies $\tilde{u} \leq \psi + \sup_X u$ letting $k \rightarrow +\infty$. Thus, $\tilde{u} \leq P_\omega[\psi](u)$ as the inequality $\tilde{u} \leq u$ is trivial. Moreover, since $\psi \leq \psi_k$ we also have $P_\omega[\psi](u) \leq P_\omega[\psi_k](u)$, which clearly yields $P_\omega[\psi](u) \leq \tilde{u}$ and concludes this part.

Suppose $\psi_k \nearrow \psi$. Then the inequality $\tilde{u} \leq P_\omega[\psi](u)$ is immediate. Next, combining Theorem 2.2 and Proposition 2.10 we have

$$\begin{aligned} MA_\omega(P_\omega[\psi_k](u)) &= MA_\omega\left(P_\omega[\psi_k](P_\omega[\psi](u))\right) \leq \mathbb{1}_{\{P_\omega[\psi_k](u) = P_\omega[\psi](u)\}} MA_\omega(P_\omega[\psi](u)) \leq \\ &\leq \mathbb{1}_{\{\tilde{u} = P_\omega[\psi](u)\}} MA_\omega(P_\omega[\psi](u)) \end{aligned}$$

where the last inequality follows from $P_\omega[\psi_k](u) \leq \tilde{u} \leq P_\omega[\psi](u)$. Thus, as $MA_\omega(P_\omega[\psi_k](u)) \rightarrow MA_\omega(\tilde{u})$ weakly by Theorem 2.3 in [13], we deduce that $\tilde{u} \in \mathcal{E}(X, \omega, \psi)$ and that $MA_\omega(\tilde{u}) \leq \mathbb{1}_{\{\tilde{u} = P_\omega[\psi](u)\}} MA_\omega(P_\omega[\psi](u))$. Moreover we also have $P_\omega[\psi](u) \in \mathcal{E}(X, \omega, \psi)$. Indeed $P_\omega[\psi](u) \leq P_\omega[\psi](\sup_X u) = \psi + \sup_X u$, i.e. $P_\omega[\psi](u) \preccurlyeq \psi$, while $P_\omega[\psi](u) \geq P_\omega[\psi](\psi_k - C_k) = \psi_k - C_k$ for non-negative constants C_k and for any $k \gg 1$ big enough as u, ψ are less singular than ψ_k . Thus $P_\omega[\psi](u) \succcurlyeq \psi_k$ for any k , which yields $\int_X MA_\omega(P_\omega[\psi](u)) \geq V_\psi > 0$ and gives $P_\omega[\psi](u) \in \mathcal{E}(X, \omega, \psi)$. Hence

$$0 \leq \int_X (P_\omega[\psi](u) - \tilde{u}) MA_\omega(\tilde{u}) \leq \int_{\{\tilde{u} = P_\omega[\psi](u)\}} (P_\omega[\psi](u) - \tilde{u}) MA_\omega(P_\omega[\psi](u)) = 0,$$

which by the domination principle of Proposition 3.11 in [13] implies $\tilde{u} \geq P_\omega[\psi](u)$ and concludes the proof. \square

3 Tools.

In this section we collect some uniform estimates on $\mathcal{E}^1(X, \omega, \psi)$ for $\psi \in \mathcal{M}^+$, we recall the ψ -relative capacity and we will prove the upper semicontinuity of $E(\cdot)$ on $X_{\mathcal{A}}$.

3.1 Uniform estimates.

Let $\psi \in \mathcal{M}^+$.

We first define in the ψ -relative setting the analogous of some well-known functionals of the variational approach (see [4] and reference therein).

We introduce respectively the ψ -relative I -functional and the ψ -relative J -functional (see also [1]) $I_\psi, J_\psi : \mathcal{E}^1(X, \omega, \psi) \times \mathcal{E}^1(X, \omega, \psi) \rightarrow \mathbb{R}$ where $\psi \in \mathcal{M}^+$ as

$$I_\psi(u, v) := \int_X (u - v)(MA_\omega(v) - MA_\omega(u)),$$

$$J_\psi(u, v) := J_u^\psi(v) := E_\psi(u) - E_\psi(v) + \int_X (v - u)MA_\omega(u).$$

They assume non-negative values by Proposition 2.4, I_ψ is clearly symmetric while J_ψ is convex again by Proposition 2.4. Moreover the ψ -relative I and J functionals are related each other by the following result.

Lemma 3.1. *Let $u, v \in \mathcal{E}^1(X, \omega, \psi)$. Then*

- (i) $\frac{1}{n+1}I_\psi(u, v) \leq J_u^\psi(v) \leq \frac{n}{n+1}I_\psi(u, v)$;
- (ii) $\frac{1}{n}J_u^\psi(v) \leq J_v^\psi(u) \leq nJ_u^\psi(v)$.

In particular

$$d(\psi, u) \leq nJ_u^\psi(\psi) + (\|\psi\|_{L^1} + \|u\|_{L^1})$$

for any $u \in \mathcal{E}^1(X, \omega, \psi)$ such that $u \leq \psi$.

Proof. By Proposition 2.4 it follows that

$$n \int_X (u - v)MA_\omega(u) + \int_X (u - v)MA_\omega(v) \leq$$

$$\leq (n+1)(E_\psi(u) - E_\psi(v)) \leq \int_X (u - v)MA_\omega(u) + n \int_X (u - v)MA_\omega(v)$$

for any $u, v \in \mathcal{E}^1(X, \omega, \psi)$, which yields (i) and (ii).

Next considering $v = \psi$ and assuming $u \leq \psi$ from the second inequality in (ii) we obtain

$$d(u, \psi) = -E_\psi(u) \leq nJ_u^\psi(\psi) + \int_X (\psi - u)MA_\omega(\psi),$$

which implies the assertion since $MA_\omega(\psi) \leq MA_\omega(0)$ by Theorem 2.2. \square

We can now proceed showing the uniform estimates, adapting some results in [4].

Lemma 3.2 (Lemma 3.7, [23]). *Let $\psi \in \mathcal{M}^+$. Then there exists positive constants $A > 1, B > 0$ depending only on n, ω such that for any $u \in \mathcal{E}^1(X, \omega, \psi)$*

$$-d(\psi, u) \leq V_\psi \sup_X (u - \psi) = V_\psi \sup_X u \leq Ad(\psi, u) + B$$

Remark 3.3. As a consequence of Lemma 3.2 if $d(\psi, u) \leq C$ then $\sup_X u \leq (AC+B)/V_\psi$ while $-E_\psi(u) = d(\psi + (AC+B)/V_\psi, u) - (AC+B) \leq d(\psi, u) \leq C$, i.e. $u \in \mathcal{E}_D^1(X, \omega, \psi)$ where $D := \max(C, (AC+B)/V_\psi)$. Conversely, it is easy to check that $d(u, \psi) \leq C(2V_\psi + 1)$ for any $u \in \mathcal{E}_C^1(X, \omega, \psi)$ using the definitions and the triangle inequality.

Proposition 3.4. *Let $C \in \mathbb{R}_{>0}$. Then there exists a continuous increasing function $f_C : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ depending only on C, ω, n with $f_C(0) = 0$ such that*

$$\left| \int_X (u - v)(MA_\omega(\varphi_1) - MA_\omega(\varphi_2)) \right| \leq f_C(d(u, v)) \quad (7)$$

for any $u, v, \varphi_1, \varphi_2 \in \mathcal{E}^1(X, \omega, \psi)$ with $d(u, \psi), d(v, \psi), d(\varphi_1, \psi), d(\varphi_2, \psi) \leq C$.

Proof. As said in Remark 3.3 if $w \in \mathcal{E}^1(X, \omega, \psi)$ with $d(\psi, w) \leq C$ then $\tilde{w} := w - (AC + B)/V_\psi$ satisfies $\sup_X \tilde{w} \leq 0$ and

$$-E_\psi(\tilde{w}) = d(\psi, \tilde{w}) \leq d(\psi, w) + d(w, \tilde{w}) \leq C + AC + B =: D.$$

Therefore setting $\tilde{u} := u - (AC + B)/V_\psi$, $\tilde{v} := v - (AC + B)/V_\psi$ we can proceed exactly as in Lemma 5.8 in [4] using the integration by parts formula in [25] (see also Theorem 1.14 in [7]) to get

$$\left| \int_X (\tilde{u} - \tilde{v})(MA_\omega(\varphi_1) - MA_\omega(\varphi_2)) \right| \leq I_\psi(\tilde{u}, \tilde{v}) + h_D(I_\psi(\tilde{u}, \tilde{v})) \quad (8)$$

where $h_D : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ is an increasing continuous function depending only on D such that $h_D(0) = 0$. Furthermore, by definition

$$d(\psi, P_\omega(\tilde{u}, \tilde{v})) \leq d(\psi, \tilde{u}) + d(\tilde{u}, P_\omega(\tilde{u}, \tilde{v})) \leq d(\psi, \tilde{u}) + d(\tilde{u}, \tilde{v}) \leq 3D,$$

so, by the triangle inequality and (8), we have

$$\begin{aligned} \left| \int_X (u - v)(MA_\omega(\varphi_1) - MA_\omega(\varphi_2)) \right| &\leq \\ &\leq I_\psi(\tilde{u}, P_\omega(\tilde{u}, \tilde{v})) + I_\psi(\tilde{v}, P_\omega(\tilde{u}, \tilde{v})) + h_{3D}(I_\psi(\tilde{u}, P_\omega(\tilde{u}, \tilde{v}))) + h_{3D}(I_\psi(\tilde{v}, P_\omega(\tilde{u}, \tilde{v}))). \end{aligned} \quad (9)$$

On the other hand, if $w_1, w_2 \in \mathcal{E}^1(X, \omega, \psi)$ with $w_1 \geq w_2$ then by Proposition 2.4

$$I_\psi(w_1, w_2) \leq \int_X (w_1 - w_2)MA_\omega(w_2) \leq (n+1)d(w_1, w_2).$$

Hence from (9) it is sufficient to set $f_C(x) := (n+1)x + 2h_{3D}((n+1)x)$ to conclude the proof since clearly $d(\tilde{u}, \tilde{v}) = d(u, v)$. \square

Corollary 3.5. *Let $\psi \in \mathcal{M}^+$ and let $C \in \mathbb{R}_{>0}$. Then there exists a continuous increasing functions $f_C : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ depending only on C, ω, n with $f_C(0) = 0$ such that*

$$\int_X |u - v|MA_\omega(\varphi) \leq f_C(d(u, v))$$

for any $u, v, \varphi \in \mathcal{E}^1(X, \omega, \psi)$ with $d(\psi, u), d(\psi, v), d(\psi, \varphi) \leq C$.

Proof. Since $d(\psi, P_\omega(u, v)) \leq 3C$, letting $g_{3C} : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be the map (7) of Proposition 3.4, it follows that

$$\begin{aligned} \int_X (u - P_\omega(u, v))MA_\omega(\varphi) &\leq \int_X (u - P_\omega(u, v))MA_\omega(P_\omega(u, v)) + g_{3C}(d(u, P_\omega(u, v))) \leq \\ &\leq (n+1)d(u, P_\omega(u, v)) + g_{3C}(d(u, v)), \end{aligned}$$

where in the last inequality we used Proposition 2.4. Hence by the triangle inequality we get

$$\begin{aligned} \int_X |u - v|MA_\omega(\varphi) &\leq (n+1)d(u, P_\omega(u, v)) + (n+1)d(v, P_\omega(u, v)) + 2g_{3C}(d(u, v)) = \\ &= (n+1)d(u, v) + 2g_{3C}(d(u, v)). \end{aligned}$$

Defining $f_C(x) := (n+1)x + 2g_{3C}(x)$ concludes the proof. \square

As first important consequence we obtain that the strong convergence in $\mathcal{E}^1(X, \omega, \psi)$ implies the weak convergence.

Proposition 3.6. *Let $\psi \in \mathcal{M}^+$ and let $C \in \mathbb{R}_{>0}$. Then there exists a continuous increasing function $f_{C, \psi} : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ depending on C, ω, n, ψ with $f_{C, \psi}(0) = 0$ such that*

$$\|u - v\|_{L^1} \leq f_{C, \psi}(d(u, v))$$

for any $u, v \in \mathcal{E}^1(X, \omega, \psi)$ with $d(\psi, u), d(\psi, v) \leq C$. In particular $u_k \rightarrow u$ weakly if $u_k \rightarrow u$ strongly.

Proof. Theorem A in [14] (see also Theorem 1.4 in [13]) implies that there exists $\phi \in \mathcal{E}^1(X, \omega, \psi)$ with $\sup_X \phi = 0$ such that

$$MA_\omega(\phi) = cMA_\omega(0)$$

where $c := V_\psi/V_0 > 0$. Therefore it follows that

$$\|u - v\|_{L^1} \leq \frac{1}{c} g_{\hat{C}}(d(u, v))$$

where $\hat{C} := \max(d(\psi, \phi), C)$ and $g_{\hat{C}}$ is the continuous increasing function with $g_{\hat{C}}(0) = 0$ given by Corollary 3.5. Setting $f_{C, \psi} := \frac{1}{c} g_{\hat{C}}$ concludes the proof. \square

Finally we also get the following useful estimate.

Proposition 3.7. *Let $\psi \in \mathcal{M}^+$ and let $C \in \mathbb{R}_{>0}$. Then there exists a constant \tilde{C} depending only on C, ω, n such that*

$$\left| \int_X (u - v)(MA_\omega(\varphi_1) - MA_\omega(\varphi_2)) \right| \leq \tilde{C} I_\psi(\varphi_1, \varphi_2)^{\frac{1}{2}} \quad (10)$$

for any $u, v, \varphi_1, \varphi_2 \in \mathcal{E}^1(X, \omega, \psi)$ with $d(u, \psi), d(v, \psi), d(\varphi_1, \psi), d(\varphi_2, \psi) \leq C$.

Proof. As seen during the proof of Proposition 3.4 and with the same notations, the function $\tilde{u} := u - (AC + B)/V_\psi$ satisfy $\sup_X \tilde{u} \leq 0$ (by Lemma 3.2) and $-E_\psi(\tilde{u}) \leq C + AC + B =: D$ (and similarly for v, φ_1, φ_2). Therefore by integration by parts and using Lemma 3.8 below, it follows exactly as in Lemma 3.13 in [4] that there exists a constant \tilde{C} depending only on D, n such that

$$\left| \int_X (\tilde{u} - \tilde{v})(MA_\omega(\tilde{\varphi}_1) - MA_\omega(\tilde{\varphi}_2)) \right| \leq \tilde{C} I_\psi(\tilde{\varphi}_1, \tilde{\varphi}_2)^{\frac{1}{2}},$$

which clearly implies (10). \square

Lemma 3.8. *Let $C \in \mathbb{R}_{>0}$. Then there exists a constant \tilde{C} depending only on C, ω, n such that*

$$\int_X |u_0 - \psi|(\omega + dd^c u_1) \wedge \cdots \wedge (\omega + dd^c u_n) \leq \tilde{C}$$

for any $u_0, \dots, u_n \in \mathcal{E}^1(X, \omega, \psi)$ with $d(u_j, \psi) \leq C$ for any $j = 0, \dots, n$.

Proof. As in Proposition 3.4 and with the same notations $v_j := u_j - (AC + B)/V_\psi$ satisfies $\sup_X v_j \leq 0$, and setting $v := \frac{1}{n+1}(v_0 + \cdots + v_n)$ we obtain $\psi - u_0 \leq (n+1)(\psi - v)$. Thus by Proposition 2.4 it follows that

$$\begin{aligned} \int_X (\psi - v_0)MA_\omega(v) &\leq (n+1) \int_X (\psi - v)MA_\omega(v) \leq (n+1)^2 |E_\psi(v)| \leq \\ &\leq (n+1) \sum_{j=0}^n |E_\psi(v_j)| \leq (n+1) \sum_{j=0}^n (d(\psi, u_j) + D) \leq (n+1)^2(C + D) \end{aligned}$$

where $D := AC + B$. On the other hand $MA_\omega(v) \geq E(\omega + dd^c u_1) \wedge \cdots \wedge (\omega + dd^c u_n)$ where the constant E depends only on n . Finally we get

$$\int_X |u_0 - \psi|(\omega + dd^c u_1) \wedge \cdots \wedge (\omega + dd^c u_n) \leq D + \frac{1}{E} \int_X (\psi - v_0)MA_\omega(v) \leq D + \frac{(n+1)^2(C + D)}{E},$$

which concludes the proof. \square

3.2 ψ -relative Monge-Ampère capacity.

Definition 3.9 (Section §4.1, [13]; Definition 3.1, [14]). *Let $B \subset X$ be a Borel set, and let $\psi \in \mathcal{M}^+$. Then its ψ -relative Monge-Ampère capacity is defined as*

$$\text{Cap}_\psi(B) := \sup \left\{ \int_B MA_\omega(u) : u \in PSH(X, \omega), \psi - 1 \leq u \leq \psi \right\}.$$

In the absolute setting the Monge-Ampère capacity is very useful to study the existence and the regularity of solutions of degenerate complex Monge-Ampère equation ([20]), and analog holds in the relative setting ([13], [14]). We refer to these articles just cited to many properties of the Monge-Ampère capacity.

For any fixed constant A , $\mathcal{C}_{A,\psi}$ denotes the set of all probability measures μ on X such that

$$\mu(B) \leq A \text{Cap}_\psi(B)$$

for any Borel set $B \subset X$ (Section §4.3, [13]).

Proposition 3.10. *Let $u \in \mathcal{E}^1(X, \omega, \psi)$ with ψ -relative minimal singularities. Then $MA_\omega(u)/V_\psi \in \mathcal{C}_{A,\psi}$ for a constant $A > 0$.*

Proof. Let $j \in \mathbb{R}$ such that $u \geq \psi - j$ and assume without loss of generality that $u \leq \psi$ and that $j \geq 1$. Then the function $v := j^{-1}u + (1 - j^{-1})\psi$ is a candidate in the definition of Cap_ψ , which implies that $MA_\omega(v) \leq \text{Cap}_\psi$. Hence, since $MA_\omega(u) \leq j^n MA_\omega(v)$ we get that $MA_\omega(u) \in \mathcal{C}_{A,\psi}$ for $A = j^n$ and the result follows. \square

We also need to quote the following result.

Lemma 3.11 (Lemma 4.18, [13]). *If $\mu \in \mathcal{C}_{A,\psi}$ then there is a constant $B > 0$ depending only on A, n such that*

$$\int_X (u - \psi)^2 \mu \leq B(|E_\psi(u)| + 1)$$

for any $u \in PSH(X, \omega, \psi)$ such that $\sup_X u = 0$.

Similarly to the case $\psi = 0$ (see [19]), we say that a sequence $u_k \in PSH(X, \omega)$ converges to $u \in PSH(X, \omega)$ in ψ -relative capacity for $\psi \in \mathcal{M}$ if

$$\text{Cap}_\psi(\{|u_k - u| \geq \delta\}) \rightarrow 0$$

as $k \rightarrow \infty$ for any $\delta > 0$.

By Theorem 10.37 in [19] (see also Theorem 5.7 in [4]) the convergence in $(\mathcal{E}^1(X, \omega), d)$ implies the convergence in capacity. The analogous holds for $\psi \in \mathcal{M}^+$, i.e. that the strong convergence in $\mathcal{E}^1(X, \omega, \psi)$ implies the convergence in ψ -relative capacity. Indeed in Proposition 5.7 we will prove the the strong convergence implies the convergence in ψ' -relative capacity for any $\psi' \in \mathcal{M}^+$.

3.3 (Weak) Upper Semicontinuity of $u \rightarrow E_{P_\omega[u]}(u)$ over $X_{\mathcal{A}}$.

One of the main feature of E_ψ for $\psi \in \mathcal{M}$ is its upper semicontinuity with respect to the weak topology. Here we prove the analogous for $E(\cdot)$ over $X_{\mathcal{A}}$.

Lemma 3.12. *The map $MA_\omega : \overline{\mathcal{A}} \rightarrow MA_\omega(\overline{\mathcal{A}}) \subset \{\mu \text{ positive measure on } X\}$ is a homeomorphism considering the weak topologies. In particular $\overline{\mathcal{A}}$ is homeomorphic to a closed set contained in $[0, \int_X MA_\omega(0)]$ through the map $\psi \rightarrow V_\psi$.*

Proof. The map is well-defined and continuous by Lemma 2.6 in [23]. Moreover the injectivity follows from the fact that $V_{\psi_1} = V_{\psi_2}$ for $\psi_1, \psi_2 \in \overline{\mathcal{A}}$ implies $\psi_1 = \psi_2$ using Theorem 2.1 and the fact that $\mathcal{A} \subset \mathcal{M}^+$. Finally to conclude the proof it is enough to prove that $\psi_k \rightarrow \psi$ weakly assuming $V_{\psi_k} \rightarrow V_\psi$ and it is clearly sufficient to show that any subsequence of $\{\psi_k\}_{k \in \mathbb{N}}$ admits a subsequence weakly convergent to ψ . Moreover since $\overline{\mathcal{A}}$ is totally ordered and \succcurlyeq coincides with \geq on \mathcal{M} , we may assume $\{\psi_k\}_{k \in \mathbb{N}}$ monotonic sequence. Then, up to considering a further subsequence, ψ_k converges almost everywhere to an element $\psi' \in \overline{\mathcal{A}}$ by compactness, and Lemma 2.12 implies that $V_{\psi'} = V_\psi$, i.e $\psi = \psi'$. \square

In the case $\mathcal{A} := \{\psi_k\}_{k \in \mathbb{N}} \subset \mathcal{M}^+$, we say that $u_k \in \mathcal{E}^1(X, \omega, \psi_k)$ converges weakly to $P_{\psi_{\min}}$ where $\psi_{\min} \in \mathcal{M} \setminus \mathcal{M}^+$ if $|\sup_X u_k| \leq C$ for any $k \in \mathbb{N}$ and any weak accumulation point u of $\{u_k\}_{k \in \mathbb{N}}$ satisfies $u \preccurlyeq \psi_{\min}$. This definition is the most natural since $PSH(X, \omega, \psi) = \mathcal{E}^1(X, \omega, \psi_{\min})$.

Lemma 3.13. *Let $\{u_k\}_{k \in \mathbb{N}} \subset X_{\mathcal{A}}$ be a sequence converging weakly to $u \in X_{\mathcal{A}}$. If $E_{P_\omega[u_k]}(u_k) \geq C$ uniformly, then $P_\omega[u_k] \rightarrow P_\omega[u]$ weakly.*

Proof. By Lemma 3.12 the convergence requested is equivalent to $V_{\psi_k} \rightarrow V_\psi$, where we set $\psi_k := P_\omega[u_k]$, $\psi := P_\omega[u]$.

Moreover by a simple contradiction argument it is enough to show that any subsequence $\{\psi_{k_h}\}_{h \in \mathbb{N}}$ admits a subsequence $\{\psi_{k_{h_j}}\}_{j \in \mathbb{N}}$ such that $V_{\psi_{k_{h_j}}} \rightarrow V_\psi$. Thus up to considering a subsequence, by abuse of notations and by the lower semicontinuity $\liminf_{k \rightarrow \infty} V_{\psi_k} \geq V_\psi$ of Theorem 2.3. in [13], we may suppose by contradiction that $\psi_k \searrow \psi'$ for $\psi' \in \mathcal{M}$ such that $V_{\psi'} > V_\psi$. In particular $V_{\psi'} > 0$ and $\psi' \succneq \psi$. Then by Proposition 2.10 and Remark 3.3 the sequence $\{P_\omega[\psi'](u_k)\}_{k \in \mathbb{N}}$ is bounded in $(\mathcal{E}^1(X, \omega, \psi'), d)$ and it belongs to $\mathcal{E}_{C'}^1(X, \omega, \psi')$ for some $C' \in \mathbb{R}$. Therefore, up to considering a subsequence, we have that $\{u_k\}_{k \in \mathbb{N}}$ converges weakly to an element $v \in \mathcal{E}^1(X, \omega, \psi)$ (which is the element u itself when $u \neq P_{\psi_{\min}}$) while the sequence $P_\omega[\psi'](u_k)$ converges weakly to an element $w \in \mathcal{E}^1(X, \omega, \psi')$. Thus the contradiction follows from $w \leq v$ since $\psi' \succneq \psi$, $V_{\psi'} > 0$ and $\mathcal{E}^1(X, \omega, \psi') \cap \mathcal{E}^1(X, \omega, \psi) = \emptyset$. \square

Proposition 3.14. *Let $\{u_k\}_{k \in \mathbb{N}} \subset X_{\mathcal{A}}$ be a sequence converging weakly to $u \in X_{\mathcal{A}}$. Then*

$$\limsup_{k \rightarrow \infty} E_{P_\omega[u_k]}(u_k) \leq E_{P_\omega[u]}(u). \quad (11)$$

Proof. Let $\psi_k := P_\omega[u_k]$, $\psi := P_\omega[u] \in \overline{\mathcal{A}}$. We may clearly assume $\psi_k \neq \psi_{\min}$ for any $k \in \mathbb{N}$ if $\psi = \psi_{\min}$ and $V_{\psi_{\min}} = 0$.

Moreover we can also suppose that $E_{\psi_k}(u_k)$ is bounded from below, which implies that $u_k \in \mathcal{E}_C^1(X, \omega, \psi_k)$ for a uniform constant C and that $\psi_k \rightarrow \psi$ weakly by Lemma 3.13. Thus since $E_{\psi_k}(u_k) = E_{\psi_k}(u_k - C) + CV_{\psi_k}$ for any $k \in \mathbb{N}$, Lemma 3.12 implies that we may assume that $\sup_X u_k \leq 0$. Furthermore since \mathcal{A} is totally ordered, it is enough to show (11) when $\psi_k \rightarrow \psi$ a.e. monotonically.

If $\psi_k \searrow \psi$, setting $v_k := (\sup\{u_j : j \geq k\})^* \in \mathcal{E}^1(X, \omega, \psi_k)$, we easily have

$$\limsup_{k \rightarrow \infty} E_{\psi_k}(u_k) \leq \limsup_{k \rightarrow \infty} E_{\psi_k}(v_k) \leq \limsup_{k \rightarrow \infty} E_\psi(P_\omega[\psi](v_k))$$

using the monotonicity of E_{ψ_k} and Proposition 2.10. Hence if $\psi = \psi_{\min}$ and $V_{\psi_{\min}} = 0$ then $E_\psi(P_\omega[\psi](v_k)) = 0 = E_\psi(u)$, while otherwise the conclusion follows from Proposition 2.4 since $P_\omega[\psi](v_k) \searrow u$ by construction.

If instead $\psi_k \nearrow \psi$, fix $\epsilon > 0$ and for any $k \in \mathbb{N}$ let $j_k \geq k$ such that

$$\sup_{j \geq k} E_{\psi_j}(u_j) \leq E_{\psi_{j_k}}(u_{j_k}) + \epsilon.$$

Thus again by Proposition 2.10, $E_{\psi_{j_k}}(u_{j_k}) \leq E_{\psi_l}(P_\omega[\psi_l](u_{j_k}))$ for any $l \leq j_k$. Moreover, assuming $E_{\psi_{j_k}}(u_{j_k})$ bounded from below, $-E_{\psi_l}(P_\omega[\psi_l](u_{j_k})) = d(\psi_l, P_\omega[\psi_l](u_{j_k}))$ is uniformly bounded in l, k , which implies that $\sup_X P_\omega[\psi_l](u_{j_k})$ is uniformly bounded by Remark 3.3 since $V_{\psi_{j_k}} \geq a > 0$ for $k \gg 0$ big enough. By compactness, up to considering a subsequence, we obtain $P_\omega[\psi_l](u_{j_k}) \rightarrow v_l$ weakly where $v_l \in \mathcal{E}^1(X, \omega, \psi_l)$ by the upper semicontinuity of $E_{\psi_l}(\cdot)$ on $\mathcal{E}^1(X, \omega, \psi_l)$. Hence

$$\limsup_{k \rightarrow \infty} E_{\psi_k}(u_k) \leq \limsup_{k \rightarrow \infty} E_{\psi_l}(P_\omega[\psi_l](u_{j_k})) + \epsilon = E_{\psi_l}(v_l) + \epsilon$$

for any $l \in \mathbb{N}$. Moreover by construction $v_l \leq P_\omega[\psi_l](u)$ since $P_\omega[\psi_l](u_{j_k}) \leq u_{j_k}$ for any k such that $j_k \geq l$ and $u_{j_k} \rightarrow u$ weakly. Therefore by the monotonicity of $E_{\psi_l}(\cdot)$ and by Proposition 2.11.(ii) we conclude that

$$\limsup_{k \rightarrow \infty} E_{\psi_k}(u_k) \leq \lim_{l \rightarrow \infty} E_{\psi_l}(P_\omega[\psi_l](u)) + \epsilon = E_\psi(u) + \epsilon$$

letting $l \rightarrow \infty$. \square

As a consequence, defining

$$X_{\mathcal{A}, C} := \bigsqcup_{\psi \in \overline{\mathcal{A}}} \mathcal{E}_C^1(X, \omega, \psi),$$

we get the following compactness result.

Proposition 3.15. *Let $C, a \in \mathbb{R}_{>0}$. The set*

$$X_{\mathcal{A}, C}^a := X_{\mathcal{A}, C} \cap \left(\bigsqcup_{\psi \in \overline{\mathcal{A}}: V_\psi \geq a} \mathcal{E}^1(X, \omega, \psi) \right)$$

is compact with respect to the weak topology.

Proof. It follows directly from the definition that

$$X_{\mathcal{A},C}^a \subset \left\{ u \in PSH(X, \omega) : \sup_X u \leq C' \right\}$$

where $C' := \max(C, C/a)$. Therefore by Proposition 8.5 in [19], $X_{\mathcal{A},C}^a$ is weakly relatively compact. Finally Proposition 3.14 and Hartogs' Lemma imply that $X_{\mathcal{A},C}^a$ is also closed with respect to the weak topology, concluding the proof. \square

Remark 3.16. The whole set $X_{\mathcal{A},C}$ may not be weakly compact. Indeed assuming $V_{\psi_{\min}} = 0$ and letting $\psi_k \in \overline{\mathcal{A}}$ such that $\psi_k \searrow \psi_{\min}$, the functions $u_k := \psi_k - 1/\sqrt{V_{\psi_k}}$ belong to $X_{\mathcal{A},V}$ for $V = \int_X MA_\omega(0)$ since $E_{\psi_k}(u_k) = -\sqrt{V_{\psi_k}}$ but $\sup_X u_k = -1/\sqrt{V_{\psi_k}} \rightarrow -\infty$.

4 The action of measures on $PSH(X, \omega)$.

In this section we want to replace the action on $PSH(X, \omega)$ defined in [4] given by a probability measure μ with an action which assume finite values on elements $u \in PSH(X, \omega)$ with ψ -relative minimal singularities where $\psi = P_\omega[u]$ for almost all $\psi \in \mathcal{M}$. On the other hand for any $\psi \in \mathcal{M}$ we want that there exists many measures μ whose action over $\{u \in PSH(X, \omega) : P_\omega[u] = \psi\}$ is well-defined. The problem is that μ varies among *all* probability measures while ψ among *all* model type envelopes. So it may happen that μ takes mass on non-pluripolar sets and that the unbounded locus of $\psi \in \mathcal{M}$ is very nasty.

Definition 4.1. Let μ be a probability measure on X . Then μ acts on $PSH(X, \omega)$ through the functional $L_\mu : PSH(X, \omega) \rightarrow \mathbb{R} \cup \{-\infty\}$ defined as $L_\mu(u) = -\infty$ if μ charges $\{P_\omega[u] = -\infty\}$, as

$$L_\mu(u) := \int_X (u - P_\omega[u]) \mu$$

if u has $P_\omega[u]$ -relative minimal singularities and μ does not charge $\{P_\omega[u] = -\infty\}$ and as

$$L_\mu(u) := \inf\{L_\mu(v) : v \in PSH(X, \omega) \text{ with } P_\omega[u]\text{-relative minimal singularities, } v \geq u\}$$

otherwise.

Proposition 4.2. The following properties hold:

- (i) L_μ is affine, i.e. it satisfies the scaling property $L_\mu(u+c) = L_\mu(u) + c$ for any $c \in \mathbb{R}$, $u \in PSH(X, \omega)$;
- (ii) L_μ is non-decreasing on $\{u \in PSH(X, \omega) : P_\omega[u] = \psi\}$ for any $\psi \in \mathcal{M}$;
- (iii) $L_\mu(u) = \lim_{j \rightarrow \infty} L_\mu(\max(u, P_\omega[u] - j))$ for any $u \in PSH(X, \omega)$;
- (iv) if μ is non-pluripolar then L_μ is convex;
- (v) if μ is non-pluripolar and $u_k \rightarrow u$ and $P_\omega[u_k] \rightarrow P_\omega[u]$ weakly as $k \rightarrow \infty$ then $L_\mu(u) \geq \limsup_{k \rightarrow \infty} L_\mu(u_k)$;
- (vi) if $u \in \mathcal{E}^1(X, \omega, \psi)$ for $\psi \in \mathcal{M}^+$ then $L_{MA_\omega(u)/V_\psi}$ is finite on $\mathcal{E}^1(X, \omega, \psi)$.

Proof. The first two points follow by definition.

For the third point, setting $\psi := P_\omega[u]$, clearly $L_\mu(u) \leq \lim_{j \rightarrow \infty} L_\mu(\max(u, \psi - j))$. Conversely, for any $v \geq u$ with ψ -relative minimal singularities $v \geq \max(u, \psi - j)$ for $j \gg 0$ big enough, hence by (ii) we get $L_\mu(v) \geq \lim_{j \rightarrow \infty} L_\mu(\max(u, \psi - j))$ which implies (iii) by definition.

Next, we prove (iv). Let $v = \sum_{l=1}^m a_l u_l$ be a convex combination of elements $u_l \in PSH(X, \omega)$, and without loss of generality we may assume $\sup_X v, \sup_X u_l \leq 0$. In particular we have $L_\mu(v), L_\mu(u_l) \leq 0$.

Suppose $L_\mu(v) > -\infty$ (otherwise it is trivial) and let $\psi := P_\omega[v]$, $\psi_l := P_\omega[u_l]$. Then for any $C \in \mathbb{R}_{>0}$ it is easy to see that

$$\sum_{l=1}^m a_l P_\omega(u_l + C, 0) \leq P_\omega(v + C, 0) \leq \psi,$$

which leads to $\sum_{l=1}^m a_l \psi_l \leq \psi$ letting $C \rightarrow \infty$. Hence (iii) yields

$$-\infty < L_\mu(v) = \int_X (v - \psi) \mu \leq \sum_{l=1}^m a_l \int_X (u_l - \psi_l) \mu = \sum_{l=1}^m a_l L_\mu(u_l).$$

The point (v) easily follows from $\limsup_{k \rightarrow \infty} \max(u_k, P_\omega[u_k] - j) \leq \max(u, P_\omega[u] - j)$ and (iii), while the last point is a consequence of Lemma 3.8. \square

Next, since for any $t \in [0, 1]$ and any $u, v \in \mathcal{E}^1(X, \omega, \psi)$

$$\begin{aligned} \int_X (u - v)MA_\omega(tu + (1 - t)v) &= \\ &= (1 - t)^n \int_X (u - v)MA_\omega(v) + \sum_{j=1}^n \binom{n}{j} t^j (1 - t)^{n-j} \int_X (u - v)MA_\omega(u^j, v^{n-j}) \geq \\ &\geq (1 - t)^n \int_X (u - v)MA_\omega(v) + (1 - (1 - t)^n) \int_X (u - v)MA_\omega(u), \end{aligned}$$

we can proceed exactly as in Proposition 3.4 in [4] (see also Lemma 2.11. in [18]), replacing V_θ with ψ , to get the following result.

Proposition 4.3. *Let $A \subset PSH(X, \omega)$ and let $L : A \rightarrow \mathbb{R} \cup \{-\infty\}$ be a convex and non-decreasing function satisfying the scaling property $L(u + c) = L(u) + c$ for any $c \in \mathbb{R}$. Then*

(i) *if L is finite valued on a weakly compact convex set $K \subset A$, then $L(K)$ is bounded;*

(ii) *if $\mathcal{E}^1(X, \omega, \psi) \subset A$ and L is finite valued on $\mathcal{E}^1(X, \omega, \psi)$ then $\sup_{\{u \in \mathcal{E}_C^1(X, \omega, \psi) : \sup_X u \leq 0\}} |L| = O(C^{1/2})$ as $C \rightarrow \infty$.*

4.1 When is L_μ continuous?

The continuity of L_μ is a hard problem. However we can characterize its continuity on some weakly compact sets as the next Theorem shows.

Theorem 4.4. *Let μ be a non-pluripolar probability measure, and let $K \subset PSH(X, \omega)$ be a compact convex set such that L_μ is finite on K , the set $\{P_\omega[u] : u \in K\} \subset \mathcal{M}$ is totally ordered and its closure in $PSH(X, \omega)$ has at most one element in $\mathcal{M} \setminus \mathcal{M}^+$. Suppose also that there exists $C \in \mathbb{R}$ such that $|E_{P_\omega[u]}(u)| \leq C$ for any $u \in K$. Then the following properties are equivalent:*

(i) *L_μ is continuous on K ;*

(ii) *the map $\tau : K \rightarrow L^1(\mu)$, $\tau(u) := u - P_\omega[u]$ is continuous;*

(iii) *the set $\tau(K) \subset L^1(\mu)$ is uniformly integrable, i.e.*

$$\int_{t=m}^{\infty} \mu\{u \leq P_\omega[u] - t\} \rightarrow 0$$

as $m \rightarrow \infty$, uniformly for $u \in K$.

Proof. We first observe that if $u_k \in K$ converges to $u \in K$ then by Lemma 3.13 $\psi_k \rightarrow \psi$ where we set $\psi_k := P_\omega[u_k]$, $\psi := P_\omega[u]$.

Then we can proceed exactly as in Theorem 3.10 in [4] to get the equivalence between (i) and (ii), (ii) \Rightarrow (iii) and the fact that the graph of τ is closed. It is important to underline that (iii) is equivalent to say that $\tau(K)$ is *weakly* relative compact by Dunford-Pettis Theorem, i.e. with respect to the weak topology on $L^1(\mu)$ induced by $L^\infty(\mu) = L^1(\mu)^*$.

Finally assuming that (iii) holds, it remains to prove (i). So, letting $u_k, u \in K$ such that $u_k \rightarrow u$, we have to show that $\int_X \tau(u_k)\mu \rightarrow \int_X \tau(u)\mu$. Since $\tau(K) \subset L^1(\mu)$ is bounded, unless considering a subsequence, we may suppose $\int_X \tau(u_k)\mu \rightarrow L \in \mathbb{R}$. By Fatou's Lemma,

$$L = \lim_{k \rightarrow \infty} \int_X \tau(u_k)\mu \leq \int_X \tau(u)\mu. \quad (12)$$

Then for any $k \in \mathbb{N}$ the closed convex envelope

$$C_k := \overline{\text{Conv}\{\tau(u_j) : j \geq k\}},$$

is weakly closed in $L^1(\mu)$ by Hahn-Banach Theorem, which implies that C_k is weakly compact since it is contained in $\tau(K)$. Thus since C_k is a decreasing sequence of non-empty weakly compact sets, there exists $f \in \bigcap_{k \geq 1} C_k$ and there exist elements $v_k \in \text{Conv}(u_j : j \geq k)$ given as finite convex combination such that

$\tau(v_k) \rightarrow f$ in $L^1(\mu)$. Moreover by the closed graph property $f = \tau(u)$ since $v_k \rightarrow u$ as a consequence of $u_k \rightarrow u$. On the other hand by Proposition 4.2.(iv) we get

$$\int_X \tau(v_k) \mu \leq \sum_{l=1}^{m_k} a_{l,k} \int_X \tau(u_{k_l}) \mu$$

if $v_k = \sum_{l=1}^{m_k} a_{l,k} u_{k_l}$. Hence $L \geq \int_X \tau(u) \mu$, which together (12) implies $L = \int_X \tau(u) \mu$ and concludes the proof. \square

Corollary 4.5. *Let $\psi \in \mathcal{M}^+$ and $\mu \in \mathcal{C}_{A,\psi}$. Then L_μ is continuous on $\mathcal{E}_C^1(X, \omega, \psi)$ for any $C \in \mathbb{R}_{>0}$. In particular if $\mu = MA_\omega(u)/V_\psi$ for $u \in \mathcal{E}^1(X, \omega, \psi)$ with ψ -relative minimal singularities then L_μ is continuous on $\mathcal{E}_C^1(X, \omega, \psi)$ for any $C \in \mathbb{R}_{>0}$.*

Proof. With the notations of Theorem 4.4, $\tau(\mathcal{E}_C^1(X, \omega, \psi))$ is bounded in $L^2(\mu)$ by Lemma 3.11. Hence by Holder's inequality $\tau(\mathcal{E}_C^1(X, \omega, \psi))$ is uniformly integrable and Theorem 4.4 yields the continuity of L_μ on $\mathcal{E}_C^1(X, \omega, \psi)$ for any $C \in \mathbb{R}_{>0}$.

The last assertion follows directly from Proposition 3.10. \square

The following Lemma will be essential to prove Theorem A, Theorem B.

Lemma 4.6. *Let $\varphi \in \mathcal{H}_\omega$ and let $\mathcal{A} \subset \mathcal{M}$ be a totally ordered subset. Set also $v_\psi := P_\omega[\psi](\varphi)$ for any $\psi \in \mathcal{A}$. Then the actions $\{V_\psi L_{MA_\omega(v_\psi)/V_\psi}\}_{\psi \in \mathcal{A}}$ take finite values and they are equicontinuous on any compact set $K \subset PSH(X, \omega)$ such that $\{P_\omega[u] : u \in K\}$ is a totally ordered set whose closure in $PSH(X, \omega)$ has at most one element in $\mathcal{M} \setminus \mathcal{M}^+$ and such that $|E_{P_\omega[u]}(u)| \leq C$ uniformly for any $u \in K$. If $\psi \in \mathcal{M} \setminus \mathcal{M}^+$, for the action $V_\psi L_{MA_\omega(v_\psi)/V_\psi}$ we mean the null action. In particular if $\psi_k \rightarrow \psi$ monotonically almost everywhere and $\{u_k\}_{k \in \mathbb{N}} \subset K$ converges weakly to $u \in K$, then*

$$\int_X (u_k - P_\omega[u_k]) MA_\omega(v_{\psi_k}) \rightarrow \int_X (u - P_\omega[u]) MA_\omega(v_\psi). \quad (13)$$

Proof. By Theorem 2.2, $|V_\psi L_{MA_\omega(v_\psi)/V_\psi}(u)| \leq \int_X |u - P_\omega[u]| MA_\omega(\varphi)$ for any $u \in PSH(X, \omega)$ and any $\psi \in \mathcal{A}$, so the actions in the statement assume finite values. Then the equicontinuity on any weak compact set $K \subset PSH(X, \omega)$ satisfying the assumptions of the Lemma follows from

$$V_\psi \left| L_{MA_\omega(v_\psi)/V_\psi}(w_1) - L_{MA_\omega(v_\psi)/V_\psi}(w_2) \right| \leq \int_X |w_1 - P_\omega[w_1] - w_2 + P_\omega[w_2]| MA_\omega(\varphi)$$

for any $w_1, w_2 \in PSH(X, \omega)$ since $MA_\omega(\varphi)$ is a volume form on X and $P_\omega[w_k] \rightarrow P_\omega[w]$ if $\{w_k\}_{k \in \mathbb{N}} \subset K$ converges to $w \in K$ under our hypothesis by Lemma 3.13.

For the second assertion, if $\psi_k \searrow \psi$ (resp. $\psi_k \nearrow \psi$ almost everywhere), letting $f_k, f \in L^\infty$ such that $MA_\omega(v_{\psi_k}) = f_k MA_\omega(\varphi)$ and $MA_\omega(v_\psi) = f MA_\omega(\varphi)$ (Theorem 2.2), we have $0 \leq f_k \leq 1$, $0 \leq f \leq 1$ and $\{f_k\}_{k \in \mathbb{N}}$ is a monotone sequence. Therefore $f_k \rightarrow f$ in L^p for any $p > 1$ as $k \rightarrow \infty$ which implies

$$\int_X (u - P_\omega[u]) MA_\omega(v_{\psi_k}) \rightarrow \int_X (u - P_\omega[u]) MA_\omega(v_\psi)$$

as $k \rightarrow \infty$ since $MA_\omega(\varphi)$ is a volume form. Hence (13) follows since by the first part of the proof

$$\int_X (u_k - P_\omega[u_k] - u + P_\omega[u]) MA_\omega(v_{\psi_k}) \rightarrow 0.$$

\square

5 Theorem A

In this section we fix $\psi \in \mathcal{M}^+$ and using a variational approach we first prove the bijectivity of the Monge-Ampère operator between $\mathcal{E}_{norm}^1(X, \omega, \psi)$ and $\mathcal{M}^1(X, \omega, \psi)$, and then we prove that it is actually a homeomorphism considering the strong topologies.

5.1 Degenerate complex Monge–Ampère equations.

Letting μ be a probability measure and $\psi \in \mathcal{M}$, we define the functional $F_{\mu,\psi} : \mathcal{E}^1(X, \omega, \psi) \rightarrow \mathbb{R} \cup \{-\infty\}$ as

$$F_{\mu,\psi}(u) := (E_\psi - V_\psi L_\mu)(u)$$

where we recall that $L_\mu(u) = \lim_{j \rightarrow \infty} L_\mu(\max(u, \psi - j)) = \lim_{j \rightarrow \infty} \int_X (\max(u, \psi - j) - \psi) \mu$ (see section 4). $F_{\mu,\psi}$ is clearly a translation invariant functional and $F_{\mu,\psi} \equiv 0$ for any μ if $V_\psi = 0$.

Proposition 5.1. *Let μ be a probability measure, $\psi \in \mathcal{M}^+$ and let $F := F_{\mu,\psi}$. If L_μ is continuous then F is upper semicontinuous on $\mathcal{E}^1(X, \omega, \psi)$. Moreover if L_μ is finite valued on $\mathcal{E}^1(X, \omega, \psi)$ then there exist $A, B > 0$ such that*

$$F(v) \leq -Ad(\psi, v) + B$$

for any $v \in \mathcal{E}_{norm}^1(X, \omega, \psi)$, i.e. F is d -coercive. In particular F is upper semicontinuous on $\mathcal{E}^1(X, \omega, \psi)$ and d -coercive on $\mathcal{E}_{norm}^1(X, \omega, \psi)$ if $\mu = MA_\omega(u)/V_\psi$ for $u \in \mathcal{E}^1(X, \omega, \psi)$.

Proof. If L_μ is continuous then F is easily upper semicontinuous by Proposition 2.4.

Then, since $d(\psi, v) = -E_\psi(v)$ on $\mathcal{E}_{norm}^1(X, \omega, \psi)$, it is easy to check that the coercivity requested is equivalent to

$$\sup_{\mathcal{E}_C^1(X, \omega, \psi) \cap \mathcal{E}_{norm}^1(X, \omega, \psi)} |L_\mu| \leq \frac{(1-A)}{V_\psi} C + O(1),$$

which holds by Proposition 4.3.(ii).

Next assuming $\mu = MA_\omega(u)/V_\psi$ it is sufficient to check the continuity of L_μ since L_μ is finite valued on $\mathcal{E}^1(X, \omega, \psi)$ by Proposition 4.2. We may suppose without loss of generality that $u \leq \psi$. By Proposition 3.7 and Remark 3.3, for any $C \in \mathbb{R}_{>0}$, L_μ restricted to $\mathcal{E}_C^1(X, \omega, \psi)$ is the uniform limit of L_{μ_j} , where $\mu_j := MA_\omega(\max(u, \psi - j))$, since $I_\psi(\max(u, \psi - j), u) \rightarrow 0$ as $j \rightarrow \infty$. Therefore L_μ is continuous on $\mathcal{E}_C^1(X, \omega, \psi)$ since uniform limit of continuous functionals L_{μ_j} (Corollary 4.5). \square

As a consequence of the concavity of E_ψ if $\mu = MA_\omega(u)/V_\psi$ for $u \in \mathcal{E}^1(X, \omega, \psi)$ where $V_\psi > 0$ then

$$J_u^\psi(\psi) = F_{\mu,\psi}(u) = \sup_{\mathcal{E}^1(X, \omega, \psi)} F_{\mu,\psi},$$

i.e. u is a maximizer for $F_{\mu,\psi}$. The other way around also holds as the next result shows.

Proposition 5.2. *Let $\psi \in \mathcal{M}^+$ and let μ be a probability measure such that L_μ is finite valued on $\mathcal{E}^1(X, \omega, \psi)$. Then $\mu = MA_\omega(u)/V_\psi$ for $u \in \mathcal{E}^1(X, \omega, \psi)$ if and only if u is a maximizer of $F_{\mu,\psi}$.*

Proof. As said before, it is clear that $\mu = MA_\omega(u)/V_\psi$ implies that u is a maximizer for $F_{\mu,\psi}$. Conversely, if u is a maximizer of $F_{\mu,\psi}$ then by Theorem 4.22 in [13] $\mu = MA_\omega(u)/V_\psi$. \square

Similarly to [4] we, thus, define the ψ -relative energy for $\psi \in \mathcal{M}$ of a probability measure μ as

$$E_\psi^*(\mu) := \sup_{u \in \mathcal{E}^1(X, \omega, \psi)} F_{\mu,\psi}(u)$$

i.e. essentially as the Legendre transform of E_ψ . It takes non-negative values ($F_{\mu,\psi}(\psi) = 0$) and it is easy to check that E_ψ^* is a convex function.

Moreover defining

$$\mathcal{M}^1(X, \omega, \psi) := \{V_\psi \mu : \mu \text{ is a probability measure satisfying } E_\psi^*(\mu) < \infty\},$$

we note that $\mathcal{M}^1(X, \omega, \psi)$ consists only of the null measure if $V_\psi = 0$ while in $V_\psi > 0$ any probability measure μ such that $V_\psi \mu \in \mathcal{M}^1(X, \omega, \psi)$ is non-pluripolar as the next Lemma shows.

Lemma 5.3. *Let $A \subset X$ be a (locally) pluripolar set. Then there exists $u \in \mathcal{E}^1(X, \omega, \psi)$ such that $A \subset \{u = -\infty\}$. In particular if $V_\psi \mu \in \mathcal{M}^1(X, \omega, \psi)$ for $\psi \in \mathcal{M}^+$ then μ is non-pluripolar.*

Proof. By Corollary 2.11 in [4] there exists $\varphi \in \mathcal{E}^1(X, \omega)$ such that $A \subset \{\varphi = -\infty\}$. Therefore setting $u := P_\omega[\psi](\varphi)$ proves the first part.

Next let $V_\psi \mu \in \mathcal{M}^1(X, \omega, \psi)$ for $\psi \in \mathcal{M}^+$ and μ probability measure and assume by contradiction that μ takes mass on a pluripolar set A . Then by the first part of the proof there exists $u \in \mathcal{E}^1(X, \omega, \psi)$ such that $A \subset \{u = -\infty\}$. On the other hand, since $V_\psi \mu \in \mathcal{M}^1(X, \omega, \psi)$ by definition μ does not charge $\{\psi = -\infty\}$. Thus by Proposition 4.2.(iii) we obtain $L_\mu(u) = -\infty$, which is a contradiction. \square

We can now prove that the Monge-Ampère operation is a bijection between $\mathcal{E}^1(X, \omega, \psi)$ and $\mathcal{M}^1(X, \omega, \psi)$.

Lemma 5.4. *Let $\psi \in \mathcal{M}^+$ and let $\mu \in \mathcal{C}_{A, \psi}$ where $A \in \mathbb{R}$. Then there exists $u \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ maximizing $F_{\mu, \psi}$.*

Proof. By Lemma 3.11 L_μ is finite valued on $\mathcal{E}^1(X, \omega, \psi)$, and it is continuous on $\mathcal{E}_C^1(X, \omega, \psi)$ for any $C \in \mathbb{R}$ thanks to Corollary 4.5. Therefore it follows from Proposition 5.1 that $F_{\mu, \psi}$ is upper semicontinuous and d -coercive on $\mathcal{E}_{norm}^1(X, \omega, \psi)$. Hence $F_{\mu, \psi}$ admits a maximizer $u \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ as easy consequence of the weak compactness of $\mathcal{E}_C^1(X, \omega, \psi)$. \square

Proposition 5.5. *Let $\psi \in \mathcal{M}^+$. Then the Monge-Ampère map $MA : \mathcal{E}_{norm}^1(X, \omega, \psi) \rightarrow \mathcal{M}^1(X, \omega, \psi)$, $u \rightarrow MA_\omega(u)$ is bijective. Furthermore if $V_\psi \mu = MA_\omega(u) \in \mathcal{M}^1(X, \omega, \psi)$ for $u \in \mathcal{E}^1(X, \omega, \psi)$ then any maximizing sequence $u_k \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ for $F_{\mu, \psi}$ necessarily converges weakly to u .*

Proof. The proof is inspired by Theorem 4.7 in [4].

The map is well-defined as a consequence of Proposition 5.1, i.e. $MA_\omega(u) \in \mathcal{M}^1(X, \omega, \psi)$ for any $u \in \mathcal{E}^1(X, \omega, \psi)$. Moreover the injectivity follows from Theorem 4.8 in [14].

Let $u_k \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ be a sequence such that $F_{\mu, \psi}(u_k) \nearrow \sup_{\mathcal{E}^1(X, \omega, \psi)} F_{\mu, \psi}$ where $\mu = MA_\omega(u)/V_\psi$ is a probability measure and $u \in \mathcal{E}_{norm}^1(X, \omega, \psi)$. Up to considering a subsequence, we may also assume that $u_k \rightarrow v \in PSH(X, \omega)$. Then, by the upper semicontinuity and the d -coercivity of $F_{\mu, \psi}$ (Proposition 5.1) it follows that $v \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ and $F_{\mu, \psi}(v) = \sup_{\mathcal{E}^1(X, \omega, \psi)} F_{\mu, \psi}$. Thus by Proposition 5.2 we get $\mu = MA_\omega(v)/V_\psi$. Hence $v = u$ since $\sup_X v = \sup_X u = 0$.

Then let μ be a probability measure such that $V_\psi \mu \in \mathcal{M}^1(X, \omega, \psi)$. Again by Proposition 5.2, to prove the existence of $u \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ such that $\mu = MA_\omega(u)/V_\psi$ it is sufficient to check that $F_{\mu, \psi}$ admits a maximum over $\mathcal{E}_{norm}^1(X, \omega, \psi)$. Moreover by Proposition 5.1 we also know that $F_{\mu, \psi}$ is d -coercive on $\mathcal{E}_{norm}^1(X, \omega, \psi)$. Thus if there exists a constant $A > 0$ such that $\mu \in \mathcal{C}_{A, \psi}$ then Corollary 4.5 leads to the upper semicontinuity of $F_{\mu, \psi}$ which clearly implies that $V_\psi \mu = MA_\omega(u)$ for $u \in \mathcal{E}^1(X, \omega, \psi)$ since $\mathcal{E}_C^1(X, \omega, \psi) \subset PSH(X, \omega)$ is compact for any $C \in \mathbb{R}_{>0}$.

In the general case by Lemma 4.26 in [13] (see also [8]) μ is absolutely continuous with respect to $\nu \in \mathcal{C}_{1, \psi}$ using also that μ is a non-pluripolar measure (Lemma 5.3). Therefore letting $f \in L^1(\nu)$ such that $\mu = f\nu$, we define for any $k \in \mathbb{N}$

$$\mu_k := (1 + \epsilon_k) \min(f, k)\nu$$

where $\epsilon_k > 0$ are chosen so that μ_k is a probability measure, noting that $(1 + \epsilon_k) \min(f, k) \rightarrow f$ in $L^1(\nu)$. Then by Lemma 5.4 it follows that $\mu_k = MA_\omega(u_k)/V_\psi$ for $u_k \in \mathcal{E}_{norm}^1(X, \omega, \psi)$.

Moreover by weak compactness, without loss of generality, we may also assume that $u_k \rightarrow u \in PSH(X, \omega)$. Note that $u \leq \psi$ since $u_k \leq \psi$ for any $k \in \mathbb{N}$. Then by Lemma 2.8 in [14] we obtain

$$MA_\omega(u) \geq V_\psi f \nu = V_\psi \mu,$$

which implies $MA_\omega(u) = V_\psi \mu$ by [24] since u is more singular than ψ and μ is a probability measure. It remains to prove that $u \in \mathcal{E}^1(X, \omega, \psi)$.

It is not difficult to see that $\mu_k \leq 2\mu$ for $k \gg 0$, thus Proposition 4.3 implies that there exists a constant $B > 0$ such that

$$\sup_{\mathcal{E}_C^1(X, \omega, \psi)} |L_{\mu_k}| \leq 2 \sup_{\mathcal{E}_C^1(X, \omega, \psi)} |L_\mu| \leq 2B(1 + C^{1/2})$$

for any $C \in \mathbb{R}_{>0}$. Therefore

$$J_{u_k}^\psi(\psi) = E_\psi(u_k) + V_\psi |L_{\mu_k}(u_k)| \leq \sup_{C>0} (2V_\psi B(1 + C^{1/2}) - C)$$

and Lemma 3.1 yields $d(\psi, u_k) \leq D$ for a uniform constant D , i.e. $u_k \in \mathcal{E}_{D'}^1(X, \omega, \psi)$ for any $k \in \mathbb{N}$ for a uniform constant D' (Remark 3.3). Hence since $\mathcal{E}_{D'}^1(X, \omega, \psi)$ is weakly compact we obtain $u \in \mathcal{E}_{D'}^1(X, \omega, \psi)$. \square

5.2 Proof of Theorem A.

We first need to explore further the properties of the strong topology on $\mathcal{E}^1(X, \omega, \psi)$.

By Proposition 3.6 the strong convergence implies the weak convergence. Moreover the strong topology is the coarsest refinement of the weak topology such that $E_\psi(\cdot)$ becomes continuous.

Proposition 5.6. *Let $\psi \in \mathcal{M}^+$ and $u_k, u \in \mathcal{E}^1(X, \omega, \psi)$. Then $u_k \rightarrow u$ strongly if and only if $u_k \rightarrow u$ weakly and $E_\psi(u_k) \rightarrow E_\psi(u)$.*

Proof. Assume that $u_k \rightarrow u$ weakly and that $E_\psi(u_k) \rightarrow E_\psi(u)$. Then $w_k := (\sup\{u_j : j \geq k\})^* \in \mathcal{E}^1(X, \omega, \psi)$ and it decreases to u . Thus by Proposition 2.4 $E_\psi(w_k) \rightarrow E_\psi(u)$ and

$$d(u_k, u) \leq d(u_k, w_k) + d(w_k, u) = 2E_\psi(w_k) - E_\psi(u_k) - E_\psi(u) \rightarrow 0.$$

Conversely, assuming that $d(u_k, u) \rightarrow 0$, we immediately get that $u_k \rightarrow u$ weakly as said above (Proposition 3.6). Moreover $\sup_X u_k, \sup_X u \leq A$ uniformly for a constant $A \in \mathbb{R}$. Thus

$$|E_\psi(u_k) - E_\psi(u)| = |d(\psi + A, u_k) - d(\psi + A, u)| \leq d(u_k, u) \rightarrow 0,$$

which concludes the proof. \square

Then we also observe that the strong convergence implies the convergence in ψ' -capacity for any $\psi' \in \mathcal{M}^+$.

Proposition 5.7. *Let $\psi \in \mathcal{M}^+$ and $u_k, u \in \mathcal{E}^1(X, \omega, \psi)$ such that $d(u_k, u) \rightarrow 0$. Then there exists a subsequence $\{u_{k_j}\}_{j \in \mathbb{N}}$ such that $w_j := (\sup\{u_{k_h} : h \geq j\})^*$, $v_j := P_\omega(u_{k_j}, u_{k_{j+1}}, \dots)$ belong to $\mathcal{E}^1(X, \omega, \psi)$ and converge monotonically almost everywhere to u . In particular $u_k \rightarrow u$ in ψ' -capacity for any $\psi' \in \mathcal{M}^+$ and $MA_\omega(u_{k_j}^j, \psi^{n-j}) \rightarrow MA_\omega(u^j, \psi^{n-j})$ weakly for any $j = 0, \dots, n$.*

Proof. Since the strong convergence implies the weak convergence by Proposition 5.6 it is clear that $w_k \in \mathcal{E}^1(X, \omega, \psi)$ and that it decreases to u . In particular up to considering a subsequence we may assume that $d(u_k, w_k) \leq 1/2^k$ for any $k \in \mathbb{N}$.

Next for any $j \geq k$ we set $v_{k,j} := P_\omega(u_k, \dots, u_j) \in \mathcal{E}^1(X, \omega, \psi)$ and $v_{k,j}^u := P_\omega(v_{k,j}, u) \in \mathcal{E}^1(X, \omega, \psi)$. Then it follows from Proposition 2.4 and Lemma 3.7 in [13] that

$$\begin{aligned} d(u, v_{k,j}^u) &\leq \int_X (u - v_{k,j}^u) MA_\omega(v_{k,j}^u) \leq \int_{\{v_{k,j}^u = v_{k,j}\}} (u - v_{k,j}) MA_\omega(v_{k,j}) \leq \\ &\leq \sum_{s=k}^j \int_X (w_s - u_s) MA_\omega(u_s) \leq (n+1) \sum_{s=k}^j d(w_s, u_s) \leq \frac{(n+1)}{2^{k-1}}. \end{aligned}$$

Therefore by Proposition 3.15 $v_{k,j}^u$ decreases (hence converges strongly) to a function $\phi_k \in \mathcal{E}^1(X, \omega, \psi)$ as $j \rightarrow \infty$. Similarly we also observe that

$$d(v_{k,j}, v_{k,j}^u) \leq \int_{\{v_{k,j}^u = u\}} (v_{k,j} - u) MA_\omega(u) \leq \int_X |v_{k,1} - u| MA_\omega(u) \leq C$$

uniformly in j by Corollary 3.5. Hence by definition $d(u, v_{k,j}) \leq C + \frac{(n+1)}{2^{k-1}}$, i.e. $v_{k,j}$ decreases and converges strongly as $j \rightarrow \infty$ to the function $v_k = P_\omega(u_k, u_{k+1}, \dots) \in \mathcal{E}^1(X, \omega, \psi)$ again by Proposition 3.15. Moreover by construction $u_k \geq v_k \geq \phi_k$ since $v_k \leq v_{k,j} \leq u_k$ for any $j \geq k$. Hence

$$d(u, v_k) \leq d(u, \phi_k) \leq \frac{(n+1)}{2^{k-1}} \rightarrow 0$$

as $k \rightarrow \infty$, i.e. $v_k \nearrow u$ strongly.

The convergence in ψ' -capacity for $\psi' \in \mathcal{M}^+$ is now clearly an immediate consequence. Indeed by an easy contradiction argument it is enough to prove that any arbitrary subsequence, which we will keep denoting with $\{u_k\}_{k \in \mathbb{N}}$ for the sake of simplicity, admits a further subsequence $\{u_{k_j}\}_{j \in \mathbb{N}}$ converging in ψ' -capacity to u . Thus taking the subsequence satisfying $v_j \leq u_{k_j} \leq w_j$ where v_j, w_j are the monotonic sequence of the first part of the Proposition, the convergence in ψ' -capacity follows from the inclusions

$$\{|u - u_{k_j}| > \delta\} = \{u - u_{k_j} > \delta\} \cup \{u_{k_j} - u > \delta\} \subset \{u - v_j > \delta\} \cup \{w_j - u > \delta\}$$

for any $\delta > 0$. Finally Lemma 2.12 gives the weak convergence of the measures. \square

We can now endow the set $\mathcal{M}^1(X, \omega, \psi) = \{V_\psi \mu : \mu \text{ is a probability measure satisfying } E_\psi^*(\mu) < +\infty\}$ (subsection 5.1) with its natural strong topology given as the coarsest refinement of the weak topology such that $E_\psi^*(\cdot)$ becomes continuous, and prove our Theorem A.

Theorem A. *Let $\psi \in \mathcal{M}^+$. Then*

$$MA_\omega : (\mathcal{E}_{norm}^1(X, \omega, \psi), d) \rightarrow (\mathcal{M}^1(X, \omega, \psi), \text{strong})$$

is a homeomorphism.

Proof. The map is bijective as immediate consequence of Proposition 5.5.

Next, letting $u_k \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ converging strongly to $u \in \mathcal{E}_{norm}^1(X, \omega, \psi)$, Proposition 5.7 gives the weak convergence of $MA_\omega(u_k) \rightarrow MA_\omega(u)$ as $k \rightarrow \infty$. Moreover since $E_\psi^*(MA_\omega(v)/V_\psi) = J_v^\psi(\psi)$ for any $v \in \mathcal{E}^1(X, \omega, \psi)$, we get

$$\begin{aligned} & \left| E_\psi^*(MA_\omega(u_k)/V_\psi) - E_\psi^*(MA_\omega(u)/V_\psi) \right| \leq \\ & \leq |E_\psi(u_k) - E_\psi(u)| + \left| \int_X (\psi - u_k) MA_\omega(u_k) - \int_X (\psi - u) MA_\omega(u) \right| \leq \\ & \leq |E_\psi(u_k) - E_\psi(u)| + \left| \int_X (\psi - u_k)(MA_\omega(u_k) - MA_\omega(u)) \right| + \int_X |u_k - u| MA_\omega(u). \end{aligned} \quad (14)$$

Hence $MA_\omega(u_k) \rightarrow MA_\omega(u)$ strongly in $\mathcal{M}^1(X, \omega, \psi)$ since each term on the right-hand side of (14) goes to 0 as $k \rightarrow +\infty$ combining Proposition 5.6, Proposition 3.7 and Corollary 3.5 recalling that by Proposition 3.4 $I_\psi(u_k, u) \rightarrow 0$ as $k \rightarrow \infty$.

Conversely, suppose that $MA_\omega(u_k) \rightarrow MA_\omega(u)$ strongly in $\mathcal{M}^1(X, \omega, \psi)$ where $u_k, u \in \mathcal{E}_{norm}^1(X, \omega, \psi)$. Then, letting $\{\varphi_j\}_{j \in \mathbb{N}} \subset \mathcal{H}_\omega$ such that $\varphi_j \searrow u$ ([6]) and setting $v_j := P_\omega[\psi](\varphi_j)$, by Lemma 3.1

$$\begin{aligned} (n+1)I_\psi(u_k, v_j) & \leq E_\psi(u_k) - E_\psi(v_j) + \int_X (v_j - u_k) MA_\omega(u_k) = \\ & = E_\psi^*(MA_\omega(u_k)/V_\psi) - E_\psi^*(MA_\omega(v_j)/V_\psi) + \int_X (v_j - \psi)(MA_\omega(u_k) - MA_\omega(v_j)). \end{aligned} \quad (15)$$

By construction and the first part of the proof, it follows that $E_\psi^*(MA_\omega(u_k)/V_\psi) - E_\psi^*(MA_\omega(v_j)/V_\psi) \rightarrow 0$ as $k, j \rightarrow \infty$. While setting $f_j := v_j - \psi$ we want to prove that

$$\limsup_{k \rightarrow \infty} \int_X f_j MA_\omega(u_k) = \int_X f_j MA_\omega(u),$$

which would imply $\limsup_{j \rightarrow \infty} \limsup_{k \rightarrow \infty} I_\psi(u_k, v_j) = 0$ since $\int_X f_j (MA_\omega(u) - MA_\omega(v_j)) \rightarrow 0$ as a consequence of Propositions 3.7 and 3.4.

We observe that $\|f_j\|_{L^\infty} \leq \|\varphi_j\|_{L^\infty}$ by Proposition 2.10 and we denote by $\{f_j^s\}_{s \in \mathbb{N}} \subset C^\infty$ a sequence of smooth functions converging in capacity to f_j such that $\|f_j^s\|_{L^\infty} \leq 2\|f_j\|_{L^\infty}$. We recall here briefly how to construct such sequence. Let $\{g_j^s\}_{s \in \mathbb{N}}$ be the sequence of bounded functions converging in capacity to f_j defined as $g_j^s := \max(v_j, -s) - \max(\psi, -s)$. We have that $\|g_j^s\|_{L^\infty} \leq \|f_j\|_{L^\infty}$ and that $\max(v_j, -s), \max(\psi, -s) \in PSH(X, \omega)$. Therefore by a regularization process (see for instance [6]) and a diagonal argument we can now construct a sequence $\{f_j^s\}_{j \in \mathbb{N}} \subset C^\infty$ converging in capacity to f_j such that $\|f_j^s\|_{L^\infty} \leq 2\|g_j^s\|_{L^\infty} \leq 2\|f_j\|_{L^\infty}$ where $f_j^s = v_j^s - \psi^s$ with v_j^s, ψ^s quasi-psh functions decreasing respectively to v_j, ψ .

Then letting $\delta > 0$ we have

$$\int_X (f_j - f_j^s) MA_\omega(u_k) \leq \delta V_\psi + 3\|\varphi_j\|_{L^\infty} \int_{\{f_j - f_j^s > \delta\}} MA_\omega(u_k) \leq \delta V_\psi + 3\|\varphi_j\|_{L^\infty} \int_{\{\psi^s - \psi > \delta\}} MA_\omega(u_k)$$

from the trivial inclusion $\{f_j - f_j^s > \delta\} \subset \{\psi^s - \psi > \delta\}$. Therefore

$$\begin{aligned} \limsup_{s \rightarrow \infty} \limsup_{k \rightarrow \infty} \int_X (f_j - f_j^s) MA_\omega(u_k) & \leq \delta V_\psi + \limsup_{s \rightarrow \infty} \limsup_{k \rightarrow \infty} \int_{\{\psi^s - \psi \geq \delta\}} MA_\omega(u_k) \leq \\ & \leq \delta V_\psi + \limsup_{s \rightarrow \infty} \int_{\{\psi^s - \psi \geq \delta\}} MA_\omega(u) = \delta V_\psi, \end{aligned}$$

where we used that $\{\psi^s - \psi \geq \delta\}$ is a closed set in the plurifine topology. Hence since $f_j^s \in C^\infty$ we obtain

$$\begin{aligned} \limsup_{k \rightarrow \infty} \int_X f_j MA_\omega(u_k) & = \limsup_{s \rightarrow \infty} \limsup_{k \rightarrow \infty} \left(\int_X (f_j - f_j^s) MA_\omega(u_k) + \int_X f_j^s MA_\omega(u_k) \right) \leq \\ & \leq \limsup_{s \rightarrow \infty} \int_X f_j^s MA_\omega(u) = \int_X f_j MA_\omega(u), \end{aligned}$$

which as said above implies $I_\psi(u_k, v_j) \rightarrow 0$ letting $k, j \rightarrow \infty$ in this order.

Next, again by Lemma 3.1, we obtain $u_k \in \mathcal{E}_C^1(X, \omega, \psi)$ for some $C \in \mathbb{N}$ big enough since $J_{u_k}^\psi(\psi) = E_\psi^*(MA_\omega(u_k)/V_\psi)$. In particular, up to considering a subsequence, $u_k \rightarrow w \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ weakly by Proposition 3.15. Observe also that by Proposition 3.7

$$\left| \int_X (\psi - u_k)(MA_\omega(v_j) - MA_\omega(u_k)) \right| \rightarrow 0 \quad (16)$$

as $k, j \rightarrow \infty$ in this order. Moreover by Proposition 3.14 and Lemma 4.6

$$\begin{aligned} \limsup_{k \rightarrow \infty} \left(E_\psi^*(MA_\omega(u_k)/V_\psi) + \int_X (\psi - u_k)(MA_\omega(v_j) - MA_\omega(u_k)) \right) &= \\ &= \limsup_{k \rightarrow \infty} \left(E_\psi(u_k) + \int_X (\psi - u_k)MA_\omega(v_j) \right) \leq E_\psi(w) + \int_X (\psi - w)MA_\omega(v_j). \end{aligned} \quad (17)$$

Therefore combining (16) and (17) with the strong convergence of v_j to u we obtain

$$\begin{aligned} E_\psi(u) + \int_X (\psi - u)MA_\omega(u) &= \lim_{k \rightarrow \infty} E_\psi^*(MA_\omega(u_k)/V_\psi) \leq \\ &\leq \limsup_{j \rightarrow \infty} \left(E_\psi(w) + \int_X (\psi - w)MA_\omega(v_j) \right) = E_\psi(w) + \int_X (\psi - w)MA_\omega(u), \end{aligned}$$

i.e. w is a maximizer of $F_{MA_\omega(u)/V_\psi, \psi}$. Hence $w = u$ (Proposition 5.5), i.e. $u_k \rightarrow u$ weakly. Furthermore again by Lemma 3.1 and Lemma 4.6

$$\begin{aligned} \limsup_{k \rightarrow \infty} (E_\psi(v_j) - E_\psi(u_k)) &\leq \limsup_{k \rightarrow \infty} \left(\frac{n}{n+1} I_\psi(u_k, v_j) + \left| \int_X (u_k - v_j)MA_\omega(v_j) \right| \right) \leq \\ &\leq \left| \int_X (u - v_j)MA_\omega(v_j) \right| + \limsup_{k \rightarrow \infty} \frac{n}{n+1} I_\psi(u_k, v_j). \end{aligned} \quad (18)$$

Finally letting $j \rightarrow \infty$, since $v_j \searrow u$ strongly, we obtain $\liminf_{j \rightarrow \infty} E_\psi(u_k) \geq \lim_{j \rightarrow \infty} E_\psi(v_j) = E_\psi(u)$ which implies that $E_\psi(u_k) \rightarrow E_\psi(u)$ and that $u_k \rightarrow u$ strongly by Proposition 5.6. \square

The main difference between the proof of Theorem A with respect to the same result in the absolute setting, i.e. when $\psi = 0$, is that for fixed $u \in \mathcal{E}^1(X, \omega, \psi)$ the action $\mathcal{M}^1(X, \omega, \psi) \ni MA_\omega(v) \rightarrow \int_X (u - \psi)MA_\omega(v)$ is not a priori continuous with respect to the weak topologies of measures even if we restrict the action on $\mathcal{M}_C^1(X, \omega, \psi) := \{V_\psi \mu : E_\psi^*(\mu) \leq C\}$ for $C \in \mathbb{R}$ while in the absolute setting this is given by Proposition 1.7. in [3] where the authors used the fact that any $u \in \mathcal{E}^1(X, \omega)$ can be approximated inside the class $\mathcal{E}^1(X, \omega)$ by a sequence of continuous functions.

6 Strong Topologies.

In this section we investigate the strong topology on $X_{\mathcal{A}}$ in detail, proving that it is the coarsest refinement of the weak topology such that $E(\cdot)$ becomes continuous (Theorem 6.2) and proving that the strong convergence implies the convergence in ψ -capacity for any $\psi \in \mathcal{M}^+$ (Theorem 6.3), i.e. we extend all the typical properties of the L^1 -metric geometry to the bigger space $X_{\mathcal{A}}$, justifying further the construction of the distance $d_{\mathcal{A}}$ ([23]) and its naturality. Moreover we define the set $Y_{\mathcal{A}}$, and we prove Theorem B.

6.1 About $(X_{\mathcal{A}}, d_{\mathcal{A}})$.

First we prove that the strong convergence in $X_{\mathcal{A}}$ implies the weak convergence, recalling that for weak convergence of $u_k \in \mathcal{E}^1(X, \omega, \psi_k)$ to $P_{\psi_{\min}}$ where $\psi_{\min} \in \mathcal{M}$ with $V_{\psi_{\min}} = 0$ we mean that $|\sup_X u_k| \leq C$ and that any weak accumulation point of $\{u_k\}_{k \in \mathbb{N}}$ is more singular than ψ_{\min} .

Proposition 6.1. *Let $u_k, u \in X_{\mathcal{A}}$ such that $u_k \rightarrow u$ strongly. If $u \neq P_{\psi_{\min}}$ then $u_k \rightarrow u$ weakly. If instead $u = P_{\psi_{\min}}$ the following dichotomy holds:*

- (i) $u_k \rightarrow P_{\psi_{\min}}$ weakly;
- (ii) $\limsup_{k \rightarrow \infty} |\sup_X u_k| = +\infty$.

Proof. The dichotomy for the case $u = P_{\psi_{\min}}$ follows by definition. Indeed if $|\sup_X u_k| \leq C$ and $d_{\mathcal{A}}(u_k, u) \rightarrow 0$ as $k \rightarrow \infty$, then $V_{\psi_k} \rightarrow V_{\psi_{\min}} = 0$ by Proposition 2.11.(iv) which implies that $\psi_k \rightarrow \psi_{\min}$ by Lemma 3.12. Hence any weak accumulation point u of $\{u_k\}_{k \in \mathbb{N}}$ satisfies $u \leq \psi_{\min} + C$. Thus, let $\psi_k, \psi \in \mathcal{A}$ such that $u_k \in \mathcal{E}^1(X, \omega, \psi_k)$ and $u \in \mathcal{E}^1(X, \omega, \psi)$ where $\psi \in \mathcal{M}^+$. Observe that

$$d(u_k, \psi_k) \leq d_{\mathcal{A}}(u_k, u) + d(u, \psi) + d_{\mathcal{A}}(\psi, \psi_k) \leq A$$

for a uniform constant $A > 0$ by Proposition 2.11.(iv)

On the other hand for any $j \in \mathbb{N}$ by [6] there exists $h_j \in \mathcal{H}_{\omega}$ such that $h_j \geq u$, $\|h_j - u\|_{L^1} \leq 1/j$ and $d(u, P_{\omega}[\psi](h_j)) \leq 1/j$. In particular by the triangle inequality and Proposition 2.11 we have

$$\limsup_{k \rightarrow \infty} d(P_{\omega}[\psi_k](h_j), \psi_k) \leq \limsup_{k \rightarrow \infty} \left(d_{\mathcal{A}}(P_{\omega}[\psi_k](h_j), P_{\omega}[\psi](h_j)) + \frac{1}{j} + d(u, \psi) + d(\psi, \psi_k) \right) \leq d(u, \psi) + \frac{1}{j}, \quad (19)$$

Similarly again by the triangle inequality and Proposition 2.11

$$\limsup_{k \rightarrow \infty} d(u_k, P_{\omega}[\psi_k](h_j)) \leq \limsup_{k \rightarrow \infty} \left(d_{\mathcal{A}}(P_{\omega}[\psi_k](h_j), P_{\omega}[\psi](h_j)) + \frac{1}{j} + d_{\mathcal{A}}(u, u_k) \right) \leq \frac{1}{j} \quad (20)$$

and

$$\begin{aligned} \limsup_{k \rightarrow \infty} \|u_k - u\|_{L^1} &\leq \limsup_{k \rightarrow \infty} \left(\|u_k - P_{\omega}[\psi_k](h_j)\|_{L^1} + \|P_{\omega}[\psi_k](h_j) - P_{\omega}[\psi](h_j)\|_{L^1} + \|P_{\omega}[\psi](h_j) - u\|_{L^1} \right) \leq \\ &\leq \frac{1}{j} + \limsup_{k \rightarrow \infty} \|u_k - P_{\omega}[\psi_k](h_j)\|_{L^1} \quad (21) \end{aligned}$$

where we also used Lemma 2.14. In particular from (19) and (20) we deduce that $d(\psi_k, P_{\omega}[\psi_k](h_j)), d(\psi_k, u_k) \leq C$ for a uniform constant $C \in \mathbb{R}$. Next let $\phi_k \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ the unique solution of $MA_{\omega}(\phi_k) = \frac{V_{\psi_k}}{V_0} MA_{\omega}(0)$ and observe that by Proposition 2.4

$$d(\psi_k, \phi_k) = -E_{\psi_k}(\phi_k) \leq \int_X (\psi_k - \phi_k) MA_{\omega}(\phi_k) \leq \frac{V_{\psi_k}}{V_0} \int_X |\phi_k| MA_{\omega}(0) \leq \|\phi_k\|_{L^1} \leq C'$$

since ϕ_k belongs to a compact (hence bounded) subset of $PSH(X, \omega) \subset L^1$. Therefore, since $V_{\psi_k} \geq a > 0$ for $k \gg 0$ big enough, by Proposition 3.6 it follows that there exists a continuous increasing function $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ with $f(0) = 0$ such that

$$\|u_k - P_{\omega}[\psi_k](h_j)\|_{L^1} \leq f(d(u_k, P_{\omega}[\psi_k](h_j)))$$

for any k, j big enough. Hence combining (20) and (21) the convergence requested follows letting $k, j \rightarrow +\infty$ in this order. \square

We can now prove the important characterization of the strong convergence as the coarsest refinement of the weak topology such that $E(\cdot)$ becomes continuous.

Theorem 6.2. *Let $u_k \in \mathcal{E}^1(X, \omega, \psi_k), u \in \mathcal{E}^1(X, \omega, \psi)$ for $\{\psi_k\}_{k \in \mathbb{N}}, \psi \in \overline{\mathcal{A}}$. If $\psi \neq \psi_{\min}$ or $V_{\psi_{\min}} > 0$ then the followings are equivalent:*

- i) $u_k \rightarrow u$ strongly;
- ii) $u_k \rightarrow u$ weakly and $E_{\psi_k}(u_k) \rightarrow E_{\psi}(u)$.

In the case $\psi = \psi_{\min}$ and $V_{\psi_{\min}} = 0$, if $u_k \rightarrow P_{\psi_{\min}}$ weakly and $E_{\psi_k}(u_k) \rightarrow 0$ then $u_k \rightarrow P_{\psi_{\min}}$ strongly. Finally if $d_{\mathcal{A}}(u_k, P_{\psi_{\min}}) \rightarrow 0$ as $k \rightarrow \infty$, then the following dichotomy holds:

- a) $u_k \rightarrow P_{\psi_{\min}}$ weakly and $E_{\psi_k}(u_k) \rightarrow 0$;
- b) $\limsup_{k \rightarrow \infty} |\sup_X u_k| = \infty$.

Proof. Implication (ii) \Rightarrow (i).

Assume that (ii) holds where we include the case $u = P_{\psi_{\min}}$ setting $E_{\psi}(P_{\psi_{\min}}) := 0$. Clearly it is enough to prove that any subsequence of $\{u_k\}_{k \in \mathbb{N}}$ admits a subsequence which is $d_{\mathcal{A}}$ -convergent to u . For the sake of simplicity we denote by $\{u_k\}_{k \in \mathbb{N}}$ the arbitrary initial subsequence, and since \mathcal{A} is totally ordered by Lemma 3.13 we may also assume either $\psi_k \searrow \psi$ or $\psi_k \nearrow \psi$ almost everywhere. In particular even if $u = P_{\psi_{\min}}$

we may suppose that u_k converges weakly to a proper element $v \in \mathcal{E}^1(X, \omega, \psi)$ up to considering a further subsequence by definition of weak convergence to the point $P_{\psi_{\min}}$. In this case by abuse of notation we denote the function v , which depends on the subsequence chosen, by u . Note also that by Hartogs' Lemma we have $u_k \leq \psi_k + A$, $u \leq \psi + A$ for a uniform constant $A \in \mathbb{R}_{\geq 0}$ since $|\sup_X u_k| \leq A$.

In the case $\psi_k \searrow \psi$, $v_k := (\sup\{u_j : j \geq k\})^* \in \mathcal{E}^1(X, \omega, \psi_k)$ decreases to u . Thus $w_k := P_\omega[\psi](v_k) \in \mathcal{E}^1(X, \omega, \psi)$ decreases to u , which implies $d(u, w_k) \rightarrow 0$ as $k \rightarrow \infty$ (if $u = P_{\psi_{\min}}$ we immediately have $w_k = P_{\psi_{\min}}$).

Moreover by Propositions 2.4 and 2.10 it follows that

$$\begin{aligned} E_\psi(u) &= \lim_{k \rightarrow \infty} E_\psi(w_k) = AV_\psi - \lim_{k \rightarrow \infty} d(\psi + A, w_k) \geq \lim_{k \rightarrow \infty} (AV_{\psi_k} - d(\psi_k + A, v_k)) = \\ &= \limsup_{k \rightarrow \infty} E_{\psi_k}(v_k) \geq \lim_{k \rightarrow \infty} E_{\psi_k}(u_k) = E_\psi(u) \end{aligned}$$

since $\psi_k + A = P_\omega[\psi_k](A)$. Hence $\limsup_{k \rightarrow \infty} d(v_k, u_k) = \limsup_{k \rightarrow \infty} d(\psi_k + A, u_k) - d(v_k, \psi_k + A) = \lim_{k \rightarrow \infty} E_{\psi_k}(v_k) - E_{\psi_k}(u_k) = 0$. Thus by the triangle inequality it is sufficient to show that $\limsup_{k \rightarrow \infty} d_{\mathcal{A}}(u, v_k) = 0$.

Next for any $C \in \mathbb{R}$ we set $v_k^C := \max(v_k, \psi_k - C)$, $u^C := \max(u, \psi - C)$ and we observe that $d(\psi_k + A, v_k^C) \rightarrow d(\psi + A, u^C)$ by Proposition 2.11 since $v_k^C \searrow u^C$. This implies that

$$\begin{aligned} d(v_k, v_k^C) &= d(\psi_k + A, v_k) - d(\psi_k + A, v_k^C) = AV_{\psi_k} - E_{\psi_k}(v_k) - d(\psi_k + A, v_k^C) \longrightarrow \\ &\longrightarrow AV_\psi - E_\psi(u) - d(\psi + A, u^C) = d(\psi + A, u) - d(\psi + A, u^C) = d(u, u^C). \end{aligned}$$

Thus, since $u^C \rightarrow u$ strongly, again by the triangle inequality it remains to estimate $d_{\mathcal{A}}(u, v_k^C)$. Fix $\epsilon > 0$ and $\phi_\epsilon \in \mathcal{P}_{\mathcal{H}_\omega}(X, \omega, \psi)$ such that $d(\phi_\epsilon, u) \leq \epsilon$ (by Lemma 2.13). Then letting $\varphi \in \mathcal{H}_\omega$ such that $\phi_\epsilon = P_\omega[\psi](\varphi)$ and setting $\phi_{\epsilon, k} := P_\omega[\psi_k](\varphi)$ by Proposition 2.11 we have

$$\limsup_{k \rightarrow \infty} d_{\mathcal{A}}(u, v_k^C) \leq \limsup_{k \rightarrow \infty} (d(u, \phi_\epsilon) + d_{\mathcal{A}}(\phi_\epsilon, \phi_{\epsilon, k}) + d(\phi_{\epsilon, k}, v_k^C)) \leq \epsilon + d(\phi_\epsilon, u^C) \leq 2\epsilon + d(u, u^C),$$

which concludes the first case of (ii) \Rightarrow (i) by the arbitrariness of ϵ since $u^C \rightarrow u$ strongly in $\mathcal{E}^1(X, \omega, \psi)$. Next assume that $\psi_k \nearrow \psi$ almost everywhere. In this case we clearly may assume $V_{\psi_k} > 0$ for any $k \in \mathbb{N}$. Then $v_k := (\sup\{u_j : j \geq k\})^* \in \mathcal{E}^1(X, \omega, \psi)$ decreases to u . Moreover setting $w_k := P_\omega[\psi_k](v_k) \in \mathcal{E}^1(X, \omega, \psi_k)$ and combining the monotonicity of $E_{\psi_k}(\cdot)$, the upper semicontinuity of $E(\cdot)$ (Proposition 3.14) and the contraction property of Proposition 2.10 we obtain

$$\begin{aligned} E_\psi(u) &= \lim_{k \rightarrow \infty} E_\psi(v_k) = AV_\psi - \lim_{k \rightarrow \infty} d(v_k, \psi + A) \leq \liminf_{k \rightarrow \infty} (AV_{\psi_k} - d(w_k, \psi_k + A)) = \\ &= \liminf_{k \rightarrow \infty} E_{\psi_k}(w_k) \leq \limsup_{k \rightarrow \infty} E_{\psi_k}(w_k) \leq E_\psi(u), \end{aligned}$$

i.e. $E_{\psi_k}(w_k) \rightarrow E_\psi(u)$ as $k \rightarrow \infty$. As a easy consequence we also get $d(w_k, u_k) = E_{\psi_k}(w_k) - E_{\psi_k}(u_k) \rightarrow 0$, thus it is sufficient to prove that

$$\limsup_{k \rightarrow \infty} d_{\mathcal{A}}(u, w_k) = 0.$$

Similarly to the previous case, fix $\epsilon > 0$ and let $\phi_\epsilon = P_\omega[\psi](\varphi_\epsilon)$ for $\varphi_\epsilon \in \mathcal{H}_\omega$ such that $d(u, \phi_\epsilon) \leq \epsilon$. Again Proposition 2.10 and Proposition 2.11 yield

$$\begin{aligned} \limsup_{k \rightarrow \infty} d_{\mathcal{A}}(u, w_k) &\leq \epsilon + \limsup_{k \rightarrow \infty} (d_{\mathcal{A}}(\phi_\epsilon, P_\omega[\psi_k](\phi_\epsilon)) + d(P_\omega[\psi_k](\phi_\epsilon), w_k)) \leq \\ &\leq \epsilon + \limsup_{k \rightarrow \infty} (d_{\mathcal{A}}(\phi_\epsilon, P_\omega[\psi_k](\phi_\epsilon)) + d(\phi_\epsilon, v_k)) \leq 2\epsilon, \end{aligned}$$

which concludes the first part.

Implication (i) \Rightarrow (ii) if $u \neq P_{\psi_{\min}}$ while (i) implies the dichotomy if $u = P_{\psi_{\min}}$.

If $u \neq P_{\psi_{\min}}$, Proposition 6.1 implies that $u_k \rightarrow u$ weakly and in particular that $|\sup_X u_k| \leq A$. Thus it remains to prove that $E_{\psi_k}(u_k) \rightarrow E_\psi(u)$.

If $u = P_{\psi_{\min}}$ then again by Proposition 6.1 it remains to show that $E_{\psi_k}(u_k) \rightarrow 0$ assuming $u_{k_h} \rightarrow P_{\psi_{\min}}$ strongly and weakly. Note that we also have $|\sup_X u_k| \leq A$ for a uniform constant $A \in \mathbb{R}$ by definition of weak convergence to $P_{\psi_{\min}}$.

So, since by an easy contradiction argument it is enough to prove that any subsequence of $\{u_k\}_{k \in \mathbb{N}}$ admits a further subsequence such that the convergence of the energies holds, without loss of generality we may

assume that $u_k \rightarrow u \in \mathcal{E}^1(X, \omega, \psi)$ weakly even in the case $V_\psi = 0$ (i.e. when, with abuse of notation, $u = P_{\psi_{\min}}$).

Therefore we want to show the existence of a further subsequence $\{u_{k_h}\}_{h \in \mathbb{N}}$ such that $E_{\psi_{k_h}}(u_{k_h}) \rightarrow E_\psi(u)$ (note that if $V_\psi = 0$ then $E_\psi(u) = 0$). It easily follows that

$$|E_{\psi_k}(u_k) - E_\psi(u)| \leq |d(\psi_k + A, u_k) - d(\psi + A, u)| + A|V_{\psi_k} - V_\psi| \leq d_{\mathcal{A}}(u, u_k) + d(\psi_k + A, \psi + A) + A|V_{\psi_k} - V_\psi|,$$

and this leads to $\lim_{k \rightarrow \infty} E_{\psi_k}(u_k) = E_\psi(u)$ by Proposition 2.11 since $\psi_k + A = P_\omega[\psi_k](A)$ and $\psi + A = P_\omega[\psi](A)$. Hence $E_{\psi_k}(u_k) \rightarrow E_\psi(u)$ as requested. \square

Note that in Theorem 6.2 the case (b) may happen (Remark 3.16) but obviously one can consider

$$X_{\mathcal{A}, norm} = \bigsqcup_{\psi \in \overline{\mathcal{A}}} \mathcal{E}_{norm}^1(X, \omega, \psi)$$

to exclude such pathology.

The strong convergence also implies the convergence in ψ' -capacity for any $\psi' \in \mathcal{M}^+$ as our next result shows.

Theorem 6.3. *Let $\psi_k, \psi \in \mathcal{A}$, and let $u_k \in \mathcal{E}^1(X, \omega, \psi_k)$ strongly converging to $u \in \mathcal{E}^1(X, \omega, \psi)$. Assuming also that $V_\psi > 0$. Then there exists a subsequence $\{u_{k_j}\}_{j \in \mathbb{N}}$ such that the sequences $w_j := (\sup\{u_{k_s} : s \geq j\})^*$, $v_j := P_\omega(u_{k_j}, u_{k_{j+1}}, \dots)$ belong to $X_{\mathcal{A}}$, satisfy $v_j \leq u_{k_j} \leq w_j$ and converge strongly and monotonically to u . In particular $u_k \rightarrow u$ in ψ' -capacity for any $\psi' \in \mathcal{M}^+$ and $MA_\omega(u_k^j, \psi_k^{n-j}) \rightarrow MA_\omega(u^k, \psi^{n-j})$ weakly for any $j \in \{0, \dots, n\}$.*

Proof. We first observe that by Theorem 6.2 $u_k \rightarrow u$ weakly and $E_{\psi_k}(u_k) \rightarrow E_\psi(u)$. In particular $\sup_X u_k$ is uniformly bounded and the sequence of ω -psh $w_k := (\sup\{u_j : j \geq k\})^*$ decreases to u .

Up to considering a subsequence we may assume either $\psi_k \searrow \psi$ or $\psi_k \nearrow \psi$ almost everywhere. We treat the two cases separately.

Assume first that $\psi_k \searrow \psi$. Since clearly $w_k \in \mathcal{E}^1(X, \omega, \psi_k)$ and $E_{\psi_k}(w_k) \geq E_{\psi_k}(u_k)$, Theorem 6.2 and Proposition 3.14 yields

$$E_\psi(u) = \lim_{k \rightarrow \infty} E_{\psi_k}(u_k) \leq \limsup_{k \rightarrow \infty} E_{\psi_k}(w_k) \leq E_\psi(u),$$

i.e. $w_k \rightarrow u$ strongly. Thus up to considering a further subsequence we can suppose that $d(u_k, w_k) \leq 1/2^k$ for any $k \in \mathbb{N}$.

Next similarly as during the proof of Proposition 5.7 we define $v_{j,l} := P_\omega(u_j, \dots, u_{j+l})$ for any $j, l \in \mathbb{N}$, observing that $v_{j,l} \in \mathcal{E}^1(X, \omega, \psi_{j+l})$. Thus the function $v_{j,l}^u := P_\omega(u, v_{j,l}) \in \mathcal{E}^1(X, \omega, \psi)$ satisfies

$$\begin{aligned} d(u, v_{j,l}^u) &\leq \int_X (u - v_{j,l}^u) MA_\omega(v_{j,l}^u) \leq \int_{\{v_{j,l}^u = v_{j,l}\}} (u - v_{j,l}) MA_\omega(v_{j,l}) \leq \\ &\leq \sum_{s=j}^{j+l} \int_X (w_s - u_s) MA_\omega(u_s) \leq (n+1) \sum_{s=j}^{j+l} d(w_s, u_s) \leq \frac{(n+1)}{2^{j-1}}, \end{aligned} \quad (22)$$

where we combined Proposition 2.4 and Lemma 3.7 in [13]. Therefore by Proposition 3.15 $v_{j,l}^u$ converges decreasingly and strongly in $\mathcal{E}^1(X, \omega, \psi)$ to a function ϕ_j which satisfies $\phi_j \leq u$.

Similarly $\int_{\{P_\omega(u, v_{j,l}^u) = u\}} (v_{j,l}^u - u) MA_\omega(u) \leq \int_X |v_{j,l}^u - u| MA_\omega(u) < \infty$ by Corollary 3.5, which implies that $v_{j,l}$ converges decreasingly to $v_j \in \mathcal{E}^1(X, \omega, \psi)$ such that $u \geq v_j \geq \phi_j$ since $v_j \leq u_s$ for any $s \geq j$ and $v_{j,l} \geq v_{j,l}^u$. Hence from (22) we obtain

$$d(u, v_j) \leq d(u, \phi_j) = \lim_{l \rightarrow \infty} d(u, v_{j,l}^u) \leq \frac{(n+1)}{2^{j-1}},$$

i.e. v_j converges increasingly and strongly to u as $j \rightarrow \infty$.

Next assume $\psi_k \nearrow \psi$ almost everywhere. In this case $w_k \in \mathcal{E}^1(X, \omega, \psi)$ for any $k \in \mathbb{N}$, and clearly w_k converges strongly and decreasingly to u . On the other hand, letting $w_{k,k} := P_\omega[\psi_k](w_k)$ we observe that $w_{k,k} \rightarrow u$ weakly since $w_k \geq w_{k,k} \geq u_k$ and

$$E_\psi(u) = \lim_{k \rightarrow \infty} E_{\psi_k}(u_k) \leq \limsup_{k \rightarrow \infty} E_{\psi_k}(w_{k,k}) \leq E_\psi(u)$$

by Theorem 6.2 and Proposition 3.14, i.e. $w_{k,k} \rightarrow u$ strongly again by Theorem 6.2. Thus, similarly to the previous case, we may assume that $d(u_k, w_{k,k}) \leq 1/2^k$ up to considering a further subsequence. Therefore setting $v_{j,l} := P_\omega(u_j, \dots, u_{j+l}) \in \mathcal{E}^1(X, \omega, \psi_j)$, $u^j := P_\omega[\psi_j](u)$ and $v_{j,l}^{u^j} := P_\omega(v_{j,l}, u^j)$ we obtain

$$d(u^j, v_{j,l}^{u^j}) \leq \int_X (u^j - v_{j,l}^{u^j}) MA_\omega(v_{j,l}^{u^j}) \leq \sum_{s=j}^{j+l} \int_X (w_{s,s} - u_s) MA_\omega(u_s) \leq \frac{(n+1)}{2^{j-1}} \quad (23)$$

proceeding similarly as before. This implies that $v_{j,l}^{u^j}$ and $v_{j,l}$ converge decreasingly and strongly respectively to functions $\phi_j, v_j \in \mathcal{E}^1(X, \omega, \psi_j)$ as $l \rightarrow +\infty$ which satisfy $\phi_j \leq v_j \leq u^j$. Therefore combining (23), Proposition 2.11 and the triangle inequality we get

$$\limsup_{j \rightarrow \infty} d_{\mathcal{A}}(u, v_j) \leq \limsup_{j \rightarrow \infty} \left(d_{\mathcal{A}}(u, u^j) + d(u^j, \phi_j) \right) \leq \limsup_{j \rightarrow \infty} \left(d_{\mathcal{A}}(u, u^j) + \frac{(n+1)}{2^{j-1}} \right) = 0.$$

Hence v_j converges strongly and increasingly to u , so $v_j \nearrow u$ almost everywhere (Proposition 6.1) and the first part of the proof is concluded.

The convergence in ψ' -capacity and the weak convergence of the mixed Monge-Ampère measures follow exactly as seen during the proof of Proposition 5.7. \square

We observe that the assumption $u \neq P_{\psi_{\min}}$ if $V_{\psi_{\min}} = 0$ in Theorem 6.3 is obviously necessary as the counterexample of Remark 3.16 shows. On the other hand if $d_{\mathcal{A}}(u_k, P_{\psi_{\min}}) \rightarrow 0$ then trivially $MA_\omega(u_k^j, \psi_k^{n-j}) \rightarrow 0$ weakly as $k \rightarrow \infty$ for any $j \in \{0, \dots, n\}$ as a consequence of $V_{\psi_k} \searrow 0$.

6.2 Proof of Theorem B

Definition 6.4. We define $Y_{\mathcal{A}}$ as

$$Y_{\mathcal{A}} := \bigsqcup_{\psi \in \overline{\mathcal{A}}} \mathcal{M}^1(X, \omega, \psi),$$

and we endow it with its natural strong topology given as the coarsest refinement of the weak topology such that E^* becomes continuous, i.e. $V_{\psi_k} \mu_k$ converges strongly to $V_{\psi} \mu$ if and only if $V_{\psi_k} \mu_k \rightarrow V_{\psi} \mu$ weakly and $E_{\psi_k}^*(\mu_k) \rightarrow E_{\psi}^*(\mu)$ as $k \rightarrow \infty$.

Observe that $Y_{\mathcal{A}} \subset \{\text{non-pluripolar measures of total mass belonging to } [V_{\psi_{\min}}, V_{\psi_{\max}}]\}$ where clearly $\psi_{\max} := \sup \mathcal{A}$. As stated in the Introduction, the denomination is coherent with [3] since if $\psi = 0 \in \overline{\mathcal{A}}$ then the induced topology on $\mathcal{M}^1(X, \omega)$ coincides with the strong topology as defined in [3].

We also recall that

$$X_{\mathcal{A}, \text{norm}} := \bigsqcup_{\psi \in \overline{\mathcal{A}}} \mathcal{E}_{\text{norm}}^1(X, \omega, \psi)$$

where $\mathcal{E}_{\text{norm}}^1(X, \omega, \psi) := \{u \in \mathcal{E}^1(X, \omega, \psi) \text{ such that } \sup_X u = 0\}$ (if $V_{\psi_{\min}} = 0$ then we clearly assume $P_{\psi_{\min}} \in X_{\mathcal{A}, \text{norm}}$).

Theorem B. The Monge-Ampère map

$$MA_\omega : (X_{\mathcal{A}, \text{norm}}, d_{\mathcal{A}}) \rightarrow (Y_{\mathcal{A}}, \text{strong})$$

is a homeomorphism.

Proof. The map is a bijection as a consequence of Lemma 3.12 and Proposition 5.5 defining clearly $MA_\omega(P_{\psi_{\min}}) := 0$, i.e. to be the null measure.

Step 1: Continuity. Assume first that $V_{\psi_{\min}} = 0$ and that $d_{\mathcal{A}}(u_k, P_{\psi_{\min}}) \rightarrow 0$ as $k \rightarrow \infty$. Then easily $MA_\omega(u_k) \rightarrow 0$ weakly. Moreover, assuming $u_k \neq P_{\psi_{\min}}$ for any k , it follows from Proposition 2.4 that

$$\begin{aligned} E_{\psi_k}^*(MA_\omega(u_k)/V_{\psi_k}) &= E_{\psi_k}(u_k) + \int_X (\psi_k - u_k) MA_\omega(u_k) \leq \\ &\leq \frac{n}{n+1} \int_X (\psi_k - u_k) MA_\omega(u_k) \leq -n E_{\psi_k}(u_k) \rightarrow 0 \end{aligned}$$

as $k \rightarrow \infty$ where the convergence is given by Theorem 6.2. Hence $MA_\omega(u_k) \rightarrow 0$ strongly in $Y_{\mathcal{A}}$.

We can now assume that $u \neq P_{\psi_{\min}}$.

Theorem 6.3 immediately gives the weak convergence of $MA_\omega(u_k)$ to $MA_\omega(u)$. Fix $\varphi_j \in \mathcal{H}_\omega$ be a decreasing sequence converging to u such that $d(u, P_\omega[\psi](\varphi_j)) \leq 1/j$ for any $j \in \mathbb{N}$ ([6]) and set $v_{k,j} := P_\omega[\psi_k](\varphi_j)$ and $v_j := P_\omega[\psi](\varphi_j)$. Observe also that as a consequence of Proposition 2.11 and Theorem 6.2, for any $j \in \mathbb{N}$ there exists $k_j \gg 0$ big enough such that $d(\psi_k, v_{k,j}) \leq d_{\mathcal{A}}(\psi_k, \psi) + d(\psi, v_j) + d_{\mathcal{A}}(v_j, v_{k,j}) \leq d(\psi, v_j) + 1 \leq C$ for any $k \geq k_j$, where C is a uniform constant independent on $j \in \mathbb{N}$. Therefore combining again Theorem 6.2 with Lemma 4.6 and Proposition 3.7 we obtain

$$\begin{aligned} & \limsup_{k \rightarrow \infty} \left| E_{\psi_k}^* (MA_\omega(u_k)/V_{\psi_k}) - E_{\psi_k}^* (MA_\omega(v_{k,j})/V_{\psi_k}) \right| \leq \\ & \leq \limsup_{k \rightarrow \infty} \left(|E_{\psi_k}(u_k) - E_{\psi_k}(v_{k,j})| + \left| \int_X (\psi_k - u_k)(MA_\omega(u_k) - MA_\omega(v_{k,j})) \right| + \left| \int_X (v_{k,j} - u_k) MA_\omega(v_{k,j}) \right| \right) \leq \\ & \leq |E_\psi(u) - E_\psi(v_j)| + \limsup_{k \rightarrow \infty} CI_{\psi_k}(u_k, v_{k,j})^{1/2} + \int_X (v_j - u) MA_\omega(v_j) \quad (24) \end{aligned}$$

since clearly we may assume that either $\psi_k \searrow \psi$ or $\psi_k \nearrow \psi$ almost everywhere, up to considering a subsequence. On the other hand, if $k \geq k_j$, Proposition 3.4 implies $I_{\psi_k}(u_k, v_{k,j}) \leq 2f_{\tilde{C}}(d(u_k, v_{k,j}))$ where \tilde{C} is a uniform constant independent of j, k and $f_{\tilde{C}} : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ is a continuous increasing function such that $f_{\tilde{C}}(0) = 0$. Hence continuing the estimates in (24) we get

$$(24) \leq |E_\psi(u) - E_\psi(v_j)| + 2Cf_{\tilde{C}}(d(u, v_j)) + d(v_j, u) \quad (25)$$

using also Propositions 2.4 and 2.11. Letting $j \rightarrow \infty$ in (25), it follows that

$$\limsup_{j \rightarrow \infty} \limsup_{k \rightarrow \infty} \left| E_{\psi_k}^* (MA_\omega(u_k)/V_{\psi_k}) - E_{\psi_k}^* (MA_\omega(v_{k,j})/V_{\psi_k}) \right| = 0$$

since $v_j \searrow u$. Furthermore it is easy to check that $E_{\psi_k}^* (MA_\omega(v_{k,j})/V_{\psi_k}) \rightarrow E_\psi^* (MA_\omega(v_j)/V_\psi)$ as $k \rightarrow \infty$ for j fixed by Lemma 4.6 and Proposition 2.11. Therefore the convergence

$$E_\psi^* (MA_\omega(v_j)/V_\psi) \rightarrow E_\psi^* (MA_\omega(u)/V_\psi) \quad (26)$$

as $j \rightarrow \infty$ given by Theorem A concludes this step.

Step 2: Continuity of the inverse. Assume $u_k \in \mathcal{E}_{norm}^1(X, \omega, \psi_k), u \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ such that $MA_\omega(u_k) \rightarrow MA_\omega(u)$ strongly. Note that when $\psi = \psi_{\min}$ and $V_{\psi_{\min}} = 0$ the assumption does not depend on the function u chosen. Clearly this implies $V_{\psi_k} \rightarrow V_\psi$ which leads to $\psi_k \rightarrow \psi$ as $k \rightarrow \infty$ by Lemma 3.12 since $\mathcal{A} \subset M^+$ is totally ordered. Hence, up to considering a subsequence, we may assume that $\psi_k \rightarrow \psi$ monotonically almost everywhere. We keep the same notations of the previous step for $v_{k,j}, v_j$. We may also suppose that $V_{\psi_k} > 0$ for any $k \in \mathbb{N}$ big enough otherwise it would be trivial.

The strategy is to proceed similarly as during the proof of Theorem A, i.e. we want first to prove that $I_{\psi_k}(u_k, v_{k,j}) \rightarrow 0$ as $k, j \rightarrow \infty$ in this order. Then we want to use this to prove that the unique weak accumulation point of $\{u_k\}_{k \in \mathbb{N}}$ is u . Finally we will deduce also the convergence of the ψ_k -relative energies to conclude that $u_k \rightarrow u$ strongly thanks to Theorem 6.2.

By Lemma 3.1

$$\begin{aligned} (n+1)^{-1} I_{\psi_k}(u_k, v_{k,j}) & \leq E_{\psi_k}(u_k) - E_{\psi_k}(v_{k,j}) + \int_X (v_{k,j} - u_k) MA_\omega(u_k) = \\ & = E_{\psi_k}^* (MA_\omega(u_k)/V_{\psi_k}) - E_{\psi_k}^* (MA_\omega(v_{k,j})/V_{\psi_k}) + \int_X (v_{k,j} - \psi_k)(MA_\omega(u_k) - MA_\omega(v_{k,j})) \quad (27) \end{aligned}$$

for any j, k . Moreover by Step 1 and Proposition 2.11 we know that $E_{\psi_k}^* (MA_\omega(v_{k,j})/V_{\psi_k})$ converges, as $k \rightarrow +\infty$, respectively to 0 if $V_\psi = 0$ and to $E_\psi^* (MA_\omega(v_j)/V_\psi)$ if $V_\psi > 0$. Next by Lemma 4.6

$$\int_X (v_{k,j} - \psi_k) MA_\omega(v_{k,j}) \rightarrow \int_X (v_j - \psi) MA_\omega(v_j)$$

letting $k \rightarrow \infty$. So if $V_\psi = 0$ then from $\lim_{k \rightarrow \infty} \sup_X (v_{k,j} - \psi_k) = \sup_X (v_j - \psi) = \sup_X v_j$ we easily get $\limsup_{k \rightarrow \infty} I_{\psi_k}(u_k, v_{k,j}) = 0$. Thus we may assume $V_\psi > 0$ and it remains to estimate $\int_X (v_{k,j} - \psi_k) MA_\omega(u_k)$ from above.

We set $f_{k,j} := v_{k,j} - \psi_k$ and analogously to the proof of Theorem A we construct a sequence of smooth

functions $f_j^s := v_j^s - \psi^s$ converging in capacity to $f_j := v_j - \psi$ and satisfying $\|f_j^s\|_{L^\infty} \leq 2\|f_j\|_{L^\infty} \leq 2\|\varphi_j\|_{L^\infty}$. Here v_j^s, ψ^s are sequences of ω -psh functions decreasing respectively to v_j, ψ . Then we write

$$\int_X f_{k,j} MA_\omega(u_k) = \int_X (f_{k,j} - f_j^s) MA_\omega(u_k) + \int_X f_j^s MA_\omega(u_k) \quad (28)$$

and we observe that $\limsup_{s \rightarrow \infty} \limsup_{k \rightarrow \infty} \int_X f_j^s MA_\omega(u_k) = \int_X f_j MA_\omega(u)$ since $MA_\omega(u_k) \rightarrow MA_\omega(u)$ weakly, $f_j^s \in C^\infty$, f_j^s converges to f_j in capacity and $\|f_j^s\|_{L^\infty} \leq 2\|f_j\|_{L^\infty}$. While we claim that the first term on the right-hand side of (28) goes to 0 letting $k, s \rightarrow \infty$ in this order. Indeed for any $\delta > 0$

$$\int_X (f_{k,j} - f_j) MA_\omega(u_k) \leq \delta V_{\psi_k} + 2\|\varphi_j\|_{L^\infty} \int_{\{|f_{k,j} - f_j| > \delta\}} MA_\omega(u_k) \leq \delta V_{\psi_k} + 2\|\varphi_j\|_{L^\infty} \int_{\{|h_{k,j} - h_j| > \delta\}} MA_\omega(u_k) \quad (29)$$

where we set $h_{k,j} := v_{k,j}, h_j := v_j$ if $\psi_k \searrow \psi$ and $h_{k,j} := \psi_k, h_j := \psi$ if instead $\psi_k \nearrow \psi$ almost everywhere. Moreover since $\{|h_{k,j} - h_j| > \delta\} \subset \{|h_{l,j} - h_j| > \delta\}$ for any $l \leq k$, from (29) we obtain

$$\begin{aligned} \limsup_{k \rightarrow \infty} \int_X (f_{k,j} - f_j) MA_\omega(u_k) &\leq \delta V_\psi + \limsup_{l \rightarrow \infty} \limsup_{k \rightarrow \infty} 2\|\varphi_j\|_{L^\infty} \int_{\{|h_{l,j} - h_j| \geq \delta\}} MA_\omega(u_k) \leq \\ &\leq \delta V_\psi + \limsup_{l \rightarrow \infty} 2\|\varphi_j\|_{L^\infty} \int_{\{|h_{l,j} - h_j| \geq \delta\}} MA_\omega(u) = \delta V_\psi \end{aligned}$$

where we also used that $\{|h_{l,j} - h_j| \geq \delta\}$ is a closed set in the plurifine topology since it is equal to $\{v_{l,j} - v_j \geq \delta\}$ if $\psi_l \searrow \psi$ and to $\{\psi - \psi_l \geq \delta\}$ if $\psi_l \nearrow \psi$ almost everywhere. Hence $\limsup_{k \rightarrow \infty} \int_X (f_{k,j} - f_j) MA_\omega(u_k) \leq 0$. Similarly we also get $\limsup_{s \rightarrow \infty} \limsup_{k \rightarrow \infty} \int_X (f_j - f_j^s) MA_\omega(u_k) \leq 0$. (see also the proof of Theorem A).

Summarizing from (27), we obtain

$$\begin{aligned} \limsup_{k \rightarrow \infty} (n+1)^{-1} I_{\psi_k}(u_k, v_{k,j}) &\leq E_{\psi}^*(MA_\omega(u)/V_\psi) - E_{\psi}^*(MA_\omega(v_j)/V_\psi) + \\ &+ \int_X (v_j - \psi) MA_\omega(u) - \int_X (v_j - \psi) MA_\omega(v_j) =: F_j, \quad (30) \end{aligned}$$

and $F_j \rightarrow 0$ as $j \rightarrow \infty$ by Step 1 and Proposition 3.7 since $\mathcal{E}^1(X, \omega, \psi) \ni v_j \searrow u \in \mathcal{E}_{norm}^1(X, \omega, \psi)$, hence strongly.

Next by Lemma 3.1 $u_k \in X_{A,C}$ for $C \gg 1$ since $E^*(MA_\omega(u_k)/V_{\psi_k}) = J_{u_k}^\psi(\psi)$ and $\sup_X u_k = 0$, thus up to considering a further subsequence $u_k \rightarrow w \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ weakly where $d(w, \psi) \leq C$. Indeed if $V_\psi > 0$ this follows from Proposition 3.15 while it is trivial if $V_\psi = 0$. In particular by Lemma 4.6

$$\int_X (\psi_k - u_k) MA_\omega(v_{k,j}) \rightarrow \int_X (\psi - w) MA_\omega(v_j) \quad (31)$$

$$\int_X (v_{k,j} - u_k) MA_\omega(v_{k,j}) \rightarrow \int_X (v_j - w) MA_\omega(v_j) \quad (32)$$

as $j \rightarrow \infty$. Therefore if $V_\psi = 0$ then combining $I_{\psi_k}(u_k, v_{k,j}) \rightarrow 0$ as $k \rightarrow \infty$ with (32) and Lemma 3.1, we obtain

$$\limsup_{k \rightarrow \infty} \left(-E_{\psi_k}(u_k) + E_{\psi_k}(v_{k,j}) \right) \leq \limsup_{k \rightarrow \infty} \left(\frac{n}{n+1} I_{\psi_k}(u_k, v_{k,j}) + \left| \int_X (v_{k,j} - u_k) MA_\omega(v_{k,j}) \right| \right) = 0.$$

This implies that $d(\psi_k, u_k) = -E_{\psi_k}(u_k) \rightarrow 0$ as $k \rightarrow \infty$, i.e. that $d_{\mathcal{A}}(P_{\psi_{\min}}, u_k) \rightarrow 0$ using Theorem 6.2. Thus we may assume from now until the end of the proof that $V_\psi > 0$.

By (31) and Proposition 3.14 it follows that

$$\begin{aligned} \limsup_{k \rightarrow \infty} \left(E_{\psi_k}^*(MA_\omega(u_k)/V_{\psi_k}) + \int_X (\psi_k - u_k) (MA_\omega(v_{k,j}) - MA_\omega(u_k)) \right) &= \\ = \limsup_{k \rightarrow \infty} \left(E_{\psi_k}(u_k) + \int_X (\psi_k - u_k) MA_\omega(v_{k,j}) \right) &\leq E_\psi(w) + \int_X (\psi - w) MA_\omega(v_j). \quad (33) \end{aligned}$$

On the other hand by Proposition 3.7 and (30),

$$\limsup_{k \rightarrow \infty} \left| \int_X (\psi_k - u_k) (MA_\omega(v_{k,j}) - MA_\omega(u_k)) \right| \leq CF_j^{1/2}. \quad (34)$$

In conclusion by the triangle inequality combining (33) and (34) we get

$$\begin{aligned} E_\psi(u) + \int_X (\psi - u)MA_\omega(u) &= \lim_{k \rightarrow \infty} E^*(MA_\omega(u_k)/V_{\psi_k}) \leq \\ &\leq \limsup_{j \rightarrow \infty} \left(E_\psi(w) + \int_X (\psi - w)MA_\omega(v_j) + CF_j^{1/2} \right) = E_\omega(w) + \int_X (\psi - w)MA_\omega(u) \end{aligned}$$

since $F_j \rightarrow 0$, i.e. $w \in \mathcal{E}_{norm}^1(X, \omega, \psi)$ is a maximizer of $F_{MA_\omega(u)/V_{\psi, \psi}}$. Hence $w = u$ (Proposition 5.5), i.e. $u_k \rightarrow u$ weakly. Furthermore, similarly to the case $V_\psi = 0$, Lemma 3.1 and (32) imply

$$\begin{aligned} E_\psi(v_j) - \liminf_{k \rightarrow \infty} E_{\psi_k}(u_k) &= \limsup_{k \rightarrow \infty} \left(-E_{\psi_k}(u_k) + E_{\psi_k}(v_{k,j}) \right) \leq \\ &\leq \limsup_{k \rightarrow \infty} \left(\frac{n}{n+1} I_{\psi_k}(u_k, v_{k,j}) + \left| \int_X (u_k - v_{j,k})MA_\omega(v_{k,j}) \right| \right) \leq \frac{n}{n+1} F_j + \left| \int_X (u - v_j)MA_\omega(v_j) \right| \end{aligned}$$

Finally letting $j \rightarrow \infty$, since $v_j \rightarrow u$ strongly, we obtain $\liminf_{k \rightarrow \infty} E_{\psi_k}(u_k) \geq \lim_{j \rightarrow \infty} E_\psi(v_j) = E_\psi(u)$. Hence $E_{\psi_k}(u_k) \rightarrow E_\psi(u)$ by Proposition 3.14 which implies $d_{\mathcal{A}}(u_k, u) \rightarrow 0$ by Theorem 6.2 and concludes the proof. \square

7 Stability of Complex Monge-Ampère equations.

As stated in the Introduction, we want to use the homeomorphism of Theorem B to deduce the strong stability of solutions of complex Monge-Ampère equations with prescribed singularities when the measures have uniformly bounded L^p density for $p > 1$.

Theorem C. *Let $\mathcal{A} := \{\psi_k\}_{k \in \mathbb{N}} \subset \mathcal{M}^+$ be totally ordered, and let $\{f_k\}_{k \in \mathbb{N}} \subset L^1$ a sequence of non-negative functions such that $f_k \rightarrow f \in L^1 \setminus \{0\}$ and such that $\int_X f_k \omega^n = V_{\psi_k}$ for any $k \in \mathbb{N}$. Assume also that there exists $p > 1$ such that $\|f_k\|_{L^p}, \|f\|_{L^p}$ are uniformly bounded. Then $\psi_k \rightarrow \psi \in \overline{\mathcal{A}} \subset \mathcal{M}^+$, and the sequence of solutions of*

$$\begin{cases} MA_\omega(u_k) = f_k \omega^n \\ u_k \in \mathcal{E}_{norm}^1(X, \omega, \psi_k) \end{cases} \quad (35)$$

converges strongly to $u \in X_{\mathcal{A}}$ which is the unique solution of

$$\begin{cases} MA_\omega(u) = f \omega^n \\ u \in \mathcal{E}_{norm}^1(X, \omega, \psi). \end{cases} \quad (36)$$

In particular $u_k \rightarrow u$ in capacity.

Proof. We first observe that the existence of the unique solutions of (35) follows by Theorem A in [14]. Moreover letting u any weak accumulation point for $\{u_k\}_{k \in \mathbb{N}}$ (there exists at least one by compactness), Lemma 2.8 in [14] yields $MA_\omega(u) \geq f \omega^n$ and by the convergence of f_k to f we also obtain $\int_X f \omega^n = \lim_{k \rightarrow \infty} \int_X f_k \omega^n = \lim_{k \rightarrow \infty} V_{\psi_k}$. Moreover since $u_k \leq \psi_k$ for any $k \in \mathbb{N}$, by [24] we obtain $\int_X MA_\omega(u) \leq \lim_{k \rightarrow \infty} \int_X V_{\psi_k}$. Hence $MA_\omega(u) = f \omega^n$ which in particular means that there is a unique weak accumulation point for $\{u_k\}_{k \in \mathbb{N}}$ and that $\psi_k \rightarrow \psi$ as $k \rightarrow \infty$ since $V_{\psi_k} \rightarrow V_\psi$ (by Lemma 3.12). Then it easily follows combining Fatou's Lemma with Proposition 2.10 and Lemma 2.12 that for any $\varphi \in \mathcal{H}_\omega$

$$\begin{aligned} \liminf_{k \rightarrow \infty} E_{\psi_k}^*(MA_\omega(u_k)/V_{\psi_k}) &\geq \liminf_{k \rightarrow \infty} \left(E_{\psi_k}(P_\omega[\psi_k](\varphi)) + \int_X (\psi_k - P_\omega[\psi_k](\varphi)) f_k \omega^n \right) \geq \\ &\geq E_\psi(P_\omega[\psi](\varphi)) + \int_X (\psi - P_\omega[\psi](\varphi)) f \omega^n \quad (37) \end{aligned}$$

since $(\psi_k - P_\omega[\psi_k](\varphi)) f_k \rightarrow (\psi - P_\omega[\psi](\varphi)) f$ almost everywhere by Lemma 2.14. Thus, for any $v \in \mathcal{E}^1(X, \omega, \psi)$ letting $\varphi_j \in \mathcal{H}_\omega$ be a decreasing sequence converging to v ([6]), from the inequality (37) we get

$$\liminf_{k \rightarrow \infty} E_{\psi_k}^*(MA_\omega(u_k)/V_{\psi_k}) \geq \limsup_{j \rightarrow \infty} \left(E_\psi(P_\omega[\psi](\varphi_j)) + \int_X (\psi - P_\omega[\psi](\varphi_j)) f \omega^n \right) = E_\psi(v) + \int_X (\psi - v) f \omega^n$$

using Proposition 2.4 and the Monotone Convergence Theorem. Hence by definition

$$\liminf_{k \rightarrow \infty} E_{\psi_k}^*(MA_\omega(u_k)/V_{\psi_k}) \geq E_\psi^*(f\omega^n/V_\psi). \quad (38)$$

On the other hand since $\|f_k\|_{L^p}, \|f\|_{L^p}$ are uniformly bounded where $p > 1$ and $u_k \rightarrow u, \psi_k \rightarrow \psi$ in L^q for any $q \in [1, +\infty)$ (see Theorem 1.48 in [19]), we also have

$$\int_X (\psi_k - u_k) f_k \omega^n \rightarrow \int_X (\psi - u) f \omega^n < +\infty,$$

which implies that $\int_X (\psi - u) MA_\omega(u) < +\infty$, i.e. $u \in \mathcal{E}^1(X, \omega, \psi)$ by Proposition 2.4. Moreover by Proposition 3.14 we also get

$$\limsup_{k \rightarrow \infty} E_{\psi_k}^*(MA_\omega(u_k)/V_{\psi_k}) \leq E_\psi^*(MA_\omega(u)/V_\psi),$$

which together with (38) leads to $MA_\omega(u_k) \rightarrow MA_\omega(u)$ strongly in $Y_{\mathcal{A}}$ by definition (observe that $MA_\omega(u_k) = f_k \omega^n \rightarrow MA_\omega(u) = f \omega^n$ weakly). Hence $u_k \rightarrow u$ strongly by Theorem B while the convergence in capacity follows from Theorem 6.3. \square

Remark 7.1. As said in the Introduction, the convergence in capacity of Theorem C was already obtained in Theorem 1.4 in [15]. Indeed under the hypothesis of Theorem C it follows from Lemma 2.12 and Lemma 3.4 in [15] that $d_S(\psi_k, \psi) \rightarrow 0$ where d_S is the pseudometric on $\{[u] : u \in PSH(X, \omega)\}$ introduced in [15] where the class $[u]$ is given by the partial order \preceq .

References

- [1] AUBIN, T. Réduction du cas positif de l'équation de Monge-Ampère sur les variétés Kählériennes compactes à la démonstration d'une inégalité. *Journal of Functional Analysis* 57 (1984), 143–153.
- [2] BERMAN, R., AND BOUCKSOM, S. Growth of balls of holomorphic sections and energy at equilibrium. *Inventiones mathematicae* 181, 2 (2010), 337–394.
- [3] BERMAN, R. J., BOUCKSOM, S., EYSSIDIEUX, P., GUEDJ, V., AND ZERIAHI, A. Kähler–einstein metrics and the Kähler–Ricci flow on log Fano varieties. *Journal für die reine und angewandte Mathematik* 2019, 751 (2019), 27–89.
- [4] BERMAN, R. J., BOUCKSOM, S., GUEDJ, V., AND ZERIAHI, A. A variational approach to complex Monge-Ampère equations. *Publications mathématiques de l'IHES* 117, 1 (2013), 179–245.
- [5] BERMAN, R. J., DARVAS, T., AND LU, C. H. Regularity of weak minimizers of the K-energy and applications to properness and K-stability. *arXiv preprint arXiv:1602.03114* (2016).
- [6] BLOCKI, Z., AND KOŁODZIEJ, S. On regularization of plurisubharmonic functions on manifolds. *Proceedings of the American Mathematical Society* 135, 7 (2007), 2089–2093.
- [7] BOUCKSOM, S., EYSSIDIEUX, P., GUEDJ, V., AND ZERIAHI, A. Monge-Ampère equations in big cohomology classes. *Acta mathematica* 205, 2 (2010), 199–262.
- [8] CEGRELL, U. Pluricomplex energy. *Acta mathematica* 180, 2 (1998), 187–217.
- [9] CHEN, X., AND CHENG, J. On the constant scalar curvature Kähler metrics (I) - A priori estimates. *Journal of the American Mathematical Society* (2021).
- [10] CHEN, X., AND CHENG, J. On the constant scalar curvature Kähler metrics (II) - existence results. *Journal of the American Mathematical Society* (2021).
- [11] CHEN, X., AND CHENG, J. On the constant scalar curvature Kähler metrics (II) - existence results. *Journal of the American Mathematical Society* (2021).
- [12] DARVAS, T. The Mabuchi geometry of finite energy classes. *Advances in Mathematics* 285 (2015), 182–219.
- [13] DARVAS, T., DI NEZZA, E., AND LU, C. H. Monotonicity of nonpluripolar products and complex Monge-Ampère equations with prescribed singularity. *Analysis & PDE* 11, 8 (2018), 2049–2087.
- [14] DARVAS, T., DI NEZZA, E., AND LU, C. H. Log-concavity of volume and complex Monge-Ampère equations with prescribed singularity. *Mathematische Annalen* (2019), 1–38.
- [15] DARVAS, T., NEZZA, E. D., AND LU, C. H. The metric geometry of singularity types. *Journal für die reine und angewandte Mathematik (Crelles Journal)* 2021, 771 (2021), 137–170.
- [16] DARVAS, T., AND RUBINSTEIN, Y. Tian's properness conjectures and Finsler geometry of the space of Kähler metrics. *Journal of the American Mathematical Society* 30, 2 (2017), 347–387.

- [17] DI NEZZA, E., AND TRAPANI, S. Monge-Ampère measures on contact sets. *arXiv preprint arXiv:1912.12720* (2019).
- [18] GUEDJ, V., AND ZERIAHI, A. The weighted Monge–Ampère energy of quasisubharmonic functions. *Journal of Functional Analysis* 250, 2 (2007), 442–482.
- [19] GUEDJ, V., AND ZERIAHI, A. Degenerate complex Monge-Ampère equations. *EMS Tracts in Mathematics* 26 (2017).
- [20] KOŁODZIEJ, S. The complex Monge-Ampère equation. *Acta mathematica* 180, 1 (1998), 69–117.
- [21] MABUCHI, T. K-energy maps integrating Futaki invariants. *Tohoku Mathematical Journal, Second Series* 38, 4 (1986), 575–593.
- [22] ROSS, J., AND WITT NYSTRÖM, D. Analytic test configurations and geodesic rays. *Journal of Symplectic Geometry* 12, 1 (2014), 125–169.
- [23] TRUSIANI, A. L^1 metric geometry of potentials with prescribed singularities on compact Kähler manifolds. *The Journal of Geometric Analysis* 32, 2 (2022), 1–37.
- [24] WITT NYSTRÖM, D. Monotonicity of non-pluripolar Monge-Ampère masses. *Indiana University Mathematics Journal* 68 (2019), 579–591.
- [25] XIA, M. Integration by parts formula for non-pluripolar product. *arXiv preprint arXiv:1907.06359* (2019).
- [26] YAU, S.-T. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, i. *Communications on pure and applied mathematics* 31, 3 (1978), 339–411.