

Well-posedness of SVI solutions to singular-degenerate stochastic porous media equations arising in self-organised criticality

Marius Neuß

Abstract

We consider a class of generalised stochastic porous media equations with multiplicative Lipschitz continuous noise. These equations can be related to physical models exhibiting self-organised criticality. We show that these SPDEs have unique SVI solutions which depend continuously on the initial value. In order to formulate this notion of solution and to prove uniqueness in the case of a slowly growing nonlinearity, the arising energy functional is analysed in detail.

KEYWORDS: singular-degenerate SPDE, stochastic variational inequalities, generalised porous media, self-organised criticality

MSC 2010: 60H15, 76S05

1 Introduction

We consider a class of singular-degenerate generalised stochastic porous media equations

$$\begin{aligned} dX_t &\in \Delta(\phi(X_t)) dt + B(t, X_t) dW_t, \\ X_0 &= x_0, \end{aligned} \tag{1.1}$$

on a bounded, smooth domain $\mathcal{O} \subseteq \mathbb{R}^d$ with zero Dirichlet boundary conditions and $x_0 \in H^{-1}$, where H^{-1} is the dual of $H_0^1(\mathcal{O})$. In the following, W is a cylindrical Wiener process on some separable Hilbert space U , and the diffusion coefficients $B : [0, T] \times H^{-1} \times \Omega \rightarrow L_2(U, H^{-1})$ take values in the space of Hilbert-Schmidt operators $L_2(U, H^{-1})$. The nonlinearity $\phi : \mathbb{R} \rightarrow 2^{\mathbb{R}}$ is the subdifferential of a convex lower-semicontinuous symmetric function $\psi : \mathbb{R} \rightarrow \mathbb{R}$ (sometimes called “potential”), which grows at least linearly and at most quadratically for $|x| \rightarrow \infty$. As paradigmatic examples, we mention the maximal monotone extensions of

$$\phi_1(x) = \operatorname{sgn}(x) (1 - \mathbb{1}_{(-1,1)}(x)) \quad \text{and} \quad \phi_2(x) = x (1 - \mathbb{1}_{(-1,1)}(x)), \tag{1.2}$$

which are encountered in the context of self-organised criticality. Indeed, equation (1.1) with the first nonlinearity in (1.2) is related to a particle model which was first introduced by Bak, Tang and Wiesenfeld in their celebrated works [3] and [4]. We refer to Section 1.2 below for details and references.

The main merits of this article are as follows. First, we give a meaning to (1.1) by defining a suitable notion of solution and proving the existence and uniqueness of such solutions. Second, we extend the applicability of the framework of SVI solutions, which features several properties which are desirable independently of the specific equation presented above. For instance, it applies to stochastic partial differential equations (SPDE) with a very general nonlinear drift term, and solutions for general initial data can be identified by means of the equation and not only in a limiting sense.

We briefly outline the strategy that we are going to apply. First, we rewrite (1.1) into the form

$$dX_t \in -\partial\varphi(X_t) dt + B(t, X_t) dW_t, \tag{1.3}$$

which incorporates the multivalued function ϕ into an energy functional $\varphi : H^{-1} \rightarrow [0, \infty]$. For example, in case of the nonlinearity ϕ_1 in (2.5), we define

$$\varphi(u) = \begin{cases} \|\psi(u)\|_{TV}, & \text{if } u \text{ is a finite Radon measure on } \mathcal{O}, \\ +\infty, & \text{else,} \end{cases} \tag{1.4}$$

where ψ is the anti-derivative of ϕ , i.e. $\partial\psi = \phi$, with $\psi(0) = 0$. For the precise definition of a convex function of a measure, we refer to Section 3 below. We then derive a stochastic variational inequality (SVI) from (1.3) and define a corresponding notion of solution, see Definition 2.4 below. In order to construct such a solution we first show that φ as defined above is lower-semicontinuous, which then allows to show the convergence of an approximating sequence gained by a Yosida approximation of the nonlinearity and the addition of a viscosity term. Furthermore, in the proof of uniqueness, it is crucial to show that φ can be well approximated by its values on L^2 , which we ensure by showing that it coincides with the lower-semicontinuous hull of $\varphi|_{L^2}$ in H^{-1} . To this end, we will construct approximating sequences by an interplay of mollification and shifts, inspired by the construction of [1, Lemma A6.7]. This constitutes one technical focus of this work.

The structure of this article is as follows: In the subsequent sections of the introduction, we will give a brief overview on the mathematical literature concerning the solution theory of generalised stochastic porous media equations, and we will point out how equation (1.1) is motivated by the physics literature. In Section 2 we state the precise assumptions and formulate the first main result of this article, in which the well-posedness of Equation (1.1) is established (see Theorem 2.6 below). We prove the lower-semicontinuity of the abovementioned energy functional φ and the property of φ being the lower-semicontinuous hull of $\varphi|_{L^2}$ in H^{-1} in Section 3, the latter of which is the second main result (see Theorem 3.8 below). In Section 4, the well-posedness result will be proved, following the arguments of [37, Section 2].

1.1 Mathematical Literature

In the recent decades, stochastic porous media equations have been very present in the mathematical literature. For the original case

$$dX_t = \Delta\phi(X_t)dt + B(t, X_t)dW_t, \quad (1.5)$$

where $\phi(r) = r^{[m]} := |r|^{m-1}r$ for $r \in \mathbb{R}$ and $m \geq 1$ ($m = 1$ representing the stochastic heat equation), a concisely summarised well-posedness analysis can be found in [53], which goes back to the work of Krylov and Rozovskii [45] and Pardoux [51]. In [54], the theory is extended to the fast diffusion case $m \in (0, 1)$, and other nonlinear functions ϕ are considered. A setting with a more general monotone and differentiable nonlinearity is considered in [9].

A severe additional difficulty arises when one considers the limit case $m = 0$, in which ϕ becomes multivalued. The first articles treating this type of porous medium equations, [10] and [8], either require ϕ to be surjective or more restrictions on the initial state or the noise. In [41], the $m = 0$ limit of (1.5) can be treated, but one has to restrict to more regular initial data or to the concept of limiting solutions. For general initial conditions, this notion of solution contains no characterisation in terms of the equation, which is often necessary for further work such as stability results (see e.g. [39]).

In [7] and later in [13, 36], the concept of stochastic variational inequalities (SVIs) and a corresponding notion of solution have been used to overcome these issues. We note that in [36], an identification of a functional as a lower-semicontinuous hull was needed in the context of p -Laplace type equations with a C^2 potential, going back to results from [2, 27]. In [37], the existence and uniqueness of SVI solutions was proven for the $m = 0$ limit of (1.5), for which a refinement of previous methods became necessary, because the naive choice for the energy functional does not lead to an energy space with adequate compactness properties. The arising difficulties when setting up the energy functional are similar to the ones mentioned above for φ from (1.4). They have been overcome in [37] by using the specific shape of the nonlinearity, which allows to set the energy functional to

$$\varphi(u) = \begin{cases} \|u\|_{TV}, & \text{if } u \text{ is a finite Radon measure on } \mathcal{O}, \\ +\infty, & \text{else} \end{cases}$$

for $u \in H^{-1}$, which then allows to use structural properties of the TV norm. With more regularity or structural assumptions on the noise and/or the initial state, more regularity for SVI solutions or the existence of strong solutions can be proved, as e.g. in [37, 36, 13, 32]. For the regularisation by noise of quasi-linear SPDE with possibly singular drift terms, we also mention the works [31, 43].

We next mention several different approaches to stochastic porous media equations. The article [14] considers the equation on an unbounded domain, the works [6, 18] use an approach via Kolmogorov

equations. In [12], an operatorial approach to SPDE is introduced which can be applied to generalised stochastic porous media equations with continuous nonlinearities. In [35, 21] and [19], stochastic porous media equations are solved in the sense of kinetic or entropy solutions, respectively. Previous works in those directions are, e.g., [16, 22] and [17, 26, 44]. [38] makes use of a rough path approach leading to pathwise rough kinetic/entropy solutions and including regularity results, with [30, 46] as some of the related preceding works.

Regarding the construction and analysis of the energy functional arising in the context of SVIs, we rely on techniques from [23, 55] on convex functionals of Radon measures. For the deterministic theory on porous medium equations, we refer to [50] and [56]. Regarding results on the long-time behaviour of singular-degenerate SPDE, see e.g. [28, 33] for the existence of random attractors, [40, 20, 48] for ergodicity and [34, 11] for finite-time extinction in the case of purely multiplicative noise.

1.2 Self-organised criticality (SOC)

The model (1.1) can to some extent be associated with processes exhibiting self-organised criticality (SOC). This concept postulates that many randomly driven processes featuring a critical threshold, at which relaxation events are triggered, possess a non-equilibrium statistical invariant state, in which intermittent events can be observed, the size of which is distributed by a power law. SOC has been initially discussed in view of certain cellular automaton models, which are introduced and explained in much detail in [3] and [4], as well as later by [52]. In these models, particles can be interpreted as units of granular material piling up, which coined the notion of “sandpile models”. Other applications, where self-organised critical behaviour has been observed, are the size of landslides [49], earthquakes (the famous Gutenberg-Richter law, see [42]) and stock prices [47].

In [25] and [24], the abovementioned sandpile models are related to a model similar to (1.1), i.e. a stochastic process in a continuous function space where mass of a continuously distributed size is both added and subtracted. In contrast to the assumptions mentioned above, the potential in [25] is only one-sided. As this leads to a process just forced towards $-\infty$, where no avalanches would occur, we consider symmetric potentials instead.

The underlying mechanisms of SOC have been a matter of lively discussion in the literature, see e.g. [57] for a review. The present work is supposed to contribute to this question by noting that SPDEs with singular-degenerate drift and additive noise incorporate several characteristic properties of the original sandpile models, such as deterministic dynamics which are locally switched on at a certain threshold. However, they also differ from them in other perspectives, such as the non-discrete structure. By setting up a theory for those processes, we ultimately hope to gain insight into their long-time statistics, see e.g. [48]. Thereby, we aim to investigate whether SOC extends to the continuous setting and potentially set the stage for new ways of explaining this statistical effect.

1.3 General notation

Unless specified differently, function or measure spaces will be understood to be defined on a smooth, bounded domain $\mathcal{O} \subset \mathbb{R}^d$, $d \in \mathbb{N}$. We write $L^p = L^p(\mathcal{O})$ for the usual Lebesgue spaces with norm $\|\cdot\|_{L^p}$ and scalar product $\langle \cdot, \cdot \rangle_{L^2}$ if $p = 2$. The Lebesgue measure is denoted by dx , and a measure with density $h \in L^1$ with respect to dx is denoted by $h dx$. Furthermore, $H_0^1 = H_0^1(\mathcal{O})$ denotes the Sobolev space of L^2 functions whose first-order weak derivatives exist and are in L^2 , and which have zero trace, with norm $\|u\|_{H_0^1} = \|\nabla u\|_{L^2}$. The full space analogues $L^2(\mathbb{R}^d)$, $H^1(\mathbb{R}^d)$ are defined correspondingly. Furthermore, let H^{-1} denote the topological dual of H_0^1 . We use $-\Delta$ to denote the corresponding Riesz isomorphism, which gives rise to the inner product

$$\langle u, v \rangle_{H^{-1}} = {}_{H^{-1}} \langle u, (-\Delta)^{-1} v \rangle_{H_0^1} \quad \text{for all } u, v \in H^{-1},$$

where the notation ${}_V \langle u, v \rangle_V = {}_V \langle v, u \rangle_V$ denotes evaluating a functional u belonging to the dual space V' of a Banach space V at a vector $v \in V$.

Moreover, we let $\mathcal{C}_0^0 = \mathcal{C}_0^0(\mathcal{O})$ denote the set of all continuous functions on \mathcal{O} vanishing at the boundary, while we write $\mathcal{C}_c^0 = \mathcal{C}_c^0(\mathcal{O})$ for continuous functions with compact support. The same notation applies to spaces \mathcal{C}^k of k times continuously differentiable functions.

For $m \in [0, 1]$ we define the set

$$L^{m+1} \cap H^{-1} := \left\{ v \in L^{m+1} : \exists C \geq 0 \text{ such that } \int v \eta \, dx \leq C \|\eta\|_{H_0^1} \text{ for all } \eta \in C_c^1 \right\}.$$

Note that $L^2 = L^2 \cap H^{-1}$ by the Cauchy-Schwarz and Poincaré inequalities. To each $v \in L^{m+1} \cap H^{-1}$ one can injectively assign a map

$$\mathcal{C}_c^1 \ni \eta \mapsto \int v \eta \, dx. \quad (1.6)$$

By continuity, (1.6) can be injectively extended to a bounded linear functional on H_0^1 , which we call $\iota_m(v)$. The resulting map $\iota_m : L^{m+1} \cap H^{-1} \rightarrow H^{-1}$ is thus injective, which allows to identify v with $\iota_m(v)$.

Let $\mathcal{M} = \mathcal{M}(\mathcal{O})$ be the space of all signed Radon measures on \mathcal{O} with finite total variation, which is isomorphic to the dual space $(\mathcal{C}_0^0)'$ via

$$\mathcal{M} \ni \mu \mapsto \tilde{\mu} \in (\mathcal{C}_0^0)', \quad \tilde{\mu}(f) = \int f \, d\mu. \quad (1.7)$$

This allows us to use $(\mathcal{C}_0^0)'$ and \mathcal{M} , as well as $\tilde{\mu}$ and μ interchangeably. The variation measure of $\mu \in \mathcal{M}$ is denoted by $|\mu| := \mu_+ + \mu_-$ and the total variation of μ is given by

$$\|\mu\|_{TV} = |\mu|(\mathcal{O}).$$

Note that the total variation is also the operator norm if the measure is interpreted as an element of $(\mathcal{C}_0^0)'$ by the Riesz-Markov representation theorem (see e. g. [29, Theorem 1.200]). We define the space of *measures of bounded energy* by

$$\mathcal{M} \cap H^{-1} := \left\{ \mu \in \mathcal{M} : \exists C \geq 0 \text{ such that } \int \eta(x) \, d\mu(x) \leq C \|\eta\|_{H_0^1} \text{ for all } \eta \in C_c^1(\mathcal{O}) \right\}.$$

By a density argument, restricting a measure $\mu \in \mathcal{M} \cap H^{-1}$ to a function on \mathcal{C}_c^1 is an injective operation. Moreover, by continuity $\mu|_{\mathcal{C}_c^1}$ can be injectively extended to a bounded linear functional on H_0^1 , which we call $\iota(\mu)$. The resulting map $\iota : \mathcal{M} \cap H^{-1} \rightarrow H^{-1}$ is thus injective, which allows to identify μ with $\iota(\mu)$.

In general, constants may vary from line to line, but are always positive and finite.

1.4 Acknowledgements

I would like to thank my advisor Benjamin Gess for his support throughout my work on this article. Support by the International Max Planck Research School in Leipzig and by the ‘‘Cusanuswerk – Bischöfliche Studienförderung’’ is gratefully acknowledged.

2 Assumptions and main result

Assumptions 2.1. We require the following assumptions throughout this article.

(A1) W is a cylindrical Id-Wiener process in some separable Hilbert space U defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with normal filtration $(\mathcal{F}_t)_{t \geq 0}$, which means the following: There is a Hilbert-Schmidt embedding J from U to another Hilbert space U_1 , which can be chosen to be bijective (see e. g. [53, Remark 2.5.1]). Defining $Q_1 := JJ^*$, Q_1 is linear, bounded, non-negative definite, symmetric and has finite trace, so that we obtain a classical Q_1 -Wiener process \tilde{W} on U_1 . Moreover, for an operator $\tilde{B} : U \rightarrow H^{-1}$ we have

$$\tilde{B} \in L_2(U, H^{-1}) \Leftrightarrow \tilde{B} \circ J^{-1} \in L_2\left(Q_1^{\frac{1}{2}}(U_1), H^{-1}\right), \quad (2.1)$$

such that if (2.1) is satisfied, we can define

$$\int_0^T \tilde{B} \, dW_t := \int_0^T \tilde{B} \circ J^{-1} \, d\tilde{W}_t.$$

(A2) The diffusion coefficients $B : [0, T] \times H^{-1} \times \Omega \rightarrow L_2(U, H^{-1})$ take values in the space of Hilbert-Schmidt operators, are progressively measurable and satisfy

$$\|B(t, v) - B(t, w)\|_{L_2(U, H^{-1})}^2 \leq C \|v - w\|_{H^{-1}}^2 \quad \text{for all } v, w \in H^{-1}, \quad (2.2)$$

$$\|B(t, v)\|_{L_2(U, L^2)}^2 \leq C(1 + \|v\|_{L^2}^2) \quad \text{for all } v \in L^2, \quad (2.3)$$

$$\|B(t, 0)\|_{L_2(U, H^{-1})}^2 \leq C, \quad (2.4)$$

for some constant $C > 0$ and all $(t, \omega) \in [0, T] \times \Omega$.

(A3) The so-called *potential* $\psi : \mathbb{R} \rightarrow [0, \infty)$ is convex and lower-semicontinuous, and we assume $\psi(0) = 0$, which then implies $0 \in \partial\psi(0)$. For simplicity, we furthermore impose the symmetry assumption $\psi(x) = \psi(-x)$ for all $x \in \mathbb{R}$.

(A4) Define $\phi = \partial\psi : \mathbb{R} \rightarrow 2^{\mathbb{R}}$, the subdifferential of ψ , and assume for all $r \in \mathbb{R}$

$$\inf\{|\eta|^2 : \eta \in \phi(r)\} \leq C(1 + |r|^2). \quad (2.5)$$

In case that

$$\lim_{|x| \rightarrow \infty} \frac{\psi(x)}{|x|} \rightarrow \infty, \quad (2.6)$$

i. e. ψ is superlinear, we require

(A5) There exists $m \in (0, 1]$, such that $\psi(v) \in L^1(\mathcal{O})$ if and only if $v \in L^{m+1}(\mathcal{O})$.

In case that the potential is sublinear, i. e. that there exists a constant $C > 0$ such that

$$\psi(x) \leq C(1 + |x|) \quad \text{for all } x \in \mathbb{R}, \quad (2.7)$$

we require

(A5') There exists $y > 0$ such that $\psi(y) > 0$.

Note that by convexity, Assumption (A5') implies that

$$\psi(x) \geq \frac{\psi(y)}{y} |x| - \psi(y) \quad \text{for all } x \in \mathbb{R}.$$

Next, we define the energy functional for the notion of solution we are going to consider.

Definition 2.2. Let Assumptions 2.1 be satisfied.

(i) In the case of a superlinear potential, i. e. if (2.6) is satisfied, we define for $u \in H^{-1}$ the functional

$$\varphi(u) = \begin{cases} \int \psi(u) \, dx, & \text{if } u \in L^{m+1} \cap H^{-1}, \\ +\infty, & \text{else,} \end{cases} \quad (2.8)$$

where m is the exponent from (A5).

(ii) In the case of a sublinear potential, i. e. if (2.7) is satisfied, we define for $u \in H^{-1}$ the functional

$$\varphi(u) = \begin{cases} \|\psi(u)\|_{TV}, & \text{if } u \in \mathcal{M} \cap H^{-1}, \\ +\infty, & \text{else,} \end{cases} \quad (2.9)$$

where the construction of a nonlinear functional of a measure, which is needed in (2.9), is given in Definition 3.3 below.

Remark 2.3. The choice of the energy functional in Definition 2.2 allows us to reformulate (1.1) as a gradient flow, i. e. to rewrite it in the form

$$\begin{aligned} dX_t &\in -\partial\varphi(X_t)dt + B(t, X_t)dW_t, \\ X_0 &= x_0, \end{aligned} \tag{2.10}$$

where the subdifferential is well-defined due to Proposition 3.7 below. More precisely, let a “classical” solution to (1.1) with $x_0 \in H^{-1}$ be defined as an $(\mathcal{F}_t)_{t \geq 0}$ -adapted process $X \in L^2(\Omega; \mathcal{C}([0, T]; H^{-1}))$ with the following properties: \mathbb{P} -almost surely, for all $t \in [0, T]$ we have $X_t \in L^2$, there is a choice $v_t \in \phi(X_t)$ such that $v_t \in H_0^1$, and

$$X_t = x_0 + \int_0^t \Delta v_r dr + \int_0^t B(r, X_r) dW_r.$$

Furthermore, we impose $\Delta v \in L^2([0, T] \times \Omega; H^{-1})$. Then, one can check that if X is a classical solution in this sense, $(X, \Delta v)$ is a strong solution to

$$\begin{aligned} dX_t &\in -\partial\varphi(X_t)dt + B(t, X_t)dW_t \\ X_0 &= x_0 \end{aligned} \tag{2.11}$$

in the sense of [37, Appendix A].

Now we are in the position to formulate the notion of solution we want to consider.

Definition 2.4 (SVI solution). Given Assumptions 2.1, let $x_0 \in L^2(\Omega, \mathcal{F}_0; H^{-1})$, $T > 0$ and φ be defined as in Definition 2.2. We say that an \mathcal{F}_t -adapted process $X \in L^2(\Omega; \mathcal{C}([0, T]; H^{-1}))$ is an SVI solution to (1.1) if the following conditions are satisfied:

(i) (Regularity)

$$\varphi(X) \in L^1([0, T] \times \Omega).$$

(ii) (Variational inequality) For each \mathcal{F}_t -progressively measurable process $G \in L^2([0, T] \times \Omega; H^{-1})$, and each \mathcal{F}_t -adapted process $Z \in L^2(\Omega; \mathcal{C}([0, T]; H^{-1})) \cap L^2([0, T] \times \Omega; L^2)$ solving the equation

$$Z_t - Z_0 = \int_0^t G_s ds + \int_0^t B(s, Z_s) dW_s \quad \text{for all } t \in [0, T],$$

we have

$$\begin{aligned} \mathbb{E} \|X_t - Z_t\|_{H^{-1}}^2 + 2\mathbb{E} \int_0^t \varphi(X_r) dr &\leq \mathbb{E} \|x_0 - Z_0\|_{H^{-1}}^2 + 2\mathbb{E} \int_0^t \varphi(Z_r) dr \\ &\quad - 2\mathbb{E} \int_0^t \langle G_r, X_r - Z_r \rangle_{H^{-1}} dr \\ &\quad + C \mathbb{E} \int_0^t \|X_r - Z_r\|_{H^{-1}}^2 dr \quad \text{for all } t \in [0, T] \end{aligned} \tag{2.12}$$

for some $C > 0$.

Remark 2.5. It is shown in [37, Remark 2.2] that if (X, η) is a strong solution to (2.10) in H^{-1} , as defined in [37, Appendix A], then X is an SVI solution to (1.1).

The main result of this article is as follows.

Theorem 2.6. *Given Assumptions 2.1, let $x_0 \in L^2(\Omega, \mathcal{F}_0; H^{-1})$ and $T > 0$. Then there is a unique SVI solution X to (1.1). For two SVI solutions X, Y with initial conditions $x_0, y_0 \in L^2(\Omega, \mathcal{F}_0; H^{-1})$, we have*

$$\sup_{t \in [0, T]} \mathbb{E} \|X_t - Y_t\|_{H^{-1}}^2 \leq C \mathbb{E} \|x_0 - y_0\|_{H^{-1}}^2. \tag{2.13}$$

The proof of this theorem will be given in Section 4 below.

3 Properties of the energy functional

The aim of this section is to make Definition 2.2 rigorous by recalling the concept of convex functionals on measures, and to prove certain properties of the energy functional defined in Definition 2.2, which are needed for the proof of the main theorem. We start with some basic concepts concerning convex functions.

Definition 3.1. Let $f : \mathbb{R} \rightarrow [0, \infty]$ be a convex and lower-semicontinuous function with $f(0) = 0$. We then define its convex conjugate $f^* : \mathbb{R} \rightarrow [0, \infty]$ by

$$f^*(x) = \sup_{y \in \mathbb{R}} (xy - f(y)), \quad (3.1)$$

and its recession function $f_\infty : \mathbb{R} \rightarrow [0, \infty]$ by

$$f_\infty(x) = \lim_{t \rightarrow \infty} \frac{f(tx)}{t}. \quad (3.2)$$

Remark 3.2. Note that f_∞ and f^* are convex. If f is symmetric, so are f_∞ and f^* . Moreover, f_∞ is positively homogeneous.

For the notion of solution that we are aiming at, we need the concept of a convex function of a measure, which has been developed in [23].

Definition 3.3. Let ψ satisfy (2.7) as well as Assumptions 2.1 (A3), (A5'). Define the set

$$\mathcal{D}_\psi = \{v \in \mathcal{C}_c^0(\mathcal{O}) : \psi^*(v) \in L^1(\mathcal{O})\}$$

and let $\mu \in \mathcal{M}(\mathcal{O})$. We then define the positive measure $\psi(\mu) \in \mathcal{M}(\mathcal{O})$ by

$$\int_{\mathcal{O}} \eta \psi(\mu) := \mathcal{M}(\mathcal{O}) \langle \psi(\mu), \eta \rangle_{\mathcal{C}_0^0(\mathcal{O})} := \sup \left\{ \int_{\mathcal{O}} v \eta \, d\mu - \int_{\mathcal{O}} \psi^*(v) \eta \, dx : v \in \mathcal{D}_\psi \right\} \quad (3.3)$$

for $\eta \in \mathcal{C}_0^0(\mathcal{O})$, $\eta \geq 0$, and for general $\eta \in \mathcal{C}_0^0(\mathcal{O})$ we set

$$\mathcal{M}(\mathcal{O}) \langle \psi(\mu), \eta \rangle_{\mathcal{C}_0^0(\mathcal{O})} = \mathcal{M}(\mathcal{O}) \langle \psi(\mu), \eta \vee 0 \rangle_{\mathcal{C}_0^0(\mathcal{O})} - \mathcal{M}(\mathcal{O}) \langle \psi(\mu), (-\eta) \vee 0 \rangle_{\mathcal{C}_0^0(\mathcal{O})},$$

according to [23, Theorem 1.1].

Remark 3.4. As argued in [23, Lemma 1.1], one can write for $\mu \in \mathcal{M}(\mathcal{O})$

$$\int_{\mathcal{O}} \psi(\mu) = \|\psi(\mu)\|_{TV} = \sup \left\{ \int_{\mathcal{O}} v \, d\mu - \int_{\mathcal{O}} \psi^*(v) \, dx : v \in \mathcal{D}_\psi \right\}.$$

Remark 3.5. Let $\mu \in \mathcal{M}(\mathcal{O})$ with Lebesgue decomposition $\mu^a + \mu^s$, where μ^a has the density $h \in L^1(\mathcal{O})$ with respect to the Lebesgue measure. Then, by [23, Theorem 1.1], we have

$$\int_{\mathcal{O}} \eta \psi(\mu) = \int_{\mathcal{O}} \eta(x) \psi(h(x)) \, dx + \int_{\mathcal{O}} \eta \psi_\infty(\mu^s), \quad (3.4)$$

where the recession function ψ_∞ is defined as in (3.2). In particular, this formulation shows the useful fact that

$$\psi(\mu) = \psi(\mu^a) + \psi(\mu^s). \quad (3.5)$$

Our next aim is to prove the lower-semicontinuity of the energy functional defined in Definition 2.2 and Definition 3.3. First, we show that the Radon measure $\psi(\mu)$ constructed in Definition 3.3 controls the norm of its original measure μ in the following way.

Lemma 3.6. Let ψ satisfy (2.7) as well as Assumptions 2.1 (A3), (A5'). Let $\mu \in \mathcal{M}(\mathcal{O})$ and let $y > 0$ such that $\psi(y) > 0$ as demanded in Assumption 2.1 (A5'). Then

$$\|\psi(\mu)\|_{TV} \geq \frac{\psi(y)}{y} \|\mu\|_{TV} - \psi(y) |\mathcal{O}|.$$

Proof. For $\mu \in \mathcal{M}(\mathcal{O})$, denote by $\mu = \mu^a + \mu^s$ the Lebesgue decomposition of μ with respect to Lebesgue measure, and let $h = \frac{d\mu^a}{dx}$ be the Radon-Nikodym derivative of μ^a . As $\psi_\infty(\mu^s)$ is singular by [55, Theorem 4.2], we can use the decomposition (3.4) to obtain

$$\|\psi(\mu)\|_{TV} = \int_{\mathcal{O}} \psi(h) dx + \|\psi_\infty(\mu^s)\|_{TV}. \quad (3.6)$$

We now estimate the summands separately. For the absolutely continuous part we obtain using Assumption 2.1 (A5')

$$\int_{\mathcal{O}} \psi(h) dx \geq \frac{\psi(y)}{y} \int_{\mathcal{O}} |h| dx - \psi(y) |\mathcal{O}| = \frac{\psi(y)}{y} \|\mu^a\|_{TV} - \psi(y) |\mathcal{O}|.$$

For the singular part, we note by Lemma A.5 that for $v \in \mathcal{C}_c^0(\mathcal{O})$ being in $\mathcal{D}_{\psi_\infty}$ is equivalent to $-\psi_\infty(1) \leq v \leq \psi_\infty(1)$, and for such v , $\psi_\infty^*(v) \equiv 0$. Thus, we get with Corollary A.4 with $k := \frac{\psi(y)}{y}$

$$\int_{\mathcal{O}} \psi_\infty(\mu^s) = \sup_{v \in \mathcal{D}_{\psi_\infty}} \left(\int_{\mathcal{O}} v d\mu^s - \int_{\mathcal{O}} \psi_\infty^*(v) dx \right) \geq \sup_{\substack{v \in \mathcal{C}_c^0(\mathcal{O}) \\ -k \leq v \leq k}} \int_{\mathcal{O}} v d\mu^s = k \|\mu^s\|_{TV}.$$

Thus, we can continue (3.6) by

$$\|\psi(\mu)\|_{TV} \geq \frac{\psi(y)}{y} \|\mu^a\|_{TV} + k \|\mu^s\|_{TV} - \psi(y) |\mathcal{O}| = \frac{\psi(y)}{y} \|\mu\|_{TV} - \psi(y) |\mathcal{O}|,$$

as required. \square

Proposition 3.7. *In both settings of Definition 2.2, $\varphi : H^{-1} \rightarrow [0, \infty]$ is convex and lower-semicontinuous.*

Proof. In the superlinear case, i. e. Definition 2.2 (i) applies, convexity and lower-semicontinuity of φ are proved in [5, p. 68]. In the sublinear case, i. e. Definition 2.2 (ii) applies, convexity becomes clear by Remark 3.4. It remains to prove lower-semicontinuity in the sublinear case.

Step 1: As a preparatory step, we establish weak* lower-semicontinuity of the functional $\tilde{\varphi} : \mathcal{M}(\mathcal{O}) \rightarrow [0, \infty)$,

$$\tilde{\varphi}(\mu) = \|\psi(\mu)\|_{TV},$$

for which we have

$$\tilde{\varphi}|_{\mathcal{M}(\mathcal{O}) \cap H^{-1}} = \varphi.$$

Consider $\mu_n \rightarrow \mu$ weakly* for $n \rightarrow \infty$. We can assume that $\psi(\mu_n)$ contains a subsequence which is bounded in TV norm (otherwise there is nothing to show). Then we select a subsequence $(\mu_{n_k})_{k \in \mathbb{N}}$ such that $\|\psi(\mu_{n_k})\|_{TV} \rightarrow \liminf_{n \rightarrow \infty} \|\psi(\mu_n)\|_{TV}$ for $k \rightarrow \infty$, from which we can choose a nonrelabeled subsequence $(\psi(\mu_{n_k}))_{k \in \mathbb{N}}$ which converges weakly* to some $\nu \in \mathcal{M}(\mathcal{O})$ (e. g. by [1, Satz 6.5]). By [23, Lemma 2.1], we get that

$$\mathcal{M}(\mathcal{O}) \langle \psi(\mu), \eta \rangle_{\mathcal{C}_c^0(\mathcal{O})} \leq \mathcal{M}(\mathcal{O}) \langle \nu, \eta \rangle_{\mathcal{C}_c^0(\mathcal{O})} = \lim_{k \rightarrow \infty} \mathcal{M}(\mathcal{O}) \langle \psi(\mu_{n_k}), \eta \rangle_{\mathcal{C}_c^0(\mathcal{O})} \leq \lim_{k \rightarrow \infty} \|\psi(\mu_{n_k})\|_{TV} \|\eta\|_{\mathcal{C}_c^0(\mathcal{O})}$$

for $\eta \in \mathcal{C}_c^0(\mathcal{O})$, $\eta \geq 0$. Now, using that $\psi(\rho)$ is a positive measure for any $\rho \in \mathcal{M}(\mathcal{O})$ by (3.3), we obtain

$$\begin{aligned} \|\psi(\mu)\|_{TV} &= \sup_{\substack{\eta \in \mathcal{C}_c^0(\mathcal{O}) \\ \eta \in [0,1]}} \mathcal{M}(\mathcal{O}) \langle \psi(\mu), \eta \rangle_{\mathcal{C}_c^0(\mathcal{O})} \leq \sup_{\substack{\eta \in \mathcal{C}_c^0(\mathcal{O}) \\ \eta \in [0,1]}} \lim_{k \rightarrow \infty} \mathcal{M}(\mathcal{O}) \langle \psi(\mu_{n_k}), \eta \rangle_{\mathcal{C}_c^0(\mathcal{O})} \\ &\leq \sup_{\substack{\eta \in \mathcal{C}_c^0(\mathcal{O}) \\ \eta \in [0,1]}} \lim_{k \rightarrow \infty} \|\psi(\mu_{n_k})\|_{TV} = \liminf_{n \rightarrow \infty} \|\psi(\mu_n)\|_{TV}, \end{aligned}$$

as required.

Step 2: Assume now that $(u_n)_{n \in \mathbb{N}} \subset H^{-1}$, $u \in H^{-1}$, and $u_n \rightarrow u$ for $n \rightarrow \infty$. Being the only non-trivial case, we can assume that $(u_n)_{n \in \mathbb{N}}$ contains a subsequence (which we call again (u_n)) for which $(\varphi(u_n))_{n \in \mathbb{N}}$ is bounded. Thus, there are measures $\mu_n \in \mathcal{M}(\mathcal{O}) \cap H^{-1}$ such that

$$u_n(\eta) = \int_{\mathcal{O}} \eta d\mu_n \quad \text{for all } \eta \in \mathcal{C}_c^1(\mathcal{O}).$$

By definition of φ , $\varphi(u_n) = \|\psi(\mu_n)\|_{TV}$, such that Lemma 3.6 implies that $\|\mu_n\|_{TV}$ is bounded. Thus, there is $\tilde{\mu} \in \mathcal{M}(\mathcal{O})$ an again nonrelabeled subsubsequence $(\mu_n)_{n \in \mathbb{N}}$ such that $\mu_n \xrightarrow{*} \tilde{\mu}$. For $\eta \in \mathcal{C}_c^1(\mathcal{O}) \subseteq \mathcal{C}_c^0(\mathcal{O})$ we have

$$\int_{\mathcal{O}} \eta \, d\tilde{\mu} = \lim_{n \rightarrow \infty} \int_{\mathcal{O}} \eta \, d\mu_n = \lim_{n \rightarrow \infty} u_n(\eta) = u(\eta) \leq \|u\|_{H^{-1}} \|\eta\|_{H_0^1(\mathcal{O})},$$

so $\tilde{\mu} \in \mathcal{M}(\mathcal{O}) \cap H^{-1}$ and $u = \tilde{\mu}$. Using the weak* lower-semicontinuity of $\tilde{\varphi}$ from Step 1, we get

$$\varphi(u) = \tilde{\varphi}(\tilde{\mu}) \leq \liminf_{n \rightarrow \infty} \tilde{\varphi}(\mu_n) = \liminf_{n \rightarrow \infty} \varphi(u_n). \quad (3.7)$$

As this argument works for any bounded subsequence of $(u_n)_{n \in \mathbb{N}}$, (3.7) is also true for the original sequence $(u_n)_{n \in \mathbb{N}}$. \square

As one can see from the definition of the energy functional φ in the second part of Definition 2.2, it has an explicit representation on $H^{-1} \setminus \mathcal{M}(\mathcal{O})$, where it is ∞ , and on $L^1(\mathcal{O}) \cap H^{-1}$, where it is an integral. However, whenever we evaluate φ for general measures in $\mathcal{M}(\mathcal{O}) \cap H^{-1}$, e. g. in the uniqueness part of the proof of Theorem 2.6, we need an approximation reducing it to evaluations on $L^1(\mathcal{O})$ functions. This will be made precise in the following theorem, the proof of which will take the rest of this section.

Theorem 3.8. *Assume that ψ satisfies (2.7) as well as Assumptions 2.1 (A3), (A5'). Let φ be defined as in Definition 2.2 (ii) and $u \in \mathcal{M}(\mathcal{O}) \cap H^{-1}$. Then there exists a sequence $u_n \in L^2(\mathcal{O})$ such that*

$$u_n \rightharpoonup u \quad \text{in } H^{-1}, \quad \text{and} \quad (3.8)$$

$$\varphi(u_n) \rightarrow \varphi(u) \quad (3.9)$$

for $n \rightarrow \infty$.

Corollary 3.9. *Since convex functions on a real Hilbert space are lower-semicontinuous if and only if they are weakly sequentially lower-semicontinuous (see e. g. [15, Theorem 9.1]), Theorem 3.8 implies that φ is the lower-semicontinuous hull of $\varphi|_{L^2(\mathcal{O})}$ in H^{-1} , which means that*

$$\varphi = \sup \{ \beta : H^{-1} \rightarrow [0, \infty] \mid \beta \text{ convex and lower-semicontinuous, } \beta|_{L^2(\mathcal{O})} \leq \varphi|_{L^2(\mathcal{O})} \}, \quad (3.10)$$

where sup denotes the pointwise supremum.

We will approach Theorem 3.8 by giving an explicit construction for the sequence $(\mu_n)_{n \in \mathbb{N}}$, inspired by the construction in [1, Lemma A6.7]. It will rely on applying the original functional to modified functions, which is why we first introduce several modifications to functions on \mathcal{O} .

We next introduce further notation and recall some concepts relying on the regularity of the boundary.

Notations 3.10. Since the domain \mathcal{O} is bounded and smooth, its boundary is locally the graph of a smooth function. More precisely, we recall from [1, Section A6.2] that for each $y \in \partial\mathcal{O}$ there is a neighbourhood $\tilde{U} \subset \mathbb{R}^d$, an orthonormal system e_1, \dots, e_d of \mathbb{R}^d , $r, h \in \mathbb{R}$ with $r > h > 0$, and a smooth bounded function $g : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$, such that with the notation

$$x_{,d} := (x_1, \dots, x_{d-1}), \quad \text{for } x = \sum_{i=1}^d x_i e_i,$$

we have

$$\tilde{U} = \{ x \in \mathbb{R}^d : |x_{,d} - y| < r \text{ and } |x_d - g(x_{,d})| < h \},$$

and for $x \in \tilde{U}$

$$\begin{aligned} x_d &= g(x_{,d}) && \text{if and only if } x \in \partial\mathcal{O}, \\ x_d &\in (g(x_{,d}), g(x_{,d}) + h) && \text{if and only if } x \in \mathcal{O}, \text{ and} \\ x_d &\in (g(x_{,d}) - h, g(x_{,d})) && \text{if and only if } x \notin \mathcal{O}. \end{aligned}$$

For technical reasons we set

$$U = \left\{ x \in \tilde{U} : |x_{,d} - y| < \frac{r}{2} \text{ and } |x_d - g(x_{,d})| < \frac{h}{2} \right\}. \quad (3.11)$$

The boundary $\partial\mathcal{O}$ is covered by those open sets U belonging to all possible reference points y . As $\partial\mathcal{O}$ is compact, we can choose a finite subcovering $(U^j)_{j=1}^l$, and for each U^j , we denote the elements belonging to it by a superindex j , e. g. $y^j, e_d^j, g^j, h^j, \tilde{U}^j$. At last, we fix an open set U^0 with $\overline{U^0} \subset \mathcal{O}$, such that $\overline{\mathcal{O}} \subset \cup_{j=0}^l U^j$ and we set $e_d^0 := 0$.

Subordinate to the covering $\cup_{j=0}^l U^j$, let now ζ^0, \dots, ζ^l be a partition of unity on $\overline{\mathcal{O}}$, i. e. $0 \leq \zeta^j \leq 1, \zeta^j \in C_c^\infty(\mathbb{R}^d), \text{supp}(\zeta^j) \subseteq U^j$ for all $j = 0, \dots, l$, and

$$\sum_{j=0}^l \zeta^j = 1 \quad \text{on } \overline{\mathcal{O}}.$$

For $\eta : \mathcal{O} \rightarrow \mathbb{R}$ and $\mu \in \mathcal{M}(\mathcal{O})$, we define $\eta_{\text{ext}} : \mathbb{R}^d \rightarrow \mathbb{R}$ and $\mu_{\text{ext}} \in \mathcal{M}(\mathbb{R}^d)$ as the extended function (resp. measure) by zero. Finally, we define for $\rho \in C_c^\infty(\mathbb{R}^d)$ with

$$\text{supp}(\rho) \subseteq B_1(0), \quad \int_{\mathcal{O}} \rho \, dx = 1, \quad \rho(x) = \rho(-x) \quad (3.12)$$

a Dirac sequence $(\rho_\delta)_{\delta>0} \subset C_c^\infty(\mathbb{R}^d)$ of mollifiers by

$$\rho_\delta(x) = \frac{1}{\delta^d} \rho\left(\frac{x}{\delta}\right). \quad (3.13)$$

For $\eta \in L^2(\mathbb{R}^d), \mu \in \mathcal{M}(\mathbb{R}^d)$, we then define functions $\rho_\delta * \eta, \rho_\delta * \mu \in C^\infty(\mathbb{R}^d)$ by

$$\rho_\delta * \eta(x) = \int_{\mathbb{R}^d} \rho_\delta(x-y)\eta(y) \, dy \quad \text{and} \quad \rho_\delta * \mu(x) = \int_{\mathbb{R}^d} \rho_\delta(x-y) \, d\mu(y).$$

For brevity, we write $\rho_\delta * \eta := \rho_\delta * \eta_{\text{ext}}$ for $\eta \in L^2(\mathcal{O})$.

The following construction allows to shift a function ‘‘away from the boundary’’.

Definition 3.11. Let $\varepsilon > 0$ and $\eta : \mathcal{O} \rightarrow \mathbb{R}$. Then we define $\eta_\varepsilon : \mathcal{O} \rightarrow \mathbb{R}$ by

$$\eta_\varepsilon(x) = \sum_{j=0}^l \zeta^j(x) \eta_{\text{ext}}(x - \varepsilon e_d^j), \quad (3.14)$$

where we recall that e_d^0 is set to 0.

Remark 3.12. By this construction, we achieve that $\eta_\varepsilon = 0$ on a $w(\varepsilon)$ -neighbourhood of $\partial\mathcal{O}$ with

$$w(\varepsilon) := \min \left\{ \text{dist}(U^0, \mathcal{O}^c), \min_{j=1, \dots, l} \left(\min \left\{ \frac{\varepsilon}{2}, \frac{\varepsilon}{2L^j}, \frac{h^j}{4}, \frac{h^j}{4L^j} \right\} \right) \right\} > 0, \quad (3.15)$$

where L^j denotes the Lipschitz constant of g^j defined in Notations 3.10.

Proof. The number $w(\varepsilon)$ is obviously strictly positive by the construction of the covering $(U^j)_{j=0}^l$. To show the support property, let $j \in \{0, 1, \dots, l\}$ and $U_\varepsilon^j := U^j \cap ((U^j \cap \mathcal{O}) + \varepsilon e_d^j)$. By definition, $\eta_{\text{ext}}(x - \varepsilon e_d^j) = 0$ if $x \in U^j \setminus U_\varepsilon^j$. By the definition of ζ^j , we furthermore conclude that $\zeta^j(x) \eta_{\text{ext}}(x - \varepsilon e_d^j) = 0$ for $x \notin U_\varepsilon^j$. Consequently,

$$\eta_\varepsilon : x \mapsto \sum_{j=0}^l \zeta^j(x) \eta_{\text{ext}}(x - \varepsilon e_d^j)$$

is supported on

$$U_\varepsilon := \bigcup_{j=0}^l U_\varepsilon^j,$$

such that it remains to show that $\text{dist}(U_\varepsilon, \mathcal{O}^c) \geq w(\varepsilon)$, or equivalently, that $\text{dist}(U_\varepsilon^j, \mathcal{O}^c) \geq w(\varepsilon)$ for all $j \in \{0, \dots, l\}$.

For $j = 0$, this is trivial by construction of $U_\varepsilon^0 = U^0$ and $w(\varepsilon)$. For $j = 1, \dots, l$, using the coordinate system $(x_{,d}^j, x_d^j)$ we can rewrite

$$U_\varepsilon^j = \{x \in U^j : x_d^j > g^j(x_{,d}^j) + \varepsilon\}.$$

Hence, we can compute for any $x \in U_\varepsilon^j$, i. e. $x = (x_{,d}^j, g^j(x_{,d}^j) + \varepsilon')$ for some $\varepsilon' \in (\varepsilon, \frac{h^j}{2})$, and $y \in \partial\mathcal{O} \cap \tilde{U}^j$

$$\begin{aligned} \|x - y\|^2 &= \|x_{,d} - y_{,d}\|^2 + |g(x_{,d}) + \varepsilon' - g(y_{,d})|^2 \\ &\geq \|x_{,d} - y_{,d}\|^2 + (\varepsilon' - |g(x_{,d}) - g(y_{,d})|)^2, \end{aligned}$$

where $\|\cdot\|$ denotes the Euclidean norm both in \mathbb{R}^d and in \mathbb{R}^{d-1} . Letting L^j be the Lipschitz constant of g^j , we can then argue that either $\|x_{,d} - y_{,d}\| > \frac{\varepsilon}{2L^j}$ or

$$|g(x_{,d}) - g(y_{,d})| \leq L^j \frac{\varepsilon}{2L^j} = \frac{\varepsilon}{2},$$

such that $\text{dist}(U_\varepsilon^j, \partial\mathcal{O} \cap \tilde{U}^j)$ is at least $\min\{\frac{\varepsilon}{2}, \frac{\varepsilon}{2L^j}\}$. By similar arguments, we can obtain from the construction of U^j in (3.11) (note that $r^j > h^j$ by construction) that

$$\text{dist}(U_\varepsilon^j, (\tilde{U}^j)^c) \geq \min\left\{\frac{h^j}{4}, \frac{h^j}{4L^j}\right\},$$

such that we conclude

$$\begin{aligned} \text{dist}(U_\varepsilon^j, \partial\mathcal{O}) &= \min\{\text{dist}(U_\varepsilon^j, \partial\mathcal{O} \cap \tilde{U}^j), \text{dist}(U_\varepsilon^j, \partial\mathcal{O} \cap (\tilde{U}^j)^c)\} \\ &\geq \min\{\text{dist}(U_\varepsilon^j, \partial\mathcal{O} \cap \tilde{U}^j), \text{dist}(U_\varepsilon^j, (\tilde{U}^j)^c)\} \\ &\geq \min\left\{\frac{\varepsilon}{2}, \frac{\varepsilon}{2L^j}, \frac{h^j}{4}, \frac{h^j}{4L^j}\right\} \geq w(\varepsilon). \end{aligned}$$

□

This allows to define the following approximating objects for $u \in \mathcal{M}(\mathcal{O}) \cap H^{-1}$.

Definition 3.13. Let $\varepsilon > 0$, $0 < \delta \leq \frac{w(\varepsilon)}{2}$ and $u \in \mathcal{M}(\mathcal{O}) \cap H^{-1}$. We then define for $\eta \in H_0^1(\mathcal{O})$

$$\begin{aligned} \tilde{u}_\varepsilon(\eta) &=_{H^{-1}} \langle u, \eta_\varepsilon \rangle_{H_0^1(\mathcal{O})} \\ \text{and } \tilde{u}_{\varepsilon, \delta}(\eta) &=_{H^{-1}} \langle u, \rho_\delta * \eta_\varepsilon \rangle_{H_0^1(\mathcal{O})}. \end{aligned} \tag{3.16}$$

These functionals are in H^{-1} by Lemma 3.14 and Lemma 3.15 below. For $\eta \in C_0^0(\mathcal{O})$, we define

$$\begin{aligned} \mu_\varepsilon(\eta) &=_{\mathcal{M}(\mathcal{O})} \langle \mu, \eta_\varepsilon \rangle_{C_0^0(\mathcal{O})} \\ \text{and } \mu_{\varepsilon, \delta}(\eta) &=_{\mathcal{M}(\mathcal{O})} \langle \mu, \rho_\delta * \eta_\varepsilon \rangle_{C_0^0(\mathcal{O})}. \end{aligned} \tag{3.17}$$

These functionals are in $\mathcal{M}(\mathcal{O})$ by Lemma 3.16 below. By uniqueness of the linear continuation, this allows to conclude that

$$u_\varepsilon, u_{\varepsilon, \delta} \in \mathcal{M} \cap H^{-1}, \quad \text{as well as } u_\varepsilon = \tilde{u}_\varepsilon \text{ and } u_{\varepsilon, \delta} = \tilde{u}_{\varepsilon, \delta}.$$

Lemma 3.14. Let $\varepsilon > 0$ and $\eta \in H_0^1(\mathcal{O})$. Then the map $H_0^1(\mathcal{O}) \ni \eta \mapsto \eta_\varepsilon \in H_0^1(\mathcal{O})$ is linear, and

$$\|\eta_\varepsilon\|_{H_0^1(\mathcal{O})} \leq C \|\eta\|_{H_0^1(\mathcal{O})},$$

where C only depends on the localising functions $(\zeta^j)_{j=0}^l$, the number of covering sets l , the Poincaré constant of the domain \mathcal{O} and the spatial dimension d .

Proof. The proof of the linearity claim is straightforward and therefore skipped. In order to prove boundedness, let $V^j = U^j \cap \mathcal{O}$ and $U_\varepsilon^j := U^j \cap ((U^j \cap \mathcal{O}) + \varepsilon e_d^j)$ as before. We first note

$$\|\eta_\varepsilon\|_{H_0^1(\mathcal{O})} = \left\| \sum_{j=0}^l \zeta^j \eta_\varepsilon^j \right\|_{H_0^1(\mathcal{O})} \leq \sum_{j=0}^l \|\zeta^j \eta_\varepsilon^j\|_{H_0^1(\mathcal{O})}, \tag{3.18}$$

where we have written

$$\eta_\varepsilon^j \in H^1(\mathbb{R}^d), \quad \eta_\varepsilon^j(x) = \eta_{\text{ext}}(x - \varepsilon e_d^j).$$

We now analyse the summands separately, where we make use of the fact that for all $j \in \{1, \dots, l\}$, $\zeta^j \in \mathcal{C}_c^\infty(U^j)$ and $\zeta^j \eta_\varepsilon^j$ is supported on V^j . In the following, $(\partial_i)_{i=1}^d$ represent the weak partial derivatives of first order. We then compute for $i \in \{1, \dots, d\}$

$$\begin{aligned} \|\partial_i(\zeta^j \eta_\varepsilon^j)\|_{L^2(\mathcal{O})} &= \|\partial_i(\zeta^j \eta_\varepsilon^j)\|_{L^2(V^j)} \leq \|(\partial_i \zeta^j) \eta_\varepsilon^j\|_{L^2(V^j)} + \|\zeta^j \partial_i \eta_\varepsilon^j\|_{L^2(V^j)} \\ &\leq C \|\eta_\varepsilon^j\|_{L^2(V^j)} + \left(\int_{V^j} \left| \partial_i(\eta_{\text{ext}}(x - \varepsilon e_d^j)) \right|^2 dx \right)^{\frac{1}{2}} \\ &\leq C \|\eta\|_{L^2(\mathcal{O})} + \left(\int_{U_\varepsilon^j} \left| (\partial_i \eta)(x - \varepsilon e_d^j) \right|^2 dx \right)^{\frac{1}{2}} \\ &\leq C \|\eta\|_{L^2(\mathcal{O})} + \|\partial_i \eta\|_{L^2(\mathcal{O})}. \end{aligned}$$

This yields

$$\begin{aligned} \|\zeta^j \eta_\varepsilon^j\|_{H_0^1(\mathcal{O})}^2 &= \sum_{i=1}^d \|\partial_i(\zeta^j \eta_\varepsilon^j)\|_{L^2(\mathcal{O})}^2 \leq \sum_{i=1}^d \left(C \|\eta\|_{L^2(\mathcal{O})} + \|\partial_i \eta\|_{L^2(\mathcal{O})} \right)^2 \\ &\leq C \|\eta\|_{H_0^1(\mathcal{O})}^2 + 2 \sum_{i=1}^d \|\partial_i \eta\|_{L^2(\mathcal{O})}^2 \leq C \|\eta\|_{H_0^1(\mathcal{O})}^2, \end{aligned}$$

where C may depend on d, \mathcal{O} (through the Poincaré constant) and ζ^j . Thus, we can continue (3.18) by

$$\|\eta_\varepsilon\|_{H_0^1(\mathcal{O})} \leq \sum_{j=0}^l \|\zeta^j \eta_\varepsilon^j\|_{H_0^1(\mathcal{O})} \leq (l+1)C \|\eta\|_{H_0^1(\mathcal{O})},$$

as required. \square

Concerning the mollification step, we note that by Remark 3.12, $\rho_\delta * \eta_\varepsilon(x) = 0$ if $\text{dist}(x, \partial\mathcal{O}) \leq \frac{w(\varepsilon)}{2}$ and $0 < \delta \leq \frac{w(\varepsilon)}{2}$, so that in this case we can restrict $\rho_\delta * \eta_\varepsilon$ to \mathcal{O} to get a $\mathcal{C}_c^1(\mathcal{O})$ function. By a slight abuse of notation, we then write

$$(\rho_\delta * \eta_\varepsilon)|_{\mathcal{O}} = \rho_\delta * \eta_\varepsilon \in \mathcal{C}_c^1(\mathcal{O}) \subseteq H_0^1(\mathcal{O}) \cap \mathcal{C}_0^0(\mathcal{O}). \quad (3.19)$$

Also for this step, we have to ensure linearity, which is clear, and an estimate on the $H_0^1(\mathcal{O})$ norm, which is done in the following lemma.

Lemma 3.15. *Let $\varepsilon > 0$ and $0 < \delta \leq \frac{w(\varepsilon)}{2}$. Then the map $H_0^1(\mathcal{O}) \ni \eta \mapsto \eta_{\varepsilon, \delta} \in H_0^1(\mathcal{O})$ is linear, and*

$$\|\rho_\delta * \eta_\varepsilon\|_{H_0^1(\mathcal{O})} \leq C \|\eta\|_{H_0^1(\mathcal{O})} \quad \text{for all } \eta \in H_0^1(\mathcal{O}),$$

where C is the constant from Lemma 3.14.

Proof. The proof of linearity is straightforward. In order to show boundedness, for any $g \in L^2(\mathcal{O})$ such that $\rho_\delta * g = 0$ on \mathcal{O}^c we can compute

$$\begin{aligned} \|\rho_\delta * g\|_{L^2(\mathcal{O})}^2 &= \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} \rho_\delta(x-y) g_{\text{ext}}(y) dy \right)^2 dx \leq \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \rho_\delta(x-y) (g_{\text{ext}}(y))^2 dy dx \\ &= \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \rho_\delta(x-y) dx (g_{\text{ext}}(y))^2 dy = \|g_{\text{ext}}\|_{L^2(\mathbb{R}^d)}^2 = \|g\|_{L^2(\mathcal{O})}^2, \end{aligned} \quad (3.20)$$

where in the second step we could apply Jensen's inequality since $\rho_\delta(x-y) dy$ is a probability measure for each $x \in \mathbb{R}^d$. By Remark 3.12 for all $i \in \{1, \dots, d\}$, $\rho_\delta * (\partial_i \eta_\varepsilon)$ vanishes outside of \mathcal{O} if $0 < \delta \leq \frac{w(\varepsilon)}{2}$. Hence g in (3.20) can be replaced by each partial derivative $\partial_i \eta_\varepsilon$ which yields

$$\begin{aligned} \|\rho_\delta * \eta_\varepsilon\|_{H_0^1(\mathcal{O})}^2 &= \sum_{i=1}^d \|\partial_i(\rho_\delta * \eta_\varepsilon)\|_{L^2(\mathcal{O})}^2 = \sum_{i=1}^d \|\rho_\delta * \partial_i(\eta_\varepsilon)\|_{L^2(\mathcal{O})}^2 \\ &\leq \sum_{i=1}^d \|\partial_i \eta_\varepsilon\|_{L^2(\mathcal{O})}^2 = \|\eta_\varepsilon\|_{H_0^1(\mathcal{O})}^2 \leq C \|\eta\|_{H_0^1(\mathcal{O})}^2, \end{aligned}$$

where the second equality can be found e. g. in [1, Section 2.23] and the last inequality is the statement of Lemma 3.14. \square

Lemma 3.16. *Let $\varepsilon > 0$, $0 < \delta \leq \frac{w(\varepsilon)}{2}$ and $\eta \in \mathcal{C}_c^0(\mathcal{O})$. Then, the map*

$$\mathcal{C}_c^0(\mathcal{O}) \ni \eta \mapsto (\eta_\varepsilon, \rho_\delta * \eta_\varepsilon) \in (\mathcal{C}_c^0(\mathcal{O}))^2$$

is linear. Furthermore, we have

$$\|\rho_\delta * \eta_\varepsilon\|_\infty \leq \|\eta_\varepsilon\|_\infty \leq \|\eta\|_\infty, \quad (3.21)$$

where $\|\cdot\|_\infty$ denotes the supremum norm.

Proof. The proof of the linearity claim is straightforward. In order to show boundedness, we first note that for $0 < \delta \leq \frac{w(\varepsilon)}{2}$, $\rho_\delta * \eta_\varepsilon \in \mathcal{C}_c^0(\mathcal{O})$ by construction and Remark 3.12. To obtain (3.21), we estimate for arbitrary $x \in \mathcal{O}$

$$|\eta_\varepsilon(x)| \leq \sum_{j=0}^l \zeta^j(x) \left| \eta_{\text{ext}}(x - \varepsilon e_d^j) \right| \leq \sum_{j=0}^l \zeta^j(x) \|\eta\|_\infty = \|\eta\|_\infty,$$

which yields the second relation. The first one can be seen by

$$|\rho_\delta * \eta_\varepsilon(x)| \leq \int_{\mathbb{R}^d} \rho_\delta(x - y) \|\eta_\varepsilon\|_\infty \, dx = \|\eta_\varepsilon\|_\infty,$$

which concludes the proof. \square

We next analyse how φ as given in Definition 2.2 (ii) acts on the approximating measures from Definition 3.13. First, we state that if μ is absolutely continuous with respect to the Lebesgue measure, so is μ_ε , which we show by giving its density.

Lemma 3.17. *Let $\varepsilon > 0$, $h \in L^1(\mathcal{O})$ and $\mu := h \, dx \in \mathcal{M}(\mathcal{O})$. Then μ_ε has the density*

$$\mathcal{O} \ni x \mapsto \sum_{j=0}^l \zeta^j(x + \varepsilon e_d^j) h_{\text{ext}}(x + \varepsilon e_d^j)$$

with respect to the Lebesgue measure.

A more direct construction of $\mu_{\varepsilon,\delta}$ is given by the following lemma.

Lemma 3.18. *Let $\varepsilon, \delta \leq \frac{w(\varepsilon)}{2}$. Then the measure $\tilde{\mu}_{\varepsilon,\delta} \in \mathcal{M}(\mathcal{O})$ defined by*

$$\tilde{\mu}_{\varepsilon,\delta} := ((\rho_\delta * \mu_{\text{ext}})|_{\mathcal{O}} \, dx)_\varepsilon, \quad (3.22)$$

coincides with $\mu_{\varepsilon,\delta}$.

The proofs of the preceding lemmas are straightforward and therefore suppressed.

In the rest of this section, we will argue that the sequence

$$\left(\mu_{\frac{1}{n}, \frac{1}{2}w(\frac{1}{n})} \right)_{n \in \mathbb{N}}$$

is an approximation of $\mu \in \mathcal{M}(\mathcal{O}) \cap H^{-1}$ in the sense of Theorem 3.8. First we address the regularity of $\mu_{\varepsilon,\delta}$, where $\varepsilon > 0$ and $0 < \delta \leq \frac{w(\varepsilon)}{2}$.

Lemma 3.19. *For all $\varepsilon > 0$, $0 < \delta \leq \frac{w(\varepsilon)}{2}$, the approximating measures $\mu_{\varepsilon,\delta}$ have a bounded density with respect to Lebesgue measure.*

Proof. The fact that $\mu_{\varepsilon,\delta}$ has a density with respect to Lebesgue measure follows from its characterisation in Lemma 3.18 and Lemma 3.17. This density is bounded in space since

$$\left| \sum_{j=0}^l \zeta^j(x + \varepsilon e_d^j) (\rho_{\frac{1}{2}w(\varepsilon)} * \mu_{\text{ext}})|_{\mathcal{O}}(x) \right| \leq l \sup_{x \in \mathcal{O}} \left| \rho_{\frac{1}{2}w(\varepsilon)} * \mu(x) \right| \leq l \sup_{x \in \mathbb{R}^d} \left| \rho_{\frac{1}{2}w(\varepsilon)}(x) \right| \|\mu\|_{TV}.$$

\square

The first part of the following proposition allows to deduce property (3.8), while the second part is needed for the further proof of (3.9).

Proposition 3.20. *Let ρ be as in (3.13) and $0 < \delta_\varepsilon \leq \frac{w(\varepsilon)}{2}$.*

1. For $\eta \in H_0^1(\mathcal{O})$, we have

$$\rho_{\delta_\varepsilon} * \eta_\varepsilon \rightarrow \eta \quad \text{for } \varepsilon \rightarrow 0 \text{ in } H_0^1(\mathcal{O}). \quad (3.23)$$

2. For $\eta \in C_c^0(\mathcal{O})$, we have

$$\rho_{\delta_\varepsilon} * \eta_\varepsilon \rightarrow \eta \quad \text{for } \varepsilon \rightarrow 0 \text{ in } C_c^0(\mathcal{O}). \quad (3.24)$$

Proof. Throughout this proof, we will write δ instead of δ_ε , always assuming that $0 < \delta \leq \frac{w(\varepsilon)}{2}$.

Proof of part 1: It is enough to show that for all $i \in \{1, \dots, d\}$

$$\|\partial_i(\rho_\delta * \eta_\varepsilon) - \partial_i \eta\|_{L^2(\mathcal{O})} \rightarrow 0 \quad \text{for } \varepsilon \rightarrow 0. \quad (3.25)$$

By the density of $C_0^\infty(\mathcal{O})$ in $H_0^1(\mathcal{O})$, for any $\beta > 0$ we can choose $\varphi \in C_0^\infty(\mathcal{O})$ such that

$$\max \left\{ \|\varphi - \eta\|_{L^2(\mathcal{O})}, \|\partial_i \varphi - \partial_i \eta\|_{L^2(\mathcal{O})} \right\} \leq \frac{\beta}{6(l+1)\tilde{C}}, \quad (3.26)$$

where

$$\tilde{C} := \max \left\{ \max_{j=1, \dots, l} \left(\sup_{\mathbb{R}^d} |\partial_i \zeta^j| \right), 1 \right\}$$

As $\varphi_{\text{ext}}, \zeta^j \in C_b^1(\mathcal{O})$ for each $j \in \{1, \dots, l\}$, we can choose $\varepsilon_0 > 0$ small enough, such that for all $x \in \mathbb{R}^d$ and $y, z \in B_{\varepsilon_0}(x)$

$$|\partial_i \zeta^j(y) \varphi_{\text{ext}}(z) - \partial_i \zeta^j(x) \varphi_{\text{ext}}(x)| \leq \frac{\beta}{6(l+1)|\mathcal{O}|^{\frac{1}{2}}} \quad (3.27)$$

and

$$|\zeta^j(y) \partial_i \varphi_{\text{ext}}(z) - \zeta^j(x) \partial_i \varphi_{\text{ext}}(x)| \leq \frac{\beta}{6(l+1)|\mathcal{O}|^{\frac{1}{2}}}. \quad (3.28)$$

We approach (3.25) by splitting the term under consideration into the more convenient pieces

$$\begin{aligned} \|\partial_i(\rho_\delta * \eta_\varepsilon) - \partial_i \eta\|_{L^2(\mathcal{O})} &= \|\rho_\delta * \partial_i \eta_\varepsilon - \partial_i \eta_{\text{ext}}\|_{L^2(\mathbb{R}^d)} \\ &= \|\rho_\delta * \partial_i(\eta_\varepsilon - \varphi_\varepsilon) + \rho_\delta * \partial_i \varphi_\varepsilon - \partial_i \varphi_{\text{ext}} + \partial_i \varphi_{\text{ext}} - \partial_i \eta_{\text{ext}}\|_{L^2(\mathbb{R}^d)} \\ &\leq \|\rho_\delta * \partial_i(\eta_\varepsilon - \varphi_\varepsilon)\|_{L^2(\mathbb{R}^d)} + \|\rho_\delta * \partial_i \varphi_\varepsilon - \partial_i \varphi_{\text{ext}}\|_{L^2(\mathbb{R}^d)} + \|\partial_i \varphi_{\text{ext}} - \partial_i \eta_{\text{ext}}\|_{L^2(\mathbb{R}^d)} \\ &= \text{(I)} + \text{(II)} + \text{(III)}. \end{aligned}$$

We estimate the summands separately. For the first one we get with the convolution estimate (e. g. [1, Section 2.13])

$$\begin{aligned} \text{(I)} &\leq \|\partial_i(\eta_\varepsilon - \varphi_\varepsilon)\|_{L^2(\mathbb{R}^d)} \\ &= \left\| \sum_{j=0}^l \partial_i [\zeta^j(\eta_{\text{ext}}(\cdot - \varepsilon e_d^j) - \varphi_{\text{ext}}(\cdot - \varepsilon e_d^j))] \right\|_{L^2(\mathbb{R}^d)} \\ &\leq \sum_{j=0}^l \left\| \partial_i \zeta^j \left(\eta_{\text{ext}}(\cdot - \varepsilon e_d^j) - \varphi_{\text{ext}}(\cdot - \varepsilon e_d^j) \right) \right\|_{L^2(\mathbb{R}^d)} \\ &\quad + \sum_{j=0}^l \left\| \zeta^j \left(\partial_i \eta_{\text{ext}}(\cdot - \varepsilon e_d^j) - \partial_i \varphi_{\text{ext}}(\cdot - \varepsilon e_d^j) \right) \right\|_{L^2(\mathbb{R}^d)} \\ &\leq \sum_{j=0}^l \left(\sup_{\mathbb{R}^d} |\partial_i \zeta^j| \|\eta_{\text{ext}} - \varphi_{\text{ext}}\|_{L^2(\mathbb{R}^d)} + \|\partial_i \eta_{\text{ext}} - \partial_i \varphi_{\text{ext}}\|_{L^2(\mathbb{R}^d)} \right) \leq \frac{\beta}{3}, \end{aligned} \quad (3.29)$$

where we used (3.26) in the last step. For the second term, we recall that $(\zeta^j)_{j=0}^l$ is a partition of unity on the support of φ . Thus, we can compute

$$\begin{aligned}
(\text{II}) &\leq \sum_{j=0}^l \left\| \rho_\delta * \partial_i \left(\zeta^j \varphi_{\text{ext}}(\cdot - \varepsilon e_d^j) \right) - \partial_i \left(\zeta^j \varphi_{\text{ext}} \right) \right\|_{L^2(\mathbb{R}^d)} \\
&= \sum_{j=0}^l \left\| \rho_\delta * \left(\partial_i \zeta^j \varphi_{\text{ext}}(\cdot - \varepsilon e_d^j) + \zeta^j \partial_i \varphi_{\text{ext}}(\cdot - \varepsilon e_d^j) \right) - \partial_i \zeta^j \varphi_{\text{ext}} - \zeta^j \partial_i \varphi_{\text{ext}} \right\|_{L^2(\mathbb{R}^d)} \\
&\leq \sum_{j=0}^l \left\| \rho_\delta * \left(\partial_i \zeta^j \varphi_{\text{ext}}(\cdot - \varepsilon e_d^j) \right) - \partial_i \zeta^j \varphi_{\text{ext}} \right\|_{L^2(\mathbb{R}^d)} + \sum_{j=0}^l \left\| \rho_\delta * \left(\zeta^j \partial_i \varphi_{\text{ext}}(\cdot - \varepsilon e_d^j) \right) - \zeta^j \partial_i \varphi_{\text{ext}} \right\|_{L^2(\mathbb{R}^d)} \\
&=: \sum_{j=0}^l (\text{IV})_j + \sum_{j=0}^l (\text{V})_j.
\end{aligned}$$

$(\text{IV})_j$ and $(\text{V})_j$ are treated analogously, so we only show the estimate for $(\text{V})_j$, where we choose $\varepsilon < \frac{\varepsilon_0}{2}$ with ε_0 as for (3.27). Noting that ρ_δ integrates to 1 for any $\delta > 0$ and using Jensen's inequality in the second step, we obtain

$$\begin{aligned}
(\text{V})_j^2 &= \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}^d} \rho_\delta(x-y) \left(\zeta^j(y) \partial_i \varphi_{\text{ext}}(y - \varepsilon e_d^j) - \zeta^j(x) \partial_i \varphi_{\text{ext}}(x) \right) dy \right|^2 dx \\
&\leq \int_{\mathbb{R}^d} \int_{B_\delta(x)} \rho_\delta(x-y) \left| \zeta^j(y) \partial_i \varphi_{\text{ext}}(y - \varepsilon e_d^j) - \zeta^j(x) \partial_i \varphi_{\text{ext}}(x) \right|^2 dy dx. \tag{3.30}
\end{aligned}$$

As $\partial_i \varphi_{\text{ext}}$ is supported on \mathcal{O} and, for the analogous step for (IV) , so is φ_{ext} , we can argue as in the proof of Remark 3.12 to see that the integrand of the outer integral is supported on \mathcal{O} . Thus, we can restrict the integration domain to obtain

$$\begin{aligned}
(3.30) &= \int_{\mathcal{O}} \int_{B_\delta(x)} \rho_\delta(x-y) \left| \zeta^j(y) \partial_i \varphi_{\text{ext}}(y - \varepsilon e_d^j) - \zeta^j(x) \partial_i \varphi_{\text{ext}}(x) \right|^2 dy dx \\
&\leq \int_{\mathcal{O}} \frac{\beta^2}{36(l+1)^2 |\mathcal{O}|} \int_{\mathbb{R}^d} \rho_\delta(x-y) dy dx = \left(\frac{\beta}{6(l+1)} \right)^2.
\end{aligned}$$

While we have used (3.28) in the second step, the estimate for $(\text{IV})_j$ uses (3.27) instead and gets the same result. We conclude

$$(\text{II}) = \sum_{j=0}^l \left((\text{IV})_j + (\text{V})_j \right) \leq \frac{\beta}{3}. \tag{3.31}$$

Finally the estimate

$$(\text{III}) \leq \frac{\beta}{3} \tag{3.32}$$

is obvious by property (3.26). Collecting (3.29), (3.31), and (3.32), we obtain

$$\|\partial_i(\rho_\delta * \eta_\varepsilon - \eta)\|_{L^2(\mathcal{O})} \leq \beta$$

only by choosing ε small enough and adapting $0 < \delta \leq \frac{w(\varepsilon)}{2}$, which proves (3.23).

Proof of part 2: Since η is now assumed to be continuous and to have compact support, it is uniformly continuous. For arbitrary $\beta > 0$, we can thus fix $\varepsilon_0 > 0$ such that for all $x, y \in \mathbb{R}^d$

$$|x - y| \leq \varepsilon_0 \quad \text{implies} \quad |\eta_{\text{ext}}(x) - \eta_{\text{ext}}(y)| \leq \frac{\beta}{l+1}.$$

For $\varepsilon \leq \frac{1}{2}\varepsilon_0$, we use $\delta \leq \frac{w(\varepsilon)}{2} \leq \varepsilon$ by (3.15) to calculate for $x \in \mathcal{O}$

$$\begin{aligned} |\rho_\delta * \eta_\varepsilon(x) - \eta(x)| &= \left| \int_{B_\delta(x)} \rho_\delta(x-y) \left(\sum_{j=0}^l \zeta^j(y) (\eta_{\text{ext}}(y - \varepsilon e_d^j) - \eta(x)) \right) dy \right| \\ &\leq \int_{B_\delta(x)} \rho_\delta(x-y) \sum_{j=0}^l \left| \eta_{\text{ext}}(y - \varepsilon e_d^j) - \eta_{\text{ext}}(x) \right| dy \\ &\leq \int_{B_\delta(x)} \rho_\delta(x-y) \sum_{j=0}^l \frac{\beta}{l+1} dy = \beta, \end{aligned}$$

where for the second step we observe that for $y \in B_\delta(x)$, we have

$$\left| (y - \varepsilon e_d^j) - x \right| \leq \delta + \varepsilon \leq 2\varepsilon \leq \varepsilon_0.$$

This proves (3.24). \square

We now turn to prove Property (3.9). Recall the definition of a convex function of a measure from Definition 3.3. We need some more lemmas on measures obtained by this technique, the first of which can be found in [23, Equation (2.11)].

Lemma 3.21. *Let ψ satisfy (2.7) as well as conditions Assumptions 2.1 (A3), (A5'). Let $\mu \in \mathcal{M}(\mathbb{R}^d)$ and let $(\rho_\delta)_{\delta>0}$ be a family of mollifying kernels as specified in (3.12) and (3.13). Then*

$$\int_{\mathbb{R}^d} \psi(\rho_\delta * \mu) dx \leq \int_{\mathbb{R}^d} \psi(\mu) \quad \text{for all } \delta > 0. \quad (3.33)$$

Remark 3.22. Given the assumptions on ψ , the theory of Definition 3.3 indeed also applies to finite measures on \mathbb{R}^d (cf. [55, p. 202]).

Lemma 3.23. *Let ψ satisfy (2.7) as well as conditions Assumptions 2.1 (A3), (A5'). For $\mu \in \mathcal{M}(\mathcal{O})$ we have*

$$\int_{\mathbb{R}^d} \psi(\mu_{\text{ext}}) = \int_{\mathcal{O}} \psi(\mu). \quad (3.34)$$

Proof. We define

$$\mathcal{D}_1 := \left\{ \int_{\mathcal{O}} v d\mu - \int_{\mathcal{O}} \psi^*(v) dx : v \in L^1(\mu), \psi^*(v) \in L^1(\mathcal{O}) \right\}$$

and

$$\mathcal{D}_2 := \left\{ \int_{\mathbb{R}^d} v d\mu_{\text{ext}} - \int_{\mathbb{R}^d} \psi^*(v) dx : v \in L^1(\mu_{\text{ext}}), \psi^*(v) \in L^1(\mathbb{R}^d) \right\},$$

which allows us to write

$$\int_{\mathcal{O}} \psi(\mu) = \sup \mathcal{D}_1 \quad \text{and} \quad \int_{\mathbb{R}^d} \psi(\mu_{\text{ext}}) = \sup \mathcal{D}_2.$$

We note that for v satisfying the conditions of \mathcal{D}_1 , v_{ext} satisfies the conditions of \mathcal{D}_2 , while the involved integrals agree due to the definition of μ_{ext} and $\psi^*(0) = 0$. This yields “ \geq ”.

Conversely, for v satisfying the conditions of \mathcal{D}_2 we can define $\tilde{v} = v|_{\mathcal{O}}$. \tilde{v} satisfies the conditions of \mathcal{D}_1 . Furthermore, we have

$$\begin{aligned} \int_{\mathcal{O}} \tilde{v} d\mu &= \int_{\mathbb{R}^d} v d\mu_{\text{ext}} \quad \text{and} \\ \int_{\mathcal{O}} \psi^*(\tilde{v}) dx &\leq \int_{\mathbb{R}^d} \psi^*(v) dx \quad \text{due to } \psi^* \geq 0. \end{aligned}$$

Thus, we have found an element in \mathcal{D}_1 being larger than or equal to

$$\int_{\mathbb{R}^d} v d\mu_{\text{ext}} - \int_{\mathbb{R}^d} \psi^*(v) dx,$$

which yields “ \leq ”, completing the proof. \square

The key tool to prove the approximation property (3.9) is the following proposition.

Proposition 3.24. *Let $\varepsilon > 0$ and $0 < \delta \leq \frac{w(\varepsilon)}{2}$. Then,*

$$\|\psi(\mu_{\varepsilon, \delta})\|_{TV} \leq \|\psi(\mu)\|_{TV}. \quad (3.35)$$

Proof. Recall Notations 3.10 and let $V^j = U^j \cap \mathcal{O}$. Let $(\xi_\alpha)_{\alpha > 0} \subset C_c^0(\mathbb{R}^d)$ be a sequence of non-negative cut-off functions compactly supported in \mathcal{O} , which converge to 1 pointwise in \mathcal{O} for $\alpha \rightarrow 0$, and each of which is monotonically increasing on each V^j in e_d^j direction.

Let $h \in L^1(\mathcal{O})$ and $\mu = h dx$. In the following argument, we will need $\xi_\alpha(x) \geq \xi_\alpha(x - \varepsilon e_d^j)$ for $x \in V^j$, where $x - \varepsilon e_d^j$ is not a priori in \mathcal{O} . However, since $\xi_\alpha = 0$ outside of \mathcal{O} , it is clear that the statement is valid even if $x - \varepsilon e_d^j \notin \mathcal{O}$. By the convexity of ψ , the construction of $(\zeta^j)_{j=0}^l$ and Lemma 3.17, we then estimate

$$\begin{aligned} \int_{\mathcal{O}} \xi_\alpha \psi(\mu_\varepsilon) &= \int_{\mathcal{O}} \xi_\alpha(x) \psi \left(\sum_{j=0}^l \zeta^j(x + \varepsilon e_d^j) h_{\text{ext}}(x + \varepsilon e_d^j) \right) dx \\ &\leq \int_{\mathcal{O}} \xi_\alpha(x) \sum_{j=0}^l \zeta^j(x + \varepsilon e_d^j) \psi(h_{\text{ext}}(x + \varepsilon e_d^j)) dx \\ &= \int_{\mathbb{R}^d} \xi_\alpha(x) \sum_{j=0}^l \zeta^j(x + \varepsilon e_d^j) \psi(h_{\text{ext}}(x + \varepsilon e_d^j)) dx \\ &= \int_{\mathbb{R}^d} \psi(h_{\text{ext}}(x)) \sum_{j=0}^l \xi_\alpha(x - \varepsilon e_d^j) \zeta^j(x) dx. \end{aligned} \quad (3.36)$$

We note that $\sum_{j=0}^l \xi_\alpha(x - \varepsilon e_d^j) \zeta^j(x)$ is supported on \mathcal{O} by Remark 3.12. Furthermore, by the construction of ξ_α , we have

$$\xi_\alpha(x - \varepsilon e_d^j) \leq \xi_\alpha(x)$$

for all $x \in V^j$, so this holds especially for $x \in \mathcal{O}$ for which $\zeta^j(x) > 0$. Thus, we can continue

$$\begin{aligned} (3.36) &= \sum_{j=0}^l \int_{\mathcal{O}} \xi_\alpha(x - \varepsilon e_d^j) \zeta^j(x) \psi(h(x)) dx \leq \int_{\mathcal{O}} \sum_{j=0}^l \zeta^j(x) \xi_\alpha(x) \psi(h(x)) dx \\ &= \int_{\mathcal{O}} \xi_\alpha(x) \psi(h(x)) dx = \int_{\mathcal{O}} \xi_\alpha \psi(\mu). \end{aligned} \quad (3.37)$$

For a positive Radon measure μ , we have $\mu(\mathcal{O}) = \sup\{\mu(K) : K \subseteq \mathcal{O} \text{ compact}\}$. Since any such K is included in

$$K_\alpha := \{x \in \mathcal{O} : \text{dist}(x, \mathcal{O}^c) \geq \alpha\}$$

for α small enough, we can as well write $\mu(\mathcal{O}) = \lim_{\alpha \rightarrow 0} \mu(K_\alpha)$. Then, noting that $\xi_\alpha \geq \mathbb{1}_{K_\alpha}$, we can argue by definition of the Radon measure of compact sets that

$$\mu(\mathcal{O}) \geq \int_{\mathcal{O}} \xi_\alpha d\mu \geq \mu(K_\alpha) \xrightarrow{\alpha \rightarrow 0} \mu(\mathcal{O}),$$

thus $\mu(\mathcal{O}) = \lim_{\alpha \rightarrow 0} \int_{\mathcal{O}} \xi_\alpha d\mu$.

Hence, we conclude by (3.37) for $\mu = h dx$, $h \in L^1(\mathcal{O})$, that

$$\int_{\mathcal{O}} \psi(\mu_\varepsilon) = \lim_{\alpha \rightarrow 0} \int_{\mathcal{O}} \xi_\alpha \psi(\mu_\varepsilon) \leq \lim_{\alpha \rightarrow 0} \int_{\mathcal{O}} \xi_\alpha \psi(\mu) = \int_{\mathcal{O}} \psi(\mu). \quad (3.38)$$

Using (3.38), Lemma 3.21 and Lemma 3.23, we then obtain for $0 < \delta \leq \frac{w(\varepsilon)}{2}$

$$\begin{aligned} \int_{\mathcal{O}} \psi(\mu_{\varepsilon, \delta}) &= \int_{\mathcal{O}} \psi((\rho_\delta * \mu_{\text{ext}})|_{\mathcal{O}} dx)_\varepsilon \leq \int_{\mathcal{O}} \psi((\rho_\delta * \mu_{\text{ext}})|_{\mathcal{O}}) dx \\ &= \int_{\mathbb{R}^d} \psi(\rho_\delta * \mu_{\text{ext}}) \mathbb{1}_{\mathcal{O}} dx \leq \int_{\mathbb{R}^d} \psi(\rho_\delta * \mu_{\text{ext}}) dx \leq \int_{\mathbb{R}^d} \psi(\mu_{\text{ext}}) = \int_{\mathcal{O}} \psi(\mu), \end{aligned}$$

which finishes the proof. \square

Corollary 3.25. *Together with Remark 3.4, Proposition 3.24 immediately implies*

$$\limsup_{\varepsilon \rightarrow 0} \int_{\mathcal{O}} \psi(\mu_{\varepsilon, \delta_\varepsilon}) \leq \int_{\mathcal{O}} \psi(\mu)$$

as long as $0 < \delta_\varepsilon \leq \frac{w(\varepsilon)}{2}$.

Proof of Theorem 3.8. For μ as in Theorem 3.8, we show that the sequence

$$(\mu_n)_{n \in \mathbb{N}} := \left(\mu_{\frac{1}{n}, \frac{1}{2}w(\frac{1}{n})} \right)_{n \in \mathbb{N}},$$

where w was defined in Remark 3.12, meets all requirements.

By construction, $\mu_n \in \mathcal{M}(\mathcal{O}) \cap H^{-1}$ for all $n \in \mathbb{N}$, and by Lemma 3.19, the density of μ_n is bounded and thus in $L^2(\mathcal{O})$. Property (3.8) is proved in the first part of Proposition 3.20. For Property (3.9), note that Corollary 3.25 especially shows that $(\mu_n)_{n \in \mathbb{N}}$ is uniformly bounded in the TV norm, which means that it contains a subsequence that converges weakly* to $\psi(\mu)$ by Proposition 3.20 and [23, Lemma 2.1]. Since this argument can be carried out for any subsequence, we get weak* convergence for the whole sequence and, also by [23, Lemma 2.1],

$$\left\| \psi \left(\mu_{\frac{1}{n}, \frac{1}{2}w(\frac{1}{n})} \right) \right\|_{TV} := \int_{\mathcal{O}} \psi \left(\mu_{\frac{1}{n}, \frac{1}{2}w(\frac{1}{n})} \right) \rightarrow \int_{\mathcal{O}} \psi(\mu) := \|\psi(\mu)\|_{TV} \quad \text{as } n \rightarrow \infty.$$

This yields (3.9) and thereby concludes the proof. \square

4 Proof of the main result

Throughout this section, we work under Assumptions 2.1. We mostly sketch the argument, which is closely along the lines of [37, Proof of Theorem 2.3], and only give more details for the parts where additional results are needed due to the different nonlinearity.

We consider the SPDE

$$\begin{aligned} dX_t^\varepsilon &= \varepsilon \Delta X_t^\varepsilon dt + \Delta \phi^\varepsilon(X_t^\varepsilon) dt + B(t, X_t^\varepsilon) dW_t, \\ X_0^\varepsilon &= x_0, \end{aligned} \tag{4.1}$$

where we use the notation for the Yosida approximation of [37, Appendix C] and assume $x_0 \in L^2(\Omega, \mathcal{F}_0; L^2)$. Now and in the following we omit the domain \mathcal{O} when using Lebesgue and Sobolev spaces.

Lemma 4.1. *For all $T > 0$, Problem (4.1) gives rise to a solution in sense of [53, Definition 4.2.1] with respect to the Gelfand triple $V := L^2 \hookrightarrow H^{-1} \hookrightarrow (L^2)' = V'$.*

Proof. One shows that (4.1) fits into the framework of [53, Example 4.1.11] with the operator

$$A(u) = \Delta(\varepsilon u + \phi^\varepsilon(u)) \quad \text{for } u \in L^2.$$

The statement then follows by [53, Theorem 4.2.4]. \square

The following lemma provides an important estimate on the regularity of these approximate solutions and corresponds to [37, Lemma B.1]:

Lemma 4.2. *Let $\varepsilon > 0$, $x_0 \in L^2(\Omega, \mathcal{F}_0; L^2)$ and $T > 0$. Then for the solution $(X_t^\varepsilon)_{t \in [0, T]}$ to (4.1) we have*

$$\mathbb{E} \sup_{t \in [0, T]} \|X_t^\varepsilon\|_2^2 + \varepsilon \mathbb{E} \int_0^T \|X_r^\varepsilon\|_{H_0^1}^2 dr \leq C(\mathbb{E} \|x_0\|_2^2 + 1)$$

with a constant $C > 0$ independent of ε .

Proof of Lemma 4.2. Let $(e_i)_{i \in \mathbb{N}} \subset \mathcal{C}_0^2$ be an orthonormal basis in H^{-1} of smooth eigenvectors to $-\Delta$, and let $P^n : H^{-1} \rightarrow H_n := \text{span}\{e_1, \dots, e_n\}$ be the orthogonal projection onto the span of the first n eigenvectors. Recall that the unique variational solution X^ε to (4.1) is constructed in [53] as a (weak) limit in $L^2([0, T] \times \Omega; L^2)$ of the solutions to the Galerkin approximation

$$\begin{aligned} dX_t^n &= \varepsilon P^n \Delta X_t^n dt + P^n \Delta \phi^\varepsilon(X_t^n) dt + P^n B(t, X_t^n) dW_t^n \\ X_0^n &= P^n x_0, \end{aligned}$$

in H_n , where for simplicity we omit the ε -dependence of X^n , and for an orthonormal basis $(g_i)_{i \in \mathbb{N}}$ of U (as defined in Assumption 2.1 (A1)) we let

$$W_t^n = \sum_{i=1}^n \langle J^{-1}(W_t), g_i \rangle_U g_i.$$

Using the finite-dimensional Ito formula and the Burkholder-Davis-Gundy inequality, one shows the energy estimate

$$\mathbb{E} \sup_{r \in [0, T]} \|X_r^n\|_{L^2}^2 + \varepsilon \mathbb{E} \int_0^T \|X_r^n\|_{H_0^1}^2 dr \leq C(\mathbb{E} \|x_0\|_{L^2}^2 + 1).$$

Thus, $(X^n)_{n \in \mathbb{N}}$ is bounded in $L^2(\Omega; L^\infty([0, T]; L^2))$ and in $L^2(\Omega \times [0, T]; H_0^1)$. The latter is a Hilbert space, thus we can extract a weakly converging subsequence whose limit can be identified with the unique weak $L^2(\Omega \times [0, T]; L^2)$ limit X^ε . The former is the dual space of $L^2(\Omega; L^1([0, T]; L^2))$ which is separable. Thus, we can extract a weak* converging subsequence whose limit can again be identified with X^ε . By weak (respectively weak*) lower-semicontinuity of the norms, we can thus pass to the limit $n \rightarrow \infty$ to obtain the required inequality. \square

Proof of Theorem 2.6. Existence: Let $(x_0^n)_{n \in \mathbb{N}} \subset L^2(\Omega, \mathcal{F}_0; L^2)$ such that $x_0^n \rightarrow x_0$ in $L^2(\Omega; H^{-1})$, and let $X^{\varepsilon_1, n}, X^{\varepsilon_2, n}$ be the solutions to (4.1) with initial state x_0^n for $\varepsilon_1, \varepsilon_2 > 0$. By the Ito formula on $e^{-Kt} \|X_t^{\varepsilon_1, n} - X_t^{\varepsilon_2, m}\|_{H^{-1}}^2$, the Burkholder-Davis-Gundy inequality and [37, Equation (C.5)], we obtain

$$\begin{aligned} \mathbb{E} \sup_{t \in [0, T]} \left(e^{-Kt} \|X_t^{\varepsilon_1, n} - X_t^{\varepsilon_2, m}\|_{H^{-1}}^2 \right) &\leq 2 \mathbb{E} \|x_0^n - x_0^m\|_{H^{-1}}^2 \\ &+ C(\varepsilon_1 + \varepsilon_2) \left(\mathbb{E} \|x_0^n\|_{L^2}^2 + \mathbb{E} \|x_0^m\|_{L^2}^2 + 1 \right) \end{aligned} \quad (4.2)$$

for $K > 0$ large enough. Letting first $\varepsilon \rightarrow 0$ and then $n \rightarrow \infty$ yields a limit $X \in L^2(\Omega; \mathcal{C}([0, T]; H^{-1}))$ by completeness, which will be shown to be an SVI solution. To this end, define with the notation of [37, Appendix C]

$$\varphi^\varepsilon(v) = \begin{cases} \int_{\mathcal{O}} \psi^\varepsilon(v) dx, & v \in L^2, \\ +\infty, & \text{otherwise,} \end{cases} \quad (4.3)$$

for $v \in H^{-1}$. Using the Ito formula on $e^{-Kt} \|X_t^{\varepsilon, n}\|_{H^{-1}}^2$ and the fact that $-\Delta \phi^\varepsilon(x) \in \partial \varphi^\varepsilon(x)$ for $x \in H_0^1$, one obtains

$$\mathbb{E} \int_0^t \varphi^\varepsilon(X_r^{\varepsilon, n}) dr \leq C + \mathbb{E} \|x_0^n\|_{H^{-1}}^2 \leq \tilde{C} \quad (4.4)$$

for some $C, \tilde{C} > 0$ independent of ε and n . Together with Assumption 2.1 (A4), which allows to use [37, Equation (C.4)], and the lower-semicontinuity of φ from Proposition 3.7, one obtains part (i) of Definition 2.4.

For the variational inequality part, let G, Z, t be as in Definition 2.4 (ii). Using Ito's formula on $\|X_t^{\varepsilon, n} - Z_t\|_{H^{-1}}^2$, (2.2) and the weighted Young inequality we obtain

$$\begin{aligned} &\mathbb{E} \|X_t^{\varepsilon, n} - Z_t\|_{H^{-1}} + 2 \mathbb{E} \int_0^t \varphi^\varepsilon(X_r^{\varepsilon, n}) dr \\ &\leq \mathbb{E} \|x_0^n - Z_0\|_{H^{-1}}^2 + 2 \mathbb{E} \int_0^t \varphi^\varepsilon(Z_r) dr \\ &\quad - 2 \mathbb{E} \int_0^t \langle G_r, X_r^{\varepsilon, n} - Z_r \rangle_{H^{-1}} dr + C \mathbb{E} \int_0^t \|X_r^{\varepsilon, n} - Z_r\|_{H^{-1}}^2 dr \\ &\quad + 2 \mathbb{E} \int_0^t \frac{1}{2} \varepsilon^{\frac{4}{3}} \|\Delta X_r^{\varepsilon, n}\|_{H^{-1}}^2 + \frac{1}{2} \varepsilon^{\frac{2}{3}} \|X_r^{\varepsilon, n} - Z_r\|_{H^{-1}}^2 dr. \end{aligned} \quad (4.5)$$

Then using [37, Equations (C.3) and (C.4)], the lower-semicontinuity of φ from Proposition 3.7 and Lemma 4.2, we can take first $\liminf_{\varepsilon \rightarrow 0}$ and then $\liminf_{n \rightarrow \infty}$ to obtain (2.12) and therefore the remaining part (ii) of Definition 2.4.

Uniqueness: It remains to show that the solution constructed in the previous step is unique. To this end, let $x_0, y_0 \in L^2(\Omega, \mathcal{F}_0; H^{-1})$, $(y_0^n)_{n \in \mathbb{N}} \subset L^2(\Omega, \mathcal{F}_0; L^2)$ satisfying $y_0^n \rightarrow y_0$ in $L^2(\Omega; H^{-1})$ for $n \rightarrow \infty$. Let X be an arbitrary SVI solution to (1.1) with initial condition x_0 and let $(Y^{\varepsilon, n})_{\varepsilon > 0, n \in \mathbb{N}}$ be the solutions to (4.1) with respective initial conditions $(y_0^n)_{n \in \mathbb{N}}$. One can check that

$$Z = Y^{\varepsilon, n} \quad \text{and} \quad G = \varepsilon \Delta Y^{\varepsilon, n} + \Delta \phi^\varepsilon(Y^{\varepsilon, n}) \quad (4.6)$$

are admissible choices for (2.12). Then, (2.12) yields for $t \in [0, T]$

$$\begin{aligned} \mathbb{E} \|X_t - Y_t^{\varepsilon, n}\|_{H^{-1}}^2 + 2\mathbb{E} \int_0^t \varphi(X_r) dr &\leq \mathbb{E} \|x_0 - y_0^n\|_{H^{-1}}^2 + 2\mathbb{E} \int_0^t \varphi(Y_r^{\varepsilon, n}) dr \\ &\quad - 2\mathbb{E} \int_0^t \langle \varepsilon \Delta Y_r^{\varepsilon, n} + \Delta \phi^\varepsilon(Y_r^{\varepsilon, n}), X_r - Y_r^{\varepsilon, n} \rangle_{H^{-1}} dr \\ &\quad + C\mathbb{E} \int_0^t \|X_r - Y_r^{\varepsilon, n}\|_{H^{-1}}^2 dr. \end{aligned} \quad (4.7)$$

For $u \in L^2$ and φ^ε as in (4.3), we have

$$\langle -\Delta \phi^\varepsilon(Y^{\varepsilon, n}), u - Y^{\varepsilon, n} \rangle_{H^{-1}} + \varphi^\varepsilon(Y^{\varepsilon, n}) \leq \varphi^\varepsilon(u) \quad dt \otimes \mathbb{P}\text{-a. e.} \quad (4.8)$$

Since $Y^{\varepsilon, n} \in H_0^1 \subset L^2$ $dt \otimes \mathbb{P}$ -a. e. we can use [37, Equation (C.4)] to obtain $dt \otimes \mathbb{P}$ -almost everywhere

$$|\varphi^\varepsilon(Y^{\varepsilon, n}) - \varphi(Y^{\varepsilon, n})| \leq C\varepsilon \left(1 + \|Y^{\varepsilon, n}\|_{L^2}^2\right). \quad (4.9)$$

Using 4.9 and [37, Equation (C.3)], we can modify (4.8) to obtain

$$\langle -\Delta \phi^\varepsilon(Y^{\varepsilon, n}), u - Y^{\varepsilon, n} \rangle_{H^{-1}} + \varphi(Y^{\varepsilon, n}) \leq \varphi(u) + C\varepsilon \left(1 + \|Y^{\varepsilon, n}\|_{L^2}^2\right) \quad dt \otimes \mathbb{P}\text{-a. e.} \quad (4.10)$$

Note that (4.10) is trivial if $\varphi(u) = \infty$. Furthermore, (4.10) can be deduced analogously for $u \in L^{m+1} \cap H^{-1}$ in the superlinear setting, i. e. when φ is given by (2.8), with m as in Assumption 2.1 (A5). In the sublinear setting, i. e. φ is given by (2.9), and $u \in \mathcal{M}(\mathcal{O}) \cap H^{-1}$, we consider the approximating sequence $(u_j)_{j \in \mathbb{N}} \subset L^2$ given by Theorem 3.8, such that (4.10) is satisfied for all $u_j, j \in \mathbb{N}$. We then pass to the limit $j \rightarrow \infty$ and notice that by the construction of $(u_j)_{j \in \mathbb{N}}$ we have both $\varphi(u_j) \rightarrow \varphi(u)$ and

$$\begin{aligned} &\langle -\Delta \phi^\varepsilon(Y^{\varepsilon, n}), u_j - Y^{\varepsilon, n} \rangle_{H^{-1}} \\ &= {}_{H_0^1} \langle \phi^\varepsilon(Y^{\varepsilon, n}), u_j - Y^{\varepsilon, n} \rangle_{H^{-1}} \longrightarrow {}_{H_0^1} \langle \phi^\varepsilon(Y^{\varepsilon, n}), u - Y^{\varepsilon, n} \rangle_{H^{-1}} \\ &= \langle -\Delta \phi^\varepsilon(Y^{\varepsilon, n}), u - Y^{\varepsilon, n} \rangle_{H^{-1}}. \end{aligned}$$

Consequently, replacing u by X in (4.10), we have in any case

$$\langle -\Delta \phi^\varepsilon(Y^{\varepsilon, n}), X - Y^{\varepsilon, n} \rangle_{H^{-1}} + \varphi(Y^{\varepsilon, n}) \leq \varphi(X) + C\varepsilon \left(1 + \|Y^{\varepsilon, n}\|_{L^2}^2\right) \quad dt \otimes \mathbb{P}\text{-a. e.} \quad (4.11)$$

Using (4.11) and the weighted Young inequality, we can modify (4.7) to obtain for $t \in [0, T]$

$$\begin{aligned} \mathbb{E} \|X_t - Y_t^{\varepsilon, n}\|_{H^{-1}}^2 &\leq \mathbb{E} \|x_0 - y_0^n\|_{H^{-1}}^2 \\ &\quad + 2\mathbb{E} \int_0^t \varepsilon^{\frac{4}{3}} \|\Delta Y_r^{\varepsilon, n}\|_{H^{-1}}^2 dr + \varepsilon^{\frac{2}{3}} \|X_r - Y_r^{\varepsilon, n}\|_{H^{-1}}^2 dr \\ &\quad + C\mathbb{E} \int_0^t \|X_r - Y_r^{\varepsilon, n}\|_{H^{-1}}^2 dr + C\varepsilon \mathbb{E} \int_0^t \left(1 + \|Y_t^{\varepsilon, n}\|_{L^2}^2\right) dr. \end{aligned}$$

Taking $\varepsilon \rightarrow 0$ and then $n \rightarrow \infty$ yields

$$\mathbb{E} \|X_t - Y_t\|_{H^{-1}}^2 \leq \mathbb{E} \|x_0 - y_0\|_{H^{-1}}^2 + C\mathbb{E} \int_0^t \|X_r - Y_r\|_{H^{-1}}^2 dr \quad \text{for } t \in [0, T]. \quad (4.12)$$

where Y is the SVI solution constructed from $(Y^{\varepsilon, n})$ by the limiting procedure at the beginning of this proof. Gronwall's inequality then yields $X = Y$ if $x_0 = y_0$, and thus uniqueness of SVI solutions. Then, estimate (2.13) follows by applying Gronwall's inequality to (4.12) with different initial values, which concludes the proof. \square

A Generalities on convex functions

We collect and prove some statements on convex functions defined on \mathbb{R} .

Lemma A.1. *Let $f : \mathbb{R} \rightarrow [0, \infty)$ be convex with $f(0) = 0$ and $x, y \in \mathbb{R} \setminus \{0\}$ with $x < y$. Then*

$$\frac{f(x)}{x} \leq \frac{f(y)}{y}. \quad (\text{A.1})$$

In particular, for $x > 0$ this implies $f(x) \leq f(y)$.

Proof. Note that by convexity, we have for $\lambda \in (0, 1)$, $x \in \mathbb{R}$

$$f(\lambda x) = f(\lambda x + (1 - \lambda)0) \leq \lambda f(x) + (1 - \lambda)f(0) = \lambda f(x). \quad (\text{A.2})$$

If $x < 0 < y$, the statement is obvious by the nonnegativity of f . If $0 < x < y$, we use (A.2) with $\lambda = \frac{x}{y}$ to get

$$\frac{f(x)}{x} = \frac{f(\lambda y)}{\lambda y} \leq \frac{\lambda f(y)}{\lambda y} = \frac{f(y)}{y},$$

while for $x < y < 0$ we use (A.2) with $\lambda := \frac{y}{x}$ to get

$$\frac{f(y)}{y} = \frac{f(\lambda x)}{\lambda x} \geq \frac{\lambda f(x)}{\lambda x} = \frac{f(x)}{x},$$

as required. □

Lemma A.2. *Let ψ satisfy Assumptions 2.1 and $y > 0$. Then, if $\psi(y) > 0$, we have*

$$\psi^*(-x) = \psi^*(x) \leq \psi(y) \quad \text{for } x \in \left[0, \frac{\psi(y)}{y}\right],$$

where ψ^ is defined as in Definition 3.1.*

Proof. By Remark 3.2, the last part of Lemma A.1 and the nonnegativity of ψ^* , it is enough to show

$$\psi^*\left(\frac{\psi(y)}{y}\right) \leq \psi(y). \quad (\text{A.3})$$

To verify (A.3), we distinguish three cases for $y' \in \mathbb{R}$. For $y' \geq y$ we have by Lemma A.1

$$\frac{\psi(y)}{y}y' - \psi(y') = y' \left(\frac{\psi(y)}{y} - \frac{\psi(y')}{y'} \right) \leq 0,$$

for $y' \leq 0$ we have by the nonnegativity of ψ

$$\frac{\psi(y)}{y}y' - \psi(y') \leq 0,$$

and for $y' \in (0, y)$ we have

$$\frac{\psi(y)}{y}y' - \psi(y') \leq \frac{\psi(y)}{y}y = \psi(y),$$

which yields the claim. □

Lemma A.3. *Let ψ satisfy Assumptions 2.1. For $K = \text{dom}(\psi^*) := \{x \in \mathbb{R} : \psi^*(x) < \infty\}$ we have*

$$\sup K = \lim_{t \rightarrow \infty} \frac{\psi(t)}{t} \quad \text{and} \quad \sup(-K) = \lim_{t \rightarrow \infty} \frac{\psi(-t)}{t}.$$

Proof. We only prove the first statement, the second then becomes clear by symmetry. To this end, note first that the limit is actually a supremum, as $\frac{\psi(t)}{t}$ is increasing (by (A.1)). Let now $x \in K$, which means that $xt - \psi(t) \leq c_x < \infty$ and thus $\frac{\psi(t)}{t} \geq x - \frac{c_x}{t}$ for all $t \in [0, \infty)$, which yields “ \leq ” by letting $t \rightarrow \infty$.

Conversely, we have $\frac{\psi(t)}{t} \in K$ for $t > 0$, $\psi(t) > 0$ by Lemma A.2. As $\psi^*(0) = 0$, this is true also if $\psi(t) = 0$, thereby proving “ \geq ”. □

Corollary A.4. *Let ψ satisfy Assumptions 2.1. By Lemmas A.2 and A.3, we have that*

$$\psi_\infty(1) = \psi_\infty(-1) \geq \frac{\psi(y)}{y}$$

for $y > 0$ with $\psi(y) > 0$.

Lemma A.5. *Let ψ satisfy Assumptions 2.1. For the convex conjugate of the recession function, we have*

$$\psi_\infty^*(x) := (\psi_\infty)^*(x) = \chi_{[-\psi_\infty(1), \psi_\infty(1)]}(x)$$

for $x \in \mathbb{R}$, where for an Interval I we have written

$$\chi_I(x) = \begin{cases} 0, & \text{if } x \in I \\ +\infty, & \text{else.} \end{cases}$$

Proof. In the superlinear case, i. e. (2.6) is satisfied, we have $\psi_\infty = \chi_{\{0\}}$ and thus $\psi_\infty^* \equiv 0$, as required. In the sublinear case, we first note that ψ_∞ is, by definition, positively homogeneous, which by symmetry amounts to absolute homogeneity. Thus

$$\psi_\infty(x) = \psi_\infty(1) |x|,$$

where $\psi_\infty(1) > 0$ by Corollary A.4, which allows to conclude by the definition of the convex conjugate. \square

References

- [1] H. Alt. *Lineare Funktionalanalysis: Eine anwendungsorientierte Einführung*. Masterclass. Springer Berlin Heidelberg, 2012.
- [2] G. Anzellotti. The Euler equation for functionals with linear growth. *Trans. Amer. Math. Soc.*, 290(2):483–501, 1985.
- [3] P. Bak, C. Tang, and K. Wiesenfeld. Self-organized criticality: An explanation of the 1/f noise. *Phys. Rev. Lett.*, 59:381–384, 1987.
- [4] P. Bak, C. Tang, and K. Wiesenfeld. Self-organized criticality. *Phys. Rev. A*, 38:364–374, 1988.
- [5] V. Barbu. *Nonlinear Differential Equations of Monotone Types in Banach Spaces*. Springer New York, 2010.
- [6] V. Barbu, V. I. Bogachev, G. D. Prato, and M. Röckner. Weak solutions to the stochastic porous media equation via Kolmogorov equations: The degenerate case. *Journal of Functional Analysis*, 237(1):54–75, 2006.
- [7] V. Barbu, G. Da Prato, and M. Röckner. Stochastic nonlinear diffusion equations with singular diffusivity. *SIAM Journal on Mathematical Analysis*, 41(3):1106–1120, 2009.
- [8] V. Barbu, G. D. Prato, and M. Röckner. Stochastic porous media equations and self-organized criticality. *Communications in Mathematical Physics*, 285(3):901–923, Feb 2009.
- [9] V. Barbu, G. D. Prato, and M. Röckner. Existence and uniqueness of nonnegative solutions to the stochastic porous media equation. *Indiana University Mathematics Journal*, 57(1):187–211, 2008.
- [10] V. Barbu, G. D. Prato, and M. Röckner. Existence of strong solutions for stochastic porous media equation under general monotonicity conditions. *The Annals of Probability*, 37(2):428–452, 2009.
- [11] V. Barbu, G. D. Prato, and M. Röckner. Finite time extinction of solutions to fast diffusion equations driven by linear multiplicative noise. *Journal of Mathematical Analysis and Applications*, 389(1):147–164, 2012.
- [12] V. Barbu and M. Röckner. An operatorial approach to stochastic partial differential equations driven by linear multiplicative noise. *J. Eur. Math. Soc. (JEMS)*, 17(7):1789–1815, 2015.

-
- [13] V. Barbu and M. Röckner. Stochastic variational inequalities and applications to the total variation flow perturbed by linear multiplicative noise. *Archive for Rational Mechanics and Analysis*, 209(3):797–834, Sep 2013.
- [14] V. Barbu, M. Röckner, and F. Russo. Stochastic porous media equations in \mathbb{R}^d . *Journal de Mathématiques Pures et Appliquées*, 103(4):1024–1052, 2015.
- [15] H. Bauschke and P. Combettes. *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*. CMS Books in Mathematics. Springer International Publishing, 2017.
- [16] C. Bauzet, G. Vallet, and P. Wittbold. A degenerate parabolic-hyperbolic Cauchy problem with a stochastic force. *J. Hyperbolic Differ. Equ.*, 12(3):501–533, 2015.
- [17] I. H. Biswas and A. K. Majee. Stochastic conservation laws: weak-in-time formulation and strong entropy condition. *J. Funct. Anal.*, 267(7):2199–2252, 2014.
- [18] G. Da Prato and M. Röckner. Weak solutions to stochastic porous media equations. *Journal of Evolution Equations*, 4(2):249–271, May 2004.
- [19] K. Dareiotis, M. Gerencsér, and B. Gess. Entropy solutions for stochastic porous media equations. *J. Differential Equations*, 266(6):3732–3763, 2019.
- [20] K. Dareiotis, B. Gess, and P. Tsatsoulis. Ergodicity for Stochastic Porous Media Equations. *arXiv e-prints*, page arXiv:1907.04605, Jul 2019.
- [21] A. Debussche, M. Hofmanová, and J. Vovelle. Degenerate parabolic stochastic partial differential equations: quasilinear case. *Ann. Probab.*, 44(3):1916–1955, 2016.
- [22] A. Debussche and J. Vovelle. Scalar conservation laws with stochastic forcing. *J. Funct. Anal.*, 259(4):1014–1042, 2010.
- [23] F. Demengel and R. Temam. Convex functions of a measure and applications. *Indiana Univ. Math. J.*, 33:673 – 709, 1984.
- [24] A. Díaz-Guilera. Noise and dynamics of self-organized critical phenomena. *Phys. Rev. A*, 45:8551–8558, Jun 1992.
- [25] A. Díaz-Guilera. Dynamic renormalization group approach to self-organized critical phenomena. *EPL (Europhysics Letters)*, 26:177 – 182, 1994.
- [26] J. Feng and D. Nualart. Stochastic scalar conservation laws. *J. Funct. Anal.*, 255(2):313–373, 2008.
- [27] F. Ferro. Integral characterization of functionals defined on spaces of BV functions. *Rend. Sem. Mat. Univ. Padova*, 61:177–201 (1980), 1979.
- [28] F. Flandoli, B. Gess, and M. Scheutzow. Synchronization by noise. *Probability Theory and Related Fields*, 168(3):511–556, Aug 2017.
- [29] I. Fonseca and G. Leoni. *Modern Methods in the Calculus of Variations: L^p Spaces*. Springer Monographs in Mathematics. Springer New York, 2007.
- [30] P. K. Friz and B. Gess. Stochastic scalar conservation laws driven by rough paths. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 33(4):933–963, 2016.
- [31] P. Gassiat and B. Gess. Regularization by noise for stochastic Hamilton-Jacobi equations. *Probab. Theory Related Fields*, 173(3-4):1063–1098, 2019.
- [32] B. Gess. Strong solutions for stochastic partial differential equations of gradient type. *J. Funct. Anal.*, 263(8):2355–2383, 2012.
- [33] B. Gess. Random attractors for singular stochastic evolution equations. *Journal of Differential Equations*, 255(3):524 – 559, 2013.
- [34] B. Gess. Finite time extinction for stochastic sign fast diffusion and self-organized criticality. *Communications in Mathematical Physics*, 335(1):309–344, Apr 2015.

-
- [35] B. Gess and M. Hofmanová. Well-posedness and regularity for quasilinear degenerate parabolic-hyperbolic SPDE. *Ann. Probab.*, 46(5):2495–2544, 2018.
- [36] B. Gess and M. Röckner. Stochastic variational inequalities and regularity for degenerate stochastic partial differential equations. *Trans. Amer. Math. Soc.*, 369(5):3017–3045, 2017.
- [37] B. Gess and M. Röckner. Singular-degenerate multivalued stochastic fast diffusion equations. *SIAM J. Math. Anal.*, 47:4059 – 4090, 2015.
- [38] B. Gess and P. E. Souganidis. Stochastic non-isotropic degenerate parabolic-hyperbolic equations. *Stochastic Process. Appl.*, 127(9):2961–3004, 2017.
- [39] B. Gess and J. M. Tölle. Stability of solutions to stochastic partial differential equations. *J. Differential Equations*, 260(6):4973–5025, 2016.
- [40] B. Gess and J. Tölle. Ergodicity and local limits for stochastic local and nonlocal p -laplace equations. *SIAM Journal on Mathematical Analysis*, 48(6):4094–4125, 2016.
- [41] B. Gess and J. M. Tölle. Multi-valued, singular stochastic evolution inclusions. *Journal de Mathématiques Pures et Appliquées*, 101(6):789 – 827, 2014.
- [42] B. Gutenberg and C. F. Richter. *Seismicity Of The Earth And Associated Phenomena*. Princeton University Press, 1949.
- [43] I. Gyöngy and E. Pardoux. On the regularization effect of space-time white noise on quasi-linear parabolic partial differential equations. *Probab. Theory Related Fields*, 97(1-2):211–229, 1993.
- [44] J. U. Kim. On a stochastic scalar conservation law. *Indiana Univ. Math. J.*, 52(1):227–256, 2003.
- [45] N. Krylov and B. Rozovskii. Stochastic evolution equations. *J. Soviet Math.*, 16 (Transl.):1233 – 1277, 1981.
- [46] P.-L. Lions, B. Perthame, and P. E. Souganidis. Scalar conservation laws with rough (stochastic) fluxes. *Stoch. Partial Differ. Equ. Anal. Comput.*, 1(4):664–686, 2013.
- [47] B. Mandelbrot. The variation of certain speculative prices. *The Journal of Business*, 36, 1963.
- [48] M. Neuß. Ergodicity for singular-degenerate porous media equations. *arXiv e-prints*, page arXiv:1909.05161, Sep 2019.
- [49] D. A. Noever. Himalayan sandpiles. *Phys. Rev. E*, 47:724–725, 1993.
- [50] F. Otto. The geometry of dissipative evolution equations: the porous medium equation. *Comm. Partial Differential Equations*, 26(1-2):101–174, 2001.
- [51] E. Pardoux. Équations aux dérivées partielles stochastiques de type monotone. In *Séminaire sur les Équations aux Dérivées Partielles (1974–1975), III, Exp. No. 2*, page 10. Collège de France Paris, 1975.
- [52] C. J. Pérez, Á. Corral, A. Díaz-Guilera, K. Christensen, and A. Arenas. On Self-Organized Criticality and Synchronization in Lattice Models of Coupled Dynamical Systems. *International Journal of Modern Physics B*, 10:1111–1151, 1996.
- [53] C. Prévot and M. Röckner. *A Concise Course on Stochastic Partial Differential Equations*. Springer Berlin/Heidelberg, 2007.
- [54] J. Ren, M. Röckner, and F.-Y. Wang. Stochastic generalized porous media and fast diffusion equations. *Journal of Differential Equations*, 238(1):118–152, 2007.
- [55] R. Temam. *Mathematical problems in plasticity*. Gauthier-Villars, 1985.
- [56] J. Vazquez. *The Porous Medium Equation: Mathematical Theory*. Oxford Mathematical Monographs. Clarendon Press, 2006.
- [57] N. W. Watkins, G. Pruessner, S. C. Chapman, N. B. Crosby, and H. J. Jensen. 25 years of self-organized criticality: Concepts and controversies. *Space Science Reviews*, 198(1):3–44, Jan 2016.

MARIUS NEUSS
MAX-PLANCK-INSTITUT FÜR MATHEMATIK IN DEN NATURWISSENSCHAFTEN
04103 LEIPZIG, GERMANY
marius.neuss@mis.mpg.de