

GROUP-VALUED MOMENTUM MAPS FOR ACTIONS OF AUTOMORPHISM GROUPS

Tobias Diez^a and Tudor S. Ratiu^b

February 5, 2020

The space of smooth sections of a symplectic fiber bundle carries a natural symplectic structure. We provide a general framework to determine the momentum map for the action of the group of bundle automorphism on this space. Since, in general, this action does not admit a classical momentum map, we introduce the more general class of group-valued momentum maps which is inspired by the Poisson Lie setting. In this approach, the group-valued momentum map assigns to every section of the symplectic fiber bundle a principal circle-bundle with connection. The power of this general framework is illustrated in many examples: we construct generalized Clebsch variables for fluids with integral helicity; the anti-canonical bundle turns out to be the momentum map for the action of the group of symplectomorphisms on the space of compatible complex structures; the Teichmüller moduli space is realized as a symplectic orbit reduced space associated to a coadjoint orbit of $SL(2, \mathbb{R})$ and spaces related to the other coadjoint orbits are identified and studied. Moreover, we show that the momentum map for the group of bundle automorphisms on the space of connections over a Riemann surface encodes, besides the curvature, also topological information of the bundle.

KEYWORDS: symplectic structure on moduli spaces of geometric structures, infinite-dimensional symplectic geometry, momentum maps, symplectic fiber bundles, differential characters, Kähler geometry, Teichmüller space

MSC 2010: 53D20, (58D27, 53C08, 53C10, 58B99, 32G15)

^a Max Planck Institute for Mathematics in the Sciences, 04103 Leipzig, Germany and Institut für Theoretische Physik, Universität Leipzig, 04009 Leipzig, Germany and Institute of Applied Mathematics, Delft University of Technology, 2628 XE Delft, Netherlands. Partially supported by the Max Planck Institute for Mathematics in the Sciences (Leipzig), by the German National Academic Foundation (DAAD) and by the above NWO grant 639.032.734. tobias.diez@mis.mpg.de

^b School of Mathematical Sciences and Ministry of Education Laboratory of Scientific Computing (MOE-LSC), Shanghai Jiao Tong University, Minhang District, 800 Dongchuan Road, 200240 China, Section de Mathématiques, Université de Genève, 2–4 rue du Lièvre, case postale 64, 1211 Genève 4, and Ecole Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland. Partially supported by the National Natural Science Foundation of China (No. 11871334) and by NCCR SwissMAP grant of the Swiss National Science Foundation. ratiu@sjtu.edu.cn, tudor.ratiu@epfl.ch

CONTENTS

1	Introduction	2	
2	Group-Valued Momentum Maps	8	
2.1	Poisson momentum maps	8	
2.2	Dual pairs of Lie algebras	10	
2.3	Group-valued momentum maps	13	
2.4	Existence and uniqueness	16	
2.5	Equivariance and Poisson property	19	
2.6	Momentum maps for group extensions	35	
3	Global Analysis of Symplectic Fiber Bundles	38	
3.1	Symplectic form	38	
3.2	Action of the automorphism group	39	
3.2.1	Momentum map for the gauge group	41	
3.2.2	Momentum map for the diffeomorphism group	44	
3.2.3	Momentum map for volume-preserving diffeomorphisms	44	
3.2.4	Momentum map for symplectomorphisms	49	
4	Applications	52	
4.1	Trivial bundle case	52	
4.1.1	Hydrodynamics	53	
4.1.2	Lagrangian embeddings	59	
4.2	Reduction of structure group	60	
4.2.1	Pull-back of prequantum bundles as associated bundles	61	
4.2.2	Kähler geometry	64	
4.2.3	Teichmüller space and weighted Lagrangian subbundles	70	
4.3	Action of the quantomorphism group	77	
4.4	Gauge theory	79	
	Appendices	87	
A	Notations and conventions	87	
B	Fiber integration	90	
C	Hat product for fiber bundles	92	
D	Cheeger-Simons differential characters	95	
D.1	Motivation: circle bundles	96	
D.2	Differential characters	97	
D.3	Hat product of differential characters	102	

1 INTRODUCTION

Noether's theorem states that every symmetry of a given system has a corresponding conservation law. In terms of symplectic geometry, these conserved quantities are encoded in the momentum map. It was, however, quickly realized

that the momentum map geometry not only plays an important role in dynamical systems but is also a valuable tool in the study of differential geometric questions. In their seminal work, Atiyah and Bott [AB83] showed that the curvature of a connection on a principal bundle over a Riemannian surface is the momentum map for the action of the group of gauge transformations. They applied Morse theory to the norm-squared of the momentum map (the Yang–Mills functional) in order to obtain the cohomology of the moduli space of Yang–Mills solutions which, by the Narasimhan–Seshadri theorem, can be identified with the moduli space of stable holomorphic structures. Within the same circle of ideas, Fujiki [Fuj92] and Donaldson [Don97; Don00; Don03] provided a momentum map picture for the relationship between the existence of constant scalar curvature Kähler metrics and stability in the sense of geometric invariant theory.

The first main aim of this paper is to provide a framework which encompasses the gauge theory setting of Atiyah and Bott and, at the same time, the action of diffeomorphism groups of Fujiki and Donaldson. Our starting point is a symplectic fiber bundle of the form $F = P \times_G \underline{F}$ for a principal G -bundle $P \rightarrow M$, where the typical fiber \underline{F} is endowed with a G -invariant symplectic form. The fiberwise symplectic structure, combined with a volume form on the base M , induces a symplectic form Ω on the space \mathcal{F} of sections of $F \rightarrow M$. The gauge group of P acts in a natural way on \mathcal{F} , leaving the induced symplectic form Ω invariant. As we will see, this action possesses a momentum map which is completely determined by the momentum map of the G -action on the fiber \underline{F} . Suppose the bundle P is natural, i.e., it comes with a lift of $\text{Diff}(M)$ to bundle automorphisms (for example, this is the case when P is the frame bundle of M). In this case, every subgroup of the group of volume-preserving diffeomorphisms acts naturally on the space \mathcal{F} of sections and leaves Ω invariant. Our first result is summarized in Theorems 3.3 and 3.7 which determine the momentum map for the group of volume-preserving diffeomorphisms and symplectomorphisms, respectively. There are, essentially, two contributions to the momentum map. The first term is the pull-back of the fiberwise symplectic structure. The second term involves the fiber momentum map and, morally speaking, captures how much the lift of diffeomorphisms to bundle automorphisms shifts in the vertical direction. The interesting point is that the momentum map for the automorphism group on the *infinite-dimensional* space of sections is canonically constructed from the *finite-dimensional* symplectic G -manifold \underline{F} .

In contrast to the case of the action of the gauge group, the momentum map for the symplectic action of the diffeomorphism group on the space of sections does not exist in full generality. This was already pointed out in [Don00; Don03]. The obstruction has a topological character, i.e., certain cohomology groups have to vanish. To remedy this situation, one restricts attention to a certain “exact” subgroup, e.g., the subgroup of Hamiltonian diffeomorphisms in the group of all

symplectomorphisms. The action of this subgroup then usually admits a classical momentum map. Working from a completely different point of view, similar observations were made by Gay-Balmaz and Vizman [GV12] in their study of the classical dual pair in hydrodynamics. In this case, the symplectic action of volume-preserving diffeomorphisms on a symplectic manifold of mappings only has a momentum map under certain topological conditions and one is forced to work with suitable central extensions of the group of exact volume-preserving diffeomorphisms.

In this paper, we take the viewpoint that the above mentioned topological obstructions are not a bug but a feature of the theory. The action of the diffeomorphism group interacts with, and is largely determined by, the topological structure of the bundle. Thus, one would expect to capture certain topological data (like characteristic classes) that are “conserved” by the action. Such “conservation laws” should be encoded in the momentum map. Since the classical momentum map takes values in a continuous vector space, there is no space to “store” discrete topological information. Hence, whenever those classes do not vanish, a classical momentum map does not exist. Nonetheless, one could hope that a generalized momentum map exists and captures the conserved topological data.

The second main aim of the paper is to translate these philosophical remarks into explicit mathematical statements. In order to do this, we generalize the notion of momentum maps allowing them to take values in groups. Our concept of a group-valued momentum map is inspired by the Lu momentum map [Lu90; LW90] in Poisson geometry. We emphasize right away that the group-valued momentum map we introduce in this paper is a vast generalization of many notions of momentum maps appearing in the literature including circle-valued, cylinder-valued, and Lie algebra-valued momentum maps. We show that our generalized group-valued momentum map always exists for the action of the diffeomorphism group, without any topological assumptions on the base but some integrability conditions on the fiber model. The resulting momentum map captures topological invariants of the geometry, exactly in (the dual of) those cohomology classes which prevented the existence of a classical momentum map. This approach of extending the definition of the momentum map, besides the situation described above in Poisson geometry, in order to capture conservation laws not available using the classical definition, has been used successfully before in the theory of the cylinder-valued and optimal momentum maps; see [OR03] for a detailed presentation.

To see how our approach works, consider the following setting. Let (M, μ) be a closed (i.e., compact and boundaryless) n -manifold with volume form μ and (F, ω) a symplectic manifold. The space $C^\infty(M, F)$ of smooth maps from M to F

carries the weak symplectic form

$$\Omega_\phi(X, Y) = \int_M \omega_{\phi(m)}(X(m), Y(m)) \mu(m), \quad (1.1)$$

where $\phi \in C^\infty(M, F)$ and $X, Y \in T_\phi C^\infty(M, F)$, i.e., the maps $X, Y: M \rightarrow TF$ satisfy $X(m), Y(m) \in T_{\phi(m)}F$ for all $m \in M$. The natural action by precomposition of $\text{Diff}_\mu(M)$, the group of diffeomorphisms of M preserving the volume form μ , leaves Ω invariant. If ω is exact, say with primitive ϑ , then the momentum map is given by

$$\overline{\mathcal{J}}: C^\infty(M, F) \rightarrow \Omega^1(M)/d\Omega^0(M), \quad \phi \mapsto [\phi^*\vartheta], \quad (1.2)$$

where the space of volume-preserving vector fields $\mathfrak{X}_\mu(M)$ (the vector fields whose μ -divergence vanishes) is identified with the space of closed $(n-1)$ -forms, so that $\mathfrak{X}_\mu(M)^* \simeq \Omega^1(M)/d\Omega^0(M)$ with respect to the natural integration pairing. More generally, Gay-Balmaz and Vizman [GV12] showed that a (non-equivariant) momentum map also exists when the pull-back of ω by all maps $\phi \in C^\infty(M, F)$ is exact; for example, this happens when $H^2(M, \mathbb{R})$ is trivial. Our generalized group-valued momentum map takes no longer values in $\mathfrak{X}_\mu(M)^*$, but instead in the Abelian group $\hat{H}^2(M, U(1))$ that parametrizes principal circle bundles with connections modulo gauge equivalence. If (F, ω) has a prequantum bundle (L, ϑ) , then the map

$$\mathcal{J}: C^\infty(M, F) \rightarrow \hat{H}^2(M, U(1)), \quad \phi \mapsto [\phi^*(L, \vartheta)], \quad (1.3)$$

which sends ϕ to the pull-back bundle with connection $\phi^*(L, \vartheta)$, is a group-valued momentum map. We see that no (topological) restrictions have to be imposed on M and only the integrability condition of the symplectic form ω on F is needed for the existence of a group-valued momentum map. In contrast to the classical momentum map, a $\hat{H}^2(M, U(1))$ -valued momentum map contains topological information. First, the Chern class of the bundle, as a class in $H^2(M, \mathbb{Z})$, is available from the generalized momentum map. In our simple example, this is the pull-back of the Chern class of L . A second class in $H^1(M, U(1))$ is related to the equivariance of the momentum map; see Remark 3.6 for details.

In Table 1, we illustrate the conclusions implied by our general framework in several important examples. Next, we comment separately on each one of them.

Marsden and Weinstein [MW83] construct Clebsch variables for ideal fluids starting from a similar infinite-dimensional symplectic system as discussed above. It turns out, that every vector field represented in those Clebsch variables has vanishing helicity, i.e., such a fluid configuration has trivial topology and no links or knots. Thus, topological interesting configurations such as the ones

	Space	Action of	Chern class	Secondary topological class
Hydrodynamics	$C^\infty(M, F)$	$\text{Diff}_\mu(M)$	0 (total vorticity) in $H^2(M, \mathbb{Z})$	Circulations in $H^1(M, \mathbb{R})$
Lagrangian embeddings	$C^\infty(L, M)$	$\text{Diff}_\mu(L)$	Torsion class in $H^2(M, \mathbb{Z})$	Liouville class in $H^1(M, \mathbb{R})$
Kähler geometry	$\Gamma^\infty(LM \times_{\text{Sp}} \text{Sp}/U)$	$\text{Diff}_\omega(M)$	$c_1(M) \cup [\omega]^{n-1}$ in $H^{2n}(M, \mathbb{Z})$	
Quantomorphism	$\Gamma^\infty(P \times_{U(1)} \mathbb{C})$	$\text{Aut}_\Gamma(P)$	trivial	
Gauge theory	$C(P)$	$\text{Aut}(P)$	Torsion class in $H^{2n}(M, \mathbb{Z})$	

Table 1: Overview of the examples discussed in Chapter 4. Here, μ is a volume form and ω a symplectic form. Moreover, $Q \rightarrow M$ denotes a prequantum circle bundle with connection Γ and $P \rightarrow M$ is an arbitrary principal G -bundle. The frame bundle is denoted by LM . We also abbreviated the homogeneous space $\text{Sp}(2n, \mathbb{R})/U(n)$ by Sp/U .

constructed in [EP12] cannot be written in terms of classical Clebsch variables. Our more general framework allows to construct generalized Clebsch variables for vector fields with integral helicity; see Subsection 4.1.1.

When applied to the space of Lagrangian immersions, the group-valued momentum map recovers the Liouville class as the conserved topological data. Moreover, we realize moduli spaces of Lagrangian immersions (and modifications thereof) as symplectic quotients (reduced spaces).

A wide range of interesting examples with geometric significance are obtained when the typical fiber \underline{F} is a symplectic homogeneous space G/H . In this case, sections of $LM \times_G \underline{F}$ correspond to reductions of the G -frame bundle LM to H . Special focus is put on the space of almost complex structures compatible with a given symplectic structure, i.e., $\underline{F} = \text{Sp}(2n, \mathbb{R})/U(n)$. In this case, the group-valued momentum map for the group of symplectomorphisms assigns to an almost complex structure the anti-canonical bundle. It was already observed by Donaldson [Don97] and Fujiki [Fuj92] that the Hermitian scalar curvature furnishes a classical momentum map for the action of the group of Hamiltonian symplectomorphisms. Of course, the Hermitian scalar curvature is the curvature of the anti-canonical bundle. Thus, the group-valued momentum map combines the geometric curvature structure with the topological data of the anti-canonical bundle. For the case of a 2-dimensional base manifold, we realize the Teichmüller moduli space with the symplectic Weil–Petersson form as a symplectic orbit reduced space.

Finally, we revisit the classical gauge theoretic setting of Atiyah and Bott [AB83]

and extend their work in two ways. First, we generalize from 2-dimensional surfaces to arbitrary symplectic manifolds M as the base (a similar extension was already discussed by Donaldson [Don87]). Secondly, we investigate the group-valued momentum map for the action of the *full automorphism group* on the space of connections. The existence of the momentum map for this action is tightly connected to the triviality of a certain class in $H^3(M, \mathbb{Z})$ that is canonically constructed from the symplectic structure and the principal G -bundle.

We emphasize that, in contrast to most papers discussing infinite-dimensional symplectic geometry, we do not work formally, but really address the functional analytical problems arising from the transition to the infinite-dimensional setting. In particular, smoothness of maps between infinite dimensional manifolds is understood in the sense of locally convex spaces as, for example, presented in [Nee06].

Throughout the paper, integrality of certain symplectic forms plays a central role. We remark that, to a large extent, this assumption was made for convenience. Most results carry over directly to symplectic forms with discrete period groups per $\omega \subseteq \mathbb{R}$, without much technical effort. In spirit, our results also hold in the general setting without any assumptions on the period group; however, then one is forced to work in the diffeological category because the quotient $\mathbb{R}/\text{per } \omega$ may no longer be a Lie group.

Finally, we note that most of our symplectic reduced spaces are obtained as (sometimes singular) *orbit* reduced spaces, a theory that is not yet present in the literature for infinite dimensional systems, even though we state theorems using it. However, the techniques in [Die19] which completely treats infinite dimensional singular symplectic *point* reduction, combined with the strategy in [OR03] for finite dimensional singular symplectic orbit reduction, yields a general theory of infinite dimensional singular symplectic orbit reduction, which is precisely what is needed here. In the interest of the focus of this paper, as well as its length, this theory will be presented in a future article.

STRUCTURE OF THE PAPER Starting from the Poisson setting, Chapter 2 introduces the notion of a group-valued momentum map. It is shown that this generalized momentum map still satisfies Noether's theorem. Moreover, existence and uniqueness questions are addressed. Section 2.6 prepares the calculation of the momentum map for the automorphism group by giving a construction of the momentum map for a group extension (to the best of our knowledge, this construction is original, even for classical momentum maps). In Chapter 3, the symplectic geometry of the space of sections of a symplectic fiber bundle is discussed. First, the momentum map for the gauge group is determined in Subsection 3.2.1 and then for the group of volume-preserving diffeomorphisms, culminating in Theorem 3.3. The momentum map for the group of symplec-

tomorphisms, given in Theorem 3.7, is obtained by viewing it as a subgroup of the volume-preserving diffeomorphisms. Finally, Chapter 4 discusses the applications outlined above.

For the convenience of the reader, in Appendix A we summarize the notations and conventions from tensor calculus, symplectic and Poisson geometry, etc., in force throughout this article. In Appendix B and Appendix D, we recall the main properties of fiber integration and differential characters, respectively. The only original work in the appendices is Proposition D.3, Appendix C, and Appendix D.3, where we calculate the derivative of the pull-back map and generalize the hat product of [Viz11] to fiber bundles and differential characters, respectively.

ACKNOWLEDGMENTS We are indebted to F. Gay-Balmaz, J. Huebschmann, B. Janssens, J.-H. Lu, K.-H. Neeb, G. Rudolph, and C. Vizman for helpful discussions. We gratefully acknowledge support of the German National Academic Foundation (DAAD), the National Natural Science Foundation of China (NNSFC), the Swiss National Science Foundation (SNF), the Max Planck Institute for Mathematics in the Sciences in Leipzig, the Swiss Federal Institute of Technology in Lausanne, and the Shanghai Jiao Tong University which made our collaboration possible.

2 GROUP-VALUED MOMENTUM MAPS

In order to handle the full groups of volume-preserving diffeomorphisms or symplectomorphisms, we need a more general concept of a momentum map. Our starting point is the notion of a momentum map in Poisson geometry as introduced by Lu and Weinstein [Lu90; LW90]. We hasten to add that this group-valued momentum map is not built on the pattern from the theory of quasi-Hamiltonian actions and shall comment on the relation between the two later on. Standard definitions and terminology, with the specification of sign and coefficient conventions, are recalled in Appendix A.

2.1 Poisson Lie group momentum maps

All manifolds and Lie groups occurring in this subsection are assumed to be finite-dimensional. The proofs of the statements below can be found in [Lu90; LW90].

A Lie group G is a *Poisson Lie group* if it is simultaneously a Poisson manifold (relative to the underlying manifold structure) such that group multiplication and inversion are Poisson maps. Let $\omega_G \in \mathfrak{X}^2(G)$ denote the Poisson tensor of G . Let (M, ω_M) be a Poisson manifold. A *Poisson action* of the Poisson Lie group G on (M, ω_M) is a smooth (left) action $G \times M \rightarrow M$ which is, in addition, a

Poisson map (with $G \times M$ endowed with the product Poisson structure $\omega_G \times \omega_M$, i.e., the Poisson bracket of two functions depending only on G is given by the Poisson bracket on G , the Poisson bracket of two functions depending only on M is given by the Poisson bracket on M , and functions on G Poisson commute with functions on M).

The Poisson tensor ω_G of a Poisson Lie group G , with Lie algebra \mathfrak{g} , necessarily vanishes at the identity element $e \in G$, which then allows for the definition of the *intrinsic derivative* $\epsilon: \mathfrak{g} \rightarrow \mathfrak{g} \wedge \mathfrak{g}$ by $\epsilon(A) := (\mathfrak{L}_X \omega_G)_e$, where $X \in \mathfrak{X}(G)$ is an arbitrary vector field satisfying $X_e = A$ and \mathfrak{L}_X denotes the Lie derivative in the direction X . The dual map $\epsilon^*: \mathfrak{g}^* \wedge \mathfrak{g}^* \rightarrow \mathfrak{g}^*$ satisfies the Jacobi identity and thus endows \mathfrak{g}^* with a Lie algebra structure. The unique connected and simply connected Lie group G^* whose Lie algebra is \mathfrak{g}^* is called the *dual group* of G . The Lie group G^* has a unique Poisson structure ω_{G^*} relative to which G^* is a Poisson Lie group such that the intrinsic derivative of ω_{G^*} coincides with the Lie bracket on \mathfrak{g} . If G is connected and simply connected, the intrinsic derivative ϵ is a cocycle which uniquely determines both Poisson Lie tensors ω_G and ω_{G^*} .

Let $G \times M \rightarrow M$ be a left Poisson action of the Poisson Lie group (G, ω_G) on the Poisson manifold (M, ω_M) , i.e., the left action map $G \times M \rightarrow M$ is Poisson. A smooth map $J: M \rightarrow G^*$, if it exists, is called a *momentum map* of this action if

$$A^* + \omega_M(\cdot, J^* A^l) = 0, \quad \text{for all } A \in \mathfrak{g}. \quad (2.1)$$

Here, A^* denotes the fundamental (or infinitesimal generator) vector field on M induced by the infinitesimal action of $A \in \mathfrak{g}$, i.e.,

$$A^*(m) := \left. \frac{d}{dt} \right|_{t=0} \exp(tA) \cdot m, \quad \text{for all } m \in M, \quad (2.2)$$

where $g \cdot m$ denotes the action of $g \in G$ on $m \in M$. The second term in (2.1) is interpreted in the following way. Since \mathfrak{g} is the dual of \mathfrak{g}^* (which is the Lie algebra of G^*), we may think of A as a linear functional on \mathfrak{g}^* and extend it to a left invariant one-form $A^l \in \Omega^1(G^*)$, i.e., $(A^l)_a(v) = \langle A, L_{a^{-1}} v \rangle$ for every $a \in G^*$ and $v \in T_a G^*$, where $L_{a^{-1}}$ denotes both the left translation by $a^{-1} \in G^*$ in G^* and its tangent map (derivative) on TG^* .

Assume now that the Poisson manifold (M, ω_M) is symplectic with symplectic form ω and let us unwind the definition in this case. For any $X_m \in T_m M$ we have

$$\begin{aligned} \omega_m(A_m^*, X_m) &= (\omega_M)_m \left((\omega_M^\sharp)^{-1} A_m^*, (\omega_M^\sharp)^{-1} X_m \right) \stackrel{(2.1)}{=} -(J^* A^l)_m(X_m) \\ &= -A_{J(m)}^l(T_m J(X_m)) = -\langle A, L_{J(m)^{-1}} T_m J(X_m) \rangle \\ &= -\langle A, (\delta J)_m(X_m) \rangle, \end{aligned} \quad (2.3)$$

where $T_m J: T_m M \rightarrow T_{J(m)} G^*$ is the derivative (tangent map) of $J: M \rightarrow G^*$ and $\delta J \in \Omega^1(M, \mathfrak{g}^*)$, defined by the last equality, is its *left logarithmic derivative*.

Note that the identity $\omega_m(A_m^*, X_m) + \langle A, (\delta J)_m(X_m) \rangle = 0$ proved in (2.3) does not use the fact that G is a Poisson Lie group. Indeed, this identity still makes sense if the momentum map is replaced by a smooth map $J: M \rightarrow H$ with values in an arbitrary Lie group H , as long as there is a duality between the Lie algebras of G and H . This observation leads to our generalization of Lu's momentum map. In order to define this generalization, we need a few preliminary concepts that are inspired by their counterparts in the theory of Poisson Lie groups.

2.2 Dual pairs of Lie algebras

A *dual pair of Lie algebras* (not necessarily finite-dimensional) consists of two Lie algebras \mathfrak{g} and \mathfrak{h} , which are in duality through a given (weakly) non-degenerate bilinear map $\kappa: \mathfrak{g} \times \mathfrak{h} \rightarrow \mathbb{R}$. Using notation stemming from functional analysis, we often write the dual pair as $\kappa(\mathfrak{g}, \mathfrak{h})$. Intuitively, we think of \mathfrak{h} as the dual vector space of \mathfrak{g} , endowed with its own Lie bracket operation. For this reason, we often denote $\mathfrak{g}^* := \mathfrak{h}$, even though \mathfrak{g}^* is not necessarily the functional analytic dual of \mathfrak{g} . Two Lie groups G and H are said to be *dual* to each other if there exists a (weakly) non-degenerate bilinear form $\kappa: \mathfrak{g} \times \mathfrak{h} \rightarrow \mathbb{R}$ relative to which the associated Lie algebras are in duality. We use the notation $\kappa(G, H)$ in this case. As for Lie algebras, we often write $G^* := H$, intuitively thinking of G^* as the dual Lie group, as in the theory of Poisson Lie groups ([LW90]).

Note that the notion of a dual pair of Lie algebras involves only the underlying vector spaces, while the Lie brackets play no role. We introduce a more rigid concept of duality, which takes all structures into account. For a given dual pair $\kappa(\mathfrak{g}, \mathfrak{h})$ of Lie algebras, define a bilinear skew-symmetric bracket on the *double* $\mathfrak{d} := \mathfrak{g} \times \mathfrak{h}$ by

$$[(A, \mu), (B, \nu)] = ([A, B]_{\mathfrak{g}} - \text{ad}_{\mu}^* B + \text{ad}_{\nu}^* A, [\mu, \nu]_{\mathfrak{h}} - \text{ad}_A^* \nu + \text{ad}_B^* \mu), \quad (2.4)$$

for $A, B \in \mathfrak{g}$, $\mu, \nu \in \mathfrak{h}$, where the infinitesimal coadjoint actions are defined with respect to κ by¹

$$\begin{aligned} \kappa(B, \text{ad}_A^* \mu) &= \kappa([A, B]_{\mathfrak{g}}, \mu), \\ \kappa(\text{ad}_{\mu}^* A, \nu) &= \kappa(A, [\mu, \nu]_{\mathfrak{h}}). \end{aligned} \quad (2.5)$$

However, the bracket operation defined in (2.4) does not satisfy the Jacobi identity, in general. A dual pair $\kappa(\mathfrak{g}, \mathfrak{h})$ of Lie algebras is called a *Lie bialgebra*, if

¹ In infinite dimensions, equation (2.5) ensures only uniqueness of the adjoints but not their existence. In the sequel, we always assume that the coadjoint actions exist.

the bracket (2.4) on $\mathfrak{d} = \mathfrak{g} \times \mathfrak{h}$ is a Lie bracket. In this case, we denote the double by $\mathfrak{g} \bowtie \mathfrak{h}$. The notation displays the similarity to the semidirect product but also emphasizes that the bracket on $\mathfrak{g} \times \mathfrak{h}$ uses both the adjoint actions of \mathfrak{g} and \mathfrak{h} .

EXAMPLE 2.1 Let $\kappa(\mathfrak{g}, \mathfrak{h})$ be a dual pair. Assume that \mathfrak{h} is an Abelian Lie algebra. Then (2.5) implies that the coadjoint action $\text{ad}_\mu^* : \mathfrak{g} \rightarrow \mathfrak{g}$ is trivial for every $\mu \in \mathfrak{h}$. Hence the bracket on the double \mathfrak{d} simplifies to

$$[(A, \mu), (B, \nu)] = ([A, B]_{\mathfrak{g}}, -\text{ad}_A^* \nu + \text{ad}_B^* \mu). \quad (2.6)$$

Thus, \mathfrak{d} is the semidirect product $\mathfrak{g} \bowtie_{\text{ad}^*} \mathfrak{h}$ of Lie algebras, where \mathfrak{g} acts on \mathfrak{h} by the κ -coadjoint action (2.5). Furthermore, the Jacobi identity always holds and thus $\kappa(\mathfrak{g}, \mathfrak{h})$ is a Lie bialgebra. \diamond

EXAMPLE 2.2 (Group of volume-preserving diffeomorphisms) Let M be a compact manifold endowed with a volume form μ . Then the group $G = \text{Diff}_\mu(M)$ of volume-preserving diffeomorphisms is a Fréchet Lie group, see [Ham82, Theorem 2.5.3]. Its Lie algebra consists of μ -divergence-free vector fields X on M . Equivalently,

$$\mathfrak{g} = \mathfrak{X}_\mu(M) = \{X \in \mathfrak{X}(M) \mid \text{d}(X \lrcorner \mu) = 0\}, \quad (2.7)$$

where d denotes the exterior derivative on forms and $X \lrcorner \mu := i_X \mu$ is the interior product (contraction on the first index) of X with μ . Hence, we also identify $\mathfrak{X}_\mu(M)$ with $\Omega_{\text{cl}}^{\dim M - 1}(M)$ via $X \mapsto X \lrcorner \mu$, where $\Omega_{\text{cl}}^k(M)$ denotes the space of closed k -forms on M . Thus $\Omega^1(M)/\text{d}\Omega^0(M)$ is the regular dual with respect to the weakly non-degenerate integration pairing

$$\kappa(X, \alpha) := (-1)^{\dim M - 1} \int_M (X \lrcorner \mu) \wedge \alpha = \int_M (X \lrcorner \alpha) \mu. \quad (2.8)$$

We now note that a 1-form α can be interpreted as a trivial principal circle bundle with curvature $\text{d}\alpha$. From this point of view, $\Omega^1(M)/\text{d}\Omega^0(M)$ parametrizes gauge equivalence classes of connections on a trivial principal circle bundle. Thus, it is natural to think of it as the Lie algebra of the Abelian group $H := \hat{H}^2(M, \text{U}(1))$ of all principal circle bundles with connections, modulo gauge equivalence. This heuristic argument can be made rigorous using the theory of Cheeger–Simons differential characters; see Appendix D for a quick review. In summary, we get a dual pair $\kappa(\text{Diff}_\mu(M), \hat{H}^2(M, \text{U}(1)))$ of Lie groups. For later use, it is convenient to introduce the notation $\hat{\mathfrak{h}}^2(M, \text{U}(1))$ for the Lie algebra $\Omega^1(M)/\text{d}\Omega^0(M)$ of $\hat{H}^2(M, \text{U}(1))$. \diamond

EXAMPLE 2.3 (Group of symplectomorphisms) Consider a compact symplectic manifold (M, ω) . The group $G = \text{Diff}_\omega(M)$ of symplectomorphisms is a Fréchet

Lie group [KM97, Theorem 43.12.] with Lie algebra

$$\mathfrak{g} = \mathfrak{X}_\omega(M) = \{X \in \mathfrak{X}(M) \mid d(X \lrcorner \omega) = 0\}. \quad (2.9)$$

The map $X \mapsto X \lrcorner \omega$ identifies \mathfrak{g} with the space $\Omega_{\text{cl}}^1(M)$ of closed 1-forms on M . Thus the regular dual of \mathfrak{g} with respect to the natural integration pairing

$$\kappa(X, \alpha) := \frac{(-1)^{\dim M - 1}}{\left(\frac{1}{2} \dim M - 1\right)!} \int_M (X \lrcorner \omega) \wedge \alpha \quad (2.10)$$

is $\hat{\mathfrak{h}}^{\dim M}(M, U(1)) := \Omega^{\dim M - 1}(M) / d\Omega^{\dim M - 2}(M)$. The prefactor in front of the integral turns out to be a convenient choice in Subsection 3.2.4. Similarly to the Example 2.2 of volume-preserving diffeomorphisms, the Abelian Lie algebra $\hat{\mathfrak{h}}^{\dim M}(M, U(1))$ is integrated by the group $\hat{H}^{\dim M}(M, U(1))$ of Cheeger–Simons differential characters with degree $\dim M$. These can be thought of as equivalence classes of circle n -bundles with connections in the sense of higher differential geometry. \diamond

REMARK 2.4 The group of volume-preserving diffeomorphism and the group of symplectomorphisms both have an Abelian dual group. Thus they fall in the realm of Example 2.1. In particular, both dual pairs $(\mathfrak{X}_\mu(M), \hat{\mathfrak{h}}^2(M, U(1)))$ and $(\mathfrak{X}_\omega(M), \hat{\mathfrak{h}}^{\dim M}(M, U(1)))$ are actually Lie bialgebras. Ignoring the particularities of the infinite-dimensional setting for a moment, Drinfeld’s theorem [ES02, Theorem 2.2.2] states that there are essentially unique connected and simply connected Poisson Lie groups, whose Lie algebras are $\mathfrak{X}_\mu(M)$ and $\mathfrak{X}_\omega(M)$, respectively. We do not know if the groups $\text{Diff}_\mu(M)$ or $\text{Diff}_\omega(M)$ carry a non-trivial Poisson Lie structure integrating the above Lie bialgebras (this would require to find a non-trivial integration of the adjoint action, see [Dri93] and [Lu90, Theorem 2.31]). Moreover, we are not aware of any Poisson Lie structure on these groups, such that the actions discussed in Chapter 3 are Poisson maps. \diamond

EXAMPLE 2.5 (Gauge group) Let $P \rightarrow M$ be a right principal G -bundle over the compact manifold M . The group $\text{Gau}(P)$ of gauge transformations is identified with the space of sections of $P \times_G G := (P \times G)/G$ and thus is a Fréchet Lie group with Lie algebra $\mathfrak{gau}(P) = \Gamma^\infty(\text{Ad}P)$, the space of sections of the adjoint bundle $\text{Ad}P := (P \times \mathfrak{g})/G$, see [CM85]. Denote the dual of the adjoint bundle by $\text{Ad}^*P := (P \times \mathfrak{g}^*)/G$, the action of G on \mathfrak{g}^* being the left coadjoint action. The natural pairing

$$\kappa(\phi, \alpha) = \int_M \langle \phi, \alpha \rangle, \quad \phi \in \mathfrak{gau}(P), \alpha \in \Omega^{\dim M}(M, \text{Ad}^*P), \quad (2.11)$$

identifies $\Omega^{\dim M}(M, \text{Ad}^*P)$ as the regular dual to $\mathfrak{gau}(P)$. In particular, if M

is endowed with a volume form μ , then $\text{gau}^*(P) = \Gamma^\infty(\text{Ad}^*P)$ is the dual by integration against μ :

$$\langle \cdot, \cdot \rangle_{\text{Ad}}: \Gamma^\infty(\text{Ad}P) \times \Gamma^\infty(\text{Ad}^*P) \rightarrow \mathbb{R}, \quad (\phi, \varrho) \mapsto \int_M \langle \phi, \varrho \rangle \mu. \quad (2.12)$$

Moreover, an Ad_G -invariant non-degenerate bilinear form on \mathfrak{g} identifies $\text{gau}^*(P)$ with $\text{gau}(P)$ so that $\text{gau}(P)$ is self-dual in this case. \diamond

2.3 Group-valued momentum maps

With these preliminaries out of the way, we can introduce our generalization of a group-valued momentum map.

DEFINITION 2.6 Let M be G -manifold endowed with a symplectic form¹ ω . A *group-valued momentum map* is a pair (J, κ) , where $\kappa(G, G^*)$ is a dual pair of Lie groups and $J: M \rightarrow G^*$ is a smooth map satisfying

$$A^* \lrcorner \omega + \kappa(A, \delta J) = 0, \quad A \in \mathfrak{g}. \quad (2.13)$$

In this formula, A^* denotes the fundamental vector field on M induced by $A \in \mathfrak{g}$, $\delta J \in \Omega^1(M, \mathfrak{g}^*)$ is the left logarithmic derivative of J , \mathfrak{g} is the Lie algebra of G , and \mathfrak{g}^* is the Lie algebra of G^* . \diamond

EXAMPLE 2.7 (As a generalization of the usual momentum map) Let (M, ω) be a finite-dimensional, symplectic G -manifold. We view $G^* = \mathfrak{g}^*$ as an Abelian group and consider the natural pairing $\kappa: \mathfrak{g} \times \mathfrak{g}^* \rightarrow \mathbb{R}$. Thus a \mathfrak{g}^* -valued momentum map is a smooth map $J: M \rightarrow \mathfrak{g}^*$ satisfying the usual relation

$$A^* \lrcorner \omega + dJ_A = 0, \quad A \in \mathfrak{g}, \quad (2.14)$$

where $J_A: M \rightarrow \mathbb{R}$ is defined by $J_A(m) = \kappa(A, J(m))$ for $m \in M$, and $dJ_A = \kappa(A, \delta J)$ holds as a consequence of the Abelian character of \mathfrak{g}^* . \diamond

EXAMPLE 2.8 (As a generalization of the Lie algebra-valued momentum map) If $\kappa: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{R}$ is a continuous, weakly non-degenerate, Ad_G -invariant symmetric bilinear form, one identifies the dual \mathfrak{g}^* with \mathfrak{g} and the coadjoint orbits with adjoint orbits. In view of Example 2.7, we are led to the concept of a Lie algebra-valued momentum map. Although this notion has been in the literature since the mid 70s, it was recently formalized by Neeb, Sahlmann, and Thiemann [NST14, Definition 4.3]: a Lie algebra-valued momentum map is a smooth map $J: M \rightarrow \mathfrak{g}$

¹ We do not require the symplectic form ω to be invariant under the group action. In fact, we will see below in Lemma 2.14 that this will not be the case if the dual group H is non-Abelian.

such that, for all $A \in \mathfrak{g}$, the component functions $J_A = \kappa(A, J): M \rightarrow \mathbb{R}$ satisfy

$$A^* \lrcorner \omega + dJ_A = 0. \quad (2.15)$$

It is immediate from the definition, that such a Lie algebra-valued momentum map can be regarded as a group-valued momentum map with respect to the dual pair $\kappa(G, \mathfrak{g})$, where \mathfrak{g} is viewed as an Abelian Lie group. \diamond

EXAMPLE 2.9 (As a generalization of the Poisson momentum map) We now describe how our motivational example fits in the new framework. Let (M, ω) be a finite-dimensional symplectic manifold. Suppose a finite-dimensional Poisson Lie group G acts on M such that the action is Poisson. The Poisson structure of G induces a Lie bracket on the dual \mathfrak{g}^* of the Lie algebra [Lugo, Theorem 2.18]. Let G^* denote the unique connected and simply connected Lie group integrating the Lie algebra \mathfrak{g}^* . As above, let κ be the natural pairing $\mathfrak{g} \times \mathfrak{g}^* \rightarrow \mathbb{R}$. Hence a G^* -valued momentum map is a smooth map $J: M \rightarrow G^*$ satisfying

$$A^* \lrcorner \omega + \kappa(A, \delta J) = 0. \quad (2.16)$$

As we have explained above, this equation is a reformulation of the usual Lu momentum map relation in the context of symplectic geometry, see (2.3). In other words, our group-valued momentum map is the natural generalization of the Poisson momentum map if the Lie group G is not necessarily a Poisson Lie group. \diamond

EXAMPLE 2.10 (As a generalization of the circle-valued momentum map) Consider a symplectic manifold (M, ω) with a symplectic action of $G = U(1)$. We let $G^* = U(1)$ and take (minus) the Killing form $\kappa: \mathfrak{u}(1) \times \mathfrak{u}(1) \rightarrow \mathbb{R}$ as the pairing between the Lie algebras of G and G^* , i.e., $\kappa(x, y) = xy$ is the usual multiplication of real numbers under the identification $\mathfrak{u}(1) \simeq \mathbb{R}$. Thus, a map $J: M \rightarrow U(1)$ is a group-valued momentum map if and only if

$$1^* \lrcorner \omega + \delta J = 0. \quad (2.17)$$

In this way, we recover the usual definition of a circle-valued momentum map (see, for example, [PR12, Definition 1]). \diamond

EXAMPLE 2.11 (Symplectic torus) Consider a symplectic vector space (V, ω) . Let Λ be a lattice in V , i.e., a discrete subgroup of the additive group V . The symplectic structure is invariant under the natural action of Λ on V by translations and hence it descends to a symplectic form ω_T on the torus $T = V/\Lambda$. Moreover, the translation action of V on itself commutes with the lattice action and thus induces a symplectic V -action on T . The action of V on itself has the momentum map

$$J: V \rightarrow V^*, \quad v \mapsto \omega(v, \cdot). \quad (2.18)$$

However, J is not invariant under the lattice action and so does *not* descend to a momentum map for the induced action of V on the torus T . Indeed, it is well-known that, for cohomological reasons, the symplectic action on the torus does not admit a standard momentum map. Rather, J transforms as

$$J(v + \lambda) = J(v) + \omega(\lambda, \cdot), \quad \lambda \in \Lambda. \quad (2.19)$$

Thus, if $\omega(\lambda_1, \lambda_2) \in \mathbb{Z}$ holds for all $\lambda_1, \lambda_2 \in \Lambda$, then J is equivariant with respect to the dual lattice action

$$\Lambda^* = \{\alpha \in V^* \mid \alpha(\lambda) \in \mathbb{Z} \text{ for all } \lambda \in \Lambda\}. \quad (2.20)$$

In this case, J induces a V^*/Λ^* -valued momentum map J_T on the torus T . It is interesting to note that the integrality condition $\omega(\lambda_1, \lambda_2) \in \mathbb{Z}$ is equivalent to ω_T being prequantizable. \diamond

EXAMPLE 2.12 (As a generalization of the cylinder-valued momentum map) In a finite-dimensional context, Condevaux, Dazord, and Molino [CDM88] introduced a momentum map with values in the cylinder $C := \mathfrak{g}^*/H$, where H is the holonomy group of a flat connection on some bundle constructed in terms of the symplectic form and the action (in our language, α defined in (2.22) below plays the role of the connection form). If the holonomy group H is discrete, then C is a Lie group with Lie algebra \mathfrak{g}^* . Thus C is a dual group. Under the identification of the Lie algebra $\mathfrak{c} = \mathfrak{g}^*$, the cylinder-valued momentum map also satisfies (2.13), see [OR03, Theorem 5.2.8], and hence is a group-valued momentum map. The group-valued momentum map for the symplectic torus discussed in Example 2.11 is also the cylinder-valued momentum map, see [OR03, Example 5.2.5].

The case when the holonomy group $H \subseteq \mathfrak{g}^*$ has accumulation points is pathological both in the framework of cylinder- as well as group-valued momentum maps, see Example 2.17 and cf. [OR03, Example 5.2.6]. \diamond

As we have seen, the notion of a group-valued momentum map unifies many different concepts of momentum maps. Despite its general nature, a group-valued momentum map still captures conserved quantities of the dynamical system, i.e., it has the *Noether property* (see [OR03, Definition 4.3.1]).

PROPOSITION 2.13 (Noether's theorem) *Let (M, ω) be symplectic G -manifold. Assume that the action has a G^* -valued momentum map $J: M \rightarrow G^*$. Let $h \in C^\infty(M)$ be a smooth function for which the Hamiltonian vector field X^h exists and has a unique local flow¹. If h is G -invariant, then J is constant along the integral curves of X^h . \diamond*

¹ Recall that vector fields on Fréchet manifolds do not need to have flows. Having a flow is more or less equivalent to local in time solutions of the corresponding partial differential equation.

Proof. Let $A \in \mathfrak{g}$ and $m \in M$. Using the defining equation for the momentum map, we have

$$\begin{aligned} \kappa(A, (\delta J)_m X_m^h) &= -\omega_m(A_m^*, X_m^h) = -(\mathrm{d}h)_m A_m^* \\ &= -\left. \frac{\mathrm{d}}{\mathrm{d}\varepsilon} \right|_0 h(\exp(\varepsilon A) \cdot m) = 0 \end{aligned} \quad (2.21)$$

by G -invariance of h . Since $A \in \mathfrak{g}$ is arbitrary and the pairing κ is weakly non-degenerate, we conclude $(\delta J)X^h = 0$. Hence, J is constant along integral curves of X^h . \square

2.4 Existence and uniqueness

Let (M, ω) be a symplectic manifold. For a given G -action on M and a fixed dual pair $\kappa(G, G^*)$ of Lie groups, we will now search for conditions which ensure the existence of a G^* -valued momentum map.

Define a \mathfrak{g}^* -valued 1-form $\alpha \in \Omega^1(M, \mathfrak{g}^*)$ by $A^* \lrcorner \omega + \kappa(A, \alpha) = 0$, that is,

$$\kappa(A, \alpha_m(X_m)) = \omega_m(X_m, A_m^*) \quad \text{for all } X_m \in T_m M, A \in \mathfrak{g}. \quad (2.22)$$

In infinite dimensions, the dual pairing κ is often not strongly non-degenerate. In such cases, there might not exist $\alpha \in \Omega^1(M, \mathfrak{g}^*)$ satisfying (2.22), although α is unique, if it exists. We will assume for the rest of the subsection, that we have such an α .

By (2.13), the G -action on M admits a G^* -valued momentum map if and only if $\alpha \in \Omega^1(M, \mathfrak{g}^*)$ defined by (2.22) is log-exact, i.e., if $\alpha = \delta J$ for some smooth function $J: M \rightarrow G^*$. A necessary condition is that α satisfies the Maurer–Cartan equation, as the next lemma shows. Below, $[\alpha \wedge \beta]$ means the wedge product of the \mathfrak{g}^* -valued forms α and β on M associated to the bracket operation on \mathfrak{g}^* (as the Lie algebra of G^*).

LEMMA 2.14 (Corresponds to [Lu90, Theorem 3.7])

(i) *The \mathfrak{g}^* -valued one-form α on M defined by (2.22) satisfies the Maurer–Cartan equation*

$$\mathrm{d}\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0 \quad (2.23)$$

if and only if $\mathfrak{L}_{A^}\omega = \frac{1}{2}\kappa(A, [\alpha \wedge \alpha])$ holds for all $A \in \mathfrak{g}$.*

(ii) *If the G -action on M admits a G^* -valued momentum map $J: M \rightarrow G^*$, then $\delta J \in \Omega^1(M, \mathfrak{g}^*)$ satisfies the Maurer–Cartan equation (2.23). \diamond*

If the form $\alpha \in \Omega^1(M, \mathfrak{g}^*)$ defined by (2.22) exists and satisfies the Maurer–Cartan equation (2.23), then we say that the G -action on (M, ω) is G^* -symplectic.

For Abelian G^* , the notions of symplectic and G^* -symplectic group actions coincide.

Proof. (i) For $A \in \mathfrak{g}$ and $X, Y \in \mathfrak{X}(M)$, we calculate

$$\begin{aligned}
 (\mathcal{Q}_{A^*}\omega)(X, Y) &= d(A^* \lrcorner \omega)(X, Y) \\
 &= X(\omega(A^*, Y)) - Y(\omega(A^*, X)) - \omega(A^*, [X, Y]) \\
 &\stackrel{(2.22)}{=} -\kappa(A, X(\alpha(Y)) - Y(\alpha(X)) - \alpha([X, Y])) \\
 &= -\kappa(A, (d\alpha)(X, Y)),
 \end{aligned} \tag{2.24}$$

which is equivalent to $\mathcal{Q}_{A^*}\omega + \kappa(A, d\alpha) = 0$, from which the claim follows.

(ii) If the G -action admits a G^* -valued momentum map $J: M \rightarrow G^*$, then δJ satisfies the Maurer-Cartan equation (2.23) if and only if $\mathcal{Q}_{A^*}\omega + \kappa(A, d\delta J) = 0$, as we just showed in (i). This identity is obtained by taking the exterior derivative of the defining relation (2.13). \square

Although, at first glance, it may seem unnatural to consider momentum maps for actions which do not leave the symplectic form invariant, we remind the reader that this is also the case in the Poisson category. There, the action $\Upsilon: G \times M \rightarrow M$ is a Poisson map, while for fixed $g \in G$ the map $\Upsilon_g: M \rightarrow M$ might not preserve the Poisson structure. If one restricts attention to objects invariant under the action, then non-closed almost symplectic structures have to be considered. In other words, this path leads to quasi-Hamiltonian systems [AMM98; HJ94].

EXAMPLE 2.15 (Poisson Lie Momentum Map [FR96]) A Poisson action of a non-Abelian Poisson Lie group on a symplectic manifold, even if it admits a momentum map (see Example 2.9), never preserves the symplectic form. For example, consider the Poisson Lie group $SU(n)$ endowed with the Lu-Weinstein Poisson structure. Its dual Poisson Lie group is $B := AN$, where A is the subgroup of positive diagonal matrices, N is the subgroup of complex triangular matrices with 1 on the diagonal, both viewed as real subgroups of $SL(n, \mathbb{C})$, also viewed as a real Lie group. Thus, B has a natural Poisson Lie group structure whose symplectic leaves are the sets Σ_λ of matrices with fixed singular values $\lambda := \{\lambda_1 \geq \dots \geq \lambda_n > 0\}$ (the dressing orbits). The Lie group $SU(n)$ acts on the symplectic manifold $\Sigma_\lambda \times \Sigma_\mu$ in a Poisson fashion and admits the momentum map $J: \Sigma_\lambda \times \Sigma_\mu \ni (b_1, b_2) \mapsto b_1 b_2 \in B$. The Poisson action is given in the following way. First, one shows that if $b_1 \in \Sigma_\lambda$, $b_2 \in \Sigma_\mu$, and $b_1 b_2 \in \Sigma_\nu$, then there exists $b_3 \in \Sigma_\lambda$, $b_4 \in \Sigma_\mu$ such that $b_3 b_4 = \text{diag}(v_1, \dots, v_n)$. Second, since $b_1 b_2 \in \Sigma_\nu$, there exist $k_1, k_2 \in SU(n)$ such that $k_1 b_1 b_2 k_2 = \text{diag}(v_1, \dots, v_n)$. With these notations, the aforementioned Poisson action of $SU(n)$ on $\Sigma_\lambda \times \Sigma_\mu$ is given by $k_2^{-1} \cdot (b_1, b_2) := (b_3, b_4)$. See [FR96] for the proofs of these statements and their link to a convexity theorem for compact Poisson Lie group actions on

compact connected symplectic manifolds. In this case, the theorem states that $J(\Sigma_\lambda \times \Sigma_\mu) \cap A_+$ is the exponential of a convex polytope, where A_+ denotes the set of elements of A with decreasingly ordered entries. \diamond

Returning to the existence and uniqueness question, recall that the Maurer–Cartan theory involves also a topological obstruction for the existence of a primitive for α in the form of the so-called period homomorphism [Nee06, Theorem III.1.2]. To strengthen Lemma 2.14 in terms of the period map, we need to recall the notion of a regular Lie group (see [Nee06, Definition II.5.5]). A Lie group G modeled on a locally convex space is *regular* if for each curve $c \in C^\infty([0, 1], \mathfrak{g})$, the initial value problem $\delta\eta(t) := L_{\eta(t)^{-1}} \dot{\eta}(t) = c(t)$, $\eta(0) = e$, formulated in terms of the left logarithmic derivative $\delta\eta$ of η , has a solution $\eta_c \in C^\infty([0, 1], G)$ and the endpoint evaluation map $C^\infty([0, 1], \mathfrak{g}) \ni c \mapsto \eta_c(1) \in G$ is smooth. If G is regular, then it has a smooth exponential function. All Banach (so, in particular, all finite dimensional) Lie groups are regular.

Fix a point $m_0 \in M$ and consider a piece-wise smooth loop $\gamma: I \rightarrow M$ based at m_0 . The pull-back of $\alpha \in \Omega^1(M, \mathfrak{g}^*)$ by γ is a \mathfrak{g}^* -valued 1-form $\gamma^*\alpha$ on I . Denote by $\eta_\gamma \in C^\infty(I, G^*)$ the solution of the initial value problem

$$\delta\eta = \gamma^*\alpha, \quad \eta(0) = e, \quad (2.25)$$

which exists if G^* is regular. Evaluating η_γ at the endpoint 1, we obtain the period homomorphism $\text{per}_\alpha: \pi_1(M, m_0) \ni [\gamma] \mapsto \eta_\gamma(1) \in G^*$, where $[\gamma]$ is the homotopy class of the loop γ .

PROPOSITION 2.16 *Let (M, ω) be a connected symplectic manifold and $\kappa(G, G^*)$ a dual pair of Lie groups. In infinite dimensions, we additionally assume that G^* is a regular Lie group. Let G act on M in a G^* -symplectic way. Then there exists a G^* -valued momentum map if and only if the period homomorphism $\text{per}_\alpha: \pi_1(M, m_0) \rightarrow G^*$ is trivial. Moreover, the momentum map is unique up to translation by a constant element $h \in G^*$. \diamond*

Proof. This proposition follows directly from Lemma 2.14 and the general fundamental theorem for Lie group-valued functions [Nee06, Theorem III.1.2.].

For the convince of the reader, we quickly sketch how the period homomorphism enters the game. Recall that an ordinary closed 1-form has a local primitive by the Poincaré lemma. The Maurer–Cartan equation (2.23) replaces the role of closedness for Lie algebra-valued 1-forms. Since the action is G^* -symplectic, the solution α of (2.22) satisfies (2.23) by Lemma 2.14. Hence, on connected and simply connected domains, a G^* -valued momentum map exists. If M is not simply connected, then we pull back everything to the universal covering $\tilde{M} \rightarrow M$ and obtain a lifted primitive $\tilde{J}: \tilde{M} \rightarrow G^*$. A simple calculation reveals

that \tilde{J} is equivariant with respect to per_α , i.e.,

$$\tilde{J}([\gamma] \cdot \tilde{m}) = \text{per}_\alpha([\gamma]) \cdot \tilde{J}(\tilde{m}), \quad \tilde{m} \in \tilde{M}, \quad [\gamma] \in \pi_1(M, m_0). \quad (2.26)$$

Thus \tilde{J} drops to M if and only if the period homomorphism is trivial. \square

EXAMPLE 2.17 A slight modification of Example 2.11 yields a symplectic group action without a group-valued momentum map. Let $M = T^2 = (\mathbb{R}/\mathbb{Z})^4$ be the product of two tori and denote its elements by 4-tuples $(\varphi_1, \varphi_2, \psi_1, \psi_2)$. Endow M with the symplectic form

$$\omega = d\varphi_1 \wedge d\varphi_2 + \sqrt{2} d\psi_1 \wedge d\psi_2. \quad (2.27)$$

The circle action given by $\lambda \cdot (\varphi_1, \varphi_2, \psi_1, \psi_2) = (\varphi_1 - \lambda, \varphi_2, \psi_1 - \lambda, \psi_2)$ is clearly symplectic. The 1-form α defined by equation (2.22) becomes

$$\alpha = d\varphi_2 + \sqrt{2} d\psi_2. \quad (2.28)$$

As the generators of $\pi_1(M)$ we take the four natural loops γ_i , where, for $1 \leq i \leq 4$, the loop $\gamma_i: I \rightarrow M$ winds once around the i -th circle in $M = (\mathbb{R}/\mathbb{Z})^4$. The pull-back of α by γ_1 and γ_3 vanishes and we find

$$\gamma_2^* \alpha = dt \quad \text{and} \quad \gamma_4^* \alpha = \sqrt{2} dt, \quad (2.29)$$

where t denotes the canonical variable in $I = [0, 1]$. Since there are only two connected one-dimensional Lie groups, the only possible choices for the dual group are $G^* = \mathbb{R}$ and $G^* = \mathbb{R}/\mathbb{Z}$. In both cases, the initial value problem (2.25) has the unique solutions $\eta_{\gamma_2}(t) = t$ and $\eta_{\gamma_4}(t) = \sqrt{2}t$. Thus neither for $G^* = \mathbb{R}$ nor for $G^* = \mathbb{R}/\mathbb{Z}$ the period homomorphism is trivial and hence no group-valued momentum map exists for this action.

This example exhibits another phenomenon that is particular for group-valued momentum maps: the action of a subgroup may not possess a group-valued momentum map even if the bigger group has a group-valued momentum map. In fact, the action $(\lambda_1, \lambda_2) \cdot (\varphi_1, \varphi_2, \psi_1, \psi_2) = (\varphi_1 - \lambda_1, \varphi_2, \psi_1 - \lambda_2, \psi_2)$ by $G = S^1 \times S^1$ has a group-valued momentum map (which is the product of two copies of the one discussed in Example 2.11) but the action of the diagonally embedded circle has no group-valued momentum map as we have just seen. \diamond

2.5 Equivariance and Poisson property

Fix a dual pair $\kappa(G, G^*)$ of Lie groups. Let G act on the symplectic manifold (M, ω) such that the action has a group-valued momentum map $J: M \rightarrow G^*$. A natural question to ask is in which sense J is equivariant. For classical momentum

maps with values in \mathfrak{g}^* , the equivariance of interest is with respect to the coadjoint action. Before we give the definition of the analogue of the coadjoint action in the group-valued setting, let us recall a few basic notations and conventions; see Appendix A for details. The coadjoint action of G on \mathfrak{g}^* is defined with respect to the duality pairing κ by¹ $\kappa(A, \text{Ad}_g^* \mu) = \kappa(\text{Ad}_g A, \mu)$ for $g \in G, A \in \mathfrak{g}$ and $\mu \in \mathfrak{g}^*$. The coadjoint action of G^* on \mathfrak{g} is similarly defined and will be denoted by Ad^* as well. For an action $\Upsilon: G \times G^* \rightarrow G^*$, we introduce the partial maps $\Upsilon_g: G^* \rightarrow G^*$ and $\Upsilon_\eta: G \rightarrow G^*$ by $\Upsilon_g(\eta) \equiv \Upsilon_\eta(g) \equiv \Upsilon(g, \eta)$ for $g \in G$ and $\eta \in G^*$.

DEFINITION 2.18 A *coconjugation action* on the dual pair $\kappa(G, G^*)$ is a pair (Υ, Υ^*) consisting of a left action $\Upsilon: G \times G^* \rightarrow G^*$ of G on G^* and a left action $\Upsilon^*: G^* \times G \rightarrow G$ of G^* on G which integrate the coadjoint actions in the sense that $\Upsilon_g(e) = e$, $\Upsilon_\eta^*(e) = e$ and

$$T_e \Upsilon_g(\mu) = \text{Ad}_{g^{-1}}^* \mu, \quad T_e \Upsilon_\eta^*(A) = \text{Ad}_{\eta^{-1}}^* A \quad (2.30)$$

holds for all $g \in G, \eta \in G^*, \mu \in \mathfrak{g}^*$, and $A \in \mathfrak{g}$. If, moreover,

$$\Upsilon_g(\eta_1 \eta_2) = \Upsilon_g(\eta_1) \Upsilon_{\Upsilon_{\eta_1^{-1}}(g)}(\eta_2), \quad \Upsilon_\eta^*(g_1 g_2) = \Upsilon_{\Upsilon_{g_2}(\eta^{-1})}^*(g_1) \Upsilon_\eta^*(g_2) \quad (2.31)$$

for all $g_1, g_2 \in G$ and $\eta_1, \eta_2 \in G^*$, then we say that the tuple $(G, G^*, \Upsilon, \Upsilon^*)$ (or shortly (Υ, Υ^*)) is a *coconjugation matched pair*. \diamond

A pair of groups endowed with two actions on each other that are compatible in the sense of (2.31) is called a *matched pair of groups*. The notion of a matched pair of groups appears in [Tak81, Section 2] and has been connected to the theory of Poisson Lie groups in [LW90; Maj90]. The compatibility condition (2.31) allows to define a (non-trivial) Lie group structure on the product $G \times G^*$ by

$$(g_1, \eta_1) \cdot (g_2, \eta_2) = (g_1 \Upsilon_{\eta_1}^*(g_2^{-1})^{-1}, \Upsilon_{g_2^{-1}}(\eta_1^{-1})^{-1} \eta_2), \quad (2.32)$$

see [Tak81, Proposition 2.2]. Conversely, every factorization $D = GG^*$ of a Lie group D into two Lie subgroups G and G^* with $G \cap G^* = \{e\}$ yields actions of G and G^* on each other that satisfy (2.31) by declaring

$$\eta g^{-1} = (\Upsilon_\eta^*(g))^{-1} (\Upsilon_g(\eta^{-1}))^{-1} \in GG^*. \quad (2.33)$$

The class of coconjugation matched pairs is rather rigid, especially if one of the coconjugation actions is predetermined.

LEMMA 2.19 Let $\kappa(G, G^*)$ be a dual pair of Lie groups with a matched pair (Υ, Υ^*) . If $\tilde{\Upsilon}: G \times G^* \rightarrow G^*$ is another action of G on G^* such that $(\tilde{\Upsilon}, \Upsilon^*)$ forms a coconjugation

¹ In infinite dimensions, κ is, in general, only weakly non-degenerate and thus this relation only ensures uniqueness of Ad^* but not its existence. In the sequel, we assume that Ad^* exists.

matched pair, then there exists a map $c: G \times \pi_0(G^*) \rightarrow G^*$ with $c(g, [e]) = e$ such that $\tilde{\Upsilon}(g, \eta) = c(g, [\eta]) \Upsilon(g, \eta)$. In particular, if G^* is connected, then coconjugation actions Υ compatible with Υ^* are unique. \diamond

Proof. Using the defining properties (2.30) and (2.31), for $g \in G$, $\eta \in G^*$ and $\mu \in \mathfrak{g}^*$, we calculate

$$T_\eta \Upsilon_g(\eta \cdot \mu) = \left. \frac{d}{d\varepsilon} \right|_0 \Upsilon_g(\eta \mu_\varepsilon) = \left. \frac{d}{d\varepsilon} \right|_0 \Upsilon_g(\eta) \Upsilon_{\Upsilon_{\eta^{-1}}(g)}(\mu_\varepsilon) = \Upsilon_g(\eta) \cdot (\text{Ad}_{\Upsilon_{\eta^{-1}}(g)}^* \mu), \quad (2.34)$$

where $\varepsilon \mapsto \mu_\varepsilon$ is a curve in G^* starting at the identity with $\left. \frac{d}{d\varepsilon} \right|_0 \mu_\varepsilon = \mu$. A similar result holds, of course, also for Υ replaced by $\tilde{\Upsilon}$. In particular, the left-logarithmic derivatives of $\Upsilon_g: G^* \rightarrow G^*$ and $\tilde{\Upsilon}_g: G^* \rightarrow G^*$ coincide for every $g \in G$. The uniqueness lemma for Lie group-valued functions [Nee06, Remark II.3.6.(c)] hence implies that on every connected component of G^* the maps Υ_g and $\tilde{\Upsilon}_g$ differ by a constant. Thus, there exists a map $c: G \times \pi_0(G^*) \rightarrow G^*$ such that $\tilde{\Upsilon}(g, \eta) = c(g, [\eta]) \Upsilon(g, \eta)$. Finally, $\tilde{\Upsilon}(g, e) = e = \Upsilon(g, e)$ implies $c(g, [e]) = e$. \square

EXAMPLE 2.20 Let G be a Lie group with Lie algebra \mathfrak{g} and let $\kappa(\mathfrak{g}, \mathfrak{g}^*)$ be a dual pair of (locally convex) vector spaces. Considering \mathfrak{g}^* as an Abelian Lie group, we obtain the dual pair $\kappa(G, \mathfrak{g}^*)$ of Lie groups. Then the coadjoint representation $\Upsilon(g, \mu) = \text{Ad}_{g^{-1}}^* \mu$ and the trivial action $\Upsilon^*(\mu, g) = g$ form a coconjugation matched pair on $\kappa(G, \mathfrak{g}^*)$. \diamond

Since \mathfrak{g}^* is connected, the previous proposition implies that the coadjoint representation is the only coconjugation action compatible with the trivial action of \mathfrak{g}^* on G . However, there are interesting examples where the dual group does not act trivially.

EXAMPLE 2.21 Let D be a finite-dimensional connected complex semisimple Lie group and let K be its compact real form. Consider the Iwasawa decomposition $D = KAN$ and set $B = AN$. Then B is a (solvable) Lie subgroup of D with Lie algebra \mathfrak{b} . Let $\text{pr}_K: D \rightarrow K$ and $\text{pr}_B: D \rightarrow B$ be the natural projections, and $\text{pr}_{\mathfrak{k}}: \mathfrak{d} \rightarrow \mathfrak{k}$ and $\text{pr}_{\mathfrak{b}}: \mathfrak{d} \rightarrow \mathfrak{b}$ be the associated projections on the level of Lie algebras. By (2.33), the decomposition $D = KB$ induces the following actions:

$$\Upsilon_k(b) = \text{pr}_B(b^{-1}k^{-1})^{-1}, \quad \Upsilon_b^*(k) = \text{pr}_K(bk^{-1})^{-1}. \quad (2.35)$$

Going back to the construction of the Iwasawa decomposition, one can show that \mathfrak{k} and \mathfrak{b} are isotropic subspaces of \mathfrak{d} relative to $\text{Im } \kappa_{\mathfrak{d}}$, see [LW90, p. 520]. Thus, we obtain a dual pair $\kappa(K, B)$ of Lie groups. We have $T_e \Upsilon_k(\mu) = \text{pr}_{\mathfrak{b}}(\text{Ad}_k \mu)$ and $T_e \Upsilon_b^*(A) = \text{pr}_{\mathfrak{k}}(\text{Ad}_b A)$ for $\mu \in \mathfrak{b}$ and $A \in \mathfrak{k}$. These identities show that the actions integrate the coadjoint actions and thus (Υ, Υ^*) is a coconjugation action on

$\kappa(K, B)$. As $D = KB$ is a decomposition in subgroups, (Y, Y^*) is a coconjugation matched pair. \diamond

Recall that, in the classical setting $G^* \equiv \mathfrak{g}^*$, the coadjoint action is closely connected to the natural Poisson structure of Kostant–Kirillov–Souriau. Moreover, the Cartan and Iwasawa decompositions are well-known constructions in the theory of Poisson Lie groups. Thus, one would expect a close interplay between coconjugation actions and Poisson structures. As we will see, this is indeed the case.

Poisson structures in infinite dimensions need to be treated with caution. For example, according to [BGT18] not every Poisson bracket is induced by a Poisson bivector field. We refer the reader to [NST14] for a recent discussion of the problems concerning infinite-dimensional Poisson geometry. In the present case, we can use the group structure to circumvent most of these issues. Let us start by recalling the necessary terminology and notation from the finite-dimensional theory of Poisson Lie groups. Let H be a finite-dimensional Lie group, \mathfrak{h} be its Lie algebra with dual \mathfrak{h}^* . We recall that for a smooth map $F: M \rightarrow H$ the *left logarithmic derivative* is the \mathfrak{h} -valued 1-form δF on M defined by $\delta_m F(X) = F(m)^{-1} \cdot T_m F(X)$ for $X \in T_m M$. In the following, we also need the concept of the *left derivative*. This time, the domain of the map is a Lie group, i.e., we are considering a smooth map $f: H \rightarrow N$. The left derivative $T_h^L f: \mathfrak{h} \rightarrow T_{f(h)} N$ of f at $h \in H$ in the direction $A \in \mathfrak{h}$ is defined by

$$T_h^L f(A) := T_h f(h \cdot A), \quad (2.36)$$

that is, as the usual derivative $Tf: TH \rightarrow TN$ precomposed with the left trivialization $H \times \mathfrak{h} \simeq TH$. In particular, for a real-valued function $f: H \rightarrow \mathbb{R}$, the left derivative $T_h^L f: \mathfrak{h} \rightarrow \mathbb{R}$ can be viewed as an element of \mathfrak{h}^* .

In finite dimensions, a Poisson Lie group is a Lie group H endowed with a Poisson structure $\omega_H \in \mathfrak{X}^2(H)$ such that the multiplication $H \times H \rightarrow H$ is a Poisson map (with respect to the product Poisson structure on $H \times H$). By [Lu90, Theorem 2.18], the linearization of ω_H at the identity $e \in H$ induces a Lie bracket on \mathfrak{h}^* by

$$[A, B]_{\omega_H} = d_e(\omega_H(\bar{A}, \bar{B})), \quad (2.37)$$

where $A, B \in \mathfrak{h}^*$ and \bar{A}, \bar{B} are arbitrary 1-forms on H with $\bar{A}_e = A$ and $\bar{B}_e = B$. In the left-trivialization $TH \simeq H \times \mathfrak{h}$, the bivector field ω_H is identified with a smooth map $\Lambda_H: H \rightarrow \Lambda^2 \mathfrak{h}$ and we have $[A, B]_{\omega_H} = T_e(\Lambda_H(\cdot)(A, B))$.

Let $\kappa(\mathfrak{g}, \mathfrak{h})$ be a dual pair of finite-dimensional Lie algebras. For every \mathfrak{g} -module M , consider the Chevalley–Eilenberg complex $\text{Hom}(\Lambda^\bullet \mathfrak{g}, M)$ consisting

of continuous alternating multilinear maps $\lambda: \Lambda^k \mathfrak{g} \rightarrow M$ with differential

$$\begin{aligned} d^{\text{CE}} \lambda(A_1, \dots, A_{k+1}) &= \sum_i (-1)^{i+1} A_i \cdot \lambda(A_1, \dots, \hat{i} \dots A_{k+1}) \\ &\quad + \sum_{i < j} (-1)^{i+j} \lambda([A_i, A_j], A_1, \dots, \hat{i} \dots \hat{j} \dots A_{k+1}). \end{aligned} \quad (2.38)$$

In particular, for $M = \mathfrak{h}$ with (minus) the coadjoint action, we have

$$\begin{aligned} d_{\mathfrak{h}}^{\text{CE}} \lambda(A_1, \dots, A_{k+1}) &= \sum_i (-1)^i \text{ad}_{A_i}^* \lambda(A_1, \dots, \hat{i} \dots A_{k+1}) \\ &\quad + \sum_{i < j} (-1)^{i+j} \lambda([A_i, A_j], A_1, \dots, \hat{i} \dots \hat{j} \dots A_{k+1}). \end{aligned} \quad (2.39)$$

Moreover, using the identification $\text{Hom}(\Lambda^\bullet \mathfrak{g}, \mathbb{R}) \simeq \Lambda^\bullet \mathfrak{g}^* \simeq \Lambda^\bullet \mathfrak{h}$, we can transfer the Chevalley–Eilenberg differential to $\Lambda^\bullet \mathfrak{h}$. Explicitly, it is given by

$$\begin{aligned} \kappa(A_1 \wedge \dots \wedge A_{k+1}, d_{\mathfrak{g}}^{\text{CE}} \nu) \\ = \sum_{i < j} (-1)^{i+j} \kappa([A_i, A_j]_{\mathfrak{g}} \wedge A_1 \wedge \dots \wedge \hat{i} \dots \hat{j} \dots \wedge A_{k+1}, \nu) \end{aligned} \quad (2.40)$$

for every $\nu \in \Lambda^k \mathfrak{h}$ and $A_i \in \mathfrak{g}$. Moreover, given $T \in \text{Hom}(\mathfrak{h}, \Lambda^2 \mathfrak{h}) \simeq \mathfrak{g} \otimes \Lambda^2 \mathfrak{h}$ and $\beta \in \Lambda^2 \mathfrak{h}$, we define the contraction $T \lrcorner \beta \in \Lambda^3 \mathfrak{h}$ by

$$(A \otimes \alpha) \lrcorner \beta := \alpha \wedge (A \lrcorner \beta) = (A \lrcorner \beta) \wedge \alpha \quad (2.41)$$

for $A \in \mathfrak{g}$ and $\alpha \in \Lambda^2 \mathfrak{h}$. Finally, recall that the (algebraic) Schouten–Nijenhuis bracket $[\cdot, \cdot]: \Lambda^k \mathfrak{h} \times \Lambda^l \mathfrak{h} \rightarrow \Lambda^{k+l-1} \mathfrak{h}$ is defined by

$$[v_1 \wedge \dots \wedge v_k, \mu_1 \wedge \dots \wedge \mu_l] = \sum_{i,j} (-1)^{i+j} [v_i, \mu_j]_{\mathfrak{h}} \wedge v_1 \wedge \dots \wedge \hat{i} \dots \wedge v_k \wedge \mu_1 \wedge \dots \wedge \hat{j} \dots \wedge \mu_l. \quad (2.42)$$

Let us introduce a related bracket as follows. For linear maps $\phi: \Lambda^k \mathfrak{g} \rightarrow \mathfrak{h}$ and $\varphi: \Lambda^l \mathfrak{g} \rightarrow \mathfrak{h}$, define $[\phi, \varphi]: \Lambda^{k+l} \mathfrak{g} \rightarrow \mathfrak{h}$ by

$$\begin{aligned} [\phi, \varphi](A) &= \sum_{\sigma} (-1)^{\text{sgn} \sigma} \left(\phi(\text{ad}_{\varphi(\sigma_l(A))}^* \sigma_k(A)) - \varphi(\text{ad}_{\phi(\sigma_k(A))}^* \sigma_l(A)) \right. \\ &\quad \left. + [\phi(\sigma_k(A)), \varphi(\sigma_l(A))]_{\mathfrak{h}} \right), \end{aligned} \quad (2.43)$$

where the sum is over (unordered) partitions σ of $A \in \Lambda^{k+l} \mathfrak{g}$ in pairs $\sigma_k(A) \in \Lambda^k \mathfrak{g}$

and $\sigma_l(A) \in \Lambda^l \mathfrak{g}$. In particular, for a linear map $\phi: \mathfrak{g} \rightarrow \mathfrak{h}$, we get

$$\frac{1}{2}[\phi, \phi](A, B) = \phi(\text{ad}_{\phi(A)}^* B) - \phi(\text{ad}_{\phi(B)}^* A) + [\phi(A), \phi(B)]_{\mathfrak{h}} \quad (2.44)$$

for all $A, B \in \mathfrak{g}$.

The following result is the first step towards a reformation of the notion of a Poisson Lie structure that is suitable for the infinite-dimensional setting.

PROPOSITION 2.22 *Let $\kappa(G, H)$ be a dual pair of finite-dimensional Lie groups. Let $\Lambda_H: H \rightarrow \Lambda^2 \mathfrak{h}$ be a smooth map and let $\omega_H \in \mathfrak{X}^2(H)$ be the associated bivector field. Then ω_H is a Poisson Lie structure on H whose linearization $[\cdot, \cdot]_{\omega_H}$ coincides with the Lie bracket on $\mathfrak{g} \simeq \mathfrak{h}^*$ if and only if Λ_H possesses the following properties:*

(i) (multiplicativity) For all $\eta, \zeta \in H$,

$$\Lambda_H(\eta \zeta) = \text{Ad}_{\zeta^{-1}} \Lambda_H(\eta) + \Lambda_H(\zeta), \quad (2.45)$$

where Ad denotes the natural extension to $\Lambda^2 \mathfrak{h}$ of the adjoint action of H on \mathfrak{h} .

(ii) (Poisson property)

$$d_{\mathfrak{g}}^{\text{CE}} \Lambda_H - \frac{1}{2}[\Lambda_H, \Lambda_H] = 0. \quad (2.46)$$

(iii) (compatibility) For all $\mu \in \mathfrak{h}$ and $A, B \in \mathfrak{g}$,

$$\kappa(A \wedge B, T_e \Lambda_H(\mu)) = \kappa([A, B]_{\mathfrak{g}}, \mu). \quad (2.47)$$

◇

Note that the Poisson property is reformulated in terms of the purely algebraic condition (2.46). For the particular case where H is Abelian, (2.46) holds if and only if $\Lambda_H(\eta)$ is a 2-cocycle on \mathfrak{g} for all $\eta \in H$.

Proof. By [Ale+98, Lemma 4.1], multiplication $H \times H \rightarrow H$ is a Poisson map if and only if Λ_H is multiplicative in the sense that (2.45) holds.

If $\bar{A}, \bar{B} \in \Omega^1(H)$ are left-invariant extensions of $A, B \in \mathfrak{g} \simeq \mathfrak{h}^*$, then by definition $(\omega_H)_{\eta}(\bar{A}, \bar{B}) = \Lambda_H(\eta)(A, B)$ for every $\eta \in H$. Thus, the Lie bracket on \mathfrak{g} induced by ω_H is given by

$$\kappa([A, B]_{\omega_H}, \mu) = \kappa(d_e(\omega_H(\bar{A}, \bar{B})), \mu) = \kappa(A \wedge B, T_e \Lambda_H(\mu)). \quad (2.48)$$

Thus, $[\cdot, \cdot]_{\omega_H}$ coincides with the original Lie bracket on \mathfrak{g} if and only if (2.47) holds.

By [Ale+98, Proposition 3.4], the Schouten–Nijenhuis bracket of ω_H with itself vanishes if and only if

$$(T^L \Lambda_H) \lrcorner \Lambda_H = \frac{1}{2}[\Lambda_H, \Lambda_H], \quad (2.49)$$

where, for every $\eta \in H$, the left derivative $T_\eta^L \Lambda_H$ is an element of $\text{Hom}(\mathfrak{h}, \Lambda^2 \mathfrak{h})$. By differentiating (2.45), we obtain

$$T_\eta^L \Lambda_H(\mu) = T_\eta \Lambda_H(\eta \cdot \mu) = -\text{ad}_\mu \Lambda_H(\eta) + T_e \Lambda_H(\mu). \quad (2.50)$$

Using the compatibility relation (2.47), Lemmas 2.23 and 2.24 below imply that

$$(T^L \Lambda_H) \lrcorner \Lambda_H = [\Lambda_H, \Lambda_H] - d_{\mathfrak{g}}^{\text{CE}} \Lambda_H. \quad (2.51)$$

Thus, (2.49) is equivalent to (2.46). \square

LEMMA 2.23 *Let $\kappa(\mathfrak{g}, \mathfrak{h})$ be a dual pair of finite-dimensional Lie algebras. For every $\Lambda \in \Lambda^2 \mathfrak{h}$, the operator $T: \mathfrak{h} \rightarrow \Lambda^2 \mathfrak{h}$ defined by $T(\mu) = \text{ad}_\mu \Lambda$ satisfies*

$$T \lrcorner \Lambda = -[\Lambda, \Lambda]. \quad (2.52)$$

\diamond

Proof. Choose a basis $\{e_i\}$ in \mathfrak{g} and the dual basis $\{\varepsilon^i\}$ in \mathfrak{h} . Write

$$T = e_i \otimes \text{ad}_{e_i} \Lambda \quad \text{and} \quad \Lambda = \frac{1}{2} \Lambda_{kl} \varepsilon^k \wedge \varepsilon^l. \quad (2.53)$$

Then, we have

$$e_i \lrcorner \Lambda = \frac{1}{2} \Lambda_{kl} (\delta_i^k \varepsilon^l - \delta_i^l \varepsilon^k) = \Lambda_{il} \varepsilon^l \quad (2.54)$$

and

$$\text{ad}_{e_i} \Lambda = \frac{1}{2} \Lambda_{kl} ([\varepsilon^i, \varepsilon^k]_{\mathfrak{h}} \wedge \varepsilon^l + \varepsilon^k \wedge [\varepsilon^i, \varepsilon^l]_{\mathfrak{h}}) = \Lambda_{kl} [\varepsilon^i, \varepsilon^k]_{\mathfrak{h}} \wedge \varepsilon^l. \quad (2.55)$$

Thus, using (2.41), we obtain

$$T \lrcorner \Lambda = \text{ad}_{e_i} \Lambda \wedge (e_i \lrcorner \Lambda) = \Lambda_{kl} \Lambda_{ij} [\varepsilon^i, \varepsilon^k]_{\mathfrak{h}} \wedge \varepsilon^l \wedge \varepsilon^j. \quad (2.56)$$

On the other hand, we find

$$\begin{aligned} [\Lambda, \Lambda] &= \frac{1}{4} \Lambda_{kl} \Lambda_{ij} [\varepsilon^k \wedge \varepsilon^l, \varepsilon^i \wedge \varepsilon^j] \\ &= \frac{1}{4} \Lambda_{kl} \Lambda_{ij} ([\varepsilon^k, \varepsilon^i]_{\mathfrak{h}} \wedge \varepsilon^l \wedge \varepsilon^j - [\varepsilon^k, \varepsilon^j]_{\mathfrak{h}} \wedge \varepsilon^l \wedge \varepsilon^i \\ &\quad - [\varepsilon^l, \varepsilon^i]_{\mathfrak{h}} \wedge \varepsilon^k \wedge \varepsilon^j + [\varepsilon^l, \varepsilon^j]_{\mathfrak{h}} \wedge \varepsilon^k \wedge \varepsilon^i) \\ &= \Lambda_{kl} \Lambda_{ij} [\varepsilon^k, \varepsilon^i]_{\mathfrak{h}} \wedge \varepsilon^l \wedge \varepsilon^j. \end{aligned} \quad (2.57)$$

Comparing the last two identities yields the claim. \square

LEMMA 2.24 *Let $\kappa(\mathfrak{g}, \mathfrak{h})$ be a dual pair of finite-dimensional Lie algebras. For every*

$\Lambda \in \Lambda^2 \mathfrak{h}$,

$$\omega \lrcorner \Lambda = -d_{\mathfrak{g}}^{\text{CE}} \Lambda, \quad (2.58)$$

where the operator $\omega: \mathfrak{h} \rightarrow \Lambda^2 \mathfrak{h}$ is defined by $\kappa(A \wedge B, \omega(\mu)) = \kappa([A, B]_{\mathfrak{g}}, \mu)$ for all $A, B \in \mathfrak{g}$. \diamond

Proof. Fix a basis $\{e_k\}$ of \mathfrak{g} . Denote by $[\cdot, \cdot]^k \in \Lambda^2 \mathfrak{h}$ the k -th component of the commutator, i.e., $[A, B]^k e_k = [A, B]_{\mathfrak{g}}$ for all $A, B \in \mathfrak{g}$. With this notation, we have $\omega = e_k \otimes [\cdot, \cdot]^k$, because

$$\kappa(A \wedge B, \omega(\mu)) = \kappa([A, B]_{\mathfrak{g}}, \mu) = \kappa(e_k, \mu) [A, B]^k. \quad (2.59)$$

Thus we find

$$\begin{aligned} (\omega \lrcorner \Lambda)(A, B, C) &= ((e_k \otimes [\cdot, \cdot]^k) \lrcorner \Lambda)(A, B, C) \\ &= ((e_k \lrcorner \Lambda) \wedge [\cdot, \cdot]^k)(A, B, C) \\ &= \Lambda(e_k, A) [B, C]^k - \Lambda(e_k, B) [A, C]^k + \Lambda(e_k, C) [A, B]^k \\ &= \Lambda([B, C], A) - \Lambda([A, C], B) + \Lambda([A, B], C) \\ &= -\Lambda(A, [B, C]) + \Lambda(B, [A, C]) - \Lambda(C, [A, B]) \\ &= -d_{\mathfrak{g}}^{\text{CE}} \Lambda(A, B, C). \quad \square \end{aligned}$$

Proposition 2.22 yields a convenient reformulation of what it means for a bivector field $\omega_H \in \mathfrak{X}^2(H)$ to be a Poisson Lie structure on H in terms of the associated map $\Lambda_H: H \rightarrow \Lambda^2 \mathfrak{h}$. In the sequel, it will be convenient to view the latter as a map

$$\pi_H: H \times \mathfrak{g} \rightarrow \mathfrak{h}, \quad (2.60)$$

that is, $\Lambda_H(\eta)(A, B) = \kappa(B, \pi_H(\eta, A))$. From this viewpoint, we have the following equivalent description of a Poisson Lie structure on H .

PROPOSITION 2.25 *Let $\kappa(G, H)$ be a dual pair of finite-dimensional Lie groups. Let $\Lambda_H: H \rightarrow \Lambda^2 \mathfrak{h}$ be a smooth map satisfying the properties of Proposition 2.22 (so that its extension $\omega_H \in \mathfrak{X}^2(H)$ is a Poisson Lie structure on H). Then the map $\pi_H: H \times \mathfrak{g} \rightarrow \mathfrak{h}$ defined by $\pi_H(\eta, A) = A \lrcorner \Lambda_H$ satisfies the following conditions:*

(i) *(linearity and skew-symmetry) π_H is linear in the second argument and satisfies*

$$\kappa(A, \pi_H(\eta, B)) = -\kappa(B, \pi_H(\eta, A)) \quad (2.61)$$

for all $A, B \in \mathfrak{g}$ and $\eta \in H$.

(ii) *(multiplicativity) For all $\eta, \zeta \in H$ and $A \in \mathfrak{g}$,*

$$\pi_H(\eta \zeta, A) = \text{Ad}_{\zeta^{-1}} \pi_H(\eta, \text{Ad}_{\zeta^{-1}}^* A) + \pi_H(\zeta, A). \quad (2.62)$$

(iii) (Poisson property)

$$d_{\mathfrak{h}}^{\text{CE}} \pi_H - \frac{1}{2}[\pi_H, \pi_H] = 0. \quad (2.63)$$

(iv) (compatibility) For all $\mu \in \mathfrak{h}$ and $A \in \mathfrak{g}$,

$$T_e(\pi_H(\cdot, A))(\mu) = \text{ad}_A^* \mu, \quad (2.64)$$

where $\pi_H(\cdot, A): H \rightarrow \mathfrak{h}$.

Conversely, to every map $\pi_H: H \times \mathfrak{g} \rightarrow \mathfrak{h}$ satisfying these conditions, the map $\Lambda_H: H \rightarrow \Lambda^2 \mathfrak{h}$ defined by $\pi_H(\eta, A) = A \lrcorner \Lambda_H$ posses the properties of Proposition 2.22. \diamond

Proof. Linearity in the second argument and skew-symmetry of π_H is clear as Λ_H takes values in $\Lambda^2 \mathfrak{h}$. Moreover, the multiplicativity and compatibility conditions are simple reformulations of the corresponding properties in Proposition 2.22.

Thus, it is left to show that (2.46) and (2.63) are equivalent. Since this a completely algebraic question (for fixed $\eta \in H$), it suffices to consider a linear map $\pi: \mathfrak{g} \rightarrow \mathfrak{h}$ with associated $\Lambda \in \Lambda^2 \mathfrak{h}$. First note that we have

$$\begin{aligned} \kappa(C, d_{\mathfrak{h}}^{\text{CE}} \pi(A, B)) &= \kappa(C, -\text{ad}_A^* \pi(B) + \text{ad}_B^* \pi(A) - \pi([A, B])) \\ &= -\kappa([A, C], \pi(B)) + \kappa([B, C], \pi(A)) - \kappa(C, \pi([A, B])) \\ &= \kappa([A, C] \wedge B - [B, C] \wedge A - [A, B] \wedge C, \Lambda) \\ &= \kappa(A \wedge B \wedge C, d_{\mathfrak{g}}^{\text{CE}} \Lambda). \end{aligned} \quad (2.65)$$

Choose a basis $\{e_i\}$ in \mathfrak{g} and the dual basis $\{\varepsilon^i\}$ in \mathfrak{h} . According to (2.57), we have $[\Lambda, \Lambda] = \Lambda_{kl} \Lambda_{pq} [\varepsilon^k, \varepsilon^p]_r \varepsilon^r \wedge \varepsilon^l \wedge \varepsilon^q$, where $\Lambda = \frac{1}{2} \Lambda_{kl} \varepsilon^k \wedge \varepsilon^l$ and $[\varepsilon^k, \varepsilon^p]_r \varepsilon^r = [\varepsilon^k, \varepsilon^p]_{\mathfrak{h}}$. Thus,

$$\begin{aligned} e_j \lrcorner e_i \lrcorner [\Lambda, \Lambda] &= \Lambda_{kl} \Lambda_{pq} [\varepsilon^k, \varepsilon^p]_r (\delta_i^r \delta_j^l \varepsilon^q + \delta_i^l \delta_j^q \varepsilon^r + \delta_i^q \delta_j^r \varepsilon^l \\ &\quad - \delta_i^r \delta_j^q \varepsilon^l - \delta_i^q \delta_j^l \varepsilon^r - \delta_i^l \delta_j^r \varepsilon^q) \\ &= 2 \Lambda_{kl} \Lambda_{pq} [\varepsilon^k, \varepsilon^p]_r (\delta_i^r \delta_j^l \varepsilon^q + \delta_i^l \delta_j^q \varepsilon^r + \delta_i^q \delta_j^r \varepsilon^l) \\ &= 2(\Lambda_{kj} \Lambda_{lr} [\varepsilon^k, \varepsilon^l]_i + \Lambda_{ki} \Lambda_{lj} [\varepsilon^k, \varepsilon^l]_r + \Lambda_{kr} \Lambda_{li} [\varepsilon^k, \varepsilon^l]_j) \varepsilon^r, \end{aligned} \quad (2.66)$$

because $\Lambda_{kl} \Lambda_{pq} [\varepsilon^k, \varepsilon^p]_r$ is skew-symmetric in the indices l and q . On the other hand, using (2.44), the identities $\pi(e_i) = e_i \lrcorner \Lambda = \Lambda_{il} \varepsilon^l$ and $\text{ad}_{\varepsilon^l}^* e_i = [\varepsilon^l, \varepsilon^k]_i e_k$

imply

$$\begin{aligned} \frac{1}{2}[\pi, \pi](e_i, e_j) &= \pi(\text{ad}_{\pi(e_i)}^* e_j) - \pi(\text{ad}_{\pi(e_j)}^* e_i) + [\pi(e_i), \pi(e_j)]_{\mathfrak{h}} \\ &= \Lambda_{il}\Lambda_{kr}[\varepsilon^l, \varepsilon^k]_j \varepsilon^r - \Lambda_{jl}\Lambda_{kr}[\varepsilon^l, \varepsilon^k]_i \varepsilon^r + \Lambda_{il}\Lambda_{jk}[\varepsilon^l, \varepsilon^k]_r \varepsilon^r. \end{aligned} \quad (2.67)$$

In summary, we obtain

$$\kappa\left(A \wedge B \wedge C, d_{\mathfrak{g}}^{\text{CE}} \Lambda - \frac{1}{2}[\Lambda, \Lambda]\right) = \kappa\left(C, d_{\mathfrak{h}}^{\text{CE}} \pi(A, B) - \frac{1}{2}[\pi, \pi](A, B)\right), \quad (2.68)$$

so that (2.46) and (2.63) are equivalent indeed. \square

It is this form that we adopt as the definition of a Poisson Lie structure in the infinite-dimensional setting.

DEFINITION 2.26 Let $\kappa(G, H)$ be a dual pair of (possibly infinite-dimensional) Lie groups. A smooth map $\pi_H: H \times \mathfrak{g} \rightarrow \mathfrak{h}$ is said to be a *Poisson Lie structure on H compatible with the Lie bracket on \mathfrak{g}* if it satisfies the properties stated in Proposition 2.25. \diamond

In [NST14] it has been observed that a Poisson tensor on an infinite-dimensional manifold does *not* induce a Poisson bracket on the space of all smooth functions. This continues to be true in our particular setting of Poisson Lie structures. Indeed, a Poisson Lie structure π_H on H defines a Lie bracket

$$\{f, g\}(\eta) = \kappa(T_{\eta}^{\text{L}} f, \pi_H(\eta, T_{\eta}^{\text{L}} g)) \quad (2.69)$$

only on the subalgebra of $C^{\infty}(H)$ consisting of functions f for which the functional $T^{\text{L}} f: \mathfrak{h} \rightarrow \mathbb{R}$ can be represented by an element of \mathfrak{g} . Conversely, we note that, in infinite dimensions, the Poisson bracket on the space of smooth functions may depend on higher-order derivatives of functions and thus may not be induced by a Poisson tensor, see [BGT18].

After this excursion into the theory of infinite-dimensional Poisson Lie groups, let us return to our original quest of establishing a relation between coconjugation actions and induced Poisson Lie structures. In finite dimensions, every Poisson Lie group with complete dressing vector fields can be integrated to matched pair of Lie groups, see [LW90, Theorem 3.12]. The following result establishes the corresponding inverse “derivation” operation.

THEOREM 2.27 Let $\kappa(G, G^*)$ be a dual pair of Lie groups. For every coconjugation matched pair (Υ, Υ^*) on $\kappa(G, G^*)$, define $\pi_G: G \times \mathfrak{g}^* \rightarrow \mathfrak{g}$ and $\pi_{G^*}: G^* \times \mathfrak{g} \rightarrow \mathfrak{g}^*$ by

$$\pi_G(g, \mu) := T_e \Upsilon_{g^{-1}}^* (\text{Ad}_{g^{-1}}^* \mu) \cdot g, \quad \pi_{G^*}(\eta, A) := -\eta^{-1} \cdot T_e \Upsilon_{\eta} (\text{Ad}_{\eta^{-1}}^* A). \quad (2.70)$$

If π_G and π_{G^*} are skew-symmetric in the sense of (2.61), then they define Poisson Lie structures on G and on G^* , respectively. \diamond

Proof. We will give the proof only for π_{G^*} as similar arguments show that π_G is a Poisson Lie structure as well. We have to verify that π_{G^*} satisfies the conditions of Proposition 2.25.

First, note that π_{G^*} is clearly linear in the second argument and it is skew-symmetric by assumption.

The compatibility condition (2.31) implies

$$\begin{aligned}
 \pi_{G^*}(\eta \zeta, A) &= -(\zeta^{-1} \eta^{-1}) \cdot \mathrm{T}_e \Upsilon_{\eta \zeta}(\mathrm{Ad}_{\eta^{-1}}^* \mathrm{Ad}_{\zeta^{-1}}^* A) \\
 &= -(\zeta^{-1} \eta^{-1}) \cdot \mathrm{T}_e (\Upsilon_{\eta} \cdot (\Upsilon_{\zeta} \circ \Upsilon_{\eta^{-1}}^*))(\mathrm{Ad}_{\eta^{-1}}^* \mathrm{Ad}_{\zeta^{-1}}^* A) \\
 &= -\mathrm{Ad}_{\zeta^{-1}}(\eta^{-1} \cdot \mathrm{T}_e \Upsilon_{\eta}(\mathrm{Ad}_{\eta^{-1}}^* \mathrm{Ad}_{\zeta^{-1}}^* A)) \\
 &\quad - \zeta^{-1} \cdot \mathrm{T}_e \Upsilon_{\zeta}(\mathrm{T}_e \Upsilon_{\eta^{-1}}^*(\mathrm{Ad}_{\eta^{-1}}^* \mathrm{Ad}_{\zeta^{-1}}^* A)) \\
 &= \mathrm{Ad}_{\zeta^{-1}}(\pi_{G^*}(\eta, \mathrm{Ad}_{\zeta^{-1}}^* A)) + \pi_{G^*}(\zeta, A).
 \end{aligned} \tag{2.71}$$

This verifies the multiplicativity of π_{G^*} , cf. (2.62).

Let us now turn to the compatibility relation (2.64). Note that we have $(\delta \Upsilon)_{(e, \eta)}(A, o) = -\pi_{G^*}(\eta, \mathrm{Ad}_{\eta}^* A)$ and $(\delta \Upsilon)_{(g, \eta)}(o, \eta \cdot \mu) = \mathrm{Ad}_{\Upsilon_{\eta^{-1}}^*(g)^{-1}}^* \mu$, see (2.34). Since the 1-form $\delta \Upsilon \in \Omega^1(G \times G^*, \mathfrak{g}^*)$ satisfies the Maurer–Cartan equation, we thus obtain

$$\begin{aligned}
 o &= \left(d \delta \Upsilon + \frac{1}{2} [\delta \Upsilon, \delta \Upsilon] \right)_{(e, e)}((o, \mu), (A, o)) \\
 &= (o, \bar{\mu})_{(e, e)}(\delta \Upsilon(\bar{A}, o)) - (\bar{A}, o)_{(e, e)}(\delta \Upsilon(o, \bar{\mu})) \\
 &\quad + [(\delta \Upsilon)_{(e, e)}(o, \mu), (\delta \Upsilon)_{(e, e)}(A, o)] \\
 &= -\mathrm{T}_e(\eta \mapsto \pi_{G^*}(\eta, \mathrm{Ad}_{\eta}^* A))(\mu) - \mathrm{T}_e(g \mapsto \mathrm{Ad}_{g^{-1}}^* \mu)(A) + o \\
 &= -\mathrm{T}_e(\pi_{G^*}(\cdot, A))(\mu) + \mathrm{ad}_A^* \mu,
 \end{aligned} \tag{2.72}$$

where \bar{A} and $\bar{\mu}$ denote the left-invariant extensions of A and μ , respectively. Thus, $\mathrm{T}_e(\pi_{G^*}(\cdot, A))(\mu) = \mathrm{ad}_A^* \mu$, which is the compatibility relation (2.64).

For the Poisson property (2.63), we need a bit of preparation. First, the fact that Υ is a left G -action implies

$$\begin{aligned}
 \delta_g \Upsilon_{\eta}(g \cdot A) &= \Upsilon(g, \eta)^{-1} \cdot \mathrm{T}_{\eta} \Upsilon_g(\mathrm{T}_e \Upsilon_{\eta}(A)) \\
 &= -\Upsilon(g, \eta)^{-1} \cdot \mathrm{T}_{\eta} \Upsilon_g(\eta \cdot \pi_{G^*}(\eta, \mathrm{Ad}_{\eta}^* A)) \\
 &= -\mathrm{Ad}_{\Upsilon_{\eta^{-1}}^*(g)^{-1}}^* \pi_{G^*}(\eta, \mathrm{Ad}_{\eta}^* A),
 \end{aligned} \tag{2.73}$$

where, for the last equality, we used (2.34). Since $\delta \Upsilon_{\eta} \in \Omega^1(G, \mathfrak{g}^*)$ satisfies the

Maurer–Cartan equation for every $\eta \in G^*$, we thus find

$$\begin{aligned}
\circ &= \left(d \delta \Upsilon_\eta + \frac{1}{2} [\delta \Upsilon_\eta, \delta \Upsilon_\eta] \right)_e (A, B) \\
&= \bar{A}_e (\delta \Upsilon_\eta(\bar{B})) - \bar{B}_e (\delta \Upsilon_\eta(\bar{A})) - \delta_e \Upsilon_\eta([\bar{A}, \bar{B}]) \\
&\quad + [\delta_e \Upsilon_\eta(A), \delta_e \Upsilon_\eta(B)] \\
&= \text{ad}_{\text{Ad}_\eta^* A}^* \pi_{G^*}(\eta, \text{Ad}_\eta^* B) - \text{ad}_{\text{Ad}_\eta^* B}^* \pi_{G^*}(\eta, \text{Ad}_\eta^* A) \\
&\quad + \pi_{G^*}(\eta, \text{Ad}_\eta^* [A, B]) + [\pi_{G^*}(\eta, \text{Ad}_\eta^* A), \pi_{G^*}(\eta, \text{Ad}_\eta^* B)].
\end{aligned} \tag{2.74}$$

Similarly, the twisted multiplicativity (2.31) of Υ_η^* implies

$$\begin{aligned}
\delta_g \Upsilon_\eta^*(g \cdot A) &= \Upsilon^*(g, \eta)^{-1} \cdot T_g \Upsilon_\eta^*(g \cdot A) \\
&= \Upsilon^*(g, \eta)^{-1} \cdot (T_\eta \Upsilon_g^*(\pi_{G^*}(\eta^{-1}, \text{Ad}_{\eta^{-1}}^* A) \cdot \eta) + \Upsilon_\eta^*(g) \cdot T_e \Upsilon_\eta^*(A)) \\
&= \text{Ad}_{\Upsilon_\eta^*(g)^{-1}} \pi_G(\Upsilon_\eta^*(g)^{-1}, \text{Ad}_{\Upsilon_\eta^*(g)^{-1}}^* \pi_{G^*}(\eta^{-1}, \text{Ad}_{\eta^{-1}}^* A)) + \text{Ad}_{\eta^{-1}}^* A,
\end{aligned} \tag{2.75}$$

where we used $T_e(g \mapsto \Upsilon_g(\eta^{-1})^{-1})(A) = \pi_{G^*}(\eta^{-1}, \text{Ad}_{\eta^{-1}}^* A) \cdot \eta$ and $T_\eta \Upsilon_g^*(\mu \cdot \eta) = T_e \Upsilon_{\Upsilon_\eta^*(g)}^*(\mu) = \pi_G(\Upsilon_\eta^*(g)^{-1}, \text{Ad}_{\Upsilon_\eta^*(g)^{-1}}^* \mu) \cdot \Upsilon_\eta^*(g)$. Taking the derivative with respect to g at the identity in the direction $B \in \mathfrak{g}$ yields

$$\begin{aligned}
\bar{B}_e (\delta \Upsilon_\eta^*(\bar{A})) &= -T_e(\pi_G(\cdot, \pi_{G^*}(\eta^{-1}, \text{Ad}_{\eta^{-1}}^* A))) (T_e \Upsilon_\eta^*(B)) \\
&= -\text{ad}_{\pi_{G^*}(\eta^{-1}, \text{Ad}_{\eta^{-1}}^* A)}^* (\text{Ad}_{\eta^{-1}}^* B)
\end{aligned} \tag{2.76}$$

Since $\delta \Upsilon_\eta^* \in \Omega^1(G, \mathfrak{g})$ satisfies the Maurer–Cartan equation for every $\eta \in G^*$, we thus find

$$\begin{aligned}
\circ &= \left(d \delta \Upsilon_\eta^* + \frac{1}{2} [\delta \Upsilon_\eta^*, \delta \Upsilon_\eta^*] \right)_e (A, B) \\
&= \bar{A}_e (\delta \Upsilon_\eta^*(\bar{B})) - \bar{B}_e (\delta \Upsilon_\eta^*(\bar{A})) - \delta_e \Upsilon_\eta^*([\bar{A}, \bar{B}]) \\
&\quad + [\delta_e \Upsilon_\eta^*(A), \delta_e \Upsilon_\eta^*(B)] \\
&= -\text{ad}_{\pi_{G^*}(\eta^{-1}, \text{Ad}_{\eta^{-1}}^* B)}^* (\text{Ad}_{\eta^{-1}}^* A) + \text{ad}_{\pi_{G^*}(\eta^{-1}, \text{Ad}_{\eta^{-1}}^* A)}^* (\text{Ad}_{\eta^{-1}}^* B) - \text{Ad}_{\eta^{-1}}^* [A, B] \\
&\quad + [\text{Ad}_{\eta^{-1}}^* A, \text{Ad}_{\eta^{-1}}^* B]
\end{aligned} \tag{2.77}$$

Using (2.74) and (2.77), we now obtain

$$\begin{aligned}
 d_{\mathfrak{g}}^{\text{CE}} \pi_{G^*}(\eta, A, B) &- \frac{1}{2}[\pi_{G^*}, \pi_{G^*}](\eta, A, B) \\
 &= -\text{ad}_A^* \pi_{G^*}(\eta, B) + \text{ad}_B^* \pi_{G^*}(\eta, A) - \pi_{G^*}(\eta, [A, B]) \\
 &\quad - \pi_{G^*}(\eta, \text{ad}_{\pi_{G^*}(\eta, A)}^* B) + \pi_{G^*}(\eta, \text{ad}_{\pi_{G^*}(\eta, B)}^* A) \\
 &\quad - [\pi_{G^*}(\eta, A), \pi_{G^*}(\eta, B)] \\
 &= \pi_{G^*}(\eta, \text{Ad}_\eta^*[\text{Ad}_{\eta^{-1}}^* A, \text{Ad}_{\eta^{-1}}^* B]) - \pi_{G^*}(\eta, [A, B]) \\
 &\quad - \pi_{G^*}(\eta, \text{ad}_{\pi_{G^*}(\eta, A)}^* B) + \pi_{G^*}(\eta, \text{ad}_{\pi_{G^*}(\eta, B)}^* A) \\
 &= 0.
 \end{aligned} \tag{2.78}$$

This verifies the Poisson property (2.63). \square

Note that, by construction, the coconjugation action Υ and the Poisson bivector field π_{G^*} are connected by

$$-\eta \cdot \pi_{G^*}(\eta, \text{Ad}_\eta^* A) = T_e \Upsilon_\eta(A) \in T_\eta G^*, \quad \eta \in G^*, A \in \mathfrak{g}. \tag{2.79}$$

In Poisson geometry, actions satisfying this relation with respect to a Poisson structure on a Lie group are called *Dressing transformations* (see, e.g., [Lu90, Definition 2.38] or [FR96]). For questions concerning the equivariance of momentum maps, it feels however more natural to put the action first and then derive the Poisson structure from it, than the other way around. In summary, we obtain the following hierarchy:

$$\begin{array}{ccc}
 \text{Coconjugation actions } (\Upsilon, \Upsilon^*) & & \\
 \text{dressing transformations } \left(\begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \right) & \text{Theorem 2.27} & \\
 \text{Dual pair of Poisson Lie groups } \kappa(G, G^*) & & (2.80) \\
 [\text{LW90, Theorem 1.8}] \left(\begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \right) & [\text{LW90, Theorem 1.8}] & \\
 \text{Lie bialgebra } \kappa(\mathfrak{g}, \mathfrak{g}^*) & &
 \end{array}$$

The dotted arrows indicate that the integration of structure is not always possible. The integration of a Lie bialgebra to a dual pair of Poisson Lie groups, [LW90, Theorem 1.8.], requires the groups to be connected and simply connected. Moreover, to obtain dressing transformations one needs that the dressing vector fields are complete, see [LW90, Theorem 3.12].

EXAMPLE 2.28 For the coadjoint action dual pair $\kappa(G, \mathfrak{g}^*)$ discussed in Exam-

ple 2.20, the induced Poisson Lie structure on \mathfrak{g}^* is given by

$$\pi_{\mathfrak{g}^*}(\mu, A) = \text{ad}_A^* \mu. \quad (2.81)$$

This is the usual Kostant–Kirillov–Souriau Poisson structure. \diamond

EXAMPLE 2.29 Continuing with Example 2.21, let D be a finite-dimensional connected complex semisimple Lie group and let $D = KB$ be its Iwasawa decomposition. The action (2.35) of K on B induces the following Poisson Lie structure on B :

$$\pi_B(b, A) = -\text{pr}_b(\text{Ad}_{b^{-1}} \text{Ad}_{b^{-1}}^* A). \quad (2.82)$$

The corresponding bivector field is given by

$$\Lambda_b(A, B) = \text{Im } \kappa_{\mathfrak{b}}(\text{pr}_b(\text{Ad}_b A), \text{pr}_{\mathfrak{f}}(\text{Ad}_b B)), \quad (2.83)$$

which coincides with the one given in [LR91, Definition 4.2]. Similarly, the action (2.35) of B on K induces the Poisson Lie structure on K

$$\pi_K(k, \mu) = -\text{pr}_{\mathfrak{f}}(\text{Ad}_{k^{-1}} \text{Ad}_{k^{-1}}^* \mu) \quad (2.84)$$

with bivector field

$$\Lambda_k(\mu, \nu) = \text{Im } \kappa_{\mathfrak{b}}(\text{pr}_{\mathfrak{f}}(\text{Ad}_k \mu), \text{pr}_{\mathfrak{b}}(\text{Ad}_k \nu)). \quad (2.85)$$

This coincides with the Poisson structure of [LW90, Theorem 4.3]. \diamond

EXAMPLE 2.30 Consider the dual pair of Lie groups $\kappa(\text{Diff}_{\mu}(M), \hat{\mathbb{H}}^2(M, \text{U}(1)))$ from Example 2.2. The coadjoint action of a diffeomorphism is given by pull-back. Thus a natural coconjugation action is given by

$$\text{Diff}_{\mu}(M) \times \hat{\mathbb{H}}^2(M, \text{U}(1)) \rightarrow \hat{\mathbb{H}}^2(M, \text{U}(1)), \quad (\phi, h) \mapsto (\phi^{-1})^* h. \quad (2.86)$$

This indeed endows $\kappa(\text{Diff}_{\mu}(M), \hat{\mathbb{H}}^2(M, \text{U}(1)))$ with the structure of a coconjugation matched pair (relative to the trivial action of $\hat{\mathbb{H}}^2(M, \text{U}(1))$ on $\text{Diff}_{\mu}(M)$). Using Proposition D.3, we see that the induced Poisson Lie structure on $\hat{\mathbb{H}}^2(M, \text{U}(1))$ is defined by

$$\hat{\mathbb{H}}^2(M, \text{U}(1)) \times \mathfrak{X}_{\mu}(M) \rightarrow \Omega^1(M)/d\Omega^0(M), \quad (h, X) \mapsto [X \lrcorner \text{curv } h]. \quad (2.87)$$

Under the integration pairing, this corresponds to the map

$$\hat{\mathbb{H}}^2(M, \text{U}(1)) \times \mathfrak{X}_{\mu}(M) \times \mathfrak{X}_{\mu}(M) \rightarrow \mathbb{R} \quad (h, X, Y) \mapsto \int_M \text{curv } h(X, Y) \mu. \quad (2.88)$$

Note that, for fixed $h \in \hat{H}^2(M, U(1))$, this is precisely the Lichnerowicz cocycle on $\mathfrak{X}_\mu(M)$ defined by the 2-form curv h . In other words, the Lichnerowicz cocycle is derived from the pull-back action. \diamond

For the rest of the section, we consider the case where G^* is Abelian in more detail. To emphasize the Abelian nature we write group operation in G^* as addition. In this case, the coadjoint action of G^* on \mathfrak{g} is trivial so that we can choose $\Upsilon^*: G^* \times G \rightarrow G$ to be the trivial action. Accordingly, we refer to the G -action Υ on G^* as the conjugation action. Moreover, the matched pair condition (2.31) simplifies to

$$\Upsilon_g(\eta_1 + \eta_2) = \Upsilon_g(\eta_1) + \Upsilon_g(\eta_2), \quad (2.89)$$

and we will call Υ standard if this relation is satisfied.

The following is a refinement of Lemma 2.19 in the Abelian setting.

PROPOSITION 2.31 *Let $\Upsilon: G \times G^* \rightarrow G^*$ be a standard coconjugation action. A map $c: G \rightarrow G^*$ defines a coconjugation action by $\tilde{\Upsilon} := \Upsilon + c$ if and only if c satisfies $c(gh) = c(g) + \Upsilon_g(c(h))$ for all $g, h \in G$. That is, $\tilde{\Upsilon}$ is a coconjugation action if and only if c is a 1-cocycle with respect to Υ . \diamond*

Proof. Suppose that $c: G \rightarrow G^*$ is a map satisfying $c(gh) = c(g) + \Upsilon_g(c(h))$. Define the map $\tilde{\Upsilon}: G \times G^* \rightarrow G^*$ by $\tilde{\Upsilon}_g(\eta) = \Upsilon_g(\eta) + c(g)$. It is clear that $\tilde{\Upsilon}$ still integrates the coadjoint action. Thus it is left to show that $\tilde{\Upsilon}$ is a left action indeed. This follows from the following straightforward calculation:

$$\begin{aligned} (\tilde{\Upsilon}_g \circ \tilde{\Upsilon}_h)(\eta) &= \Upsilon_g(\Upsilon_h(\eta) + c(h)) + c(g) \\ &= \Upsilon_g(\Upsilon_h(\eta)) + \Upsilon_g(c(h)) + c(g) \\ &= \Upsilon_{gh}(\eta) + c(gh) \\ &= \tilde{\Upsilon}_{gh}(\eta). \end{aligned} \quad (2.90)$$

Reading the argument backwards establishes the converse claim in the “if and only if” statement. \square

EXAMPLE 2.32 The coadjoint representation is a standard coconjugation action of G on $G^* := \mathfrak{g}^*$. Moreover, since \mathfrak{g}^* is connected, the previous proposition in combination with Lemma 2.19 establishes a bijection between coconjugation actions on \mathfrak{g}^* and 1-cocycles $c: G \rightarrow \mathfrak{g}^*$. The coconjugation action corresponding to a 1-cocycle c is the affine action $(g, \mu) \mapsto \text{Ad}_{g^{-1}}^* \mu + c(g)$, which plays an important role for classical non-equivariant momentum maps (see [OR03, Definiton 4.5.23]). \diamond

According to Theorem 2.27, every standard coconjugation action yields an associated Lie Poisson structure π_{G^*} on G^* (the induced Lie Poisson structure π_G

on G vanishes, because Υ^* is trivial). For the Poisson structure induced on G^* we have the following affine version.

PROPOSITION 2.33 *Let $\kappa(G, G^*)$ be a dual pair of Lie groups with Abelian G^* . For every standard coconjugation action Υ and for every cocycle $c: G \rightarrow G^*$, let $\tilde{\Upsilon} = \Upsilon + c$ be the associated coconjugation action and define $\tilde{\pi}_{G^*}: G^* \times \mathfrak{g} \rightarrow \mathfrak{g}^*$ by*

$$\tilde{\pi}_{G^*}(\eta, A) := -\eta^{-1} \cdot T_e \tilde{\Upsilon}_\eta(A). \quad (2.91)$$

If π_{G^*} is skew-symmetric in the sense of (2.61), then it defines an affine Poisson structures on G^* . \diamond

Proof. Since we have $\tilde{\Upsilon}_{\zeta+\eta} = \tilde{\Upsilon}_\zeta + \tilde{\Upsilon}_\eta - c$, a similar calculation as in the proof of Theorem 2.27 shows that

$$\begin{aligned} \tilde{\pi}_{G^*}(\zeta + \eta, A) &= \tilde{\pi}_{G^*}(\zeta, A) + \tilde{\pi}_{G^*}(\eta, A) - T_e c(A) \\ &= \tilde{\pi}_{G^*}(\zeta, A) + \tilde{\pi}_{G^*}(\eta, A) - \tilde{\pi}_{G^*}(o, A). \end{aligned} \quad (2.92)$$

Thus $\tilde{\pi}_{G^*}$ is an affine bivector field. The Poisson property follows as in Theorem 2.27. \square

For a given coconjugation action Υ of G on G^* , we say that the group-valued momentum map is *equivariant* if it is G -equivariant as a map $J: M \rightarrow G^*$. It is well-known that a classical momentum map is equivariant if and only if it is a Poisson map. This relation carries over to our more general setting.

As above, we first have to reformulate what it means for J to be a Poisson map in terms of π_{G^*} . For a smooth function $f: G^* \rightarrow \mathbb{R}$, the left derivative $T_\eta^L f: \mathfrak{g}^* \rightarrow \mathbb{R}$ at $\eta \in G^*$ is an element of the double dual \mathfrak{g}^{**} . In finite dimensions, we can view $T_\eta^L f$ thus as an element of \mathfrak{g} . The Poisson bracket of $f, g \in C^\infty(G^*)$ is given by $\{f, g\}_{G^*}(\eta) = \kappa(T_\eta^L f, \pi_{G^*}(\eta, T_\eta^L g))$ according to (2.69). Given a group-valued momentum map $J: M \rightarrow G^*$, we calculate for $X_m \in T_m M$

$$\begin{aligned} T_m(f \circ J)(X) &= T_{J(m)} f(J(m) \cdot \delta_m J(X)) = \kappa(T_{J(m)}^L f, \delta_m J(X)) \\ &= -\omega_m((T_{J(m)}^L f) \cdot m, X) \end{aligned} \quad (2.93)$$

by (2.13). Hence, we obtain (see (2.1))

$$\begin{aligned} \{f \circ J, g \circ J\}_M(m) &= \omega_m(T_m(f \circ J), T_m(g \circ J)) \\ &= \omega_m((T_{J(m)}^L f) \cdot m, (T_{J(m)}^L g) \cdot m). \end{aligned} \quad (2.94)$$

Thus, in summary, J is a Poisson map, i.e., $\{f \circ J, g \circ J\}_M = \{f, g\}_{G^*} \circ J$, if and only if

$$\omega_m(A \cdot m, B \cdot m) = \kappa(A, \pi_{G^*}(J(m), B)) \quad (2.95)$$

holds for all $A, B \in \mathfrak{g}$ and $m \in M$. This equation no longer relies on reflexivity to make sense and so we adopt it as the definition for J to be a Poisson map in infinite dimensions. Recall that the left-hand side of (2.95) defines the so called *non-equivariance* cocycle $\sigma_m(A, B) = \omega_m(A \cdot m, B \cdot m)$. Thus, J is a Poisson map if and only if the Lie algebra cocycles σ_m and $\kappa(\cdot, \pi_{G^*}(J(m), \cdot))$ coincide for all $m \in M$.

PROPOSITION 2.34 *Let $\kappa(G, G^*)$ be a dual pair of Lie groups where G^* is Abelian. Assume that G acts on the symplectic manifold (M, ω) with a group-valued momentum map $J: M \rightarrow G^*$. If J is equivariant with respect to a given coconjugation action $\tilde{Y} = Y + c$, then it is a Poisson map relative to the induced Poisson structure $\tilde{\pi}_{G^*}$ of Proposition 2.33. \diamond*

Proof. If $J: M \rightarrow G^*$ is equivariant, then $J(g \cdot m) = \tilde{Y}_g J(m)$ relative to the coconjugation action $\tilde{Y}: G \times G^* \rightarrow G^*$. Thus,

$$\delta_m J(B \cdot m) = J(m)^{-1} \cdot T_e \tilde{Y}_{J(m)}(B) = -\tilde{\pi}_{G^*}(J(m), B). \quad (2.96)$$

On the other hand, $\kappa(A, \delta_m J(B \cdot m)) = -\omega_m(A \cdot m, B \cdot m)$ by the definition of the momentum map, and the claim follows. \square

2.6 Momentum maps for group extensions

Consider the following set-up. Let H, K, G be Lie groups which fit in the exact sequence

$$e \longrightarrow H \xrightarrow{\iota} K \xrightarrow{\pi} G \longrightarrow e, \quad (2.97)$$

where ι and π are Lie group morphisms. Suppose that K acts on a symplectic manifold (M, ω) such that the diffeomorphism defined by each group element is symplectic. We are interested in the momentum map for this action. Starting from the extension sequence (2.97), it is natural to seek an expression of the momentum map for the K -action in terms of the momentum maps for the groups H and G , assuming they exist. Similar questions occur in the context of symplectic reduction by stages (see [Mar+07]).

The corresponding sequence on the infinitesimal level

$$0 \longrightarrow \mathfrak{h} \xrightarrow{\iota} \mathfrak{k} \xrightarrow{\pi} \mathfrak{g} \longrightarrow 0 \quad (2.98)$$

always splits as vector spaces but not necessarily as Lie algebras. Let us fix a splitting $\sigma: \mathfrak{g} \rightarrow \mathfrak{k}$ in the category of locally convex vector spaces and write $\mathfrak{h} \oplus_{\sigma} \mathfrak{g} = \mathfrak{k}$ for the corresponding direct sum. Thus, every $A \in \mathfrak{k}$ can be uniquely written as the sum $A = \iota(A_H) + \sigma(A_G)$ with $A_H \in \mathfrak{h}$ and $A_G = \pi(A) \in \mathfrak{g}$.

PROPOSITION 2.35 *Let $\kappa(G, G^*)$ be a dual pair of Lie groups and let $\langle \mathfrak{h}, \mathfrak{h} \rangle$ be a pairing of \mathfrak{h} with itself. Assume that the induced action of H on M has a standard momentum map $J_H: M \rightarrow \mathfrak{h}$ with respect to the pairing $\langle \mathfrak{h}, \mathfrak{h} \rangle$ and that there exists a map $J_\sigma: M \rightarrow G^*$ satisfying*

$$\sigma(A_G)^* \lrcorner \omega + \kappa(A_G, \delta J_\sigma) = 0, \quad A_G \in \mathfrak{g}. \quad (2.99)$$

Then, $J_K = (J_H, J_\sigma): M \rightarrow \mathfrak{h} \times G^$ is a group-valued momentum map for the K -action with respect to the pairing $(\mathfrak{h} \oplus_\sigma \mathfrak{g}, \mathfrak{h} \oplus \mathfrak{g}^*) = \langle \mathfrak{h}, \mathfrak{h} \rangle + \kappa(\mathfrak{g}, \mathfrak{g}^*)$. \diamond*

Proof. The momentum map equation

$$A^* \lrcorner \omega + (A, \delta J_K) = 0, \quad A \in \mathfrak{k} \quad (2.100)$$

is linear in $A = \iota(A_H) + \sigma(A_G)$ and thus decomposes in

$$A_H^* \lrcorner \omega + \langle A_H, dJ_H \rangle = 0 \quad (2.101)$$

and in

$$\sigma(A_G)^* \lrcorner \omega + \kappa(A_G, \delta J_\sigma) = 0. \quad (2.102)$$

Both equations hold by assumption. \square

Formally, (2.99) looks like a momentum map relation. However, we do not assume that σ is a splitting on the level of Lie algebras and, hence G , or its Lie algebra \mathfrak{g} , does not act on M via σ . However, if G happens to act on M through a different splitting $\chi: G \rightarrow K$ which is a group section of π (see (2.97)), then J_σ is the momentum map up to some twisting by J_H .

PROPOSITION 2.36 *In the setting above, let $\chi: \mathfrak{g} \rightarrow \mathfrak{k}$ be a Lie algebra homomorphism splitting the exact sequence (2.98) and hence there is an induced (infinitesimal) \mathfrak{g} -action on M . Define $\tau_{\sigma\chi} := \chi - \sigma: \mathfrak{g} \rightarrow \mathfrak{h}$ and denote its dual map with respect to the chosen pairings by $\tau_{\sigma\chi}^*: \mathfrak{h}^* \rightarrow \mathfrak{g}^*$. Moreover, assume that the dual group G^* of G is Abelian. Then,*

$$J_\chi: M \rightarrow G^*, \quad m \mapsto J_\sigma(m) \cdot \exp(\tau_{\sigma\chi}^* J_H(m)) \quad (2.103)$$

is a group-valued momentum map for the G -action on M . Moreover, J_χ does not depend on the splitting σ . \diamond

Proof. For $B \in \mathfrak{g}$, the infinitesimal generator vector field is $B^* = (\chi(B))^* = \sigma(B)^* + \tau_{\sigma\chi}(B)^*$. Since G^* is by assumption Abelian, we have the ordinary product rule $\delta J_\chi = \delta J_\sigma + \tau_{\sigma\chi}^* dJ_H$. Calculating, using the defining equations for J_H and J_σ ,

yields

$$\begin{aligned}
& B^* \lrcorner \omega + \kappa(B, \delta J_\chi) \\
&= \sigma(B)^* \lrcorner \omega + \kappa(B, \delta J_\sigma) + \tau_{\sigma\chi}(B)^* \lrcorner \omega + \kappa(B, \tau_{\sigma\chi}^* dJ_H) \\
&= \sigma(B)^* \lrcorner \omega + \kappa(B, \delta J_\sigma) - \langle \tau_{\sigma\chi}(B), dJ_H \rangle + \kappa(B, \tau_{\sigma\chi}^* dJ_H) \\
&= 0,
\end{aligned} \tag{2.104}$$

which shows that J_χ is a group-valued momentum map.

It is left to show that the momentum map J_χ does not depend on the splitting that was used in the construction. Let σ_1 and σ_2 be different splittings. First, we start with the observation that if J_{σ_1} satisfies (2.99) for σ_1 , then the map defined by $J_{\sigma_2} := J_{\sigma_1} \cdot \exp(-(\sigma_1 - \sigma_2)^* J_H)$ satisfies the same relation with σ replaced by σ_2 . Indeed, since G^* is assumed to be Abelian, the chain rule for the logarithmic derivative simplifies and is not twisted by the adjoint representation. We thus obtain

$$\delta J_{\sigma_2} = \delta J_{\sigma_1} - (\sigma_1 - \sigma_2)^* dJ_H. \tag{2.105}$$

Using (2.99) and the momentum relation for J_H , we get for all $A_G \in \mathfrak{g}$

$$\begin{aligned}
\kappa(A_G, \delta J_{\sigma_2}) &= \kappa(A_G, \delta J_{\sigma_1}) - \kappa(A_G, (\sigma_1 - \sigma_2)^* dJ_H) \\
&= \kappa(A_G, \delta J_{\sigma_1}) - \langle (\sigma_1 - \sigma_2)A_G, dJ_H \rangle \\
&= -(\sigma_1 A_G)^* \lrcorner \omega + ((\sigma_1 - \sigma_2)A_G)^* \lrcorner \omega \\
&= -(\sigma_2 A_G)^* \lrcorner \omega.
\end{aligned} \tag{2.106}$$

On the other hand, we clearly have $\tau_{\sigma_2\chi}^* = \tau_{\sigma_1\chi}^* + (\sigma_1 - \sigma_2)^*$. Hence, in summary,

$$\begin{aligned}
J_{\sigma_2} \cdot \exp(\tau_{\sigma_2\chi}^* J_H) &= J_{\sigma_1} \cdot \exp(-(\sigma_1 - \sigma_2)^* J_H) \exp(\tau_{\sigma_1\chi}^* J_H) \exp((\sigma_1 - \sigma_2)^* J_H) \\
&= J_{\sigma_1} \cdot \exp(\tau_{\sigma_1\chi}^* J_H)
\end{aligned} \tag{2.107}$$

and thus J_χ is independent of the chosen splitting σ . \square

In a similar spirit, we can determine the momentum map for the action of a subgroup.

PROPOSITION 2.37 *Let G act on the symplectic manifold (M, ω) with group-valued momentum map $J: M \rightarrow G^*$. Let $\iota: H \rightarrow G$ be a Lie group homomorphism, so that H acts through G on M . Fix a dual group H^* of H . Assume¹ that there is a Lie group homomorphism $\rho: G^* \rightarrow H^*$ whose associated Lie algebra homomorphism $\rho: \mathfrak{g}^* \rightarrow \mathfrak{h}^*$ is the dual of $\iota: \mathfrak{h} \rightarrow \mathfrak{g}$ with respect to $\kappa(\mathfrak{g}, \mathfrak{g}^*)$ and $\langle \mathfrak{h}, \mathfrak{h}^* \rangle$. Then $J^H := \rho \circ J: M \rightarrow H^*$ is a group-valued momentum map for the induced H -action. \diamond*

¹ Note that these assumptions are automatically satisfied in finite dimensions if G is 1-connected,

Proof. Let $B \in \mathfrak{h}$. Then using the fact that taking the logarithmic derivative commutes with Lie group homomorphisms [Nee06, Proposition II.4.1.1], we have

$$\iota(B)^* \lrcorner \omega = -\kappa(\iota(B), \delta J) = -\langle B, \rho \circ \delta J \rangle = -\langle B, \delta(\rho \circ J) \rangle, \quad (2.108)$$

which proves the proposition. \square

3 GLOBAL ANALYSIS OF SYMPLECTIC FIBER BUNDLES

3.1 Symplectic form on the space of sections

The prime example of an infinite-dimensional manifold is the space of sections of a fiber bundle. So, for the study of infinite-dimensional symplectic geometry, it is suggestive to search for a way to construct a symplectic form on the space of sections using some geometric structure on the fiber bundle. As we will see, a symplectic form along the fibers is what we are looking for.

Let $\pi: P \rightarrow M$ be a finite-dimensional right principal G -bundle. Given a left G -manifold \underline{E} , where the action is denoted by $\underline{l}: G \times \underline{E} \rightarrow \underline{E}$, form the associated fiber bundle $F = P \times_G \underline{E} \rightarrow M$ (see Appendix A for definitions and conventions). If, in addition, \underline{E} carries a G -invariant symplectic form $\underline{\omega}$, then there exists a unique symplectic structure ω_m on each fiber F_m such that the map

$$\iota_p: \underline{E} \rightarrow F_m, \quad \underline{f} \mapsto [p, \underline{f}] \quad (3.1)$$

is a symplectomorphism for all $p \in P_m$. The fiber bundle F with the induced fiberwise symplectic structure ω is called a *symplectic fiber bundle* and we denote it by (F, ω) . We emphasize that ω is not a 2-form on F but is merely a notation for the fiberwise symplectic structure.

A 2-form $\hat{\omega}$ on F is called an *extension* of the fiberwise symplectic structure if its pull-back to each fiber F_m coincides with ω_m . Such extensions always exist (for example, choose a connection in F and construct the extension $\hat{\omega}$ by letting it vanish on horizontal vectors). In the theory of symplectic fiber bundles one usually tries to construct extensions that are again symplectic forms [Wei80]. We emphasize that we do not require the extension $\hat{\omega}$ to be non-degenerate. An extension $\hat{\omega}$ is called *compatible with a given connection* on P if the vertical and horizontal subbundles of the associated connection on F are orthogonal with respect to $\hat{\omega}$.

i.e., connected and $\pi_1(G, e) = 1$. In infinite dimensions, however, the adjoint $\rho: \mathfrak{g}^* \rightarrow \mathfrak{h}^*$ of the linear map ι does not need to exist, and even if it exists, it does not necessarily integrate to a Lie group homomorphism (for this, we would need some regularity assumptions on the dual group H^* , cf. [Nee06, Theorem III.1.5]).

In what follows, we assume that the base manifold M is closed (i.e., compact without boundary) so that the space of sections $\mathcal{F} = \Gamma^\infty(F)$ is a Fréchet manifold. The tangent space to \mathcal{F} at a section ϕ consists of vertical vector fields along ϕ , i.e., sections of ϕ^*VF , where $VF \subset TF$ is the vector subbundle of vertical vectors. Fix a volume form μ on M and choose an extension $\hat{\omega}$ of the fiberwise symplectic structure ω on F . Using the evaluation map $\text{ev}: M \times \mathcal{F} \rightarrow F$ and the projection $\text{pr}_M: M \times \mathcal{F} \rightarrow M$, we define a 2-form Ω on \mathcal{F} by

$$\Omega = \mu \hat{*} \omega \equiv \int_M \text{pr}_M^* \mu \wedge \text{ev}^* \hat{\omega}, \quad (3.2)$$

which uses the hat product introduced by Vizman [Viz11], here generalized to sections of fiber bundles. Details about the resulting hat calculus are collected in Appendix C. The form Ω depends only on the fiber symplectic form ω and not on the extension $\hat{\omega}$, since only the vertical part of the tangent vector plays a role; see Remark C.1 (which also justifies the notation $\mu \hat{*} \omega$ instead of $\mu \hat{*} \hat{\omega}$). In fact, by (C.4), we have the explicit description

$$\Omega_\phi(Y_1, Y_2) = \int_M \omega_\phi(Y_1, Y_2) \mu, \quad Y_1, Y_2 \in \Gamma^\infty(\phi^*VF). \quad (3.3)$$

This identity also shows that Ω is non-degenerate, because ω is fiberwise non-degenerate, by assumption. The exterior differential of Ω is given by $d\Omega = (-1)^{\dim M} \mu \hat{*} d_\pi \omega$, where d_π denotes the vertical differential along the fibers; see (C.6). Thus Ω is closed. Let us record this observation.

PROPOSITION 3.1 *Let M be a closed manifold endowed with volume form μ and let (F, ω) be a symplectic fiber bundle over M . Then $\Omega = \mu \hat{*} \omega$ is a symplectic structure on the space of smooth sections \mathcal{F} of F . \diamond*

Note that (3.3) shows that Ω coincides with the symplectic form considered in a slightly more restricted setting by Donaldson [Don03, Equation 7].

3.2 Action of $\text{Aut}(P)$

Every bundle automorphism ψ of F induces a diffeomorphism $\check{\psi}$ on the base M . Consider the natural smooth left action of $\text{Aut}(F)$ on the section space \mathcal{F} defined by

$$\Upsilon: \text{Aut}(F) \times \mathcal{F} \rightarrow \mathcal{F}, \quad \psi \cdot \phi = \psi \circ \phi \circ \check{\psi}^{-1}. \quad (3.4)$$

Let

$$\text{Aut}_{\mu, \omega}(F) = \{\psi \in \text{Aut}(F) \mid \check{\psi} \in \text{Diff}_\mu(M), \psi^* \omega = \omega\} \quad (3.5)$$

be the subgroup of automorphisms of F preserving the fiber symplectic structure and inducing volume-preserving diffeomorphisms on the base. Equation (C.9) implies that the induced action of $\text{Aut}_{\mu,\omega}(F)$ on \mathcal{F} leaves the symplectic form $\Omega = \mu \hat{*} \omega$ invariant.

Although it is thus perfectly natural to seek a momentum map for the $\text{Aut}_{\mu,\omega}(F)$ action on \mathcal{F} , one is confronted with serious difficulties in this quest. We will further comment on these matters below in Remark 4.1 and for now concentrate on bundle automorphisms of $F = P \times_G \underline{E}$ coming from the principal bundle. Every principal bundle automorphism φ of P defines a bundle automorphism φ_F of F by $\varphi_F = \varphi \times_G \text{id}_{\underline{E}}$. Hence, by definition, $\varphi_F \circ \iota_p = \iota_{\varphi(p)}$ for every $p \in P$, where $\iota_p: \underline{E} \rightarrow F$ was introduced in (3.1). This relation implies $\iota_p^* \varphi_F^* \omega_{\check{\varphi}(m)} = \iota_{\varphi(p)}^* \omega_{\check{\varphi}(m)} = \underline{\omega}$, or equivalently $\varphi_F^* \omega_{\check{\varphi}(m)} = \omega_m$ for $m = \pi(p)$. Thus, φ_F leaves the fiber symplectic structure invariant and we get a group homomorphism $\text{Aut}_{\mu}(P) \rightarrow \text{Aut}_{\mu,\omega}(F)$, where

$$\text{Aut}_{\mu}(P) = \{\varphi \in \text{Aut}(P) \mid \check{\varphi} \in \text{Diff}_{\mu}(M)\} \quad (3.6)$$

is the subgroup of automorphisms of P that induce volume-preserving diffeomorphisms on the base. Therefore, we seek a momentum map for the $\text{Aut}_{\mu}(P)$ action on \mathcal{F} . We will divide this problem into its vertical and horizontal parts using Proposition 2.35.

Assigning to a principal bundle automorphism ψ of P the induced diffeomorphism $\check{\psi}$ on the base M , yields a group homomorphism $\text{Aut}_{\mu}(P) \rightarrow \text{Diff}_{\mu}(M)$, whose kernel is the gauge group $\text{Gau}(P)$. Thus we obtain a short exact sequence of Lie groups

$$\text{id} \rightarrow \text{Gau}(P) \rightarrow \text{Aut}_{\mu}(P) \rightarrow \text{Diff}_{\mu,P}(M) \rightarrow \text{id}, \quad (3.7)$$

where $\text{Diff}_{\mu,P}(M)$ denotes the subgroup of volume-preserving diffeomorphisms which is the range of the projection given by the third arrow in (3.7). Note that $\text{Diff}_{\mu,P}(M)$ consists of those diffeomorphisms that preserve the equivalence class of P under pull-back, and thus it is an open subgroup of $\text{Diff}_{\mu}(M)$.

So we find ourselves in the setting discussed in Section 2.6. Proposition 2.35 shows that the momentum map for $\text{Aut}_{\mu}(P)$ is obtained from the momentum map for the gauge group as well as the ‘partial’ momentum map for $\text{Diff}_{\mu,P}(M)$. Before we determine these two ingredients, let us record the common starting point. By (C.8), the infinitesimal generator vector field on \mathcal{F} induced by $Y \in \text{aut}_{\mu}(P)$ is given by

$$Y_{\check{\phi}}^* = Y_F \circ \phi - T\phi(Y_M) \in \Gamma^{\infty}(\phi^*VF), \quad (3.8)$$

where $Y_M = T\pi(Y)$ denotes the induced vector field on M and where Y_F is the vector field on F induced by the natural map $\text{aut}_{\mu}(P) \rightarrow \text{aut}_{\mu,\omega}(F)$. Using (C.11),

we obtain

$$Y^* \lrcorner \Omega = (Y_M \lrcorner \mu) \hat{*} \hat{\omega} + (-1)^{\dim M} \mu \hat{*} (Y_F \lrcorner \hat{\omega}), \quad Y \in \text{aut}_\mu(P). \quad (3.9)$$

Hence, $Y^* \lrcorner \Omega$ decomposes into vertical and horizontal terms, so that we can determine the momentum map for the gauge group separately from the momentum map for the diffeomorphism group.

3.2.1 Momentum map for the gauge group

Since for every $Y \in \text{gau}(P)$ the induced vector field Y_M on the base vanishes, (3.9) simplifies to

$$Y^* \lrcorner \Omega + (-1)^{\dim M+1} \mu \hat{*} (Y_F \lrcorner \hat{\omega}) = 0. \quad (3.10)$$

To study the second term $(-1)^{\dim M} \mu \hat{*} (Y_F \lrcorner \hat{\omega}) \in \Omega^1(\mathcal{F})$, we evaluate it on $Z \in T_\phi \mathcal{F}$ and, using (C.2), obtain

$$(-1)^{\dim M} (\mu \hat{*} (Y_F \lrcorner \hat{\omega}))_\phi(Z) = \int_M \hat{\omega}_{\phi(m)}(Y_F \circ \phi(m), Z(m)) \mu. \quad (3.11)$$

In order to shift the calculation from section spaces to the fiber model \underline{E} , we introduce the smooth bundle map¹ over M

$$l: \text{Conj } P \times_M F \rightarrow F, \quad ([p, g], [p, \underline{f}]) \mapsto [p, g \cdot \underline{f}], \quad (3.12)$$

where $\text{Conj } P = P \times_G G \rightarrow M$ is the conjugation bundle associated to the principal G -bundle $P \rightarrow M$ (see Appendix A). The fiber derivative of l is given by

$$V_1 l: \text{Ad } P \times_M F \rightarrow VF = P \times_G T\underline{E}, \quad ([p, A], [p, \underline{f}]) \mapsto [p, A \underline{f}], \quad (3.13)$$

where $\text{Ad } P \rightarrow M$ is the adjoint bundle of the principal G -bundle $P \rightarrow M$. Let $\mathfrak{X}^v(F)$ denote the set of vertical vector fields on the associated fiber bundle F . Note that l induces a natural map $l_*: \text{Gau}(P) \rightarrow \text{Gau}(F)$ on sections and that the map $\text{gau}(P) \rightarrow \mathfrak{X}^v(F)$ sending a vertical vector field Y on P to its partner Y_F on F is identified with $(V_1 l)_*: \Gamma^\infty(\text{Ad } P) \rightarrow \mathfrak{X}^v(F)$. In other words, $Y_F = (V_1 l)_* Y$.

To calculate the contraction $Y_F \lrcorner \omega = (V_1 l)_* Y \lrcorner \omega$ of the vertical vector field Y_F with the fiber symplectic form ω we need a G -equivariant momentum map $J: \underline{E} \rightarrow \mathfrak{g}^*$ for the symplectic G -action on (\underline{E}, ω) . By G -equivariance, J yields a bundle map $J: F \rightarrow \text{Ad}^* P$, where $\text{Ad}^* P = P \times_G \mathfrak{g}^*$ is the coadjoint bundle. Taking the derivative in the fiber direction yields a bundle map $VJ: VF \rightarrow \text{Ad}^* P$. Note that the derivative of $J_*: \mathcal{F} \rightarrow \Gamma^\infty(\text{Ad}^* P)$ at $\phi \in \mathcal{F}$ equals $(VJ)_* \circ \phi$. Let $l'A$ be

¹ Morally, $l = \text{id}_P \times_G \underline{l}$.

	Fiber model \underline{F}	Bundle F
Symplectic form	$\underline{\omega}$	ω
Action	$\underline{l}: G \times \underline{F} \rightarrow \underline{F}$	$l: \text{Conj} P \times_M F \rightarrow F$
Infinitesimal action	$\underline{l}': \mathfrak{g} \times \underline{F} \rightarrow \underline{TF}$	$V_1 l: \text{Ad} P \times_M F \rightarrow VF$
Momentum map	$\underline{J}: \underline{F} \rightarrow \mathfrak{g}^*$	$J: F \rightarrow \text{Ad}^* P$

Table 2: Overview of the objects of the fiberwise Hamiltonian system and their companions in the bundle picture.

the infinitesimal generator vector field for the G -action $l: G \times \underline{F} \rightarrow \underline{F}$ associated to $A \in \mathfrak{g}$. Unravelling the definitions shows that the momentum map relation $(\underline{l}'A) \lrcorner \underline{\omega} + \langle A, d\underline{J} \rangle = 0$ on \underline{F} translates in the bundle picture to

$$((V_1 l)_* Y) \lrcorner \omega + \langle Y, VJ \rangle = 0 \quad (3.14)$$

on vertical vectors. Indeed, let $[p, \underline{u}] \in P \times_G \underline{TF}$ be a vertical vector at the point $[p, f] \in F$ over $m \in M$ and represent $Y(m) \in \text{Ad} P$ by the pair $[p, A]$ with $A \in \mathfrak{g}$. Then, using (3.13), we obtain

$$\begin{aligned} \omega_m(V_1 l(Y(m)), [p, \underline{u}]) &= \omega_m([p, A_f^*], [p, \underline{u}]) \\ &= \underline{\omega}_f(A_f^*, \underline{u}) \\ &= -\langle A, T_f \underline{J}(\underline{u}) \rangle. \end{aligned} \quad (3.15)$$

On the other hand, we have

$$\langle Y(m), VJ([p, \underline{u}]) \rangle = \langle [p, A], [p, T_f \underline{J}(\underline{u})] \rangle = \langle A, T_f \underline{J}(\underline{u}) \rangle, \quad (3.16)$$

which verifies (3.14). The additional structure on the bundle F derived from the fiberwise Hamiltonian system on \underline{F} is summarized in Table 2.

Returning to the original problem of computing the momentum map of the

gauge group, we can now rewrite (3.11) as

$$\begin{aligned}
 (-1)^{\dim M} (\mu \hat{*} (Y_F \lrcorner \hat{\omega}))_{\phi}(Z) &= \int_M \hat{\omega}_{\phi(m)}(Y_F \circ \phi(m), Z(m)) \mu \\
 &= \int_M \hat{\omega}_{\phi(m)}((V_1 l)_* Y \circ \phi(m), Z(m)) \mu \\
 &= - \int_M \langle Y(m), (VJ)_{\phi(m)}(Z(m)) \rangle \mu \\
 &= - \langle Y, (VJ)_{\phi}(Z) \rangle_{\text{Ad}'},
 \end{aligned} \tag{3.17}$$

where in the last step we used the natural pairing $\langle \cdot, \cdot \rangle_{\text{Ad}}: \Gamma^{\infty}(\text{Ad}P) \times \Gamma^{\infty}(\text{Ad}^*P) \rightarrow \mathbb{R}$ discussed in Example 2.5. Summarizing, we get

$$Y^* \lrcorner \Omega + \langle Y, (VJ)_* \rangle_{\text{Ad}} = 0. \tag{3.18}$$

Keeping in mind that $(VJ)_* \circ \phi$ is the derivative of $J_*: \mathcal{F} \rightarrow \text{gau}^*(P)$ at $\phi \in \mathcal{F}$, we read off the momentum map for the group of gauge transformation.

PROPOSITION 3.2 *Let $P \rightarrow M$ be a principal G -bundle over the closed manifold M endowed with a volume form μ . Let $F = P \times_G \underline{E}$ be a symplectic fiber bundle modeled on the Hamiltonian G -manifold $(\underline{E}, \underline{\omega}, \underline{J})$. The momentum map for the action of the group $\text{Gau}(P)$ of gauge transformations on the space \mathcal{F} of sections of F is given by*

$$\mathcal{J}_{\text{Gau}}: \mathcal{F} \rightarrow \text{gau}^*(P), \quad \phi \mapsto J_* \phi. \tag{3.19}$$

Here, $\text{gau}^*(P) = \Gamma^{\infty}(\text{Ad}^*P)$ is dual to $\text{gau}(P)$ by the integration pairing $\langle \cdot, \cdot \rangle_{\text{Ad}}$ introduced in Example 2.5 and, moreover, $J = \text{id}_P \times_G \underline{J}: F \rightarrow \text{Ad}^*P$ is the bundle map induced by \underline{J} . That is,

$$\langle Y, \mathcal{J}_{\text{Gau}}(\phi) \rangle_{\text{Ad}} = \int_M \langle Y(m), (\text{id}_P \times_G \underline{J}) \circ \phi(m) \rangle \mu(m) \tag{3.20}$$

for all $Y \in \text{gau}(P)$. ◇

In special cases, similar expressions for the momentum map of the gauge group were found in [CGSoo, Remark 3.1] (on Riemann surfaces) and in [Munoo, p. 4.2.2] (on Kähler manifolds), both in the context of Yang–Mills equations coupled to matter fields.

3.2.2 Momentum map for the diffeomorphism group

Let us now determine the momentum map for the action of the diffeomorphism group of the base. In the computations below we use a dual pair $\kappa(\text{Diff}_{\mu,P}(M), \mathcal{H})$ but we do not specify \mathcal{H} for the general case. The situations interesting to us are treated explicitly later. In light of Proposition 2.35, it is enough to split the infinitesimal sequence corresponding to (3.7) and subsequently determine the ‘partial’ momentum map. In the present setting, the associated sequence is

$$0 \rightarrow \text{gau}(P) \rightarrow \text{aut}_{\mu}(P) \rightarrow \mathfrak{X}_{\mu}(M) \rightarrow 0, \quad (3.21)$$

which naturally splits (in the category of locally convex vector spaces, not as Lie algebras) using a principal connection in P . Given a principal connection Γ in P , let $\Gamma: \mathfrak{X}_{\mu}(M) \rightarrow \text{aut}_{\mu}(P)$ be the vector space homomorphism that assigns to a vector field $X \in \mathfrak{X}_{\mu}(M)$ its horizontal lift $X^h \in \text{aut}_{\mu}(P)$. Recall that the curvature of Γ is the obstruction for the lift $\Gamma: \mathfrak{X}_{\mu}(M) \rightarrow \text{aut}_{\mu}(P)$ to be a Lie algebra homomorphism. By Proposition 2.35 applied to (3.21) with $\Gamma: \mathfrak{X}_{\mu}(M) \rightarrow \text{aut}_{\mu}(P)$ as the splitting, the ‘partial’ momentum map $\mathcal{J}_{\Gamma}: \mathcal{F} \rightarrow \mathcal{H}$ has to satisfy

$$(X^h)^* \lrcorner \Omega + \kappa(X, \delta \mathcal{J}_{\Gamma}) = 0, \quad X \in \mathfrak{X}_{\mu}(M) \quad (3.22)$$

for a suitable dual pair $\kappa(\text{Diff}_{\mu,P}(M), \mathcal{H})$. If we choose an extension $\hat{\omega}$ compatible with the connection Γ , then the second term in (3.9) vanishes and thus (3.22) simplifies to

$$(X \lrcorner \mu) \hat{*} \hat{\omega} + \kappa(X, \delta \mathcal{J}_{\Gamma}) = 0, \quad X \in \mathfrak{X}_{\mu}(M). \quad (3.23)$$

In general, it is hard to determine the ‘partial’ momentum map for the diffeomorphism group, mainly because one has to describe the dual group of $\text{Diff}_{\mu,P}(M)$ rather concretely. For this reason, we will only study the momentum map for the most important cases of volume-preserving and symplectic diffeomorphisms.

3.2.3 Momentum map for volume-preserving diffeomorphisms

Recall the exact sequence (3.7) and assume, for simplicity, that $\text{Diff}_{\mu,P}(M) = \text{Diff}_{\mu}(M)$. In Example 2.2 we showed that the dual group to $\text{Diff}_{\mu}(M)$ is the group of Cheeger–Simons differential characters $\hat{H}^2(M, \text{U}(1))$, which parametrizes principal $\text{U}(1)$ -bundles over M . Let $\kappa(\text{Diff}_{\mu}(M), \hat{H}^2(M, \text{U}(1)))$ be the resulting dual pair of Lie groups.

We continue to work in the general setting where $F = P \times_G \underline{F}$ is an associated fiber bundle endowed with a fiber symplectic structure ω induced by a symplectic form $\underline{\omega}$ on \underline{F} . Assume that $(\underline{F}, \underline{\omega})$ has a G -equivariant prequantization $\text{U}(1)$ -bundle $\underline{L} \rightarrow \underline{F}$ with connection $\underline{\vartheta}$. That is, G acts on \underline{L} by principal bundle automorphisms covering the action on \underline{F} , and the connection $\underline{\vartheta}$ is G -invariant.

Since G acts by bundle automorphisms, the actions of $U(1)$ and G commute. Consider the associated $U(1)$ -bundle $L := P \times_G \underline{L} \rightarrow F = P \times_G \underline{F}$. Since $L \rightarrow M$ is a fiber bundle with typical fiber \underline{L} , the vertical subbundle $VL = P \times_G T\underline{L}$ naturally projects onto $VF = P \times_G T\underline{F}$. The G -equivariant $U(1)$ -connection $\underline{\vartheta}$ in $\underline{L} \rightarrow \underline{F}$ yields an G -equivariant lift $T\underline{F} \rightarrow T\underline{L}$ and, thereby, induces a lift $\vartheta: VF \rightarrow VL$. The principal G -connection Γ on $P \rightarrow M$ completes ϑ to a $U(1)$ -connection ϑ^Γ in $L \rightarrow F$. Indeed, the associated connection 1-form on L is, under the identification $TL = TP \times_{TG} T\underline{L}$, given by

$$\vartheta_{[p, \underline{l}]}^\Gamma([Z_p^\Gamma + B_p^*, u_{\underline{l}}]) := \underline{\vartheta}_{\underline{l}}(B_{\underline{l}}^* + u_{\underline{l}}), \quad (3.24)$$

where Z_p^Γ is a Γ -horizontal vector at the point $p \in P$, $B \in \mathfrak{g}$, and $u_{\underline{l}} \in T\underline{L}$. By construction, ϑ^Γ coincides with ϑ on the vertical part and hence the curvature F_{ϑ^Γ} of ϑ^Γ , seen as a 2-form on F , is an extension of the fiber symplectic structure ω . In other words, (L, ϑ^Γ) yields a prequantization of the fiber symplectic bundle (F, ω) . We will set $\hat{\omega} = F_{\vartheta^\Gamma}$ in the general theory discussed above. Moreover, as discussed in Appendix D.1, the holonomy of (L, ϑ^Γ) yields a differential character $h_\Gamma \in \hat{H}^2(F, U(1))$.

The pull-back map

$$\text{pb}_h: \mathcal{F} \rightarrow \hat{H}^2(M, U(1)), \quad \phi \mapsto \phi^* h_\Gamma \quad (3.25)$$

has logarithmic derivative $(\delta \text{pb}_h)_\phi Y = \phi^*((Y \lrcorner F_{\vartheta^\Gamma}) \circ \phi) = \phi^*((Y \lrcorner \hat{\omega}) \circ \phi)$ according to Proposition D.3. On the other hand, using (C.2), we evaluate $(X \lrcorner \mu) \hat{*} \hat{\omega}$ on $Y \in T_\phi \mathcal{F}$ to obtain

$$((X \lrcorner \mu) \hat{*} \hat{\omega})_\phi(Y) = (-1)^{\dim M - 1} \int_M (X \lrcorner \mu) \wedge \phi^*((Y \lrcorner \hat{\omega}) \circ \phi). \quad (3.26)$$

Thus we have

$$(X \lrcorner \mu) \hat{*} \hat{\omega} = \kappa(X, \delta \text{pb}_h), \quad (3.27)$$

so that $\mathcal{J}_\Gamma := -\text{pb}_h: \mathcal{F} \rightarrow \hat{H}^2(M, U(1))$ satisfies the ‘partial’ momentum map relation (3.23).

We finally gathered all the necessary ingredients to apply Propositions 2.35 and 2.36.

THEOREM 3.3 (Momentum map in the volume-preserving case) *Let $\pi: P \rightarrow M$ be a finite-dimensional principal G -bundle over the closed volume manifold (M, μ) and assume that $\text{Diff}_{\mu, P}(M) = \text{Diff}_\mu(M)$. Then the following holds:*

- (i) *For every Hamiltonian G -manifold $(\underline{F}, \underline{\omega}, J)$, the space \mathcal{F} of sections of the associated symplectic fiber bundle $(F = P \times_G \underline{F}, \omega)$ is a symplectic Fréchet manifold with weak symplectic form $\Omega = \mu \hat{*} \omega$.*

- (ii) Choose a principal connection Γ in P . Assume that $(\underline{F}, \underline{\omega})$ has a G -equivariant prequantization $U(1)$ -bundle \underline{L} and denote the resulting prequantization differential character of (F, ω) by $h_\Gamma \in \hat{H}^2(F, U(1))$. The group $\text{Aut}_\mu(P)$ of bundle automorphisms, whose induced base diffeomorphisms are volume-preserving, acts symplectically on (\mathcal{F}, Ω) and has a group-valued momentum map

$$\mathcal{J}_{\text{Aut}}: \mathcal{F} \rightarrow \text{gau}^*(P) \times \hat{H}^2(M, U(1)), \quad \phi \mapsto (J_*\phi, -\phi^*h_\Gamma). \quad (3.28)$$

This momentum map is equivariant relative to the natural actions of $\text{Aut}_\mu(P)$ (see Example 2.30).

- (iii) A lift $\chi: \text{Diff}_\mu(M) \rightarrow \text{Aut}_\mu(P)$ of the group of volume-preserving diffeomorphisms to principal bundle automorphisms (this is a section of the projection in (3.7)) yields a symplectic group action of $\text{Diff}_\mu(M)$ on \mathcal{F} with $\hat{H}^2(M, U(1))$ -valued momentum map

$$\mathcal{J}_{\text{Diff}}: \mathcal{F} \rightarrow \hat{H}^2(M, U(1)), \quad \phi \mapsto -\phi^*h_\Gamma + \iota \circ \tau_{\Gamma, \chi}^* \circ J_*(\phi). \quad (3.29)$$

Here, $\iota: \hat{\mathfrak{h}}^2(M, U(1)) \rightarrow \hat{H}^2(M, U(1))$ denotes the natural inclusion of topological trivial bundles (see Appendix D.2). Moreover, $\tau_{\Gamma, \chi}^*: \text{gau}^*(P) \rightarrow \hat{\mathfrak{h}}^2(M, U(1))$ denotes the adjoint of $\tau_{\Gamma, \chi} := \chi' - \Gamma: \mathfrak{X}_\mu(M) \rightarrow \text{gau}(P)$ with respect to the dual pairs $\kappa(\mathfrak{X}_\mu(M), \hat{\mathfrak{h}}^2(M, U(1)))$ and $\langle \text{gau}(P), \text{gau}^*(P) \rangle_{\text{Ad}}$. Furthermore, $\mathcal{J}_{\text{Diff}}$ does not depend on the connection Γ that was used in the construction. \diamond

REMARK 3.4 In order to illustrate the meaning of the map $\tau_{\Gamma, \chi}$ and its adjoint, we compute it for the $\text{SL}(n, \mathbb{R})$ -frame bundle $\pi: LM \rightarrow M$. Every volume-preserving diffeomorphism ϕ lifts in a natural way to a diffeomorphism

$$\chi(\phi) = \hat{\phi}: LM \rightarrow LM, \quad u_m \mapsto T_m\phi \circ u_m, \quad (3.30)$$

where we view a frame $u_m \in L_m M$ as a linear isomorphism $u_m: \mathbb{R}^n \rightarrow T_m M$. Thus the frame bundle comes equipped with a natural lift $\chi: \text{Diff}_\mu(M) \rightarrow \text{Aut}_\mu(LM)$. Of course, we also get an induced lifting map for vector fields, which we write as $\chi': \mathfrak{X}_\mu(M) \ni X \mapsto \hat{X} \in \text{aut}_\mu(LM)$.

Recall that there exists a natural 1-form $\theta \in \Omega^1(LM, \mathbb{R}^n)$ defined by (see [KN63, Section III.2])

$$\theta(Z_u) = u^{-1} \circ T_u \pi(Z_u), \quad Z_u \in T_u(LM). \quad (3.31)$$

The form θ is invariant under naturally lifted vector fields, i.e., $\mathfrak{L}_{\hat{X}}\theta = 0$.

Let $\alpha \in \Omega^1(LM, \mathfrak{sl}(n, \mathbb{R}))$ be a principal connection 1-form on the bundle $LM \rightarrow M$ and denote by Γ the associated horizontal lift. The associated covariant derivative on the tangent bundle $TM \rightarrow M$ is defined by (see [KN63, Lemma on p. 133])

$$\nabla_Y X := u \circ \hat{Y}_u^\Gamma(\theta(\hat{X})), \quad u \in LM, \quad (3.32)$$

where \hat{Y}^Γ denotes the Γ -horizontal lift of $Y \in \mathfrak{X}(M)$. The torsion $t_\Gamma \in \Omega^2(LM, \mathbb{R}^n)$ of Γ is given as the covariant derivative of θ . The structure equation (see [KN63, Theorem III.2.4]) allows us to write

$$t_\Gamma(Z_1, Z_2) = d\theta(Z_1, Z_2) + \alpha(Z_1) \cdot \theta(Z_2) - \alpha(Z_2) \cdot \theta(Z_1), \quad (3.33)$$

where the \cdot in the last two terms denotes the natural action of $\mathfrak{sl}(n, \mathbb{R})$ on \mathbb{R}^n . We evaluate this expression for the case where $Z_1 = \hat{X} \in \mathfrak{X}(LM)$ is the natural lift of $X \in \mathfrak{X}(M)$ and $Z_2 = \hat{Y}^\Gamma \in \mathfrak{X}(LM)$ is the horizontal lift of $Y \in \mathfrak{X}(M)$:

$$\begin{aligned} t_\Gamma(\hat{X}, \hat{Y}^\Gamma) &= d\theta(\hat{X}, \hat{Y}^\Gamma) + \alpha(\hat{X}) \cdot \theta(\hat{Y}^\Gamma) \\ &= -\hat{Y}^\Gamma \lrcorner d(\hat{X} \lrcorner \theta) + \alpha(\hat{X}) \cdot u^{-1}(Y) \\ &= -u^{-1}(\nabla_Y X) + \alpha(\hat{X}) \cdot u^{-1}(Y), \end{aligned} \quad (3.34)$$

where we used $\mathfrak{L}_{\hat{X}}\theta = 0$. On the other hand, for $\tau_{\Gamma, \chi} = \chi' - \Gamma$, we have

$$\tau_{\Gamma, \chi}(X) = \alpha(\tau_{\Gamma, \chi}(X)) = \alpha(\hat{X} - \hat{X}^\Gamma) = \alpha(\hat{X}). \quad (3.35)$$

Combining both equations, we get

$$\begin{aligned} u(\tau_{\Gamma, \chi}(X) \cdot u^{-1}(Y)) &= u \circ t_\Gamma(\hat{X}, \hat{Y}^\Gamma) + \nabla_Y X \\ &= \nabla_X Y - [X, Y]. \end{aligned} \quad (3.36)$$

Thus, for torsion-free connections, we identify $\tau_{\Gamma, \chi}$ with the covariant derivative

$$\tau_{\Gamma, \chi}: \mathfrak{X}_\mu(M) \rightarrow \Gamma^\infty(T^*M \otimes TM), \quad X \mapsto \nabla X. \quad (3.37)$$

For a section s of $TM \otimes T^*M \rightarrow M$, the covariant derivative ∇s is a section of $(TM \otimes T^*M) \otimes T^*M \rightarrow M$. We obtain a 1-form $c(\nabla s)$ by contracting the first with the last component (i.e., the vector component with the "derivative" component; in Penrose's abstract index notation $c(\nabla s)_i = \nabla_j s^j{}_i$). The identity

$$\operatorname{div}(X \lrcorner s) = \langle \nabla X, s \rangle + X \lrcorner c(\nabla s) \quad (3.38)$$

yields the following expression for the adjoint operator of $\tau_{\Gamma, \chi}$:

$$\tau_{\Gamma, \chi}^*: \Gamma^\infty(TM \otimes T^*M) \rightarrow \Omega^1(M)/d\Omega^0(M), \quad s \mapsto -[c(\nabla s)]. \quad (3.39)$$

◇

REMARK 3.5 (Momentum map in the classical sense) If we compose the momentum map $\mathcal{J}_{\text{Diff}}: \mathcal{F} \rightarrow \hat{\mathcal{H}}^2(M, U(1))$ with the curvature map $\hat{\mathcal{H}}^2(M, U(1)) \rightarrow \Omega^2(M, \mathbb{R})$, then we recover Donaldson's expression for the "momentum map"

[Dono3, Equation 8]¹

$$\mathcal{F} \rightarrow \Omega^2(M, \mathbb{R}), \quad \phi \mapsto -\phi^* \hat{\omega} + d(\tau_{\Gamma, \chi}^* \circ J_*(\phi)). \quad (3.40)$$

Note, however, that $\Omega^2(M, \mathbb{R})$ is not the dual of the Lie algebra $\mathfrak{X}_\mu(M)$. In fact, it is only the dual of the Lie algebra of *exact* volume-preserving vector fields. Thus, (3.40) does not yield a bona-fide momentum map for the action of $\text{Diff}_\mu(M)$.

Although we were able to determine a group-valued momentum map for the diffeomorphism action under rather weak assumptions on the fiber model $(\underline{F}, \underline{\omega})$, a momentum map in the classical sense exists only under certain topological conditions on the fiber bundle F . Indeed, a necessary and sufficient condition for the existence of a momentum map $\mathcal{J}: \mathcal{F} \rightarrow (\mathfrak{X}_\mu(M))^*$ is that the obstruction map $\mathfrak{X}_\mu(M)/[\mathfrak{X}_\mu(M), \mathfrak{X}_\mu(M)] \rightarrow H^1(\mathcal{F}, \mathbb{R})$ sending the class of $X \in \mathfrak{X}_\mu(M)$ to the cohomology class of $X^* \lrcorner \Omega$ is identically zero (see, for example, [ORo3, Proposition 4.5.17]).

Suppose we have a lift $\chi: \text{Diff}_\mu(M) \rightarrow \text{Aut}_\mu(P)$. Then there is an induced lifting map for vector fields, which we write as $\mathfrak{X}_\mu(M) \ni X \mapsto \hat{X} \in \text{aut}_\mu(P)$. Let us assume² that \hat{X} is horizontal with respect to a principal connection on $P \rightarrow M$ compatible with the extension $\hat{\omega}$ and that $\text{Diff}_{\mu, P}(M) = \text{Diff}_\mu(M)$. In this setting, the obstruction to the existence of a classical momentum map takes the form

$$H^{\dim M - 1}(M, \mathbb{R}) \rightarrow H^1(\mathcal{F}, \mathbb{R}), \quad [\alpha] \mapsto [\alpha] \hat{*} [\hat{\omega}], \quad (3.41)$$

where we used the isomorphism $\mathfrak{X}_\mu(M)/[\mathfrak{X}_\mu(M), \mathfrak{X}_\mu(M)] \ni [X] \mapsto [X \lrcorner \mu] \in H^{\dim M - 1}(M, \mathbb{R})$, see [Ban97, Theorem 1.4.3]. We do not know what are the minimal conditions guaranteeing that the above obstruction (3.41) vanishes. Clearly, this is the case if the corresponding cohomology class of M vanishes. However, this is a rather strong topological assumption. \diamond

REMARK 3.6 (Topological data contained in the group-valued momentum map) The $\hat{H}^2(M, U(1))$ -valued momentum map contains two types of information. First, the curvature as a geometric datum essentially corresponds to the classical momentum map as discussed in Remark 3.5 above.

Second, the group-valued momentum map also yields topological classes that are not present in the classical momentum map. The Chern class

$$c(\mathcal{J}_{\text{Diff}}(\phi)) = -\phi^* c(h_\Gamma) \quad (3.42)$$

can be viewed as an obstruction to the existence of a classical momentum map. Indeed, if a classical momentum map with values in $\Omega^1(M)/d\Omega^0(M)$ exists, then

¹ Actually, Donaldson's expression for the momentum map differs by a sign, which comes from the fact that in [Dono3] the diffeomorphism group acts via a right action on the section space.

² These assumptions are satisfied, for example, for the trivial bundle $F = M \times \underline{F} \rightarrow M$.

its composition with the inclusion $\iota: \Omega^1(M)/d\Omega^0(M) \rightarrow \hat{H}^2(M, U(1))$ (see (D.6)) is a group-valued momentum map whose Chern class vanishes, because the image of ι lies in the kernel of the Chern class map c .

If the Chern class is trivial, then a secondary topological class related to the equivariance of the momentum map appears. This secondary class has its origin in the short exact sequence from (D.6)

$$0 \longrightarrow \Omega^1(M)/\Omega_{\text{cl},\mathbb{Z}}^1(M) \xrightarrow{\iota} \hat{H}^2(M, U(1)) \xrightarrow{c} H^2(M, \mathbb{Z}) \longrightarrow 0. \quad (3.43)$$

Exactness implies that the kernel of the characteristic class map c is isomorphic to $\Omega^1(M)/\Omega_{\text{cl},\mathbb{Z}}^1(M)$. However, this isomorphism is not canonical and involves a choice. For example, we may choose a basis of the singular homology $H_1(M, \mathbb{Z})$ given by piece-wise smooth loops γ_i . The curvature of a differential character $h \in \hat{H}^2(M, U(1))$ with trivial Chern class is an exact 2-form, say $\text{curv } h = d\alpha_h$. The primitive α_h is uniquely determined, modulo $\Omega_{\text{cl},\mathbb{Z}}^1(M)$, by the requirement that $\exp\left(2\pi i \int_{\gamma_i} \alpha_h\right) = h(\gamma_i)$ for all i . By construction, the map $\iota_\gamma^{-1}: h \mapsto [\alpha_h]$ is inverse to ι and thus yields the claimed isomorphism $\text{Ker } c \rightarrow \Omega^1(M)/\Omega_{\text{cl},\mathbb{Z}}^1(M)$ of Abelian Lie groups.

Given an $\hat{H}^2(M, U(1))$ -valued momentum map \mathcal{J} , whose image is contained in $\text{Ker } c$, the composition $\mathcal{J}_\gamma = \iota_\gamma^{-1} \circ \mathcal{J}$ is a $\Omega^1(M)/\Omega_{\text{cl},\mathbb{Z}}^1(M)$ -valued momentum map, i.e., almost a classical momentum map. Note that, even though \mathcal{J} is equivariant with respect to pull-backs by diffeomorphisms, the reduced momentum map \mathcal{J}_γ is no longer equivariant, because the loops γ_i are also affected by the diffeomorphism. The obstruction for the equivariance is the class in $H^1(M, U(1))$ given by evaluating h on the generators γ_i . We note that the same construction goes through in the slightly more general case, where the Chern class does not vanish but where it is a torsion class¹ (because then the curvature is an exact form as well).

We return to this secondary topological class in Chapter 4, where we recover, as an example, the Liouville class of a Lagrangian embedding.

In summary, if an equivariant $\hat{H}^2(M, U(1))$ -valued momentum map exists, then the existence of a classical equivariant momentum map is obstructed by two topological classes: the Chern class in $H^2(M, \mathbb{Z})$ and a class in $H^1(M, U(1))$. \diamond

3.2.4 Momentum map for symplectomorphisms

Let M be a closed $2n$ -dimensional manifold endowed with a symplectic form σ . The Liouville volume form is denoted by $\mu_\sigma = \sigma^n/n!$. Instead of the exact

¹An element $c \in H^k(M, \mathbb{Z})$ is called a torsion class if it belongs to the subgroup $\text{Ext}_{\text{Ab}}(H_{k-1}(M, \mathbb{Z}), \mathbb{Z}) \subseteq H^k(M, \mathbb{Z})$.

sequence (3.7), we consider the exact sequence

$$\text{id} \rightarrow \text{Gau}(P) \rightarrow \text{Aut}_\sigma(P) \rightarrow \text{Diff}_{\sigma,P}(M) \rightarrow \text{id}, \quad (3.44)$$

where $\text{Aut}_\sigma(P)$ is the group of automorphisms of the bundle $P \rightarrow M$ whose induced diffeomorphisms on the base p reserve σ . We assume that $\text{Diff}_{\sigma,P}(M) = \text{Diff}_\sigma(M)$. The momentum map is determined similarly to the volume-preserving case. Indeed, we only need to replace the dual pair $\kappa(\text{Diff}_\mu(M), \hat{H}^2(M, U(1)))$ by the dual pair $\kappa(\text{Diff}_\sigma(M), \hat{H}^{2n}(M, U(1)))$ appearing in Example 2.3.

Assume that σ is prequantizable and choose a prequantization circle bundle with connection $h_\sigma \in \hat{H}^2(M, U(1))$. As above, we assume that (\underline{F}, ω) has a G -equivariant prequantization bundle \underline{L} . A principal connection Γ in $P \rightarrow M$ yields a prequantization $h_\Gamma \in \hat{H}^2(F, U(1))$ of the symplectic fiber bundle (F, ω) . We claim that the ‘partial’ momentum map $\mathcal{J}_\Gamma: \mathcal{F} \rightarrow \hat{H}^{2n}(M, U(1))$ assigns to a section ϕ the degree $2n$ -bundle $-\phi^* h_\Gamma \star h_\sigma^{n-1}$, where the star product extends the wedge product to differential characters and the power $n - 1$ should also be understood in this sense, see (D.23). In order to verify that the ‘partial’ momentum map relation, cf. (3.23),

$$(X \lrcorner \mu_\sigma) \hat{\star} \hat{\omega} + \kappa(X, \delta \mathcal{J}_\Gamma) = 0, \quad X \in \mathfrak{X}_\sigma(M) \quad (3.45)$$

holds, we first observe that \mathcal{J}_Γ is the composition of the pull-back map pb_{h_Γ} with the Lie group homomorphism

$$\hat{H}^2(M, U(1)) \rightarrow \hat{H}^{2n}(M, U(1)), \quad h \mapsto h \star h_\sigma^{n-1}. \quad (3.46)$$

The corresponding Lie algebra homomorphism is given by $\alpha \mapsto \alpha \wedge \sigma^{n-1}$, because, using the properties listed in Appendix D.2, we have

$$\iota(\alpha) \star h_\sigma^{n-1} = \iota(\alpha \wedge \text{curv } h_\sigma^{n-1}). \quad (3.47)$$

Using the distribution law of the logarithmic derivative over Lie group homomorphisms [Nee06, Proposition II.4.1.1], we thus get

$$\begin{aligned} \kappa(X, \delta_\phi \mathcal{J}_\Gamma(Y)) &= - \frac{(-1)^{\dim M - 1}}{(n-1)!} \int_M (X \lrcorner \sigma) \wedge \phi^*((Y \lrcorner \hat{\omega}) \circ \phi) \wedge \sigma^{n-1} \\ &= - (-1)^{\dim M - 1} \int_M (X \lrcorner \sigma) \wedge \frac{\sigma^{n-1}}{(n-1)!} \wedge \phi^*((Y \lrcorner \hat{\omega}) \circ \phi) \\ &= - (-1)^{\dim M - 1} \int_M (X \lrcorner \mu_\sigma) \wedge \phi^*((Y \lrcorner \hat{\omega}) \circ \phi), \\ &= - ((X \lrcorner \mu_\sigma) \hat{\star} \hat{\omega})_\phi(Y), \end{aligned} \quad (3.48)$$

where the last equality follows from (C.2). This completes the proof that \mathcal{J}_Γ satisfies the ‘partial’ momentum map relation (3.45). Thus, Propositions 2.35 and 2.36 yield the following.

THEOREM 3.7 (Momentum map in the symplectic case) *Let $\pi: P \rightarrow M$ be a finite-dimensional principal G -bundle over the closed $2n$ -dimensional symplectic manifold (M, σ) and assume that $\text{Diff}_{\sigma, P}(M) = \text{Diff}_\sigma(M)$ in the exact sequence (3.44). Let $\mu_\sigma := \sigma^n / n!$ be the Liouville volume form on M . Then the following holds:*

- (i) *For every Hamiltonian G -manifold $(\underline{F}, \underline{\omega}, J)$, the space of sections \mathcal{F} of the associated symplectic fiber bundle $(F = P \times_G \underline{F}, \omega)$ is a symplectic Fréchet manifold with symplectic form $\Omega = \mu_\sigma \hat{*} \omega$.*
- (ii) *Choose a principal connection Γ in $P \rightarrow M$. Assume that $(\underline{F}, \underline{\omega})$ has a G -equivariant prequantization $U(1)$ -bundle \underline{L} and denote the resulting prequantization differential character of (F, ω) by $h_\Gamma \in \hat{H}^2(F, U(1))$. Assume that σ has a prequantization $h_\sigma \in \hat{H}^2(M, U(1))$. The group $\text{Aut}_\sigma(P)$ of bundle automorphisms whose induced base diffeomorphisms preserve the symplectic form σ acts symplectically on (\mathcal{F}, Ω) and has a group-valued momentum map*

$$\mathcal{J}_{\text{Aut}}: \mathcal{F} \rightarrow \text{gau}^*(P) \times \hat{H}^{2n}(M, U(1)), \quad \phi \mapsto (J_*\phi, -\phi^* h_\Gamma \star h_\sigma^{n-1}). \quad (3.49)$$

This momentum map is equivariant relative to the natural actions of $\text{Aut}_\sigma(P)$ (compare Example 2.30).

- (iii) *A lift $\chi: \text{Diff}_\sigma(M) \rightarrow \text{Aut}_\sigma(P)$ of the group of symplectomorphisms to principal bundle automorphisms (this is a section of the projection in (3.44)) yields a symplectic group action of $\text{Diff}_\sigma(M)$ on \mathcal{F} with $\hat{H}^{2n}(M, U(1))$ -valued momentum map*

$$\mathcal{J}_{\text{Diff}}: \mathcal{F} \rightarrow \hat{H}^{2n}(M, U(1)), \quad \phi \mapsto -\phi^* h_\Gamma \star h_\sigma^{n-1} + \iota \circ \tau_{\Gamma, \chi}^* \circ J_*(\phi). \quad (3.50)$$

Here, $\iota: \hat{h}^{2n}(M, U(1)) \rightarrow \hat{H}^{2n}(M, U(1))$ denotes the natural inclusion of topological trivial bundles (see Appendix D.2). Moreover, $\tau_{\Gamma, \chi}^: \text{gau}^*(P) \rightarrow \hat{h}^{2n}(M, U(1))$ denotes the adjoint of $\tau_{\Gamma, \chi} = \chi' - \Gamma: \mathfrak{X}_\sigma(M) \rightarrow \text{gau}(P)$ with respect to the dual pairs $\kappa(\mathfrak{X}_\sigma(M), \hat{h}^{2n}(M, U(1)))$ and $\langle \text{gau}(P), \text{gau}^*(P) \rangle_{\text{Ad}}$. Furthermore, $\mathcal{J}_{\text{Diff}}$ does not depend on the connection Γ that was used in the construction. \diamond*

REMARK 3.8 (Momentum map in the classical sense) *Consider the case of a trivial bundle $F = M \times \underline{F} \rightarrow M$. In this setting the term involving $\tau_{\Gamma, \chi}$ vanishes. After composing the momentum map $\mathcal{J}_{\text{Diff}}: \mathcal{F} \rightarrow \hat{H}^{2n}(M, U(1))$ with the curvature $\hat{H}^{2n}(M, U(1)) \rightarrow \Omega^{2n}(M, \mathbb{R})$ we get the map*

$$\mathcal{F} \rightarrow \Omega^{2n}(M, \mathbb{R}), \quad \phi \mapsto -\phi^* \hat{\omega} \wedge \sigma^{n-1}. \quad (3.51)$$

Donaldson [Don00, Proposition 3] found that this map is the momentum map for the action of the symplectomorphism group *under certain topological conditions* on M . However, we remark again that $\Omega^{2n}(M, \mathbb{R})$ is not the dual of the Lie algebra of symplectic vector fields $\mathfrak{X}_\sigma(M)$ but that it is only the dual of the space of *Hamiltonian* vector fields.

The existence of a momentum map in the classical sense is again obstructed by a map on the Abelianization of $\mathfrak{X}_\sigma(M)$. The map $\mathfrak{X}_\sigma(M)/[\mathfrak{X}_\sigma(M), \mathfrak{X}_\sigma(M)] \ni [X] \mapsto [X \lrcorner \sigma] \in H^1(M, \mathbb{R})$ is an isomorphism; see, [Ban97, Theorem 1.4.3]. In particular, if $H^1(M, \mathbb{R})$ is trivial, then a classical momentum map exists. \diamond

4 APPLICATIONS

4.1 Trivial bundle case

Let M be a closed manifold endowed with a volume form μ . We are interested in the space of smooth maps $\mathcal{F} = C^\infty(M, F)$ from M to a given Hamiltonian G -manifold (F, ω, J) . Of course, this set-up fits in the general theory developed in Chapters 2 and 3 by thinking of \mathcal{F} as the space of sections of the trivial bundle $M \times F$. The symplectic form on \mathcal{F} is expressed in this case as

$$\Omega_\phi(Y_1, Y_2) = \int_M \omega_\phi(Y_1, Y_2) \mu. \quad (4.1)$$

We consider the two natural actions

$$\begin{aligned} C^\infty(M, G) \times \mathcal{F} &\rightarrow \mathcal{F}, & \psi \cdot \phi &= \psi \cdot_G \phi, \\ \text{Diff}_\mu(M) \times \mathcal{F} &\rightarrow \mathcal{F}, & \varphi \cdot \phi &= \phi \circ \varphi^{-1}, \end{aligned} \quad (4.2)$$

where \cdot_G denotes the G -action on F . If (F, ω) has a G -equivariant prequantum bundle (L, ϑ) , then, by Theorem 3.3, the above actions have group-valued momentum maps

$$\begin{array}{ccc} & \mathcal{F} & \\ \mathcal{J}_G \swarrow & & \searrow \mathcal{J}_{\text{Diff}_\mu} \\ C^\infty(M, \mathfrak{g}^*) & & \hat{H}^2(M, U(1)) \end{array} \quad (4.3)$$

given by

$$\mathcal{J}_G(\phi) = J \circ \phi, \quad \mathcal{J}_{\text{Diff}_\mu}(\phi) = -\phi^*[L, \vartheta]. \quad (4.4)$$

Moreover, if the symplectic form is exact, say $\omega = d\vartheta$, then the Diff_μ -action admits an ordinary momentum map

$$\mathcal{J}_{\text{Diff}_\mu}: \mathcal{F} \rightarrow \Omega^1(M)/d\Omega^0(M), \quad \phi \mapsto -[\phi^*\vartheta]. \quad (4.5)$$

REMARK 4.1 (Extension to the action of $\text{Diff}_\omega(F)$) The group $\text{Diff}_\omega(F)$ of symplectomorphisms leaves the symplectic form on \mathcal{F} invariant and thus it is natural to seek a momentum map for it. This corresponds to replacing $\text{Aut}_\mu(P)$ by $\text{Aut}_{\mu,\omega}(F)$ in the general theory outlined in the previous chapter. However, we are confronted with serious problems if we try to determine the momentum map in this generality: Such a momentum map would take values in the *distributional* dual. As shown in [GV12], the action of the subgroup of Hamiltonian diffeomorphisms of F admits a momentum map $\mathcal{J}: \mathcal{F} \rightarrow \mathfrak{ham}(F, \omega)^*$ satisfying $\langle X_h, \mathcal{J}(\phi) \rangle = \int_M (\phi^*h) \mu$, after choosing a basis in $H^0(F, \mathbb{R})$ to identify Hamiltonian vector fields with smooth functions. However, the functional

$$T_\phi: C^\infty(F, \mathbb{R}) \rightarrow \mathbb{R}, \quad h \mapsto \int_M (\phi^*h) \mu \quad (4.6)$$

is usually not regular, i.e. not represented by integration against a smooth function on F . For example, if M is the one-point space $\{x\}$, then T_ϕ is the delta distribution at the point $\phi(x)$. The distributional character of the momentum map for the $\text{Diff}_\omega(F)$ -action is not surprising. Conceptually, the momentum map is the push-forward of μ , and the push-forward of a differential form into a higher-dimensional manifold results in a distributional functional.

In order to apply the same strategy as in the general theory above, we would need a Lie group whose Lie algebra consists of distributions; or, even better, we would need “distributional line bundles”, i.e., singular bundles whose curvature is a distribution. \diamond

4.1.1 Hydrodynamics

The Lie group-valued momentum map can be used to construct a generalized Clebsch representation of a vector field that cannot be expressed in terms of classical Clebsch variables.

Let us start by recalling the usual setting. Let M be a compact 3-dimensional orientable boundaryless Riemannian manifold, oriented by the Riemannian volume form μ . For simplicity, we also assume that M is connected. Euler’s equation for an incompressible perfect fluid on M is

$$\partial_t v + \nabla_v v = -\text{grad } p \quad (4.7)$$

for $v \in \mathfrak{X}(M)$ subject to the incompressibility condition $\mathfrak{L}_v \mu = 0$ (i.e., $\operatorname{div}_\mu v = 0$). By taking the Riemannian dual and using the identity

$$\mathfrak{L}_v(v^b) = (\nabla_v v)^b + \frac{1}{2} d\|v\|^2 \quad (4.8)$$

we can express Euler's equation as

$$\begin{cases} \partial_t \alpha + \mathfrak{L}_v \alpha = d\tilde{p}, \\ v^b = \alpha, \end{cases} \quad (4.9)$$

where $\tilde{p} := -p + \frac{1}{2}\|v\|^2$. The Hodge decomposition $\Omega^1(M) = \operatorname{Ker} d^* \oplus \operatorname{Im} d$ (which is L^2 -orthogonal) shows that the projection $\operatorname{pr}_{\operatorname{Ker} d^*}$ onto the kernel of the codifferential induces an isomorphism between divergence free vector fields and $\Omega^1(M)/d\Omega^0(M)$. The resulting isomorphism

$$A: \mathfrak{X}_\mu(M) \ni v \mapsto [v^b] \in \Omega^1(M)/d\Omega^0(M) \quad (4.10)$$

is called the *inertia operator*. This allows us to view Euler's equation as a dynamical system on $\Omega^1(M)/d\Omega^0(M)$:

$$\begin{cases} \partial_t [\alpha] + \mathfrak{L}_v [\alpha] = 0 \\ A(v) = [\alpha]. \end{cases} \quad (4.11)$$

It turns out that these equations are a Lie-Poisson system for the Hamiltonian $\mathcal{H}([\alpha]) = \frac{1}{2} \int_M \|A^{-1}[\alpha]\|^2 \mu$, see [MW83].

Clebsch variables give a parametrization of $[\alpha] \in \Omega^1(M)/d\Omega^0(M)$ in such a way that Euler's equation is cast into a classical (i.e., symplectic) Hamiltonian form. We refer the reader to [GV12; MW83] for more details on the following geometric construction of the classical Clebsch representation. Let $(F, \omega = d\vartheta)$ be an exact symplectic manifold of dimension 2 and let \mathcal{F} denote the space of smooth functions from M to F . As we have seen in (4.5), the map

$$\mathcal{J}: \mathcal{F} \rightarrow \Omega^1(M)/d\Omega^0(M), \quad \phi \mapsto -[\phi^* \vartheta] \quad (4.12)$$

is an equivariant momentum map for the action of $\operatorname{Diff}_\mu(M)$ on \mathcal{F} . Since an equivariant momentum map is Poisson, \mathcal{J} maps solutions of the Hamiltonian system $(\mathcal{F}, \Omega, \mathcal{H} \circ \mathcal{J})$ to solutions of Euler's equation. For the velocity field, \mathcal{J} gives the following representation in terms of Clebsch variables:

$$v^b = -\operatorname{pr}_{\operatorname{Ker} d^*} \phi^* \vartheta = -\phi^* \vartheta + df, \quad (4.13)$$

where $f \in C^\infty(M)$ ensures that the resulting vector field is divergence free (such

a function always exists by the Hodge decomposition $\Omega^1(M) = \text{Ker } d^* \oplus \text{Im } d$. For example, in the case $F = \mathbb{R}^2$ with $\vartheta = -x \, dy$ we obtain

$$v^b = \phi_1 \, d\phi_2 + df, \quad (4.14)$$

which is the classical Clebsch representation for the velocity field [Cle57; Cle59].

If a vector field admits a representation in terms of Clebsch variables, then its helicity has to vanish. Indeed, since $\dim F = 2$ and $\partial M = \emptyset$, we have

$$\text{Hel}(v) := \int_M v^b \wedge dv^b = \int_M \phi^*(\vartheta \wedge d\vartheta) = 0. \quad (4.15)$$

Helicity measures the linkage and/or knottedness of vortex lines. Hence the helicity constraint implies that topological non-trivial vector fields do not admit a classical Clebsch representation.

We show in the rest of this subsection how the group-valued momentum map gives a generalized Clebsch representation for vector fields with integral helicity. We continue to work in the general setting where (M, μ) is a compact 3-dimensional manifold endowed with a volume form μ and where (F, ω) is a 2-dimensional symplectic manifold. However, instead of requiring that ω is exact, we now only assume that it has integral periods. Thus, there exists a prequantum $U(1)$ -bundle $L \rightarrow F$ with connection ϑ and curvature ω . By (4.4) in Section 4.1, the group-valued momentum map for the $\text{Diff}_\mu(M)$ -action is given by

$$\mathcal{J}: \mathcal{F} \rightarrow \hat{H}^2(M, U(1)) \quad \phi \mapsto -\phi^*[L, \vartheta]. \quad (4.16)$$

Suppose there exists $\phi_o \in \mathcal{F}$ such that the pull-back bundle ϕ_o^*L over M is trivial. Then also for every $\phi \in \mathcal{F}$ in the connected component \mathcal{F}_{ϕ_o} of ϕ_o , the bundle ϕ^*L is trivial, because the composition

$$\mathcal{F} \xrightarrow{\mathcal{J}} \hat{H}^2(M, U(1)) \xrightarrow{c} H^2(M, \mathbb{Z}) \quad (4.17)$$

is a continuous map. Hence the restriction $\mathcal{J}|_{\mathcal{F}_{\phi_o}}$ of \mathcal{J} to \mathcal{F}_{ϕ_o} takes values in $c^{-1}(0) \simeq \Omega^1(M)/\Omega_{\text{cl},\mathbb{Z}}^1(M)$. By duality, we get the following *generalized Clebsch representation* of a vector field

$$v^b = -\phi^*\vartheta + v, \quad (4.18)$$

where $v \in \Omega_{\text{cl},\mathbb{Z}}^1(M)$ and, with a slight abuse of notation, $\phi^*\vartheta$ denotes the 1-form on M that is induced by the pull-back connection $\phi^*\vartheta$ in the trivial bundle ϕ^*L .

We note that Euler's equation descend to $\Omega^1(M)/\Omega_{\text{cl},\mathbb{Z}}^1(M)$. In fact, the Lie

derivative of a 1-form β with integral periods is exact since

$$\int_{\gamma} \mathfrak{L}_X \beta = \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \underbrace{\int_{\text{FL}_{\varepsilon}^X(\gamma)} \beta}_{\in \mathbb{Z}} = 0 \quad (4.19)$$

for every closed curve γ in M . Thus Euler's equation (4.11) implies that the class $[\beta] \in \Omega_{\text{cl}, \mathbb{Z}}^1(M)/d\Omega^0(M) \simeq H^1(M, \mathbb{Z})$ is constant in time, and that thus the interesting part of the motion takes place in $\Omega^1(M)/\Omega_{\text{cl}, \mathbb{Z}}^1(M)$. We note, however, that it is only possible to define an inertia operator $\Omega^1(M)/\Omega_{\text{cl}, \mathbb{Z}}^1(M) \rightarrow \mathfrak{X}_{\mu}(M)$ for small times since the principal bundle $H^1(M, \mathbb{R}) \rightarrow H^1(M, \mathbb{R})/H^1(M, \mathbb{Z})$ is only locally trivial.

In order to show that the helicity of a vector field admitting a generalized Clebsch representation has to be an integer, we start with the following general remarks about the helicity. On abstract grounds, the helicity functional is defined by

$$\text{Hel}: \Omega^1(M) \ni \alpha \mapsto \int_M \alpha \wedge d\alpha \in \mathbb{R}. \quad (4.20)$$

Note that Hel vanishes on closed forms. If one forgets the hydrodynamical context for a moment, then one would mistake such an expression as the Abelian Chern–Simons functional of a trivial bundle. In fact, the similarity between helicity and Chern–Simons theory was already pointed out previously in [LR12]. The Chern–Simons functional admits a well-known generalization to non-trivial circle bundles; see, e.g., [FMS07, Eq. 1.28]. In fact, for a differential character $h \in \hat{H}^2(M, U(1))$, the square $h \star h \in \hat{H}^4(M, U(1))$ is a group homomorphism $Z_3(M, \mathbb{Z}) \rightarrow U(1)$. We can evaluate it, in particular, on the 3-cycle M . Writing the evaluation as integration in order to keep the similarity with (4.20), we obtain the *generalized helicity* (or generalized Chern–Simons functional)

$$\text{Hel}: \hat{H}^2(M, U(1)) \ni h \mapsto \int_M h \star h \in U(1). \quad (4.21)$$

The properties of the star product (see Appendix D.2) yield the following commutative diagram connecting the ordinary helicity with its generalized version:

$$\begin{array}{ccc} \Omega^1(M)/\Omega_{\text{cl}, \mathbb{Z}}^1(M) & \xrightarrow{\text{Hel}} & \mathbb{R} \\ \downarrow \iota & & \downarrow \exp \\ \hat{H}^2(M, U(1)) & \xrightarrow{\text{Hel}} & U(1). \end{array} \quad (4.22)$$

We finally observe that the generalized helicity of elements in the image of the group-valued momentum map \mathcal{J} vanishes, due to a dimension count similar to the one leading to the identity (4.15), namely,

$$\begin{aligned} (\text{Hel} \circ \mathcal{J})(\phi) &= \text{Hel}(-\phi^*[L, \vartheta]) = \int_M (\phi^*[L, \vartheta]) \star (\phi^*[L, \vartheta]) \\ &= \int_M \phi^*([L, \vartheta] \star [L, \vartheta]) = 1. \end{aligned} \quad (4.23)$$

Hence, the commutative diagram (4.22) implies that the ordinary helicity of $\phi^*\vartheta$ has to be an integer. In other words, a vector field admitting a generalized Clebsch representation of the form (4.18) must have integral helicity.

We thus arrive at the following.

THEOREM 4.2 *Let (M, μ) be a compact 3-dimensional manifold endowed with a volume form μ . A solution $v(t) \in \mathfrak{X}_\mu(M)$ of Euler's equation (4.7) for an incompressible perfect fluid on M projects via $\mathfrak{X}_\mu(M) \ni v \mapsto [v^b] \in \Omega^1(M)/\Omega_{\text{cl},\mathbb{Z}}^1(M)$ to a solution of the system*

$$\begin{cases} \partial_t[\alpha] + \mathfrak{L}_v[\alpha] = 0, \\ [\alpha] = [v^b]. \end{cases} \quad (4.24)$$

on $\Omega^1(M)/\Omega_{\text{cl},\mathbb{Z}}^1(M)$.

Let (F, ω) be a 2-dimensional symplectic manifold that has a prequantum $U(1)$ -bundle $L \rightarrow F$ with connection ϑ . The natural action of $\text{Diff}_\mu(M)$ on $\mathcal{F} = C^\infty(M, F)$ preserves the symplectic form $\mu \hat{*} \omega$ and has a group-valued momentum map

$$\mathcal{J}: \mathcal{F} \rightarrow \hat{H}^2(M, U(1)), \quad \phi \mapsto -\phi^*[L, \vartheta]. \quad (4.25)$$

Moreover, assume there exists $\phi_0 \in \mathcal{F}$ such that the pull-back bundle ϕ_0^*L over M is trivial. Then the restriction of \mathcal{J} to the connected component \mathcal{F}_{ϕ_0} of ϕ_0 takes values in $\Omega^1(M)/\Omega_{\text{cl},\mathbb{Z}}^1(M)$ and yields generalized Clebsch variables for the projected Euler equation (4.24) in the sense of (4.18). Every vector field admitting such a generalized Clebsch representation has integral helicity. \diamond

EXAMPLE 4.3 For concreteness, let $M = S^3$, and consider $F = S^2$ endowed with its usual normalized volume ω . We take the Hopf bundle (L, ϑ) as the prequantization of (S^2, ω) . Since S^3 has trivial cohomology in degree 2, the pull-back of L by every map $\phi: S^3 \rightarrow S^2$ is trivial. Many interesting and topological non-trivial knots on S^3 can be constructed from the pull-back of ϑ by a special choice of ϕ . For example, the Hopf vector field [AK98, Example III.1.9] corresponds to the classical Hopf fibration

$$\phi: \mathbb{C}^2 \supseteq S^3 \ni (z_1, z_2) \mapsto [z_1: z_2] \in \mathbb{C}P^1 \simeq S^2. \quad (4.26)$$

Other knots that can be realized in this way include the figure of eight knot, linked rings, and the trefoil knot, among others; see [Ked+16] for details. The helicity of such vector fields coincides with the Hopf invariant of ϕ . Hence, these knots have a non-vanishing but integral helicity. Hence they *do not* admit a classical Clebsch representation, but have, by construction, a generalized Clebsch representation. \diamond

The following example suggests that not every vector field with integral helicity admits a generalized Clebsch representation and that one might need to replace the symplectic manifold F by a singular symplectic stratified space to ensure such a representation for every vector field with integral helicity.

EXAMPLE 4.4 Consider the vector field $v = A \sin z \partial_x + A \cos z \partial_y$ on \mathbb{R}^3 . Since v is 2π -periodic, it naturally lives on the torus $M = T^3$. Note that v is a special case of an Arnold–Beltrami–Childress (ABC) flow with $B = 0 = C$ in the usual notation. In particular, v provides an example for a simple steady-state solution of Euler’s equation. The vorticity of v is given by $\omega = A \cos z dz \wedge dx + A \sin z dy \wedge dz$, so that its helicity is

$$\text{Hel}(v) = \int_{T^3} v^b \wedge \omega = \int_{T^3} A^2 \text{vol} = A^2 \text{vol}(T^3). \quad (4.27)$$

Seen as a vector field on \mathbb{R}^3 , we can represent v as

$$v^b = f dg + dh, \quad \begin{cases} f = y \sin z - x \cos z, \\ g = z, \\ h = x \sin z + y \cos z. \end{cases} \quad (4.28)$$

The pair (f, g) , seen as a map $\mathbb{R}^3 \rightarrow \mathbb{R}^2$, is \mathbb{Z}^3 -equivariant with respect to the natural action on \mathbb{R}^3 and the twisted action on \mathbb{R}^2 , where the group element $(n_1, n_2, n_3) \in \mathbb{Z}^3$ acts by

$$\mathbb{R}^2 \ni (u, v) \mapsto (2\pi n_2 \sin v - 2\pi n_1 \cos v, v + 2\pi n_3) \in \mathbb{R}^2. \quad (4.29)$$

Let $F = \mathbb{R}^2/\mathbb{Z}^3$ be the orbit space. The representation (4.28) suggests that the induced map $(f, g): T^3 \rightarrow F$ yields generalized Clebsch variables for v . Note, however, that the \mathbb{Z}^3 -action on \mathbb{R}^2 is not free and that the orbit space F thus has singularities. \diamond

REMARK 4.5 The classical Clebsch representation admits a natural generalization (sometimes called a Weber–Clebsch representation)

$$v^b = \sum_i \phi_i d\psi_i + df, \quad (4.30)$$

where one allows for multiple potentials, see e.g. [GHoo]. In our setting, this corresponds to considering higher-dimensional symplectic manifolds F . We claim that every vector field admits a generalized Clebsch representation if we allow higher-dimensional targets F . To see this, let v be a vector field on M and let $v^b \in \Omega^1(M)$ be the associated 1-form. We may view v^b as a connection 1-form on the trivial bundle $L = M \times U(1) \rightarrow M$. The theory of universal bundles [NR61, Theorem 1] entails that there exists a smooth map $\phi: M \rightarrow \mathbb{C}P^N$ for sufficient large N such that the bundle (L, v^b) with connection is isomorphic to the pull-back of the canonical bundle over $\mathbb{C}P^N$ endowed with the canonical connection ϑ_{can} . Hence, there exists a gauge transformation $f \in C^\infty(M, U(1))$ of L such that

$$v^b = \phi^* \vartheta_{\text{can}} + \delta f, \quad (4.31)$$

where we again identified the connection $\phi^* \vartheta_{\text{can}}$ in the trivial bundle L with a 1-form on M . \diamond

4.1.2 Isotropic and Lagrangian embeddings

By a slight modification of the previous set-up, the moduli space of Lagrangian embeddings can be realized as a symplectic quotient. Let (S, μ) be a closed manifold with volume form, and let (M, ω) be a symplectic manifold. Since the space of smooth embeddings of S into M is an open subset of $C^\infty(S, M)$ [KM97, Theorem 44.1.], the symplectic form $\Omega = \omega \hat{*} \mu$ on $C^\infty(S, M)$ restricts to a symplectic form on $\mathcal{E}mb(S, M)$. We assume that (M, ω) has a prequantization bundle (L, ϑ) . By (4.4), the momentum map for the action of $\text{Diff}_\mu(S)$ is given by

$$\mathcal{J}_{\text{Diff}_\mu}: \mathcal{E}mb(S, M) \rightarrow \hat{H}^2(S, U(1)), \quad \iota \mapsto -\iota^*[L, \vartheta]. \quad (4.32)$$

Thus, the inverse image under $\mathcal{J}_{\text{Diff}_\mu}$ of the normal subgroup of all flat circle bundles on S consists precisely of isotropic embeddings. For $\dim(S) = \dim M/2$, the set $\mathcal{J}_{\text{Diff}_\mu}^{-1}(\text{flat})$ corresponds to Lagrangian embeddings. More generally, the curvature $-\iota^*\omega$ of $\mathcal{J}_{\text{Diff}_\mu}(\iota)$ vanishes if and only if ι is an isotropic embedding. Nonetheless, the Chern class of $\iota^*[L, \vartheta]$ may still be non-trivial in this case. The exact sequence (D.6) implies that $c(\iota^*[L, \vartheta])$ has to be completely torsion. We are not aware that this torsion class in $H^2(S, \mathbb{Z})$ is discussed anywhere in the literature.

If the symplectic form is exact, say $\omega = d\vartheta$, then the Chern class of $\iota^*[L, \vartheta]$ vanishes. Nonetheless, the group-valued momentum map contains secondary topological data which corresponds to the well-known Liouville class as we will see now. Indeed, then the line bundle (L, ϑ) is trivial so that the momentum map takes values in the subgroup of topological trivial bundles. Hence $\iota^*[L, \vartheta]$ is completely characterized by its curvature $\iota^*\omega$ and a class in the torus

$H^1(S, \mathbb{R})/H^1(S, \mathbb{Z})$; see (D.6). For an isotropic embedding ι , the curvature $\iota^*\omega$ vanishes and the topological class equals $[\iota^*\vartheta]$ and hence it coincides with the *Liouville class* of the embedding ι [BW97, p. 30].

Recall that an isotropic embedding is called *prequantizable* if its Liouville class vanishes; see [BW97, Definition 4.4]. More generally, we say that an isotropic embedding is *torsion-free* if the torsion class $c(\iota^*[L, \vartheta])$ in $H^2(S, \mathbb{Z})$ discussed above vanishes. Summarizing, we arrive at the following correspondence:

Moduli space of	Symplectic reduction $\mathcal{J}_{\text{Diff}_\mu}^{-1}(\cdot)/\text{Diff}_\mu(S)$ at
Lagrangian embeddings	flat bundles
Torsion-free Lagrangian embeddings	flat and trivial bundles
Prequantizable Lagrangian embeddings	0

Here, we are a bit sloppy with the notion “moduli space”. As we only take the quotient by the group $\text{Diff}_\mu(S) \subseteq \text{Diff}(S)$, the resulting reduced space still remembers the volume form. Hence, being more precise, the symplectic quotient $\mathcal{J}_{\text{Diff}_\mu}^{-1}(\cdot)/\text{Diff}_\mu(S)$ yields the moduli space of Lagrangian (or isotropic) weighted submanifolds of type (S, μ) .

In the case of an exact symplectic form, a non-equivariant momentum map for the action of the diffeomorphism group $\text{Diff}_\mu(S)$ is discussed in [MS99, Example 5.32]. Recently, Gay-Balmaz and Vizman [GV19] showed that the quotient of isotropic embeddings modulo volume-preserving diffeomorphisms is indeed a symplectic manifold and that its connected components correspond to coadjoint orbits of $\text{Ham}(M, \omega)$. These results are obtained under the additional assumptions that the first cohomology group of S vanishes. It would be of interest to study the symplectic quotient without this topological assumption and search for relations with coadjoint orbits of the whole symplectomorphism group. In light of our refined results, one would like to conclude:

CONJECTURE 4.6 The moduli space of weighted prequantizable Lagrangian embeddings is a symplectic manifold whose connected components are related to the coadjoint orbits of the symplectomorphism group. \diamond

Moreover, it is an interesting question whether the moduli space of (torsion-free) Lagrangian embeddings is also symplectic. A systematic study along the same lines as [GV19] requires to develop the dual pair picture for group-valued momentum maps.

4.2 Reduction of structure group

Geometric objects usually correspond to the reduction of a principal G -bundle to a Lie subgroup $H \subseteq G$. In this setting, the fiber model \underline{F} is the homogeneous

space G/H . A section ϕ of the associated bundle $F = P \times_G (G/H)$ results in a reduction of $\pi: P \rightarrow M$ to the principal H -bundle

$$Q_\phi := \{p \in P \mid \phi(\pi(p)) = [p, eH]\} \rightarrow M. \quad (4.33)$$

Conversely, every reduction of P to a H -subbundle defines a section of F .

Let μ be a volume form on the base manifold M of P . If $\underline{F} = G/H$ is endowed with a G -invariant symplectic form $\underline{\omega}$ then, according to Proposition 3.1, the space of H -reductions of P carries a natural symplectic structure $\Omega = \mu \hat{*} \omega$ with ω being the induced fiber symplectic structure on F . In this subsection, we discuss the content of Theorem 3.3 for this particular setting.

4.2.1 Pull-back of prequantum bundles as associated bundles

As in the general case, assume that $(\underline{F} = G/H, \underline{\omega}, J)$ is a Hamiltonian G -space. Let $\underline{L} \rightarrow \underline{F}$ be a G -equivariant prequantum bundle, and let $L = P \times_G \underline{L}$. As a preparation, we now discuss how the pull-back bundle ϕ^*L can be identified with an associated bundle to Q_ϕ for every section ϕ of F . Evaluating the momentum map $J: \underline{F} \rightarrow \mathfrak{g}^*$ at the point eH yields a Lie algebra homomorphism

$$J(eH)|_{\mathfrak{h}}: \mathfrak{h} \rightarrow \mathbb{R}. \quad (4.34)$$

It is a well-known fact in prequantization theory [Kos70, Theorem 5.7.1] that G -equivariant prequantizations over G/H are in one-to-one correspondence with Lie group homomorphisms $\rho: H \rightarrow U(1)$ integrating $J(eH)|_{\mathfrak{h}}$. Indeed, given an equivariant prequantization bundle \underline{L} , such a homomorphism can be extracted from the circle action in the fiber over eH , that is

$$h \cdot \underline{l} = \underline{l} \cdot \rho(h), \quad \underline{l} \in \underline{L}_{eH}. \quad (4.35)$$

Conversely, a Lie group homomorphism $\rho: H \rightarrow U(1)$ yields a G -equivariant prequantization bundle with connection as the associated bundle $\underline{L} = G \times_\rho U(1)$ to the canonical principal H -bundle $G \rightarrow G/H$. The G -action on \underline{L} is given by left translation in the G -factor, i.e., $g \cdot [a, z] = [ga, z]$, and the connection 1-form $\underline{\vartheta}$ on \underline{L} is defined by

$$\underline{\vartheta}_{[a, z]}([a \cdot A, v]) = \langle J(eH), A \rangle + v, \quad (4.36)$$

where $A \in \mathfrak{g}$ and $v \in T_z U(1) \simeq \mathbb{R}$.

Now we establish a natural isomorphism $\phi^*L \simeq Q_\phi \times_\rho U(1)$. Since the left G - and the right H -actions on G commute, we can identify the bundle $L = P \times_G \underline{L} \rightarrow F$

with the quotient of $P \times G \times U(1)$ by the simultaneous actions of G and H , namely,

$$(p, a, z) \cdot (g, h) := (p \cdot g, g^{-1}ah, \rho(h^{-1})z). \quad (4.37)$$

From this viewpoint, the projection $L \rightarrow F$ maps $[p, a, z]$ to $[p, aH]$. Thus, for a section ϕ of $F \rightarrow M$, the pull-back bundle is identified with

$$\phi^*L = \{(m, [p, a, z]) \in M \times L \mid \phi(m) = [p, aH]\}. \quad (4.38)$$

Recall that Q_ϕ denotes the reduced H -bundle defined in (4.33).

LEMMA 4.7 *The map*

$$\phi^*L \rightarrow Q_\phi \times_\rho U(1), \quad (m, [p, a, z]) \mapsto [p \cdot a, z] \quad (4.39)$$

is a principal bundle isomorphism over M . \diamond

Proof. First, note that $p \cdot a \in Q_\phi$, because

$$\phi(\pi(p \cdot a)) = \phi(m) = [p, aH] = [p \cdot a, eH]. \quad (4.40)$$

Second, the definition of the map in (4.39) does not depend on the chosen representative. Indeed, the point $(m, [p \cdot g, g^{-1}ah, \rho(h^{-1})z])$ is sent under (4.39) to $[p \cdot g g^{-1}ah, \rho(h^{-1})z] = [p \cdot a, z]$. Third, the vector bundle map

$$Q_\phi \times_\rho U(1) \rightarrow \phi^*L, \quad [q, z] \mapsto (\pi(q), [q, e, z]) \quad (4.41)$$

is clearly inverse to (4.39), so that we obtain a principal bundle isomorphism $\phi^*L \simeq Q_\phi \times_\rho U(1)$, as claimed. \square

The next lemma shows that this isomorphism also identifies the natural connections on both bundles. Recall from (3.24) the connection ϑ^Γ on L constructed from a connection Γ in P and the connection $\underline{\vartheta}$ on \underline{L} .

LEMMA 4.8 *Let Γ be a principal connection on P and assume that it reduces to a connection on Q_ϕ . Then the pull-back of the connection ϑ^Γ from L to ϕ^*L coincides with the induced connection $\rho_*\Gamma$ on the associated bundle $Q_\phi \times_\rho U(1)$ under the identification $\phi^*L \simeq Q_\phi \times_\rho U(1)$ of Lemma 4.7. \diamond*

Proof. We have seen that L can be realized as the quotient of $P \times G \times U(1)$ by the simultaneous actions of G and H . From this point of view, the connection ϑ^Γ on L takes the form

$$\begin{aligned} \vartheta_{[p,a,z]}^\Gamma([Z_p^\Gamma + p \cdot B, a \cdot A, v]) &= \underline{\vartheta}_{[a,z]}(B \cdot [a, z] + [a \cdot A, v]) \\ &= \underline{\vartheta}_{[a,z]}([a \cdot \text{Ad}_a^{-1} B + a \cdot A, v]) \\ &= \langle \underline{J}(eH), \text{Ad}_a^{-1} B + A \rangle + v, \end{aligned} \quad (4.42)$$

where $p \in P$, $a \in G$, $z \in U(1)$, $Z_p^\Gamma \in T_p P$ is horizontal, $B \in \mathfrak{g}$ and $z \in T_z U(1) \simeq \mathbb{R}$. Note that the following diagram of fiber bundle morphisms is commutative

$$\begin{array}{ccccc}
 Q_\phi \times_\rho U(1) & \longrightarrow & \phi^* L & \longrightarrow & L \\
 [q, z] & \longmapsto & (\pi(q), [q, e, z]) & \longmapsto & [q, e, z] \\
 \downarrow & & \downarrow & & \downarrow \\
 \pi(q) & \longmapsto & \pi(q) & \xrightarrow{\phi} & [q, eH] \\
 M & \xrightarrow{\text{id}} & M & \xrightarrow{\phi} & F
 \end{array} \tag{4.43}$$

Thus the pull-back of ϑ^Γ to $Q_\phi \times_\rho U(1)$ is given by

$$(\phi^* \vartheta^\Gamma)_{[q, z]}([Z_q^\Gamma + q \cdot B, v]) = \vartheta_{[q, e, z]}^\Gamma([Z_q^\Gamma + q \cdot B, v]) = \langle \underline{J}(eH), B \rangle + v. \tag{4.44}$$

On the other hand, recall that $\underline{J}(eH)|_{\mathfrak{h}}$ is, by definition, the derivative of the Lie group homomorphism ρ . Hence, the connection $\rho_* \Gamma$ on $Q_\phi \times_\rho U(1)$ induced by Γ satisfies

$$(\rho_* \Gamma)_{[q, z]}([Z_q^\Gamma + q \cdot B, v]) = T_e \rho(B) + v = \langle \underline{J}(eH), B \rangle + v. \tag{4.45}$$

Thus, the connections $\phi^* \vartheta^\Gamma$ and $\rho_* \Gamma$ coincide. \square

Thus, we get the following as a special case of Theorem 3.3.

THEOREM 4.9 (Momentum map for reductions of structure group) *Let $\pi: P \rightarrow M$ be a finite-dimensional principal G -bundle over the closed volume manifold (M, μ) and assume that $\text{Diff}_{\mu, P}(M) = \text{Diff}_\mu(M)$. Let H be a Lie subgroup of G and let $\underline{\omega}$ be a G -invariant symplectic form on G/H with equivariant momentum map $\underline{J}: G/H \rightarrow \mathfrak{g}^*$. Then the following holds:*

- (i) *The space \mathcal{F} of reductions of P to a principal H -subbundle is a symplectic Fréchet manifold with weak symplectic form $\Omega = \mu \hat{*} \omega$, where ω is the fiber symplectic form on $P \times_G (G/H)$ induced by $\underline{\omega}$.*
- (ii) *The group $\text{Aut}_\mu(P)$ of bundle automorphisms, whose induced base diffeomorphisms are volume-preserving, acts symplectically on (\mathcal{F}, Ω) . Moreover, for a principal connection Γ in P that is reducible to each Q_ϕ and for a Lie group homomorphism $\rho: H \rightarrow U(1)$ integrating $\underline{J}(eH)|_{\mathfrak{h}}$, the $\text{Aut}_\mu(P)$ -action on \mathcal{F} has a group-valued momentum map $\mathcal{J}_{\text{Aut}}: \mathcal{F} \rightarrow \mathfrak{gau}^*(P) \times \hat{H}^2(M, U(1))$ given by*

$$\mathcal{J}_{\text{Aut}}(\phi) = \left(J_* \phi, [Q_\phi \times_{\rho^{-1}} U(1), (\rho^{-1})_* \Gamma] \right), \tag{4.46}$$

where $J: P \times_G (G/H) \rightarrow \text{Ad}^* P$ is the bundle map induced by \underline{J} . The momentum map \mathcal{J}_{Aut} is equivariant relative to the natural actions of $\overline{\text{Aut}}_\mu(P)$ (see

Example 2.30).

- (iii) A lift $\chi: \text{Diff}_\mu(M) \rightarrow \text{Aut}_\mu(P)$ of the group of volume-preserving diffeomorphisms to principal bundle automorphisms yields a symplectic group action of $\text{Diff}_\mu(M)$ on \mathcal{F} with group-valued momentum map $\mathcal{J}_{\text{Diff}}: \mathcal{F} \rightarrow \hat{\mathcal{H}}^2(M, \text{U}(1))$ given by

$$\mathcal{J}_{\text{Diff}}(\phi) = [Q_\phi \times_{\rho^{-1}} \text{U}(1), (\rho^{-1})_* \Gamma] + \iota \circ \tau_{\Gamma, \chi}^* \circ J_*(\phi), \quad (4.47)$$

where $\iota: \hat{\mathfrak{h}}^2(M, \text{U}(1)) \rightarrow \hat{\mathcal{H}}^2(M, \text{U}(1))$ and $\tau_{\Gamma, \chi}^*: \mathfrak{gau}^*(P) \rightarrow \hat{\mathfrak{h}}^2(M, \text{U}(1))$ are the same maps as in Theorem 3.3. \diamond

Clearly there is also a version of this theorem for symplectic base manifolds similar to Theorem 3.7. We leave it to the interested reader to spell out the details.

In the following, we are interested in the special case where P is the frame bundle of M so that there is a canonical lift of diffeomorphisms of M to bundle automorphisms of $P = LM$. Thus, this setting falls in the realm of the last point in Theorem 4.9.

4.2.2 Kähler geometry

Let (M, σ) be a symplectic $2n$ -dimensional manifold. We are interested in the space \mathcal{I} of almost complex structures compatible with σ , i.e., almost complex structures J for which $\sigma(\cdot, J\cdot)$ is a Riemannian metric on M . Each compatible almost complex structure on M gives rise to an almost Kähler manifold structure on M and thus reduces the symplectic frame bundle $LM \rightarrow M$ to an $\text{U}(n)$ -bundle. Hence \mathcal{I} is identified with the space of sections of $LM \times_{\text{Sp}(2n, \mathbb{R})} \underline{F}$ with fiber model $\underline{F} = \text{Sp}(2n, \mathbb{R})/\text{U}(n)$. This homogeneous space \underline{F} can be identified with the Siegel upper half space and thus is a symplectic manifold.

To get some additional insight in the problem, we will follow [Ohs15; SV19] and realize the Siegel upper half space as a symplectic quotient. Consider the vector space $\mathfrak{M}(2n \times 2n, \mathbb{R})$ of real $2n \times 2n$ matrices endowed with the constant symplectic form

$$\omega(X, Y) = \text{Tr}(X^T \mathbb{J} Y), \quad (4.48)$$

where \mathbb{J} denotes the standard symplectic matrix (whose right upper $n \times n$ -corner is the identity). Left multiplication by $\text{Sp}(2n, \mathbb{R})$ leaves the symplectic form invariant. Similarly, $\text{O}(2n)$ acts by right multiplication by the inverse (so it is also a left action), and this action is symplectic as well. These actions possess the momentum maps

$$\begin{array}{ccc} \mathfrak{sp}(2n, \mathbb{R}) & \xleftarrow{J_{\text{Sp}}} & \mathfrak{M}(2n \times 2n, \mathbb{R}) & \xrightarrow{J_{\text{O}}} & \mathfrak{o}(2n) \\ -XX^T \mathbb{J} & \longleftarrow & X & \longrightarrow & X^T \mathbb{J} X, \end{array} \quad (4.49)$$

where we identified the duals with the Lie algebras itself using the bilinear form $\frac{1}{2} \text{Tr}(XY^T)$. Note that $U(n)$ is the stabilizer of \mathbb{J} under the (co)adjoint action of either $\text{Sp}(2n, \mathbb{R})$ or $O(2n)$. Hence, we identify the symplectic quotients at \mathbb{J} with the spaces

$$J_{\text{Sp}}^{-1}(\mathbb{J})/\text{Sp}(2n, \mathbb{R})_{\mathbb{J}} = O(2n)/U(n), \quad J_{O}^{-1}(\mathbb{J})/O(2n)_{\mathbb{J}} = \text{Sp}(2n, \mathbb{R})/U(n). \quad (4.50)$$

It is intriguing that this duality on the level of momentum maps and symplectic quotients corresponds to the two ways to get an almost Kähler manifold: start either from a symplectic manifold or a Riemannian manifold, and then choose a compatible almost complex structure.

Using the momentum maps, we can determine the prequantizations. For this, recall from Subsection 4.2.1 that equivariant prequantization bundles over the homogeneous space G/H are in bijective correspondence with Lie group homomorphisms $\rho: H \rightarrow U(1)$ integrating the Lie algebra homomorphism $J(eH)_{\mathfrak{h}}: \mathfrak{h} \rightarrow \mathbb{R}$. In the present setting, we have $J_{\text{Sp}}(e U(n)) = -\mathbb{J} = -J_O(e U(n))$. The restriction of the corresponding functionals to $\mathfrak{u}(n)$ yields the Lie algebra homomorphism $\mp \frac{1}{2} \text{Tr}(\mathbb{J} \cdot): \mathfrak{u}(n) \rightarrow \mathbb{R}$, which equals $\pm \text{Tr}_{\mathbb{C}}: \mathfrak{u}(n) \rightarrow i\mathbb{R}$ when $\mathfrak{u}(n)$ is viewed as a subalgebra of complex matrices. Thus the integrating group homomorphism is given by

$$\rho_{\pm}: U(n) \rightarrow U(1), \quad A \mapsto (\det_{\mathbb{C}} A)^{\pm 1}. \quad (4.51)$$

In our sign convention, ρ_+ corresponds to J_O while ρ_- integrates J_{Sp} .

Let us now return to the non-linear setting given by a symplectic manifold (M, σ) with associated symplectic frame bundle $LM \rightarrow M$. For every almost complex structure I on M , let $L_I M$ be the corresponding $U(n)$ -reduction of LM and let Γ_I be the Levi-Civita connection, viewed as a connection on $L_I M$. With a slight abuse of notation, we also denote the extended connection on LM by Γ_I . The set \mathcal{I} of almost complex structures compatible with σ is identified with the space of sections of $F = LM \times_{\text{Sp}(2n, \mathbb{R})} \underline{F}$ with $\underline{F} = \text{Sp}(2n, \mathbb{R})/U(n)$. As discussed above, the fiber model \underline{F} is a symplectic manifold and hence our general theory implies that \mathcal{I} is a symplectic manifold and that the group of symplectomorphisms acts symplectically on it. According to Theorem 4.9, the momentum map for the action of the group of symplectomorphisms involves the associated bundle $(L_I M \times_{\rho^{-1}} U(1), (\rho^{-1})_* \Gamma_I)$. The group homomorphism $\rho^{-1}: U(n) \rightarrow U(1)$ is the determinant and thus the associated bundle $L_I M \times_{\rho^{-1}} U(1)$ coincides with the canonical circle bundle¹ $K_I M$ endowed with its natural Chern connection determined by the almost complex structure I . Moreover, note that the adjoint

¹ Given an almost complex structure I on M , the complex line bundle $\Lambda^{n,0} M$ of holomorphic forms is called the *canonical line bundle* and its dual $\Lambda^{0,n} M$ the *anti-canonical line bundle*. The volume form $\frac{\sigma^n}{n!}$ is a non-vanishing section of $\Lambda^{n,n} M = \Lambda^{n,0} M \otimes \Lambda^{0,n} M$ and thus can be viewed as a

of $\tau_{\Gamma_I, \chi}: \mathfrak{X}_\sigma(M) \rightarrow \mathfrak{gau}(LM)$ appears in the construction of the momentum map. As shown in Remark 3.4, $\tau_{\Gamma_I, \chi}^*(s)$ involves the covariant derivative of $s = J_*(I)$ and thus this term vanishes, because I is parallel with respect to Γ_I . To summarize, we obtain an important special case of Theorem 4.9 (in its symplectic version analogous to Theorem 3.7).

THEOREM 4.10 *Let (M, σ) be a compact symplectic manifold of dimension $2n$. Then the set \mathcal{I} of almost complex structures compatible with σ is naturally endowed with the structure of a symplectic Fréchet manifold. If (M, σ) has a prequantum bundle $h_\sigma \in \hat{H}^2(M, \mathbb{U}(1))$, then the symplectic action of $\text{Diff}_\sigma(M)$ on \mathcal{I} has a group-valued momentum map*

$$\mathcal{J}_{\text{Diff}}: \mathcal{I} \rightarrow \hat{H}^{2n}(M, \mathbb{U}(1)), \quad I \mapsto K_I M \star h_\sigma^{n-1}, \quad (4.52)$$

where $K_I M$ is the canonical bundle, viewed as an element of $\hat{H}^2(M, \mathbb{U}(1))$. \diamond

By definition, the Chern–Ricci form Ric_I is the curvature of the Chern connection on the anti-canonical bundle. If I is integrable, then Ric_I coincides with the Ricci form of the associated Kähler metric. We hence have

$$\text{curv} \circ \mathcal{J}_{\text{Diff}}: \mathcal{I} \rightarrow \Omega^{2n}(M, \mathbb{U}(1)), \quad I \mapsto -\text{Ric}_I \wedge \sigma^{n-1} = -\frac{S_I}{n} \sigma^n, \quad (4.53)$$

where $S_I = \text{Tr}_\sigma \text{Ric}_I$ is the Chern scalar curvature. It has been shown by Donaldson [Don97, Proposition 9] that, under rather restrictive topological assumptions, the map $I \mapsto S_I$ yields a momentum map for the $\text{Diff}_\sigma(M)$ -action on \mathcal{I} . As in the examples above, the curvature momentum map is, in general, not the momentum map for the full symplectomorphism group but only for the subgroup of *Hamiltonian symplectomorphisms*. Moreover, the group-valued momentum map contains additional topological information: the Chern class of the bundle $K_I^{-1} M \star h_\sigma^{n-1}$, which is given as the cup-product $c_I(M) \cup c(h_\sigma)^{n-1}$ of the Chern class $c(h_\sigma)$ with the first Chern class $c_I(M) := c(K_I M) \in H^2(M, \mathbb{Z})$ of the almost complex manifold (M, I) . In real cohomology, this class simplifies to $c_I(M) \cup [\sigma]^{n-1}$.

Equation (4.53) shows that the curvature of the momentum map is essentially the scalar curvature. Hence, taking different symplectic quotients yield various moduli spaces of almost Kähler metrics with prescribed scalar curvature. The following cases are of special interest.

- (i) For $c \in \hat{H}^{2n}(M, \mathbb{U}(1))$, the level set $\mathcal{J}_{\text{Diff}}^{-1}(c)$ consists of all almost complex structures I whose canonical bundle satisfies $K_I M \star h_\sigma^{n-1} = c$.

Hermitian metric on the (anti-)canonical bundle. The associated Hermitian frame bundles $K_I M$ and $K_I^{-1} M$ are principal $\mathbb{U}(1)$ -bundles and are called the *canonical*, respectively, *anti-canonical circle bundles*.

- (ii) Recall from the exact sequence (D.6) that the set $c + H^{2n-1}(M, U(1))$ comprises all differential characters with the same curvature as c . Thus the inverse image $\mathcal{J}_{\text{Diff}}^{-1}(c + H^{2n-1}(M, U(1)))$ is the set of all almost complex structures I with prescribed scalar curvature $S_I = -n \frac{\text{curv } c}{\sigma^n}$.
- (iii) Similarly, the set $c + \text{Jac}^{n-1}(M)$ parametrizes differential characters having the same curvature and topological class as c . Thus the inverse image $\mathcal{J}_{\text{Diff}}^{-1}(c + \text{Jac}^{n-1}(M))$ is identified with the set of all almost complex structures whose scalar curvature is prescribed by the above relation and, in addition, for which $c_I(M) \cup c(h_\sigma)^{n-1}$ coincides with the Chern class of c .

Ignoring the particularities of the infinite-dimensional setting, one would thus conjecture that all the corresponding moduli spaces are symplectic manifolds or at least symplectic stratified spaces.

REMARK 4.11 (Kähler–Einstein metrics) If the curvature of c is a constant multiple of σ^n , then the inverse image $\mathcal{J}_{\text{Diff}}^{-1}(c + \text{Jac}^{n-1}(M))$ is equal to the set of Kähler–Einstein metrics. Indeed, an (almost) complex structure I whose Ricci form satisfies the Einstein relation $\text{Ric}_I = \lambda \sigma$ has constant scalar curvature $S_I = n\lambda$. Conversely, every *integrable* complex structure I with constant scalar curvature and whose Chern class satisfies $c_I(M) = \lambda[\sigma]$ in real cohomology is automatically Kähler–Einstein. Indeed, in this case, Ric_I and $\lambda\sigma$ are two harmonic forms representing the same cohomology class and thus are equal. The issue of existence of Kähler–Einstein metrics has been an important subject of study for decades and was recently settled with the resolution of the Yau–Tian–Donaldson conjecture (we refer the reader to the recent review [Don15] and references therein for more details). The infinite-dimensional symplectic picture we have just developed suggests that Kähler metrics with constant scalar curvature are actually a more natural object to study, especially since their moduli spaces are symplectic quotients. However, the existence problem is harder than for Kähler–Einstein metrics, and thus it is not clear how the analytic question concerning the infinite-dimensional symplectic reduction pan out. \diamond

REMARK 4.12 (Relation to GIT quotient) It is a recurrent theme in Kähler geometry that the existence of certain special metrics (Kähler–Einstein, positive Ricci curvature, . . .) has an algebraic stability obstruction. Donaldson [Don97; Don01] related this geometric-algebraic equivalence to the interplay of the symplectic momentum map geometry with geometric invariant theory. The fundamental idea is to detect zeros of the momentum map using an algebraic condition phrased in terms of the prequantum bundle. However, this approach relies on the existence of a classical momentum map and thus requires certain topological classes to vanish. In light of our refined results, it would be interesting to see how the group-valued momentum map interacts with geometric invariant theory. In this direction, [Die+19] shows how the hat product of differential

characters can be used to construct a prequantization bundle over certain infinite-dimensional function spaces. However, it is not yet clear what the existence of a group-valued momentum map means in terms of the prequantum geometry. \diamond

REMARK 4.13 The anti-canonical bundle often plays the role of the algebraic counterpart to a geometric property. For example, a compact complex manifold admits a Kähler metric of positive Ricci curvature only if it has an ample anti-canonical bundle (by Yau’s proof of the Calabi conjecture combined with Kodaira’s theorem). Our general group-valued momentum map picture explains, at least in principle, the importance of the (anti-)canonical bundle in many important questions concerning the geometry of (almost) Kähler manifolds: $K_I M$ is the “conserved quantity” of the group of symplectomorphisms. \diamond

Recall that we realized the Siegel upper half space $\mathrm{Sp}(2n, \mathbb{R})/\mathrm{U}(n)$ as a symplectic quotient. This additional structure transfers to the space of sections, and puts the moduli space of constant scalar curvature metrics in the framework of reduction by stages. For the moment, let us return to the following general set-up. Let $(\underline{F}, \underline{\omega})$ be a finite-dimensional symplectic manifold. Consider two Lie groups G and K that act on \underline{F} from the left and preserve the symplectic form. In the Kähler setting, we have $\underline{F} = \mathrm{GL}(2n, \mathbb{R})$, $G = \mathrm{O}(2n)$, and $K = \mathrm{Sp}(2n, \mathbb{R})$. Assume that the two actions are free, proper, commute, and have momentum maps

$$\mathfrak{g}^* \xleftarrow{J_G} \underline{F} \xrightarrow{J_K} \mathfrak{k}^*. \tag{4.54}$$

Moreover, we assume that $J_K : \underline{F} \rightarrow \mathfrak{k}^*$ is invariant under G and that, vice-versa, J_G is invariant under K . Hence we are in the setting of commuting reduction by stages [Mar+07, Section 15.5]. Let $\mu \in \mathfrak{g}^*$. The general theory [Mar+07, Lemma 4.1.2] tells us that the symplectic reduction by G yields the first stage reduced Hamiltonian K -space $(\underline{F}_\mu, \underline{\omega}_\mu, J_\mu)$. The momentum map of the reduced space is induced by $J_K : \underline{F} \rightarrow \mathfrak{k}^*$ in the obvious way. Reducing once more by the remaining K -action gives a symplectic quotient that is symplectically diffeomorphic to the corresponding symplectic quotient of the product group $G \times K$.

Let us now discuss the corresponding story in the bundle picture and for the spaces of sections. For this, let $\pi : P \rightarrow M$ be a principal K -bundle. The spaces occurring in the symplectic reduction diagram

$$\begin{array}{ccc} J_G^{-1}(\mu) & \hookrightarrow & \underline{F} \\ \downarrow / G & & \\ \underline{F}_\mu & & \end{array} \tag{4.55}$$

all carry a natural K -action such that the inclusion and projection maps are equivariant. Thus we obtain a similar diagram for the section spaces of the associated bundles

$$\begin{array}{ccc} \Gamma^\infty(P \times_K J_G^{-1}(\mu)) & \hookrightarrow & \mathcal{F} = \Gamma^\infty(P \times_K \underline{E}) \\ \downarrow / C^\infty(M, G) & & \\ \Gamma^\infty(P \times_K \underline{E}_\mu) & & \end{array} \quad (4.56)$$

The reduced space $\Gamma^\infty(P \times_K \underline{E}_\mu)$ is symplectic by our general theory, since it is the section space of a symplectic fiber bundle. Hence it is natural to suppose that $\Gamma^\infty(P \times_K \underline{E}_\mu)$ is the symplectic reduction starting from \mathcal{F} . This is indeed the case as we will see now. Consider the action of $C^\infty(M, G)$ on $\mathcal{F} = \Gamma^\infty(P \times_K \underline{E})$ induced by the G -action on \underline{E} . This action is well-defined since the G - and K -actions on \underline{E} commute. Although this setting is not completely the same as the action of a gauge group on the space of sections of an associated bundle — $C^\infty(M, G)$ is the gauge group of the trivial bundle not of P —, the general strategy of Proposition 3.2 carries through and yields the momentum map

$$\mathcal{J}_G: \mathcal{F} \rightarrow C^\infty(M, \mathfrak{g}^*), \quad \phi \mapsto J_G \circ \phi, \quad (4.57)$$

where $J_G: P \times_K \underline{E} \ni [p, \underline{f}] \mapsto (\pi(p), J_G(\underline{f})) \in M \times \mathfrak{g}^*$ is the bundle map induced by the K -invariant map $J_G: \underline{E} \rightarrow \mathfrak{g}^*$. From this expression, it is easy to see that $\Gamma^\infty(P \times_K \underline{E}_\mu)$ is the symplectic reduced space of \mathcal{F} by $C^\infty(M, G)$ at the constant map $\mu \in C^\infty(M, \mathfrak{g}^*)$.

In the Kähler setting, we have $\underline{E} = \text{GL}(2n, \mathbb{R})$, $G = \text{O}(2n)$, and $K = \text{Sp}(2n, \mathbb{R})$ so that the G -reduced space \underline{E}_μ is the upper Siegel half space for $\mu = \mathbb{J}$. The space \mathcal{F} of sections is identified with the space $\mathcal{T}^2(M)$ of invertible 2-tensors, and the first reduced stage $\Gamma^\infty(P \times_K \underline{E}_\mu)$ coincides with the space \mathcal{I} of almost complex structures compatible with the symplectic form σ . As discussed above, the second reduction by $\text{Diff}_\sigma(M)$ at the subset $c + \hat{H}^{2n-1}(M, \text{U}(1))$ yields the moduli space \mathcal{SC} of almost complex structures I with prescribed scalar curvature. Thus we get the following reduction by stages diagram:

$$\begin{array}{ccc} \mathcal{T}^2(M) & \xrightarrow{\parallel_{\mathbb{J}} C^\infty(M, \text{O}(2n))} & \mathcal{I} & \xrightarrow{\parallel_c \text{Diff}_\sigma(M)} & \mathcal{SC}. \\ & & \text{?} & & \\ & \searrow & & \nearrow & \\ & & \parallel_{\mathbb{J}, c} C^\infty(M, \text{O}(2n)) \times \text{Diff}_\sigma(M) & & \end{array} \quad (4.58)$$

We believe that this diagram commutes, i.e., the two-stage reduced space \mathcal{SC} can be obtained by the one-shot reduction from $\mathcal{T}^2(M)$ by the product group

$C^\infty(M, \mathcal{O}(2n)) \times \text{Diff}_\sigma(M)$. In the finite-dimensional setting, such questions about commuting reductions are well-understood [Mar+07]. However, the framework of reduction by stages is completely missing in infinite dimensions. Once the general theory of reduction by stages in infinite dimensions is formulated, one can state that the curved arrow is a one-shot reduction. This general theory is the subject of a future paper.

4.2.3 Teichmüller space and weighted Lagrangian subbundles

Coadjoint orbits are the canonical examples of symplectic homogeneous spaces. Let G be a finite-dimensional Lie group and $\mu \in \mathfrak{g}^*$. The coadjoint orbit $\mathcal{O}_\mu \simeq G/G_\mu$ through μ is endowed with the Kostant–Kirillov–Souriau symplectic form

$$\omega_v(\text{ad}_A^* v, \text{ad}_B^* v) = \langle v, [A, B] \rangle, \quad v \in \mathcal{O}_\mu \subset \mathfrak{g}^*, \quad A, B \in \mathfrak{g}. \quad (4.59)$$

Let M be a finite-dimensional closed manifold carrying a G -structure, i.e., the structure group of the frame bundle is reduced to G . A section of $LM \times_G (G/G_\mu)$ corresponds to a further reduction to G_μ . Thus, our general theory implies that the space of all such G_μ -reductions carries a symplectic structure, which is naturally induced by the coadjoint orbit symplectic form (4.59). The momentum map for the action of the group of diffeomorphisms of M on the space of all such G_μ -structures can be calculated with the help of Theorem 4.9. Recall that the momentum map involved the prequantization of the fiber model. In the present setting, prequantizations of G/G_μ bijectively correspond to Lie group homomorphisms $\rho_\mu: G_\mu \rightarrow \text{U}(1)$ that integrate the Lie algebra homomorphism (see, Subsection 4.2.1)

$$\check{\rho}_\mu = \underline{J}(eG_\mu)|_{\mathfrak{g}_\mu}: \mathfrak{g}_\mu \rightarrow \mathbb{R}, \quad (4.60)$$

where $\underline{J}: G/G_\mu \rightarrow \mathfrak{g}^*$ is the momentum map for the G -action on the coadjoint orbit. It is a straightforward calculation to verify that the momentum map is just the canonical embedding of the coadjoint orbit in \mathfrak{g}^* . Thus we are looking for group homomorphisms ρ_μ that integrate the pairing with μ , that is,

$$\check{\rho}_\mu: \mathfrak{g}_\mu \rightarrow \mathbb{R}, \quad A \mapsto \langle \mu, A \rangle. \quad (4.61)$$

In summary, starting from a G -structure we can derive different geometric moduli spaces that arise from an infinite-dimensional symplectic reduction by performing the following steps:

- (i) Classify all coadjoint orbits \mathcal{O}_μ of G .
- (ii) For each $\mu \in \mathfrak{g}^*$ in this classification, determine the stabilizer G_μ and find an identification of G/G_μ in terms of a more geometric object. In practice,




Figure 1: Coadjoint orbits of $SL(2, \mathbb{R})$ projected onto the $e_- - h$ plane.

this comes down to finding a transitive G -action on some space having stabilizer G_μ .

- (iii) Calculate $\check{\rho}_\mu: \mathfrak{g}_\mu \rightarrow \mathbb{R}$ and integrate it to a Lie group homomorphism $\rho_\mu: G_\mu \rightarrow U(1)$. This integration is usually possible only for certain values of λ . These are the so-called prequantizable coadjoint orbits.
- (iv) Give a geometric interpretation of the associated bundle $L_\mu M \times_{\rho_\mu} U(1)$, where $L_\mu M$ is a given G_μ -reduction of the G -frame bundle LM .

In this subsection, we carry out this program for the coadjoint orbits of $SL(2, \mathbb{R})$. The low dimensionality of the spaces involved allows to explicitly determine the coadjoint orbits and the stabilizer subgroups G_μ . Even in this relatively simple example, the geometric structures involved turn out to be of special interest: we recover the Teichmüller moduli space as a symplectic reduction; another coadjoint orbit yields a hyperbolic cousin of the classical Teichmüller space and yet another orbit is related to Lagrangian distributions endowed with a density. Throughout this subsection, let (M, σ) be closed surface endowed with a symplectic form σ (which can, equivalently, be viewed as a volume form).

We begin with a systematic discussion of the coadjoint orbits of $SL(2, \mathbb{R})$. The Lie algebra $\mathfrak{sl}(2, \mathbb{R})$ consists of real 2×2 matrices with vanishing trace. We choose the following generators

$$e_+ = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad e_- = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad h = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (4.62)$$

subject to the commutation relations $[h, e_\pm] = \mp 2e_\mp$ and $[e_+, e_-] = 2h$. The Killing form $\kappa(A, B) = \frac{1}{2} \text{Tr}(AB)$ is non-degenerate and thus yields an identification of \mathfrak{g} with \mathfrak{g}^* . The generators form an orthonormal basis of \mathfrak{g} (and thus also of \mathfrak{g}^*) with signature $(++-)$. The Killing form is Ad-invariant and so identifies the coadjoint action with the adjoint action. Every non-vanishing element $A \in \mathfrak{sl}(2, \mathbb{R})$ can be

brought into one of the following four normal forms under the adjoint action:

$$N_e^\lambda = \lambda \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad N_h^\lambda = \lambda \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad N_p^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad N_p^- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \quad (4.63)$$

The orbit through N_e^λ is an elliptic hyperboloid (or two-sheeted hyperboloid). Positive λ parametrize the family of the upper sheets and negative λ correspond to the lower sheets. The stabilizer of N_e^λ is independent of λ and is isomorphic to $\mathrm{SO}(2)$. The coadjoint orbit for $\lambda = 1$ is symplectomorphic to the reduced space $J_{\mathbb{O}(2)}^{-1}(\mathbb{J})/\mathbb{O}(2)_{\mathbb{J}} = \mathrm{Sp}(2, \mathbb{R})/\mathrm{U}(1)$ discussed in the previous section (see (4.50) with $n = 1$). Sections of $\mathrm{LM} \times_{\mathrm{SL}(2, \mathbb{R})} (\mathrm{SL}(2, \mathbb{R})/\mathrm{SO}(2))$ correspond to reductions of the $\mathrm{SL}(2, \mathbb{R})$ -frame bundle to $\mathrm{SO}(2)$, i.e., Riemannian metrics g compatible with the prescribed volume form σ . To make this interpretation in terms of Riemannian metrics more explicit, we identify the coadjoint orbit $\mathcal{O}_{N_e^\lambda} \simeq \mathrm{SL}(2, \mathbb{R})/\mathrm{SO}(2)$ with the space $\mathrm{SSym}^+(2, \mathbb{R})$ of symmetric positive definite matrices having determinant 1 via

$$\mathcal{O}_{N_e^\lambda} \ni \mathrm{Ad}_g N_e^\lambda \mapsto gg^T \in \mathrm{SSym}^+(2, \mathbb{R}), \quad g \in \mathrm{SL}(2, \mathbb{R}). \quad (4.64)$$

Using this diffeomorphism, the coadjoint orbit symplectic form can be pushed forward to yield the following symplectic form on $\mathrm{SSym}^+(2, \mathbb{R})$:

$$\omega_B(C_1, C_2) = -\frac{1}{4} \mathrm{Tr}(B^{-1}C_1 B^{-1}N_e^\lambda B^{-1}C_2), \quad (4.65)$$

where $B \in \mathrm{SSym}^+(2, \mathbb{R})$, and C_1, C_2 are traceless symmetric matrices.

The orbit through N_h^λ is a hyperbolic hyperboloid (or one-sheeted hyperboloid). The stabilizer of N_h^λ is again independent of λ , consists of elements of the form

$$S = \begin{pmatrix} \pm \cosh r & \sinh r \\ \sinh r & \pm \cosh r \end{pmatrix}, \quad r \in \mathbb{R}, \quad (4.66)$$

and hence is isomorphic to $\mathrm{SO}(1, 1)$. Sections of $\mathrm{LM} \times_{\mathrm{SL}(2, \mathbb{R})} (\mathrm{SL}(2, \mathbb{R})/\mathrm{SO}(1, 1))$ are again reductions of the $\mathrm{SL}(2, \mathbb{R})$ -frame bundle, this time, to $\mathrm{SO}(1, 1)$. Geometrically, these are semi- (or pseudo-, in another terminology) Riemannian metrics η . Hence the space of semi-Riemannian metrics inducing the given volume form σ is an infinite-dimensional symplectic manifold.

The parabolic orbits, that is, the two light-cones, are orbits through N_p^+ and N_p^- . As the notation suggests, N_p^+ corresponds to the upper cone. Its stabilizer P^+ consists of elements of the form

$$S = \begin{pmatrix} \pm 1 & r \\ 0 & \pm 1 \end{pmatrix} \quad (4.67)$$

for $r \in \mathbb{R}$. Similarly, the lower cone is the orbit through N_p^- and the elements of

its stabilizer P_- are of the form

$$S = \begin{pmatrix} \pm 1 & 0 \\ r & \pm 1 \end{pmatrix} \quad (4.68)$$

for $r \in \mathbb{R}$.

In order to illuminate the geometric structure that leads to a reduction to the stabilizer subgroups P^\pm , it is actually helpful to consider the higher dimensional case. Thus, let (M, σ) be, for the moment, a $2n$ -dimensional symplectic manifold. An n -dimensional distribution D on M yields a reduction of the symplectic frame bundle to the group consisting of elements of the form

$$\begin{pmatrix} A & B \\ 0 & C \end{pmatrix}, \quad (4.69)$$

where $A, C \in \text{GL}(n, \mathbb{R})$ and B is an arbitrary $n \times n$ matrix. Indeed, the reduced bundle consists only of frames $u_m: \mathbb{R}^{2n} \rightarrow T_m M$ that map the last n standard vectors to vectors spanning D_m . If we would, instead, require that the first n vectors span D_m , we obtain a reduction to elements of the form

$$\begin{pmatrix} A & 0 \\ B & C \end{pmatrix}. \quad (4.70)$$

If the distribution carries an additional structure, then this is reflected in further conditions on the matrix C . For example, a distribution endowed with a density $|\nu|$ leads to the condition $|\det C| = 1$. We then say that the distribution is weighted.

Specializing again to the case $n = 1$ at hand, we see that a reduction of the symplectic frame bundle to P^\pm corresponds to a weighted Lagrangian subbundle $(L, |\nu|)$ of TM . Thus the space of all weighted Lagrangian distributions carries two natural symplectic structures (depending on the point N_p^\pm chosen for the identification).

In order to determine the prequantizations of the orbits, we have to calculate

$$\check{\rho}_\mu: \mathfrak{g}_\mu \rightarrow \mathbb{R}, \quad A \mapsto \langle \mu, A \rangle \quad (4.71)$$

for each orbit. Note that we always have $\mathfrak{g}_\mu = \mathbb{R}\mu$ since the stabilizers are 1-dimensional. We obtain

$$\check{\rho}_{N_e^\lambda}(a \cdot N_e^\lambda) = -a\lambda, \quad (4.72)$$

$$\check{\rho}_{N_h^\lambda}(a \cdot N_h^\lambda) = a\lambda, \quad (4.73)$$

$$\check{\rho}_{N_p^\pm}(a \cdot N_p^\pm) = 0. \quad (4.74)$$

If λ satisfies the prequantization condition $\lambda \in \mathbb{Z}$, then $\check{\rho}_{N_e^\lambda}$ integrates to the Lie

μ	N_e^λ	N_h^λ	N_p^\pm
Type	elliptic	hyperbolic	parabolic
G_μ	SO(2)	SO(1, 1)	P^\pm
Quantizable	$\lambda \in \mathbb{Z}$	always	always
ρ_μ	$\rho_{N_e^\lambda}$	$\rho_{N_h^\lambda}, \sqrt{\rho_{N_h^{2\lambda}}}$	$1, \rho_{N_p^\pm}$
\mathcal{F}_μ	Riemannian metric g	semi-Riemannian metric η	weighted Lagrangian distribution (L, v)
P_μ	$K_g^{-1}M$	$K_\eta M, K_\eta^{\frac{1}{2}}M$	OrL
Characteristic class	$c_1(M)$?	$w_1(L)$
Moduli space	Teichmüller space	hyperbolic Teichmüller space	?

Table 3: Overview over the properties of the coadjoint orbits. The information concerning the geometric structures induced by the coadjoint orbits will become clear after the statement of Theorem 4.14 and the comments following it. Here, \mathcal{F}_μ denotes the space of sections of $LM \times_{\text{SL}(2, \mathbb{R})} (\text{SL}(2, \mathbb{R})/G_\mu)$. Furthermore, P_μ is the associated circle bundle of the reduced G_μ -frame bundle using the group homomorphism ρ_μ .

group homomorphism

$$\rho_{N_e^\lambda}: \text{SO}(2) \rightarrow \text{U}(1), \quad \begin{pmatrix} \cos \vartheta & \sin \vartheta \\ -\sin \vartheta & \cos \vartheta \end{pmatrix} \mapsto e^{-\vartheta \lambda i}, \quad (4.75)$$

Note that for $\lambda = 1$, this is just the inverse of the canonical identification of $\text{SO}(2)$ with $\text{U}(1)$. Thus, the associated bundle $L_{\text{SO}(2)}M \times_{\rho_{N_e^1}} \text{U}(1)$ is the anti-canonical bundle $K_g^{-1}M$ constructed from the Riemannian metric g . Other values of $\lambda \in \mathbb{Z}$ correspond to higher powers $K_g^{-\lambda}M$.

In contrast, $\text{SO}(1, 1)$ is not connected and $\check{\rho}_{N_h^\lambda}$ admits two types of integrating group homomorphisms for all $\lambda \in \mathbb{R}$

$$\rho_{N_h^\lambda}: \text{SO}(1, 1) \rightarrow \text{U}(1), \quad \begin{pmatrix} \pm \cosh r & \sinh r \\ \sinh r & \pm \cosh r \end{pmatrix} \mapsto e^{\pm r \lambda i}, \quad (4.76)$$

$$\sqrt{\rho_{N_h^{2\lambda}}}: \text{SO}(1, 1) \rightarrow \text{U}(1), \quad \begin{pmatrix} \pm \cosh r & \sinh r \\ \sinh r & \pm \cosh r \end{pmatrix} \mapsto \begin{cases} e^{r \lambda i} & \text{for } + \\ -e^{-r \lambda i} & \text{for } - \end{cases} \quad (4.77)$$

As the notation suggests, we have $\sqrt{\rho_{N_h^{2\lambda}}} = \rho_{N_h^\lambda}$. But, of course, also $\rho_{N_h^\lambda}^2 = \rho_{N_h^{2\lambda}}$ holds, so that $\sqrt{\rho_{N_h^{2\lambda}}}$ is the “non-canonical square root”. In analogy to the previous case, the associated bundle $L_{\text{SO}(1,1)}M \times_{\rho_{N_h^2}} \text{U}(1)$ is the canonical bundle $K_\eta M$ induced by the semi-Riemannian metric η . The bundles $\rho_{N_h^\lambda}$ and $\sqrt{\rho_{N_h^{2\lambda}}}$

yield the two inequivalent square roots $K_\eta^{+\frac{1}{2}}M$ and $K_\eta^{-\frac{1}{2}}M$ of $K_\eta M$. Thus, in contrast to the Riemannian case, also the square roots of the canonical bundle are of particular importance based on the coadjoint orbit geometry. Analogously, for $\lambda \in \mathbb{R}$, we use the notations $K_\eta^{+\frac{\lambda}{2}}M$ and $K_\eta^{-\frac{\lambda}{2}}M$ for the associated bundles $L_{\text{SO}(1,1)}M \times_{\rho_{N_h^\lambda}} U(1)$ and $L_{\text{SO}(1,1)}M \times_{\sqrt{\rho_{N_h^{2\lambda}}}} U(1)$, respectively.

Since $\check{\rho}_{N_p^\pm} = 0$, the integrating group homomorphism $\rho_{N_p^\pm}: P^\pm \rightarrow U(1)$ may be chosen to be the trivial homomorphism. However, P^\pm is also not connected and thus there is a second non-trivial integrating homomorphisms $\rho_{N_p^\pm}$, which sends $\begin{pmatrix} -1 & r \\ 0 & -1 \end{pmatrix}$, respectively $\begin{pmatrix} -1 & 0 \\ r & -1 \end{pmatrix}$, to $-1 \in U(1)$. Note that the latter homomorphism factors through a homomorphism with values in \mathbb{Z}_2 .

In order to illuminate the significance of the associated \mathbb{Z}_2 -bundle $L_{P^\pm}M \times_{\rho_{N_p^\pm}} \mathbb{Z}_2$, it is again helpful to leave the safe 2-dimensional harbor and move to higher dimensional terrain. Thus, let (M, σ) be, temporarily, a $2n$ -dimensional symplectic manifold. Recall that a weighted Lagrangian distribution on M yields a reduction of the symplectic frame bundle to the structure group consisting of symplectic block matrices $\begin{pmatrix} A & B \\ 0 & C \end{pmatrix}$ with $C \in \text{SL}^\pm(n, \mathbb{R}) := \{C \in \text{SL}(n, \mathbb{R}) \mid \det C = \pm 1\}$. The higher dimensional analogue of $\rho_{N_p^\pm}$ is the group homomorphism

$$\rho_{N_p^\pm}: \begin{pmatrix} A & B \\ 0 & C \end{pmatrix} \mapsto \det C \in \mathbb{Z}_2. \quad (4.78)$$

Thus a global section of the associated flat bundle $LM \times_{\rho_{N_p^\pm}} \mathbb{Z}_2$ corresponds to a consistent choice of orientation of the Lagrangian subspaces. We hence refer to this associated \mathbb{Z}_2 -bundle as the *orientation bundle* $\text{Or}L$ of the Lagrangian distribution L . The orientation bundle is trivial if and only if the first Stiefel–Whitney class $w_1(L)$ of the Lagrangian distribution vanishes.

We have thus assembled all ingredients to apply Theorem 4.9 (see also Table 3 for a summary).

THEOREM 4.14 *Let M be a closed surface endowed with a symplectic form σ . Then the following hold:*

- (i) *The space $\text{Metr}_\sigma(M)$ of Riemannian metrics on M compatible with σ is endowed with the symplectic form*

$$\Omega_g(h_1, h_2) = -\frac{\lambda}{4} \int_M \text{Tr}(g^{-1}h_1 \cdot g^{-1}\sigma \cdot g^{-1}h_2) \sigma. \quad (4.79)$$

If $\lambda \in \mathbb{Z}$, then the action of $\text{Diff}_\sigma(M)$ has a group-valued momentum map given by

$$\text{Metr}_\sigma(M) \rightarrow \hat{\text{H}}^2(M, U(1)), \quad g \mapsto K_g M. \quad (4.80)$$

(ii) The space $\text{Metr}_\sigma^{+-}(M)$ of semi-Riemannian metrics with signature $(+-)$ and compatible with σ carries a symplectic form defined as in the Riemannian case by (4.79). For all $\lambda \in \mathbb{R}$, the action of $\text{Diff}_\sigma(M)$ has two group-valued momentum maps given by

$$\text{Metr}_\sigma^{+-}(M) \rightarrow \hat{H}^2(M, \mathbb{U}(1)), \quad \eta \mapsto \mathbb{K}_\eta^{+\frac{\lambda}{2}} M \quad (4.81)$$

and

$$\text{Metr}_\sigma^{+-}(M) \rightarrow \hat{H}^2(M, \mathbb{U}(1)), \quad \eta \mapsto \mathbb{K}_\eta^{-\frac{\lambda}{2}} M. \quad (4.82)$$

(iii) The space of weighted Lagrangian distributions is a symplectic manifold and the group-valued momentum map for the action of $\text{Diff}_\sigma(M)$ is the assignment of the orientation bundle $\text{Or}L$ to each Lagrangian distribution L . \diamond

The curvature of the canonical bundle $\mathbb{K}_g M$ is given by $-S_I \sigma$, where S_I denotes the scalar curvature. Hence, symplectic reduction at the subset $\text{curv}^{-1}(\sigma)$ of all bundles with constant curvature σ yields the Riemann moduli space (for $\lambda = 1$):

$$\mathcal{J}_{\text{Diff}}^{-1}(\text{curv}^{-1}(\sigma))/\text{Diff}_\sigma(M) = \{I \in \mathcal{I} \mid S_I = -1\}/\text{Diff}_\sigma(M). \quad (4.83)$$

Instead of taking the quotient with respect to $\text{Diff}_\sigma(M)$, we could restrict attention to the connected component of the identity $\text{Diff}_\sigma(M)^\circ$. The action of $\text{Diff}_\sigma(M)^\circ$ is free and its momentum map is given by the same formula (4.80). Thus, the reduced space coincides with $\{I \in \mathcal{I} \mid S_I = -1\}/\text{Diff}_\sigma(M)^\circ$ and hence yields the Teichmüller space. Formula (4.79) for the symplectic form on the space of Riemannian metrics shows that the reduced symplectic form is proportional to the Weil–Petersson symplectic form on the Teichmüller space. However, in contrast to classical symplectic reduction, we take the inverse image of a set and not just of a point. We show now that $\text{Diff}_\sigma(M)$ acts (infinitesimally) transitively on $\text{curv}^{-1}(\sigma)$, so that the reduction is a *symplectic orbit reduction* [OR03, Section 6.3]. Suppose that σ is normalized to have total volume 1. Then there exists a differential character $h_\sigma \in \hat{H}^2(M, \mathbb{U}(1))$ with curvature σ . Consider the flux homomorphism¹ relative to h_σ

$$\text{Flux}_{h_\sigma}: \text{Diff}_\sigma(M) \rightarrow \mathbb{H}^1(M, \mathbb{U}(1)), \quad \phi \mapsto (\phi^{-1})^* h_\sigma - h_\sigma. \quad (4.84)$$

By Proposition D.3, the corresponding Lie algebra homomorphism is

$$\text{Flux}_\sigma: \mathfrak{X}_\sigma(M) \rightarrow \mathbb{H}^1(M, \mathbb{R}), \quad X \mapsto [X \lrcorner \sigma], \quad (4.85)$$

¹ Flux is actually only a cocycle and not necessarily a group homomorphism on $\text{Diff}_\sigma(M)$. Only when restricted to $\text{Diff}_\sigma(M) \cap \text{Diff}(M)^\circ$ is the flux a bona-fide group homomorphism.

which is clearly surjective. Thus $\text{Diff}_\sigma(M)$ acts infinitesimally transitive on $\text{curv}^{-1}(\sigma)$. In summary, we obtain the following.

THEOREM 4.15 *The Riemann moduli space*

$$\{I \in \mathcal{I} \mid S_I = -1\} / \text{Diff}_\sigma(M) \quad (4.86)$$

and the Teichmüller space

$$\{I \in \mathcal{I} \mid S_I = -1\} / \text{Diff}_\sigma(M)^\circ \quad (4.87)$$

are symplectic orbit reduced spaces. ◇

An analogous discussion for the coadjoint orbit $\text{SL}(2, \mathbb{R}) / \text{SO}(1, 1)$ yields a hyperbolic version of the Teichmüller space. Instead of Riemannian metrics, the moduli space then consists of semi-Riemannian metrics with prescribed curvature modulo symplectomorphisms. It also carries a symplectic form analogous to the one induced by the Weil–Petersson metric, and it is a symplectic orbit reduced space. Formally, we just have to replace ‘elliptic’ by ‘hyperbolic’, ‘Riemannian’ by ‘semi-Riemannian’ and ‘complex structure’ by ‘para-complex structure’. The analysis of hyperbolic operators is, of course, more delicate than elliptic operators and thus it is not clear that the hyperbolic Teichmüller space is indeed a smooth manifold. From a physics point of view, elements of the hyperbolic Teichmüller space parametrize equivalence classes of solutions of Einstein’s equation in two dimensions. In the physics literature [Stroo] one finds arguments that such moduli spaces of semi-Riemannian metrics with prescribed curvature are in fact finite dimensional, similar to the Riemannian case. We are not aware of any rigorous mathematical proof of this statement.

4.3 Action of the quantomorphism group

Let (M, ω) be a compact $2n$ -dimensional symplectic manifold and let $\mu_\omega = \omega^n / n!$ be the Liouville volume form. Assume that ω has integral periods so that there exists a prequantum circle bundle $\pi: P \rightarrow M$ with connection Γ having curvature ω . As the fiber model we take $\underline{F} = \mathbb{C}$ with the usual symplectic structure $\omega_{\mathbb{C}}$ endowed with k -times the natural $\text{U}(1)$ -action for some $k \in \mathbb{N}$. The associated line bundle $F^k = P \times_{k \cdot \text{U}(1)} \mathbb{C}$ is the k -th tensor product of the line bundle F^1 . As in Section 3.2, the connection Γ in P and the symplectic structure $\omega_{\mathbb{C}}$ on \mathbb{C} combine to an extension $\hat{\omega}_{\mathbb{C}}^\Gamma \in \Omega^2(F^k)$ of the induced fiber symplectic structure on F^k .

Theorem 3.7 determines the momentum map of the action on $\Gamma^\infty(F^k)$ of the group $\text{Aut}_\omega(P)$ of automorphisms of P that preserve the symplectic structure on the base. In this section, we put the focus on the subgroup $\text{Aut}_\Gamma(P) \subseteq \text{Aut}_\omega(P)$ consisting of automorphisms that even preserve the connection Γ . Due to its

importance in the theory of geometric quantization, $\text{Aut}_\Gamma(P)$ is often called the *quantomorphism group* of (P, Γ) . Recall that the Lie algebra $\text{aut}_\Gamma(P)$ of $\text{Aut}_\Gamma(P)$ is identified with the Poisson algebra $C^\infty(M)$ using the Koszul prescription

$$C^\infty(M) \ni f \mapsto X_f^\Gamma + \pi^* f \partial_{\mathfrak{g}} \in \text{aut}_\Gamma(P), \quad (4.88)$$

where $\partial_{\mathfrak{g}}$ is the canonical vector field along the fiber and X_f^Γ is the horizontal lift of the Hamiltonian vector field X_f . Accordingly, it is natural to choose $\text{aut}_\Gamma(P)^* = \Omega^{2n}(M)$ as the dual of $\text{aut}_\Gamma(P)$. On the level of vector spaces, the Lie algebra $\text{aut}_\omega(P)$ of $\text{Aut}_\omega(P)$ is identified with $C^\infty(M) \times \mathfrak{X}_\omega(M)$. Hence, a natural choice for the dual group of $\text{Aut}_\omega(P)$ is the Abelian Lie group $\text{Aut}_\omega(P)^* = C^\infty(M) \times \hat{H}^{2n}(M, U(1))$, whose Lie algebra is $C^\infty(M) \times \Omega^{2n-1}(M)/d\Omega^{2n-2}(M)$. Under the above identification, the inclusion of $\text{aut}_\Gamma(P)$ into $\text{aut}_\omega(P)$ is given by

$$C^\infty(M) \ni f \mapsto (f, X_f) \in C^\infty(M) \times \mathfrak{X}_\omega(M). \quad (4.89)$$

The dual projection, with respect to the pairings (2.10) and (2.11), has the form

$$C^\infty(M) \times \Omega^{2n-1}(M)/d\Omega^{2n-2}(M) \ni (g, [\alpha]) \mapsto g\mu_\omega - \frac{1}{(n-1)!} d\alpha \in \Omega^{2n}(M) \quad (4.90)$$

and lifts to the dual group $\text{Aut}_\omega(P)^*$ as follows:

$$C^\infty(M) \times \hat{H}^{2n}(M, U(1)) \ni (g, h) \mapsto g\mu_\omega - \frac{1}{(n-1)!} \text{curv } h \in \Omega^{2n}(M). \quad (4.91)$$

Clearly, the momentum map $\mathcal{J}_{\text{Aut}_\Gamma}$ for the quantomorphism group is the composition of this “projection” with the momentum map for the automorphism group. Using Theorem 3.7, we thus obtain

$$\mathcal{J}_{\text{Aut}_\Gamma}: \Gamma^\infty(F^k) \rightarrow \Omega^{2n}(M), \quad \phi \mapsto -k\|\phi\|^2\mu_\omega + \frac{1}{(n-1)!}\phi^*\hat{\omega}_\mathbb{C}^\Gamma \wedge \omega^{n-1}. \quad (4.92)$$

Using the identity $\phi^*\hat{\omega}_\mathbb{C}^\Gamma = \omega_\mathbb{C}(\nabla^\Gamma\phi, \nabla^\Gamma\phi) + (J_*\phi) \text{curv } \Gamma$ of [Don03, Lemma 13], we can rewrite the momentum map as follows:

$$\begin{aligned} \mathcal{J}_{\text{Aut}_\Gamma}(\phi) &= -k\|\phi\|^2\mu_\omega + \frac{1}{(n-1)!} \left(\omega_\mathbb{C}(\nabla^\Gamma\phi, \nabla^\Gamma\phi) - k\|\phi\|^2\omega \right) \wedge \omega^{n-1} \\ &= -k(n+1)\|\phi\|^2\mu_\omega + \frac{i}{2}\nabla^\Gamma\phi \wedge \nabla^\Gamma\bar{\phi} \wedge \frac{\omega^{n-1}}{(n-1)!}. \end{aligned} \quad (4.93)$$

Thus, in summary, we obtain the following.

THEOREM 4.16 *Let (M, ω) be a compact $2n$ -dimensional symplectic manifold. Assume that ω has integral periods so that there exists a prequantum circle bundle $\pi: P \rightarrow M$*

with connection Γ having curvature ω . Then the space $\Gamma^\infty(F^k)$ of smooth sections of the associated line bundle $F^k = P \times_{k \cdot \mathbb{U}(1)} \mathbb{C}$ carries a natural symplectic structure $\Omega = \mu_\omega \hat{*} \omega_{\mathbb{C}}$. Moreover, the quantomorphism group $\text{Aut}_\Gamma(P)$ of P acts symplectically on $\Gamma^\infty(F^k)$ and has a classical momentum map $\mathcal{J}_{\text{Aut}_\Gamma} : \Gamma^\infty(F^k) \rightarrow \Omega^{2n}(M)$ given by

$$\mathcal{J}_{\text{Aut}_\Gamma}(\phi) = -k(n+1)\|\phi\|^2\mu_\omega + \frac{i}{2}\nabla^\Gamma\phi \wedge \nabla^\Gamma\bar{\phi} \wedge \frac{\omega^{n-1}}{(n-1)!}. \quad (4.94)$$

◇

A similar expression for the momentum map has been found by Donaldson [Don01, eq. 11], but with a few typos in the relative prefactors (in the derivation, [Don01, eq. 13] misses a factor of n). We emphasize that $\mathcal{J}_{\text{Aut}_\Gamma}$ is a classical momentum map with values in the dual vector space of $\text{aut}_\Gamma(P)$. The existence of a classical momentum map should not come as a surprise, because the quantomorphism group plays the role of the group of exact symplectomorphisms and as such is not sensitive to the same topological properties of M that the group of all symplectomorphisms sees.

4.4 Gauge theory

Atiyah and Bott [AB83] showed that the curvature furnishes a momentum map for the action of the group of gauge transformations on the space of connections over a surface. In this section, we outline how this setting fits in our general framework. Moreover, we also discuss the action of the whole group of bundle automorphisms, whose existence is tightly connected to the triviality of a certain class in the third cohomology of the base.

As we have already seen in Subsection 3.2.1, the group of gauge transformations admits an ordinary momentum map, and thus the concept of a group-valued momentum map is not needed for this action. Nonetheless, even in this case, our general framework provides the interesting insight that the momentum map geometry of the infinite-dimensional space of connection is induced by a finite-dimensional symplectic system.

Originally, Atiyah and Bott studied the Yang–Mills equation over a Riemannian surface. We consider the more general situation where the base is a compact $2n$ -dimensional symplectic manifold (M, σ) ; see also [Don87] for an almost identical setting.

Let $\pi : P \rightarrow M$ be a principal G -bundle, and let $\text{Ad}P \rightarrow M$ be the adjoint bundle of P . Recall that a connection Γ on P is a splitting of the exact sequence of vector bundles over M

$$0 \longrightarrow \text{Ad}P \longrightarrow (TP)/G \xrightarrow{\text{T}\pi} TM \longrightarrow 0, \quad (4.95)$$

where $\text{T}\pi$ also denotes the natural map on $(TP)/G$ induced by $\text{T}\pi : TP \rightarrow TM$.

The bundle $CP \rightarrow M$ of principal connections on P is, by definition, the subbundle of $L(TM, (TP)/G) \rightarrow M$ whose fiber $C_m P$ at $m \in M$ consists of all linear maps $\Gamma_m: T_m M \rightarrow (TP/G)_m$ satisfying $T\pi(\Gamma_m(X_m)) = X_m$ for all $X_m \in T_m M$. A smooth section $m \mapsto \Gamma_m$ of CP determines a principal G -connection on P as follows. The image of Γ_m yields a consistent choice of a subspace of $T_p P$ which is complementary to the vertical subspace for any $p \in \pi^{-1}(m)$. This induces a complementary subbundle of TP to the vertical subbundle that is invariant under the G -action, i.e., a horizontal G -equivariant distribution, and hence a principal connection. Conversely, the horizontal lift of a principal connection on P yields a splitting of (4.95) and hence a section of CP . Thus, the space $C(P)$ of principal connections on P is identified with the space of sections of CP . Note that $CP \rightarrow M$ is an affine bundle modeled over the vector bundle $L(TM, AdP) \rightarrow M$.

In order to apply our general framework, we need to realize CP as an associated fiber bundle with a symplectic fiber. This will require the notion of the principal jet prolongation WP of P . We now recall this construction (mostly suppressing the jet-theoretic background) and refer the reader to [KMS93, Section 15.6 and 17.5] for more details.

- The principal jet prolongation of the Lie group G is by definition the semidirect product

$$WG = (\mathrm{Sp}(2n, \mathbb{R}) \times G) \rtimes L(\mathbb{R}^{2n}, \mathfrak{g}) \quad (4.96)$$

with multiplication

$$(a, g, S) \cdot (b, h, T) = (a \cdot b, g \cdot h, \mathrm{Ad}_{h^{-1}} S \circ b + T) \quad (4.97)$$

and inversion

$$(a, g, S)^{-1} = (a^{-1}, g^{-1}, -\mathrm{Ad}_g S \circ a^{-1}). \quad (4.98)$$

The Lie algebra of WG is the semidirect product

$$W\mathfrak{g} = (\mathfrak{sp}(2n, \mathbb{R}) \times \mathfrak{g}) \rtimes L(\mathbb{R}^{2n}, \mathfrak{g}). \quad (4.99)$$

- The principal jet prolongation WP is the product fiber bundle $LM \times_M J^1 P$ over M , where LM denotes the symplectic frame bundle. Equivalently, the fiber $W_m P$ over $m \in M$ consists of pairs (u_m, v_p) of a linear symplectomorphism $u_m: \mathbb{R}^{2n} \rightarrow T_m M$ of \mathbb{R}^{2n} endowed with its standard symplectic form ω_o and $(T_m M, \sigma_m)$, and a linear map $v_p: T_m M \rightarrow T_p P$ satisfying $T_p \pi \circ v_p = \mathrm{id}_{T_m M}$. The right action of WG on WP defined by

$$(u_m, v_p) \cdot (a, g, S) := (u_m \circ a, v_p \cdot g + (p \cdot g) \cdot (S \circ a^{-1} \circ u_m^{-1})) \quad (4.100)$$

endows WP with the structure of a principal WG -bundle. Here, the lower

dot notation stands for the natural action of G on TP and of \mathfrak{g} on P .

- WG acts affinely on $L(\mathbb{R}^{2n}, \mathfrak{g})$ by

$$(a, g, S) \cdot T = \text{Ad}_g(S + T) \circ a^{-1}. \quad (4.101)$$

The associated bundle $WP \times_{WG} L(\mathbb{R}^{2n}, \mathfrak{g})$ is the bundle CP of connections. The horizontal lift Γ_m corresponding to the equivalence class $[u_m, v_p, T] \in WP \times_{WG} L(\mathbb{R}^{2n}, \mathfrak{g})$ is given by

$$\Gamma_m = v_p - p \cdot (T \circ u_m^{-1}): T_m M \rightarrow T_p P. \quad (4.102)$$

Assume that \mathfrak{g} is endowed with an Ad-invariant scalar product. We denote by $\text{Tr}(\cdot \wedge \cdot)$ the induced wedge product of linear maps with values in \mathfrak{g} . The linear symplectic form $\underline{\omega}$ on $L(\mathbb{R}^{2n}, \mathfrak{g})$ is defined by the identity

$$\underline{\omega}(T_1, T_2) \omega_0^n = \text{Tr}(T_1 \wedge T_2) \wedge \omega_0^{n-1}, \quad (4.103)$$

where ω_0 denotes the standard symplectic form on \mathbb{R}^{2n} . Clearly, $\underline{\omega}$ is invariant under the action of WG , because $a \in \text{Sp}(2n, \mathbb{R})$ leaves ω_0 invariant and because the scalar product on \mathfrak{g} is Ad-invariant. The associated momentum map $\underline{J}: L(\mathbb{R}^{2n}, \mathfrak{g}) \rightarrow \mathfrak{sp}(2n, \mathbb{R})^* \times \mathfrak{g}^* \times L(\mathbb{R}^{2n}, \mathfrak{g})^*$ is given by

$$\langle \underline{J}(T), (A, \xi, L) \rangle \omega_0^n = \left(\frac{1}{2} \text{Tr}(\xi, [T \wedge T]) - \text{Tr}(L \wedge T) + \frac{1}{2} \text{Tr}((T \circ A) \wedge T) \right) \wedge \omega_0^{n-1} \quad (4.104)$$

for $A \in \mathfrak{sp}(2n, \mathbb{R})$, $\xi \in \mathfrak{g}$ and $L \in L(\mathbb{R}^{2n}, \mathfrak{g})$. Thus the typical fiber $L(\mathbb{R}^{2n}, \mathfrak{g})$ of the connection bundle is a symplectic system with WG -symmetry. However, this system does not quite fit into the general setting discussed in Chapter 3, because the momentum map \underline{J} is not equivariant. In fact, a straightforward calculation shows that the non-equivariance 1-cocycle $\sigma(a, g, S) = \underline{J}((a, g, S) \cdot T) - \text{Ad}_{(a, g, S)^{-1}}^* \underline{J}(T)$ is given by

$$\sigma(a, g, S) = \text{Ad}_{(a, g, 0)^{-1}}^* \underline{J}(S). \quad (4.105)$$

The associated Lie algebra 2-cocycle $\Sigma: W\mathfrak{g} \times W\mathfrak{g} \rightarrow \mathbb{R}$ is

$$\Sigma((A_1, \xi_1, L_1), (A_2, \xi_2, L_2)) = \underline{\omega}(L_1, L_2). \quad (4.106)$$

For the same reason, we need to pass to a central extension of WG to obtain an equivariant prequantization bundle. Endow the trivial circle bundle $\underline{L} =$

$L(\mathbb{R}^{2n}, \mathfrak{g}) \times U(1)$ with the connection 1-form

$$\underline{\theta}_{(T,z)}(U, u) = \frac{1}{2}\underline{\omega}(T, U) + u, \quad (4.107)$$

where $T, U \in L(\mathbb{R}^{2n}, \mathfrak{g})$, $z \in U(1)$ and $u \in \mathbb{R} \simeq \mathfrak{u}(1)$. The curvature of $\underline{\theta}$ is $\underline{\omega}$, so that the trivial bundle \underline{L} endowed with the connection $\underline{\theta}$ is a prequantum bundle of the symplectic vector space $(L(\mathbb{R}^{2n}, \mathfrak{g}), \underline{\omega})$.

LEMMA 4.17 *The group of connection-preserving bundle automorphisms of the prequantum bundle $(\underline{L}, \underline{\theta})$ covering the action of WG on $L(\mathbb{R}^{2n}, \mathfrak{g})$ is identified with $\widehat{WG} = WG \times U(1)$ endowed with the multiplication*

$$\begin{aligned} (a, g, S, u) \star (b, h, T, v) \\ = \left(a \cdot b, g \cdot h, \text{Ad}_{h^{-1}} S \circ b + T, \exp\left(-\frac{i}{2}\underline{\omega}(S, \text{Ad}_h T \circ b^{-1})\right)uv \right), \end{aligned} \quad (4.108)$$

where $(a, g, S), (b, h, T) \in WG$ and $u, v \in U(1)$. The group \widehat{WG} is a non-trivial central extension of WG by $U(1)$ whose induced Lie algebra 2-cocycle is Σ . \diamond

Roughly speaking, one can view \widehat{WG} as the principal jet prolonged Heisenberg group of the symplectic vector space $(L(\mathbb{R}^{2n}, \mathfrak{g}), \underline{\omega})$.

Proof. Every bundle automorphism ϕ of $L(\mathbb{R}^{2n}, \mathfrak{g}) \times U(1)$ covering the action of $(a, g, S) \in WG$ is of the form

$$(T, z) \mapsto ((a, g, S) \cdot T, \phi_{(a,g,S)}(T)z) \quad (4.109)$$

for some smooth map $\phi_{(a,g,S)}: L(\mathbb{R}^{2n}, \mathfrak{g}) \rightarrow U(1)$. The automorphism ϕ preserves the connection $\underline{\theta}$ if and only if

$$\begin{aligned} \delta_T \phi_{(a,g,S)}(U) &= \frac{1}{2}\underline{\omega}(T, U) - \frac{1}{2}\underline{\omega}((a, g, S) \cdot T, (a, g, S) \cdot U) \\ &= \frac{1}{2}\underline{\omega}(T, U) - \frac{1}{2}\underline{\omega}(\text{Ad}_g(S + T) \circ a^{-1}, \text{Ad}_g U \circ a^{-1}) \\ &= -\frac{1}{2}\underline{\omega}(S, U). \end{aligned} \quad (4.110)$$

Consequently, $\phi_{(a,g,S)}(T) = \exp\left(-\frac{i}{2}\underline{\omega}(S, T)\right)u$ for some $u \in U(1)$. One easily verifies that the assignment of the bundle automorphism

$$\phi_{(a,g,S,u)}(T, z) = \left((a, g, S) \cdot T, \exp\left(-\frac{i}{2}\underline{\omega}(S, T)\right)uz \right) \quad (4.111)$$

to $(a, g, S, u) \in \widehat{WG}$ is a group isomorphism with respect to the product (4.108). \square

Finally, observe that the action of \widehat{WG} on $L(\mathbb{R}^{2n}, \mathfrak{g})$ is transitive, and that the stabilizer of o is the direct product group $\mathrm{Sp}(2n, \mathbb{R}) \times G \times \mathrm{U}(1)$. We can thus view $L(\mathbb{R}^{2n}, \mathfrak{g})$ as the homogeneous space $\widehat{WG}/(\mathrm{Sp}(2n, \mathbb{R}) \times G \times \mathrm{U}(1))$. The Lie group homomorphism that characterizes the \widehat{WG} -equivariant prequantization bundle $(\underline{L}, \underline{\theta})$ according to Section 4.2 is given by

$$\rho: \mathrm{Sp}(2n, \mathbb{R}) \times G \times \mathrm{U}(1) \rightarrow \mathrm{U}(1), \quad (a, g, u) \mapsto u. \quad (4.112)$$

In summary, one needs to pass to the central extension \widehat{WG} of WG to obtain a Hamiltonian system $(L(\mathbb{R}^{2n}, \mathfrak{g}), \underline{\omega})$ with equivariant momentum map and equivariant prequantum bundle $(\underline{L}, \underline{\theta})$.

The issue of non-equivariance extends to the bundle picture as well. For example, the action of the gauge group of WP on the space $C(P)$ of principal connections on P has a non-equivariant momentum map and one needs to pass to an extension to get an equivariant momentum map. Surprisingly perhaps, these problems disappear under the jet prolongation, and we obtain the following generalization of the result of Atiyah and Bott [AB83, p. 587] to higher dimensional base manifolds.

THEOREM 4.18 *Let (M, σ) be a compact $2n$ -dimensional symplectic manifold, and let $\pi: P \rightarrow M$ be a principal G -bundle with compact structure group G . The 2-form*

$$\Omega(\alpha, \beta) = \int_M \mathrm{Tr}(\alpha \wedge \beta) \wedge \sigma^{n-1}, \quad \alpha, \beta \in \Omega^1(M, \mathrm{Ad}P) \quad (4.113)$$

defines a symplectic structure on the space $C(P)$ of principal connections on P . Moreover, the natural action of the group $\mathrm{Gau}(P)$ of gauge transformations of P on $C(P)$ leaves Ω invariant, and possesses an equivariant momentum map

$$\mathcal{J}_{\mathrm{Gau}}: C(P) \rightarrow \Omega^{2n}(M, \mathrm{Ad}P), \quad \Gamma \mapsto \mathrm{curv} \Gamma \wedge \sigma^{n-1}, \quad (4.114)$$

where $\mathrm{curv} \Gamma \in \Omega^2(M, \mathrm{Ad}P)$ is the curvature of Γ . ◇

Proof. The claim follows from a simple direct calculation along the lines of [AB83, p. 587]. We now sketch how the same result can be obtained in our framework utilizing the fiberwise symplectic system.

Theorem 3.7 shows that the momentum map for the action of the gauge group of WP is the push-forward of the fiber momentum map (4.104), that is,

$$\langle \mathcal{J}_{\mathrm{Gau}(WP)}(\Gamma), \eta \rangle = \int_M \langle J \circ T_\Gamma, \eta \rangle \sigma^n, \quad (4.115)$$

where $\eta \in \mathrm{gau}(WP)$, and where $T_\Gamma: WP \rightarrow L(\mathbb{R}^{2n}, \mathfrak{g})$ denotes the equivariant

map corresponding to the connection Γ ; that is, $\Gamma_p = v_p - p \cdot (T_\Gamma(u_m, v_p) \circ u_m^{-1})$, see (4.102). The momentum map for $\text{Gau}(P)$ is obtained by composing with the projection $\mathfrak{gau}(WP)^* \rightarrow \mathfrak{gau}(P)^*$ dual to the derivative of the natural prolongation map $W: \text{Gau}(P) \rightarrow \text{Gau}(WP)$, cf. [KMS93, p. 151]. If $\chi \in \text{Gau}(P)$ is viewed as an equivariant map $\chi: P \rightarrow G$, then we find for the prolongation $W\chi: WP \rightarrow WP$

$$\begin{aligned} W\chi(u_m, v_p) &= (u_m, v_p \cdot \chi(p) + p \cdot (T_p\chi \circ v_p)) \\ &= (u_m, v_p) \cdot (\text{id}, \chi(p), \chi(p)^{-1} \cdot (T_p\chi \circ v_p \circ u_m)), \end{aligned} \quad (4.116)$$

where we used (4.100) for the second equality. Thus, changing viewpoints again, we read off

$$W\chi: WP \rightarrow WG, \quad (u_m, v_p) \mapsto (\text{id}, \chi(p), \delta_p^L \chi \circ v_p \circ u_m). \quad (4.117)$$

On the Lie algebra level, $\xi \in \mathfrak{gau}(P)$ seen as a map $\xi: P \rightarrow \mathfrak{g}$ is prolonged to

$$W\xi: WP \rightarrow W\mathfrak{g}, \quad (u_m, v_p) \mapsto (0, \xi(p), T_p\xi \circ v_p \circ u_m). \quad (4.118)$$

The momentum map $\mathcal{J}_{\text{Gau}(P)}$ for the action of $\text{Gau}(P)$ satisfies, by definition,

$$\langle \mathcal{J}_{\text{Gau}(P)}(\Gamma), \xi \rangle = \langle \mathcal{J}_{\text{Gau}(WP)}(\Gamma), W\xi \rangle = \int_M \langle J \circ T_\Gamma, W\xi \rangle \sigma^n. \quad (4.119)$$

For the integrand we find using (4.104)

$$\begin{aligned} &\langle J \circ T_\Gamma, W\xi \rangle(u_m, v_p) \sigma_m^n \\ &= \left(\frac{1}{2} \text{Tr}(\xi(p), [T_\Gamma(u_m, v_p) \wedge T_\Gamma(u_m, v_p)]) \right. \\ &\quad \left. - \text{Tr}((T_p\xi \circ v_p \circ u_m) \wedge T_\Gamma(u_m, v_p)) \right) \wedge \omega_0^{n-1} \\ &= \left(\frac{1}{2} \text{Tr}(\xi(p), [T_\Gamma(v_p) \wedge T_\Gamma(v_p)]) - \text{Tr}((T_p\xi \circ v_p) \wedge T_\Gamma(v_p)) \right) \wedge \sigma^{n-1}, \end{aligned} \quad (4.120)$$

where we abbreviated $T_\Gamma(v_p) \equiv T_\Gamma(u_m, v_p) \circ u_m^{-1}$. After partial integration on the second term, we use the structure equation $\text{curv } \Gamma = \frac{1}{2}[\Gamma \wedge \Gamma] + d\Gamma$ to obtain

$$\langle \mathcal{J}_{\text{Gau}(P)}(\Gamma), \xi \rangle = \int_M \text{Tr}(\xi, \text{curv } \Gamma) \wedge \sigma^{n-1}. \quad (4.121)$$

Thus $\Gamma \mapsto \text{curv } \Gamma \wedge \sigma^{n-1}$ is the momentum map for the action of $\text{Gau}(P)$, as claimed. \square

The discussion of the momentum map for the action of the full automorphism group of P is more involved and requires the following additional structure.

Recall the central extension

$$U(1) \longrightarrow \widehat{WG} \longrightarrow WG. \quad (4.122)$$

A lift of WP to \widehat{WG} is a principal \widehat{WG} -bundle \widehat{WP} together with a vertical bundle morphism $\chi: \widehat{WP} \rightarrow WP$ intertwining the principal actions of \widehat{WG} and WG . The bundle morphism χ induces an isomorphism of the bundles $\widehat{WP}/U(1)$ and WP . We will usually suppress this isomorphism and directly view \widehat{WP} as a principal $U(1)$ -bundle over WP . Note, however, that such a lift \widehat{WP} does not need to exist. In fact, the existence and uniqueness question is governed by the following exact sequence in Čech cohomology

$$\begin{array}{ccccccc} \hat{H}^1(M, U(1)) & \longrightarrow & \hat{H}^1(M, \widehat{WG}) & \longrightarrow & \hat{H}^1(M, WG) & \longrightarrow & \hat{H}^2(M, U(1)) \\ \parallel & & & & & & \parallel \\ H^2(M, \mathbb{Z}) & & & & & & H^3(M, \mathbb{Z}) \end{array} \quad (4.123)$$

induced by the exact sequence (4.122), see [Bry07, Theorem 4.1.4]. The vertical isomorphisms are induced by the short exact sequence $\mathbb{Z} \rightarrow \mathbb{R} \rightarrow U(1)$. As the Čech cohomology $\hat{H}^1(M, K)$ parameterizes equivalence classes of K -bundles, this exact sequence shows that there exists a lift of WP to \widehat{WG} given that a certain class in $H^3(M, \mathbb{Z})$ vanishes. In fact, this class admits the following description, cf. [Bry07, p. 161]. Choose an open covering U_i of M , and let $g_{ij}: U_i \cap U_j \rightarrow G$ be the transition functions of P . By jet prolongation, we obtain the transition functions $Wg_{ij}: U_i \cap U_j \rightarrow WG$ of WP . Choose a lift \widehat{Wg}_{ij} of Wg_{ij} to \widehat{WG} . These are the candidate transition functions for \widehat{WP} . However, they satisfy the cocycle condition of triple overlaps $U_i \cap U_j \cap U_k$ only in the form

$$\widehat{Wg}_{ij} \widehat{Wg}_{jk} = u_{ijk} \widehat{Wg}_{ik} \quad (4.124)$$

for some $U(1)$ -valued map u_{ijk} . The class of u_{ijk} in $\hat{H}^2(M, U(1)) \simeq H^3(M, \mathbb{Z})$ is the obstruction for the lift. We emphasize that this class in the third cohomology can be constructed for each principal bundle over a symplectic manifold in a natural and canonical way. Although being canonical, we are not aware that this class has been discussed before in the literature. Recall that elements in $H^3(M, \mathbb{Z})$ parameterize equivalence classes of bundle gerbes (in the same way $H^2(M, \mathbb{Z})$ parameterizes principal circle bundles). The bundle gerbe corresponding to $[u_{ijk}]$ is known as the so-called lifting gerbe associated to the lifting problem of WP to \widehat{WG} , see [Mur96, Section 4]. It comes equipped with a natural connection and curving [Gom03]. In this context, the following open question is of interest.

PROBLEM 4.19 Find an explicit description of the lifting gerbe (and the class

in $H^3(M, \mathbb{Z})$) that obstructs the lifting of WP to \widehat{WG} . In particular, clarify the geometric meaning of this gerbe that is naturally associated to each principal bundle over a symplectic manifold. \diamond

In the following, we assume that there exists a lift \widehat{WP} of WP to \widehat{WG} . According to the sequence (4.123), such a lift is unique up to a principal circle bundle on M . The natural identifications

$$\begin{aligned} \widehat{WP} \times_{\widehat{WG}} (\widehat{WG}/(\mathrm{Sp}(2n, \mathbb{R}) \times G \times \mathrm{U}(1))) &\simeq \widehat{WP} \times_{\widehat{WG}} \mathrm{L}(\mathbb{R}^{2n}, \mathfrak{g}) \\ &\simeq WP \times_{WG} \mathrm{L}(\mathbb{R}^{2n}, \mathfrak{g}) \\ &\simeq CP \end{aligned} \quad (4.125)$$

allow us to view a connection Γ on P as a reduction Q_Γ of \widehat{WP} to the structure group $\mathrm{Sp}(2n, \mathbb{R}) \times G \times \mathrm{U}(1)$. We are thus in the setting discussed in Section 4.2. Assume that there exists a principal connection on \widehat{WP} that is reducible to each reduction Q_Γ . Then, by Theorem 4.9, the group of bundle automorphism of \widehat{WP} has an equivariant group-valued momentum map, whose second component is given by the principal $\mathrm{U}(1)$ -bundle $Q_\Gamma \times_{\rho^{-1}} \mathrm{U}(1)$. We would like to proceed as in the proof of Theorem 4.18, and pull back everything along the jet prolongation $W: \mathrm{Aut}(P) \rightarrow \mathrm{Aut}(WP)$. However, we are faced with the problem that W takes values in $\mathrm{Aut}(WP)$ and not in $\mathrm{Aut}(\widehat{WP})$. Clearly, we have the sequence

$$e \longrightarrow C^\infty(M, \mathrm{U}(1)) \longrightarrow \mathrm{Aut}(\widehat{WP}) \longrightarrow \mathrm{Aut}(WP). \quad (4.126)$$

However, the last map in this sequence is in general not surjective. If the $\mathrm{U}(1)$ -bundle $\widehat{WP} \rightarrow WP$ has a \widehat{WG} -invariant connection, then we can use its horizontal lift to show exactness of the infinitesimal sequence:

$$0 \longrightarrow C^\infty(M, \mathbb{R}) \longrightarrow \mathrm{aut}(\widehat{WP}) \longrightarrow \mathrm{aut}(WP) \longrightarrow 0. \quad (4.127)$$

In this case, we can pull-back (4.127) along the jet prolongation $W: \mathrm{aut}(P) \rightarrow \mathrm{aut}(WP)$ to obtain an Abelian extension

$$0 \longrightarrow C^\infty(M, \mathbb{R}) \longrightarrow \widehat{\mathrm{aut}}(P) \longrightarrow \mathrm{aut}(P) \longrightarrow 0. \quad (4.128)$$

Under these assumptions, we conclude that the Lie algebra $\widehat{\mathrm{aut}}(P)$ acts on the space $C(P)$ of connections and that it has an (infinitesimal) equivariant momentum map. Comparing with the usual non-equivariance picture, we conjecture that the $\mathrm{Aut}(P)$ -action has a non-equivariant momentum map and that the passage to the extension is only necessary to obtain an equivariant momentum map. In this context, one would also expect that an extension by \mathbb{R} is sufficient instead of an

extension by the much larger space $C^\infty(M, \mathbb{R})$. Moreover, it is not clear whether the extension (4.128) splits. As the general methods and techniques developed in Chapter 3 are tailored to symplectic fiber models with *equivariant* momentum maps, these questions go beyond the scope of this paper.

APPENDICES

A Notations and conventions

Because notations and, especially, sign and coefficient conventions are not uniform in the literature, we provide below those used in this article. We also comment on some peculiarities of the infinite-dimensional setting.

By a manifold we understand a possibly infinite-dimensional smooth manifold without boundary modeled on a locally convex space as, for example discussed in [Nee06]. If $f: M \rightarrow N$ is a smooth map between manifolds, its derivative, or tangent map, at $m \in M$ is denoted by $T_m f: T_m M \rightarrow T_{f(m)} N$. The Lie algebra of vector fields on a smooth manifold M is denoted by $\mathfrak{X}(M)$, the Jacobi-Lie bracket on $\mathfrak{X}(M)$ by $[\cdot, \cdot]$, the space of k -exterior differential forms by $\Omega^k(M)$, the space of k -contravariant tensors by $\mathfrak{X}^k(M)$, the exterior differential by $d: \Omega^k(M) \rightarrow \Omega^{k+1}(M)$, the Lie derivative along a vector field $Z \in \mathfrak{X}(M)$ by \mathfrak{L}_Z (acting on any tensor field), the interior product (contraction on the first index) of a form $\alpha \in \Omega^k(M)$ with $Z \in \mathfrak{X}(M)$ by $i_Z \alpha = Z \lrcorner \alpha \in \Omega^{k-1}(M)$. In infinite dimensions, the cotangent bundle is no longer a *smooth* bundle [Nee06, Remark 1.3.9] and hence smoothness of differential forms has to be specified explicitly. A differential form is defined as a set-theoretic section of the exterior bundle such that all chart representations are smooth as maps $U \times E^k \rightarrow \mathbb{R}$, where U is an open subset of M and E denotes the model space of M . For the wedge product $\alpha \wedge \beta \in \Omega^{k+l}(M)$ of the differential forms $\alpha \in \Omega^k(M)$ and $\beta \in \Omega^l(M)$ we use the Bourbaki convention

$$(\alpha \wedge \beta)(X_1, \dots, X_{k+l}) = \sum_{\sigma} (\text{sign } \sigma) \alpha(X_{\sigma(1)}, \dots, X_{\sigma(k)}) \beta(X_{\sigma(k+1)}, \dots, X_{\sigma(k+l)}),$$

where $X_1, \dots, X_{k+l} \in \mathfrak{X}(M)$ and the sum is taken over all (k, l) -shuffles σ , i.e., permutations of $\{1, 2, \dots, k+l\}$ such that $\sigma(1) < \dots < \sigma(k)$ and $\sigma(k+1) < \dots < \sigma(k+l)$. Evaluation of a tensor field ξ on M at $m \in M$ is denoted by $\xi_m = \xi(m)$, yielding a tensor of the same type on $T_m M$. The vector space of skew-symmetric k -linear maps on $T_m^* M$ and $T_m M$ is denoted by $\Lambda^k(T_m^* M)$ and $\Lambda^k(T_m M)$, respectively; thus if $\alpha \in \Omega^k(M)$ and $m \in M$, then $\alpha_m \in \Lambda^k(T_m^* M)$.

Given topological vector spaces V and V^* (not necessarily the functional analytic dual), $\langle \cdot, \cdot \rangle: V^* \times V \rightarrow \mathbb{R}$ denotes a weakly non-degenerate pairing. If W is another topological vector space in weak duality with some topological vector

space W^* and $T: V \rightarrow W$ is a linear continuous map, then its dual $T^*: W^* \rightarrow V^*$ is defined by $\langle T^*\alpha, v \rangle := \langle \alpha, Tv \rangle$, for any $v \in V$ and $\alpha \in W^*$. Weak non-degeneracy of the pairings guarantees only uniqueness of the linear dual map T^* , not its existence, so in concrete situations, its existence needs to be checked.

Given a Lie group G , left and right translations by $a \in G$ are denoted by $b \ni G \mapsto L_a b := ab \in G$ and $b \ni G \mapsto R_a b := ba \in G$, respectively. The tangent map $T_b L_a: T_b G \rightarrow T_{ab} G$ of L_a at an arbitrary $b \in G$ is denoted, for simplicity, by the same symbol L_a ; similar conventions are used for right translation. To shorten notation even further, we will abbreviate $T_b L_a(A_b)$ for $A_b \in T_b G$ by $a \cdot A_b \in T_{ab} G$. Similarly, we write $A_b \cdot a \equiv T_b R_a(A_b)$ for the right multiplication. The Lie algebra of G is denoted by \mathfrak{g} and the (left) adjoint representation by $\text{Ad}_a := T_e(L_a \circ R_{a^{-1}}): \mathfrak{g} \rightarrow \mathfrak{g}$; Ad_a is a Lie algebra isomorphism for any $a \in G$. The (left) coadjoint representation of G on the dual \mathfrak{g}^* of \mathfrak{g} is denoted by $\text{Ad}_{a^{-1}}^*: \mathfrak{g}^* \rightarrow \mathfrak{g}^*$ for any $a \in G$. The adjoint and coadjoint Lie algebra representations of \mathfrak{g} on \mathfrak{g} and \mathfrak{g}^* , respectively, are denoted by $\text{ad}_A B := [A, B]$ and $\text{ad}_A^*: \mathfrak{g}^* \rightarrow \mathfrak{g}^*$, respectively, for any $A, B \in \mathfrak{g}$. We have $T_e(\text{Ad} \cdot B)(A) = [A, B] = \text{ad}_A B$ for any $A, B \in \mathfrak{g}$.

The space of \mathfrak{g} -valued k -forms on a manifold M is denoted by $\Omega^k(M, \mathfrak{g})$. The wedge product associated to the Lie bracket $[\cdot, \cdot]$ on \mathfrak{g} , denoted by $[\alpha \wedge \beta] \in \Omega^{k+l}(M, \mathfrak{g})$ for any $\alpha \in \Omega^k(M, \mathfrak{g})$ and $\beta \in \Omega^l(M, \mathfrak{g})$, is defined with the same coefficient conventions as in the scalar case, i.e.,

$$[\alpha \wedge \beta](X_1, \dots, X_{k+l}) = \sum_{\sigma} (\text{sign } \sigma) [\alpha(X_{\sigma(1)}, \dots, X_{\sigma(k)}), \beta(X_{\sigma(k+1)}, \dots, X_{\sigma(k+l)})],$$

the sum being over all (k, l) -shuffles σ and $X_1, \dots, X_{k+l} \in \mathfrak{X}(M)$ arbitrary.

If $\Upsilon: G \times M \rightarrow M$ is a left Lie group action and $A \in \mathfrak{g}$, the infinitesimal generator, or fundamental, vector field defined by A is denoted by

$$A_m^* := \left. \frac{d}{dt} \right|_{t=0} \Upsilon(\exp(tA), m) \in T_m M$$

for any $m \in M$, where $\exp: \mathfrak{g} \rightarrow G$ is the exponential map of G . We have $[A, B]^* = -[A^*, B^*]$ for all $A, B \in \mathfrak{g}$. If the G -action on M is on the right, then $[A, B]^* = [A^*, B^*]$. If the action is clear from the context, we also use the “dot notation” for the action and its derivatives. So, for $m \in M$ and $g \in G$, we abbreviate $g \cdot m = \Upsilon(g, m)$. On the infinitesimal level, for $A \in \mathfrak{g}$, we will write the fundamental vector field as $A_m^* = A \cdot m$. Similarly, $g \cdot X_m = T_m \Upsilon_g(X_m)$.

If $f: M \rightarrow G$ is a smooth function, the left logarithmic derivative $\delta_m f: T_m M \rightarrow \mathfrak{g}$ of f at $m \in M$ is defined by $\delta_m f := L_{f(m)}^{-1} \circ T_m f$. Therefore $\delta f \in \Omega^1(M, \mathfrak{g})$.

A Poisson structure on a smooth finite-dimensional manifold M is a Lie algebra structure $\{\cdot, \cdot\}$ on the vector space $C^\infty(M)$ of smooth functions satisfying the Leibniz identity $\{fg, h\} = f\{g, h\} + \{f, h\}g$ for any $f, g, h \in C^\infty(M)$. Thus, $\{h, \cdot\}: C^\infty(M) \rightarrow C^\infty(M)$ is a derivation and hence defines a vector field $X_h \in$

$\mathfrak{X}(M)$ on M , i.e., $\langle df, X_h \rangle := \{h, f\}$ for any $f \in C^\infty(M)$, called the Hamiltonian vector field of h . Hamilton's equations $\frac{d}{dt}m(t) = X_h(m(t))$ can be written in Poisson bracket form as $\frac{d}{dt}f(m(t)) = \{h, f\}(m(t))$ for any $f \in C^\infty(M)$. The Poisson tensor $\omega \in \mathfrak{X}^2(M)$ is defined by $\omega(df, dg) := \{f, g\}$ for any $f, g \in C^\infty(M)$ and the Schouten bracket of ω with itself vanishes. Conversely, any $\omega \in \mathfrak{X}^2(M)$ with this property defines a Poisson bracket by the formula above. Denote by $\omega^\sharp: T^*M \rightarrow TM$ the vector bundle map covering the identity defined by $\langle \alpha, \omega^\sharp \beta \rangle := \omega(\alpha, \beta)$ for any $\alpha, \beta \in \Omega^1(M)$. If the map ω^\sharp is invertible, its inverse, $\omega^\flat: TM \rightarrow T^*M$ defines a closed two form $\omega \in \Omega^2(M)$ by $\omega(X_1, X_2) := \langle \omega^\flat(X_1), X_2 \rangle = \omega\left((\omega^\sharp)^{-1}X_1, (\omega^\sharp)^{-1}X_2\right)$ for any $X_1, X_2 \in \mathfrak{X}(M)$. In this case, (M, ω) is a symplectic manifold. The Hamiltonian vector field X_h of $h \in C^\infty(M)$ is given then, alternatively, by the identity $X_h \lrcorner \omega = -dh$. Any symplectic manifold (M, ω) is Poisson by setting $\{f, h\} := \omega(X_f, X_h)$.

If Q is finite dimensional, the cotangent bundle is naturally an exact symplectic manifold $(T^*Q, \omega_{\text{can}})$, where $\omega_{\text{can}} = d\theta_{\text{can}}$, where $\theta_{\text{can}} \in \Omega^1(T^*Q)$ is given by $\langle \theta_{\text{can}}(\alpha_q), V_{\alpha_q} \rangle = \langle \alpha_q, T_q \pi(V_{\alpha_q}) \rangle$, for all $\alpha_q \in T^*Q$, $V_{\alpha_q} \in T_{\alpha_q}(T^*Q)$, and $\pi: T^*Q \ni \alpha_q \mapsto q \in Q$ is the cotangent bundle projection. In canonical cotangent bundle coordinates $(q^1, \dots, q^n, p_1, \dots, p_n)$, we have $\theta_{\text{can}} = p_i dq^i$ and $\omega_{\text{can}} = dp_i \wedge dq^i$. Thus, the Hamiltonian vector field is given by $X_f = \frac{\partial f}{\partial p_i} \partial_{q^i} - \frac{\partial f}{\partial q^i} \partial_{p_i}$ and the Poisson bracket takes the form $\{f, h\} = \frac{\partial f}{\partial p_i} \frac{\partial h}{\partial q^i} - \frac{\partial f}{\partial q^i} \frac{\partial h}{\partial p_i}$.

Let (M, ω) be a symplectic G -manifold. A momentum map for the action, if it exists, is a smooth map $J: M \rightarrow \mathfrak{g}^*$ satisfying

$$A^* \lrcorner \omega + dJ_A = 0, \quad A \in \mathfrak{g},$$

where $J_A := \langle A, J \rangle: M \rightarrow \mathbb{R}$.

A principal G -bundle is a fiber bundle $P \rightarrow M$ together with a smooth right action of G on P , which is fiber-preserving, free, and transitive on each fiber. In particular, $M = P/G$. Given a left G -manifold \underline{E} , we form the associated bundle $F = P \times_G \underline{E} := (P \times \underline{E})/G$ as the quotient of $P \times \underline{E}$ by the natural free right action of G , i.e., $(p, \underline{f}) \cdot g := (p \cdot g, g^{-1} \cdot \underline{f})$, for any $p \in P$, $\underline{f} \in \underline{E}$, and $g \in G$. Denoting by $[p, \underline{f}]$ points in $P \times_G \underline{E}$, we have the identity

$$[p \cdot g, \underline{f}] = [p, g \cdot \underline{f}],$$

often used in the text. The smooth projection map $P \times_G \underline{E} \ni [p, \underline{f}] \mapsto [p] \in M = P/G$ defines a locally trivial fiber bundle with total space $P \times_G \underline{E}$, base M , and typical fiber \underline{E} .

An important particular case of this construction occurs by taking $\underline{E} = \mathfrak{g}$, where \mathfrak{g} is the Lie algebra of G , and the G -action on \mathfrak{g} is the adjoint action. In this case,

the associated bundle is called the adjoint bundle and is denoted by $\text{Ad}P$; it is a vector bundle all of whose fibers are Lie algebras and the bracket operation depends smoothly on the base point, i.e., $\text{Ad}P$ is a Lie algebra bundle. Taking $\underline{F} = \mathfrak{g}^*$ and the coadjoint action on \mathfrak{g}^* yields as associated bundle the coadjoint bundle $\text{Ad}^*P \rightarrow M$. Each fiber of this vector bundle carries the Lie-Poisson bracket which depends smoothly on the base, i.e., Ad^*P is a Lie-Poisson vector bundle. A third useful example is the conjugation bundle $\text{Conj}P \rightarrow M$ which is the associated bundle obtained by taking $\underline{F} = G$ with the G -action being the conjugation on G . The fibers of this bundle carry a natural group structure, depending smoothly on the base, so this is a group fiber bundle.

B Fiber integration

In this appendix, we review the method of fiber integration in the special case of trivial bundles, because it appears often in formulas and computations in the text. All the proofs of the following statements can be found in [GHV72, Section VII.5], where also the more general situation of locally trivial fiber bundles is discussed.

Let $\pi: M \times F \rightarrow M$ be a finite-dimensional trivial fiber bundle. Assume that the fiber F is oriented and closed (i.e., F is compact and without boundary). Integrating a differential form $\alpha \in \Omega^k(M \times F)$ over the fiber F yields a $(k - \dim F)$ -form $\int_F \alpha$ on the base M by defining

$$\left(\int_F \alpha \right)_m := \int_F \alpha_m \in \Lambda^{k-\dim F}(\mathbb{T}_m^*M), \quad (\text{B.1})$$

where, for $m \in M$, the *retrenchment* $\alpha_m \in \Omega^{\dim F}(F, \Lambda^{k-\dim F}(\mathbb{T}_m^*M))$, a $(\dim F)$ -form on F with values in the vector space $\Lambda^{k-\dim F}(\mathbb{T}_m^*M)$, is defined by

$$\begin{aligned} & \langle (\alpha_m)_f(Z_1, \dots, Z_{\dim F}), X_1 \wedge \dots \wedge X_{k-\dim F} \rangle \\ & = \alpha_{(m,f)}(X_1, \dots, X_{k-\dim F}, Z_1, \dots, Z_{\dim F}) \end{aligned} \quad (\text{B.2})$$

for $m \in M$, $f \in F$, $X_i \in \mathbb{T}_mM$, and $Z_j \in \mathbb{T}_fF$. This defines a linear map

$$\int_F : \Omega^k(M \times F) \rightarrow \Omega^{k-\dim F}(M). \quad (\text{B.3})$$

REMARK B.1 Note that, on a formal level, the discussion above also applies to the case of an infinite-dimensional manifold M , because in the end, we are just integrating over the finite-dimensional fiber F . However, a direct and carefree translation of the previous approach to an infinite-dimensional manifold M

is impeded by the fact that the cotangent bundle T^*M is not a *smooth* bundle in this case [Nee06, Remark I.3.9]. Thus, it is then not clear in which sense the retraction depends smoothly on $m \in M$. Moreover, to integrate a form $\beta \in \Omega^{\dim F}(F, V)$ on F with values in a locally convex space V , one would need at least Mackey completeness of V (see [Nee06, Remark I.4.4]). To circumvent these problems for infinite-dimensional base manifolds, we directly define the fiber integral by

$$\left(\int_F \alpha \right)_m (X_1, \dots, X_{k-\dim F}) = \int_F \alpha_{(m, \cdot)}(X_1, \dots, X_{k-\dim F}, \cdot). \quad (\text{B.4})$$

Here, the integrand is a real valued, top-degree form on F . We then have to check that $\int \alpha$ is indeed a *smooth* form on M . But this follows from the local expression as in the finite-dimensional case; compare [GHV72, Proof of Proposition VII.5.7.11.VII]. \diamond

Fiber integration enjoys the following properties:

(i) Equivariance with respect to maps on the base:

$$\phi^* \int_F \alpha = \int_F (\phi \times \text{id}_F)^* \alpha \quad (\text{B.5})$$

holds for all smooth maps $\phi: M \rightarrow M$, see [GHV72, Proposition VII.5.7.12.VIII].

(ii) Invariance under fiber transformations: for all orientation-preserving diffeomorphisms ψ of F ,

$$\int_F (\text{id}_M \times \psi)^* \alpha = \int_F \alpha, \quad (\text{B.6})$$

see [GHV72, Proposition VII.5.7.12.VIII].

(iii) Contraction with vector fields:

$$X \lrcorner \int_F \alpha = \int_F (X \times \text{id}_F) \lrcorner \alpha \quad (\text{B.7})$$

holds for all vector fields $X \in \mathfrak{X}(M)$. Furthermore, for $Z \in \mathfrak{X}(F)$, we have

$$\int_F (\text{id}_M \times Z) \lrcorner \alpha = 0, \quad (\text{B.8})$$

see [GHV72, Proposition VII.5.7.13.X].

(iv) Fiber integration commutes with the exterior differential on M and $M \times F$:

$$d \int_F \alpha = \int_F d\alpha, \quad (\text{B.9})$$

see [GHV72, Proposition VII.5.7.13.X]. In this formula, the fact that F is boundaryless is essential. If F has a non-empty boundary, the formula above has contributions from the boundary (see [GHV72, Problem VII.4]):

$$d \int_F \alpha = \int_F d\alpha + (-1)^{\deg \alpha + \dim F} \int_{\partial F} \alpha. \quad (\text{B.10})$$

(v) Up-down formula: for any $\beta \in \Omega^k(M)$ and $\alpha \in \Omega^l(M \times F)$, we have

$$\int_F \pi^* \beta \wedge \alpha = \beta \wedge \int_F \alpha, \quad (\text{B.11})$$

see [GHV72, Proposition VII.5.7.13.IX].

C Hat product for fiber bundles

Vizman [Viz11] introduced the hat product of differential forms as a method to construct differential forms on the mapping space $C^\infty(M, F)$ starting from differential forms on M and F . We generalize these ideas to induce differential forms on the space of smooth sections of a fiber bundle.

Let M be a compact, oriented, finite-dimensional manifold without boundary and let $\pi: F \rightarrow M$ be a finite-dimensional fiber bundle over M . Then the space \mathcal{F} of smooth sections of F is a Fréchet manifold. The evaluation map $\text{ev}: M \times \mathcal{F} \rightarrow F$ assigns to a section $\phi \in \mathcal{F}$ and a point $m \in M$ the value $\phi(m) \in F$. Furthermore, let $\text{pr}_M: M \times \mathcal{F} \rightarrow M$ denote the projection on the first factor. We define the *hat product* as the bilinear map¹

$$\hat{*}: \Omega^k(M) \times \Omega^l(F) \rightarrow \Omega^{k+l-\dim M}(\mathcal{F}), \quad \alpha \hat{*} \omega = \int_M \text{pr}_M^* \alpha \wedge \text{ev}^* \omega, \quad (\text{C.1})$$

where the fiber integration over the trivial bundle $M \times \mathcal{F} \rightarrow \mathcal{F}$ is defined as explained in Remark B.1. Unravelling the definitions using (B.4) gives the explicit

¹ We use a different sign convention than the one used in [Viz11].

formula

$$(\alpha \hat{*} \omega)_\phi(Y_1, \dots, Y_r) = (-1)^{kr} \int_M \alpha \wedge \phi^*(Y_r \lrcorner \cdots Y_1 \lrcorner (\omega \circ \phi)), \quad (\text{C.2})$$

where $r = k + l - \dim M$ denotes the degree of the hat product, $\alpha \in \Omega^k(M)$, $\omega \in \Omega^l(F)$, $Y_i \in \Gamma^\infty(\phi^*VF)$, and $VF \subset TF$ is the vertical subbundle. Moreover, we used the abbreviation $\phi^*(Y_r \lrcorner \cdots Y_1 \lrcorner (\omega \circ \phi)) \in \Omega^{l-r}(M)$ for the partial pull-back

$$\begin{aligned} & \phi^*(Y_r \lrcorner \cdots Y_1 \lrcorner (\omega \circ \phi))_m(X_1, \dots, X_{l-r}) \\ &= \omega_{\phi(m)}(Y_1(m), \dots, Y_r(m), T_m\phi(X_1), \dots, T_m\phi(X_{l-r})) \\ &= (\text{ev}^*\omega)_{m,\phi}(Y_1, \dots, Y_r, X_1, \dots, X_{l-r}). \end{aligned} \quad (\text{C.3})$$

REMARK C.1 In the special case where α is a volume form μ on M , we get the following l -form on \mathcal{F}

$$(\mu \hat{*} \omega)_\phi(Y_1, \dots, Y_l) = (-1)^{l \dim M} \int_M \phi^*(Y_l \lrcorner \cdots Y_1 \lrcorner (\omega \circ \phi)) \mu. \quad (\text{C.4})$$

Since $Y_i(m)$ is a vertical tangent vector at $\phi(m)$, we see that $\mu \hat{*} \omega$ only depends on the vertical component of the form ω . Hence, in this case, we can also consider the hat product of μ with a differential form ω along the fibers of F by using (C.4) as the definition. \diamond

Let us now explore the resulting calculus. Since the exterior differential commutes with fiber integration by (B.9), we have

$$\begin{aligned} d(\alpha \hat{*} \omega) &= \int_M d(\text{pr}_M^* \alpha \wedge \text{ev}^* \omega) \\ &= \int_M \text{pr}_M^* d\alpha \wedge \text{ev}^* \omega + (-1)^k \int_M \text{pr}_M^* \alpha \wedge \text{ev}^* d\omega \\ &= d\alpha \hat{*} \omega + (-1)^k \alpha \hat{*} d\omega. \end{aligned} \quad (\text{C.5})$$

In particular, if $\alpha = \mu$ is a volume form on M , then, by Remark C.1,

$$d(\mu \hat{*} \omega) = (-1)^{\dim M} \mu \hat{*} d_\pi \omega, \quad (\text{C.6})$$

where d_π denotes the vertical differential along the fibers.

For every bundle automorphism ψ of F , we denote the induced diffeomorphism on the base M by $\check{\psi}$. With this notation, the canonical smooth left action of

$\text{Aut}(F)$ on the section space \mathcal{F} takes the form

$$\Upsilon: \text{Aut}(F) \times \mathcal{F} \rightarrow \mathcal{F}, \quad \psi \cdot \phi = \psi \circ \phi \circ \check{\psi}^{-1}. \quad (\text{C.7})$$

On the infinitesimal level, we obtain

$$Y_\phi^* = T_{\text{id}} \Upsilon_\phi(Y) = \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \psi_\varepsilon \circ \phi \circ \check{\psi}_\varepsilon^{-1} = Y \circ \phi - T\phi(\check{Y}) \in \Gamma^\infty(\phi^*VF), \quad (\text{C.8})$$

where, for $Y \in \text{aut}(F)$, the induced vector field on M is denoted by \check{Y} and the curve $\varepsilon \mapsto \psi_\varepsilon \in \text{Aut}(F)$ is tangent to Y at $\varepsilon = 0$, i.e., $\psi_0 = \text{id}$ and $\left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \psi_\varepsilon = Y$. The next proposition shows how the action of $\text{Aut}(F)$ on \mathcal{F} interacts with the hat product.

PROPOSITION C.2 *Let $\alpha \in \Omega^k(M)$ and $\omega \in \Omega^l(F)$. For every $\psi \in \text{Aut}(F)$, whose induced diffeomorphism $\check{\psi}$ on M is orientation-preserving, and $Y \in \text{aut}(F)$ we have*

$$\Upsilon_\psi^*(\alpha \hat{*} \omega) = (\check{\psi}^* \alpha) \hat{*} (\psi^* \omega), \quad (\text{C.9})$$

$$\mathfrak{L}_{Y^*}(\alpha \hat{*} \omega) = (\mathfrak{L}_{\check{Y}} \alpha) \hat{*} \omega + \alpha \hat{*} (\mathfrak{L}_Y \omega). \quad (\text{C.10})$$

Moreover,

$$Y^* \lrcorner (\alpha \hat{*} \omega) = (\check{Y} \lrcorner \alpha) \hat{*} \omega + (-1)^k \alpha \hat{*} (Y \lrcorner \omega). \quad (\text{C.11})$$

◇

Proof. The following identities clearly hold:

$$\text{pr}_M \circ (\text{id}_M \times \Upsilon_\psi) = \text{pr}_M, \quad (\text{C.12})$$

$$\text{ev} \circ (\text{id}_M \times \Upsilon_\psi) = \psi \circ \text{ev} \circ (\check{\psi}^{-1} \times \text{id}_{\mathcal{F}}), \quad (\text{C.13})$$

$$\text{pr}_M \circ (\check{\psi} \times \text{id}_{\mathcal{F}}) = \check{\psi} \circ \text{pr}_M. \quad (\text{C.14})$$

Using these relations, we calculate

$$\begin{aligned} \Upsilon_\psi^*(\alpha \hat{*} \omega) &\stackrel{(\text{B.5})}{=} \int_M (\text{id}_M \times \Upsilon_\psi)^*(\text{pr}_M^* \alpha \wedge \text{ev}^* \omega) \\ &= \int_M \text{pr}_M^* \alpha \wedge (\psi \circ \text{ev} \circ (\check{\psi}^{-1} \times \text{id}_{\mathcal{F}}))^* \omega \\ &= \int_M (\check{\psi}^{-1} \times \text{id}_{\mathcal{F}})^* ((\check{\psi} \times \text{id}_{\mathcal{F}})^* \text{pr}_M^* \alpha \wedge \text{ev}^*(\psi^* \omega)) \\ &\stackrel{(\text{B.6})}{=} \int_M \text{pr}_M^*(\check{\psi}^* \alpha) \wedge \text{ev}^*(\psi^* \omega) \\ &= (\check{\psi}^* \alpha) \hat{*} (\psi^* \omega). \end{aligned} \quad (\text{C.15})$$

The infinitesimal version (C.10) of (C.9) follows by differentiating using the product law. For the contraction with the Killing vector field, we have

$$\begin{aligned}
Y^* \lrcorner (\alpha \hat{*} \omega) &\stackrel{(B.7)}{=} \int_M (o_M \times Y^*) \lrcorner (\text{pr}_M^* \alpha \wedge \text{ev}^* \omega) \\
&= (-1)^k \int_M \text{pr}_M^* \alpha \wedge ((o_M \times Y^*) \lrcorner \text{ev}^* \omega) \\
&= (-1)^k \int_M \text{pr}_M^* \alpha \wedge \text{ev}^*(Y \lrcorner \omega) \\
&\quad + (-1)^k \int_M \text{pr}_M^* \alpha \wedge ((-\check{Y} \times o_{\mathcal{F}}) \lrcorner \text{ev}^* \omega) \\
&= (-1)^k \alpha \hat{*} (Y \lrcorner \omega) - \int_M (\check{Y} \times o_{\mathcal{F}}) \lrcorner (\text{pr}_M^* \alpha \wedge \text{ev}^* \omega) \\
&\quad + \int_M ((\check{Y} \times o_{\mathcal{F}}) \lrcorner \text{pr}_M^* \alpha) \wedge \text{ev}^* \omega \\
&\stackrel{(B.8)}{=} (-1)^k \alpha \hat{*} (Y \lrcorner \omega) + 0 + \int_M \text{pr}_M^*(\check{Y} \lrcorner \alpha) \wedge \text{ev}^* \omega \\
&= (\check{Y} \lrcorner \alpha) \hat{*} \omega + (-1)^k \alpha \hat{*} (Y \lrcorner \omega).
\end{aligned} \tag{C.16}$$

Here, the general identity $Y \lrcorner f^* \beta = f^*(TY \lrcorner \beta \circ f)$ for the partial pull-back, see (C.3), was utilized in the form

$$(o_M \times Y^*) \lrcorner \text{pr}_M^* \alpha = \text{pr}_M^*(\text{Tpr}_M(o_M \times Y^*) \lrcorner \alpha \circ \text{pr}) = 0, \tag{C.17}$$

$$\begin{aligned}
(o_M \times Y^*) \lrcorner \text{ev}^* \omega &= \text{ev}^*(\text{Tev}(o_M \times Y^*) \lrcorner \omega \circ \text{ev}) \\
&= \text{ev}^*(Y \lrcorner \omega) - \text{ev}^*(\text{Tev}(\check{Y} \times o_{\mathcal{F}}) \lrcorner \omega \circ \text{ev}) \\
&= \text{ev}^*(Y \lrcorner \omega) - (\check{Y} \times o_{\mathcal{F}}) \lrcorner \text{ev}^* \omega,
\end{aligned} \tag{C.18}$$

$$(\check{Y} \times o_{\mathcal{F}}) \lrcorner \text{pr}_M^* \alpha = \text{pr}_M^*(\text{Tpr}_M(\check{Y} \times o_{\mathcal{F}}) \lrcorner \alpha) = \text{pr}_M^*(\check{Y} \lrcorner \alpha), \tag{C.19}$$

where, for the second identity, we used the infinitesimal version of (C.13). \square

D Cheeger-Simons differential characters

Cheeger–Simons differential characters model connections on higher circle n -bundles. As such, they generalize circle bundles to “bundles” whose curvature is not a 2-form but a n -form. Differential characters were introduced by Cheeger and Simons [CS85]. The idea is to adopt and generalize the representation

of principal circle bundles with connection via their holonomy map. First we summarize the main constructions and operations for differential characters and then we introduce the hat product of differential characters. All details and proofs can be found in [BB14] if no other references are given. The only original work in this section are Proposition D.3 and Appendix D.3 where we calculate the derivative of the pull-back map and where we generalize the hat product to differential characters, respectively.

D.1 Motivation: circle bundles

Let us first recall some basic facts about ordinary principal $U(1)$ -bundles in a language which will motivate their generalization to higher bundles.

We will denote the set of equivalence classes of principal $U(1)$ -bundles with connections over M by $\hat{H}^2(M, U(1))$. The reason for this notation will become clear from the considerations below. We have four canonical maps:

CURVATURE The curvature $\text{curv } \Gamma$ of the connection Γ is a 2-form on M . By the Bianchi identity, $\text{curv } \Gamma$ is closed. Furthermore, the periods of the curvature are integral (after dividing by 2π). Thus, in summary, we get a curvature map to the space of closed 2-forms with integral periods

$$\text{curv}: \hat{H}^2(M, U(1)) \rightarrow \Omega_{\text{cl}, \mathbb{Z}}^2(M, \mathbb{R}). \quad (\text{D.1})$$

FLAT BUNDLE The holonomy of a flat connection yields a homomorphism $\text{Hol}: \pi_1(M) \rightarrow U(1)$ from the fundamental group to the circle group. Since the target is an Abelian group, the commutator subgroup $[\pi_1(M), \pi_1(M)]$ lies in the kernel of Hol . On the other hand, the Abelianization

$$\pi_1(M)/[\pi_1(M), \pi_1(M)] \text{ is isomorphic to } H_1(M, \mathbb{Z})$$

by the Hurewicz theorem. The universal coefficient theorem now gives the isomorphism $\text{Hom}(H_1(M, \mathbb{Z}), U(1)) = H^1(M, U(1))$ since $U(1)$ is divisible. Thus, flat line bundles are represented by $H^1(M, U(1))$ and we get a natural injection

$$j: H^1(M, U(1)) \simeq \text{Hom}(\pi_1(M)/[\pi_1(M), \pi_1(M)], U(1)) \rightarrow \hat{H}^2(M, U(1)). \quad (\text{D.2})$$

CHARACTERISTIC CLASS Assigning the Chern class to a bundle furnishes a map

$$c: \hat{H}^2(M, U(1)) \rightarrow H^2(M, \mathbb{Z}). \quad (\text{D.3})$$

TRIVIAL BUNDLE Connections on the trivial bundle $M \times U(1)$ are canonically identified with real valued 1-forms. So, we can view elements of $\Omega^1(M, \mathbb{R})$ as trivial bundles with connection, and thus get a map $\iota: \Omega^1(M, \mathbb{R}) \rightarrow \hat{H}^2(M, U(1))$.

The kernel of ι consists of all $\alpha \in \Omega^1(M, \mathbb{R})$ which are gauge-equivalent to 0, i.e., there exists a gauge transformation $\phi: M \rightarrow U(1)$ such that $\alpha = \delta\phi$. For connected M , this is exactly the case if α is closed and integral. Thus we get an injection

$$\iota: \Omega^1(M, \mathbb{R}) / \Omega_{\text{cl}, \mathbb{Z}}^1(M, \mathbb{R}) \rightarrow \hat{H}^2(M, U(1)). \quad (\text{D.4})$$

D.2 Differential characters

Let M be smooth manifold. In the following, we assume that all singular simplices are smooth, i.e., we work with smooth singular (co-)homology. Let $Z_{k-1}(M, \mathbb{Z})$ denote the group of smooth integral-valued singular cycles of degree $k-1$.

DEFINITION D.1 ([BB14, Section 5.1]) A group homomorphism $h: Z_{k-1}(M, \mathbb{Z}) \rightarrow U(1)$ is called a *differential character* if there exists a differential form $\text{curv } h \in \Omega^k(M, \mathbb{R})$ satisfying

$$h(\partial c) = \exp\left(2\pi i \int_c \text{curv } h\right) \quad (\text{D.5})$$

for every smooth singular k -chain c . The k -th Cheeger–Simons differential cohomology group $\hat{H}^k(M, U(1))$ is the subgroup of $\text{Hom}(Z_{k-1}(M, \mathbb{Z}), U(1))$ consisting of differential characters. \diamond

Cheeger–Simons differential characters provide a model for differential cohomology theory [BB14, Definition 5.9] in the sense that we get a functor $\hat{H}^*(\cdot, U(1))$ from the category of smooth manifolds to the category of \mathbb{Z} -graded Abelian groups, together with four natural transformations

- curvature $\text{curv}: \hat{H}^*(\cdot, U(1)) \rightarrow \Omega_{\text{cl}, \mathbb{Z}}^*(\cdot, \mathbb{R})$
- inclusion of flat classes $j: H^{*-1}(\cdot, U(1)) \rightarrow \hat{H}^*(\cdot, U(1))$
- characteristic class $c: \hat{H}^*(\cdot, U(1)) \rightarrow H^*(\cdot, \mathbb{Z})$
- topological trivialization $\iota: \Omega^{*-1}(\cdot, \mathbb{R}) / \Omega_{\text{cl}, \mathbb{Z}}^{*-1}(\cdot, \mathbb{R}) \rightarrow \hat{H}^*(\cdot, U(1))$

These transformations generalize the maps discussed in Appendix D.1. Moreover, for every smooth manifold M and $k \in \mathbb{Z}$ the following diagram commutes and

all rows and columns are exact:

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \text{Jac}^{k-1}(M) & \longrightarrow & \frac{\Omega^{k-1}(M, \mathbb{R})}{\Omega_{\text{cl}, \mathbb{Z}}^{k-1}(M, \mathbb{R})} & \longrightarrow & d\Omega^{k-1}(M, \mathbb{R}) \longrightarrow 0 \\
& & \downarrow & & \downarrow \iota & & \downarrow \\
0 & \longrightarrow & H^{k-1}(M, U(1)) & \xrightarrow{j} & \hat{H}^k(M, U(1)) & \xrightarrow{\text{curv}} & \Omega_{\text{cl}, \mathbb{Z}}^k(M, \mathbb{R}) \longrightarrow 0 \\
& & \downarrow & & \downarrow c & & \downarrow \\
0 & \longrightarrow & \text{Ext}_{\text{Ab}}(H_{k-1}(M, \mathbb{Z}), \mathbb{Z}) & \longrightarrow & H^k(M, \mathbb{Z}) & \longrightarrow & \text{Hom}(H_k(M, \mathbb{Z}), \mathbb{Z}) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & 0 & & 0 & & 0
\end{array} \tag{D.6}$$

Here $\text{Jac}^l(M)$ denotes the Jacobian torus $H^l(M, \mathbb{R})/H^l(M, \mathbb{Z})$, and the Abelian group $\text{Ext}_{\text{Ab}}(H_{k-1}(M, \mathbb{Z}), \mathbb{Z})$ of equivalence classes of \mathbb{Z} -extensions of $H_{k-1}(M, \mathbb{Z})$ is canonically identified with the torsion subgroup of $H^k(M, \mathbb{Z})$. The horizontal sequence in the middle captures information mainly about the connection. For ordinary principal $U(1)$ -bundle it states that every closed integer 2-form can be realized as the curvature of a connection and that flat bundles are parametrized by their holonomy on generators of $\pi_1(M)$. The topological information is encoded in the vertical sequences. The surjective map $c: \hat{H}^k(M, U(1)) \rightarrow H^k(M, \mathbb{Z})$ shows that differential cohomology refines integer-valued cohomology.

Let us finally comment on the group structure for the case of ordinary principal $U(1)$ -bundles. As far as we know, the group structure was first observed by Kobayashi [Kob56]. The addition of two principal $U(1)$ -bundles P and \tilde{P} over M is defined as follows. First, consider the fiber product $P \times_M \tilde{P}$ and then identify the points which differ by an $U(1)$ -action, i.e., we set

$$P + \tilde{P} := P \times_M \tilde{P} \quad / \quad (p, \tilde{p}) \sim (p \cdot z, \tilde{p} \cdot z^{-1}) \tag{D.7}$$

The so-defined object $P + \tilde{P}$ is indeed a smooth principal $U(1)$ -bundle, where the $U(1)$ -action is the translation in the first factor. The trivial bundle constitutes the identity element, i.e., $P + (M \times U(1))$ is isomorphic to P . For a given principal bundle P , we let $-P$ denote the $U(1)$ bundle that has the same underlying bundle structure as P but carries the modified $U(1)$ -action

$$p * z := p \cdot z^{-1}, \tag{D.8}$$

where on the right side $U(1)$ -acts as in P . Then $P + (-P)$ is isomorphic to the trivial bundle. We leave it to the reader to verify that the addition of bundles is

commutative and associative. Connections A and \tilde{A} on the bundles combine to a connection $\text{pr}_1^*A + \text{pr}_2^*\tilde{A}$ on $P \times_M \tilde{P}$, which descends to a connection $A + \tilde{A}$ on $P + \tilde{P}$. The curvature of $A + \tilde{A}$ is the sum of the corresponding curvatures.

Coming back to the general case, $\hat{H}^k(M, U(1))$ is not just a group but even a Lie group.

PROPOSITION D.2 (Lie group structure [BSS17, Section A]) *Let M be a compact manifold. For every $k \geq 1$, the group $\hat{H}^k(M, U(1))$ is an infinite-dimensional Fréchet Lie group diffeomorphic to*

$$H^k(M, \mathbb{Z}) \times \left(\Omega^{k-1}(M, \mathbb{R}) / \Omega_{\text{cl}, \mathbb{Z}}^{k-1}(M, \mathbb{R}) \right). \quad (\text{D.9})$$

Its Lie algebra $\hat{\mathfrak{h}}^k(M, U(1))$ is isomorphic to the Abelian Fréchet Lie algebra

$$\Omega^{k-1}(M, \mathbb{R}) / \text{d}\Omega^{k-2}(M, \mathbb{R}). \quad (\text{D.10})$$

With respect to this differentiable structure, all natural maps curv , j , c , ι are smooth and hence the defining diagram (D.6) is a commutative diagram of Lie groups. \diamond

Let us now describe the different operations that constitute the calculus of differential cohomology.

PULL-BACK A smooth map $f: M \rightarrow N$ induces a pull-back operation

$$f^*: \hat{H}^k(N, U(1)) \rightarrow \hat{H}^k(M, U(1)) \quad (\text{D.11})$$

on the level of differential characters. Here f^*h is defined by $f^*h = h \circ f_*$, i.e., as the dual of the induced map $f_*: Z_{k-1}(M, \mathbb{Z}) \rightarrow Z_{k-1}(N, \mathbb{Z})$ on cycles. Pull-back commutes with the curvature and the characteristic map.

PROPOSITION D.3 (Smoothness of the pull-back operation) *Let $h \in \hat{H}^k(N, U(1))$ be a differential character on the finite-dimensional manifold N . For every closed manifold M , the pull-back map*

$$\text{pb}_h: C^\infty(M, N) \rightarrow \hat{H}^k(M, U(1)), \quad \phi \mapsto \phi^*h \quad (\text{D.12})$$

is smooth with logarithmic derivative

$$\delta_\phi \text{pb}_h(X) = [\phi^*(X \lrcorner \text{curv } h \circ \phi)] \in \Omega^{k-1}(M, \mathbb{R}) / \text{d}\Omega^{k-2}(M, \mathbb{R}) \quad (\text{D.13})$$

for all $X \in T_\phi C^\infty(M, N)$. \diamond

Proof. Let $\phi_\varepsilon: M \rightarrow N$ be a smooth family of maps whose derivative at $\varepsilon = 0$ is the

tangent vector $X \in T_\phi C^\infty(M, N)$. The homotopy formula [BB14, Example 7.19]

$$\phi_\varepsilon^* h - \phi^* h = (-1)^{k-1} \int_{[0,1]} \tilde{\phi}_\varepsilon^* \text{curv } h \quad (\text{D.14})$$

reduces the problem to the case of differential forms. Here, $\tilde{\phi}_\varepsilon: M \times [0, 1] \rightarrow N$ denotes the smooth homotopy between ϕ and ϕ_ε (for a fixed ε). The following calculation using Lemma D.4 below, proves the identity (D.13):

$$\begin{aligned} \frac{d}{d\varepsilon} \Big|_0 \int_{[0,1]} \tilde{\phi}_\varepsilon^* \text{curv } h &= \int_{[0,1]} \frac{d}{d\varepsilon} \Big|_0 \tilde{\phi}_\varepsilon^* \text{curv } h \\ &\stackrel{(\text{D.20})}{=} \int_{[0,1]} d \tilde{\phi}_0^* \left(\frac{d}{d\varepsilon} \Big|_0 \tilde{\phi}_\varepsilon \lrcorner \text{curv } h \circ \tilde{\phi}_0 \right) \\ &= \int_{[0,1]} d \left(\text{pr}_M^* (\phi^* (X \lrcorner \text{curv } h \circ \phi)) \wedge \text{pr}_{[0,1]} \right) \\ &\stackrel{(\text{B.10})}{=} \int_{[0,1]} \text{pr}_M^* (\phi^* (X \lrcorner \text{curv } h \circ \phi)) \wedge \text{pr}_{[0,1]} \\ &\quad + (-1)^{k-1} \int_{\partial[0,1]} \text{pr}_M^* (\phi^* (X \lrcorner \text{curv } h \circ \phi)) \wedge \text{pr}_{[0,1]} \\ &\stackrel{(\text{B.11})}{=} 0 + (-1)^{k-1} \phi^* (X \lrcorner \text{curv } h \circ \phi) \wedge \int_{\partial[0,1]} \text{pr}_{[0,1]} \\ &= (-1)^{k-1} \phi^* (X \lrcorner \text{curv } h \circ \phi), \end{aligned} \quad (\text{D.15})$$

where we used the notation of the partial pull-back (C.3). \square

Alternative proof. In the case $k = 2$ (where $\hat{H}^2(N, U(1))$ parametrizes principal circle bundles with connection), we can give a more direct and geometric proof. Let $P \rightarrow N$ be principal $U(1)$ -bundle with connection Γ . Let η_ε be a family of smooth closed curves in N representing the tangent vector $Y \in T_\eta C^\infty(S^1, N)$. The derivative of the holonomy map $\text{Hol}: C^\infty(S^1, N) \rightarrow U(1)$ in the direction of Y can be calculated, heuristically, by

$$\frac{d}{d\varepsilon} \Big|_0 \text{Hol}(\eta_\varepsilon) = \frac{d}{d\varepsilon} \Big|_0 \exp \left(\int_{\eta_\varepsilon} \Gamma \right) = \text{Hol}(\eta) + \int_{S^1} (\text{curv } \Gamma)_{\eta(\tau)} (Y(\tau), \dot{\eta}(\tau)) d\tau. \quad (\text{D.16})$$

This calculation can be made rigorous and even generalizes to non-Abelian bundles; see [Die] for details. Now let γ be a closed curve in M and consider a family of smooth maps $\phi_\varepsilon : M \rightarrow N$ representing $X \in T_\phi C^\infty(M, N)$. The above calculation implies

$$\begin{aligned} \frac{d}{d\varepsilon} \Big|_0 \text{pb}_h(\phi_\varepsilon)(\gamma) &= \frac{d}{d\varepsilon} \Big|_0 h(\phi_\varepsilon \circ \gamma) \\ &= h(\phi \circ \gamma) + \int_{S^1} (\text{curv } h)_{\phi \circ \gamma(\tau)}(X_{\gamma(\tau)}, T\phi(\dot{\gamma}(\tau))) d\tau. \end{aligned} \quad (\text{D.17})$$

Thus, the derivative of the pull-back map is

$$\delta_\phi \text{pb}_h(X)(\gamma) = \int_\gamma \phi^*(X \lrcorner \text{curv } h \circ \phi). \quad (\text{D.18})$$

□

LEMMA D.4 *Let α be a differential k -form on the finite-dimensional manifold N . For every closed finite-dimensional manifold M , the pull-back map*

$$\text{pb}_\alpha : C^\infty(M, N) \rightarrow \Omega^k(M), \quad \phi \mapsto \phi^* \alpha \quad (\text{D.19})$$

is smooth and has derivative

$$T_\phi \text{pb}_\alpha(X) = \mathfrak{L}_X^\phi \alpha = d(\phi^*(X \lrcorner \alpha \circ \phi)) + \phi^*(X \lrcorner (d\alpha \circ \phi)) \quad (\text{D.20})$$

for $X \in T_\phi C^\infty(M, N)$. Here $\mathfrak{L}_X^\phi \alpha := \frac{d}{d\varepsilon} \Big|_0 \phi_\varepsilon^* \alpha$ denotes the Lie derivative along the map ϕ , so that the second equality in (D.20) generalizes Cartan's formula. Moreover, we used the notation of the partial pull-back (C.3). \diamond

Proof. Let $\phi_\varepsilon : M \rightarrow N$ be a family of maps representing the tangent vector $X \in T_\phi C^\infty(M, N)$. We have $T_\phi \text{pb}_\alpha(X) = \frac{d}{d\varepsilon} \Big|_0 \phi_\varepsilon^* \alpha = \mathfrak{L}_X^\phi \alpha$ by definition. For the generalized Cartan formula, we need to check the identity only for functions and for exact 1-forms, because these generate the exterior algebra. On functions $\alpha = f \in C^\infty(N)$, we get

$$(\mathfrak{L}_X^\phi f)_m = \frac{d}{d\varepsilon} \Big|_0 (\phi_\varepsilon^* f)(m) = T_{\phi(m)} f(X_m) = (\phi^*(X \lrcorner df \circ \phi))_m. \quad (\text{D.21})$$

Similarly, for exact one forms $\alpha = df$, we find

$$\mathfrak{L}_X^\phi(df) = \frac{d}{d\varepsilon} \Big|_0 \phi_\varepsilon^* df = d \frac{d}{d\varepsilon} \Big|_0 \phi_\varepsilon^* f = d(\phi^*(X \lrcorner df \circ \phi)). \quad (\text{D.22})$$

This completes the proof. \square

PRODUCT There exists an associative and \mathbb{Z} -bilinear product of differential characters denoted by

$$\star: \hat{H}^k(M, \mathbb{U}(1)) \times \hat{H}^l(M, \mathbb{U}(1)) \rightarrow \hat{H}^{k+l}(M, \mathbb{U}(1)). \quad (\text{D.23})$$

It satisfies the following properties [BB14, Definition 6.1]:

- (i) (Ring structure) The product \star is associative and bilinear.
- (ii) (Graded commutativity) $h \star g = (-1)^{kl} g \star h$.
- (iii) (Naturality under pull-back) For a smooth map $f: N \rightarrow M$, we have $f^*(h \star g) = (f^*h) \star (f^*g)$.
- (iv) (Compatibility with curvature) $\text{curv}(h \star g) = \text{curv } h \wedge \text{curv } g$.
- (v) (Compatibility with characteristic class) $c(h \star g) = c(h) \cup c(g)$.
- (vi) (Compatibility with trivializations) $\iota(\alpha) \star h = \iota(\alpha \wedge \text{curv } h)$.

As shown in [BB14], these properties completely characterize the product (D.23); we refer to this book for the lengthy proof of this statement. In this paper, we use only these properties without appealing to the concrete construction of \star .

FIBER INTEGRATION Let $\pi: F \rightarrow M$ be a smooth fiber bundle whose typical fiber \underline{F} is a closed, finite-dimensional oriented manifold. Then fiber integration [BB14, Definition 7.1] yields a group homomorphism

$$\int: \hat{H}^k(F, \mathbb{U}(1)) \rightarrow \hat{H}^{k-\dim \underline{F}}(M, \mathbb{U}(1)) \quad (\text{D.24})$$

such that the curvature map intertwines fiber integration of differential characters with ordinary fiber integration of differential forms $f: \Omega^k(F) \rightarrow \Omega^{k-\dim \underline{F}}(M)$, i.e., the following diagram commutes:

$$\begin{array}{ccc} \hat{H}^k(F, \mathbb{U}(1)) & \xrightarrow{f} & \hat{H}^{k-\dim \underline{F}}(M, \mathbb{U}(1)) \\ \downarrow \text{curv} & & \downarrow \text{curv} \\ \Omega^k(F) & \xrightarrow{f} & \Omega^{k-\dim \underline{F}}(M). \end{array} \quad (\text{D.25})$$

D.3 Hat product of differential characters

Vizman [Viz11] introduced the hat product of differential forms as a means to construct differential forms on the mapping space $C^\infty(M, F)$ from differential forms on the finite-dimensional manifolds M and F . We now generalize these

ideas to the level of differential characters. A similar construction has been used also in [Die+19]. The hat product of differential characters yields a method that combines higher bundles on M and F to bundles on the mapping space. In the process, we also generalize the procedure of higher dimensional transgression [BB14, Chapter 9.2].

As in Appendix C, we consider the more general setup of sections of a fiber bundle. Let M be a compact, oriented, finite-dimensional manifold without boundary, and let $\pi: F \rightarrow M$ be a finite-dimensional fiber bundle over M . Recall from [Viz11] that the hat product of differential forms used the evaluation and projection map to construct differential forms on $M \times \mathcal{F}$ and then integrate out the M -factor. We use the same strategy to define the hat product of differential characters. The evaluation map $\text{ev}: M \times \mathcal{F} \rightarrow F$ yields a pull-back map of differential characters $\text{ev}^*: \hat{H}^l(F, \mathbb{U}(1)) \rightarrow \hat{H}^l(M \times \mathcal{F}, \mathbb{U}(1))$. Similarly, the projection on the first factor induces a homomorphism $\text{pr}_M^*: \hat{H}^k(M, \mathbb{U}(1)) \rightarrow \hat{H}^k(M \times \mathcal{F})$. Taking the product of these differential forms yields an element of $\hat{H}^{k+l}(M \times \mathcal{F})$. Finally, we integrate over M by viewing $M \times \mathcal{F}$ as a trivial bundle over \mathcal{F} . Summarizing, we have constructed a map

$$\begin{aligned} \hat{*}: \hat{H}^k(M, \mathbb{U}(1)) \times \hat{H}^l(F, \mathbb{U}(1)) &\rightarrow \hat{H}^{k+l-\dim M}(\mathcal{F}, \mathbb{U}(1)), \\ h \hat{*} g &:= \int_M \text{pr}_M^* h \star \text{ev}^* g. \end{aligned} \quad (\text{D.26})$$

Using the compatibility with the curvature map, we calculate

$$\text{curv}(h \hat{*} g) = \int_M \text{pr}_M^* \text{curv } h \wedge \text{ev}^* \text{curv } g = \text{curv } h \hat{*} \text{curv } g, \quad (\text{D.27})$$

that is, the curvature of the hat product of differential characters equals the hat product of the corresponding curvatures. Let us now discuss a few special cases.

PREQUANTIZATION Fix a differential character μ on M whose degree coincides with the dimension of M . Then we define a μ -dependent map by

$$\hat{H}^l(F, \mathbb{U}(1)) \rightarrow \hat{H}^l(\mathcal{F}, \mathbb{U}(1)), \quad h \mapsto \mu \hat{*} h. \quad (\text{D.28})$$

In particular, for $l = 2$, principal $\mathbb{U}(1)$ -bundles on F yield principal $\mathbb{U}(1)$ -bundles on the section space \mathcal{F} . By (D.27), the differential character $\mu \hat{*} h$ is a prequantization of $(\mathcal{F}, \text{curv } \mu \hat{*} \text{curv } g)$.

TRANSGRESSION The *transgression map* [BB14, Definition 9.1] uses the hat product

with the identity element $e \in \hat{H}^0(M, U(1))$:

$$\hat{H}^l(F, U(1)) \rightarrow \hat{H}^{l-\dim M}(\mathcal{F}, U(1)), \quad h \mapsto e \hat{*} h = \int_M \text{ev}^* h. \quad (\text{D.29})$$

For the special case of a trivial fiber bundle $F = \underline{F} \times M$ over $M = S^1$, the space of smooth sections is identified with the space $\mathcal{L}^\infty(\underline{F})$ of smooth loops in \underline{F} , and transgression yields a map $\hat{H}^l(\underline{F}, U(1)) \rightarrow \hat{H}^{l-1}(\mathcal{L}^\infty(\underline{F}), U(1))$. In degree $l = 2$, it assigns to a principal $U(1)$ -bundle over \underline{F} its holonomy map $\mathcal{L}^\infty(\underline{F}) \rightarrow U(1)$.

REFERENCES

- [AMM98] A. Alekseev, A. Malkin, and E. Meinrenken. “Lie group valued moment maps”. *J. Differential Geom.* 48.3 (1998), pp. 445–495. arXiv: dg-ga/9707021 [dg-ga].
- [Ale+98] D. Alekseevsky, J. Grabowski, G. Marmo, and P. W. Michor. “Poisson structures on double Lie groups”. *J. Geom. Phys.* 26.3-4 (1998), pp. 340–379.
- [AK98] V. I. Arnold and B. A. Khesin. *Topological Methods in Hydrodynamics*. Springer, 1998.
- [AB83] M. F. Atiyah and R. Bott. “The Yang-Mills equations over Riemann surfaces”. *Philos. Trans. Roy. Soc. London Ser. A* 308.1505 (1983), pp. 523–615.
- [Ban97] A. Banyaga. *The Structure of Classical Diffeomorphism Groups*. Vol. 400. Mathematics and Its Applications. Springer, 1997.
- [BB14] C. Bär and C. Becker. *Differential Characters*. Vol. 2112. Lecture Notes in Mathematics. Springer, 2014, pp. 1–90. arXiv: 1303.6457 [math.DG].
- [BW97] S. Bates and A. Weinstein. *Lectures on the Geometry of Quantization*. Vol. 8. Berkeley Mathematical Lecture Notes. American Mathematical Society, 1997.
- [BSS17] C. Becker, A. Schenkel, and R. J. Szabo. “Differential cohomology and locally covariant quantum field theory”. *Rev. Math. Phys.* 29.1 (2017). arXiv: 1406.1514 [hep-th].
- [BGT18] D. Beltiță, T. Goliński, and A.-B. Tumpach. “Queer Poisson brackets”. *Journal of Geometry and Physics* 132 (2018), pp. 358–362. arXiv: 1710.03057.
- [Bry07] J.-L. Brylinski. *Loop Spaces, Characteristic Classes and Geometric Quantization*. Birkhäuser Boston, 2007.

- [CS85] J. Cheeger and J. Simons. “Differential characters and geometric invariants”. In: *Geometry and Topology*. Vol. 1167. Lecture Notes in Mathematics. Springer, Berlin, 1985, pp. 50–80.
- [CGS00] K. Cieliebak, A. R. Gaio, and D. A. Salamon. “J-holomorphic curves, moment maps, and invariants of Hamiltonian group actions”. *Internat. Math. Res. Notices* 16 (2000), pp. 831–882. arXiv: math/9909122v1 [math.SG].
- [CM85] R. Cirelli and A. Manià. “The group of gauge transformations as a Schwartz–Lie group”. *J. Math. Phys.* 26.12 (1985), pp. 3036–3041.
- [Cle57] A. Clebsch. “Über eine allgemeine Transformation der hydrodynamischen Gleichungen”. *J. Reine Angew. Math.* (1857), pp. 293–312.
- [Cle59] A. Clebsch. “Ueber die Integration der hydrodynamischen Gleichungen”. *J. Reine Angew. Math.* 56 (1859), pp. 1–10.
- [CDM88] M. Condevaux, P. Dazord, and P. Molino. “Géométrie du moment”. In: *Travaux du Séminaire Sud-Rhodanien de Géométrie, I*. Vol. 88. Publ. Dép. Math. 1B. Université Claude Bernard - Lyon 1, 1988, pp. 131–160.
- [Die19] T. Diez. “Singular symplectic reduction for Fréchet group actions and the Cauchy problem for the Yang–Mills equation”. PhD thesis. Universität Leipzig, 2019. arXiv: 1909.00744 [math.SG].
- [Die] T. Diez. “Smoothness of the holonomy map”. In preparation.
- [Die+19] T. Diez, B. Janssens, K.-H. Neeb, and C. Vizman. “Central extensions of Lie groups preserving a differential form” (2019). arXiv: 1906.03234 [math.DG]. Submitted.
- [Don87] S. K. Donaldson. “Infinite determinants, stable bundles and curvature”. *Duke Math. J.* 54.1 (1987), pp. 231–247.
- [Don97] S. K. Donaldson. “Remarks on gauge theory, complex geometry and 4-manifold topology”. In: *Fields Medallists’ Lectures*. Ed. by M. Atiyah and D. Iagolnitzer. Vol. 5. World Sci. Ser. 20th Century Math. 5. World Sci. Publ., River Edge, NJ, 1997, pp. 384–403.
- [Don00] S. K. Donaldson. “Moment maps and diffeomorphisms”. In: *Surveys in differential geometry*. Vol. 3. Surv. Differ. Geom. 1. Int. Press, Somerville, MA, 2000, pp. 1–16.
- [Don01] S. K. Donaldson. “Scalar curvature and projective embeddings, I”. *J. Differential Geom.* 59.3 (2001), pp. 479–522.
- [Don03] S. K. Donaldson. “Moment maps in differential geometry”. In: *Surveys in differential geometry*. Vol. 8. Surv. Differ. Geom. 2003, pp. 171–189.

- [Don15] S. Donaldson. “Kähler-Einstein metrics and algebraic geometry”. In: *Current developments in mathematics*. Vol. 2015. 1. International Press of Boston, 2015, pp. 1–25. arXiv: 1702.05748v2 [math.DG].
- [Dri93] V. G. Drinfeld. “On Poisson homogeneous spaces of Poisson-Lie groups”. *Teoret. Mat. Fiz.* 95.2 (1993), pp. 226–227.
- [EP12] A. Enciso and D. Peralta-Salas. “Knots and links in steady solutions of the Euler equation”. *Annals of Mathematics* 175.1 (2012), pp. 345–367. arXiv: 1003.3122 [math-ph].
- [ES02] P. Etingof and O. Schiffmann. *Lectures on Quantum Groups*. 2nd ed. Lectures in Mathematical Physics. International Press, 2002.
- [FR96] H. Flaschka and T. Ratiu. “A convexity theorem for Poisson actions of compact Lie groups”. *Ann. Sci. Éc. Norm. Supér. (4)* 29.6 (1996), pp. 787–809.
- [FMS07] D. S. Freed, G. W. Moore, and G. Segal. “The uncertainty of fluxes”. *Commun. Math. Phys.* 271.1 (2007), pp. 247–274. arXiv: hep-th/0605198v3 [hep-th].
- [Fuj92] A. Fujiki. “Moduli space of polarized algebraic manifolds and Kähler metrics”. *Sugaku Expositions* 5.2 (1992), pp. 173–191.
- [GV12] F. Gay-Balmaz and C. Vizman. “Dual pairs in fluid dynamics”. *Ann. Global Anal. Geom.* 41.1 (2012), pp. 1–24. arXiv: 1007.1347 [math.SG].
- [GV19] F. Gay-Balmaz and C. Vizman. “Isotropic submanifolds and coadjoint orbits of the Hamiltonian group”. *Journ. Symplectic. Geom.* 17.3 (2019), pp. 663–702. arXiv: 1606.07994v2 [math.SG].
- [Gom03] K. Gomi. “Connections and curvings on lifting bundle gerbes”. *J. Lond. Math. Soc.* 67.2 (2003), pp. 510–526.
- [GH00] C. R. Graham and F. S. Henyey. “Clebsch representation near points where the vorticity vanishes”. *Phys. Fluids* 12.4 (2000), pp. 744–746.
- [GHV72] W. Greub, S. Halperin, and R. Vanstone. *Connections, curvature, and cohomology. De Rham Cohomology of Manifolds and Vector Bundles*. Vol. 1. Academic Press, 1972.
- [Ham82] R. S. Hamilton. “The inverse function theorem of Nash and Moser”. *Bull. Amer. Math. Soc* 7 (1982), pp. 65–223.
- [HJ94] J. Huebschmann and L. C. Jeffrey. “Group cohomology construction of symplectic forms on certain moduli spaces”. *Internat. Math. Res. Notices* 1994.6 (1994), pp. 245–249.
- [Ked+16] H. Kedia, D. Foster, M. R. Dennis, and W. T. M. Irvine. “Weaving knotted vector fields with tunable helicity”. *Phys. Rev. Lett.* 117.27 (2016). arXiv: 1610.02463v1 [math-ph].

- [Kob56] S. Kobayashi. “Principal fibre bundles with the 1-dimensional toroidal group”. *Tohoku Math. J.* 2nd ser. 8.1 (1956), pp. 29–45.
- [KN63] S. Kobayashi and K. Nomizu. *Foundations of Differential Geometry*. Vol. 1. Wiley, 1963.
- [KMS93] I. Kolář, P. W. Michor, and J. Slovák. *Natural Operations in Differential Geometry*. Springer-Verlag, Berlin, 1993.
- [Kos70] B. Kostant. “Quantization and unitary representations”. In: *Lectures in Modern Analysis and Applications III*. Vol. 170. Lecture Notes in Mathematics. Springer Berlin / Heidelberg, 1970, pp. 87–208.
- [KM97] A. Kriegl and P. W. Michor. *The Convenient Setting of Global Analysis*. Vol. 53. Mathematical Surveys and Monographs. American Mathematical Society, 1997.
- [LR12] X. Liu and R. L. Ricca. “The Jones polynomial for fluid knots from helicity”. *J. Phys. A* 45.20 (2012), pp. 205501, 14.
- [Lu90] J.-H. Lu. “Multiplicative and affine Poisson structures on Lie groups”. PhD thesis. University of California, Berkeley, 1990.
- [LR91] J.-H. Lu and T. Ratiu. “On the nonlinear convexity theorem of Kostant”. *Journal of the American Mathematical Society* 4.2 (1991), pp. 349–349.
- [LW90] J.-H. Lu and A. Weinstein. “Poisson Lie groups, dressing transformations, and Bruhat decompositions”. *J. Differential Geom.* 31.2 (1990), pp. 501–526.
- [Maj90] S. Majid. “Matched pairs of Lie groups associated to solutions of the Yang-Baxter equations”. *Pacific Journal of Mathematics* 141.2 (1990), pp. 311–332.
- [Mar+07] J. E. Marsden, G. Misiolek, J.-P. Ortega, M. Perlmutter, and T. S. Ratiu. *Hamiltonian Reduction by Stages*. Vol. 1913. Lecture Notes in Mathematics. Springer, 2007.
- [MW83] J. Marsden and A. Weinstein. “Coadjoint orbits, vortices, and Clebsch variables for incompressible fluids”. *Phys. D* 7.1–3 (1983), pp. 305–323.
- [MS99] D. McDuff and D. Salamon. *Introduction to Symplectic Topology*. 2nd ed. Oxford Graduate Texts in Mathematics. Oxford University Press, Oxford, 1999.
- [Mun00] I. Mundet i Riera. “A Hitchin-Kobayashi correspondence for Kähler fibrations”. *J. Reine Angew. Math.* (528 2000), pp. 41–80. arXiv: math/9901076v2 [math.DG].
- [Mur96] M. K. Murray. “Bundle Gerbes”. *J. Lond. Math. Soc.* 54.2 (1996), pp. 403–416. arXiv: dg-ga/9407015.

- [NR61] M. S. Narasimhan and S. Ramanan. “Existence of Universal Connections”. *Amer. J. Math.* 83.3 (1961), pp. 563–572.
- [Nee06] K.-H. Neeb. “Towards a Lie theory of locally convex groups”. *Japan. J. Math.* 1.2 (2006), pp. 291–468.
- [NST14] K.-H. Neeb, H. Sahlmann, and T. Thiemann. “Weak Poisson structures on infinite dimensional manifolds and Hamiltonian actions”. In: *Lie theory and its applications in physics*. Ed. by V. Dobrev. Vol. 111. Springer Proceedings in Mathematics & Statistics. Springer, Tokyo, 2014, pp. 105–135. arXiv: 1402.6818 [math.DG].
- [Ohs15] T. Ohsawa. “The Siegel upper half space is a Marsden-Weinstein quotient: symplectic reduction and Gaussian wave packets”. *Lett. Math. Phys.* 105.9 (2015), pp. 1301–1320. arXiv: 1504.03963 [math-ph].
- [OR03] J.-P. Ortega and T. S. Ratiu. *Momentum Maps and Hamiltonian Reduction*. Birkhäuser Boston, 2003.
- [PR12] Á. Pelayo and T. S. Ratiu. “Circle-valued momentum maps for symplectic periodic flows”. *Enseign. Math.* 2nd ser. 58.1/2 (2012), pp. 205–219.
- [SV19] P. Skerritt and C. Vizman. “Dual pairs for matrix Lie groups”. *J. Geom. Mech.* 11.2 (2019), pp. 255–275. arXiv: 1805.01519 [math].
- [Stro0] T. Strobl. “Gravity in two spacetime dimensions”. Habilitation thesis. 2000. arXiv: hep-th/0011240v1 [hep-th].
- [Tak81] M. Takeuchi. “Matched pairs of groups and bismash products of hopf algebras”. *Communications in Algebra* 9.8 (1981), pp. 841–882.
- [Viz11] C. Vizman. “Induced differential forms on manifolds of functions”. *Arch. Math. (Brno)* 47.3 (2011), pp. 201–215. arXiv: 1111.3889 [math.DG].
- [Wei80] A. Weinstein. “Fat bundles and symplectic manifolds”. *Adv. Math.* 37.3 (1980), pp. 239–250.