

Multivariate normal approximation for traces of random unitary matrices

Kurt Johansson * Gaultier Lambert †

February 6, 2020

Abstract

In this article, we obtain a super-exponential rate of convergence in total variation between the traces of the first m powers of an $n \times n$ random unitary matrices and a $2m$ -dimensional Gaussian random variable. This generalizes previous results in the scalar case to the multivariate setting, and we also give the precise dependence on the dimensions m and n in the estimate with explicit constants. We are especially interested in the regime where m grows with n and our main result basically states that if $m \ll \sqrt{n}$, then the rate of convergence in the Gaussian approximation is $\Gamma(\frac{n}{m} + 1)^{-1}$ times a correction. We also show that the Gaussian approximation remains valid for all $m \ll n^{2/3}$ without a fast rate of convergence.

1 Introduction and main results

1.1 Introduction

Let \mathbf{U} be a random unitary matrix distributed according to the normalized Haar measure \mathbb{P}_n on the unitary group $U(n)$ of size $n \in \mathbb{N}$. In random matrix theory this is known as the circular unitary ensemble or CUE. The joint law of the eigenvalues $(e^{i\theta_1}, \dots, e^{i\theta_n})$ of \mathbf{U} , $i = \sqrt{-1}$, $\theta_j \in [-\pi, \pi]$, under this probability measure has an explicit density given by the Weyl integration formula,

$$\frac{1}{(2\pi)^{nn!}} \prod_{1 \leq k < j \leq n} |e^{i\theta_k} - e^{i\theta_j}|^2 = \frac{\mathfrak{U}_n}{(2\pi)^n} \prod_{1 \leq k < j \leq n} \sin^2 \left(\frac{\theta_k - \theta_j}{2} \right), \quad (1.1)$$

where $\mathfrak{U}_n = 2^{n(n-1)}/n!$. Consider the random variable

$$Z = \sum_{k=1}^m \xi_{2k-1} \sqrt{\frac{2}{k}} \Re \operatorname{Tr} \mathbf{U}^k + \xi_{2k} \sqrt{\frac{2}{k}} \Im \operatorname{Tr} \mathbf{U}^k,$$

where $\xi = (\xi_1, \dots, \xi_{2m}) \in \mathbb{R}^{2m}$, built from the traces of the unitary matrix \mathbf{U} . It is a well-known consequence of the Strong Szegő theorem ((Szegő, 1952) – see Theorem 2.1 below) that for any fixed $m \in \mathbb{N}$, $Z \rightarrow \|\xi\| \mathcal{N}$ weakly as $n \rightarrow \infty$, where \mathcal{N} is a standard Gaussian random variable. This is a surprising result since the trace is the sum of n random variables and there is no normalization in n .

*KTH Royal Institute of Technology, Matematiska institutionen, 100 44 Stockholm, Sweden.

K.J. was supported by the grant KAW 2015.0270 from the Knut and Alice Wallenberg Foundation and the Swedish Research Council grant 2015-04872.

Email: <mailto:kurtj@kth.se>

†University of Zurich, Winterthurerstrasse 190, 8057 Zürich, Switzerland.

G.L. research is supported by the SNSF Ambizione grant S-71114-05-01.

Email: gaultier.lambert@math.uzh.ch

This limit theorem is also a consequence of the striking fact proved by (Diaconis and Shahshahani, 1994) that all joint moments of $\sqrt{\frac{2}{k}}\Re \operatorname{Tr} \mathbf{U}^k$ and $\sqrt{\frac{2}{k}}\Im \operatorname{Tr} \mathbf{U}^k$ up to a certain order are identical to those of independent standard Gaussian random variables (see Theorem 7.1 below). Based on this result, Persi Diaconis (Diaconis, 1994) conjectured that the rate of convergence in total variation norm of Z to a normal random variable should be very fast, even super-exponential. This was proved in (Johansson, 1997), where it was shown that there are positive constants C and δ so that

$$d_{\text{TV}}(Z, \|\xi\|\mathcal{N}) \leq Cn^{-\delta n}, \quad (1.2)$$

where d_{TV} denotes the total variation distance (see (1.7) below for a definition). No explicit expression for C or δ or their dependence on m and the parameters was given.

A related but separate problem is to consider the multivariate convergence of the random variables

$$X_{2k-1} := \sqrt{\frac{2}{k}}\Re \operatorname{Tr} \mathbf{U}^k \quad \text{and} \quad X_{2k} := \sqrt{\frac{2}{k}}\Im \operatorname{Tr} \mathbf{U}^k, \quad (1.3)$$

$1 \leq k \leq m$. We are interested in the law of the random vector $\mathbf{X} = (X_1, \dots, X_{2m})$ when the dimension of the matrix \mathbf{U} is large. Let $\mathbf{G} = (G_1, \dots, G_{2m})$ be i.i.d. standard Gaussian random variables. For a fixed $m \in \mathbb{N}$ it again follows from the Strong Szegő theorem that $\mathbf{X} \rightarrow \mathbf{G}$ weakly as $n \rightarrow \infty$. Peter Sarnak (Sarnak, 2019) raised the following problem in connection with his work with M. Rubinstein on computing zeros of L-functions and under-determined matrix moment problems. How close is \mathbf{X} to \mathbf{G} in total variation distance as a function of m for a given n ? Here, m can depend on n , e.g. be a power of n . Is \mathbf{X} still very close to \mathbf{G} ? This is the main problem investigated in the present paper. Theorem 1.1 and Theorem 1.3 give our results. We get a statement for m almost up to \sqrt{n} . The other classical groups can also be considered, see (Courteaut and Johansson). Since we are mainly interested in the case when m is large we assume that $m \geq 3$ throughout this paper. In the case $m = 1$ it is possible to get a more precise result and this together with results on single traces will be considered for all the classical compact groups in a forthcoming publication, (Courteaut et al.). (A bound for $m = 2$ can be directly inferred from the case $m = 3$; a special treatment of this case would only give a slight improvement.) An important aspect of the present work is that, in contrast to (Johansson, 1997), we keep explicit track of the constants and the dependence on m . We have also made an effort to optimize in the argument and get reasonable numerical constants.

Since $Z = \mathbf{X} \cdot \xi$, as a consequence of our multivariate results we can improve (1.2), for a fixed m and uniformly for all ξ , to

$$d_{\text{TV}}(Z, \|\xi\|\mathcal{N}) = \mathcal{O}\left(\frac{e^{\frac{n}{m}(\log(1+\log m)+\frac{1}{2})}}{\sqrt{n} \Gamma(\frac{n}{m}+1)}\right), \quad (1.4)$$

where the implied constant has an explicit dependence in $m \in \mathbb{N}$. Broadly speaking, we expect that the best possible estimate for the RHS of (1.4) is $\Gamma(\frac{n}{m}+1)^{-1}$ times some sub-exponential corrections. We can also let the degree m grow as $n \rightarrow +\infty$. From Proposition 1.6 we deduce the following estimate:

$$\sup_{m \leq \frac{\sqrt{n}}{6.45(\log n)^{1/4}}} d_{\text{TV}}(Z, \|\xi\|\mathcal{N}) \leq \sqrt{n} \exp(19.4 - 0.83\sqrt{n}(\log n)^{5/4}).$$

uniformly for all ξ when n is large enough.

Using Stein's method and the exact moment identities from (Diaconis and Shahshahani, 1994), one can infer the following rate of convergence in the multivariate problem: for any $m \leq 2n$,

$$W_1(\mathbf{X}, \mathbf{G}) = \mathcal{O}(m^2/n), \quad (1.5)$$

where W_1 denotes the Wasserstein 1 distance between two probability measures on \mathbb{R}^{2m} – see (Döbler and Stolz, 2011, Theorem 3.1). By relying on the recent techniques from (Lambert et al., 2019), we

can improve on (1.5) – see Theorem 1.5 below. See also (Webb, 2016) for an analogous multivariate result that applies to more general circular β -ensembles and Remark 1.1 below. Recently, rates of convergence to the Gaussian law have also been obtained for $\text{Tr } f(\mathbf{M})$ where $f : \mathbb{R} \rightarrow \mathbb{R}$ is a real-analytic function and \mathbf{M} is a random matrix from the Gaussian, Laguerre or Jacobi unitary ensembles by Berezin and Bufetov (2019) using Riemann–Hilbert techniques. In contrast to the CUE, in these cases, the optimal rates of convergence are expected to be polynomial in the dimension of the random matrix.

The fast rate of convergence of \mathbf{X} to \mathbf{G} holds for $m \ll \sqrt{n}$ by Theorem 1.1, but we see from Theorem 1.5 below that we have convergence to the multivariate Gaussian for $m \ll n^{2/3}$. We have no conjecture concerning the threshold $m \in \mathbb{N}$ at which the Gaussian approximation fails. Also, we do not know whether there is some transition when varying m where we go from a fast convergence rate to some other rate of convergence.

1.2 Main results

For any $m \in \mathbb{N}$, we denote by $\Omega_m = \frac{\pi^m}{m!}$ the volume of the unit ball and by $\|x\| = \sqrt{x_1^2 + \dots + x_{2m}^2}$ the Euclidean norm in \mathbb{R}^{2m} . It is straightforward to see that for any $m, n \in \mathbb{N}$, the random vector \mathbf{X} has a density on \mathbb{R}^{2m} that we denote by $\mathcal{P}_{n,m}$. For any $n, m \in \mathbb{N}$ and $k \in \mathbb{N}$, we define

$$\Delta_{n,m}^{(k)} := \left(\int_{\mathbb{R}^{2m}} \left| \mathcal{P}_{n,m}(x) - \frac{e^{-\|x\|^2/2}}{(2\pi)^m} \right|^k dx \right)^{1/k}. \quad (1.6)$$

In this paper, we focus on getting (non-asymptotic) estimates for $\Delta_{n,m}^{(1)}$ and $\Delta_{n,m}^{(2)}$ with explicit constants which hold for large $n \in \mathbb{N}$ when $m \ll \sqrt{n}$. Let us observe that $\Delta_{n,m}^{(1)}$ controls the *total variation* distance between \mathbf{X} and \mathbf{G} (a standard Gaussian random variable on \mathbb{R}^{2m}). Namely, we have

$$d_{\text{TV}}(\mathbf{X}, \mathbf{G}) := \sup_{A \subset \mathbb{R}^{2m}} |\mathbb{P}_n[\mathbf{X} \in A] - \mathbb{P}[\mathbf{G} \in A]| \leq \Delta_{n,m}^{(1)}, \quad (1.7)$$

where the supremum is taken over all Borel subsets $A \subset \mathbb{R}^{2m}$. Our main result, which is a quantitative generalization of the estimates (1.2) from (Johansson, 1997) in a multi-dimensional setting can be summarized as follows.

Theorem 1.1. *For all $n, m \in \mathbb{N}$ such that $n \geq 1911$ and $N = n/m \geq 146.5m\sqrt{1 + \log m}$, we have the following estimate in total variation distance,*

$$d_{\text{TV}}(\mathbf{X}, \mathbf{G}) \leq 16\sqrt{\Omega_m} m^{\frac{5}{2}} 4^m e^{\frac{N}{2} + \frac{m^2}{4N}} \frac{(N\sqrt{\log N})^m (1 + \log m)^N}{\sqrt{N} \Gamma(N + 1)}.$$

We expect that, up to corrections, the factor $\Gamma(\frac{n}{m} + 1)^{-1}$ is actually the correct order for the statistical distance between the random vectors \mathbf{X} and \mathbf{G} as long as $m \ll \sqrt{n}$. To clarify the meaning of this estimate in the regime where m grows with n , let us also give the following consequence when m is like n^α , $\alpha < 1/2$.

Proposition 1.2. *Let $m = \lfloor n^\alpha \rfloor$ with $0 < \alpha < 1/2$, then for all $n \geq n_\alpha$,*

$$d_{\text{TV}}(\mathbf{X}, \mathbf{G}) \leq \frac{18e^{8\pi}}{(2\pi)^{\frac{3}{4}}} n^{3\alpha - \frac{3}{2}} \exp\left(- (1 - \epsilon_n) n^{1-\alpha} \log(n^{1-\alpha})\right),$$

where $n_\alpha := \inf \{n \geq 18^{1/\alpha} : n^{1-2\alpha} \geq 20.4\sqrt{\log n}\}$, $1 - \epsilon_n \geq 87 \cdot 10^{-3}$ and $\epsilon_n \rightarrow 0$ as $n \rightarrow \infty$; see (3.17) for a more precise bound on ϵ_n .

The proof of Theorem 1.1 and Proposition 1.2 are given in Section 3.3. According to (1.7), these results are consequences of the following more precise bounds. We postpone the definition of $\Theta_{N,m}$ to the Appendix B since it is rather involved.

Theorem 1.3. Let $\Theta_{N,m}$ be given by (B.10). For any $n, m \in \mathbb{N}$ such that $m \geq 3$ and $N = n/m > 4m$, we have

$$\Delta_{n,m}^{(2)} \leq 8\sqrt{\Omega_m N^{\frac{m}{2}}} \Theta_{N,m}. \quad (1.8)$$

If we assume that $\Delta_{n,m}^{(2)} \leq 5 \cdot 2^{-m} m^{1-\frac{m}{2}} e^{-\frac{m}{2}}$, then

$$\Delta_{n,m}^{(1)} \leq 2(8 \log \Delta_{n,m}^{(2)-1})^{\frac{m}{2}} \Delta_{n,m}^{(2)}. \quad (1.9)$$

The proof of Theorem 1.3 is explained in Section 2 and it is given in Section 3. This shows that the parameter $\Theta_{N,m}$ controls the statistical distance between the random vectors \mathbf{X} and \mathbf{G} . We have made significant efforts to keep track carefully of the dependency in n, m of our estimate with reasonable numerical constants. Unfortunately, this leads to an expression for $\Theta_{N,m}$ which is rather involved – see Section B.1. In particular, there are several regimes depending on n and m where different contributions are relevant. Let us just point out that in the cases we are most interested in, that is when m is large and $N = \frac{n}{m}$ is sufficiently large compared to m , we obtain the following bounds which allow us to verify the second assumption in the formulation of Theorem 1.3.

Proposition 1.4. Fix an integer $M \geq 3$. For all $m \geq M$ and $N = n/m \geq c(M)m\sqrt{1 + \log m}$,

$$\Theta_{N,m} \leq (1 + \epsilon) m^{\frac{5}{2}} 2^{\frac{m}{2}} e^{\frac{m^2}{4N}} \frac{e^{\frac{N}{2}} (1 + \log m)^N}{\sqrt{N} \Gamma(N + 1)},$$

with $\epsilon \leq 25 \cdot 10^{-5}$ and $c(M)$ are explicit constants given in the table (B.22). We emphasize that $c(M)$ is non-increasing in $M \in \mathbb{N}$ with $c(3) = 146.5$ as in Theorem 1.1 and $c(M) = 19.4$ for $M \geq 70$.

The proof of Proposition 1.4 involves rather technical numerical estimates (which have been obtained with *Mathematica*) and it is given in the Appendix B.2.

Our next result shows that it is still possible to approximate \mathbf{X} by a Gaussian random vector when $m \gg \sqrt{n}$. It is an interesting question whether the approximation also holds for the total variation distance. Recall that the Kantorovich or Wasserstein distances between the random vectors \mathbf{X} and the Gaussian \mathbf{G} are defined by for any $q \geq 1$,

$$W_q(\mathbf{X}, \mathbf{G}) = \inf_{\mathbb{P}} \left(\mathbb{E}[\|\mathbf{x} - \mathbf{g}\|^q] \right)^{1/q}, \quad (1.10)$$

where the infimum is taken over all probability measures on $\mathbb{R}^{2m} \times \mathbb{R}^{2m}$ such that the first marginal of \mathbb{P} , \mathbf{x} has the same law as \mathbf{X} and the second marginal of \mathbb{P} , \mathbf{g} is a standard Gaussian on \mathbb{R}^{2m} .

Theorem 1.5. For any $n, m \in \mathbb{N}$ such that $n \geq 2m$, it holds

$$W_2(\mathbf{X}, \mathbf{G}) \leq (\sqrt{8} + \sqrt{2}) \frac{(m+1)\sqrt{m}}{3n}.$$

This shows that if $m \rightarrow +\infty$ in such a way that $m = o(n^{2/3})$, then the Kantorovich distance between the random vector \mathbf{X} and a standard Gaussian \mathbf{G} on \mathbb{R}^{2m} converges to 0 as $n \rightarrow +\infty$. The proof of Theorem 1.5 is given in Section 7 and it relies on the normal approximation method from (Lambert et al., 2019), see Proposition 7.2 below. This result allows to turn the moments' identities of (Diaconis and Shahshahani, 1994) into a quantitative statement about the rate of convergence to the normal distribution in the Kantorovich distance. Let us emphasize that the result from (Lambert et al., 2019) which is used to prove Theorem 1.5 is inspired by Stein's method and is therefore completely unrelated to the techniques that we develop in Sections 2–6 to prove our main result.

Remark 1.1. If we let for $k \geq 1$,

$$X_{2k-1} = \frac{2}{\sqrt{\beta k}} \sum_{j=1}^n \cos(k\theta_j) \quad \text{and} \quad X_{2k} = \frac{2}{\sqrt{\beta k}} \sum_{j=1}^n \sin(k\theta_j),$$

then, the counterpart of Theorem 1.5 also holds for the circular β -ensembles $\{\theta_1, \dots, \theta_n\}$. That is, for any $\beta > 0$, there exists a constant $C_\beta > 0$ such that for all $n, m \in \mathbb{N}$ with $n \geq 2m$,

$$W_2(\mathbf{X}, \mathbf{G}) \leq C_\beta \frac{m^{3/2}}{n}.$$

The proof is similar to that of Theorem 1.5 and it relies on Proposition 7.2 and Lemma 7.3. The only differences lie in that instead of using the moments' identities of (Diaconis and Shahshahani, 1994), one can make use of the estimates from (Jiang and Matsumoto, 2015, Theorem 1). These estimates for the joint moments of \mathbf{X} corresponds to the analogue for general $\beta > 0$ of Theorem 7.1 with constants which are not sharp as they are obtained by using the Jack functions instead of Schur functions as in the case of the unitary group ($\beta = 2$). Then, it is straightforward to control the errors as in Lemmas 7.4 and 7.5. Likewise, a similar result also holds for the other classical compact groups (that is for the circular orthogonal and symplectic ensembles) with the appropriate normalization. \blacksquare

Let us give a final application of Theorem 1.3 when m is close to \sqrt{n} and the dimension n of the random matrix \mathbf{U} is large. Namely, we obtain the following corollary.

Proposition 1.6. *Let us assume that $n \geq 4322$. Then, it holds for any integer $m \leq \sqrt{\frac{n}{41.5\sqrt{\log n}}}$,*

$$d_{\text{TV}}(\mathbf{X}, \mathbf{G}) \leq \sqrt{n} \exp(19.4 - 0.93\sqrt{n}(\log n)^{5/4}).$$

The proof of Proposition 1.6 is also given in Section 3.3. We verify numerically that under the assumptions of Proposition 1.6, $d_{\text{TV}}(\mathbf{X}, \mathbf{G}) \leq 10^{-367}$ which is far below *Machine Epsilon* (of order 10^{-33} for quad(ruple) precision decimal). In the Appendix B.3, we present further numerical plots which illustrate our estimates in the case $m = 3$.

2 Overview of the proof of Theorem 1.3

The core of the proof of Theorem 1.3 is to obtain the estimate (1.8) for the L^2 distance $\Delta_{n,m}^{(2)}$ between the density $\mathcal{P}_{n,m}$ of the random vector \mathbf{X} and the standard Gaussian density on \mathbb{R}^{2m} . Observe that by Parseval's formula, we can rewrite for any $n, m \in \mathbb{N}$,

$$\Delta_{n,m}^{(2)} = \left(\int_{\mathbb{R}^{2m}} \left| F_{n,m}(\xi) - e^{-\|\xi\|^2/2} \right|^2 d\xi \right)^{1/2}, \quad (2.1)$$

where $F_{n,m}$ denotes the characteristic function of the random vector \mathbf{X} . Like in the proof of (Johansson, 1997), the general strategy is to obtain precise estimates for $F_{n,m}$ and we need to distinguish different *regimes* depending the parameters ξ , m and $N = n/m$. These *regimes* are explained in Section 2.4 and we use different methods to treat them. Compared with the arguments of (Johansson, 1997) considerable improvement is needed. There are two new challenges that come up since we allow the degree $m \in \mathbb{N}$ to grow with n and we want to keep track carefully of the constants. Let us also point out that the improvements of Theorem 1.3 come from new techniques, especially from using the *Borodin–Okounkov formula* that we recall in the next section. We also make a more careful use of the *change of variables method* from (Johansson, 1997) that we review in Section 2.3. The main steps of the proof of the estimate (1.8) are presented in Section 2.4, while the details of the proof are given in Section 3.

2.1 Notation

In this section, we collect the main notation that will be use throughout the rest of this paper.

We let $\mathbb{T} = \mathbb{R}/[2\pi]$ and view the CUE measure (1.1) as a probability measure on \mathbb{T}^n . For any $f : \mathbb{T} \rightarrow \mathbb{C}$ which is integrable, the random variable $\text{Tr } f(\mathbf{U}) = \sum_{k=1}^n f(\theta_k)$ is well-defined with $\mathbb{E}_n[\text{Tr } f(\mathbf{U})] = \widehat{f}_0$. Then, for any $\xi \in \mathbb{R}^{2m}$, we have $\mathbf{X} \cdot \xi = \text{Tr } g(\mathbf{U})$ where g is a real-valued trigonometric polynomial:

$$g(\theta) = \sum_{\substack{|k| \leq m \\ k \neq 0}} \frac{\zeta_k}{\sqrt{2|k|}} e^{ik\theta}, \quad (2.2)$$

with $\zeta_k = \xi_{2k-1} - i\xi_{2k}$ and $\zeta_{-k} = \overline{\zeta_k}$ for all $k = 1, \dots, m$. In particular the characteristic function of the random vector \mathbf{X} can be written as

$$\begin{aligned} F_{n,m}(\xi) &:= \int_{\mathbb{R}^{2m}} e^{i\xi \cdot x} \mathcal{P}_{n,m}(x) dx \\ &= \mathbb{E}_n[e^{i \text{Tr } g(\mathbf{U})}]. \end{aligned} \quad (2.3)$$

For any function $f \in L^1$, we define its Fourier coefficients for all $k \in \mathbb{Z}$,

$$\widehat{f}_k = \int_{\mathbb{T}} f(\theta) e^{-ik\theta} \frac{d\theta}{2\pi}.$$

Then, we define the following (semi)-norm

$$\|f\|_{H^{1/2}}^2 = \sum_{k \in \mathbb{Z}} |k| |\widehat{f}_k|^2.$$

If $f \in H^{1/2}$, that is if $f \in L^1$ and $\|f\|_{H^{1/2}}^2 < +\infty$, we let

$$\mathcal{A}(f) = \sum_{k \geq 1} k \widehat{f}_k \widehat{f}_{-k}. \quad (2.4)$$

If the real-valued function f lies in the Sobolev space H^1 , we also verify that

$$\|f\|_{H^{1/2}}^2 = - \int f'(\theta) \mathcal{U} f(\theta) \frac{d\theta}{2\pi},$$

where $\mathcal{U} f = - \sum_{k \in \mathbb{Z}} i \text{sgn}(k) \widehat{f}_k e^{ik\theta}$ denotes the Hilbert transform of f .

2.2 Preliminaries: Toeplitz determinants and the Borodin–Okounkov formula

Recall that the CUE refers to a random matrix \mathbf{U} which is distributed according to the Haar measure on the unitary group $U(n)$ and that the eigenvalues of \mathbf{U} have a joint law which is explicitly given by (1.1). One of the most remarkable feature of the CUE is the connection with Toeplitz determinants. Namely, for any integrable function $w = e^f$, $f : \mathbb{T} \rightarrow \mathbb{C}$ and $n \in \mathbb{N}$, if $\text{Tr } f(\mathbf{U}) = \sum_{j=1}^n f(\theta_j)$, then we have

$$\mathbb{E}_n[e^{\text{Tr } f(\mathbf{U})}] = \det_{n \times n} [\widehat{w}_{i-j}]. \quad (2.5)$$

Formula (2.5) implies that we can obtain the asymptotics of the Laplace transform of the random variable $\text{Tr } f(\mathbf{U})$ by using the Strong Szegő limit theorem.

Theorem 2.1. *If $f \in H^{1/2}$, then as $n \rightarrow +\infty$,*

$$\mathbb{E}_n [e^{\text{Tr } f(\mathbf{U})}] = \exp \left(n\widehat{f}_0 + \mathcal{A}(f) + o(1) \right), \quad (2.6)$$

where $\mathcal{A}(f) = \sum_{k \geq 1} k \widehat{f}_k \widehat{f}_{-k} \in \mathbb{C}$.

The first version of Theorem 2.1 was first proved by (Szegő, 1952) when $f \in C^{1,\alpha}$ is real-valued. The hypothesis from Theorem 2.1 are optimal and this version was first obtained for real-valued f by (Ibragimov, 1968) and (Golinskii and Ibragimov, 1971). We refer to the survey paper of (Deift et al., 2013) for a history of the Szegő Strong Limit theorem and its later generalizations and to the book (Simon, 2005b, Chapter 6) for a detailed presentation of several proofs. A proof of Theorem 2.1 which holds for complex-valued f can be found in (Johansson, 1988).

Actually, one can also obtain Theorem 2.1 as a consequence of the *Borodin–Okounkov formula*. This formula expresses the Toeplitz determinant (2.5) in terms of Fredholm determinant which is more amenable for asymptotic analysis. If $f : \mathbb{T} \rightarrow \mathbb{C}$ is an L^2 function, we denote

$$f^+(\theta) = \sum_{k \geq 1} \widehat{f}_k e^{ik\theta}, \quad f^-(\theta) = \sum_{k \geq 1} \widehat{f}_{-k} e^{-ik\theta}.$$

Let $w : \mathbb{T} \rightarrow \mathbb{C}$ be an integrable function such that $\sum_{k \in \mathbb{Z}} |k| |\widehat{w}_k|^2 < +\infty$, and define two Hankel operators:

$$H_+(w) = \begin{pmatrix} \widehat{w}_1 & \widehat{w}_2 & \widehat{w}_3 & \dots \\ \widehat{w}_2 & \widehat{w}_3 & \widehat{w}_4 & \dots \\ \widehat{w}_3 & \widehat{w}_4 & \widehat{w}_5 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad \text{and} \quad H_-(w) = \begin{pmatrix} \widehat{w}_{-1} & \widehat{w}_{-2} & \widehat{w}_{-3} & \dots \\ \widehat{w}_{-2} & \widehat{w}_{-3} & \widehat{w}_{-4} & \dots \\ \widehat{w}_{-3} & \widehat{w}_{-4} & \widehat{w}_{-5} & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}. \quad (2.7)$$

Note that the condition $\sum_{k \in \mathbb{Z}} |k| |\widehat{w}_k|^2 < +\infty$ guarantees that these operators are Hilbert–Schmidt on $L^2(\mathbb{N})$. We also denote by (e_1, e_2, \dots) the standard basis of $L^2(\mathbb{N})$.

Theorem 2.2. *Let $f : \mathbb{T} \rightarrow \mathbb{C}$ be a L^∞ function such that $\sum_{k \in \mathbb{Z}} |k| |\widehat{f}_k|^2 < +\infty$ and $\widehat{f}_0 = 0$. Let us also define*

$$K_f = H_+(e^{f^- - f^+}) H_-(e^{f^+ - f^-}). \quad (2.8)$$

The operator K_f is trace-class and for any $n \in \mathbb{N}$,

$$\mathbb{E}_n [e^{\text{Tr } f(\mathbf{U})}] = e^{\mathcal{A}(f)} \det[\mathbf{I} - K_f Q_n], \quad (2.9)$$

where Q_n denotes the orthogonal projection with kernel $\text{span}(e_1, \dots, e_{n-1})$ and the RHS is a Fredholm determinant on $L^2(\mathbb{N})$.

Since the operator K_f is trace class, by definition of Q_n , we have $\det[\mathbf{I} - K_f Q_n] \rightarrow 1$ as $n \rightarrow +\infty$, so that Theorem 2.2 implies the Szegő Strong Limit theorem. The *Borodin–Okounkov formula* (2.9) (sometimes also known as *Geronimo–Case formula*) first appeared (formally) in (Geronimo and Case, 1979). (Borodin and Okounkov, 2000) proved formula (2.9) in a different form when f is analytic using Gessel’s Theorem which allows to express Toeplitz determinants as series in Schur functions, (Gessel, 1990). The version from Theorem 2.2 is due to (Basor and Widom, 2000) – see also (Böttcher, 2002) for a different proof. It is possible to remove the condition $f \in L^\infty$ from Theorem 2.2, see e.g. (Simon, 2005b, Chapter 6.2).

Concerning our method, let us point out that in order to obtain the *super-exponential* rate of convergence in (1.2), (Johansson, 1997) relied on exact formulae for Toeplitz determinants with certain specific symbols which are due to (Baxter, 1961) and relates to the original proof of the Strong Szegő

theorem. Observe that according to (2.2), we have $\mathcal{A}(\text{ig}) = -\|\zeta\|^2/2 = -\|\xi\|^2/2$ so that by (2.3) and (2.9), we can rewrite for all $n, m \in \mathbb{N}$ and $\xi \in \mathbb{R}^{2m}$,

$$F_{n,m}(\xi) = e^{-\|\xi\|^2/2} \det[\mathbb{I} - K_{\text{ig}} Q_n]. \quad (2.10)$$

Hence, by controlling precisely how close the Fredholm determinant $\det[\mathbb{I} - K_{\text{ig}} Q_n]$ is to 1, we are able to significantly improve the rate of convergence from (Johansson, 1997). Even though this might be difficult to verify, it is natural to expect that modulo corrections, $1/\Gamma(N+1)$ should be the true rate of convergence in Theorem 1.3 in the regime where $m \ll N = \frac{n}{m}$.

Throughout this article, we also make crucial use of the following bound.

Lemma 2.3. *Suppose that $f \in \mathcal{C}(\mathbb{T})$ is real-valued with $\mathcal{A}(f) < +\infty$ where \mathcal{A} is as in (2.4). Then for any $n \in \mathbb{N}$,*

$$\mathbb{E}_n[e^{\text{Tr} f(\mathbf{U})}] \leq \exp(n\widehat{f}_0 + \mathcal{A}(f)). \quad (2.11)$$

Let us recall that $\mathbb{E}_n[\text{Tr} f(\mathbf{U})] = n\widehat{f}_0$ and that by Theorem 2.1, $\text{Var}[\text{Tr} f(\mathbf{U})] \rightarrow 2\mathcal{A}(f)$ as $n \rightarrow +\infty$, so that the estimate (2.11) is sharp. The upper-bound (2.11) is classical and it follows for instance from the monotonicity of Toeplitz determinants, (Simon, 2005a). For completeness, we show in the appendix (Section A.1), how one can immediately deduce Lemma 2.3 from the *Borodin–Okounkov formula*.

2.3 Change of variables

In addition to the *Borodin–Okounkov formula* (Theorem 2.2) and Lemma 2.3, our main tool to prove Theorem 1.3 is the change of variables method introduced in (Johansson, 1988). More specifically we rely on an estimate from the proof of (Johansson, 1997, Proposition 2.8). Recall that according to (2.3), $F_{n,m}$ denotes the characteristic function of the random variable $\text{Tr} g(\mathbf{U})$.

Lemma 2.4. *Let $\nu > 0$ and $h : \mathbb{T} \rightarrow \mathbb{R}$ be a C^1 function. Then, for any $n, m \in \mathbb{N}$ and $\xi \in \mathbb{R}^{2m}$,*

$$|F_{n,m}(\xi)| \leq \mathbb{E}_n \left[\prod_{1 \leq i < j \leq n} \left| \frac{\sin\left(\frac{\theta_i - \theta_j}{2} + i\nu \frac{h(\theta_i) - h(\theta_j)}{2n}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)} \right|^2 \prod_{j=1}^n |1 + i\frac{\nu}{n} h'(\theta_j)| e^{-\Im g(\theta_j + i\frac{\nu}{n} h(\theta_j))} \right].$$

Proof. For completeness, let us give the proof of Lemma 2.4. Using the explicit formula (1.1) for the joint law of the eigenvalues of the random matrix \mathbf{U} , we obtain

$$F_{n,m}(\xi) = \mathcal{U}_n \int_{[-\pi, \pi]^n} \prod_{1 \leq i < j \leq n} \sin^2\left(\frac{\theta_i - \theta_j}{2}\right) \prod_{k=1}^n e^{ig(\theta_k)} \frac{d\theta_k}{2\pi}.$$

If we regard θ_k as complex variables in the previous integral, since the integrand is an entire function, we can deform the contours of integration in the complex plane. Let γ be a positively oriented curve given by

$$\gamma = \left\{ \theta + i\frac{\nu}{n} h(\theta) : \theta \in [-\pi, \pi] \right\}.$$

Since the functions g and $\sin^2(\cdot/2)$ are also 2π -periodic, we have by Cauchy's theorem,

$$\begin{aligned} F_{n,m}(\xi) &= \mathcal{U}_n \int_{\gamma^n} \prod_{1 \leq i < j \leq n} \sin^2\left(\frac{\theta_i - \theta_j}{2}\right) \prod_{k=1}^n e^{ig(\theta_k)} \frac{d\theta_k}{2\pi} \\ &= \mathcal{U}_n \int_{[-\pi, \pi]^n} \prod_{1 \leq i < j \leq n} \left(\sin\left(\frac{\theta_i - \theta_j}{2} + i\nu \frac{h(\theta_i) - h(\theta_j)}{2n}\right) \right)^2 \prod_{j=1}^n e^{ig(\theta_j + i\frac{\nu}{n} h(\theta_j))} (1 + i\frac{\nu}{n} h'(\theta_j)) \frac{d\theta_j}{2\pi}. \end{aligned} \quad (2.12)$$

Hence, by (1.1), this implies that

$$\begin{aligned} |F_{n,m}(\xi)| &\leq \mathcal{U}_n \int_{[-\pi,\pi]^n} \prod_{1 \leq i < j \leq n} \left| \sin \left(\frac{\theta_i - \theta_j}{2} + i\nu \frac{h(\theta_i) - h(\theta_j)}{2n} \right) \right|^2 \prod_{k=1}^n e^{-\Im g(\theta_j + i\frac{\nu}{n}h(\theta_j))} \left| 1 + i\frac{\nu}{n}h'(\theta_j) \right| \frac{d\theta_j}{2\pi} \\ &= \mathbb{E}_n \left[\prod_{1 \leq i < j \leq n} \left| \frac{\sin \left(\frac{\theta_i - \theta_j}{2} + i\nu \frac{h(\theta_i) - h(\theta_j)}{2n} \right)}{\sin \left(\frac{\theta_i - \theta_j}{2} \right)} \right|^2 \prod_{j=1}^n \left| 1 + i\frac{\nu}{n}h'(\theta_j) \right| e^{-\Im g(\theta_j + i\frac{\nu}{n}h(\theta_j))} \right]. \end{aligned}$$

□

The key idea underlying this change of variables is that the eigenvalues of \mathbf{U} are almost uniformly distributed on the unit circle (like the vertices of a regular n -gon). This means that at first order, we can approximate the empirical measure $\sum_{j=1}^n \delta_{\theta_j} \simeq n \frac{d\theta}{2\pi}$. Choose $h = \mathcal{U}g$ where \mathcal{U} is the Hilbert transform:

$$h(\theta) = - \sum_{\substack{|k| \leq m \\ k \neq 0}} \operatorname{sgn}(k) \frac{i\zeta_k}{\sqrt{2|k|}} e^{ik\theta}. \quad (2.13)$$

By making the change of variables θ_j by $\theta_j + i\frac{\nu}{n}h(\theta_j)$ in (2.12), we expect that using first order Taylor approximations:

$$\begin{aligned} F_{n,m}(\xi) &\simeq \mathcal{U}_n \int_{[-\pi,\pi]^n} \prod_{1 \leq i < j \leq n} \sin^2 \left(\frac{\theta_i - \theta_j}{2} \right) e^{\frac{\nu^2}{n^2} H(\theta_i, \theta_j)} \prod_{j=1}^n e^{i\mathfrak{g}(\theta_j) - \frac{\nu}{n} \mathfrak{g}'(\theta_j)h(\theta_j) + \frac{\nu^2}{n^2} h'(\theta_j)^2} \frac{d\theta_j}{2\pi} \\ &= \mathbb{E}_n \left[e^{\frac{\nu^2}{2n^2} \sum_{i,j=1}^n H(\theta_i, \theta_j)} e^{i \sum_{j=1}^n \mathfrak{g}(\theta_j) - \frac{\nu}{n} \sum_{j=1}^n \mathfrak{g}'(\theta_j)h(\theta_j)} \right], \end{aligned}$$

where

$$H(\theta, x) = \begin{cases} \left(\frac{h(\theta) - h(x)}{2 \sin \left(\frac{\theta - x}{2} \right)} \right)^2 & x \neq \theta \\ h'(\theta)^2 & x = \theta \end{cases}. \quad (2.14)$$

Then, since $\widehat{\mathfrak{g}}_0 = 0$, we expect that

$$F_{n,m}(\xi) \simeq \exp \left(-\nu \int_{[0,2\pi]} g'(\theta)h(\theta) \frac{d\theta}{2\pi} + \frac{\nu^2}{2} \iint_{[0,2\pi]^2} H(\theta, x) \frac{d\theta}{2\pi} \frac{dx}{2\pi} \right).$$

Then, by Devinatz's formula (Simon, 2005a, Proposition 6.1.10), since $h = -\mathcal{U}g$, we have

$$\|h\|_{H^{1/2}}^2 = \int_{[0,2\pi]} g'(\theta)h(\theta) \frac{d\theta}{2\pi} = \iint_{[0,2\pi]^2} H(\theta, x) \frac{d\theta}{2\pi} \frac{dx}{2\pi} \quad (2.15)$$

and

$$\|h\|_{H^{1/2}}^2 = \sum_{k \in \mathbb{Z}} |k| |h_k|^2 = \sum_{k=1}^m |\zeta_k|^2 = \|\xi\|^2. \quad (2.16)$$

Whence it follows from this heuristic with $\nu = 1$ that

$$F_{n,m}(\xi) \simeq e^{-\|\xi\|^2/2}.$$

To turn this heuristics rigorous, one needs to justify the approximation $\sum_{j=1}^n \delta_{\theta_j} \simeq n \frac{d\theta}{2\pi}$ and to control the errors coming from the Taylor expansions. This can be done by using *rigidity estimates* for the CUE eigenvalues, see (Lambert, 2019), but we present a different approach below (see **iii**) Intermediate regime in the next section).

2.4 Estimates for the function $F_{n,m}(\xi)$ in the different regimes

Recall that we let $N = \frac{n}{m}$ and that our main goal is to obtain the following bound.

Proposition 2.5. *For any $n, m \in \mathbb{N}$ such that $m \geq 3$ and $N = n/m > 4m$, we have*

$$\Delta_{n,m}^{(2)} = \left(\int_{\mathbb{R}^{2m}} \left| F_{n,m}(\xi) - e^{-\|\xi\|^2/2} \right|^2 d\xi \right)^{1/2} \leq c_3 \sqrt{\Omega_m} N^{\frac{m}{2}} \Theta_{N,m}, \quad (2.17)$$

where $\Theta_{N,m}$ is as in (B.7)–(B.10).

In this section, we present the main estimates for the characteristic function $F_{n,m}(\xi)$ that are required to prove Proposition 2.5. We postpone the technical details of the arguments to Sections 4–6. All the constants c_j used below, which can depend on m are defined in the Appendix B. Let us define

$$\Lambda_1 = \frac{c_4 N}{\sqrt{1 + \log m}}. \quad (2.18)$$

The proof consists in splitting the integral on the LHS of (2.17) in three different regimes depending on whether **i)** $\|\xi\| \leq \Lambda_1$, **ii)** $\|\xi\| \geq \Lambda_3$ or **iii)** $\Lambda_1 \leq \|\xi\| \leq \Lambda_3$ where $\Lambda_3 \gg \Lambda_1$ is a parameter that we will choose later.

i) Gaussian approximation for $\|\xi\| \leq \Lambda_1$. In this regime, our goal is to compare the characteristic function $F_{n,m}$ with that of a $2m$ -dimensional standard Gaussian by using the *Borodin–Okounkov formula* from Theorem 2.2. We obtain the following estimates.

Proposition 2.6. *Under the assumptions of Proposition 2.5, we have for all $\xi \in \mathbb{R}^{2m}$ such that $\|\xi\| \leq \Lambda_1$,*

$$\left| F_{n,m}(\xi) - e^{-\|\xi\|^2/2} \right|^2 \leq c_8^2 m^4 e^{2\sqrt{2(1+\log m)}\|\xi\|} \left(\frac{1 + \log m}{2} \right)^{2N} \frac{\|\xi\|^{4N}}{\Gamma(N+1)^4} e^{-\|\xi\|^2}.$$

Let us point out that Proposition 2.6 gives the main contribution $\Theta_{N,m}^0$ to $\Delta_{n,m}^{(2)}$. We expect that the main error in the normal approximation should come from the regime where $\|\xi\|$ is not too large. The proof of Proposition 2.6 is given in Section 4. Let us observe that according to formula (2.10), we have

$$\left| F_{n,m}(\xi) - e^{-\|\xi\|^2/2} \right|^2 = \left| 1 - \det[\mathbf{I} - K_{\text{ig}} Q_n] \right|^2 e^{-\|\xi\|^2},$$

and we expect that if both the degree m and $\|g\|_{H^{1/2}}^2 = \|\xi\|^2$ are sufficiently small (depending on the dimension $n \in \mathbb{N}$ of the random matrix \mathbf{U}), then by definition of the projection Q_n , the operator $K_{\text{ig}} Q_n$ is also small (in trace norm) so that $\det[\mathbf{I} - K_{\text{ig}} Q_n] \simeq 1$. This can be quantified by using the bound for Fredholm determinant from (Simon, 2005b, Theorem 3.4),

$$\left| 1 - \det[\mathbf{I} - K_{\text{ig}} Q_n] \right| \leq \|K_{\text{ig}} Q_n\|_{\mathcal{S}_1} e^{1 + \|K_{\text{ig}} Q_n\|_{\mathcal{S}_1}}, \quad (2.19)$$

where $\|\cdot\|_{\mathcal{S}_1}$ denotes the Schatten 1-norm or trace norm of an operator. Then, in order to compute $\|K_{\text{ig}} Q_n\|_{\mathcal{S}_1}$, we use the product structure of the operator K_{ig} , (2.8), and the Cauchy–Schwartz inequality (for the Hilbert–Schmidt norm $\|\cdot\|_{\mathcal{S}_2}$):

$$\|K_{\text{ig}} Q_n\|_{\mathcal{S}_1} \leq \|Q_n H_+(e^{2\Im g^+})\|_{\mathcal{S}_2} \|H_-(e^{-2\Im g^+}) Q_n\|_{\mathcal{S}_2}.$$

Moreover, since $H_{\pm}(\cdot)$ are Hankel operators (2.7), we can estimate the norms $\|H_{\pm}(e^{-2\Im g^{\pm}}) Q_n\|_{\mathcal{S}_2}$ by obtaining bounds for the Fourier coefficients of the symbols $e^{-2\Im g^{\pm}}$, see Lemma 4.1 below. To sum up, we show in Section 4 that $\|Q_n H_{\pm}(e^{2\Im g^{\pm}})\|_{\mathcal{S}_2} \ll 1/\Gamma(1+N)$ provided that $\|\xi\| \ll N$ and we use

this estimate to deduce Proposition 2.6. Let us emphasize again that we expect that these estimates are of the right order and hold only in the regime where $\|\xi\| \ll N$.

ii) Tail bound for large $\|\xi\|$. If $\|\xi\|$ is very large, we are not looking to compare $F_{n,m}$ with the characteristic function of a standard Gaussian, but rather aiming at obtaining a *good* tail bound for $F_{n,m}$. By *good*, we mean that we aim for estimates which yield errors that are smaller than $\Theta_{N,m}^0$ when N is sufficiently large. (Johansson, 1997, Proposition 2.13) used the Hadamard's inequality

$$|F_{n,m}(\xi)|^2 \leq \prod_{j=1}^n \sum_{i=1}^n |(e^{i\mathbf{g}})_{j-i}|^2 \quad (2.20)$$

and an estimate for the Fourier coefficients of the function $e^{i\mathbf{g}}$ to obtain the tail bound $|F_{n,m}(\xi)|^2 \leq \frac{C^n n^{\frac{3n}{2}}}{\|\xi\|^{\frac{N}{2}}}$ for a constant $C > 0$. By using (2.20) and a (sharp) *Van der Corput's inequality*, this estimate can be improved and we obtain for all $m, n \geq 3$ and $\xi \in \mathbb{R}^{2m}$,

$$|F_{n,m}(\xi)|^2 \leq \frac{c^n n^n}{\|\xi\|^{\frac{n}{m+1}}}, \quad c = 4\pi e(1 + 1/\sqrt{3}). \quad (2.21)$$

To obtain a good multi-dimensional approximation for a growing number of traces we would like to have a better estimate that does not contain the very large factor n^n . We can obtain a different tail bound by relying on Lemma 2.4 with $h = g'$. Choosing $\nu > 0$ appropriately, we obtain

$$|F_{n,m}(\xi)| \leq e^{cn} \mathbb{E}_n \left[e^{-\gamma \sum_{j=1}^n g'(\theta_j)^2} \right], \quad (2.22)$$

for a constant $c > 0$ and $\gamma \rightarrow 0$ as $n \rightarrow +\infty$ – see Proposition 5.1 below for further details. Then, to estimate the RHS of (2.22), we use that

$$\mathbb{E}_n \left[e^{-\gamma \sum_{j=1}^n g'(\theta_j)^2} \right] \leq \frac{e^n}{\sqrt{2\pi n}} \left(\int_{\mathbb{T}} e^{-\gamma g'(\theta)^2} \frac{d\theta}{2\pi} \right)^n, \quad (2.23)$$

and since $g' : \mathbb{T} \rightarrow \mathbb{R}$ is a trigonometric polynomial of degree $m \in \mathbb{N}$,

$$\int_{\mathbb{T}} e^{-\gamma g'(\theta)^2} \frac{d\theta}{2\pi} \leq \frac{2e}{(2\gamma \|g'\|_{L^2}^2)^{1/4m}}. \quad (2.24)$$

The estimate (2.23) is rather classical and its proof is given in the appendix – Lemma 5.3 – for completeness. On the other-hand, (2.24) relies on an estimate of the measure of the set where a trigonometric polynomial is small by its L^2 norm which is taken from (Chahkiev, 2008) – see Lemma 5.4 below. By combining these estimates, we obtain the following bound in Section 5.

Proposition 2.7. *Fix $n, m \in \mathbb{N}$ and suppose that $N \geq 4m$. For any $\xi \in \mathbb{R}^{2m}$, we have*

$$|F_{n,m}(\xi)|^2 \leq \Upsilon_3(m)^{N/2} \frac{c_{15}^{2n} N^{N/4}}{\|\xi\|^{N/2}}.$$

While this tail bound has a worse decay in $\|\xi\|$ than (2.21), the factor N^N is significantly better than n^n when the degree $m \in \mathbb{N}$ is large. Moreover, we see that this estimate will be useful in the proof of Proposition 2.5 in the regime where $\|\xi\| \gg N^c$ for a sufficiently large constant c . In the proof, we will actually choose $\Lambda_3 = e^{4c_1 \frac{N}{1+\log m}}$ times some corrections – see formula (3.1) below.

iii) Intermediate regime. It remains to deal with the intermediate regime where $\Lambda_1 \leq \|\xi\| \leq \Lambda_3$. As we already pointed out, when $\|\xi\| \gg N$, we do not expect that the Fredholm determinant $\det[\mathbf{I} - K_{\text{ig}} Q_n]$ is close to 1. However, we still expect that $|F_{n,m}(\xi)|^2 \ll 1/\Gamma(1+N)^2$ for all such $\xi \in \mathbb{R}^{2m}$. From a

technical perspective, this intermediate regime is the most challenging one because the direct estimates (e.g. the method used in (Johansson, 1997, Section 2.2)) lead to errors which are bigger than that of Proposition 2.6 – see the estimate (2.30) below. Our final bounds are summarized in the next proposition. Define

$$\Lambda_2 = \frac{c_0^{-1}(1 - c_{10})N\sqrt{m+1}}{8(1 + \log m)^{3/4}c_{11}}, \quad (2.25)$$

where $c_1(m)$, $c_2(m)$, $c_{10}(m)$ and $c_{11}(m)$ are as in (B.3). We verify that both $c_{10}(m)$ and $c_{11}(m)$ are decreasing for $m \geq 3$. Since $c_{10}(3) \approx 0.0124$ and $c_{11}(3) \approx 1.583$, this shows that $\Lambda_2 \geq \Lambda_1$ and Λ_2 is increasing as a function of m for all $m \geq 3$.

Proposition 2.8. *Fix $m, n \in \mathbb{N}$ with $m \geq 3$. We have for all $\xi \in \mathbb{R}^{2m}$,*

$$|F_{n,m}(\xi)| \leq \exp\left(c_9 - \frac{c_1(m)N^2}{1 + \log m}\right) \quad \text{if } \|\xi\| \geq \Lambda_2, \quad (2.26)$$

and

$$|F_{n,m}(\xi)| \leq \exp\left(c_9 - \frac{c_2(m)N^2}{\sqrt{m+1}(1 + \log m)^{3/4}}\right) \quad \text{if } \Lambda_1 \leq \|\xi\| \leq \Lambda_2. \quad (2.27)$$

Let us observe that these bounds directly relate to the error terms $\Theta_{N,m}^1$ and $\Theta_{N,m}^2$ from (B.9) and (B.8) respectively. The proof of Proposition 2.8 is given in Section 6. The starting point of this proof is the change of variables and the heuristics described in Section 2.3. Namely, using Lemma 2.4 with $h = -\mathcal{U}g$ as in (2.13), we obtain the following bound.

Lemma 2.9. *Let $n, m \in \mathbb{N}$ and $\xi \in \mathbb{R}^{2m}$. We have for any $\nu > 0$,*

$$|F_{n,m}(\xi)|^2 \leq \mathbb{E}_n \left[\exp\left(\frac{\nu^2}{n^2} \sum_{i,j=1}^n H(\theta_i, \theta_j)\right) \right] \mathbb{E}_n \left[e^{-2 \sum_{j=1}^n \Im g(\theta_j + i \frac{\nu}{n} h(\theta_j))} \right], \quad (2.28)$$

where the function H is given by (2.14).

The proof of Lemma 2.9 is postponed to Section 6.2. Using Lemma 2.3, we can easily control the second factor in the RHS of (2.28). We obtain that there exists a constant $c > 0$ such that if $\|\xi\| \ll \Lambda_2$,

$$\mathbb{E}_n \left[e^{-2 \sum_{j=1}^n \Im g(\theta_j + i \frac{\nu}{n} h(\theta_j))} \right] \leq e^{-c\nu \|\xi\|^2} \quad (2.29)$$

see Lemma 6.1 below for further details. Moreover, using the deterministic bound $H(\theta_i, \theta_j) \leq \|h'\|_\infty^2$ in (2.28) combined with $\|h'\|_\infty \leq \sqrt{2m}\|\xi\|$, this implies that

$$|F_{n,m}(\xi)| \leq \exp(-c\nu \|\xi\|^2 + 2\nu^2 m^2 \|\xi\|^2).$$

If we optimize over $\nu > 0$, this leads to

$$|F_{n,m}(\xi)| \leq \exp\left(-\frac{c^2 \|\xi\|^2}{8m^2}\right). \quad (2.30)$$

In the regime where the degree $m \in \mathbb{N}$ depends on the dimension $n \in \mathbb{N}$ with $N \geq 4m$, the naive estimate (2.30) is not precise enough to lead to errors which are small compared with that of Proposition 2.6. Therefore, to prove Proposition 2.8, we need to introduce a new idea. One approach would be to use precise rigidity estimates from (Lambert, 2019) to obtain a better estimate for the first term on the RHS of (2.28). But, the method that we use consists in writing $\sum_{i,j=1}^n H(\theta_i, \theta_j)$ as a quadratic form in the random variables $T_k = \text{Tr } \mathbf{U}^k$,

$$\sum_{i,j=1}^n H(\theta_i, \theta_j) = n^2 \iint_{[0,2\pi]^2} H(\theta, x) \frac{d\theta}{2\pi} \frac{dx}{2\pi} + n(\mathbf{a}^* \mathbf{T} + \mathbf{T}^* \mathbf{a}) + \mathbf{T}^* \mathbf{M} \mathbf{T}, \quad \text{where } \mathbf{T} = \begin{pmatrix} T_1 \\ \vdots \\ T_{2m-1} \end{pmatrix},$$

$\mathbf{a}(\xi) \in \mathbb{C}^{2m}$ is a deterministic vector and $\mathbf{M}(\xi) \in \mathbb{C}^{2m \times 2m}$ is a deterministic matrix which depend on $\xi \in \mathbb{R}^{2m}$, see Lemma 6.3 below. This allows us to express the Laplace transform of the random variable $\sum_{i,j=1}^n H(\theta_i, \theta_j)$ as an integral against a Gaussian measure on \mathbb{C}^{2m} . It is not at all clear that the matrix $\mathbf{M}(\xi)$ is positive definite, but *if* it were (see the Remark 6.1), by formulae (2.15)–(2.16), we would obtain for any $\nu > 0$,

$$\mathbb{E}_n \left[e^{\frac{\nu^2}{n^2} \sum_{i,j=1}^n H(\theta_i, \theta_j)} \right] = \frac{e^{\nu^2 \|\xi\|^2}}{\pi^{2m} \det(\mathbf{M})} \int_{\mathbb{C}^{2m-1}} e^{-\mathbf{z}^* \mathbf{M}^{-1} \mathbf{z}} \mathbb{E}_n \left[e^{\frac{\nu}{n} (\mathbf{z}^* \mathbf{T} + \mathbf{T}^* \mathbf{z}) + \frac{\nu^2}{n} (\mathbf{a}^* \mathbf{T} + \mathbf{T}^* \mathbf{a})} \right] d\mathbf{z},$$

where $d\mathbf{z}$ is the Lebesgue measure on \mathbb{C}^{2m-1} . The idea is now to use Lemma 2.3 to estimate the expectation on the RHS of the previous formula and then to evaluate the Gaussian integral. The details in the implementation of this idea, which requires a modification of \mathbf{M} , is somewhat involved and we refer to the proof in Section 6 for further details.

3 Proof of the main result

3.1 Proof of Proposition 2.5

In this section, we give the proof of Proposition 2.5 relying on the estimates from Propositions 2.6, 2.7 and 2.8. Recall that we assume that $m \geq 3$ and $N = \frac{n}{m} > 4m$. Then, using the notation (B.3) and (B.11), we define

$$\Lambda_3 = e^{-4c_9/N} c_{15}^{4m} \left(\frac{N}{4m} - 1 \right)^{2/N} \Upsilon_3(m) \sqrt{N} \exp \left(\frac{4c_1(m)N}{1 + \log m} \right). \quad (3.1)$$

Let us also recall that Λ_1 is given by (2.18) and Λ_2 is given by (2.25), so that $\Lambda_3 \gg \Lambda_2 \gg \Lambda_1$ as $n \rightarrow +\infty$ (and possibly $m \rightarrow +\infty$). We will also need the following bound which is proved in the appendix (Section A.2).

Lemma 3.1. *For any $m \in \mathbb{N}$, if $\Lambda > \sqrt{m}$,*

$$\int_{\|\xi\| \geq \Lambda} e^{-\|\xi\|^2} d\xi \leq \Omega_m \frac{e^{-\Lambda^2}}{\Lambda^2 - m}.$$

Since $\Lambda_1^2 \geq 2m$ for all $m \geq 3$, it follows from Lemma 3.1 that

$$\int_{\|\xi\| \geq \Lambda_1} e^{-\|\xi\|^2} d\xi \leq \Omega_m \frac{e^{-\Lambda_1^2}}{\Lambda_1^2 - m} \leq \frac{\Omega_m}{m} \exp \left(-\frac{N^2}{8(1 + \log m)} \right). \quad (3.2)$$

Set $\Lambda_4 = +\infty$ and let us denote for $k = 1, 2, 3$,

$$\mathcal{J}_0^2 = \int_{\substack{\mathbb{R}^{2m} \\ \|\xi\| \leq \Lambda_1}} \left| F_{n,m}(\xi) - e^{-\|\xi\|^2/2} \right|^2 d\xi, \quad \mathcal{J}_k^2 = \int_{\substack{\mathbb{R}^{2m} \\ \Lambda_k \leq \|\xi\| \leq \Lambda_{k+1}}} |F_{n,m}(\xi)|^2 d\xi.$$

Then, by splitting the integral in (2.1) and using the estimate (3.2), we obtain

$$\Delta_{n,m}^{(2)} \leq \mathcal{J}_0 + \mathcal{J}_1 + \mathcal{J}_2 + \mathcal{J}_3 + \sqrt{\frac{\Omega_m}{m}} \exp \left(-\frac{N^2}{16(1 + \log m)} \right). \quad (3.3)$$

As we explain in Section 2.4, we expect that the main contribution in (3.3) comes from \mathcal{J}_0 . By Proposition 2.6, we obtain

$$\mathcal{J}_0^2 = \int_{\|\xi\| \leq \Lambda_1} \left| e^{-\|\xi\|^2/2} - F_{n,m}(\xi) \right|^2 d\xi \leq \frac{c_8^2 m^4 e^{2\sqrt{2(1+\log m)}\Lambda_1} \left(\frac{1+\log m}{2} \right)^{2N}}{\Gamma(N+1)^4} \int_{\|\xi\| \leq \Lambda_1} \|\xi\|^{4N} e^{-\|\xi\|^2} d\xi.$$

Moreover, by going to polar coordinates, we have

$$\begin{aligned} \int_{\|\xi\| \leq \Lambda_1} \|\xi\|^{4N} e^{-\|\xi\|^2} d\xi &= m\Omega_m \int_0^{\Lambda_1^2} u^{2N+m-1} e^{-u} du \\ &\leq m\Omega_m \Gamma(2N+m). \end{aligned}$$

For $3 \leq m \leq N$, we also have

$$\begin{aligned} \frac{\Gamma(2N+m)}{\Gamma(N+1)^2} &\leq e^{-m} \frac{(2N+m)^{2N+m}}{\sqrt{2\pi} N^{2N+1}} = \frac{4^N}{\sqrt{2\pi}} N^{m-1} \left(1 + \frac{m}{2N}\right)^{2N+m} \left(\frac{2}{e}\right)^m \\ &\leq \frac{4^N 2^m e^{\frac{m^2}{2N}}}{\sqrt{2\pi}} N^{m-1}. \end{aligned}$$

Here we used the upper-bound $\Gamma(k) \leq \sqrt{2\pi} k^k e^{-k}$ which holds for all integer $k \geq 9$ and the lower-bound $\Gamma(N+1) \geq \sqrt{2\pi} N^N e^{-N}$ which holds for all $N \in \mathbb{N}$. For the last step, we used that $(1+x)^\alpha \leq e^{\alpha x}$ for any $x, \alpha \geq 0$. By (2.18), it holds that $2\sqrt{2}(1+\log m)\Lambda_1 = N$, so we obtain

$$\begin{aligned} \mathcal{J}_0^2 &\leq \frac{c_8^2}{\sqrt{2\pi}} m^5 \Omega_m 2^m \frac{e^{N+\frac{m^2}{2N}} (1+\log m)^{2N} N^{m-1}}{\Gamma(N+1)^2} \\ &= (c_3 \sqrt{\Omega_m} N^{\frac{m}{2}} \Theta_{N,m}^0)^2, \end{aligned} \tag{3.4}$$

according to formula (B.7) and (B.3).

In the rest of the proof, we give bounds for the integrals \mathcal{J}_k for $k = 1, 2, 3$. First, using the tail bound from Proposition 2.7, we have

$$\mathcal{J}_3^2 = \int_{\|\xi\| \geq \Lambda_3} |F_{n,m}(\xi)|^2 d\xi \leq \Upsilon_3(m)^{N/2} c_{15}^{2n} N^{N/4} \int_{\|\xi\| \geq \Lambda_3} \|\xi\|^{-N/2} d\xi.$$

Hence, since we assume that $N > 4m$, the previous integral is finite and we obtain

$$\mathcal{J}_3^2 \leq \frac{2m\Omega_m}{N/2 - 2m} \frac{\Upsilon_3^{N/2} c_{15}^{2n} N^{N/4}}{\Lambda_3^{N/2-2m}}. \tag{3.5}$$

Second, by using the estimate (2.26) from Proposition 2.8, we also have

$$\mathcal{J}_2^2 = \int_{\Lambda_2 \leq \|\xi\| \leq \Lambda_3} |F_{n,m}(\xi)|^2 d\xi \leq \Omega_m \Lambda_3^{2m} \exp\left(2c_9 - \frac{2c_1 N^2}{1+\log m}\right) \tag{3.6}$$

Hence, by combining the estimates (3.5) and (3.6), this implies that

$$\mathcal{J}_2 + \mathcal{J}_3 \leq \sqrt{\Omega_m} \Lambda_3^m \left(\exp\left(c_9 - \frac{c_1 N^2}{1+\log m}\right) + \frac{N^{N/8} c_{15}^n \Upsilon_3^{N/4}}{\sqrt{N/4m - 1} \Lambda_3^{N/4}} \right). \tag{3.7}$$

Our choice of Λ_3 consists in optimizing¹ the RHS of (3.7). Namely, by choosing Λ_3 according to (3.1), we obtain for all $N > 4m$,

$$\begin{aligned} \mathcal{J}_2 + \mathcal{J}_3 &\leq \frac{\sqrt{\Omega_m}}{1 - 4m/N} \Lambda_3^m \exp\left(c_9 - \frac{c_1 N^2}{1+\log m}\right) \\ &\leq \left(\frac{N}{4m}\right)^{\frac{2m}{N}} \frac{e^{c_9(1-\frac{4m}{N})} \sqrt{\Omega_m}}{(1 - 4m/N)^{1-\frac{2m}{N}}} N^{\frac{m}{2}} c_{15}^{4m^2} \Upsilon_3^m \exp\left(-\frac{c_1 N(N-4m)}{1+\log m}\right). \end{aligned}$$

¹If $\alpha > m$, the minimum of the function $\Lambda^m \epsilon + C\Lambda^{m-\alpha}$ over all $\Lambda > 0$ is attained when $\Lambda^\alpha = (\frac{\alpha}{m} - 1)\epsilon^{-1}C$ and equals to $\frac{\epsilon \Lambda^m}{1-m/\alpha}$.

Then, by using the estimate (B.13) and that $\left(\frac{N}{4m}\right)^{\frac{2m}{N}} \leq \sqrt{e}$, this implies that

$$\mathcal{J}_2 + \mathcal{J}_3 \leq \frac{e^{3/2} e^{c_9(1-\frac{4m}{N})} \sqrt{\Omega_m} N^{\frac{m}{2}} c_7^m c_{15}^{4m^2} m^{\frac{5m}{2}} \exp\left(-\frac{c_1 N(N-4m)}{1+\log m}\right)}{(1-4m/N)^{1-\frac{2m}{N}}}.$$

According to the notation (B.3), (B.4) and (B.11), since $c_6 = 4 \log c_{15}$, we have

$$e^{\Upsilon_1(m)} = e^{3/2} c_7^m c_{15}^{4m^2} m^{\frac{5m}{2}}.$$

Hence, according to formula (B.9), we have shown that for all $N > 4m$,

$$\begin{aligned} \mathcal{J}_2 + \mathcal{J}_3 &\leq \frac{e^{c_9(1-\frac{4m}{N})} \sqrt{\Omega_m} N^{\frac{m}{2}} \exp\left(\Upsilon_1(m) - \frac{c_1 N(N-4m)}{1+\log m}\right)}{(1-4m/N)^{1-\frac{2m}{N}}} \\ &= c_3 \sqrt{\Omega_m} N^{\frac{m}{2}} \Theta_{N,m}^1. \end{aligned} \quad (3.8)$$

Third, by using the estimate (2.27) from Proposition 2.8, we also have the bound

$$\mathcal{J}_1^2 = \int_{\Lambda_1 \leq \|\xi\| \leq \Lambda_2} |F_{n,m}(\xi)|^2 d\xi \leq \Omega_m \Lambda_2^{2m} \exp\left(2c_9 - \frac{2c_2 N^2}{\sqrt{m+1}(1+\log m)^{3/4}}\right)$$

and according to (2.25),

$$\Lambda_2^m \leq \sqrt{e} \frac{(8c_0)^{-m} N^m m^{m/2}}{(1+\log m)^{3m/4}},$$

where used that by (B.3), $0 < c_{10}(m) < 1$, $c_{11}(m) \geq 1$ and $(m+1)^m \leq \sqrt{e} m^{m/2}$ for all $m \geq 3$. According to (B.4), this shows that $\Lambda_2^m \leq e^{\Upsilon_2(m)}$, so that for all $m \geq 3$,

$$\begin{aligned} \mathcal{J}_1 &\leq e^{c_9} \sqrt{\Omega_m} N^m \exp\left(\Upsilon_2(m) - \frac{c_2 N^2}{\sqrt{m+1}(1+\log m)^{3/4}}\right) \\ &= c_3 \sqrt{\Omega_m} N^{\frac{m}{2}} \Theta_{N,m}^2, \end{aligned} \quad (3.9)$$

according to formula (B.8).

Finally, by collecting the estimates (3.4), (3.8) and (3.9), we deduce from the decomposition (3.3) that for any $m \geq 3$ and $N > 4m$,

$$\Delta_{n,m}^{(2)} \leq c_3 \sqrt{\Omega_m} N^{\frac{m}{2}} \left(\Theta_{N,m}^0 + \Theta_{N,m}^1 + \Theta_{N,m}^2 + \frac{c_3^{-1}}{\sqrt{m}} N^{-\frac{m}{2}} \exp\left(-\frac{N^2}{16(1+\log m)}\right) \right).$$

After replacing the last term by $\Theta_{N,m}^3$ according to (B.7), this completes the proof. \square

3.2 Proof of Theorem 1.3

First, the estimate (1.8) follows directly from Proposition 2.5 and the fact that $c_3 \leq 8$. Then, in order to prove the estimate (1.9), we need the following Gaussian tail-bound which is a straightforward consequence of Lemma 2.3.

Lemma 3.2 (Large deviation estimates). *For any $L > 0$, let $\square_L = [-\frac{L}{2}, \frac{L}{2}]^{2m}$. Then, we have for any $n, m \in \mathbb{Z}_+$,*

$$\mathbb{P}_n[\mathbf{X} \notin \square_L] \leq 4me^{-L^2/8}$$

and

$$\int_{\mathbb{R}^{2m} \setminus \square_L} \frac{e^{-\|x\|^2/2}}{(2\pi)^m} dx \leq \frac{8m}{\sqrt{2\pi}L} e^{-L^2/8}.$$

Proof. For any $k \geq 1$, it follows from Lemma 2.3 that for any $t \in \mathbb{R}$,

$$\mathbb{E}_n[e^{tX_{2k}}], \mathbb{E}_n[e^{tX_{2k-1}}] \leq e^{t^2/2}.$$

By Markov inequality, this implies that for any $k \geq 1$ and $t > 0$,

$$\mathbb{P}_n[|X_k| \geq L] \leq 2e^{-tL+t^2/2}.$$

Choosing $t = L$, we obtain

$$\mathbb{P}_n[|X_k| \geq L] \leq 2e^{-L^2/2}.$$

Hence, by a union bound, we obtain

$$\mathbb{P}_n[(X_1, \dots, X_{2m}) \notin \square_L] \leq \sum_{k \leq 2m} \mathbb{P}_n[|X_k| \geq L/2] \leq 4me^{-L^2/8}.$$

By a similar union bound, an analogous estimate holds in the Gaussian case. \square

Recall the definitions (1.6) and let us split

$$\begin{aligned} \Delta_{n,m}^{(1)} &= \left(\int_{\square_L} + \int_{\mathbb{R}^{2m} \setminus \square_L} \right) \left| \mathcal{P}_{n,m}(x) - \frac{e^{-\|x\|^2/2}}{(2\pi)^m} \right| dx \\ &\leq L^m \Delta_{n,m}^{(2)} + \int_{\mathbb{R}^{2m} \setminus \square_L} \left(\mathcal{P}_{n,m}(x) + \frac{e^{-\|x\|^2/2}}{(2\pi)^m} \right) dx, \end{aligned}$$

where we used the Cauchy–Schwartz inequality to bound the first integral. By Lemma 3.2, this implies that for any $L \geq 2\sqrt{3}$,

$$\Delta_{n,m}^{(1)} \leq L^m \Delta_{n,m}^{(2)} + 5me^{-L^2/8}, \quad (3.10)$$

We choose the parameter L which minimizes the RHS of (3.10), that is the (unique) solution of the equation:

$$\Delta_{n,m}^{(2)} = \frac{5}{4}L^{2-m}e^{-L^2/8}. \quad (3.11)$$

Since $m \geq 3$, the function $L \mapsto L^{2-m}e^{-L^2/8}$ is decreasing and it is bounded from below by $2^{2-m}m^{1-\frac{m}{2}}e^{-\frac{m}{2}}$ for $L \leq 2\sqrt{m}$, under the assumption that $\Delta_{n,m}^{(2)} \leq 5 \cdot 2^{-m}m^{1-\frac{m}{2}}e^{-\frac{m}{2}}$, the solution of the equation (3.11) satisfies

$$2\sqrt{m} \leq L \leq \sqrt{8 \log \Delta_{n,m}^{(2)-1}}. \quad (3.12)$$

Hence, by (3.10), this implies that

$$\Delta_{n,m}^{(1)} \leq L^m \left(1 + \frac{4m}{L^2} \right) \Delta_{n,m}^{(2)}.$$

Finally, using the conditions (3.12) for L , we conclude that

$$\Delta_{n,m}^{(1)} \leq 2(8 \log \Delta_{n,m}^{(2)-1})^{\frac{m}{2}} \Delta_{n,m}^{(2)}.$$

This completes the proof. \square

3.3 Proof of Theorem 1.1, Proposition 1.2 and Proposition 1.6

Throughout the proof, we fix an integer $M \geq 3$. By (B.17), we have $\sqrt{\Omega_m} \leq \frac{m^{-\frac{m}{2}} (e\pi)^{\frac{m}{2}}}{(2\pi)^{1/4}}$ and it follows from the estimate (1.8) that the condition $\Delta_{n,m}^{(2)} \leq 5 \cdot 2^{-m} m^{1-\frac{m}{2}} e^{-\frac{m}{2}}$ from Theorem 1.3 is satisfied if

$$\Theta_{n,m} \leq \frac{m}{2} c_{16}^{-m} N^{-\frac{m}{2}}. \quad (3.13)$$

Then we immediately deduce from the estimates of Proposition B.3 that for all $m \geq M$ and $N \geq c(M)m\sqrt{1+\log m}$,

$$\Theta_{n,m} \leq (1+\epsilon)\Theta_{N,m}^0 \leq N^{-\frac{m}{2}} \frac{\exp(-12m(\log m - 0.26))}{c_{16}^m}.$$

This shows that under the hypothesis of Theorem 1.1, the condition (3.13) holds. Accordingly, by (1.9), we obtain

$$\begin{aligned} \Delta_{n,m}^{(1)} &\leq 2(8 \log \Delta_{n,m}^{(2)})^{\frac{m}{2}} \Delta_{n,m}^{(2)} \\ &\leq 2c_3 \sqrt{\Omega_m} (8N \log (c_3 \sqrt{\Omega_m} N^{\frac{m}{2}} \Theta_{N,m})^{-1})^{\frac{m}{2}} \Theta_{N,m}, \end{aligned}$$

where we used that the function $x \mapsto x(\log x^{-1})^{\frac{m}{2}}$ is non-decreasing for $x \in [0, e^{-\frac{m}{2}}]$ as well as Proposition 2.5 to get the second bound. By (B.17), we also have $\Omega_m N^m \geq \frac{e^m \pi^m (N/m)^m}{3\sqrt{m}}$, so that according to formula (B.7), we obtain the (crude) lower-bound

$$c_3 \sqrt{\Omega_m} N^{\frac{m}{2}} \Theta_{N,m}^0 \geq N^{-N}.$$

This implies that if $m \geq M$ and $N \geq c(M)m\sqrt{1+\log m}$, then

$$\Delta_{n,m}^{(1)} \leq 2c_3 \sqrt{\Omega_m} (N\sqrt{8\log N})^m \Theta_{N,m}. \quad (3.14)$$

Using Corollary B.3 once more, we obtain

$$\Delta_{n,m}^{(1)} \leq 2c_3(1+\epsilon)\sqrt{\Omega_m} (N\sqrt{8\log N})^m \Theta_{N,m}^0.$$

Since $2c_3(1+\epsilon) \leq 16$, by (B.7) and (1.7), we conclude that for all $m \geq M$ which satisfies the condition $N \geq c(M)m\sqrt{1+\log m}$,

$$d_{\text{TV}}(\mathbf{X}, \mathbf{G}) \leq \Delta_{n,m}^{(1)} \leq 16\sqrt{\Omega_m} m^3 4^m e^{\frac{m^2}{4N}} (N\sqrt{\log N})^m e^{\frac{N}{2}} \frac{(1+\log m)^N}{\sqrt{n} \Gamma(N+1)}. \quad (3.15)$$

This completes the proof of Theorem 1.1. \square

Now, let us choose $m = \lfloor n^\alpha \rfloor$ with $0 < \alpha < 1/2$ and let us assume that $m \geq 17$. By (3.15), this implies that for all integer $n \in \mathbb{N}$ such that $n^{1-2\alpha} \geq 20.4\sqrt{\log n}$,

$$\Delta_{n,m}^{(1)} \leq 16\sqrt{\Omega_m} n^{3\alpha-1} 4^m e^{\frac{n^{3\alpha-1}}{4}} (N\sqrt{\log N})^{n^\alpha} \frac{e^{\frac{N}{2}} (\log N)^N}{\Gamma(N+1)}$$

where we used that $N \geq e^2 m$ and that $c(17) = 28.8 \leq 20.4\sqrt{2}$ – see the Table (B.22). First, observe that by (B.17), it holds for all $m \in \mathbb{N}$,

$$\sqrt{\Omega_m} 4^m \leq \frac{(4\sqrt{\pi e})^m}{(2\pi)^{\frac{1}{4}} \sqrt{m}} m^{-\frac{m}{2}} \leq \frac{e^{8\pi}}{\sqrt{m\sqrt{2\pi}}}, \quad (3.16)$$

where we used that the $\max_{m \geq 0} \{(16\pi e)^m m^{-m}\} = \exp(16\pi)$. Second, let us observe that the function $\frac{e^{\frac{N}{2}} (\log N)^N}{\Gamma(N+1)}$ is decreasing for $N \geq e^2$ so that by (B.17),

$$\frac{e^{\frac{N}{2}} (\log N)^N}{\Gamma(N+1)} \leq \frac{n^{\frac{\alpha-1}{2}}}{\sqrt{2\pi}} \exp\left(-n^{1-\alpha} \log(n^{1-\alpha}) \left(1 - \frac{\log(\log \sqrt{n}) + 3/2}{\log(n^{1-\alpha})}\right)\right).$$

Since $1/\sqrt{m} \leq n^{-\frac{\alpha}{2}} \sqrt{18/17}$ for $n \geq 18^{\alpha^{-1}}$, these estimates imply that

$$\Delta_{n,m}^{(1)} \leq C n^{3\alpha - \frac{3}{2}} e^{\frac{n^{3\alpha-1}}{4}} (N\sqrt{\log N})^{n^\alpha} \exp\left(-n^{1-\alpha} \log(n^{1-\alpha}) \left(1 - \frac{\log(\log \sqrt{n}) + 3/2}{\log(n^{1-\alpha})}\right)\right),$$

with $C = \frac{18e^{8\pi}}{(2\pi)^{\frac{3}{4}}}$. Now, let us also observe that $N \leq e^{0.0572} n^{1-\alpha}$ for $n \geq 18^{\alpha^{-1}}$, so that

$$(N\sqrt{\log N})^{n^\alpha} \leq \exp\left(n^\alpha \log(n^{1-\alpha}) \left(1 + \frac{\log \log n + 0.1144}{2 \log(n^{1-\alpha})}\right)\right).$$

This shows that if $n \geq 18^{\alpha^{-1}}$ (so that $m \geq 17$) and $n^{1-2\alpha} \geq 20.4\sqrt{\log n}$,

$$\Delta_{n,m}^{(1)} \leq C n^{3\alpha - \frac{3}{2}} \exp\left(- (1 - \epsilon_n) n^{1-\alpha} \log(n^{1-\alpha})\right)$$

with

$$\epsilon_n := \frac{\log \log(\sqrt{n}) + 3/2}{\log(n^{1-\alpha})} + n^{-(1-2\alpha)} \left(1 + \frac{\log \log n + 0.1144}{2 \log(n^{1-\alpha})}\right) + \frac{n^{-2(1-2\alpha)}}{4 \log(n^{1-\alpha})} \quad (3.17)$$

$$\leq \frac{2(\log \log n + 0.8069)}{\log n} + \frac{0.0649}{\sqrt{\log n}} + \frac{0.0012}{(\log n)^2}, \quad (3.18)$$

where we have used that $n^{1-2\alpha} \geq 20.4\sqrt{\log n}$, $\alpha \leq 1/2$ and the numerical bound $1 + \frac{\log \log n + 0.1144}{\log n} \leq 20.4 \cdot 0.0649$ for all $n \geq 18^2$ to obtain the estimate (3.18). We also deduce from (3.18) that $\epsilon_n \leq 1 - 87 \cdot 10^{-3}$ for all $n \geq 18^2$. Since $d_{\text{TV}}(\mathbf{X}, \mathbf{G}) \leq \Delta_{n,m}^{(1)}$, this completes the proof of Proposition 1.2. \square

We now turn to the proof of Proposition 1.6. Let us choose $m = \left\lfloor \sqrt{\frac{n}{41.5\sqrt{\log n}}} \right\rfloor$ and suppose that $n \geq 4322$ so that $m \geq 6$ and we can use the estimate (3.15) – we have $c(6) = 58.66 \leq 41.5\sqrt{2}$ according to the Table (B.22). As $N = \frac{n}{m} \geq e^2 m$, this implies that

$$\Delta_{n,m}^{(1)} \leq 16\sqrt{\Omega_m} m^{\frac{7}{2}} 4^m e^{\frac{m^3}{4n}} (N\sqrt{\log N})^m \frac{e^{\frac{3}{2}N} (\log N)^N}{\sqrt{2\pi n N^N}}.$$

Moreover, we verify that as $N \geq \sqrt{41.5n\sqrt{\log n}}$,

$$(\log N)^N N^{m-N} e^{\frac{3}{2}N} \leq \exp\left(-\frac{\sqrt{41.5}}{2} \sqrt{n} (\log n)^{5/4} \left(1 - \frac{1}{41.5\sqrt{\log n}} - \frac{2 \log(\log \sqrt{n}) + 3}{\log n}\right)\right)$$

and using the estimate (3.16), this shows that

$$\Delta_{n,m}^{(1)} \leq \frac{16e^{8\pi} m^3}{(2\pi)^{\frac{3}{4}} n} e^{\frac{m^3}{4n}} (\log N)^{\frac{m}{2}} \exp\left(-\frac{\sqrt{41.5}}{2} \sqrt{n} (\log n)^{5/4} \left(1 - \frac{1}{41.5\sqrt{\log n}} - \frac{2 \log(\log \sqrt{n}) + 3}{\log n}\right)\right).$$

Moreover, since $(\log N)^{\frac{m}{2}} e^{\frac{m^3}{4n}} \leq \exp\left(\frac{\sqrt{n}}{4(41.5\sqrt{\log n})^{3/2}} + \frac{1}{2} \sqrt{\frac{n}{41.5\sqrt{\log n}}} \log \log n\right)$, we obtain

$$\Delta_{n,m}^{(1)} \leq \frac{16e^{8\pi} \sqrt{n}}{(2\pi \log n)^{\frac{3}{4}} (41.5)^{\frac{3}{2}}} \exp\left(-\frac{\sqrt{41.5}}{2} \sqrt{n} (\log n)^{5/4} (1 - \epsilon_n)\right).$$

where

$$\epsilon_n = \frac{1}{41.5\sqrt{\log n}} + \frac{3 - 2\log 2 + 2\log \log n}{\log n} + \frac{\log \log n}{41.5(\log n)^{\frac{3}{2}}} + \frac{1/2}{(41.5 \log n)^2}$$

We verify numerically that $\epsilon_n \leq 0.711$ for all $n \geq 4322$. In particular, this implies that

$$\Delta_{n,m}^{(1)} \leq \sqrt{n} \exp(19.4 - 0.93\sqrt{n}(\log n)^{5/4}).$$

Since $\Delta_{n,m}^{(1)}$ is non-decreasing in $m \in \mathbb{N}$, this completes the proof of Proposition 1.6. \square

4 Gaussian approximation: Proof of Proposition 2.6

Recall that $F_{n,m}$ denotes the characteristic function of the random vector \mathbf{X} and that it is given by formula (2.10). In particular, it holds for any $\xi \in \mathbb{R}^{2m}$,

$$|e^{-\|\xi\|^2/2} - F_{n,m}(\xi)|^2 = e^{-\|\xi\|^2} |1 - \det[\mathbb{I} - K_{\text{ig}} Q_n]|^2, \quad (4.1)$$

where g is a trigonometric polynomial (2.2), Q_n is the orthogonal projection with kernel $\text{span}(e_1, \dots, e_{n-1})$ and according to formula (2.8),

$$K_{\text{ig}} = H_+(e^{2\Im g^+}) H_-(e^{2\Im g^-}). \quad (4.2)$$

Recall that the operator K_{ig} is trace-class, but observe that it is not self-adjoint since by (2.7), $K_{\text{ig}}^* = H_+(e^{2\Im g^-}) H_-(e^{2\Im g^+})$ with $\Im g^- = -\Im g^+$ because the function g is real-valued. As we explained in Section 2.4, in order to prove Proposition 2.6, we provide estimates for the Fredholm determinant on the RHS of (4.1) in the regime where $\|\xi\| \ll N$ in order to guarantee that the Schatten norm $\|K_{\text{ig}} Q_n\|_{\mathcal{S}_1}$ remains small.

The first step of the proof consists in obtaining a priori estimates on Fourier coefficients of the functions $e^{2\Im g^\pm}$.

Lemma 4.1. *Fix $m \in \mathbb{N}$ and $\xi \in \mathbb{R}^{2m}$. Let $\rho = \sqrt{\frac{1+\log m}{2}} \|\xi\|$. We have for all integers $k > 2m\rho$,*

$$\left| \widehat{(e^{\pm 2\Im g^\pm})}_k \right| \leq 2e^\rho \frac{\rho^{\lceil k/m \rceil}}{\lceil k/m \rceil!}.$$

Proof. Let us define $\phi_M(w) = \sum_{k=0}^M \frac{w^k}{k!}$ for $M \geq 1$. Since $g^+(\theta) = \sum_{k=1}^m \frac{\zeta_k}{\sqrt{2k}} e^{ik\theta}$ and $g^- = \overline{g^+}$, we have for all integers $k > Mm$,

$$\int_{\mathbb{T}} \phi_M(-ig^+(\theta)) e^{ig^-(\theta) - ik\theta} \frac{d\theta}{2\pi} = 0.$$

This implies that any $k > Mm$,

$$\begin{aligned} \left| \widehat{(e^{2\Im g^+})}_k \right| &= \left| \int_{\mathbb{T}} e^{-ig^+(\theta) + ig^-(\theta) - ik\theta} \frac{d\theta}{2\pi} \right| \\ &\leq \int_{\mathbb{T}} \left| e^{-ig^+(\theta)} - \phi_M(-ig^+(\theta)) \right| e^{-\Im g^-(\theta)} \frac{d\theta}{2\pi}. \end{aligned} \quad (4.3)$$

Now, let us observe that for any $|w| \leq M/2$,

$$\begin{aligned} |e^w - \phi_M(w)| &\leq \frac{|w|^{M+1}}{(M+1)!} \sum_{j \geq 0} \left(\frac{|w|}{M+2} \right)^j \\ &\leq 2 \frac{|w|^{M+1}}{(M+1)!}. \end{aligned} \quad (4.4)$$

Moreover, by (2.2) and since $\sum_{k=1}^m |\zeta_k|^2 = \|\xi\|^2$, we also have

$$\|g^+\|_\infty \leq \sum_{k=1}^m \frac{|\zeta_k|}{\sqrt{2k}} \leq \rho = \sqrt{\frac{1 + \log m}{2}} \|\xi\|, \quad (4.5)$$

where we used that $\sum_{k=1}^m k^{-1} \leq 1 + \log m$ for any $m \geq 1$ and the Cauchy–Schwartz inequality. Then, using the estimates (4.3), (4.4) and (4.5), we obtain if both $M \geq 2\rho$ and $k > Mm$,

$$\left| \widehat{(e^{2\Im g^+})}_k \right| \leq 2e^\rho \frac{\rho^{M+1}}{(M+1)!}.$$

By choosing $M = \lfloor k/m \rfloor$, this implies the claim. Indeed, by the same argument, we obtain the same bound for $\left| \widehat{(e^{-2\Im g^+})}_k \right|$. \square

Now, let us use these estimates to bound the Schatten norm $\|K_{\text{ig}} Q_n\|_{\mathcal{J}_1}$.

Lemma 4.2. *Fix $m \in \mathbb{N}$, $\xi \in \mathbb{R}^{2m}$ and let $\rho = \sqrt{\frac{1+\log m}{2}} \|\xi\|$. If we assume that $N = \frac{n}{m} \geq c_*^{-1} \rho$ with $c_* < \frac{1}{2}$, then*

$$\|K_{\text{ig}} Q_n\|_{\mathcal{J}_1} \leq \frac{4m^2 e^{2\rho}}{(1 - c_*^2)^2} \frac{\rho^{2N}}{\Gamma(N+1)^2}.$$

Proof. Let us recall that the operators $H_\pm(e^{2\Im g^\pm})$ are Hilbert–Schmidt and that by (2.7), we have

$$\|Q_n H_\pm(e^{2\Im g^\pm})\|_{\mathcal{J}_2}^2 \leq \sum_{k \geq n} (k - n + 1) \left| \widehat{(e^{2\Im g^\pm})}_k \right|^2.$$

Moreover, by formula (4.2) and the Cauchy–Schwartz inequality (for the Schatten norms), since Q_n is a projection, we have

$$\|K_{\text{ig}} Q_n\|_{\mathcal{J}_1} \leq \|H_+(e^{2\Im g^+}) Q_n\|_{\mathcal{J}_2} \|H_-(e^{2\Im g^-}) Q_n\|_{\mathcal{J}_2}.$$

Using the estimates from Lemma 4.1, this implies that if the dimension $n > 2m\rho$, then

$$\|K_{\text{ig}} Q_n\|_{\mathcal{J}_1} \leq 4e^{2\rho} \sum_{k \geq n} (k - n + 1) \frac{\rho^{2\lceil k/m \rceil}}{(\lceil k/m \rceil!)^2}. \quad (4.6)$$

Under the condition $N = \frac{n}{m} \geq c_*^{-1} \rho$, since $j! \geq \Gamma(N+1) N^{j-N}$ for all $j \geq N$, we obtain

$$\begin{aligned} \sum_{k \geq n} (k - n + 1) \frac{\rho^{2\lceil k/m \rceil}}{(\lceil k/m \rceil!)^2} &\leq m^2 \sum_{j \geq N} (j + 1 - N) \frac{\rho^{2j}}{(j!)^2} \\ &\leq m^2 \frac{\rho^{2N}}{\Gamma(N+1)^2} \sum_{j \geq 0} (j + 1) \left(\frac{\rho}{N} \right)^{2j} \\ &\leq (1 - c_*^2)^{-2} m^2 \frac{\rho^{2N}}{\Gamma(N+1)^2}. \end{aligned}$$

Note that for the last bound, it suffices that $c_* < 1$. However, we impose that $c_* < \frac{1}{2}$ to guarantee that $n > 2m\rho$. Then, by combining the previous estimate with (4.6), this completes the proof. \square

We are now ready to finish the proof of Proposition 2.6. First, let us observe that by Lemma 4.2 and using formula (B.17) for the Γ function, we obtain that if $N \geq c_*^{-1}\rho$,

$$\|K_{\text{ig}}Q_n\|_{\mathcal{J}_1} \leq \frac{2/\pi}{(1-c_*^2)^2} m^2 e^{2\rho} \frac{(\rho e)^{2N}}{N^{2N+1}} \leq \frac{2/\pi}{(1-c_*^2)^2} \frac{m^2}{N} (c_* e^{c_*+1})^{2N}. \quad (4.7)$$

If we choose $c_* = 1/4$, then $c_* e^{c_*+1} \leq 0.873$ so that the RHS of (4.7) is very small for large N . Actually, in the regime where $N > 4m$ (in particular when $N \geq 13$), this implies that

$$\|K_{\text{ig}}Q_n\|_{\mathcal{J}_1} \leq \frac{32}{225 \cdot \pi} N (c_* e^{c_*+1})^{2N} \leq \frac{416}{225 \cdot \pi} (0.873)^{26} \leq \log(2.766) - 1,$$

where we obtained the last two bounds numerically. Hence, using the inequality (2.19) from (Simon, 2005b, Theorem 3.4), we deduce from Lemma 4.2 with $c_* = 1/4$ and the previous estimate that if $N \geq 4(\rho \vee m)$,

$$|1 - \det[\mathbf{I} - K_{\text{ig}}Q_n]|^2 \leq \|K_{\text{ig}}Q_n\|_{\mathcal{J}_1}^2 e^{2(1+\|K_{\text{ig}}Q_n\|_{\mathcal{J}_1})} \leq c_8^2 m^4 e^{4\rho} \frac{\rho^{4N}}{\Gamma(N+1)^4},$$

where $c_8 = 2.766 \frac{4}{(1-c_*^2)^2}$ according to (B.3). If we combine this estimate with formula (4.1) and replace $\rho = \sqrt{(1 + \log m)/2} \|\xi\|$, this implies that for any $N \geq 4m$ and all $\|\xi\| \leq \Lambda_1 = \frac{N}{4\sqrt{(1+\log m)/2}}$,

$$|e^{-\|\xi\|^2/2} - F_{n,m}(\xi)|^2 \leq c_8^2 m^4 e^{4\rho} \frac{\rho^{4N}}{\Gamma(N+1)^4} e^{-\|\xi\|^2}.$$

This completes the proof. \square

5 Tail bound for large $\|\xi\|$: Proof of Proposition 2.7

Recall that the function g is given by (2.2) and let us observe that by choosing $h = g'$ in Lemma 2.4, we obtain the following bound.

Proposition 5.1. *Fix $m, n \in \mathbb{N}$ and let $N = \frac{n}{m}$. For any $\eta > 0$ and any $\xi \in \mathbb{R}^{2m}$, we have*

$$|F_{n,m}(\xi)| \leq \exp\left(c_{20}\left(n + \frac{2}{\pi^2}\right)\right) \mathbb{E}_n \left[e^{-\gamma \sum_{j=1}^n g'(\theta_j)^2} \right],$$

where $\gamma = \frac{\eta}{\sqrt{nm(m+1)\|\xi\|}} \left(1 - \frac{\eta^2 c_{21}}{n}\right)$ and $c_{20} = \frac{\pi^2 \eta^2}{8}$.

In order to prove Proposition 5.1, we need the following basic estimate which is proved in the Appendix A.3.

Lemma 5.2. *For any $y \in [-1, 1]$, $y \neq 0$ and $x \in \mathbb{R}$, we have*

$$1 + \left(\frac{\sinh(x)}{y}\right)^2 \leq \exp\left(\frac{x}{y}\right)^2.$$

Proof. We apply Lemma 2.4 with $h = g'$ and $\nu = \frac{\eta\sqrt{n}}{m(m+1)\|\xi\|}$ where $\eta > 0$. We obtain

$$|F_{n,m}(\xi)| \leq \mathbb{E}_n \left[\prod_{i < j} \left| \frac{\sin\left(\frac{\theta_i - \theta_j}{2} + i\nu \frac{g'(\theta_i) - g'(\theta_j)}{2n}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)} \right|^2 \prod_{j=1}^n \left| 1 + i \frac{\nu}{n} g''(\theta_j) \right| e^{-\Im g\left(\theta_j + i \frac{\nu}{n} g'(\theta_j)\right)} \right]. \quad (5.1)$$

Moreover, by the Cauchy–Schwartz, we have

$$\|g'\|_\infty \leq \sum_{k=1}^m \sqrt{2k} |\zeta_k| \leq \sqrt{2 \sum_{k=1}^m |\zeta_k|^2 \sum_{k=1}^m k} = \sqrt{m(m+1)} \|\xi\|. \quad (5.2)$$

Observe that with these choices, we have $\frac{\nu}{n} \|g'\|_\infty \leq \frac{\eta/m}{\sqrt{n(1+1/m)}}$, so that by Taylor’s theorem, since g is real–valued, we have for $j \in \{1, \dots, n\}$,

$$\left| \Im g\left(\theta_j + i \frac{\nu}{n} g'(\theta_j)\right) - \frac{\nu}{n} g'(\theta_j)^2 \right| \leq \frac{1}{6} \left| \frac{\nu}{n} g'(\theta_j) \right|^3 \sup_{|\Re z| \leq \pi, |\Im z| \leq \frac{\eta/m}{\sqrt{n(1+1/m)}}} |g'''(z)|.$$

We also have

$$|g'''(z)| \leq \sum_{|k| \leq m} |\zeta_k| \frac{|k|^{5/2}}{\sqrt{2}} e^{k|\Im z|} \quad \text{so that} \quad \sup_{|\Re z| \leq \pi, |\Im z| \leq \frac{\eta/m}{\sqrt{n(1+1/m)}}} |g'''(z)| \leq 6c_{21} (m(m+1))^{3/2} \|\xi\|,$$

where $c_{21} = \frac{\exp(\eta/\sqrt{n(1+1/m)})}{6\sqrt{3}}$ and we used that $\sum_{k=1}^m k^5 \leq \frac{m^3(m+1)^3}{6}$. Then using the estimate (5.2), the previous bounds imply that

$$\begin{aligned} \left| \Im g\left(\theta_j + i \frac{\nu}{n} g'(\theta_j)\right) - \frac{\nu}{n} g'(\theta_j)^2 \right| &\leq c_{21} \frac{\nu^3 m^2 (m+1)^2 \|\xi\|^2}{n^3} g'(\theta_j)^2 \\ &= \frac{\nu}{n} \frac{\eta^2 c_{21}}{n} g'(\theta_j)^2, \end{aligned}$$

where we used our choice for ν . This shows that

$$\prod_{j=1}^n e^{-\Im g\left(\theta_j + i \frac{\nu}{n} g'(\theta_j)\right)} \leq \exp\left(-\frac{\nu}{n} \left(1 - \frac{\eta^2 c_{21}}{n}\right) \sum_{j=1}^n g'(\theta_j)^2\right). \quad (5.3)$$

Moreover, by Lemma 5.2 and since by convexity, $\sin(u/2) \geq u/\pi$ for all $u \in [0, \pi]$, we obtain for any $u \in [-\pi, \pi]$ and $\alpha > 0$,

$$1 + \left(\frac{\sinh(\alpha u/2)}{\sin(u/2)}\right)^2 \leq \exp\left(\frac{\alpha u}{2 \sin(u/2)}\right)^2 \leq \exp\left(\frac{\pi \alpha}{2}\right)^2.$$

This estimate implies that for all $i, j \in \{1, \dots, n\}$,

$$\begin{aligned} \left| \frac{\sin\left(\frac{\theta_i - \theta_j}{2} + i \nu \frac{g'(\theta_i) - g'(\theta_j)}{2n}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)} \right|^2 &= 1 + \left(\frac{\sinh\left(\nu \frac{g'(\theta_i) - g'(\theta_j)}{2n}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)}\right)^2 \\ &\leq 1 + \left(\frac{\sinh\left(\frac{\nu \|g''\|_\infty (\theta_i - \theta_j)}{2}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)}\right)^2 \\ &\leq \exp\left(\frac{\nu \pi \|g''\|_\infty}{2n}\right)^2. \end{aligned}$$

Moreover, by the Cauchy–Schwartz inequality, we have

$$\|g''\|_\infty \leq \sum_{|k| \leq m} \frac{k^{3/2}}{\sqrt{2}} |\zeta_k| \leq \sqrt{\sum_{k=1}^m k^3 \sum_{|k| \leq m} |\zeta_k|^2} = \frac{m(m+1)}{\sqrt{2}} \|\xi\|.$$

Hence, this shows that

$$\prod_{1 \leq i < j \leq n} \left| \frac{\sin\left(\frac{\theta_i - \theta_j}{2} + i\nu \frac{g'(\theta_i) - g'(\theta_j)}{2n}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)} \right|^2 \leq \exp\left(\frac{\nu m(m+1)\|\xi\|}{2\sqrt{2}/\pi}\right)^2 = e^{c_{20}n}, \quad (5.4)$$

where we used the definition of ν and set $c_{20} = \frac{\pi^2 \eta^2}{8}$. Similarly, we have

$$\begin{aligned} \prod_{j=1}^n |1 + i\frac{\nu}{n}g''(\theta_j)| &\leq \left(1 + \frac{\nu^2 \|g''\|_\infty^2}{n^2}\right)^{n/2} \\ &\leq \exp\left(\frac{(\nu m(m+1)\|\xi\|)^2}{4n}\right) = \exp\left(\frac{2c_{20}}{\pi^2}\right), \end{aligned} \quad (5.5)$$

where we used that $1 + x \leq e^x$ for all $x \in \mathbb{R}$ to obtain the second estimate. By combining the estimates (5.3), (5.4), (5.5) with (5.1), we obtain that for all $\xi \in \mathbb{R}^{2m}$,

$$|F_{n,m}(\xi)| \leq \mathbb{E}_n \left[\exp\left(c_{20}\left(n + \frac{2}{\pi^2}\right) - \gamma \sum_{j=1}^n g'(\theta_j)^2\right) \right],$$

where $\gamma = \frac{\nu}{n} \left(1 - \frac{\eta^2 c_{21}}{n}\right) = \frac{\eta}{\sqrt{nm(m+1)}\|\xi\|} \left(1 - \eta^2 \frac{\exp\left(\frac{\eta}{\sqrt{n(1+1/m)}}\right)}{6\sqrt{3n}}\right)$. \square

Thus, in order to estimate $|F_{n,m}(\xi)|$ using Proposition 5.1, we need a bound for $\mathbb{E}_n[e^{-\gamma \sum_{j=1}^n g'(\theta_j)^2}]$. Let us point out that in the regime where $\|\xi\|$ is large compared with N , we cannot use the bound from Lemma 2.3 to estimate this quantity. Indeed, we have $\|g'\|_{L^2}^2 \geq \|\xi\|^2$, while our basic estimate for $\mathcal{A}(g'^2)$ is of the form $\mathcal{A}(g'^2) \leq cm^5 \|\xi\|^4$ for a numerical constant $c > 0$. Then, by optimizing over all $\gamma > 0$, we would obtain

$$\mathbb{E}_n [e^{-\gamma \sum_{j=1}^n g'(\theta_j)^2}] \leq \exp\left(-\gamma n \|g'\|_{L^2}^2 + \gamma^2 \mathcal{A}(g'^2)\right) \leq \exp\left(-\frac{N^2}{4cm^3}\right).$$

This estimate is similar to those from Proposition 2.8 but it not as good for large $m \in \mathbb{N}$. More importantly, it does not yield any decay as $\|\xi\| \rightarrow +\infty$. So, instead of Lemma 2.3, we will use the bound (2.23) which follows from the next Lemma.

Lemma 5.3. *For any function $f : \mathbb{T} \rightarrow \mathbb{R}$ such that e^{-f} is integrable, we have for any $n \geq 2$,*

$$\mathbb{E}_n \left[e^{-\sum_{j=1}^n f(\theta_j)} \right] \leq \frac{e^n}{\sqrt{2\pi n}} \left(\int_{\mathbb{T}} e^{-f(\theta)} \frac{d\theta}{2\pi} \right)^n. \quad (5.6)$$

The proof of Lemma 5.3 is given in the appendix (Section A.4) and it relies on the fact that the configurations which minimize the *energy* associated with the probability measure \mathbb{P}_n are uniformly distributed on \mathbb{T} (like the vertices of a regular n -gon) so that we know explicitly the minimal energy as well as the partition function.

To complete the proof of Proposition 2.7, we also need (Chahkiev, 2008, Lemma 2) in order to give an estimate for the integral on the RHS of (5.6).

Lemma 5.4 ((Chahkiev, 2008)). *Let $f : \mathbb{T} \rightarrow \mathbb{R}$ be a trigonometric polynomial of degree $m \in \mathbb{N}$ and let $\|f\|_{L^2} = \sqrt{\int_{\mathbb{T}} f(\theta)^2 d\mu}$ where $d\mu = \frac{d\theta}{2\pi}$ denotes the uniform measure on \mathbb{T} . If we let $\mathcal{T}_\lambda = \{\theta \in \mathbb{T} : |f(\theta)| \leq \lambda\}$, then we have for any $\lambda > 0$,*

$$\mu(\mathcal{T}_\lambda) \leq 2e \left(\frac{\lambda}{\sqrt{2}\|f\|_{L^2}} \right)^{1/2m}.$$

From Lemma 5.4, we deduce that for any trigonometric polynomial $f : \mathbb{T} \rightarrow \mathbb{R}$ of degree at most $m \in \mathbb{N}$, we have

$$\begin{aligned} \int_{\mathbb{T}} e^{-f(\theta)^2} \frac{d\theta}{2\pi} &= \int_{\mathbb{T}} \left(\int_0^{+\infty} e^{-\lambda} \mathbf{1}_{\{|f(\theta)| \leq \sqrt{\lambda}\}} d\lambda \right) \mu(d\theta) \\ &= \int_0^{+\infty} e^{-\lambda} \mu(\mathcal{T}_{\sqrt{\lambda}}) d\lambda \\ &\leq 2e \int_0^{+\infty} e^{-\lambda} \left(\frac{\lambda}{2\|f\|_{L^2}^2} \right)^{1/4m} d\lambda \\ &\leq \frac{2e}{(2\|f\|_{L^2}^2)^{1/4m}}, \end{aligned}$$

where we used that $\Gamma(1 + 1/4m) = \int_0^{+\infty} e^{-\lambda} \lambda^{1/4m} d\lambda \leq 1$ for any $m \in \mathbb{N}$ in the last step. Hence, by combining this estimate with (5.6), we obtain the following general bound.

Proposition 5.5. *Let $f : \mathbb{T} \rightarrow \mathbb{R}$ be a trigonometric polynomial for degree $m \in \mathbb{N}$. We have for any $n \geq 2$,*

$$\mathbb{E}_n \left[e^{-\sum_{j=1}^n f(\theta_j)^2} \right] \leq \frac{c_{15}^n}{\sqrt{2\pi n} (2\|f\|_{L^2}^2)^{N/4}},$$

where $N = \frac{n}{m}$, $c_{15} = 2e^2$ and $\|f\|_{L^2} = \sqrt{\int_{\mathbb{T}} f(\theta)^2 \frac{d\theta}{2\pi}}$.

We are now ready to complete the proof of Proposition 2.7. By combining the estimates from Proposition 5.1 and Proposition 5.5 with $f = \sqrt{\gamma}g'$ which is a real-valued² trigonometric polynomial of degree $m \in \mathbb{N}$, we obtain that for any $n \geq 2$ and any $\eta \in (0, 1]$,

$$\begin{aligned} |F_{n,m}(\xi)| &\leq \exp\left(c_{20}\left(n + \frac{2}{\pi^2}\right)\right) \frac{c_{15}^n}{\sqrt{2\pi n} (2\gamma\|g'\|_{L^2}^2)^{N/4}} \\ &\leq \frac{1}{\sqrt{2\pi n}} \left(\frac{e^{m\eta^2(\frac{\pi^2}{2} + \frac{1}{n})}}{2\gamma\|\xi\|} \right)^{N/4} \frac{c_{15}^n}{\|\xi\|^{N/4}}, \end{aligned} \quad (5.7)$$

where we used that by definition we have $\|g'\|_{L^2}^2 = \sum_{k=1}^m k|\zeta_k|^2 \geq \|\xi\|^2$ and we replaced $c_{20} = \frac{\pi^2\eta^2}{8}$. We still have the freedom to choose the parameter $\eta \in (0, 1]$ in the estimate (5.7) and we choose it in such a way to minimize $\eta^{-1}e^{m\eta^2\frac{\pi^2}{2}}$. That is, we choose $\eta = \frac{1/\pi}{\sqrt{m}}$ and since $2\gamma\|\xi\| = \frac{2\eta}{\sqrt{nm(m+1)}} \left(1 - \frac{\eta^2 c_{21}}{n}\right)$, this implies that

$$|F_{n,m}(\xi)| \leq \left(\frac{\pi e^{\frac{1}{2}(1 + \frac{2}{\pi^2 n})} \sqrt{nm}^{3/2} (m+1)}{2\left(1 - \frac{c_{21}}{nm}\right)} \right)^{N/4} \frac{c_{15}^n}{\|\xi\|^{N/4}}.$$

Finally, let us observe that in the regime where $n \geq 4m^2$ (note that it is the only place where we use this condition), this implies that

$$|F_{n,m}(\xi)|^2 \leq \Upsilon_3(m)^{N/2} \frac{c_{15}^{2n} n^{N/4}}{\|\xi\|^{N/2}}.$$

where $\Upsilon_3(m) = \frac{\pi m^{3/2} (m+1) e^{\frac{1}{2}\left(1 + \frac{1/2}{(\pi m)^2}\right)}}{2\left(1 - \frac{c_{21}}{4\pi^2 m^3}\right)}$ according to (B.11). This completes the proof. \square

²We verify that for any $n, m \in \mathbb{N}$ and $\eta \in (0, 1]$, $\gamma > 0$.

6 Intermediate regime

The goal of this section is to prove Proposition 2.8. Recall that the polynomial g is given by (2.2) and that $h = -\mathcal{U}g$ is the Hilbert transform of the function $-g$ – see (2.13). We will make use of the following basic estimates. We have for any $\xi \in \mathbb{R}^{2m}$,

$$\|g\|_\infty, \|h\|_\infty \leq \sqrt{2(1 + \log m)} \|\xi\|. \quad (6.1)$$

Similarly, for any $\xi \in \mathbb{R}^{2m}$ and any integer $\kappa \geq 0$,

$$\|h^{(\kappa+1)}\|_\infty \leq \sum_{k \leq m} |\zeta_k| k^\kappa \sqrt{2k} \leq C_\kappa \|\xi\| (m(m+1))^{\frac{\kappa+1}{2}}, \quad (6.2)$$

with $C_0 = 1$, $C_1 = 1/\sqrt{2}$, $C_2 = 1/\sqrt{3}$ and

$$\|h^{(\kappa+1)}\|_{L^2} = \sqrt{\sum_{k \leq m} |\zeta_k|^2 k^{2\kappa+1}} \leq m^{\kappa+1/2} \|\xi\|. \quad (6.3)$$

We will also make use of Lemma 2.9 which is proved in Section 6.2 and we fix (throughout this section) the parameter

$$\nu = \frac{\nu_* N}{\sqrt{m+1}(1 + \log m)^{1/4} \|\xi\|}, \quad (6.4)$$

where $N = \frac{n}{m}$ and $0 < \nu_* \leq c_0$ as in (B.1). This last condition is necessary for our proof of Proposition 6.2 below and we will optimize over the parameter ν_* in the proof of Proposition 2.8 which is given in the next section. This proof relies crucially on the following two estimates.

Proposition 6.1. *Let $n, m \in \mathbb{Z}_+$ and $\xi \in \mathbb{R}^{2m}$. If $\nu > 0$ is given by (6.4), then*

$$\mathbb{E}_n \left[e^{-2 \sum_{j=1}^n \Im g(\theta_j + i \frac{\nu}{n} h(\theta_j))} \right] \leq \exp \left(-2\nu \|\xi\|^2 \left(1 - c_{10} - \frac{4c_{11}\nu_* \|\xi\|}{N\sqrt{m+1}} (1 + \log m)^{3/4} \right) \right).$$

Proposition 6.2. *Let $n, m \in \mathbb{Z}_+$ with $m \geq 3$, $\xi \in \mathbb{R}^{2m}$ and suppose that the parameter ν is given by (6.4) with $0 < \nu_* \leq c_0$. If H is given by (2.14), we have*

$$\mathbb{E}_n \left[\exp \left(\frac{\nu^2}{n^2} \sum_{i,j=1}^n H(\theta_i, \theta_j) \right) \right] \leq \exp \left(2c_9 + \frac{\nu_*^2 N^2 (1 + \epsilon_0)}{(m+1)\sqrt{1 + \log m}} \right).$$

The proof of Proposition 6.1 is given in Section 6.3 while the proof of Proposition 6.2 is given in Section 6.4. Now that we are equipped with these two estimates, we can proceed with the proof of Proposition 2.8.

6.1 Proof of Proposition 2.8

Let us recall that the parameter ν is chosen according to (6.4) and we assume that $0 < \nu_* \leq c_0 = \sqrt{\frac{1}{6\sqrt{2}}}$. By combining Lemma 2.9, Lemma 6.1 and Proposition 6.2, we obtain

$$|F_{n,m}(\xi)|^2 \leq \exp \left(2c_9 + \frac{\nu_*^2 N^2 (1 + \epsilon_0)}{(m+1)\sqrt{1 + \log m}} - \frac{2\nu_* N \|\xi\|}{\sqrt{m+1}(1 + \log m)^{1/4}} \left(1 - c_{10} - \frac{4c_{11}\nu_* \|\xi\|}{N\sqrt{m+1}} (1 + \log m)^{3/4} \right) \right). \quad (6.5)$$

Let Λ_2 be as in (2.25), that is

$$\Lambda_2 = \frac{c_0^{-1}(1 - c_{10})N\sqrt{m+1}}{8(1 + \log m)^{3/4}c_{11}}.$$

In order to maximize the polynomial $\nu_*(1 - c_{10} - 4\nu_*c_{11}\frac{\|\xi\|}{N\sqrt{m+1}}(1 + \log m)^{3/4})$, we choose $\nu_* = \frac{(1-c_{10})N\sqrt{m+1}}{8\|\xi\|(1+\log m)^{3/4}c_{11}}$. Then, we verify that in the regime where $\|\xi\| \geq \Lambda_2$, we have $\nu_* \leq c_0$ so that we are allowed to use the estimate (6.5). We obtain

$$|F_{n,m}(\xi)|^2 \leq \exp\left(2c_9 + \frac{c_0^2 N^2(1 + \epsilon_0)}{(m+1)\sqrt{1 + \log m}} - \frac{(1 - c_{10})^2 N^2}{8(1 + \log m)c_{11}}\right)$$

If $c_1 = \frac{(1-c_{10})^2}{16c_{11}} - c_0^2(1 + \epsilon_0)\frac{\sqrt{1+\log m}}{2(m+1)}$ according to (B.3), it follows from the previous formula that in the regime where $\|\xi\| \geq \Lambda_2$,

$$|F_{n,m}(\xi)|^2 \leq \exp\left(2c_9 - \frac{2c_1 N^2}{1 + \log m}\right).$$

This proves the estimate (2.26).

On the other hand, in the regime where $\|\xi\| \leq \Lambda_2$ if we choose $\nu_* = c_0$ in the estimate (6.5), by (2.25), we verify that

$$\begin{aligned} |F_{n,m}(\xi)|^2 &\leq \exp\left(2c_9 + c_0^2\left(\frac{N^2(1 + \epsilon_0)}{(m+1)\sqrt{1 + \log m}} - 8c_{11}\frac{\sqrt{1 + \log m}}{m+1}\|\xi\|(2\Lambda_2 - \|\xi\|)\right)\right) \\ &\leq \exp\left(2c_9 + c_0^2\left(\frac{N^2(1 + \epsilon_0)}{(m+1)\sqrt{1 + \log m}} - 8c_{11}\frac{\sqrt{1 + \log m}}{m+1}\Lambda_1(2\Lambda_2 - \Lambda_1)\right)\right), \end{aligned}$$

where we used that the minimum of the function $\xi \mapsto \|\xi\|(2\Lambda_2 - \|\xi\|)$ for $\Lambda_1 \leq \|\xi\| \leq \Lambda_2$ equals

$$\begin{aligned} \Lambda_1(2\Lambda_2 - \Lambda_1) &= \frac{c_4 N}{\sqrt{1 + \log m}}\left(\frac{c_0^{-1}(1 - c_{10})N\sqrt{m+1}}{4c_{11}(1 + \log m)^{3/4}} - \frac{c_4 N}{\sqrt{1 + \log m}}\right) \\ &= \frac{c_0^{-1}c_4\sqrt{m+1}}{4c_{11}(1 + \log m)^{5/4}}N^2\left(1 - c_{10} - \frac{4c_4c_0c_{11}(1 + \log m)^{1/4}}{\sqrt{m+1}}\right). \end{aligned}$$

Hence, if $c_2 = c_0c_4\left(1 - c_{10} - \frac{4c_4c_0c_{11}(1+\log m)^{1/4}}{\sqrt{m+1}}\right) - c_0^2\frac{(1+\epsilon_0)(1+\log m)^{1/4}}{2\sqrt{m+1}}$ according to (B.3), it follows from the previous formulae that in the regime where $\Lambda_1 \leq \|\xi\| \leq \Lambda_2$,

$$|F_{n,m}(\xi)|^2 \leq \exp\left(2c_9 - \frac{2c_2(m)N^2}{\sqrt{m+1}(1 + \log m)^{3/4}}\right).$$

This proves the estimate (2.27) and it completes the proof. It just remains to prove Lemma 2.9 as well as Propositions 6.1 and 6.2 which is the task that we undertake in the next sections. \square

6.2 Proof of Lemma 2.9

Let us recall that by Lemma 2.4, we have for any $\nu > 0$,

$$|F_{n,m}(\xi)| \leq \mathbb{E}_n \left[\prod_{1 \leq i < j \leq n} \left| \frac{\sin\left(\frac{\theta_i - \theta_j}{2} + i\nu\frac{h(\theta_i) - h(\theta_j)}{2n}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)} \right|^2 \prod_{j=1}^n \left| 1 + i\frac{\nu}{n}h'(\theta_j) \right| e^{-\Im_{\mathbf{g}}(\theta_j + i\frac{\nu}{n}h(\theta_j))} \right].$$

By Lemma 5.2, we obtain for all $\theta_i, \theta_j \in \mathbb{T}$ with $\theta_i \neq \theta_j$,

$$\begin{aligned} \left| \frac{\sin\left(\frac{\theta_i - \theta_j}{2} + i\nu\frac{h(\theta_i) - h(\theta_j)}{2n}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)} \right|^2 &= 1 + \left(\frac{\sinh\left(\nu\frac{h(\theta_i) - h(\theta_j)}{2n}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)} \right)^2 \\ &\leq \exp\left(\nu\frac{h(\theta_i) - h(\theta_j)}{2n\sin\left(\frac{\theta_i - \theta_j}{2}\right)}\right)^2 \\ &= \exp\left(\frac{\nu^2}{n^2}H(\theta_i, \theta_j)\right), \end{aligned}$$

where the function H is as in (2.14). Moreover, we also have

$$\prod_{j=1}^n |1 + i\frac{\nu}{n}h'(\theta_j)|^2 \leq \exp\left(\frac{\nu^2}{n^2}\sum_{j=1}^n H(\theta_j, \theta_j)\right).$$

Combining these bounds, we obtain for any $\theta_1, \dots, \theta_n \in \mathbb{T}$ distinct and any $\nu > 0$,

$$\prod_{1 \leq i < j \leq n} \left| \frac{\sin\left(\frac{\theta_i - \theta_j}{2} + i\nu\frac{h(\theta_i) - h(\theta_j)}{2n}\right)}{\sin\left(\frac{\theta_i - \theta_j}{2}\right)} \right|^2 \prod_{j=1}^n |1 + i\frac{\nu}{n}h'(\theta_j)| \leq \exp\left(\frac{1}{2}\frac{\nu^2}{n^2}\sum_{i,j=1}^n H(\theta_i, \theta_j)\right).$$

Hence, by the Cauchy–Schwartz inequality, this implies that

$$|F_{n,m}(\xi)|^2 \leq \mathbb{E}_n \left[\exp\left(\frac{\nu^2}{n^2}\sum_{i,j=1}^n H(\theta_i, \theta_j)\right) \right] \mathbb{E}_n \left[e^{-2\sum_{j=1}^n \Im g(\theta_j + i\frac{\nu}{n}h(\theta_j))} \right].$$

□

6.3 Proof of Proposition 6.1

Recall that according to (6.4), we assume that $\nu = \frac{\nu_* n}{m\sqrt{m+1}(1+\log m)^{1/4}\|\xi\|}$ for a constant $\nu_* > 0$. Using the estimate (6.1), this implies that $\frac{\nu}{n}\|h\|_\infty \leq \sqrt{2}\nu_*\frac{(1+\log m)^{1/4}}{m\sqrt{m+1}}$. Then, since both functions g, h are real-valued on \mathbb{T} and g is an analytic function, we have

$$\left| \Im g\left(\theta_j + i\frac{\nu}{n}h(\theta_j)\right) - \frac{\nu}{n}g'(\theta_j)h(\theta_j) \right| \leq \frac{\nu^3}{6n^3}|h(\theta_j)|^3 \sup_{\substack{z \in \mathbb{C}: \\ |\Re z| \leq \pi, |\Im z| \leq \sqrt{2}\nu_*\frac{(1+\log m)^{1/4}}{m\sqrt{m+1}}}} |g'''(z)|.$$

Moreover, by (2.2), we have for any $z \in \mathbb{C}$,

$$g'''(z) = \frac{-i}{\sqrt{2}} \sum_{|k| \leq m} |k|^{5/2} \zeta_k e^{ikz}$$

so that if $|\Re z| \leq \pi, |\Im z| \leq \sqrt{2}\nu_*\frac{(1+\log m)^{1/4}}{m\sqrt{m+1}}$, then

$$|g'''(z)| \leq \sqrt{2}e^{\sqrt{2}\nu_*\frac{(1+\log m)^{1/4}}{\sqrt{m+1}}} \sum_{k=1}^m |\zeta_k| k^{5/2} \leq 3\sqrt{2}c_{19}\|\xi\| m^{3/2}(m+1)^{3/2},$$

where $c_{19}(m) = \frac{1}{3\sqrt{6}}e^{\sqrt{2}c_0\frac{(1+\log m)^{1/4}}{\sqrt{m+1}}}$ and we used that $\sum_{k=1}^m k^5 \leq \frac{m^3(m+1)^3}{6}$. These bounds and the estimate (6.1) show that

$$\begin{aligned} & \mathbb{E}_n \left[\exp\left(-2\sum_{j=1}^n \Im g\left(\theta_j + i\frac{\nu}{n}h(\theta_j)\right)\right) \right] \\ & \leq \mathbb{E}_n \left[\exp\left(-\frac{2\nu}{n}\sum_{j=1}^n g'(\theta_j)h(\theta_j) + 2c_{19}\|\xi\|^2 \frac{\nu^3 m^{3/2}(m+1)^{3/2}}{n^3} \sqrt{1+\log m} \sum_{j=1}^n h(\theta_j)^2\right) \right]. \end{aligned}$$

Let us denote $\gamma = c_{19}\|\xi\|^2 \frac{\nu^2 m^{3/2}(m+1)^{3/2}}{n^2} \sqrt{1+\log m}$ and $f = g' - \gamma h$. By Lemma 2.3, this implies that

$$\begin{aligned} \mathbb{E}_n \left[\exp\left(-2\sum_{j=1}^n \Im g\left(\theta_j + i\frac{\nu}{n}h(\theta_j)\right)\right) \right] & \leq \mathbb{E}_n \left[\exp\left(-\frac{2\nu}{n}\sum_{j=1}^n f(\theta_j)h(\theta_j)\right) \right] \\ & \leq \exp\left(-2\nu \int_{\mathbb{T}} f(\theta)h(\theta) \frac{d\theta}{2\pi} + \frac{4\nu^2}{n^2} \mathcal{A}(fh)\right). \end{aligned}$$

First observe that since we have chosen $h = -\mathcal{U}g$, we have

$$\int_{\mathbb{T}} h(\theta)^2 \frac{d\theta}{2\pi} \leq \|\xi\|^2 \quad (6.6)$$

and by formulae (2.15)–(2.16), we obtain

$$\mathbb{E}_n \left[\exp \left(-2 \sum_{j=1}^n \Im g(\theta_j + i \frac{\nu}{n} h(\theta_j)) \right) \right] \leq \exp \left(-2\nu(1-\gamma)\|\xi\|^2 + \frac{4\nu^2}{n^2} \mathcal{A}(fh) \right). \quad (6.7)$$

It remains to estimate the quantities $\mathcal{A}(fh)$ where the seminorm \mathcal{A} is given by (2.4) and $f = g' - \gamma h$. To that end, we may use the bound $\mathcal{A}(u) \leq \|u\|_{L^2} \|u'\|_{L^2}$ which holds for any smooth function $u : \mathbb{T} \rightarrow \mathbb{C}$. First, we have

$$\|fh\|_{L^2} \leq \|h\|_{\infty} \|f\|_{L^2} \leq \sqrt{2(1+\log m)} (\sqrt{m} + \gamma) \|\xi\|^2$$

where we used the estimates (6.1), (6.3) and (6.6). Second, we have

$$\begin{aligned} \|(fh)'\|_{L^2} &\leq \|h\|_{\infty} \|f'\|_{L^2} + \|f\|_{\infty} \|h'\|_{L^2} \\ &\leq \left(\sqrt{2m(1+\log m)} (m + \gamma) + \left(\sqrt{m(m+1)} + \gamma \sqrt{2(1+\log m)} \right) \sqrt{m} \right) \|\xi\|^2 \\ &= m \sqrt{2m(1+\log m)} \left(1 + \frac{2\gamma}{m} + \sqrt{\frac{1+1/m}{2(1+\log m)}} \right) \|\xi\|^2 \end{aligned}$$

Here we used that $\|g^{(\kappa)}\|_{L^2} = \|h^{(\kappa)}\|_{L^2}$ for any $\kappa \geq 0$ since h is the Hilbert transform of g and the estimates (6.1)–(6.3). Combining all these estimates, we deduce from formula (6.7) that

$$\begin{aligned} &\mathbb{E}_n \left[\exp \left(-2 \sum_{j=1}^n \Im \left\{ g(\theta_j + i \frac{\nu}{n} h(\theta_j)) \right\} \right) \right] \\ &\leq \exp \left(-2\nu\|\xi\|^2 \left(1 - \gamma - \frac{4\nu\|\xi\|^2 m^2}{n^2} (1+\log m) \left(1 + \frac{\gamma}{\sqrt{m}} \right) \left(1 + \frac{2\gamma}{m} + \sqrt{\frac{1+1/m}{2(1+\log m)}} \right) \right) \right). \end{aligned}$$

To complete the proof, it remains to observe that by (6.4) and (B.3), we have

$$\gamma = c_{19} \|\xi\|^2 \frac{\nu^2 (m+1)^{3/2}}{N^2 \sqrt{m}} \sqrt{1+\log m} = c_{19} \nu_*^2 \sqrt{1+1/m} \leq c_{10}(m) = c_0^2 \frac{\sqrt{1+1/m}}{3\sqrt{6}} e^{c_0 \frac{(1+\log m)^{1/4}}{\sqrt{(m+1)^{1/2}}}}$$

after replacing $c_{19} = \frac{1}{3\sqrt{6}} e^{c_0 \frac{(1+\log m)^{1/4}}{\sqrt{(m+1)^{1/2}}}}$ and using that $\nu_* \leq c_0$. Moreover, by (6.4), we also have $\frac{\nu\|\xi\|^2 m^2}{n^2} (1+\log m) = \frac{\nu_* \|\xi\|}{N \sqrt{m+1}} (1+\log m)^{3/4}$, so as $c_{11} = \left(1 + \frac{c_{10}}{\sqrt{m}} \right) \left(1 + \frac{2c_{10}}{m} + \sqrt{\frac{1+1/m}{2(1+\log m)}} \right)$, this proves the claimed bound. \square

6.4 Proof of Proposition 6.2

Let us denote for any $k \in \mathbb{Z}$,

$$\mathbf{T}_k = \text{Tr } \mathbf{U}^k = \sum_{j=1}^n e^{ik\theta_j} = \sqrt{\frac{k}{2}} (X_{2k-1} + iX_{2k}).$$

The idea of the proof is to view $\sum_{i,j=1}^n H(\theta_i, \theta_j)$ as a quadratic form in the random variables $(\mathbf{T}_k)_{k \in \mathbb{Z}}$ and to use this observation to express the Laplace transform of the random variable $\sum_{i,j=1}^n H(\theta_i, \theta_j)$ as a (multivariate) Gaussian integral as explained at the end of Section 2.4.

Lemma 6.3. *We have the identity*

$$\sum_{i,j=1}^n H(\theta_i, \theta_j) = \frac{1}{2} \Re \left\{ \sum_{p,q \in \mathbb{Z}} A_{pq} \mathbb{T}_p \mathbb{T}_q + \sum_{p,q \in \mathbb{Z}} B_{pq} \mathbb{T}_p \mathbb{T}_q \right\},$$

where

$$A_{pq} = \sum_{1 \leq k \leq \ell \leq m} (\mathbf{1}_{1 \leq p-k+1 \leq p+q-\ell \leq m} + \mathbf{1}_{1 \leq q-k+1 \leq p+q-\ell \leq m}) \frac{\zeta_\ell \zeta_{p+q-\ell}}{\sqrt{\ell(p+q-\ell)}}$$

and

$$B_{pq} = \sum_{1 \leq k \leq \ell \leq m} (\mathbf{1}_{1 \leq k-p \leq \ell-p-q \leq m} + \mathbf{1}_{1 \leq k-q \leq \ell-p-q \leq m}) \frac{\zeta_\ell \zeta_{p+q-\ell}}{\sqrt{\ell(\ell-p-q)}}.$$

Proof. An elementary computation gives that for any $\ell \in \mathbb{Z}$,

$$\frac{e^{i\ell\theta} - e^{i\ell x}}{2i \sin(\frac{\theta-x}{2})} = \sum_{k=1}^{\ell} e^{i(k-1/2)\theta} e^{i(\ell-k+1/2)x}, \quad x, \theta \in \mathbb{T}.$$

By (2.13) – (2.14), this directly implies that for any $i, j = 1, \dots, n$,

$$\begin{aligned} H(\theta_i, \theta_j) &= \Re \left\{ \sum_{1 \leq k \leq \ell \leq m} \sum_{1 \leq r \leq s \leq m} \frac{\zeta_\ell \zeta_s}{\sqrt{\ell s}} e^{i(k-1/2)\theta_i} e^{i(\ell-k+1/2)\theta_j} e^{i(r-1/2)\theta_i} e^{i(s-r+1/2)\theta_j} \right\} \\ &+ \Re \left\{ \sum_{1 \leq k \leq \ell \leq m} \sum_{1 \leq r \leq s \leq m} \frac{\zeta_\ell \bar{\zeta}_s}{\sqrt{\ell s}} e^{i(k-1/2)\theta_i} e^{i(\ell-k+1/2)\theta_j} e^{-i(r-1/2)\theta_i} e^{-i(s-r+1/2)\theta_j} \right\}. \end{aligned}$$

Then summing over all variables θ_i, θ_j , we obtain

$$\sum_{1 \leq i, j \leq n} H(\theta_i, \theta_j) = \Re \left\{ \sum_{1 \leq k \leq \ell \leq m} \sum_{1 \leq r \leq s \leq m} \frac{\zeta_\ell \zeta_s}{\sqrt{\ell s}} \mathbb{T}_{k+r-1} \mathbb{T}_{\ell+s-k-r+1} \right\} \quad (6.8)$$

$$+ \Re \left\{ \sum_{1 \leq k \leq \ell \leq m} \sum_{1 \leq r \leq s \leq m} \frac{\zeta_\ell \bar{\zeta}_s}{\sqrt{\ell s}} \mathbb{T}_{k-r} \mathbb{T}_{\ell-s+r-k} \right\}. \quad (6.9)$$

In (6.8) we make the change of variables $(r, s) \leftrightarrow (p, q)$ given by $r = p - k + 1$ and $s = q + p - \ell$. Similarly, in (6.9) we make the change of variables $(r, s) \leftrightarrow (p, q)$ given by $r = k - p$ and $s = \ell - q - p$. This implies that

$$\sum_{1 \leq i, j \leq n} H(\theta_i, \theta_j) = \Re \left\{ \sum_{1 \leq k \leq \ell \leq m} \sum_{p, q \in \mathbb{Z}} \frac{\zeta_\ell \zeta_{q+p-\ell}}{\sqrt{\ell(q+p-\ell)}} \mathbf{1}_{1 \leq p-k+1 \leq q+p-\ell \leq m} \mathbb{T}_p \mathbb{T}_q \right\} \quad (6.10)$$

$$+ \Re \left\{ \sum_{1 \leq k \leq \ell \leq m} \sum_{p, q \in \mathbb{Z}} \frac{\zeta_\ell \bar{\zeta}_{\ell-q-p}}{\sqrt{\ell(\ell-q-p)}} \mathbf{1}_{1 \leq k-p \leq \ell-q-p \leq m} \mathbb{T}_p \mathbb{T}_q \right\}. \quad (6.11)$$

To finish the proof, it remains to symmetrize the previous formula over (p, q) and use that $\bar{\zeta}_{-j} = \zeta_j$ for all $j = 1, \dots, m$. Then (6.10) corresponds to $\frac{1}{2} \Re \{ \sum_{p, q \in \mathbb{Z}} A_{pq} \mathbb{T}_p \mathbb{T}_q \}$ and (6.11) corresponds to $\frac{1}{2} \Re \{ \sum_{p, q \in \mathbb{Z}} B_{pq} \mathbb{T}_p \mathbb{T}_q \}$. \square

Let us observe that in the notation of Lemma 6.3, $A_{pq} \neq 0$ only if $1 \leq p, q \leq 2m - 1$ and $B_{pq} \neq 0$ only if $|p|, |q| \leq m - 1$, so we may view $\mathbf{A} = (A_{pq})_{p, q=1}^{2m-1}$ and $\mathbf{B} = (B_{pq})_{1 \leq |p|, |q| < m}$ as symmetric matrix-valued functions of the parameters $(\zeta_k)_{k=1}^m$. In the following, we denote

$$\mathfrak{Q}_{\mathbf{A}} = \Re \left\{ \sum_{p, q \in \mathbb{Z}} A_{pq} \mathbb{T}_p \mathbb{T}_q \right\}, \quad \mathfrak{Q}_{\mathbf{B}} = \Re \left\{ \sum_{\substack{p, q \in \mathbb{Z} \\ p, q \neq 0}} B_{pq} \mathbb{T}_p \mathbb{T}_q \right\} \quad \text{and} \quad \mathfrak{L} = \Re \left\{ n B_{00} + 2 \sum_{\substack{p \in \mathbb{Z} \\ p \neq 0}} B_{p0} \mathbb{T}_p \right\}.$$

We introduce this decomposition because $T_0 = n$ is not a random variable and should be treated individually. By Cauchy–Schwartz inequality, Lemma 6.3 implies that

$$\mathbb{E}_n \left[\exp \left(\delta \sum_{i,j=1}^n H(\theta_i, \theta_j) \right) \right] \leq \left(\mathbb{E}_n [\exp(2\delta\mathfrak{Q}_\mathbf{A})] \mathbb{E}_n [\exp(2\delta\mathfrak{Q}_\mathbf{B})] \right)^{1/4} \sqrt{\mathbb{E}_n [\exp(\delta n\mathfrak{L})]}, \quad (6.12)$$

where $\delta = (\frac{\nu}{n})^2$. Our first observation is that \mathfrak{L} is a linear statistic associated with the trigonometric polynomial $f = nB_{00} + 2\Re\{\sum_{p \neq 0} B_{p0} e^{ip\theta}\}$ so that by Lemma 2.3, we have the estimate

$$\mathbb{E}_n [\exp(\delta n\mathfrak{L})] \leq \exp \left(\delta n^2 B_{00} + (\delta n)^2 \sum_{0 < p < m} p |B_{p0} + \overline{B_{-p0}}|^2 \right).$$

In combination with Lemma 6.4 below, this implies that

$$\mathbb{E}_n [\exp(\delta n\mathfrak{L})] \leq \exp \left(2\delta n^2 \|\xi\|^2 + (\delta n)^2 \frac{4m^3}{3} \|\xi\|^4 \right). \quad (6.13)$$

Lemma 6.4. *In the notation of Lemma 6.3, we have $B_{00} = 2\|\xi\|^2$ and*

$$\sum_{0 < p < m} p |B_{p0} + \overline{B_{-p0}}|^2 \leq \frac{4m^3}{3} \|\xi\|^4.$$

Proof. First of all, by definition, we have

$$\frac{B_{00}}{2} = \sum_{1 \leq k \leq \ell \leq m} \frac{\zeta_k \zeta_{-\ell}}{\ell} = \sum_{1 \leq k \leq m} |\zeta_k|^2 = \|\xi\|^2.$$

Secondly, we also have for any $p \in \mathbb{Z}$,

$$B_{p0} = \sum_{1 \leq k \leq \ell \leq m} (\mathbf{1}_{1 \leq k-p \leq \ell-p \leq m} + \mathbf{1}_{1 \leq k \leq \ell-p \leq m}) \frac{\zeta_k \zeta_{p-\ell}}{\sqrt{\ell(\ell-p)}}.$$

This shows that for $p \geq 1$,

$$\begin{aligned} |B_{p0}| &\leq \|\xi\| \sum_{1 \leq k \leq \ell \leq m} (\mathbf{1}_{p+1 \leq k \leq \ell \leq m} + \mathbf{1}_{1 \leq k \leq \ell-p \leq m}) \frac{|\zeta_\ell|}{\sqrt{\ell(\ell-p)}} \\ &= 2\|\xi\| \sum_{p+1 \leq \ell \leq m} \sqrt{1-p/\ell} |\zeta_\ell|, \end{aligned}$$

where at the second step we computed the sum over k . By Cauchy–Schwartz inequality, this shows that

$$|B_{p0}| \leq 2\sqrt{m-p} \|\xi\|^2$$

This estimate implies that

$$\sum_{p=1}^{m-1} p |B_{p0}|^2 \leq 4\|\xi\|^4 \sum_{p=1}^{m-1} p(m-p) \leq \frac{2m^3}{3} \|\xi\|^4.$$

Similarly, we can show that $|B_{-p0}| \leq 2\sqrt{m-p} \|\xi\|^2$ for any $p \geq 1$ so that we also have $\sum_{p=1}^{m-1} p |B_{-p0}|^2 \leq \frac{2m^3}{3} \|\xi\|^4$. This completes the proof. \square

In the remainder of this section, our task is to bound the terms which involve the quadratic forms $\mathfrak{Q}_{\mathbf{A}}$ and $\mathfrak{Q}_{\mathbf{B}}$ on the RHS of (6.12). In order to do this, we need a priori estimates for the norms of the corresponding matrices \mathbf{A} and \mathbf{B} .

Lemma 6.5. *Let $\|\mathbf{A}\| = \max_{1 \leq p < 2m} \sum_{q=1}^{2m-1} |A_{pq}|$ and $\|\mathbf{B}\| = \max_{1 \leq |p| < m} \sum_{|q|=1}^{m-1} |B_{pq}|$. We have*

$$\|\mathbf{A}\|, \|\mathbf{B}\| \leq \sqrt{2m(m+1)(1+\log m)} \|\xi\|^2.$$

Proof. By definition, we have

$$\sum_{q=1}^{2m-1} |A_{pq}| \leq 2 \sum_{1 \leq k \leq \ell \leq m} \frac{|\zeta_\ell|}{\sqrt{\ell}} \sum_{q=1}^{2m-1} \mathbf{1}_{1 \leq p+q-\ell \leq m} \frac{|\zeta_{p+q-\ell}|}{\sqrt{p+q-\ell}}.$$

The last sum is bounded by $\sum_{r=1}^m \frac{|\zeta_r|}{\sqrt{r}}$, so we obtain

$$\sum_{q=1}^{2m-1} |A_{pq}| \leq 2 \sum_{k,r=1}^m \sqrt{\frac{\ell}{r}} |\zeta_\ell| |\zeta_r|.$$

By the Cauchy–Schwartz inequality, this shows that

$$\sum_{q=1}^{2m-1} |A_{pq}| \leq 2 \|\zeta\|^2 \sqrt{\sum_{k,r=1}^m \frac{\ell}{r}} \leq \sqrt{2m(m+1)(1+\log m)} \|\zeta\|^2.$$

Since $\|\zeta\| = \|\xi\|$, this gives the estimate for $\|\mathbf{A}\|$ – the argument for $\|\mathbf{B}\|$ is exactly the same. \square

Let us define new objects. For $\delta_1, \delta_2 > 0$, we set

$$\mathbf{M} = \begin{pmatrix} \mathbf{I}_{2m-1} & \delta_2 \mathbf{A}^* \\ \delta_2 \mathbf{A} & \mathbf{I}_{2m-1} \end{pmatrix} \quad \text{and} \quad \mathbf{v} = \sqrt{\delta_1} \begin{pmatrix} \mathbf{T}_1 \\ \vdots \\ \mathbf{T}_{2m-1} \end{pmatrix}. \quad (6.14)$$

Remark 6.1. As explained in Section 2.4, it is not clear whether the matrices \mathbf{A} and \mathbf{B} are positive definite. This issue is resolved by bounding the quadratic form $\mathfrak{Q}_{\mathbf{A}}$ using the matrix \mathbf{M} (see the estimate (6.16)) and by choosing the parameter δ_2 small enough to guarantee that \mathbf{M} is positive definite and the Gaussian integral (6.17) is convergent. \blacksquare

Since \mathbf{A} is a symmetric matrix, we have $\|\mathbf{A}^*\| = \|\mathbf{A}\|$ and

$$\|\mathbf{M} - \mathbf{I}_{4m-2}\| = \max_{1 \leq p < 4m-1} \sum_{q=1}^{4m-2} |M_{pq} - \mathbf{1}_{pq}| = \delta_2 \|\mathbf{A}\|.$$

Hence, by Lemma 6.5, if $\delta_2 \leq \frac{1}{3\sqrt{2m(m+1)(1+\log m)} \|\xi\|^2}$, then

$$\|\mathbf{M} - \mathbf{I}_{4m-2}\| \leq \frac{1}{3}, \quad (6.15)$$

so that the matrix \mathbf{M} is invertible with $\mathbf{M}^{-1} = \sum_{k=0}^{+\infty} (\mathbf{I}_{4m-2} - \mathbf{M})^k$ (convergent Neumann series). This also implies that \mathbf{M} is positive definite and we have

$$(\mathbf{v}^* \quad \bar{\mathbf{v}}^*) \begin{pmatrix} \mathbf{I} & \delta_2 \mathbf{A}^* \\ \delta_2 \mathbf{A} & \mathbf{I} \end{pmatrix} \begin{pmatrix} \mathbf{v} \\ \bar{\mathbf{v}} \end{pmatrix} = 2\delta_2 \Re \{ \mathbf{v}^t \mathbf{A} \mathbf{v} \} + 2|\mathbf{v}|^2.$$

Thus, if we set $\delta = \delta_1 \delta_2$ and use the notation (6.14), this shows that

$$2\delta\Omega_{\mathbf{A}} = 2\delta\Re\left\{\sum_{p,q\in\mathbb{Z}} A_{pq}\mathbb{T}_p\mathbb{T}_q\right\} \leq \left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right)^* \mathbf{M} \left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right). \quad (6.16)$$

Then, in order to estimate the quantity $\mathbb{E}_n[\exp(2\delta\Omega_{\mathbf{A}})]$, we may use the identity

$$\pi^{4m-2} \det(\mathbf{M}) \exp\left(\left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right)^* \mathbf{M} \left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right)\right) = \int_{\mathbb{C}^{4m-2}} \exp\left(-\mathbf{z}^* \mathbf{M}^{-1} \mathbf{z} + \mathbf{z}^* \left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right) + \left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right)^* \mathbf{z}\right) d^2 \mathbf{z}, \quad (6.17)$$

where $\mathbf{z} = \begin{pmatrix} z_1 \\ \vdots \\ z_{4m-2} \end{pmatrix}$ and $d^2 \mathbf{z} = \prod_{k=1}^{4m-2} d\Re(z_k) d\Im(z_k)$ denotes the Lebesgue measure on \mathbb{C}^{4m-2} .

Formula (6.17) is a simple Gaussian integration on \mathbb{C}^{4m-2} and it makes sense since we have seen that the matrix \mathbf{M} is positive definite by (6.15). Moreover, it is useful since

$$\left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right)^* \mathbf{z} = \sqrt{\delta_1} \sum_{k=1}^{2m-1} (z_k \bar{\Gamma}_k + z_{2m-1+k} \Gamma_k)$$

is a (mean-zero) linear statistic of a trigonometric polynomial, so that by Lemma 2.3, we have

$$\begin{aligned} \mathbb{E}_n \left[\exp\left(\mathbf{z}^* \left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right) + \left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right)^* \mathbf{z}\right) \right] &\leq \exp\left(\delta_1 \sum_{k=1}^{2m-1} k |\bar{z}_k + z_{2m-1+k}|^2\right) \\ &\leq \exp(2\delta_1 \mathbf{z}^* \mathbf{C} \mathbf{z}) \end{aligned} \quad (6.18)$$

where \mathbf{C} is a diagonal matrix given by

$$\mathbf{C} = \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & 2m-1 & & \\ & & & 1 & \\ & & & & \ddots & \\ & & & & & 2m-1 \end{pmatrix}$$

Hence, taking expectation in formula (6.17) and using the bound (6.18), we obtain

$$\begin{aligned} \mathbb{E}_n \left[\exp\left(\left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right)^* \mathbf{M} \left(\frac{\mathbf{v}}{\bar{\mathbf{v}}}\right)\right) \right] &\leq \frac{1}{\pi^{4m-2} \det(\mathbf{M})} \int_{\mathbb{C}^{4m-2}} \exp(-\mathbf{z}^* (\mathbf{M}^{-1} - 2\delta_1 \mathbf{C}) \mathbf{z}) d^2 \mathbf{z} \\ &= \frac{\det(\mathbf{M}^{-1} - 2\delta_1 \mathbf{C})^{-1}}{\det(\mathbf{M})} = \frac{1}{\det(\mathbf{I} - 2\delta_1 \mathbf{M} \mathbf{C})}. \end{aligned} \quad (6.19)$$

Here we used that the matrix $\mathbf{M}^{-1} - 2\delta_1 \mathbf{C}$ is also positive definite. Indeed, it follows from the above discussion (in particular from the estimate (6.15)) that if $\delta_2 \leq \frac{1}{3\sqrt{2m(m+1)(1+\log m)} \|\xi\|^2}$ and $\delta_1 \leq \frac{1}{2m^{3/2}\sqrt{m+1}}$, then for any $m \geq 3$,

$$\|\mathbf{M}^{-1} - \mathbf{I}_{4m-2}\| \leq \sum_{k=1}^{+\infty} \|\mathbf{M} - \mathbf{I}_{4m-2}\|^k \leq \frac{1}{2} \quad \text{and} \quad 2\delta_1 \|\mathbf{C}\| \leq \frac{2m-1}{m^{3/2}\sqrt{m+1}} \leq \frac{5}{6\sqrt{3}} < \frac{1}{2}.$$

Note that the condition $m \geq 3$ is crucial in order to obtain the second estimate. Moreover, since \mathbf{M}, \mathbf{C} are Hermitian matrices with $\|\mathbf{M}\| \leq 4/3$, it follows that for all $m \geq 3$,

$$\det(\mathbf{I} - 2\delta_1 \mathbf{M} \mathbf{C}) \geq \left(1 - \frac{4(2m-1)}{3m^{3/2}\sqrt{m+1}}\right)^{2(2m-1)} \geq e^{-c_{17}},$$

where $c_{17} = \frac{32}{3}(1 + \frac{(2-1/m)^3}{3(m+1)})$ and we used that $1 - x \geq e^{-x-x^2}$ for $0 \leq x \leq 2/3$. Hence, by formula (6.16) and (6.19), we obtain for $m \geq 3$,

$$\mathbb{E}_n[\exp(2\delta\Omega_{\mathbf{A}})] \leq \mathbb{E}_n\left[\exp\left(\left(\frac{\mathbf{v}}{\overline{\mathbf{v}}}\right)^* \mathbf{M} \left(\frac{\mathbf{v}}{\overline{\mathbf{v}}}\right)\right)\right] \leq e^{c_{17}}. \quad (6.20)$$

In an analogous way, let us denote

$$\mathbf{N} = \begin{pmatrix} \mathbf{I} & \delta_2 \mathbf{B}^* \\ \delta_2 \mathbf{B} & \mathbf{I} \end{pmatrix}, \quad \mathbf{D} = \begin{pmatrix} m-1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & 1 & & \\ & & & & \ddots & \\ & & & & & m-1 \end{pmatrix} \quad \text{and} \quad \mathbf{w} = \sqrt{\delta_1} \begin{pmatrix} \mathbf{T}_{-m+1} \\ \vdots \\ \mathbf{T}_{-1} \\ \mathbf{T}_1 \\ \vdots \\ \mathbf{T}_{m-1} \end{pmatrix}.$$

Then we have

$$2\delta\Omega_{\mathbf{B}} = 2\delta\Re\left\{\sum_{\substack{p,q \in \mathbb{Z} \\ p,q \neq 0}} B_{pq} \mathbf{T}_p \mathbf{T}_q\right\} \leq \left(\frac{\mathbf{w}}{\overline{\mathbf{w}}}\right)^* \mathbf{N} \left(\frac{\mathbf{w}}{\overline{\mathbf{w}}}\right).$$

By Lemma 6.5, if $\delta_2 \leq \frac{1}{3\sqrt{2m(m+1)(1+\log m)}\|\xi\|^2}$, then $\|\mathbf{N} - \mathbf{I}_{4m-2}\| \leq 1/3$ so that both \mathbf{N} and $\mathbf{N}^{-1} - 2\delta_1 \mathbf{D}$ are positive definite matrices. Like in our previous computations, this implies that

$$\begin{aligned} \mathbb{E}_n\left[\exp\left(\left(\frac{\mathbf{w}}{\overline{\mathbf{w}}}\right)^* \mathbf{N} \left(\frac{\mathbf{w}}{\overline{\mathbf{w}}}\right)\right)\right] &= \frac{1}{\pi^{4m-4} \det(\mathbf{N})} \int_{\mathbb{C}^{4m-4}} \exp(-\mathbf{z}^* \mathbf{N}^{-1} \mathbf{z}) \mathbb{E}_n\left[\exp\left(\mathbf{z}^* \left(\frac{\mathbf{w}}{\overline{\mathbf{w}}}\right) + \left(\frac{\mathbf{w}}{\overline{\mathbf{w}}}\right)^* \mathbf{z}\right)\right] d^2 \mathbf{z} \\ &\leq \frac{1}{\pi^{4m-4} \det(\mathbf{N})} \int_{\mathbb{C}^{4m-4}} \exp(-\mathbf{z}^* (\mathbf{N}^{-1} - 2\delta_1 \mathbf{D}) \mathbf{z}) d^2 \mathbf{z} \\ &= \frac{1}{\det(\mathbf{I} - 2\delta_1 \mathbf{N} \mathbf{D})}, \end{aligned}$$

where at the second step we used an estimate analogous to (6.18). Moreover, since \mathbf{N}, \mathbf{D} are Hermitian matrices with $\|\mathbf{N}\| \leq 4/3$ and $\|\mathbf{D}\| = m-1$ for $m \geq 3$, if $\delta_1 \leq \frac{1}{2m^{3/2}\sqrt{m+1}}$, we have

$$\det(\mathbf{I} - 2\delta_1 \mathbf{N} \mathbf{D}) \geq \left(1 - \frac{4(m-1)}{3m^{3/2}\sqrt{m+1}}\right)^{2(m-1)} \geq e^{-c_{18}},$$

where $c_{18} = \frac{8}{3}(1 + \frac{4(1-1/m)^3}{3(m+1)})$ and we used that $1 - x \geq e^{-x-x^2}$ for $0 \leq x \leq 2/3$. Combining these estimates, this implies that for $m \geq 3$,

$$\mathbb{E}_n[\exp(2\delta\Omega_{\mathbf{B}})] \leq \mathbb{E}_n\left[\exp\left(\left(\frac{\mathbf{w}}{\overline{\mathbf{w}}}\right)^* \mathbf{N} \left(\frac{\mathbf{w}}{\overline{\mathbf{w}}}\right)\right)\right] \leq e^{c_{18}}. \quad (6.21)$$

Now, let us recall that we must have $\delta = \left(\frac{\nu}{n}\right)^2 = \delta_1 \delta_2$. Hence, if we choose

$$\delta_1 = \frac{1}{2m^{3/2}\sqrt{m+1}} \quad \text{and} \quad \delta_2 = \frac{2\nu^2 m^{3/2} \sqrt{m+1}}{n^2} = \frac{2\nu_*^2}{\sqrt{m(m+1)(1+\log m)}\|\xi\|^2}$$

according to (6.4), then we have $\delta_2 \leq \frac{1}{3\sqrt{2m(m+1)(1+\log m)}\|\xi\|^2}$ as required provided that $\nu_*^2 \leq \frac{1}{6\sqrt{2}}$. Observe that this explains our choice for c_0 as the maximum admissible value for ν_* . In the end, if we combine all our estimates (6.12), (6.13), (6.20) and (6.21), if the parameter ν is given by (6.4) and $m \geq 3$, then we obtain

$$\mathbb{E}_n \left[\exp \left(\delta \sum_{i,j=1}^n H(\theta_i, \theta_j) \right) \right] \leq \exp \left(2 \frac{c_{17} + c_{18}}{8} + \delta n^2 \|\xi\|^2 + (\delta n)^2 \frac{2m^3}{3} \|\xi\|^4 \right). \quad (6.22)$$

By definitions, we have $\frac{c_{17}+c_{18}}{8} = \frac{1}{3} \left(5 + 4 \frac{(1-1/m)^3 + (2-1/m)^3}{3(m+1)} \right)$. This function attains its maximum over the positive integers for $m = 3$, so that $\frac{c_{17}+c_{18}}{4} \leq c_9 = \frac{538}{243}$. Hence, if we replace $\delta = \frac{\nu^2}{n^2}$ in formula (6.22) and use (6.4), we conclude that

$$\begin{aligned} \mathbb{E}_n \left[\exp \left(\frac{\nu^2}{n^2} \sum_{i,j=1}^n H(\theta_i, \theta_j) \right) \right] &\leq \exp \left(2c_9 + \nu^2 \|\xi\|^2 + \frac{2\nu^4 m^3}{3n^2} \|\xi\|^4 \right) \\ &= \exp \left(2c_9 + \frac{\nu_*^2 N^2}{(m+1)\sqrt{1+\log m}} + \frac{2\nu_*^4 N^2}{3(m+1)(1+1/m)(1+\log m)} \right). \end{aligned}$$

By definition of $\epsilon_0(m) \geq 0$, (B.2), this completes the proof. \square

7 Proof of Theorem 1.5

The method used in this section relies on the formalism introduced in (Lambert et al., 2019) which provides a normal approximation result for certain observable of a Gibbs-type distribution and the following moment identities from (Diaconis and Shahshahani, 1994). According to (1.3), we let for any $k \geq 1$,

$$\mathbb{T}_k = \sqrt{\frac{2}{k}} \operatorname{Tr} \mathbf{U}^k = \mathbf{X}_{2k} + i\mathbf{X}_{2k+1}. \quad (7.1)$$

Theorem 7.1 ((Diaconis and Shahshahani, 1994)). *Fix $m \in \mathbb{N}$ and let $\mathbf{a}, \mathbf{b} \in \{0, 1, \dots\}^m$. Then, for all $n \geq \sum_{k=1}^m ka_k \vee \sum_{k=1}^m kb_k$,*

$$\mathbb{E}_n \left[\prod_{k=1}^m \mathbb{T}_k^{a_k} \overline{\mathbb{T}_k}^{b_k} \right] = \mathbb{E} \left[\prod_{k=1}^m Z_k^{a_k} \overline{Z_k}^{b_k} \right].$$

where $Z_k = \mathbf{G}_{2k} + i\mathbf{G}_{2k+1}$ for all $k \geq 1$ and \mathbf{G}_k are i.i.d. standard Gaussian random variables.

Note that the hypothesis of Theorem 7.1 are incorrectly stated in (Diaconis and Shahshahani, 1994) and we refer instead to (Diaconis and Evans, 2001) for a correct version of this Theorem as well as several applications to the asymptotic distributions of linear statistics of the eigenvalues of the CUE.

One can interpret the law (1.1) of the eigenvalues of the CUE as a Gibbs distribution³ on \mathbb{T}^n with energy $\Phi(\boldsymbol{\theta}) := \sum_{1 \leq i < j \leq n} \log |2 \sin(\frac{\theta_i - \theta_j}{2})|^{-2}$. This implies that formally, (1.1) is the stationary measure of a diffusion with generator

$$\mathbb{L} = -\Delta + \nabla \Phi \cdot \nabla = - \sum_{j=1}^n \left(\partial_{jj} + \sum_{i \neq j} \frac{\partial_j}{\tan\left(\frac{\theta_j - \theta_i}{2}\right)} \right). \quad (7.2)$$

³This means that the probability measure (1.1) also describes a 2d Coulomb gas of N point charges confined on the unit circle at inverse temperature $\beta = 2$.

We view the vector $\mathbf{X} : \mathbb{T}^n \rightarrow \mathbb{R}^{2m}$ as a smooth function in $L^\infty(\mathbb{P}_n)$, so that we can define the vector \mathbf{LX} and the $2m \times 2m$ matrix

$$\mathbf{\Gamma}_{k,\ell} = \nabla X_k \cdot \nabla X_\ell. \quad (7.3)$$

Recall also the definition of the Kantorovich or Wasserstein distance (1.10). Then, by applying (Lambert et al., 2019, Corollary 2.4) to the random variable \mathbf{X} we obtain the following result.

Proposition 7.2. *For all $n, m \in \mathbb{N}$ and for any positive definite diagonal matrix \mathbf{K} of size $2m \times 2m$, we have*

$$W_2(\mathbf{X}, \mathbf{G}) \leq \sqrt{\mathbb{E}_n[|\mathbf{K}^{-1}\mathbf{LX} - \mathbf{X}|^2]} + \sqrt{\mathbb{E}_n[\|\mathbf{I} - \mathbf{K}^{-1}\mathbf{\Gamma}\|^2]}, \quad (7.4)$$

where $\|\cdot\|$ denotes the Hilbert–Schmidt norm.

The reason the RHS of (7.4) is small is because the random variables T_1, T_2, \dots are *approximate eigenfunctions* of the generator L and the matrix \mathbf{K} records the corresponding eigenvalues. The following Lemma makes this claim precise. Observe that $\boldsymbol{\xi}$ is small compared to \mathbf{K} which is of order n .

Lemma 7.3. *For all $n, m \in \mathbb{N}$, we have $\mathbf{LX} = \mathbf{KX} + \boldsymbol{\xi}$ where*

$$\begin{aligned} \mathbf{K} &= n \cdot \text{diag}(1, 1, 2, 2, \dots, m, m) \\ \boldsymbol{\xi} &= (\Re\zeta_1, \Im\zeta_1, \Re\zeta_2, \Im\zeta_2, \dots, \Re\zeta_m, \Im\zeta_m) \end{aligned}$$

and for all $k \geq 1$,

$$\zeta_k = \sqrt{\frac{k}{2}} \sum_{\ell=1}^{k-1} \sqrt{\ell(k-\ell)} T_\ell T_{k-\ell}.$$

Proof. The Lemma follows from the fact that $T_k = \sqrt{\frac{2}{k}} \sum_{j=1}^n e^{ik\theta_j}$ and explicit computations. Let us fix $k \in \mathbb{N}$ and observe that

$$\Delta T_k = -k^2 T_k. \quad (7.5)$$

Second, since $\tan\left(\frac{\theta_j - \theta_i}{2}\right) = -i \frac{e^{i\theta_j} - e^{i\theta_i}}{e^{i\theta_j} + e^{i\theta_i}}$ for any $i, j = 1, \dots, n$, we have

$$\sum_{i \neq j} \frac{\partial_j T_k}{\tan\left(\frac{\theta_j - \theta_i}{2}\right)} = -\sqrt{2k} \sum_{i \neq j} \frac{e^{i\theta_j} + e^{i\theta_i}}{e^{i\theta_j} - e^{i\theta_i}} e^{ik\theta_j}.$$

By symmetry, this implies that

$$\begin{aligned} \sum_{i \neq j} \frac{\partial_j T_k}{\tan\left(\frac{\theta_j - \theta_i}{2}\right)} &= -\sqrt{2k} \sum_{i \neq j} \frac{e^{ik\theta_j} - e^{ik\theta_i}}{e^{i\theta_j} - e^{i\theta_i}} e^{i\theta_j} \\ &= -\sqrt{2k} \sum_{i \neq j} \sum_{\ell=1}^k e^{i\ell\theta_j} e^{i(k-\ell)\theta_i} \\ &= -\sqrt{2k} \sum_{i,j} \sum_{\ell=1}^k e^{i\ell\theta_j} e^{i(k-\ell)\theta_i} + k^2 T_k, \end{aligned} \quad (7.6)$$

where we have used that $\frac{e^{ik\theta_j} - e^{ik\theta_i}}{e^{i\theta_j} - e^{i\theta_i}} = \sum_{\ell=1}^k e^{i(\ell-1)\theta_j} e^{i(k-\ell)\theta_i}$. Note that in the sum on the RHS of (7.6), the term $\ell = k$ equals to $-nkT_k$ while the other terms can be expressed in terms of the variables $(T_\ell)_{\ell=1}^{k-1}$. Hence, according to formula (7.2) and by combining formulae (7.5) and (7.6), this shows that for any $k \geq 1$,

$$LT_k = nkT_k + \zeta_k \quad (7.7)$$

Taking real and imaginary parts of the equation (7.7), this completes the proof. \square

Remarkably with Lemma 7.3 and Theorem 7.1, we can exactly compute the error terms on the RHS of the estimate (7.4). We obtain the following results.

Lemma 7.4. *For any $n, m \in \mathbb{N}$ such that $m \leq n$,*

$$\mathbb{E}_n [|\mathbf{K}^{-1}\boldsymbol{\xi}|^2] = \frac{(2m+5)m(m-1)}{9n^2}$$

Lemma 7.5. *For any $n, m \in \mathbb{N}$ such that $m \leq n/2$, we have*

$$\mathbb{E}_n [\|\mathbf{I} - \mathbf{K}^{-1}\boldsymbol{\Gamma}\|^2] = \frac{(8m+7)(m+1)m}{6n^2}.$$

Using Lemmas 7.4 and 7.5, we can complete the proof of Theorem 1.5. According to Lemma 7.3, we have

$$\mathbf{K}^{-1}\mathbf{L}\mathbf{X} - \mathbf{X} = \mathbf{K}^{-1}\boldsymbol{\xi},$$

so that for all $m, n \in \mathbb{N}$ such that $m \leq n/2$,

$$\begin{aligned} \sqrt{\mathbb{E}_n [|\mathbf{K}^{-1}\mathbf{L}\mathbf{X} - \mathbf{X}|^2]} + \sqrt{\mathbb{E}_n [\|\mathbf{I} - \mathbf{K}^{-1}\boldsymbol{\Gamma}\|^2]} &\leq \frac{\sqrt{(2m+5)m(m-1)} + \sqrt{(8m+7)(m+1)9m/6}}{3n} \\ &\leq (\sqrt{8} + \sqrt{2}) \frac{(m+1)\sqrt{m}}{3n}. \end{aligned}$$

By Proposition 7.2, we obtain the required bound for the Kantorovich distance $W_2(\mathbf{X}, \mathbf{G})$ between the random vector \mathbf{X} and a standard Gaussian random variable on \mathbb{R}^{2m} . Thus, to complete the proof, it remains to prove Lemmas 7.4 and 7.5.

Proof of Lemma 7.4. According to the notation of Lemma 7.3, we have

$$|\mathbf{K}^{-1}\boldsymbol{\xi}|^2 = \sum_{k=1}^m \frac{|\zeta_k|^2}{n^2 k^2} = \sum_{1 \leq \ell, \ell' < k \leq m} \frac{\sqrt{\ell(k-\ell)}\sqrt{\ell'(k-\ell')}}{2kn^2} \mathbf{T}_\ell \mathbf{T}_{k-\ell} \overline{\mathbf{T}_{\ell'} \mathbf{T}_{k-\ell'}}.$$

Moreover, according to Theorem 7.1, if $m \leq n$, then it holds for any integers $1 \leq \ell, \ell' < k \leq m$,

$$\begin{aligned} \mathbb{E}_n [\mathbf{T}_\ell \mathbf{T}_{k-\ell} \overline{\mathbf{T}_{\ell'} \mathbf{T}_{k-\ell'}}] &= (\mathbf{1}_{\{\ell=\ell', \ell \neq k/2\}} + \mathbf{1}_{\{\ell=k-\ell', \ell \neq k/2\}}) \mathbb{E}_n [|\mathbf{Z}_\ell|^2 |\mathbf{Z}_{k-\ell}|^2] + \mathbf{1}_{\ell=\ell'=k/2} \mathbb{E}_n [|\mathbf{Z}_\ell|^4] \\ &= 4(\mathbf{1}_{\{\ell=\ell', \ell \neq k/2\}} + \mathbf{1}_{\{\ell=k-\ell', \ell \neq k/2\}}) + 8\mathbf{1}_{\ell=\ell'=k/2}. \end{aligned}$$

This implies that

$$\mathbb{E}_n [|\mathbf{K}^{-1}\boldsymbol{\xi}|^2] = \frac{4}{n^2} \sum_{1 \leq \ell < k \leq m} \frac{\ell(k-\ell)}{k} = \frac{(2m+5)m(m-1)}{9n^2},$$

where we have used that $\sum_{1 \leq \ell < k} \frac{\ell(k-\ell)}{k} = \frac{k^2-1}{6}$. \square

Proof of Lemma 7.5. Let us decompose $\boldsymbol{\Gamma} = \tilde{\boldsymbol{\Gamma}} + \boldsymbol{\Delta}$ where $\boldsymbol{\Delta} = \text{diag}(\boldsymbol{\Gamma})$. The point is that

$$\|\mathbf{I} - \mathbf{K}^{-1}\boldsymbol{\Gamma}\|^2 = \|\mathbf{I} - \mathbf{K}^{-1}\boldsymbol{\Delta}\|^2 + \|\mathbf{K}^{-1}\tilde{\boldsymbol{\Gamma}}\|^2. \quad (7.8)$$

Since $\mathbf{X}_{2k-1} = \sqrt{\frac{2}{k}} \sum_{j=1}^n \cos(k\theta_j)$ and $\mathbf{X}_{2k} = \sqrt{\frac{2}{k}} \sum_{j=1}^n \sin(k\theta_j)$, by (7.3), we have for any $k = 1, \dots, m$,

$$\begin{aligned} \boldsymbol{\Gamma}_{2k-1, 2k-1} &= 2k \sum_{j=1}^n \sin^2(k\theta_j) = nk - \frac{k^{3/2}}{\sqrt{2}} \Re(\mathbf{T}_{2k}) \\ \boldsymbol{\Gamma}_{2k, 2k} &= 2k \sum_{j=1}^n \cos^2(k\theta_j) = nk + \frac{k^{3/2}}{\sqrt{2}} \Re(\mathbf{T}_{2k}). \end{aligned} \quad (7.9)$$

According to the notation of Lemma 7.3, this shows that

$$\|\mathbf{I} - \mathbf{K}^{-1}\mathbf{\Delta}\|^2 = \sum_{k=1}^m \frac{k}{n^2} \Re(\mathbf{T}_{2k})^2. \quad (7.10)$$

It remains to compute the second term on the RHS of (7.8). Let $\mathbf{K}^{1/2}$ be the positive square-root of the diagonal matrix \mathbf{K} and observe that by definition of the Hilbert Schmidt norm:

$$\|\mathbf{K}^{-1}\tilde{\mathbf{\Gamma}}\| = \|\mathbf{K}^{-1/2}\tilde{\mathbf{\Gamma}}\mathbf{K}^{-1/2}\|^2 = \frac{2}{n^2} \left(\sum_{1 \leq k < \ell \leq m} \frac{\mathbf{\Gamma}_{2\ell,2k}^2 + \mathbf{\Gamma}_{2\ell-1,2k-1}^2}{k\ell} + \sum_{1 \leq k \leq \ell \leq m} \frac{\mathbf{\Gamma}_{2\ell-1,2k}^2 + \mathbf{\Gamma}_{2\ell,2k-1}^2}{k\ell} \right). \quad (7.11)$$

Like (7.9), we can compute the coefficients on the RHS of (7.11). We check that for any $1 \leq k \leq \ell \leq m$,

$$\begin{aligned} \mathbf{\Gamma}_{2\ell,2k} &= 2\sqrt{k\ell} \sum_{j=1}^n \cos(k\theta_j) \cos(\ell\theta_j) = \sqrt{k\ell \left(\frac{\ell-k}{2}\right)} \Re(\mathbf{T}_{\ell-k}) + \sqrt{k\ell \left(\frac{\ell+k}{2}\right)} \Re(\mathbf{T}_{\ell+k}), \\ \mathbf{\Gamma}_{2\ell-1,2k-1} &= 2\sqrt{k\ell} \sum_{j=1}^n \sin(k\theta_j) \sin(\ell\theta_j) = \sqrt{k\ell \left(\frac{\ell-k}{2}\right)} \Re(\mathbf{T}_{\ell-k}) - \sqrt{k\ell \left(\frac{\ell+k}{2}\right)} \Re(\mathbf{T}_{\ell+k}), \\ \mathbf{\Gamma}_{2\ell-1,2k} &= -2\sqrt{k\ell} \sum_{j=1}^n \cos(k\theta_j) \sin(\ell\theta_j) = -\sqrt{k\ell \left(\frac{\ell-k}{2}\right)} \Im(\mathbf{T}_{\ell-k}) - \sqrt{k\ell \left(\frac{\ell+k}{2}\right)} \Im(\mathbf{T}_{\ell+k}), \\ \mathbf{\Gamma}_{2\ell,2k-1} &= -2\sqrt{k\ell} \sum_{j=1}^n \sin(k\theta_j) \cos(\ell\theta_j) = +\sqrt{k\ell \left(\frac{\ell-k}{2}\right)} \Im(\mathbf{T}_{\ell-k}) - \sqrt{k\ell \left(\frac{\ell+k}{2}\right)} \Im(\mathbf{T}_{\ell+k}). \end{aligned}$$

By (7.11), this implies that

$$\begin{aligned} \|\mathbf{K}^{-1}\tilde{\mathbf{\Gamma}}\|^2 &= \frac{2}{n^2} \left(\sum_{1 \leq k < \ell \leq m} ((\ell-k)\Re(\mathbf{T}_{\ell-k})^2 + (\ell+k)\Re(\mathbf{T}_{\ell+k})^2) + \sum_{1 \leq k \leq \ell \leq m} ((\ell-k)\Im(\mathbf{T}_{\ell-k})^2 + (\ell+k)\Im(\mathbf{T}_{\ell+k})^2) \right) \\ &= \frac{2}{n^2} \left(\sum_{1 \leq k < \ell \leq m} ((\ell-k)|\mathbf{T}_{\ell-k}|^2 + (\ell+k)|\mathbf{T}_{\ell+k}|^2) + 2 \sum_{k=1}^m k \Im(\mathbf{T}_{2k})^2 \right). \end{aligned}$$

Combining the previous formula with (7.8) and (7.10), we obtain

$$\mathbb{E}_n [\|\mathbf{I} - \mathbf{K}^{-1}\mathbf{\Gamma}\|^2] = \frac{2}{n^2} \left(\sum_{1 \leq k < \ell \leq m} (\ell-k)\mathbb{E}_n [|\mathbf{T}_{\ell-k}|^2] + (\ell+k)\mathbb{E}_n [|\mathbf{T}_{\ell+k}|^2] \right) + \frac{5}{n^2} \sum_{k=1}^m k \mathbb{E}_n [\Re(\mathbf{T}_{2k})^2],$$

where we used that the random variables $\Re(\mathbf{T}_k)$ and $\Im(\mathbf{T}_k)$ have the same law for all $k \geq 1$. Hence, by Theorem 7.1, we conclude that if $m \leq n/2$,

$$\mathbb{E}_n [\|\mathbf{I} - \mathbf{K}^{-1}\mathbf{\Gamma}\|^2] = \sum_{1 \leq k < \ell \leq m} \frac{4\ell}{n^2} + \sum_{k=1}^m \frac{5k}{n^2} = \frac{(8m+7)(m+1)m}{6n^2}.$$

This completes the proof. \square

A Additional proofs

A.1 Proof of Lemma 2.3

Without loss of generality, we assume that $\hat{f}_0 = 0$, then by (2.9), we have

$$\mathbb{E}_n [\exp \operatorname{Tr} f(\mathbf{U})] = e^{\mathcal{A}(f)} \det[\mathbf{I} - K_f Q_n], \quad (A.1)$$

where according to (2.8), if we let $w = e^{-2i\Im(f^+)}$, the kernel K_f is given by

$$K_f = H_+(w)H_-(\bar{w}) = H(w)H(w)^*,$$

where $H(w)^*$ is the adjoint of $H(w)$. Therefore, $K_f > 0$ as a trace-class operator and this implies that for any $n \in \mathbb{Z}_+$,

$$0 < \det[\mathbf{I} - K_f Q_n] \leq 1.$$

Then, the claim follows directly from (A.1).

A.2 Proof of Lemma 3.1

Recall that for any $m \in \mathbb{N}$ and $\Lambda > 0$, we let

$$g_m(\Lambda) = e^{-\Lambda^2} \sum_{0 \leq k < m} m^k \Lambda^{-2(k+1)}.$$

First, by going to polar coordinates and making the change of variable $u = \|\xi\|^2$, we have for any $\Lambda > 0$,

$$\int_{\substack{\mathbb{R}^{2m} \\ \|\xi\| \geq \Lambda}} e^{-\|\xi\|^2} d\xi = \Omega_m \int_{\Lambda^2}^{+\infty} e^{-u} d(u^m)$$

The integral on the RHS corresponds to the incomplete Gamma function – see (DLMF, Formula (8.2.2)) – and repeated integrations by parts give for any $\lambda > 0$,

$$\int_{\lambda}^{+\infty} e^{-u} d(u^m) = (m-1)! e^{-\lambda} \sum_{0 \leq k < m} \frac{\lambda^k}{k!}.$$

Using the bound $(m-k)! \geq \frac{(m-1)!}{m^{k-1}}$ valid for all $k = 1, \dots, m$, this implies that

$$\int_{\substack{\mathbb{R}^{2m} \\ \|\xi\| \geq \Lambda}} e^{-\|\xi\|^2} d\xi \leq \Omega_m e^{-\Lambda^2} \sum_{1 \leq k \leq m} m^{k-1} \Lambda^{2k} = \Omega_m g_m(\Lambda).$$

Finally, if $\Lambda^2 > m$, by summing the geometric sum, we obtain $g_m(\Lambda) \leq \frac{e^{-\Lambda^2}}{\Lambda^2 - m}$.

A.3 Proof of Lemma 5.2

By symmetry, it suffices to prove that for all $y \in (0, 1]$ and $x \geq 0$,

$$1 + \left(\frac{\sinh(x)}{y} \right)^2 \leq \exp\left(\frac{x^2}{y} \right).$$

We have for any fixed $y \in (0, 1]$ and $x \geq 0$,

$$\begin{aligned} \frac{d}{dx} \left(\left(1 + \left(\frac{\sinh(x)}{y} \right)^2 \right) \exp\left(-\frac{x^2}{y^2} \right) \right) &= -\frac{2}{y^2} \exp\left(-\frac{x^2}{y^2} \right) \left(x \left(1 + \left(\frac{\sinh(x)}{y} \right)^2 \right) - \sinh(x) \cosh(x) \right) \\ &\leq -\frac{2}{y^2} \exp\left(-\frac{x^2}{y^2} \right) \left(x (1 + \sinh(x)^2) - \sinh(x) \cosh(x) \right) \\ &\leq -\frac{2}{y^2} \exp\left(-\frac{x^2}{y^2} \right) \cosh(x) \left(x \cosh(x) - \sinh(x) \right). \end{aligned}$$

Since $x \cosh(x) - \sinh(x) \geq 0$, this shows that for any fixed $y \in (0, 1]$ and $x \geq 0$

$$\frac{d}{dx} \left(\left(1 + \left(\frac{\sinh(x)}{y} \right)^2 \right) \exp\left(-\frac{x^2}{y^2} \right) \right) \leq 0.$$

This implies that for any $y \in (0, 1]$

$$\max_{x>0} \left\{ \left(1 + \left(\frac{\sinh(x)}{y} \right)^2 \right) \exp \left(-\frac{x^2}{y^2} \right) \right\} = 1.$$

Since the RHS is independent of $y \in (0, 1]$, this completes the proof.

A.4 Proof of Lemma 5.3

Let us define the function $\Phi(\boldsymbol{\theta}) = \sum_{1 \leq i < j \leq n} \log |e^{i\theta_i} - e^{i\theta_j}|^{-2}$ for $\boldsymbol{\theta} \in \Delta$ where $\Delta := \{\boldsymbol{\theta} \in \mathbb{R}^n : \theta_1 = 0 < \theta_2 < \dots < \theta_n < 2\pi\}$ is a convex set. Observe that by symmetry, we have

$$\max_{\theta_1, \dots, \theta_n \in \mathbb{T}} \left(\prod_{1 \leq i < j \leq n} |e^{i\theta_i} - e^{i\theta_j}|^2 \right) = \max_{\boldsymbol{\theta} \in \Delta} (e^{-\Phi(\boldsymbol{\theta})}) = e^{-\min_{\boldsymbol{\theta} \in \Delta} \Phi(\boldsymbol{\theta})}.$$

Since function Φ is smooth on Δ , by computing its Hessian (with respect to $\theta_2, \dots, \theta_n$), we verify that Φ is strictly convex⁴. Moreover, if we let $\boldsymbol{\vartheta} = (0, \frac{2\pi}{n}, \dots, \frac{2\pi(n-1)}{n})$, we see that by symmetry for any $j = 2, \dots, n$,

$$\nabla_j \Phi(\boldsymbol{\vartheta}) = \sum_{i \neq j} \frac{1}{\tan(\frac{\vartheta_i - \vartheta_j}{2})} = \sum_{i \neq j} \frac{1}{\tan(\pi \frac{i-j}{n})} = 0.$$

This implies that $\boldsymbol{\vartheta}$ is the only critical point of Φ inside Δ and since $\Phi = +\infty$ on $\partial\Delta$, we have

$$\min_{\boldsymbol{\theta} \in \Delta} \Phi(\boldsymbol{\theta}) = \Phi(\boldsymbol{\vartheta}).$$

Moreover, by definition of the Vandermonde determinant,

$$e^{-\Phi(\boldsymbol{\vartheta})} = \prod_{1 \leq i < j \leq n} |e^{i\vartheta_i} - e^{i\vartheta_j}|^2 = \left| \det_{n \times n} (e^{i(j-1)\vartheta_i}) \right|^2 = \left| \det_{n \times n} (e^{i2\pi \frac{(j-1)(i-1)}{n}}) \right|^2.$$

This shows that for any $n \geq 2$,

$$\max_{\theta_1, \dots, \theta_n \in \mathbb{T}} \prod_{1 \leq i < j \leq n} |e^{i\theta_i} - e^{i\theta_j}|^2 = e^{-\Phi(\boldsymbol{\vartheta})} = n^n \left| \det_{n \times n} A \right|^2,$$

where $A_{ij} = \frac{e^{i2\pi \frac{(j-1)(i-1)}{n}}}{\sqrt{n}}$. We easily verify that the columns of the matrix A are orthonormal so that A is a unitary matrix and $|\det_{n \times n} A| = 1$. This proves that for any integer $n \geq 2$,

$$\max_{\theta_1, \dots, \theta_n \in \mathbb{T}} \prod_{1 \leq i < j \leq n} |e^{i\theta_i} - e^{i\theta_j}|^2 = n^n.$$

We immediately deduce from this fact and formula (1.1) for the joint density of \mathbb{P}_n that

$$\begin{aligned} \mathbb{E}_n \left[e^{-\sum_{j=1}^n f(\theta_j)} \right] &= \frac{1}{n!} \int_{\mathbb{T}^n} \prod_{1 \leq i < j \leq n} |e^{i\theta_i} - e^{i\theta_j}|^2 e^{-\sum_{j=1}^n f(\theta_j)} \frac{d\theta_1}{2\pi} \dots \frac{d\theta_n}{2\pi} \\ &\leq \frac{e^n}{\sqrt{2\pi n}} \left(\int_{\mathbb{T}} e^{-f(\theta)} \frac{d\theta}{2\pi} \right)^n, \end{aligned}$$

where we used that $\frac{n^n}{n!} \leq \frac{e^n}{\sqrt{2\pi n}}$ by (B.17).

⁴This follows from the fact that the Hessian $\nabla^2 \Phi$ has a strictly dominant diagonal with positive entries on Δ .

B Constants and numerical approximations

As we pointed out in the introduction, one of the main challenge of the proof of Theorem 1.3 is to try to optimize and keep track of all the constants involved in our different estimates. For the convenience of the readers, these constants as well as the error terms in Theorem 1.3 are collected in this section. The constants are denoted by $c_j = c_j(m)$, $\epsilon_j = \epsilon_j(m)$ and $\Upsilon_j = \Upsilon_j(m)$ for $j \in \mathbb{N}_0$ since they are allowed to depend on m but not on the dimension n the random matrix. They are positive for all $m \geq 3$ and we use the following conventions:

- $c_j(m) \rightarrow \widehat{c}_j$ as $m \rightarrow +\infty$ where $\widehat{c}_j > 0$.
- $\epsilon_j(m) \rightarrow 0$ as $m \rightarrow +\infty$.
- $\Upsilon_j(m) \rightarrow +\infty$ as $m \rightarrow +\infty$ and $\Upsilon_j(m)$ is a regularly varying function.

When relevant, we also provide a numerical approximation or an estimate for these constants.

B.1 Errors in Theorem 1.3

We let

$$c_0 = \sqrt{\frac{1}{6\sqrt{2}}} \approx 0.343 \quad (\text{B.1})$$

and

$$\epsilon_0(m) = \frac{2c_0^2}{3(1+1/m)\sqrt{1+\log m}}. \quad (\text{B.2})$$

We note that $\epsilon_0(m) \leq 0.041$ for all $m \geq 3$. The constants which are directly involved in $\Theta_{N,m}$ from Theorem 1.3 are given by

$$\begin{aligned} c_1(m) &= \frac{(1-c_{10})^2}{16c_{11}} - c_0^2(1+\epsilon_0)\frac{\sqrt{1+\log m}}{2(m+1)} \\ c_2(m) &= c_0c_4 \left(1 - c_{10} - \frac{4c_4c_0c_{11}(1+\log m)^{1/4}}{\sqrt{m+1}} \right) - c_0^2 \frac{(1+\epsilon_0)(1+\log m)^{1/4}}{2\sqrt{m+1}} \\ c_3 &= c_8/(2\pi)^{1/4} \approx 7.98 \\ c_4 &= \frac{1}{2\sqrt{2}} \approx 0.354 \\ c_5 &= c_3^{-1}e^{c_9} \approx 1.147 \\ c_6 &= 4(2+\log 2) \approx 10.78 \\ c_7 &= \frac{\pi\sqrt{e}}{2} \approx 2.59 \\ c_8 &= \frac{4 \cdot 2.766}{(1-1/16)^2} \approx 12.63 \\ c_9 &= \frac{538}{243} \approx 2.21 \\ c_{10}(m) &= c_0^2c_{19}(m)\sqrt{1+1/m} = c_0^2 \frac{\sqrt{1+1/m}}{3\sqrt{6}} e^{c_0 \frac{(1+\log m)^{1/4}}{\sqrt{(m+1)/2}}}, \quad \widehat{c}_{10} = \frac{c_0^2}{3\sqrt{6}} \approx 0.016 \\ c_{11}(m) &= \left(1 + \frac{c_{10}}{\sqrt{m}} \right) \left(1 + \frac{2c_{10}}{m} + \sqrt{\frac{1+1/m}{2(1+\log m)}} \right). \end{aligned} \quad (\text{B.3})$$

We verify that both $c_1(m)$ and $c_2(m)$ are increasing for $m \geq 3$ and we have

$$\widehat{c}_1 = (1 - \widehat{c}_{10})^2/16 \approx 0.0605, \quad \widehat{c}_2 = 4c_0c_4\sqrt{\widehat{c}_1} \approx 0.119.$$

Note also that the convergence is slow since $c_j = \widehat{c}_j + \mathcal{O}\left(\frac{(1+\log m)^{1/4}}{\sqrt{(m+1)}}\right)$ for $j = 1, 2$ as $m \rightarrow +\infty$.

Let us also define for all $m \geq 1$,

$$\begin{aligned}\Upsilon_1(m) &= c_6 m^2 + \frac{5}{2} m \log m + \log(c_7) m + \frac{3}{2} \\ \Upsilon_2(m) &= \frac{1}{2} m \log m - \frac{3}{4} m \log(1 + \log m) - m \log(8c_0) + \frac{1}{2}.\end{aligned}\tag{B.4}$$

It will turn out that we need the following estimates for the functions $\Upsilon_1(m)$ and $\Upsilon_2(m)$. We have for all $m \geq 3$,

$$\sqrt{c_1(m)^{-1}(1 + \log m)\Upsilon_1(m)} \geq 34m\tag{B.5}$$

and

$$\frac{c_1(m)\Upsilon_2(m)\sqrt{m+1}}{c_2(m)(1 + \log m)^{1/4}} \leq \frac{\Upsilon_1(m)}{1500}.\tag{B.6}$$

The numerical constants in (B.5) and (B.6) are not optimal but they suffice for our applications.

For any $N, m \geq 1$, let us define the following functions:

$$\Theta_{N,m}^0 = m^{\frac{5}{2}} 2^{\frac{m}{2}} e^{\frac{m^2}{4N}} \frac{e^{\frac{N}{2}} (1 + \log m)^N}{\sqrt{N} \Gamma(N+1)}, \quad \Theta_{N,m}^3 = \frac{c_3^{-1}}{\sqrt{m}} N^{-\frac{m}{2}} \exp\left(-\frac{N^2}{16(1 + \log m)}\right),\tag{B.7}$$

$$\Theta_{N,m}^2 = c_5 N^{\frac{m}{2}} \exp\left(\Upsilon_2(m) - \frac{c_2(m)N^2}{\sqrt{m+1}(1 + \log m)^{\frac{3}{4}}}\right),\tag{B.8}$$

and if $N > 4m$,

$$\Theta_{N,m}^1 = \frac{c_5 e^{-c_9 \frac{4m}{N}}}{(1 - \frac{4m}{N})^{1 - \frac{2m}{N}}} \exp\left(\Upsilon_1(m) - c_1(m) \frac{N(N-4m)}{(1 + \log m)}\right).\tag{B.9}$$

Then, the error in Theorem 1.3 is given by

$$\Theta_{N,m} = \Theta_{N,m}^0 + \Theta_{N,m}^1 + \Theta_{N,m}^2 + \Theta_{N,m}^3.\tag{B.10}$$

One should keep in mind that $\Theta_{N,m}^0$ is the main term, the term $\Theta_{N,m}^3$ is always negligible, while $\Theta_{N,m}^1$ and $\Theta_{N,m}^2$ are corrections which become negligible when $m \ll N$. This is quantified by Proposition B.1 below.

In Sections 2–6, the following constants will come in play.

$$\begin{aligned}c_{12} &= \frac{1 + \sqrt{290}}{17} \approx 1.06 \\ c_{13} &= \frac{(\log 108)\sqrt{1 + \log 3}}{68\sqrt{108}c_2(m)} \\ c_{14} &= \frac{\sqrt{13}}{1500} \approx 0.0024 \\ c_{15} &= 2e^2 \approx 14.78 \\ c_{16} &= 2e\sqrt{\pi} \approx 9.64 \\ c_{17} &= \frac{32}{3} \left(1 + \frac{(2 - 1/m)^3}{3(m+1)}\right) \\ c_{18} &= \frac{8}{3} \left(1 + \frac{4(1 - 1/m)^3}{3(m+1)}\right)\end{aligned}\tag{B.11}$$

$$\begin{aligned}
c_{19}(m) &= \frac{1}{3\sqrt{6}} e^{\sqrt{2}c_0 \frac{(1+\log m)^{1/4}}{\sqrt{m+1}}} \\
c_{20}(\eta) &= \frac{\pi^2 \eta^2}{8}, \quad \eta = \frac{1/\pi}{\sqrt{m}} \\
c_{21}(m, \eta) &= \frac{\exp\left(\frac{\eta}{2\sqrt{m(m+1)}}\right)}{6\sqrt{3}}, \quad \eta = \frac{1/\pi}{\sqrt{m}} \\
\Upsilon_3(m) &= \frac{\pi m^{3/2}(m+1)e^{\frac{1}{2}\left(1+\frac{1/2}{(\pi m)^2}\right)}}{2\left(1-\frac{c_{21}}{4\pi^2 m^3}\right)}.
\end{aligned} \tag{B.12}$$

Observe that $\Upsilon_3(m) = c_7 m^{\frac{5}{2}}(1 + \mathcal{O}(m^{-1}))$ as $m \rightarrow +\infty$. Moreover, as $c_{21} \leq \frac{1}{12}$, we verify that for all $m \geq 3$,

$$\Upsilon_3^m \leq c_7^m m^{\frac{5m}{2}} \frac{e^{\frac{1}{4\pi^2 m}(1+1/m)^m}}{\left(1-\frac{m^{-3}}{48\pi^2}\right)^m} \leq e c_7^m m^{\frac{5m}{2}}. \tag{B.13}$$

B.2 Estimates for errors – Proof of Proposition 1.4.

Proposition B.1. Fix $\gamma > c_{12} = \frac{1+\sqrt{290}}{17}$. For all $N, m \geq 3$ such that $N \geq \gamma\sqrt{c_1(m)^{-1}(1+\log m)\Upsilon_1(m)}$, we have the estimates

$$\Theta_{N,m}^1 \leq c_5 \exp\left(-\left(1-\frac{2\gamma^{-1}}{17}-\gamma^{-2}\right)\frac{c_1(m)N^2}{1+\log m}\right) \tag{B.14}$$

and

$$\Theta_{N,m}^2 \leq c_5 N^{\frac{m}{2}} \exp\left(-\frac{\left(1-\frac{\gamma^{-2}}{1500}\right)c_2(m)N^2}{\sqrt{m+1}(1+\log m)^{\frac{3}{4}}}\right) \leq c_5 \exp\left(-\left(\sqrt{13}\gamma-c_{13}\gamma^{-1}-c_{14}\gamma^{-\frac{3}{2}}\right)\frac{c_2(m)N^{\frac{3}{2}}}{\sqrt{1+\log m}}\right). \tag{B.15}$$

Moreover, we have the lower-bounds: $c_1(m) \geq 0.0148$, $c_2(m) \geq 0.077$ and $c_{13} \leq 0.125$ for all $m \geq 3$.

Proof. Since $e^{c_9} \geq 4$, we verify that the function $x \mapsto e^{-c_9 x}(1-x)^{1-2x}$ is decreasing on $[0, \frac{1}{2}]$ so that we deduce from (B.9) that for all $N \geq 8m$,

$$\Theta_{N,m}^1 \leq c_5 \exp\left(\Upsilon_1(m) - c_1(m)\frac{N(N-4m)}{(1+\log m)}\right).$$

Then, we verify that if the condition $N \geq \gamma\sqrt{c_1(m)^{-1}(1+\log m)\Upsilon_1(m)}$ holds with $\gamma > 0$,

$$c_1(m)\frac{N(N-4m)}{(1+\log m)} - \Upsilon_1(m) \geq \frac{c_1(m)N^2}{(1+\log m)}\left(1-\frac{2\gamma^{-1}}{17}-\gamma^{-2}\right), \tag{B.16}$$

where we used the lower-bound (B.5). The RHS of (B.16) is positive so long as $\gamma > c_{12} = \frac{1+\sqrt{290}}{17}$ and this yields the estimate (B.14). For the estimate (B.15), let us also observe that according to (B.6), we have for all $N, m \geq 3$ such that $N \geq \gamma\sqrt{c_1(m)^{-1}(1+\log m)\Upsilon_1(m)}$,

$$c_2(m)^{-1}\Upsilon_2(m)\sqrt{m+1}(1+\log m)^{3/4} \leq \frac{\gamma^{-2}}{1500}N^2.$$

This implies that

$$\Theta_{N,m}^2 \leq c_5 N^{\frac{m}{2}} \exp\left(-\left(1-\frac{\gamma^{-2}}{1500}\right)\frac{c_2(m)N^2}{\sqrt{m+1}(1+\log m)^{\frac{3}{4}}}\right).$$

Moreover, we also verify that for all $m \geq 3$,

$$(m+1)^2 \leq \frac{c_1(m)^{-1} \Upsilon_1(m)}{13^2}$$

so that under our hypothesis,

$$\frac{N^2}{\sqrt{m+1}(1+\log m)^{\frac{3}{4}}} \geq \frac{\sqrt{13} N^2}{\sqrt{1+\log m} (c_1(m)^{-1}(1+\log m) \Upsilon_1(m))^{\frac{1}{4}}} \geq \frac{\sqrt{13\gamma} N^{3/2}}{\sqrt{1+\log m}}.$$

Hence, if we agree to loose the Gaussian decay in N of $\Theta_{N,m}^2$, we obtain that for all $N, m \geq 3$ such that $N \geq \gamma \sqrt{c_1(m)^{-1}(1+\log m) \Upsilon_1(m)}$,

$$\Theta_{N,m}^2 \leq c_5 N^{\frac{m}{2}} \exp\left(-\left(\sqrt{13\gamma} - \frac{\sqrt{13}}{1500} \gamma^{-\frac{3}{2}}\right) \frac{c_2(m) N^{\frac{3}{2}}}{\sqrt{1+\log m}}\right).$$

Finally, it follows from (B.5) that under our hypothesis, $N \geq 34\gamma m$ with $\gamma > c_{12}$ so that $N \geq 108$ and

$$N^{\frac{m}{2}} \leq \exp\left(\frac{N \log N}{68\gamma}\right) \leq \exp\left(\frac{N^{\frac{3}{2}}}{\sqrt{1+\log m}} \frac{\log N \sqrt{1+\log(N/36)}}{68\gamma\sqrt{N}}\right) \leq \exp\left(\frac{c_{13} c_2(m) N^{\frac{3}{2}}}{\gamma \sqrt{1+\log m}}\right),$$

where $c_{13} = \frac{(\log 108)\sqrt{1+\log 3}}{68\sqrt{108}c_2(m)}$. This yields the estimate (B.15). Since c_1, c_2 are increasing functions for $m \geq 3$, we obtain the numerical estimates for c_1, c_2 and c_{13} by evaluating these functions for $m = 3$ on *Mathematica*. This completes the proof. \square

We will also need the following basic estimates for the main error term $\Theta_{N,m}^0$.

Lemma B.2. *For all $m, N \in \mathbb{N}$ such that $m \geq 3$ and $N \geq 5m$, it holds*

$$\Theta_{N,m}^0 \leq \frac{1}{\sqrt{\pi}} \exp\left(-N \log m \left(1 - \frac{\log(1+\log m)}{\log m}\right)\right).$$

Proof. Let us recall from (DLMF, Formula (5.6.1)) that for any $x > 0$,

$$\Gamma(x+1) = \sqrt{2\pi x} x^x \exp\left(-x + \frac{\theta_x}{12x}\right) \quad \text{where } \theta_x \in (0, 1). \quad (\text{B.17})$$

In addition, since $e^{\frac{3}{2}} \leq 5$, let us observe that we have for all $N \geq 5m$,

$$\frac{m^{\frac{3}{2}} 2^{\frac{m}{2}} e^{\frac{m^2}{4N}} e^{\frac{3N}{2}}}{\sqrt{2\pi} 5^{N+1}} \leq \frac{m^{\frac{3}{2}} e^{-cm}}{5\sqrt{2\pi}} \leq \frac{1}{\sqrt{\pi}},$$

where we used that $c = 5 \log 5 - \frac{15}{2} - \frac{1}{20} - \frac{\log 2}{2} \geq 0.15$. By (B.17), this implies that for all $N \geq 5m$,

$$\Theta_{N,m}^0 \leq \frac{m^{\frac{3}{2}} 2^{\frac{m}{2}} e^{\frac{m^2}{4N}} e^{\frac{3N}{2}}}{\sqrt{2\pi} 5^{N+1}} \frac{(1+\log m)^N}{m^N} \leq \frac{1}{\sqrt{\pi}} \exp\left(-N \log m \left(1 - \frac{\log(1+\log m)}{\log m}\right)\right).$$

\square

Using the previous estimates, we are now ready to prove Proposition 1.4.

Proposition B.3. *Fix $M \geq 3$. For all $m \geq M$ and $N \geq c(M)m\sqrt{1+\log m}$, we have*

$$\Theta_{N,m}^1 + \Theta_{N,m}^2 + \Theta_{N,m}^3 \leq 0.011 \frac{(1+\log m)^N e^{\frac{N}{2}}}{\sqrt{N} \Gamma(N+1)} \leq \epsilon \Theta_{N,m}^0 \quad (\text{B.18})$$

where $\epsilon \leq 25 \cdot 10^{-5}$ and the constant $c(M)$ are explicitly given by the Table (B.22) below. Moreover, under the same conditions, we also have $\Theta_{N,m}^0 \leq N^{-\frac{m}{2}} \frac{\exp(-12m(\log m - 0.26))}{\sqrt{\pi c_{16}^m}}$.

Proof. First, observe that for all $N \geq \gamma \sqrt{c_1(m)^{-1}(1 + \log m) \Upsilon_1(m)}$, if $\theta(m) := \sqrt{\frac{c_1(m) \Upsilon_1(m)}{1 + \log m}} \geq (\gamma - \frac{2}{17} - \gamma^{-1})^{-1}$, then it holds that

$$\frac{e^{-3/2} \cdot N}{1 + \log m} \exp\left(-\frac{(1 - \frac{2\gamma^{-1}}{17} - \gamma^{-2})c_1 N}{1 + \log m}\right) \leq c_1^{-1} \gamma \theta e^{-3/2 - (\gamma - 2/17 - \gamma^{-1})\theta}.$$

Let us suppose that $\gamma \leq 5.12$. This shows that if we choose γ depending on $m \geq 3$ in such a way that

$$(\gamma - 2/17 - \gamma^{-1})\theta \geq \log(5.12c_1^{-1}\theta) - 1.48 > 0, \quad (\text{B.19})$$

then we have

$$\frac{e^{-3/2} \cdot N}{1 + \log m} \exp\left(-\frac{(1 - \frac{2\gamma^{-1}}{17} - \gamma^{-2})c_1 N}{1 + \log m}\right) \leq e^{-0.02}.$$

Any solution of (B.19) satisfies $\gamma > c_{12}$ and we can choose a (numerical) solution $\gamma(m)$ which is non-increasing in the following way:

m	3	4	5	6	8	12	17	23	30	40	≥ 70		(B.20)
$\gamma(m)$	5.119	3.806	3.149	2.754	2.30	1.882	1.65	1.507	1.413	1.334	1.230		

Observe that the function $m^{-1} \sqrt{c_1(m)^{-1} \Upsilon_1(m)}$ is also decreasing for $m \geq 3$. By (B.14), this implies that for any $M \geq 3$, if $m \geq M$ and $N \geq c(M)m\sqrt{1 + \log m}$, then we obtain

$$\begin{aligned} \Theta_{N,m}^1 &\leq c_5 \exp\left(-\frac{(1 - \frac{2\gamma^{-1}}{17} - \gamma^{-2})c_1 N^2}{1 + \log m}\right) \\ &\leq c_5 e^{-0.02N} \left(\frac{e^{-3/2} \cdot N}{1 + \log m}\right)^{-N} \end{aligned} \quad (\text{B.21})$$

where $c(M) = \gamma(M)M^{-1} \sqrt{c_1(M)^{-1} \Upsilon_1(M)}$ is a decreasing function

M	3	4	5	6	8	12	17	23	30	40	70		(B.22)
$c(M)$	146.5	93.8	71.1	58.66	45.5	34.5	28.8	25.5	23.4	21.64	19.4		

Moreover, since $N \geq 600$ in the regime that we consider, we also verify that

$$N e^{-0.02N} \leq 600 e^{-12} \leq 3.7 \cdot 10^{-3},$$

so that by (B.21), this implies that for $m \geq M$ and $N \geq c(M)m\sqrt{1 + \log m}$,

$$\Theta_{N,m}^1 \leq 0.0107 \frac{(1 + \log m)^N e^{\frac{N}{2}}}{\sqrt{N} \Gamma(N + 1)} \quad (\text{B.23})$$

where we used that according to (B.17), $\Gamma(N + 1) \leq 2.52 N^{N+1/2} e^{-N}$. By a similar argument, we have $\Theta_{N,m}^3 \leq 0.073 \exp\left(-\frac{0.0625 \cdot N^2}{1 + \log m}\right)$ and $\frac{N}{1 + \log m} \geq \min_{M \geq 3} \frac{c(M)M}{\sqrt{1 + \log M}} \geq 200$, so that

$$\frac{e^{-3/2} \cdot N}{1 + \log m} \exp\left(-\frac{0.0625 \cdot N}{1 + \log m}\right) \leq 200 e^{-11} \leq 3.4 \cdot 10^{-3}$$

and

$$\Theta_{N,m}^3 \leq 0.2N (3.4 \cdot 10^{-3})^N \frac{(1 + \log m)^N e^{\frac{N}{2}}}{\sqrt{N} \Gamma(N + 1)}. \quad (\text{B.24})$$

Using the estimate (B.15) and the fact that according to the Table (B.20) $\min_{m \geq 3} \{(\sqrt{13\gamma(m)} - c_{13}/\gamma(m) - c_{14}/\gamma(m)^{\frac{3}{2}})c_2(m)\} \geq 0.422$, we obtain the estimate

$$\Theta_{N,m}^2 \leq c_5 \exp\left(-\frac{0.422 \cdot N^{\frac{3}{2}}}{\sqrt{1 + \log m}}\right).$$

Since we have seen that $\frac{N}{1 + \log m} \geq 200$, this implies that

$$\frac{N}{1 + \log m} \exp\left(-\frac{0.422\sqrt{N}}{\sqrt{1 + \log m}}\right) \leq \max_{x \geq \sqrt{200}} \{x^2 e^{-0.422x}\} \leq 1.$$

So, using the same argument once more, we obtain that for any $m \geq M$ and $N \geq c(M)m\sqrt{1 + \log m}$,

$$\Theta_{N,m}^2 \leq 3N e^{-3N/2} \frac{(1 + \log m)^N e^{\frac{N}{2}}}{\sqrt{N}\Gamma(N+1)}. \quad (\text{B.25})$$

By combining the estimates (B.23), (B.24) and (B.25), we easily verify that for any $m \geq M$ and for all $N \geq c(M)m\sqrt{1 + \log m}$,

$$\Theta_{N,m}^1 + \Theta_{N,m}^2 + \Theta_{N,m}^3 \leq 0.011 \frac{(1 + \log m)^N e^{\frac{N}{2}}}{\sqrt{N}\Gamma(N+1)}.$$

Then, from (B.7), we deduce the bound (B.18) with $\epsilon \leq 0.011 \cdot 3^{-\frac{5}{2}} 2^{-\frac{3}{2}} \leq 25 \cdot 10^{-5}$. Finally, it remains to obtain the upper-bound for $\Theta_{N,m}^0$. According to Lemma B.2, we have for all $m \geq 3$ and $N \geq 5m$,

$$N^{\frac{m}{2}} \Theta_{N,m}^0 \leq \frac{1}{\sqrt{\pi}} \exp(-0.3N \log m + 0.5m \log N).$$

Since the function $N \mapsto 0.3N \log m - 0.5m \log N$ is increasing and $\min_{M \geq 3} c(M)\sqrt{1 + \log M} \geq 42$, this implies that for $N \geq 42m$,

$$N^{\frac{m}{2}} \Theta_{N,m}^0 \leq \frac{\exp(-12m \log m + \frac{\log 42}{2}m)}{\sqrt{\pi}} \leq \frac{\exp(-12m(\log m - 0.26))}{\sqrt{\pi}c_{16}^m},$$

where we used that $\frac{\log(42c_{16})}{24} \leq 0.26$. □

B.3 Numerics for $m = 3$

Figure 1: Log-Log plot of the errors (B.7)–(B.8) for $m = 3$ as functions of $N = n/3$ where n is the dimension of the random unitary matrix. We observe that $\Theta_{N,3}^0 \geq \Theta_{N,3}^1$ when $N \geq 631$ which is consistent with the threshold $3c(3)\sqrt{1 + \log 3} \approx 637$ from Proposition B.3.





Figure 2: Plot of $\log(\Delta_{n,3}^{(2)})$ as a function of the dimension n of the random unitary matrix. By Theorem 1.3, this quantity controls the total variation distance between \mathbf{X} and a standard Gaussian vector in \mathbb{R}^6 . We observe that our estimates become relevant as soon as $n \geq 400$ which can still be considered a small size random matrix.



References

- E. L. Basor and H. Widom. On a Toeplitz determinant identity of Borodin and Okounkov. *Integral Equations Operator Theory*, 37(4):397–401, 2000. ISSN 0378-620X. doi: 10.1007/BF01192828. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=1780119>.
- G. Baxter. Polynomials defined by a difference system. *J. Math. Anal. Appl.*, 2:223–263, 1961. ISSN 0022-247x. doi: 10.1016/0022-247X(61)90033-6. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=0126125>.
- S. Berezin and A. I. Bufetov. On the rate of convergence in the central limit theorem for linear statistics of Gaussian, Laguerre, and Jacobi ensembles. 2019. URL <https://arxiv.org/pdf/1904.09685.pdf>.
- A. Borodin and A. Okounkov. A Fredholm determinant formula for Toeplitz determinants. *Integral Equations Operator Theory*, 37(4):386–396, 2000. ISSN 0378-620X. doi: 10.1007/BF01192827. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=1780118>.
- A. Böttcher. On the determinant formulas by Borodin, Okounkov, Baik, Deift and Rains. In *Toeplitz matrices and singular integral equations (Pobershau, 2001)*, volume 135 of *Oper. Theory Adv. Appl.*, pages 91–99. Birkhäuser, Basel, 2002. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=1935759>.
- M. A. Chahkiev. On oscillatory integral Hilbert transformation with trigonometric polynomial phase. *Anal. Math.*, 34(3):177–185, 2008. ISSN 0133-3852. doi: 10.1007/s10476-008-0302-7. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=2434672>.
- K. Courteaut and K. Johansson. Multivariate normal approximation for traces of orthogonal and symplectic matrices. in preparation.
- K. Courteaut, K. Johansson, and G. Lambert. in preparation.
- P. Deift, A. Its, and I. Krasovsky. Toeplitz matrices and Toeplitz determinants under the impetus of the Ising model: some history and some recent results. *Comm. Pure Appl. Math.*, 66(9):1360–1438, 2013. ISSN 0010-3640. doi: 10.1002/cpa.21467. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=3078693>.
- P. Diaconis. personal communication, 1994.
- P. Diaconis and S. N. Evans. Linear functionals of eigenvalues of random matrices. *Trans. Amer. Math. Soc.*, 353(7):2615–2633, 2001. ISSN 0002-9947. doi: 10.1090/S0002-9947-01-02800-8. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=1828463>.

- P. Diaconis and M. Shahshahani. On the eigenvalues of random matrices. *J. Appl. Probab.*, 31A:49–62, 1994. ISSN 0021-9002. doi: 10.2307/3214948. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=1274717>. Studies in applied probability.
- DLMF. Nist digital library of mathematical functions. URL <http://dlmf.nist.gov/>.
- C. Döbler and M. Stolz. Stein’s method and the multivariate CLT for traces of powers on the classical compact groups. *Electron. J. Probab.*, 16:no. 86, 2375–2405, 2011. ISSN 1083-6489. doi: 10.1214/EJP.v16-960. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=2861678>.
- J. S. Geronimo and K. M. Case. Scattering theory and polynomials orthogonal on the unit circle. *J. Math. Phys.*, 20(2):299–310, 1979. ISSN 0022-2488. doi: 10.1063/1.524077. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=519213>.
- I. M. Gessel. Symmetric functions and P-recursiveness. *J. Combin. Theory Ser. A*, 53(2):257–285, 1990. ISSN 0097-3165. doi: 10.1016/0097-3165(90)90060-A. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=1041448>.
- B. L. Golinskii and I. A. Ibragimov. A limit theorem of G. Szegő. *Izv. Akad. Nauk SSSR Ser. Mat.*, 35:408–427, 1971. ISSN 0373-2436. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=0291713>.
- I. A. Ibragimov. A theorem of Gabor Szegő. *Mat. Zametki*, 3:693–702, 1968. ISSN 0025-567X. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=231114>.
- T. Jiang and S. Matsumoto. Moments of traces of circular beta-ensembles. *Ann. Probab.*, 43(6):3279–3336, 2015. ISSN 0091-1798. doi: 10.1214/14-AOP960. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=3433582>.
- K. Johansson. On Szegő’s asymptotic formula for Toeplitz determinants and generalizations. *Bull. Sci. Math. (2)*, 112(3):257–304, 1988. ISSN 0007-4497. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=975365>.
- K. Johansson. On random matrices from the compact classical groups. *Ann. of Math. (2)*, 145(3):519–545, 1997. ISSN 0003-486X. doi: 10.2307/2951843. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=1454702>.
- G. Lambert. Mesoscopic central limit theorem for the circular β -ensembles and applications. [arXiv:1902.06611](https://arxiv.org/abs/1902.06611), 2019.
- G. Lambert, M. Ledoux, and C. Webb. Quantitative normal approximation of linear statistics of β -ensembles. *Ann. Probab.*, 47(5):2619–2685, 2019. ISSN 0091-1798. doi: 10.1214/18-AOP1314. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=4021234>.
- P. Sarnak. Memorial conference for I. Piatetski-Shapiro: An underdetermined matrix moment problem and its applications to computing zeros of l-functions, 2019. URL <http://publications.ias.edu/sarnak/paper/2705>.
- B. Simon. *Orthogonal polynomials on the unit circle. Part 1*, volume 54 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2005a. ISBN 0-8218-3446-0. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=2105088>. Classical theory.
- B. Simon. *Trace ideals and their applications*, volume 120 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2005b. ISBN 0-8218-3581-5. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=2154153>.
- G. Szegő. On certain Hermitian forms associated with the Fourier series of a positive function. *Comm. Sém. Math. Univ. Lund [Medd. Lunds Univ. Mat. Sem.]*, 1952(Tome Supplémentaire):228–238, 1952. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=0051961>.
- C. Webb. Linear statistics of the circular β -ensemble, Stein’s method, and circular Dyson Brownian motion. *Electron. J. Probab.*, 21:Paper No. 25, 16, 2016. ISSN 1083-6489. doi: 10.1214/16-EJP4535. URL <https://mathscinet.ams.org/mathscinet-getitem?mr=3485367>.