

Adjoining roots in homotopy theory

Tyler Lawson

February 7, 2020

Abstract

We use a “twisted group algebra” method to constructively adjoin formal radicals $\sqrt[n]{\alpha}$, for α a unit in a commutative ring spectrum or an invertible object in a symmetric monoidal ∞ -category. We show that this construction is classified by maps from Eilenberg–Mac Lane objects to the unit spectrum \mathfrak{gl}_1 , the Picard spectrum pic , and the Brauer spectrum br .

1 Introduction

Given a commutative ring R and an element $\alpha \in R$, we can adjoin a formal radical $\sqrt[n]{\alpha}$ to R by embedding R into the extension ring $R[x]/(x^n - \alpha)$. These ring extensions come equipped with a ready-made basis $\{1, x, \dots, x^{n-1}\}$ over R and are fundamental constructions in algebra. In the derived setting, however, it is less clear when these types of constructions are possible. Given a commutative ring spectrum R and an element $\alpha \in \pi_0 R$, one can construct a commutative R -algebra with an n 'th root of α using the same type of presentation in terms of generators and relations. However, away from characteristic zero the universal property enjoyed by this construction is not as strong and its coefficient ring can be unpredictable.

We can try instead to lift the algebra directly by constructing a commutative R -algebra S with a map $R \rightarrow S$ such that, on coefficient rings, we have the algebraic extension: $\pi_* S \cong (\pi_* R)[x]/(x^n - \alpha)$. Depending on α , such an extension may not be possible or may not be unique. It is always possible to adjoin roots of 1, because those algebras are realized by the group algebras $R[C_n]$ of finite cyclic groups. If both α and n are units in $\pi_0 R$ then the resulting extension on coefficient rings is *étale*, and the obstruction theory of Robinson [Rob03] or Goerss–Hopkins [GH04] can be used to show that the extension ring S exists, is essentially unique, and has a universal property among R -algebras with such a root adjoined. If 2 is invertible, this means that we can adjoin $\sqrt{-1}$. However, Schwänzl–Vogt–Waldhausen showed in [SVW99], using topological Hochschild homology, that it is impossible to adjoin a square root of -1 to the sphere spectrum. A different argument of Hopkins with $K(1)$ -local power operations shows that the p -complete K -theory spectrum cannot admit p 'th roots of unity [LN14, A.6.iii], and this was further generalized by Devalapurkar to $K(n)$ -local theory [Dev17]. Further difficulties appear when attempting to adjoin a root that appears in a nonzero degree.

Our starting observation is that these formal radicals are a special case of twisted group algebras. Given an abelian group extension $0 \rightarrow G \rightarrow E \rightarrow A \rightarrow 0$ and a group homomorphism $G \rightarrow R^\times$, we can form the relative tensor product $\mathbb{Z}[E] \otimes_{\mathbb{Z}[G]} R$. The elements of A lift to a basis over R , and the resulting algebra differs from the group algebra $R[A]$ by this central extension. Moreover, all such extensions can be constructed in a universal case by pushing forward the extension from G to R^\times .

This process can be applied to ring spectra. We will begin by showing that, when it is possible to construct similar extensions of the spectrum of units $\mathrm{gl}_1(R)$, we can give systematic constructions of formal radicals and other twisted group algebras, and obstructions are detectable with sufficient knowledge of $\mathrm{gl}_1(R)$. By then reinterpreting these constructions as Thom spectra for maps to $\mathrm{pic}(R)$, we can use recent work of Antolín-Camarena–Barthel [AB14] to both generalize this and allow us to identify these algebras as having a universal property. This recovers several constructions: adjoining n 'th roots of elements to a ring spectrum where n is invertible, usually carried out using obstruction theory, and adjoining similar roots to elements in gradings outside zero. There are also new constructions: we find that we can extend the first Postnikov stage $\tau_{\leq 1} \mathbb{S}_{(2)}$ of the 2-local sphere by adjoining \sqrt{D} for $D \equiv 1 \pmod{4}$.

Once we have accomplished this, our second goal will be to dig one categorical level down.

The same methods can be applied to adjoin formal radicals of elements in the Picard group. This allows us to take an extension of the Picard spectrum $\mathrm{pic}(R)$ and use it to embed the category of R -modules into a graded category with a larger group of invertible objects. This formalism recovers algebraic examples, such as Rezk's ω -twisted tensor product for $\mathbb{Z}/2$ -graded modules [Rez09]. There are also new topological examples: if R is an MU -algebra we can embed the category LMod_R of R -modules, where integer suspensions are possible, into a larger category $\prod_{\mathbb{Q}/\mathbb{Z}} \mathrm{LMod}_R$ with a symmetric monoidal structure that allows suspensions by elements of $\mathbb{Q} \times \mathbb{Z}/2$. This is also possible for modules over the topological K -theory spectrum ku and the algebraic K -theory spectrum $K\mathbb{C}$. Although our focus is on ring spectra, many of the results are proved in the generality of presentable symmetric monoidal ∞ -categories.

We finally close with the observation that algebraic K -theory relates these two processes.

Further directions

A first issue is that our discussion of adjoining roots is less satisfying for units outside degree zero. In particular, the identification of such units is somewhat roundabout. Ideally a solution to this problem would make use of a spectrum of graded units similar to those developed by Sagave [Sag16], and in particular his construction of $\mathrm{bgl}_1^* R$.

Second, we restrict our attention to strictly commutative objects (meaning E_∞ -ring objects). The constructions in this paper should have interesting and useful E_n -variants, making use of the iterated classifying spaces $B^{(n)}GL_1 R$.

Finally, we study unit groups because they are more easily analyzed via their associated spectrum. This means that we lose any ability to extract formal radicals of nonunit elements. We are hopeful that the future will bring a better understanding of the structure theory of E_∞ -spaces, allowing us to move beyond unit groups to effectively study multiplicative monoids.

Conventions and background

Our paper is written homotopically, and in particular we use the phrase “commutative ring spectrum” to mean an E_∞ -ring spectrum.

For expedience, we will use the same name for both an abelian group A and the associated Eilenberg–Mac Lane spectrum, regarding the category of abelian groups as embedded fully faithfully into the higher category of spectra. A commutative ring k gives rise to a commutative ring spectrum.

For a ring spectrum R , the unit group $\mathrm{GL}_1(R) \subset \Omega^\infty R$ is the space of units under the multiplicative monoidal product, or equivalently the space of self-equivalences of R as a left R -module. If R has a commutative ring structure we write $\mathfrak{gl}_1(R)$ for the associated spectrum of units [May77, ABG⁺14]. There is a unit map $\mathbb{S}[\mathrm{GL}_1(R)] \rightarrow R$ for the adjunction between unit groups and spherical group algebras.

For a monoidal ∞ -category \mathcal{C} , the Picard space $\mathrm{Pic}(\mathcal{C}) \subset \mathcal{C}^\simeq$ is the space of invertible objects and equivalences between them [Cla11, MS16].¹ If \mathcal{C} has a symmetric monoidal structure then $\mathrm{Pic}(\mathcal{C})$ has an E_∞ -structure and we write $\mathrm{pic}(\mathcal{C})$ for the associated Picard spectrum. If $\mathcal{C} = \mathrm{LMod}_R$ is the category of modules over R , we simply write $\mathrm{Pic}(R)$ and $\mathrm{pic}(R)$ instead.

We will require known identifications of the groups $[A, \Sigma^k B]$ of homotopy classes of maps between Eilenberg–Mac Lane spectra, which we will simply state [EML54].²

1. The group $[A, B]$ is isomorphic to the group $\mathrm{Hom}(A, B)$.
2. The group $[A, \Sigma B]$ is isomorphic to the group $\mathrm{Ext}(A, B)$: this extension is identified with the fiber of a map $A \rightarrow \Sigma B$.
3. The group $[A, \Sigma^2 B]$ is isomorphic to the group $\mathrm{Hom}(A, B[2])$ of 2-torsion homomorphisms $A \rightarrow B$.
4. The group $[A, \Sigma^3 B]$ is part of a short exact sequence

$$0 \rightarrow \mathrm{Ext}(A, B[2]) \rightarrow [A, \Sigma^3 B] \rightarrow \mathrm{Hom}(A, B/2) \rightarrow 0.$$

5. These identifications respect composition. In particular, the composition map $[B, \Sigma^2 C] \times [A, \Sigma B] \rightarrow [A, \Sigma^3 C]$ is the Yoneda pairing

$$\mathrm{Hom}(B, C[2]) \times \mathrm{Ext}(A, B) \rightarrow \mathrm{Ext}(A, C[2]).$$

¹(cf. [Mat14, §2.2]) If the unit of \mathcal{C} is κ -compact for some cardinal κ , then the objects of $\mathrm{Pic}(\mathcal{C})$ are also κ -compact, and if \mathcal{C} is presentable then the full subcategory of κ -compact objects is essentially small. Therefore, for presentable monoidal ∞ -categories it is possible to identify $\mathrm{Pic}(\mathcal{C})$ with a small space even though \mathcal{C} is large.

²All of these can be determined by first calculating that the groups of maps $\mathbb{Z} \rightarrow \Sigma^k \mathbb{Z}$ are $\mathbb{Z}, 0, 0, \mathbb{Z}/2, 0$ for $k = 0 \dots 5$, and then using free resolutions of the source and target. The generator in degree 3 is the composite of mod-2 reduction, the Steenrod square Sq^2 , and the Bockstein.

As a result, a pair (g, Γ) representing a homomorphism $B \rightarrow C[2]$ and an extension $0 \rightarrow B \rightarrow \Gamma \rightarrow A \rightarrow 0$ maps to zero if and only if the homomorphism g extends from B to all of Γ .

Given an ordinary symmetric monoidal category \mathcal{D} , the Picard space $\text{Pic}(\mathcal{D})$ is the nerve of an ordinary groupoid, consisting of the invertible objects and isomorphisms between them. This means there is a fiber sequence

$$\text{pic}(\mathcal{D}) \rightarrow \pi_0 \text{pic}(\mathcal{D}) \xrightarrow{k} \Sigma^2 \pi_1 \text{pic}(\mathcal{D}).$$

Here $\pi_0 \text{pic}(\mathcal{D})$ is the classical Picard group of the category \mathcal{D} , and $\pi_1 \text{pic}(\mathcal{D})$ is the automorphism group $\text{Aut}_{\mathcal{D}}(\mathbb{1})$ of the monoidal unit. This first k -invariant is always expressed in terms of the twist map. Namely, given an invertible module γ , the twist-self-isomorphism $\gamma \otimes \gamma \rightarrow \gamma \otimes \gamma$ is multiplication by a 2-torsion automorphism $\tau(\gamma)$ of the monoidal unit, an element of $\pi_1 \text{pic}(\mathcal{D})$ that satisfies $\tau(\gamma) \circ \tau(\gamma) = \text{id}$. The k -invariant $\pi_0 \text{pic}(\mathcal{D}) \xrightarrow{k} \Sigma^2 \pi_1 \text{pic}(\mathcal{D})$ is identified with τ .

Acknowledgements

The author would like to thank Tobias Barthel, Clark Barwick, Sanath Devalapurkar, Fabien Hebestreit, Lars Hesselholt, and Charles Rezk for discussions related to this paper, and Benjamin Antieau for comments on an earlier draft. The author would also like to thank the Max Planck Institute for their hospitality and financial support while this paper was being written. The author was partially supported by NSF grant 1610408.

2 Formal radicals

Let $\alpha \in \pi_0(R)$ be a unit; α is then also represented by an element of $\pi_0(\mathfrak{gl}_1(R))$. Fix a positive integer n . From this, we can construct an extension of abelian groups

$$0 \rightarrow \pi_0(\mathfrak{gl}_1 R) \rightarrow E \rightarrow \mathbb{Z}/n \rightarrow 0$$

such that the generator 1 of \mathbb{Z}/n has a chosen lift to an element $x \in E$ with $x^n = \alpha$. As a set with an action of $\pi_0 \mathfrak{gl}_1(R)$, $E \cong \coprod_{i=0}^{n-1} \pi_0 \mathfrak{gl}_1(R) \cdot \{x^i\}$.

This extension is determined by an extension class $\text{Ext}^1(\mathbb{Z}/n, \pi_0 \mathfrak{gl}_1(R))$, or equivalently by a map

$$\bar{\rho}: \mathbb{Z}/n \rightarrow \Sigma \pi_0 \mathfrak{gl}_1(R)$$

between Eilenberg–Mac Lane spectra.

Definition 1. A *formal n 'th root of α* is a lift ρ of $\bar{\rho}$ to $\mathfrak{gl}_1(R)$:

$$\begin{array}{ccc} & \Sigma \mathfrak{gl}_1(R) & \\ & \nearrow \rho & \downarrow \\ \mathbb{Z}/n & \xrightarrow{\bar{\rho}} & \Sigma \pi_0 \mathfrak{gl}_1(R) \end{array}$$

We refer to the fiber of ρ as the *extended unit spectrum* $\mathfrak{gl}_1(R, \rho)$ associated to ρ , and the associated infinite loop space as *extended unit group* $GL_1(R, \rho)$.

The extended unit spectrum is part of a fiber sequence

$$\mathfrak{gl}_1(R) \rightarrow \mathfrak{gl}_1(R, \rho) \rightarrow \mathbb{Z}/n,$$

and hence we get a decomposition $GL_1(R, \rho) \cong \coprod_{i=0}^{n-1} GL_1(R) \cdot \{x^i\}$ as spaces with an action of $GL_1(R)$.

Remark 2. The map ρ determines $\bar{\rho}$: the map $\mathfrak{gl}_1(R) \rightarrow \mathfrak{gl}_1(R, \rho)$ is an isomorphism on π_k except when $k = 0$, when it is the inclusion $\pi_0 \mathfrak{gl}_1(R) \rightarrow E$. Therefore, ρ determines the extension E and hence determines α up to n 'th powers.

Definition 3. Suppose that ρ is a formal n 'th root of α . Then the algebra obtained by *adjoining this root* is the relative smash product

$$R[\rho] = \mathbb{S}[GL_1(R, \rho)] \otimes_{\mathbb{S}[GL_1(R)]} R.$$

Proposition 4. *The coefficient ring of $R[\rho]$ is*

$$\pi_* R[\rho] \cong \pi_* R[x]/(x^n - \alpha).$$

Proof. The decomposition $GL_1(R, \rho) \cong \coprod_{i=0}^{n-1} GL_1(R) \cdot \{x^i\}$ means that the spherical group algebra $\mathbb{S}[GL_1(R, \rho)]$ decomposes as $\bigoplus_{i=0}^{n-1} \mathbb{S}[GL_1(R)] \cdot \{x^i\}$, a free left $\mathbb{S}[GL_1(R)]$ -module. Therefore, there is a simple Künneth formula that gives us an isomorphism of modules:

$$\begin{aligned} \pi_* R[\rho] &\cong \pi_* \mathbb{S}[GL_1(R, \rho)] \otimes_{\pi_* \mathbb{S}[GL_1(R)]} \pi_* R \\ &\cong \bigoplus_{i=0}^{n-1} \pi_* R \cdot \{x^i\}. \end{aligned}$$

Moreover, the identity $x^n = \alpha$ for the element $x \in E = \pi_0 GL_1(R, \rho)$ completely determines the multiplication in $\pi_* R[\rho]$. \square

Remark 5. It is clear that, other than the calculation of the structure of the coefficient ring, there is nothing special about the group \mathbb{Z}/n in the above discussion. Given a map $\rho: A \rightarrow \Sigma \mathfrak{gl}_1(R)$ for some abelian group A , lifting an extension

$$0 \rightarrow \pi_0 \mathfrak{gl}_1(R) \rightarrow E \rightarrow A \rightarrow 0,$$

there is an associated algebra $R[\rho]$ whose coefficient ring is a twisted central extension:

$$\begin{aligned} \pi_* R[\rho] &\cong \mathbb{Z}[E] \otimes_{\mathbb{Z}[\pi_0 \mathfrak{gl}_1(R)]} \pi_* R \\ &\cong \bigoplus_{a \in A} \pi_* R \cdot \{a\}. \end{aligned}$$

We will see similar algebras in later sections.

Example 6. Suppose that n is a unit in $\pi_0 R$. Then n acts invertibly on the homotopy groups $\pi_k \mathfrak{gl}_1(R) \cong \pi_k R$ for $k > 0$: therefore the spectrum of maps from \mathbb{Z}/n to the 0-connected cover $\tau_{\geq 1} \mathfrak{gl}_1(R)$ is trivial. Using the fiber sequence

$$\Sigma \mathfrak{gl}_1(R) \rightarrow \Sigma \pi_0 \mathfrak{gl}_1(R) \rightarrow \Sigma^2 \tau_{\geq 1} \mathfrak{gl}_1(R),$$

we find that, for any unit α , the map $\mathbb{Z}/n \rightarrow \Sigma \pi_0 \mathfrak{gl}_1(R)$ lifts essentially uniquely to a formal n 'th root $\mathbb{Z}/n \rightarrow \Sigma \mathfrak{gl}_1(R)$.

Example 7. Let K be the p -complete K -theory spectrum. Then it is possible to show that the group $[\mathbb{Z}/p, \Sigma \mathfrak{gl}_1 K]$ is trivial, and thus this method does not allow us to adjoin $x = \sqrt[p]{\alpha}$ for any nontrivial element α of $\mathbb{Z}_p^\times / (\mathbb{Z}_p^\times)^p$. To give a proof, however, we need to know structural properties of the multiplication on K . The straight-line proof we know uses Rezk's $K(1)$ -local logarithm $\ell_1: \mathfrak{gl}_1 K \rightarrow K$ [Rez06], together with nontrivial knowledge of low-degree k -invariants for $\mathfrak{gl}_1 K$. In rough, the fact that Rezk's logarithm gives us an equivalence in degrees greater than 2 implies that there is a diagram of fiber sequences:

$$\begin{array}{ccccc} \Sigma^4 (ku)_p^\wedge & \longrightarrow & \tau_{\geq 2} \mathfrak{gl}_1 K & \longrightarrow & \mathfrak{gl}_1 K \\ & & \downarrow & \swarrow & \downarrow \\ & & \Sigma^2 \mathbb{Z}_p & & \mathbb{Z}_p^\times \end{array}$$

Applying $[\mathbb{Z}/p, -]$ gives us a spectral sequence that computes $[\mathbb{Z}/p, \Sigma \mathfrak{gl}_1 K]$; in the critical group the k -invariants of $\mathfrak{gl}_1 K$ give this spectral sequence one nontrivial differential for $p > 2$ and two nontrivial differentials for $p = 2$.

However, the impossibility of adjoining these radicals can be shown directly using $K(1)$ -local power operations, in a manner exactly analogous to the proof that one cannot adjoin roots of unity; in this form it generalizes. Let us sketch this argument.

If we had such an E_∞ -ring K -algebra L , it would be p -complete and thus $K(1)$ -local. Its coefficient ring $K_*[x]/(x^p - \alpha)$ would then have a $K(1)$ -local power operation ψ^p , a ring endomorphism that agrees with the Frobenius mod p [Hop14]. The element $\zeta = \psi^p(x)/x$ would then be a p 'th root of unity. If ζ is not in \mathbb{Z}_p^\times , then L is a $K(1)$ -local E_∞ -ring containing a nontrivial p 'th root of unity and Devalapurkar has shown this to be impossible [Dev17]. If ζ is in \mathbb{Z}_p^\times , then $\alpha \equiv \psi^p x = \zeta^{-1} x \pmod{p}$, which contradicts the fact that $1, x, \dots, x^{p-1}$ are a basis of this ring mod p .

3 Strict units

One source of formal roots is the theory of strictly commutative elements.

Definition 8. The space $\mathbb{G}_m(R)$ of *strictly commutative units* of R is the space of maps $\mathbb{Z} \rightarrow \mathfrak{gl}_1(R)$, or equivalently the space of E_∞ -maps $\mathbb{Z} \rightarrow \mathrm{GL}_1(R)$. The generator $1 \in \mathbb{Z}$ induces forgetful maps $\mathbb{G}_m(R) \rightarrow \mathrm{GL}_1(R) \rightarrow \pi_0 \mathfrak{gl}_1(R)$.

In particular, a strictly commutative unit of R has an underlying unit in $\pi_0(R)$.

Proposition 9. *Suppose α is a strictly commutative unit of R . Then, for any $n > 0$, α has a canonical lift to a formal n 'th root.*

Proof. The canonical Bockstein map $\mathbb{Z}/n \rightarrow \Sigma\mathbb{Z}$ can be composed with the map $\mathbb{Z} \rightarrow \mathfrak{gl}_1(R)$ classifying α . \square

Example 10. Let $R = S^{-1}(\tau_{\leq 1}\mathbb{S})$ be the localization of the first Postnikov truncation of the sphere spectrum with respect to some set S of primes (not containing 2). Then there is a fiber sequence

$$\mathfrak{gl}_1(R) \rightarrow (S^{-1}\mathbb{Z})^\times \xrightarrow{k} \Sigma^2\mathbb{Z}/2.$$

This k -invariant corresponds to a (2-torsion) homomorphism $(S^{-1}\mathbb{Z})^\times \rightarrow \mathbb{Z}/2$. This homomorphism is a classical calculation of orientation theory: it is the map

$$n \mapsto \begin{cases} 1 & \text{if } n \equiv +1 \pmod{4}, \\ -1 & \text{if } n \equiv -1 \pmod{4}. \end{cases}$$

As a result, one can determine the homotopy groups of the space of strictly commuting elements, and in particular there is an exact sequence

$$0 \rightarrow [\mathbb{Z}, \mathfrak{gl}_1(R)] \rightarrow [\mathbb{Z}, (S^{-1}\mathbb{Z})^\times] \rightarrow [\mathbb{Z}, \Sigma^2\mathbb{Z}/2].$$

We find that any unit in $S^{-1}\mathbb{Z}$ which is congruent to 1 mod 4 lifts, essentially uniquely, to a strictly commutative unit of R . This allows us to construct commutative algebras such as $R[\sqrt{5}]$ and $R[\sqrt{-3}]$, even though these are ramified extensions on the level of coefficient rings.

Remark 11. In the case of strictly commutative units, we obtain a second description of the algebra obtained by adjoining this root ρ . A strictly commutative element determines a composite map

$$\mathbb{S}[\mathbb{Z}] \rightarrow \mathbb{S}[\mathrm{GL}_1(R)] \rightarrow R,$$

and so we can construct the algebra $R[\rho]$ as $R \otimes_{\mathbb{S}[\mathbb{Z}]} \mathbb{S}[\frac{1}{n}\mathbb{Z}]$.

This has the benefit that it readily lifts to a *nonunit* version. If we define the *strictly commutative multiplicative monoid* $\mathbb{M}_m(R)$ to be the space of E_∞ -maps

$$\mathbb{N} \rightarrow M_1(R) = \Omega_\otimes^\infty R$$

to the multiplicative monoid of R , then a strictly commutative element α can have an n 'th root adjoined via the construction

$$\mathbb{S}[\frac{1}{n}\mathbb{N}] \otimes_{\mathbb{S}[\mathbb{N}]} R.$$

Remark 12. A further property possessed by strict units is that it is possible to trivialize them. Using the natural augmentation $\mathbb{S}[\mathbb{Z}] \rightarrow \mathbb{S}$ of the spherical group

algebra, any strictly commutative unit of R with underlying unit $x \in \pi_0(R)$ gives rise to a new commutative ring spectrum

$$R/(x-1) = \mathbb{S} \otimes_{\mathbb{S}[\mathbb{Z}]} R,$$

whose underlying R -module is equivalent to the cofiber of the map $R \rightarrow R$ induced by $(x-1) \in \pi_0(R)$. This algebra has the following property: it is the universal commutative R -algebra with a chosen commuting diagram

$$\begin{array}{ccc} \mathbb{Z} & \longrightarrow & \mathfrak{gl}_1(R) \\ \downarrow & & \downarrow \\ * & \longrightarrow & \mathfrak{gl}_1(R/(x-1)). \end{array}$$

We will see similarly themed universal properties in later sections.

4 Strict gradings

The shift-by-1 in our definition of formal roots is strongly suggestive: the suspended unit spectrum $\Sigma \mathfrak{gl}_1(R)$ is a connective cover of the Picard spectrum $\mathrm{pic}(R)$. In this section we will begin exploring Picard-graded analogues of our constructions.

Definition 13. Suppose that A is an abelian group and \mathcal{C} is a symmetric monoidal ∞ -category. The space of *strict A -gradings for \mathcal{C}* is the space of maps $A \rightarrow \mathrm{pic}(\mathcal{C})$, or equivalently the space of E_∞ -maps $A \rightarrow \mathrm{Pic}(\mathcal{C})$. There is a composite $A \rightarrow \pi_0 \mathrm{pic}(\mathcal{C})$, which we refer to as the *underlying A -grading*.

Remark 14. Suppose that we have any symmetric monoidal functor $A \rightarrow \mathcal{C}$. The space A could be regarded as a discrete groupoid, so its image lies in \mathcal{C}^{\times} ; the objects in A have inverses under the monoidal product, so monoidality of ρ implies that its image lies inside $\mathrm{Pic}(\mathcal{C})$. We will not distinguish between symmetric monoidal functors $A \rightarrow \mathcal{C}$ and symmetric monoidal functors $A \rightarrow \mathrm{Pic}(\mathcal{C})$.

Example 15. Let \mathcal{C} be a symmetric monoidal ∞ -category. The *strict Picard space* of \mathcal{C} , denoted by $\mathbb{P}\mathrm{ic}(\mathcal{C})$, is the space of strict \mathbb{Z} -gradings: maps $\mathbb{Z} \rightarrow \mathrm{pic}(\mathcal{C})$, or equivalently E_∞ -maps $\mathbb{Z} \rightarrow \mathrm{Pic}(\mathcal{C})$. The generator $1 \in \mathbb{Z}$ induces forgetful maps $\mathbb{P}\mathrm{ic}(\mathcal{C}) \rightarrow \mathrm{Pic}(\mathcal{C}) \rightarrow \pi_0 \mathrm{Pic}(\mathcal{C})$, and we refer to the image as the *underlying object*.

Example 16. The space of *strict n -torsion objects* of \mathcal{C} , denoted by $\mathbb{P}\mathrm{ic}^{[n]}(\mathcal{C})$, is the space of strict \mathbb{Z}/n -gradings: maps $\rho: \mathbb{Z}/n \rightarrow \mathrm{pic}(\mathcal{C})$, or equivalently maps $\mathbb{Z}/n \rightarrow \mathrm{Pic}(\mathcal{C})$ of E_∞ -spaces. The cofiber sequence $\mathbb{Z} \xrightarrow{n} \mathbb{Z} \rightarrow \mathbb{Z}/n$ of spectra gives rise to the following maps, where each double composite is a fiber sequence:

$$\mu_n(R) \rightarrow \mathbb{G}_m(R) \xrightarrow{n} \mathbb{G}_m(R) \xrightarrow{\partial} \mathbb{P}\mathrm{ic}^{[n]}(R) \rightarrow \mathbb{P}\mathrm{ic}(R) \xrightarrow{n} \mathbb{P}\mathrm{ic}(R)$$

In particular, the map ∂ sends a strictly commutative element $\alpha: \mathbb{Z} \rightarrow \mathfrak{gl}_1(R)$ to the image

$$\mathbb{Z}/n \rightarrow \Sigma \mathbb{Z} \xrightarrow{\alpha} \Sigma \mathfrak{gl}_1(R) \rightarrow \mathrm{pic}(R)$$

of the formal n 'th root associated to α .

Remark 17. For a commutative ring spectrum R , a strict n -torsion R -module with underlying left R -module L has a choice of equivalence $L^{\otimes_R n} \rightarrow R$. If the module L is equivalent to R , then the map $\mathbb{Z}/n \rightarrow \pi_0 \text{pic}(R)$ is trivial and so the map lifts to a map $\mathbb{Z}/n \rightarrow \Sigma \mathfrak{gl}_1(R)$: a formal n 'th root. We can detect which root (up to n 'th powers) by making a choice of an equivalence $R \rightarrow L$; this determines a composite equivalence $R \simeq R^{\otimes_R n} \rightarrow I^{\otimes_R n} \rightarrow R$, and hence a unit in $\pi_0(R)$.

Example 18. Suppose that \mathcal{C} is a symmetric monoidal stable ∞ -category such that n is a unit in the ring $\pi_0 \text{End}_{\mathcal{C}}(\mathbb{1})$. Then for $k > 0$ there are isomorphisms

$$\pi_k \text{pic}(\mathcal{C}) \cong \pi_{k+1} \text{Aut}_{\mathcal{C}}(\mathbb{1}) \cong \pi_{k+1} \text{End}_{\mathcal{C}}(\mathbb{1}),$$

and the latter are acted on invertibly by n . Therefore, the fiber sequence

$$\tau_{\geq 2} \text{pic}(\mathcal{C}) \rightarrow \text{pic}(\mathcal{C}) \rightarrow \text{pic}(h\mathcal{C})$$

induces an equivalence $\text{Map}(\mathbb{Z}/n, \text{pic}(\mathcal{C})) \rightarrow \text{Map}(\mathbb{Z}/n, \text{pic}(h\mathcal{C}))$. In this case, strict n -torsion objects in \mathcal{C} are equivalent to strict n -torsion objects in the homotopy category $h\mathcal{C}$.

Example 19. For a local ring k , there is a fiber sequence

$$\text{pic}(k) \rightarrow \mathbb{Z} \rightarrow \Sigma^2 k^\times,$$

where Σk is a generator of $\pi_0 \text{pic}(k)$. The k -invariant is the twist permutation of Σk , and is represented by the homomorphism $\mathbb{Z} \twoheadrightarrow \{\pm 1\} \rightarrow k^\times$. This k -invariant becomes trivial on $2\mathbb{Z}$, and so the category of k -modules has a strict $2\mathbb{Z}$ -grading. The spaces $\mathbb{G}_m(k)$ are connected but not contractible, so the $2\mathbb{Z}$ -gradings are unique up to isomorphism but not canonical. We can make them canonical by choosing a $2\mathbb{Z}$ -grading of \mathbb{Z} .³

Example 20 ([Law18, 1.3.7]). For any commutative ring spectrum R , the element ΣR in $\text{Pic}(R)$ determines a map $\mathbb{Z} = \pi_0 \text{pic}(\mathbb{S}) \rightarrow \pi_0 \text{pic}(R)$. For any $d > 0$, we get a composite $d\mathbb{Z} \rightarrow \pi_0 \text{pic}(R)$. If this lifts to a strict $d\mathbb{Z}$ -grading, we could call this an E_∞^d -structure on R : it should be a strengthening of the notion of an H_∞^d -structure from [BMMS86]. However, our enthusiasm for extending this notational convention is very low.

The universal property of the real bordism spectrum MO is that it is initial among commutative ring spectra with a nullhomotopy of the map $BO \rightarrow \text{Pic}(\mathbb{S}) \rightarrow \text{Pic}(MO)$ of E_∞ -spaces. Equivalently, it is initial among commutative rings with a commutative diagram of spectra

$$\begin{array}{ccc} ko & \longrightarrow & \text{pic}(\mathbb{S}) \\ \downarrow & & \downarrow \\ \mathbb{Z} & \longrightarrow & \text{pic}(MO). \end{array}$$

³If $2 = 0$ in k , this k -invariant is trivial and so the category of k -modules has a strict \mathbb{Z} -grading. Moreover, $\text{pic}(\mathbb{F}_2) \simeq \mathbb{Z}$, and so this \mathbb{Z} -grading is canonical.

In particular, this gives the real bordism spectrum MO a strict \mathbb{Z} -grading and any commutative MO -algebra inherits it. Similar considerations with complex or spin structures give the complex bordism spectrum MU a strict $2\mathbb{Z}$ -grading and the spin bordism spectrum $MSpin$ a strict $4\mathbb{Z}$ -grading.

Example 21. The Atiyah–Bott–Shapiro orientation lifts to give the complex K -theory spectrum ku a commutative MU -algebra structure, and the real K -theory spectrum ko a commutative $MSpin$ -algebra structure, due to work of Joachim [Joa04]. Therefore, ku admits a strict $2\mathbb{Z}$ -grading and ko admits a strict $4\mathbb{Z}$ -grading.

Example 22. Let $K\mathbb{C}$ be the algebraic K -theory spectrum of the complex numbers, which comes equipped with a map $f: K\mathbb{C} \rightarrow ku$ to the complex topological K -theory spectrum. Work of Suslin showed that the map f is an equivalence after profinite completion, and hence the fiber of f is rational. We would like to show that $K\mathbb{C}$ has a strict $2\mathbb{Z}$ -grading. (A similar argument applies to show that the strict $4\mathbb{Z}$ -grading of the real K -theory spectrum ko lifts to the algebraic K -theory $K\mathbb{R}$.)

Consider the arithmetic square:

$$\begin{array}{ccc} K\mathbb{C} & \longrightarrow & K\mathbb{C}_{\mathbb{Q}} \\ \downarrow & & \downarrow \\ K\mathbb{C}^{\wedge} & \longrightarrow & (K\mathbb{C}^{\wedge})_{\mathbb{Q}} \end{array}$$

The functor GL_1 preserves this pullback. When we apply pic we get a diagram of connective spectra:

$$\begin{array}{ccc} \text{pic}(K\mathbb{C}) & \longrightarrow & \text{pic}(K\mathbb{C}_{\mathbb{Q}}) \\ \downarrow & & \downarrow \\ \text{pic}(K\mathbb{C}^{\wedge}) & \longrightarrow & \text{pic}((K\mathbb{C}^{\wedge})_{\mathbb{Q}}) \end{array}$$

On π_0 this is the constant square \mathbb{Z} , and on π_1 we get a bicartesian square

$$\begin{array}{ccc} \mathbb{Z}^{\times} & \longrightarrow & \mathbb{Q}^{\times} \\ \downarrow & & \downarrow \\ \widehat{\mathbb{Z}}^{\times} & \longrightarrow & (\widehat{\mathbb{Z}}_{\mathbb{Q}})^{\times}. \end{array}$$

Together these show that the diagram of Picard spectra is a homotopy pullback diagram. Let C be the cofiber of $\text{pic}(K\mathbb{C}) \rightarrow \text{pic}(K\mathbb{C}^{\wedge})$; its homotopy groups are then rational above degree 2, and equal to the torsion-free group $\widehat{\mathbb{Z}}^{\times}/\mathbb{Z}^{\times}$ in degree one.

The obstruction to lifting the strict $2\mathbb{Z}$ -grading

$$2\mathbb{Z} \rightarrow \text{pic}(ku) \rightarrow \text{pic}(ku^{\wedge}) \simeq \text{pic}(K\mathbb{C}^{\wedge})$$

to a $2\mathbb{Z}$ -grading of $\text{pic}(K\mathbb{C})$ is the map $2\mathbb{Z} \rightarrow C$. However, consider the fiber sequence

$$\tau_{\geq 2}C \rightarrow C \rightarrow \Sigma\widehat{\mathbb{Z}}^{\times}/\mathbb{Z}^{\times}.$$

Since the left term is 1-connected and rational and the right term is torsion-free, there are no nontrivial homotopy classes of maps from $2\mathbb{Z}$ into either term, and hence $[\mathbb{Z}, C] = 0$. Therefore, the strict $2\mathbb{Z}$ -grading of ku can be extended to $K\mathbb{C}$.

5 Trivializing algebras

The ring spectrum constructed by adjoining formal radicals turns out to be a special case of a more general construction associated to strict gradings. From this point forward we will need to make heavier use of [Lur09, Lur17].

Definition 23. Suppose that \mathcal{C} is a presentable symmetric monoidal ∞ -category, A is an abelian group regarded as a discrete symmetric monoidal category, and that $\rho: A \rightarrow \mathcal{C}$ is a strict A -grading. We define the *trivializing algebra* T_ρ to be the homotopy colimit of ρ .

Remark 24. Since A is a discrete category, there is an equivalence in \mathcal{C} of the form

$$T_\rho \simeq \coprod_{a \in A} \rho(a).$$

This homotopy colimit is a very special case of the Thom construction. As such, work of Antolín-Camarena-Barthel gives the trivializing algebra a universal property, generalizing the results of [May77, ABG⁺] from the category of spectra.

Proposition 25. *Suppose that A is an abelian group and that $\rho: A \rightarrow \mathcal{C}$ is a symmetric monoidal functor with trivializing algebra T_ρ .*

1. T_ρ has a natural lift to a commutative algebra object: $T_\rho \in \text{CAlg}(\mathcal{C})$.
2. The algebra T_ρ is universal among commutative algebras in $\text{CAlg}(\mathcal{C})$ with a chosen commuting diagram

$$\begin{array}{ccc} A & \xrightarrow{\rho} & \text{pic}(\mathcal{C}) \\ \downarrow & & \downarrow \\ * & \longrightarrow & \text{pic}(\text{LMod}_{T_\rho}). \end{array}$$

In particular, for any $a \in A$ the algebra T_ρ has a chosen equivalence of left T_ρ -modules $\phi_a: T_\rho \rightarrow T_\rho \otimes \rho(a)$, and there are chosen coherences $\phi_a \otimes 1 \circ \phi_b \simeq \phi_{ab}$.

Proof. The colimit has a commutative algebra structure by [AB14, Theorem 2.8].

We will now prove the universal property essentially, following the same line of argument in [AB14, Lemma 3.15]. Applying [AB14, Theorem 2.13] to the functor $A \rightarrow \mathcal{C}$ of symmetric monoidal ∞ -categories, we find the following: maps $T_\rho \rightarrow R$ in $\text{CAlg}(\mathcal{C})$ are equivalent to lax symmetric monoidal lifts in the diagram

$$\begin{array}{ccc} & & \mathcal{C}/R \\ & \nearrow \text{dotted} & \downarrow \\ A & \longrightarrow \text{Pic}(\mathcal{C}) & \longrightarrow \mathcal{C}. \end{array}$$

The objects of $\mathcal{C}_{/R}$ are maps $N \rightarrow R$, with symmetric monoidal product given by

$$(N \rightarrow R) \otimes (M \rightarrow R) \simeq (N \otimes M \rightarrow R \otimes R \rightarrow R).$$

The monoidal unit is the map $\mathbb{I} \rightarrow R$. There is a symmetric monoidal functor $\mathcal{C}_{/R} \rightarrow \mathcal{C}$, given by forgetting the structure map, and a symmetric monoidal functor $\mathcal{C}_{/R} \rightarrow (\mathrm{LMod}_R)_{/R}$, given by $(L \rightarrow R) \mapsto (R \otimes L \rightarrow R)$.

However, since A is grouplike the image $L \rightarrow R$ of any object must be contained in the invertible objects of $\mathcal{C}_{/R}$. This implies first that L is an invertible object of \mathcal{C} . This implies second that $R \otimes L \rightarrow R$ is an invertible object of $(\mathrm{LMod}_R)_{/R}$; this happens only when this adjoint structure map is an equivalence. Conversely, if $L \rightarrow R$ is a map whose adjoint $R \otimes L \rightarrow R$ is an equivalence, tensoring with L^{-1} gives an equivalence of R -modules $R \rightarrow R \otimes L^{-1}$ of left R -modules, whose inverse is adjoint to a map $L^{-1} \rightarrow R$. \square

Remark 26. Suppose that the functor $\rho: A \rightarrow \mathcal{C}$ maps entirely to the unit component: it is a map $A \rightarrow B\mathrm{Aut}_{\mathcal{C}}(\mathbb{I})$ of E_{∞} -spaces, classifying an extension G of the automorphism space $\mathrm{Aut}_{\mathcal{C}}(\mathbb{I})$. Then the trivializing algebra T_{ρ} is universal among commutative algebras R in \mathcal{C} where the map $\mathrm{Aut}_{\mathcal{C}}(\mathbb{I}) \rightarrow \mathrm{Aut}_{\mathcal{C}}(R)$ extends to a map of E_{∞} -spaces $G \rightarrow \mathrm{Aut}_{\mathcal{C}}(R)$.

In particular, when \mathcal{C} is the category Mod_R of modules over a commutative ring spectrum, this universal property recovers the relative tensor product defined earlier.

Example 27. The trivializing algebra for the map $2\mathbb{Z} \rightarrow \mathrm{pic}(MU)$ is a periodic MU -spectrum MUP , whose coefficient ring is $\pi_*MU[u^{\pm 1}]$ for a generator u in degree 2. It is universal among commutative algebras R with a nullhomotopy of the composite $ku \rightarrow \mathrm{pic}(\mathbb{S}) \rightarrow \mathrm{pic}(R)$. The algebra MUP is often useful for translating between even-periodic and $\mathbb{Z}/2$ -graded interpretations in chromatic theory.

6 Root obstructions

The symmetric monoidal functor from \mathcal{C} to its homotopy category $h\mathcal{C}$ induces a map of Picard spectra $\mathrm{pic}(\mathcal{C}) \rightarrow \mathrm{pic}(h\mathcal{C})$, identifying $\mathrm{pic}(h\mathcal{C})$ with the first nontrivial stage $\tau_{\leq 1} \mathrm{pic}(\mathcal{C})$ in the Postnikov tower for $\mathrm{pic}(\mathcal{C})$. For us to lift a map $\bar{\rho}: A \rightarrow \pi_0 \mathrm{pic}(\mathcal{C})$ to the first Postnikov stage $\mathrm{pic}(h\mathcal{C})$, it is necessary and sufficient that the composite map

$$A \xrightarrow{\bar{\rho}} \pi_0 \mathrm{pic}(\mathcal{C}) \xrightarrow{k} \Sigma^2 \pi_1 \mathrm{pic}(\mathcal{C})$$

is trivial. The result is an obstruction class: an element of $[A, \Sigma^2 \pi_1 \mathrm{pic}(\mathcal{C})]$, and a lift exists if and only if this obstruction vanishes. Two different choices of lift to a map $A \rightarrow \mathrm{pic}(h\mathcal{C})$ are represented by homotopy classes of maps $A \rightarrow \Sigma \pi_1 \mathrm{pic}(\mathcal{C})$.

The identification of the first k -invariant with the twist homomorphism τ leads us to the following definition [Rez09].

Definition 28. An element $\gamma \in \pi_0 \mathrm{pic}(\mathcal{C})$ is *symmetric* if $\tau(\gamma) = \mathrm{id}$ in $\pi_1 \mathrm{pic}(\mathcal{C})$.

This allows us to conclude the following.

Proposition 29. *A map $\bar{\rho}: A \rightarrow \pi_0 \text{pic}(\mathcal{C})$ lifts to a map $A \rightarrow \text{pic}(h\mathcal{C})$ if and only if the elements $\bar{\rho}(a)$ are symmetric for all $a \in A$. Two such lifts determine a difference class in $\text{Ext}(A, \pi_1 \text{pic}(h\mathcal{C}))$.*

Example 30. Suppose that L is an invertible object such that there is an equivalence $v: \mathbb{I} \rightarrow L^{\otimes n}$. The object L determines a map $\mathbb{S} \rightarrow \text{pic}(\mathcal{C})$, and the equivalence v determines a nullhomotopy of n times this map. Alternatively, there is a commutative diagram of symmetric monoidal ∞ -categories

$$\begin{array}{ccc} F(L^{\otimes n}) & \longrightarrow & F(L) \\ \downarrow & & \downarrow \\ * & \longrightarrow & \text{Pic}(\mathcal{C}), \end{array}$$

where $F(x)$ is the free E_∞ -space on an object named x . Taking associated spectra gives a diagram

$$\begin{array}{ccc} \mathbb{S} & \xrightarrow{n} & \mathbb{S} \\ \downarrow & & \downarrow \\ * & \longrightarrow & \text{pic}(\mathcal{C}), \end{array}$$

or equivalently a map $\mathbb{S}/n \rightarrow \text{pic}(\mathcal{C})$. Conversely, there is a short exact sequence

$$0 \rightarrow \pi_1 \text{pic}(\mathcal{C}) / [\pi_1 \text{pic}(\mathcal{C})]^n \rightarrow [\mathbb{S}/n, \text{pic}(\mathcal{C})] \rightarrow \pi_0 \text{pic}(\mathcal{C})[n] \rightarrow 0.$$

The quotient expresses that a map from \mathbb{S}/n determines an underlying n -torsion object L in $\pi_0 \text{pic}(\mathcal{C})$. The kernel expresses that two different maps representing the same object L may differ in their choice of equivalence $v: \mathbb{I} \rightarrow L^{\otimes n}$ (modulo self-equivalences of L).

If there is a symmetric monoidal functor $f: \mathcal{C} \rightarrow \mathcal{D}$ such that there is an equivalence $u: \mathbb{I} \rightarrow f(L)$ in \mathcal{D} , then the map $\mathbb{S} \rightarrow \text{pic}(\mathcal{D})$ determining $f(L)$ becomes trivial. However, the extended map $\mathbb{S}/n \rightarrow \text{pic}(\mathcal{D})$ does not always become trivial: it becomes trivial precisely when there exists a choice of $u: \mathbb{I} \rightarrow f(L)$ in \mathcal{D} such that $u^n = f(v)$.

The bottom homotopy group of \mathbb{S}/n is \mathbb{Z}/n . The map $\mathbb{S}/n \rightarrow \text{pic}(h\mathcal{C})$ extends to a map $\mathbb{Z}/n \rightarrow \text{pic}(h\mathcal{C})$ if and only if L is symmetric, and if so it extends uniquely. A strict \mathbb{Z}/n -grading would be an extension to a map $\mathbb{Z}/n \rightarrow \text{pic}(\mathcal{C})$. In rough, we can think of this in the following way. If we have a strict \mathbb{Z}/n -grading, then it is a rigid version of choosing an object L with an equivalence $v: \mathbb{I} \rightarrow \mathbb{I}^{\otimes n}$; the trivializing algebra T then extracts an n 'th root of this *chosen* equivalence v .

Example 31. Suppose that R is a commutative ring spectrum which is $2n$ -periodic: there is a unit $v \in \pi_{2n}R$. This determines a symmetric element $\Sigma^2 R$, and a lift of $\Sigma^2 R$ to a strict n -torsion object allows us to construct an R -algebra whose \mathbb{Z} -graded coefficient ring is

$$\pi_* R[x] / (x^n - av)$$

for some well-defined $a \in (\pi_0 R)^\times / [(\pi_0 R)^\times]^n$. If n is a unit in $\pi_0 R$ then we also find that such algebras can be constructed, essentially uniquely, for any value of a . These extensions appear, for example, when relating completed Johnson–Wilson spectra to Lubin–Tate spectra [LN12, §4].

7 Grading extensions

In this section, we will begin the process of extending gradings by adjoining formal radicals to elements in the Picard group. We fix a symmetric monoidal presentable ∞ -category \mathcal{C} , and let $\text{Pic}(\mathcal{C})$ be the Picard space of invertible elements in \mathcal{C} .

Definition 32. Let

$$0 \rightarrow \pi_0 \text{pic}(\mathcal{C}) \rightarrow \Gamma \rightarrow B \rightarrow 0$$

be an extension of abelian groups, represented by a map

$$\bar{\rho}: B \rightarrow \Sigma \pi_0 \text{pic}(\mathcal{C})$$

between Eilenberg–Mac Lane spectra. A *extension to Γ -grading* is a lift of $\bar{\rho}$ to $\text{pic}(\mathcal{C})$:

$$\begin{array}{ccc} & & \Sigma \text{pic}(\mathcal{C}) \\ & \nearrow \rho & \downarrow \\ B & \xrightarrow{\bar{\rho}} & \Sigma \pi_0 \text{pic}(\mathcal{C}) \end{array}$$

We refer to the fiber of ρ as the *extended Picard spectrum* $\text{pic}(\mathcal{C}, \rho)$ associated to ρ , and the associated infinite loop space as the *extended Picard group* $\text{Pic}(\mathcal{C}, \rho)$.

The extended Picard spectrum is part of a fiber sequence

$$\text{pic}(\mathcal{C}) \rightarrow \text{pic}(\mathcal{C}, \rho) \rightarrow B,$$

and on π_0 this realizes the extension $0 \rightarrow \pi_0 \text{pic}(\mathcal{C}) \rightarrow \Gamma \rightarrow B \rightarrow 0$. We get a decomposition $\text{Pic}(\mathcal{C}, \rho) \cong \coprod_{b \in B} \text{Pic}(\mathcal{C}) \cdot \{b\}$ as spaces with an action of $\text{Pic}(\mathcal{C})$.

Remark 33. Again, the map ρ determines $\bar{\rho}$ and the extension Γ .

By definition, there is an inclusion $i: \text{Pic}(\mathcal{C}) \subset \mathcal{C}$ of symmetric monoidal ∞ -categories. The latter is presentable, whereas the former is (essentially) small. By formally adjoining colimits to \mathcal{C} , we obtain a factorization

$$\text{Pic}(\mathcal{C}) \rightarrow \mathcal{P}(\text{Pic}(\mathcal{C})) \rightarrow \mathcal{C}$$

through the presheaf ∞ -category, where the second functor preserves colimits.

Proposition 34. *For a symmetric monoidal ∞ -category \mathcal{C} , there is a diagram*

$$\mathcal{P}(\text{Pic}(\mathcal{C}, \rho)) \leftarrow \mathcal{P}(\text{Pic}(\mathcal{C})) \rightarrow \mathcal{C}$$

of symmetric monoidal presentable ∞ -categories.

Proof. Fix a regular cardinal κ such that the unit of \mathcal{C} is κ -compact. Then all of the objects of $\text{Pic}(\mathcal{C})$ are contained inside the essentially small subcategory \mathcal{C}^κ of κ -compact objects.

The functor \mathcal{P} is the left adjoint in an adjunction between κ -small ∞ -categories and κ -presentable ∞ -categories; $(-)^{\kappa}$ is the right adjoint. Moreover, the tensor product of presentable ∞ -categories is universal with respect to functors that are colimit-preserving in each variable separately; in particular, this gives us canonical identifications

$$\text{Fun}^{PrL}(\mathcal{P}(\prod S_i), \mathcal{C}) \simeq \text{Fun}^{PrL}(\otimes \mathcal{P}(S_i), \mathcal{C})$$

natural in \mathcal{C} . This makes the functor \mathcal{P} strong symmetric monoidal, and lifts it to a left adjoint to the functor taking a κ -presentable symmetric monoidal ∞ -category to the symmetric monoidal subcategory of κ -compact objects. For a small symmetric monoidal ∞ -category S , the induced symmetric monoidal structure on the category $\mathcal{P}(S)$ is given by left Kan extension: this is the Day convolution monoidal structure [Gla16, Lur17]. It is colimit-preserving in each variable, and for objects s and t of S with associated presheaves j_s and j_t there is a natural isomorphism $j_s \otimes j_t \cong j_{s \otimes t}$.

The Day convolution makes the functor $\mathcal{P}(\text{Pic}(\mathcal{C})) \rightarrow \mathcal{P}(\text{Pic}(\mathcal{C}, \rho))$ symmetric monoidal, and the adjunction gives us a composite symmetric monoidal functor

$$\mathcal{P}(\text{Pic}(\mathcal{C})) \rightarrow \mathcal{P}(\mathcal{C}^\kappa) \rightarrow \mathcal{C}$$

as desired. \square

Definition 35. Suppose that ρ is an extension to Γ -grading. We define the category obtained by *extending gradings to Γ* to be the symmetric presentable ∞ -category

$$\mathcal{C}[\rho] = \mathcal{P}(\text{Pic}(\mathcal{C}, \rho)) \otimes_{\mathcal{P}(\text{Pic}(R))} \mathcal{C}.$$

Proposition 36. *As a presentable category left-tensored over \mathcal{C} , we have*

$$\mathcal{C}[\rho] \cong \prod_{b \in B} \mathcal{C}.$$

In particular, $\mathcal{C}[\rho]$ is isomorphic to the category of B -graded objects of \mathcal{C} .

Proof. Since $\text{Pic}(\mathcal{C}, \rho) \cong \coprod_{b \in B} \text{Pic}(\mathcal{C})$ as categories left-tensored over $\text{Pic}(\mathcal{C})$,

$$\mathcal{P}(\text{Pic}(\mathcal{C}, \rho)) \cong \prod_{b \in B}^{pres} \mathcal{P}(\text{Pic}(\mathcal{C}))$$

as presentable ∞ -categories left-tensored over $\mathcal{P}(\mathcal{C})$ —here the coproduct taking place within presentable ∞ -categories. The relative tensor product preserves colimits in each variable, and thus we have

$$\mathcal{C}[\rho] \simeq \prod_{b \in B}^{pres} \mathcal{C}$$

as categories left-tensored over \mathcal{C} . However, within presentable ∞ -categories, coproducts and products over a small index set coincide. \square

One source of grading extensions is the theory of strict gradings.

Proposition 37. *Suppose that $0 \rightarrow G \rightarrow \Gamma \rightarrow B \rightarrow 0$ is an extension of abelian groups. Then every strict G -grading of \mathcal{C} has a naturally associated extension to a Γ -grading.*

Proof. The extension Γ is classified by a map $B \rightarrow \Sigma G$, which can be composed with the strict G -grading $G \rightarrow \text{pic}(\mathcal{C})$. \square

Example 38. Since MO has a strict \mathbb{Z} -grading, for any MO -algebra R we can then adjoin invertible objects L to the category of MO -modules such that $L^{\otimes n} \simeq \Sigma MO$, or extend to any grading $\mathbb{Z} \subset \Gamma$. For example, we can embed the category of MO -modules into the category of \mathbb{Q}/\mathbb{Z} -graded MO -modules, giving the latter a symmetric monoidal structure where shifts by integers are extended to shifts by rational numbers.

Example 39. Similarly, the strict $2\mathbb{Z}$ -grading on MU allow us to adjoin invertible objects L such that $L^{\otimes n} \cong \Sigma^2 MU$. (Note that if we take $n = 2$ in this construction, we find that the object $\Sigma^{-1}L$ is a nontrivial object with $(\Sigma^{-1}L)^{\otimes 2} \cong MU$.) This allows us to extend from a \mathbb{Z} -grading on MU -modules to a grading over $\mathbb{Z} \oplus_{2\mathbb{Z}} \mathbb{Q} \cong \mathbb{Q} \times \mathbb{Z}/2$. Similar constructions are possible with MU -algebras like ku or with the algebraic K -theory spectrum $K\mathbb{C}$.

8 Grading obstructions

As in §6, we can identify $\text{pic}(h\mathcal{C})$ with the first nontrivial stage $\tau_{\leq 1} \text{pic}(\mathcal{C})$ in the Postnikov tower for $\text{pic}(\mathcal{C})$. For us to lift a map $\bar{\rho}: B \rightarrow \Sigma \pi_0 \text{pic}(\mathcal{C})$ to the first Postnikov stage, it is necessary and sufficient that the associated obstruction

$$B \xrightarrow{\bar{\rho}} \Sigma \pi_0 \text{pic}(\mathcal{C}) \xrightarrow{k} \Sigma^3 \pi_1 \text{pic}(\mathcal{C})$$

is trivial. Two different choices of lift to a map $B \rightarrow \Sigma \text{pic}(h\mathcal{C})$ differ by an element of $[B \rightarrow \Sigma^2 \pi_1 \text{pic}(\mathcal{C})]$.

This can be concisely packaged into the following result.

Proposition 40. *Given an extension $0 \rightarrow \pi_0 \text{pic}(\mathcal{C}) \rightarrow \Gamma \rightarrow B \rightarrow 0$, the lifts of the map $\bar{\rho}: B \rightarrow \Sigma \pi_0 \text{pic}(\mathcal{C})$ to a map $\rho_{\leq 1}: B \rightarrow \Sigma \text{pic}(h\mathcal{C})$ are in bijective correspondence with extensions of the twist homomorphism $\tau: \pi_0 \text{pic}(\mathcal{C}) \rightarrow \pi_1 \text{pic}(\mathcal{C})[2]$ to all of Γ .*

Example 41. This construction can recover the twisted $\mathbb{Z}/2$ -graded categories of Rezk [Rez09, §2].⁴ Let \mathcal{C} be an ordinary presentable symmetric monoidal category and ω an invertible object of \mathcal{C} . We would like to construct a larger category $\mathcal{C}[\sqrt{\omega}]$ where ω is the square of another invertible module. This would be a symmetric monoidal category of $\mathbb{Z}/2$ -graded objects (A_0, A_1) of \mathcal{C} , representing $A_0 \oplus (\sqrt{\omega} \otimes A_1)$, with a tensor product satisfying

$$(A_0, A_1) \otimes (B_0, B_1) \cong \left((A_0 \otimes B_0) \oplus (\omega \otimes A_1 \otimes B_1), (A_0 \otimes B_1) \oplus (A_1 \otimes B_0) \right).$$

⁴Rezk gives these constructions in the case where \mathcal{C} is merely additive, which is not covered by our assumption that \mathcal{C} is presentable.

In this case, the underlying category is ordinary and so $\text{pic}(\mathcal{C}) = \text{pic}(h\mathcal{C})$. For us to extend gradings, it is necessary and sufficient that ω be symmetric. If this is the case, the possible choices of extension represent choices of 2-torsion element $\tau(\sqrt{\omega}) \in \text{Aut}_{\mathcal{C}}(\mathbb{I})$, a Koszul sign rule for the twist isomorphism on the square root of ω . If \mathcal{C} is additive, choosing $\tau(\sqrt{\omega}) = -1$ recovers Rezk's ω -twisted tensor product.

9 Brauer roots

Just as the unit spectrum is extended by the Picard spectrum, the Picard spectrum is extended by the Brauer spectrum [GL16, Hau17]. Fix a symmetric monoidal presentable ∞ -category \mathcal{C} and let $\text{Cat}_{\mathcal{C}}$ be the category of presentable ∞ -categories left-tensored over \mathcal{C} . This has a symmetric monoidal under $\otimes_{\mathcal{C}}$. We write $\text{Br}(\mathcal{C}) = \text{Pic}(\text{Cat}_{\mathcal{C}})$ for the Brauer space parametrizing invertible objects of $\text{Cat}_{\mathcal{C}}$ that admit a compact generator, and $\text{br}(\mathcal{C})$ for the associated spectrum.

The unit of $\text{Cat}_{\mathcal{C}}$ is \mathcal{C} itself, and all \mathcal{C} -linear functors are of the form $X \mapsto X \otimes B$ for some $B \in \mathcal{C}$; in particular, this identifies the \mathcal{C} -linear functors $\mathcal{C} \rightarrow \mathcal{C}$ with \mathcal{C}^{\simeq} itself, with composition given by the tensor in \mathcal{C} . As a result, the space of self-equivalences of the unit is $\text{Pic}(\mathcal{C})$, and so there is a fiber sequence

$$\Sigma \text{pic}(\mathcal{C}) \rightarrow \text{br}(\mathcal{C}) \rightarrow \pi_0 \text{br}(\mathcal{C}).$$

Definition 42. Suppose that \mathcal{C} is a presentable symmetric monoidal ∞ -category, B is an abelian group regarded as a discrete symmetric monoidal category, and that $\rho: B \rightarrow \text{Cat}(\mathcal{C})$ is a symmetric monoidal functor. We define the *trivializing category* \mathcal{T}_{ρ} to be the homotopy colimit of ρ , calculated in $\text{Cat}_{\mathcal{C}}$.

Remark 43. Since B is a discrete category, there is an equivalence in $\text{Cat}_{\mathcal{C}}$ of the form

$$\mathcal{T}_{\rho} \simeq \coprod_{b \in B}^{\text{Cat}_{\mathcal{C}}} \rho(b) \simeq \prod_{b \in B} \rho(b).$$

Proposition 44. *Suppose that B is an abelian group and that $\rho: B \rightarrow \mathcal{C}$ is a symmetric monoidal functor with trivializing category \mathcal{T}_{ρ} .*

1. *The map ρ has a natural lift to a map $B \rightarrow \text{Br}(\mathcal{C})$.*
2. *\mathcal{T}_{ρ} has a natural lift to a symmetric monoidal ∞ -category under \mathcal{C} .*
3. *The symmetric monoidal ∞ -category \mathcal{T}_{ρ} is universal among symmetric monoidal presentable ∞ -categories under \mathcal{C} with a chosen commutative diagram*

$$\begin{array}{ccc} B & \longrightarrow & \text{br}(\mathcal{C}) \\ \downarrow & & \downarrow \\ * & \longrightarrow & \text{br}(\mathcal{T}_{\rho}). \end{array}$$

In particular, for any $b \in B$ there is an equivalence $\phi_b: \mathcal{T}_\rho \rightarrow \mathcal{T}_\rho \otimes_{\mathcal{C}} \rho(b)$ of presentable ∞ -categories left-tensored over \mathcal{T}_ρ , and there are chosen equivalences of functors $\phi_a \otimes 1 \circ \phi_b \xrightarrow{\sim} \phi_{ab}$.

Proof. This is Proposition 25, applied to the (large) ∞ -category of presentable ∞ -categories. \square

Example 45. Let $\mathcal{C} = \text{LMod}_R$ where R is a commutative ring spectrum. Then each object of $\text{Br}(R)$ is represented by the category of right modules over an Azumaya R -algebra Q which is well-defined up to Morita equivalence [AG14, Toë12, BRS], see also [GL16, 5.13]. Given a symmetric monoidal functor $\rho: B \rightarrow \text{Br}(R)$, we can therefore choose algebras $Q(b)$ so that there is an equivalence

$$\mathcal{T}_\rho \simeq \prod_{b \in B} \text{RMod}_{Q(b)}.$$

The symmetric monoidal structure on this category takes more work to describe. The symmetric monoidal structure on ρ gives Morita equivalences between $Q(p) \otimes_R Q(q)$ and $Q(p+q)$, which are expressed by $Q(p) \otimes_R Q(q) \otimes_R Q(p+q)^{op}$ -modules $M_{p,q}$. The symmetric monoidal structure is given by a formula of the form

$$(X_b)_{b \in B} \otimes (Y_b)_{b \in B} \cong \left(\bigoplus_{p+q=b} (X_p \otimes Y_q) \otimes_{Q(p) \otimes_R Q(q)} M_{p,q} \right)_{b \in B}.$$

However, describing the full symmetric monoidal structure in this fashion would require us to carefully express coherence relations between tensor products of the bimodules $M_{p,q}$. We can think of these objects as “coefficients for the multiplication” that give the symmetric monoidal structure on this product category.

Remark 46. The first k -invariant in the Brauer spectrum is a map $\pi_0 \text{br}(\mathcal{C}) \rightarrow \Sigma^2 \pi_0 \text{pic}(\mathcal{C})$, and this is still expressed by the twist self-equivalence. However, now this twist self-equivalence occurs on the level of module categories. For a commutative ring spectrum R with an Azumaya R -algebra Q , the twist equivalence of $\text{RMod}_{Q \otimes_R Q}$ is expressed by tensoring with the $Q \otimes_R Q$ -bimodule $Q \otimes_R Q^\tau$, where the action on the left is the standard one and on the right factors through the twist automorphism. The fact that $Q \otimes_R Q$ is Azumaya means that this bimodule must be of the form $Q \otimes_R Q \otimes_R \tau(Q)$ for some $\tau(Q) \in \text{Pic}(R)$.⁵

More concretely, we can identify $\tau(Q)$ with the R -module

$$F_{Q \otimes_R Q\text{-bimod}}(Q \otimes_R Q, (Q \otimes_R Q)^\tau).$$

This assignment remains relatively mysterious to the author. In algebraic examples it is trivial, which is shown by descent-theoretic methods in [Vis].

⁵The algebraic version of this assignment, taking k -algebra automorphisms of an Azumaya algebra T to elements in the Picard group, is part of the Rosenberg-Zelinsky exact sequence

$$1 \rightarrow k^\times \rightarrow T^\times \rightarrow \text{Aut}_{\text{Alg}(k)}(T) \rightarrow \pi_0 \text{pic}(k)$$

that expresses the potential failure of algebra automorphisms to be inner. If k is a field this recovers the Noether-Skolem theorem.

10 Remarks on algebraic K -theory

We close with some conjectural remarks on the relationship with algebraic K -theory.

We recall the following from [GL16, 5.11]. The functor Mod is symmetric monoidal by [Lur17, 4.8.5.16]: for A and B R -algebras, the tensor product in Mod_R -linear ∞ -categories has a natural equivalence

$$\text{Mod}_A \otimes \text{Mod}_B \simeq \text{Mod}_{A \otimes_R B}.$$

Moreover, by [GGN15, 8.6] the algebraic K -theory functor is lax symmetric monoidal, and hence induces a functor

$$\text{Cat}_R \rightarrow \text{Mod}_{KR}.$$

While this functor is only lax symmetric monoidal, it does send the unit Mod_R to the unit KR , and hence the restriction to the unit component is a symmetric monoidal functor

$$BPic(R) \rightarrow BGL_1(KR).$$

In other words, there is a spectrum map $\text{pic}(R) \rightarrow \text{gl}_1(KR)$, inducing a map $\Sigma \text{pic}(R) \rightarrow \text{pic}(KR)$. (The author sees no evidence that this extends to a map $\text{br}(R) \rightarrow \text{pic}(KR)$.)

A map $\rho: B \rightarrow \Sigma \text{pic}(R)$, which can be used to embed Mod_R into a symmetric monoidal category of B -graded modules, therefore also induces a map $B \rightarrow \Sigma \text{gl}_1 KR$, which can be used to extend the unit group of the algebraic K -theory spectrum. Moreover, because B is discrete the additive structure of the extension of Mod_R and the additive structure of the extension of KR are straightforward. The universal property can be used to show that the algebraic K -theory functor takes the category of compact B -graded R -modules, with the symmetric monoidal structure determined by ρ , to the induced extension of KR .

References

- [AB14] Omar Antolin-Camarena and Tobias Barthel, *A simple universal property of Thom ring spectra*, arXiv e-prints (2014), arXiv:1411.7988.
- [ABG⁺] Matthew Ando, Andrew Blumberg, David Gepner, Michael Hopkins, and Charles Rezk, *Units of ring spectra and thom spectra*, arXiv:0810.4535.
- [ABG⁺14] Matthew Ando, Andrew J. Blumberg, David Gepner, Michael J. Hopkins, and Charles Rezk, *Units of ring spectra, orientations and Thom spectra via rigid infinite loop space theory*, J. Topol. **7** (2014), no. 4, 1077–1117. MR 3286898
- [AG14] Benjamin Antieau and David Gepner, *Brauer groups and étale cohomology in derived algebraic geometry*, Geom. Topol. **18** (2014), no. 2, 1149–1244. MR 3190610
- [BMMS86] R. R. Bruner, J. P. May, J. E. McClure, and M. Steinberger, *H_∞ ring spectra and their applications*, Lecture Notes in Mathematics, vol. 1176, Springer-Verlag, Berlin, 1986. MR 836132 (88e:55001)

- [BRS] Andrew Baker, Birgit Richter, and Markus Szymik, *Brauer groups for commutative S -algebras*, arXiv:1005.5370.
- [Cla11] Dustin Clausen, *p -adic J -homomorphisms and a product formula*, arXiv e-prints (2011), arXiv:1110.5851.
- [Dev17] Sanath K. Devalapurkar, *Roots of unity in $K(n)$ -local E_∞ -rings*, arXiv e-prints (2017), arXiv:1707.09957.
- [EML54] Samuel Eilenberg and Saunders Mac Lane, *On the groups $H(\Pi, n)$. II. Methods of computation*, Ann. of Math. (2) **60** (1954), 49–139. MR 65162
- [GGN15] David Gepner, Moritz Groth, and Thomas Nikolaus, *Universality of multiplicative infinite loop space machines*, Algebr. Geom. Topol. **15** (2015), no. 6, 3107–3153. MR 3450758
- [GH04] Paul G. Goerss and Michael J. Hopkins, *Moduli spaces of commutative ring spectra*, Structured ring spectra, London Math. Soc. Lecture Note Ser., vol. 315, Cambridge Univ. Press, Cambridge, 2004, pp. 151–200. MR 2125040 (2006b:55010)
- [GL16] D. Gepner and T. Lawson, *Brauer groups and Galois cohomology of commutative ring spectra*, ArXiv e-prints (2016).
- [Gla16] Saul Glasman, *Day convolution for ∞ -categories*, Math. Res. Lett. **23** (2016), no. 5, 1369–1385. MR 3601070
- [Hau17] Rune Haugseng, *The higher Morita category of \mathbb{E}_n -algebras*, Geom. Topol. **21** (2017), no. 3, 1631–1730. MR 3650080
- [Hop14] Michael J. Hopkins, *$K(1)$ -local E_∞ -ring spectra*, Topological modular forms, Math. Surveys Monogr., vol. 201, Amer. Math. Soc., Providence, RI, 2014, pp. 287–302. MR 3328537
- [Joa04] Michael Joachim, *Higher coherences for equivariant K -theory*, Structured ring spectra, London Math. Soc. Lecture Note Ser., vol. 315, Cambridge Univ. Press, Cambridge, 2004, pp. 87–114. MR 2122155
- [Law18] Tyler Lawson, *Secondary power operations and the Brown–Peterson spectrum at the prime 2*, Ann. of Math. (2) **188** (2018), no. 2, 513–576. MR 3862946
- [LN12] Tyler Lawson and Niko Naumann, *Commutativity conditions for truncated Brown–Peterson spectra of height 2*, J. Topol. **5** (2012), no. 1, 137–168. MR 2897051
- [LN14] ———, *Strictly commutative realizations of diagrams over the Steenrod algebra and topological modular forms at the prime 2*, Int. Math. Res. Not. IMRN (2014), no. 10, 2773–2813. MR 3214285

- [Lur09] Jacob Lurie, *Higher topos theory*, Annals of Mathematics Studies, vol. 170, Princeton University Press, Princeton, NJ, 2009. MR 2522659 (2010j:18001)
- [Lur17] ———, *Higher Algebra*, Draft version available at: <http://www.math.harvard.edu/~lurie/papers/higheralgebra.pdf>, 2017.
- [Mat14] Akhil Mathew, *The Galois group of a stable homotopy theory*, 2014.
- [May77] J. Peter May, *E_∞ ring spaces and E_∞ ring spectra*, Lecture Notes in Mathematics, Vol. 577, Springer-Verlag, Berlin-New York, 1977, With contributions by Frank Quinn, Nigel Ray, and Jørgen Tornehave. MR 0494077 (58 #13008)
- [MS16] Akhil Mathew and Vesna Stojanoska, *The Picard group of topological modular forms via descent theory*, Geom. Topol. **20** (2016), no. 6, 3133–3217. MR 3590352
- [Rez06] Charles Rezk, *The units of a ring spectrum and a logarithmic cohomology operation*, J. Amer. Math. Soc. **19** (2006), no. 4, 969–1014. MR 2219307 (2007h:55006)
- [Rez09] ———, *The congruence criterion for power operations in Morava E -theory*, Homology, Homotopy Appl. **11** (2009), no. 2, 327–379. MR 2591924 (2011e:55021)
- [Rob03] Alan Robinson, *Gamma homology, Lie representations and E_∞ multiplications*, Invent. Math. **152** (2003), no. 2, 331–348. MR 1974890
- [Sag16] Steffen Sagave, *Spectra of units for periodic ring spectra and group completion of graded E_∞ spaces*, Algebr. Geom. Topol. **16** (2016), no. 2, 1203–1251. MR 3493419
- [SVW99] R. Schwänzl, R. M. Vogt, and F. Waldhausen, *Adjoining roots of unity to E_∞ ring spectra in good cases—a remark*, Homotopy invariant algebraic structures (Baltimore, MD, 1998), Contemp. Math., vol. 239, Amer. Math. Soc., Providence, RI, 1999, pp. 245–249. MR 1718085 (2001b:55026)
- [Toë12] Bertrand Toën, *Derived Azumaya algebras and generators for twisted derived categories*, Invent. Math. **189** (2012), no. 3, 581–652. MR 2957304
- [Vis] Angelo Vistoli, *On a morphism from the brauer group to the picard group*, MathOverflow, <https://mathoverflow.net/q/334293>.