Probability distribution connected to stationary flow of substance in a channel of network containing finite number of arms

Roumen Borisov¹, Zlatinka I. Dimitrova², Nikolay K. Vitanov¹

¹Institute of Mechanics, Bulgarian Academy of Sciences, Acad. G. Bonchev Str., Bl. 4, 1113 Sofia, Bulgaria ²"G. Nadjakov" Institute of Solid State Physics, Bulgarian Academy of Sciences, Blvd. Tzarigradsko Chaussee 72, 1784, Sofia, Bulgaria

Abstract

We discuss a channel consisting of nodes of a network and lines which connect these nodes and form ways for motion of a substance through the channel. We study stationary flow of substance for channel which arms contain finite number of nodes each and obtain probability distribution for substance in arms of this channel. Finally we calculate Shannon information measure for the case of stationary flow of substance in a simple channel consisting of a single arm having just three nodes.

1 Introduction

In the course of time researchers began to study systems with increased complexity in different areas of sciences such as, for example, natural sciences [2] -[8], [12], [13] population dynamics [14] - [17], [26], [32],[45],[47] or social sciences [27],[44],[46]. Large number of structures and processes in these complex systems can be modeled by networks and differential equations [11], [19], [22], [25], [28] - [43], [54]. Our interest is concentrated on motions of substance through channels of complicated structures. These motions are theoretically interesting and of significant practical importance [1], [18] as they are connected to transportation problems [21], [23], migration flows [24], [20], [48] or other kinds of flows in networks [10]. In this article flow

of substance in a channel which has have arbitrary number of arms. The channel has special nodes where split of an arm happens. More than one arm may arise by this split. The substance can move only in one direction along the channel. The substance can also move out of the channel - "leakage" of substance.

The organization of text is as follows. In Sect. 2 we discuss a mathematical model for flow of substance in studied channel for the case when each channel arm has finite number of nodes. We obtain probability distribution connected to amounts of substance in nodes of arms of this channel for the case of stationary flow of substance. In Sect. 3 we discuss information measures connected to the obtained probability distributions.

2 Flow of substance in a channel with finite number of nodes in each arm

The discussed model of motion of substance is an extension of model discussed in [48], [35]. The channel contains chains of nodes of the network and the convention for numbering of nodes of channel is as follows. We assign 4 indexes to each node: $\mathcal{V}_{i,j}^{a,b}$. Lower indexes specify position of node in current arm. *i* is associated with current arm. *j* is the number of node of the *i*-th arm. Upper indexes specify the origin of arm *i*. Index *a* denotes number of arm from which arm *i* splits. Index *b* denotes number of node of the arm *a* where this split happens. Then $\mathcal{V}_{4,6}^{8,4}$ means the 6-th node of arm 4 which splits at node 4 of channel's arm 8. The node where arm *i* begins is labeled as the 0-th node of *i*-th arm. This node is the next one after splitting at node *b* of arm *a*.

The motion of substance in arms of studied channel is as follows. Substance enters the channel from external environment only through channel's main arm (labeled by q = 0 below in the text). The substance can move only in one direction in any arm: from nodes labeled by smaller number to nodes labeled by larger numbers. The nodes of each arm are connected by lines and each node is connected only to two neighboring nodes of the arm exclusive for nodes where a split of an arm happens. Last nodes can be connected to one or more additional nodes. We assume that substance can quit the channel and can move to environment - "leakage" of substance. As substance can enter the channel only through 0-th node of main arm then leakage is possible only in direction from channel to environment and not in opposite direction.

We consider each arm to be an array of of cells indexed in succession by

non-negative integers. We assume that an amount $x_q^{a_q,b_q}$ of some substance is distributed among cells of the arm q which splits at node (a_q, b_q) of the network. This substance can move from one cell to next cell. Let $x_{q,i}^{a_q,b_q}$ be the amount of substance in *i*-th cell of q-th arm of channel. We consider first a channel containing number of nodes $N_q + 1$ in each of its arms (the

number of arms is M). For this case $x_q^{a_q,b_q} = \sum_{i=0}^{N_q} x_{q,i}^{a_q,b_q}$. The fractions $y_{q,i}^{a_q,b_q} =$

 $x_{q,i}^{a_q,b_q}/x_q^{a_q,b_q}$ are considered as probability values of distribution of a discrete random variable ζ in corresponding arm of channel: $y_{q,i}^{a_q,b_q} = p_q^{a_q,b_q}(\zeta = i)$, $i = 0, 1, \ldots$ The amount $x_{q,i}^{a_q,b_q}$ of substance in *i*-th node of *q*-th arm of channel can change because of the following processes. First of all some amount s_q^{a,b_q} of substance enters arm *q* from external environment through 0-th cell of the arm. We consider two kinds of external environments. For the root of the channel (arm with label q = 0): substance $s_0^{0,0}$ enters the root through environment of the channel. For the arms of the channel that are not root $(q \neq 0)$: substance $s_q^{a,b}$ is part of substance presented in node (a, b) of parent arm.

The substance $s_q^{a,b}$ is presented only in node 0 of q-th arm of channel. For other nodes of channel there is no substance that enters the node from environment or from other arms of the channel. Next amount $f_{q,i}^{a,b}$ from $x_{q,i}^{a,b}$ is transferred from the *i*-th cell into i + 1-th cell of q-th arm and amount $g_{q,i}^{a,b}$ of $x_{q,i}^{a,b}$ leaks out the *i*-th cell of q-th arm into environment of the arm of the channel. This leakage can be of two kinds. First of all there can be leakage to the environment of the channel. This leads to loss of substance for the channel. In addition there can be a leakage to other arms of the channel which begin from node *b* of arm *a*. This leakage is connected to the substance $s_q^{a,b}$ that enters corresponding child arm of the channel which splits from node *b* of arm *a*.

The process of motion of substance is continuous in time and motion of substance among nodes of q-th channel is modeled mathematically by a system of ordinary differential equations which contains an equation for 0th node, equations for nodes $1, \ldots, N_{q-1}$ and equation for node N_q . This system is $\frac{dx_{q,0}^{a,b}}{dt} = s_q^{a,b} - f_{q,0}^{a,b} - g_{q,0}^{a,b}$; $\frac{dx_{q,i}^{a,b}}{dt} = f_{q,i-1}^{a,b} - f_{q,i}^{a,b} - g_{q,i}^{a,b}$, $i = 1, 2, N_q 1 \frac{dx_{q,N_q}^{a,b}}{dt} = f_{q,N_q-1}^{a,b} - g_{q,N_q}^{a,b}$. We shall discuss below the stationary regime of functioning of channel where $dx_{i,q}^{a,b}/dt = 0$, $i = 0, 1, N_q$. We mark the quantities for the stationary case with *. Then from equations above one obtains: $f_{q,0}^{*a,b} = s_q^{*a,b} - g_{q,0}^{*a,b}$; $f_{q,i}^{*a,b} = f_{q,i-1}^{*a,b} - g_{q,i}^{*a,b}$; $f_{q,N_q}^{*a,b} = f_{q,N_q-1}^{*a,b} - g_{q,N_q}^{*a,b}$. We assume the following forms of amounts of moving substances in above equations $(\alpha_i, \beta_i, \gamma_i, \sigma \text{ are parameters})$: $s_0^{0,0} = \sigma_0 x_{0,0}^{0,0} > 0$; $s_q^{a,b} = \delta_q x_{a,b}^{c,d}$ $(1 \ge \delta_q \ge 0)$; $f_{q,i}^{a,b} = \alpha_{q,i}^{a,b} x_{q,i}^{a,b}$ $(1 > \alpha_{q,i}^{a,b} > 0)$; $g_{q,i}^{a,b} = \gamma_{q,i}^{*a,b} x_{q,i}^{a,b}$; $1 \ge \gamma_{q,i}^{*a,b} \ge 0 \rightarrow$ non-uniform leakage in the nodes. Indexes c and d above describe parent arm (numbered by c) and parent node of the arm c (numbered by d) for the arm q. $\gamma_{q,i}^{*a,b} = \gamma_{q,i}^{a,b} + \sum_{p \in (q,i)} \delta_{p,q,i}^{a,b}$ describes the situation with leakages in cells

where new arm arises and part of substance flows to this new arm. $\gamma_{q,i}^{a,b}$ is the leakage to environment from *i*-th node of *q*-th arm. Notation $p \in (q, i)$ in the sum denotes all the arms which arise from node (q, i). The terms in the sum describe leakages because of outflow of substance from parent arm to child arms. The model system of equations for node 0 and for nodes $1, \ldots, N_{q-1}$ of *q*-th arm of the channel is

$$\frac{dx_{q,0}^{a,b}}{dt} = s_q^{a,b} - \alpha_{q,0}^{a,b} x_{q,0}^{a,b} - \gamma_{q,0}^{*a,b} x_{q,0}^{a,b},
\frac{dx_{q,i}^{a,b}}{dt} = \alpha_{q,i-1}^{a,b} x_{q,i-1}^{a,b} - (\alpha_{q,i}^{a,b} + \gamma_{q,i}^{*a,b}) x_{q,i}^{a,b}; \quad i = 1, 2, \dots, N_q - 1.$$
(1)

For node N_q of q-th arm there is no outflow to next mode of the arm (as the node N_q is the last node of q-th arm) and the equation for motion of substance for this node is

$$\frac{dx_{q,N}^{a,b}}{dt} = \alpha_{q,N_q-1}^{a,b} x_{q,N_q-1}^{a,b} - \gamma_{q,N_q}^{*a,b} x_{q,N_q}^{a,b}.$$
 (2)

We discuss stationary motion of substance through arms of studied channel. Then $dx_{q,0}^{a,b}/dt = 0$ in the first of Eqs. (1) and

$$x_{q,0}^{*a,b} = \frac{s_q^{a,b}}{\alpha_{0,q}^{a,b} + \gamma_{0,q}^{*,a,b}} \quad . \tag{3}$$

For root of the channel (arm 0) we substitute $s_0^{0,0}$ in (3) and obtain that $x_{0,0}^{0,0}$ is a free parameter and in addition $\sigma_0 = \alpha_{0,0}^{0,0} + \gamma_{0,0}^{*,0,0}$. For arm r which arises from node m of arm q, $dx_{r,0}^{q,m}/dt = 0$ and thus we obtain

$$x_{r,0}^{*q,m} = \frac{\delta_{r,q,m}^{a,m}}{\alpha_{r,0}^{q,m} + \gamma_{r,0}^{q,m}} x_{q,m}^{*c,d}.$$
(4)

For $x_{q,i}^{*a,b}$ one obtains the relationship (just set $dx_{q,i}^{a,b}/dt = 0$ in second of Eqs. (1))

$$x_{q,i}^{*a,b} = \frac{\alpha_{q,i-1}^{a,b}}{\alpha_{q,i}^{a,b} + \gamma_{q,i}^{*a,b}} x_{q,i-1}^{*a,b}, \ i = 1, 2, \dots, N_q - 1;$$
(5)

In order to calculate $x_{q,N_q}^{*a,b}$ we have to use (2). The result is

$$x_{q,N_q}^{*a,b} = \frac{\alpha_{q,N_q-1}^{a,b}}{\gamma_{q,N_q}^{*a,b}} x_{q,N_q-1}^{*a,b}.$$
 (6)

From (5) we obtain

$$x_{q,i}^{*a,b} = \frac{\prod_{j=0}^{i-1} \alpha_{q,i-j-1}^{a,b}}{\prod_{j=0}^{i-1} \left[\alpha_{q,i-j}^{a,b} + \gamma_{q,i-j}^{*a,b} \right]} x_{q,0}^{*a,b}, \quad i = 1, \dots, N_q - 1.$$
(7)

From Eq.(6) one obtains


$$x_{q,N_q}^{*a,b} = \frac{\alpha_{q,N_q-1}^{a,b}}{\gamma_{q,N_q}^{*a,b}} \frac{\prod_{j=0}^{N_q-2} \alpha_{q,N_q-j-2}^{a,b}}{\prod_{j=0}^{N_q-2} \left[\alpha_{q,N_q-j-1}^{a,b} + \gamma_{q,N_q-j-1}^{*a,b}\right]} x_{q,0}^{*a,b}.$$
(8)


The total amount of the substance in q-th arm of the channel is

$$\begin{pmatrix}
x_{q}^{*,a,b} = x_{q,0}^{*a,b} + x_{q,i}^{*,a,b} + x_{q,N}^{*,a,b} = x_{q,0}^{*a,b} \times \\
x_{q}^{*,a,b} = x_{q,0}^{*a,b} + x_{q,i}^{*,a,b} + x_{q,N}^{*,a,b} = x_{q,0}^{*a,b} \times \\
\begin{pmatrix}
1 + \sum_{i=1}^{N_{q-1}} \prod_{j=0}^{i-1} \alpha_{q,i-j-1}^{a,b} \\
\prod_{j=0}^{i-1} \left[\alpha_{q,i-j}^{a,b} + \gamma_{q,i-j}^{*a,b} \right] \\
+ \frac{\alpha_{q,N_{q-1}}^{a,b}}{\gamma_{q,N_{q}}^{*a,b}} \prod_{j=0}^{N_{q-2}} \left[\alpha_{q,N_{q-j-1}}^{a,b} + \gamma_{q,N_{q-j-1}}^{*a,b} \right] \\
\end{pmatrix}$$
(9)

The distribution of substance in nodes of q-th arm of the channel is

$$y_{q,0}^{*a,b}=1 \Bigg/$$





$$y_{q,N_q}^{*a,b} = \left(\frac{\alpha_{q,N_q-1}^{a,b}}{\gamma_{q,N_q}^{*a,b}} \frac{\prod_{j=0}^{N_q-2} \alpha_{q,N_q-j-2}^{a,b}}{\prod_{j=0}^{N_q-2} \left[\alpha_{q,N_q-j-1}^{a,b} + \gamma_{q,N_q-j-1}^{*a,b} \right]} \right) /$$

$$\left(1 + \sum_{i=1}^{N_{q}-1} \frac{\prod_{j=0}^{i-1} \alpha_{q,i-j-1}^{a,b}}{\prod_{j=0}^{i-1} \left[\alpha_{q,i-j}^{a,b} + \gamma_{q,i-j}^{*a,b}\right]} + \frac{\alpha_{q,N_{q}-1}^{a,b}}{\gamma_{q,N_{q}}^{*a,b}} \frac{\prod_{j=0}^{N_{q}-2} \alpha_{q,N_{q}-j-2}^{a,b}}{\prod_{j=0}^{N_{q}-2} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b}\right]}\right)$$
(12)

3 Information measures for studied channel

We have discussed probability distributions corresponding to a channel with single arm in [9], [49] - [53]. Particular cases of these distributions are Waring distribution, Zipf distribution, Yule-Simon distribution, Binomial distribution, etc. We can study also other kinds of distributions on the basis of the model discussed in this text. One example is probability distribution of the substance in a part of the studied network which contains several channels for motion of substance. We can calculate various characteristics of the distributions obtained above and let us consider an information problem connected to the flow of substance in studied channel. Each node of the channel is numbered and we consider the nodes as letters of an alphabet. Some kind of an event happens in any of nodes of the channel and let probability of happening of this event be proportional to amount of substance in corresponding node. Thus the probability of happening of event in a node of channel will be equal to the probability from corresponding probability distribution obtained above in the text. The channel (the source) will generate events with corresponding probability and we can calculate measure of information and Shannon measure of information for these sequences as probability distribution is known. The information measure connected to an event with probability p is $I(p) = -\log(p)$, and Shannon information measure (average information we get from a symbol in a stream) connected to probability distribution $P = (p_0, \ldots, p_N)$ is $H(P) = -\sum_{i=0}^N p_i \log(p_i)$. Let us consider distribution P^* of substance in the q-th arm of the channel given by (10) - (12). The information connected to event with probability p_i from *i*-th node of this arm is

$$I(p_{i}) = -\log\left(\frac{\prod_{j=0}^{i-1} \alpha_{q,i-j-1}^{a,b}}{\prod_{j=0}^{i-1} \left[\alpha_{q,i-j}^{a,b} \gamma_{q,i-j}^{*a,b}\right]}\right) + \log\left(1 + \sum_{i=1}^{N_{q}-1} \frac{\prod_{j=0}^{i-1} \alpha_{q,i-j-1}^{a,b}}{\prod_{j=0}^{i-1} \left[\alpha_{q,i-j}^{a,b} + \gamma_{q,i-j}^{*a,b}\right]} + \frac{\alpha_{q,N_{q}-1}^{a,b}}{\frac{\alpha_{q,N_{q}-1}^{a,b}}{\gamma_{q,N_{q}}^{*a,b}}} \frac{\prod_{j=0}^{N_{q}-2} \alpha_{q,N_{q}-j-2}^{a,b}}{\prod_{j=0}^{N_{q}-2} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b}\right]}\right), \ i = 1, \dots, N_{q} - 1$$

$$(13)$$

For the 0-th node and for the $N_q\mbox{-th}$ node we obtain the following relationships for the information measures

$$I(p_{0}) = \log \left(1 + \sum_{i=1}^{N_{q}-1} \frac{\prod_{j=0}^{i-1} \alpha_{q,i-j-1}^{a,b}}{\prod_{j=0}^{i-1} \left[\alpha_{q,i-j}^{a,b} + \gamma_{q,i-j}^{*a,b} \right]} + \frac{\alpha_{q,N_{q}-1}^{a,b}}{\gamma_{q,N_{q}}^{*a,b}} \frac{\prod_{j=0}^{N_{q}-2} \alpha_{q,N_{q}-j-2}^{a,b}}{\prod_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right)$$
$$I(p_{N}) = -\log \left(\frac{\alpha_{q,N_{q}-1}^{a,b}}{\gamma_{q,N_{q}}^{*a,b}} \frac{\prod_{j=0}^{N_{q}-2} \alpha_{q,N_{q}-j-2}^{a,b}}{\prod_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}} \left(\frac{\alpha_{q,N_{q}-1}^{a,b}}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}} \left(\frac{\alpha_{q,N_{q}-1}^{a,b}}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}} \left(\frac{\alpha_{q,N_{q}-1}^{a,b}}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}} \left(\frac{\alpha_{q,N_{q}-1}^{a,b}}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}} \left(\frac{\alpha_{q,N_{q}-1}^{a,b}}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}} \left(\frac{\alpha_{q,N_{q}-1}^{a,b}}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}} \left(\frac{\alpha_{q,N_{q}-1}^{a,b}}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}} \left(\frac{\alpha_{q,N_{q}-j}^{a,b}}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}} \left(\frac{\alpha_{q,N_{q}-j}^{a,b}}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]}{\sum_{j=0}^{I} \left[\alpha_{q,N_{q}-j-1}^{a,b} + \gamma_{q,N_{q}-j-1}^{*a,b} \right]} \right) + \frac{1}{N_{q}$$

$$\log\left(1+\sum_{i=1}^{N_{q}-1}\frac{\prod_{j=0}^{i-1}\alpha_{q,i-j-1}^{a,b}}{\prod_{j=0}^{i-1}\left[\alpha_{q,i-j}^{a,b}+\gamma_{q,i-j}^{*a,b}\right]}+\frac{\alpha_{q,N_{q}-1}^{a,b}}{\gamma_{q,N_{q}}^{*a,b}}\frac{\prod_{j=0}^{N_{q}-2}\alpha_{q,N_{q}-j-2}^{a,b}}{\prod_{j=0}^{N_{q}-2}\left[\alpha_{q,N_{q}-j-1}^{a,b}+\gamma_{q,N_{q}-j-1}^{*a,b}\right]}\right).(14)$$

The general expression for the Shannon information measure is very long and we will not write it here. Instead of this we present the Shannon information measure for a simple case of a channel arm that has just three nodes. We shall omit the indices a, b and q. For discussed case $N_q = N = 2$ as labels of nodes of the arm are 0, 1 and 2. The parameters below account for following processes in the studied arm. α_0 ($0 < \alpha_0 < 1$) accounts for flow between first and second node; α_1 ($0 < \alpha_1 < 1$) accounts for flow between second and third node; β_1 ($0 < \beta_1 < 1 - \alpha_1$) accounts for preference for the third node; γ_1^* ($0 \le \gamma_1^* < 1 - \alpha_1 - \beta_1$) accounts for leakage from the second node; γ_2^* ($0 < \gamma_2^* \le 1$) accounts for leakage from the third node. The Shannon information measure is

$$H = \frac{1}{\left[1 + \frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}} \left(1 + \frac{\alpha_{1} + \beta_{1}}{\gamma_{2}^{*}}\right)\right]} \log \left[1 + \frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}} \left(1 + \frac{\alpha_{1} + \beta_{1}}{\gamma_{2}^{*}}\right)\right] + \frac{\left(\frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}}\right)}{\left[1 + \frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}} \left(1 + \frac{\alpha_{1} + \beta_{1}}{\gamma_{2}^{*}}\right)\right]} \left\{\log \left[1 + \frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}} \left(1 + \frac{\alpha_{1} + \beta_{1}}{\gamma_{2}^{*}}\right)\right] - \log \left(\frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}}\right)\right\} + \frac{\left(\frac{\alpha_{1} + \beta_{1}}{\gamma_{2}^{*}} + \frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}}\right)}{\left[1 + \frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}} \left(1 + \frac{\alpha_{1} + \beta_{1}}{\gamma_{2}^{*}}\right)\right]} \left\{\log \left[1 + \frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}} \times \left(1 + \frac{\alpha_{1} + \beta_{1}}{\gamma_{2}^{*}}\right)\right] - \log \left(\frac{\alpha_{1} + \beta_{1}}{\gamma_{2}^{*}} + \frac{\alpha_{0}}{\alpha_{1} + \beta_{1} + \gamma_{1}^{*}}\right)\right\}.$$
(15)

The studies show that H can have a maximum value for some value of the parameters of the problem. This means that average information we can obtain from an events happening in the channel can have local maxima for some values of problem parameters.

References

 Albert, R., Barabasi, A.-L.: Statistical mechanics of complex networks. Rev. Mod. Phys. 74, 47 – 97 (2002).

- [2] Amaral, L. A. N., Ottino, J. M.: Complex networks. Augmenting and framework for the study of complex systems. Eur. Phys. J. B 38, 147 – 162 (2004).
- [3] Ashenfelter, K.T., Boker, S.M., Waddell, J.R., Vitanov, N.: Spatiotemporal symmetry and multifractal structure of head movements during dyadic conversation. Journal of Experimental Psychology: Human Perception and Performance 35 1072 – 1091 (2009)
- [4] Ausloos, M., Cloots, R., Gadomski, A., Vitanov, N.K.: Ranking structures and rankrank correlations of countries: The FIFA and UEFA cases. International Journal of Modern Physics C 25, 1450060 (2014)
- [5] Basuchoudhary, A., Bang, J. T., Sen, T.: Machine-learning techniques in economics. New tools for predicting economic growth. Springer, Cham (2017)
- [6] Bo Wang, Dimitrova, Z. I., Vitanov, N. K.: Statistical analysis of the water level of Huang He river (Yellow river) in China. Journal of Theoretical and Applied Mechanics 49, 136 – 148 (2019).
- [7] Boccaletti, S., Latora, Moreno, V. Y., Chavez, M., Hwang, D. U.. Complex networks: Structure and dynamics. Physics Reports, 424, 175 – 308 (2006).
- Boeck, T., Vitanov, N.K.: Low-dimensional chaos in zero-Prandtlnumber Benard - Marangoni convection. Physical Review E 65, 037203 (2002).
- [9] Borisov, R., Vitanov, N. K.: Human migration: Model of a migration channel with a secondary and a tertiary arm. AIP Conference Proceedings 2075, 150001 (2019).
- [10] Chan, W. -K.: Theory of Nets: Flows in Networks. Wiley, New York (1990).
- [11] Chen, G., Wang, X., Li, X.: Fundamentals of complex networks. Wiley, Singapore (2015).
- [12] Dimitrova, Z.: Fluctuations and dynamics of the chaotic attractor connected to an instability in heating from below fluid layer. Comptes Rendus de lAcademie bulgare des Sciences 60, No. 10, 1065 – 1071 (2007)

- [13] Dimitrova, Z.: On traveling waves in lattices: The case of Riccati lattices. Journal of Theoretical and Applied Mechanics 42, No.3, 3 – 22 (2012).
- [14] Dimitrova, Z.I., Vitanov, N.K.: Influence of adaptation on the nonlinear dynamics of a system of competing populations. Physics Letters A 272, 368 – 380 (2000).
- [15] Dimitrova, Z.I., Vitanov, N.K.: Dynamical consequences of adaptation of the growth rates in a system of three competing populations. Journal of Physics A: Mathematical and General 34, 7459 – 7473 (2001)
- [16] Dimitrova, Z.I., Vitanov, N.K.: Adaptation and its impact on the dynamics of a system of three competing populations. Physica A 300, 91 - 115 (2001)
- [17] Dimitrova, Z.I., Vitanov, N.K.: Chaotic pairwise competition. Theoretical Population Biology 66, 1 – 12 (2004)
- [18] Dorogovtsev, S. N., Mendes, J. F. F. Evolution of networks. Advances in Physics, 51, 1079 – 1187 (2002).
- [19] Estrada, E. The structure of complex networks.Oxford Unversity Press, Oxford, UK (2011).
- [20] Fawcet, J. T.: Networks, linkages, and migration systems. International Migration Review 23, 671 – 680 (1989).
- [21] Ford Jr., L. D., Fulkerson, D. R.: Flows in Networks. Princeton University Press, Princeton, NJ (1962).
- [22] Fu, X., Small, M., Chen, G. Propagation dynamics on complex networks. Wiley, Chichester, UK (2014).
- [23] Gartner, N. H., Imrota, G. (Eds.) Urban Traffic Networks. Dynamic Flow Modeling and Control. Springer, Berlin (1995).
- [24] Harris, J. R., Todaro, M. O.: Migration, unemployment and development: A two-sector analysis. The American Economic Review, 60, 126 – 142 (1970).
- [25] Kantz, H., Holstein, D., Ragwitz, M., Vitanov, N.K.: Markov chain model for turbulent wind speed data. Physica A 342, 315 – 321 (2004).

- [26] May, R.M., Leonard, W.J. Nonlinear aspects of competition between three species. SIAM Journal of Applied Mathematics 29, 243 – 253 (1975).
- [27] Marsan, G. A., Bellomo, N., Tosin, A.: Complex Systems and Society: Modeling and Simulation. Springer, New York (2013).
- [28] Martinov, N.K., Vitanov, N.K.: New class of running-wave solutions of the (2+ 1)-dimensional sine-Gordon equation. Journal of Physics A:Mathematical and General 27, 4611 – 4617 (1994)
- [29] Nikolova, E.V.: On nonlinear waves in a blood-filled artery with an aneurysm. AIP Conference Proceedings 1978, 470050 (2018).
- [30] Nikolova, E.V., Serbezov, D.Z., Jordanov, I.P.: On the spatio-temporal dynamics of interacting economic agents: Application of the modified method of simplest equation. AIP Conference Proceedings 2075, 150003 (2019)
- [31] Panchev, S., Spassova, T., Vitanov, N.K.: Analytical and numerical investigation of two families of Lorenz-like dynamical systems. Chaos, Solitons & Fractals 33, 1658 – 1671 (2007).
- [32] Pastor-Sattoras, Vespignani, R. A.: Epidemic dynamics and endemic states in complex networks. Physical Review E **63**, 066117 (2001).
- [33] Scharnhorst, A., Borner, K., Van den Besselaar, P. (Eds.). Models for Science Dynamics. Springer, Berlin, (2012)
- [34] Silva, T. C., Zhao, L. Machine learning in complex networks. Springer, Cham (2016).
- [35] Schubert, A., Glänzel, W.: A dynamic look at a class of skew distributions. A model with scientometric application. Scientometrics 6, 149 – 167 (1984).
- [36] Vitanov, N.K.: On travelling waves and double-periodic structures in two-dimensional sine-Gordon systems. Journal of Physics A: Mathematical and General 29, 5195 – 5207 (1996)
- [37] Vitanov, N.K., Sakai, K.: Upper bounds on the number of significant degrees of freedom of noise influenced oscillations of moving machines. Systems Analysis Modelling Simulation 43, 815 – 828 (2003)

- [38] Vitanov, N.K., Dimitrova, Z.I., Kantz, H.: On the trap of extinction and its elimination. Physics Letters A 349, 350 – 355 (2006)
- [39] Vitanov, N.K., Jordanov, I.P., Dimitrova, Z.I.: On nonlinear dynamics of interacting populations: Coupled kink waves in a system of two populations. Communications in Nonlinear Science and Numerical Simulation 14, 2379 – 2388 (2009)
- [40] Vitanov, N.K., Jordanov, I.P., Dimitrova, Z.I.: On nonlinear population waves. Applied Mathematics and Computation 215, 2950 - 2964 (2009)
- [41] Vitanov, N.K.: Application of simplest equations of Bernoulli and Riccati kind for obtaining exact traveling-wave solutions for a class of PDEs with polynomial nonlinearity. Communications in Nonlinear Science and Numerical Simulation 15, 2050 – 2060 (2010)
- [42] Vitanov, N.K.: Modified method of simplest equation: powerful tool for obtaining exact and approximate traveling-wave solutions of nonlinear PDEs. Communications in Nonlinear Science and Numerical Simulation 16, 1176 - 1185 (2011)
- [43] Vitanov, N.K.: On modified method of simplest equation for obtaining exact and approximate solutions of nonlinear PDEs: the role of the simplest equation. Communications in Nonlinear Science and Numerical Simulation 16, 4215 - 4231 (2011).
- [44] Vitanov, N.K.: Science Dynamics and Research Production. Indicators, Indexes, Statistical Laws and Mathematical Models, Springer, Cham (2016).
- [45] Vitanov, N. K., Dimitrova, Z. I., Ausloos, M.: Verhulst-Lotka-Volterra model of ideological struggle. *Physica A* 389, 4970 – 4980 (2010).
- [46] Vitanov, N. K., Ausloos, M., Rotundo, G.. Discrete model of ideological struggle accounting for migration. Advances in Complex Systems, 15 (supp01), 1250049 (2012).
- [47] Vitanov, N. K., Dimitrova, Z. I., Vitanov, K. N.: Traveling waves and statistical distributions connected to systems of interacting populations. Computers & Mathematics with Applications 66, 1666 – 1684 (2013).
- [48] Vitanov, N. K., Vitanov, K. N.: Box model of migration channels. Mathematical Social Sciences 80, 108 – 114 (2016).

- [49] Vitanov, N. K., Vitanov, K. N.: Discrete-time model for a motion of substance in a channel of a network with application to channels of human migration. Physica A 509, 635 – 650 (2018).
- [50] Vitanov, N. K., Vitanov, K. N.: On the motion of substance in a channel of a network and human migration. Physica A **490**, 1277 1294 (2018).
- [51] Vitanov, N. K., Vitanov, K. N.: Statistical distributions connected to motion of substance in a channel of a network. Physica A 527, 121174 (2019).
- [52] Vitanov, N. K., Vitanov, K. N., Ivanova, T.: Box model of migration in channels of migration networks. In K. Georgiev, Todorov, M., Georgiev, I. (eds.), Advanced Computing in Industrial Mathematics, Studies in Computational Intelligence, **728**, 203 – 215 (2017).
- [53] Vitanov, N. K., Borisov, R: A model of a motion of substance in a channel of a network. Journal of Theoretical and Applied Mechanics 48, No. 3, 74 – 84 (2018).
- [54] Wu, J., Wang, Y. Opportuistic mobile social networks. CRC Press, Boca Raton, FL (2015).