

RUELLE ZETA FUNCTION FROM FIELD THEORY

CHARLES HADFIELD, SANTOSH KANDEL, AND MICHELE SCHIAVINA

ABSTRACT. We propose a field-theoretic interpretation of Ruelle zeta function, and show how it can be seen as the partition function for BF theory when an unusual gauge fixing condition on contact manifolds is imposed. This suggests an alternative rephrasing of a conjecture due to Fried on the equivalence between Ruelle zeta function and analytic torsion, in terms of homotopies of Lagrangian submanifolds.

CONTENTS

Introduction	2
What to expect from this paper	3
1. Lagrangian field theory, the Batalin–Vilkovisky formalism and regularised determinants	5
1.1. Classical field theory, symmetries and quantisation	5
1.2. Cohomological approach and the BV complex	6
1.3. Partition functions	8
2. Geometric Setting	10
2.1. Flat vector bundle	10
2.2. Riemannian structure and analytic torsion	10
2.3. Contact structure	12
2.4. Anosov dynamics	12
3. Ruelle Zeta function	13
3.1. Differential forms decomposition	14
3.2. Ruelle zeta function as regularized determinant	15
3.3. Meromorphic extension of the resolvent	16
4. BF theory on contact manifolds	17
4.1. Analytic torsion from resolutions of de Rham differential	18
4.2. BV interpretation: metric gauge for BF theory	20
4.3. Contact gauge fixing for BF theory	22
5. Lagrangian homotopies, Fried’s conjecture and gauge-fixing independence	23
5.1. A sphere bundle construction	24

Acknowledgements	27
References	27

INTRODUCTION

Quantum field theory is a useful tool in many areas of pure and applied mathematics. It provides a number of precise answers, often involving insight coming from statements that are theorems in finite dimensions, and that need to be appropriately checked and generalised in infinite dimensions.

A positive example of this is the interpretation by Schwarz of the Ray–Singer analytic torsion in terms of a partition function for a degenerate functional [Sch78, Sch79]. The main ingredient in Schwarz’s construction is a topological field theory involving differential forms, which enjoys a symmetry given by the shift of closed forms by exact ones [BBRT91, CCRFM95, CR01]. This is known nowadays with the name of BF theory.

From a field-theoretic point of view, such symmetry needs to be removed, or gauge fixed, as it represents a fundamental redundancy in the description. One possible way to do this is by choosing a reference metric g and enforcing a g -dependent condition on fields¹. It allows to compute the partition function of the theory - the starting point for quantum considerations on the system - and one is left to show that the choice of metric is immaterial. The proof that such choice of metric is irrelevant was given by Schwarz for the partition function of abelian BF theory, and it is tantamount to the statement of independence of the analytic torsion on the metric used to define a Laplacian on the underlying manifold.

There are several ways of encoding a choice of gauge fixing within the framework of field theory, starting from the original idea of Faddeev and Popov [FP16], later understood in terms of Lie algebra cohomology by Becchi, Rouet, Stora and Tyutin [BRS74, BRS75, BRS76, Tyu75]. A more general approach follows the ideas of by Batalin and Vilkovisky [BV83, BV84], and implements the choice of a gauge as the choice of a Lagrangian submanifold in an appropriate (graded)-symplectic manifold of fields \mathcal{F} . In this context, gauge-fixing independence is phrased in terms of isotopies of embedded Lagrangian submanifolds, and needs to be proven in some appropriate regularisations scheme. In finite dimensions this is a theorem: the partition function for an action functional S that satisfies the quantum master equation (a differential condition on S) does not depend on the choice of a particular Lagrangian submanifold inside a smooth family $\mathbb{L}_t \subset \mathcal{F}$. Observe that this statement can be phrased as local constancy of the partition function w.r.t a parametrisation of the Lagrangian homotopy.

Leaving field theory aside for a moment, consider a closed manifold M endowed with an Anosov vector field (Definition 15). The flow associated with the vector field is a typical example of a dynamical system displaying “hard chaos” [Gut91]. An important example of such a dynamical system is obtained from a Riemannian manifold (Σ, g) whose sectional

¹This is often called Lorenz gauge fixing.

curvature is negative; then $M = S_g^*\Sigma$, the unit-cotangent bundle of Σ (a sphere bundle), is such that the Reeb vector field X associated with the natural contact structure is an Anosov vector field and its flow coincides with the geodesic flow.

If an Anosov flow (generated by the vector field X) admits closed orbits, one defines a (dynamical) Ruelle zeta function $\zeta_X(\lambda)$ to count lengths of closed orbits associated with the flow in a similar spirit to how the Riemann zeta function counts prime numbers [Rue76, Rue86]. The zeta function also may be defined in the presence of a representation ρ of $\pi_1(M)$ which provides a flat vector bundle over M , this leads to a twisted zeta function. The chaotic nature of the dynamical system ensures that the zeta function is well-defined for $\text{Re}(\lambda) \gg 1$, however work is required to show that the function extends meromorphically to the whole complex plane [GLP13]. Conjecture 21, due to Fried [Fri86], proposes that when $M = S_g^*\Sigma$, the Ruelle zeta function (evaluated at zero) exactly computes the analytic torsion of the associated sphere bundle. To connect this to field theory, we observe that this means that Ruelle zeta function is expected to compute - in Schwarz's terms - the partition function of BF theory in a given (metric dependent) gauge fixing.

Fried's conjecture has received considerable attention recently. The proposed equality was confirmed in [Fri86] for Σ a hyperbolic manifold, and conjectured in [Fri87] that it also holds for compact locally symmetric spaces with non positive curvature. Conjecture 21, as we state it, appears in [Fri95]. A more precise version for locally symmetric manifolds has been proved in [She17] following [MS91]. In the variable curvature case a perturbative result has been obtained in [DGRS20] and extended in [CD19]. A surprising result in the case of surfaces with variable negative curvature, but without reference to an acyclic representation, showed the zeta function at zero is determined by the topology of the surface [DZ17]. This has been extended to the case of surfaces with boundary [Had18] and to higher dimensional closed manifolds perturbatively close to hyperbolic space [KW20]. However Fried's conjecture, along with its three star bounty [Zwo17, Section 3, footnote 6], remains open.

What to expect from this paper. We present a new class of gauge fixings for BF theory on contact manifolds based on the Reeb vector field associated with the contact structure; we call this the contact gauge in Definition 34. We then go on to show that, on sphere bundles with an Anosov–Reeb vector field, the Ruelle zeta function can be interpreted as an appropriately regularised determinant for the Lie derivative operator \mathcal{L}_X on k -forms in the kernel of the contraction ι_X . Taking this regularised determinant as the definition for the partition function of BF theory in the contact gauge allows us to conclude that this coincides with the Ruelle zeta function. This point of view is analogous to Schwarz's calculation of the partition function of BF theory (in the metric gauge), whose output is the analytic torsion, and to the recent proof of Chern–Gauss–Bonnet Theorem that has been given with similar techniques in [BE15] (see also [CMR95]).

As a consequence, we relate the expected gauge-fixing independence of the partition function of BF theory to Fried's conjecture; in particular, we show how modern proofs of the conjecture for certain classes of manifolds (see [DGRS20]) can be taken as a proof of gauge-fixing independence. On the other hand, we believe that the field theoretic presentation of

the Ruelle zeta function provided in this paper will allow the problem to be tackled from a different angle: by means of homotopies of Lagrangian submanifolds.

To this aim, we setup a convenient construction to compare Anosov vector fields that are related to a choice of a metric on a base manifold Σ (Section 5). By means of a natural construction for sphere bundles, we map smooth paths of metrics into smooth paths of Anosov vector fields, effectively constructing an isotopy between their associated Lagrangian submanifolds. This, together with the crucial local-constancy results of [DGRS20], allow us to test our approach to the known case of 2d surfaces — Theorem 47 provides an alternative proof of Fried’s conjecture on surfaces — and interprets it as gauge-fixing independence for BF theory.

From the point of view of algebraic topology, this result suggests that, under certain assumptions, ι_X can be made into a chain contraction for the de Rham complex, namely one can construct $\eta_X = (\mathcal{L}_X)^{-1}\iota_X$ with the appropriate conditions of nondegeneracy of \mathcal{L}_X . This interpretation appears to be related with the notion of a dynamical torsion introduced in [CD19]. There appears to be a sweet spot at the intersection of Anosov and Reeb vector fields where the independence of the “torsion” of the de Rham complex on the choice of a chain contraction, and independence of the partition function of BF theory on a choice of gauge fixing appear to be aspects of the same statement, expressed by Fried’s conjecture.

This work is mostly addressed to the mathematical physics community working with or closely related to field theory in the Batalin–Vilkovisky formalism, but it is also aimed at the community interested in the microlocal analysis of Anosov/geodesic flows and Fried’s conjecture. Therefore, we will present some basic background on field theory with symmetries to set the stage, terminology and expectations, but we will not present a complete treatment of the mathematics behind it. Results and constructions that will be somewhat assumed in this exposition of field theory can be found, e.g., in [And92, DF99, CMR14, Del17].

Our main goal is to present a novel link between field theory and geometric and microlocal analysis, that will hopefully allow to import techniques across research fields, and stimulate fruitful interaction between scientific communities.

Section 1 is an overview on Lagrangian field theory aimed at introducing the Batalin–Vilkovisky formalism and the problem of gauge fixing. It sets the stage for the field-theoretic interpretations that will follow.

Section 2 establishes the geometric conventions and notations, clarifying what incremental/alternative data one needs at different stages, and briefly describes the analytic torsion and Anosov dynamics.

In Section 3 we introduce Ruelle zeta function and its k -form decomposition, and state Fried’s conjecture. We interpret the zeta function as a regularised (super)determinant.

In Section 4 we describe a field theory called BF theory, we summarise the famous interpretation (due to Schwarz) of the analytic torsion in terms of the partition function of BF theory, and introduce a new gauge fixing condition on contact manifolds. We show how, with that gauge-fixing, the partition function of BF theory computes the Ruelle zeta function of the associated geodesic/Anosov flow.

Finally, in Section 5 we interpret Fried's conjecture in terms of gauge-fixing independence of BF theory in the BV formalism, and suggest a construction for sphere bundles that allows to present explicit homotopies between Lagrangian submanifolds.

1. LAGRANGIAN FIELD THEORY, THE BATALIN–VILKOVISKY FORMALISM AND REGULARISED DETERMINANTS

In this section we will review the basics of the Batalin–Vilkovisky (BV) formalism [BV83, BV84] for Lagrangian field theories and how it handles gauge fixing. We will use a particular kind of regularisation based on the notion of flat traces to define determinants of operators and partition functions of quadratic functionals.

1.1. Classical field theory, symmetries and quantisation. The standard framework for Lagrangian field theories is as follows. To a compact manifold M , possibly endowed with extra geometric data, like a Riemannian metric or a contact structure, we associate a space of classical fields F_M , which is usually modelled on the space of sections of some vector bundle² $E \rightarrow M$, together with a local functional S_M , called action functional. Local here means that it has the form of an integral over M of a density-valued functional of the fields and a finite number of jets³:

$$S_M = \int_M L_M[\phi, \partial^I \phi], \quad (1)$$

where I is a finite multi-index and L_M is called Lagrangian density. For simplicity we will consider compact manifolds without boundary, although it is possible to adapt the construction to non-compact ones or manifolds with boundary (see e.g. [FR13, CMR14]).

The dynamical content of the theory is encoded in the Euler–Lagrange locus $EL[S_M]$, the space of solution of the Euler–Lagrange equations coming from the variational problem for S_M . In other words, the Euler–Lagrange locus is the set of critical points of S_M .

The action functional might enjoy a symmetry. That is, it might be invariant under some transformation of the fields, for example when considering Lie algebra actions on fields taking values in Lie algebra modules. Symmetries are usually described by a (smooth) distribution $D_M \subset TF_M$, and they make the critical points of the action functional degenerate⁴ and will become an issue when dealing with perturbative quantisation of the theory (see below). In what follows we will only consider symmetry distributions that are involutive.

Quantisation, loosely speaking, is meant to replace the (commutative) algebra of functions over the space of physical configurations of the system with some (noncommutative) algebra of operators over a suitable vector space, also called the space of quantum states. Without delving too much into how this is achieved in general, for our purposes it will be important

²More generally, a sheaf.

³There is an equivalent formulation of this in the variational bi-complex [And92, DF99], where the full jet bundle is taken into account.

⁴We will only be concerned with continuous symmetries.

to mention that one possible procedure starts by making sense of the following expression:

$$Z = \int \exp\left(\frac{i}{\hbar} S_M\right), \quad (2)$$

usually called the partition function, where the integral sign should ideally represent actual integration over F_M , with some measure. However, an appropriate integration theory for such (infinite dimensional) spaces of fields is generally not available, and one defines the previous expression as a formal power-series expansion in the parameter \hbar . This approach, however, requires the critical points of S_M to be isolated, as it involves a saddle point or stationary phase approximation around critical points. It therefore automatically fails in the presence of symmetries, unless appropriate prescriptions are enforced.

We choose to deal with this problem by means of the Batalin–Vilkovisky (BV) formalism.

1.2. Cohomological approach and the BV complex. Degenerate functionals are usually accompanied by involutive (symmetry) distributions D_M , and the space of inequivalent field configurations is the quotient $EL[S_M]/D_M$. Most of the times the quotient is singular and one looks for a replacement for it.

A resolution of $EL[S_M]/D_M$ is given by a complex (C^\bullet, d_C) such that⁵ for all $i > 0$

$$H^{-i}(C^\bullet) = 0, \quad H^0(C^\bullet) \simeq C^\infty(EL[S_M]/D_M). \quad (3)$$

One way to obtain a resolution is by first localising to the submanifold $EL[S_M]$ constructing the Koszul–Tate complex, and then following the Chevalley–Eilenberg procedure to describe D_M -invariant functions on it (see [Sta98] for a “geometric” jet-bundle explanation of this and [FR13] for a more “algebraic” one).

The Batalin–Vilkovisky formalism is essentially the interpretation of said complex as the space of functions over a (-1) -symplectic graded manifold $(\mathcal{F}_{BV}, \Omega_{BV})$ [Hen90, Sta98, CS11, CMR14, CMR18], whose degree-0 part coincides with the original space of fields F_M , endowed with an odd vector field of degree-1 $Q \in C^\infty(\mathcal{F}_{BV}, T[1]\mathcal{F}_{BV})$ such that⁶ $[Q, Q] = 0$ and a degree-0 functional $\mathcal{S}_M: \mathcal{F}_{BV} \rightarrow \mathbb{R}$ satisfying

$$\iota_Q \iota_Q \Omega_{BV} = \{\mathcal{S}_M, \mathcal{S}_M\}_{\Omega_{BV}} = 0, \quad (4)$$

with $\{\cdot, \cdot\}_{\Omega_{BV}}$ the Poisson bracket associated with the symplectic structure Ω_{BV} , and the compatibility condition

$$\iota_Q \Omega_{BV} = d\mathcal{S}_M. \quad (5)$$

Remark 1. In infinite dimensions one models \mathcal{F}_{BV} on some appropriate space of sections of the jet bundle of a vector bundle $E \rightarrow M$. Then, the de Rham differential in Equation (5) is replaced with δ , the variation operator interpreted as the vertical differential on local functionals over \mathcal{F}_{BV} (see [And92]). Observe that we could take Equation (4) as a definition of Poisson brackets in infinite dimensions.

⁵In some practical cases the vanishing of the negative cohomology is not guaranteed. We will anyway not need this condition in what follows.

⁶ Q is essentially the derivation d_C interpreted as a vector field.

On $C^\infty(\mathcal{F}_{BV})$ one constructs another (second order) differential ($\Delta_{BV}^2 = 0$) called BV-Laplacian and defines gauge fixing to be the choice of a Lagrangian submanifold $\mathbb{L} \subset \mathcal{F}_{BV}$. The main results are as follows:

Theorem 2. *Let $(\mathcal{F}_{BV}, \Omega_{BV})$ be a finite dimensional (-1) -symplectic graded manifold, with a measure μ and the BV Laplacian Δ_μ , a coboundary operator defined on $C^\infty(\mathcal{F}_{BV})$ such that for all $f \in C^\infty(\mathcal{F}_{BV})$ we have*

$$\Delta_\mu f = -\frac{1}{2} \operatorname{div}_\mu(X_f) \quad (6)$$

with X_f the Hamiltonian vector field of f with respect to Ω_{BV} . Assuming that $\Delta_{\mu_{BV}} f = 0$ and $g = \Delta_{\mu_{BV}} h$, with $f, g, h \in C^\infty(\mathcal{F}_{BV})$ then:

- for any Lagrangian submanifold $\mathbb{L} \subset \mathcal{F}_{BV}$

$$\int_{\mathbb{L}} g \mu_{BV}|_{\mathbb{L}} = 0; \quad (7)$$

- given a continuous family of Lagrangian submanifolds \mathbb{L}_t

$$\frac{d}{dt} \int_{\mathbb{L}_t} f \mu_{BV}|_{\mathbb{L}_t} = 0. \quad (8)$$

Remark 3. The definition of the BV Laplacian in (6) implies the relations

$$\Delta_\mu(fg) = (\Delta_\mu f)g + (-1)^{|f|} f(\Delta_\mu g) + (-1)^{|f|} \{f, g\}_{\Omega_{BV}} \quad (9)$$

$$\Delta_\mu \{f, g\}_{\Omega_{BV}} = \{\Delta_\mu f, g\}_{\Omega_{BV}} + (-1)^{|f|+1} \{f, \Delta_\mu g\}_{\Omega_{BV}}, \quad (10)$$

making the tuple $(C^\infty(\mathcal{F}_{BV}), \cdot, \{\cdot, \cdot\}_{\Omega_{BV}}, \Delta_\mu)$ into a BV algebra [CFL06].

Remark 4. Theorem 2 is stated for finite dimensional manifolds. In this case Δ_μ always exists. On infinite dimensional manifolds a number of complications arise. One needs an appropriate regularisation of Δ_μ , and the corresponding adaptations of statements in Theorem 2 must be checked. In this paper we are interested in abelian BF theory, which is a non-interacting topological field theory whose partition function (cf. Equation (2)) is expected to be independent of the gauge fixing, and has been computed to be the Reidemeister (or equivalently) Analytic torsion of the de Rham complex [Sch78, Sch79]. More recent work generalised the BV theorem for certain classes of field theories (and gauge fixings) [Cos11, CG16], while a perturbative approach has been shown to work for the theory at hand in the presence of boundaries [CMR18], and with a regularisation coming from cellular decompositions in [CMR20].

Remark 5. Notice that the notion of Lagrangian submanifold has to be appropriately adapted in infinite dimensions and when dealing with \mathbb{Z} -grading. The Lagrangian submanifolds L we will consider in this paper are such that, locally, the symplectic space looks like $L \oplus K$, with the symplectic form given by a nondegenerate pairing between L and K . This notion of a Lagrangian submanifold coincides with the one used in [CC18]. Often L can be seen as the vanishing locus of a Poisson subalgebra \mathcal{I} of the Poisson algebra of functions on \mathcal{F}_{BF} , which

is also isotropic, i.e. $\Omega_{BF}|_L = 0$. This means that L is isotropic and coisotropic⁷. For a more in depth analysis of Lagrangian submanifolds in infinite dimensions and the symplectic category we refer to [Wei10], building on [Wei71].

Remark 6. For concreteness, in what follows we will discuss field theories where fields are given by differential forms on a manifold. If needed, one can think of the space of fields as a Fréchet vector space, but indeed this specification will not be necessary for our purposes.

1.3. Partition functions. If we look at the quadratic part of the action functional, we can interpret the partition function as a (formal) Gaussian integral. Assume from now on that the action functional is at most quadratic.

In finite dimensions the result of said integral would be the determinant of the operator featured in the action functional S_M . In infinite dimensions, this requires defining an appropriate regularisation of determinants.

The standard approach to partition functions for degenerate quadratic functionals follows from Schwarz [Sch78, Sch79], where the resolution of (the kernel of) an elliptic differential operator on a closed Riemannian manifold is presented, which outputs a (co-)chain complex, and the partition function is given in terms of products of (regularised) determinants of operators associated with the resolving complex. The explicit example for BF theory is given in Section 4.1. In short, the insights from Schwarz allow us to interpret partition functions of quadratic degenerate functionals as (a product of) regularised determinants, provided a suitable resolution can be found such that the regularised determinants of the associated operators exist.

For the purposes of this paper, instead of the standard zeta-function regularisation, we will use the notion of a flat-determinant, based on flat-traces, inspired by Atiyah and Bott's constructions [AB64, AB68]. Details on the definition of flat traces and determinant can be found in several places: [BV84, Definition 3.12], a microlocal version in [DZ16, Section 2.4], and a mollifier approach in [Bal18, Section 3.2.2]. Since we will not be concerned with the microlocal analysis of operators, we will avoid discussing the necessary tools to define flat traces, and will only work up to the requirements that they exist for the operators we will consider. In this spirit, we give the following definition.

Definition 7. *Let $A: V \rightarrow V$ be an operator on an appropriate inner product space, such that the flat trace $\text{tr}^b(\exp(-t(A + \lambda)))$ exists for $\lambda \in \mathbb{C}$. We define the flat determinant of $A + \lambda$ to be*

$$\log \det^b(A + \lambda) := - \left. \frac{d}{ds} \right|_{s=0} \left[\frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \text{tr}^b(\exp(-t(A + \lambda)) - \Pi_\lambda) dt \right] \quad (11)$$

where Π_λ is the spectral projector on the kernel of $(A + \lambda)$, whenever the integrals converge.

Remark 8. Observe that if A is such that $e^{-t(A+\lambda)}$ is trace-class, then $\text{tr}(e^{-t(A+\lambda)}) = \text{tr}^b(e^{-t(A+\lambda)})$. As a consequence, if the zeta-regularized determinant of $A + \lambda$ exists, it

⁷This notion coincides with requiring L to be maximal isotropic.

coincides with the flat-determinant: $\det^b(A + \lambda) = \det(A + \lambda)$. See e.g. [Bal18, Proposition 6.8] for details.

Let $A = \text{diag}(B, C)$ be a graded, degree-preserving (block-diagonal) linear map on a finite dimensional graded vector space. A graded Gaussian integral for $\exp(-\langle y, Ax \rangle)$ returns $\text{sdet}(A)^{-1} = \frac{\det(C)}{\det(B)}$. More generally, if A_k is the k -th component of A acting on a graded space with a finite number of nonzero components, each A_k acting on vectors of degree k , we get⁸

$$\text{sdet}(A) = \prod_{k=0}^n \det(A)^{(-1)^k}.$$

For an introduction to Berezinians and odd integration see, for example, [Mne19, Section 3.8] and [Vor91], while the original notion was introduced in [Ber83, BL75].

In infinite dimensions, to an operator A on a graded space we can associate a regularised superdeterminant in the same way, but replacing the determinants on the block operators with their flat-regularised versions. We will use this notion to define the partition function of a degenerate quadratic functional, as follows:

Definition 9. Let $S: V \rightarrow \mathbb{R}$ of the form $S = \frac{1}{2} \int_M(x, Ax)$ for some operator $A: V \rightarrow V$, with V a (possibly graded) vector space endowed with an inner product (\cdot, \cdot) . We define the partition function Z of S to be the square root of the flat (super)-determinant of the (graded) operator A :

$$Z(S) = \int_V e^{iS_M} := |\text{sdet}^b(A)|^{-\frac{1}{2}}. \quad (12)$$

When the field theory has symmetries, we assume that V is further endowed with a (-1) symplectic form. A choice of a Lagrangian submanifold $\mathbb{L} \subset V$ will be called gauge-fixing, and we define the partition function of S in the gauge-fixing \mathbb{L} to be

$$Z(S, \mathbb{L}) := Z(S|_{\mathbb{L}}). \quad (13)$$

Remark 10. To obtain a true generalisation of finite dimensional Gaussian integrals one should append to formula (12) the phase factor $\exp(-i\frac{\pi}{4}\text{sg}(A))$ where $\text{sg}(A)$ is the signature of the operator A , appropriately regularised. Since in what follows we will not discuss the phase of partition functions, we will omit this term from the definition.

Remark 11. Observe that often one encounters the situation in which $S_M = \int_M(y, Bx)$ for some operator B and $x, y \in V'$ for some space V' . Define⁹ $V = V' \oplus V'$ and

$$A = \begin{pmatrix} 0 & B \\ B^t & 0 \end{pmatrix}$$

so that $(y, Bx) = \frac{1}{2}(z, Az)$ with $z = (x, y) \in V$. Then $Z = |\text{sdet}^b(A)|^{-\frac{1}{2}} = |\text{sdet}^b(B)|^{-1}$.

⁸Observe that we are considering parities modulo 2. In principle a graded determinant would return $\prod_{k=0}^n \det(A)^{k(-1)^k}$. We will not make such a distinction in what follows.

⁹Note that the V' components in V are not considered to have different degrees, i.e. A is an even matrix of graded operators.

Remark 12. Let V be a graded vector space and $V[k]$ its k -shift, so that $(V[k])^i := V^{i+k}$. In particular, if $z \in V[1]$ is a homogeneous element of degree k it will be parametrised by homogeneous elements in V of opposite parity. In particular, if A is a graded linear map on V , the Gaussian integral

$$\int_{V[1]} e^{-\frac{1}{2}\langle z, Az \rangle} := |\text{sdet}^b(A)|^{\frac{1}{2}}. \quad (14)$$

2. GEOMETRIC SETTING

This section establishes the geometry and notations. In the following three sections we progressively introduce more structure to our initial setup of a flat vector bundle over a manifold, whose twisted cohomology is trivial. The plainest setting involves a differentiable manifold endowed with a flat vector bundle. On top of that we consider the introduction of either a Riemannian structure or a contact structure. The intersection of the two will require the base manifold to display Anosov dynamics.

It is useful to distinguish between these geometric settings as when we will only need certain geometric properties when discussing different field theories. This distinction between geometric data with which a differentiable manifold is endowed reflects the practice of complementing topological theories with additional geometric structures (e.g. Riemannian or contact) for the sake of gauge fixing.

2.1. Flat vector bundle. Let M be an N -dimensional compact manifold without boundary which is oriented and connected. Let $\rho : \pi_1(M) \rightarrow U(\mathbb{C}^r)$ denote a unitary representation. This representation endows M with a Hermitian vector bundle (E, h) of rank r with flat connection ∇ . We collectively denote this data by (M, E) .

Let $\Omega^\bullet(M; E)$ denote the space of (smooth) differential forms on M taking values in E , and let

$$d_\nabla \equiv d_k : \Omega^k(M, E) \rightarrow \Omega^{k+1}(M, E) \quad (15)$$

be two notations for the twisted de Rham differential. We denote by $H^\bullet(M; E)$ the cohomology associated to the twisted de Rham complex $(\Omega^\bullet(M; E), d_\nabla)$, with Betti numbers $\beta_k := \dim H^k(M; E)$.

From now on we will assume that (M, E) is such that its twisted de Rham complex is acyclic, i.e. $\beta_k = 0$ for all $0 \leq k \leq N$.

2.2. Riemannian structure and analytic torsion. Suppose (M, E) is supplemented with a Riemannian metric g_M . We collectively denote this data (M, E, g_M) . Let $d\text{Vol}$ denote the associated volume form and let $\langle \cdot, \cdot \rangle$ be the inherited inner product on $\bigwedge^\bullet T^*M$. The Hodge star $\star : \bigwedge^k T^*M \rightarrow \bigwedge^{N-k} T^*M$ is defined through

$$u \wedge \star v = \langle u, v \rangle d\text{Vol} \quad (16)$$

for $u, v \in \bigwedge^k T^*M$. This lifts to E -valued forms by identifying E with its dual via the Hermitian metric h . An inner product is then placed on $\Omega^\bullet(M; E)$ by declaring

$$(u, v) = \int_M [u \wedge \star v]^{\text{top}} \quad (17)$$

for $u, v \in \Omega^\bullet(M; E)$ with top refers to only taking the top-form (degree N) part of $u \wedge \star v$.

The de Rham differential has an adjoint

$$d_{\nabla}^* \equiv d_k^*: \Omega^{k+1}(M, E) \rightarrow \Omega^k(M, E) \quad (18)$$

which provides the twisted Laplace–de Rham operator (henceforth simply Laplacian)

$$\Delta_k := (d_{\nabla}^* + d_{\nabla})^2 : \Omega^k(M; E) \rightarrow \Omega^k(M; E). \quad (19)$$

When acting on $L^2(M; \bigwedge^k T^*M \otimes E)$ the Laplacian has nonnegative eigenvalues $\lambda_n \geq 0$. Denote by Π_λ the L^2 spectral projector onto the kernel of $\Delta_k + \lambda$ and consider the function

$$F_{\Delta_k}(\lambda, s) := \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} \text{tr}^b \left(e^{-t(\Delta_k + \lambda)} - \Pi_\lambda \right) dt. \quad (20)$$

The (flat) regularised determinant of the operator $\Delta_k + \lambda$ is

$$\log \det^b(\Delta_k + \lambda) = - \left. \frac{\partial}{\partial s} \right|_{s=0} F_{\Delta_k}(\lambda, s). \quad (21)$$

We also write $f_{\Delta_k}(s) := F_{\Delta_k}(0, s)$ and observe that $f_{\Delta_k}(s) = \sum_{\lambda_j > 0} \lambda_j^{-s}$.

Definition 13. *The analytic torsion of M [RS71] is defined to be*

$$\tau_\rho(M) := \prod_{k=1}^N \det^b(\Delta_k)^{\frac{k}{2}(-1)^{k+1}}. \quad (22)$$

Alternatively, we can write

$$2 \log \tau_\rho(M) = \sum_{k=1}^N (-1)^k k \left. \frac{d}{ds} \right|_{s=0} f_{\Delta_k}(s). \quad (23)$$

The Hodge decomposition provides an orthogonal decomposition of $L^2(M; \bigwedge^\bullet T^*M \otimes E)$ into exact and coexact forms (no harmonic forms are present due to acyclicity). Introducing $d_k^* d_k := \Delta_k|_{\text{coexact}}$ we can also write the analytic torsion as [Sch79] (see also [Mne14, CMR20])

$$\tau_\rho(M) = \prod_{k=0}^{N-1} \det^b(d_k^* d_k)^{\frac{1}{2}(-1)^k}, \quad (24)$$

which alternatively reads

$$2 \log \tau_\rho(M) = \sum_{k=0}^{N-1} (-1)^{k+1} \left. \frac{d}{ds} \right|_{s=0} f_{d_k^* d_k}(s). \quad (25)$$

2.3. Contact structure. Suppose that (M, E) is supplemented with a contact form $\alpha \in \Omega^1(M)$, and $\dim(M) = N = 2n + 1$. We will denote $d\text{Vol} = \alpha \wedge (d\alpha)^n$. Let X be the associated Reeb vector field, defined by the relations

$$\iota_X \alpha = 1, \quad \iota_X d\alpha = 0. \quad (26)$$

We collectively denote this data (M, E, X) .

Denote by T_0^*M the $2n$ -rank subbundle of T^*M defined as the conormal of X , so that pointwise $T^*M = \mathbb{R}\alpha + T_0^*M$. Transferring this to the space of E -valued differential forms, we write

$$\Omega^k(M, E) = \Omega_0^k(M, E) \oplus \alpha \wedge \Omega_0^{k-1}(M, E). \quad (27)$$

where $\Omega_0^k(M, E) = \ker \iota_X|_{\Omega^k(M)}$.

2.3.1. Contact-Riemannian structure. Let (M, E, X) as above and introduce a metric g_M on M of the form $g_M = \alpha^2 + g_0$ such that $T^*M = \mathbb{R}\alpha + T_0^*M$ becomes an orthogonal decomposition. Let \star and \star_0 denote the Hodge stars associated with (T^*M, g_M) and (T_0^*M, g_0) respectively. We may choose g_0 such that $\star\alpha = (d\alpha)^n = \star_0 1$ whence the Hodge star behaves nicely with respect to the splitting of T^*M . Specifically, we have maps

$$\star : \Omega_0^k(M, E) \rightarrow \alpha \wedge \Omega_0^{N-k-1}(M, E) \quad \star : \alpha \wedge \Omega_0^k(M, E) \rightarrow \Omega_0^{N-k-1}(M, E) \quad (28a)$$

$$\varphi \mapsto (-1)^k \alpha \wedge \star_0 \varphi \quad \alpha \wedge \psi \mapsto \star_0 \psi \quad (28b)$$

Moreover, noting that $\langle \alpha \wedge \varphi, \alpha \wedge \varphi \rangle = \langle \varphi, \varphi \rangle$ on $\Omega_0^\bullet(M, E)$ we conclude $(\iota_X)^T = \alpha \wedge$. When a metric is chosen in this way, compatible with the contact structure, we collectively denote this data (M, E, X, g_M) .

Definition 14. *Considering the maps*

$$\alpha \wedge : \Omega_0^\bullet(M, E) \rightarrow \alpha \wedge \Omega_0^\bullet(M, E), \quad \iota_X : \alpha \wedge \Omega_0^\bullet(M, E) \rightarrow \Omega_0^\bullet(M, E), \quad (29)$$

we set

$$\text{sdet}^b(\alpha \wedge) = \text{sdet}^b(\iota_X) := \left| \text{sdet}^b(\iota_X \circ \alpha \wedge)^{\frac{1}{2}} \right| = 1. \quad (30)$$

2.4. Anosov dynamics. Suppose M is supplemented with a flow $\varphi_t : M \rightarrow M$ for $t \in \mathbb{R}$. We will reuse $X \in C^\infty(M; TM)$ to denote the vector field which generates φ_t .

Definition 15. *The flow is Anosov if there exists a $d\varphi_t$ -invariant continuous splitting of the tangent bundle:*

$$T_x M = E_n(x) \oplus E_s(x) \oplus E_u(x), \quad E_n(x) = \mathbb{R}X_x, \quad (31)$$

and for a given norm $\|\cdot\|$ on TM , there exist constants $C, \lambda > 0$ so that for all $t \geq 0$,

$$\forall v \in E_s(x), \quad \|d\varphi_t(x)v\| \leq Ce^{-\lambda t}\|v\|, \quad \forall v \in E_u(x), \quad \|d\varphi_{-t}(x)v\| \leq Ce^{-\lambda t}\|v\|. \quad (32)$$

The subbundles E_n, E_s, E_u are respectively called *neutral, stable, unstable*.

Remark 16. We will prefer to work with the cotangent bundle T^*M due to the Lie derivative acting naturally on differential forms. The cotangent bundle also has a decomposition $T_x^*M = E_n^*(x) \oplus E_s^*(x) \oplus E_u^*(x)$ whose stable and unstable bundles are understood through the action of $(d\varphi_{-t})^T$ (rather than $d\varphi_t$).

Henceforth we will always assume that the stable and unstable bundles are orientable and each have rank n .

2.4.1. *A guiding example.* We provide an example of the geometric setting discussed in the previous subsections. Let (Σ, g) be a compact manifold without boundary which is oriented, connected, and of dimension $n + 1$. Suppose that Σ has sectional curvature which is everywhere strictly negative. Let $M := S_g^*\Sigma$ be the unit cotangent bundle of Σ . Set $\alpha \in \Omega^1(M)$ to be the pull-back of the canonical one-form on $T^*\Sigma$. Then (M, α) is a contact manifold, and the Reeb vector field $X_g \in C^\infty(M; TM)$ generates the geodesic flow φ_t which is Anosov [Ano67, AS67, AA68].

If we consider $M = S_g^*\Sigma$ within the geometric setting (M, E) of Subsection 2.1 (in particular the representation ρ is unitary and ∇ is flat), we denote the resulting contact, Anosov, Riemannian data on $S_g^*\Sigma$ by

$$(M = S_g^*\Sigma, E, X_g, g).$$

Remark 17. $M \rightarrow \Sigma$ is an \mathbb{S}^n -bundle, and if $n \geq 2$ then $\pi_1(\mathbb{S}^n) = 0$ whence representations of $\pi_1(\Sigma)$ are in one-to-one correspondence with representations of $\pi_1(M)$.

Remark 18. Observe that in the case of a unitary representation and a flat vector bundle, and when $n \geq 2$, the preceding remark implies that $\tau_\rho(M) = (\tau_\rho(\Sigma))^2$ [Fri86, Section 1, p. 526]. For the rest of this article, particularly the announcements of theorems and conjectures, we have chosen to emphasise the role of $M = S_g^*\Sigma$ rather than Σ .

3. RUELLE ZETA FUNCTION

Consider the geometric data (M, E, X) of Section 2.3 and assume the flow φ_t associated to X is Anosov (see Section 2.4). We denote by \mathcal{P} the set of primitive orbits of the flow φ_t and by $\ell(\gamma)$ the period of any given $\gamma \in \mathcal{P}$.

Definition 19. *The Ruelle zeta function (associated with the trivial representation of $\pi_1(M)$) is defined as*

$$\zeta(\lambda) := \prod_{\gamma \in \mathcal{P}} (1 - e^{-\lambda \ell(\gamma)}) \quad (33)$$

whose convergence is assured for $\operatorname{Re} \lambda \gg 1$. The Ruelle zeta function twisted by an arbitrary representation ρ is

$$\zeta_\rho(\lambda) := \prod_{\gamma \in \mathcal{P}} \det(I - \rho([\gamma])e^{-\lambda \ell(\gamma)}). \quad (34)$$

whose convergence is again assured for $\operatorname{Re} \lambda \gg 1$.

Here $[\gamma]$ represents the conjugacy class of γ in $\pi_1(M)$. It has been shown that the zeta functions continue meromorphically to \mathbb{C} [BL07, Mar11, GLP13, DZ16]. We have the following:

Theorem 20 ([Fri86]). *Let (M, E, X, g) be the geometric data of Subsection 2.4.1, where $M = S_g^*\Sigma$, with Σ a closed, oriented hyperbolic manifold $\Sigma = \Gamma \backslash \mathbb{H}^{n+1}$, and g the induced hyperbolic metric. Then, the Ruelle zeta function, defined for $\operatorname{Re}(s) > n$ by equation (34) extends meromorphically to \mathbb{C} and*

$$|\zeta_\rho(0)|^{(-1)^n} = \tau_\rho(M).$$

This fact has inspired a conjecture [Fri87]:

Conjecture 21 (Fried). *Let (M, E, X, g) be the geometric data of Subsection 2.4.1 with $\dim(M) = 2n + 1$. Then the (twisted) Ruelle zeta function computes the analytic torsion:*

$$|\zeta_\rho(0)|^{(-1)^n} = \tau_\rho(M). \quad (35)$$

3.1. Differential forms decomposition. This section shows how the (twisted) Ruelle zeta function may be written as an alternating product of zeta functions associated with $(\bigwedge^k T_0^*M) \otimes E$ for $0 \leq k \leq 2n$.

Given a closed orbit γ , of length $\ell(\gamma)$, and a point $x \in M$ in the orbit, let $P(\gamma, x)$ denote the linearised Poincaré map on the fibre of T_0^*M above x :

$$P(\gamma, p) := (d\varphi_{-\ell(\gamma)})^T : T_0^*M(x) \rightarrow T_0^*M(x) \quad (36)$$

This map is conjugate to $P(\gamma, x')$ for other x' in the same orbit γ , and as we need only evaluate this map's trace and determinant, we will refer to all maps as $P(\gamma)$. Recall T_0^*M is the subbundle of T^*M conormal to the vector field X .

We start with the basic linear algebra identity

$$\det(I - A) = \sum_{k=0}^m (-1)^k \operatorname{tr}(\bigwedge^k A)$$

for an endomorphism A on an m -dimensional vector space and the observation that $P(\gamma)$ has precisely n eigenvalues greater than 1 (where n is the rank of E_u^*). Therefore for $j \geq 1$

$$(-1)^n |\det(I - P(\gamma)^j)| = \sum_{k=0}^{2n} (-1)^k \operatorname{tr}(\bigwedge^k P(\gamma)^j). \quad (37)$$

Using identity (37) to obtain equality (38c) below, we derive

$$\log \zeta_\rho(\lambda) = \sum_{\gamma \in \mathcal{P}} \operatorname{tr} \log(I - \rho([\gamma])e^{-\lambda\ell(\gamma)}) \quad (38a)$$

$$= - \sum_{\gamma \in \mathcal{P}} \operatorname{tr} \sum_{j=1}^{\infty} \frac{1}{j} e^{-\lambda j \ell(\gamma)} \rho([\gamma])^j \quad (38b)$$

$$= - \sum_{\gamma \in \mathcal{P}} \sum_{j=1}^{\infty} \frac{1}{j} e^{-\lambda j \ell(\gamma)} \operatorname{tr}(\rho([\gamma])^j) \left((-1)^n \sum_{k=0}^{2n} \frac{(-1)^k \operatorname{tr}(\bigwedge^k P(\gamma)^j)}{|\det(I - P(\gamma)^j)|} \right) \quad (38c)$$

$$= (-1)^n \sum_{k=0}^{2n} (-1)^k \log \zeta_{\rho, k}(\lambda). \quad (38d)$$

Equation (38d) in the preceding display defines implicitly the Ruelle zeta function twisted by ρ upon restriction to k -forms

$$\zeta_{\rho,k}(\lambda) := \exp \left(- \sum_{\gamma \in \mathcal{P}} \sum_{j=1}^{\infty} \frac{1}{j} \frac{e^{-\lambda j \ell(\gamma)} \operatorname{tr}(\rho([\gamma])^j) \operatorname{tr}(\wedge^k P(\gamma)^j)}{|\det(I - P(\gamma)^j)|} \right) \quad (39)$$

so in the form of a compact equality, we have

$$\zeta_{\rho}(\lambda)^{(-1)^n} = \prod_{k=0}^{2n} \zeta_{\rho,k}(\lambda)^{(-1)^k}. \quad (40)$$

3.2. Ruelle zeta function as regularized determinant. We aim to link the function $\zeta_{\rho,k}$ with the operator $\mathcal{L}_{X,k}$ acting on $\Omega_0^k(M; E)$. This is done with the help of the Atiyah–Bott–Guillemin trace formula [Gui77]. We use the notation

$$e^{-t\mathcal{L}_{X,k}} = \varphi_{-t}^* : L^2(M; \bigwedge^k T_0^* M \otimes E) \rightarrow L^2(M; \bigwedge^k T_0^* M \otimes E). \quad (41)$$

and write the Schwartz kernel $K_k(t, \cdot, \cdot)$ such that

$$(e^{-t\mathcal{L}_{X,k}} \psi)(x) = \int_M K_k(t, x, y) \psi(y) dy. \quad (42)$$

Due to the microlocal structure of K_k we may take its flat trace, which leads to the Atiyah–Bott–Guillemin trace formula:

$$\operatorname{tr}^b e^{-t\mathcal{L}_{X,k}} = \sum_{\gamma \in \mathcal{P}} \sum_{j=1}^{\infty} \ell(\gamma) \delta(t - j\ell(\gamma)) \frac{\operatorname{tr}(\rho([\gamma])^j) \operatorname{tr}(\wedge^k P(\gamma)^j)}{|\det(I - P(\gamma)^j)|} \quad (43)$$

as a distribution on \mathbb{R}_+ . Integrating this distribution against the function $-t^{-1}e^{-t\lambda}$ provides an integral representation of $\zeta_{\rho,k}(\lambda)$:

$$\log \zeta_{\rho,k}(\lambda) = - \int_0^{\infty} t^{-1} e^{-t\lambda} \operatorname{tr}^b e^{-t\mathcal{L}_{X,k}} dt. \quad (44)$$

Consider now the following function, dependent on two variables:

$$F_{\mathcal{L}_{X,k}}(\lambda, s) := \frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-1} \operatorname{tr}^b e^{-t(\mathcal{L}_{X,k} + \lambda)} dt. \quad (45)$$

This function is holomorphic for small s (near $s = 0$) and for $\operatorname{Re}(\lambda) \gg 1$ (see Subsection 3.3). Naively, at $s = 0$ the integrand poses a problem due to the t^{-1} structure as $t \rightarrow 0$, however the trace formula (Equation (43)) provides a natural cut-off in small t so that we avoid this problem. Moreover, for small s we expand $1/\Gamma(s) = s + O(s^2)$ showing

$$\partial_s|_{s=0} F_{\mathcal{L}_{X,k}}(\lambda, s) = \left(\partial_s|_{s=0} \frac{1}{\Gamma(s)} \right) \cdot \int_0^{\infty} t^{s-1} \operatorname{tr}^b e^{-t(\mathcal{L}_{X,k} + \lambda)} dt \Big|_{s=0} \quad (46)$$

$$= -\log \zeta_{\rho,k}(\lambda). \quad (47)$$

Recalling Definition 7 for the flat determinant of an operator now indicates that, for $\operatorname{Re} \lambda \gg 1$

$$\log \det^b(\mathcal{L}_{X,k} + \lambda) \equiv -\partial_s|_{s=0} F_{\mathcal{L}_{X,k}}(\lambda, s) = \log \zeta_{\rho,k}(\lambda). \quad (48)$$

Since $\zeta_{\rho,k}(\lambda)$ has a meromorphic extension to the complex plane, so does the function $\det^b(\mathcal{L}_{X,k} + \lambda)$. Assuming there are no poles at $\lambda = 0$, we sensibly have $\det^b(\mathcal{L}_{X,k})$ as the value at zero of the meromorphic extension of the Ruelle zeta function (restricted to k -forms) at zero:

$$\det^b(\mathcal{L}_{X,k}) = \zeta_{\rho,k}(0). \quad (49)$$

The decomposition in Equation (40) of the Ruelle zeta function now gives an operator interpretation of the Ruelle zeta function:

$$\zeta_{\rho}(\lambda)^{(-1)^n} = \prod_{k=0}^{2n} \det^b(\mathcal{L}_{X,k} + \lambda)^{(-1)^k}. \quad (50)$$

Remark 22. Notice that, due to the decomposition (40) and the trace formula (43), we can consider ζ_{ρ} as directly dependent on an Anosov vector field X . To stress this fact we will use the notation $\zeta(X, \lambda)$ (resp. $\zeta_k(X, \lambda)$) instead of $\zeta_{\rho}(\lambda)$ (resp. $\zeta_{\rho,k}(\lambda)$).

3.3. Meromorphic extension of the resolvent. In the preceding section we related the resolvent $(\mathcal{L}_{X,k} + \lambda)^{-1}$ with $\zeta_{\rho,k}(\lambda)$ for $\lambda \gg 1$. Here we announce a more precise statement for the meromorphic extension of $(\mathcal{L}_{X,k} + \lambda)^{-1}$ [DZ16].

The operator norm of $e^{-t\mathcal{L}_{X,k}}$ is bounded by $e^{C_0 t}$ for some $C_0 > 0$. Therefore the resolvent $(\mathcal{L}_{X,k} + \lambda)^{-1}$, as an operator on L^2 sections, exists for $\text{Re } \lambda > C_0$ and is given by the formula

$$(\mathcal{L}_{X,k} + \lambda)^{-1} = \int_0^{\infty} e^{-t(\mathcal{L}_{X,k} + \lambda)} dt. \quad (51)$$

The restricted resolvent

$$R_k(\lambda) = (\mathcal{L}_{X,k} + \lambda)^{-1} : C^{\infty}(M; \bigwedge^k T_0^* M \otimes E) \rightarrow \mathcal{D}'(M; \bigwedge^k T_0^* M \otimes E) \quad (52)$$

has a nowhere-vanishing meromorphic continuation to \mathbb{C} whose poles are of finite rank, and are called Pollicott-Ruelle resonances. For each $\lambda_0 \in \mathbb{C}$, we have the expansion

$$R_k(\lambda) = R_k^H(\lambda) + \sum_{j=1}^{J(\lambda_0)} \frac{(-1)^{j-1} (\mathcal{L}_{X,k} + \lambda_0)^{j-1} \Pi_{\lambda_0}}{(\lambda - \lambda_0)^j} \quad (53)$$

where R_k^H is holomorphic near λ_0 and $\Pi_{\lambda_0} : C^{\infty}(M; \bigwedge^k T_0^* M \otimes E) \rightarrow \mathcal{D}'(M; \bigwedge^k T_0^* M \otimes E)$ is a finite rank projector. The range of Π_{λ_0} defines (generalised) resonant states. They are characterised as

$$\text{Range } \Pi_{\lambda_0} = \left\{ \varphi \in \mathcal{D}'(M; \bigwedge^k T_0^* M \otimes E) : \text{WF}(u) \subset E_u^*, (\mathcal{L}_{X,k} + \lambda_0)^{J(\lambda_0)} \varphi = 0 \right\}. \quad (54)$$

where WF refers to the wave-front of a distribution (or current), E_u^* is the unstable bundle referenced in Remark 16, and $J(\lambda_0)$ denotes the multiplicity. The adjective ‘‘generalised’’ refers to the possibility that the pole may not be simple (and is superfluous in the case $J(\lambda_0) = 1$).

Finally, the poles of the meromorphic continuation $R_k(\lambda)$ correspond to zeros of the zeta function $\zeta_k(\lambda)$ (which is entire for each k), and the rank of the projector Π_{λ} equals the multiplicity of the zero.

4. BF THEORY ON CONTACT MANIFOLDS

In this section we will analyse a field theory that goes under the name of BF theory¹⁰, in the Batalin-Vilkovisky (BV) formalism. Consider the basic geometric data (M, E) of Subsection 2.1. The classical version of abelian BF theory (i.e. without BV Formalism) is given by the following assignment:

Definition 23. *Define the space of classical fields to be $F_{BF} := \Omega^1(M, E) \oplus \Omega^{N-2}(M, E)$, and the BF action functional:*

$$S_{BF} = \int_M B \wedge d_{\nabla} A. \quad (55)$$

Then we call BF theory the assignment $(M, E) \rightsquigarrow (F_{BF}, S_{BF})$.

Remark 24. Recall that Subsection 2.1 equips E with a Hermitian metric h . This metric has been implicitly used to couple B and A . Specifically, both B and $d_{\nabla} A$ take values in E whence h must be used so that $B \wedge d_{\nabla} A \in \Omega^N(M)$.

Remark 25. It is easy to see that shifting either B or A by a d_{∇} -exact form leaves the action functional unchanged.¹¹ This goes under the name of reducible symmetry, and it is conveniently treated by means of the BV formalism.

Let us consider the space of differential forms $\Omega^{-\bullet}(M, E)$ as a \mathbb{Z} -graded vector space, such that homogeneous forms $\omega^{(k)} \in \Omega^k(M, E)$ will have degree $|\omega^{(k)}| := -k$. We define the space of Batalin-Vilkovisky fields for BF theory to be the graded vector space

$$\mathcal{F}_{BF} := \Omega^{-\bullet}(M, E)[1] \oplus \Omega^{-\bullet}(M, E)[N-2] \ni (\mathbb{A}, \mathbb{B}) \quad (56)$$

where the degree shift means that a k -form in $\Omega^{-\bullet}(M, E)[1]$ will have degree $|\mathbb{A}| = 1 - k$. The symplectic structure reads

$$\Omega_{BF} = \int_M [\delta \mathbb{B} \delta \mathbb{A}]^{\text{top}}, \quad (57)$$

and we define an action functional on \mathcal{F}_{BF} as

$$\mathbb{S}_{BF} = \int_M [\mathbb{B} d_{\nabla} \mathbb{A}]^{\text{top}}. \quad (58)$$

Observe that $|\mathbb{S}_{BF}| = 0$ and $|\Omega_{BF}| = -1$.

Remark 26. We stress that, although the functional form of S_{BF} in Equation (55) and \mathbb{S}_{BF} in (58) is the same, in the latter \mathbb{B} and \mathbb{A} are inhomogeneous forms with an additional shift in degree. Such a degree shift effectively switches the total parity of inhomogeneous forms, so that, if N is odd, even forms will have odd parity and vice-versa. This will have a crucial impact in the partition-function interpretation of quantities such as the analytic torsion and Ruelle zeta function.

¹⁰The name BF comes from the tradition of denoting fields with B and A , and Lagrangian density $B \wedge F_A$.

¹¹Strictly speaking this is true only up to boundary terms. We will assume that M has no boundary, but otherwise the analysis of boundary terms is relevant, and can be performed with a version of the BV formalism [CMR18].

Definition 27. *The assignment $(M, E) \rightsquigarrow (\mathcal{F}_{BF}, \Omega_{BF}, \mathbb{S}_{BF}, Q_{BF})$, with*

$$Q_{BF}\mathbb{B} = d_{\nabla}\mathbb{B}, \quad Q_{BF}\mathbb{A} = d_{\nabla}\mathbb{A} \quad (59)$$

is called BF theory in the Batalin–Vilkovisky formalism.

4.1. Analytic torsion from resolutions of de Rham differential. In this section we will discuss the relation of analytic torsion with degenerate action functionals, following Schwarz [Sch78, Sch79]. This is related to BF, as we will highlight in what follows. In Section 4.2 we will interpret this relation in terms of a gauge-fixing for BF theory in the Batalin–Vilkovisky framework.

Consider the geometric data (M, E, g_M) of Subsection 2.2. Note that the only importance of the vector bundle is to ensure acyclicity of the twisted de Rham complex. This requirement will be necessary in Section 5, however, what we will say here can be extended to nontrivial cohomology along the lines of [CMR20, CMR18].

We can define the partition function associated to abelian BF theory as follows. We first need a resolution of the kernel of the operator featuring in the action functional, represented then as the 0th-cohomology of a chain complex. For abelian BF theory it is given by the following:

$$0 \rightarrow V_{N-2} \xrightarrow{T_{N-2}} V_{N-1} \xrightarrow{T_{N-1}} \dots \xrightarrow{T_3} V_2 \xrightarrow{T_2} V_1 \xrightarrow{T_1} V_0 \xrightarrow{T} V_0 \rightarrow 0 \quad (60)$$

where

$$V_0 = \Omega^1(M) \oplus \Omega^{N-2}(M), \quad T = \begin{bmatrix} 0 & \star d_{N-2} \\ \star d_1 & 0 \end{bmatrix}; \quad (61a)$$

$$V_1 = \Omega^0(M) \oplus \Omega^{N-3}(M), \quad T_1 = \begin{bmatrix} d_0 & 0 \\ 0 & d_{N-3} \end{bmatrix}; \quad (61b)$$

$$V_2 = \Omega^{N-4}(M), \quad T_2 = \begin{bmatrix} 0 \\ d_{N-4} \end{bmatrix}; \quad (61c)$$

$$V_k = \Omega^{N-(k+2)}(M), \quad T_k = d_{N-(k+2)} \quad (61d)$$

for $3 \leq k \leq N-2$. This implies $T_k T_k^* = d_{N-(k+2)} d_{N-(k+2)}^*$ and

$$T^2 = \begin{bmatrix} d_1^* d_1 & 0 \\ 0 & d_{N-2}^* d_{N-2} \end{bmatrix}, \quad T_1 T_1^* = \begin{bmatrix} d_0 d_0^* & 0 \\ 0 & d_{N-3} d_{N-3}^* \end{bmatrix}, \quad T_2 T_2^* = \begin{bmatrix} 0 & 0 \\ 0 & d_{N-4} d_{N-4}^* \end{bmatrix}. \quad (62)$$

In fact, denoting $C := (A, B) \in V_0$, the operator T allows us to rewrite

$$S_{BF} = \int_M B \wedge dA = \frac{1}{2} (C, TC) \quad (63)$$

where (\cdot, \cdot) is the inner product (17) on V_0 as discussed in Section 2.2. Then, the partition function of the degenerate action functional (63), with respect to the resolution (60), is

defined by Schwarz in [Sch79] to be:

$$Z_{\text{Sch}}[T, T_i] := \det^b(T^2)^{-\frac{1}{4}} \prod_{k=1}^{N-2} \det^b(T_k T_k^*)^{(-1)^{k+1} \frac{1}{2}}. \quad (64)$$

Remark 28. The procedure outlined above recursively produces spaces V_k at every stage in order to parametrise the relevant quotients $V_k/\ker(T_k)$, on which integration makes sense. Indeed, said quotients coincide with $\text{coker}(T_{k+1}) \simeq \ker(T_{k+1}^*)$, and the localisation to the correct integration subspaces is obtained by restriction to $\ker(T_k^*)$. Thus, Schwarz's definition of the partition function goes through an extension to a larger space of fields, and the choice of a subspace where integration is well defined: the kernel of the map $\mathbb{T} := \bigoplus_{j=1}^{N-2} T_j^*$. Notice that this construction produces a resolution in the sense of Equation (3).

In the case of abelian BF theory this leads to

$$\begin{aligned} Z_{\text{Sch}}[T, T_i] &= \det^b(d_1^* d_1)^{-\frac{1}{4}} \det^b(d_{N-2}^* d_{N-2})^{-\frac{1}{4}} \det^b(d_0^* d_0)^{\frac{1}{2}} \\ &\quad \times \det^b(d_{N-3}^* d_{N-3})^{\frac{1}{2}} \prod_{k=2}^{N-2} \det^b(d_{N-(k+2)} d_{N-(k+2)}^*)^{\frac{(-1)^{k+1}}{2}}. \end{aligned} \quad (65)$$

It is possible rewrite the expression for the partition function in a familiar form using the following lemma.

Lemma 29. *Let (M, E, ∇, ρ) be as in Subsection 2.1, then the following holds:*

- (1) $\det^b(d_k^* d_k) = \det^b(d_k d_k^*)$
- (2) $\det^b(d_{k-1} d_{k-1}^*) = \det^b(d_{N-k}^* d_{N-k})$
- (3) $\det^b(\Delta_k) = \det^b(d_{k-1} d_{k-1}^*) \det^b(d_k^* d_k) = \det^b(d_{k-1}^* d_{k-1}) \det^b(d_k d_k^*)$.

Sketch of proof. The proof is immediate from the analysis of the spectra of the operators under the consideration. More precisely, note that in this case \det^b 's are spectral invariants as they are the usual zeta-regularized determinants.

(1) follows from the fact that the operators $d_k^* d_k$ and $d_k d_k^*$ are isospectral, a property one can check by acting with d on a coexact eigenform of Δ .

Similarly, we observe that $(*_{N-k})^{-1} \circ d_{k-1} d_{k-1}^* \circ *_N = d_{N-k}^* d_{N-k}$, which implies that $d_{k-1} d_{k-1}^*$ and $d_{N-k}^* d_{N-k}$ are isospectral and (2) follows as well.

Finally (3) follows from the observation that the spectrum of $\Delta_k = d_k^* d_k + d_{k-1} d_{k-1}^*$ is the union of spectrum of $d_{k-1} d_{k-1}^*$ and $d_k^* d_k$. \square

Thanks to the previous lemma, we get:

Proposition 30. *The partition function $Z_{\text{Sch}}[T, T_i]$ for the resolution (60) yields*

$$Z_{\text{Sch}}[T, T_i] = \tau_\rho(M). \quad (66)$$

Proof. From (1) and (2) of Lemma 29, we have

$$\det^b(d_{N-(k+2)} d_{N-(k+2)}^*) = \det^b(d_{k+1}^* d_{k+1}).$$

Using this relation and Lemma 29 again, we can rewrite (65) as

$$Z_{\text{Sch}}[T, T_i] = \det^b(d_1^* d_1)^{-\frac{1}{4}} \det^b(d_{N-2}^* d_{N-2})^{-\frac{1}{4}} \det^b(d_0^* d_0)^{\frac{1}{2}} \\ \det^b(d_{N-3}^* d_{N-3})^{\frac{1}{2}} \prod_{k=2}^{N-2} \det^b(d_{k+1}^* d_{k+1})^{\frac{(-1)^{k+1}}{2}}. \quad (67)$$

Now, we know again by Lemma 29 that $\det^b(d_1^* d_1) = \det^b(d_{N-2}^* d_{N-2})$ and $\det^b(d_{N-3}^* d_{N-3}) = \det^b(d_2^* d_2)$.

Finally, using (3) of the Lemma 29, we can write

$$\prod_{k=0}^{N-1} \det^b(d_k^* d_k)^{(-1)^k \frac{1}{2}} = \prod_{k=1}^N \det^b(\Delta_k)^{(-1)^{k+1} \frac{k}{2}}$$

In summary, we have

$$Z_{\text{Sch}}[T, T_i] = \prod_{k=1}^N \det^b(\Delta_k)^{(-1)^{k+1} \frac{k}{2}},$$

showing that the partition function $Z_{\text{Sch}}[T, T_i]$ for the resolution (60) of the operator d coincides with the analytic torsion $\tau_\rho(M)$ of Definition 13. \square

4.2. BV interpretation: metric gauge for BF theory. We would like to phrase the procedure outlined in Section 4.1 in the Batalin–Vilkovisky framework.

First of all we need to observe that, when the complex is acyclic, the submanifold

$$\mathbb{L}_g : \{d_\nabla^* \mathbb{B} = d_\nabla^* \mathbb{A} = 0\} \quad (68)$$

is Lagrangian in the space of BV fields \mathcal{F}_{BF} (see Remark 5), and can be therefore used as a gauge fixing for BF theory. This is a consequence of Hodge decomposition (on closed manifolds without boundary). In fact one can easily check that the images of d and d^* are both isotropic in \mathcal{F}_{BF} , and complementary due to Hodge decomposition. We refer to this gauge fixing as the metric gauge.

Due to the vanishing of the cohomology of $\Omega^\bullet(M, E)$, the metric gauge Lagrangian submanifold \mathbb{L}_g can be parametrised as follows. The condition $d_\nabla^* \mathbb{B} = 0$ can be equivalently written as $\mathbb{B} = d_\nabla^* \eta$, and if we specify the form degree of the various fields with a subscript, we write $\mathbb{B}_{N-k-1} = d_\nabla^* \eta_{N-k}$. Finally, writing $\eta_{N-k} = \star \tau_k$, together with the inner product given in Equation 17 provides the following formula:

$$\mathbb{S}_{BF}|_{\mathbb{L}_g} \equiv \int_M [\mathbb{B} d_\nabla \mathbb{A}]_{\mathbb{L}_g}^{\text{top}} = \sum_{k=0}^{N-1} \int_M \star d_\nabla \tau_k d_\nabla \mathbb{A}_k |_{d^* \mathbb{A} = 0} = \sum_{k=1}^N (\tau_k, d_\nabla^* d_\nabla |_{\text{coexact } \mathbb{A}_k}). \quad (69)$$

Remark 31. We consider an interpretation of Equation 66 in terms of the BV construction we just outlined¹². We will see how the analytic torsion can be seen as a way of making sense of the (super) determinant of the (graded) operator $d: \Omega_{\text{coexact}}^\bullet \rightarrow \Omega_{\text{exact}}^{\bullet+1}$. Let us define first

$$\text{sdet}^b(d_\nabla^*) \equiv \text{sdet}^b(d_\nabla) := \text{sdet}^b(d_\nabla^* d_\nabla)^{\frac{1}{2}}. \quad (70)$$

¹²The authors would like to thank P. Mnev and A.S. Cattaneo for valuable insight on this interpretation.

Definition 9 interprets partition functions of quadratic functionals in terms of the super determinant of the operator that features in the functional. However, when an explicit gauge fixing is considered, one needs to take into account the parametrisation of the gauge-fixing Lagrangian. In our case this introduces a ‘‘Jacobian superdeterminant’’ for the change of coordinates $\mathbb{B} = d_{\nabla}^* \eta$. Moreover, one should pay attention to the shift in degree $\Omega^{-\bullet}(M, E)[1]$ and $\Omega^{-\bullet}(M, E)[N - 2]$. As a matter of fact, in this circumstance the change of variables operator d^* is acting on k -form of parity $k + 1 \pmod 2$ (instead of $k \pmod 2$), so that the Jacobian super determinant will appear with the ‘‘opposite’’ power. For the same reason, recalling Remark 12, the output of a Gaussian integration over a 1-shifted graded vector space will yield the superdeterminant of the relevant operator (instead of its inverse).

Hence, one recovers the analytic torsion formally as a Gaussian integral on a super vector space by defining the partition function for gauge-fixed BF theory to be a flat-regularised Berezinian/super determinant, times the Jacobian (super) determinant $\text{sdet}^b(d^*)^{-1}$ of the change of variables operator:

$$Z(\mathcal{S}_{BF}, \mathbb{L}_g) = \int_{\mathbb{L}_g} e^{i\mathbb{S}_{BF}}|_{\mathbb{L}_g} = |\text{sdet}^b(d_{\nabla}^*)|^{-1} \int \exp\left(i \sum_{k=0}^{N-1} (\tau_k, (d_k^* d_k) \mathbb{A}_k)\right) \quad (71a)$$

$$:= |\text{sdet}^b(d_{\nabla})|^{-1} \left| \text{sdet}^b(d_{\nabla}^* d_{\nabla}) \right| = \left| \text{sdet}^b(d_{\nabla}^* d_{\nabla}) \right|^{\frac{1}{2}} \quad (71b)$$

$$= \prod_{k=0}^{N-1} \det^b(d_k^* d_k)^{(-1)^k \frac{1}{2}} = \tau_{\rho}(M), \quad (71c)$$

which, comparing with (70), means (cf. [CMR20, Lemma 5.5])

$$\text{sdet}^b(d_{\nabla}) \equiv \text{sdet}^b(d_{\nabla}^*) := \tau_{\rho}(M).$$

Observe that, as mentioned in Remark 10, our definition of partition function discards a potential phase factor in Equations (71), which depends on the signature of the operator. However, as observed in [CMR20, Proposition 5.7 and Remarks 5.8, 5.9], in the case of acyclic complexes the phase drops out. The analysis of the partition function of BF theory in the metric gauge fixing is discussed in detail in [CMR20], where BF theory is cast on triangulated manifolds with boundary (cellular decompositions of cobordisms) and its partition function is shown to coincide with the Reidemeister torsion. The triangulation of a manifold can also be seen as a way of regularising the infinite dimensional BV Laplacian Δ_{BV} , of which \mathbb{S}_{BF} is a cocycle.

Remark 32. In Schwarz’s formalism the computation of the partition function is done by performing integration over $\ker(\mathbb{T}) \subset \bigoplus_{j=0}^{N-2} V_j$ (see Remark 28). It is tantamount to the 0-th Koszul–Tate cohomology, i.e. the resolution presented in Equation (3). This can be compared with the metric gauge in the BV interpretation, where $\mathcal{F}_{BF} \simeq T^*[-1] \bigoplus_{k=0}^{N-2} V_k[k]$. In order to represent integration over $\ker(\mathbb{T})$ within the BV formalism, one considers \mathbb{L}_g , which is essentially the same condition — i.e. restriction to coclosed forms — extended to the shifted cotangent bundle.

4.3. Contact gauge fixing for BF theory. This subsection shows that if BF theory is cast on a manifold that admits a contact structure, it is possible to find an alternative gauge fixing condition. Adopting the same point of view on the partition function, we show how one recovers the Ruelle zeta function.

Consider the geometric data¹³ from Subsection 2.3, (M, E, X) , and consider BF theory cast in the Batalin–Vilkovisky formalism as in Definition 27.

Proposition 33. *The submanifold*

$$\mathbb{L}_X := \{(\mathbb{A}, \mathbb{B}) \in \mathcal{F}_{BF} \mid \iota_X \mathbb{B} = 0; \iota_X \mathbb{A} = 0\}$$

is Lagrangian in \mathcal{F}_{BF} .

Proof. We show that the condition $\mathbb{L}_X : \iota_X \mathbb{B} = \iota_X \mathbb{A} = 0$ defines an isotropic submanifold with an isotropic complement. Indeed, in virtue of the splitting in Eq. (27), we can decompose $\Omega^\bullet(M) = \ker(\iota_X) \oplus \ker(\alpha \wedge)$, so that $\mathbb{A} = \varphi + \alpha \wedge \eta$ and $\mathbb{B} = \psi + \alpha \wedge \xi$. We observe that $\eta = \xi = 0$ on \mathbb{L}_X and, defining $\mathbb{L}_X^\perp := \{(\mathbb{A}, \mathbb{B}) \in \mathcal{F}_{BF} \mid \alpha \wedge \mathbb{A} = \alpha \wedge \mathbb{B} = 0\}$, also that $\psi = \phi = 0$ on \mathbb{L}_X^\perp . Then we have (we understand the top-form part of all the integrands)

$$\Omega_{BF}|_{\mathbb{L}_X} = \int_M [\delta\psi\delta(\alpha \wedge \eta) + \delta(\alpha \wedge \xi)\delta\varphi]_{\mathbb{L}_X} \equiv 0 \quad (72)$$

$$\Omega_{BF}|_{\mathbb{L}_X^\perp} = \int_M [\delta\psi\delta(\alpha \wedge \eta) + \delta(\alpha \wedge \xi)\delta\varphi]_{\mathbb{L}_X^\perp} \equiv 0. \quad (73)$$

This shows that both \mathbb{L}_X and \mathbb{L}_X^\perp are isotropic. \square

Definition 34. *We shall refer to the choice of gauge fixing proposed in Proposition 33 for BF theory as the contact gauge.*

Theorem 35. *Consider the geometric data (M, E, X) . Suppose that the Ruelle zeta function for X has a meromorphic extension which does not vanish at zero. Then, the Ruelle zeta function at zero $\zeta_\rho(0)$ computes the partition function of BF theory in the contact gauge:*

$$Z(\mathcal{S}_{BF}, \mathbb{L}_X) = |\zeta_\rho(0)|^{(-1)^{n+1}}. \quad (74)$$

Proof. We want to compute $Z(\mathcal{S}_{BF}, \mathbb{L}_X) \equiv Z(\mathbb{S}_{BF}|_{\mathbb{L}_X})$. Because of the decomposition (27) we have that $\iota_X \mathbb{B} = 0 \iff \mathbb{B} = (-1)^{|\tau|+1} \iota_X \tau$, for some τ , hence:

$$\mathbb{S}_{BF}|_{\mathbb{L}_X} = \int_M [\mathbb{B} d\nabla \mathbb{A}]_{\mathbb{L}_X}^{\text{top}} = \int_M [(-1)^{|\tau|} \iota_X \tau d\nabla \mathbb{A}]_{\iota_X \mathbb{A}=0}^{\text{top}} \quad (75)$$

$$= \int_M [\tau \iota_X d\nabla \mathbb{A}]_{\iota_X \mathbb{A}=0}^{\text{top}} = \int_M [\tau \mathcal{L}_X |_{\Omega_0^\bullet \mathbb{A}}] \quad (76)$$

Observe that τ and \mathbb{A} are inhomogeneous forms in $\alpha \wedge \Omega_0^\bullet(M, E)[N-2]$ and $\Omega_0^\bullet(M, E)[1]$, respectively. Then, according to Definition 9, recalling that \mathcal{L}_X acts on a 1-shifted graded

¹³Again, the importance of E is to ensure acyclicity.

vector space and referring to Definition 14 for $\text{sdet}^b(\iota_X)$, we have

$$Z(\mathcal{S}_{BF}, \mathbb{L}_X) = \left| \text{sdet}^b \mathcal{L}_X |_{\Omega_0^\bullet} \right| = \left| \prod_{k=0}^{N-1} \left(\det^b \mathcal{L}_{X,k} |_{\Omega_0^k} \right)^{(-1)^k} \right|. \quad (77)$$

This, compared with Equation (50) and (40), allows us to conclude the proof. \square

Remark 36. We would like to stress how the crucial element in both metric and contact gauges is the existence of a ‘‘Hodge decomposition’’ for the space of k -forms. In the metric case it is given, in particular, by the kernel of the de Rham differential and its dual, but truly it can be considered independently of it, as in the contact case, where the maps ι_X and $\alpha\lambda$ define the splitting.

Remark 37. It is tempting to consider a field theory with action functional $S_X := \int_M [\phi \mathcal{L}_X \psi]^{\text{top}}$ on differential forms $\phi, \psi \in \Omega^\bullet(M)$, and call it Ruelle theory. Observe that this is what one gets out of Theorem 35, but with a nontrivial shift in degree. This is akin to considering the theory $\int_M [\mathcal{B}d\mathcal{A}]^{\text{top}}$, for unshifted inhomogeneous differential forms $\mathcal{A}, \mathcal{B} \in \Omega^\bullet$, but - to the best of our knowledge - that does not seem to have a clear interpretation up to now.

5. LAGRANGIAN HOMOTOPIES, FRIED’S CONJECTURE AND GAUGE-FIXING INDEPENDENCE

In this section we outline a strategy to interpret the recent results of [DGRS20] on Fried’s conjecture as invariance of gauge-fixing in the BV formalism, and vice-versa. We will setup a general geometric framework to discuss perturbation of the Anosov vector field as an argument for the Ruelle zeta function, and argue how this can be used to construct homotopies for Lagrangian submanifolds. Indeed, consider the following:

Claim 38. *Assume Theorem 2 holds on \mathcal{F}_{BF} for some appropriate choice of BV Laplacian Δ_{BV} , and assume there exists a Lagrangian homotopy between \mathbb{L}_X and \mathbb{L}_g for any X contact and Anosov. Then, up to phase,*

$$\tau_\rho(M) = Z(\mathcal{S}_{BF}, \mathbb{L}_g) = Z(\mathcal{S}_{BF}, \mathbb{L}_X) = |\zeta_\rho(0)|^{(-1)^n}, \quad (78)$$

which is the statement of (Fried’s) Conjecture 21.

Remark 39. We stress here that a universal generalisation of Theorem 2 to infinite dimensions, i.e. one that works independently of the details of the field theory (and gauge fixing), is currently not yet available. The results of [CMR20] suggest that one possible regularisation scheme for the BV Laplacian might arise as a limit of finite-dimensional cellular analogues. The works of [Cos11] and [Gwi, GH18] assume instead a somewhat different setup, which would also need to be adapted to the case at hand. Ideally, a scheme should be found that makes the regularisation of the BV Laplacian compatible with the context of Definition 7. Observe that the recent results of [CD19] suggest that a number of subtleties might arise, already in acyclic cases.

The central notions for this section, and the main theorem we will need are as follows.

Definition 40. Let X be an Anosov vector field on a manifold M . We will say that X is regular whenever the restricted resolvents $R_k(\lambda) = (\lambda_{X,k} + \lambda)^{-1}$ of Equation (52) do not have poles at $\lambda = 0$ for all $0 \leq k \leq N - 1$.

Recalling Remark 22, we report now a result by Dang, Guillarmou, Rivière and Shen on the properties of Ruelle zeta function seen as a function on Anosov vector fields:

Theorem 41 ([DGRS20]). Let (M, E) denote the geometric data of Subsection 2.1. Consider the set $U \subset C^\infty(M, TM)$ of regular smooth Anosov vector fields X . Then this set is open, and the map $\zeta: U \rightarrow \mathbb{C}$, sending X to $\zeta(X, 0)$ is locally constant and nonzero.

5.1. A sphere bundle construction. Let Σ be a compact manifold. We denote by $\mathcal{R}(\Sigma)$ the space of all Riemannian metrics on Σ , by $\mathcal{R}_<(\Sigma)$ the space of negative sectional curvature metrics, and by $\mathcal{R}_h(\Sigma)$ the space of hyperbolic metrics.

Definition 42. Let us fix a reference metric g_0 . We denote the sphere bundle associated to g_0 by $S_0^*\Sigma := S_{g_0}^*\Sigma$. Moreover, let $g, \tilde{g} \in \mathcal{R}(\Sigma)$ and consider the diffeomorphism

$$\sigma_{\tilde{g}}^g: S_g^*\Sigma \longrightarrow S_{\tilde{g}}^*\Sigma \quad (79)$$

obtained by rescaling lengths in $T\Sigma$. Then, we denote the diffeomorphisms $\sigma_g: S_g^*\Sigma \rightarrow S_0^*\Sigma$, with $\sigma_g := \sigma_{g_0}^{g_0}$ for all $g \in \mathcal{R}(\Sigma)$. Denoting by φ_g the geodesic flow on $S_g^*\Sigma$, and by X_g the associated vector field. One can transfer the geodesic flow φ_g to $S_0^*\Sigma$ by $\varphi_g^0 = \sigma_g \circ \varphi_g \circ \sigma_g^{-1}$; we denote the associated vector field on $S_0^*\Sigma$ by X_g^0 .

Since we are interested in computing the Ruelle zeta function on sphere bundles, we would like to ensure that the geodesic (Reeb) vector field X_g is Anosov in $S_g^*\Sigma$. This won't be true in general, but it will be true for metrics of negative sectional curvature $\mathcal{R}_<$. The map σ_g allows us to consider all Anosov-geodesic vector fields on the same reference space $S_0^*\Sigma$.

Lemma 43. The Anosov property of is preserved under pushforward by σ_g .

Proof. This is an adaptation of a result proved in [Mat13]. Given $S_g^*\Sigma$ with Anosov flow φ_g and associated vector field X_g , we also have a decomposition

$$T(S_g^*\Sigma) = \mathbb{R}X_g + E_s(X_g) + E_u(X_g) \quad (80)$$

with constants C_g, λ_g so that for all $v_s \in E_s(X_g)$, $v_u \in E_u(X_g)$ and all $t > 0$:

$$\|d\varphi_{g,t}v_s\|_g \leq C_g e^{-\lambda_g t} \|v_s\|_g, \quad \|d\varphi_{g,-t}v_u\|_g \leq C_g e^{-\lambda_g t} \|v_u\|_g. \quad (81)$$

Using $d\sigma_g$ we push the decomposition of $T(S_g^*\Sigma)$ to $T(S_0^*\Sigma)$ providing

$$T(S_0^*\Sigma) = \mathbb{R}X_g^0 + E_s(X_g^0) + E_u(X_g^0) \quad (82)$$

where $E_s(X_g^0) := d\sigma_g(E_s(X_g))$ and $E_u(X_g^0) := d\sigma_g(E_u(X_g))$. Now $d\sigma_g$ and $d\sigma_g^{-1}$ are both bounded so for $v_s^0 \in E_s(X_g^0)$, there is $v_s \in E_s$ such that $d\sigma_g v_s = v_s^0$, whence for $t > 0$:

$$\|d\varphi_{g,t}^0 v_s^0\|_0 = \|d\sigma_g d\varphi_{g,t} v_s\|_0 \leq \|d\sigma_g\| \|d\varphi_{g,t} v_s\|_g \leq \|d\sigma_g\| C_g e^{-\lambda_g t} \|v_s\|_g \quad (83)$$

$$= \|d\sigma_g\| C_g e^{-\lambda_g t} \|d\sigma_g^{-1} v_s^0\|_g \leq \|d\sigma_g\| \|d\sigma_g^{-1}\| C_g e^{-\lambda_g t} \|v_s^0\|_0 = C'_g e^{-\lambda_g t} \|v_s^0\|_0 \quad (84)$$

A similar result holds for vectors in $E_u(X_g^0)$. \square

Definition 44. We define the assignment:

$$\mathbb{X}: \mathcal{R}(\Sigma) \longrightarrow C^\infty(S_0^*\Sigma; T(S_0^*\Sigma)); \quad \mathbb{X}(g) = d\sigma_g X_g = X_g^0 \quad (85)$$

where X_g is the geodesic vector field induced by g on $S_g^*\Sigma$.

Proposition 45. Let us denote by $\mathcal{A}(\Sigma)$ the set of Anosov vector fields on $S_0^*\Sigma$, and by $\mathcal{R}_<(\Sigma)$ the space of negative sectional curvature metrics on Σ . Then

$$\mathcal{A}_<(\Sigma) := \text{Im}(\mathbb{X}|_{\mathcal{R}_<(\Sigma)}) \subset \mathcal{A}(\Sigma). \quad (86)$$

Proof. This follows from the general fact that all negative sectional curvature metrics have Anosov geodesic flows, and application of Lemma 43. \square

Lemma 46. For Σ a 2-dimensional surface of negative Euler characteristic $\chi(\Sigma) < 0$, the image of \mathbb{X} restricted to $\mathcal{R}_h(\Sigma)$, the space of hyperbolic metrics on Σ , is a deformation retract of $\mathcal{A}_<(\Sigma)$.

Proof. On 2-dimensional surfaces, the space of negative sectional curvature metrics coincides with the space of negative-Ricci-curvature metrics, which is a contractible subset of \mathcal{R} . In fact, via the Ricci flow one canonically deforms any metric g into a hyperbolic one and, in particular, negatively curved metrics are deformed in such a way along a path of negatively curved metrics [TW15]. Thus, the space of hyperbolic metrics on Σ is a deformation retract of the space of all negatively curved metrics $\mathcal{R}_<(\Sigma)$, and so will be its image under \mathbb{X} with respect to the space $\mathcal{A}_<(\Sigma)$. \square

Theorem 47. Consider the contact, Anosov-Riemannian structure (M, E, X_g, g) of Subsection 2.4.1, where $M = S_g^*\Sigma$ with $\dim(\Sigma) = 2$, the Euler characteristic $\chi(\Sigma) < 0$ and such that $g \in \mathcal{R}_<(\Sigma)$. Then,

$$|\zeta(X_g, 0)| = \tau_\rho(M)^{-1}. \quad (87)$$

Proof. In virtue of Lemma 46 there exists a smooth path $X(t)$ of Anosov vector fields in $\mathcal{A}_<(\Sigma)$ connecting X_g to X_{g_h} for some g_h hyperbolic, where Theorem 20 holds: let $g(t)$ be a retraction such that $g(0) = g_0$ and $g(1) = g$, and set $X(t) = \mathbb{X}(g(t))$. If we can show that each $X(t)$ is regular then local constancy of the zeta function at zero (Theorem 41) provides the result.

The idea for showing regularity is essentially in [DZ17] even though that setting is using the trivial representation, so that the de Rham complex is not acyclic. A more explicit version which adapts the spirit of [DZ17] is in [DGRS20]. Below we cite the necessary results of [DGRS20] to conclude that $X(t)$ is regular.

Since $g \in \mathcal{R}_<(\Sigma)$, its associated geodesic vector field X_g on $S_g^*\Sigma$ is Anosov [Ano67, AS67, AA68]. For regularity, one shows that no k -form resonant states $\varphi^{(k)}$ associated with the spectral parameter $\lambda = 0$ exist for $k \in \{0, 1, 2\}$ (recall Equation (54)). By [DGRS20, Lemma 7.4], 0-form resonant states $\varphi^{(0)}$ are closed and smooth in this setting so may be identified with degree-0 de Rham cohomology. Acyclicity implies such states are necessarily the 0-section. For 2-form resonant states $\varphi^{(k)}$, there is an isomorphism with 0-form resonant states

upon wedging with $d\alpha$ [DGRS20, Lemma 7.2]. Effectively $\varphi^{(2)} = \varphi^{(0)}d\alpha$ so in this case also, no such resonant states exist. Finally, for 1-form resonant states, we require degree-1 cohomology to vanish and appeal to [DGRS20, Lemma 7.2 Hypothesis (1)] to conclude no non-trivial resonant states exist. \square

Corollary 48. *Under the assumptions of Theorem 47, denoting by*

$$\mathbb{L}_X = \{(\mathbb{A}, \mathbb{B}) \in \mathcal{F}_{BF} \mid \iota_{\mathbb{X}(g)}\mathbb{A} = \iota_{\mathbb{X}(g)}\mathbb{B} = 0\}$$

the Lagrangian submanifold defined by the Anosov vector field $\mathbb{X}(g)$, we have

$$Z(\mathcal{S}_{BF}, \mathbb{L}_X) = \tau_\rho(M)^{-1}, \quad (88)$$

for every $g \in \mathcal{R}_<(\Sigma)$.

Proof. This follows from Theorems 35 and 47. \square

Remark 49. Observe that the above construction can be generalised to some extent. In general, the space of negative sectional curvature metric will not be path connected, so let us consider the connected components that contain a hyperbolic metric. The image under \mathbb{X} of the disjoint union of such connected components in $\mathcal{A}(\Sigma)$ will be a union of islands of Anosov vector fields, path connected to an anosov vector fields in $\text{Im}(\mathbb{X}|_{\mathcal{R}_h})$. In a neighborhood of a hyperbolic metric Anosov vector field, the requirements of Theorem 41 are satisfied, and Fried's conjecture might be extended to open subsets of $\mathcal{A}_<(\Sigma)$. We defer the development of such a generalisation to a subsequent work.

Remark 50. The spirit of gauge-fixing homotopies is that of replacing an ill-defined integral with a well-defined one, and can be considered as providing a family of integral representations of the same quantity, only some of which are directly computable. From this point of view, the Ruelle zeta function at zero might not be computable for a generic (Anosov) vector field, but it will be once deformed away from an invalid point in $\mathcal{A}(\Sigma)$.

We wish to interpret this result in terms of homotopies of Lagrangian submanifolds in \mathcal{F}_{BF} and gauge fixing independence for BF theory.

Theorem 51. *Consider the geometric data (M, E, X_g, g) of Subsection 2.4.1, a smooth path $g_t : [0, 1] \rightarrow \mathcal{R}(\Sigma)$ such that $\mathbb{X}(g_0)$ is regular, and let*

$$\mathbb{L}_t := \{(\mathbb{A}, \mathbb{B}) \in \mathcal{F}_{BF} \mid \iota_{\mathbb{X}(g_t)}\mathbb{A} = \iota_{\mathbb{X}(g_t)}\mathbb{B} = 0\}$$

be the associated smooth family of Lagrangian submanifolds in \mathcal{F}_{BF} . Then, $Z(\mathcal{S}_{BF}, \mathbb{L}_0) \neq 0$ and

$$\left. \frac{d}{dt} \right|_{t=0} Z(\mathcal{S}_{BF}, \mathbb{L}_t) = 0. \quad (89)$$

Moreover, if g_0 is hyperbolic, we have that

$$Z(\mathcal{S}_{BF}, \mathbb{L}_0) = \tau_\rho(M). \quad (90)$$

Proof. Lemma 43 ensures that $\mathbb{X}(g_t) = d\sigma_{g_t}X_{g_t}$ is a smooth path of Anosov vector fields in $S_0^*\Sigma$, and in virtue of Theorem 35 we have that

$$Z(\mathcal{S}_{BF}, \mathbb{L}_t) \equiv Z(\mathbb{S}_R|_{\mathbb{L}_t}) = |\zeta(\mathbb{X}(g_t), 0)|^{(-1)^n}.$$

By assumption, at g_0 the k -th Ruelle zeta factors $\zeta_k(\mathbb{X}(g_0), 0)$ are well defined and different from zero. Then, in virtue of Theorem 41 $\zeta(\mathbb{X}(g_0), 0)$ is constant in a open neighborhood of $\mathbb{X}(g_0)$, hence it is on the whole path g_t for $t \in [0, T)$ for some appropriately chosen T and in particular its derivative at t vanishes. If we choose g_0 hyperbolic, using Theorem 20 we can conclude that

$$Z(\mathcal{S}_{BF}, \mathbb{L}_0) = |\zeta(\mathbb{X}(g_0), 0)|^{(-1)^n} \equiv |\zeta_\rho(M)|^{(-1)^n} = \tau_\rho(M).$$

□

ACKNOWLEDGEMENTS

This research was (partly) supported by the NCCR SwissMAP, funded by the Swiss National Science Foundation. MS acknowledges partial support from Swiss National Science Foundation grants P2ZHP2_164999 and P300P2_177862. SK acknowledges partial support from Deutsche Forschungs Gemeinschaft (DFG) under Graduiertenkolleg 1821 (Cohomological Methods in Geometry). This project began, and was mostly developed, during the period when the first and last author were at UC Berkeley, to which we are grateful for having provided such a stimulating research environment. M.S. would like to thank the University of Freiburg for hospitality. We are indebted to A.S. Cattaneo and P. Mnev for their support and feedback at different stages of this work. Since the preprint appeared on arXiv, we have received several useful comments from Y. Chaubet and N.V. Dang whom we warmly thank. We also thank the anonymous referee for providing useful comments that helped improve this paper.

REFERENCES

- [AA68] V.I. Arnold and A. Avez. Problèmes ergodiques de la mécanique classique, Paris, Gauthier-Villars (1967); (English translation: Ergodic problems in classical mechanics) New York, 1968.
- [AB64] M.F. Atiyah and R. Bott. *Notes on the Lefschetz fixed point theorem for elliptic complexes*. Harvard University, 1964.
- [AB68] M.F. Atiyah and R. Bott. A Lefschetz fixed point formula for elliptic complexes: II. Applications. *Annals of Mathematics*, pages 451–491, 1968.
- [And92] I.M. Anderson. Introduction to the variational bicomplex. *Contemporary Mathematics*, 132, 1992.
- [Ano67] D.V. Anosov. Geodesic flows on closed Riemannian manifolds of negative curvature. *Trudy Mat. Inst. Steklov*, 90(5), 1967.
- [AS67] D.V. Anosov and Y.G. Sinai. Some smooth ergodic systems. *Russian Mathematical Surveys*, 22(5):103, 1967.
- [Bal18] V. Baladi. *Dynamical zeta functions and dynamical determinants for hyperbolic maps*. Springer, 2018.
- [BBRT91] D. Birmingham, M. Blau, M. Rakowski, and G. Thompson. Topological field theory. *Physics Reports*, 209(4):129 – 340, 1991.
- [BE15] D. Berwick-Evans. The Chern–Gauss–Bonnet theorem via supersymmetric Euclidean field theories. *Communications in Mathematical Physics*, 335(3):1121–1157, May 2015.
- [Ber83] F.A. Berezin. Introduction to algebra and analysis with anticommuting variables. *Moscow Univ*, 1983.
- [BL75] F.A. Berezin and D.A. Leites. Supermanifolds. In *Doklady Akademii Nauk*, volume 224, pages 505–508. Russian Academy of Sciences, 1975.

- [BL07] O. Butterley and C. Liverani. Smooth Anosov flows: correlation spectra and stability. *J. Mod. Dyn.*, 1(2):301–322, 2007.
- [BRS74] C. Becchi, A. Rouet, and R. Stora. The abelian higgs kibble model, unitarity of the s-operator. *Physics Letters B*, 52(3):344–346, 1974.
- [BRS75] C. Becchi, A. Rouet, and R. Stora. Renormalization of the abelian Higgs-Kibble model. *Communications in Mathematical Physics*, 42(2):127–162, 1975.
- [BRS76] C. Becchi, A. Rouet, and R. Stora. Renormalization of gauge theories. *Annals of Physics*, 98(2):287–321, 1976.
- [BV83] I.A. Batalin and G.A. Vilkovisky. Quantization of gauge theories with linearly dependent generators. *Physical Review D*, 28(10):2567, 1983.
- [BV84] I.A. Batalin and G.A. Vilkovisky. Gauge algebra and quantization. In *Quantum Gravity*, pages 463–480. Springer, 1984.
- [CC18] A.S. Cattaneo and I. Contreras. Split canonical relations. *arXiv preprint arXiv:1811.10107*, 2018.
- [CCRFM95] A.S. Cattaneo, P. Cotta-Ramusino, J. Fröhlich, and M. Martellini. Topological BF theories in 3 and 4 dimensions. *Journal of Mathematical Physics*, 36(11):6137–6160, 1995.
- [CD19] Y. Chaubet and N.V. Dang. Dynamical torsion for contact Anosov flows. *arXiv preprint arXiv:1911.09931*, 2019.
- [CFL06] A.S. Cattaneo, D. Fiorenza, and R. Longoni. Graded Poisson algebras. In J.P. Francoise, G.L. Naber, and T.S. Tsun, editors, *Encyclopedia of Mathematical Physics*, pages 560 – 567. Academic Press, Oxford, 2006.
- [CG16] K. Costello and O. Gwilliam. *Factorization algebras in quantum field theory*, volume 1. Cambridge University Press, 2016.
- [CMR95] S. Cordes, G. Moore, and S. Ramgoolam. Lectures on 2d Yang-Mills theory, equivariant cohomology and topological field theories. *Nuclear Physics B - Proceedings Supplements*, 41(1):184 – 244, 1995.
- [CMR14] A. S. Cattaneo, P. Mnev, and N. Reshetikhin. Classical BV theories on manifolds with boundary. *Communications in Mathematical Physics*, 332(2):535–603, 2014.
- [CMR18] A.S. Cattaneo, P. Mnev, and N. Reshetikhin. Perturbative quantum gauge theories on manifolds with boundary. *Communications in Mathematical Physics*, 357(2):631–730, 2018.
- [CMR20] Alberto S. Cattaneo, Pavel Mnev, and Nicolai Reshetikhin. A cellular topological field theory. *Communications in Mathematical Physics*, 374(2):1229–1320, 2020.
- [Cos11] K. Costello. *Renormalization and effective field theory*, volume 170. American Mathematical Soc., 2011.
- [CR01] A.S. Cattaneo and C.A. Rossi. Higher-dimensional BF theories in the Batalin–Vilkovisky formalism: The BV action and generalized Wilson loops. *Communications in Mathematical physics*, 221(3):591–657, 2001.
- [CS11] A.S. Cattaneo and F. Schätz. Introduction to supergeometry. *Reviews in Mathematical Physics*, 23(06):669–690, 2011.
- [Del17] N.L. Delgado. *Lagrangian field theories: ind/pro-approach and L-infinity algebra of local observables*. PhD thesis, Max Planck Institute for Mathematics, 2017.
- [DF99] P. Deligne and D.S. Freed. Classical field theory. *Quantum fields and strings: a course for mathematicians*, 2:137–226, 1999.
- [DGRS20] Nguyen Viet Dang, Colin Guillarmou, Gabriel Rivière, and Shu Shen. The Fried conjecture in small dimensions. *Inventiones mathematicae*, 220(2):525–579, 2020.
- [DZ16] S. Dyatlov and M. Zworski. Dynamical zeta functions for Anosov flows via microlocal analysis. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(3):543–577, 2016.
- [DZ17] S. Dyatlov and M. Zworski. Ruelle zeta function at zero for surfaces. *Inventiones mathematicae*, 210(1):211–229, 2017.
- [FP16] L.D. Faddeev and V.N. Popov. Feynman diagrams for the Yang-Mills field. In *Fifty Years of Mathematical Physics: Selected Works of Ludwig Faddeev*, pages 157–158. World Scientific, 2016.

- [FR13] K. Fredenhagen and K. Rejzner. Batalin-Vilkovisky formalism in perturbative algebraic quantum field theory. *Communications in Mathematical Physics*, 317(3):697–725, 2013.
- [Fri86] D. Fried. Analytic torsion and closed geodesics on hyperbolic manifolds. *Inventiones mathematicae*, 84(3):523–540, 1986.
- [Fri87] D. Fried. Lefschetz formulas for flows. *Contemporary Mathematics*, 58:19–69, 1987.
- [Fri95] D. Fried. Meromorphic zeta functions for analytic flows. *Communications in Mathematical Physics*, 174(1):161–190, 1995.
- [GH18] Owen Gwilliam and Rune Haugseng. Linear Batalin–Vilkovisky quantization as a functor of ∞ -categories. *Selecta Mathematica*, 24(2):1247–1313, 2018.
- [GLP13] P. Giulietti, C. Liverani, and M. Pollicott. Anosov flows and dynamical zeta functions. *Annals of Mathematics*, pages 687–773, 2013.
- [Gui77] V. Guillemin. Lectures on spectral theory of elliptic operators. *Duke Mathematical Journal*, 44(3):485–517, 1977.
- [Gut91] M.C. Gutzwiller. *Chaos in Classical and Quantum Mechanics*, volume 1. Springer Science & Business Media, 1991.
- [Gwi] Owen Gwilliam. *Factorization Algebras and Free Field Theories*. PhD thesis, Northwestern University.
- [Had18] C.S. Hadfield. Zeta function at zero for surfaces with boundary. *arXiv preprint arXiv:1803.10982*, 2018.
- [Hen90] M. Henneaux. Lectures on the antifield-BRST formalism for gauge theories. *Nuclear Physics B-Proceedings Supplements*, 18(1):47–105, 1990.
- [KW20] Benjamin Küster and Tobias Weich. Pollicott-ruelle resonant states and betti numbers. *Communications in Mathematical Physics*, 378(2):917–941, 2020.
- [Mar11] J. Marklof. Selberg’s trace formula: an introduction. In J. Bolte and F. Steiner, editors, *Hyperbolic Geometry and Applications in Quantum Chaos and Cosmology*, pages 83–119. Cambridge University Press, 2011.
- [Mat13] S. Matsumoto. The space of (contact) Anosov flows on 3-manifolds. *J. Math. Sci. Univ. Tokyo*, 20:445–460, 2013.
- [Mne14] P. Mnev. Lecture notes on torsions. *arXiv preprint arXiv:1406.3705*, 2014.
- [Mne19] P. Mnev. *Quantum Field Theory: Batalin–Vilkovisky Formalism and Its Applications*, volume 72. American Mathematical Soc., 2019.
- [MS91] H. Moscovici and R.J. Stanton. R-torsion and zeta functions for locally symmetric manifolds. *Inventiones mathematicae*, 105(1):185–216, 1991.
- [RS71] D.B. Ray and I.M. Singer. R-torsion and the Laplacian on riemannian manifolds. *Advances in Mathematics*, 7(2):145–210, 1971.
- [Rue76] D. Ruelle. Zeta-functions for expanding maps and Anosov flows. *Inventiones mathematicae*, 34(3):231–242, 1976.
- [Rue86] D. Ruelle. Resonances of chaotic dynamical systems. *Physical review letters*, 56(5):405, 1986.
- [Sch78] A.S. Schwarz. The partition function of degenerate quadratic functional and Ray-Singer invariants. *Letters in Mathematical Physics*, 2(3):247–252, 1978.
- [Sch79] A.S. Schwarz. The partition function of a degenerate functional. *Communications in Mathematical Physics*, 67(1):1–16, 1979.
- [She17] S. Shen. Analytic torsion, dynamical zeta functions, and the Fried conjecture. *Analysis & PDE*, 11(1):1–74, 2017.
- [Sta98] J. Stasheff. The (secret?) homological algebra of the Batalin-Vilkovisky approach. In *Conference Secondary Calculus and Cohomological Physics*, volume 219, pages 195–210. Moscow, August 1998.
- [TW15] W. Tuschmann and D.J. Wraith. *Moduli spaces of Riemannian metrics*. Springer, 2015.
- [Tyu75] I.V. Tyutin. Gauge invariance in field theory and statistical physics in operator formalism. *Lebedev Physics Institute preprint arXiv:0812.0580*, 1975.

- [Vor91] T. Voronov. *Geometric integration theory on supermanifolds*, volume 1. CRC Press, 1991.
- [Wei71] A. Weinstein. Symplectic manifolds and their Lagrangian submanifolds. *Advances in Mathematics*, 6(3):329–346, 1971.
- [Wei10] A. Weinstein. Symplectic categories. *Portugaliae Mathematica*, 67(2):261–278, 2010.
- [Zwo17] M. Zworski. Mathematical study of scattering resonances. *Bulletin of Mathematical Sciences*, 7(1):1–85, 2017.

IBM T.J. WATSON RESEARCH CENTER, 1101 KITCHAWAN RD, YORKTOWN HEIGHTS, NY 10598, USA

E-mail address: `charles.hadfield@ibm.com`

MATHEMATICS INSTITUTE, UNIVERSITY OF FREIBURG, FREIBURG, 79104, GERMANY

E-mail address: `skandel1@alumni.nd.edu`

ETH ZÜRICH

DEPARTMENT OF MATHEMATICS, RÄMISTRASSE 101, 8092 ZÜRICH, SWITZERLAND, &

INSTITUTE FOR THEORETICAL PHYSICS, WOLFGANG-PAULI STRASSE 27, 8093 ZÜRICH, SWITZERLAND

E-mail address: `micschia@phys.ethz.ch`