# Invariant measures for multidimensional fractional stochastic volatility models \*

Balázs Gerencsér† Miklós Rásonyi‡

August 30, 2021

Dedicated to István Gyöngy on the occasion of his 70th birthday

#### **Abstract**

We establish convergence to an invariant measure as time tends to infinity, for a large class of (possibly non-Markovian) stochastic volatility models. Our arguments are based on a novel coupling idea for Markov chains which also extends to Markov chains in random environments in an efficient way.

**Keywords:** Markov chain in random environment; coupling; stochastic volatility; invariant measure; fractional volatility

**MSC 2010:** 60J05, 60J25, 91G80

#### **1 Introduction**

Stochastic volatility models (in the simplest one-dimensional case) are of the form

<span id="page-0-0"></span>
$$
dS_t = v_1(S_t, V_t)S_t dt + V_t S_t d\overline{W}_t, \qquad (1)
$$

where  $\overline{W}$  is a Brownian motion,  $v_1$  is a suitable function and S describes the (discounted) price of an asset with volatility process *V*.

The present paper is about the long-term behaviour of *S*. In the Markovian case, *V* satisfies a stochastic differential equation (SDE),

$$
dV_t = v_2(V_t)dt + \sigma(V_t)dB_t, \qquad (2)
$$

where *B* is another Brownian motion, possibly correlated with  $\overline{W}$ ;  $v_2$ ,  $\sigma$  are suitable functions. In such diffusion models there is an arsenal of techniques from Markov process theory to show that the law of  $S_t$  tends to a limit as  $t \to \infty$ , see e.g. [\[41,](#page-22-0) [32,](#page-21-0) [33,](#page-21-1) [12,](#page-20-0) [38,](#page-22-1) [42,](#page-22-2) [25\]](#page-21-2), Chapter 20 of [\[31\]](#page-21-3) and Subsection 7.1 of [\[22\]](#page-21-4).

Recently, however, fractional stochastic volatility models have become popular (see [\[9,](#page-20-1) [15,](#page-20-2) [45\]](#page-22-3)), where the process *V* is not Markovian. For instance,

<span id="page-0-1"></span>
$$
V_t = \exp(J_t), \quad J_t := \int_{-\infty}^t K(t-s) dB_s,\tag{3}
$$

with some (two-sided) Brownian motion *B*, and a suitable function  $K: \mathbb{R}_+ \to \mathbb{R}$ . In such a setting the question of stochastic stability becomes difficult, one cannot rely on the usual Markovian techniques and there seems to be no results in the literature that would imply the convergence of the law of  $S_t$ as  $t \rightarrow \infty$  at this level of generality.

<sup>\*</sup>Both authors enjoyed the support of the "Lendület" grant LP 2015-6 of the Hungarian Academy of Sciences. The first author was also supported by NRDI (National Research, Development and Innovation Office) grant FK 135711, the János Bolyai Research Scholarship of the Hungarian Academy of Sciences and by the ÚNKP-20-5 New National Excellence Program of the Ministry for Innovation and Technology from the source of the National Research, Development and Innovation Fund.

<sup>†</sup>Alfréd Rényi Institute of Mathematics and Eötvös Loránd University, Budapest, Hungary

<sup>‡</sup>Alfréd Rényi Institute of Mathematics, Budapest, Hungary; rasonyi@renyi.hu

We now explain our motivations for studying such models. Asset price processes often show mean-reversion (for instance, commodities or commodity futures, see [\[6,](#page-20-3) [8\]](#page-20-4)). Optimal investment problems for such assets were considered in [\[17\]](#page-21-5), see also the study [\[13\]](#page-20-5) on asymptotic arbitrage. Long-term investments may also be studied in the framework of ergodic, risk-sensitive or adaptive control (see e.g. [\[24,](#page-21-6) [11,](#page-20-6) [36,](#page-21-7) [5\]](#page-20-7)). All these approaches require that the law of  $S_t$  should converge to a steady state as  $t \to \infty$ . Long-term investment problems for fractional processes were treated in [\[18,](#page-21-8) [34\]](#page-21-9), but these studies do not cover fractional stochastic volatility models.

The present paper proves that – under mean-reversion and smoothness conditions on the drift of *S* and integrability assumptions on  $V_0$ ,  $S_0$  – the stochastic system  $(S_t, V_t)$  converges to an invariant probability, independent of the initialization  $S_0$ . A multi-asset framework is treated and  $B$ ,  $\overline{W}$  will be allowed to have a stochastic correlation. Our arguments are based on a new coupling construction for (discrete-time) Markov processes in random environments.

In the extant literature on fractional volatility, asset dynamics is most often considered for purposes of derivative pricing; [\[4,](#page-20-8) [14,](#page-20-9) [20\]](#page-21-10) are early examples. These papers thus work under the riskneutral measure, which corresponds to taking  $v_1 \equiv 0$  in [\(1\)](#page-0-0). As we have in mind a different class of problems (portfolio optimization), we work under the physical probability, where  $v_1$  is non-zero.

In Section [2](#page-1-0) we rigorously formulate our main results, Theorems [2.9](#page-2-0) and [2.10.](#page-3-0) A novel (discretetime) coupling method is introduced in Section [3.](#page-3-1) As a warm-up, it is first presented for (ordinary) Markov chains in Subsection [3.1.](#page-3-2) Subsection [3.2](#page-7-0) develops the same ideas in the more involved setting of Markov chains in random environments. In Section [4](#page-11-0) we prove the main results, combining advanced Malliavin calculus techniques with the discrete-time construction of Subsection [3.2.](#page-7-0)

#### <span id="page-1-0"></span>**2 Results**

Scalar product in finite-dimensional Euclidean spaces is denoted by  $\langle \cdot, \cdot \rangle$ , the coresponding norm is | · |, where the dimension of the space may vary. For a matrix *A*, *A* ∗ denotes its transpose. For matrices *A*, |*A*| denotes the operator norm.

All the random objects in the present paper will live on a fixed probability space  $(\Omega, \mathscr{F}, P)$ . For a Polish space  $\mathcal{I}$ , its Borel sigma-algebra is denoted  $\mathscr{B}(\mathcal{I})$ . If  $Z:\Omega\to\mathcal{I}$  is  $\mathscr{F}/\mathscr{B}(\mathcal{I})$  is measurable (that is, if *Z* is a *Z*-valued random variable) then  $\mathcal{L}(Z)$  denotes its law on  $\mathcal{B}(\mathcal{Z})$ .

Fix  $d, m \in \mathbb{N} \setminus \{0\}$  with  $d \leq m$ . The number of assets will be *d* and the dimension of the driving noise *m*. For every  $k \geq 1$ , let  $\mathcal{W}^k$  denote the set of continuous  $\mathbb{R}^k$ -valued functions on  $\mathbb R$  which is a Polish space under the metric

$$
\mathbf{d}_k(f,g) := \sum_{i=-\infty}^{\infty} \frac{1}{2^{|i|}} \left[ 1 \wedge \sup_{u \in [i,i+1]} |f(u) - g(u)| \right], \quad f, g \in \mathcal{W}^k.
$$

Let  $B_t$ ,  $t \in \mathbb{R}$  be a two-sided *m*-dimensional Brownian motion (i.e.  $B_t$ ,  $B_{-t}$ ,  $t \in \mathbb{R}_+$  are independent standard *m*-dimensional Brownian motions), let  $\mathscr{G}_t$ ,  $t \in \mathbb{R}$  denote its completed natural filtration. Let V denote the set of  $d \times d$  non-singular matrices,  $\mathcal{R}$  the set of  $d \times m$  matrices r satisfying  $rr^* < I$ , where *I* is the *d*-dimensional identity matrix and  $A < B$  for symmetric, positive semidefinite  $d \times d$ matrices *A*,*B* means that *B* − *A* is positive definite. Similarly,  $A \leq B$  means that  $B - A$  is positive semindefinite and  $\sqrt{A}$  denotes the usual square root of semidefinite matrices.

Let  $V_t$ ,  $t \in \mathbb{R}$  (resp.  $\rho_t$ ,  $t \in \mathbb{R}$ ) be  $\mathcal{V}$ -valued (resp.  $\mathcal{R}$ -valued) processes with continuous trajectories. Notice that  $\mathbf{B}_t := (B_t - B_{t+s})_{s \in \mathbb{R}}$  (resp.  $\mathbf{V}_t := (V_{t+s})_{s \in \mathbb{R}}$  and  $\mathbf{R}_t := (\rho_{t+s})_{s \in \mathbb{R}}$ ) can be naturally regarded as a  $\mathcal{W}^m$ -valued (resp.  $\mathcal{W}^{d \times d}$ -valued and  $\mathcal{W}^{d \times m}$ -valued) random process indexed by  $t \in \mathbb{R}$ .

<span id="page-1-1"></span>**Assumption 2.1.** *There are measurable functions*  $F_1: W^m \to W^{d \times d}$ ,  $F_2: W^m \to W^{d \times m}$  such that  $\mathbf{V}_t = F_1(\mathbf{B}_t)$ ,  $\mathbf{R}_t = F_2(\mathbf{B}_t)$ . Furthermore,  $(V_t, \rho_t)$ ,  $t \in \mathbb{R}$  is adapted to  $\mathscr{G}_t$ ,  $t \in \mathbb{R}$ .

In plain English,  $(V_t, \rho_t)$  is a nonanticipative functional of the increments of the Brownian motion *B* up to *t*. A specification like [\(3\)](#page-0-1) is a typical example. Under Assumption [2.1,](#page-1-1)  $(\mathbf{V}_t, \mathbf{R}_t, \mathbf{B}_t)$ ,  $t \in \mathbb{R}$  is a stationary process in the strong sense.

Let  $W_t$ ,  $t \in \mathbb{R}_+$  be another, *d*-dimensional standard Brownian motion with (completed) natural filtration  $\mathcal{F}_t$ ,  $t \in \mathbb{R}_+$ . Instead of prices, it is more convenient to work with log-prices. Hence we

consider *d* financial assets whose log-price is given by the *d*-dimensional process  $L_t$ ,  $t \in \mathbb{R}_+$  which is the solution of the stochastic differential equation

<span id="page-2-4"></span>
$$
dL_t = \zeta(L_t, V_t)dt + V_t \rho_t dB_t + V_t \sqrt{I - \rho_t \rho_t^*} dW_t,
$$
\n(4)

where  $L_0$  is a random variable and  $\zeta : \mathbb{R}^d \times \mathcal{V} \to \mathbb{R}^d$  is a measurable function.

 $\Delta$ ssumptions [2.2,](#page-2-1) [2.4](#page-2-2) and [2.7,](#page-2-3) stipulated below, guarantee a unique  $(\mathscr{F}_t\vee\mathscr{G}_t)_{t\in\mathbb{R}_+}$ -adapted solution to [\(4\)](#page-2-4), by Theorem 7 on page 82 of [\[28\]](#page-21-11).

<span id="page-2-1"></span>**Assumption 2.2.** *Let*  $\mathscr{G}_{\infty}$  *be independent of*  $\mathscr{F}_{\infty}$ *. Let*  $L_0 = l(R, V_0, R_0)$  *for some measurable l* :  $[0,1] \times W^{d \times d} \times W^{d \times m} \to \mathbb{R}$  and  $[0,1]$ *-uniformly distributed random variable R, which is assumed to be independent of*  $\mathscr{G}_{\infty} \vee \mathscr{F}_{\infty}$ *.* 

**Remark 2.3.** An arbitrary joint law for  $(L_0, V_0, R_0)$  can be realized for suitable *l*, hence Assumption [2.2](#page-2-1) is not restrictive at all. For practical applications, actually, one may assume  $L_0$  to be constant.

<span id="page-2-2"></span>**Assumption 2.4.** *The function*  $\zeta(x, v)$  *is twice continuously differentiable in its first variable,*  $\partial_x \zeta, \partial_{xx}\zeta$ *are bounded. Furthermore, there is*  $K > 0$  *such that*  $|\zeta(x, v_1) - \zeta(x, v_2)| \leq K(1 + |v_1| + |v_2|)|v_1 - v_2|$  for all  $x \in \mathbb{R}^d$ ,  $v_1, v_2 \in V$  (polynomial Lipschitz condition in v).

The following mean-reversion (or dissipativity) condition is rather standard, also in a non-Markovian context, see e.g. [\[21\]](#page-21-12).

<span id="page-2-5"></span>**Assumption 2.5.** *There exist*  $\alpha, \beta > 0$ ,  $\xi \ge 2$  *such that* 

$$
\langle x,\zeta(x,v)\rangle\leq -\alpha|x|^2+\beta(1+|v|^{\zeta}),\ x\in\mathbb{R}^d,\ v\in\mathcal{V}.
$$

**Example 2.6.** We briefly comment on the meaning of Assumptions [2.4](#page-2-2) and [2.5](#page-2-5) in a simple case with one asset  $(d = 1)$  whose price satisfies

$$
dS_t = v_1(S_t)S_t dt + V_t S_t d\overline{W}_t
$$

with some  $S_0 > 0$ , with a  $(\mathbb{R} \setminus \{0\}) \times (-1, 1)$ -valued stationary process  $(V_t, \rho_t)$  and Brownian motion  $\overline{W}_t = \rho_t dB_t + \sqrt{1-\rho_t^2} \, dW_t.$  Let the function  $v_1$  be such that  $\bar v_1(x) := v_1(\exp(x))$  is twice continuously differentiable with  $\bar{v}'_1$ ,  $\bar{v}''_1$  bounded and satisfying

$$
x\bar{v}_1(x) \leq -\bar{\alpha}|x|^2 + \bar{\beta}, \quad x \in \mathbb{R}
$$

with some  $\bar{\alpha}, \bar{\beta} > 0$ . Then  $L_t := \ln(S_t)$  has dynamics

$$
dL_t = \left[\bar{v}_1(L_t) + \frac{V_t^2}{2}\right]dt + V_t d\overline{W}_t
$$

and  $\zeta(x, v) := \bar{v}_1(x) + v^2/2$  satisfies Assumption [2.5](#page-2-5) (with  $\zeta = 2$  and with suitable  $\alpha, \beta$ ). Assumption [2.4](#page-2-2) also holds true. This example shows how the Lipschitz-continuity condition on *v* naturally arises in Assumption [2.4.](#page-2-2) It also shows that the most relevant case is where  $\zeta = 2$ .

Finally, we stipulate moment conditions on the volatility process and on the initial condition.

<span id="page-2-3"></span>**Assumption 2.7.** *Let*  $E[|V_0|^{\max{\{\xi,4\}}}] < \infty$  *holds for the ξ of Assumption [2.5.](#page-2-5)* 

<span id="page-2-6"></span>**Assumption 2.8.** *Let*  $E[|L_0|^2] < \infty$  *hold.* 

Our principal result is now presented.

<span id="page-2-0"></span>**Theorem 2.9.** *Let Assumptions [2.1,](#page-1-1) [2.2,](#page-2-1) [2.4,](#page-2-2) [2.5,](#page-2-5) [2.7](#page-2-3) and [2.8](#page-2-6) be in force. Then*

$$
\mathscr{L}(L_t, \mathbf{V}_t, \mathbf{R}_t) \to \mu_{\sharp}, \ t \to \infty \tag{5}
$$

*holds for some probability µ*<sup>♯</sup> *on* B(R *<sup>d</sup>*×<sup>W</sup> *<sup>d</sup>*×*d*×<sup>W</sup> *<sup>d</sup>*×*m*)*, in the sense of weak convergence of probability measures. The probability*  $\mu$ <sup> $\parallel$ </sup> *does not depend on*  $L_0$  *and it is invariant in the following sense: if*  $\mathscr{L}(L_0, \mathbf{V}_0, \mathbf{R}_0) = \mu_{\sharp}$  then  $\mathscr{L}(L_t, \mathbf{V}_t, \mathbf{R}_t) = \mu_{\sharp}$  for every  $t > 0$ .

In the following theorem, instead of Assumption [2.5](#page-2-5) one assumes the weaker condition [\(6\)](#page-3-3) below. This comes at the price of strengthening Assumptions [2.7](#page-2-3) and [2.8](#page-2-6) to [\(7\)](#page-3-4) below.

<span id="page-3-0"></span>**Theorem 2.10.** *Let Assumptions [2.2](#page-2-1) and [2.4](#page-2-2) hold, let*

<span id="page-3-3"></span>
$$
\langle x, \zeta(x, v) \rangle \le -\alpha |x|^{1+\gamma} + \beta (1+|v|^{\zeta}), \ x \in \mathbb{R}^d, \ v \in \mathcal{V}
$$
 (6)

*hold for some*  $\alpha, \beta > 0$ ,  $\xi \geq 2$  *and*  $0 < \gamma < 1$ *. Let us assume* 

<span id="page-3-4"></span>
$$
E\left[e^{\kappa_0|L_0|}\right]<\infty,\quad E\left[e^{\kappa_0|V_0|^{\xi/\gamma}}\right]<\infty\tag{7}
$$

<span id="page-3-1"></span>*for some*  $\kappa_0 > 0$ *. Then the conclusions of Theorem [2.9](#page-2-0) hold.* 

## **3 Coupling constructions**

Following the conventions of measure theory, the total variation norm of a finite signed measure *µ* on  $\mathscr{B}(\mathcal{I})$  is defined as

$$
||\mu||_{TV} := \sup_{\phi \in \Phi_1} \left| \int_{\mathcal{Z}} \phi(z) \mu(dz) \right|,
$$

where  $\Phi_1$  denotes the family of measurable functions  $\phi : \mathcal{Z} \to [-1,1]$ . The underlying  $\mathcal{Z}$  may vary but it will always be clear from the context. Note that for  $\mathcal{Z}\text{-value}$  random variables  $Z_1, Z_2$  we always have

<span id="page-3-5"></span>
$$
||\mathcal{L}(Z_1) - \mathcal{L}(Z_2)||_{TV} \le 2P(Z_1 \ne Z_2). \tag{8}
$$

#### <span id="page-3-2"></span>**3.1 Markov chains**

First we will work in the setting of general state space discrete-time Markov chains. Our main ideas will be explained in this simple context before turning to Markov chains in random environments in the next subsection.

Proofs for the stochastic stability of Markov chains are usually based on two ingredients, see e.g. [\[31\]](#page-21-3). First, it is checked (using Lyapunov functions) that the chain returns often enough to a fixed set *C*. Second, a minorization condition holds on *C* for the transition kernel so couplings occur whose probabilities can be estimated. Such *C* are called "small sets".

When the state space is  $\mathbb{R}^d$ , it happens often that *all* compact sets are small. This is the case for both discretized and discretely sampled non-degenerate diffusions. The coupling method of the present subsection exploits the latter property, formulated in more abstract terms. Otherwise we rely on standard "coupling from the past" ideas, see e.g. [\[37,](#page-21-13) [10\]](#page-20-10).

Although Theorem [3.4](#page-4-0) below seems to be new, its statement contains little revelation. Its proof, on the contrary, presents original ideas which will become fruitful in the more general setting of the next subsection where existing results do not apply. We will construct couplings on *a sequence of* small sets and then exploit (assuming a certain form of tightness) that the chain stays in these sets with large enough probabilities. The crucial methodological contribution of this approach is that, instead of analysing return times to a set *C* (which have a complicated dependence structure due to the random environment), one can repeatedly use simple, one-step estimates.

Another approach based on one-step estimates was presented in [\[23\]](#page-21-14), using a contraction argument in a suitable metric. When applying it in the presence of the random environment, however, the metric to be used becomes dependent on that environment which sets limitations to the use of that method, see [\[16\]](#page-21-15).

Let X be a Polish space. Let  $Q(\cdot, \cdot)$  be a probabilistic kernel, i.e.  $Q(\cdot, A)$  is measurable for each  $A \in \mathcal{B}(\mathcal{X})$  and  $Q(x, \cdot)$  is a probability law for each  $x \in \mathcal{X}$ . Let  $X_t$ ,  $t \in \mathbb{N}$  denote a Markov chain with transition kernel  $Q$ , started from some  $X_0$ . We now define the set of initial laws starting from which the chain satisfies a tightness-like assumption. We assume in the sequel that we are given a non-decreasing sequence of sets  $\mathscr{X}_n \in \mathscr{B}(\mathscr{X})$ ,  $n \in \mathbb{N}$  with  $\mathscr{X}_0 \neq \emptyset$ .

**Definition 3.1.** Let  $\mathcal{P}_b$  denote the set of probabilities  $\mu$  on  $\mathcal{B}(\mathcal{X})$  such that if  $X_0$  has law  $\mu$  then

$$
\lim_{n\to\infty}\sup_{t\in\mathbb{N}}P(X_t\notin\mathscr{X}_n)=0.
$$

Notice that  $\mathscr{P}_b$  might well be empty. We will write  $X_0 \in \mathscr{P}_b$  when we indeed mean  $\mathscr{L}(X_0) \in \mathscr{P}_b$ . We stipulate next that minorization conditions should hold on *each* of the sets  $\mathscr{X}_n$ .

<span id="page-4-2"></span>**Assumption 3.2.** *There exists a sequence*  $\alpha_n \in (0,1]$ *,*  $n \in \mathbb{N}$  and a sequence of probability measures *<sup>ν</sup>n, <sup>n</sup>* <sup>∈</sup> <sup>N</sup> *such that*

<span id="page-4-1"></span>
$$
Q(x,A) \ge \alpha_n v_n(A), \ A \in \mathcal{B}(\mathcal{X}), \ x \in \mathcal{X}_n, \ n \in \mathbb{N}.
$$

We recall a representation result for kernels satisfying the minorization condition [\(9\)](#page-4-1), in terms of random mappings that are constant on the respective  $\mathscr{X}_n$  with probability at least  $\alpha_n$ .

<span id="page-4-3"></span>**Lemma 3.3.** *Let Assumption [3.2](#page-4-2) be in force. Let* **U** *be a uniform random variable on* [0,1]*. For each*  $n \in \mathbb{N}$ , there exists a mapping  $T^n(\cdot,\cdot): [0,1] \times \mathcal{X} \to \mathcal{X}$  satisfying

$$
Q(x, A) = P(T(\mathbf{U}, x) \in A), \ x \in \mathcal{X}, \ A \in \mathcal{B}(\mathcal{X}),
$$

*such that for all*  $u \in [0, \alpha_n]$ ,

$$
T^n(u, x_1) = T^n(u, x_2) \quad \text{for all} \quad x_1, x_2 \in \mathcal{X}_n. \tag{10}
$$

*Proof.* Such a representation is well-known, see page 228 in [\[7\]](#page-20-11).

<span id="page-4-0"></span>**Theorem 3.4.** *Let Assumption* [3.2](#page-4-2) *hold. Then there exists a probability*  $\mu_*$  *on*  $\mathcal{B}(\mathcal{X})$  *such that* 

$$
||\mathcal{L}(X_t) - \mu_*||_{TV} \to 0, \ t \to \infty \tag{11}
$$

 $\Box$ 

*holds for every*  $X_0 \in \mathcal{P}_b$ .

*Proof.* Theorem [3.4](#page-4-0) follows from Theorem [3.10](#page-8-0) below (choosing  $\mathcal Y$  a singleton). Nonetheless we provide a proof in the present, simple setting, in order to elucidate the main ideas.

Fix  $\varepsilon > 0$  and choose  $n = n(\varepsilon)$  so large that

<span id="page-4-4"></span>
$$
\sup_{t \in \mathbb{N}} P(X_t \notin \mathcal{X}_n) \le \varepsilon. \tag{12}
$$

We will estimate coupling probabilities on  $\mathscr{X}_n$ , using independent copies of the random mappings constructed in Lemma [3.3](#page-4-3) above.

Let **,**  $k \in -N$  **be an independent sequence of uniform random variables on [0,1], independent** of  $X_0$ . Let  $T^n(\cdot, \cdot)$  be the mapping constructed in Lemma [3.3.](#page-4-3) Define the process

 $\tilde{X}_t$  := [ $T^n(\mathbf{U}_0, \cdot) \circ \cdots \circ T^n(\mathbf{U}_{-t+1}, \cdot)](X_0), t \in \mathbb{N}$ 

where we mean  $\tilde{X}_0 = X_0$ . Notice that  $\mathscr{L}(\tilde{X}_t) = \mathscr{L}(X_t)$  for each  $t \in \mathbb{N}$ .

Fix integers  $1 \leq s \leq t$ . For each  $j = 0, \ldots, s$ , define the following disjoint events:

$$
A_j^{s,t} := \{ [T^n(\mathbf{U}_{-j}, \cdot) \circ \cdots \circ T^n(\mathbf{U}_{-t+1}, \cdot)](X_0) = [T^n(\mathbf{U}_{-j}) \circ \dots \circ T^n(\mathbf{U}_{-s+1}, \cdot)](X_0) \},
$$
  
\n
$$
B_j^{s,t} := \{ [T^n(\mathbf{U}_{-j}, \cdot) \circ \cdots \circ T^n(\mathbf{U}_{-t+1}, \cdot)](X_0) \neq [T^n(\mathbf{U}_{-j}, \cdot) \circ \dots \circ T^n(\mathbf{U}_{-s+1}, \cdot)](X_0),
$$
  
\n
$$
[T^n(\mathbf{U}_{-j}, \cdot) \circ \cdots \circ T^n(\mathbf{U}_{-t+1}, \cdot)](X_0) \in \mathcal{X}_n, [T^n(\mathbf{U}_{-j}, \cdot) \circ \dots \circ T^n(\mathbf{U}_{-s+1}, \cdot)](X_0) \in \mathcal{X}_n \},
$$
  
\n
$$
C_j^{s,t} := \Omega \setminus (A_j^{s,t} \cup B_j^{s,t}),
$$

where we mean

$$
A_s^{s,t} := \{ [T^n(\mathbf{U}_{-s}, \cdot) \circ \cdots \circ T^n(\mathbf{U}_{-t+1}, \cdot)](X_0) = X_0 \},
$$
  
\n
$$
B_s^{s,t} := \{ [T^n(\mathbf{U}_{-s}, \cdot) \circ \cdots \circ T^n(\mathbf{U}_{-t+1}, \cdot)](X_0) \neq X_0, [T^n(\mathbf{U}_{-s}, \cdot) \circ \cdots \circ T^n(\mathbf{U}_{-t+1}, \cdot)](X_0) \in \mathcal{X}_n, X_0 \in \mathcal{X}_n \}.
$$

Define also  $p^{s,t}$  $j^{s,t} := P(A_j^{s,t})$  $j^{s,t}$ ). We aim to show that, for *s* large,  $p_0^{s,t}$  $\int_0^{s,t}$  is close to 1 for each  $t > s$ , which means that  $\tilde{X}_t$  very likely equals  $\tilde{X}_s$ . We will estimate  $p_j^{s,t}$  $j_j^{s,t}$  by backward recursion. Notice that

<span id="page-4-5"></span>
$$
P(C_j^{s,t}) \leq P([T^n(\mathbf{U}_{-j},\cdot)\circ\cdots\circ T^n(\mathbf{U}_{-t+1},\cdot)](X_0) \notin \mathcal{X}_n) + P([T^n(\mathbf{U}_{-j},\cdot)\circ\ldots\circ T^n(\mathbf{U}_{-s+1},\cdot)](X_0) \notin \mathcal{X}_n)
$$
  
= 
$$
P(X_{t-j} \notin \mathcal{X}_n) + P(X_{s-j} \notin \mathcal{X}_n) \leq 2\varepsilon,
$$
 (13)

by [\(12\)](#page-4-4). Define  $\mathcal{H}_{j,t} := \sigma(X_0, \mathbf{U}_{-j}, \dots, \mathbf{U}_{-t+1})$ . On the event  $B_j^{s,t}$  $j^{s,t} \in \mathscr{H}_{j,t}$  we have

$$
P\left(A_{j-1}^{s,t} \mid \mathcal{H}_{j,t}\right) \ge P\left(\mathbf{U}_{-j+1} \in [0,\alpha_n] \mid \mathcal{H}_{j,t}\right) = P(\mathbf{U}_{-j+1} \in [0,\alpha_n]) = \alpha_n \text{ a.s.}
$$

since  $T^n(\mathbf{U}_{-j+1},\cdot)$  is a constant mapping on  $\mathcal{X}_n$  when  $\mathbf{U}_{-j+1}^1 \in [0,\alpha_n]$ , and  $\mathbf{U}_{-j+1}$  is independent of  $\mathscr{H}_{j,t}$ . On the other hand, on the event  $A_j^{s,t}$  $\mathcal{F}^{s,t}_j \in \mathcal{H}_{j,t}$  we have  $P\left(A^{s,t}_{j-1}\right)$  $j_{j-1}^{s,t}$  *√* $H_{j,t}$  = 1 a.s. for trivial reasons. Hence

<span id="page-5-0"></span>
$$
p_{j-1}^{s,t} \ge p_j^{s,t} + \alpha_n P(B_j^{s,t}) \ge p_j^{s,t} + \alpha_n (1 - p_j^{s,t} - 2\varepsilon),\tag{14}
$$

using [\(13\)](#page-4-5). We get by backward recursion using [\(14\)](#page-5-0), starting from the trivial  $p_s^{s,t} \ge 0$ , that

$$
p_0^{s,t} \ge (1 - 2\varepsilon)\alpha_n \frac{1 - (1 - \alpha_n)^s}{1 - (1 - \alpha_n)} = (1 - 2\varepsilon)[1 - (1 - \alpha_n)^s],
$$

remembering also the formula for the sum of a geometric series. It follows from [\(8\)](#page-3-5) that for all integers  $1 \leq s \leq t$ .

<span id="page-5-3"></span>
$$
||\mathcal{L}(X_t) - \mathcal{L}(X_s)||_{TV} \le 2P(\tilde{X}_t \ne \tilde{X}_s) = 2(1 - p_0^{s,t}) \le 4\varepsilon + 2(1 - \alpha_n)^s,
$$
\n(15)

which is smaller than 5*ε* for *s* large enough. As *ε* was arbitrary, the sequence  $\mathscr{L}(X_t)$ ,  $t \in \mathbb{N}$  is shown to be Cauchy for the total variation distance hence it converges to some probability  $\mu_*$ .

Let  $X_t$ ,  $X'_t$ ,  $t \in \mathbb{N}$  denote Markov chains with transition kernel *Q*, started from  $X_0, X'_0 \in \mathcal{P}_b$ , respectively. Then, using  $U_k$ ,  $k \in -\mathbb{N}$  independent of  $\sigma(X_0, X'_0)$ , we get  $||\mathcal{L}(X_t) - \mathcal{L}(X'_t)||_{TV} \to 0$ as *t* → ∞ analogously to the argument above. This shows that  $\mu_*$  is independent of the choice of  $X_0 \in \mathcal{P}_h$ .  $X_0 \in \mathscr{P}_b$ .

<span id="page-5-2"></span>**Remark 3.5.** Assume  $\mathcal{X} := \mathbb{R}^d$  and  $\mathcal{X}_n := \{x \in \mathcal{X} : |x| \le n\}$ ,  $n \in \mathbb{N}$ . Let  $V(x) := g(|x|)$  for some nondecreasing  $g : \mathbb{R}_+ \to \mathbb{R}_+$  with  $g(\infty) = \infty$ . If the initial state  $X_0$  is such that

<span id="page-5-5"></span>
$$
\sup_{k \in \mathbb{N}} E[V(X_k)] < \infty,\tag{16}
$$

then  $X_0 \in \mathcal{P}_b$ , as seen from Markov's inequality.

The argument for proving Theorem [3.4](#page-4-0) above, in fact, provides us with a convergence rate estimate, too. For each *t*, [\(17\)](#page-5-1) below allows to optimize over *n* and to choose  $n = n(t)$  that gives the best estimate.

<span id="page-5-4"></span>**Corollary 3.6.** *Under Assumption* [3.2,](#page-4-2) *in the setting of Remark* [3.5,](#page-5-2) *for each*  $n \in \mathbb{N}$  and  $t \in \mathbb{N}$ ,

<span id="page-5-1"></span>
$$
||\mathcal{L}(X_t) - \mu_*||_{TV} \le 4 \frac{\sup_{k \in \mathbb{N}} E[V(X_k)]}{g(n)} + 2(1 - \alpha_n)^t.
$$
 (17)

*Proof.* This follows from [\(12\)](#page-4-4), [\(15\)](#page-5-3) and from Markov's inequality.

$$
\Box
$$

We demonstrate the application of Corollary [3.6](#page-5-4) and the resulting rate through a simple example.

**Example 3.7.** Consider a stable scalar AR(1) process, where  $\mathcal{X} = \mathbb{R}$  and the dynamics is

<span id="page-5-6"></span>
$$
X_{t+1} = \gamma X_t + \varepsilon_{t+1},\tag{18}
$$

where  $0 < \gamma < 1$ ,  $\varepsilon_t$  is an independent series of standard Gaussian variables, and  $X_0$  is a constant initialization.

In order to apply Corollary [3.6,](#page-5-4) we choose  $V(x) = g(|x|) = e^{\beta x^2}$  with  $\beta < \frac{1-\gamma^2}{2}$  $\frac{y}{2}$ . To confirm [\(16\)](#page-5-5), expanding the dynamics equation [\(18\)](#page-5-6) we see

$$
X_t = \gamma^t X_0 + \sum_{s=1}^t \gamma^{t-s} \varepsilon_s \quad \sim \quad \mathcal{N}\left(\gamma^t X_0, \frac{1-\gamma^{2t}}{1-\gamma^2}\right).
$$

Consequently,

$$
EV(X_t) = \frac{1}{\sqrt{2\pi \frac{1-\gamma^{2t}}{1-\gamma^2}}} \int_{-\infty}^{\infty} e^{-\frac{1-\gamma^2}{2(1-\gamma^{2t})}(z-\gamma^t X_0)^2} e^{\beta z^2} dz
$$
  

$$
\leq \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{1-\gamma^2}{2}(z-\gamma^t X_0)^2} e^{\beta z^2} dz < \infty,
$$

and this quantity is also bounded above uniformly in *t* by some  $c(\gamma, \beta, X_0)$  since  $|\gamma^t X_0|$  decreases as  $t \rightarrow \infty$ .

We also need Assumption [3.2,](#page-4-2) the minorization condition for a sequence of small sets. Let

$$
\mathscr{X}_n = [-n, n], \qquad v = \frac{1}{2} Leb|_{[-1,1]},
$$

for all *n*. In order to acquire  $\alpha_n$ , we need to find the infimum of  $\frac{dQ(x)}{dy}$  on the appropriate sets, and now that they are both absolutely continuous distributions, this boils down to comparing the densities, therefore

$$
\alpha_n = \inf_{x \in [-n,n], z \in [-1,1]} \frac{Q(x,dz)}{\frac{1}{2}dz} = \sqrt{\frac{2}{\pi}} e^{-\frac{(\gamma n + 1)^2}{2}}.
$$
\n(19)

Substituting the computed expressions Corollary [3.6](#page-5-4) provides

<span id="page-6-0"></span>
$$
||\mathcal{L}(X_t) - \mu_*||_{TV} \le \frac{4c(\gamma, \beta, X_0)}{\exp(\beta n^2)} + 2\left(1 - \sqrt{\frac{2}{\pi}}e^{-\frac{(\gamma n + 1)^2}{2}}\right)^t.
$$
 (20)

It remains to choose *n* depending on *t* to get the best bound possible. Clearly there is a tradeoff: for small values of *n*, the first term is weak while for large values of *n* the second term increases and can remain bounded away from 0.

Let us present the heuristics to find a near-optimal *n*. The second term in [\(20\)](#page-6-0) is approximately

$$
2\exp\left(-t\sqrt{\frac{2}{\pi}}\exp\left(-\frac{(\gamma n+1)^2}{2}\right)\right).
$$

We get the optimal bounds if the two terms agree (ignoring constants):

$$
\exp(-\beta n^2) = \exp\left(-t\sqrt{\frac{2}{\pi}}\exp\left(-\frac{(\gamma n+1)^2}{2}\right)\right),\
$$

$$
\log \beta + 2\log n = \log t + \frac{1}{2}\log\frac{2}{\pi} - \frac{(\gamma n+1)^2}{2}.
$$

It is easy to see that the value of  $\sqrt{2 \log t}$  $\frac{f_{\text{log}t}}{\gamma}$  is slightly too high for *n*. Still, inspired by this option we choose  $\lceil \sqrt{2} \rceil$ !

$$
n = \left\lceil \left( \frac{\sqrt{2}}{\gamma} - \eta \right) \sqrt{\log t} \right\rceil
$$

with some small  $\eta > 0$ . Using this choice in our bound [\(20\)](#page-6-0) and noting

$$
(\gamma n + 1)^2 \le \left(\gamma \left(\frac{\sqrt{2}}{\gamma} - \eta\right) \sqrt{\log t} + 2\right)^2
$$

we get

$$
||\mathcal{L}(X_t) - \mu_*||_{TV} \leq \frac{4c(\gamma,\beta,X_0)}{\exp\left(\beta\left(\frac{\sqrt{2}}{\gamma} - \eta\right)^2\log t\right)} + 2\left(1 - \sqrt{\frac{2}{\pi}}\exp\left[-\left(\left(1 - \frac{\gamma\eta}{\sqrt{2}}\right)\sqrt{\log t} + \sqrt{2}\right)^2\right]\right)^t.
$$

In the exponent of the first term we could choose the coefficient of the logarithm arbitrarily close to  $\frac{1-\gamma^2}{2}$  $\frac{(-\gamma^2)}{2} \left(\frac{\sqrt{2}}{\gamma}\right)^2 = \frac{1}{\gamma^2}$  $\frac{1}{\gamma^2}$  – 1. Although the second term looks daunting, observe that it has the order of  $(1 - t^{-1+\eta'})^t$  with some  $\eta' > 0$  therefore it has subpolynomial decay and is negligible compared to the first term.

Summing up, for a rate estimate we get that for any  $h > 0$  there is some constant  $C_h > 0$  such that

<span id="page-7-1"></span>
$$
||\mathcal{L}(X_t) - \mu_*||_{TV} \le \frac{C_h}{t^{\frac{1}{\gamma^2} - 1 - h}}.\tag{21}
$$

In the model [\(18\)](#page-5-6),  $||\mathcal{L}(X_t) - \mu_*||_{TV}$  decreases geometrically in *t* so only a suboptimal rate can be achieved by our method. Nevertheless, the estimates leading to [\(21\)](#page-7-1) are of great interest since they can serve as a basis for similar results for certain non-Markovian models, where power convergence rates are common, see e.g. [\[21\]](#page-21-12). One can thus treat models like [\(32\)](#page-10-0) below (which are not covered by current literature). Then, using technology from [\[16,](#page-21-15) [29\]](#page-21-16), various mixing properties and laws of large numbers (with rate estimates) can be established for functionals of the process  $X_t$ ,  $t \in \mathbb{N}$ . Central limit theorems can also be derived from mixing conditions, see [\[44\]](#page-22-4). These developments, however, are out of the scope of the present article.

#### <span id="page-7-0"></span>**3.2 Markov chains in random environments**

We now extend Theorem [3.4](#page-4-0) to Markov chains in random environments. These processes will still evolve in  $\mathscr{X} = \cup_{n \in \mathbb{N}} \mathscr{X}_n$  but their dynamics will be influenced by another random process we are just about to introduce. Let  $\mathscr Y$  be another Polish space and let  $Y_t$ ,  $t \in \mathbb Z$  be a (strict sense) stationary process in  $\mathscr Y$ . We assume that a non-decreasing sequence  $\mathscr Y_n \in \mathscr B(\mathscr Y)$ ,  $n \in \mathbb N$  is given with  $\mathscr Y_0 \neq \emptyset$ . Let  $Q : \mathcal{X} \times \mathcal{Y} \times \mathcal{B}(\mathcal{X}) \to [0,1]$  be a parametrized family of transition kernels, i.e.  $Q(\cdot,\cdot,A)$  is measurable for all  $A \in \mathcal{B}(\mathcal{X})$  and  $Q(x, y, \cdot)$  is a probability for all  $(x, y) \in \mathcal{X} \times \mathcal{Y}$ . We say that the process  $X_t$ ,  $t \in \mathbb{N}$ is a Markov chain in a random environment with transition kernel  $Q$  if it is an  $\mathscr X$ -valued stochastic process such that

$$
P(X_{t+1} \in A \mid \sigma(Y_j, \ j \in \mathbb{Z}; \ X_j, \ 0 \le j \le t)) = Q(X_t, Y_t, A), \ t \in \mathbb{N}.
$$
 (22)

Denote by  $\mathscr{M}_0$  the set of probability laws on  $\mathscr{X}\times\mathscr{Y}^{\mathbb{Z}}$  such that their second marginal equals the law of  $(Y_k)_{k\in\mathbb{Z}}$ . Let  $\mathcal{M}_b$  denote the set of those  $\mu \in \mathcal{M}_0$  for which the process  $X_t$ ,  $t \in \mathbb{N}$  started from  $X_0$  with  $L(X_0, (Y_k)_{k \in \mathbb{Z}}) = \mu$  satisfies

<span id="page-7-4"></span>
$$
\sup_{t \in \mathbb{N}} P(X_t \notin \mathcal{X}_n) \to 0, \ n \to \infty.
$$
 (23)

We will write  $X_0 \in \mathcal{M}_b$  in the sequel when we really mean  $\mathcal{L}(X_0,(Y_k)_{k \in \mathbb{Z}}) \in \mathcal{M}_b$ .

<span id="page-7-2"></span>**Assumption 3.8.** *Let*  $P(Y_0 \notin \mathcal{Y}_n) \to 0$  *hold as*  $n \to \infty$ *. There exists a sequence*  $\alpha_n \in (0,1]$ *,*  $n \in \mathbb{N}$  *and a sequence of probability measures*  $v_n$ ,  $n \in \mathbb{N}$  *such that for all*  $n \in \mathbb{N}$ *,* 

$$
Q(x,y,A)\geq \alpha_n v_n(A),\ A\in \mathcal{B}(\mathcal{X}),\ y\in \mathcal{Y}_n,\ x\in \mathcal{X}_n.
$$

A parametric version of Lemma [3.3](#page-4-3) comes next.

<span id="page-7-3"></span>**Lemma 3.9.** *Let Assumption [3.8](#page-7-2) be in force. Let* **U** *be a uniform random variable on* [0,1]*. For each*  $n \in \mathbb{N}$ , there exists a measurable mapping  $T^n(\cdot,\cdot,\cdot): [0,1] \times \mathcal{X} \times \mathcal{Y} \to \mathcal{X}$  satisfying  $Q(x, y, A) =$  $P(T^n(\mathbf{U}, x, y) \in A)$ ,  $x \in \mathcal{X}$ ,  $y \in \mathcal{Y}$ ,  $A \in \mathcal{B}(\mathcal{X})$  such that for all  $u \in [0, a_n]$ ,

$$
T^n(u,x_1,y)=T^n(u,x_2,y)\quad\text{for all}\quad x_1,x_2\in\mathscr{X}_n,\ y\in\mathscr{Y}_n.
$$

*Proof.* This is a straightforward extension of the case with  $\mathcal{Y}$  a singleton, that is, of Lemma [3.3](#page-4-3) above. See Lemma 7.1 of [\[29\]](#page-21-16).  $\Box$ 

The following abstract result serves as the basis of Section [4](#page-11-0) below. We do not know of any similar results in the literature. Existing papers have fairly restrictive assumptions: either Doeblin-like conditions (as in [\[26,](#page-21-17) [27,](#page-21-18) [39\]](#page-22-5)) or strong contractivity hypotheses (as in [\[40\]](#page-22-6)).

<span id="page-8-0"></span>**Theorem 3.10.** *Let* Assumption [3.8](#page-7-2) hold and let  $\mathcal{M}_b \neq \emptyset$ . Let  $X_t$ ,  $t \in \mathbb{N}$  denote a Markov chain in *a random environment with transition kernel Q, started from some*  $X_0 \in \mathcal{M}_b$ *. Then there exists a*  $p$ robability  $\mu_\sharp$  on  $\mathscr{B}( \mathscr{X} \times \mathscr{Y}^{\mathbb{N}})$  such that

$$
||\mathcal{L}(X_t,(Y_{t+k})_{k\in\mathbb{Z}}) - \mu_{\sharp}||_{TV} \to 0, \ t \to \infty.
$$
 (24)

*If*  $X'_t$ ,  $t \in \mathbb{N}$  *is another such Markov chain in random environment started from*  $X'_0$  ∈  $\mathcal{M}_b$  *then* 

<span id="page-8-2"></span>
$$
||\mathcal{L}(X_t,(Y_{t+k})_{k\in\mathbb{Z}}) - \mathcal{L}(X'_t,(Y_{t+k})_{k\in\mathbb{Z}})||_{TV} \to 0, t \to \infty.
$$
 (25)

*In particular,*  $\mu_{\sharp}$  does not depend on the choice of  $X_0 \in \mathcal{M}_b$ . The probability  $\mu_{\sharp}$  is invariant in the following sense: if  $X_0$  is such that  $\mathscr{L}(X_0,(Y_k)_{k\in\mathbb{Z}})=\mu_\sharp$  then  $\mathscr{L}(X_t,(Y_{t+k})_{k\in\mathbb{Z}})=\mu_\sharp$  for each  $t\in\mathbb{N}$ .

*Proof.* The core idea of the proof is identical to that of Theorem [3.4,](#page-4-0) with the extra task of checking whether the process  $Y$  stays in  $\mathcal{Y}_n$  for some suitable  $n$ . In order to prove invariance, however, here we need to construct  $\tilde X_\infty$  such that  $\tilde X_t$  (to be defined soon) converges to  $\tilde X_\infty$  a.s. in a stationary way (along a suitable subsequence). This requires a more complicated setup.

There exists a measurable function  $g: \mathscr{Y}^{\mathbb{Z}} \times [0,1] \to \mathscr{X}$  and a uniform [0,1]-valued random variable R, independent of  $\sigma(Y_k, k \in \mathbb{Z})$ , such that  $\mathscr{L}(X_0, (Y_k)_{k \in \mathbb{Z}}) = \mathscr{L}(g((Y_k)_{k \in \mathbb{Z}}, R), (Y_k)_{k \in \mathbb{Z}})$ . Let  $U_k$ , *k* ∈ −N be an independent family of uniform random variables on [0,1], independent of  $\sigma(R,(Y_k)_{k\in\mathbb{Z}})$ . Let  $T^n(\cdot,\cdot,\cdot)$ ,  $n \in \mathbb{N}$  be the mappings constructed in Lemma [3.9.](#page-7-3)

For each integer  $m \geq 1$  choose  $n(m) \in \mathbb{N}$  so large that

<span id="page-8-1"></span>
$$
P(Y_0 \notin \mathcal{Y}_{n(m)}) + \sup_{k \in \mathbb{N}} P(X_k \notin \mathcal{X}_{n(m)}) \le 1/2^m.
$$
 (26)

Let  $N(m) \ge 1$  be so large that  $(1 - a_{n(m)})^{N(m)} \le 1/2^m$ . Define  $M_0 := 0$ ,  $M_m := \sum_{j=1}^m N(j)$ . Define the following random mappings from  $\mathcal{X} \to \mathcal{X}$ , for each  $m \geq 1$ :

$$
\tilde{T}_m(\cdot) := T^{n(m)}(\mathbf{U}_{-M_{m-1}}, \cdot, Y_{-M_{m-1}-1}) \circ \dots \circ T^{n(m)}(\mathbf{U}_{-M_m+1}, \cdot, Y_{-M_m})
$$

and

$$
\mathbf{T}_m(\cdot) := \tilde{T}_1(\cdot) \circ \ldots \circ \tilde{T}_m(\cdot).
$$

Let  $\mathbf{T}_0$  be the identity mapping of  $\mathcal{X}$ .

Let  $\tilde{X}_0 := g((Y_k)_{k \in \mathbb{Z}}, R)$  and for each  $m \in \mathbb{N}$  and each  $M_m + 1 \le t \le M_{m+1}$ , define the process

$$
\tilde{X}_t := \mathbf{T}_m(\cdot) \circ T^{n(m+1)}(\mathbf{U}_{-M_m}, \cdot, Y_{-M_m-1}) \circ \dots \circ T^{n(m+1)}(\mathbf{U}_{-t+1}, \cdot, Y_{-t})(g((Y_{-t+k})_{k \in \mathbb{Z}}, R)).
$$

Notice that  $\mathscr{L}(\tilde{X}_t, (Y_k)_{k \in \mathbb{Z}}) = \mathscr{L}(X_t, (Y_{t+k})_{k \in \mathbb{Z}})$  by construction, for each  $t \in \mathbb{N}$ .

Fix  $m \geq 2$  and let  $M_m + 1 \leq t \leq M_{m+1}$  be arbitrary. For each  $j = M_{m-1},...,M_m$  we will define the following random variables:

$$
V_{j,t} := [T^{n(m)}(\mathbf{U}_{-j}, \cdot, Y_{-j-1}) \circ \cdots \circ T^{n(m)}(\mathbf{U}_{-M_m+1}, \cdot, Y_{-M_m}) \circ T^{n(m+1)}(\mathbf{U}_{-M_m}, \cdot, Y_{-M_m-1}) \circ \cdots \circ T^{n(m+1)}(\mathbf{U}_{-t+1}, \cdot, Y_{-t})](g((Y_{-t+k})_{k \in \mathbb{Z}}, R)),
$$
  
\n
$$
W_{j,t} := [T^{n(m)}(\mathbf{U}_{-j}, \cdot, Y_{-j-1}) \circ \cdots \circ T^{n(m)}(\mathbf{U}_{-M_m+1}, \cdot, Y_{-M_m})](g((Y_{-M_m+k})_{k \in \mathbb{Z}}, R)),
$$

with the understanding that

$$
W_{M_m,t}=g((Y_{-M_m+k})_{k\in\mathbb{Z}},R)
$$

and

$$
V_{M_m,t} := T^{n(m+1)}(\mathbf{U}_{-M_m}, \cdot, Y_{-M_m-1}) \circ \cdots \circ T^{n(m+1)}(\mathbf{U}_{-t+1}, \cdot, Y_{-t})(g((Y_{-t+k})_{k \in \mathbb{Z}}, R)).
$$

Consider the corresponding disjoint events

$$
A_{j,t} := \{V_{j,t} = W_{j,t}\},
$$
  
\n
$$
B_{j,t} := \{V_{j,t} \neq W_{j,t}, V_{j,t} \in \mathcal{X}_{n(m)}, W_{j,t} \in \mathcal{X}_{n(m)}, Y_{-j} \in \mathcal{Y}_{n(m)}\},
$$
  
\n
$$
C_{j,t} := \Omega \setminus (A_{j,t} \cup B_{j,t}).
$$

Define also  $p_{j,t} := P(A_{j,t}), j = M_{m-1},...,M_m$ . Notice that

<span id="page-9-0"></span>
$$
P(C_{j,t}) \leq P(V_{j,t} \notin \mathcal{X}_{n(m)}) + P(W_{j,t} \notin \mathcal{X}_{n(m)}) + P(Y_{-j} \notin \mathcal{Y}_{n(m)})
$$
  
=  $P(X_{t-j} \notin \mathcal{X}_{n(m)}) + P(X_{M_m - j} \notin \mathcal{X}_{n(m)}) + P(Y_{-j} \notin \mathcal{Y}_{n(m)})$   
 $\leq 1/2^{m-1},$  (27)

by the stationarity of the process *Y* and by [\(26\)](#page-8-1).

Define  $\mathcal{H}_{j,t} := \sigma((Y_k)_{k \in \mathbb{Z}}, R, \mathbf{U}_{-j}, \ldots, \mathbf{U}_{-t+1})$ . On  $B_{j,t} \in \mathcal{H}_{j,t}$  we have

 $P(A_{j-1,t} | \mathcal{H}_{j,t}) \ge P(\mathbf{U}_{-j+1} \in [0, \alpha_{n(m)}] | \mathcal{H}_{j,t}) = P(\mathbf{U}_{-j+1} \in [0, \alpha_{n(m)}]) = \alpha_{n(m)}$  a.s.

since  $T^{n(m)}(\mathbf{U}_{-j+1},\cdot,y)$  is a constant mapping on  $\mathscr{X}_{n(m)}$ , for each  $y \in \mathscr{Y}_{n(m)}$  when  $\mathbf{U}_{-j+1} \in [0,\alpha_{n(m)}],$ and  $\mathbf{U}_{-j+1}$  is independent of  $\mathcal{H}_{j,t}$ . On the other hand, on  $A_{j,t} \in \mathcal{H}_{j,t}$  we have  $P(A_{j-1,t}|\mathcal{H}_{j,t}) = 1$  a.s., trivially. Hence

$$
p_{j-1,t} \ge p_{j,t} + \alpha_{n(m)} P(B_{j,t}) \ge p_{j,t} + \alpha_{n(m)} (1 - p_{j,t} - 1/2^{m-1}),
$$
\n(28)

using [\(27\)](#page-9-0), which leads (by backward induction starting from  $p_{M_m,t} \ge 0$ ) to

$$
p_{M_{m-1},t} \ge (1 - 1/2^{m-1})[1 - (1 - \alpha_{n(m)})^{N(m)}],
$$

and eventually to

<span id="page-9-1"></span>
$$
P(\tilde{X}_t \neq \tilde{X}_{M_m}) \le P(V_{M_{m-1},t} \neq W_{M_{m-1},t}) = 1 - p_{M_{m-1},t} \le 1/2^{m-1} + (1 - \alpha_{n(m)})^{N(m)} \le 1/2^{m-2},\tag{29}
$$

remembering the choice of  $N(m)$ . These relations establish, in particular, that for the event

$$
A_m := \left\{ \tilde{X}_{M_j} = \tilde{X}_{M_m} \text{ for all } j \geq m \right\},\
$$

we have

<span id="page-9-2"></span>
$$
P(\Omega \setminus A_m) \le \sum_{j=m} \frac{1}{2^{j-2}} \le 1/2^{m-3}.\tag{30}
$$

We can thus define unambiguously  $\tilde{X}_{\infty} := \tilde{X}_{M_m}$  on  $A_m$  and, doing this for all  $m \geq 2$ , a random variable  $\tilde{X}_{\infty}$  gets almost surely defined. Clearly, for all  $M_m + 1 \le t \le M_{m+1}$ ,

$$
P(\tilde{X}_t \neq \tilde{X}_{\infty}) \le P(\tilde{X}_t \neq \tilde{X}_{M_m}) + P(\tilde{X}_{M_m} \neq \tilde{X}_{\infty}) \le 1/2^{m-4},
$$

by [\(29\)](#page-9-1) and [\(30\)](#page-9-2). Denoting by  $\mu_{\parallel}$  the law of  $(\tilde{X}_{\infty}, (Y_k)_{k \in \mathbb{Z}})$ ,

$$
||\mathcal{L}(X_t,(Y_{t+k})_{k\in\mathbb{Z}})-\mu_{\sharp}||_{TV}\leq 2P(\tilde{X}_t\neq \tilde{X}_{\infty})\rightarrow 0,~t\rightarrow\infty.
$$

Now we turn to proving [\(25\)](#page-8-2). In addition to  $\tilde{X}_t$ , let us also define  $\tilde{X}^\prime_t$  in the same manner with *g* replaced by  $g' : \mathscr{Y}^{\mathbb{Z}} \times [0,1] \to \mathscr{X}$  such that  $\mathscr{L}(X'_{0},(Y_{k})_{k \in \mathbb{Z}}) = \mathscr{L}(g'((Y_{k})_{k \in \mathbb{Z}}^{\prime},R), (Y_{k})_{k \in \mathbb{Z}})$ . We get by analogous arguments that

$$
||\mathcal{L}(X_t,(Y_{t+k})_{k\in\mathbb{Z}})-\mathcal{L}(X_t',(Y_{t+k})_{k\in\mathbb{Z}})||_{TV}=||\mathcal{L}(\tilde{X}_t,(Y_k)_{k\in\mathbb{Z}})-\mathcal{L}(\tilde{X}_t',(Y_k)_{k\in\mathbb{Z}})||_{TV}\rightarrow 0,\; t\rightarrow\infty.
$$

To see invariance, fix  $\varepsilon > 0$  and notice that for  $m = m(\varepsilon)$  large enough,

<span id="page-9-3"></span>
$$
P(\tilde{X}_{M_m} \neq \tilde{X}_{\infty}) + P(\tilde{X}_{M_m+1} \neq \tilde{X}_{\infty}) \leq \varepsilon.
$$
\n(31)

Let us take **U**∗ uniform on [0,1], independent of all the random objects that have appeared so far. We will use the mapping  $T^0(\cdot,\cdot,\cdot)$  below but  $T^n(\cdot,\cdot,\cdot)$  for any  $n$  would do equally well. Notice that

$$
\mathcal{L}(T^0(\mathbf{U}^*, \tilde{X}_{M_m}, Y_0), (Y_{1+k})_{k \in \mathbb{Z}}) = \mathcal{L}(\tilde{X}_{M_m+1}, (Y_k)_{k \in \mathbb{Z}})
$$

and then from [\(31\)](#page-9-3), necessarily,

$$
||\mathcal{L}(T^0(\mathbf{U}^*, \tilde{X}_{M_m}, Y_0), (Y_{1+k})_{k \in \mathbb{Z}}) - \mu_{\sharp}||_{TV} \leq 2\varepsilon.
$$

Now employing the limiting random variable representing  $\mu_\sharp,$ 

$$
\| \mathcal{L}(T^0(\mathbf{U}^*, \tilde{X}_{M_m}, Y_0), (Y_{1+k})_{k \in \mathbb{Z}}) - \mathcal{L}(T^0(\mathbf{U}^*, \tilde{X}_{\infty}, Y_0), (Y_{1+k})_{k \in \mathbb{Z}})) \|_{TV}
$$
  
\n
$$
\leq 2P(T^0(\mathbf{U}^*, \tilde{X}_{M_m}, Y_0) \neq T^0(\mathbf{U}^*, \tilde{X}_{\infty}, Y_0))
$$
  
\n
$$
\leq 2P(\tilde{X}_{M_m} \neq \tilde{X}_{\infty}) \leq 2\varepsilon.
$$

Thus we have

$$
||\mathcal{L}(T^0(\mathbf{U}^*, \tilde{X}_{\infty}, Y_0), (Y_{1+k})_{k \in \mathbb{Z}}) - \mu_{\sharp}||_{TV} \leq 4\varepsilon
$$

and, as *ε* was arbitrary,  $\mathscr{L}(T^0(\mathbf{U}^*, \tilde{X}_{\infty}, Y_0), (Y_{1+k})_{k \in \mathbb{Z}}) = \mu_{\sharp}$  follows. Clearly, this means that

$$
\mathcal{L}(X_0,(Y_k)_{k\in\mathbb{Z}})=\mu_\sharp \text{ implies } \mathcal{L}(X_1,(Y_{1+k})_{k\in\mathbb{Z}})=\mu_\sharp
$$

and the latter extends immediately to  $\mathscr{L}(X_t,(Y_{t+k})_{k\in\mathbb{Z}}) = \mu_{\sharp}$  for all  $t \geq 2$ , too. The proof is complete.

Before transitioning to the analysis of continuous-time processes, let us demonstrate the application of Theorem [3.10](#page-8-0) on a benchmark model: the discrete-time counterpart of [\(1\)](#page-0-0) with log-Gaussian *Vt* . We take the simplest mean-reverting drift, but the same argument applies under more general dissipativity conditions.

**Example 3.11.** Consider the following model for financial time series. Let  $\eta_t, t \in \mathbb{Z}$  be independent standard Gaussian random variables and

$$
Z_t = \sum_{k=0}^{\infty} a_k \eta_{t-k},
$$

a causal moving average with constants  $a_k$ ,  $k \in \mathbb{N}$  satisfying  $\sum_k a_k^2 < \infty$ . Therefore  $Z_t$  is almost surely well defined and is a stationary Gaussian process. *Z<sup>t</sup>* represents the log-volatility of an asset's logprice  $X_t$  which in turn is defined as

<span id="page-10-0"></span>
$$
X_{t+1} = \gamma X_t + \rho e^{Z_t} \eta_{t+1} + \sqrt{1 - \rho^2} e^{Z_t} \varepsilon_{t+1},
$$
\n(32)

where  $\gamma \in (0,1)$ ,  $\rho \in (-1,1)$  and  $\varepsilon_k$ ,  $k \in \mathbb{N}$  is an i.i.d. series of zero-mean random variables, also independent of  $\eta_t, t \in \mathbb{Z}$ .

For the  $\varepsilon_k$  we assume they have finite variance and have a positive density  $f(x)$  such that for all *n* ∈ N, inf<sub>*x*∈[-*n*,*n*]</sub>  $f(x) = c(n) > 0$ . Additionally, we assume the initial price  $X_0$  has finite variance and is independent of  $\eta_t, t \in \mathbb{Z}, \varepsilon_k, k \in \mathbb{N}$ .

We claim that under these natural assumptions Theorem [3.10](#page-8-0) is applicable to the model [\(32\)](#page-10-0). First of all, the random environment is defined as  $Y_t := (Z_t, \eta_{t+1})$ . We choose

 $\mathscr{X}_n = \{x \in \mathbb{R} : |x| \leq n\}$   $\mathscr{Y}_n = \{(z, \eta) \in \mathbb{R}^2 : |z|, |\eta| \leq n\}.$ 

We first verify Assumption [3.8,](#page-7-2) fix some  $n \in \mathbb{N}$  and  $v_n = \frac{1}{2} Leb|_{[-1,1]}$ . Now that we are working with absolutely continuous distributions, we have to find a lower bound of the transition density to  $[-1,1]$  from any departure point  $X_t \in \mathcal{X}_n$ ,  $(Z_t, \eta_{t+1}) \in \mathcal{Y}_n$ .

Rearranging [\(32\)](#page-10-0), we get

$$
\varepsilon_{t+1} = \frac{X_{t+1} - \gamma X_t}{\sqrt{1 - \rho^2} e^{Z_t}} + \frac{\rho}{\sqrt{1 - \rho^2}} \eta_{t+1}.
$$

Requiring  $X_{t+1}$  to arrive in [−1,1], knowing  $X_t, \eta_{t+1} \in [-n, n]$ ,  $e^{Z_t} \in [e^{-n}, e^n]$ , the possible needed values of  $\varepsilon_{t+1}$  are restricted within some bounded interval  $[-d(n),d(n)]$ . Using the condition on the bounded positivity of the density  $f(x)$  of  $\varepsilon_{t+1}$  we get a valid minorization with

$$
\alpha_n = 2 \inf_{x \in [-d(n), d(n)]} f(x) = 2c(d(n)) > 0.
$$

It is left to confirm that  $X_0 \in \mathcal{M}_b$ , so that  $X_t$  uniformly rarely leaves the small sets. By recursively using [\(32\)](#page-10-0) we may express  $X_t$  as follows:

$$
X_{t} = \sum_{s=1}^{t} \gamma^{t-s} e^{Z_{s-1}} \left( \rho \eta_{s} + \sqrt{1 - \rho^{2}} \varepsilon_{s} \right) + \gamma^{t} X_{0}.
$$
 (33)

To bound  $X_t$ , we compute  $E[X_t^2]$ . Observe that when evaluating the square of this sum, all crossterms cancel when taking expectation, even the ones only involving *Z* and *η*. Consequently,

$$
E[X_t^2] = \sum_{s=1}^t \gamma^{2t-2s} E\left[e^{2Z_{s-1}}\right] (\rho^2 E[\eta_s^2] + (1-\rho^2)E[\varepsilon_s^2]) + \gamma^{2t} E[X_0^2].
$$

Regarding these terms, remember that *Z<sup>t</sup>* was Gaussian thus it has finite exponential moments and all appearing variables had finite variances. Moreover, due to the stationarity of all components appearing, we have the time-independent bound

$$
E[X_t^2] \le \frac{1}{1-\gamma^2} E\left[e^{2Z_0}\right] (\rho^2 E[\eta_1^2] + (1-\rho^2)E[\varepsilon_1^2]) + E[X_0^2] =: K < \infty.
$$

From here we can conveniently bound

$$
\sup_{t\in\mathbb{N}} P(|X_t|>n) \leq \frac{K}{n^2},
$$

which indeed converges to 0 as  $n \to \infty$ . This reasoning shows that  $\mathscr{L}(X_0,(Z_k,\eta_{k+1})_{k \in \mathbb{Z}}) \in \mathscr{M}_b$ . We have verified the minorization Assumption [3.8](#page-7-2) just before so Theorem [3.10](#page-8-0) applies, ensuring convergence in total variation. The present example complements Example 3.4 of [\[16\]](#page-21-15) where convergence in total variation was established under stronger assumptions (but with a rate estimate).

## <span id="page-11-0"></span>**4 Proofs in continuous time**

Until finishing the proof of Theorem [2.9,](#page-2-0) we assume that all the hypotheses of that theorem are in force. Let us first establish a simple continuity property.

<span id="page-11-2"></span>**Lemma 4.1.** When  $s \to 0$ ,  $\sup_{t \in \mathbb{R}} \{E[\mathbf{d}_{d \times d}(\mathbf{V}_{t+s}, \mathbf{V}_t)] + E[\mathbf{d}_{d \times m}(\mathbf{R}_{t+s}, \mathbf{R}_t)]\} \to 0$  holds true.

*Proof.* By stationarity of *V*, *ρ*, this amounts to proving  $E[\mathbf{d}_{d \times d}(\mathbf{V}_s, \mathbf{V}_0)] + E[\mathbf{d}_{d \times m}(\mathbf{R}_s, \mathbf{R}_0)] \to 0$ . The process *V* has trajectories that are uniformly continuous on compacts hence  $\sup_{u \in [i,j+1]} |V_{u+s} - V_u| \to$ 0 almost surely as *s* → 0, for each *i* ∈ Z. Then  $E[1 \wedge \sup_{u \in [i, i+1]} |V_{u+s} - V_u|] \to 0$  for each *i* and, finally,  $E[\mathbf{d}_{d \times d}(\mathbf{V}_s, \mathbf{V}_0)] \to 0$  by the definition of **d**. We argue in the same manner for **R**.  $E[\mathbf{d}_{d \times d}(\mathbf{V}_s, \mathbf{V}_0)] \rightarrow 0$  by the definition of **d**. We argue in the same manner for **R**.

Now let us prove a moment estimate.

<span id="page-11-1"></span>**Lemma 4.2.** *We have*  $\tilde{L} := \sup_{t \in \mathbb{R}_+} E[|L_t|^2] < \infty$ *.* 

*Proof.* Fix  $k \in \mathbb{N}$ , for the moment. Define the stopping times  $\tau_l := \inf\{t > k : |L_t| > l\}$  for  $l \in \mathbb{N}$ . Itô's formula and Assumption [2.5](#page-2-5) imply that, for all  $k \le t \le k + 1$ ,

$$
e^{2\alpha(t\wedge\tau_{l}-k)}|L_{t\wedge\tau_{l}}|^{2} = |L_{k}|^{2} + \int_{k}^{t\wedge\tau_{l}} 2e^{2\alpha(s-k)}\langle L_{s},\zeta(L_{s},V_{s})\rangle ds + \int_{k}^{t\wedge\tau_{l}} 2e^{2\alpha(s-k)}L_{s}^{*}V_{s}d\overline{W}_{s}
$$
  
+ 
$$
\int_{k}^{t\wedge\tau_{l}} e^{2\alpha(s-k)}tr(V_{s}^{*}V_{s})ds + \int_{k}^{t\wedge\tau_{l}} 2\alpha e^{2\alpha(s-k)}|L_{s}|^{2}ds
$$
  

$$
\leq |L_{k}|^{2} - \int_{k}^{t\wedge\tau_{l}} 2\alpha e^{2\alpha(s-k)}|L_{s}|^{2}ds + \int_{k}^{k+1} 2e^{2\alpha(s-k)}\beta(1+|V_{s}|^{\zeta})ds
$$
  
+ 
$$
\int_{k}^{t\wedge\tau_{l}} 2e^{2\alpha(s-k)}L_{s}^{*}V_{s}d\overline{W}_{s} + e^{2\alpha} \int_{k}^{k+1} d|V_{s}|^{2}ds
$$
  
+ 
$$
\int_{k}^{t\wedge\tau_{l}} 2\alpha e^{2\alpha(s-k)}|L_{s}|^{2}ds
$$
  

$$
\leq |L_{k}|^{2} + \int_{k}^{k+1} 2e^{2\alpha}\beta(1+|V_{s}|^{\zeta})ds
$$
  
+ 
$$
\int_{k}^{t\wedge\tau_{l}} 2e^{2\alpha(s-k)}L_{s}^{*}V_{s}d\overline{W}_{s} + e^{2\alpha} \int_{k}^{k+1} d|V_{s}|^{2}ds,
$$

where  $\overline{W}_t = \int_0^t \rho_s dB_s + \int_0^t \sqrt{I - \rho_s \rho_s^*} dW_s$  is a *d*-dimensional standard Brownian motion. Taking expectations and noting the martingale property of the stochastic integral,

$$
E[e^{2\alpha(t\wedge\tau_l-k)}|L_{t\wedge\tau_l}|^2] \leq E[|L_k|^2] + \int_k^{k+1} e^{2\alpha}(2\beta+d)(1+E[|V_s|^{\xi}])ds.
$$

Noting stationarity of *V* and applying Fatou's lemma,

<span id="page-12-0"></span>
$$
E[|L_t|^2] \le e^{-2\alpha(t-k)}E[|L_k|^2] + e^{2\alpha}(2\beta + d)(1 + E[|V_0|^{\xi}]).
$$
\n(34)

Setting  $t = k + 1$ , Assumption [2.8](#page-2-6) and an induction on *k* show that  $E[|L_k|^2] < \infty$  for all  $k \in \mathbb{N}$ , in fact,  $\sup_k E[|L_k|^2] < \infty$ . Finally, also  $\sup_{t \geq 0} E[|L_t|^2] < \infty$  follows from [\(34\)](#page-12-0).

Let  $\mathbf{C}_k$  denote the Banach space of continuous  $\mathbb{R}^k$ -valued functions on [0,1] equipped with the usual maximum norm  $|| \cdot ||_{\mathbf{C}_k}$ . The family of functions in  $\mathbf{C}_{d \times d}$  whose values are non-singular is denoted by **C**+. We further define

$$
\mathbf{C}^1 := \{ \mathbf{r} \in \mathbf{C}_{d \times m} : \mathbf{r}_t \mathbf{r}_t^* \leq I, \ t \in [0,1] \}
$$

as well as

$$
\mathbf{C}^{1+} := \{\mathbf{r} \in \mathbf{C}_{d \times m} : \mathbf{r}_t \mathbf{r}_t^* < I, \ t \in [0,1]\}.
$$

The auxiliary process to be defined in [\(35\)](#page-12-1) below plays a key role in our arguments. The parameters **v**,**r** represent the "frozen" values of trajectories of the volatility and correlation processes, while **z** will be a generic value of the stochastic integral of  $V\rho$  with respect to *B*.

For each  $\mathbf{v} \in \mathbf{C}_{d \times d}$ ,  $\mathbf{z} \in \mathbf{C}_d$ ,  $\mathbf{r} \in \mathbf{C}^1$  and  $x \in \mathbb{R}^d$ , let  $\tilde{X}_t(\mathbf{v}, \mathbf{z}, \mathbf{r}, x)$ ,  $t \in [0, 1]$  denote the unique  $\mathscr{F}_t$ . adapted solution of the SDE

<span id="page-12-1"></span>
$$
d\tilde{X}_t(\mathbf{v},\mathbf{z},\mathbf{r},x) = \zeta \left( \tilde{X}_t(\mathbf{v},\mathbf{z},\mathbf{r},x) + \mathbf{z}_t, \mathbf{v}_t \right) dt + \mathbf{v}_t \sqrt{I - \mathbf{r}_t \mathbf{r}_t^*} dW_t, \ \tilde{X}_0(\mathbf{v},\mathbf{z},\mathbf{r},x) = x,\tag{35}
$$

which exists e.g. by Theorem 7 on page 82 of [\[28\]](#page-21-11). We shall use the shorthand notation  $\mathbf{q} := (\mathbf{v}, \mathbf{z}, \mathbf{r}, x)$ in the sequel. Introduce also the space  $\mathcal{Y} := \mathbf{C}_{d \times d} \times \mathbf{C}_d \times \mathbf{C}^1$  where the random environment (to be defined in [\(37\)](#page-13-0) below) will evolve.

In line with the notations of the standard reference work [\[35\]](#page-21-19),  $\mathbb{D}^{k,p}$  denotes the  $p$ -Sobolev space of *k* times Malliavin differentiable functionals. The first and second Malliavin derivative of a functional *F* will be denoted by  $DF$  ,  $D^2F$  or  $D_rF$  ,  $D^2_{r_1,r_2}F$  when we need to emphasize that these are random processes/fields indexed by *r*, *r*1, *r*2. The Skorokhod integral operator (the adjoint of *D*) is denoted by  $\delta$ . The notation *H* refers to the Hilbert-space of square-integrable  $\mathbb{R}^d$ -valued functions on [0,1].

For  $F = (F^1, \ldots, F^d)$  with  $F^i \in \mathbb{D}^{1,2}$ ,  $i = 1, \ldots, d$  the corresponding Malliavin matrix  $\sigma(F)$  is defined as

$$
\sigma(F)_{ij} = \sum_{l=1}^d \int_0^1 D_s^{(l)} F^i D_s^{(l)} F^j ds,
$$

where  $D^{(l)}F^i$  denotes the *l*th coordinate of  $DF^i$ . In the sequel, the notation  $L^p$  refers to the usual space of *p*-integrable real-valued random variables, for  $p \ge 1$ . We define  $\gamma := \sigma^{-1}$  on the event where  $\sigma$  is invertible and 0 otherwise.

<span id="page-12-3"></span>**Lemma 4.3.** For each  $\mathbf{q} \in \mathcal{Y} \times \mathbb{R}^d$ , we have  $\tilde{X}_1(\mathbf{q}) \in \cap_{p \geq 1} \mathbb{D}^{2,p}$ ,  $D\tilde{X}_1(\mathbf{q})$  and  $D^2\tilde{X}_1(\mathbf{q})$  are bounded. *Furthermore, if*  $\mathbf{v} \in \mathbf{C}^+$  *and*  $\mathbf{r} \in \mathbf{C}^{1+}$  *then*  $\gamma$  *is uniformly bounded, in particular,*  $1/\text{det}(\sigma(\tilde{X}_1(\mathbf{q}))) \in$  $∩_{p≥1}L^p$  *holds.* 

*Proof.* The first statement follows from the proof of Theorem 2.2.2 of [\[35\]](#page-21-19) which applies in the cases  $N = 1,2$  by Assumption [2.4.](#page-2-2)

To see the last statement, recall from Theorem 2.2.1 of [\[35\]](#page-21-19) that the matrix-valued process  $(M_t(u))_{ij} := D^{(j)} \tilde{X}^i_t(\mathbf{q}), \, t \in [u,1] \text{ satisfies the (random) ordinary differential equation}$ 

<span id="page-12-2"></span>
$$
dM_t(u) = \partial_x \zeta \left( \tilde{X}_t(\mathbf{q}) + \mathbf{z}_t, \mathbf{v}_t \right) M_t(u) dt, \ M_u(u) = \mathbf{v}_u \sqrt{I - \mathbf{r}_u \mathbf{r}_u^*},
$$
(36)

for each  $u \in [0,1]$ . Assumption [\(2.4\)](#page-2-2) gives us  $K'$  with  $|\partial_x \zeta| \le K'$ . We see that

$$
M_1(u)M_1^*(u) = \mathbf{v}_u \sqrt{I - \mathbf{r}_u \mathbf{r}_u^*} \exp\left(\int_u^1 \partial_x \zeta(\tilde{X}_s(\mathbf{q}) + \mathbf{z}_s, \mathbf{v}_s) + \partial_x \zeta^*(\tilde{X}_s(\mathbf{q}) + \mathbf{z}_s, \mathbf{v}_s) ds\right) \sqrt{I - \mathbf{r}_u \mathbf{r}_u^*} \mathbf{v}_u^*
$$
  
\n
$$
\geq \exp(-2K') \mathbf{v}_u (I - \mathbf{r}_u \mathbf{r}_u^*) \mathbf{v}_u^*
$$
  
\n
$$
\geq \epsilon I,
$$

for some  $\epsilon = \epsilon(\mathbf{v}, \mathbf{r}) > 0$  since  $\mathbf{v} \in \mathbb{C}^+$  and  $\mathbf{r} \in \mathbb{C}^{1+}$ . This implies our claim since

$$
\sigma(\tilde{X}_s(\mathbf{q})) = \int_0^1 M_1(u) M_1(u)^* du.
$$

It follows also from [\(36\)](#page-12-2) that  $D\tilde{X}_{1}(\bold{q})$  is bounded, for each  $\bold{q}$ . Finally, Lemma 2.2.2 of [\[35\]](#page-21-19) implies that, for each pair of indices  $i, l$ , the second derivative  $DD^{(l)}\tilde{X}_t^i(\mathbf{q})$  satisfies for all indices  $j$  and for all  $s \le u \le t$ ,

$$
\begin{array}{rcl} |D_s^{(j)}D_u^{(l)}\tilde{X}_t^i(\mathbf{q})|&=&\displaystyle \left|\int_u^t(D_s^{(j)}\partial_x\zeta(\tilde{X}_r(\mathbf{q})+\mathbf{z}_r,\mathbf{v}_r))D_u^{(l)}\tilde{X}_r^i(\mathbf{q})+\partial_x\zeta(\tilde{X}_r(\mathbf{q})+\mathbf{z}_r,\mathbf{v}_r)D_s^{(l)}\tilde{X}_r^i(\mathbf{q})dr\right|\\&\leq&\displaystyle \int_u^tK_1+K_2|D_s^{(j)}D_u^{(l)}\tilde{X}_r^i(\mathbf{q})|dr\end{array}
$$

for constants  $K_1,K_2$  since  $\partial_x\zeta$  ,  $\partial_{xx}\zeta$  ,  $D\tilde X_t^i(\bf q)$  are all bounded. Gronwall's lemma now guarantees that also  $D_s^{(j)} D_u^{(l)} \tilde{X}_1^i(\mathbf{q})$  is bounded.  $\Box$ 

We now set up a discrete-time machinery so that we can invoke the results of Subsection [3.2.](#page-7-0) Set  $\mathcal{X}$ : = ℝ<sup>d</sup> and  $\mathcal{X}_n$ : = { $x \in \mathbb{R}^d$ : | $x | ≤ n$ },  $n ∈ \mathbb{N}$ . Define, for  $k ∈ \mathbb{Z}$ , the  $\mathcal{Y}$ -valued random variables

<span id="page-13-0"></span>
$$
Y_k := \left( (V_{k+t})_{t \in [0,1]}, (Z_{k+t} - Z_k)_{t \in [0,1]}, (\rho_{k+t})_{t \in [0,1]} \right),\tag{37}
$$

where we denote  $Z_u := \int_0^u V_s \rho_s dB_s$ ,  $u \in \mathbb{R}_+$ . *Y* is a stationary process by Assumption [2.1.](#page-1-1)

By Prokhorov's theorem, there exist an increasing sequence of compact sets  $\mathbf{D}_n \subset \mathbf{C}_{d \times d} \times \mathbf{C}_d \times \mathbf{C}^1$ ,  $n \in \mathbb{N}$  such that  $P(Y_0 \notin \mathbf{D}_n) \leq 1/n$ . As  $V \in \mathcal{V}$  and  $\rho \in \mathcal{R}$ ,  $P(Y_0 \in \mathbf{C}^+ \times \mathbf{C}_d \times \mathbf{C}^{1+}) = 1$  holds. Thus there is an increasing N-valued sequence  $l(n) \to \infty$ ,  $n \to \infty$  such that the sets

$$
\mathcal{Y}_n := \{ (\mathbf{v}, \mathbf{z}, \mathbf{r}) \in \mathbf{D}_n : \ \mathbf{r}_s \mathbf{r}_s^* \le (1 - 1/l(n))I, \ \mathbf{v}_s \mathbf{v}_s^* \ge I/l(n), \ s \in [0, 1] \}
$$

satisfy  $P(Y_0 \notin \mathcal{Y}_n) \le 2/n$ ,  $n \in \mathbb{N}$ . Being closed subsets of the respective  $\mathbf{D}_n$ , they are compact and satisfy  $P(Y_0 \notin \mathcal{Y}_n) \to 0, n \to \infty$ .

We define a metric on  $\mathcal{Q} := \mathcal{Y} \times \mathbb{R}^d$  by setting, for  $\mathbf{q}_i = (\mathbf{v}^i, \mathbf{z}^i, \mathbf{r}^i, x^i), i = 1, 2,$ 

$$
\rho(\mathbf{q}_1, \mathbf{q}_2) := ||\mathbf{v}^1 - \mathbf{v}^2||_{\mathbf{C}_{d \times d}} + ||\mathbf{z}^1 - \mathbf{z}^2||_{\mathbf{C}_d} + ||\mathbf{r}^1 - \mathbf{r}^2||_{\mathbf{C}_{d \times m}} + |x^1 - x^2|.
$$

Continuity of  $\tilde{X}_t(\mathbf{q})$  and its Malliavin derivatives with respect to the parameter  $\mathbf{q}$  is established next.

<span id="page-13-1"></span>**Lemma 4.4.** For each  $n \in \mathbb{N}$  and  $p \ge 2$  there exists  $C(n, p) > 0$  such that for all  $\mathbf{q}^1, \mathbf{q}^2 \in \mathcal{Y}_n \times \mathcal{X}_n$  we *have*

$$
E^{1/p}[\sup_{t\in[0,1]}|\tilde{X}_t(\mathbf{q}^1)-\tilde{X}_t(\mathbf{q}^2)|^p] \leq C(n,p)\rho(\mathbf{q}^1,\mathbf{q}^2),
$$
  

$$
E^{1/p}[\|\tilde{D}\tilde{X}_1(\mathbf{q}^1)-\tilde{X}_1(\mathbf{q}^2)\|_H^p] \leq C(n,p)\rho(\mathbf{q}^1,\mathbf{q}^2),
$$
  

$$
E^{1/p}[\|\tilde{D}^2\tilde{X}_1(\mathbf{q}^1)-\tilde{D}^2\tilde{X}_1(\mathbf{q}^2)\|_{H\otimes H}^p] \leq C(n,p)\rho(\mathbf{q}^1,\mathbf{q}^2).
$$

*Proof.* For  $i = 1, 2$ , define the Picard iterates  $Z_0^i(t) := x^i, t \in [0, 1]$  and

$$
Z_{l+1}^i(t) := x^i + \int_0^t \zeta \left( Z_l^i(s) + \mathbf{z}_s^i, \mathbf{v}_s^i \right) ds + \int_0^t \mathbf{v}_s^i \sqrt{I - \mathbf{r}_s^i \mathbf{r}_s^{i*}} dW_s, \tag{38}
$$

for  $t \in [0,1]$  and  $l \in \mathbb{N}$ . Let  $K_0$  denote a bound for  $|\partial_x \zeta|$  such that  $K_0 \ge K$  where  $K$  is as in Assumption [2.4.](#page-2-2) Clearly,

$$
\sup_{u \in [0,t]} |Z_{l+1}^1(u) - Z_{l+1}^2(u)|
$$
\n
$$
\leq |x^1 - x^2| + K_0 \int_0^t \sup_{u \in [0,s]} |Z_{l+1}^1(u) - Z_{l+1}^2(u)| + (1 + |\mathbf{v}_s^1| + |\mathbf{z}_s^1 - \mathbf{z}_s^2| + |\mathbf{v}_s^2|) |\mathbf{v}_s^1 - \mathbf{v}_s^2| ds
$$
\n
$$
+ \sup_{s \in [0,t]} \left| \int_0^s \mathbf{v}_u^1 \sqrt{I - \mathbf{r}_u^1 \mathbf{r}_u^{1*}} - \mathbf{v}_u^2 \sqrt{I - \mathbf{r}_u^2 \mathbf{r}_u^{2*}} dW_u \right|.
$$

Note that there is  $C > 0$  such that

$$
(x_1 + x_2 + x_3 + x_4 + x_5)^2 \le C(x_1^2 + x_2^2 + x_3^2 + x_4^2 + x_5^2), \ x_i \in \mathbb{R}, \ i = 1, 2, 3, 4, 5.
$$

Taking squares and using Cauchy's inequality we arrive at

$$
\sup_{u \in [0,t]} |Z_{l+1}^1(u) - Z_{l+1}^2(u)|^2
$$
\n
$$
\leq C|x^1 - x^2|^2 + CtK_0^2 \int_0^t \sup_{u \in [0,s]} |Z_{l+1}^1(u) - Z_{l+1}^2(u)|^2 + |\mathbf{z}_s^1 - \mathbf{z}_s^2|^2 + (1 + |\mathbf{v}_s^1| + |\mathbf{v}_s^2|)^2 |\mathbf{v}_s^1 - \mathbf{v}_s^2|^2 ds
$$
\n
$$
+ C \sup_{s \in [0,t]} \left| \int_0^s \mathbf{v}_u^1 \sqrt{I - \mathbf{r}_u^1 \mathbf{r}_u^{1*}} - \mathbf{v}_u^2 \sqrt{I - \mathbf{r}_u^2 \mathbf{r}_u^{2*}} dW_u \right|^2.
$$

Taking expectations, applying Doob's inequality and noting  $t \leq 1$ ,

$$
E \sup_{u \in [0,t]} |Z_{l+1}^1(u) - Z_{l+1}^2(u)|^2
$$
  
\n
$$
\leq C|x^1 - x^2|^2 + CK_0^2 \int_0^t E \sup_{u \in [0,s]} |Z_{l+1}^1(u) - Z_{l+1}^2(u)|^2 + (1 + |\mathbf{v}_s^1| + |\mathbf{z}_s^1 - \mathbf{z}_s^2|^2 + |\mathbf{v}_s^2|)^2 |\mathbf{v}_s^1 - \mathbf{v}_s^2|^2 ds
$$
  
\n+ 
$$
4CE \int_0^1 (\mathbf{v}_u^1 \sqrt{I - \mathbf{r}_u^1 \mathbf{r}_u^1} - \mathbf{v}_u^2 \sqrt{I - \mathbf{r}_u^2 \mathbf{r}_u^2})^2 du
$$
  
\n
$$
\leq C'_n \left[ \rho^2 (\mathbf{q}^1, \mathbf{q}^2) + \int_0^t E \sup_{u \in [0,s]} |Z_{l+1}^1(u) - Z_{l+1}^2(u)|^2 ds \right].
$$

 $C'_n$  because  $z \to \sqrt{I - zz^*}$  is Lipschitz-continuous on the set  $\{z \in \mathbf{C}^1 : zz^* \leq (1-\epsilon)I\}$ , for all  $\epsilon > 0$ . Grönwall's lemma implies that for some constant  $C''_n$ , independent of *l*,

$$
E\left[\sup_{t\in[0,1]}|Z_{l+1}^1(t)-Z_{l+1}^2(t)|^2\right]\leq C''_n\rho^2(\mathbf{q}^1,\mathbf{q}^2).
$$

Since Picard iterates converge, (see e.g. Lemma 2.2.1 in [\[35\]](#page-21-19)), we get

$$
E^{1/2}\left[\sup_{t\in[0,1]}|\tilde{X}(\mathbf{q}^1)-\tilde{X}(\mathbf{q}^2)|^2\right]\leq\sqrt{C''_n}\rho(\mathbf{q}^1,\mathbf{q}^2).
$$

A similar argument works in  $L^p$  with  $p > 2$ , too. Now recall that  $D\tilde{X}(\mathbf{q}), D^2\tilde{X}(\mathbf{q})$  also satisfy similar (even simpler) equations, see Theorem 2.2.1 of [\[35\]](#page-21-19), so analogous arguments apply to them, proving the remaining two inequalities.  $\Box$ 

We continue with some more technical material. In the following lemma, we will rely on the powerful techniques presented in [\[3\]](#page-20-12).

<span id="page-14-0"></span>**Lemma 4.5.** The random variables  $\tilde{X}_1(q)$  have densities  $p_q(u)$ ,  $u \in \mathbb{R}^d$  with respect to Leb<sub>d</sub>, the d*dimensional Lebesgue measure, for each*  $q \in \mathcal{Y}_n \times \mathcal{X}_n$ , for each *n*. These densities have versions such *that the mapping*  $(u, \mathbf{q}) \to p_{\mathbf{q}}(u)$  *is continuous on*  $\mathcal{Y}_n \times \mathcal{X}_n$ *, for each n.* 

*Proof.* Fix *n*, let  $\mathbf{q}_k \in \mathcal{Y}_n \times \mathcal{X}_n$  and  $u_k \in \mathbb{R}^d$ ,  $k \in \mathbb{N}$  be such that

$$
(u_n, \mathbf{q}_n) := (u_n, \mathbf{v}_n, \mathbf{r}_n, \mathbf{z}_n, x_n) \rightarrow (u, \mathbf{q}) := (u, \mathbf{v}, \mathbf{r}, \mathbf{z}, x)
$$

hold as  $n \to \infty$ . Let  $\partial_i Q$ ,  $i = 1, ..., d$  denote the *i*th partial derivative of the Poisson kernel on  $\mathbb{R}^d$ , see page 14 of [\[3\]](#page-20-12). We rely on the Malliavin-Thalmaier formula for the density of functionals on the Wiener space [\[30\]](#page-21-20), as presented in [\[3\]](#page-20-12). By Theorem 2.3.1 of [\[3\]](#page-20-12), the representation

$$
p_{\mathbf{q}}(u) = \sum_{i=1}^{d} \sum_{j=1}^{d} E\left[\partial_i Q\left(\tilde{X}_1(\mathbf{q}) - u\right) \delta\left(\gamma^{i,j} (\tilde{X}_1(\mathbf{q})) D\tilde{X}_1(\mathbf{q})\right)\right]
$$
(39)

provides the density function of  $\tilde{X}_1(\mathbf{q})$  (with respect to the  $d$ -dimensional Lebesgue-measure).

Fix *i*, *j*. By Lemma [4.3,](#page-12-3) the sequence  $\tilde{X}_1(q_n)$  is bounded in  $\mathbb{D}^{2,p}$  for all *p* and  $\gamma(\tilde{X}_1(q_n))$  is uniformly bounded, hence  $\partial_i Q\left(\tilde{X}_1(\mathbf{q}_n) - u\right)$  is bounded in  $L^{(d+1)/d}$  by (2.86) in Theorem 2.3.1 of [\[3\]](#page-20-12). By Corollary 2.2.12 in [\[3\]](#page-20-12), the sequence  $\delta(\gamma^{i,j}(\tilde{X}_1(\mathbf{q}))D\tilde{X}_1(\mathbf{q}_n))$  is bounded in  $L^p$  for all  $p \ge 1$ , in particular in  $L^{(d+1/2)(2d+2)/d}$ . But then Hölder's inequality implies

<span id="page-15-0"></span>
$$
\sup_{n} E\left[\left|\partial_{i} Q\left(\tilde{X}_{1}(\mathbf{q}_{n})-u\right) \delta\left(\gamma^{i,j}(\tilde{X}_{1}(\mathbf{q}_{n}))D\tilde{X}_{1}(\mathbf{q}_{n})\right)\right|^{(d+1/2)/d}\right]
$$
\n
$$
\leq \sup_{n} E\left[\left|\partial_{i} Q\left(\tilde{X}_{1}(\mathbf{q}_{n})-u\right)\right|^{(d+1)/d}\right]^{(d+1/2)/(d+1)} \tag{40}
$$

$$
\times \quad \sup_{n} E\left[\left|\delta^{(d+1/2)(2d+2)/d}\left(\gamma^{i,j}(\tilde{X}_1(\mathbf{q}_n))D\tilde{X}_1(\mathbf{q}_n)\right)\right|\right]^{1/(2d+2)} < \infty. \tag{41}
$$

This means that the sequence  $\partial_iQ\left(\tilde X_1(\mathbf q_n)-u\right)\delta\left(\gamma^{i,j}(\tilde X_1(\mathbf q_n))D\tilde X_1(\mathbf q_n)\right),\,n\in\mathbb N$  is uniformly integrable hence it suffices to prove that

<span id="page-15-1"></span>
$$
\partial_i Q\left(\tilde{X}_1(\mathbf{q}_n) - u_n\right) \delta\left(\gamma^{i,j}(\tilde{X}_1(\mathbf{q}_n))D\tilde{X}_1(\mathbf{q}_n)\right) \to \partial_i Q\left(\tilde{X}_1(\mathbf{q}) - u\right) \delta\left(\gamma^{i,j}(\tilde{X}_1(\mathbf{q}))D\tilde{X}_1(\mathbf{q})\right), \ n \to \infty \qquad (42)
$$

in probability. We remark that in [\(41\)](#page-15-0) we can also take a supremum in  $u \in C$  for any compact  $C \subset \mathbb{R}^d$ and this also implies

<span id="page-15-2"></span>
$$
\sup_n \sup_{u \in C} p_{\mathbf{q}_n}(u) < \infty. \tag{43}
$$

We start by treating the factor  $\partial_i Q$  in [\(42\)](#page-15-1). Let  $\varepsilon > 0$  be given. Notice that each  $\partial_i Q$  is Lipschitzcontinuous outside a ball of radius  $\varepsilon$ , with Lipschitz-constant, say,  $K_{\varepsilon}$ . Let M be such that  $\sup_n |u_n| \le$  $M$  and denote by  $A_d$  the volume of the unit ball in  $\mathbb{R}^d.$  We thus have, by the Markov inequality,

$$
P\left(|\partial_i Q\left(\tilde{X}_1(\mathbf{q}_n) - u_n\right) - \partial_i Q\left(\tilde{X}_1(\mathbf{q}) - u\right)\right| \geq \varepsilon\right)
$$
  
\n
$$
\leq P\left(|\tilde{X}_1(\mathbf{q}_n) - u_n| \leq \varepsilon\right) + P\left(|\tilde{X}_1(\mathbf{q}) - u| \leq \varepsilon\right)
$$
  
\n+  $K_{\varepsilon} E\left[|\tilde{X}_1(\mathbf{q}_n) - u_n - \tilde{X}_1(\mathbf{q}) + u|\right] / \varepsilon$   
\n
$$
\leq A_d \varepsilon^d \sup_{|v| \leq M + \varepsilon} \sup_{k \in \mathbb{N}} p_{\mathbf{q}_k}(v) + A_d \varepsilon^d \sup_{|v| \leq M + \varepsilon} p_{\mathbf{q}}(v) + K_{\varepsilon} E\left[|\tilde{X}_1(\mathbf{q}_n) - u_n - \tilde{X}_1(\mathbf{q}) + u|\right] / \varepsilon.
$$

Here the third term is smaller than *ε* for *n* large enough, by Lemma [4.4.](#page-13-1) Then all three terms can be made arbitrarily small by choosing *ε* small enough and *n* large enough, the suprema being finite by [\(43\)](#page-15-2). This shows convergence in probability for the first factor in [\(42\)](#page-15-1).

Now we turn to the second factor in [\(42\)](#page-15-1). By Proposition 1.5.4 of [\[35\]](#page-21-19), convergence of

<span id="page-15-3"></span>
$$
\gamma^{i,j}(\tilde{X}_1(\mathbf{q}_n))D\tilde{X}_1(\mathbf{q}_n) \to \gamma^{i,j}(\tilde{X}_1(\mathbf{q}))D\tilde{X}_1(\mathbf{q}) \quad \text{in } \mathbb{D}^{1,2} \tag{44}
$$

implies the  $L^2$ -convergence (hence also convergence in probability) of

$$
\delta\left(\gamma^{i,j}(\tilde{X}_1(\mathbf{q}_n))D\tilde{X}_1(\mathbf{q}_n)\right) \to \delta\left(\gamma^{i,j}(\tilde{X}_1(\mathbf{q}))D\tilde{X}_1(\mathbf{q})\right).
$$

So it remains to establish [\(44\)](#page-15-3).

Since  $\gamma^{i,j}(\tilde{X}_1(\mathbf{q}_n))$  are uniformly bounded and  $D\tilde{X}_1(\mathbf{q}_n) \to D\tilde{X}_1(\mathbf{q})$  in  $L^2([0,1] \times \Omega)$ , we clearly have that  $\gamma^{i,j}(\tilde{X}_1(\mathbf{q}_n))D\tilde{X}_1(\mathbf{q}_n) \to \gamma^{i,j}(\tilde{X}_1(\mathbf{q}))D\tilde{X}_1(\mathbf{q})$  in  $L^2([0,1] \times \Omega)$  by Lemma [4.4](#page-13-1) and by the fact that matrix inversion is a continuous operation. Let us now have a closer look at the Malliavin derivative of  $\gamma^{i,j}(\tilde{X}_1(\mathbf{q}_n))D\tilde{X}_1(\mathbf{q}_n)$ .

First, let us recall that  $D^2 \tilde{X}_1(\mathbf{q}_n) \to D^2 \tilde{X}_1(\mathbf{q})$  in  $L^p([0,1]^2 \times \Omega)$  for all  $p \ge 1$ , by Lemma [4.4.](#page-13-1) Also,  $\gamma^{i,j}(\tilde{X}_1(\mathbf{q}_n)) \to \gamma^{i,j}(\tilde{X}_1(\mathbf{q}))$  in  $L^p$  and  $D\tilde{X}_1(\mathbf{q}_n) \to D\tilde{X}_1(\mathbf{q})$  in  $L^p$ . It remains to establish

<span id="page-16-0"></span>
$$
D\gamma^{i,j}(\tilde{X}_1(\mathbf{q}_n)) \to D\gamma^{i,j}(\tilde{X}_1(\mathbf{q}))
$$
\n(45)

 $\operatorname{in} L^p$ .

Denote by  $G: V \to \mathbb{R}^{d \times d}$  the operation of matrix inversion and by  $G'$  its derivative. We find that, on the set of positive definite matrices bounded away from 0, *G*′ is bounded. Now [\(45\)](#page-16-0) follows from Lemma [4.4](#page-13-1) and from our previous observations. As these arguments work for all *i*, *j*, we indeed get  $p_{\mathbf{q}_n}(u_n) \rightarrow p_{\mathbf{q}}(u).$  $\Box$ 

Proving the positivity of densities is an evergreen topic in Malliavin calculus. We will rely on the deep study [\[2\]](#page-20-13) in the next lemma.

<span id="page-16-1"></span>**Lemma 4.6.** We have  $p_{q}(u) > 0$  for every  $q$ ,  $u$ .

*Proof.* Fix **q**, *u*. We will choose  $0 < \eta < 1/2$  later. We will apply Theorem 3.3 of [\[2\]](#page-20-13) with the choice  $y = u, r = 1, T = 1, \delta = n$ 

$$
F = \tilde{X}_1(\mathbf{q}), F_{1-\eta} = \tilde{X}_{1-\eta}(\mathbf{q}),
$$
  
\n
$$
G_{\eta} = \mathbf{v}_{1-\eta} \sqrt{I - \mathbf{r}_{1-\eta} \mathbf{r}_{1-\eta}^*} (W_1 - W_{1-\eta}),
$$
  
\n
$$
R_{\eta} = R_{\eta}^1 + R_{\eta}^2 = \int_{1-\eta}^1 (\mathbf{v}_s \sqrt{I - \mathbf{r}_s \mathbf{r}_s^*} - \mathbf{v}_{1-\eta} \sqrt{I - \mathbf{r}_{1-\eta} \mathbf{r}_{1-\eta}^*}) dW_s
$$
  
\n+ 
$$
\int_{1-\eta}^1 \zeta(\tilde{X}_s(\mathbf{q}) + \mathbf{z}_s, \mathbf{v}_s) ds.
$$

We are now checking the conditions of that theorem.

First,  $F_{1-\eta}$  is  $\mathscr{F}_{1-\eta}$ -measurable. Second, as all coordinates of  $\tilde{X}_1(\mathbf{q})$  are in  $\mathbb{D}^{2,\infty}$  by Lemma [4.3,](#page-12-3) so are those of  $R_\delta$ . Third, since  $s \to (\mathbf{r}_s, \mathbf{v}_s)$  is continuous and  $\mathbf{r}_s \in \mathbb{C}^{1+}$ ,  $\mathbf{v}_s \in \mathbb{C}^+$ , there is  $\varepsilon > 0$  such that  $\mathbf{v}_s \sqrt{I - \mathbf{r}_s \mathbf{r}_s^*} \geq \varepsilon I$  for all *s*. It follows that

$$
C_{\eta} := \int_{1-\eta}^{1} \mathbf{v}_s^*[I - \mathbf{r}_s \mathbf{r}_s^*] \mathbf{v}_s ds \geq \eta \varepsilon^2 I.
$$

Clearly,  $\det(C_n) \neq 0$ .

In order to apply Theorem 3.3 of [\[2\]](#page-20-13), it remains to check that the event

$$
\widetilde{\Gamma}_{\eta,1}=\{|F_{1-\eta}-y|\leq 1/2\}\cap\{||C_{\eta}^{-1/2}R_{\eta}||_{\eta,2,q}\leq a\mathrm{e}^{-1}\}
$$

has positive probability for a suitable *η*. Here *q* and *a* are explicit constants whose precise form can be found in [\[2\]](#page-20-13). For  $U \in \mathbb{R}^d$  with all coordinates in  $\mathbb{D}^{2,q}$  the norm  $||U||_{\delta,2,q}$  is defined as the random quantity

$$
\left(E_{T-\eta}[\left|U\right|^q\right] + E_{T-\eta}\left[\left(\int_{T-\eta}^T |D_s(U)|^2 ds\right)^{q/2}\right] + E_{T-\eta}\left[\left(\int_{T-\eta}^T \int_{T-\eta}^T |D^2_{s_1,s_2}(U)|^2 ds_1 ds_2\right)^{q/2}\right]\right)^{1/q}, \quad (46)
$$

with  $E_{T-n}$  denoting conditional expectation with respect to  $\mathscr{F}_{T-n}$ . As we have already seen,

$$
C_{\eta}^{-1/2}\leq \frac{1}{\varepsilon \sqrt{\eta}}I,
$$

so it suffices to show that

$$
\hat{\Gamma}_{\eta,1} := \{ |F_{1-\eta} - u| \le 1/2 \} \cap \{ ||R_{\eta}||_{\eta,2,q} \le a e^{-1} \varepsilon \sqrt{\eta} \}
$$

has positive probability.

By an easy extension of the support theorem for diffusions, see [\[19\]](#page-21-21), the process  $\tilde{X}({\bf q})$  has full support on the space of continuous functions starting from *x*, so we clearly have that

$$
P(A_{\eta}) > 0 \text{ for } A_{\eta} := \{|F_{1-\eta} - u| \le 1/2\},\
$$

for each  $0 < \eta < 1/2$ . A standard argument (like Lemma 19 of [\[1\]](#page-20-14)) shows that

$$
||R_\eta||_{\eta,2,q} \leq (1+|F_{1-\eta}|)\eta
$$

almost surely. But then on  $A_\eta$  we have  $||R_\eta||_{\delta,2,q} \leq (1+|u|+1/2)\eta$ . Clearly, this is smaller than  $a e^{-1} ε \sqrt{η}$  for *η* small enough. We conclude that the set  $\hat{\Gamma}_{\eta,q}$  contains  $A_\eta$  for *η* small enough, conse-quently it has positive probability. Now Theorem 3.3 of [\[2\]](#page-20-13) implies that  $p_{\mathbf{q}}(\cdot) \geq c$  Lebesgue-a.s. with some  $c > 0$  in a neighbourhood of *u*. As  $p_{\mathbf{q}}(u)$  is continuous in *u*,  $p_{\mathbf{q}}(u) \ge c$ , showing our lemma.  $\Box$ 

<span id="page-17-0"></span>**Corollary 4.7.** *There exist constants*  $\tilde{c}_n > 0$ ,  $n \in \mathbb{N}$  *such that for each*  $A \in \mathcal{B}(\mathbb{R})$  *with*  $A \subset [-1,1]$  *and for all*  $(\mathbf{v}, \mathbf{z}, \mathbf{r}) \in \mathcal{Y}_n$ ,  $x \in \mathcal{X}_n$ ,

$$
P(\tilde{X}_1(\mathbf{v}, \mathbf{z}, \mathbf{r}, x) \in A) \ge \tilde{c}_n \text{Leb}(A).
$$

*Proof.* Compactness of  $[-1,1] \times \mathcal{Y}_n \times \mathcal{X}_n$ , Lemmas [4.5](#page-14-0) and [4.6](#page-16-1) imply that

$$
\inf_{(u,\mathbf{q})\in[-1,1]\times\mathcal{Y}_n\times\mathcal{X}_n}p_{\mathbf{q}}(u)>0.
$$



Define  $\hat{X}_t(\mathbf{v}, \mathbf{z}, \mathbf{r}, x) := \tilde{X}_t(\mathbf{v}, \mathbf{z}, \mathbf{r}, x) + \mathbf{z}_t, t \in [0, 1].$  This process satisfies the integral equation

$$
\hat{X}_t(\mathbf{v}, \mathbf{z}, \mathbf{r}, x) = x + \int_0^t \zeta(\hat{X}_s, \mathbf{v}_s) ds + \mathbf{z}_t + \int_0^t \mathbf{v}_s \sqrt{I - \mathbf{r}_s \mathbf{r}_s^*} dW_s, \ t \in [0, 1]
$$

hence it will serve as the "parametric version" of [\(4\)](#page-2-4).

**Corollary 4.8.** There exist constants  $\hat{c}_n > 0$ ,  $n \in \mathbb{N}$  such that for each  $A \in \mathcal{B}(\mathbb{R})$  with  $A \subset [-1,1]$  and *for all*  $(\mathbf{v}, \mathbf{z}, \mathbf{r}) \in \mathcal{Y}_n$ ,  $x \in \mathcal{X}_n$ ,

$$
P(\hat{X}_1(\mathbf{v}, \mathbf{z}, \mathbf{r}, x) \in A) \ge \hat{c}_n \text{Leb}(A).
$$

*Proof.* Note that  $(v, z, r) \rightarrow ||z||_{C_d}$  is bounded on each  $\mathcal{Y}_n$ . Hence there is  $N \ge n$  such that whenever *x* ∈  $\mathcal{X}_n$  one has  $x + z_1 \in \mathcal{X}_N$  for all  $q \in \mathcal{Y}_n$ . Now Corollary [4.7](#page-17-0) readily implies the statement.

<span id="page-17-1"></span>**Lemma 4.9.** *There exists a measurable mapping*  $\Xi : \Omega \times \cup_n \mathcal{Y}_n \times \mathbb{R} \to \mathbb{C}_d$  *such that it satisfies for all* **q** ∈ ∪<sub>*n*</sub> $\mathcal{Y}_n$  × ℝ *the equation* 

$$
\Xi_t(\mathbf{q}) = x + \int_0^t \zeta(\Xi_s(\mathbf{q}), \mathbf{v}_s) ds + \mathbf{z}_t + \int_0^t \mathbf{v}_s \sqrt{I - \mathbf{r}_s \mathbf{r}_s^*} dW_t, \ t \in [0, 1].
$$

*For almost all*  $\omega$ ,  $\Xi(\omega, \cdot, \cdot)$  *is continuous. Furthermore*,  $\Xi_t(\cdot, Y_k, L_k)$ ,  $t \in [0,1]$  *is a version of*  $L_{k+t}$ , *t* ∈ [0,1]*. From now on we always take this version of L.*

*Proof.* Let us take an increasing sequence of sets  $B_n \subset \mathcal{Y}_n \times \mathcal{X}_n$ ,  $n \in \mathbb{N}$  which are countable and dense in  $\mathscr{Y}_n \times \mathscr{X}_n$ . By Lemma [4.4,](#page-13-1) there is a common *P*-null set  $N \in \mathscr{F}$  such that for  $\omega \in \Omega \setminus N$ the mapping  $\mathbf{q} \to (\hat{X}_u(\mathbf{q})(\omega))_{u \in [0,1]} \in \mathbf{C}_d$  is uniformly continuous on  $B_n$  for each *n* hence it has a continuous extension to  $\mathcal{Y}_n \times \mathcal{X}_n$  which coincides with the respective extensions on  $\mathcal{Y}_l \times \mathcal{X}_l$  for  $l \leq n$ . Hence we eventually get a function  $\Xi$ :  $(\Omega \setminus N) \times \cup_n \mathcal{Y}_n \times \mathbb{R} \to \mathbb{C}_d$  that is measurable in its first variable and jointly continuous in its second and third, hence jointly measurable in all three variables. (We set  $\Xi := 0$  on  $N$ .)

For any  $\mathscr{G}_{\infty} \vee \mathscr{F}_{k}$ -measurable step function  $\mathbf{Q}: \Omega \to \cup_{n} \mathscr{Y}_{n} \times \mathbb{R}$  with  $\mathbf{Q} = (\mathbf{V}, \mathbf{Z}, \mathbf{R}, X)$  it clearly holds that

$$
\Xi_t(\mathbf{Q}) = X + \int_0^t \zeta(\Xi_s(\mathbf{Q})), \mathbf{v}_s) ds + \mathbf{Z}_t + \int_0^t \mathbf{v}_s \sqrt{I - \mathbf{R}_s \mathbf{R}_s^*} dW_t, \ t \in [0, 1],
$$

and then this extends by continuity to all  $\cup_n \mathcal{Y}_n \times \mathbb{R}$ -valued  $\mathcal{G}_{\infty} \vee \mathcal{F}_k$ -measurable random variables **Q**, by continuity. In particular, it holds for **Q** :=  $(Y_k, L_k)$ , which proves the second statement. **Q**, by continuity. In particular, it holds for  $\mathbf{Q} := (Y_k, L_k)$ , which proves the second statement.

Let us define the parametrized kernel *Q* as follows: for each  $(x, y) \in \mathbb{R} \times \bigcup_{n} \mathcal{Y}_n$  and for all continu- $\text{ous and bounded } \phi : \mathbb{R}^d \to \mathbb{R}^d \text{ we let }$ 

$$
\int_{\mathbb{R}^d} \phi(z) Q(x,y,dz) := E[\phi(\Xi_1(y,x))].
$$

This clearly defines a probability for all  $(x, y)$ , and for a fixed  $\phi$  it is measurable in  $(x, y)$  by Lemma [4.9.](#page-17-1) Now we can recursively generate

$$
X_0:=L_0,\ X_{t+1}:=\Xi_{t+1}(Y_t,X_t), t\in \mathbb{N}\setminus\{0\}
$$

and see that *X* is a Markov chain in random environment with kernel *Q* which satisfies  $X_t = L_t$ ,  $t \in \mathbb{N}$ . Notice that [\(23\)](#page-7-4) holds by Lemma [4.2](#page-11-1) above.

Let  $\mu, \nu$  be probabilities on  $\mathscr{B}(\mathbb{R}^d \times \mathbb{W}^{d \times d} \times \mathbb{W}^{d \times m})$ . Let  $\mathscr{C}(\mu, \nu)$  denote the set of probabilities  $\pi$  on  $\mathscr{B}(\mathbb{R}^d \times \mathbb{W}^{d \times d} \times \mathbb{W}^{d \times m} \times \mathbb{R} \times \mathbb{W}^{d \times d} \times \mathbb{W}^{d \times m})$  such that their respective marginals are  $\mu, \nu$ . Define

$$
\mathbf{w}(\mu, v) := \inf_{\zeta \in \mathscr{C}(\mu, v)} \int_{(\mathbb{R}^d \times \mathscr{W}^{d \times d} \times \mathscr{W}^{d \times m})^2} ([1 \wedge |x_1 - x_2|] + \mathbf{d}_{d \times d}(v_1, v_2) + \mathbf{d}_{d \times m}(w_1, w_2)) \pi(dx_1, dv_1, dw_1, dx_2, dv_2, dw_2).
$$

This bounded Wasserstein distance metrizes weak convergence of probabilities on  $\mathscr{B}(\mathbb{R}^d \times \mathbb{W}^{d \times d} \times$  $W^{d \times m}$ ) and satisfies  $\mathbf{w}(\mu, \nu) \leq C ||\mu - \nu||_{TV}$  for some  $C > 0$ , see Theorem 6.15 of [\[43\]](#page-22-7).

*Proof of Theorem [2.9.](#page-2-0)* The letter *C* refers to various constants in this proof. Invoking Theorem [3.10,](#page-8-0) we can establish the existence of  $\mu_{\sharp}$  such that

$$
\mathscr{L}(L_l,\mathbf{V}_l,\mathbf{R}_l)\to\mu_{\sharp},\ l\to\infty,\ l\in\mathbb{N}
$$

holds in  $|| \cdot ||_{TV}$ . Working on a finer time grid, we similarly obtain that, for each  $k \in \mathbb{N}$ , the sequence of laws  $\mathscr{L}(L_{l/2^k}, \mathbf{V}_{l/2^k}, \mathbf{R}_{l/2^k})$ ,  $l \in \mathbb{N}$  converge in  $\lVert \cdot \rVert_{TV}$  as  $l \to \infty$  and all these limits necessarily equal *µ*♯ .

Assumption [2.4](#page-2-2) implies Lipschitz-continuity of *ζ* in its first variable and local Lipschitz-continuity with linearly growing Lipschitz-continuity in its second variable. In particular,  $|\zeta(x,v)| \leq C(1+|x|+1)$  $|v|^2$ ), hence for  $0 < h \le 1$ ,

<span id="page-18-0"></span>
$$
E[|L_{t+h} - L_t|^2]
$$
\n
$$
\leq 3E \left[ \left( \int_t^{t+h} \zeta(L_s, V_s) ds \right)^2 \right] + 3E \left[ \left( \int_t^{t+h} V_s \rho_s dB_s \right)^2 \right] + 3E \left[ \left( \int_t^{t+h} \sqrt{I - \rho_s \rho_s^*} V_s dW_s \right)^2 \right]
$$
\n
$$
\leq \int_t^{t+h} C \left[ E[|L_s|^2] + E[|V_0|^4] + 1 \right] ds + 3 \int_t^{t+h} E[|V_0|^2] ds + 3 \int_t^{t+h} E[|V_0|^2] ds
$$
\n
$$
\leq hC[\tilde{L} + E[|V_0|^4] + 1 + E[|V_0|^2]] \leq Ch,
$$
\n(47)

by Assumption [2.4](#page-2-2) and Lemma [4.2.](#page-11-1) It is only at this point that we need  $E[|V_0|^4] < \infty$ .

For each  $t \in \mathbb{R}_+$  and  $k \in \mathbb{N}$ , let  $l(k,t)$  denote the integer satisfying  $l(k,t)/2^k \le t < [l(k,t)+1]/2^k$ . Notice that, for *k* fixed,  $l(k, t) \rightarrow \infty$  as  $t \rightarrow \infty$ . We estimate, using [\(47\)](#page-18-0),

$$
\mathbf{w}(\mathcal{L}(L_t, \mathbf{V}_t, \mathbf{R}_t), \mu_{\sharp})
$$
\n
$$
\leq \mathbf{w}(\mathcal{L}(L_t, \mathbf{V}_t, \mathbf{R}_t), \mathcal{L}(L_{l(k,t)/2^k}, \mathbf{V}_{l(k,t)/2^k}, \mathbf{R}_{l(k,t)/2^k})) + \mathbf{w}(\mathcal{L}(L_{l(k,t)/2^k}, \mathbf{V}_{l(k,t)/2^k}, \mathbf{R}_{l(k,t)/2^k}), \mu_{\sharp})
$$
\n
$$
\leq E|L_t - L_{l(k,t)/2^k}| + E[\mathbf{d}_{d \times d}(\mathbf{V}_t, \mathbf{V}_{l(k,t)/2^k})]
$$
\n+  $E[\mathbf{d}_{d \times m}(\mathbf{R}_t, \mathbf{R}_{l(k,t)/2^k}] + C||\mathcal{L}(L_{l(k,t)/2^k}, \mathbf{V}_{l(k,t)/2^k}, \mathbf{R}_{l(k,t)/2^k}) - \mu_{\sharp}||_{TV}$   
\n
$$
\leq \sqrt{C/2^k} + \sup_{t \in \mathbb{R}} \{E[\mathbf{d}_{d \times d}(\mathbf{V}_t, \mathbf{V}_{l(k,t)/2^k})] + E[\mathbf{d}_{d \times m}(\mathbf{R}_t, \mathbf{R}_{l(k,t)/2^k}]]\}
$$
\n+  $C||\mathcal{L}(L_{l(k,t)/2^k}) - \mu_{\sharp}||_{TV}$ .

Noting Lemma [4.1](#page-11-2) and Theorem [3.10,](#page-8-0) the latter expression can be made arbitrarily small by first choosing *k* large enough and then choosing *t* large enough.

Now we turn to proving stationarity. Theorem [3.10](#page-8-0) implies that, if  $\mathcal{L}(L_0, \mathbf{V}_0, \mathbf{R}_0) = \mu_{\sharp}$  then

<span id="page-19-0"></span>
$$
\mathscr{L}(L_t, \mathbf{V}_t, \mathbf{R}_t) = \mu_{\sharp} \tag{48}
$$

 $\Box$ 

holds for all dyadic rationals  $t \ge 0$ . For an arbitrary  $t \in \mathbb{R}$ , take dyadic rationals  $t_n \to t$ ,  $n \to \infty$  and estimate

$$
\mathbf{w}(\mathcal{L}(L_t, \mathbf{V}_t, \mathbf{R}_t), \mathcal{L}(L_{t_n}, \mathbf{V}_{t_n}, \mathbf{R}_{t_n}))
$$
  
\n
$$
\leq E|L_t - L_{t_n}| + E[\mathbf{d}_{d \times d}(\mathbf{V}_t, \mathbf{V}_{t_n})] + E[\mathbf{d}_{d \times m}(\mathbf{R}_t, \mathbf{R}_{t_n})],
$$

which tends to 0 as  $n \to \infty$ , by Lemma [4.1](#page-11-2) and by [\(47\)](#page-18-0). Hence [\(48\)](#page-19-0) holds for all  $t \in \mathbb{R}$ .

*Proof of Theorem [2.10.](#page-3-0)* Notice that, in the proof of Theorem [2.9,](#page-2-0) we used Assumption [2.5](#page-2-5) only in Lemma [4.2.](#page-11-1) Under our current assumptions, we will verify

$$
\sup_{t\geq 0} E[e^{\kappa|L_t|}] < \infty
$$

for some  $\kappa > 0$ . This trivially entails

$$
\tilde{L} := \sup_{t \geq 0} E[|L_t|^2] < \infty,
$$

and the rest of the proof follows verbatim that of Theorem [2.9.](#page-2-0)

 $\text{We will use the Lyapunov-function } g(x) := \exp\left(\kappa\sqrt{1+|x|^2}\right)\!, x\in\mathbb{R}^d\!, \text{ where } 0<\kappa\leq\kappa_0 \text{ will be chosen}$ later. Note that

$$
\partial_i g(x) = \exp\left(\kappa \sqrt{1+|x|^2}\right) \frac{\kappa x^i}{\sqrt{1+|x|^2}}, i=1,\ldots,d,
$$

and  $|\partial_{ij} g(x)| \leq C_0 \kappa g(x)$  for all *x*, with some constant  $C_0 > 0$ , for all  $1 \leq i, j \leq d$ .

Fix  $k \in \mathbb{N}$ . Define the stopping times  $\tau_l := \inf\{t > k : |L_t| > l\}$  for  $l \in \mathbb{N}$ . Apply Itô's lemma to obtain

$$
e^{\alpha(t\wedge\tau_{l}-k)}e^{\kappa\sqrt{1+|L_{t\wedge\tau_{l}}|^2}} \leq e^{\kappa\sqrt{1+|L_{k}|^2}} + \int_{k}^{t\wedge\tau_{l}} e^{\alpha(s-k)}\kappa \frac{e^{\kappa\sqrt{1+|L_{s}|^2}}}{\sqrt{1+|L_{s}|^2}}\langle L_{s},\zeta(L_{s},V_{s})\rangle ds + \int_{k}^{t\wedge\tau_{l}} e^{\alpha(s-k)}\kappa \frac{e^{\kappa\sqrt{1+|L_{s}|^2}}}{\sqrt{1+|L_{s}|^2}}L_{s}^{*}V_{s}d\overline{W}_{s} + \int_{k}^{t\wedge\tau_{l}}C_{1}\kappa e^{\alpha(s-k)}e^{\kappa\sqrt{1+|L_{s}|^2}}|V_{s}|^2 ds + \int_{k}^{t\wedge\tau_{l}} \alpha e^{\alpha(s-k)}e^{\kappa\sqrt{1+|L_{s}|^2}}ds, t \geq k,
$$

for some  $C_1 > 0$ . Taking expectations, using the martingale property of stochastic integrals and [\(6\)](#page-3-3), we arrive at

$$
E\left[e^{\alpha(t\wedge\tau_l-k)}e^{\kappa\sqrt{1+|L_{t\wedge\tau_l}|^2}}\right] \leq E\left[e^{\kappa\sqrt{1+|L_k|^2}}\right] + E\left[\int_k^{t\wedge\tau_l} \kappa e^{\alpha(s-k)} \frac{e^{\kappa\sqrt{1+|L_s|^2}}}{\sqrt{1+|L_s|^2}} (-\alpha|L_s|^{1+\gamma} + \beta(1+|V_s|^{\zeta}))ds\right] + E\left[\int_k^{t\wedge\tau_l} C_1 \kappa e^{\alpha(s-k)} e^{\kappa\sqrt{1+|L_s|^2}} |V_s|^2 ds\right] + E\left[\int_k^{t\wedge\tau_l} \alpha e^{\alpha(s-k)} e^{\kappa\sqrt{1+|L_s|^2}} ds\right].
$$

Set  $C_2 := C_1 + \beta$ . Let us notice that, on the event

$$
A := \left\{ |L_s| \ge \max\left\{1, \left(\frac{2\sqrt{2}}{\kappa}\right)^{1/\gamma} + \left(\frac{2\sqrt{2}C_2}{\alpha}\right)^{1/\gamma}(1+|V_s|^{\zeta})^{1/\gamma}\right\} \right\}
$$

we have

$$
\frac{-\alpha\kappa}{\sqrt{1+|L_s|^2}}|L_s|^{1+\gamma} + C_2\kappa(1+|V_s|^{\xi}) + \alpha \leq 0
$$

On the complement of *A*,

$$
\exp\left(\kappa\sqrt{1+|L_s|^2}\right) \le \exp\left(\kappa+\kappa\left(1+\left(\frac{2\sqrt{2}}{\kappa}\right)^{1/\gamma}+\left(\frac{2\sqrt{2}C_2}{\alpha}\right)^{1/\gamma}\left(1+|V_s|^{\xi}\right)^{1/\gamma}\right)\right).
$$

Tending  $l \rightarrow \infty$  and applying Fatou's lemma, our estimate takes the form

$$
E\left[e^{\alpha(t-k)}e^{\kappa\sqrt{1+|L_t|^2}}\right] \leq E\left[e^{\kappa\sqrt{1+|L_k|^2}}\right]
$$
  
+ 
$$
C_3(\kappa)\int_k^{k+1} E\left[(1+|V_s|^{\xi})\exp\left(\kappa\left(\frac{2\sqrt{2}C_2}{\alpha}\right)^{1/\gamma}(1+|V_s|^{\xi})^{1/\gamma}\right)\right] ds
$$
  

$$
\leq E\left[e^{\kappa\sqrt{1+|L_k|^2}}\right] + C_4(\kappa)E\left[(1+|V_0|^{\xi})\exp\left(\kappa\left(\frac{2\sqrt{2}C_2}{\alpha}\right)^{1/\gamma}2^{1/\gamma}|V_0|^{\xi/\gamma}\right)\right].
$$

 $\int \int_0^{\pi} e^{i(x-x)} \, dx \leq f(x) + \int_0^{\pi} e^{i(x-x)} \, dx$  and  $\int e^{i(x-x)} \, dx \leq f(x) + \int_0^{\pi} e^{i(x-x)} \, dx$  and  $\int_0^{\pi} 2^{i(x)} \, dx < \kappa_0$ , the second integral is finite, by  $(6)$ . Now we can easily conclude, as in Lemma [4.2](#page-11-1) above.  $\Box$ 

## <span id="page-20-14"></span>**References**

- [1] V. Bally. Lower bounds for the density of locally elliptic Itô processes. *Annals of Probability*, 34:2406–2440, 2006.
- <span id="page-20-13"></span><span id="page-20-12"></span>[2] V. Bally and L. Caramellino. Positivity and lower bounds for the density of Wiener functionals. *Potential Analysis*, 39:141–168, 2013.
- [3] V. Bally and L. Caramellino. Integration by parts formulas, Malliavin calculus and regularity of probability laws. *In: V. Bally, L. Caramellino and R. Cont: Stochastic integration by parts and functional Itô calculus*, 1–114, Birkhäuser, 2016.
- <span id="page-20-8"></span>[4] Ch. Bayer, P. Friz and J. Gatheral. Pricing under rough volatility. *Quantitative Finance*, 16:887– 904, 2016.
- <span id="page-20-7"></span>[5] A. Benveniste, M. Métivier and P. Priouret. *Adaptive algorithms and stochastic approximations.* Springer, 1990.
- <span id="page-20-3"></span>[6] H. Bessembinder, J. F. Coughenor, P. J. Seguin and M. M. Smoller. Mean-reversion in equilibrium asset prices: evidence from the futures term structure. *Journal of Finance*, 50:361–375, 1995.
- <span id="page-20-11"></span>[7] R. Bhattacharya and E. Waymire. *Stochastic Processes with Applications*. Wiley & Sons, New York, 1990.
- <span id="page-20-4"></span>[8] J. Casassus and P. Collin-Dufresne. Stochastic convenience yield implied from commodity futures and interest rates. *Journal of Finance*, 60:2283–2331, 2005.
- <span id="page-20-1"></span>[9] F. Comte and É. Renault. Long memory in continuous-time stochastic volatility models. *Math. Finance*, 8:291–323, 1998.
- <span id="page-20-10"></span><span id="page-20-6"></span>[10] P. Diaconis and D. Freedman. Iterated random functions. *SIAM Review*, 41:45–76, 1999.
- [11] G.B. Di Masi and L. Stettner. Risk sensitive control of discrete time Markov processes with infinite horizon. *SIAM J. Control Optimiz.*, 38:61–78, 2000.
- <span id="page-20-0"></span>[12] D. Down, S. P. Meyn and R. L. Tweedie. Exponential and uniform ergodicity of Markov processes. *Ann. Probab.*, 23:1671–1691, 1995.
- <span id="page-20-5"></span>[13] H. Föllmer and W. Schachermayer. Asymptotic Arbitrage and large deviations. *Math. Financ. Econ.*, 1:213–249, 2008.
- <span id="page-20-9"></span>[14] M. Fukasawa. Short-time at-the-money skew and rough fractional volatility. *Quantitative Finance*, 17:189–198, 2017.
- <span id="page-20-2"></span>[15] J. Gatheral, T. Jaisson and M. Rosenbaum. Volatility is rough. *Quantitative Finance*, 18:933– 949, 2018.
- <span id="page-21-15"></span><span id="page-21-5"></span>[16] B. Gerencsér and M. Rásonyi. On the ergodicity of certain Markov chains in random environments. *Submitted*, 2020. [arXiv:1807.03568v](http://arxiv.org/abs/1807.03568)3
- <span id="page-21-8"></span>[17] P. Guasoni, L. Nagy, M. Rásonyi. Young, timid and risk seekers. *To appear in Mathematical Finance*, 2021. SSRN:3763151
- <span id="page-21-21"></span>[18] P. Guasoni, Zs. Nika and M. Rásonyi. Trading fractional Brownian motion. *SIAM J. Financial Mathematics*, vol. 10, 769–789, 2019.
- <span id="page-21-10"></span>[19] P. Guasoni and M. Rásonyi. Fragility of arbitrage and bubbles in local martingale diffusion models. *Finance Stoch.*, 19:215–231, 2015.
- <span id="page-21-12"></span>[20] H. Guennoun, A. Jacquier, P. Roome and F. Shi. Asymptotic behavior of the fractional Heston model. *SIAM J. Finan. Math.*, 9:1017–1045, 2018.
- <span id="page-21-4"></span>[21] M. Hairer. Ergodicity of stochastic differential equations driven by fractional Brownian motion. *Annals of Probability*, 703–758, 2005.
- <span id="page-21-14"></span>[22] M. Hairer. *Convergence of Markov processes.* Lecture notes, 2016[.http://hairer.org](http://hairer.org)
- [23] M. Hairer and J. Mattingly. Yet another look at Harris' ergodic theorem for Markov chains. *In: Seminar on stochastic analysis, random fields and applications VI (eds. R. Dalang, M. Dozzi and F. Russo F.)*, Progress in Probability, vol. 63, 109–117, 2011.
- <span id="page-21-6"></span><span id="page-21-2"></span>[24] O. Hernandez-Lerma and J. B. Lasserre. *Discrete-time Markov control processes*. Springer, 1996.
- <span id="page-21-17"></span>[25] R. Z. Khasminskii. *Stochastic stability of differential equations.* 2nd edition. Springer, 2012.
- [26] Y. Kifer. Perron-Frobenius theorem, large deviations, and random perturbations in random environments. *Math. Zeitschrift*, 222:677–698, 1996.
- <span id="page-21-18"></span>[27] Y. Kifer. Limit theorems for random transformations and processes in random environments. *Trans. American Math. Soc.*, 350:1481–1518, 1998.
- <span id="page-21-16"></span><span id="page-21-11"></span>[28] N. V. Krylov. *Controlled diffusion processes.* Springer, 1980.
- [29] A. Lovas and M. Rásonyi. Markov chains in random environment with applications in queuing theory and machine learning. *To appear in Stochastic Processes and their Applications*, 2021. [arXiv:1911.04377v](http://arxiv.org/abs/1911.04377)2
- <span id="page-21-20"></span><span id="page-21-3"></span>[30] P. Malliavin and A. Thalmaier. *Malliavin calculus for financial applications.* Springer, 2010.
- [31] S. P. Meyn and R. L. Tweedie. *Markov chains and stochastic stability.* 2nd edition. Cambridge University Press, 2009.
- <span id="page-21-0"></span>[32] S. P. Meyn and R. L. Tweedie. Stability of Markovian processes II. Continuous time processes and sampled chains. *Adv. Appl. Prob.* 25:487–517, 1993.
- <span id="page-21-1"></span>[33] S. P. Meyn and R. L. Tweedie. Stability of Markovian processes. III. Foster-Lyapunov criteria for continuous-time processes. *Adv. Appl. Probab.*, 25:518–548, 1993.
- <span id="page-21-9"></span>[34] L. Nagy, M. Rásonyi. Optimal long-term investment in illiquid markets when prices have negative memory. *To appear in Electronic Communications in Probability*, 2021. [arXiv:2005.07080](http://arxiv.org/abs/2005.07080)
- <span id="page-21-19"></span><span id="page-21-7"></span>[35] D. Nualart. *The Malliavin calculus and related topics.* 2nd edition, Springer, 2006.
- [36] M. Pitera and L. Stettner. Long run risk-sensitive portfolio with general factors. *Math. Method. Oper. Res.* 83:265–293, 2016.
- <span id="page-21-13"></span>[37] J. G. Propp and D. B. Wilson. Exact sampling with coupled Markov chains and applications to statistical mechanics. *Random Structures and Algorithms*, 9:223–252, 1996.
- <span id="page-22-5"></span><span id="page-22-1"></span>[38] G. O. Roberts and R. L. Tweedie. Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, 2:341–363, 1996.
- <span id="page-22-6"></span>[39] T. Seppäläinen. Large deviations for Markov chains with random transitions. *Ann. Probab.*, 22:713–748, 1994.
- <span id="page-22-0"></span>[40] Ö. Stenflo. Markov chains in random environments and random iterated function systems. *Trans. American Math. Soc.*, 353:3547–3562, 2001.
- [41] A. Yu. Veretennikov. Bounds for the mixing rate in the theory of stochastic equations. *Theory Probab. Appl.*, 32:273–281, 1987.
- <span id="page-22-2"></span>[42] A. Yu. Veretennikov. On polynomial mixing bounds for stochastic differential equations. *Stoch. Processes Appl.*, 70:115–127, 1997.
- <span id="page-22-7"></span><span id="page-22-4"></span>[43] C. Villani. *Optimal transport. Old and new.* Springer, 2009.
- [44] C. S. Withers. Central limit theorems for dependent variables I. *Z. Wahsch. verw. Gebiete*, 57:509–534, 1981.
- <span id="page-22-3"></span>[45] https://sites.google.com/site/roughvol/home/risks-1

**Data availability statement:** there is no associated data to this manuscript.