

UNIQUENESS OF COMMUNITIES IN REGULAR STOCHASTIC BLOCK MODELS

SAYAR KARMAKAR, MOUMANTI PODDER

ABSTRACT. This paper studies the regular stochastic block model comprising *several* communities: each of the k non-overlapping communities, for $k \geq 3$, possesses n vertices, each of which has total degree d . The values of the intra-cluster degrees (i.e. the number of neighbours of a vertex inside the cluster it belongs to) and the inter-cluster degrees (i.e. the number of neighbours of a vertex inside a cluster different from its own) are allowed to vary across clusters. We discuss two main results: the first compares the probability measure induced by our model with the uniform measure on the space of d -regular graphs on kn vertices, and the second establishes that the clusters, under rather weak assumptions, are unique asymptotically almost surely as $n \rightarrow \infty$.

1. INTRODUCTION

This paper concerns itself with the *regular stochastic block model*, henceforth abbreviated as RSBM, that is used to study *clustered networks*. These networks exhibit *community structure*, whereby the individuals participating in the network, typically indicated as nodes or vertices of a graph, are split into overlapping or non-overlapping groups, usually with dense connections internally and sparser connections between different groups. Community structure is common in many complex networks such as computer and information networks ([28]), online social networks and biological networks ([20, 11, 17, 35]) that include protein-protein and gene-gene interactions ([31]), biological neural networks ([19]), metabolic networks ([47]) etc. Detecting communities in clustered networks has been pursued with fervour ([41, 42, 43, 6, 40, 25]), since communities often act as meta-nodes in a network and individuals within the same community tend to exhibit behavioural and functional similarities, simplifying the analysis of the underlying features of the network. The characteristics displayed by each distinct community may also vary greatly from the average properties of the network. The existence of communities may also significantly affect the spreading of rumours, epidemics etc. within the network.

The *stochastic block model*, henceforth abbreviated as SBM, (introduced in [22], surveyed in [1]), has been the most popular model, so far, in studying clustered networks. In its most simplified form, this model comprises $2n$ vertices that are partitioned into two equi-sized clusters. Edges between all pairs of vertices appear mutually independently, with probability p if both vertices belong to the same cluster, and probability q if they belong to different clusters. Letting the intra-cluster average degree be $a \sim pn$ and the inter-cluster average degree be $b \sim qn$, [34] and [3] studied the SBM in the regime where $a, b = O(\log n)$, whereas [12], [14], [15], [36], [39], [37], [38] and [30] studied SBM

SAYAR KARMAKAR, DEPARTMENT OF STATISTICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FL 32611, UNITED STATES.

MOUMANTI PODDER, DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH (IISER) PUNE, DR. HOMI BHABHA ROAD, PASHAN, PUNE 411008, INDIA.

E-mail address: sayarkarmakar@ufl.edu, moumantip3@gmail.com.

2020 *Mathematics Subject Classification.* 60C05, 94C15, 68R05, 68R01.

Key words and phrases. regular stochastic block model, uniqueness of clusters, configuration models.

The second author was partially supported by the grant NSF DMS-1444084.

in the regime where $a, b = O(1)$, as n grows to ∞ . Other variants of this model studied are the Bayesian SBM ([46]), degree-corrected block models ([48]), labeled SBM ([27, 21]), SBM in sparse hypergraphs ([45]) etc. We also refer the reader to [16], [24] and [13] for discussions on relations between community detection in SBM and the minimum bisection problem that seeks to partition a graph of $2n$ vertices into two equi-sized parts such that the number of edges across the parts is minimized.

Since the essence of our paper is to focus on the case where the given model constitutes *several* underlying clusters, we emphasize on the following developments in the literature. In [14], it was conjectured that if the signal-to-noise ratio of a given SBM is strictly higher than 1, then it is possible to detect communities in polynomial time, or, in other words, the well-known Kesten-Stigum threshold is achieved; moreover, if the number of underlying communities in the model exceeds 4, it is possible to detect the communities information-theoretically for some signal-to-noise ratio strictly lower than 1. It was shown in [8] that the Kesten-Stigum threshold is achieved in SBM's with multiple communities satisfying certain asymmetry assumptions, whereas the full conjecture of [14], for several clusters, was established in [2]. The extension of SBM from two to several communities has proven to be a veritable challenge during the course of development of this field.

The RSBM was introduced in [9] (note that two regular versions of the SBM in the sparse regime was proposed in [36], and *equitable random graphs* have been studied in [44] and [4]). As in [9], we assume that each intra-cluster degree and each inter-cluster degree exceeds 3, ensuring that the resulting graph is connected with high probability. The RSBM differs from the SBM with constant average intra-cluster and inter-cluster degrees in that the latter has a positive probability of possessing isolated vertices. In RSBM, the imposition of the constraint that each vertex has a constant number of neighbours in each given cluster gives more structure to the graph, but at the same time, robs the model of the edge-independence that is present in SBM.

We now highlight the novelties as well as describe the organization of our paper. We emphasize that our model is more general than that of [9] in that it takes into account multiple communities as well as intra-cluster and inter-cluster degree values that differ across communities. We answer questions similar to those in [9], but the proof techniques are more involved and require careful analysis. To set the stage, in §1.1, we describe the notations and terminology used throughout the paper; in §1.2, we describe the model and its underlying measure in details; in §1.3, we describe the well-known configuration model and the associated exploration process, and their importance in the generation of uniformly random regular or bipartite-regular graphs.

In §2, we show that the measure induced by RSBM on kn vertices, each with degree d , where k denotes the number of communities and n the number of members in each community, is distinct from the measure that makes a uniformly random selection out of the collection of all d -regular graphs on kn vertices. In §3, we show that under rather weak assumptions, the underlying clusters of the model are unique almost surely as n approaches ∞ . We draw attention of the reader to a key difference between our analysis and the analysis in [[9], §3.2.2]: while they had the symmetry, around $1/2$, of the binary entropy function $H(\alpha) = -\alpha \log_2 \alpha - (1 - \alpha) \log_2 (1 - \alpha)$ in their favour due to the presence of only two clusters, we require a somewhat different strategy to handle the higher number of clusters in our model. Even in the homogeneous case, where all the intra-cluster degrees are the same, there is need for a thorough case-by-case analysis that is much more intricate than in [9]. We emphasize here that in the homogeneous case, our analysis allows for the inter-cluster degrees to exceed, by far, the intra-cluster degrees. This is a significant generalization over the much more usual assumption of denser intra-cluster connections and sparser inter-cluster

connections found in the literature. In the heterogeneous scenario, we need a more restricted range of intra-cluster and inter-cluster degrees, as described in [(3.1) and (3.2), Theorem 3.1].

1.1. Notations. Given $n, d \in \mathbb{N}$, we denote by \mathcal{R}_d^n the set of all d -regular graphs on n labeled vertices, and by \mathcal{B}_d^n the set of all d -bipartite-regular graphs on $2n$ labeled vertices where each cluster comprises n vertices. We shall denote by μ_d^n the uniform measure on \mathcal{R}_d^n .

Given a graph G , we denote by $V(G)$ its vertex set and by $E(G)$ its edge set. Given $S \subset V(G)$ and $v \in V(G)$, we let $\deg_S(v)$ denote the number of edges $\{u, v\}$ where $u \in S$. We denote by $G|_S$ the subgraph of G that is induced on S . For disjoint subsets S_1 and S_2 of $V(G)$, we denote by $\deg(S_1, S_2)$ the number of edges $\{u_1, u_2\}$ where $u_1 \in S_1$ and $u_2 \in S_2$. We let $G|_{S_1, S_2}$ denote the subgraph with vertex set $S_1 \cup S_2$ and edge set $\{\{u_1, u_2\} : u_1 \in S_1, u_2 \in S_2\}$. Given $v \in V(G)$ and $r \in \mathbb{N}$, we let $B(v, r) = \{u \in V(G) : \rho(u, v) \leq r\}$ be the neighbourhood of radius r around v , where ρ is the usual graph metric. We let $\delta B(v, r) = \{u \in V(G) : \rho(u, v) = r\}$ denote the boundary of $B(v, r)$.

Given an infinite sequence of graphs $\{G_n\}$ and a graph property A , we say that A holds *asymptotically almost surely* (a.a.s.) for this sequence if $\mathbf{P}[G_n \text{ satisfies property } A] \rightarrow 1$ as $n \rightarrow \infty$. Given any two functions $f : \mathbb{N} \rightarrow \mathbb{R}^+$ and $g : \mathbb{N} \rightarrow \mathbb{R}^+$, we write $f(n) \sim g(n)$ to indicate that $f(n)/g(n) \rightarrow 1$ as $n \rightarrow \infty$.

For any $\alpha \in (0, 1)$, recall that the Shannon entropy for a Bernoulli(α) distribution is given by $H(\alpha) = -\alpha \log_2 \alpha - (1 - \alpha) \log_2 (1 - \alpha)$. This will be used in §3.

1.2. Description of the model. Our k -cluster RSBM, denoted \mathcal{G}_A^n , has the following parameters:

- (i) n denotes the number of vertices in each cluster,
- (ii) k denotes the number of clusters,
- (iii) and $A = (A_{i,j})_{1 \leq i, j \leq k}$ is a $k \times k$ symmetric matrix of strictly positive integers such that, for some $d \in \mathbb{N}$,

$$\sum_{j=1}^k a_{i,j} = d \text{ for all } i = 1, \dots, k. \quad (1.1)$$

Starting with kn labeled vertices, we uniformly randomly partition them into k clusters $\mathcal{C}_1, \dots, \mathcal{C}_k$, each of size n . Independent of each other, we now place on the vertices of \mathcal{C}_i a uniformly random member of $\mathcal{R}_{a_{i,i}}^n$, and across the clusters \mathcal{C}_i and \mathcal{C}_j a uniformly random member of $\mathcal{B}_{a_{i,j}}^n$, for all $i, j \in \{1, \dots, k\}$ with $i \neq j$. The criterion in (1.1) ensures that any realization of our model will be d -regular. All the parameters except n remain fixed throughout our analysis. We analyze the asymptotic behaviour of the model as $n \rightarrow \infty$.

1.3. Configuration model and exploration process. The *configuration model* plays a crucial role as a tool in our arguments in §3. Given $d, n \in \mathbb{N}$ such that dn is even, this model (see [5, 7]) allows us to generate a d -regular random graph on n labeled vertices v_1, \dots, v_n (possibly with self-loops and parallel edges) according to the following procedure, also known as the *exploration process*:

- (i) Fix a total order $v_1 < v_2 < \dots < v_n$ on the vertex set, and let $\Xi_i = \{\xi_{i,j} : 1 \leq j \leq d\}$ denote the set of *half-edges* emanating from v_i , for all $i = 1, \dots, n$. Let $\Xi = \bigcup_{i=1}^n \Xi_i$. We define a total ordering on Ξ as follows: all half-edges in Ξ_i come before every half-edge in Ξ_{i+1} for all $i = 1, \dots, n-1$, and within each Ξ_i , we have $\xi_{i,j} < \xi_{i,j+1}$ for all $1 \leq j \leq d-1$.
- (ii) We first choose $\hat{\xi}$ uniformly randomly from the set $\Xi \setminus \{\xi_{1,1}\}$ and form the edge $\{\hat{\xi}, \xi_{1,1}\}$. Having constructed the first k edges, we find the smallest half-edge $\xi_{i,j}$ yet unmatched

with another half-edge, and choosing a $\tilde{\xi}$ uniformly randomly from the remaining subset of half-edges, we form the edge $\{\xi_{i,j}, \tilde{\xi}\}$. Thus we form a perfect matching on Ξ .

We also describe here the exploration process aimed at generating a random d -regular bipartite graph in which each cluster contains n vertices. In this case, we label the vertices of one partition as $u_1 < u_2 < \dots < u_n$ and the other as $v_1 < v_2 < \dots < v_n$. We let $\Xi_i = \{\xi_{i,j} : 1 \leq j \leq d\}$ denote the set of half-edges emanating from u_i and $\Gamma_i = \{\gamma_{i,j} : 1 \leq j \leq n\}$ the set of half-edges emanating from v_i , and the total orderings on $\Xi = \bigcup_{i=1}^n \Xi_i$ and on $\Gamma = \bigcup_{i=1}^n \Gamma_i$ are analogous to the one described above. We first choose, uniformly randomly, a $\hat{\gamma}$ out of Γ , and form the edge $\{\hat{\gamma}, \xi_{1,1}\}$. After having constructed the k -th edge, we find the smallest $\xi_{i,j}$ in Ξ that is yet to be matched with a half-edge from Γ . We choose, uniformly randomly, a half-edge $\tilde{\gamma}$ from Γ that has not yet been matched, and form the edge $\{\tilde{\gamma}, \xi_{i,j}\}$. This leads to a perfect matching between Ξ and Γ .

It has been shown in [7] that in either of the cases above, the probability that the generated random graph is simple, i.e. devoid of self-loops and parallel edges, stays bounded away from 0 as d stays bounded and $n \rightarrow \infty$. We can thus condition on the event that the generated random graph is simple, which in turn allows us to prove the results in §3 using the exploration process.

Henceforth, we call a half-edge emanating from a vertex in \mathcal{C}_i and matched with a half-edge from a vertex in \mathcal{C}_j , a *half-edge of type* $\{i, j\}$, for all $i, j \in \{1, \dots, k\}$.

2. COMPARING RSBM WITH UNIFORM MEASURE ON d -REGULAR GRAPHS ON kn VERTICES

We state here our first main result. Let μ_A^n denote the probability measure of \mathcal{G}_n^A and \mathcal{S}_A^n the support of μ_A^n . Recall from §1.1 that μ_d^{kn} denotes the uniform random measure on \mathcal{R}_d^{kn} .

Theorem 2.1. *Under the above set-up, keeping the matrix A fixed, we have*

$$\lim_{n \rightarrow \infty} \left\| \mu_A^n, \mu_d^{kn} \right\|_{\text{TV}} = 1, \quad (2.1)$$

where TV denotes the total variation distance between two probability measures.

The proof begins with stating two well-known results. For given n and d with $1 \leq d = o(n^{1/2})$, [[33], Corollary 5.3] states that

$$|\mathcal{R}_d^n| = C \frac{(nd)!}{(nd/2)! 2^{nd/2} (d!)^n}, \quad (2.2)$$

where $C = C(n, d)$ remains bounded as n grows. Similarly, [[32], Theorem 2] states that

$$|\mathcal{B}_d^n| = C' \frac{(dn)!}{(d!)^{2n}}, \quad (2.3)$$

where $C' = C'(n, d)$ remains bounded as n grows. We now use these estimates to count the total number of possible realizations of \mathcal{G}_A^n . Given kn labeled vertices, we choose the vertex sets for the clusters $\mathcal{C}_1, \dots, \mathcal{C}_k$ in

$$\binom{kn}{n} \binom{(k-1)n}{n} \dots \binom{2n}{n} \binom{n}{n} = \frac{(kn)!}{(n!)^k} \sim \frac{\sqrt{2\pi(kn)} \left(\frac{kn}{e}\right)^{kn}}{\left\{ \sqrt{2\pi n} \left(\frac{n}{e}\right)^n \right\}^k} = \sqrt{\frac{k}{(2\pi n)^{k-1}}} k^{kn} = \Theta\left(\frac{k^{kn}}{n^{(k-1)/2}}\right)$$

many ways. The number of possible $a_{i,i}$ -regular graphs on \mathcal{C}_i , for each $i = 1, \dots, k$, equals, by (2.2),

$$\Theta\left(\frac{(na_{i,i})!}{(na_{i,i}/2)! 2^{na_{i,i}/2} (a_{i,i}!)^n}\right) = \Theta\left(\frac{\sqrt{2\pi(a_{i,i}n)} \left(\frac{a_{i,i}n}{e}\right)^{a_{i,i}n}}{\sqrt{\pi(a_{i,i}n)} \left(\frac{a_{i,i}n}{2e}\right)^{a_{i,i}n/2} 2^{na_{i,i}/2} \left\{ \sqrt{2\pi a_{i,i}} \left(\frac{a_{i,i}}{e}\right)^{a_{i,i}} \right\}^n}\right)$$

$$\begin{aligned}
&= \Theta \left(\frac{a_{i,i}^{a_{i,i}n} n^{a_{i,i}n} e^{-a_{i,i}n}}{a_{i,i}^{a_{i,i}n/2} n^{a_{i,i}n/2} 2^{-a_{i,i}n/2} e^{-a_{i,i}n/2} 2^{a_{i,i}n/2} (2\pi a_{i,i})^{n/2} a_{i,i}^{a_{i,i}n} e^{-a_{i,i}n}} \right) \\
&= \Theta \left(\frac{n^{a_{i,i}n/2} e^{a_{i,i}n/2}}{a_{i,i}^{(a_{i,i}+1)n/2} (2\pi)^{n/2}} \right).
\end{aligned}$$

Similarly, by (2.3), the number of possible $a_{i,j}$ -bipartite-regular graphs across clusters \mathcal{C}_i and \mathcal{C}_j , for each $i \neq j$ and $i, j \in \{1, \dots, k\}$, is given by

$$\Theta \left(\frac{(a_{i,j}n)!}{(a_{i,j}!)^{2n}} \right) = \Theta \left(\frac{\sqrt{2\pi(a_{i,j}n)} \left(\frac{a_{i,j}n}{e}\right)^{a_{i,j}n}}{\left\{ \sqrt{2\pi a_{i,j}} \left(\frac{a_{i,j}}{e}\right)^{a_{i,j}} \right\}^{2n}} \right) = \Theta \left(\frac{\sqrt{n} a_{i,j}^{a_{i,j}n} n^{a_{i,j}n} e^{-a_{i,j}n}}{(2\pi a_{i,j})^n a_{i,j}^{2a_{i,j}n} e^{-2a_{i,j}n}} \right) = \Theta \left(\frac{\sqrt{n} n^{a_{i,j}n} e^{a_{i,j}n}}{(2\pi)^n a_{i,j}^{(a_{i,j}+1)n}} \right).$$

Therefore, combining these estimates, the total number of possible realizations of \mathcal{G}_A^n on a given set of kn labeled vertices becomes

$$\begin{aligned}
&\Theta \left(\frac{k^{kn}}{n^{(k-1)/2}} \prod_{i=1}^k \frac{n^{a_{i,i}n/2} e^{a_{i,i}n/2}}{a_{i,i}^{(a_{i,i}+1)n/2} (2\pi)^{n/2}} \prod_{i < j} \frac{\sqrt{n} n^{a_{i,j}n} e^{a_{i,j}n}}{(2\pi)^n a_{i,j}^{(a_{i,j}+1)n}} \right) \\
&= \Theta \left(\frac{k^{kn} n^{\sum_{i=1}^k n/2 \sum_{j=1}^k a_{i,j}} e^{\sum_{i=1}^k n/2 \sum_{j=1}^k a_{i,j}} n^{k(k-1)/4}}{n^{(k-1)/2} \prod_{i,j} a_{i,j}^{(a_{i,j}+1)n/2} (2\pi)^{nk/2} (2\pi)^{k(k-1)n/2}} \right) = \Theta \left(\frac{k^{kn} n^{knd/2} e^{knd/2} n^{(k^2-3k+2)/4}}{\prod_{i,j} a_{i,j}^{(a_{i,j}+1)n/2} (2\pi)^{nk^2/2}} \right). \tag{2.4}
\end{aligned}$$

On the other hand, from (2.2), the number of d -regular graphs on kn vertices is

$$\Theta \left(\frac{(knd)!}{(knd/2)! 2^{knd/2} (d!)^{kn}} \right) = \Theta \left(\frac{\sqrt{2\pi(knd)} \left(\frac{knd}{e}\right)^{knd}}{\sqrt{2\pi(knd/2)} \left(\frac{knd}{2e}\right)^{knd/2} 2^{knd/2} \left(\sqrt{2\pi d} \left(\frac{d}{e}\right)^d\right)^{kn}} \right) = \Theta \left(\frac{(knd)^{knd/2} e^{knd/2}}{(2\pi)^{kn/2} d^{(d+1/2)kn}} \right). \tag{2.5}$$

We recall here the well-known weighted geometric mean – harmonic mean inequality. Given $m \in \mathbb{N}$ and positive reals x_i and α_i for all $i \in \{1, \dots, m\}$ with $\sum_{i=1}^m \alpha_i = 1$, we have

$$\prod_{i=1}^m x_i^{\alpha_i} \geq \left(\sum_{i=1}^m \frac{\alpha_i}{x_i} \right)^{-1}. \tag{2.6}$$

Setting $x_j = a_{i,j}$ and $\alpha_j = \frac{a_{i,j}}{d}$ for all $j \in \{1, \dots, k\}$, from (2.6), we get

$$\prod_{j=1}^k a_{i,j}^{a_{i,j}/d} \geq \frac{d}{k} \implies \prod_{j=1}^k a_{i,j}^{a_{i,j}} \geq \frac{d^d}{k^d}$$

for each $i = 1, \dots, k$. This in turn yields

$$\left\{ \prod_{i,j} a_{i,j}^{a_{i,j}} \right\}^{n/2} \geq \frac{d^{knd/2}}{k^{knd/2}}. \tag{2.7}$$

We also observe that given positive integers x_1, \dots, x_k for any $k \in \mathbb{N}$, the following inequality holds:

$$k \prod_{i=1}^k x_i \geq \sum_{i=1}^k x_i. \tag{2.8}$$

From (2.4), (2.5), (2.7) and (2.8), we see that

$$\begin{aligned}
\mu_d^{kn}(\mathcal{S}_A^n) &\leq \Theta \left(\frac{k^{kn} n^{knd/2} e^{knd/2} n^{(k^2-3k+2)/4}}{\prod_{i,j} a_{i,j}^{(a_{i,j}+1)n/2}} \frac{(2\pi)^{kn/2} d^{(d+1/2)kn}}{(knd)^{knd/2} e^{knd/2}} \right) \\
&\leq \Theta \left(\frac{k^{kn} n^{(k^2-3k+2)/4} d^{knd/2} d^{kn/2}}{\left\{ \prod_{i,j} a_{i,j}^{a_{i,j}} \right\}^{n/2} \left\{ \prod_{i,j \in [k]} a_{i,j} \right\}^{n/2} (2\pi)^{k(k-1)n/2} k^{knd/2}} \right) \\
&\leq \Theta \left(\frac{k^{kn} n^{(k^2-3k+2)/4} d^{kn/2}}{\left\{ \prod_{i,j} a_{i,j} \right\}^{n/2} (2\pi)^{k(k-1)n/2}} \right) = \Theta \left(\left\{ \frac{k^3}{(2\pi)^{k-1}} \right\}^{kn/2} \cdot n^{(k^2-3k+2)/4} \right). \quad (2.9)
\end{aligned}$$

The ratio $\frac{k^3}{(2\pi)^{k-1}}$ is strictly less than 1 for all k that satisfy $\frac{k-1}{\log k} > \frac{3}{\log(2\pi)}$. Now, the function $f(x) = \frac{x-1}{\log x}$ is strictly increasing in x for all $x \geq 2$, and $f(3) = \frac{2}{\log 3} > \frac{3}{\log(2\pi)}$. This shows that the ratio $\frac{k^3}{(2\pi)^{k-1}}$ is strictly less than 1 for all $k \geq 3$, thus showing that the bound in (2.9) is $o(1)$.

3. ALMOST SURE UNIQUENESS OF CLUSTERS IN RSBM

This section is devoted to establishing fairly general sufficient conditions under which the communities involved in the regular stochastic block model are unique asymptotically almost surely as the size of each community approaches infinity. This is what constitutes our second main result.

Theorem 3.1. *Suppose the model described in §1.2 satisfies the following conditions:*

- (i) *There exists a positive constant C such that $Ca_{i,i} > B_i$ for each $i \in [k]$, where $B_i = \max \{a_{i,\ell} : \ell \in \{1, \dots, k\} \setminus \{i\}\}$.*
- (ii) *When not all intra-cluster degrees are equal, for $i, j \in \{1, \dots, k\}$ with $a_{i,i} < a_{j,j}$, there exist constants $\delta_{i,j} \in (0, \frac{1}{4})$, independent of all entries of A , such that*

$$a_{i,i} \geq \left(\frac{1}{2} + 2\delta_{i,j} \right) a_{j,j}. \quad (3.1)$$

Moreover, there exist constants $\epsilon_{i,j} \in (0, \frac{1}{4})$, independent of the entries of A , such that for all $i, j \in \{1, \dots, k\}$ with $a_{i,i} > a_{j,j}$ and $B_j > a_{i,i} - a_{j,j}$,

$$\left(\frac{1}{2} - 2\epsilon_{i,j} \right) a_{i,i} \geq B_j. \quad (3.2)$$

Then, for all sufficiently large values of the entries of the matrix A , the clusters \mathcal{C}_i , $i = 1, \dots, k$, are a.a.s. unique as $n \rightarrow \infty$ while the matrix A stays fixed.

From the discussion in §1.3, it suffices to establish Theorem 3.1 on the random multigraph in which each of the intra-cluster regular graphs and inter-cluster bipartite-regular graphs is generated via the configuration model. Fix *any* non-negative $\alpha_1, \dots, \alpha_k$ such that $\sum_{i=1}^k \alpha_i = 1$, and subsets C_i of \mathcal{C}_i such that $|C_i| = \alpha_i n$ for $i = 1, \dots, k$. Let $D = \bigcup_{i=1}^k C_i$. To prove Theorem 3.1, it is enough to establish Proposition 3.2. We note here that although Proposition 3.2 is stated for $a_{1,1}$, its proof will be analogous if we replace $a_{1,1}$ by any $a_{i,i}$ for $i \in \{2, \dots, k\}$.

Proposition 3.2. *Assume that the hypotheses of Theorem 3.1 hold, and that the entries of A are sufficiently large. Suppose there exist at least two distinct $i, j \in \{1, \dots, k\}$ such that α_i and α_j are strictly positive. Then a.a.s. the following cannot be true simultaneously:*

- (i) the subgraph $\mathcal{G}|_D$ is $a_{1,1}$ -regular;
- (ii) there exists a partition of $V(G) \setminus D$ into subsets D_2, \dots, D_k , each of size n , such that $\mathcal{G}|_{D_i}$ is $a_{i,i}$ -regular, $\mathcal{G}|_{D_i, D_j}$ is $a_{i,j}$ -bipartite-regular and $\mathcal{G}|_{D, D_i}$ is $a_{1,i}$ -bipartite-regular for all distinct $i, j \in \{2, \dots, k\}$.

In order to prove Proposition 3.2, we start with the assumption that $\mathcal{G}|_D$ is $a_{1,1}$ -regular. The proof requires consideration of a few different cases depending on the values of the α_i 's, and these are addressed in Lemma 3.3, §3.1, §3.2 and §3.3.

Lemma 3.3. *If $a_{i,i} > a_{1,1}$ and $\alpha_i > \frac{B_1}{a_{i,i} - a_{1,1} + B_1}$, then the conclusion of Proposition 3.2 holds.*

Proof. Assume that $\alpha_i > 0$ for some i such that $a_{i,i} > a_{1,1}$, and that Proposition 3.2 does not hold. As $\mathcal{G}|_D$ is $a_{1,1}$ -regular, we have $\deg_{\mathcal{C}_i \setminus C_i}(v) \geq (a_{i,i} - a_{1,1})$ for every $v \in C_i$, yielding $\deg(C_i, \mathcal{C}_i \setminus C_i) \geq (a_{i,i} - a_{1,1})\alpha_i n$. On the other hand, each u in $\mathcal{C}_i \setminus C_i$ belongs to precisely one of the remaining clusters D_2, \dots, D_k . If $u \in D_j$, then $\deg_{\mathcal{C}_i}(u) \leq \deg_D(u) = a_{1,j}$. Thus $\deg(C_i, \mathcal{C}_i \setminus C_i) \leq B_1(1 - \alpha_i)n$. These two inequalities together yield

$$(a_{i,i} - a_{1,1})\alpha_i \leq B_1(1 - \alpha_i) \quad \implies \quad \alpha_i \leq \frac{B_1}{a_{i,i} - a_{1,1} + B_1},$$

thus completing the proof. \square

From here onward, we only consider those $\ell \in \{1, \dots, k\}$ such that $\alpha_\ell > 0$, without mentioning so every time. We shall let i denote that index in $\{1, \dots, k\}$ (if this is not unique, we choose any such i and fix it) for which $\alpha_i \geq \alpha_\ell$ for all $\ell \in \{1, \dots, k\} \setminus \{i\}$. Note that this guarantees, by the pigeon hole principle, that $\alpha_i \geq \frac{1}{k}$.

Under the assumption that $\mathcal{G}|_D$ is $a_{1,1}$ -regular, we have $\sum_{\ell=1}^k \deg_{C_j}(v) = a_{1,1}$ for every $v \in C_i$. On the other hand, $\deg_{C_i}(v) + \deg_{\mathcal{C}_i \setminus C_i}(v) = a_{i,i}$ for each $v \in C_i$. These together imply

$$\sum_{\ell \in \{1, \dots, k\} \setminus \{i\}} \deg_{C_\ell}(v) = a_{1,1} - a_{i,i} + \deg_{\mathcal{C}_i \setminus C_i}(v). \quad (3.3)$$

To prove Proposition 3.2, we first condition on the σ -field \mathcal{F} comprising the following information:

- (i) the vertex sets of \mathcal{C}_ℓ and C_ℓ for all $\ell = 1, \dots, k$,
- (ii) the subgraph $\mathcal{G}|_{\mathcal{C}_i}$ induced on \mathcal{C}_i .

Given \mathcal{F} , we enumerate the vertices of \mathcal{C}_i as v_1, \dots, v_n such that $C_i = \{v_1, \dots, v_{\alpha_i n}\}$. From (3.3), we set

$$g_s = a_{1,1} - a_{i,i} + \deg_{\mathcal{C}_i \setminus C_i}(v_s) \quad (3.4)$$

for all $s = 1, \dots, \alpha_i n$. The random variables g_s are measurable with respect to \mathcal{F} . The conditional probability of the event that $\mathcal{G}|_D$ is $a_{1,1}$ -regular is bounded above by the conditional probability of the event

$$A = \left\{ \sum_{\ell \in \{1, \dots, k\} \setminus \{i\}} \deg_{C_\ell}(v_s) = g_s \text{ for all } s = 1, \dots, \alpha_i n \right\}.$$

We show that the probability of the event A is $o(1)$ as $n \rightarrow \infty$.

First, we express A as the union of pairwise disjoint events. For $g \in \mathbb{N}$, let us define the following subset of ordered $(k-1)$ -tuples of non-negative integers:

$$\mathcal{S}_g = \left\{ (m_1, \dots, m_{i-1}, m_{i+1}, \dots, m_k) : 0 \leq m_\ell \leq a_{i,\ell} \text{ for all } \ell \in \{1, \dots, k\} \setminus \{i\}, \sum_{\ell \in \{1, \dots, k\} \setminus \{i\}} m_\ell = g \right\}.$$

Then A can be written as the union of the events

$$A \left(m^{(s)} : s = 1, \dots, \alpha_i n \right) = \left\{ \deg_{C_\ell}(v_s) = m_\ell^{(s)} \text{ for all } \ell \in \{1, \dots, k\} \setminus \{i\} \text{ and } s = 1, \dots, \alpha_i n \right\}$$

where $m^{(s)} = \left(m_\ell^{(s)} : \ell \in \{1, \dots, k\} \setminus \{i\} \right)$ belongs to \mathcal{S}_{g_s} for all $s = 1, \dots, \alpha_i n$. Note, from the mutual independence of the subgraphs $\mathcal{G}_{|C_i, C_\ell}$ for $\ell \in \{1, \dots, k\} \setminus \{i\}$, that

$$\begin{aligned} \mathbf{P} \left[A \left(m^{(s)} : s = 1, \dots, \alpha_i n \right) \middle| \mathcal{F} \right] &= \prod_{\ell \in \{1, \dots, k\} \setminus \{i\}} \mathbf{P} \left[\deg_{C_\ell}(v_s) = m_\ell^{(s)} \text{ for all } s = 1, \dots, \alpha_i n \middle| \mathcal{F} \right] \\ &\leq \mathbf{P} \left[\deg_{C_\ell}(v_s) = m_\ell^{(s)} \text{ for all } s = 1, \dots, \alpha_i n \middle| \mathcal{F} \right] \end{aligned} \quad (3.5)$$

for each $\ell \in \{1, \dots, k\} \setminus \{i\}$. The goal now is to fix *any* $m^{(s)} \in \mathcal{S}_{g_s}$ for each s and establish that the probability of the event $\left\{ \deg_{C_\ell}(v_s) = m_\ell^{(s)} \text{ for all } s = 1, \dots, \alpha_i n \right\}$ for at least one $\ell \in \{1, \dots, k\} \setminus \{i\}$ is $o(n^{-1})$ as $n \rightarrow \infty$.

Since $m^{(s)} \in \mathcal{S}_{g_s}$ for each s , from (3.4), we have

$$\sum_{\ell \in \{1, \dots, k\} \setminus \{i\}} \sum_{s=1}^{\alpha_i n} m_\ell^{(s)} = \sum_{s=1}^{\alpha_i n} \sum_{\ell \in \{1, \dots, k\} \setminus \{i\}} m_\ell^{(s)} = \sum_{s=1}^{\alpha_i n} g_s = \deg(C_i, C_i \setminus C_i) + (a_{1,1} - a_{i,i})\alpha_i n. \quad (3.6)$$

For G uniformly randomly chosen from \mathcal{R}_d^n , [[18], Theorem 1.1] showed that $\gamma \geq 1 - \frac{2}{\sqrt{d}}$ a.a.s. as $n \rightarrow \infty$, where γ is the spectral gap for the adjacency matrix of G . Given a d -regular graph G on n vertices and a subset S of $V(G)$ with $|S| \leq \frac{n}{2}$, [[29], Theorem 13.14] (see also [23], [26], and [[10], Theorem 6]) established that

$$\frac{\gamma}{2} \leq \frac{\deg(S, V(G) \setminus S)}{d|S|}.$$

Combining these, we get

$$\deg(C_i \setminus C_i, C_i) \geq \min\{\alpha_i, 1 - \alpha_i\} \left(\frac{1}{2} - \frac{1}{\sqrt{a_{i,i}}} \right) a_{i,i} n. \quad (3.7)$$

From (3.6) and (3.7), we get

$$\sum_{\ell \in \{1, \dots, k\} \setminus \{i\}} \sum_{s=1}^{\alpha_i n} m_\ell^{(s)} \geq \min\{\alpha_i, 1 - \alpha_i\} \left(\frac{1}{2} - \frac{1}{\sqrt{a_{i,i}}} \right) a_{i,i} n + (a_{1,1} - a_{i,i})\alpha_i n.$$

By the pigeon-hole principle, there exists at least one $j \in \{1, \dots, k\} \setminus \{i\}$ such that

$$\sum_{s=1}^{\alpha_i n} m_j^{(s)} \geq \frac{1}{k-1} \left\{ \min\{\alpha_i, 1 - \alpha_i\} \left(\frac{1}{2} - \frac{1}{\sqrt{a_{i,i}}} \right) a_{i,i} n + (a_{1,1} - a_{i,i})\alpha_i n \right\}. \quad (3.8)$$

For the rest of the proof, we fix such a j , and establish the following lemma:

Lemma 3.4. *Let A_j denote the event that there exist $C_i = \{v_1, \dots, v_{\alpha_i n}\} \subset \mathcal{C}_i$ and $C_j \subset \mathcal{C}_j$ such that $\deg_{C_j}(v_s) = m_j^{(s)}$ for all $s = 1, \dots, \alpha_i n$. Then $\mathbf{P}[A_j] = o(n^{-1})$.*

The proof of this lemma is accomplished through the consideration of three different cases, in §3.1, §3.2 and §3.3.

3.1. **When** $a_{1,1} \geq a_{i,i}$. We note at the very outset that the analysis of §3.1 is enough for the special and commonly studied situation where all intra-cluster degrees are the same. We set $G_j = \sum_{s=1}^{\alpha_i n} m_j^{(s)}$, so that from (3.8), we have, for all $a_{i,i}$ sufficiently large,

$$\frac{G_j}{n} \geq \frac{\min\{\alpha_i, 1 - \alpha_i\} a_{i,i}}{4(k-1)}. \quad (3.9)$$

We refer the reader to [[9], Lemma 2] for the following inequality:

$$\begin{aligned} & \mathbf{P} \left[\deg_{C_j}(v_s) = m_j^{(s)} \text{ for all } s = 1, \dots, \alpha_i n \mid \mathcal{F} \right] \\ & \leq \mathbf{P} \left[\deg_{C_j}(v_s) \in \{\eta_j, \eta_j + 1\} \text{ for all } s = 1, \dots, \alpha_i n, \sum_{s=1}^{\alpha_i n} \deg_{C_j}(v_s) = \sum_{s=1}^{\alpha_i n} m_j^{(s)} \mid \mathcal{F} \right], \end{aligned} \quad (3.10)$$

where $\eta_j = \lfloor \frac{G_j}{\alpha_i n} \rfloor$. Notice that we have the trivial bound $G_j = \deg(C_i, C_j) \leq \deg(C_j, C_i) = \alpha_j a_{i,j} n$, so that

$$\eta_j \leq \frac{\alpha_j a_{i,j}}{\alpha_i}. \quad (3.11)$$

We invoke the configuration model discussed in §1.3, and outline, in the next paragraph, some foundational aspects of the argument that resemble [[9], Lemma 5]. Let $\xi_{a_{i,j}(s-1)+1}, \dots, \xi_{a_{i,j}s}$ denote the half-edges of type $\{i, j\}$ emanating from vertex v_s , for each $s = 1, \dots, \alpha_i n$. Let B_t denote the indicator random variable of the event that ξ_t is matched with a half-edge of type $\{i, j\}$ emanating from C_j . Conditioned on B_1, \dots, B_t , the random variable B_{t+1} is Bernoulli with probability

$$\hat{p}_t = \frac{\alpha_j a_{i,j} n - \sum_{t' \leq t} B_{t'}}{a_{i,j} n - t}.$$

For all $1 \leq t \leq \alpha_i n$, we see that $|\hat{p}_t - \hat{p}_{t-1}| \leq O(n^{-1})$, so that for all $s = 1, \dots, \alpha_i n$, there exists $p_s \in (0, 1)$ such that

$$\left| \deg_{C_j}(v_s) \mid \mathcal{G}_{s-1}, \text{Bin}(a_{i,j}, p_s) \right|_{\text{TV}} = O\left(\frac{1}{n}\right), \quad (3.12)$$

where \mathcal{G}_s denotes the σ -field generated by $\{\xi_t, 1 \leq t \leq a_{i,j}s\}$, for each s . Given that each of $\deg_{C_j}(v_1), \dots, \deg_{C_j}(v_{s-1})$ takes values in $\{\eta_j, \eta_j + 1\}$, the number of half-edges emanating from C_j that have not yet been matched is at least $\alpha_j a_{i,j} n - (\eta_j + 1)(s-1)$ and at most $\alpha_j a_{i,j} n - \eta_j(s-1)$. The number of half-edges from C_j that are left to be matched is $a_{i,j}(n - s + 1)$. Thus

$$\frac{\alpha_j a_{i,j} n - (\eta_j + 1)(s-1)}{a_{i,j}(n - s + 1)} \leq p_s \leq \frac{\alpha_j a_{i,j} n - \eta_j(s-1)}{a_{i,j}(n - s + 1)}. \quad (3.13)$$

We shall now consider three different ranges of values of α_j , where j is as chosen by (3.8). First, consider

$$\frac{c}{a_{i,j}} < \alpha_j \leq \frac{1}{2}, \quad (3.14)$$

where

$$\log_2 c > \max \left\{ 16k + 1, \frac{2(k-1)C}{4^{8k-1}} \right\}, \quad (3.15)$$

where C is as in i of Theorem 3.1. We first consider the case of $\eta_j \neq 0$. For each $s = 1, \dots, \lfloor \frac{\alpha_i n}{4} \rfloor$, from (3.11) and (3.13), we have

$$p_s \geq \frac{\alpha_j a_{i,j} n - 2\eta_j \cdot \frac{\alpha_i n}{4}}{a_{i,j} n} \geq \frac{\alpha_j a_{i,j} n - 2 \cdot \frac{\alpha_j a_{i,j}}{\alpha_i} \cdot \frac{\alpha_i n}{4}}{a_{i,j} n} = \frac{\alpha_j}{2} > \frac{c}{2a_{i,j}}. \quad (3.16)$$

Using (3.12), (3.16), the fact that the mode of the $\text{Bin}(n, p)$ distribution is $\lfloor (n+1)p \rfloor$, and the same argument as in [[9], Lemma 4], we conclude that, for all $s = 1, \dots, \lfloor \frac{\alpha_i n}{4} \rfloor$,

$$\begin{aligned} \mathbf{P} \left[\deg_{C_j}(v_s) \in \{\eta_j, \eta_j + 1\} \middle| \mathcal{G}_{s-1} \right] &\leq \mathbf{P} [\text{Bin}(a_{i,j}, p_s) = \lfloor (a_{i,j} + 1)p_s \rfloor] + O\left(\frac{1}{n}\right) \\ &\leq O\left(\sqrt{\frac{1}{a_{i,j}p_s}}\right) \leq O\left(\sqrt{\frac{2}{c}}\right). \end{aligned} \quad (3.17)$$

We now use union bounds and Stirling's approximation to bound above $\mathbf{P}[A_j]$ by

$$\begin{aligned} \binom{n}{\alpha_i n} \binom{n}{\alpha_j n} \left\{ O\left(\sqrt{\frac{2}{c}}\right) \right\}^{\frac{\alpha_i n}{4}} &\sim O\left(\frac{2^{H(\alpha_i)n + H(\alpha_j)n}}{\sqrt{\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j)n}} \left(\frac{2}{c}\right)^{\frac{\alpha_i n}{8}}\right) \\ &\leq O\left(\frac{2^{2n}}{\sqrt{\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j)n}} \left(\frac{2}{c}\right)^{\frac{n}{8k}}\right) \end{aligned}$$

which is $o(n^{-1})$ by (3.15).

Now we consider the case of $\eta_j = 0$. We enumerate the vertices of C_j so that $\deg_{C_j}(v_s) = 1$ for $s = 1, \dots, G_j$ and $\deg_{C_j}(v_s) = 0$ for all $G_j + 1 \leq s \leq \alpha_i n$. Inspired by the lower bound in (3.13), set

$$f(s-1) = \frac{\alpha_j a_{i,j} n - s + 1}{a_{i,j}(n - s + 1)}, \text{ for all } s \in [G_j].$$

The function f being strictly increasing for α_j as in (3.14), a uniform lower bound on p_s for all $s = 1, \dots, G_j$ is $f(0) = \alpha_j > \frac{c}{a_{i,j}}$. By similar computations as used in deriving in (3.17), we get

$$\mathbf{P} \left[\deg_{C_j}(v_s) = 1 \middle| \mathcal{G}_{s-1} \right] \leq O\left(\sqrt{\frac{1}{c}}\right) \quad (3.18)$$

for all $s = 1, \dots, G_j$. Note that $\alpha_j \leq \alpha_i$ by our choice of i , and $\alpha_i + \alpha_j \leq 1$ implies that $\alpha_j \leq 1 - \alpha_i$. Hence, from (3.9), (3.14) and (3.15) we conclude that

$$\frac{G_j}{n} \geq \frac{\alpha_j a_{i,i}}{4(k-1)} > \frac{c a_{i,i}}{4(k-1)a_{i,j}} > \frac{2 \cdot 4^{8k-1}}{(k-1)C}. \quad (3.19)$$

In this case, the upper bound on $\mathbf{P}[A_j]$ is given by

$$\binom{n}{\alpha_i n} \binom{n}{\alpha_j n} \left\{ O\left(\sqrt{\frac{1}{c}}\right) \right\}^{G_j} \leq O\left(\frac{2^{2n}}{\sqrt{\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j)n}} \left(\frac{1}{c}\right)^{\frac{4^{8k-1}n}{(k-1)C}}\right)$$

which is $o(n^{-1})$ due to (3.15).

Next, we consider the following range of values of α_j :

$$\frac{1}{a_{i,j}^2} < \alpha_j \leq \frac{c}{a_{i,j}}, \quad (3.20)$$

where c is as in (3.15). At the very outset of this case, we note that, if $\alpha_i \leq \frac{1}{2}$, we must have

$$c \geq \alpha_j a_{i,j} \geq \frac{G_j}{n} \geq \frac{\alpha_i a_{i,i}}{4(k-1)},$$

implying that $\alpha_i \leq \frac{4(k-1)c}{\alpha_i a_{i,i}}$. For all $a_{i,i}$ sufficiently large, this upper bound is smaller than $\frac{1}{k}$, giving us a contradiction.

Remark 3.5. The above reasoning shows that for the range $\alpha_j \leq \frac{c}{a_{i,j}}$, we need not consider $\alpha_i \leq \frac{1}{2}$.

When $\alpha_i > \frac{1}{2}$, we have, by similar reasoning as above,

$$c \geq \frac{(1 - \alpha_i)a_{i,i}}{4(k-1)} \implies \alpha_i \geq 1 - \frac{4c(k-1)}{a_{i,i}}. \quad (3.21)$$

For all $a_{i,i}$ sufficiently large, this yields:

$$\alpha_j < 1 - \alpha_i \leq \frac{4c(k-1)}{a_{i,i}} < \frac{1}{2} < 1 - \frac{4c(k-1)}{a_{i,i}} \leq \alpha_i < 1 - \alpha_j, \quad (3.22)$$

and by the concave nature of the entropy function and its symmetry around $\frac{1}{2}$, we conclude that

$$H(\alpha_i) + H(\alpha_j) \leq 2H\left(\frac{4c(k-1)}{a_{i,i}}\right). \quad (3.23)$$

We first address the case of $\eta_j \neq 0$. From (3.13) and (3.20), for all $s = 1, \dots, \lfloor \frac{\alpha_i n}{2} \rfloor$, we get

$$p_s \leq \frac{\alpha_j a_{i,j} n}{a_{i,j} \left(n - \frac{\alpha_i n}{2}\right)} < \frac{2c}{a_{i,j}}, \quad (3.24)$$

so that $\deg_{C_j}(v_s)$, conditioned on \mathcal{G}_{s-1} , is stochastically dominated by $\text{Bin}\left(a_{i,j}, \frac{2c}{a_{i,j}}\right)$, which in turn can be approximated by the $\text{Poisson}(2c)$ distribution. Thus

$$\mathbf{P}\left[\deg_{C_j}(v_s) \in \{\eta_j, \eta_{j+1}\} \mid \mathcal{G}_{s-1}\right] \leq \mathbf{P}[\text{Poisson}(2c) \geq 1] = \gamma \quad (3.25)$$

where γ is a constant that depends only on c . Using (3.23) and (3.25), and $\frac{\alpha_i n}{2} > \frac{n}{4}$, we get the following upper bound on $\frac{1}{n} \log_2 \mathbf{P}[A_j]$:

$$-\frac{\log_2(\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j))}{2n} - \frac{\log_2 n}{n} + 2H\left(\frac{4c(k-1)}{a_{i,i}}\right) + \frac{\log_2 \gamma}{4}. \quad (3.26)$$

The first two terms approach 0 as $n \rightarrow \infty$. The last term is a strictly negative constant, and as $a_{i,i}$ grows, the third term goes to 0. Hence (3.26) is strictly negative for all sufficiently large $a_{i,i}$, as $n \rightarrow \infty$.

Remark 3.6. Observe that, in the above argument, nowhere has the lower bound on α_j from (3.20) been used. This shows that as far as the case of $\eta_j \neq 0$ is concerned, our proof of Lemma 3.4 for the regime of §3.1 ends here. In the rest of §3.1, we only consider $\eta_j = 0$.

Now, we consider $\eta_j = 0$ and α_j in the range given by (3.20). If $G_j \geq \frac{\alpha_i n}{2}$, then the same argument as above will be enough.

Remark 3.7. This shows that for all $\alpha_j \leq \frac{c}{a_{i,j}}$ and $\eta_j = 0$, as long as $G_j \geq \frac{\alpha_i n}{2}$, our proof of Lemma 3.4 is already complete. Henceforth, we only consider $\eta_j = 0$ and $G_j < \frac{\alpha_i n}{2}$.

If $G_j < \frac{\alpha_i n}{2}$, then for each $s = 1, \dots, G_j$, the bound in (3.24) holds, and hence so does (3.25). Together with (3.9) and (3.22), this yields the following upper bound on $\frac{1}{n} \log_2 \mathbf{P}[A_j]$:

$$-\frac{\log_2(\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j))}{2n} - \frac{\log_2 n}{n} + 2H(\alpha_i) + \frac{(1 - \alpha_i)a_{i,i}}{4(k-1)} \log_2 \gamma.$$

Again, the first two terms approach 0 as $n \rightarrow \infty$. We focus on the last two terms. Using the lower bound on α_j from (3.20) and the fact that $x \log x > (1 - x) \log(1 - x)$ for all $x \in (\frac{1}{2}, 1)$, we get:

$$2H(\alpha_i) + \frac{(1 - \alpha_i)a_{i,i}}{4(k-1)} \log_2 \gamma \leq -4(1 - \alpha_i) \log_2(1 - \alpha_i) + \frac{(1 - \alpha_i)a_{i,i} \log_2 \gamma}{4(k-1)}$$

$$\begin{aligned}
&\leq 8(1 - \alpha_i) \log_2 a_{i,j} + \frac{(1 - \alpha_i) a_{i,i} \log_2 \gamma}{4(k-1)} \\
&\leq 8(1 - \alpha_i) \{ \log_2 C + \log_2 a_{i,i} \} + \frac{(1 - \alpha_i) a_{i,i} \log_2 \gamma}{4(k-1)} \\
&= (1 - \alpha_i) \left\{ 8 \log_2 C + 8 \log_2 a_{i,i} + \frac{a_{i,i} \log_2 \gamma}{4(k-1)} \right\}.
\end{aligned}$$

As $a_{i,i}$ grows to ∞ much faster than $\log_2 a_{i,i}$, and the coefficient of $a_{i,i}$ is a strictly negative constant whereas that of $\log_2 a_{i,i}$ is a positive one, hence this is strictly negative for all $a_{i,i}$ sufficiently large.

Finally, we consider

$$\alpha_j \leq \frac{1}{a_{i,j}^2}, \quad (3.27)$$

and by Remarks 3.5 and 3.6, we need only consider $\alpha_i > \frac{1}{2}$ and $\eta_j = 0$. From (3.13), for all $s = 1, \dots, \lfloor \frac{\alpha_i n}{2} \rfloor$, we have

$$p_s \leq \frac{\alpha_j a_{i,j} n}{a_{i,j} (n - \frac{\alpha_i n}{2})} < 2\alpha_j \leq 2(1 - \alpha_i),$$

so that

$$\mathbf{P} \left[\deg_{C_j}(v_s) \in \{\eta_j, \eta_j + 1\} \mid \mathcal{F}_{s-1} \right] \leq \mathbf{P} [\text{Bin}(a_{i,j}, p_s) \geq 1] + O(n^{-1}) \leq a_{i,j} p_s < 2(1 - \alpha_i) a_{i,j}. \quad (3.28)$$

By Remark 3.7, we need only consider $G_j < \frac{\alpha_i n}{2}$, so that (3.28) holds for all $s = 1, \dots, G_j$. By (3.9) and (3.27), we get:

$$\frac{1}{a_{i,j}} \geq \alpha_j a_{i,j} \geq \frac{G_j}{n} \geq \frac{(1 - \alpha_i) a_{i,i}}{4(k-1)} \implies 1 - \alpha_i \leq \frac{4(k-1)}{a_{i,i} a_{i,j}}. \quad (3.29)$$

For any fixed positive integer $r > 2$, for all $a_{i,i}$ sufficiently large, by i of Theorem 3.1, we have

$$a_{i,i} > C^{\frac{1}{r-2}} \{4(k-1)\}^{\frac{r-1}{r-2}} 2^{\frac{r}{r-2}} \implies a_{i,i}^{r-1} \geq \{4(k-1)\}^{r-1} 2^r C a_{i,i} \geq \{4(k-1)\}^{r-1} 2^r a_{i,j},$$

so that by (3.29) we get

$$\log_2 \{2(1 - \alpha_i) a_{i,j}\} \leq \frac{\log_2(1 - \alpha_i)}{r}. \quad (3.30)$$

By (3.9) and (3.28), we get the following upper bound on $\frac{1}{n} \log_2 \mathbf{P}[A_j]$:

$$-\frac{\log_2(\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j))}{2n} - \frac{\log_2 n}{n} + 2H(\alpha_i) + \frac{(1 - \alpha_i) a_{i,i}}{4(k-1)} \log_2 \{2(1 - \alpha_i) a_{i,j}\}.$$

Again, it suffices to focus on the last two terms, and by (3.30), we get the following bound:

$$\begin{aligned}
2H(\alpha_i) + \frac{(1 - \alpha_i) a_{i,i}}{4(k-1)} \log_2 \{2(1 - \alpha_i) a_{i,j}\} &\leq -4(1 - \alpha_i) \log_2(1 - \alpha_i) + \frac{a_{i,i}}{4(k-1)} (1 - \alpha_i) \frac{\log_2(1 - \alpha_i)}{r} \\
&= (1 - \alpha_i) \log_2(1 - \alpha_i) \left\{ -4 + \frac{a_{i,i}}{4(k-1)r} \right\},
\end{aligned}$$

which is strictly negative for all $a_{i,i}$ sufficiently large. This brings us to the end of §3.1.

3.2. **When** $a_{1,1} < a_{i,i}$ **and** $\alpha_i > \frac{1}{2}$. Note that, by Lemma 3.3, this situation arises only when $B_1 > a_{i,i} - a_{1,1}$, and we need only consider $\alpha_i \leq \frac{B_1}{a_{i,i} - a_{1,1} + B_1}$. From (3.8) and the hypothesis of Theorem 3.1, for all sufficiently large $a_{i,i}$, we get:

$$\begin{aligned} \frac{G_j}{n} &\geq \frac{1 - \alpha_i}{k - 1} \left(\frac{1}{2} - \frac{1}{\sqrt{a_{i,i}}} \right) a_{i,i} - \frac{\alpha_i(a_{i,i} - a_{1,1})}{k - 1} \\ &\geq \frac{1 - \alpha_i}{k - 1} \left(\frac{1}{2} - \frac{1}{\sqrt{a_{i,i}}} \right) a_{i,i} - \frac{(1 - \alpha_i)(a_{i,i} - a_{1,1} + B_1)\alpha_i}{k - 1} \\ &= \frac{1 - \alpha_i}{k - 1} \left\{ \frac{a_{i,i}}{2} - \sqrt{a_{i,i}} - B_1 \right\} \\ &\geq \frac{1 - \alpha_i}{k - 1} \left\{ \left(\frac{1}{2} - \epsilon_{i,1} \right) a_{i,i} - B_1 \right\} > \frac{(1 - \alpha_i)\epsilon_{i,1}a_{i,i}}{k - 1}, \end{aligned} \quad (3.31)$$

where the last inequality follows from (3.2).

We again split the analysis into three parts depending on the ranges of values of α_j as given in (3.14), (3.20) and (3.27), with c satisfying the following condition:

$$\log_2 c > \max \left\{ 16k + 1, \frac{2C(k-1)}{\epsilon_{i,1}4^{8k}} \right\}. \quad (3.32)$$

We first consider the case of $\eta_j \neq 0$ and then the case of $\eta_j = 0$ in each of these ranges.

When we are in the regime of (3.14) and $\eta_j \neq 0$, we note that the bounds in (3.16) and (3.17) hold, and therefore the same analysis as before goes through. When $\alpha_j \leq \frac{c}{a_{i,j}}$ and $\eta_j \neq 0$, the bounds in (3.24) and (3.25) hold for all $s = 1, \dots, \lfloor \frac{\alpha_j n}{2} \rfloor$. Now, from (3.31), we have:

$$c \geq \alpha_j a_{i,j} \geq \frac{G_j}{n} \geq \frac{(1 - \alpha_i)\epsilon_{i,1}a_{i,i}}{k - 1} \implies \alpha_i \geq 1 - \frac{c(k-1)}{\epsilon_{i,1}a_{i,i}}. \quad (3.33)$$

Then $\frac{1}{n} \log_2 \mathbf{P}[A_j]$ can be bounded above by

$$-\frac{\log_2 \{ \alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j) \}}{2n} - \frac{\log_2 n}{n} + H(\alpha_i) + H(\alpha_j) + \frac{\alpha_i}{2} \log_2 \gamma$$

of which the first two terms approach 0 as $n \rightarrow \infty$. The remaining terms can be bounded above by

$$2H \left(\frac{c(k-1)}{\epsilon_{i,1}a_{i,i}} \right) + \frac{\log_2 \gamma}{4},$$

of which the first term can be made arbitrarily small by choosing $a_{i,i}$ sufficiently large, and the last term is a strictly negative constant. Hence the above is strictly negative for all $a_{i,i}$ sufficiently large.

Now, we consider $\eta_j = 0$ for the various regimes of α_j . By the same reasoning as Remark 3.7, the only interesting case is where $G_j < \frac{\alpha_j n}{2}$. First, we consider the range of α_j as in (3.14). The bound of (3.18) holds for all $s = 1, \dots, G_j$. By (3.31) and since $\alpha_j \leq 1 - \alpha_i$, we get

$$\frac{G_j}{n} \geq \frac{\alpha_j \epsilon_{i,1} a_{i,i}}{k - 1} > \frac{c \epsilon_{i,1} a_{i,i}}{(k - 1) a_{i,j}} > \frac{2 \cdot 4^{8k} \epsilon_{i,1}}{C(k - 1)} \quad (3.34)$$

by (3.31), (3.32) and (i). Thus an upper bound on $\mathbf{P}[A_j]$ is given by

$$O \left\{ \frac{2^{H(\alpha_i)n + H(\alpha_j)n}}{\sqrt{\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j)} n} \left(\frac{1}{c} \right)^{G_j/2} \right\} \leq O \left\{ \frac{2^{2n}}{\sqrt{\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j)} n} \left(\frac{1}{c} \right)^{\frac{4^{8k} \epsilon_{i,1} n}{C(k-1)}} \right\}$$

which is $o(n^{-1})$ for c as in (3.32).

Next, we consider α_j as in (3.20). Again, the bounds in (3.24) and (3.25) hold for all $s = 1, \dots, G_j$. From (3.31), we get the following upper bound on $\frac{1}{n} \log_2 \mathbf{P}[A_j]$:

$$-\frac{\log_2 \{\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j)\}}{2n} - \frac{\log_2 n}{n} + 2H(\alpha_i) + \frac{(1 - \alpha_i) \epsilon_{i,1} a_{i,i}}{k-1} \log_2 \gamma.$$

The sum of the last two terms can be bounded above by

$$\begin{aligned} -4(1 - \alpha_i) \log_2(1 - \alpha_i) + \frac{(1 - \alpha_i) \epsilon_{i,1} a_{i,i}}{k-1} \log_2 \gamma &< 8(1 - \alpha_i) \log_2 a_{i,j} + \frac{(1 - \alpha_i) \epsilon_{i,1} a_{i,i}}{k-1} \log_2 \gamma \\ &\leq (1 - \alpha_i) \left\{ 8 \log_2 C + 8 \log_2 a_{i,i} + \frac{\epsilon_{i,1} a_{i,i}}{k-1} \log_2 \gamma \right\}, \end{aligned}$$

which is strictly negative for all $a_{i,i}$ sufficiently large.

Finally, we consider α_j in the range given in (3.27). The bound in (3.28) holds for all $s = 1, \dots, G_j$. From (3.31), we have

$$\frac{1}{a_{i,j}} \geq \alpha_j a_{i,j} \geq \frac{G_j}{n} \geq \frac{(1 - \alpha_i) \epsilon_{i,1} a_{i,i}}{k-1} \implies \alpha_i \geq 1 - \frac{k-1}{\epsilon_{i,1} a_{i,i} a_{i,j}}. \quad (3.35)$$

For any fixed positive integer $r > 2$ and all $a_{i,i}$ sufficiently large,

$$a_{i,i} > C^{\frac{1}{r-2}} \left(\frac{k-1}{\epsilon_{i,1}} \right)^{\frac{r-1}{r-2}} 2^{\frac{r}{r-2}},$$

and by the same reasoning as in (3.30), using (3.35) we conclude that

$$\log_2 \{2(1 - \alpha_i) a_{i,j}\} \leq \frac{\log_2(1 - \alpha_i)}{r}.$$

An upper bound on $\frac{1}{n} \log_2 \mathbf{P}[A_j]$ is given by

$$-\frac{\log_2 \{\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j)\}}{2n} - \frac{\log_2 n}{n} + 2H(\alpha_i) + \frac{(1 - \alpha_i) \epsilon_{i,1} a_{i,i}}{k-1} \log_2 \{2(1 - \alpha_i) a_{i,j}\}$$

of which the first two terms approach 0 as $n \rightarrow \infty$, and the sum of the last two terms can be bounded by

$$-4(1 - \alpha_i) \log_2(1 - \alpha_i) + \frac{(1 - \alpha_i) \epsilon_{i,1} a_{i,i}}{k-1} \frac{\log_2(1 - \alpha_i)}{r} = (1 - \alpha_i) \log_2(1 - \alpha_i) \left\{ -4 + \frac{\epsilon_{i,1} a_{i,i}}{r(k-1)} \right\}$$

which is strictly negative for all $a_{i,i}$ sufficiently large. This brings us to the end of §3.2.

3.3. When $a_{i,i} > a_{1,1}$ and $\alpha_i \leq \frac{1}{2}$. By (3.8), we have

$$\begin{aligned} \frac{G_j}{n} &\geq \frac{\alpha_i}{k-1} \left(\frac{1}{2} - \frac{1}{\sqrt{a_{i,i}}} \right) a_{i,i} - \frac{\alpha_i (a_{i,i} - a_{1,1})}{k-1} \\ &\geq \frac{\alpha_i}{k-1} \left\{ a_{1,1} - \frac{a_{i,i}}{2} - \sqrt{a_{i,i}} \right\} \geq \frac{\alpha_i \delta_{1,i} a_{i,i}}{k-1}, \end{aligned} \quad (3.36)$$

by (3.1). Note that, if $\alpha_j \leq \frac{c}{a_{i,j}}$ for any constant $c > 1$, then

$$c \geq \alpha_j a_{i,j} \geq \frac{G_j}{n} \geq \frac{\alpha_i \delta_{1,i} a_{i,i}}{k-1} \implies \alpha_i \leq \frac{c(k-1)}{\delta_{1,i} a_{i,i}},$$

which is strictly less than $\frac{1}{k}$ for all $a_{i,i}$ sufficiently large, contradicting our choice of i . Hence we need only consider the range of (3.14) for values of α_j .

When $\eta_j \neq 0$, the argument is the same as the corresponding case in §3.1. When $\eta_j = 0$, the bound in (3.18) holds, and from (3.36) and $\alpha_i \geq \frac{1}{k}$, we get the following upper bound on $\frac{1}{n} \log_2 \mathbf{P}[A_j]$:

$$-\frac{\log_2 \{\alpha_i \alpha_j (1 - \alpha_i)(1 - \alpha_j)\}}{2n} - \frac{\log_2 n}{n} + H(\alpha_i) + H(\alpha_j) - \frac{\delta_{1,i} a_{i,i}}{2k(k-1)} \log_2 c,$$

and as $H(\alpha_i) + H(\alpha_j) \leq 2$, the above expression is strictly negative for all $a_{i,i}$ sufficiently large.

REFERENCES

- [1] Emmanuel Abbe. Community detection and stochastic block models: recent developments. *The Journal of Machine Learning Research*, 18(1):6446–6531, 2017.
- [2] Emmanuel Abbe and Colin Sandon. Detection in the stochastic block model with multiple clusters: proof of the achievability conjectures, acyclic bp, and the information-computation gap. *arXiv preprint arXiv:1512.09080*, 2015.
- [3] Emmanuel Abbe, Afonso S Bandeira, and Georgina Hall. Exact recovery in the stochastic block model. *IEEE Transactions on Information Theory*, 62(1):471–487, 2015.
- [4] Paolo Barucca. Spectral partitioning in equitable graphs. *Physical Review E*, 95(6):062310, 2017.
- [5] Edward A Bender. The asymptotic number of non-negative integer matrices with given row and column sums. *Discrete Mathematics*, 10(2):217–223, 1974.
- [6] GUILLAUMEJL BLONDELVD et al. Fast unfolding of community hierarchies in large networks, 2008.
- [7] Béla Bollobás and Bollobás Béla. *Random graphs*. Number 73. Cambridge university press, 2001.
- [8] Charles Bordenave, Marc Lelarge, and Laurent Massoulié. Non-backtracking spectrum of random graphs: community detection and non-regular ramanujan graphs. In *2015 IEEE 56th Annual Symposium on Foundations of Computer Science*, pages 1347–1357. IEEE, 2015.
- [9] Gerandy Brito, Ioana Dumitriu, Shirshendu Ganguly, Christopher Hoffman, and Linh V Tran. Recovery and rigidity in a regular stochastic block model. In *Proceedings of the twenty-seventh annual ACM-SIAM symposium on Discrete algorithms*, pages 1589–1601. Society for Industrial and Applied Mathematics, 2016.
- [10] Gerandy Brito, Ioana Dumitriu, and Kameron Decker Harris. Spectral gap in random bipartite biregular graphs and applications. *arXiv preprint arXiv:1804.07808*, 2018.
- [11] Deepjyoti Choudhury and Arnab Paul. Community detection in social networks: an overview. *International Journal of Research in Engineering and Technology*, 2(2):6–13, 2013.
- [12] Amin Coja-Oghlan. Graph partitioning via adaptive spectral techniques. *Combinatorics, Probability and Computing*, 19(2):227–284, 2010.
- [13] Anne Condon and Richard M Karp. Algorithms for graph partitioning on the planted partition model. *Random Structures & Algorithms*, 18(2):116–140, 2001.
- [14] Aurelien Decelle, Florent Krzakala, Cristopher Moore, and Lenka Zdeborová. Asymptotic analysis of the stochastic block model for modular networks and its algorithmic applications. *Physical Review E*, 84(6):066106, 2011.
- [15] Aurelien Decelle, Florent Krzakala, Cristopher Moore, and Lenka Zdeborová. Inference and phase transitions in the detection of modules in sparse networks. *Physical Review Letters*, 107(6):065701, 2011.
- [16] Martin E. Dyer and Alan M. Frieze. The solution of some random np-hard problems in polynomial expected time. *Journal of Algorithms*, 10(4):451–489, 1989.

- [17] Hossein Fani and Ebrahim Bagheri. Community detection in social networks. *Encyclopedia with Semantic Computing and Robotic Intelligence*, 1(01):1630001, 2017.
- [18] Joel Friedman. *A proof of Alon's second eigenvalue conjecture and related problems*. American Mathematical Soc., 2008.
- [19] Javier O Garcia, Arian Ashourvan, Sarah Muldoon, Jean M Vettel, and Danielle S Bassett. Applications of community detection techniques to brain graphs: Algorithmic considerations and implications for neural function. *Proceedings of the IEEE*, 106(5):846–867, 2018.
- [20] Michelle Girvan and Mark EJ Newman. Community structure in social and biological networks. *Proceedings of the national academy of sciences*, 99(12):7821–7826, 2002.
- [21] Simon Heimlicher, Marc Lelarge, and Laurent Massoulié. Community detection in the labelled stochastic block model. *arXiv preprint arXiv:1209.2910*, 2012.
- [22] Paul W Holland, Kathryn Blackmond Laskey, and Samuel Leinhardt. Stochastic blockmodels: First steps. *Social networks*, 5(2):109–137, 1983.
- [23] Mark Jerrum and Alistair Sinclair. Approximating the permanent. *SIAM journal on computing*, 18(6):1149–1178, 1989.
- [24] Mark Jerrum and Gregory B Sorkin. The metropolis algorithm for graph bisection. *Discrete Applied Mathematics*, 82(1-3):155–175, 1998.
- [25] Brian Karrer and Mark EJ Newman. Stochastic blockmodels and community structure in networks. *Physical review E*, 83(1):016107, 2011.
- [26] Gregory F Lawler and Alan D Sokal. Bounds on the l^2 spectrum for markov chains and markov processes: a generalization of cheeger's inequality. *Transactions of the American mathematical society*, 309(2):557–580, 1988.
- [27] Marc Lelarge, Laurent Massoulié, and Jiaming Xu. Reconstruction in the labelled stochastic block model. *IEEE Transactions on Network Science and Engineering*, 2(4):152–163, 2015.
- [28] Jure Leskovec, Kevin J Lang, Anirban Dasgupta, and Michael W Mahoney. Statistical properties of community structure in large social and information networks. In *Proceedings of the 17th international conference on World Wide Web*, pages 695–704, 2008.
- [29] David A. Levin, Yuval Peres, and Elizabeth L. Wilmer. *Markov chains and mixing times*, volume 107. American Mathematical Soc., 2017.
- [30] Laurent Massoulié. Community detection thresholds and the weak ramanujan property. In *Proceedings of the forty-sixth annual ACM symposium on Theory of computing*, pages 694–703. ACM, 2014.
- [31] Marwa Ben M'barek, Amel Borgi, Walid Bedhiafi, and Sana Ben Hmida. Genetic algorithm for community detection in biological networks. *Procedia Computer Science*, 126:195–204, 2018.
- [32] Brendan D McKay and Xiaoji Wang. Asymptotic enumeration of 0–1 matrices with equal row sums and equal column sums. *Linear algebra and its applications*, 373:273–287, 2003.
- [33] Brendan D McKay and Nicholas C Wormald. Asymptotic enumeration by degree sequence of graphs with degrees $(n^{1/2})$. *Combinatorica*, 11(4):369–382, 1991.
- [34] Frank McSherry. Spectral partitioning of random graphs. In *Proceedings 42nd IEEE Symposium on Foundations of Computer Science*, pages 529–537. IEEE, 2001.
- [35] Seyed Ahmad Moosavi, Mehrdad Jalali, Negin Misaghian, Shahaboddin Shamshirband, and Mohammad Hossein Anisi. Community detection in social networks using user frequent pattern mining. *Knowledge and Information Systems*, 51(1):159–186, 2017.
- [36] Elchanan Mossel, Joe Neeman, and Allan Sly. Stochastic block models and reconstruction. *arXiv preprint arXiv:1202.1499*, 2012.
- [37] Elchanan Mossel, Joe Neeman, and Allan Sly. Belief propagation, robust reconstruction and optimal recovery of block models. In *Conference on Learning Theory*, pages 356–370, 2014.

- [38] Elchanan Mossel, Joe Neeman, and Allan Sly. Consistency thresholds for the planted bisection model. In *Proceedings of the forty-seventh annual ACM symposium on Theory of computing*, pages 69–75, 2015.
- [39] Elchanan Mossel, Joe Neeman, and Allan Sly. A proof of the block model threshold conjecture. *Combinatorica*, 38(3):665–708, 2018.
- [40] Raj Rao Nadakuditi and Mark EJ Newman. Graph spectra and the detectability of community structure in networks. *Physical review letters*, 108(18):188701, 2012.
- [41] Mark EJ Newman. Detecting community structure in networks. *The European Physical Journal B*, 38(2):321–330, 2004.
- [42] Mark EJ Newman. Fast algorithm for detecting community structure in networks. *Physical review E*, 69(6):066133, 2004.
- [43] Mark EJ Newman. Finding community structure in networks using the eigenvectors of matrices. *Physical review E*, 74(3):036104, 2006.
- [44] MEJ Newman and Travis Martin. Equitable random graphs. *Physical Review E*, 90(5):052824, 2014.
- [45] Soumik Pal and Yizhe Zhu. Community detection in the sparse hypergraph stochastic block model. *arXiv preprint arXiv:1904.05981*, 2019.
- [46] Tiago P Peixoto. Bayesian stochastic blockmodeling. *Advances in network clustering and blockmodeling*, pages 289–332, 2019.
- [47] Sara Rahiminejad, Mano R Maurya, and Shankar Subramaniam. Topological and functional comparison of community detection algorithms in biological networks. *BMC bioinformatics*, 20(1):212, 2019.
- [48] Xiaoran Yan, Cosma Shalizi, Jacob E Jensen, Florent Krzakala, Cristopher Moore, Lenka Zdeborová, Pan Zhang, and Yaojia Zhu. Model selection for degree-corrected block models. *Journal of Statistical Mechanics: Theory and Experiment*, 2014(5):P05007, 2014.