Guangying Lv^a, Jinlong Wei^b, Guang-an Zou^c

^a College of Mathematics and Statistics, Nanjing University of Information Science and Technology, Nanjing 210044, China

gylvmaths@126.com

^bSchool of Statistics and Mathematics, Zhongnan University of Economics and Law, Wuhan 430073, China

weijinlong.hust@gmail.com

^cInstitute of Applied Mathematics, Henan University Kaifeng 475001, China zouguangan@henu.edu.cn

February 18, 2020

Abstract

We study the stability of reaction-diffusion equations in presence of noise. The relationship of stability of solutions between the stochastic ordinary different equations and the corresponding stochastic reaction-diffusion equation is firstly established. Then, by using the Lyapunov method, sufficient conditions for mean square and stochastic stability are given. The results show that the multiplicative noise can make the solution stable, but the additive noise will be not.

Keywords: Stochastic stability; Mean square stability; Noise; Lyapunov method. AMS subject classifications (2010): 35B35, 60H15.

1 Introduction

The stability of solutions is an important issue in the theory of PDEs (partial differential equations), which has been studied by many authors [24]. There are a lot of sufficient conditions to assure that the solutions are stable or unstable. We note that noise always exists in the real world. The reasons may be that the parameter is obtained by different measurement, and we can not get the real value, so we consider the ordinary (partial) differential equations with noise perturbation is available. In other words, in microscopic world, to describe the particle moving law must be stochastic ordinary (partial) differential equations. In macroscopic world, we often consider the case that the coefficient in equations is random or stochastic. Usually, if we consider the role of noise, we have two cases. First case: the noise is regarded as a small perturbation. In this case, the structure of solutions will not be changed and the biggest possible change is long-time behavior of the solutions, that is to say, the conditions of stability or un-stability may be different from the deterministic case. Second case: the noise is strong, such as $u^{\gamma}dW_t$, where $\gamma > 1$, u is the unknown function and W_t is the noise. In this case, the structure of solutions will be changed. More precisely, the noise can induce the solutions blow up in finite time. When the noise appears in a deterministic PDE, the impact of noise on solutions will be the first thing to be considered. In the present paper, we aim to study impact of noise on stability of solutions.

Initially, we recall some known results about the impact of noise. Flandoli et al. [5, 6] proved that the noise can make the transport equations well-posedness and the noise can prevent the singularities in linear transport equations. Chow [2] obtained that the noise can induce singularities (finite time blow up of solutions), also see [14, 17]. There are a lot of work about the impact of noise on different PDEs, for example, Hamilton-Jacobi equations [7], conservation law [8, 10], porous media equations [4], quasilinear degenerate parabolic-hyperbolic equations [9], and so on. Besides, the impact of noise on regularity of solutions of parabolic equations has been studied by Lv et al. [16].

Forty years ago, there are a lot of works about the impact of noise. The main issue is that the noise can stabilize the solution of ordinary differential equations, see the book [12, 19, 20]. Meanwhile, the stochastic stability of functional differential equations is also considered by Mackey-Nechaeva [18]. In the book [3], the long time behavior of solutions was considered in Chapter 11 and sufficient condition of mean square stable is given, see Theorem 11.14. More precisely, Da Prato- Zabczyk studied the following equation

$$\begin{cases} dX = AXdt + B(X)dW_t, \\ X(0) = x \in H, \end{cases}$$
(1.1)

where H is a Hilbert space, A generates a C_0 semigroup and $B \in L(H; L_2^0)$ (see p309 of [3] for more details). They proved that the following statements are equivalent

(i) There exists $M > 0, \gamma > 0$ such that

$$\mathbb{E}|X(t,x)|^2 \le Me^{-\gamma t}|x|^2, \quad t \ge 0.$$

(ii) For any $x \in H$ we have

$$\mathbb{E}\int_0^\infty |X(t,x)|^2 dt < \infty.$$

It is remarked that the nonlinear term is not considered in the book [3]. Moreover, the example in [3] is the stochastic reaction-diffusion equation on the bounded domain. Liu-Mao [13] also considered the stability of trivial solution 0 on the bounded domain. Wang-Li [23] considered the stability and moment boundedness of the stochastic linear age-structured model. Although in the book [1], Chow gave a abstract result to study the stability of null solution (see page 233), the concrete form was not given. In the present paper, we consider the concrete model and generalized the classical results. What's more, we find some difference between stochastic partial differential equations (SPDEs) and stochastic differential equations (SDEs).

We discuss the impact of different kinds of noise on stability. Some interesting results are obtained: the additive noise will have a "bad" effect and some multiplicative noise has "good" effect. In other words, the multiplicative noise can make the solution stable, but the additive noise will not, which is new for SPDEs. This is different from SDEs, see Remark 3.2. What's more, we obtain a new result for stability theory of SDEs, see Theorem 3.6.

This paper is arranged as follows. In Sections 2, we will present some preliminaries. Sections 3 and 4 are concerned with the bounded domain and the whole space, respectively.

2 Preliminaries

In the present paper, we always assume W_t is a one-dimensional standard Wiener process and $W_t(x)$ is a Wiener random field which are defined on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Firstly, we recall the definitions of stochastic stability. We only consider the stability of constant equilibrium of stochastic reaction-diffusion equations. Consider the following stochastic reaction-diffusion equations

$$\begin{cases} du = (\Delta u + f(u))dt + \sigma(u)dW, & t > 0, \quad x \in \mathbb{R}^d, \\ u(x,0) = u_0(x), & x \in \mathbb{R}^d, \end{cases}$$
(2.1)

where $W = W_t$ or $W_t(x)$. The Wiener random field can be chosen to have the following properties: $\mathbb{E}W_t(x) = 0$ and its covariance function q(x, y) is given by

$$\mathbb{E}W_t(x)W_s(y) = (t \wedge s)q(x,y), \quad x, y \in \mathbb{R}^d,$$

where $t \wedge s = \min\{t, s\}$ for $0 \leq t, s \leq T$, or can be chosen as one-dimension Brownian motion. Without loss of generality, we suppose that $f(0) = \sigma(0) = 0$, that is to say, 0 is a trivial solution to the first equation of problem (2.1). Let $\|\cdot\|$ denote as some norm with respect to the spatial variable.

Definition 2.1 The trivial solution 0 is called mean square stable if for any $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that for any initial data u_0 ,

$$||u_0|| \le \delta$$
 implies $\mathbb{E}||u(\cdot, t)||^2 < \varepsilon, \ \forall \ t \ge 0,$

and exponentially mean square stable, if there exists two positive constants c_1 and c_2 such that

$$\mathbb{E} \|u(\cdot,t)\|^2 < c_1 e^{-c_2 t} \|u_0\|^2, \ \forall \ t \ge 0.$$

Definition 2.2 The trivial solution 0 is called stochastically stable if for any $\varepsilon_1 > 0$ and $\varepsilon_2 > 0$, there exists $\delta = \delta(\varepsilon_1, \varepsilon_2) > 0$ such that for t > 0 the solution u satisfies

$$\mathbb{P}\left\{\sup_{t>0} \|u(\cdot,t)\| \le \varepsilon_1\right\} \ge 1 - \varepsilon_2 \quad \text{for} \quad \|u_0\| \le \delta.$$

Before ending this section, we consider the relationship of the stability of solutions between stochastic differential equations and stochastic reaction-diffusion equations.

Assume that u = 0 is a trivial solution to the first equation of problem (2.1), then u = 0 will be a trivial solution to the following equation

$$du = f(u)dt + \sigma(u)dW_t. \tag{2.2}$$

Problem (2.2) means equation (2.2) with the initial data u_0 (independent of x). Initially, we introduce a definition for (2.2).

Definition 2.3 The trivial solution 0 is called mean square stable for problem (2.2) if for any $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that for any initial data u_0 ,

$$|u_0| \le \delta$$
 implies $\mathbb{E}|u(\cdot, t)|^2 < \varepsilon, \ \forall \ t \ge 0,$

and exponentially mean square stable, if there exists two positive constants c_1 and c_2 such that

$$\mathbb{E}|u(\cdot,t)|^2 < c_1 e^{-c_2 t} |u_0|^2, \ \forall \ t \ge 0,$$

4

and stochastically stable if for any $\varepsilon_1 > 0$ and $\varepsilon_2 > 0$, there exists $\delta = \delta(\varepsilon_1, \varepsilon_2) > 0$ such that for t > 0 the solution u satisfies

$$\mathbb{P}\left\{\sup_{t>0}|u(\cdot,t)|\leq\varepsilon_1\right\}\geq 1-\varepsilon_2\quad\text{for}\quad|u_0|\leq\delta.$$

In order to establish the relationship between problems (2.1) and (2.2), we need the following lemma. Let $\eta(r) = r^-$ denote the negative part of r for $r \in \mathbb{R}$. Set

$$k(r) = \eta^2(r),$$

so that k(r) = 0 for $r \ge 0$ and $k(r) = r^2$ for r < 0. For $\epsilon > 0$, let $k_{\epsilon}(r)$ be a C^2 -regularization of k(r) defined by

$$k_{\epsilon}(r) = \begin{cases} r^2 - \frac{\epsilon^2}{6}, & r < -\epsilon, \\ -\frac{r^3}{\epsilon} \left(\frac{r}{2\epsilon} + \frac{4}{3}\right), & -\epsilon \le r < 0, \\ 0, & r \ge 0. \end{cases}$$

Then one can check that $k_{\epsilon}(r)$ has the following properties.

Lemma 2.1 [14, Lemma 3.1] The first two derivatives k'_{ϵ} , k''_{ϵ} of k_{ϵ} are continuous and satisfy the conditions: $k'_{\epsilon}(r) = 0$ for $r \ge 0$; $k'_{\epsilon} \le 0$ and $k''_{\epsilon} \ge 0$ for any $r \in \mathbb{R}$. Moreover, as $\epsilon \to 0$, we have

$$k_{\epsilon}(r) \to k(r)$$
 and $k'_{\epsilon}(r) \to -2\eta(r)$,

and the convergence is uniform for $r \in \mathbb{R}$.

Under the condition that both problems (2.1) and (2.2) have a unique strong solution, we have the following result. Here we focus on the stability of solutions and we do not talk about the existence of solutions.

Theorem 2.1 Assume that W_t is a one-dimension Wiener process, f and σ satisfy the global Lipschitz condition, and $f(0) = \sigma(0) = 0$. Then 0 is a stable (exponentially mean square stable or stochastically stable) trivial solution of (2.1) (with the L^{∞} norm in spatial variable) if and only if 0 is a stable (exponentially mean square sable or stochastically stable) trivial solution of (2.2).

Proof. Clearly, if 0 is a stable (exponentially mean square stable or stochastically stable) trivial solution of (2.1), then 0 is also a stable trivial solution of (2.2) since we can choose the initial data u_0 which is spatial variable independent.

Conversely, we will check if if 0 is a stable (exponentially mean square stable or stochastically stable) trivial solution of (2.2), then 0 is a stable trivial solution of (2.1) as well.

Assume 0 is a stable (exponentially mean square stable or stochastically stable) trivial solution of (2.2), then for any $\varepsilon > 0$, there exists constant $\delta = \delta(\varepsilon) > 0$ such that

$$\mathbb{E}|u_{\pm}(t)|^2 < \varepsilon_1$$

where $u_{\pm}(t)$ is the solution of (2.2) with initial data $\pm 2\delta$.

Let u(x,t) be the unique solution of (2.1) with initial data $u_0(x)$ which satisfies $|u_0(x)| \leq \delta$. For any fixed T > 0, we will prove that

$$u_{-}(t) \le u(x,t) \le u_{+}(t), \ \forall \ (x,t) \in \mathbb{R}^{d} \times [0,T], \ a.s..$$
 (2.3)

If the above inequality holds, then we get the desired result. In order to the inequality (2.3), we let $v = u(x,t) - u_{-}(t)$, then v satisfies

$$\begin{cases} dv = (\Delta v + f(u) - f(u_{-}))dt + (\sigma(u) - \sigma(u_{-}))dW_t, & t > 0, \quad x \in \mathbb{R}^d, \\ v(x,0) = u_0(x) + \delta \ge 0, & x \in \mathbb{R}^d. \end{cases}$$
(2.4)

Note that ∇u makes sense in $L^{\infty}(\mathbb{R}^d)$, thus we can not take $\Phi_{\varepsilon}(v(t)) = (1, k_{\varepsilon}(v(t)))$, which is different from those in [2, 17]. We need introduce a new test function.

For any R > 0, $B_R(0)$ denotes a ball centered in 0 with radius R. Let ϕ_1 be the eigenvalue function of Laplacian operator on $B_R(0)$ with respect to the first eigenvalue λ_1 , i.e.,

$$\begin{cases} -\Delta \phi_1 = \lambda_1 \phi_1, & \text{in } B_R(0), \\ \phi_1 = 0, & \text{on } \partial B_R(0). \end{cases}$$

Denote $\psi \in C^2(\mathbb{R}^d)$ satisfying

$$\psi(x) = \begin{cases} \phi_1(x), & x \in \bar{B}_R(0), \\ 0, & x \in \mathbb{R}^d \setminus B_R(0). \end{cases}$$

Define

$$\Phi_{\varepsilon}(v(t)) = (\psi, k_{\epsilon}(v(t))) = \int_{\mathbb{R}^d} \psi(x) k_{\epsilon}(v(x, t)) dx.$$

By Itô's formula, we have

$$\begin{split} \Phi_{\varepsilon}(v(t)) &= \Phi_{\varepsilon}(v_0) + \int_0^t \int_{\mathbb{R}^d} \psi(x) k'_{\epsilon}(v(x,s)) \Delta v(x,s) dx ds \\ &+ \int_0^t \int_{\mathbb{R}^d} \psi(x) k'_{\epsilon}(v(x,s)) (f(u(x,s)) - f(u_-(s))) dx ds \\ &+ \int_0^t \int_{\mathbb{R}^d} \psi(x) k'_{\epsilon}(v(x,s)) (\sigma(u(x,s)) - \sigma(u_-(s))) dx dW_s \\ &+ \frac{1}{2} \int_0^t \int_{\mathbb{R}^d} \psi(x) k''_{\epsilon}(v(x,s)) (\sigma(u(x,s)) - \sigma(u_-(s)))^2 dx ds \end{split}$$

By using the facts $k_{\epsilon}'' \ge 0, \, \psi(x) \ge 0$, we have

$$\begin{split} & \int_{\mathbb{R}^d} \psi(x) k'_{\epsilon}(v(x,s)) \Delta v(x,s) dx \\ &= -\int_{\mathbb{R}^d} \psi(x) k''_{\epsilon}(v(x,s)) |\nabla v(x,s)|^2 dx - \int_{\mathbb{R}^d} k'_{\epsilon}(v(x,s)) \nabla \psi(x) \cdot \nabla v(x,s) dx \\ &\leq -\int_{\mathbb{R}^d} \nabla k_{\epsilon}(v(x,s)) \cdot \nabla \psi(x) dx \\ &= \int_{\mathbb{R}^d} k_{\epsilon}(v(x,s)) \Delta \psi(x) dx \\ &= -\lambda_1 \int_{\mathbb{R}^d} k_{\epsilon}(v(x,s)) \psi(x) dx \leq 0. \end{split}$$

Consequently,

$$\begin{split} \Phi_{\varepsilon}(v(t)) &\leq \Phi_{\varepsilon}(v_0) + \int_0^t \int_{\mathbb{R}^d} \psi(x) k_{\epsilon}''(v(x,s)) \left(\frac{1}{2} (\sigma(u(x,s)) - \sigma(u_-(s)))^2 \right) dx ds \\ &+ \int_0^t \int_{\mathbb{R}^d} \psi(x) k_{\epsilon}'(v(x,s)) (f(u(x,s)) - f(u_-(s))) dx ds \\ &+ \int_0^t \int_{\mathbb{R}^d} \psi(x) k_{\epsilon}'(v(x,s)) (\sigma(u(x,s)) - \sigma(u_-(s))) dx dW_s. \end{split}$$

Taking expectation over the above equality and using Lemma 2.1, we get

$$\begin{split} \mathbb{E}\Phi_{\varepsilon}(v(t)) &= \mathbb{E}\Phi_{\varepsilon}(v_{0}) + \mathbb{E}\int_{0}^{t}\int_{\mathbb{R}^{d}}\psi(x)k_{\epsilon}''(v(x,s)) \\ &\times \left(\frac{1}{2}(\sigma(u(x,s)) - \sigma(u_{-}(s)))^{2}\right)dxds \\ &+ \mathbb{E}\int_{0}^{t}\int_{\mathbb{R}^{d}}\psi(x)k_{\epsilon}'(v(x,s))(f(u(x,s)) - f(u_{-}(s)))dxds \\ &\leq \mathbb{E}\Phi_{\varepsilon}(v_{0}) + \frac{L_{\sigma}}{2}\mathbb{E}\int_{0}^{t}\int_{\mathbb{R}^{d}}\psi(x)k_{\epsilon}''(v(x,s))v(x,s)^{2}dxds \\ &+ L_{f}\mathbb{E}\int_{0}^{t}\int_{\mathbb{R}^{d}}\psi(x)|k_{\epsilon}'(v(x,s))|v(x,s)|dxds. \end{split}$$

Note that $\lim_{\varepsilon \to 0} \mathbb{E}\Phi_{\varepsilon}(v(t)) = \mathbb{E}(\eta(v(t))^2, \psi)$, by taking the limits termwise as $\varepsilon \to 0$ and using Lemma 2.1, we have

$$\mathbb{E}(\eta(v(t))^2,\psi) \le (L_{\sigma}^2 + 2L_f) \int_0^t \mathbb{E}(\eta(v(s))^2,\psi) ds,$$

which, by means of Gronwall's inequality, implies that

$$\mathbb{E}(\eta(v(t))^2,\psi) = 0, \quad \forall \ t \in [0,T]$$

Note that for any R > 0, the above inequality always holds. It follows from $\psi \ge 0$ that $\eta(v(t)) = v^-(x,t) = 0$ a.s. for a.e. $x \in \mathbb{R}^d$, which implies that $v^- = 0$ a.s. for a.e. $x \in \mathbb{R}^d$, and for any $t \in [0,T]$.

Similarly, if we let $v = u_+ - u(x, t)$, then we can prove that $v^- = 0$ a.s. for a.e. $x \in \mathbb{R}^d$, and for any $t \in [0, T]$. That is to say, (2.3) holds. \Box

Remark 2.1 It is remarked that in Theorem 2.1 the norm of solution of problem (2.1) is maximum in \mathbb{R}^d . The advantage of this norm is that the estimates we obtained hold point-wise. Under this norm, Theorem 2.1 also holds if the equation (2.1) is replaced by the equation (2.5).

Now, we consider a special case on a bounded domain $D \subset \mathbb{R}^d$ $(d \ge 1)$

$$\begin{cases} du = (\Delta u + f(u))dt + udW_t, & t > 0, \quad x \in D, \\ u(x,0) = u_0(x), & x \in D, \\ u(x,t) = 0, & t > 0, \quad x \in \partial D. \end{cases}$$
(2.5)

We will establish the relationship between (2.5) and the following SDE

$$\begin{cases} dX_t = f(X_t)dt + X_t dW_t, & t > 0, \\ X(0) = u_0. \end{cases}$$
(2.6)

In order to do that, we consider the eigenvalue problem for the elliptic equation

$$\begin{cases} -\Delta \phi = \lambda \phi, & \text{in } D, \\ \phi = 0, & \text{on } \partial D. \end{cases}$$

Then, all the eigenvalues are strictly positive, increasing and the eigenfunction ϕ_1 corresponding to the smallest eigenvalue λ_1 does not change sign in domain D, as shown in [11]. Therefore, we normalize it in such a way that

$$\phi_1(x) \ge 0, \quad \int_D \phi_1(x) dx = 1.$$

Theorem 2.2 Assume that

$$(f(u),\phi_1) \le f((u,\phi_1)).$$

If 0 is a stable (exponentially mean square stable or stochastically stable) trivial solution of (2.6), then 0 is also a stable (exponentially mean square sable or stochastically stable) trivial solution of (2.5), where we take the spatial norm as $||u||_{\phi_1} = (u, \phi_1)$.

Proof. It follows from [2, Theorem 2.1] and [14] that the solutions of (2.5) keep non-negative almost surely, i.e., $u(x,t) \ge 0$, a.s. for almost every $x \in D$ and for all $t \in [0,T]$. Moreover, the solutions exist globally. Let

$$v(t) = (u, \phi_1) = \int_D u(x, t)\phi_1(x)dx$$

Then v satisfies

$$\begin{cases} dv(t) \leq [-\lambda_1 v(t) + f(v(t))]dt + v(t)dW_t, & t > 0, \quad x \in D, \\ v(0) = v_0 = (u_0, \phi_1). \end{cases}$$
(2.7)

It is easy to prove that v(t) is a sub-solution of the following problem

$$\begin{cases} dY_t = [-\lambda_1 Y_t + f(Y_t)]dt + Y_t dW_t, & t > 0, \quad x \in D, \\ Y(0) = v_0. \end{cases}$$
(2.8)

Since the solutions of (2.8) keep non-negative, we obtain that v(t) is also a sub-solution of (2.7). Set $Z_t = Y_t - v(t)$, similar to the proof of Theorem 2.1, we can prove $Z_t \ge 0$ almost surely. Indeed, one can first prove that $Y_t \ge 0$ almost surely by using the same method to the proof of Theorem 2.1. Then it follows from the definition of v that $v(t) = (u, \phi_1) > 0$ almost surely. Lastly, noting that

$$-(Y_t^r - v(t)^r) = -r\xi^{r-1}Z_t \ge 0, \text{ when } Z_t \le 0,$$

one can use the same method to the proof of Theorem 2.1 to get $Z_t \ge 0$ almost surely. Similarly, one can prove $Y_t \le X_t$ almost surely. Therefore, if 0 is a stable (exponentially mean square stable or stochastically stable) trivial solution of (2.6), then 0 is also a stable trivial solution of (2.5). The proof is complete. \Box

Example Let $f(u) = au - ku^r$, where $a \in \mathbb{R}, k \ge 0, r \ge 1$. $-u^r \ge 0$ for $u \le 0$. It follows from [2, Theorem 2.1] and [14] that the solutions of (2.5) with $f(u) = au - ku^r$ keep non-negative almost surely. By using the Hölder inequality, we have

$$(u,\phi_1)^r = \left(\int_D u(x,t)\phi_1(x)dx\right)^r \le \int_D u^r(x,t)\phi_1(x)dx = (u^r,\phi_1).$$

Therefore, all the assumptions of Theorem 2.2 are satisfied.

Remark 2.2 In particular, if f(u) = u, then the stability of the trivial solution 0 for (2.5) and the following problem

$$\begin{cases} dY_t = (1 - \lambda_1) Y_t dt + Y_t dW_t, & t > 0, \ x \in D, \\ Y(0) = (u_0, \phi_1), \end{cases}$$

are equivalent. The above problem is different from (2.6) because of the Laplacian operator.

3 Bounded domain

In this section, we consider the stability results on a bounded domain $D \subset \mathbb{R}^d$. We focus on the conditions which induce the solution stable. Meanwhile, we are interested in the difference between the stochastic partial differential equations and partial differential equations. We will consider the impact of different noise. Throughout this section, $\|\cdot\|$ means the norm of $L^2(D)$.

We first consider the following initial boundary problem

$$\begin{cases} du = \mu \Delta u dt + \sigma dW_t, & t > 0, \quad x \in D, \\ u(x,0) = u_0(x), & x \in D, \\ u(x,t) = 0, & t > 0, \quad x \in \partial D, \end{cases}$$
(3.1)

where μ and σ are positive constants. Then we obtain

Theorem 3.1 If $\sigma^2 |D| < 2\mu\lambda_1 \mathbb{E} ||u_0||^2$, then the trivial solution 0 is mean square stable.

Proof. We take the Lyapunov function as $||u||^2$. By using Itô's formula, we have

$$\begin{aligned} \frac{d}{dt} \mathbb{E} \|u(\cdot,t)\|^2 &= 2\mu \mathbb{E} \int_D u\Delta u(x,t) dx + \sigma^2 |D| \\ &= -2\mu \mathbb{E} \int_D |\nabla u(x,t)|^2 dx + \sigma^2 |D| \\ &\leq -2\mu \lambda_1 \mathbb{E} \|u(\cdot,t)\|^2 + \sigma^2 |D|, \end{aligned}$$

where we used the Poincare inequality. Solving the above inequality, we have

$$\mathbb{E}\|u(\cdot,t)\|^2 < \left[\mathbb{E}\|u_0\|^2 - \frac{\sigma^2|D|}{2\lambda_1\mu}\right]e^{-2\lambda_1\mu t} + \frac{\sigma^2|D|}{2\lambda_1\mu}.$$

Note that

$$\left[\mathbb{E}\|u_0\|^2 - \frac{\sigma^2|D|}{2\lambda_1\mu}\right]e^{-2\lambda_1\mu t} + \frac{\sigma^2|D|}{2\lambda_1\mu} \le \mathbb{E}\|u_0\|^2$$

is equivalent to

$$\left[\mathbb{E}\|u_0\|^2 - \frac{\sigma^2|D|}{2\lambda_1\mu}\right] \left[e^{-2\lambda_1\mu t} - 1\right] \le 0.$$

Therefore, if $\mathbb{E} \|u_0\|^2 - \frac{\sigma^2 |D|}{2\lambda_1 \mu} > 0$, that is, $\sigma^2 |D| < 2\mu \lambda_1 \mathbb{E} \|u_0\|^2$, we have

$$\mathbb{E}\|u(\cdot,t)\|^2 \le \mathbb{E}\|u_0\|^2.$$

The proof is complete. \Box

Remark 3.1 It is easy to see that if $\sigma = 0$, then the trivial solution 0 is stable; and if $\mu = 0$, then the trivial solution 0 will be unstable; and when $\sigma^2 > 0$, 0 will be a stable trivial solution under some more assumptions. Possibly one can say that the additive noise will have a "bad" effect on the stability of trivial solution.

By using the Chebyshev inequality, one can easily prove the trivia solution 0 is stochastic stable without any more assumption, see the next theorem for the proof.

We study the impact of additive noise. Consider the following equation

$$\begin{cases} du = (\Delta_p u + f(u))dt + \sigma dW_t, & t > 0, \quad x \in D, \\ u(x,0) = u_0(x), & x \in D, \\ u(x,t) = 0, & t > 0, \quad x \in \partial D, \end{cases}$$
(3.2)

where $\Delta_p u = \nabla \cdot (|\nabla u|^{p-2} \nabla u)$. Because we only consider the stability of solutions, throughout this paper we will assume the problem we consider admits a unique global solution. Let C_{∞} be the Sobolev embedding constant satisfying

$$\|u\|_{L^{\infty}(D)} \le C_{\infty} \|u\|_{W^{1,p}(D)}, \quad p > d.$$
(3.3)

It is noted that the solution u of (3.2) satisfies that $||u||_{W^{1,p}(D)} = ||\nabla u||_{L^p(D)}$.

Theorem 3.2 Assume the nonlinear term satisfies

$$uf(u) \le au^2 + bu^{2m}, \quad m \ge 1,$$
(3.4)

where $a, b \in \mathbb{R}$. Assume further that $p > \max\{2m, d\}$.

If

$$a + \frac{2-\gamma}{2} \left(\frac{\gamma |D| |b| C_{\infty}^p}{2}\right)^{\frac{\gamma}{2-\gamma}} + \frac{\sigma^2 |D|}{2\mathbb{E} \|u_0\|^2} < 0, \tag{3.5}$$

where $\gamma \in (0,2)$ satisfies

$$(2m-2+\gamma)\cdot\frac{2}{\gamma} = p.$$

Then the trivial solution 0 of (3.2) is mean square stable.

Proof. We remark that when m = 1, Theorem 3.2 will become easier, see [13] for similar results. We pick a Lyapunov function $V(u) = ||u||^2$. By Itô's formula, taking expectation and integrating with respect to t, we have

$$\frac{d}{dt}\mathbb{E}\|u(\cdot,t)\|^2 = 2\mathbb{E}\int_D u(\Delta_p u(x,t) + f(u))dx + \sigma^2|D|$$

$$\leq -2\mathbb{E}\int_D |\nabla u(x,t)|^p dx + 2\mathbb{E}\int_D (au^2 + bu^{2m})dx + \sigma^2|D|.$$
(3.6)

By the Sobolev embedding inequality (3.3), we have

$$\begin{aligned} \|u\|_{L^{2m}}^{2m} &= \int_{D} |u|^{2m}(x,t) dx &\leq \|u\|_{L^{\infty}}^{2m-2} \int_{D} u^{2}(x,t) dx \\ &= \|u\|_{L^{\infty}}^{2m-2} \|u\|_{L^{2}}^{2} \\ &\leq |D|^{\frac{\gamma}{2}} \|u\|_{L^{\infty}}^{2m-2+\gamma} \|u\|_{L^{2}}^{2-\gamma} \\ &\leq C_{\infty}^{2m-2+\gamma} |D|^{\frac{\gamma}{2}} \|u\|_{W^{1,p}}^{2m-2+\gamma} \|u\|_{L^{2}}^{2-\gamma} \\ &\leq \frac{2-\gamma}{2} (C_{\infty})^{\frac{2(2m-2+\gamma)}{2-\gamma}} \left(\frac{\gamma |b| |D|}{2}\right)^{\frac{\gamma}{2-\gamma}} \|u\|_{L^{2}}^{2} \\ &\quad + \frac{1}{|b|} \|u\|_{W^{1,p}}^{(2m-2+\gamma)\cdot\frac{\gamma}{\gamma}}. \end{aligned}$$
(3.7)

Noting that $p > \max\{2m, d\}$, there exists a constant $\gamma \in (0, 2)$ such that

$$(2m-2+\gamma)\cdot\frac{2}{\gamma} = p$$

Submitting (3.7) into (3.6), we have

$$\frac{d}{dt}\mathbb{E}\|u(\cdot,t)\|^2 \le 2\left(a+\hat{C}\right)\mathbb{E}\|u\|_{L^2}^2 + \sigma^2|D|,\tag{3.8}$$

where

$$\hat{C} = \frac{2 - \gamma}{2} \left(\frac{\gamma |D| |b| C_{\infty}^p}{2} \right)^{\frac{\gamma}{2 - \gamma}}.$$

Solving the differential inequality (3.8) gives

$$\mathbb{E}\|u(\cdot,t)\|^{2} \leq \left(\mathbb{E}\|u_{0}\|^{2} + \frac{\sigma^{2}|D|}{2(a+\hat{C})}\right)e^{2(a+\hat{C})t} - \frac{\sigma^{2}|D|}{2(a+\hat{C})}.$$
(3.9)

Note that $a + \hat{C} < 0$ implies that $e^{2(a+\hat{C})t} < 1$ for all t > 0. Furthermore, the assumption

$$\mathbb{E}||u_0||^2 + \frac{\sigma^2|D|}{2(a+\hat{C})} > 0$$

yields that

$$\mathbb{E}\|u(\cdot,t)\|^2 \le \mathbb{E}\|u_0\|^2,$$

which completes the proof. \Box

Example Consider

$$\begin{cases} du = (\Delta_4 u - u + \frac{1}{C_{\infty}^4} u^2) dt + \sigma dW_t, & t > 0, \quad x \in (0, 1), \\ u(x, 0) = u_0(x), & x \in (0, 1), \\ u(x, t) = 0, & t > 0, \quad x \in \partial(0, 1). \end{cases}$$

It is easy to check that $\gamma = 1$ satisfies $(2m - 2 + \gamma) \cdot \frac{2}{\gamma} = p$, where m = 3/2, p = 4. Then if the initial data satisfies

$$\frac{\sigma^2}{2\mathbb{E}\|u_0\|^2} < \frac{3}{4},$$

then Theorem 3.2 shows that the trivial solution 0 of the above problem will be mean square stable.

Remark 3.2 (1) In Theorem 3.2, we assume the constant b satisfies (3.5). Note that the constant C_{∞} depends on the domain D, the dimension d and the constant p, and thus it is hard to give a concrete constant in an example. The reason is that we used the embedding inequality (3.3). On the other hand, we can use the following embedding inequality replaced (3.3):

$$W^{1,p}(D) \hookrightarrow W^{1,2m}(D) \hookrightarrow L^{2m}(D), \quad p > 2m.$$

Let $C_{2m,2m}$ be the Sobolev embedding constant, i.e., $\|u\|_{L^{2m}(D)} \leq C_{2m,2m} \|u\|_{W^{1,2m}(D)}$. Then under the assumptions that $b \leq \frac{2}{C_{2m,2m}}$ and

$$2a\mathbb{E}||u_0||^2 + \sigma^2|D| < 0,$$

the trivial solution 0 of (3.2) is mean square stable.

(2) We now explain why we did not get the results of stochastic stability. Like the case of stochastic differential equations, we try to use a Lyapunov function $V(u) = ||u||^{2r}$ with 0 < r < 1 to prove the stochastic stability. For additive noise, we can not prove that $||u(\cdot,t)||^2 > 0$ for all t > 0. In order to use the Itô formula, we consider the following Lyapunov functional $V(u) = (||u|| + \kappa)^{2r}$ with 0 < r < 1 and $0 < \kappa \ll 1$. This leads to the expression

$$\frac{d}{dt}\mathbb{E}(\|u\|^{2}+\kappa)^{r} = 2r\mathbb{E}\left[(\|u\|^{2}+\kappa)^{r-1}\int_{D}u(\Delta_{p}u(x,t)+f(u))dx\right] \\
+r\sigma^{2}|D|\mathbb{E}\left[(\|u\|^{2}+\kappa)^{r-1}\right] \\
+2\sigma^{2}r(r-1)\mathbb{E}(\|u\|^{2}+\kappa)^{r-2}\left(\int_{D}udx\right)^{2} \\
\leq \mathbb{E}\left[r(\|u\|^{2}+\kappa)^{r-1}(a+\hat{C})\|u(\cdot,t)\|^{2} \\
+r\sigma^{2}(\|u\|^{2}+\kappa)^{r-1}\left(|D|+2(r-1)\frac{\left(\int_{D}udx\right)^{2}}{\|u\|^{2}+\kappa}\right)\right].$$
(3.10)

Due to the difference $(\int_D u dx)^2$ and $(\int_D |u| dx)^2$, we can not get any help to control the term |D|. Note that

$$\int_D u dx = 0$$

maybe happen, so we can not use this term. Even though the term $(\int_D u dx)^2$ is replaced by $(\int_D |u| dx)^2$, we can not get the desired result. The reason is the followings. The Hölder inequality implies that

$$\int_{D} |u| dx \le |D|^{\frac{1}{2}} \left(\int_{D} |u|^{2} dx \right)^{\frac{1}{2}}.$$

Consequently,

$$\frac{\|u\|_{L^1}^2}{\|u\|_{L^2}^2} \le |D|$$

Hence we can not use the above inequality in (3.10). The aim of the above discussion is to show the last two terms of right-side hand of (3.10) are in the same level, which are different from the first term for SPDEs.

But for SDEs, there will be another case. In this case, let |D| = 1, then we have

$$|D| + 2(r-1)\frac{\left(\int_D u dx\right)^2}{\|u\|^2 + \kappa} = 1 + \frac{2(r-1)}{1+\kappa}.$$

Taking 0 < r < 1/2 such that $2r + \kappa < 1$, we get

$$\frac{d}{dt}\mathbb{E}(\|u\|^2 + \kappa)^r \le 0.$$

Letting $\kappa \to 0$, and using the Chebyshev inequality, we obtain the stochastic stability. In all, we find there is a significant difference between SPDEs and SDEs in the stability theory.

We remark that the noise can be easily generalized the cylindrical Wiener process. We first generalize the classical results of deterministic reaction-diffusion equation [24, Theorem 4.2.1, p 166] to the following equation

$$\begin{cases} du(x,t) = (\Delta u(x,t) + f(x,t,u))dt + \sigma u dW_t, & t > 0, \quad x \in D, \\ u(x,0) = u_0(x), & x \in D, \\ u(x,t) = 0, & t > 0, \quad x \in \partial D. \end{cases}$$
(3.11)

Theorem 3.3 Assume that f(x,t,0) = 0, $f \in C^1(D \times [0,\infty) \times (-\infty,\infty))$.

(i) If there exists a constant $\alpha > 0$ such that for all $(x,t) \in D \times [0,\infty)$, $\eta \in \mathbb{R}$, we have

$$f(x,t,\eta) \le (\lambda_1 - \alpha)\eta,$$

then for the initial data u_0 satisfying $0 \le u_0(x) \le \rho \phi_1(x)$ with $\rho > 0$, problem (3.11) admits a unique positive solution u(x,t) and the following estimate holds almost surely

$$0 \le u(x,t) \le \rho e^{-(\alpha + \frac{\sigma^2}{2})t + \sigma W_t} \phi_1(x), \quad (x,t) \in D \times [0,\infty).$$

Consequently,

$$\mathbb{E}u(x,t) \le \rho e^{-\alpha t} \phi_1(x). \tag{3.12}$$

Assume further that the initial data u_0 is a deterministic function, we have

$$\mathbb{P}\left\{\int_{D} u(x,t)dx \le \rho e^{-(\alpha + \frac{\sigma^2}{2})t}\right\} \ge \frac{1}{2}.$$
(3.13)

(ii) If there exists a constant $\alpha > 0$ such that for all $(x,t) \in D \times [0,\infty)$, $\eta \ge 0$, we have

$$f(x,t,\eta) \ge (\lambda_1 + \alpha)\eta,$$

then for every $\delta > 0$, when $u_0(x) \ge \delta \phi_1(x)$, problem (3.11) admits a unique positive solution u(x,t), which exists globally or finite time blowup. On the lifespan, the following estimate holds almost surely

$$u(x,t) \ge \delta e^{(\alpha - \frac{\sigma^2}{2})t + \sigma W_t} \phi_1(x), \quad (x,t) \in D \times [0,\infty).$$

Consequently, $\mathbb{E} \| u(t) \|^2 \ge \delta e^{\alpha t} \mathbb{E} \| u_0 \|^2$.

Proof. We first change the stochastic reaction-diffusion equation into random reaction-diffusion, then by using comparison principle, the desired results are obtained. More precisely, let $v(x,t) = e^{-\sigma W_t} u(x,t)$, then v(x,t) satisfies that

$$\begin{cases} \frac{\partial}{\partial t}v(x,t) = \Delta v(x,t) - \frac{\sigma^2}{2}v(x,t) + e^{-\sigma W_t}f(x,t,e^{\sigma W_t}v(x,t)), & t > 0, \ x \in D, \\ v(x,0) = u_0(x), & x \in D, \\ v(x,t) = 0, & t > 0, \ x \in \partial D. \end{cases}$$
(3.14)

By using the assumptions, we get

$$\frac{\partial}{\partial t}v(x,t) \le \Delta v(x,t) - \frac{\sigma^2}{2}v(x,t) + (\lambda_1 - \alpha)v(x,t).$$

It is easy to check that $\bar{v} = \rho e^{-(\alpha + \frac{\sigma^2}{2})t} \phi_1(x)$ is an upper solution of (3.14) and 0 is a lower solution to (3.14), which implies that for $(x,t) \in D \times [0,\infty)$

$$0 \le v(x,t) \le \rho e^{-(\alpha + \frac{\sigma^2}{2})t} \phi_1(x) \Longleftrightarrow 0 \le u(x,t) \le \rho e^{-(\alpha + \frac{\sigma^2}{2})t + \sigma W_t} \phi_1(x), \quad a.s.,$$

which implies that

$$0 \le \int_D u(x,t) dx \le \rho e^{-(\alpha + \frac{\sigma^2}{2})t + \sigma W_t}, \quad a.s.$$

By using $\mathbb{E}[e^{\sigma W_t}] = e^{\frac{\sigma^2}{2}t}$, we have

$$\mathbb{E}u(x,t) \le e^{(\alpha-\lambda_1)t}\phi_1(x).$$

Note that

$$\left\{\int_D u(x,t)dx \le \rho e^{-(\alpha + \frac{\sigma^2}{2})t}\right\} \Longleftrightarrow \left\{e^{\sigma W_t} \le 1\right\} \supset \{W_t \le 0\},$$

thus we have

$$\mathbb{P}\left\{\int_{D} u(x,t)dx \le \rho e^{-(\alpha + \frac{\sigma^2}{2})t}\right\} \ge \mathbb{P}\{W_t \le 0\} = \frac{1}{2},$$

which proves (3.13).

Next, we prove (ii). Note that

$$\frac{\partial}{\partial t}v(x,t) \ge \Delta v(x,t) - \frac{\sigma^2}{2}v(x,t) + (\lambda_1 + \alpha)v(x,t).$$

It follows from that $\underline{v} = \delta e^{(\alpha - \frac{\sigma^2}{2})t}$ is a lower solution of (3.14), thus we have the desired inequality. The proof is complete. \Box

Remark 3.3 Following Theorem 3.3, it is easy to see that in mean square sense, the solution of (3.11) keeps the same properties as the deterministic case, which is different from the additive noise, see Theorems 3.1 and 3.2. Of course, the big difference between the stochastic and deterministic cases is that there exists an event such that whose probability is large than 0, where the event is that the solution of stochastic case maybe have exponentially decay. In other words, in (3.13), if $-\frac{\sigma^2}{2} < \alpha < 0$, then the solution u of (3.11) satisfies $||u(t)||^2 \leq ||u_0||^2 e^{\left(\alpha - \lambda_1 - \frac{\sigma^2}{2}\right)t}$ with probability $\frac{1}{2}$. Maybe from here we can say the noise can stabilize the solutions.

The method we used in Theorem 3.3 is comparison principle, which is different from the Lyapunov functional method. The inequality (3.12) holds pointwise, which is different from the earlier results. What's more, the index $\alpha - \lambda_1$ is different from that obtained by Lyapunov method, see the next theorem. In part (ii) of Theorem 3.3 implies the unstable condition of the trivial solution 0, which is new in this field.

Comparing with the stochastic ordinary different equations, the role of the Laplacian operator in stochastic reaction-diffusion equations gives a help with λ_1 in the stability of trivial solutions. Indeed, the reason is the Poincare inequality.

The impact of multiplicative noise in Theorem 3.3 is not satisfied. We give the next result.

Theorem 3.4 Assume that f(x,t,0) = 0 and there exists a constant K > 0 such that for all $(x,t) \in D \times [0,\infty)$, $uf(x,t,u) \leq Ku^2$. If $K - \lambda_1 + \frac{\sigma^2}{2} \leq 0$, then the trivial solution 0 is mean square stable and if

$$K - \lambda_1 - \frac{\sigma^2}{2} < 0, \tag{3.15}$$

then the trivial solution 0 is stochastically stable.

Proof. Taking Lyapunov function $V(u) = ||u||^2$, we have

$$\begin{aligned} \frac{d}{dt} \mathbb{E} \|u(t)\|^2 &= -\int_D |\nabla u(x,t)|^2 dx + \frac{\sigma^2}{2} \mathbb{E} \|u(t)\|^2 + \mathbb{E} \int_D uf(x,t,u) dx \\ &\leq \left(K - \lambda_1 + \frac{\sigma^2}{2}\right) \mathbb{E} \|u(t)\|^2, \end{aligned}$$

which yields that

$$\mathbb{E}\|u(t)\|^2 \le \mathbb{E}\|u_0\|^2 e^{\left(K-\lambda_1+\frac{\sigma^2}{2}\right)t}$$

Next, we use a Lyapunov function $V(u) = ||u||^{2r}$ with 0 < r < 1 to prove the stochastic stability. Note that in Theorem 3.3, we proved that the solution $u \ge 0$ almost surely and thus we can choose $||u||^{2r}$ as Lyapunov functional. This leads to the expression

$$\frac{d}{dt}\mathbb{E}\|u(\cdot,t)\|^{2r} = 2r\mathbb{E}\left[\|u(\cdot,t)\|^{2r-2}\int_{D}u(\Delta u(x,t)+f(x,t,u))dx\right]
+r\sigma^{2}\mathbb{E}\|u(\cdot,t)\|^{2r}+2\sigma^{2}r(r-1)\mathbb{E}\|u(\cdot,t)\|^{2r}
\leq \mathbb{E}\left[2r\|u(\cdot,t)\|^{2r}\left(K-\lambda_{1}+\sigma^{2}r-\frac{\sigma^{2}}{2}\right)\right].$$
(3.16)

If $K - \lambda_1 - \frac{\sigma^2}{2} < 0$, we can choose $0 < r < \frac{1}{2}$ such that

$$K - \lambda_1 + \sigma^2 r - \frac{\sigma^2}{2} \le 0,$$

then from the Chebyshev inequality, stochastic stability for the solution of (3.2) follows from (3.16). The proof is complete. \Box

Remark 3.4 It follows from Theorem 3.4 that the multiplicative noise can make solution stable in sense of stochastically stable. Comparing Theorem 3.4 with 3.3, we can take $K = \lambda_1 + \sigma^2/2 - \varepsilon > \lambda_1 - \alpha$, where $0 < \varepsilon \ll 1$.

In the above Theorems, we assume that the noise term satisfies the global Lipschitz condition. In the following theorem, we will see that the assumption can be weaken as local Lipschitz condition. In order to do this, we consider the following equation

$$\begin{cases} du = (\Delta u - k_1 u^r) dt + k_2 u^m dW_t(x), & t > 0, \ x \in D, \\ u(x,0) = u_0(x), & x \in D, \\ u(x,t) = 0, & t > 0, \ x \in \partial D, \end{cases}$$
(3.17)

where k_1, k_2, r and m are positive constants. Moreover, $W_t(x)$ is a Wiener random field with the covariance function q. In our paper [14], we proved the existence of global solution of (3.17) under the assumptions of Theorem 3.5. Moreover, we proved the solutions keep non-negative almost surely.

Theorem 3.5 Assume that r is an odd number and $1 < m < \frac{1+r}{2}$. Assume further that there exists a positive constant q_0 such that the covariance function q(x,y) satisfies the condition $\sup_{x,y\in\bar{D}}q(x,y) \leq q_0$. Then if $\hat{\lambda} < \lambda_1$, then the trivial solution 0 is exponentially mean square stable with the index $\lambda_1 - \hat{\lambda}$, where

$$\hat{\lambda} := \frac{r+1-2m}{r-1} \left(\frac{k_1(r-1)}{2m-2}\right)^{-\frac{2m-2}{r+1-2m}} (q_0 k_2)^{\frac{r-1}{r+1-2m}}.$$

In particular, when m = 2 and r > 3, we assume further that there exists a positive constant q_1 such that the covariance function q(x, y) satisfies the condition $\sup_{x,y\in \overline{D}} q(x, y) \ge q_1$. If

$$\lambda_1 > \frac{r-3}{r-1} \left(\frac{k_1(r-1)}{2}\right)^{-\frac{2}{r-3}} (q_0 k_2)^{\frac{r-1}{r-3}} - 2k_2^2 q_1,$$

then the trivial solution 0 is stochastic stable.

Proof. The proof is similar to that of Theorem 3.4 and we give outline of the proof for completeness. Taking Lyapunov function $V(u) = ||u||^2$, we have the following inequality

$$\frac{d}{dt}\mathbb{E}\|u(t)\|^{2} = -2\int_{D}|\nabla u(x,t)|^{2}dx + k_{2}^{2}\mathbb{E}\int_{D}u^{2m}(x,t)q(x,x)dx - 2k_{1}\mathbb{E}\int_{D}u^{r+1}(x,t)dx \\
\leq -2\lambda_{1}\mathbb{E}\|u(t)\|^{2} - 2k_{1}\mathbb{E}\|u(t)\|_{L^{1+r}}^{1+r} + k_{2}^{2}q_{0}\mathbb{E}\|u(t)\|_{L^{2m}}^{2m}.$$

By using the interpolation inequality

$$||u||_{L^r} \le ||u||_{L^p}^{\theta} ||u||_{L^q}^{1-\theta},$$

with r = 2m, p = 2 and q = 4, we have

$$\begin{aligned} q_0 \|u\|_{L^{2m}}^{2m} &\leq q_0 \|u\|_{L^2}^{2m\theta} \|u\|_{L^{1+r}}^{2m(1-\theta)} \\ &\leq k_1 \|u\|_{L^{1+r}}^{2m(1-\theta)\frac{1}{1-m\theta}} + m\theta \left(\frac{k_1}{1-m\theta}\right)^{-\frac{1-m\theta}{m\theta}} (q_0 k_2)^{\frac{1}{m\theta}} \|u\|_{L^2}^2 \\ &= k_1 \|u\|_{L^{1+r}}^{1+r} + \hat{\lambda} \|u\|_{L^2}^2, \end{aligned}$$

where

$$\theta = \frac{r+1-2m}{mr-m}, \quad \hat{\lambda} := \frac{r+1-2m}{r-1} \left(\frac{k_1(r-1)}{2m-2}\right)^{-\frac{2m-2}{r+1-2m}} (q_0k_2)^{\frac{r-1}{r+1-2m}}.$$

Consequently, we have

$$\mathbb{E}\|u(t)\|^2 \le \mathbb{E}\|u_0\|^2 e^{-(\lambda_1 - \hat{\lambda})t},$$

where implies that the trivial solution 0 is exponentially mean square stable.

Next, we use a Lyapunov function $V(u) = ||u||^{2\gamma}$ with $0 < \gamma < 1$ to prove the stochastic stability (note that the solutions of (3.17) is non-negative function, see [14]). This leads to the expression

$$\begin{split} \frac{d}{dt} \mathbb{E} \|u(\cdot,t)\|^{2\gamma} &= \gamma \mathbb{E} \left[\|u(\cdot,t)\|^{2\gamma-2} \int_D u(\Delta u(x,t) - k_1 u^r(x,t)) dx \right] \\ &+ \gamma k_2^2 \mathbb{E} \|u(\cdot,t)\|^{2\gamma-2} \int_D u^{2m} dx \\ &+ 2k_2^2 \gamma(\gamma-1) \mathbb{E} \|u(\cdot,t)\|^{2\gamma-4} \int_D \int_D q(x,y) u^m(x,t) u^m(y,t) dx \\ &\leq \gamma \mathbb{E} \left[\|u(\cdot,t)\|^{2\gamma-2} \int_D u(\Delta u(x,t) - k_1 u^r(x,t)) dx \right] \\ &+ \gamma k_2^2 \mathbb{E} \|u(\cdot,t)\|^{2\gamma-2} \int_D u^{2m} dx + 2k_2^2 q_0 \gamma(\gamma-1) \mathbb{E} \|u(\cdot,t)\|^{2\gamma}. \end{split}$$

Then by using similar method in proving mean square stable, we can choose 0 < r < 1 such that

$$\frac{d}{dt}\mathbb{E}\|u(\cdot,t)\|^{2\gamma} \le 0.$$

From the Chebyshev inequality, stochastic stability for the solution of (3.17) is obtained. The proof is complete. \Box

In paper [21], the authors considered the following problem

$$\begin{cases} dX_t = X_t(b(X_t) + k_1 X_t^{m-1})dt + k_2 X_t^{\frac{m+1}{2}} \phi(X_t) dW_t, & t > 0, \\ X_0 = x > 0, \end{cases}$$
(3.18)

where $k_1, k_2 \in \mathbb{R}, m \geq 1$. In [15], we considered the competition between the nonlinear term and noise term. The result of [21] generalized the results of [15]. Now we first recall the main results of [21].

Proposition 3.1 [21, Theorem 1.1] Let k_1 be a real number which is not zero. Assume $rb(r) \in C^1(\mathbb{R}_+)$ and there exist two positive numbers c_0 and $m_0(< m)$ such that

$$|b(r)| \le c_0(1+r^{m_0-1}), \quad r \in \mathbb{R}_+.$$

Assume in addition that $r^{\frac{m+1}{2}}\phi(r) \in C^1(\mathbb{R}_+)$. Let $\beta \in (0,1)$ and suppose there is a $r_0 > 0$ such that

$$\inf_{r \ge r_0} \phi(r) > \sqrt{\frac{2|k_1|}{(1-\beta)k_2^2}}.$$

There is a unique solution $X_t(x)$ for (3.18) on $t \ge 0$ and the solution is positive for all $t \ge 0$ almost surely. Moreover, for every T > 0

$$\sup_{0 \le t \le T} \mathbb{E} X_t^\beta(x) < +\infty.$$

Moreover, [21, Theorem 1.2] shows that the result in proposition is sharp. More precisely, if there exists $\gamma \in (\beta, 1)$ such that

$$\sup_{r \ge r_0} \phi(r) < \sqrt{\frac{2|k_1|}{(1-\gamma)k_2^2}}$$

then there is a real number $T_0 > 0$ such that

$$\sup_{0 \le t \le T} \mathbb{E}X_t^{\gamma}(x) = +\infty.$$

The above results implies that the trivial solution 0 is not mean square stable. But the stochastic stability would be possible. In the following result, we will give a positive answer.

Theorem 3.6 Let all the assumptions of Proposition 3.1 hold. Assume further that

$$rb(r) \le c_1 r + c_2 r^{m_0}, \quad 1 < m_0 < m, \quad r \in \mathbb{R}_+, \ c_1 < 0 < c_2.$$

(i) If $\inf_{r>0} \phi(r) \ge 1$ and

$$c_1 + \frac{\left[p(k_1 - \frac{k_2^2}{2})\right]^{-p/q}}{q} < 0,$$

where

$$p = \frac{m-1}{m_0 - 1}, \quad q = \frac{m-1}{m - m_0}.$$

Then the trivial solution 0 is stochastic stable.

(ii) If $\phi(r) = r^{\frac{\alpha}{2}}$ with $\alpha \ge 0$ and

$$\frac{m-1}{\alpha+m-1} \left(\frac{-c_1(\alpha+m-1)}{2\alpha}\right)^{-\frac{\alpha}{m-1}} k_1^{\frac{\alpha+m-1}{m-1}} + \frac{m_0-1}{\alpha+m-1} \left(\frac{-c_1(\alpha+m-1)}{2(\alpha+m-m_0)}\right)^{-\frac{\alpha+m-m_0}{m_0-1}} c_2^{\frac{\alpha+m-1}{m_0-1}} < \frac{k_2^2}{2}$$

Then the trivial solution 0 is stochastic stable.

Proof. (i) We use a Lyapunov function $V(X) = |X|^{\beta}$ with $0 < \beta < 1$ being fixed later to prove the stochastic stability (noting that the solutions is positive almost surely, or one can use $(|X| + \kappa)^{\beta}$ to replace $|X|^{\beta}$ and then let $\kappa \to 0$). This leads to the expression

$$\frac{d}{dt}\mathbb{E}|X|^{\beta} = \beta\mathbb{E}|X|^{\beta}(b(X_{t}) + k_{1}X_{t}^{m-1}) + \frac{1}{2}k_{2}^{2}\beta(\beta-1)\mathbb{E}|X_{t}|^{\beta+m-1}\phi^{2}(X_{t}) \\
\leq \beta\mathbb{E}|X|^{\beta}\left[c_{1} + c_{2}X_{t}^{m_{0}-1} + \left(k_{1} + (\beta-1)\frac{k_{2}^{2}}{2}\right)|X_{t}|^{m-1}\right].$$

Set $0 < \beta \ll 1$ such that

$$c_1 + \frac{\left[p(k_1 - \frac{(1-\beta)k_2^2}{2})\right]^{-p/q}}{q} \le 0.$$

By using the $\varepsilon\text{-}\mathrm{Young}$ inequality, we have

$$\frac{d}{dt}\mathbb{E}|X|^{\beta} \le 0.$$

From the Chebyshev inequality, stochastic stability for the solution of (3.18) is obtained.

(ii) In this case: $\phi(r) = r^{\frac{\alpha}{2}}$ with $\alpha \ge 0$. For every $\beta \in (0,1)$, if we take $r_0 = \left(\sqrt{\frac{3|k_1|}{k_2(1-\beta)}}\right)^{\frac{2}{\alpha}}$, then

$$\inf_{r \ge r_0} \phi(r) = \inf_{r \ge r_0} |r|^{\alpha/2} = \sqrt{\frac{3|k_1|}{k_2(1-\beta)}} > \sqrt{\frac{2|k_1|}{(1-\beta)k_2^2}}.$$

Hence Proposition 3.1 holds for this case.

Similar to case (i), we use a Lyapunov function $V(X) = |X|^{\beta}$ with $0 < \beta < 1$. This leads to the expression

$$\frac{d}{dt}\mathbb{E}|X|^{\beta} = \beta\mathbb{E}|X|^{\beta}(b(X_{t}) + k_{1}X_{t}^{m-1}) + \frac{1}{2}k_{2}^{2}\beta(\beta-1)\mathbb{E}|X_{t}|^{\beta+\alpha+m-1} \\
\leq \beta\mathbb{E}|X|^{\beta}\left[c_{1} + c_{2}X_{t}^{m_{0}-1} + k_{1}|X_{t}|^{m-1} + (\beta-1)\frac{k_{2}^{2}}{2}|X_{t}|^{\alpha+m-1}\right].$$

Set $0<\beta\ll 1$ such that

$$\frac{m-1}{\alpha+m-1} \left(\frac{-c_1(\alpha+m-1)}{2\alpha}\right)^{-\frac{\alpha}{m-1}} k_1^{\frac{\alpha+m-1}{m-1}} + \frac{m_0-1}{\alpha+m-1} \left(\frac{-c_1(\alpha+m-1)}{2(\alpha+m-m_0)}\right)^{-\frac{\alpha+m-m_0}{m_0-1}} c_2^{\frac{\alpha+m-1}{m_0-1}} \le \frac{k_2^2}{2}(1-\beta).$$

By using the ε -Young inequality, we have

$$\frac{d}{dt}\mathbb{E}|X|^{\beta} \le 0.$$

From the Chebyshev inequality, stochastic stability for the solution of (3.18) is obtained. \Box

Remark 3.5 Theorem 3.6 is new for SDEs. When $k_2 = 0$, the solution of (3.18) will blow up in finite time, and thus Theorem 3.6 implies that the multiplicative noise can make the solution stable.

Unfortunately, for SPDEs, we can not get the similar result to Theorem 3.6. Before we end this section, we give the reason. For simplicity, we consider the following problem

$$\begin{cases} du = (\Delta u + k_1 u^r) dt + k_2 u^m dW_t, & t > 0, \ x \in D, \\ u(x, 0) = u_0(x), & x \in D, \\ u(x, t) = 0, & t > 0, \ x \in \partial D, \end{cases}$$
(3.19)

where $k_1, k_2, r, m \in \mathbb{R}$. Under the condition r < m, the existence of global solution was established by [17]. In the following, we set forth the reason why we can not get that the trivial solution 0 is stochastic stable.

We use a Lyapunov function $V(u) = ||u||^{2\gamma}$ with $0 < \gamma < 1/2$ to prove the stochastic stability (if we worry about ||u|| = 0, we can use $(||u||^2 + \kappa)^{\gamma}$ instead and then let $\kappa \to 0$). This leads to the expression

$$\frac{d}{dt}\mathbb{E}\|u(\cdot,t)\|^{2\gamma} = \gamma \mathbb{E}\left[\|u(\cdot,t)\|^{2\gamma-2} \int_{D} u(\Delta u(x,t) + k_{1}u^{r}(x,t))dx\right]
+ \gamma \mathbb{E}\left[\|u(\cdot,t)\|^{2\gamma-2} \int_{D} q(x,x)u^{2m}dx\right]
+ 2k_{2}^{2}\gamma(\gamma-1)\mathbb{E}\|u(\cdot,t)\|^{2\gamma-4} \int_{D} \int_{D} q(x,y)u^{m+1}(x,t)u^{m+1}(y,t)dxdy
\leq -\gamma\lambda_{1}\mathbb{E}\|u(\cdot,t)\|^{2\gamma} + k_{1}\gamma\mathbb{E}\left[\|u(\cdot,t)\|^{2\gamma-2}\|u(\cdot,t)\|^{1+r}_{L^{1+r}}\right]
+ \gamma \mathbb{E}\left[\|u(\cdot,t)\|^{2\gamma-2} \left(q_{0}\|u\|^{2m}_{L^{2m}} + 2q_{1}k_{2}^{2}(\gamma-1)\frac{\|u(\cdot,t)\|^{2(m+1)}_{L^{m+1}}}{\|u(\cdot,t)\|^{2}}\right)\right]. \quad (3.20)$$

Hölder inequality implies that

$$\int_{D} |u|^{m+1}(x,t) dx \le \left(\int_{D} |u|^{2m}(x,t) dx \right)^{\frac{1}{2}} \left(\int_{D} |u|^{2}(x,t) dx \right)^{\frac{1}{2}}.$$

For the last term of right-side hand of (3.20), we want to get

$$\frac{\|u(\cdot,t)\|_{L^m}^{2m}}{\|u(\cdot,t)\|^2} \ge \|u\|_{L^{2m}}^{2m},$$

which is a contradiction with respect to the above Hölder inequality. This shows no difference from Remark 3.2.

4 Whole space

In this section, we consider the impact of noise in the whole space.

$$\begin{cases} du = (\Delta u + f(t, u))dt + \sigma(t, u)dW(x, t) & t > 0, \quad x \in \mathbb{R}, \\ u(x, 0) = u_0(x), & x \in \mathbb{R}, \end{cases}$$
(4.1)

where W(x,t) is the space-time white noise. Throughout this section, we assume that $f(t,0) = \sigma(t,0) = 0$. A mild solution to (4.2) in sense of Walsh [22] is any u which is adapted to the filtration generated by the white noise and satisfies the following evolution equation

$$u(x,t) = \int_{\mathbb{R}} K(x-y,t)u_0(y)dy + \int_0^t \int_{\mathbb{R}} K(x-y,t-s)\sigma(u,y,s)dW(y,s)dy,$$

where K(x,t) denotes the heat kernel of Laplacian operator.

Theorem 4.1 Assume that there exist non-negative constants α , β and γ such that

$$|f(u) + \alpha u| \le \beta(t)|u|, \quad |\sigma(t,u)| \le \gamma(t)|u|, \quad 2\int_0^t \beta(s)ds + \frac{1}{2\sqrt{\pi}}\int_0^t \frac{\gamma(s)}{(t-s)^{-\frac{1}{2}}}ds < 1.$$
(4.2)

Then the trivial solution 0 is exponentially mean square stable with index 2α .

Proof. Note that f and σ satisfy the global Lipschitz condition, one can prove that (4.2) admits a unique global solution. Let $v(x,t) = e^{\alpha t}u(x,t)$. Then v satisfies

$$\begin{cases} dv = (\Delta v + f(e^{-\alpha t}v) + \alpha e^{-\alpha t}v)dt + \sigma(t, e^{-\alpha t}v)dW(x, t), & t > 0, \quad x \in \mathbb{R}, \\ v(x, 0) = u_0(x), & x \in \mathbb{R}. \end{cases}$$
(4.3)

By taking the second moment and using the Walsh isometry, we get for $t \in [0, T]$

$$\begin{split} \mathbb{E}|v(x,t)|^2 &= \left(\int_{\mathbb{R}} K(x-y,t)u_0(y)dy + \int_0^t \int_{\mathbb{R}} K(x-y,t-s)[f(e^{-\alpha s}v) + \alpha e^{-\alpha s}v]dyds\right)^2 \\ &+ \int_0^t \int_{\mathbb{R}} K^2(x-y,t-s)\mathbb{E}\sigma^2(s,e^{-\alpha t}v)dyds \\ &\leq 2\max_{x\in\mathbb{R}}|u_0|^2(x) + \left(2\int_0^t \beta(s)ds + \frac{1}{2\sqrt{\pi}}\int_0^t \frac{\gamma(s)}{(t-s)^{-\frac{1}{2}}}ds\right)\max_{(x,t)\in\mathbb{R}\times[0,T]} \mathbb{E}v^2(x,t), \end{split}$$

which implies that

$$\mathbb{E}|v(x,t)|^2 \le C \max_{x \in \mathbb{R}} |u_0|^2(x),$$

where C is independent of t. And thus we complete the proof. \Box

The reason why we used the properties of heat kernel is that we did not know the existence of local strong solution of problem (4.2) with space-time white noise. For general d, we have the following result.

Theorem 4.2 Consider the Cauchy problem

$$\begin{cases} du = (\Delta u + f(t, u))dt + \sigma(t, u)dW_t & t > 0, \quad x \in \mathbb{R}^d, \\ u(x, 0) = u_0(x), & x \in \mathbb{R}^d. \end{cases}$$
(4.4)

Assume that there exists a constant K > 0 such that for all $(x, t) \in \mathbb{R}^d \times [0, \infty)$,

$$uf(t,u) \le Ku^2$$
, $\sigma^2(t,u) \le \sigma u^2$, $\sigma > 0$.

Then if $K + \frac{\sigma^2}{2} \leq 0$, then the trivial solution 0 is mean square stable and if $K - \frac{\sigma^2}{2} < 0$, then the trivial solution 0 is stochastically stable.

The proof of Theorem 4.2 is similar to Theorem 3.4 and we omit it here. Meanwhile, we remark that in the whole space the operator Δ will have no help to the stability of trivial solution.

Acknowledgment This research was partly supported by the NSF of China grants 11771123, 11501577, 11626085.

References

- P-L. Chow, Stochastic partial differential equations, Chapman Hall/CRC Applied Mathematics and Nonlinear Science Series. Chapman Hall/CRC, Boca Raton, FL, 2007. x+281 pp. ISBN: 978-1-58488-443-9.
- P-L. Chow, Explosive solutions of stochastic reaction-diffusion equations in mean L^p-norm, J. Differential Equations 250 (2011) 2567-2580.
- [3] G. Da Prato and J. Zabczyk, Stochastic equations in infinite dimensions, Encyclopedia of Mathematics and its Applications, 44. Cambridge University Press, Cambridge, 1992. xviii+454 pp. ISBN: 0-521-38529-6
- [4] K. Dareiotis, M. Gerencsér and B. Gess, Entropy solutions for stochastic porous media equations, J. Differential Equations 266 (2019) 3732-3763.
- [5] E. Fedrizzi, F. Flandoli, Noise prevents singularities in linear transport equations, J. Funct. Anal. 264 (6) (2012) 1329-1354.
- [6] F. Flandoli, M. Gubinelli, E. Priola, Well-posedness of the transport equation by stochastic perturbation, Invent. Math. 180 (1) (2010) 1-53.
- [7] P. Gassiat and B. Gess, Regularization by noise for stochastic Hamilton-Jacobi equations, Probab. Theory Related Fields 173 (2019) 1063-1098.
- [8] B. Gess and P. Souganidis, Long-time behavior, invariant measures, and regularizing effects for stochastic scalar conservation laws, Comm. Pure Appl. Math. 70 (2017) 1562-1597.
- B. Gess and M. Hofmanová, Well-posedness and regularity for quasilinear degenerate parabolichyperbolic SPDE, Ann. Probab. 46 (2018) 2495-2544.
- B. Gess and X. Lamy, Regularity of solutions to scalar conservation laws with a force, Ann. Inst. H. Poincare Anal. Non Lineaire 36 (2019) 505-521.
- [11] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, 2nd Ed., Springer-Verlag, New York, 1983.
- [12] R. Khasminskii, Stochastic stability of differential equations, Springer Heidelberg Dordrecht London New York, 2011.
- [13] K. Liu and X. Mao, Exponential stability of non-linear stochastic evolution equations, Stochastic Process. Appl. 78 (1998) 173-193.
- [14] G. Lv and J. Duan, Impacts of noise on a class of partial differential equations, J. Differential Equations, 258 (2015) 2196-2220.

- [15] G. Lv and J. Duan, L. Wang and J. Wu, Impact of noise on ordinary differential equations, Dynamic system and Applications, 27 (2018) 225-236.
- [16] G. Lv, H. Gao, J. Wei and J-L. Wu, BMO and Morrey-Campanato estimates for stochastic convolutions and Schauder estimates for stochastic parabolic equations, J. Differential Equations 266 (2019) 2666-2717.
- [17] G. Lv and J. Wei, Global existence and non-existence of stochastic parabolic equations, J. Differential Equations in press (arXiv:1902.07389).
- [18] M. Mackey and G. Nechaeva, Noise and stability in Differential Delay Equations, J. Dynam. Differential Equations 6 (1994) 395-426.
- [19] X. Mao, Stability of stochastic differential equations with respect to semimartingales, Longman, 1991.
- [20] X. Mao, Exponential stability of stochastic differential equations, Marcel Dekker, 1994.
- [21] R. Tian, J. Wei and J. Wu Generalized population dynamics equations with environmental noises, submitted
- [22] John B. Walsh, An introduction to stochastic partial differential equations, volume 1180 of Lecture Notes in Math., pages 265-439, Springer Berlin, 1986.
- [23] Z. Wang and X. Li, Stability and moment boundedness of the stochastic linear age-structured model, J. Dynam. Differential Equations 31 (2019) 2109-2125.
- [24] Q. Ye and Z. Li, Introduction to reaction-diffusion equations, Science Press, Beijing, 1990.