Least squares estimation for non-ergodic weighted fractional Ornstein-Uhlenbeck process of general parameters

Abdulaziz Alsenafi¹* Mishari Al-Foraih² Khalifa Es-Sebaiy³

Kuwait University

Abstract

Let $B^{a,b}:=\{B^{a,b}_t, t\geq 0\}$ be a weighted fractional Brownian motion of parameters a>-1, |b|<1, |b|<a+1. We consider a least square-type method to estimate the drift parameter $\theta>0$ of the weighted fractional Ornstein-Uhlenbeck process $X:=\{X_t,t\geq 0\}$ defined by $X_0=0;\ dX_t=\theta X_t dt+dB^{a,b}_t$. In this work, we provide least squares-type estimators for θ based continuous-time and discrete-time observations of X. The strong consistency and the asymptotic behavior in distribution of the estimators are studied for all (a,b) such that $a>-1,\ |b|<1,\ |b|<1,\ |b|<1$. Here we extend the results of [18, 19] (resp. [4]), where the strong consistency and the asymptotic distribution of the estimators are proved for $-\frac{1}{2}< a<0,\ -a< b< a+1$ (resp. $-1< a<0,\ -a< b< a+1$). Simulations are performed to illustrate the theoretical results.

Key words: Drift parameter estimation; Weighted fractional Ornstein-Uhlenbeck process; Strong consistency; Asymptotic distribution.

2010 Mathematics Subject Classification: 62F12; 60F05; 60G15; 60H05

1 Introduction

Parameter estimation for non-ergodic type diffusion processes has been developed in several papers. For motivation and further references, we refer the reader to Basawa and Scott [1], Dietz and Kutoyants [5], Jacod [14] and Shimizu [20].

Let $B^{a,b} := \{B_t^{a,b}, t \geq 0\}$ be a weighted fractional Brownian motion (wfBm) with parameters (a,b) such that a > -1, |b| < 1 and |b| < a + 1, that is, $B^{a,b}$ is defined as a centered Gaussian process starting from zero with covariance

$$R^{a,b}(t,s) = E\left(B_t^{a,b}B_s^{a,b}\right) = \int_0^{s \wedge t} u^a \left[(t-u)^b + (s-u)^b \right] du, \quad s, t \ge 0.$$
 (1.1)

¹Department of Mathematics, Faculty of Science, Kuwait University, Kuwait. E-mail: abdulaziz.alsenafi@ku.edu.kw

^{*}Corresponding author

²Department of Mathematics, Faculty of Science, Kuwait University, Kuwait. E-mail: mishari.alforaih@ku.edu.kw

³Department of Mathematics, Faculty of Science, Kuwait University, Kuwait. E-mail: khalifa.essebaiy@ku.edu.kw

For a=0, -1 < b < 1, the wfBm is a fractional Brownian motion (fBm). The process $B^{a,b}$ was introduced by [3] as an extension of fBm. Moreover, it shares several properties with fBm, such as self-similarity, path continuity, behavior of increments, long-range dependence, non-semimartingale, and others. But, unlike fBm, the wfBm does not have stationary increments for $a \neq 0$. For more details about the subject, we refer the reader to [3].

In this work we consider the non-ergodic Ornstein-Uhlenbeck process $X := \{X_t, t \ge 0\}$ driven by a wfBm $B^{a,b}$, that is the unique solution of the following linear stochastic differential equation

$$X_0 = 0; dX_t = \theta X_t dt + dB_t^{a,b}, (1.2)$$

where $\theta > 0$ is an unknown parameter.

An example of interesting problem related to (1.2) is the statistical estimation of θ when one observes X. In recent years, several researchers have been interested in studying statistical estimation problems for Gaussian Ornstein-Uhlenbeck processes. Let us mention some works in this direction in this case of Ornstein-Uhlenbeck process driven by a fractional Brownian motion $B^{0,b}$, that is, the solution of (1.2), where a=0. In the ergodic case corresponding to $\theta < 0$, the statistical estimation for the parameter θ has been studied by several papers, for instance [12, 8, 13, 11, 6] and the references therein. Further, in the non-ergodic case corresponding to $\theta > 0$, the estimation of θ has been considered by using least squares method, for example in [7, 2, 9, 10] and the references therein.

Here our aim is to estimate the drift parameter θ based on continuous-time and discrete-time observations of X, by using least squares-type estimators (LSEs) for θ . First we will consider the following LSE

$$\widetilde{\theta}_t = \frac{X_t^2}{2\int_0^t X_s^2 ds}, \quad t \ge 0, \tag{1.3}$$

as statistic to estimate θ based on the continuous-time observations $\{X_s, s \in [0, t]\}$ of (1.2), as $t \to \infty$. We will prove the strong consistency and the asymptotic behavior in distribution of the estimator $\tilde{\theta}_t$ for all parameters a > -1, |b| < 1 and |b| < a + 1. Our results extend those proved in [18, 19], where $-\frac{1}{2} < a < 0$, -a < b < a + 1 only.

Further, from a practical point of view, in parametric inference, it is more realistic and interesting to consider asymptotic estimation for (1.2) based on discrete observations. So, we will assume that the process X given in (1.2) is observed equidistantly in time with the step size Δ_n : $t_i = i\Delta_n$, i = 0, ..., n, and $T_n = n\Delta_n$ denotes the length of the "observation window". Then we will consider the following estimators

$$\hat{\theta}_n = \frac{\sum_{i=1}^n X_{t_{i-1}} (X_{t_i} - X_{t_{i-1}})}{\Delta_n \sum_{i=1}^n X_{t_{i-1}}^2},$$
(1.4)

and

$$\check{\theta}_n = \frac{X_{T_n}^2}{2\Delta_n \sum_{i=1}^n X_{t_{i-1}}^2}$$
(1.5)

as statistics to estimate θ based on the sampling data $X_{t_i}, i = 0, \ldots, n$, as $\Delta_n \longrightarrow 0$ and $n \longrightarrow \infty$. We will study the asymptotic behavior and the rate consistency of the estimators $\hat{\theta}_n$ and $\check{\theta}_n$ for all parameters a > -1, |b| < 1 and |b| < a + 1. In this case, our results extend those proved in [4], where -1 < a < 0, -a < b < a + 1 only.

The rest of the paper is organized as follows. In Section 2, we present auxiliary results that are used in the calculations of the paper. In Section 3, we prove the consistency and the asymptotic distribution of the estimator $\tilde{\theta}_t$ given in (1.3), based on the continuous-time observations of X. In Section 3, we study the asymptotic behavior and the rate consistency of the estimators $\hat{\theta}_n$ and $\check{\theta}_n$ defined in (1.4) and (1.5), respectively, based on the discrete-time observations of X. Our theoretical study is completed with simulations. We end the paper with a short review on some results from [7, 9] needed for the proofs of our results.

2 Auxiliary Results

This section is devoted to prove some technical ingredients, which will be needed throughout this paper.

In the following lemma we provide a useful decomposition of the covariance function $R^{a,b}(t,s)$ of $B^{a,b}$.

Lemma 2.1. Suppose that a > -1, |b| < 1 and |b| < a + 1. Then we can rewrite the covariance $R^{a,b}(t,s)$ of $B^{a,b}$, given in (2.1) as

$$R^{a,b}(t,s) = \beta (a+1,b+1) [t^{a+b+1} + s^{a+b+1}] - m(t,s),$$
 (2.1)

where $\beta(c,d) = \int_0^1 x^{c-1} (1-x)^{d-1}$ denotes the usual beta function, and the function m(t,s) is defined by

$$m(t,s) := \int_{s\wedge t}^{s\vee t} u^a (t\vee s - u)^b du. \tag{2.2}$$

Proof. We have for every $s, t \ge 0$,

$$R^{a,b}(t,s)$$

$$= E\left(B_t^{a,b}B_s^{a,b}\right)$$

$$= \int_0^{s\wedge t} u^a \left[(t-u)^b + (s-u)^b \right] du$$

$$= \int_0^{s\wedge t} u^a \left[(t\vee s-u)^b + (t\wedge s-u)^b \right] du$$

$$= \int_0^{s\wedge t} u^a (t\vee s-u)^b du + \int_0^{s\wedge t} u^a (t\wedge s-u)^b du$$

$$= \int_0^{s\vee t} u^a (t\vee s-u)^b du - \int_{s\wedge t}^{s\vee t} u^a (t\vee s-u)^b du + \int_0^{s\wedge t} u^a (t\wedge s-u)^b du.$$
 (2.4)

Further, making change of variables x = u/t, we have for every $t \ge 0$,

$$\int_{0}^{t} u^{a} (t - u)^{b} du = t^{b} \int_{0}^{t} u^{a} \left(1 - \frac{u}{t}\right)^{b} du$$

$$= t^{a+b+1} \int_{0}^{1} x^{a} (1 - x)^{b} du$$

$$= t^{a+b+1} \beta (a+1, b+1). \tag{2.5}$$

Therefore, combining (2.4) and (2.5), we deduce that

$$R^{a,b}(t,s) = \beta (a+1,b+1) \left[(t \vee s)^{a+b+1} + (t \wedge s)^{a+b+1} \right] - \int_{s \wedge t}^{s \vee t} u^{a}(t \vee s - u)^{b} du$$

$$= \beta (a+1,b+1) \left[t^{a+b+1} + s^{a+b+1} \right] - \int_{s \wedge t}^{s \vee t} u^{a}(t \vee s - u)^{b} du, \qquad (2.6)$$

which proves (2.1).

We will also need the following technical lemma.

Lemma 2.2. We have as $t \longrightarrow \infty$,

$$I_t := t^{-a} e^{-\theta t} \int_0^t e^{\theta s} m(t, s) ds \longrightarrow \frac{\Gamma(b+1)}{\theta^{b+2}}, \tag{2.7}$$

$$J_t := t^{-a} e^{-2\theta t} \int_0^t \int_0^t e^{\theta s} e^{\theta r} m(s, r) dr ds \longrightarrow \frac{\Gamma(b+1)}{\theta^{b+3}}, \tag{2.8}$$

where $\Gamma(.)$ is the standard gamma function, whereas the function m(t,s) is defined in (2.2).

Proof. We first prove (2.7). We have,

$$t^{-a}e^{-\theta t} \int_{0}^{t} e^{\theta s} m(t,s) ds = t^{-a}e^{-\theta t} \int_{0}^{t} e^{\theta s} \int_{s}^{t} u^{a}(t-u)^{b} du ds$$

$$= t^{-a}e^{-\theta t} \int_{0}^{t} du u^{a}(t-u)^{b} \int_{0}^{u} ds e^{\theta s}$$

$$= t^{-a}e^{-\theta t} \int_{0}^{t} du u^{a}(t-u)^{b} \frac{(e^{\theta u} - 1)}{\theta}$$

$$= \frac{t^{-a}e^{-\theta t}}{\theta} \int_{0}^{t} u^{a}(t-u)^{b}e^{\theta u} du - \frac{t^{-a}e^{-\theta t}}{\theta} \int_{0}^{t} u^{a}(t-u)^{b} du.$$

On the other hand, by the change of variables x = t - u, we get

$$\frac{t^{-a}e^{-\theta t}}{\theta} \int_0^t u^a (t-u)^b e^{\theta u} du = \frac{t^{-a}}{\theta} \int_0^t (t-x)^a x^b e^{-\theta x} dx$$

$$= \frac{1}{\theta} \int_0^t \left(1 - \frac{x}{t}\right)^a x^b e^{-\theta x} dx$$

$$\longrightarrow \frac{1}{\theta} \int_0^\infty x^b e^{-\theta x} dx = \frac{\Gamma(b+1)}{\theta^{b+2}}$$

as $t \longrightarrow \infty$. Moreover, by the change of variables x = u/t,

$$\frac{t^{-a}e^{-\theta t}}{\theta} \int_0^t u^a (t-u)^b du = \frac{e^{-\theta t}}{\theta} t^b \int_0^t (u/t)^a (1-\frac{u}{t})^b dx$$

$$= \frac{e^{-\theta t}}{\theta} t^{b+1} \int_0^t x^a (1-x)^b dx$$

$$= \frac{e^{-\theta t}}{\theta} t^{b+1} \beta (a+1,b+1)$$

$$\longrightarrow 0$$

as $t \longrightarrow \infty$. Thus the proof of the convergence (2.7) is done. For (2.8), using L'Hôpital's rule, we obtain

$$\lim_{t \to \infty} t^{-a} e^{-2\theta t} \int_0^t \int_0^t e^{\theta s} e^{\theta r} m(s,r) dr ds = \lim_{t \to \infty} \frac{2 \int_0^t \int_0^s e^{\theta s} e^{\theta r} m(s,r) dr ds}{t^a e^{2\theta t}}$$

$$= \lim_{t \to \infty} \frac{2 \int_0^t e^{\theta t} e^{\theta r} m(t,r) dr}{t^a e^{2\theta t} \left(2\theta + \frac{a}{t}\right)}$$

$$= \lim_{t \to \infty} \frac{2}{\left(2\theta + \frac{a}{t}\right)} t^{-a} e^{-\theta t} \int_0^t e^{\theta r} m(t,r) dr$$

$$= \frac{\Gamma(b+1)}{\theta^{b+3}},$$

where the latter equality comes from (2.7). Therefore the convergence (2.8) is proved. \square

3 LSE based on continuous-time observation

In this section we will establish the consistency and the asymptotic distribution of the least square-type estimator $\tilde{\theta}_t$ given in (1.3), based on the continuous-time observation $\{X_s, s \in [0, t]\}$ given by (1.2), as $t \to \infty$.

Recall that if $X \sim \mathcal{N}(m_1, \sigma_1)$ and $Y \sim \mathcal{N}(m_2, \sigma_2)$ are two independent random variables, then X/Y follows a Cauchy-type distribution. For a motivation and further references, we refer the reader to [17], as well as [15]. Notice also that if $N \sim \mathcal{N}(0, 1)$ is independent of $B^{a,b}$, then N is independent of Z_{∞} , since $Z_{\infty} := \int_0^{\infty} e^{-\theta s} B_s^{a,b} ds$ is a functional of $B^{a,b}$.

Theorem 3.1. Assume that a > -1, |b| < 1, |b| < a + 1, and let $\widetilde{\theta}_t$ be the estimator given in (1.3). Then, as $t \longrightarrow \infty$,

$$\widetilde{\theta}_t \longrightarrow \theta$$
 almost surely.

Moreover, as $t \to \infty$,

$$t^{-a/2}e^{\theta t}\left(\widetilde{\theta}_t-\theta\right) \xrightarrow{\text{law}} \frac{2\sigma_{B^{a,b}}}{\sqrt{E\left(Z_\infty^2\right)}}\mathcal{C}(1),$$

where $\sigma_{B^{a,b}} = \frac{\Gamma(b+1)}{\theta^{b+1}}$, $Z_{\infty} := \int_0^{\infty} e^{-\theta s} B_s^{a,b} ds$, whereas $\mathcal{C}(1)$ is the standard Cauchy distribution with the probability density function $\frac{1}{\pi(1+x^2)}$; $x \in \mathbb{R}$.

Proof. In order to prove this Theorem 3.1, using Theorem 6.1, it suffices to check that the assumptions $(\mathcal{H}1)$, $(\mathcal{H}2)$, $(\mathcal{H}3)$, $(\mathcal{H}4)$ hold.

It follows from (2.1) that for every $0 < s \le t$,

$$E\left(B_t^{a,b} - B_s^{a,b}\right)^2 = 2\int_s^t u^a (t-u)^b du$$
$$= 2(t-s)^{b+1} \int_0^1 (t(1-x) + sx)^a x^b dx,$$

where the latter inequality comes from the change of variables x = (t - u)/(t - s). Hence, it is easy to see that there exists a constant $C_{a,b}$ depending only on a, b, such that

$$E\left(B_t^{a,b} - B_s^{a,b}\right)^2 \le C_{a,b}t^a(t-s)^{b+1}.$$

Thus for all fixed T there exists a constant $C_{a,b}(T)$ depending only on a, b, T such that for every $0 < s \le t \le T$

$$E\left(B_t^{a,b} - B_s^{a,b}\right)^2 \le C_{a,b}(T)|t - s|^{(a+b+1)\wedge(b+1)},$$

where we used the fact that a + b + 1 > 0.

Therefore, using the fact that $B^{a,b}$ is Gaussian, and Kolmogorov's continuity criterion, we

deduce that $B^{a,b}$ has a version with $((a+b+1) \wedge (b+1) - \varepsilon)$ -Hölder continuous paths for every $\varepsilon \in (0, (a+b+1) \wedge (b+1))$. Thus $(\mathcal{H}1)$ holds for any δ in $(0, (a+b+1) \wedge (b+1))$. On the other hand, according to (2.1) we have for every $t \geq 0$,

$$E\left(B_t^{a,b}\right)^2 = 2\beta(1+a,1+b)t^{a+b+1},$$

which proves that $(\mathcal{H}2)$ holds for $\gamma = (a+b+1)/2$.

Now it remains to check that the assumptions ($\mathcal{H}3$) and ($\mathcal{H}4$) hold for $\nu = -a/2$ and $\sigma_{B^{a,b}} = \frac{\Gamma(b+1)}{\theta^{b+1}}$. Let us first compute the limiting variance of $t^{-a/2}e^{-\theta t}\int_0^t e^{\theta s}dB_s^{a,b}$ as $t \to \infty$. By (2.1) we obtain

$$E\left[\left(t^{-a/2}e^{-\theta t}\int_{0}^{t}e^{\theta s}dB_{s}^{a,b}\right)^{2}\right] = E\left[\left(t^{-a/2}e^{-\theta t}\left(e^{\theta t}B_{t}^{a,b} - \theta\int_{0}^{t}e^{\theta s}B_{s}^{a,b}ds\right)\right)^{2}\right]$$

$$= t^{-a}\left(R^{a,b}(t,t) - 2\theta e^{-\theta t}\int_{0}^{t}e^{\theta s}R^{a,b}(t,s)ds + \theta^{2}e^{-2\theta t}\int_{0}^{t}\int_{0}^{t}e^{\theta s}e^{\theta r}R^{a,b}(s,r)dsdr\right)$$

$$= t^{-a}\Delta_{g_{B^{a,b}}}(t) + 2\theta I_{t} - \theta^{2}J_{t}, \tag{3.1}$$

where I_t , J_t and $\Delta_{g_{B^a,b}}(t)$ are defined in (2.7), (2.8) and Lemma 6.1, respectively, whereas $g_{B^{a,b}}(s,r) = \beta(a+1,b+1) \left(s^{a+b+1} + r^{a+b+1}\right)$.

On the other hand, since $\frac{\partial g_{Ba,b}}{\partial s}(s,0) = \beta(a+1,b+1)(a+b+1)s^{a+b}$, and $\frac{\partial^2 g_{Ba,b}}{\partial s \partial r}(s,r) = 0$, it follows from (6.2) that

$$t^{-a}\Delta_{g_{B^{a,b}}}(t) = 2\beta(a+1,b+1)(a+b+1)t^{-a}e^{-2\theta t}\int_{0}^{t}s^{a+b}e^{\theta s}ds$$

$$\leq 2\beta(a+1,b+1)e^{-\theta t}t^{a+b+1}$$

$$\to 0 \text{ as } t \to \infty.$$
(3.2)

Combining (3.1), (3.2), (2.7) and (2.8), we get

$$E\left[\left(t^{-a/2}e^{-\theta t}\int_0^t e^{\theta s}dB_s^{a,b}\right)^2\right] \longrightarrow \frac{\Gamma(b+1)}{\theta^{b+1}} \quad \text{as } t \to \infty,$$

which implies that $(\mathcal{H}3)$ holds.

Hence, to finish the proof it remains to check that $(\mathcal{H}4)$ holds, that is, for all fixed $s \geq 0$

$$\lim_{t\to\infty} E\left(B_s^{a,b} t^{-a/2} e^{-\theta t} \int_0^t e^{\theta r} dB_r^{a,b}\right) = 0.$$

Let us consider s < t. According to (6.4), we can write

$$\begin{split} &E\left(B_{s}^{a,b}t^{-a/2}e^{-\theta t}\int_{0}^{t}e^{\theta r}dB_{r}^{a,b}\right)\\ &=\ t^{-a/2}\left(R^{a,b}(s,t)-\theta e^{-\theta t}\int_{0}^{t}e^{\theta r}R^{a,b}(s,r)dr\right)\\ &=\ t^{-a/2}\left(R^{a,b}(s,t)-\theta e^{-\theta t}\int_{s}^{t}e^{\theta r}R^{a,b}(s,r)dr-\theta e^{-\theta t}\int_{0}^{s}e^{\theta r}R^{a,b}(s,r)dr\right)\\ &=\ t^{-a/2}\left(e^{-\theta(t-s)}R^{a,b}(s,s)+e^{-\theta t}\int_{s}^{t}e^{\theta r}\frac{\partial R^{a,b}}{\partial r}(s,r)dr-\theta e^{-\theta t}\int_{0}^{s}e^{\theta r}R^{a,b}(s,r)dr\right). \end{split}$$

It is clear that $t^{-a/2}\left(e^{-\theta(t-s)}R^{a,b}(s,s)-\theta e^{-\theta t}\int_0^s e^{\theta r}R^{a,b}(s,r)dr\right)\longrightarrow 0$ as $t\to\infty$. Let us now prove that

$$t^{-a/2}e^{-\theta t}\int_{s}^{t}e^{\theta r}\frac{\partial R^{a,b}}{\partial r}(s,r)dr\longrightarrow 0$$

as $t \to \infty$. Using (1.1) we have for s < r

$$\frac{\partial R^{a,b}}{\partial r}(s,r) = b \int_0^s u^a (r-u)^{b-1} du$$

Applying L'Hôspital's rule we obtain

$$\lim_{t \to \infty} t^{-a/2} e^{-\theta t} \int_{s}^{t} e^{\theta r} \frac{\partial R^{a,b}}{\partial r}(s,r) dr = \lim_{t \to \infty} \frac{bt^{-a/2}}{\theta + \frac{a}{2t}} \int_{0}^{s} u^{a} (t-u)^{b-1} du$$

$$= \lim_{t \to \infty} \frac{bt^{b-1-\frac{a}{2}}}{\theta + \frac{a}{2t}} \int_{0}^{s} u^{a} (1-u/t)^{b-1} du$$

$$\longrightarrow 0 \text{ as } t \to \infty,$$

due to $b-1-\frac{a}{2}<0$. In fact, if -1< a<0, we use b< a+1, then $b< a+1<\frac{a}{2}+1$. Otherwise, if a>0, we use b<1, then $b-1-\frac{a}{2}< b-1-<0$. Therefore the proof of Theorem 3.1 is complete.

4 LSEs based on discrete-time observations

In this section, our purpose is to study the asymptotic behavior and the rate consistency of the estimators $\hat{\theta}_n$ and $\check{\theta}_n$ based on the sampling data $X_{t_i}, i = 0, \ldots, n$ of (1.2), where $t_i = i\Delta_n, i = 0, \ldots, n$, and $T_n = n\Delta_n$ denotes the length of the "observation window".

Definition 4.1. Let $\{Z_n\}$ be a sequence of random variables defined on a probability space (Ω, \mathcal{F}, P) . We say $\{Z_n\}$ is tight (or bounded in probability), if for every $\varepsilon > 0$, there exists $M_{\varepsilon} > 0$ such that,

$$P(|Z_n| > M_{\varepsilon}) < \varepsilon$$
, for all n .

4.1 The asymptotic behavior and the rate consistency of LSEs

Theorem 4.1. Assume that a > -1, |b| < 1, |b| < a + 1. Let $\hat{\theta}_n$ and $\check{\theta}_n$ be the estimators given in (1.4) and (1.5), respectively. Suppose that $\Delta_n \to 0$ and $n\Delta_n^{1+\alpha} \to \infty$ for some $\alpha > 0$. Then, as $n \to \infty$,

$$\hat{\theta}_n \longrightarrow \theta, \quad \check{\theta}_n \longrightarrow \theta \quad almost \ surely,$$

and for any $q \geq 0$,

$$\Delta_n^q e^{\theta T_n} (\hat{\theta}_n - \theta)$$
 and $\Delta_n^q e^{\theta T_n} (\check{\theta}_n - \theta)$ are not tight.

In addition, if we assume that $n\Delta_n^3 \to 0$ as $n \to \infty$, the estimators $\hat{\theta}_n$ and $\check{\theta}_n$ are $\sqrt{T_n}$ – consistent in the sense that the sequences

$$\sqrt{T_n}(\hat{\theta}_n - \theta)$$
 and $\sqrt{T_n}(\check{\theta}_n - \theta)$ are tight.

Proof. In order to prove this Theorem 4.1, using Theorem 6.2, it suffices to check that the assumptions $(\mathcal{H}1)$, $(\mathcal{H}2)$, $(\mathcal{H}5)$ hold.

From the proof of Theorem 3.1, the assumptions $(\mathcal{H}1)$, $(\mathcal{H}2)$ hold. Now it remains to check that $(\mathcal{H}5)$ holds. In this case, the process ζ is defined as

$$\zeta_t := \int_0^t e^{-\theta s} dB_s^{a,b}, \qquad t \ge 0,$$

whereas the integral is interpreted in the Young sense (see Appendix). Using the formula (6.4) and (6.3), we can write

$$E\left[\left(\zeta_{t_{i}} - \zeta_{t_{i-1}}\right)^{2}\right] = E\left[\left(\int_{t_{i-1}}^{t_{i}} e^{-\theta s} dB_{s}^{a,b}\right)^{2}\right]$$

$$= E\left[\left(e^{-\theta t_{i}} B_{t_{i}}^{a,b} - e^{-\theta t_{i-1}} B_{t_{i-1}}^{a,b} + \theta \int_{t_{i-1}}^{t_{i}} e^{-\theta s} B_{s}^{a,b} ds\right)^{2}\right]$$

$$= \lambda_{g_{Ba,b}}(t_{i}, t_{i-1}) - \lambda_{m}(t_{i}, t_{i-1})$$

$$= \int_{t_{i-1}}^{t_{i}} \int_{t_{i-1}}^{t_{i}} e^{-\theta(r+u)} \frac{\partial^{2} g_{Ba,b}}{\partial r \partial u}(r, u) dr du - \lambda_{m}(t_{i}, t_{i-1})$$

$$= -\lambda_{m}(t_{i}, t_{i-1}),$$

where $\lambda_{\cdot}(t_i,t_{i-1})$ is defined in Lemma 6.2, $g_{B^{a,b}}(s,r)=\beta(a+1,b+1)\left(s^{a+b+1}+r^{a+b+1}\right)$ and $\frac{\partial^2 g_{B^{a,b}}}{\partial s \partial r}(s,r)=0$, whereas the term $\lambda_m(t_i,t_{i-1})$ is equal to

$$\lambda_m(t_i, t_{i-1}) = -2m(t_i, t_{i-1})e^{-2\theta(t_{i-1}+t_i)} + 2\theta e^{-\theta t_i} \int_{t_{i-1}}^{t_i} m(r, t_i)e^{-\theta r} dr$$
$$-2\theta e^{-\theta t_{i-1}} \int_{t_{i-1}}^{t_i} m(r, t_{i-1})e^{-\theta r} dr + \theta^2 \int_{t_{i-1}}^{t_i} \int_{t_{i-1}}^{t_i} m(r, u)e^{-\theta(r+u)} dr du.$$

Combining this with the fact for every $t_{i-1} \le u \le r \le t_i$, $i \ge 2$,

$$|m(r,u)| = \left| \int_{u}^{r} x^{a} (r-x)^{b} dx \right|$$

$$\leq \begin{cases} \left| r^{a} \int_{u}^{r} (r-x)^{b} dx \right| & \text{if } -1 < a < 0 \\ \left| u^{a} \int_{u}^{r} (r-x)^{b} dx \right| & \text{if } a > 0 \end{cases}$$

$$\leq \begin{cases} \frac{\Delta_{n}^{a+b+1}}{b+1} & \text{if } -1 < a \leq 0 \\ \frac{(n\Delta_{n})^{a} \Delta_{n}^{b+1}}{b+1} & \text{if } a > 0 \end{cases}$$

together with $\Delta_n \longrightarrow 0$, we deduce that there is a positive constant C such that

$$E\left[\left(\zeta_{t_i} - \zeta_{t_{i-1}}\right)^2\right] \leq C \begin{cases} \frac{\Delta_n^{a+b+1}}{b+1} & \text{if } -1 < a \leq 0\\ \frac{(n\Delta_n)^a \Delta_n^{b+1}}{b+1} & \text{if } a > 0, \end{cases}$$

which proves that the assumption $(\mathcal{H}5)$ holds. Therefore the desired result is obtained. \square

4.2 Numerical Results

Here we simulate 100 sample paths of the process X, given by (1.2), using a regular partition of n=2000 intervals. The tables below report the means and standard deviations of the proposed estimators $\hat{\theta}_n$ and $\check{\theta}_n$ defined, respectively, by (1.4) and (1.5) of the true value of the parameter θ . The tables confirm that the estimators $\hat{\theta}_n$ and $\check{\theta}_n$ are strongly consistent even for small values of n and have small standard deviations for different true values of θ .

	$\theta = 0.7$	$\theta = 0.9$
Mean	0.7223197	0.9075169
Median	0.7286532	0.9131851
Std. dev.	0.1066532	0.08772779

Table 1: The means and standard deviations of the estimator $\hat{\theta}_n$ for a = 0.5 and b = 0.9.

	$\theta = 0.7$	$\theta = 0.9$
Mean	0.6764275	0.8892152
Median	0.7006374	0.9028244
Std. dev.	0.1260278	0.08042517

Table 2: The means and standard deviations of the estimator $\hat{\theta}_n$ for a = 0.1 and b = 0.4.

	$\theta = 0.7$	$\theta = 0.9$
Mean	0.7234658	0.9091066
Median	0.7294589	0.9144367
Std. dev.	0.1064266	0.0879878

Table 3: The means and standard deviations of the estimator $\check{\theta}_n$ for a=0.5 and b=0.9.

	$\theta = 0.7$	$\theta = 0.9$
Mean	0.6802641	0.8911407
Median	0.7015955	0.9041038
Std. dev.	0.120526	0.07755756

Table 4: The means and standard deviations of the estimator $\check{\theta}_n$ for a=0.1 and b=0.4.

5 Conclusion

To conclude, in this paper we provide least squares-type estimators for the drift parameter θ of the weighted fractional Ornstein-Uhlenbeck process X, given by (1.2), based continuous-time and discrete-time observations of X. The novelty of our approach is that it allows, comparing with the literature on statistical inference for X discussed in [18, 19, 4], to consider the general case a > -1, |b| < 1 and |b| < a + 1. More precisely,

- We estimate the drift parameter θ of (1.2) based on the continuous-time observations $\{X_s, s \in [0, t]\}$, as $t \to \infty$. We prove the strong consistency and the asymptotic behavior in distribution of the estimator $\widetilde{\theta}_t$ for all parameters a > -1, |b| < 1 and |b| < a + 1. Our results extend those proved in [18, 19], where $-\frac{1}{2} < a < 0$, -a < b < a + 1 only.
- Suppose that the process X given in (1.2) is observed equidistantly in time with the step size Δ_n : $t_i = i\Delta_n, i = 0, \ldots, n$. We estimate the drift parameter θ of (1.2) on the sampling data $X_{t_i}, i = 0, \ldots, n$, as $\Delta_n \longrightarrow 0$ and $n \longrightarrow \infty$. We study the asymptotic behavior and the rate consistency of the estimators $\hat{\theta}_n$ and $\check{\theta}_n$ for all parameters a > -1, |b| < 1 and |b| < a + 1. In this case, our results extend those proved in [4], where -1 < a < 0, -a < b < a + 1 only.

The proofs of the asymptotic behavior of the estimators are based on a new decomposition of the covariance function $R^{a,b}(t,s)$ of the wfBm $B^{a,b}$ (see Lemma 2.1), and slight extensions of results [7] and [9] (see Theorem 6.1 and Theorem 6.2 in Appendix).

Acknowledgments

The authors would like to thank the two anonymous referees for their careful reading of the manuscript and for their valuable suggestions and remarks.

6 Appendix

Here we present some ingredients needed in the paper.

Let $G = (G_t, t \ge 0)$ be a continuous centered Gaussian process defined on some probability space (Ω, \mathcal{F}, P) (Here, and throughout the text, we assume that \mathcal{F} is the sigma-field generated by G). In this section we consider the non-ergodic case of Gaussian Ornstein-Uhlenbeck processes $X = \{X_t, t \ge 0\}$ given by the following linear stochastic differential equation

$$X_0 = 0; \quad dX_t = \theta X_t dt + dG_t, \quad t \ge 0,$$
 (6.1)

where $\theta > 0$ is an unknown parameter. It is clear that the linear equation (6.1) has the following explicit solution

$$X_t = e^{\theta t} \zeta_t, \qquad t \ge 0,$$

where

$$\zeta_t := \int_0^t e^{-\theta s} dG_s, \qquad t \ge 0,$$

whereas this latter integral is interpreted in the Young sense.

Let us introduce the following required assumptions.

- $(\mathcal{H}1)$ The process G has Hölder continuous paths of some order $\delta \in (0,1]$.
- ($\mathcal{H}2$) For every $t \geq 0$, $E\left(G_t^2\right) \leq ct^{2\gamma}$ for some positive constants c and γ .
- ($\mathcal{H}3$) There is constant ν in \mathbb{R} such that the limiting variance of $t^{\nu}e^{-\theta t}\int_0^t e^{\theta s}dG_s$ exists as $t\to\infty$, that is, there exists a constant $\sigma_G>0$ such that

$$\lim_{t \to \infty} E\left[\left(t^{\nu} e^{-\theta t} \int_0^t e^{\theta s} dG_s \right)^2 \right] = \sigma_G^2.$$

 $(\mathcal{H}4)$ For ν given in $(\mathcal{H}3)$, we have all fixed $s \geq 0$

$$\lim_{t \to \infty} E\left(G_s t^{\nu} e^{-\theta t} \int_0^t e^{\theta r} dG_r\right) = 0.$$

(H5) There exist positive constants ρ , C and a real constant μ such that

$$E\left[\left(\zeta_{t_i} - \zeta_{t_{i-1}}\right)^2\right] \le C(n\Delta_n)^{\mu}\Delta_n^{\rho}e^{-2\theta t_i} \text{ for every } i = 1, \dots, n, n \ge 1.$$

The following theorem is a slight extension of the main result in [7], and it can be established following the same arguments as in [7].

Theorem 6.1. Assume that $(\mathcal{H}1)$ and $(\mathcal{H}2)$ hold and let $\widetilde{\theta_t}$ be the estimator of the form (1.3). Then, as $t \longrightarrow \infty$,

$$\widetilde{\theta}_t \longrightarrow \theta$$
 almost surely.

Moreover, if $(\mathcal{H}1)$ - $(\mathcal{H}4)$ hold, then, as $t \to \infty$,

$$t^{\nu}e^{\theta t}\left(\widetilde{\theta}_{t}-\theta\right) \stackrel{\text{law}}{\longrightarrow} \frac{2\sigma_{G}}{\sqrt{E\left(Z_{\infty}^{2}\right)}}\mathcal{C}(1),$$

where $Z_{\infty} := \int_0^{\infty} e^{-\theta s} G_s ds$, whereas $\mathcal{C}(1)$ is the standard Cauchy distribution with the probability density function $\frac{1}{\pi(1+x^2)}$; $x \in \mathbb{R}$.

The following theorem is also a slight extension of the main result in [9], and it can be proved following line by line the proofs given in [9].

Theorem 6.2. Assume that $(\mathcal{H}1)$, $(\mathcal{H}2)$ and $(\mathcal{H}5)$ hold. Let $\hat{\theta}_n$ and $\check{\theta}_n$ be the estimators of the forms (1.4) and (1.5), respectively. Suppose that $\Delta_n \to 0$ and $n\Delta_n^{1+\alpha} \to \infty$ for some $\alpha > 0$. Then, as $n \to \infty$,

$$\hat{\theta}_n \longrightarrow \theta, \quad \check{\theta}_n \longrightarrow \theta \quad almost \ surely,$$

and for any $q \geq 0$,

$$\Delta_n^q e^{\theta T_n} (\hat{\theta}_n - \theta)$$
 and $\Delta_n^q e^{\theta T_n} (\check{\theta}_n - \theta)$ are not tight.

In addition, if we assume that $n\Delta_n^3 \to 0$ as $n \to \infty$, the estimators $\hat{\theta}_n$ and $\check{\theta}_n$ are $\sqrt{T_n}$ – consistent in the sense that the sequences

$$\sqrt{T_n}(\hat{\theta}_n - \theta)$$
 and $\sqrt{T_n}(\check{\theta}_n - \theta)$ are tight.

Lemma 6.1 ([7]). Let $g:[0,\infty)\times[0,\infty)\longrightarrow\mathbb{R}$ be a symmetric function such that $\frac{\partial g}{\partial s}(s,r)$ and $\frac{\partial^2 g}{\partial s\partial r}(s,r)$ integrable on $(0,\infty)\times[0,\infty)$. Then, for every $t\geq 0$,

$$\Delta_{g}(t) := g(t,t) - 2\theta e^{-\theta t} \int_{0}^{t} g(s,t)e^{\theta s}ds + \theta^{2}e^{-2\theta t} \int_{0}^{t} \int_{0}^{t} g(s,r)e^{\theta(s+r)}drds$$

$$= 2e^{-2\theta t} \int_{0}^{t} e^{\theta s} \frac{\partial g}{\partial s}(s,0)ds + 2e^{-2\theta t} \int_{0}^{t} ds e^{\theta s} \int_{0}^{s} dr \frac{\partial^{2} g}{\partial s \partial r}(s,r)e^{\theta r}. \tag{6.2}$$

Lemma 6.2 ([9]). Let $g:[0,\infty)\times[0,\infty)\longrightarrow\mathbb{R}$ be a symmetric function such that $\frac{\partial g}{\partial s}(s,r)$ and $\frac{\partial^2 g}{\partial s\partial r}(s,r)$ integrable on $(0,\infty)\times[0,\infty)$. Then, for every $t\geq s\geq 0$,

$$\lambda_{g}(t,s) := g(t,t)e^{-2\theta t} + g(s,s)e^{-2\theta s} - 2g(s,t)e^{-2\theta(s+t)} + 2\theta e^{-\theta t} \int_{s}^{t} g(r,t)e^{-\theta r} dr$$

$$-2\theta e^{-\theta s} \int_{s}^{t} g(r,s)e^{-\theta r} dr + \theta^{2} \int_{s}^{t} \int_{s}^{t} g(r,u)e^{-\theta(r+u)} dr du$$

$$= \int_{s}^{t} \int_{s}^{t} e^{-\theta(r+u)} \frac{\partial^{2} g}{\partial r \partial u}(r,u) dr du.$$
(6.3)

Let us now recall the Young integral introduced in [21]. For any $\alpha \in (0, 1]$, we denote by $\mathcal{H}^{\alpha}([0, T])$ the set of α -Hölder continuous functions, that is, the set of functions $f:[0, T] \to \mathbb{R}$ such that

$$|f|_{\alpha} := \sup_{0 < s < t < T} \frac{|f(t) - f(s)|}{(t - s)^{\alpha}} < \infty.$$

We also set $|f|_{\infty} = \sup_{t \in [0,T]} |f(t)|$, and we equip $\mathcal{H}^{\alpha}([0,T])$ with the norm $||f||_{\alpha} := |f|_{\alpha} + |f|_{\infty}$.

Let $f \in \mathcal{H}^{\alpha}([0,T])$, and consider the operator $T_f : \mathcal{C}^1([0,T]) \to \mathcal{C}^0([0,T])$ defined as

$$T_f(g)(t) = \int_0^t f(u)g'(u)du, \quad t \in [0, T].$$

It can be shown (see, e.g., [16, Section 3.1]) that, for any $\beta \in (1 - \alpha, 1)$, there exists a constant $C_{\alpha,\beta,T} > 0$ depending only on α , β and T such that, for any $g \in \mathcal{H}^{\beta}([0,T])$,

$$\left\| \int_0^{\cdot} f(u)g'(u)du \right\|_{\beta} \le C_{\alpha,\beta,T} \|f\|_{\alpha} \|g\|_{\beta}.$$

We deduce that, for any $\alpha \in (0,1)$, any $f \in \mathcal{H}^{\alpha}([0,T])$ and any $\beta \in (1-\alpha,1)$, the linear operator $T_f : \mathcal{C}^1([0,T]) \subset \mathcal{H}^{\beta}([0,T]) \to \mathcal{H}^{\beta}([0,T])$, defined as $T_f(g) = \int_0^{\cdot} f(u)g'(u)du$, is continuous with respect to the norm $\|\cdot\|_{\beta}$. By density, it extends (in an unique way) to an operator defined on \mathcal{H}^{β} . As consequence, if $f \in \mathcal{H}^{\alpha}([0,T])$, if $g \in \mathcal{H}^{\beta}([0,T])$ and if $\alpha + \beta > 1$, then the (so-called) Young integral $\int_0^{\cdot} f(u)dg(u)$ is well-defined as being $T_f(g)$ (see [21]).

The Young integral obeys the following formula. Let $f \in \mathcal{H}^{\alpha}([0,T])$ with $\alpha \in (0,1)$ and $g \in \mathcal{H}^{\beta}([0,T])$ with $\beta \in (0,1)$ such that $\alpha + \beta > 1$. Then $\int_0^{\cdot} g_u df_u$ and $\int_0^{\cdot} f_u dg_u$ are well-defined as the Young integrals. Moreover, for all $t \in [0,T]$,

$$f_t g_t = f_0 g_0 + \int_0^t g_u df_u + \int_0^t f_u dg_u.$$
 (6.4)

References

- [1] Basawa, I. V. and Scott, D. J. (1983). Asymptotic Optimal Inference for Non-Ergodic Models. Lecture Notes in Statist. 17. New York: Springer.
- [2] Belfadli, R. Es-Sebaiy, K. and Ouknine, Y. (2011). Parameter Estimation for Fractional Ornstein-Uhlenbeck Processes: Non-Ergodic Case. Frontiers in Science and Engineering (An International Journal Edited by Hassan II Academy of Science and Technology) 1(1), 1-16.
- [3] Bojdecki, T., Gorostiza, L. and Talarczyk, A. (2007). Some extensions of fractional Brownian motion and sub-fractional Brownian motion related to particle systems. Electronic Communications in Probability, 12, 161-172.

- [4] Cheng, P., Shen, G. and Chen, Q. (2017). Parameter estimation for nonergodic Ornstein-Uhlenbeck process driven by the weighted fractional Brownian motion. Advances in Difference Equations, 2017:366. https://doi.org/10.1186/s13662-017-1420-y
- [5] Dietz, H. M. and Kutoyants, Y. A. (2003). Parameter estimation for some non-recurrent solutions of SDE. Statistics and Decisions 21, 29-46.
- [6] Douissi, S., Es-Sebaiy, K. and Viens, F. (2019). Berry-Esséen bounds for parameter estimation of general Gaussian processes. ALEA, Lat. Am. J. Probab. Math. Stat. 16, 633-664.
- [7] El Machkouri, M., Es-Sebaiy, K. and Ouknine, Y. (2016). Least squares estimator for non-ergodic Ornstein-Uhlenbeck processes driven by Gaussian processes. Journal of the Korean Statistical Society 45, 329-341.
- [8] El Onsy, B., Es-Sebaiy, K. and Viens, F. (2017). Parameter Estimation for a partially observed Ornstein-Uhlenbeck process with long-memory noise. Stochastics, 89(2), 431-468.
- [9] Es-Sebaiy, K., Alazemi, F. and Al-Foraih, M. (2019). Least squares type estimation for discretely observed non-ergodic Gaussian Ornstein-Uhlenbeck processes. Acta Mathematica Scientia, 39(4), 989-1002.
- [10] Es-Sebaiy, K. Nourdin, I. (2013). Es-Sebaiy K. and Nourdin I. (2013). Parameter Estimation for α-Fractional Bridges. In: Viens F., Feng J., Hu Y., Nualart E. (eds) Malliavin Calculus and Stochastic Analysis. Springer Proceedings in Mathematics and Statistics, 34. Springer, Boston, MA.
- [11] Es-Sebaiy, K. and Viens, F. (2019). Optimal rates for parameter estimation of stationary Gaussian processes. Stochastic Processes and their Applications, 129(9), 3018-3054.
- [12] Hu, Y. and Nualart, D. (2010). Parameter estimation for fractional Ornstein-Uhlenbeck processes. Statist. Probab. Lett. 80, 1030-1038.
- [13] Hu, Y., Nualart, D. and Zhou, H. (2019). Parameter estimation for fractional Ornstein-Uhlenbeck processes of general Hurst parameter. Statistical Inference for Stochastic Processes, 22(1), 111-142.
- [14] Jacod, J. (2006). Parametric inference for discretely observed non-ergodic diffusions. Bernoulli 12, 383-401.
- [15] Marsaglia, G. (1965). Ratios of normal variables and ratios of sums of uniform variables. J. Amer. Statist. Asso. 60: 193-204.
- [16] Nourdin, I. (2012). Selected aspects of fractional Brownian motion. Bocconi & Springer Series 4. Springer, Milan; Bocconi University Press, Milan.

- [17] Pham-Gia, T., Turkkan, N., Marchand, E. (2006). Density of the ratio of two normal random variables and applications. Communications in Statistics-Theory and Methods, 35(9), 1569-1591.
- [18] Shen, G., Yin, X., Yan, L. (2016). Least squares estimation for Ornstein-Uhlenbeck processes driven by the weighted fractional Brownian motion. Acta Math Sci, 36B(2): 394-408.
- [19] Shen, G., Yin, X., Yan, L. (2017). Erratum to: least squares estimation for Ornstein-Uhlenbeck processes driven by the weighted fractional brownian motion (Acta Mathematica Scientia 2016, 36B(2): 394-408. Acta Mathematica Scientia, 37(4), 1173-1176.
- [20] Shimizu, Y. (2009). Notes on drift estimation for certain non-recurrent diffusion from sampled data. Statistics and Probability Letters 79, 2200-2207.
- [21] Young, L. C. (1936). An inequality of the Hölder type connected with Stieltjes integration. Acta Math. 67, 251-282.