

Well-partitioned chordal graphs: obstruction set and disjoint paths*

Jungho Ahn^{1,2}, Lars Jaffke³, O-joung Kwon^{4,2}, and Paloma T. Lima³

¹Department of Mathematical Sciences, KAIST, Daejeon, South Korea

²Discrete Mathematics Group, Institute for Basic Science (IBS), Daejeon, South Korea

³Department of Informatics, University of Bergen, Bergen, Norway

⁴Department of Mathematics, Incheon National University, Incheon, South Korea

Email addresses: junghoahn@kaist.ac.kr, lars.jaffke@uib.no, ojoungkwon@gmail.com,
Paloma.Lima@uib.no

February 26, 2020

Abstract

We introduce a new subclass of chordal graphs that generalizes split graphs, which we call well-partitioned chordal graphs. Split graphs are graphs that admit a partition of the vertex set into cliques that can be arranged in a star structure, the leaves of which are of size one. Well-partitioned chordal graphs are a generalization of this concept in the following two ways. First, the cliques in the partition can be arranged in a tree structure, and second, each clique is of arbitrary size. We provide a characterization of well-partitioned chordal graphs by forbidden induced subgraphs, and give a polynomial-time algorithm that given any graph, either finds an obstruction, or outputs a partition of its vertex set that asserts that the graph is well-partitioned chordal. We demonstrate the algorithmic use of this graph class by showing that two variants of the problem of finding pairwise disjoint paths between k given pairs of vertices is in FPT parameterized by k on well-partitioned chordal graphs, while on chordal graphs, these problems are only known to be in XP. From the other end, we observe that there are problems that are polynomial-time solvable on split graphs, but become NP-complete on well-partitioned chordal graphs.

1 Introduction

A central methodology in the study of the complexity of computationally hard graph problems is to impose additional structure on the input graphs, and determine if the additional structure can be exploited in the design of an efficient algorithm. Typically, one restricts the input to be contained in a *graph class*, which is a set of graphs that share a common structural property. For example, the class of *forests* is the class of graphs that do not contain a cycle. Following the establishment of the theory of NP-hardness, numerous problems were investigated in specific classes of graphs; either providing a polynomial-time algorithm for a problem Π on a specific graph class, while Π

*J. A. and O. K. are supported by IBS-R029-C1. O. K. is also supported by the National Research Foundation of Korea (NRF) grant funded by the Ministry of Education (No. NRF-2018R1D1A1B07050294). L. J. is supported by the Bergen Research Foundation (BFS), grant number 810564. P. T. L. is supported by the Norwegian Research Council via the project “CLASSIS”, grant number 249994.




Figure 1: A well-partitioned chordal graph.

is NP-hard in a more general setting, or showing that Π remains NP-hard on a graph class. We refer to the textbooks [6, 15] for a detailed introduction to the subject. A key question in this field is to find for a given problem Π that is hard on a graph class \mathcal{A} , a subclass $\mathcal{B} \subsetneq \mathcal{A}$ such that Π is efficiently solvable on \mathcal{B} . Naturally, the goal is to narrow down the gap $\mathcal{A} \setminus \mathcal{B}$ as much as possible, and several notions of hardness/efficiency can be applied. For instance, we can require our target problem to be NP-hard on \mathcal{A} and polynomial-time solvable on \mathcal{B} ; or, from the viewpoint of parameterized complexity [9, 10], we require a target parameterized problem Π to be W[1]-hard on \mathcal{A} , while Π is in FPT on \mathcal{B} , or a separation in the kernelization complexity [12] of Π between \mathcal{A} and \mathcal{B} .

Chordal graphs are arguably one of the main characters in the algorithmic study of graph classes. They find applications for instance in computational biology [28], optimization [30], and sparse matrix computations [14]. Split graphs are an important subclass of chordal graphs. The complexities of computational problems on chordal and split graphs often coincide, see e.g. [2, 3, 11, 24]; however, this is not always the case. For instance, several variants of graph (vertex) coloring problems are polynomial-time solvable on split graphs and NP-hard on chordal graphs, see the works of Havet et al. [17], and of Silva [29]. Also, the SPAREST k -SUBGRAPH [32] and DENSEST k -SUBGRAPH [8] problems are polynomial-time solvable on split graphs and NP-hard on chordal graphs. Other problems, for instance the TREE 3-SPANNER problem [5], are easy on split graphs, while their complexity on chordal graphs is still unresolved.

In this work, we introduce the class of *well-partitioned chordal graphs*, a subclass of chordal graphs that generalizes split graphs, which can be used as a tool for narrowing down complexity gaps for problems that are hard on chordal graphs, and easy on split graphs. The definition of well-partitioned chordal graphs is mainly motivated by a property of split graphs: the vertex set of a split graph can be partitioned into sets that can be viewed as a central clique of arbitrary size and cliques of size one that have neighbors only in the central clique. Thus, this partition has the structure of a star. Well-partitioned chordal graphs relax these ideas in two ways: by allowing the parts of the partition to be arranged in a tree structure instead of a star, and by allowing the cliques in each part to have arbitrary size. The interaction between adjacent parts P and Q remains simple: it induces a complete bipartite graph between a subset of P , and a subset of Q . Such a tree structure is called a partition tree, and we give an example of a well-partitioned chordal graph in Figure 1. We formally define this class in Section 3.

That said, it is not difficult to observe that the graphs constructed in the NP-hardness proofs in the works [17, 29] are in fact well-partitioned chordal graphs — we immediately narrowed down the complexity gaps of these problems from CHORDAL \setminus SPLIT to WELL-PARTITIONED CHORDAL \setminus SPLIT.




Figure 2: The set of obstructions \mathbb{O} for well-partitioned chordal graphs.

The main structural contribution of this work is a characterization of well-partitioned chordal graphs by forbidden induced subgraphs. We also provide a polynomial-time recognition algorithm. We list the set \mathbb{O} of obstructions in Figure 2.

Theorem 1.1. *A graph is a well-partitioned chordal graph if and only if it has no induced subgraph isomorphic to a graph in \mathbb{O} . Furthermore, there is a polynomial-time algorithm that given a graph G , output either an induced subgraph of G isomorphic to a graph in \mathbb{O} , or a partition tree for each connected component which confirms that G is a well-partitioned chordal graph.*

Before we proceed with the discussion of the algorithmic results of this paper, we would like to briefly touch on the relationship of well-partitioned chordal graphs and width parameters. Each split graph is a well-partitioned chordal graph, and there are split graphs of whose *maximum induced matching width* (mim-width) depends linearly on the number of vertices [23]. This rules out the applicability of any algorithmic meta-theorem based on one of the common width parameters such as tree-width or clique-width, to the class of well-partitioned chordal graphs. It is known that mim-width is a lower bound for them [31].

Besides narrowing the complexity gap between the classes of chordal and split graphs, the class of well-partitioned chordal graphs can also be useful as a step towards determining the yet unresolved complexity of a problem Π on chordal graphs when it is known that Π is easy on split graphs. This is the case in our current work. Specifically, we study the the DISJOINT PATHS problem, formally defined as follows, and generalizations thereof. Two paths P_1 and P_2 are called *internally vertex-disjoint*, if for $i \in [2]$, no internal vertex of P_i is contained in P_{3-i} . (Note that this excludes the possibility that an endpoint of one path is used as an internal vertex in the other path.)

Graph Class	DISJOINT PATHS	SET-RESTRICTED DISJOINT PATHS
Chordal	linear FPT [20]	XP [1]
Well-partitioned chordal	$\mathcal{O}(k^3)$ kernel [T. 5.15, 5.16]	linear* FPT [T. 5.6, 5.9]
Split	$\mathcal{O}(k^2)$ kernel [18]	$\mathcal{O}(k^2)$ kernel [C. 5.11]

Table 1: Summary of some results about the parameterized complexity results of the DISJOINT PATHS and SET-RESTRICTED DISJOINT PATHS problems parameterized by the number k of terminal pairs on split, well-partitioned chordal, and chordal graphs. Size bounds for kernels are in terms of the number of *vertices* of the kernelized instances. *Assuming that we are given a partition tree.

DISJOINT PATHS

Input: A graph G , a set $\mathcal{X} = \{(s_1, t_1), \dots, (s_k, t_k)\}$ of k pairs of vertices of G , called *terminals*.

Question: Does G contain k pairwise internally vertex-disjoint paths P_1, \dots, P_k such that for all $i \in [k]$, P_i is an (s_i, t_i) -path?

This problem has already been shown by Karp to be NP-complete [21], and as a cornerstone result in the early days of fixed-parameter tractability theory, Robertson and Seymour showed that DISJOINT PATHS parameterized by k is in FPT [26]. The dependence of the runtime on the number of vertices n in the input graph is cubic in Robertson and Seymour’s algorithm, and Kawarabayashi et al. improved this to a quadratic dependence on n [22]. From the viewpoint of kernelization complexity, Bodlaender et al. showed that DISJOINT PATHS does not admit a polynomial kernel unless $\text{NP} \subseteq \text{coNP}/\text{poly}$ [4].

Restricting the problem to chordal and split graphs, Heggenes et al. showed that DISJOINT PATHS remains NP-complete on split graphs, and that it admits a polynomial kernel parameterized by k [18], and Kammer and Tholey showed that it has an FPT-algorithm with linear dependence on the size of the input chordal graph [20]. The question whether DISJOINT PATHS has a polynomial kernel on chordal graphs remains open. We go one step towards such a polynomial kernel, by showing that DISJOINT PATHS has a polynomial kernel on well-partitioned chordal graphs; generalizing the polynomial kernel on split graphs [18].

We also study a generalization of the DISJOINT PATHS problem, where in a solution, each path P_i can only use a restricted set of vertices U_i , which is specified for each terminal pair at the input. This problem was recently introduced by Belmonte et al. and given the name SET-RESTRICTED DISJOINT PATHS [1]. Since this problem contains DISJOINT PATHS as a special case (setting all domains equal to the whole vertex set), it is NP-complete. Belmonte et al. showed that SET-RESTRICTED DISJOINT PATHS parameterized by k is in XP on chordal graphs, and leave as an open question whether it is in FPT or W[1]-hard on chordal graphs. Towards showing the former, we give an FPT-algorithm on well-partitioned chordal graphs. In particular, given a partition tree, our algorithm runs in time $2^{\mathcal{O}(k \log k)} \cdot n$, so the runtime only depends linearly on the number of vertices in the input graph. While we do not settle the kernelization complexity of SET-RESTRICTED DISJOINT PATHS on well-partitioned chordal graphs, we observe that our FPT-algorithm implies a polynomial kernel on split graphs. We summarize these results in Table 1.

Finally, we also consider the SET-RESTRICTED DISJOINT CONNECTED SUBGRAPHS problem where we are given k terminal *sets* instead of pairs, and k domains, and the question is whether there are k pairwise disjoint connected subgraphs, each one connecting one of the terminal sets,

using only vertices from the specified domain. This problem was also introduced in [1] and shown to be in XP on chordal graphs, when the parameter is the total number of vertices in all terminal sets. We observe that the ideas we use in our algorithm for the path problems can be used to show that SET-RESTRICTED DISJOINT CONNECTED SUBGRAPHS on well-partitioned chordal graphs is in FPT with this parameter.

The case of coinciding terminal pairs. We want to point out that the DISJOINT PATHS is studied in several variants in the literature. According to the definition that we give here, pairs of terminal vertices may coincide, i.e. it may happen that for $i \neq j$, $\{s_i, t_i\} = \{s_j, t_j\} =: \{x, y\}$. If the edge xy is present in the input graph, then it may be used both as the path P_i and as the path P_j in the solution, without violating the definition. However, as pointed out in e.g. [18], it is natural to impose the additional condition that all paths in a solution have to be pairwise distinct. We study both variants of the (SET-RESTRICTED) DISJOINT PATHS problem, and refer to the variant that requires pairwise distinct paths in a solution as (SET-RESTRICTED) TOTALLY DISJOINT PATHS.

2 Preliminaries

For a positive integer n , we let $[n] := \{1, 2, \dots, n\}$. For a set X and an integer k , we denote by $\binom{X}{k}$ the size- k subsets of X .

A graph G is a pair of a *vertex set* $V(G)$ and an *edge set* $E(G) \subseteq \binom{V(G)}{2}$. All graphs considered in this paper are finite, i.e. their vertex sets are finite. For an edge $\{u, v\} \in E(G)$, we call u and v its *endpoints* and we use the shorthand ‘ uv ’ for ‘ $\{u, v\}$ ’. Let G and H be two graphs. We say that G is *isomorphic to* H if there is a bijection $\phi: V(G) \rightarrow V(H)$ such that for all $u, v \in V(G)$, $uv \in E(G)$ if and only if $\phi(u)\phi(v) \in E(H)$. We say that H is a *subgraph* of G , denoted by $H \subseteq G$, if $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$.

For a vertex v of a graph G , $N_G(v) := \{w \in V(G) \mid vw \in E(G)\}$ is the set of *neighbors* of v in G , and we let $N_G[v] := N_G(v) \cup \{v\}$. The *degree* of v is $\deg_G(v) := |N_G(v)|$. Given a set $X \subseteq V(G)$, we let $N_G(X) := \bigcup_{v \in X} N_G(v) \setminus X$ and $N_G[X] := N_G(X) \cup X$. In all of the above, we may drop G as a subscript if it is clear from the context. The *subgraph induced by* X , denoted by $G[X]$, is the graph $(X, E(G) \cap \binom{X}{2})$. We denote by $G - X$ the graph $G[V(G) \setminus X]$, and for a single vertex $x \in V(G)$, we use the shorthand ‘ $G - x$ ’ for ‘ $G - \{x\}$ ’. For two sets $X, Y \subseteq V(G)$, we denote by $G[X, Y]$ the graph $(X \cup Y, \{xy \in E(G) \mid x \in X, y \in Y\})$. We say that X is *complete to* Y if $X \cap Y = \emptyset$ and each vertex in X is adjacent to every vertex in Y .

Let G be a graph. We say that G is *trivial* if $|V(G)| = 1$. G is called *complete* if $E(G) = \binom{V(G)}{2}$, and *empty* if $E(G) = \emptyset$. A set $X \subseteq V(G)$ is a *clique* if $G[X]$ is complete, and an *independent set* if $G[X]$ is empty. A clique of size 3 is called a *triangle*. A graph G is called *bipartite* there is a 2-partition (A, B) of $V(G)$, called the *bipartition of* G , such that A and B are independent sets in G . A bipartite graph G on bipartition (A, B) is called *complete bipartite* if A is complete to B . For integers n and m , we denote by $K_{n,m}$ a complete bipartite graph with bipartition (A, B) such that $|A| = n$ and $|B| = m$. A graph is a *star* if it is either trivial or isomorphic to $K_{1,n}$ for some positive integer n .

A graph G is *connected* if for each 2-partition (X, Y) of $V(G)$ with $X \neq \emptyset$ and $Y \neq \emptyset$, there is a pair $x \in X, y \in Y$ such that $xy \in E(G)$. A connected component of G is a maximal connected subgraph of G . A vertex $v \in V(G)$ is a *cut vertex* if $G - v$ has more connected components than G . A graph is *2-connected* if it has no cut vertices. A *block* of a graph G is a maximal 2-connected component of G . A graph G is called *2-regular* if all vertices of G are of degree 2. A connected


2-regular graph is a cycle. A graph that has no cycle as a subgraph is called a *forest*, a connected forest is a *tree*, and a tree of maximum degree 2 is a *path*. The vertices of degree one in a tree are called *leaves* and the leaves of a path are its *endpoints*. A connected subgraph of a tree is called a *subtree*.

A *hole* in a graph G is an induced cycle of G of length at least 4. A graph is *chordal* if it has no induced subgraph isomorphic to a hole. A vertex is *simplicial* if $N_G(v)$ is a clique. We say that a graph G has a *perfect elimination ordering* v_1, \dots, v_n if v_i is simplicial in $G[\{v_i, v_{i+1}, \dots, v_n\}]$ for each $i \in [n - 1]$. It is known that a graph is chordal if and only if it has a perfect elimination ordering [13]. We will use the following hole detecting algorithm and an algorithm to generate a perfect elimination ordering of a chordal graph.

Theorem 2.1 (Nikolopoulos and Palios [25]). *Given a graph G , one can detect a hole in G in time $\mathcal{O}(|V(G)| + |E(G)|^2)$, if one exists.*

Theorem 2.2 (Rose, Tarjan, and Lueker [27]). *Given a graph G , one can generate a perfect elimination ordering of G in time $\mathcal{O}(|V(G)| + |E(G)|)$, if one exists.*

A graph G is a *split graph* if there is a 2-partition (C, I) of $V(G)$ such that C is a clique and I is an independent set. Let \mathcal{S} be a family of subsets of some set. The *intersection graph of \mathcal{S}* is the graph on vertex set \mathcal{S} and edge set $\{\{S, T\} \in \binom{\mathcal{S}}{2} \mid S \cap T \neq \emptyset\}$. It is well-known that each chordal graph is the intersection graph of vertex sets of subtrees of some tree. The following graph is called a *diamond*. Note that for all $s \in [3]$, $t \in \mathbb{N}$, the graph $W_{s,t}$ in \mathbb{O} (see Figure 2) contains two diamonds as induced subgraphs.



3 Well-partitioned chordal graphs

A connected graph G is a *well-partitioned chordal graph* if there exist a partition \mathcal{P} of $V(G)$ and a tree \mathcal{T} having \mathcal{P} as a vertex set such that the following hold.

- (i) Each part $X \in \mathcal{P}$ is a clique in G .
- (ii) For each edge $XY \in E(\mathcal{T})$, there are subsets $X' \subseteq X$ and $Y' \subseteq Y$ such that $E(G[X, Y]) = X' \times Y'$.
- (iii) For each pair of distinct $X, Y \in V(\mathcal{T})$ with $XY \notin E(\mathcal{T})$, $E(G[X, Y]) = \emptyset$.

The tree \mathcal{T} is called a *partition tree of G* , and the elements of \mathcal{P} are called its *bags*. A graph is a well-partitioned chordal graph if all of its connected components are well-partitioned chordal graphs. We remark that a well-partitioned chordal graph can have more than one partition tree. Also, observe that well-partitioned chordal graphs are closed under taking induced subgraphs.

A useful concept when considering partition trees of well-partitioned chordal graphs is that of a *boundary of a bag*. Let \mathcal{T} be a partition tree of a well-partitioned chordal graph G and let

$X, Y \in V(\mathcal{T})$ be two bags that are adjacent in \mathcal{T} . The *boundary of X with respect to Y* , denoted by $\text{bd}(X, Y)$, is the set of vertices of X that have a neighbor in Y , i.e.

$$\text{bd}(X, Y) := \{x \in X \mid N_G(x) \cap Y \neq \emptyset\}.$$

By item (ii) of the definition of the class, we know that $\text{bd}(X, Y)$ is complete to $\text{bd}(Y, X)$.

We now consider the relation between well-partitioned chordal graphs and other well-studied classes of graphs. It is easy to see that every well-partitioned chordal graph G is a chordal graph because every leaf of the partition tree of a component of G contains a simplicial vertex of G , and after removing this vertex, the remaining graph is still a well-partitioned chordal graph. Thus, we may construct a perfect elimination ordering. We show that, in fact, well-partitioned chordal graphs constitute a subclass of substar graphs. A graph is a *substar graph* [7, 19] if it is an intersection graph of substars of a tree.

Proposition 3.1. *Every well-partitioned chordal graph is a substar graph.*

Proof. Let G be a well-partitioned chordal graph with $V(G) = \{v_1, v_2, \dots, v_n\}$ and a partition tree \mathcal{T} . We will exhibit a substar intersection model for G . That is, we will show that there exists a tree \mathcal{T}' and S_1, \dots, S_n substars of \mathcal{T}' such that $v_i v_j \in E(G)$ if and only if $V(S_i) \cap V(S_j) \neq \emptyset$.

Let \mathcal{T}' be the tree obtained from \mathcal{T} by the 1-subdivision of every edge. We denote by $v_{XY} \in V(\mathcal{T}')$ the vertex originated from the 1-subdivision of the edge $XY \in E(\mathcal{T})$. Note that $N_{\mathcal{T}'}(v_{XY}) = \{X, Y\}$. For every $v_i \in V(G)$, we create a substar of \mathcal{T}' in the following way. Let $B \in V(\mathcal{T})$ be the bag containing v_i . Then S_i is a star with the center B and the leaf set $\{v_{BY} \mid v_i \in \text{bd}(B, Y)\}$.

To see that this is indeed an intersection model for G , let $v_i v_j \in E(G)$. If there exists $B \in V(\mathcal{T})$ such that $v_i, v_j \in B$, then $B \in V(S_i) \cap V(S_j)$. If v_i and v_j are not contained in the same bag, by item (ii), there exist $A, B \in V(\mathcal{T})$ such that $v_i \in A$, $v_j \in B$ and $AB \in E_{\mathcal{T}}$. Then, $v_{AB} \in V(S_i) \cap V(S_j)$. In both cases we have that $V(S_i) \cap V(S_j) \neq \emptyset$. Now suppose $V(S_i) \cap V(S_j) = \emptyset$. Note that, by construction, two stars that intersect either have the same center or they intersect in a vertex that is a leaf of both of them. If S_i and S_j have the same center B , then $v_i, v_j \in B$ and hence, by item (i), $v_i v_j \in E(G)$. If S_i and S_j have a common leaf, then this leaf is a vertex originated by the 1-subdivision of an edge. Then, there exist $A, B \in V(\mathcal{T})$ such that $v_i \in \text{bd}(A, B)$ and $v_j \in \text{bd}(B, A)$ and thus, by item (ii), $v_i v_j \in E(G)$. \square

From the definition of well-partitioned chordal graphs, one can also see that every split graph is a well-partitioned chordal graph. Indeed, if G is a split graph with clique K and independent set S , the partition tree of G will be a star, with the clique K as its central bag and each vertex of S contained in a different leaf bag. We show that, in fact, every starlike graph is a well-partitioned chordal graph. A *starlike graph* [16] is an intersection graph of substars of a star.

Proposition 3.2. *Every starlike graph is a well-partitioned chordal graph.*

Proof. Let G be a starlike graph with $V(G) = \{v_1, \dots, v_n\}$ and let \mathcal{S} be the host star of the substar intersection model of G and S_i be the substar of \mathcal{S} associated with vertex v_i . We may assume that G is connected and every vertex of \mathcal{S} is contained in some substar of the intersection model.

We know that $v_i v_j \in E(G)$ if and only if $V(S_i) \cap V(S_j) \neq \emptyset$. To show that G is a well-partitioned chordal graph, we will construct a partition tree for G . Let c be the center of the star \mathcal{S} and f_1, \dots, f_k be its leaves. The partition tree \mathcal{T} for G will be a star with center C and leaves

F_1, \dots, F_k such that $C = \{v_i \in V(G) \mid c \in S_i\}$ and $F_j = \{v_i \in V(G) \mid V(S_i) = \{f_j\}\}$. Note that this is indeed a partition of the vertex set of G , since each substar of \mathcal{S} either contains the center or consists of a single leaf and every vertex of \mathcal{S} is contained in some substar of the intersection model. Now we show this is indeed a partition tree for G . Note that, by construction, each bag is a clique, so item (i) holds. Also note that, for every i , if $v \in F_i$, then $N_G(v) \subseteq F_i \cup C$, thus item (iii) of the definition holds. Finally, note that the vertices of F_i are true twins in G , since the substars of \mathcal{S} corresponding to those vertices consist of a single vertex, namely f_i . Hence, item (ii) also holds. This concludes the proof that \mathcal{T} is a partition tree for G and thus G is a well-partitioned chordal graph. \square

We will show that the graph O_1 in Figure 2 is not a well-partitioned chordal graph. On the other hand, it is not difficult to see that O_1 is a substar graph. Also note that a path graph on 5 vertices is a well-partitioned chordal graph but not a starlike graph. These observations with Propositions 3.1 and 3.2 show that we have the following hierarchy of graph classes between split graphs and chordal graphs:

$$\begin{array}{ccccccc} \text{split} & & \text{starlike} & & \text{well-partitioned} & & \text{substar} & & \text{chordal} \\ \text{graphs} & \subsetneq & \text{graphs} & \subsetneq & \text{chordal graphs} & \subsetneq & \text{graphs} & \subsetneq & \text{graphs} \end{array}$$

4 Characterization by forbidden induced subgraphs

This section is entirely devoted to the proof of Theorem 1.1. That is, we show that the set \mathbb{O} of graphs depicted in Figure 2 is the set of all forbidden induced subgraphs for well-partitioned chordal graphs, and give a polynomial-time recognition algorithm for this graph class. For convenience, we say that an induced subgraph of a graph that is isomorphic to a graph in \mathbb{O} is an *obstruction* for well-partitioned chordal graphs, or simply an obstruction.

In Subsection 4.1, we show that the graphs in \mathbb{O} are not well-partitioned chordal graphs (Proposition 4.2). In Subsection 4.2, we introduce the notion of a boundary-crossing path which is the main tool for devising the polynomial-time recognition algorithm. We present the certifying algorithm in Subsection 4.3, which also concludes the proof of the characterization by forbidden induced subgraphs for well-partitioned chordal graphs.

It is not difficult to observe that no graph in \mathbb{O} contains another graph in \mathbb{O} as an induced subgraph. Thus, the results in this section also implies that graphs in \mathbb{O} are minimal graphs with respect to the induced subgraph relation that are not well-partitioned chordal graphs.

4.1 Graphs in \mathbb{O} are not well-partitioned chordal graphs

To argue that none of the graphs in \mathbb{O} is a well-partitioned chordal graph, we make the following observation about triangles, which follows immediately from the definition of the partition tree.

Observation 4.1. Let G be a connected well-partitioned chordal graph, and $D \subseteq V(G)$ be the vertex set of a triangle in G . In any partition tree \mathcal{T} of G , there are at most two bags whose intersection with D is non-empty.

Given a connected well-partitioned chordal graph G and a triangle with vertex set $D \subseteq V(G)$, we say that a partition tree of G *respects* D if it contains a bag that contains all the vertices of D .

For a non-empty proper subset $D' \subset D$, we say that a partition tree \mathcal{T} splits D into $(D', D \setminus D')$ if \mathcal{T} contains two distinct bags B_1 and B_2 such that $B_1 \cap D = D'$ and $B_2 \cap D = D \setminus D'$. If a partition tree splits D into $(D', D \setminus D')$ for some $D' \subset D$, then we may simply say that it splits D . By Observation 4.1, each partition tree either respects or splits each triangle.

For $s \in \{1, 2, 3\}$ and $t \geq 0$, the vertex set of a block of $W_{s,t}$ having more than 3 vertices is called a *wing* of $W_{s,t}$.

Proposition 4.2. *The graphs in \mathbb{O} are not well-partitioned chordal graphs.*

Proof. For $k \geq 4$, H_k is not a chordal graph, so it is not a well-partitioned chordal graph.

We prove an auxiliary claim that will be useful to show that the graphs O_1, O_2, O_3 , and O_4 in \mathbb{O} are not well-partitioned chordal graphs.

Claim 4.2.1. *Let H be a connected graph and $D = \{x, y, z\} \subseteq V(H)$ be a triangle in H .*

- (i) *If there are adjacent vertices $u, v \in V(H) \setminus D$ such that $D \not\subseteq N_H(u)$ and $D \not\subseteq N_H(v)$, and $\emptyset \neq N_H(u) \cap D \neq N_H(v) \cap D \neq \emptyset$, then H has no partition tree respecting D .*
- (ii) *If there exists a vertex $u \in V(H) \setminus D$ such that $N_H(u) \cap D = \{y, z\}$, then H has no partition tree splitting D into $(\{x, y\}, \{z\})$.*
- (iii) *If there exist two non-adjacent vertices $u, v \in V(H) \setminus D$ such that $N_H(u) \cap D = D = N_H(v) \cap D$, then H has no partition tree splitting D .*

Proof. In order to prove item (i), suppose there is a partition tree \mathcal{T} of H respecting D , and let B be the bag containing D . First, since $D \not\subseteq N_H(u)$ and $D \not\subseteq N_H(v)$, we have that neither v nor u is contained in B as B is a clique in H . Furthermore, since $N_H(u) \cap D \neq \emptyset$ and $N_H(v) \cap D \neq \emptyset$, and since $uv \in E(G)$, it cannot be the case that u and v are in distinct bags, otherwise there would be a triangle in \mathcal{T} . However, since $N_H(u) \cap D \neq N_H(v) \cap D$, u and v cannot be in the same bag either.

Now we proceed to the proof of item (ii). Suppose there is a partition tree \mathcal{T} of H that splits D into $(\{x, y\}, \{z\})$, and denote the two bags intersecting D by B_1 and B_2 with $B_1 \cap D = \{x, y\}$ and $B_2 \cap D = \{z\}$. Since u is not adjacent to x , $u \notin B_1$. Since $x \in N_H(z) \cap B_1$ and $x \notin N_H(u) \cap B_1$, u cannot be contained in B_2 either. However, since $uz, uy \in E(G)$, if u is in a bag other than B_1 and B_2 , then $\{u, y, z\}$ is a triangle that intersects three distinct bags of \mathcal{T} , a contradiction with Observation 4.1.

To conclude, we prove item (iii). Suppose there is a partition tree \mathcal{T} of H that splits D , and again denote the two bags intersecting D by B_1 and B_2 , with $B_1 \cap D = \{x, y\}$ and $B_2 \cap D = \{z\}$. First, since u and v are non-adjacent, they cannot be in the same bag. Furthermore, there cannot be a bag $B_3 \in V(\mathcal{T}) \setminus \{B_1, B_2\}$ such that $\{u, v\} \cap B_3 \neq \emptyset$: both u and v have neighbors in B_1 and in B_2 , so this would imply the existence of a triangle that intersects three distinct bags of \mathcal{T} (B_1, B_2 , and B_3). The last case that remains is when $u \in B_1$ and $v \in B_2$. However, in this case, B_1 contains a vertex that is adjacent to v , namely x , and a vertex that is not adjacent to v , namely u , a contradiction. \square

Now, let us consider the obstructions O_1, O_2, O_3 and O_4 and assume that their vertices are labelled as in Figure 3.

By Observation 4.1, each partition tree either respects or splits a triangle. First, consider the graph O_1 and consider the triangle $D = \{a, c, d\}$. Because of the vertices e and f , we can observe




Figure 3: Labellings of graphs O_1, O_2, O_3 , and O_4 .

that, by Claim 4.2.1(i), no partition tree of O_1 respects D . Furthermore, because of the vertices b , f , and b , we obtain by Claim 4.2.1(ii) that no partition tree splits D into $(\{a, c\}, \{d\})$, $(\{a, d\}, \{c\})$, and $(\{c, d\}, \{a\})$, respectively. Thus, no partition tree of O_1 splits D . Hence, O_1 does not admit a partition tree and therefore it is not a well-partitioned chordal graph.

For O_2 , consider again the triangle $\{a, c, d\}$. The arguments are similar to the previous ones, except that the vertex e should be used to show that no partition tree splits $\{a, c, d\}$ into $(\{a, d\}, \{c\})$.

For O_3 , consider the triangle $D = \{b, d, e\}$. Because of the vertices f and g , we observe that, by Claim 4.2.1(i), no partition tree of O_3 respects D . On the other hand, because of a and c , we observe that by Claim 4.2.1(iii), no partition tree of O_3 splits D . Hence, O_3 is not a well-partitioned chordal graph.

For O_4 , consider the triangle $D = \{b, c, g\}$. Because of the vertices d and h , we can conclude by Claim 4.2.1(i) that no partition tree of O_4 respects D . Since $N_{O_4}(d) \cap D = \{c, g\} = D \setminus \{b\}$, by Claim 4.2.1(ii), no partition tree of O_4 splits D into $(\{b, c\}, \{g\})$ or $(\{b, g\}, \{c\})$. Thus, we may assume that each partition tree splits D into $(\{c, g\}, \{b\})$. Let B_1 and B_2 be the two bags such that $B_1 \cap D = \{c, g\}$ and $B_2 \cap D = \{b\}$.

Since $hc \notin E(G)$, we have that $h \notin B_1$. Also, since $N_{O_4}(h) \cap \{g, c\} \neq N_{O_4}(d) \cap \{g, c\}$, h and d cannot be in the same bag. Thus, we conclude that $d \in B_1$, otherwise $\{d, h, g\}$ would be a triangle that intersects three distinct bags of \mathcal{T} . By considering the triangle $\{b, c, f\}$, we can conclude by symmetry that $\{a, b, f\}$ are contained in the same bag, which is B_2 . As i is adjacent to neither a nor d , the bag containing i forms a triangle with B_1 and B_2 , a contradiction. We can conclude that O_4 is not a well-partitioned chordal graph.

Next, we show that for all $s \in \{1, 2, 3\}$ and $t \geq 0$, $W_{s,t}$ is not a well-partitioned chordal graph. This will be done by induction on t , and for the base case $t = 0$, we consider these graphs with their vertices labelled as in Figure 4.

For convenience, we labelled the cut vertex of each graph d . We recall that for $s \in \{1, 2, 3\}$, the vertex set of a block of $W_{s,0}$ is a wing of $W_{s,0}$. For instance, $\{a, b, c, d\}$ and $\{d, x, y, z\}$ are the




Figure 4: Labellings of graphs $W_{1,0}, W_{2,0}$, and $W_{3,0}$.

wings of $W_{1,0}$.

Claim 4.2.2. For $s \in \{1, 2, 3\}$, $W_{s,0}$ has no partition tree having a bag whose intersection with a wing of $W_{s,0}$ consists of only the cut vertex.

Proof. Suppose there is a partition tree \mathcal{T}_1 of $W_{1,0}$ that contains a bag B such that $B \cap \{a, b, c, d\} = \{d\}$. This implies that there exists B_1 such that $\{a, b\} \subseteq B_1$, otherwise $\{a, b, d\}$ would be a triangle that intersects three distinct bags of \mathcal{T}_1 . Since c is not adjacent to a , $c \notin B_1$ and, by assumption, $c \notin B$. Thus $\{b, c, d\}$ is a triangle intersecting three bags of \mathcal{T}_1 , a contradiction.

Next, suppose there is a partition tree \mathcal{T}_2 of $W_{2,0}$ that contains a bag B whose intersection with a wing of $W_{2,0}$ consists of the cut vertex d alone. If the affected wing is $\{d, x, y, z\}$, then the argument follows from the same argument given before. If $B \cap \{a, b, c, d, e, f\} = \{d\}$, we observe the following. First, there must exist a bag B_1 containing $\{a, b, c\}$, otherwise there is a triangle violating Observation 4.1. Since neither e nor f is adjacent to a , $\{e, f\} \cap B_1 = \emptyset$. Since $N_{W_{2,0}}(e) \cap B_1 \neq N_{W_{2,0}}(f) \cap B_1$, by Claim (i), there is no partition tree respecting $\{a, b, c\}$, a contradiction.

The claim regarding $W_{3,0}$ follows as well, noting that the wings of $W_{3,0}$ are isomorphic to the one considered in the latter case. \lrcorner

Claim 4.2.3. For each $s \in \{1, 2, 3\}$ and $t \geq 0$, $W_{s,t}$ is not a well-partitioned chordal graph.

Proof. We prove the claim by induction on t . For $t = 0$, we observe that no bag of a partition tree can contain vertices from both wings of a $W_{s,0}$, unless it is the cut vertex. Hence, this case follows from Claim 4.2.2.

Now, suppose that for every $k \leq t - 1$, $W_{s,k}$ is not a well-partitioned chordal graph. Consider $W_{s,t}$, with $t \geq 1$. We may assume that besides the wings, $W_{s,t}$ has at least one triangle, call that triangle $D = \{p, q, s\}$. Suppose there is a partition tree \mathcal{T}_t for $W_{s,t}$. We will show how to transform this partition tree into a partition tree of $W_{s,t-1}$, contradicting the induction hypothesis. We know that \mathcal{T}_t either respects or splits D .

If \mathcal{T}_t respects D , let B denote the bag that contains D . Suppose q is the vertex in D that has degree 2 in $W_{s,t}$. Then, $B = D$, as no other vertex in $W_{s,t}$ is adjacent to q . Now, if we contract D to a single vertex, say p^* , then we obtain $W_{s,t-1}$. Moreover, if we replace B by $B^* := \{p^*\}$, and make B^* adjacent to all bags in $N_{\mathcal{T}_t}(B)$, then this gives a partition tree for $W_{s,t-1}$, a contradiction.

If \mathcal{T}_t splits D into $(\{p, s\}, \{q\})$, note that since no vertex other than q is adjacent to both p and s , we have that $B_1 = \{p, s\}$ and $B_2 = \{q\}$. As in the previous case, if we contract D to a single vertex p^* , we can delete B_2 and replace B_1 by $B^* := \{p^*\}$ to obtain a partition tree for $W_{s,t-1}$, a contradiction.

Suppose that \mathcal{T}_t splits D into $(\{p, q\}, \{s\})$, and let B_1 and B_2 be the bags of \mathcal{T}_t such that $B_1 \cap D = \{p, q\}$ and $B_2 \cap D = \{s\}$. Again, since no vertex other than p and s is adjacent to q , we have that $B_1 = \{p, q\}$. Now, if we simply remove B_1 from \mathcal{T}_t and make B_2 adjacent to all bags in $N_{\mathcal{T}_t}(B_1) \setminus \{B_2\}$, then we obtain a partition tree for $W_{s,t-1}$, a contradiction. The case in which \mathcal{T}_t splits D into $(\{q, s\}, \{p\})$ is symmetric to the latter case. \lrcorner

This concludes the proof of Proposition 4.2. \square

4.2 Boundary-crossing paths

In the remaining part of this section, we present the certifying algorithm for well-partitioned chordal graphs. Here, we define the main concept of a boundary-crossing path and prove some useful lemmas.

Let G be a connected well-partitioned chordal graph with a partition tree \mathcal{T} . For a bag X of \mathcal{T} and $B \subseteq X$, a vertex $z \in V(G) \setminus X$ is said to *cross B in X* , if it has a neighbor both in B and in $X \setminus B$. In this case, we also say that B has a crossing vertex. In the following definitions, a path $X_1 X_2 \dots X_\ell$ in \mathcal{T} is considered to be ordered from X_1 to X_ℓ . Let $\ell \geq 3$ be an integer. A path $X_1 X_2 \dots X_\ell$ in \mathcal{T} is called a *boundary-crossing path* if for each $1 \leq i \leq \ell - 2$, there is a vertex in X_i that crosses $\text{bd}(X_{i+1}, X_{i+2})$. If for each $1 \leq i \leq \ell - 2$, there is no bag $Y \in V(\mathcal{T}) \setminus \{X_i\}$ containing a vertex that crosses $\text{bd}(X_{i+1}, X_{i+2})$, then we say the path is *exclusive*. If for each $1 \leq i \leq \ell - 2$, $\text{bd}(X_i, X_{i+1})$ is complete to X_{i+1} , then we say the path is *complete*. If a boundary-crossing path is both complete and exclusive, then we call it *good*. For convenience, we say that any path in \mathcal{T} with at most two bags is a boundary-crossing path.

The outline of the algorithm is as follows. First we may assume that a given graph G is chordal, as we can detect a hole in polynomial time using Theorem 2.1 if it exists. We may also assume that G is connected. So, it has a simplicial vertex v , and by an inductive argument, we can assume that $G - v$ is a well-partitioned chordal graph. As v is simplicial, $G - v$ is also connected, and thus it admits a partition tree \mathcal{T} . If v has neighbors in one bag of \mathcal{T} , then we can simply put v as a new bag adjacent to that bag. Thus, we may assume that v has neighbors in two distinct bags, say C_1 and C_2 . Then our algorithm is divided into three parts:

- (i) We find a maximal good boundary-crossing path ending in $C_2 C_1$ (or $C_1 C_2$). To do this, when we currently have a good boundary-crossing path $C_i C_{i-1} \dots C_2 C_1$, find a bag C_{i+1} containing a vertex crossing $\text{bd}(C_i, C_{i-1})$. If there is no such bag, then this path is maximal. Otherwise, we argue that in polynomial time either we can find an obstruction, or verify that $C_{i+1} C_i \dots C_2 C_1$ is good.
- (ii) Assume that $C_k C_{k-1} \dots C_2 C_1$ is the obtained maximal good boundary-crossing path. Then we can in polynomial time modify \mathcal{T} so that no vertex crosses $\text{bd}(C_2, C_1)$.
- (iii) We show that if no vertex crosses $\text{bd}(C_2, C_1)$ and no vertex crosses $\text{bd}(C_1, C_2)$, then we can extend \mathcal{T} to a partition tree of G .

For the lemmas of this section, we fix that G is a connected chordal graph, v is a simplicial vertex, and $G - v$ is a connected well-partitioned chordal graph with partition tree \mathcal{T} , and furthermore assume that v has neighbors in two distinct bags C_1 and C_2 .

Regarding Step (ii), Lemma 4.3 shows that when a maximal good boundary-crossing path $C_k C_{k-1} \dots C_2 C_1$ is given, we can modify \mathcal{T} to a partition tree \mathcal{T}' such that no vertex crosses $\text{bd}(C'_2, C'_1)$, where C'_1 and C'_2 are the bags in \mathcal{T}' that correspond to C_1 and C_2 in \mathcal{T} , respectively – in particular, they are the bags containing the neighbors of v .

Lemma 4.3. *Let $C_k C_{k-1} \dots C_1$ be a good boundary-crossing path for some integer $k \geq 3$ such that no vertex crosses $\text{bd}(C_k, C_{k-1})$. One can in polynomial time output a partition tree \mathcal{T}' of $G - v$ that contains a good boundary-crossing path $C'_{k-1} C_{k-2} \dots C_1$ such that no vertex in $G - v$ crosses $\text{bd}(C'_{k-1}, C_{k-2})$.*

Proof. Since no vertex crosses $\text{bd}(C_k, C_{k-1})$, we can partition the neighbors of C_k in \mathcal{T} into \mathcal{S}_1 and \mathcal{S}_2 such that for all $S_1 \in \mathcal{S}_1$, we have that $\text{bd}(C_k, S_1) \subseteq C_k \setminus \text{bd}(C_k, C_{k-1})$, and for all $S_2 \in \mathcal{S}_2$, $\text{bd}(C_k, S_2) \subseteq \text{bd}(C_k, C_{k-1})$. Let $C'_k := C_k \setminus \text{bd}(C_k, C_{k-1})$ and $C'_{k-1} := C_{k-1} \cup \text{bd}(C_k, C_{k-1})$. We obtain \mathcal{T}' from \mathcal{T} as follows.

- Remove C_k and C_{k-1} , and add C'_k and C'_{k-1} .
- Make all bags that have been adjacent to C_{k-1} in \mathcal{T} adjacent to C'_{k-1} .
- Make all bags in \mathcal{S}_1 adjacent to C'_k , and all bags in \mathcal{S}_2 adjacent to C'_{k-1} .

Since $\text{bd}(C_k, C_{k-1})$ is complete to C'_{k-1} , C'_{k-1} is indeed a clique in $G - v$, and thus we conclude that \mathcal{T}' is a partition tree of $G - v$. Since C'_{k-1} contains C_{k-1} and there is no edge between $\text{bd}(C_k, C_{k-1})$ and C_{k-2} , we know that $C'_{k-1}C_{k-2} \dots C_1$ is a good boundary-crossing path. Clearly, \mathcal{T}' can be obtained in polynomial time.

We claim that no vertex crosses $\text{bd}(C'_{k-1}, C_{k-2})$. Suppose for a contradiction that there exists a vertex $q \in V(G - v) \setminus C'_{k-1}$ that crosses $\text{bd}(C'_{k-1}, C_{k-2})$. We consider two cases. First, we assume q also crosses $\text{bd}(C_{k-1}, C_{k-2})$. Since $C_k \dots C_1$ is exclusive, any vertex crossing $\text{bd}(C_{k-1}, C_{k-2})$ is in $\text{bd}(C_k, C_{k-1})$. This means that $q \in C'_{k-1}$, a contradiction. Now assume that q is a vertex in $V(G - v) \setminus (C_k \cup C_{k-1})$ that is adjacent to a vertex in $\text{bd}(C'_{k-1}, C_{k-2}) = \text{bd}(C_{k-1}, C_{k-2})$ and a vertex in $C'_{k-1} \setminus C_{k-1} = \text{bd}(C_k, C_{k-1})$. But this would mean that there is a triangle in \mathcal{T} , a contradiction.

We conclude that no vertex in $G - v$ crosses $\text{bd}(C'_{k-1}, C_{k-2})$. \square

With respect to Step (iii), we prove the following lemma.

Lemma 4.4. *If every vertex of $G - v$ crosses neither $\text{bd}(C_1, C_2)$ nor $\text{bd}(C_2, C_1)$, then one can output a partition tree for G in polynomial time.*

Proof. Assume that every vertex of $G - v$ crosses neither $\text{bd}(C_1, C_2)$ nor $\text{bd}(C_2, C_1)$. Let \mathcal{S}_1 denote all neighbors of C_1 in \mathcal{T} such that for each $S_1 \in \mathcal{S}_1$, $\text{bd}(C_1, S_1) \subseteq C_1 \setminus \text{bd}(C_1, C_2)$; let \mathcal{S}_2 denote the set of all neighbors of C_2 in \mathcal{T} such that for each $S_2 \in \mathcal{S}_2$, $\text{bd}(C_2, S_2) \subseteq C_2 \setminus \text{bd}(C_2, C_1)$; and let \mathcal{S}_{12} denote the set of all neighbors of C_1 or C_2 such that for each $S_1 \in \mathcal{S}_{12} \cap N_{\mathcal{T}}(C_1)$, $\text{bd}(C_1, S_1) \subseteq \text{bd}(C_1, C_2)$, and for each $S_2 \in \mathcal{S}_{12} \cap N_{\mathcal{T}}(C_2)$, $\text{bd}(C_2, S_2) \subseteq \text{bd}(C_2, C_1)$. Since no vertex of $G - v$ crosses neither $\text{bd}(C_1, C_2)$ nor $\text{bd}(C_2, C_1)$, $\mathcal{S}_1 \cup \mathcal{S}_2 \cup \mathcal{S}_{12} = N_{\mathcal{T}}(C_1) \cup N_{\mathcal{T}}(C_2) \setminus \{C_1, C_2\}$.

Now, let $C'_1 := C_1 \setminus \text{bd}(C_1, C_2)$, $C'_2 := C_2 \setminus \text{bd}(C_2, C_1)$, and $C'_{12} := \text{bd}(C_1, C_2) \cup \text{bd}(C_2, C_1)$. We obtain \mathcal{T}' from \mathcal{T} as follows.

- Remove C_1 and C_2 ; add C'_1 , C'_2 , and C'_{12} ; make C'_1 and C'_2 adjacent to C'_{12} .
- Make all bags in \mathcal{S}_1 adjacent to C'_1 , all bags in \mathcal{S}_2 adjacent to C'_2 , and all bags in \mathcal{S}_{12} adjacent to C'_{12} .
- Add a new bag $C_v := \{v\}$, and make it adjacent to C'_{12} .

This yields a partition tree for G . \square

Considering Step (i), we present some lemmas useful to find an obstruction. To describe subparts of the long obstructions $W_{s,t}$, we use the graphs $W_{1,t}^-$ and $W_{2,t}^-$ as shown in Figure 5. Note that each of them has a distinguished vertex r , that we call *terminal*. The following lemma will be useful to find a wing at the beginning of a boundary-crossing path.




Figure 5: The graphs $W_{1,t}^-$ and $W_{2,t}^-$.

Lemma 4.5. *Let XYZ be a boundary-crossing path in \mathcal{T} such that $\text{bd}(Y, Z)$ is complete to Z , and B be a non-empty proper subset of Z . If one of the following conditions does not hold, then one can in polynomial time output an induced subgraph H of $G[X \cup X' \cup Y \cup Z]$ for some neighbor X' of Y in \mathcal{T} (X' can be X) that is isomorphic to $W_{1,1}^-$ or $W_{2,0}^-$, with the terminal vertex being mapped to a vertex in B , say r_H , such that $V(H) \cap B = \{r_H\}$.*

- (i) $\text{bd}(X, Y)$ is complete to Y .
- (ii) There is no bag $X' \in V(\mathcal{T}) \setminus \{X\}$ that contains vertices crossing $\text{bd}(Y, Z)$.

Proof. Let $x \in \text{bd}(X, Y)$ be a vertex that crosses $\text{bd}(Y, Z)$. Choose a neighbor y in $\text{bd}(Y, Z)$ and a neighbor y' in $Y \setminus \text{bd}(Y, Z)$ of x . Since $\text{bd}(Y, Z)$ is complete to Z by assumption, y has a neighbor in B and a neighbor in $Z \setminus B$. Let z and z' be these neighbors, respectively. We illustrate this situation and the following arguments in Figure 6.

Suppose that (i) does not hold, i.e. that $\text{bd}(X, Y)$, in particular the vertex x is not complete to Y . Then, x has a non-neighbor, say y'' in Y . If $y'' \in Y \setminus \text{bd}(Y, Z)$, then the set $\{x, y, y', y'', z, z'\}$ induces a $W_{1,1}^-$ with the terminal being mapped to z . See Figure 6(a). On the other hand, if $y'' \in \text{bd}(Y, Z)$, then $\{x, y, y', y'', z, z'\}$ induce a $W_{2,0}^-$ with the terminal vertex being mapped to z . See Figure 6(b).

Now suppose that (ii) does not hold, and let $x' \in X'$ be a vertex crossing $\text{bd}(Y, Z)$. Then, x' has a neighbor $y \in \text{bd}(Y, Z)$ and a neighbor $y' \in Y \setminus \text{bd}(Y, Z)$. Let $x \in X$. By (i), x is adjacent to y and y' . Then $G[X \cup X' \cup Y \cup Z]$ contains a $W_{1,1}^-$ with terminal z . \square

We use the following lemmas to find an obstruction or extend a good boundary-crossing path.




Figure 6: Visual aides to the proof of Lemma 4.5.

Lemma 4.6. *Let G be a connected well-partitioned chordal graph with partition tree \mathcal{T} , and let B be a vertex set contained in some bag C_1 . If $C_k C_{k-1} \dots C_1$ is a boundary-crossing path for some $k \geq 2$ such that C_2 has a vertex that crosses B in C_1 , then one can in polynomial time either*

- (i) *output an induced subgraph H isomorphic to $W_{s,t}^-$ for some $s \in \{1, 2, 3\}$ and $t \geq 0$ with terminal v such that $V(H) \cap B = \{v\}$,*
- (ii) *output an induced subgraph H isomorphic to $W_{1,0}^-$ on $\{a, z_1, z_2, w\}$ such that both a and w have degree 2 in H , $a \in \text{bd}(D, C_1)$ for some neighbor bag D of C_1 , $z_2 \in C_1 \setminus B$, and $z_1, w \in B$, or*
- (iii) *verify that it is a good boundary-crossing path such that $\text{bd}(C_2, C_1)$ is complete to C_1 and no other bag contains a vertex crossing B in C_1 .*

Proof. We prove the lemma by induction on k . Assume that $k = 2$. We check whether $\text{bd}(C_2, C_1)$ is complete to C_1 . Suppose not. Let $a \in \text{bd}(C_2, C_1)$. Let z_1 be a neighbor of a in B , z_2 be a neighbor of a in $C_1 \setminus B$, and w be a non-neighbor of a in C_1 . If $w \in C_1 \setminus B$, then $\{a, z_1, z_2, w\}$ induces $W_{1,0}^-$ with terminal z_1 , so we have outcome (i). If $w \in B$, then $\{a, z_1, z_2, w\}$ induces a graph as in case (ii). Otherwise, we conclude that $\text{bd}(C_2, C_1)$ is complete to C_1 .

We find a bag $D \neq C_2$ in \mathcal{T} containing a vertex d crossing B in C_1 . If such a vertex d exists, then by the above procedure, we may assume that d is complete to C_1 . Then similarly to the previous case when $w \in C_1 \setminus B$, again we can find an induced subgraph isomorphic to the diamond on $\{a, d, z_1, z_2\}$. If such a vertex does not exist, then we can conclude that no other bag contains a vertex crossing B in C_1 .

Now, we assume that $k \geq 3$. By the induction hypothesis, the claim holds for the path $C_{k-1} C_{k-2} \dots C_1$. We can assume that it is good.

We check whether $\text{bd}(C_k, C_{k-1})$ is not complete to C_{k-1} , and there is a bag $D \in V(\mathcal{T}) \setminus \{C_k\}$ that has a vertex crossing $\text{bd}(C_k, C_{k-1})$. If neither of them holds, then we verified that $C_k C_{k-1} \dots C_1$ is good. Assume one of two statements holds.

Let $X := \text{bd}(C_{k-2}, C_{k-3})$ if $k \geq 4$ and $X = B$ if $k = 3$. Now, by applying Lemma 4.5 to the pair $(C_k C_{k-1} C_{k-2}, X)$, we can find an induced subgraph H isomorphic to $W_{1,1}^-$ or $W_{2,0}^-$ in $G[C_k \cup D \cup C_{k-1} \cup C_{k-2}]$ for some neighbor D of C_{k-1} in \mathcal{T} so that its terminal r is mapped to some vertex in X and $V(H) \cap X = \{r\}$. If $k = 3$, then we have outcome (i) as $X = B$.

Assume $k \geq 4$. Let $x_{k-2} := r$. We recursively choose pairs of vertices (x_i, y_i) for $i \in \{1, 2, \dots, k-3\}$ as follows. First assume $i > 1$ and x_{i+1} is defined but x_i is not defined yet. Then choose a neighbor x_i of x_{i+1} in $\text{bd}(C_i, C_{i-1})$ and a neighbor y_i of x_{i+1} in $C_i \setminus \text{bd}(C_i, C_{i-1})$. Such neighbors exists since x_{i+1} crosses $\text{bd}(C_i, C_{i-1})$. When $i = 1$, choose a neighbor x_1 of x_2 in B and y_1 of x_2 in $C_1 \setminus B$. Then it is clear that $G[\{x_1, y_1, x_2, y_2, \dots, x_{k-3}, y_{k-3}\} \cup V(H)]$ is isomorphic to $W_{s',t'}^-$ for some $s' \in \{1, 2\}$ and $t' \geq 0$ with terminal x_1 such that its intersection on B is exactly x_1 . This concludes the lemma. \square

Lemma 4.7. *Let G_1 and G_2 be two connected graphs with non-empty sets $A \subseteq V(G_1)$ and $B \subseteq V(G_2)$, and G be the graph obtained from the disjoint union of G_1 and G_2 by adding all edges between A and B such that*

- *for every $v \in B$, $G[V(G_1) \cup \{v\}]$ is isomorphic to $W_{s,t}^-$ with terminal v for some $s \in \{1, 2\}$ and $t \geq 0$,*

- G_2 is a well-partitioned chordal graph with a partition tree \mathcal{T} such that B is contained in some bag C_1 .

Then the following two statements hold.

- (1) If $C_k C_{k-1} \dots C_1$ is a boundary-crossing path in \mathcal{T} for some $k \geq 2$ such that C_2 has a vertex that crosses B in C_1 , then one can in polynomial time either output an obstruction in G , or verify that it is a good boundary-crossing path such that $\text{bd}(C_2, C_1)$ is complete to C_1 and no other bag contains a vertex crossing B in C_1 .
- (2) If $C_2 C_1$ is a boundary-crossing path in \mathcal{T} , that is, an edge in \mathcal{T} , then one can in polynomial time either output an obstruction in G , or find a maximal good boundary-crossing path ending in $C_2 C_1$ such that $\text{bd}(C_2, C_1)$ is complete to C_1 and no other bag contains a vertex crossing B in C_1 .

Proof. We prove (1). We apply Lemma 4.6 to G_2 and B , then we conclude that in polynomial time, we can either

- output an induced subgraph H isomorphic to $W_{s,t}^-$ for some $s \in \{1, 2, 3\}$ and $t \geq 0$ with terminal v such that $V(H) \cap B = \{v\}$,
- output an induced subgraph H isomorphic to the diamond on $\{a, z_1, z_2, w\}$ such that a, w have degree 2 in H , $a \in \text{bd}(D, C_1)$ for some neighbor bag D of C_1 , $z_2 \in C_1 \setminus B$, and $z_1, w \in B$, or
- verify that it is a good boundary-crossing path such that $\text{bd}(C_2, C_1)$ is complete to C_1 and no other bag contains a vertex crossing B in C_1 .

For case (i) it is clear that together with an obstruction $W_{s,t}^-$ in $G[V(G_1) \cup \{v\}]$ given by the assumption, $G[V(H) \cup V(G_1)]$ is isomorphic to $W_{s,t}$ for some $s \in \{1, 2, 3\}$ and $t \geq 0$. For case (ii), we can observe that $G[V(H) \cup V(G_1)]$ is an obstruction as follows.

- If $G[V(G_1) \cup \{z_1\}]$ is isomorphic to $W_{1,0}^-$, then $G[V(G_1) \cup \{a, w, z_1, z_2\}]$ is isomorphic to O_3 .
- If $G[V(G_1) \cup \{z_1\}]$ is isomorphic to $W_{2,0}^-$, then $G[V(G_1) \cup \{a, w, z_1, z_2\}]$ is isomorphic to O_4 .
- If $G[V(G_1) \cup \{z_1\}]$ is isomorphic to $W_{s,t}^-$ for some $s \in \{1, 2\}$ and $t \geq 1$, then $G[V(G_1) \cup \{a, w, z_1, z_2\}]$ is isomorphic to $W_{s',t-1}$ for some $s' \in \{2, 3\}$.

It shows the statement (1).

Now, we show (2). By (1), we can in polynomial time either output an obstruction, or verify that $\text{bd}(C_2, C_1)$ is complete to C_1 and no other bag crosses $\text{bd}(C_2, C_1)$. For $i \geq 3$, we recursively find a neighbor bag C_i of C_{i-1} that has a vertex crossing $\text{bd}(C_{i-1}, C_{i-2})$. If there is such a bag C_i , then by applying (1), one can in polynomial time find an obstruction or guarantee that $C_i C_{i-1} \dots C_1$ is good. As the graph is finite, this procedure terminates with some path $C_k C_{k-1} \dots C_3$ such that it is good and no vertex crosses $\text{bd}(C_k, C_{k-1})$, unless we found an obstruction. \square

4.3 A certifying algorithm

In this subsection, we prove the following.

Proposition 4.8. *Given a graph G , one can in polynomial time either output an obstruction in G or output a partition tree of G confirming that G is a well-partitioned chordal graph.*

As explained in Subsection 4.2, we mainly consider the case when G is a connected chordal graph, v is a simplicial vertex of G and $G - v$ is a connected well-partitioned chordal graph with partition tree \mathcal{T} , and v has neighbors in two distinct bags C_1 and C_2 . With these assumptions, we deal with the following three cases, and in each case, we show that either one can in polynomial time find an obstruction or output a partition tree of G .

- (Lemma 4.9) $C_1 \subseteq N_G(v)$.
- (Lemma 4.10) $\text{bd}(C_1, C_2) \setminus N_G(v) \neq \emptyset$ and $C_2 \setminus N_G(v) \neq \emptyset$.
- (Lemma 4.11) $C_1 \setminus N_G(v) \neq \emptyset$, $C_2 \setminus N_G(v) \neq \emptyset$ and $N_G(v) = \text{bd}(C_1, C_2) \cup \text{bd}(C_2, C_1)$.

We give a proof of Proposition 4.8 assuming that these lemmas hold.

Proof (Proof of Proposition 4.8). We apply Theorem 2.1 to find a hole in G if one exists. We may assume that G is chordal. Since a graph is a well-partitioned chordal graph if and only if its connected components are well-partitioned chordal graphs, it is sufficient to show it for each connected component. From now on, we assume that G is connected. Using the algorithm in Theorem 2.2, we can find a perfect elimination ordering (v_1, v_2, \dots, v_n) of G in polynomial time.

For each $i \in \{1, 2, \dots, n\}$, let $G_i := G[\{v_i, v_{i+1}, \dots, v_n\}]$. Observe that since G is connected and v_i is simplicial in G_i for all $1 \leq i \leq n - 1$, each G_i is connected. From $i = n$ to 1, we recursively find either an obstruction or a partition tree of G_i . Clearly, G_n admits a partition tree. Let $1 \leq i \leq n - 1$, and assume that we obtained a partition tree \mathcal{T} of G_{i+1} . Recall that v_i is simplicial in G_i .

Since v_i is simplicial in G_i , $N_{G_i}(v_i)$ is a clique. This implies that there are at most two bags in $V(\mathcal{T})$ that have a non-empty intersection with $N_{G_i}(v_i)$. If there is only one such bag in $V(\mathcal{T})$, say C , we can construct a partition tree for G_i by simply adding a bag consisting of v_i and making it adjacent to C .

Hence, from now on, we can assume that there are precisely two distinct adjacent bags $C_1, C_2 \in V(\mathcal{T})$ that have a non-empty intersection with $N_{G_i}(v_i)$. As $N_{G_i}(v_i)$ is a clique, we can observe that $N_{G_i}(v_i) \subseteq \text{bd}(C_1, C_2) \cup \text{bd}(C_2, C_1)$.

If $C_1 \subseteq N_{G_i}(v_i)$ or $C_2 \subseteq N_{G_i}(v_i)$, then by Lemma 4.9, we can in polynomial time either output an obstruction or output a partition tree of G_i . Thus, we may assume that $C_1 \setminus N_{G_i}(v_i) \neq \emptyset$ and $C_2 \setminus N_{G_i}(v_i) \neq \emptyset$. If $\text{bd}(C_1, C_2) \setminus N_{G_i}(v_i) \neq \emptyset$ or $\text{bd}(C_2, C_1) \setminus N_{G_i}(v_i) \neq \emptyset$, then by Lemma 4.10, we can in polynomial time either output an obstruction or output a partition tree of G_i . Thus, we may further assume that $\text{bd}(C_1, C_2) \setminus N_{G_i}(v_i) = \emptyset$ and $\text{bd}(C_2, C_1) \setminus N_{G_i}(v_i) = \emptyset$. Then by Lemma 4.11, we can in polynomial time either output an obstruction or output a partition tree of G_i , and this concludes the proposition. \square

Now, we focus on proving the three lemmas.

Lemma 4.9. *If $C_1 \subseteq N_G(v)$, then one can in polynomial time either output an obstruction in G or output a partition tree of G confirming that G is a well-partitioned chordal graph.*




Figure 7: Proof of Claim 4.9.1.

Proof. Since v is a simplicial vertex, we have that $\text{bd}(C_1, C_2) = C_1$. If $N_G(v) \cap C_2 = \text{bd}(C_2, C_1)$, then we can obtain a partition tree for G by adding v to C_1 . Thus, we may assume that $N_G(v) \cap C_2 \neq \text{bd}(C_2, C_1)$.

Assume that $C_2 = \text{bd}(C_2, C_1)$. Since $\text{bd}(C_2, C_1)$ is complete to C_1 , we have that $C_1 \cup C_2$ is a clique. Hence, we can obtain a partition tree \mathcal{T}' for G from \mathcal{T} by removing C_1 and C_2 , adding a new bag $C^* = C_1 \cup C_2$, making all neighbors of C_1 and C_2 in \mathcal{T} adjacent to C^* , and adding a new bag $C_v := \{v\}$ and making C_v adjacent to C^* . Thus, we may assume that $C_2 \setminus \text{bd}(C_2, C_1) \neq \emptyset$.

Since $C_1 = \text{bd}(C_1, C_2)$, no vertex of $G - v$ crosses $\text{bd}(C_1, C_2)$. If no vertex of $G - v$ crosses $\text{bd}(C_2, C_1)$, then by Lemma 4.4, we can obtain a partition tree for G in polynomial time. Thus, we may assume that there is a bag C_3 having a vertex that crosses $\text{bd}(C_2, C_1)$. So, $C_3 C_2 C_1$ is a boundary-crossing path. We will find either an obstruction or a maximal good boundary-crossing path ending in $C_3 C_2 C_1$. We first check that $C_3 C_2 C_1$ is good, unless some obstruction from \mathbb{O} appears.

Claim 4.9.1. Let $z_1 \in N_G(v) \cap C_1$, $z_2 \in N_G(v) \cap C_2$, $w \in \text{bd}(C_2, C_1) \setminus N_G(v)$, and $a \in C_2 \setminus \text{bd}(C_2, C_1)$.

- (i) If there is a vertex $x \in V(G) \setminus \{v, a, w, z_1, z_2\}$ such that $N(x) \cap \{v, a, w, z_1, z_2\} = \{a, w\}$, then $G[\{v, a, w, z_1, z_2, x\}]$ is isomorphic to O_1 .
- (ii) If there is a vertex $x \in V(G) \setminus \{v, a, w, z_1, z_2\}$ such that $N(x) \cap \{v, a, w, z_1, z_2\} = \{a, z_2\}$, then $G[\{v, a, w, z_1, z_2, x\}]$ is isomorphic to O_2 .
- (iii) If there is a pair of distinct non-adjacent vertices $x, y \in V(G) \setminus \{v, a, w, z_1, z_2\}$ such that $N(x) \cap \{v, a, w, z_1, z_2\} = N(y) \cap \{v, a, w, z_1, z_2\} = \{a, w, z_2\}$, then $G[\{v, a, w, z_1, z_2, x, y\}]$ is isomorphic to O_3 .

Proof. It is straightforward to check it; see Figure 7. ┘

Claim 4.9.2. One can in polynomial time output an obstruction or verify that $C_3 C_2 C_1$ is good.

Proof. We consider the bag C_3 , and first check whether $\text{bd}(C_3, C_2)$ is complete to C_2 . If so, then we are done. Otherwise, choose a vertex $p \in \text{bd}(C_3, C_2)$, and a non-neighbor q of p in C_2 . As p crosses $\text{bd}(C_2, C_1)$, p has a neighbor a in $C_2 \setminus \text{bd}(C_2, C_1)$ and a neighbor b in $\text{bd}(C_2, C_1)$. There are three possibilities; q is contained in one of $N_G(v) \cap C_2$, $\text{bd}(C_2, C_1) \setminus N_G(v)$, or $C_2 \setminus \text{bd}(C_2, C_1)$. Let $z_1 \in N_G(v) \cap C_1$.

If q and b are in distinct parts of $N_G(v) \cap C_1$ and $\text{bd}(C_2, C_1) \setminus N_G(v)$, then $G[\{p, q, z_1, a, b, v\}]$ is isomorphic to O_1 or O_2 by Claims 4.9.1(i) and (ii). Assume q and b are in the same part of

$N_G(v) \cap C_1$ or $\text{bd}(C_2, C_1) \setminus N_G(v)$. Then by the previous argument, we may assume that p is complete to the set, one of $N_G(v) \cap C_1$ and $\text{bd}(C_2, C_1) \setminus N_G(v)$, that does not contain q . Then by choosing a vertex in this set, we can again output O_1 or O_2 . Thus, we may assume that q is contained in $C_2 \setminus \text{bd}(C_2, C_1)$ and p is complete to $\text{bd}(C_2, C_1)$. Then $q \neq a$ and by using vertices from $N_G(v) \cap C_1$ and $\text{bd}(C_2, C_1) \setminus N_G(v)$ together with $\{a, p, q, v, z_1\}$, we can output O_3 by Claim 4.9.1(iii).

To verify whether $C_3C_2C_1$ is exclusive, we check if there exists another neighbor bag $D \neq C_3$ of C_2 having a vertex q that crosses $\text{bd}(C_2, C_1)$. If there is such a vertex q , then by applying the previous procedure, we may assume that q is complete to C_2 . Then by using vertices from each of $N_G(v) \cap C_2$, $\text{bd}(C_2, C_1) \setminus N_G(v)$, and $C_2 \setminus \text{bd}(C_2, C_1)$ together with $\{p, q, z_1, v\}$, we can output O_3 by Claim 4.9.1(iii). Otherwise, $C_3C_2C_1$ is a good boundary-crossing path. \lrcorner

By Claim 4.9.2, we may assume that $C_3C_2C_1$ is good. If no bag contains a vertex crossing $\text{bd}(C_3, C_2)$, then $C_3C_2C_1$ is a maximal good boundary-crossing path. So, we may assume that there is a bag C_4 containing a vertex crossing $\text{bd}(C_3, C_2)$.

We choose $z_1 \in N_G(v) \cap C_1$, $z_2 \in N_G(v) \cap C_2$, $w \in \text{bd}(C_2, C_1) \setminus N_G(v)$, and $a \in C_2 \setminus \text{bd}(C_2, C_1)$. To apply Lemma 4.7, let $G_1 = G[\{v, z_1, z_2, w, a\}]$ and G_2 be the component of $G - V(C_2)$ that contains C_3 and $G' = G[V(G_1) \cup V(G_2)]$. It is clear that G' can be obtained from the disjoint union of G_1 and G_2 by adding edges between $\text{bd}(C_3, C_2)$ and $\{w, a, z_2\}$. Also, for each vertex $p \in \text{bd}(C_3, C_2)$, $\{p\} \cup V(G_1)$ is a wing of $W_{2,0}$ with terminal p .

Thus, by (2) of Lemma 4.7, we can in polynomial time either output an obstruction, or find a maximal good boundary-crossing path ending in C_4C_3 in G_2 such that $\text{bd}(C_4, C_3)$ is complete to C_3 and no other bag contains a vertex crossing C_3 . Thus, in the latter case, we obtain a maximal boundary-crossing path ending in C_2C_1 in $G - v$. We now repeatedly apply Lemma 4.3 to modify \mathcal{T} along this path and obtain a partition tree \mathcal{T}' for $G - v$ such that no vertex crosses $\text{bd}(C_2, C_1)$. Note that, for simplicity, we call again C_1 and C_2 the bags of \mathcal{T}' containing the neighbors of v . We can now apply Lemma 4.4 to obtain a partition tree for the entire graph G in polynomial time. \square

Lemma 4.10. *If $\text{bd}(C_1, C_2) \setminus N(v) \neq \emptyset$ and $C_2 \setminus N(v) \neq \emptyset$, then one can in polynomial time either output an obstruction in G or output a partition tree of G confirming that G is a well-partitioned chordal graph.*

Proof. We choose a neighbor z_1 of v in $\text{bd}(C_1, C_2)$, a neighbor z_2 of v in $\text{bd}(C_2, C_1)$ and a non-neighbor x of v in $\text{bd}(C_1, C_2)$. We first consider the case when $\text{bd}(C_1, C_2) = C_1$.

Case 1 ($\text{bd}(C_1, C_2) = C_1$). Note that no vertex in $G - v$ crosses $\text{bd}(C_1, C_2)$. If no vertex in $G - v$ crosses $\text{bd}(C_2, C_1)$, then we can obtain a partition tree of G from \mathcal{T} by Lemma 4.4. We may assume that there is a bag C_3 containing a vertex a that crosses $\text{bd}(C_2, C_1)$.

Claim 4.10.1. *One can in polynomial time output an obstruction or verify that $C_3C_2C_1$ is good.*

Proof. As a crosses $\text{bd}(C_2, C_1)$, a has a neighbor both in $C_2 \setminus \text{bd}(C_2, C_1)$ and in $\text{bd}(C_2, C_1)$. Let b_1 and b_2 be neighbors of a in $C_2 \setminus \text{bd}(C_2, C_1)$ and $\text{bd}(C_2, C_1)$, respectively.

Assume that a and v have no common neighbors. Then b_2 is not adjacent to v and z_2 is not adjacent to a . So, $G[\{a, b_1, b_2, z_2, z_1, v\}]$ is isomorphic to O_1 , see Figure 8(a). Thus, we may assume that a and v have the common neighbor in $\text{bd}(C_2, C_1)$. We assume that z_2 is a common neighbor.




Figure 8: Illustration of some obstructions appearing in the proof of Claim 4.10.1.

Now suppose that there is a vertex $w \in C_2 \setminus \text{bd}(C_2, C_1)$ that is not adjacent to a . Recall that since $N(v) \cap C_2 \subseteq \text{bd}(C_2, C_1)$, we have that v is not adjacent to w . Thus, we can output a $W_{1,0}$ on $\{v, x, z_1, z_2, a, b_1, w\}$, see Figure 8(b). So, we may assume that a is complete to $C_2 \setminus \text{bd}(C_2, C_1)$.

Assume that there is a vertex $w \in \text{bd}(C_2, C_1)$ that is not adjacent to a . Note that v may or may not be adjacent to w . If v is adjacent to w , then G contains O_3 as an induced subgraph, and if v is not adjacent to w , then G contains O_2 as an induced subgraph; both these cases are illustrated in Figure 9. Otherwise, we can conclude that a is complete to C_2 .

To check whether $C_3C_2C_1$ is exclusive, we find a bag $D \neq C_3$ containing a vertex w crossing $\text{bd}(C_2, C_1)$. If there is no such a vertex, then it is exclusive. Assume that such a vertex w exists. By repeating the above argument, we may assume that w is complete to C_2 . Then, $G[\{v, z_1, z_2, x, a, b_1, w\}]$ is isomorphic to $W_{1,0}$ (see Figure 8(b), but note that in this case $w \notin C_2$). \square

By Claim 4.10.1, we may assume that $C_3C_2C_1$ is good. Let $a \in C_2 \setminus \text{bd}(C_2, C_1)$. If no bag contains a vertex crossing $\text{bd}(C_3, C_2)$, then $C_3C_2C_1$ is a maximal good boundary-crossing path. So, we may assume that there is a bag C_4 containing a vertex crossing $\text{bd}(C_3, C_2)$.

To apply Lemma 4.7, let $G_1 = G[\{v, x, z_1, z_2, a\}]$ and G_2 be the component of $G - V(C_2)$ containing C_3 and $G' = G[V(G_1) \cup V(G_2)]$. It is clear that G' can be obtained from the disjoint




Figure 9: Illustration of some more obstructions appearing in the proof of Claim 4.10.1. Note that the edge between v and w may or may not be present, depending on which we either have an O_3 or an O_2 as an induced subgraph in G .

union of G_1 and G_2 by adding edges between $\text{bd}(C_3, C_2)$ and $\{a, z_2\}$. Observe that for each vertex $p \in \text{bd}(C_3, C_2)$, $G[\{p, a, v, z_1, z_2, z\}]$ is isomorphic to $W_{1,1}^-$ with terminal p .

By (2) of Lemma 4.7, we can in polynomial time either output an obstruction, or find a maximal good boundary-crossing path ending in C_4C_3 in G_2 such that $\text{bd}(C_4, C_3)$ is complete to C_3 and no other bag contains a vertex crossing C_3 . Thus, in the latter case, we obtain a maximal boundary-crossing path ending in C_2C_1 in $G - v$. We can now repeatedly apply Lemma 4.3 to modify \mathcal{T} along this path and obtain a partition tree \mathcal{T}' for $G - v$ such that no vertex crosses $\text{bd}(C_2, C_1)$. Note that, for simplicity, we call again C_1 and C_2 the bags of \mathcal{T}' containing the neighbors of v . We can now apply Lemma 4.4 to obtain a partition tree for the entire graph G in polynomial time. •

Case 2 ($C_1 \setminus \text{bd}(C_1, C_2) \neq \emptyset$). If there is no vertex crossing $\text{bd}(C_1, C_2)$ and no vertex crossing $\text{bd}(C_2, C_1)$ in $G - v$, then by Lemma 4.4, one can output a partition tree of G from \mathcal{T} in polynomial time. Recall that we have neighbors of v , namely $z_1 \in \text{bd}(C_1, C_2)$ and $z_2 \in \text{bd}(C_2, C_1)$, and a non-neighbor of v , namely $x \in \text{bd}(C_1, C_2)$.

Claim 4.10.2. *If there is a vertex crossing $\text{bd}(C_1, C_2)$ or $\text{bd}(C_2, C_1)$, then one can in polynomial time output an obstruction or output a partition tree of G from \mathcal{T} confirming that G is a well-partitioned chordal graph.*

Proof. First we consider the case in which only $\text{bd}(C_1, C_2)$ has a crossing vertex. Let a be a vertex in a bag $C_3 \in V(\mathcal{T}) \setminus \{C_1, C_2\}$ that crosses $\text{bd}(C_1, C_2)$. Let $b \in C_1 \setminus \text{bd}(C_1, C_2)$ be a neighbor of a . Note that a neighbor of a in $\text{bd}(C_1, C_2)$ is either adjacent to v , as z_1 , or non-adjacent to v , as x . As in Claim 4.9.1, we can restrict the way $N(a)$ intersects $\{x, b, z_1\}$, and as we did in Claim 4.9.2, we can deduce that $\text{bd}(C_3, C_1)$ is complete to C_1 and that there is no bag other than C_3 containing a vertex that crosses $\text{bd}(C_1, C_2)$.

Observe that $\{v, z_1, z_2, x, a, b\}$ induces a $W_{2,0}^-$ with terminal vertex a . By applying Lemma 4.7 similarly in Case 1, one can in polynomial time find an obstruction or find a maximal good boundary-crossing path ending in $C_3C_1C_2$. In the latter case, we apply Lemma 4.3 to modify \mathcal{T} along this path and obtain a partition tree \mathcal{T}' for $G - v$ such that no vertex crosses $\text{bd}(C_1, C_2)$. Then, since both $\text{bd}(C_1, C_2)$ and $\text{bd}(C_2, C_1)$ have no crossing vertices, we can apply Lemma 4.4 to obtain a partition tree for G .

Now we consider the case in which only $\text{bd}(C_2, C_1)$ has a crossing vertex. Let a be a vertex in a bag C_3 that crosses $\text{bd}(C_2, C_1)$. Note that $\{v, z_1, z_2, x\}$ is a wing of $W_{1,0}$ with terminal z_2 , as in Case 1 (see (b) of Figure 8). As in Claims 4.10.1 and Lemma 4.7, we can find a maximal good boundary-crossing path ending in C_2C_1 . We apply Lemma 4.3 to modify \mathcal{T} along this path and obtain a partition tree \mathcal{T}' for $G - v$ such that no vertex crosses $\text{bd}(C_2, C_1)$. Then, since both $\text{bd}(C_1, C_2)$ and $\text{bd}(C_2, C_1)$ have no crossing vertices, we can apply Lemma 4.4 to obtain a partition tree for G .

To conclude, in the case in which both $\text{bd}(C_1, C_2)$ and $\text{bd}(C_2, C_1)$ have crossing vertices, we can first modify \mathcal{T} along a maximal boundary-crossing path ending in C_2C_1 , then along a maximal boundary-crossing path ending in C_1C_2 . In this way we obtain a partition tree for $G - v$ in which, again, both $\text{bd}(C_1, C_2)$ and $\text{bd}(C_2, C_1)$ have no crossing vertices and we proceed with Lemma 4.4. ◻

This concludes the lemma. ◻

Lemma 4.11. *If $C_1 \setminus N(v) \neq \emptyset$, $C_2 \setminus N(v) \neq \emptyset$ and $N(v) = \text{bd}(C_1, C_2) \cup \text{bd}(C_2, C_1)$, then one can in polynomial time either output an obstruction in G or output a partition tree of G confirming that G is a well-partitioned chordal graph.*

Proof. We first show that if at least one of $\text{bd}(C_1, C_2)$ and $\text{bd}(C_2, C_1)$ has no crossing vertex, then we can obtain a partition tree for G .

Claim 4.11.1. *If there is no vertex crossing $\text{bd}(C_1, C_2)$, then one can obtain a partition tree of G from \mathcal{T} in polynomial time. The same holds for $\text{bd}(C_2, C_1)$.*

Proof. We prove the claim for $\text{bd}(C_1, C_2)$ and note that the argument for $\text{bd}(C_2, C_1)$ is symmetric. Let $C'_1 := C_1 \setminus \text{bd}(C_1, C_2)$, and $C'_{12} := \text{bd}(C_1, C_2) \cup \{v\}$. Let $\mathcal{S}_1 \subseteq N_{\mathcal{T}}(C_1)$ be such that for all $S_1 \in \mathcal{S}_1$, $\text{bd}(C_1, S_1) \subseteq C_1 \setminus \text{bd}(C_1, C_2)$, and let $\mathcal{S}_2 \subseteq N_{\mathcal{T}}(C_1)$ be such that for all $S_2 \in \mathcal{S}_2$, $\text{bd}(C_1, S_2) \subseteq \text{bd}(C_1, C_2)$. We obtain a partition tree \mathcal{T}' for G from \mathcal{T} as follows.

- Remove C_1 ; add C'_1 and C'_{12} ; make C'_1 adjacent to C'_{12} , and C'_{12} adjacent to C_2 .
- Make each bag in \mathcal{S}_1 adjacent to C'_1 , and each bag in \mathcal{S}_2 adjacent to C'_{12} .

This yields a partition tree for G . ▮

From now on, we assume that both $\text{bd}(C_1, C_2)$ and $\text{bd}(C_2, C_1)$ have crossing vertices. Let C'_2 be a bag containing a vertex crossing $\text{bd}(C_1, C_2)$, and let C_3 be a bag containing a vertex crossing $\text{bd}(C_2, C_1)$. For convenience, let $C'_1 := C_1$.

Using Lemma 4.6 with $B = \text{bd}(C'_1, C_2)$, we recursively find a longer good boundary-crossing path or a partial obstruction. Starting from $C'_2 C'_1$, for a path $C'_i C'_{i-1} \dots C'_1$, we find a neighbor bag C'_{i+1} of C'_i that contains a vertex crossing $\text{bd}(C'_i, C'_{i-1})$. At the end, either we can find one of first two outcomes in Lemma 4.6, or we can find a maximal good boundary-crossing path ending in $C'_2 C'_1 C_2$. In the latter case, we can repeatedly apply Lemma 4.3 to modify \mathcal{T} along this path and obtain a partition tree \mathcal{T}' for $G - v$ such that no vertex crosses $\text{bd}(C'_1, C'_2)$. We can now apply Claim 4.11.1 to obtain a partition tree for the entire graph G . Thus, we may assume that we have an induced subgraph H_1 which is one of two outcomes in Lemma 4.6. Let v_1 be the terminal of H_1 in $\text{bd}(C'_1, C_2)$.

By applying the same argument for $C_2 C_3$, we may assume that we have an induced subgraph H_2 which is one of two outcomes in Lemma 4.6. Let v_2 be the terminal of H_2 in $\text{bd}(C_2, C_1)$.

If both H_1 and H_2 are the first outcome in Lemma 4.6, then $G[V(H_1) \cup V(H_2) \cup \{v\}]$ is isomorphic to $W_{s,t}$ for some $s \in \{1, 2, 3\}$ and $t \geq 0$. If H_1 is the first outcome and H_2 is the second outcome of Lemma 4.6, then $G[V(H_1) \cup V(H_2) \cup \{v\}]$ is isomorphic to $W_{s,t}$ for some $s \in \{2, 3\}$ and $t \geq 0$, where $G[V(H_2) \cup \{v, v_1\}]$ is isomorphic to $W_{2,0}^-$. If both are the second outcomes in Lemma 4.6, then $G[V(H_1) \cup V(H_2) \cup \{v\}]$ is isomorphic to O_4 , and this concludes the lemma. □

5 Algorithmic applications

In this section, we give several FPT-algorithms and kernels for problems on well-partitioned chordal graphs. Specifically, we consider variants of the DISJOINT PATHS problem, where each path additionally has to be from a predefined domain. Before we proceed with the algorithmic description, we review the variants of the term ‘disjoint paths’ that we use in this section. Let P_1 be an (s_1, t_1) -path and P_2 be an (s_2, t_2) -path. The paths P_1 and P_2 being disjoint is most literally translated to $V(P_1) \cap V(P_2) = \emptyset$. However, in some applications of disjoint paths problems, e.g. [1, 18], notions of *internally disjoint* paths are used, meaning that the intersection of $\{s_1, t_1\}$ and $\{s_2, t_2\}$ may be nonempty. In this section, we deal with two variants of the latter definition and note

that they in fact both generalize the setting where we require $V(P_1) \cap V(P_2) = \emptyset$. Specifically, in Section 5.1, we consider the SET-RESTRICTED DISJOINT PATHS problem, asking for *internally vertex-disjoint* paths. We say that P_1 and P_2 are internally vertex-disjoint, if for $i \in [2]$, $(V(P_i) \setminus \{s_i, t_i\}) \cap V(P_{3-i}) = \emptyset$, meaning that no internal vertex of one path is used as a vertex on the other path. However, if $\{s_1, t_1\} = \{s_2, t_2\}$ and $s_1 t_1 \in E(G)$, then according to this definition, the edge $s_1 t_1$ can be used both as the path P_1 and as the path P_2 in a solution without violating the definition. In Section 5.2, we study the SET-RESTRICTED TOTALLY DISJOINT PATHS problem that asks for internally vertex-disjoint paths that are also distinct. In Section 5.3, we sketch how to use the same methods to solve the related SET-RESTRICTED DISJOINT CONNECTED SUBGRAPHS problem on well-partitioned chordal graphs.

Moreover, in Section 5.4, we observe that the algorithms given in Sections 5.1 and 5.2 imply polynomial kernels for the SET-RESTRICTED DISJOINT PATHS and SET-RESTRICTED TOTALLY DISJOINT PATHS problems on split graphs. In Section 5.5, we show that with two more simple reduction rules regarding degree-two bags of partition trees we can obtain polynomial kernels for DISJOINT PATHS and TOTALLY DISJOINT PATHS on well-partitioned chordal graphs.

5.1 Set-Restricted Disjoint Paths

In this section we deal with the following parameterized problem, and show that it is in FPT on well-partitioned chordal graphs.

SET-RESTRICTED DISJOINT PATHS parameterized by k

Input: A graph G , a set $\mathcal{X} = \{(s_1, t_1), \dots, (s_k, t_k)\}$ of k pairs of vertices of G , called *terminals*, a set $\mathcal{U} = \{U_1, \dots, U_k\}$ of k vertex subsets of G , called *domains*.

Question: Does G contain k pairwise internally vertex-disjoint paths P_1, \dots, P_k such that for $i \in [k]$, P_i is an (s_i, t_i) -path with $V(P_i) \subseteq U_i$?

We let $V(\mathcal{X}) := \bigcup_{i \in [k]} \{s_i, t_i\}$. For a set of vertices $S \subseteq V(G)$, let $\lambda: S \rightarrow [k]$ be a labeling. We say that λ is *domain-preserving*, if for each $v \in S$, $v \in U_{\lambda(v)}$. We use domain-preserving labelings in our algorithm later to gradually build paths that only use vertices of the prescribed domains.

Remark 5.1. Let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED DISJOINT PATHS. For ease of exposition, we make the following assumptions. First, we assume that G is connected, since otherwise, we can simply solve the problem on each connected component separately. Furthermore, for each $i \in [k]$, we assume that $\{s_i, t_i\} \subseteq U_i$ and $V(\mathcal{X}) \cap (U_i \setminus \{s_i, t_i\}) = \emptyset$. This way, we ensure directly that no path P_i can use a terminal s_j or t_j ($i \neq j$) as an internal vertex.

Suppose that for some terminal pair (s_i, t_i) , we have that $s_i t_i \in E(G)$. Then, we can use the edge $s_i t_i$ as a path in a solution. Since such a path has no internal vertex, and the SET-RESTRICTED DISJOINT PATHS problem asks for *internally* vertex-disjoint paths, using this edge as the (s_i, t_i) -path cannot create any conflict with any other path. Therefore, we can safely remove the terminal pair (s_i, t_i) from the instance without changing the answer to the problem.

Reduction R.1. Let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED DISJOINT PATHS such that for some $i \in [k]$, $s_i t_i \in E(G)$. Then, reduce this instance to $(G, k - 1, \mathcal{X} \setminus \{(s_i, t_i)\}, \mathcal{U} \setminus \{U_i\})$.

Next, we observe that finding pairwise internally vertex-disjoint paths is equivalent to finding pairwise internally vertex-disjoint *induced* paths. We call a solution consisting of induced paths a *minimal* solution. The following observation is an immediate consequence by the fact that each bag of a partition tree is a clique in the underlying graph.

Observation 5.2. Let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED DISJOINT PATHS such that G is a connected well-partitioned chordal graph. Let P be a path in a minimal solution to $(G, k, \mathcal{X}, \mathcal{U})$, and let B be a bag of the partition tree of G . Then, $|V(P) \cap B| \leq 2$.

Throughout the following, let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED DISJOINT PATHS, where G is a connected well-partitioned chordal graph that is given together with a partition tree \mathcal{T} of G . Based on Observation 5.2, we now describe a marking procedure that marks at most $4k^2$ vertices of each bag $B \in V(\mathcal{T})$ such that if there is a solution to $(G, k, \mathcal{X}, \mathcal{U})$, then there is a solution whose paths use only marked vertices as internal vertices.

Lemma 5.3. *Let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED DISJOINT PATHS, such that G is a well-partitioned chordal graph with partition tree \mathcal{T} , after exhaustive application of Reduction R.1. Then, there is a $\mathcal{O}(k^2 \cdot n)$ time algorithm that computes sets $M_1, \dots, M_k \subseteq V(G)$ such that for all $i \in [k]$, $M_i \subseteq U_i$, and the following hold.*

- For each bag $B \in V(\mathcal{T})$, and each $i \in [k]$, $|B \cap M_i| \leq 4k$.
- Let \mathcal{T}' be the forest in \mathcal{T} induced by all bags with a nonempty intersection with $\bigcup_{i \in [k]} M_i$. Then, \mathcal{T}' has at most $2k$ bags of degree one.
- $(G, k, \mathcal{X}, \mathcal{U})$ is a YES-instance if and only if there exists a minimal solution (P_1, \dots, P_k) such that for each $i \in [k]$, $V(P_i) \subseteq M_i \cup \{s_i, t_i\}$.

Proof. We initialize $M_i := \emptyset$ for all $i \in [k]$. For each $i \in [k]$, we do the following. Let $B_1 B_2 \dots B_\ell$ be the path in \mathcal{T} such that $s_i \in B_1$ and $t_i \in B_\ell$. Then, for each $j \in [\ell - 1]$, we add to M_i a maximal subset of $U_i \cap \text{bd}(B_j, B_{j+1})$ of size at most $2k$, and a maximal subset of $U_i \cap \text{bd}(B_{j+1}, B_j)$ of size at most $2k$. This finishes the construction of the sets M_i , and it is not difficult to see that it can be implemented to run in time $\mathcal{O}(k^2 \cdot n)$.

We prove that M has the claimed properties. The first item is immediate. The second item follows from the observation that \mathcal{T}' consists of the union of k paths in \mathcal{T} . For the third item, suppose $(G, k, \mathcal{X}, \mathcal{U})$ is a YES-instance, and let (P_1, \dots, P_k) be a minimal solution. Suppose that for some bag $B \in V(\mathcal{T})$ and some $i \in [k]$, $V(P_i) \cap B \not\subseteq M_i \cup \{s_i, t_i\}$. Let $B_1 B_2 \dots B_\ell$ be the path in \mathcal{T} from the bag containing s_i to the bag containing t_i . We argue that we may assume that B is a bag on this path. Suppose not, and let $j \in [\ell]$ be such that B_j is the closest bag to B on the path, and let B'_j be the neighboring bag of B_j on the path from B_j to B . Then, P_i must use two vertices from $\text{bd}(B_j, B'_j)$, and hence there is a triangle in P_i , a contradiction with P_i being a path of a minimal solution.

From now on, let $j \in [\ell]$ be such that $B = B_j$, and let $Y = (V(P_i) \cap B_j) \setminus \{s_i, t_i\}$. First, if $j = 1$, then we may assume that there is only one vertex in $y \in Y \setminus M_i$, with $y \in \text{bd}(B_1, B_2)$, and that $s_i \notin \text{bd}(B_1, B_2)$ (otherwise P_i was not an induced path). Since y was not marked, there are $2k$ marked vertices in $\text{bd}(B_1, B_2) \cap M_i$. Since the paths P_j , $j \neq i$, use at most $2(k - 1)$ vertices in total from B by Observation 5.2, there is at least one vertex in $\text{bd}(B_1, B_2) \cap M_i$ that is not used by any other path, call that vertex y' . We replace y by y' in P_i , and maintain the property that

P_i is an induced (s_i, t_i) -path, since y and y' are twins in $G[B_1 \cup B_2]$. A similar argument can be given if $j = \ell$.

Now suppose that $1 < j < \ell$. Again we have that $|Y| \leq 2$. Suppose that $|Y \setminus M_i| = 2$, and let $\{y_1, y_2\} = Y \setminus M_i$. Assume wlog that $y_1 \in \text{bd}(B_j, B_{j-1})$ and that $y_2 \in \text{bd}(B_j, B_{j+1})$. By the same argument as above, there are vertices $y'_1 \in \text{bd}(B_j, B_{j-1}) \cap M_i$ and $y'_2 \in \text{bd}(B_j, B_{j+1}) \cap M_i$ such that neither y'_1 nor y'_2 are used by any other path in the solution. We can replace $\{y_1, y_2\}$ by $\{y'_1, y'_2\}$ in P_i , and P_i remains an (s_i, t_i) -path. If for $r \in [2]$, $y'_r \in \text{bd}(B_j, B_{j-1}) \cap \text{bd}(B_j, B_{j+1})$, then we remove y'_{3-r} from P_i to ensure that P_i remains an induced path. The last case, when $|Y \setminus M_i| = 1$, can be treated with similar arguments and we therefore skip the details.

We have shown how to modify the paths P_1, \dots, P_k such that they remain induced paths, and for all $i \in [k]$, $V(P_i) \subseteq M_i \cup \{s_i, t_i\}$, so the third item follows. \square

Given any bag B , we can immediately observe which paths of a solution need to use some vertices from B as internal vertices. The next definition captures the property of a bag having a set of vertices that can be used as the internal vertices of all paths that need to go through B . Note that in the algorithm of this section, we only need the special case of $[k]$ -feasible bags; however in the algorithm for SET-RESTRICTED TOTALLY DISJOINT PATHS, we need to be able to restrict this definition to a subset of $[k]$ which is why we give it in a more general form here.

Definition 5.4 (I-Feasible Bag). Let $I \subseteq [k]$. Let $B \in V(\mathcal{T})$ be a bag and $M_i \subseteq V(G)$, $i \in I$, be sets of vertices. Then, we say that B is *I-feasible w.r.t. $\{M_i \mid i \in I\}$* , if there is a set $X \subseteq B$ and a labeling $\lambda: X \rightarrow [k]$ such that the following hold. For each $i \in I$ such that B lies on the path from the bag containing s_i to the bag containing t_i in \mathcal{T} , and each neighbor C of B on that path, either $\{s_i, t_i\} \cap \text{bd}(B, C) \neq \emptyset$, or there is a vertex $x_i \in X \cap M_i \cap \text{bd}(B, C)$ such that $\lambda(x_i) = i$. We use the shorthand ‘feasible’ for ‘ $[k]$ -feasible’.

Lemma 5.5. *Let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED DISJOINT PATHS, such that G is a well-partitioned chordal graph with partition tree \mathcal{T} , after exhaustive application of Reduction R.1. Let M_1, \dots, M_k be sets of vertices given by Lemma 5.3. Then, $(G, k, \mathcal{X}, \mathcal{U})$ is a YES-instance if and only if each bag of \mathcal{T} is feasible w.r.t. M_1, \dots, M_k .*

Proof. Suppose that $(G, k, \mathcal{X}, \mathcal{U})$ is a YES-instance. By Lemma 5.3, there is a minimal solution (P_1, \dots, P_k) such that for all $i \in [k]$, $V(P_i) \subseteq M_i \cup \{s_i, t_i\}$. Let $B \in V(\mathcal{T})$ be a bag. Then, we let $X := (B \setminus V(\mathcal{X})) \cap \bigcup_{i \in [k]} V(P_i)$, and $\lambda: X \rightarrow [k]$ be such that for all $x \in X$, $\lambda(x) = i$ if $x \in V(P_i)$. Then, it is not difficult to see that X and λ show that B is a feasible bag.

For the other direction, suppose that each bag of \mathcal{T} is feasible w.r.t. M_1, \dots, M_k . Let $i \in [k]$, and denote by $B_1 B_2 \dots B_\ell$ the path in \mathcal{T} such that $s_i \in B_1$ and $t_i \in B_\ell$. Then, for $j \in [\ell]$, let X_j and λ_j be the subset of B_j and its labeling, respectively, that show that B_j is feasible. Then, for each $j \in [\ell - 1]$, there is a vertex $x_j \in \text{bd}(B_j, B_{j+1})$ such that $\lambda_j(x_j) = i$, and a vertex $y_{j+1} \in \text{bd}(B_{j+1}, B_j)$ such that $\lambda_{j+1}(y_{j+1}) = i$. Then the sequence

$$s_i, x_1, y_2, x_2, y_3, x_3, \dots, y_{\ell-1}, x_{\ell-1}, y_\ell, t_i$$

(where it may happen that $s_i = x_1$ or $y_\ell = t_i$ or for some $j \in [\ell]$, $x_j = y_j$) can be used to obtain an (s_i, t_i) -path in G , and we can take any induced subpath of it to obtain an induced (s_i, t_i) -path P_i in G . Since all labelings λ_j are domain-preserving, we have that $V(P_i) \subseteq U_i \cup \{s_i, t_i\}$. \square

Theorem 5.6. *There is an algorithm that solves each instance $(G, k, \mathcal{X}, \mathcal{U})$ of SET-RESTRICTED DISJOINT PATHS where G is a well-partitioned chordal graph given along with a partition tree \mathcal{T} , in time $2^{\mathcal{O}(k \log k)} \cdot n$.*

Proof. The algorithm works as follows.

Step 1. Apply Reduction R.1 exhaustively.

Step 2. Use Lemma 5.3 to obtain the sets M_1, \dots, M_k of marked vertices.

Step 3. For each bag $B \in V(\mathcal{T})$, check if B is feasible w.r.t. M_1, \dots, M_k . If all bags are feasible w.r.t. M_1, \dots, M_k , then report YES, and if not, report NO.

Correctness follows from Lemma 5.5. For the runtime, we have that the marking in Step 2 can be done in time $\mathcal{O}(k^2 \cdot n)$ by Lemma 5.3. Let $\mathcal{M} := \bigcup_{i \in [k]} M_i$. Since for each bag $B \in V(\mathcal{T})$, $|B \cap \mathcal{M}| = \mathcal{O}(k^2)$ by Lemma 5.3, we can try all $2^{\mathcal{O}(k \log k)}$ sets X and all $2^{\mathcal{O}(k \log k)}$ labelings of each such X to check if B is feasible. Therefore, the feasibility of B can be checked in time $2^{\mathcal{O}(k \log k)}$. Since there are at most n bags in \mathcal{T} , we have that the total runtime of verifying feasibility is $2^{\mathcal{O}(k \log k)} \cdot n$. \square

5.2 Set-Restricted Totally Disjoint Paths

As discussed above, we now adapt the algorithm of Theorem 5.6 to find *totally internally disjoint paths* instead, as it is done e.g. in the setting of finding topological minors in a graph [18]. Specifically, we are dealing with the following problem.

SET-RESTRICTED TOTALLY DISJOINT PATHS parameterized by k

Input: A graph G , a set $\mathcal{X} = \{(s_1, t_1), \dots, (s_k, t_k)\}$ of k pairs of vertices of G , called *terminals*, a set $\mathcal{U} = \{U_1, \dots, U_k\}$ of k vertex subsets of G , called *domains*.

Question: Does G contain k pairwise distinct and internally vertex-disjoint paths P_1, \dots, P_k such that for $i \in [k]$, P_i is an (s_i, t_i) -path with $V(P_i) \subseteq U_i$?

Recall that $V(\mathcal{X}) := \bigcup_{i \in [k]} \{s_i, t_i\}$, and that for a set of vertices $S \subseteq V(G)$, a labeling $\lambda: S \rightarrow [k]$ is called *domain-preserving*, if for each $v \in S$, $v \in U_{\lambda(v)}$. Furthermore, as in Remark 5.1, we assume that in any instance we consider, the input graph G is connected and for all $i \in [k]$, $\{s_i, t_i\} \subseteq U_i$ and $V(\mathcal{X}) \cap (U_i \setminus \{s_i, t_i\}) = \emptyset$.

Following the notation introduced in [18], we call an edge $xy \in E(G)$ *heavy* if for some $w \geq 2$, there are pairwise distinct indices i_1, \dots, i_w such that for each $j \in [w]$, $\{x, y\} = \{s_{i_j}, t_{i_j}\}$. In that case, we call w the *weight* of the edge xy , and we say that the indices i_1, \dots, i_w *weigh down on* xy . For each such heavy edge e , we use the notation $I(e) := \{i_1, \dots, i_w\}$. We say that paths P_{i_1}, \dots, P_{i_w} *satisfy* e if there is precisely one $a \in [w]$ such that P_{i_a} consists of the edge e , and for each $b \in [w] \setminus \{a\}$, P_{i_b} is a length-2 (x, y) -path. Furthermore, we call an index i a *heavy index* if $s_i t_i$ is a heavy edge. We say that an index is *light* if it is not heavy.

First, we observe that if for some edge $xy \in E(G)$, there is *precisely one* i such that $\{x, y\} = \{s_i, t_i\}$, then we can again remove the terminal pair (s_i, t_i) from the instance without changing the answer to the problem: We can always use the edge xy as the path connecting s_i to t_i . (Note that the following is a weaker form of Reduction R.1, and that it does not apply to heavy edges.)

Reduction R.2. Let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED TOTALLY DISJOINT PATHS such that for some edge $xy \in E(G)$ there is precisely one $i \in [k]$ such that $\{x, y\} = \{s_i, t_i\}$. Then, reduce this instance to $(G, k - 1, \mathcal{X} \setminus \{(s_i, t_i)\}, \mathcal{U} \setminus \{U_i\})$.

Due to Reduction R.2, we may from now on assume that for each $i \in [k]$, either $s_i t_i \notin E(G)$, or $s_i t_i$ is a heavy edge. The former we can handle as in the algorithm of Theorem 5.6, and we explain how to deal with the latter. The existence of a heavy edge rules out the approach of looking for *minimal* solutions: if xy is a heavy edge of weight w , then each solution contains $w - 1$ paths between x and y that are not induced due to the existence of the edge xy .

However, a slight modification of this approach works. We base our marking procedure on the existence of *minimum* solutions, i.e. solutions (P_1, \dots, P_k) such that there is no other solution that contains fewer edges: for all solutions (P'_1, \dots, P'_k) , it holds that $\sum_{i \in [k]} |E(P_i)| \leq \sum_{i \in [k]} |E(P'_i)|$. As shown in the following lemma due to Heggenes et al. [18], minimum solutions have a very restricted structure in chordal graphs as well.¹

Lemma 5.7 (Cf. Lemma 2 in [18]). *Let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED DISJOINT PATHS, such that G is a chordal graph. If (P_1, \dots, P_k) is a minimum solution to $(G, k, \mathcal{X}, \mathcal{U})$, then for each $i \in [k]$, precisely one of the following holds.*

- P_i is an induced path;
- P_i is a path of length two, and there exists some $j \in [k]$ such that P_j is of length one and $\{s_i, t_i\} = \{s_j, t_j\}$.

By the second part of the previous lemma we know that in each minimum solution, for each heavy edge e , there is a collection of paths of length at most 2 satisfying e . Hence, to accommodate the heavy edges in our marking scheme, it is enough to consider common neighbors of the endpoints of heavy edges. If our input graph G is a well-partitioned chordal graph, this gives us a lot of structure we can exploit.

We adapt the marking procedure of SET-RESTRICTED DISJOINT PATHS algorithm as follows.

Lemma 5.8. *Let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED TOTALLY DISJOINT PATHS, such that G is a well-partitioned chordal graph with partition tree \mathcal{T} , after exhaustive application of Reduction R.2. Then, there is an $\mathcal{O}(k^2 \cdot n)$ time algorithm that computes sets $M_1, \dots, M_k \subseteq V(G)$ such that for all $i \in [k]$, $M_i \subseteq U_i$, and the following hold. Let I denote the light indices.*

- For each heavy index $i \in [k] \setminus I$, $|M_i| \leq 2k$.
- For each bag $B \in V(\mathcal{T})$, and each light index $i \in I$, $|B \cap M_i| \leq 4k$.
- Let \mathcal{T}' be the forest in \mathcal{T} induced by all bags with a nonempty intersection with $\bigcup_{i \in I} M_i$. Then, \mathcal{T}' has at most $2 \cdot |I|$ bags of degree one.
- $(G, k, \mathcal{X}, \mathcal{U})$ is a YES-instance if and only if there exists a minimal solution (P_1, \dots, P_k) such that for each $i \in [k]$, $V(P_i) \subseteq M_i \cup \{s_i, t_i\}$.

¹Note that Lemma 2 in [18] is proved for the DISJOINT PATHS problem; however, the proof goes through for SET-RESTRICTED TOTALLY DISJOINT PATHS as well.

Proof. For each light index, we proceed as in Lemma 5.3. Let xy be a heavy edge. There are two cases we need to consider. First, if there are bags $B_1, B_2 \in V(\mathcal{T})$ such that $x \in \text{bd}(B_1, B_2)$ and $y \in \text{bd}(B_2, B_1)$, then the only vertices that can be used to form a length-2 (x, y) -path are in $\text{bd}(B_1, B_2) \cup \text{bd}(B_2, B_1)$. Hence, for each index i weighing down on xy , we let M_i be a maximal subset of $U_i \cap (\text{bd}(B_1, B_2) \cup \text{bd}(B_2, B_1))$ of size at most $2k$. Second, there is a bag B such that $\{x, y\} \subseteq B$. Then, the remaining vertex on a path corresponding to the terminal pair (x, y) has to be contained in B or in a neighbor bag C of B such that x and y are both in the boundary of B to C . Therefore, we let M_i be a maximal subset of

$$U_i \cap \left(B \cup \bigcup_{C \in N_{\mathcal{T}}(B), \{x, y\} \subseteq \text{bd}(B, C)} \text{bd}(C, B) \right)$$

of size at most $2k$. We can argue in the same way as in the proof of Lemma 5.3 that this is correct. \square

Theorem 5.9. *There is an algorithm that solves each instance $(G, k, \mathcal{X}, \mathcal{U})$ of SET-RESTRICTED TOTALLY DISJOINT PATHS where G is a well-partitioned chordal graph given along with a partition tree \mathcal{T} , in time $2^{\mathcal{O}(k \log k)} \cdot n$.*

Proof. Let $I \subseteq [k]$ be the light indices and $J \subseteq [k]$ the heavy indices. The algorithm works almost the same way as the one for SET-RESTRICTED DISJOINT PATHS, with an additional guessing stage of the vertices satisfying the heavy edges.

Step 1. Apply Reduction R.2 exhaustively.

Step 2. Use Lemma 5.8 to obtain the sets M_1, \dots, M_k of marked vertices.

Step 3. For each subset $Y \subseteq \bigcup_{j \in J} M_j$ of size at most $|J|$ check if the vertices in Y can be used to satisfy all heavy edges.

Step 4. If Step 3 succeeded for the set Y , then for all $i \in I$, let $M'_i := M_i \setminus Y$, and check for each bag $B \in V(\mathcal{T})$ if it is I -feasible w.r.t. $\{M'_i \mid i \in I\}$.

Correctness of this algorithm follows in a similar way as in Theorem 5.6. By Lemma 5.8, if $(G, k, \mathcal{X}, \mathcal{U})$ is a YES-instance, then there is a solution only using vertices from M_1, \dots, M_k . Therefore, if the instance has a solution, then one of the guesses in Step 3 must succeed, in such a way that all bags are I -feasible with respect to $\{M'_i \mid i \in I\}$. (Recall that I are the light indices and that they can be handled in the same way as in the algorithm of Theorem 5.6)

By Lemma 5.8, $|\bigcup_{j \in J} M_j| = \mathcal{O}(k^2)$, so there are $2^{\mathcal{O}(k \log k)}$ choices to try in Step 3. The runtime for Step 4 is again $2^{\mathcal{O}(k \log k)} \cdot n$, therefore the runtime is $2^{\mathcal{O}(k \log k)} \cdot n$. \square

5.3 Set-Restricted Disjoint Connected Subgraphs

In this section we sketch how to solve another related problem, SET-RESTRICTED DISJOINT CONNECTED SUBGRAPHS that was also introduced in [1]. For convenience, we now require the solution to be pairwise *vertex-disjoint*. However, the variants of the problem that ask for internally vertex-disjoint solutions can be handled by similar methods used in the previous sections.

SET-RESTRICTED DISJOINT CONNECTED SUBGRAPHS parameterized by $s := \sum_{i \in [k]} |S_i|$

Input: A graph G , a set $\mathcal{S} = \{S_1, \dots, S_k\}$ of pairwise disjoint vertex sets G called *terminal sets*, a set $\mathcal{U} = \{U_1, \dots, U_k\}$ of k vertex subsets of G , called *domains*.

Question: Does G contain k pairwise vertex-disjoint connected subgraphs F_1, \dots, F_k such that for $i \in [k]$, $S_i \subseteq V(F_i) \subseteq U_i$?

Let $(G, k, \mathcal{S}, \mathcal{U})$ be an instance of SET-RESTRICTED DISJOINT CONNECTED SUBGRAPHS such that G is a well-partitioned chordal graph with partition tree \mathcal{T} . Similarly to before, it is not difficult to see that if $(G, k, \mathcal{S}, \mathcal{U})$ is a YES-instance, then it has a solution that uses at most s non-terminal vertices from each bag B . To adapt the marking procedure for this problem, for each $i \in [k]$, let \mathcal{T}_i be the smallest subtree of \mathcal{T} that contains all bags that have a non-empty intersection with S_i . For each bag $B \in V(\mathcal{T}_i)$, and each $C \in N_{\mathcal{T}_i}(B)$, we mark a maximal subset of $\text{bd}(B, C)$ of size at most s . Let the marked vertices for i , M_i , be the union of all these sets.

Then, if $(G, k, \mathcal{S}, \mathcal{U})$ is a YES-instance, there is a solution such that each F_i uses only vertices from M_i . We can now define a notion of feasibility based on the subtrees \mathcal{T}_i , in analogy with Definition 5.4. Then, checking if each bag is feasible again solves the problem. Since each bag contains at most $\mathcal{O}(s^2)$ marked vertices, we obtain the following runtime bound.

Theorem 5.10. *There is an algorithm that solves each instance $(G, k, \mathcal{X}, \mathcal{U})$ of SET-RESTRICTED DISJOINT CONNECTED SUBGRAPHS where G is a well-partitioned chordal graph given along with a partition tree \mathcal{T} , in time $2^{\mathcal{O}(s \log s)} \cdot n$.*

5.4 Polynomial kernels on split graphs

We observe that Lemmas 5.3 and 5.8 imply polynomial kernels of SET-RESTRICTED DISJOINT PATHS and SET-RESTRICTED TOTALLY DISJOINT PATHS on split graphs.

Corollary 5.11. *The SET-RESTRICTED DISJOINT PATHS and SET-RESTRICTED TOTALLY DISJOINT PATHS problems on split graphs admit kernels on $\mathcal{O}(k^2)$ vertices.*

Proof. Let $(G, k, \mathcal{X}, \mathcal{U})$ be an instance of SET-RESTRICTED DISJOINT PATHS such that G is a split graph, with partition tree \mathcal{T} . Note that \mathcal{T} is a star, and assume that \mathcal{T} is rooted at the center. Then, all leaves of this star contain only one vertex. Moreover, by Lemma 5.3, we can reduce the number of vertices in the center bag to $\mathcal{O}(k^2)$, and we can remove all but $2k$ leaves, without changing the answer to the instance.

Now suppose that $(G, k, \mathcal{X}, \mathcal{U})$ is an instance of SET-RESTRICTED TOTALLY DISJOINT PATHS, and let \mathcal{T} be as above. By Lemma 5.8, we can again reduce the number of vertices in the center bag to $\mathcal{O}(k^2)$, and remove all but $\mathcal{O}(k^2)$ leaves, without changing the answer to the instance. \square

5.5 Polynomial kernel for Disjoint Paths on well-partitioned chordal graphs

We first study the variant of the problem DISJOINT PATHS as defined in the introduction, i.e. the problem that asks for pairwise *internally* vertex-disjoint paths between the terminals. Later, we consider a polynomial kernel for TOTALLY DISJOINT PATHS, which asks for pairwise distinct and internally vertex-disjoint paths.

Towards such a kernel, let (G, k, \mathcal{X}) be an instance of DISJOINT PATHS such that G is a well-partitioned chordal graph with partition tree \mathcal{T} , and let $M_1, \dots, M_k \subseteq V(G)$ be the set of marked vertices due to Lemma 5.3.² Let \mathcal{T}' be the forest in \mathcal{T} given by Lemma 5.3. If (G, k, \mathcal{X}) is a YES-instance, then there is a solution only using vertices that are contained in bags of \mathcal{T}' . We can therefore remove all vertices in bags $V(\mathcal{T}) \setminus V(\mathcal{T}')$, and continue working with the subgraph of G induced by the bags in \mathcal{T}' .

Observation 5.12. Let (G, k, \mathcal{X}) be an instance of DISJOINT PATHS such that each connected component of G is a well-partitioned chordal graph. We may assume that the forest \mathcal{F} , consisting of the partition trees of the connected components of G , has at most $2k$ bags of degree one.

For convenience, we call \mathcal{F} as in the previous definition a *partition forest*. Suppose from now on that (G, k, \mathcal{X}) is as asserted by Observation 5.12. Unless the number of vertices in G is polynomial in k , \mathcal{F} contains many bags that are of degree two, and contain no terminal. We now introduce two new reduction rules that show that either we can conclude that we are dealing with a NO-instance, or we can remove them.

Reduction R.3. Let (G, k, \mathcal{X}) be an instance of DISJOINT PATHS such that the components of G are well-partitioned chordal with partition forest \mathcal{F} . Let $B \in V(\mathcal{T})$ such that $B \cap V(\mathcal{X}) = \emptyset$ and $\deg_{\mathcal{F}}(B) = 2$. Let A and C be the two neighbors of B in \mathcal{F} . Let $I \subseteq [k]$ be the indices such that B lies on the path of the bag containing s_i to the bag containing t_i . If $|\text{bd}(B, A)| < |I|$ or $|\text{bd}(B, C)| < |I|$, then reduce (G, k, \mathcal{X}) to a trivial NO-instance.

Lemma 5.13. *If the conditions of Reduction R.3 are satisfied, then (G, k, \mathcal{X}) is a NO-instance.*

Proof. For each $i \in I$, the intersection of the vertices of the path P_i in a solution with B must be nonempty. Moreover, P_i must use a vertex from both $\text{bd}(B, C)$ and $\text{bd}(B, A)$ (which could be the same vertex). However, by the pigeonhole principle, there cannot be a solution to (G, k, \mathcal{X}) since at least one of $\text{bd}(B, A)$ and $\text{bd}(B, C)$ has less than $|I|$ vertices. \square

We now show the orthogonal to Reduction R.3: if both boundaries of B have at least $|I|$ vertices, then we can remove B and all its vertices. Note that the statement of the following reduction also tells us how to obtain a partition tree of the reduced graph.

Reduction R.4. Let (G, k, \mathcal{X}) be an instance of DISJOINT PATHS such that the components of G are well-partitioned chordal with partition forest \mathcal{F} . Let $B \in V(\mathcal{T})$ such that $B \cap V(\mathcal{X}) = \emptyset$ and $\deg_{\mathcal{F}}(B) = 2$. Let A and C be the two neighbors of B in \mathcal{F} . Let $I \subseteq [k]$ be the indices such that B lies on the path of the bag containing s_i to the bag containing t_i . If $|\text{bd}(B, A)| \geq |I|$ and $|\text{bd}(B, C)| \geq |I|$, then reduce (G, k, \mathcal{X}) to (G', k, \mathcal{X}) , where G' is obtained from G by removing B and making all vertices in $\text{bd}(A, B)$ adjacent to all vertices in $\text{bd}(C, B)$.

Lemma 5.14. *Reduction R.4 is safe, i.e. under its stated conditions, (G, k, \mathcal{X}) is a YES-instance if and only if (G', k, \mathcal{X}) is a YES-instance.*

Proof. Suppose (G, k, \mathcal{X}) is a YES-instance and let P_1, \dots, P_k be one of its solutions. For each $i \in I$, we observe that P_i contains a vertex from $\text{bd}(A, B)$, say a_i , and a vertex from $\text{bd}(C, B)$, say c_i , such that both a_i and c_i have a neighbor from B on the path P_i . Let P'_i be the path obtained

²Where for each $i \in [k]$, we let $U_i = V(G)$.

from P_i by removing the vertices $V(P_i) \cap B$, and making a_i and c_i adjacent. For each $j \in [k] \setminus I$, let $P'_j := P_j$. It is not difficult to verify that P'_1, \dots, P'_k is a solution to (G', k, \mathcal{X}) .

Conversely, suppose that (G', k, \mathcal{X}) is a YES-instance, and let P'_1, \dots, P'_k be one of its solutions, and let \mathcal{F}' be the partition forest of G' . Denote by A and C the bags in \mathcal{F}' that correspond to the neighbors of B in \mathcal{F} . Note that A and C are adjacent in \mathcal{T}' by construction. There is a pair of vertices $a_i \in \text{bd}_{\mathcal{F}'}(A, C)$, $c_i \in \text{bd}_{\mathcal{F}'}(C, A)$ such that P'_i contains the edge $a_i c_i$. Let $b_i^a \in \text{bd}_{\mathcal{F}}(B, A)$ and $b_i^c \in \text{bd}_{\mathcal{F}}(B, C)$ (possibly $b_i^a = b_i^c$). Then, the path P_i obtained from P'_i by replacing the edge $a_i c_i$ with the path $a_i b_i^a b_i^c c_i$ is an (s_i, t_i) -path in G . Moreover, since $|\text{bd}(B, A)| \geq |I|$ and $|\text{bd}(B, C)| \geq |I|$, we can assign such vertices to each of the paths P'_i , $i \in I$. Finally, for $j \in [k] \setminus I$, let $P_j := P'_j$. Then, P_1, \dots, P_k is a solution to (G, k, \mathcal{X}) . \square

We wrap up.

Theorem 5.15. *DISJOINT PATHS on well-partitioned chordal graphs parameterized by k admits a kernel on $\mathcal{O}(k^3)$ vertices.*

Proof. Let (G, k, \mathcal{X}) be an instance of DISJOINT PATHS such that G is well-partitioned chordal, and let (G, k, \mathcal{X}) be as asserted by Observation 5.12. Apply Reduction R.3 exhaustively. If at some stage the reduction returned a trivial NO-instance, then we conclude that (G, k, \mathcal{X}) is a NO-instance, which is correct by Lemma 5.13. Now apply Reduction R.4 exhaustively, and reuse (G, k, \mathcal{X}) to denote the resulting instance, and let \mathcal{T} be the partition tree of G . We count the number of bags in \mathcal{T} . By Observation 5.12, \mathcal{T} has at most $2k$ leaf nodes, therefore it also has at most $2k$ internal nodes of degree at least 3. Moreover, \mathcal{T} has at most $2k$ nodes of degree two – by Reductions R.3 and R.4, the only degree two nodes in \mathcal{T} contain terminal vertices. Therefore, $|V(\mathcal{T})| = \mathcal{O}(k)$, and since each bag contains at most $\mathcal{O}(k^2)$ vertices by Lemma 5.3, G contains at most $\mathcal{O}(k^3)$ vertices. \square

We now show that TOTALLY DISJOINT PATHS also admits a polynomial kernel on well-partitioned chordal graphs. Suppose we have an instance (G, k, \mathcal{X}) of TOTALLY DISJOINT PATHS such that G is well-partitioned chordal with partition tree \mathcal{T} . We reduce G as follows. We apply Lemma 5.8, and let \mathcal{T}' be the forest in \mathcal{T} that is described there. (Note that the construction of \mathcal{T}' is based only on the *light* indices.) It is clear that in our kernel, it suffices to take \mathcal{T}' , and the union of M_i over all *heavy* indices i . We reduce \mathcal{T}' according to Reductions R.3 and R.4. If no trivial NO-instance was returned, then we return the resulting subgraph of G as our kernelized instance. We can apply roughly the same argument as given in the proof of Theorem 5.15 to conclude that the subgraph of G induced by vertices in the bags of \mathcal{T}' after reduction, has at most $\mathcal{O}(k^3)$ vertices. Since the number of vertices that have been marked for heavy edges is at most $\mathcal{O}(k^2)$ in total, we have the following theorem.

Theorem 5.16. *TOTALLY DISJOINT PATHS on well-partitioned chordal graphs parameterized by k admits a kernel on $\mathcal{O}(k^3)$ vertices.*

Note that in resulting kernelized instance, the graph may not be connected. However, it is not difficult to see that with a slight modification of the above described procedure, one can obtain a connected graph in the kernel.

6 Conclusions

In this paper, we introduced the class of *well-partitioned chordal graphs*, a subclass of chordal graphs that generalizes split graphs. We provided a characterization by a set of forbidden induced

subgraphs which also gave a polynomial-time recognition algorithm. We showed that several parameterized variants and generalizations of the DISJOINT PATHS problem that on chordal graphs are only known to be in XP, are in FPT on well-partitioned chordal graphs. These results in some cases implied polynomial kernels on split and well-partitioned chordal graphs. It would be interesting to see for which problems well-partitioned chordal graphs can be used to narrow down complexity gaps for problems that are hard on chordal and easy on split graphs, or for which problems that are easy on split graphs and whose complexity is open on chordal graphs, well-partitioned chordal graphs can be used to obtain partial (positive) results.

Another typical characterization of (subclasses of) chordal graphs is via vertex orderings. For instance, chordal graphs are famously characterized as the graphs admitting perfect elimination orderings [13]. It would be interesting to see if well-partitioned chordal graphs admit a concise characterization in terms of vertex orderings as well. While the degree of the polynomial in the runtime of our recognition algorithm is moderate, our algorithm does not run in linear time. We therefore ask if it is possible to recognize well-partitioned chordal graphs in linear time; and note that a characterization in terms of vertex orderings can be a promising step in this direction.

References

- [1] Rémy Belmonte, Petr A. Golovach, Pinar Heggernes, Pim van ’t Hof, Marcin Kamiński, and Daniël Paulusma. Detecting fixed patterns in chordal graphs in polynomial time. *Algorithmica*, 69(3):501–521, 2014.
- [2] Alan A. Bertossi. Dominating sets for split and bipartite graphs. *Information processing letters*, 19(1):37–40, 1984.
- [3] Hans L. Bodlaender and Klaus Jansen. On the complexity of the maximum cut problem. *Nordic Journal of Computing*, 7(1):14–31, 2000.
- [4] Hans L. Bodlaender, Stéphan Thomassé, and Anders Yeo. Kernel bounds for disjoint cycles and disjoint paths. *Theoret. Comput. Sci.*, 412(35):4570–4578, 2011.
- [5] Andreas Brandstädt, Feodor F. Dragan, Hoàng-Oanh Le, and Van Bang Le. Tree spanners on chordal graphs: complexity and algorithms. *Theoret. Comput. Sci.*, 310(1-3):329–354, 2004.
- [6] Andreas Brandstädt, Van Bang Le, and Jeremy P. Spinrad. *Graph Classes: A Survey*. SIAM Monographs in Discrete Mathematics and Applications, 1999.
- [7] Yi-Wu Chang, Michael S. Jacobson, Clyde L. Monma, and Douglas B. West. Subtree and substar intersection numbers. *Discrete Appl. Math.*, 44(1-3):205–220, 1993.
- [8] D. G. Corneil and Y. Perl. Clustering and domination in perfect graphs. *Discrete Appl. Math.*, 9(1):27–39, 1984.
- [9] Marek Cygan, Fedor V. Fomin, Łukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michał Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.
- [10] Rodney G. Downey and Michael R. Fellows. *Fundamentals of Parameterized Complexity*. Texts in Computer Science. Springer, 2013.

- [11] Paola Festa, Panos M. Pardalos, and Mauricio G. C. Resende. Feedback set problems. In *Handbook of combinatorial optimization, Supplement Vol. A*, pages 209–258. Kluwer Acad. Publ., Dordrecht, 1999.
- [12] Fedor V. Fomin, Daniel Lokshtanov, Saket Saurabh, and Meirav Zehavi. *Kernelization*. Cambridge University Press, 2019.
- [13] Delbert Fulkerson and Oliver Gross. Incidence matrices and interval graphs. *Pacific Journal of Mathematics*, 15(3):835–855, 1965.
- [14] Alan George, John R. Gilbert, and Joseph W.H. Liu. *Graph theory and sparse matrix computation*, volume 56. Springer Science & Business Media, 2012.
- [15] Martin Charles Golumbic. *Algorithmic Graph Theory and Perfect Graphs*, volume 57 of *Annals of Discrete Mathematics*. North-Holland Publishing Co., 2004.
- [16] Jens Gustedt. On the pathwidth of chordal graphs. *Discrete Applied Mathematics*, 45(3):233–248, 1993.
- [17] Frédéric Havet, Claudia Linhares Sales, and Leonardo Sampaio. b-coloring of tight graphs. *Discrete Applied Mathematics*, 160(18):2709–2715, 2012.
- [18] Pinar Heggernes, Pim van't Hof, Erik Jan van Leeuwen, and Reza Saei. Finding disjoint paths in split graphs. *Theory of Computing Systems*, 57(1):140–159, 2015.
- [19] Felix Joos. A characterization of substar graphs. *Discrete Appl. Math.*, 175:115–118, 2014.
- [20] Frank Kammer and Torsten Tholey. The k-disjoint paths problem on chordal graphs. In *WG 2009*, pages 190–201. Springer, 2009.
- [21] Richard M. Karp. On the computational complexity of combinatorial problems. *Networks*, 5(1):45–68, 1975.
- [22] Ken-ichi Kawarabayashi, Yusuke Kobayashi, and Bruce Reed. The disjoint paths problem in quadratic time. *Journal of Combinatorial Theory, Series B*, 102(2):424–435, 2012.
- [23] Stefan Mengel. Lower bounds on the mim-width of some graph classes. *Discrete Applied Mathematics*, 248:28–32, 2018.
- [24] Haiko Müller. Hamiltonian circuits in chordal bipartite graphs. *Discrete Mathematics*, 156(1-3):291–298, 1996.
- [25] Stavros D. Nikolopoulos and Leonidas Palios. Detecting holes and antiholes in graphs. *Algorithmica*, 47(2):119–138, 2007.
- [26] Neil Robertson and P. D. Seymour. Graph minors. XIII. The disjoint paths problem. *J. Combin. Theory Ser. B*, 63(1):65–110, 1995.
- [27] Donald J. Rose, R. Endre Tarjan, and George S. Lueker. Algorithmic aspects of vertex elimination on graphs. *SIAM J. Comput.*, 5(2):266–283, 1976.

- [28] Charles Semple and Mike Steel. *Phylogenetics*, volume 24 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2003.
- [29] Ana Silva. Graphs with small fall-spectrum. *Discrete Applied Mathematics*, 254:183–188, 2019.
- [30] Lieven Vandenberghe and Martin S. Andersen. Chordal graphs and semidefinite optimization. *Foundations and Trends in Optimization*, 1(4):241–433, 2015.
- [31] Martin Vatshelle. *New Width Parameters of Graphs*. PhD thesis, University of Bergen, Norway, 2012.
- [32] Rémi Watrigant, Marin Bougeret, and Rodolphe Giroudeau. Approximating the Sparsest k -Subgraph in chordal graphs. *Theory Comput. Syst.*, 58(1):111–132, 2016.