

Extension of the synchrotron radiation of electrons to very high energies in clumpy environments

Dmitry Khangulyan*

Department of Physics, Rikkyo University, Nishi-Ikebukuro 3-34-1, Toshima-ku, Tokyo 171-8501, Japan

Felix Aharonian

Dublin Institute for Advanced Studies, School of Cosmic Physics, 31 Fitzwilliam Place, Dublin 2, Ireland

Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany and

National Research Nuclear University (MEPHI), Kashirskoje shosse, 31, Moscow 115409, Russia

Carlo Romoli

Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

Andrew Taylor

DESY, D-15738 Zeuthen, Germany

(Dated: March 3, 2020)

The synchrotron cooling of relativistic electrons is one of the most effective radiation mechanisms in astrophysics. It not only accompanies the process of particle acceleration but also has feedback on the formation of the energy distribution of the parent electrons. The radiative cooling time of electrons decreases with energy as $t_{\text{SYN}} \propto 1/E$; correspondingly the overall radiation efficiency increases with energy. On the other hand, this effect strictly limits the maximum energy of individual photons. Even in the so-called extreme accelerators, where the acceleration proceeds at the highest possible rate, $t_{\text{ACC}} = E/eBc$, the synchrotron radiation cannot extend well beyond the characteristic energy determined by the electron mass and the fine-structure constant: $h\nu^{\text{MAX}} \sim m_e c^2/\alpha \sim 100 \text{ MeV}$. In a highly turbulent medium with a broad magnetic field distribution, this limit is shifted towards higher energies. However, for any realistic distribution of the magnetic field, it cannot significantly exceed $h\nu^{\text{MAX}}$. In this paper, we propose a model in which the formation of synchrotron radiation takes place in compact magnetic blobs located inside the particle accelerator. We develop a formalism for calculations of synchrotron radiation emerging from such systems. We demonstrate that for certain combinations of parameters characterizing the accelerator and the magnetic blobs, the synchrotron radiation can extend into the very high energy (VHE) domain, $h\nu^{\text{MAX}} \geq 100 \text{ GeV}$. We briefly outline the possible astrophysical implications of the results in the context of VHE radiation of blazars and gamma-ray bursts.

I. INTRODUCTION

Acceleration of non-thermal particles in various astrophysical environments requires the presence of electric field. If high-energy particles are accelerated in a region with a typical magnetic field, B_0 , then the time, required for the particle to gain energy E , is $t_{\text{ACC}} = \eta E/(eB_0 c)$. Here, η is a dimensionless parameter, namely the ratio of the magnetic to the accelerating electric field, $\eta = B_0/\mathcal{E}_0$. If the acceleration proceeds in a MHD flow, then $\mathcal{E}_0 < B_0$ and the acceleration efficiency is constrained by $\eta > 1$. The acceleration is efficient only in the energy range where the acceleration time is shorter than the cooling time. For the case where synchrotron losses dominate, the cooling rate is $t_{\text{SYN}} = 6\pi m_e^2 c^3/(\sigma_T B_0^2 E)$ (here m_e , c , and σ_T are electron mass, speed of light, and Thomson cross-section, respectively). The electrons are subsequently accelerated up to a maximum energy $E_e^{\text{MAX}} = \sqrt{6\pi e m_e^2 c^4/(\sigma_T \eta B_0)}$. Correspondingly, the electrons which achieve this energy, produce synchrotron

photons of energy $h\nu^{\text{MAX}} \sim m_e c^2/(\alpha \eta) \sim 100\eta^{-1} \text{ MeV}$ [1] (here α is the fine-structure constant).

In the case of stochastic particle acceleration, the spectrum of electrons does not stop abruptly but extends beyond E_e^{MAX} , although with a rather sharp cutoff shape, e.g., exponential or faster [2]. Typically, the maximum of the spectral energy distribution (SED), νF_ν , appears at the energy exceeding $h\nu^{\text{MAX}}$. However, for any realistic spectrum of electrons, the shift does not exceed a factor of a few.

The sharp decrease in the synchrotron spectrum can be altered, however, in a medium with inhomogeneous magnetic fields. For the case of highly turbulent magnetic fields whose correlation length, λ_B , is small compared to the electron's nonrelativistic Larmor radius, $r_L = m_e c^2/eB_0$, the radiation proceeds in the so-called jitter regime. In this case, the peak in the radiation spectrum is shifted to higher frequencies by the factor of r_L/λ_B [3]. Thus, for scenarios with extremely small-scale turbulence, one can formally expect radiation in the multi-GeV or higher energies. However, the plasma physics community is quite pessimistic concerning the condition of $\lambda_B \ll r_L$, and correspondingly considers the

* d.khangulyan@rikkyo.ac.jp

realization of magnetobremstrahlung in the jitter regime as a highly unlikely process. In principle, the presence of large scale turbulence, $\lambda_B > r_L$, can also have an impact on the SED of synchrotron radiation. In particular, a broad power-law distribution of the magnetic field strength can result in a rather long power-law tail of radiation beyond $h\nu^{\text{MAX}}$ [3]. However, for more realistic magnetic field strength distributions, e.g. a Gaussian type distribution, the shift of the synchrotron peak is rather mild [4].

In this paper, we explore a new scenario for boosting the energy range of the synchrotron radiation utilising the structure of the ambient magnetic field in the acceleration region. The principle behind our proposed scheme is the separation of the region of the electron acceleration and the region(s) of effective synchrotron radiation. More specifically, we consider a scenario in which the acceleration region contains a large number of compact magnetic field condensations (blobs), whose collective volume filling factor is small. Provided that the particle-blob collision time is long compared to the acceleration time, these blobs have only a minor impact on the acceleration process. On the other hand, high-energy particles that eventually collide with these blob regions give rise to a synchrotron radiation component which can extend well beyond 100 MeV. Throughout the discussion, we do not specify the acceleration mechanism, as well as the nature of the compact magnetic condensations, but focus on the propagation and radiation of the accelerated electrons colliding with the dense magnetic blobs.

II. TOY MODEL: MATHEMATICAL FORMULATION

Although the postulation of magnetic field blobs inside the particle accelerator implies that we deal with a multi-zone scenario, mathematically one can describe the problem within a one-zone model. In this case, the description of the process is similar to the so-called *leaky-box approximation* which have been successfully applied to Galactic cosmic rays:

$$\frac{\partial n}{\partial t} + \frac{\partial(\dot{E}n)}{\partial E} + \frac{n}{\tau} = q(E). \quad (1)$$

Here $\dot{E}(E) = -E/t_{\text{SYN}}(E, B_0)$ is the energy loss rate of electrons in the main (acceleration) zone, and τ is the particle – magnetic blob collision time. Within the standard treatment, these terms are responsible for the continuous and catastrophic losses of electrons, respectively. The function $q(E)$ accounts for a phenomenological description of the acceleration process. Eq. (1) is valid for particles with energy $E > E_*$; the boundary energy is obtained from the solution of the equation:

$$t_{\text{SYN}}(E_*, B_*) = t_*, \quad (2)$$

where B_* is the magnetic field and t_* is the electron confinement time in the blobs. The latter generally depends

on the particle energy. Eq. (1) is derived in Appendix A. The relevant time-scales are shown in Fig. 1. We assume that in absence of magnetic blobs the injection spectrum contains a cutoff at energy E^{MAX} which is determined from the condition $t_{\text{ACC}}(E^{\text{MAX}}) = t_{\text{SYN}}(E^{\text{MAX}}, B_0)$. We define this timescale t^{MIN} . If the collision time is long compared to the acceleration time up to energy E^{MAX} , i.e.

$$\tau > t^{\text{MIN}}, \quad (3)$$

then the impact of the magnetic blobs on the maximum energy of accelerated electrons can be ignored. Additionally, it is also helpful to define another timescale $t^{\text{MAX}} = t_{\text{SYN}}(E_*, B_0)$. The spectral shape of electrons in the cutoff region depends on the specific acceleration process. Below, we assume that $q \propto E^{-\alpha} \exp[-(E/E^{\text{MAX}})^\beta]$, where for simplicity, but without loss of generality, we adopt $\alpha = 2$ and $\beta = 1$.

In the fast cooling regime, the solution of Eq. (1) is

$$n(E, t) = -\frac{1}{\dot{E}} \int_E^\infty dE' q(E') \exp\left[-\frac{\tau_{\text{EFF}}(E, E')}{\tau}\right], \quad (4)$$

where the auxiliary function τ_{EFF} is

$$\tau_{\text{EFF}}(E, E') = -\int_E^{E'} \frac{dE''}{\dot{E}(E'')} = \frac{6\pi m_e^2 c^3}{(\sigma_T B_0^2)} \left(\frac{1}{E} - \frac{1}{E'}\right). \quad (5)$$




FIG. 1. Characteristic time scales: the acceleration time in the main (acceleration) zone, t_{ACC} , (red line); the weak field synchrotron cooling time in the main zone, $t_{\text{SYN}}(E, B_0)$, (blue dashed line); the strong field synchrotron cooling time in the blobs, $t_{\text{SYN}}(E, B_*)$, (blue solid line); the confinement time of the electron in the blobs, t_* , (cyan line). Four characteristic collision times are considered: the purple curve for $\tau = t_*^{\text{MAX}}$, the green curve for $\tau = \sqrt{t_*^{\text{MIN}} t_*^{\text{MAX}}}$, the light blue curve for $\tau = t_*^{\text{MIN}}$, and the orange curve for $\tau < t_*^{\text{MIN}}$. Note, the timescales t_*^{MIN} and t_*^{MAX} are defined as $t_{\text{SYN}}(E^{\text{MAX}}, B_0)$ and $t_{\text{SYN}}(E_*, B_0)$, respectively.

Two distinct components of synchrotron emission are produced in this scenario. In the main zone, the synchrotron radiation produced by electrons with energy distribution given by Eq. (4), takes a spectral shape of the form

$$L_{\nu,0} = \int P_{\nu}(B_0, E)n(E)dE, \quad (6)$$

where P_{ν} is the power per unit frequency emitted by an electron with energy E (see Appendix B). The electrons that enter the blob, with its much stronger magnetic field, quickly emit their energy at a rate $1/\tau$. The time-averaged energy spectrum of their synchrotron radiation produced in the magnetic blobs is

$$L_{\nu,*} = \frac{1}{\tau} \int \left[\int_0^{t_*} dt P_{\nu}(B_*, E(t)) \right] n(E)dE, \quad (7)$$

where $E(t) = E/(1 + t/t_{\text{syn}}(E, B_*))$. In Appendix B we derive a simple analytic approximation that describes the emission spectrum produced in the blobs.

As demonstrated in Fig. 1, for the condition given by Eq. (3) two different scenarios exist: $\tau \geq t^{\text{MAX}}$ and $t^{\text{MIN}} < \tau < t^{\text{MAX}}$. In the first of these regimes, the presence of magnetic field blobs has a minor impact on the particle spectrum, i.e., $\tau \gg \tau_{\text{EFF}}$. In the second regime the particle spectrum is deformed by interactions with the magnetic blobs, although the maximum energy of the particle spectrum remains unaffected.

In Figs. 2 and 3, we show the corresponding SEDs produced in four different collision timescale scenarios illustrated in Fig. 1. In these figures two different strengths of the blob magnetic field, $B_* = 10^2 B_0$ and $B_* = 10^3 B_0$, are considered. These results show that for certain values of the collision time considered, the synchrotron radiation produced in the blobs indeed extends up to a factor B_*/B_0 beyond the limit $h\nu^{\text{MAX}}$. However, for the case of $\tau > t^{\text{MAX}}$, the overall luminosity of this high energy emitted component is significantly less than the one produced in the main zone. Only for the case of small collision timescales, $\tau < t^{\text{MAX}}$, is the SED dominated by the emission produced in the blobs. The dashed curves in these figures correspond to the emission in the accelerator's magnetic field B_0 and the solid curves correspond to the emission in the blob's magnetic field B_* . If particles enter the strong magnetic field region during the acceleration process then the maximum attainable energy can be significantly affected, decreasing the energy of the SED maximum formed in the blobs (the case of τ_4 in Figs. 2 and 3).

III. PHYSICAL JUSTIFICATION OF THE TOY MODEL

The mathematical model introduced in Sec. II formally depends on a single parameter, τ . However, there are two

hidden parameters, t_* and B_* , which determine the energy range where the leaky box approximation used is actually valid. Furthermore, the main zone is characterized by two parameters: the size, R and the magnetic field, B_0 . The question remains whether there exists a physically meaningful combination of these parameters which would allow the realization of the radiation scenario illustrated in Fig. 2.

For the sake of simplicity, we assume that there exist \mathcal{N} identical small scale large field blobs of size r_* . The interaction time of electrons with these blobs lasts a minimum duration of $t_* \simeq r_*/c$. This interaction time can be significantly longer if the electron is magnetically confined within the blob. To avoid this complication, we focus on sufficiently high electron energies (above E_*) such that the electron life-time is shorter than their minimal confinement time in the blobs $\sim r_*/c$.

For the strong field of the blobs, the geometrical size of the blob dictates the collision time of the electrons with them,

$$\tau \simeq \frac{R^3}{\mathcal{N}\pi r_*^2 c}. \quad (8)$$

On the other hand, the magnetic field strength of the blob determines the electron energy above which the leaky-box treatment remains valid:

$$E_* = \frac{6\pi m_e^2 c^4}{\sigma_T r_* B_*^2}. \quad (9)$$

The scenario introduced in Sec. II is realized when the following three conditions are satisfied: (i) $t^{\text{MIN}} < \tau <$




FIG. 2. Characteristic spectral energy distributions for four example cases considered in Fig. 1 (the same line colors for the cases considered in Fig. 1 are used here also). The dashed lines indicate the emission from particles in the accelerator magnetic field (B_0), and the solid lines indicate the emission from particles in the blob magnetic field ($B_* = 10^2 B_0$). The solid lines show the total overall emission from the two emission zones.

t^{MAX} , (ii) $E_* \ll E^{\text{MAX}}$, and (iii) $B_* \gg B_0$. The former condition determines the range for \mathcal{N} , and can be fulfilled without interference with the two other conditions. We therefore need to check just one condition, $E_* \ll E^{\text{MAX}}$, in the limit $B_* \gg B_0$:

$$r_* \gg \frac{B_0^2 E^{\text{MAX}} \eta}{B_*^2 e B_0}. \quad (10)$$

From the application of the maximum energy condition for a source (“Hillas criterion”) $\eta E^{\text{MAX}}/(e B_0) < R$, it follows that there should exist a range of r_* such that $E_* \ll E^{\text{MAX}}$ and $r_* \ll R$. However, this requires a large ratio of magnetic fields: $B_* \gg B_0$. We also note that if the acceleration efficiency is low, $\eta \gg 1$, then the allowed range of r_* shrinks. The considered radiation regime can therefore only be realized in extreme accelerators for which $\eta \approx 1$.

Let us estimate the fraction of the magnetic energy in the blobs required for strong boosting of the radiation energy. It is determined by the condition $\tau \ll t^{\text{MAX}}$:

$$\frac{\delta w_B}{w_B} = \mathcal{N} \frac{r_*^3 B_*^2}{R^3 B_0^2} = \frac{t^{\text{MAX}}}{\pi \tau} \geq 1. \quad (11)$$

This implies that significant transformation of the synchrotron spectrum beyond $h\nu^{\text{MAX}}$ is possible when the magnetic energy in the small blobs dominates over the energy in the main zone magnetic field. Note that this condition can be realized only in weakly magnetized environments, where the energy contained in the particles can provide an ample source for the formation of strong magnetic fluctuations (blobs).

IV. DISCUSSION

The synchrotron radiation of relativistic electrons is a unique process that proceeds with high efficiency in




FIG. 3. The same as Fig. 2 but for $B_* = 10^3 B_0$.

a wide variety of astrophysical environments and spans from radio to gamma-rays. The typical short time-scales characterizing the synchrotron cooling of electrons make this process as the dominant radiation channel not only at radio wavelengths but also in the relatively low (Sub-GeV) energy gamma-ray band. This is the case, for example, of gamma-ray bursts [5] or the so-called Crab flares [6]. The characteristic spectral and temporal features of this mechanism seem very attractive for interpretation of the fast and energetic phenomena in the very high energy gamma-ray band as well. One may recall, for example, the ultrafast VHE gamma-ray flares of blazars [7, 8] or extension of the gamma-ray spectra of late afterglows of GRBs [9]. However, the self-regulated cutoff in the synchrotron spectrum at ≤ 100 MeV limits the applicability of the process for energies well above 100 MeV (in the frame of the source).

Synchrotron radiation can be significantly boosted in sources where the emission region moves relativistically, $h\nu^{\text{MAX}} \simeq 100 \delta$ MeV. The Doppler boosting of the maximum of synchrotron radiation can be naturally realized in relativistic outflows like the jets of active galactic nuclei (AGN) and gamma-ray bursts (GRBs). Typically, in the AGN and late GRB afterglows, the jet’s Doppler factors do not exceed 100. Thus the Doppler boosting alone cannot shift the synchrotron radiation to very-high energies, especially given that in realistic conditions, e.g. with the inclusion of the presence of dissipative factors like adiabatic losses or plasma instabilities, the η -parameter may significantly exceed 1. The consideration of physically realistic values for η , therefore, exacerbates further the problem of accounting for emission at the $h\nu^{\text{MAX}}$ scale as being synchrotron in origin.

In this paper, we propose a scenario for the formation of synchrotron radiation in compact magnetic blobs located inside the particle accelerator. The blobs do not participate in the acceleration process but serve as passive targets where the highest energy electrons accelerated in the main zone, are instantly cooled via synchrotron losses, whose spectrum is effectively extended up to the VHE band. We have developed a simple mathematical formalism based on a “leaky box” approach, for calculations of synchrotron radiation emerging from such systems. The results of numerical calculations confirm the qualitative estimates. They demonstrate that the synchrotron radiation of such systems can be effectively released in the very-high-energy regime provided that the magnetic field in the blobs exceeds two or three orders of magnitude the magnetic field in the accelerator zone. Although the compact magnetic condensations occupy a rather small volume inside the accelerator, the total magnetic energy in these blobs should significantly exceed the electromagnetic energy contained in the acceleration zone. Remarkably, the MHD studies of magnetic amplification by turbulent dynamo [10] in relativistic outflows may result in such systems where the magnetic energy is predominantly contained in compact clumps. Moreover, this amplification of small scale fields may be a general

effect [11].

DK acknowledges support by JSPS KAKENHI Grant Numbers JP18H03722, JP24105007, and JP16H02170.

Appendix A: Leaky-box approximation

Magnetic field fluctuations play a key role in both sink and source terms in the standard injection-cooling equation:

$$\frac{\partial n}{\partial t} + \frac{\partial(\dot{E}n)}{\partial E} + \frac{n}{\tau} - \frac{n(\tilde{E})}{\tau} = q(E), \quad (\text{A1})$$

where $q(E)$ is the injection term, $n = \frac{dN}{dE}$ is energy-distribution function, \dot{E} is energy losses in the main zone magnetic field, τ is the particle – fluctuation collision time, $\tilde{E}(E)$ is the particle energy that cools down to E while interacting with the fluctuation. While the particle is interacting with the fluctuation, it loses energy via synchrotron emission in the strong magnetic field of strength B_* . If the interaction time is t_* , then the energy change is

$$\tilde{E} = \frac{E}{1 - t_*/t_{\text{SYN}}}, \quad (\text{A2})$$

where E is particle final energy and $t_{\text{SYN}}(E, B_*) (\propto E^{-1} B_*^{-2})$ is the synchrotron cooling time in the magnetic field of strength B_* . One can approximate the above function as

$$\tilde{E} = \begin{cases} E & E < E_* \\ \infty & E \geq E_* \end{cases}, \quad (\text{A3})$$

where E_* is the solution of equation $t_* = t_{\text{SYN}}(E_*, B_*)$. For the energy range $E > E_*$, the above equation takes a simple form

$$\frac{\partial n}{\partial t} + \frac{\partial(\dot{E}n)}{\partial E} + \frac{n}{\tau} = q(E), \quad (\text{A4})$$

which is identical to the equation for transport of cosmic ray under the leaky-box approximation. It allows a well-known analytical solution:

$$n(E, t) = -\frac{1}{\dot{E}} \int_E^{E_{\text{EFF}}} dE' q(E') \exp\left[-\frac{\tau_{\text{EFF}}(E, E')}{\tau}\right]. \quad (\text{A5})$$

Here the auxiliary parameters, E_{EFF} and τ_{EFF} , are

$$t = - \int_E^{E_{\text{EFF}}} \frac{dE'}{\dot{E}(E')} \quad (\text{A6})$$

and

$$\tau_{\text{EFF}}(E, E') = - \int_E^{E'} \frac{dE''}{\dot{E}(E'')}. \quad (\text{A7})$$

Since the dominant cooling mechanism is synchrotron losses the above integral can be derived analytically.

Appendix B: Approximated expression for the synchrotron spectrum

The total emission of a particle that enters a high magnetic field region needs to be computed accounting for the change of the particle energy. To be consistent with the standard textbook, we operate with the total emitted energy per frequency:

$$E_\nu = \int_0^{t_*} P_\nu(E(t)) dt = \frac{\sqrt{3}e^3 B_*}{m_e c^2} \int_0^{t_*} F_\nu(x(t)) dt, \quad (\text{B1})$$

where

$$F(x) = x \int_x^\infty K_{5/3}(\xi) d\xi \quad (\text{B2})$$

and

$$x = \frac{4\pi\nu m_e^3 c^5}{3E_0^2 e B_0} u^2 = \nu_0 u^2 \quad (\text{B3})$$

(here $u = E_0/E$ and the above equation defines ν_0). In Eq. (B1) the particle energy, $E(t)$, depends on the initial energy, E_0 , as

$$E(t) = \frac{E_0}{1 + t/t_{\text{SYN}}(E_0, B_*)} \quad (\text{B4})$$

and

$$dt = t_{\text{SYN}}(E_0, B_*) du. \quad (\text{B5})$$

Although, formally, Eq. (B1) is a four-parameter function, it is clear that

$$E_\nu(E_0, B_*, t_*) = E_\nu(E_0, B_*, \infty) - E_\nu(E(t_*), B_*, \infty), \quad (\text{B6})$$

which allows one to reduce the dependence on t_* , and in what follows we consider Eq. (B1) in the limit $t_* \rightarrow \infty$.

Combining the above equations, one obtains

$$\begin{aligned} E_\nu &= \frac{\sqrt{3}e^3 B_* t_{\text{SYN}}(E_0, B_*)}{m_e c^2} \int_1^\infty du \nu_0 u^2 \int_{\nu_0 u^2}^\infty d\xi K_{5/3}(\xi) \\ &= \frac{\sqrt{3}e^3 B_* t_{\text{SYN}}(E_0, B_*)}{m_e c^2} G(\nu_0). \end{aligned} \quad (\text{B7})$$

Changing the integral order allows one to derive one integral in the auxiliary function analytically:

$$G(\nu_0) = \frac{\nu_0}{3} \int_{\nu_0}^\infty d\xi K_{5/3}(\xi) \left(\left(\frac{\xi}{\nu_0} \right)^{3/2} - 1 \right). \quad (\text{B8})$$

Using the approach introduced in Khangulyan *et al.* [12], we approximate Eq. (B8) with a simple analytic function:

$$G(x) \simeq \tilde{G}(x) = \frac{1.04e^{-x}}{\sqrt{x}} \left(\frac{1 + 0.464x}{1 + 0.771x} \right) g(x), \quad (\text{B9})$$

where

$$g(x) = \left[1 + \frac{ax^\alpha}{1 + bx^\beta} \right]^{-1} \quad (\text{B10})$$

for $a = 0.186$, $\alpha = 0.57$, $b = 1.75$, and $\beta = 1.49$. This approximation has exact asymptotical behavior for $x \rightarrow 0$ and ∞ . In the entire range it provides an accuracy of $\simeq 0.2\%$ as shown in Fig. 4.




FIG. 4. Ratio of the exact and approximate function that describe the synchrotron spectrum accounting for change of the particle energy.

-
- [1] P. W. Guilbert, A. C. Fabian, and M. J. Rees, Spectral and variability constraints on compact sources, *MNRAS* **205**, 593 (1983).
- [2] The “100 MeV” limit does not concern the synchrotron radiation produced by secondary electrons, for example, the products of interactions of primary protons with the surrounding gas and radiation fields.
- [3] S. R. Kelner, F. A. Aharonian, and D. Khangulyan, On the Jitter Radiation, *ApJ* **774**, 61 (2013), arXiv:1304.0493 [astro-ph.HE].
- [4] E. Derishev and F. Aharonian, Exact Analytical Expression for the Synchrotron Radiation Spectrum in the Gaussian Turbulent Magnetic Field, *ApJ* **887**, 181 (2019), arXiv:1907.11663 [astro-ph.HE].
- [5] M. Ajello *et al.*, Investigating the Nature of Late-time High-energy GRB Emission through Joint Fermi/Swift Observations, *ApJ* **863**, 138 (2018).
- [6] R. Bühler and R. Blandford, The surprising Crab pulsar and its nebula: a review, *Reports on Progress in Physics* **77**, 066901 (2014), arXiv:1309.7046 [astro-ph.HE].
- [7] F. Aharonian *et al.*, An Exceptional Very High Energy Gamma-Ray Flare of PKS 2155-304, *ApJ* **664**, L71 (2007), arXiv:0706.0797 [astro-ph].
- [8] J. Aleksić *et al.*, Black hole lightning due to particle acceleration at subhorizon scales, *Science* **346**, 1080 (2014), arXiv:1412.4936 [astro-ph.HE].
- [9] H. Abdalla *et al.*, A very-high-energy component deep in the γ -ray burst afterglow, *Nature* **575**, 464 (2019), arXiv:1911.08961 [astro-ph.HE].
- [10] W. Zhang, A. MacFadyen, and P. Wang, Three-Dimensional Relativistic Magnetohydrodynamic Simulations of the Kelvin-Helmholtz Instability: Magnetic Field Amplification by a Turbulent Dynamo, *ApJ* **692**, L40 (2009), arXiv:0811.3638 [astro-ph].
- [11] A. P. Kazantsev, Enhancement of a Magnetic Field by a Conducting Fluid, *Soviet Journal of Experimental and Theoretical Physics* **26**, 1031 (1968).
- [12] D. Khangulyan, F. A. Aharonian, and S. R. Kelner, Simple Analytical Approximations for Treatment of Inverse Compton Scattering of Relativistic Electrons in the Blackbody Radiation Field, *ApJ* **783**, 100 (2014), arXiv:1310.7971 [astro-ph.HE].