

Some properties of the norm in a quaternion division algebra

Cristina FLAUT and Diana SAVIN

Abstract. In this paper we provide some applications of the norm form in some quaternion division algebras over rational field and we give some properties of Fibonacci sequence and Fibonacci sequence in connection with quaternion elements. We define a monoid structure over a finite set on which we will prove that the defined Fibonacci sequence is stationary, we provide some properties of the norm of a rational quaternion algebra, in connection to the famous Lagrange's four-square theorem and its generalizations given by Ramanujan. Moreover, we prove some results regarding the arithmetic of integer quaternions defined on some division quaternion algebras and we define and give properties of some special quaternions by using Fibonacci sequences.

Key Words: quaternion algebras; Lagrange's four-square theorem; Fibonacci quaternion elements;

2000 AMS Subject Classification: 11B39, 11R54, 17A35.

1. Introduction

Over the time, the problem of determining integer numbers a, b, c, d such that all positive integers can be written under the form $ax^2 + by^2 + cz^2 + du^2$, with $x, y, z, u \in \mathbb{Z}$, has fascinated many mathematicians. Jacobi proved that all positive integers n can be written under the form $n = x^2 + 2y^2 + 3z^2 + 6u^2$, with $x, y, z, u \in \mathbb{Z}$. Liouville and Pepin proved that for

$$(a, b, c, d) \in \{(1, 1, 1, 1), (1, 1, 2, 2), (1, 2, 2, 2), (1, 1, 1, 4), (1, 1, 2, 4), (1, 2, 2, 4)\} \\ \cup \{(1, 2, 4, 4), (1, 1, 2, 8), (1, 2, 4, 8), (1, 1, 3, 3), (1, 2, 5, 10)\},$$

all positive integers can be written under the form $n = ax^2 + by^2 + cz^2 + du^2$, with $x, y, z, u \in \mathbb{Z}$.

In the paper [Ra; 1917], S. Ramanujan has found all positive integers a, b, c, d such that all positive integers n can be expressed under the form

$$ax^2 + by^2 + cz^2 + du^2,$$

with $x, y, z, u \in \mathbb{Z}$. He proved that there are only 55 sets of values of a, b, c, d such that any positive integer n can have the form $n = ax^2 + by^2 + cz^2 + du^2$, with $x, y, z, u \in \mathbb{Z}$ ([Ra; 1917], p. 171). From these 55 sets of values, only 7 sets of values of a, b, c, d have the properties $a = 1$ and $d = bc$, namely for

$$(a, b, c, d) \in \left\{ \begin{array}{l} (1, 1, 1, 1), (1, 1, 2, 2), (1, 1, 3, 3), (1, 2, 2, 4), (1, 2, 3, 6), \\ (1, 2, 4, 8), (1, 2, 5, 10) \end{array} \right\}. \quad (1.1.)$$

In this paper we find an interesting connection between quaternion algebras and these quaternary quadratic forms. Moreover, we define a monoid structure over a finite set on which we will prove that the defined Fibonacci sequence is stationary, we prove some results regarding the arithmetic of integer quaternions defined on some division quaternion algebras and we define and give properties of some special quaternions by using Fibonacci sequences.

2. Fibonacci stationary elements

Let $(f_n)_{n \geq 0}$ be the Fibonacci sequence

$$f_n = f_{n-1} + f_{n-2}, \quad n \geq 2, f_0 = 0; f_1 = 1,$$

and $(l_n)_{n \geq 0}$ be the Lucas sequence

$$l_n = l_{n-1} + l_{n-2}, \quad n \geq 2, l_0 = 2; l_1 = 1.$$

We know that the Binet's formula for Fibonacci sequence are

$$f_n = \frac{\alpha^n - \beta^n}{\alpha - \beta} = \frac{\alpha^n - \beta^n}{\sqrt{5}}, \quad (\forall) n \in \mathbb{N}.$$

and the Binet's formula for Lucas sequence are

$$l_n = \alpha^n + \beta^n, \quad (\forall) n \in \mathbb{N}.$$

Proposition 2.1. ([Fib]). *Let $(f_n)_{n \geq 0}$ be the Fibonacci sequence and let $(l_n)_{n \geq 0}$ be the Lucas sequence. The following properties hold:*

i)

$$f_n^2 + f_{n+1}^2 = f_{2n+1}, \quad \forall n \in \mathbb{N};$$

ii)

$$f_{n+3}^2 = 2f_{n+2}^2 + 2f_{n+1}^2 - f_n^2, \quad \forall n \in \mathbb{N} - \{0\};$$

iii)

$$f_{n-1}f_m + f_n f_{m+1} = f_{n+m}, \quad \forall n, m \in \mathbb{N} - \{0\};$$

iv)

$$f_n f_m - f_{n-k} f_{m+k} = (-1)^{n-k} f_k f_{m+k-n}, \forall n, m, k \in \mathbb{N}, k \leq n.$$

It is well known that the Fibonacci sequence is periodic when it is reduced modulo m and this period is called *Pisano's period*, denoted by $\pi(m)$. Fibonacci numbers are defined by using the addition operation over the real field. Will be interesting to find other finite algebraic structures on which we will study the behavior of such a sequence. Therefore, it is interesting to find other multiplications and sets such that the behavior of such a sequence defined on to be predictable. In the following, we will define a monoid structure over a finite set on which we will prove that the defined Fibonacci sequence is stationary.

Let (M_1, \perp) and (M_2, \top) be two monoids. On their cartesian product $M_1 \times M_2$ we define the following multiplication " * ":

$$(a_1, a_2) * (b_1, b_2) = (a_1 \perp b_1, a_2 \top b_2). \quad (2.1)$$

It results that $(M_1 \times M_2, *)$ is a monoid.

Now, we consider the totally ordered sets (A_i, \leq_i) , with $A_i = \{o_i, x_i\}$, $i \in \{1, 2, \dots, k\}$. On A_i we define the multiplication " +_i "

$$x_i +_i x_i = x_i, o_i +_i o_i = o_i, x_i +_i o_i = o_i +_i x_i = x_i.$$

It results that $(A_i, +_i)$ is a commutative monoid. We consider the Cartesian product $A = A_1 \times A_2 \times \dots \times A_k$. Using relation (2.1), we obtain that $(A, *)$ is a commutative monoid. The set $A = \{y_0 \leq y_1 \leq \dots \leq y_{2^k-1}\}$ is ordered with lexicographic order and it is a totally ordered set. We denote $y_0 = 0 = (o_1, o_2, \dots, o_{2^k})$ and $y_{2^k-1} = 1 = (x_1, x_2, \dots, x_{2^k})$.

On A , the above multiplication " * " can be given by the following relations:

$$\begin{cases} y_i * y_j = 1, & \text{if } i + j > 2^k - 1; \\ y_i * y_j = y_{i+j}, & \text{if } i + j \leq 2^k - 1; \\ y * 0 = y, & \text{for all } y \in A. \end{cases}$$

Example 2.2. Let $k = 3$. We have $A = \{(y_0, y_1, \dots, y_7)\}$, where $y_0 = (o_1, o_2, o_3) = 0$, $y_1 = (o_1, o_2, x_3)$, $y_2 = (o_1, x_2, o_3)$, $y_3 = (o_1, x_2, x_3)$, $y_4 = (x_1, o_2, o_3)$, $y_5 = (x_1, o_2, x_3)$, $y_6 = (x_1, x_2, o_3)$, $y_7 = (x_1, x_2, x_3) = 1$. It results that

$$y_1 * y_2 = (o_1, x_2, x_3) = y_3, y_2 * y_5 = (x_1, x_2, x_3) = y_7 = 1, \\ y_3 * y_6 = (x_1, x_2, x_3) = 1, \text{ etc.}$$

Definition 2.3. We consider $(M, *)$ a commutative monoid. For $a, b \in M$, we define the following sequence

$$\{a, b\} = \{a, b, a * b, b * (a * b), \dots, v_n, v_{n+1}, v_{n+2}, \dots\},$$

where $v_0 = a$, $v_1 = b$ and $v_{n+2} = v_n * v_{n+1}$, for $n \in \mathbb{N}$. This sequence are called the *Fibonacci sequence attached to the elements a, b*. If we find a number $t \in \mathbb{N}$ such that $v_n = v_{n+1} = v_{n+2} = \dots$, for all $n \geq t$, then the sequence $\{a, b\}$ is called *t-stationary*.

Proposition 2.4. *With the above notations, the Fibonacci sequence defined on the set A is t -stationary.*

Proof. If $a, b \in M, a \neq 0, b \neq 0$, we consider the sequence

$$\{a, b\} = \{a, b, a * b, b * (a * b), \dots, v_n, v_{n+1}, v_{n+2}, \dots\},$$

where $v_0 = a, v_1 = b$ and $v_{n+2} = v_n * v_{n+1}$, for $n \in \mathbb{N}$. It results that $v_2 = a * b > a$ and $v_2 = a * b > b$, since $a * b = a$ or $a * b = b$ implies $b = 0$ or $a = 0$, false. If $v_2 = a * b > ax$, we have that $v_3 = b * (a * b) = (a * b) * b > a * b = v_2$, $v_4 = (a * b) * (b * (a * b)) > b * (a * b) = v_3$, etc. We get that the obtained increased sequence is stationary, since the set A is finite. Therefore, there is number $t \in \mathbb{N}$ such that $v_t = 1 = v_{t+1} = v_{t+2} = \dots$. It results that

$$\{a, b\} = \{a, b, a * b, b * (a * b), \dots, v_{t-1}, 1, 1, \dots\}.$$

Assuming now that $a, b \in A$ and $b = 0$, we obtain the sequence $\{a, b\}$ with $v_0 = a, v_1 = 0, v_2 = a, v_3 = a, v_4 = a * a > a$. Therefore, from the above, we obtain a stationary sequence, that means we get a number $t \in \mathbb{N}$ such that $v_t = 1 = v_{t+1} = v_{t+2} = \dots$.

Example 2.5. Considering the set A from Example 2.2, and the elements $y_2 = (o_1, x_2, o_3), y_4 = (x_1, o_2, o_3)$, we have $\{y_2, y_4\} = \{y_2, y_4, y_2 * y_4, y_4 * (y_2 * y_4), \dots\} = \{y_2, y_4, y_6, 1, 1, \dots\}$, therefore the sequence is stationary.

Remark 2.6. Proposition 2.4 from above generalizes results obtained in [Fl; 20], Proposition 2.6.

3. Remarks regarding the norm of a quaternion algebra

In this section, we will provide some properties of the norm of a rational quaternion algebra, in connection to the famous Lagrange's four-square theorem and its generalizations given by Ramanujan in [Ra; 1917].

First of all, we recall some results about diophantine equations, about quaternion algebras and about Fibonacci sequence, results which will be used in the following proofs.

Proposition 3.1. ([Si; 70]). *Let m be a fixed positive integer. The diophantine equation $x^2 + my^2 = z^2$ has infinitely many solutions, namely:*

$$x = a^2 - mb^2, y = 2ab, z = a^2 + mb^2, a, b \in \mathbb{Z}.$$

Proposition 3.2. ([Gi, Sz; 06]). *Let K be a field. Then, the quaternion algebra $\mathbb{H}_K(b, c)$ is a split algebra if and only if the conic*

$$C(a, b) : bx^2 + cy^2 = z^2$$

has a rational point over $K \setminus \{(0, 0, 0)\}$, i.e. there are $x_0, y_0, z_0 \in K \setminus \{(0, 0, 0)\}$ such that $bx_0^2 + cy_0^2 = z_0^2$.

Let K be an algebraic number field, let \mathcal{O}_K be the ring of integers of the field K and let $H_K(b, c)$ be a quaternion algebra over the field K . We recall that the discriminant (or reduced discriminant) $D_{H_K(b, c)}$ of the quaternion algebra $H_K(b, c)$ is the product of all prime ideals of \mathcal{O}_K that ramify in $H_K(b, c)$. If p is a prime and $K = \mathbb{Q}$, we recall the notations: $\left(\frac{\cdot}{p}\right)$ for the Legendre symbol in \mathbb{Z} and $(b, c)_p$ for the Hilbert symbol in the p -adic field \mathbb{Q}_p

$$(b, c)_p = \begin{cases} 1, & \text{if } p \text{ does not ramify in } H_K(b, c) \\ -1, & \text{if } p \text{ ramifies in } H_K(b, c). \end{cases}$$

The quaternion algebra $H_K(b, c)$ splits if and only if the discriminant $D_{H_K(b, c)}$ is equal to \mathcal{O}_K . Since a quaternion algebra is either split or a division algebra, it results that a quaternion algebra $H_K(b, c)$ is a division algebra if and only if there is a prime p such that $p | D_{H_K(b, c)}$.

Proposition 3.3. ([Ma; 07]; [Vi; 80]). *The discriminant of a quaternion algebra determines the algebra up to isomorphism. That means, if K is a field, then two quaternion algebras over the field K are isomorphic if and only if they have the same reduced discriminant.*

Let b, c be positive integer numbers and let $\mathbb{H}_{\mathbb{Q}}(-b, -c)$ be the generalized quaternion algebra over the field of rational numbers. Let $x \in \mathbb{H}_{\mathbb{Q}}(-b, -c)$, $x = x_1 \cdot 1 + x_2 e_2 + x_3 e_3 + x_4 e_4$, where $x_i \in \mathbb{Q}, i \in \{1, 2, 3, 4\}$, and the elements of the basis $\{1, e_2, e_3, e_4\}$ satisfy the following rules, given in the below multiplication table:

\cdot	1	e_2	e_3	e_4	
1	1	e_2	e_3	e_4	
e_2	e_2	$-b$	e_4	$-be_3$.
e_3	e_3	$-e_4$	$-c$	ce_2	
e_4	e_4	be_3	$-ce_2$	$-bc$	

If $x = x_1 \cdot 1 + x_2 e_2 + x_3 e_3 + x_4 e_4 \in \mathbb{H}_{\mathbb{Q}}(-b, -c)$, we denote by $\mathbf{n}(x)$ and $\mathbf{t}(x)$ the norm and the trace of a quaternion x . The conjugate of the quaternion x is $\bar{x} = x_1 \cdot 1 - x_2 e_2 - x_3 e_3 - x_4 e_4 \in \mathbb{H}_{\mathbb{Q}}(-b, -c)$. The trace of a quaternion is

$$\mathbf{t}(x) = x + \bar{x} \in \mathbb{Q}$$

and the norm has the following expression

$$\mathbf{n}(x) = x \cdot \bar{x} = x_1^2 + bx_2^2 + cx_3^2 + bcx_4^2. \quad (3.2.)$$

If, for $x \in \mathbb{H}(-b, -c)$, the relation $\mathbf{n}(x) = 0$ implies $x = 0$, then the algebra $\mathbb{H}(-b, -c)$ is called a *division algebra*, otherwise the quaternion algebra is called a *split algebra*.

Proposition 3.4. *Let b, c be positive integers and let $\mathbb{H}_{\mathbb{Q}}(-b, -c)$ be a generalized quaternion algebra. If*

$$(b, c) \in \{(1, 1), (1, 2), (1, 3), (2, 2), (2, 3), (2, 4), (2, 5)\},$$

then, the norm map

$$\mathbf{n} : \mathbb{H}_{\mathbb{Q}}(-b, -c) \rightarrow \mathbb{Q}_+$$

is a surjective function.

Proof. If we consider any positive integer m , from relation (1.1), there are some integers x_1, x_2, x_3, x_4 such that $m = x_1^2 + bx_2^2 + cx_3^2 + bcx_4^2$. Thus, there is the quaternion $x = x_1 \cdot 1 + x_2e_2 + x_3e_3 + x_4e_4 \in \mathbb{H}_{\mathbb{Q}}(-b, -c)$, such that $m = \mathbf{n}(x)$.

If we consider any positive rational number $m = \frac{m'}{l}$, where $m', l \in \mathbb{N}$, $l \neq 0$, $\text{g.c.d}(m', l) = 1$, by using the same argument as above for m'/l , there are some rational numbers $x'_1 = \frac{x_1}{l}, x'_2 = \frac{x_2}{l}, x'_3 = \frac{x_3}{l}, x'_4 = \frac{x_4}{l}$ such that

$$m = \frac{m'}{l} = \frac{m' l}{l^2} = \left(\frac{x_1}{l}\right)^2 + b \left(\frac{x_2}{l}\right)^2 + c \left(\frac{x_3}{l}\right)^2 + bc \left(\frac{x_4}{l}\right)^2 = \mathbf{n}(x'),$$

where $x' = x'_1 \cdot 1 + x'_2e_2 + x'_3e_3 + x'_4e_4 \in \mathbb{H}_{\mathbb{Q}}(-b, -c)$. From here, it results that, \mathbf{n} is a surjective function.

Proposition 3.5. *If*

$$(b, c) \in \{(1, 1), (1, 2), (1, 3), (2, 2), (2, 3), (2, 4), (2, 5)\},$$

then the obtained 7 quaternion algebras over the field of rational numbers, $\mathbb{H}_{\mathbb{Q}}(-b, -c)$, are all division algebras.

Proof. Since $b > 0, c > 0$, it result that the equation

$$-bx^2 - cy^2 = z_0^2$$

does not have solutions (x, y, z) in $(\mathbb{Q} \setminus \{(0, 0, 0)\}) \times (\mathbb{Q} \setminus \{(0, 0, 0)\}) \times (\mathbb{Q} \setminus \{(0, 0, 0)\})$.

Let K be a field. It is known that two quaternion K - algebras which split are isomorphic, but a similar result is not known in the case of two division quaternion K - algebras. These algebras may be isomorphic or may be not. We are interested to know how many of these 7 quaternions \mathbb{Q} - algebras are isomorphic and which with which is isomorphic. The answer to this question is given in the following proposition.

Proposition 3.6.

- a) The quaternion algebras $\mathbb{H}_{\mathbb{Q}}(-1, -1)$, $\mathbb{H}_{\mathbb{Q}}(-1, -2)$, $\mathbb{H}_{\mathbb{Q}}(-2, -2)$, $\mathbb{H}_{\mathbb{Q}}(-2, -3)$, $\mathbb{H}_{\mathbb{Q}}(-2, -4)$ are isomorphic.
b) The quaternion algebras $\mathbb{H}_{\mathbb{Q}}(-1, -3)$, $\mathbb{H}_{\mathbb{Q}}(-2, -5)$ are not isomorphic between them and nor with those 5 quaternions algebras from a).

Proof. It is known ([Ko]) that if a prime p divides the discriminant of the algebra $\mathbb{H}_{\mathbb{Q}}(-b, -c)$, then p must divides $2bc$. We know that the discriminant $D_{\mathbb{H}_{\mathbb{Q}}(-b, -c)}$ of the algebra $\mathbb{H}_{\mathbb{Q}}(-b, -c)$ is the product of prime ideals of \mathbb{Z} which ramify in $\mathbb{H}_{\mathbb{Q}}(-b, -c)$. Since $(\mathbb{Z}, +, \cdot)$ is a principal ideal domain, to simplify writing, an ideal of \mathbb{Z} is identified with its generator, up to units.

We consider the algebra $\mathbb{H}_{\mathbb{Q}}(-2, -3)$. Let p be a prime which divides the discriminant of $\mathbb{H}_{\mathbb{Q}}(-2, -3)$, therefore we have $p|6$.

If $p = 3$, by using the properties of the Hilbert symbol and of the Legendre symbol we obtain:

$$\begin{aligned} (-2, -3)_3 &= (-1, -3)_3(2, -3)_3 = (-1, -1)_3(-1, 3)_3(2, -1)_3(2, 3)_3 = \\ &= 1 \cdot \left(\frac{-1}{3}\right) \cdot 1 \cdot \left(\frac{2}{3}\right) = \left(\frac{2}{3}\right)^2 = 1, \end{aligned}$$

It results that 3 does not ramify in $H_{\mathbb{Q}}(-2, -3)$.

By following computation

$$\begin{aligned} (-2, -3)_2 &= (-1, -3)_2(2, -3)_2 = (-1, -1)_2(-1, 3)_2(2, -1)_2(2, 3)_2 = \\ &= (-1)(-1)^{\frac{3-1}{2} \cdot \frac{-1-1}{2}} 1(-1)^{\frac{3^2-1}{8}} = -1, \end{aligned}$$

we obtain that the only prime which ramifies in $\mathbb{H}_{\mathbb{Q}}(-2, -3)$ is 2. Therefore, the reduced discriminant of the algebra $\mathbb{H}_{\mathbb{Q}}(-2, -3)$ is 2.

In a similar way, we obtain that

$$D_{\mathbb{H}_{\mathbb{Q}}(-1, -1)} = D_{\mathbb{H}_{\mathbb{Q}}(-1, -2)} = D_{\mathbb{H}_{\mathbb{Q}}(-2, -2)} = D_{\mathbb{H}_{\mathbb{Q}}(-2, -4)} = 2.$$

Now, by using Proposition 3.3, it results that the quaternion algebras $\mathbb{H}_{\mathbb{Q}}(-1, -1)$, $\mathbb{H}_{\mathbb{Q}}(-1, -2)$, $\mathbb{H}_{\mathbb{Q}}(-2, -2)$, $\mathbb{H}_{\mathbb{Q}}(-2, -3)$, $\mathbb{H}_{\mathbb{Q}}(-2, -4)$ are isomorphic.

b) Similar to a), we calculate the discriminants for the quaternion algebras $\mathbb{H}_{\mathbb{Q}}(-1, -3)$, $\mathbb{H}_{\mathbb{Q}}(-2, -5)$ and we obtain that

$$D_{\mathbb{H}_{\mathbb{Q}}(-1, -3)} = 3, \quad D_{\mathbb{H}_{\mathbb{Q}}(-2, -5)} = 5.$$

Therefore, by using again Proposition 3.3 and a), from above, we obtain that the quaternion algebras $\mathbb{H}_{\mathbb{Q}}(-1, -3)$, $\mathbb{H}_{\mathbb{Q}}(-2, -5)$ are not isomorphic between them and nor with those 5 quaternions algebras from a).

Proposition 3.7. Let $\mathbb{Q}(i)$ be the quadratic field, where $i^2 = -1$. If

$$(b, c) \in \{(1, 1), (1, 2), (1, 3), (2, 2), (2, 3), (2, 4)\},$$

then these 6 quaternion algebras $\mathbb{H}_{\mathbb{Q}(i)}(-b, -c)$, over the quadratic field $\mathbb{Q}(i)$, are all split algebras and $\mathbb{H}_{\mathbb{Q}(i)}(-2, -5)$ is a division algebra.

Proof. For the first three cases, we use Proposition 3.1, but we are not looking for integer solutions of the equation $x^2 + my^2 = z^2$, we are looking for solutions over $\mathbb{Z}[i]$ of the equation $-x^2 - my^2 = z^2$.

Case 1. If $b = 1, c = 1$, then we remark that $(x_0, y_0, z_0) = (3i, 4i, 5)$ is a solution of the equation $-x^2 - y^2 = z^2$ over $\mathbb{Q}(i) \setminus \{(0, 0, 0)\}$.

Case 2. If $b = 1, c = 2$, then we remark that $(x_0, y_0, z_0) = (i, 2i, 3)$ is a solution of the equation $-x^2 - 2y^2 = z^2$ over $\mathbb{Q}(i) \setminus \{(0, 0, 0)\}$.

Case 3. If $b = 1, c = 3$, then we remark that $(x_0, y_0, z_0) = (i, i, 2)$ is a solution of the equation $-x^2 - 3y^2 = z^2$ over $\mathbb{Q}(i) \setminus \{(0, 0, 0)\}$.

Case 4. If $b = 2, c = 2$, then we remark that $(x_0, y_0, z_0) = (i, i, 2)$ is a solution of the equation $-2x^2 - 2y^2 = z^2$ over $\mathbb{Q}(i) \setminus \{(0, 0, 0)\}$.

Case 5. If $b = 2, c = 3$, then we remark that $(x_0, y_0, z_0) = (1, i, 1)$ is a solution of the equation $-2x^2 - 3y^2 = z^2$ over $\mathbb{Q}(i) \setminus \{(0, 0, 0)\}$.

Case 6. If $b = 2, c = 4$, then we remark that $(x_0, y_0, z_0) = (2i, 1, 2)$ is a solution of the equation $-2x^2 - 4y^2 = z^2$ over $\mathbb{Q}(i) \setminus \{(0, 0, 0)\}$.

Therefore, from Proposition 3.2, it results that the quaternion algebras $\mathbb{H}_{\mathbb{Q}(i)}(-1, -1)$, $\mathbb{H}_{\mathbb{Q}(i)}(-1, -2)$, $\mathbb{H}_{\mathbb{Q}(i)}(-1, -3)$, $\mathbb{H}_{\mathbb{Q}(i)}(-2, -2)$, $\mathbb{H}_{\mathbb{Q}(i)}(-2, -3)$, $\mathbb{H}_{\mathbb{Q}(i)}(-2, -4)$ are split algebras.

Case 7. If $b = 2, c = 5$, we use the same idea as in Proposition 3.6 and we determine the primes which ramify in $\mathbb{H}_{\mathbb{Q}(i)}(-2, -5)$. For the same purpose, we also can use the computer algebra system Magma ([Mag]) and we find that the reduced discriminant of the quaternion algebra $\mathbb{H}_{\mathbb{Q}(i)}(-b, -c)$ is $D_{\mathbb{H}_{\mathbb{Q}(i)}(-2, -5)} = 5\mathbb{Z}[i]$, therefore the quaternion algebra $\mathbb{H}_{\mathbb{Q}(i)}(-2, -5)$ is a division algebra.

4. Some aspects regarding the arithmetic of Integer Quaternions

As we remarked above, the algebras $\mathbb{H}_{\mathbb{Q}}(-1, -1)$, $\mathbb{H}_{\mathbb{Q}}(-1, -2)$, $\mathbb{H}_{\mathbb{Q}}(-2, -2)$, $\mathbb{H}_{\mathbb{Q}}(-2, -3)$, $\mathbb{H}_{\mathbb{Q}}(-2, -4)$, $\mathbb{H}_{\mathbb{Q}}(-1, -3)$, $\mathbb{H}_{\mathbb{Q}}(-2, -5)$ are division algebras and have a surjective norm form. We denote with \mathcal{H} the set of these algebras, namely $\mathcal{H} = \{\mathbb{H}_{\mathbb{Q}}(-1, -1), \mathbb{H}_{\mathbb{Q}}(-1, -2), \mathbb{H}_{\mathbb{Q}}(-2, -2), \mathbb{H}_{\mathbb{Q}}(-2, -3), \mathbb{H}_{\mathbb{Q}}(-2, -4), \mathbb{H}_{\mathbb{Q}}(-1, -3), \mathbb{H}_{\mathbb{Q}}(-2, -5)\}$. First five of these algebras are isomorphic. In $\mathbb{H}_{\mathbb{Q}}(-1, -1)$ is defined the following sets

$$\mathbb{H}(\mathbb{Z}) = \{x \in \mathbb{H}_{\mathbb{Q}}(-1, -1), x = x_1 \cdot 1 + x_2 e_2 + x_3 e_3 + x_4 e_4, x_1, x_2, x_3, x_4 \in \mathbb{Z}\},$$

called the set of *Lipschitz integers* and

$$\mathbb{H}\left(\mathbb{Z}\left[\begin{array}{c} 1 \\ 2 \end{array}\right]\right) = \left\{x \in \mathbb{H}_{\mathbb{Q}}(-1, -1), x = x_1 \cdot 1 + x_2 e_2 + x_3 e_3 + x_4 e_4, x_1, x_2, x_3, x_4 \in \mathbb{Z}\right\},$$

$$x_1, x_2, x_3, x_4 \text{ are all in } \mathbb{Z} \text{ or all in } \mathbb{Z} + \frac{1}{2},$$

called the set of *Hurwitz integers* (see [Co, Sm; 03], p.55).

In a similar way, for an algebra $A \in \mathcal{H}$, we will define the set

$$A(\mathbb{Z}) = \{x \in A, x = x_1 \cdot 1 + x_2 e_2 + x_3 e_3 + x_4 e_4, x_1, x_2, x_3, x_4 \in \mathbb{Z}\}.$$

We call the elements of this set *the integers of the algebra* $A \in \mathcal{H}$.

Proposition 4.1. *For an algebra $A \in \mathcal{H}$, for $x, y \in A(\mathbb{Z})$, there are γ and θ such that*

$$n(y)x = \gamma y + \theta n(y),$$

with $n(\theta) < n(y)$, $\gamma, \theta \in A(\mathbb{Z})$.

Proof. Since the norm n is surjective, there is a quaternion $\theta \in A(\mathbb{Z})$, such that $n(\theta) < n(y)$. Therefore, there is a quaternion $\delta \in A$ such that $\theta = x - \delta y$, with

$$\delta = (x - \theta)y^{-1} = \frac{(x - \theta)\overline{y}}{n(y)}$$

We denote $\gamma = (x - \theta)\overline{y} \in A(\mathbb{Z})$. It results

$$x = \delta y + \theta = \frac{\gamma}{n(y)}y + \theta,$$

therefore $n(y)x = \gamma y + n(y)\theta$.

Proposition 4.2. ([Da, Sa, Va; 03], Lemma 2.6.3). *Let $x, y \in \mathbb{H}(\mathbb{Z})$, with $n(y)$ an odd number. Therefore, there are γ and $\theta \in \mathbb{H}(\mathbb{Z})$ such that*

$$x = \gamma y + \theta, \text{ with } n(\theta) < n(y).$$

Remark 4.3. Proposition 4.2 was proved without using in the proof the surjectivity of the norm. Proposition 4.2 is a particular case of the Proposition 4.1, since if $n(y)$ is arbitrary, then the result obtained in Proposition 4.2 is also true in $\mathbb{H}(\mathbb{Z}[\frac{1}{2}])$.

Definition 4.4. Let $q \in \mathbb{H}(\mathbb{Z})$. If $n(q)$ is an odd integer, then the quaternion q is called an *odd* quaternion.

Definition 4.5. ([Gu; 13], Definition 2 and Theorem 1) We consider φ an odd quaternion from $\mathbb{H}(\mathbb{Z})$ and $x, y \in \mathbb{H}(\mathbb{Z})$ two arbitrary quaternions. If there is the quaternion $\theta \in \mathbb{H}(\mathbb{Z})$ such that $x - y = \theta\varphi$, therefore the quaternions x and y are called *right congruent modulo φ* and we will denote this $x \equiv_r y \pmod{\varphi}$. We obtain an equivalence relation and the quotient set is

$$\mathbb{H}(\mathbb{Z})_\varphi = \{x \pmod{\varphi}, x \in \mathbb{H}(\mathbb{Z})\}.$$

The set $\mathbb{H}(\mathbb{Z})_\varphi$ has $n^2(\varphi)$ elements and, from the above propositions, we have $n(y) < n(\varphi)$.

Definition 4.6. Let $A \in \mathcal{H}$, a quaternion $q \in A$ is a *unit* in A if $n(q) = 1$. A quaternion $q \in A(\mathbb{Z})$ is a *prime quaternion* if $q = \alpha\beta, \alpha, \beta \in A$ implies α or β is a unit element.

Remark 4.7. ([Gu; 13], Definition 5 and Remark 2)

1) We will consider the element $v \in \mathbb{H}(\mathbb{Z}[\frac{1}{2}]), v = \frac{1}{2}(1 + e_2 + e_3 + e_4)$ and we consider the set $\mathcal{A} = \{q \in \mathbb{H}(\mathbb{Z}[\frac{1}{2}]), q = \alpha + \beta v, \alpha, \beta \in \mathbb{Z}\}$. We remark that multiplication is commutative on \mathcal{A} , therefore we obtain that this set is a commutative ring. It results that we can define the same equivalence relation as above, obtaining the quotient ring \mathcal{A}_φ , with φ a prime quaternion in \mathcal{A} .

Proposition 4.8. A quaternion $q \in A(\mathbb{Z})$ is a prime quaternion if and only if its norm $\mathbf{n}(q)$ is a prime number.

Proof. We suppose that q is a prime quaternion and $\mathbf{n}(q) = m$, such that m is not a prime number. Therefore, $m = rs$, with $r \neq 1$ and $s \neq 1$. Since the norm \mathbf{n} is surjective, it results that there are the quaternions $q_1, q_2 \in A(\mathbb{Z})$ such that $\mathbf{n}(q_1) = r$ and $\mathbf{n}(q_2) = s$. We obtain that q_1 and q_2 are not unit quaternions, thus q is not a prime quaternion, which it is false. It results that $\mathbf{n}(q)$ is a prime number.

Conversely, if $\mathbf{n}(q) = p$ is a prime number and $q = q_1 q_2$, we obtain that $p = \mathbf{n}(q) = \mathbf{n}(q_1 q_2) = \mathbf{n}(q_1) \mathbf{n}(q_2)$. Therefore, $\mathbf{n}(q_1) = 1$ or $\mathbf{n}(q_2) = 1$. It results that q is a prime quaternion.

Remark 4.9. The above proposition generalized Corollary 2.6.10 from [Da, Sa, Va; 03]. In that statement the above result was proved for the algebra $\mathbb{H}_{\mathbb{Q}}(-1, -1)$ without using the surjectivity of the norm.

2) ([Gu; 13], Theorem 2) If $q \in \mathcal{A}, q = \alpha + \beta v, \alpha, \beta \in \mathbb{Z}$, with $\gcd\{\alpha, \beta\} = 1$, therefore $\mathbb{Z}_{n(q)}$ and \mathcal{A}_φ are isomorphic as rings. Therefore, if q is a prime quaternion, that means its norm $\mathbf{n}(q)$ is a prime number, we obtain that \mathcal{A}_φ is a commutative field isomorphic to $\mathbb{Z}_{n(\varphi)}$. We remark that

$$\begin{aligned} n(\varphi) &= \varphi \bar{\varphi} = (\alpha + \beta v)(\alpha + \beta \bar{v}) = \\ &= \alpha^2 + \alpha \beta \bar{v} + \alpha \beta v + \beta^2 v \bar{v} = \\ &= \alpha^2 + \alpha \beta + \beta^2. \end{aligned}$$

Example 4.10. 1) Let $q = -1 + 2v$. We obtain that $n(q) = 3$, therefore $\mathbb{Z}_3 \simeq \mathcal{A}_{-1+2v} = \{0, 1, v\}$, since we have

$$2 = (-1 + 2v)(-v) + v, \text{ with the norm } n(v) = 1 < 2.$$

2) Let $q = 1 + 2v$. We obtain that $n(q) = 7$, therefore $\mathbb{Z}_7 \simeq \mathcal{A}_{1+2v} = \{0, 1, v - 1, v, 1 + v, 2v - 1, 2v\}$.

We have:

$$2 = (1 + 2v)(1 - v) + v - 1, \text{ with the norm } n(v - 1) = 1 < 7.$$

$$3 = (1 + 2v)(1 - v) + v, \text{ with the norm } n(v) = 1 < 7.$$

$$4 = (1 + 2v)(1 - v) + 1 + v, \text{ with the norm } n(1 + v) = \frac{12}{4} = 3 < 7;$$

$$5 = (1 + 2v)(2 - 2v) + 2v - 1, \text{ with the norm } n(2v - 1) = 3 < 7;$$

$$6 = (1 + 2v)(2 - 2v) + 2v, \text{ with the norm } n(2v) = 4 < 7.$$

Remark 4.11. The isomorphism between $\mathbb{Z}_{n(q)}$ and \mathcal{A}_φ has direct applications in Coding Theory. For other details, see for example [GF; 10], [Hu; 94], [Ri; 95].

5. Quaternions with permutations

For $b = 1, c = 2$, we consider the quaternion algebra $\mathbb{H}_{\mathbb{Q}(i)}(-1, -2)$. From Proposition 3.7, this algebra is a split algebra.

Let S_4 be the set of all permutations of degree 4 on the set $\{n, n+1, n+2, n+3\}$, where n is a positive integer. We define a new element, namely, the σ -permutated n th Fibonacci-Hurtwitz quaternion

$$F_{n,\sigma} = \frac{1}{2}f_{\sigma(n)} + \frac{1}{2}f_{\sigma(n+1)}e_2 + \frac{1}{2}f_{\sigma(n+2)}e_2 + \frac{1}{2}f_{\sigma(n+3)}e_4.$$

Proposition 5.1. *Let $\mathbb{Q}(i)$ be the quadratic field where $i^2 = -1$. For the quaternion algebra $\mathbb{H}_{\mathbb{Q}(i)}(-1, -2)$ and for each positive integer n let $F_{n,\sigma}$ be the σ -permutated n th Fibonacci-Hurtwitz quaternion. Therefore, for $\sigma = \begin{pmatrix} n & n+1 & n+2 & n+3 \\ n+3 & n & n+1 & n+2 \end{pmatrix}$ or for*

$$\sigma \in \left\{ \begin{array}{l} \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n & n+3 & n+1 & n+2 \end{array} \right), \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n+3 & n & n+2 & n+1 \end{array} \right), \\ \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n & n+3 & n+2 & n+1 \end{array} \right) \end{array} \right\}$$

the norm of a such quaternion is

$$\mathbf{n}(F_{n,\sigma}) = f_{2n+3}.$$

Proof. For $\sigma = \begin{pmatrix} n & n+1 & n+2 & n+3 \\ n+3 & n & n+1 & n+2 \end{pmatrix}$, we have

$$\mathbf{n}(F_{n,\sigma}) = \frac{1}{4}(f_{n+3}^2 + f_n^2 + 2f_{n+1}^2 + 3f_{n+2}^2).$$

Using Proposition 2.1 i) and ii), we obtain

$$\mathbf{n}(F_{n,\sigma}) = \frac{1}{4}(4f_{n+1}^2 + 4f_{n+2}^2) = f_{n+1}^2 + f_{n+2}^2 = f_{2n+3}.$$

In the same way, for

$$\sigma \in \left\{ \begin{array}{l} \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n & n+3 & n+1 & n+2 \end{array} \right), \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n+3 & n & n+2 & n+1 \end{array} \right), \\ \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n & n+3 & n+2 & n+1 \end{array} \right) \end{array} \right\},$$

we obtain that $\mathbf{n}(F_{n,\sigma}) = f_{2n+3}$.

Corollary 5.2. *We consider $\mathbb{Q}(i)$ the quadratic field where $i^2 = -1$ and the quaternion algebra $\mathbb{H}_{\mathbb{Q}(i)}(-1, -2)$. Then, for $\sigma = \begin{pmatrix} n & n+1 & n+2 & n+3 \\ n+3 & n & n+1 & n+2 \end{pmatrix}$ or for*

$$\sigma \in \left\{ \begin{array}{l} \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n & n+3 & n+1 & n+2 \end{array} \right), \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n+3 & n & n+2 & n+1 \end{array} \right), \\ \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n & n+3 & n+2 & n+1 \end{array} \right) \end{array} \right\}$$

the following statements are true:

- i) All σ -permutated n th Fibonacci quaternions $F_{n,\sigma}$ are invertible;
- ii) There are not units σ -permutated n th Fibonacci quaternions $F_{n,\sigma}$.

Proof. i) Applying Proposition 5.1, for $\sigma = \begin{pmatrix} n & n+1 & n+2 & n+3 \\ n+3 & n & n+1 & n+2 \end{pmatrix}$ or for

$$\sigma \in \left\{ \begin{array}{l} \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n & n+3 & n+1 & n+2 \end{array} \right), \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n+3 & n & n+2 & n+1 \end{array} \right), \\ \left(\begin{array}{cccc} n & n+1 & n+2 & n+3 \\ n & n+3 & n+2 & n+1 \end{array} \right) \end{array} \right\},$$

we have $\mathbf{n}(F_{n,\sigma}) = f_{2n+3} \neq 0$, $(\forall)n \in \mathbb{N}$, therefore $F_{n,\sigma}$ are invertible for all $n \in \mathbb{N}$.

ii) The only Fibonacci numbers equals with 1 are f_1 and f_2 , then, by using Proposition 5.1, we have $\mathbf{n}(F_{n,\sigma}) = f_{2n+3} \neq 1$, for all $n \in \mathbb{N}$. It results that $F_{n,\sigma}$ is not a unit quaternion, for all $n \in \mathbb{N}$.

If n, k are two positive integers, from [Ra; 1917], p. 171, we get that there are $x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4 \in \mathbb{Z}$ such that $n = x_1^2 + x_2^2 + 2x_3^2 + 2x_4^2$ and $k = y_1^2 + y_2^2 + 2y_3^2 + 2y_4^2$. We want find $u_1, u_2, u_3, u_4 \in \mathbb{Z}$ such that $n \cdot k = u_1^2 + u_2^2 + 2u_3^2 + 2u_4^2$.

Proposition 5.3. *Let n, k be two positive integers, $n = x_1^2 + x_2^2 + 2x_3^2 + 2x_4^2$, $k = y_1^2 + y_2^2 + 2y_3^2 + 2y_4^2$, with $x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4 \in \mathbb{Z}$. Then*

$$n \cdot k = (x_1y_1 - x_2y_2 - 2x_3y_3 - 2x_4y_4)^2 + (x_1y_2 + x_2y_1 + 2x_3y_4 - 2x_4y_3)^2 +$$

$$+2(x_1y_3 + x_3y_1 - x_2y_4 + x_4y_2)^2 + 2(x_1y_4 + x_4y_1 + x_2y_3 - x_3y_2)^2.$$

Proof. We consider the quaternion algebra $\mathbb{H}_{\mathbb{Q}}(-1, -2)$ and $x = x_1 \cdot 1 + x_2e_2 + x_3e_3 + x_4e_4$, $y = y_1 \cdot 1 + y_2e_2 + y_3e_3 + y_4e_4 \in \mathbb{H}_{\mathbb{Q}}(-1, -2)$, where $x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4 \in \mathbb{Z}$. We have:

$$\begin{aligned} x \cdot y &= (x_1y_1 - x_2y_2 - 2x_3y_3 - 2x_4y_4) \cdot 1 + (x_1y_2 + x_2y_1 + 2x_3y_4 - 2x_4y_3) e_1 + \\ &\quad + (x_1y_3 + x_3y_1 - x_2y_4 + x_4y_2) e_2 + (x_1y_4 + x_4y_1 + x_2y_3 - x_3y_2) e_3. \end{aligned}$$

By using the fact that the norm is multiplicative, that means $\mathbf{n}(xy) = \mathbf{n}(x)\mathbf{n}(y)$, we obtain

$$\begin{aligned} n \cdot k &= (x_1y_1 - x_2y_2 - 2x_3y_3 - 2x_4y_4)^2 + (x_1y_2 + x_2y_1 + 2x_3y_4 - 2x_4y_3)^2 + \\ &\quad + 2(x_1y_3 + x_3y_1 - x_2y_4 + x_4y_2)^2 + 2(x_1y_4 + x_4y_1 + x_2y_3 - x_3y_2)^2. \end{aligned}$$

Since the numbers x_1, x_2, x_3, x_4 , obtained in the expression of n , are not unique and the numbers y_1, y_2, y_3, y_4 , obtained in the expression of k , are not unique, the representation of nk under the form $n \cdot k = u_1^2 + u_2^2 + 2u_3^2 + 2u_4^2$ is not unique.

Corollary 5.4. *Let n, k be two positive integers, $n = x_1^2 + x_2^2 + 2x_3^2 + 2x_4^2$, $k = y_1^2 + y_2^2 + 2y_3^2 + 2y_4^2$, with $x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4 \in \mathbb{Z}$. Then we have*

$$\begin{aligned} n \cdot k &= (x_1y_1 + x_2y_2 + 2x_3y_3 + 2x_4y_4)^2 + (x_1y_2 - x_2y_1 + 2x_3y_4 - 2x_4y_3)^2 + \\ &\quad + 2(x_1y_3 - x_3y_1 - x_2y_4 + x_4y_2)^2 + 2(x_1y_4 - x_4y_1 + x_2y_3 - x_3y_2)^2. \end{aligned}$$

Proof. By straightforward calculation.

Remark 5.5. The expression of nk obtained in Corollary 5.4 corresponds to the norm of the product of the following quaternions

$$x' = x_1 \cdot 1 - x_2e_2 - x_3e_3 + x_4e_4 \in \mathbb{H}_{\mathbb{Q}}(-1, -2),$$

and

$$y' = y_1 \cdot 1 + y_2e_2 + y_3e_3 - y_4e_4 \in \mathbb{H}_{\mathbb{Q}}(-1, -2),$$

where $x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4 \in \mathbb{Z}$.

From the above results, a question arise: if there is a kind of Fibonacci elements, possible similar with σ -permutated Fibonacci -Hurtwitz quaternions, such that the product of such elements is also an element of this type. In this regard, in the following, we provide a nice property involving the product of two σ -permutated Fibonacci -Hurtwitz quaternions. First of all, we give another property of the Fibonacci numbers, which we will use later in our proof.

Proposition 5.6. *Let $(f_n)_{n \geq 0}$ be the Fibonacci sequence. Then, the following property holds*

$$f_n f_l + f_{n+3} f_{l+3} = 2f_{n+l+3}, \text{ for all } n, l \in \mathbb{N}.$$

Proof. By using Binet's formula for Fibonacci sequence and Viète relations for α and β , we have

$$\begin{aligned}
f_n f_l + f_{n+3} f_{l+3} &= \frac{\alpha^n - \beta^n}{\alpha - \beta} \cdot \frac{\alpha^l - \beta^l}{\alpha - \beta} + \frac{\alpha^{n+3} - \beta^{n+3}}{\alpha - \beta} \cdot \frac{\alpha^{l+3} - \beta^{l+3}}{\alpha - \beta} = \\
&= \frac{\alpha^{n+l+6} + \beta^{n+l+6} - \alpha^{n+3} \beta^{l+3} - \alpha^{l+3} \beta^{n+3} + \alpha^{n+l} + \beta^{n+l} - \alpha^n \beta^l - \alpha^l \beta^n}{(\alpha - \beta)^2} = \\
&= \frac{\alpha^{n+l+3} (\alpha^3 + \alpha^{-3}) + \beta^{n+l+3} (\beta^3 + \beta^{-3}) - \alpha^n \beta^l (\alpha^3 \beta^3 + 1) - \alpha^l \beta^n (\alpha^3 \beta^3 + 1)}{(\alpha - \beta)^2} = \\
&= \frac{\alpha^{n+l+3} (\alpha^3 - \beta^3) + \beta^{n+l+3} (\beta^3 - \alpha^3)}{(\alpha - \beta)^2} = \frac{(\alpha^3 - \beta^3) (\alpha^{n+l+3} - \beta^{n+l+3})}{(\alpha - \beta)^2} = \\
&= \frac{(\alpha^2 + \alpha\beta + \beta^2) (\alpha^{n+l+3} - \beta^{n+l+3})}{\alpha - \beta} = 2f_{n+l+3}.
\end{aligned}$$

Proposition 5.7. Let $(f_n)_{n \geq 0}$ be the Fibonacci sequence, let $\sigma = \begin{pmatrix} n & n+1 & n+2 & n+3 \\ n+3 & n & n+1 & n+2 \end{pmatrix}$ and, for each positive integers $n < l$, let $F'_{n,\sigma}$ and $F''_{l,\sigma}$ be the σ -special permuted Fibonacci -Hurtwitz quaternions: $F'_{n,\sigma} = \frac{1}{2}f_{\sigma(n)} - \frac{1}{2}f_{\sigma(n+1)}e_2 - \frac{1}{2}f_{\sigma(n+2)}e_3 + \frac{1}{2}f_{\sigma(n+3)}e_4$, $F''_{l,\sigma} = \frac{1}{2}f_{\sigma(l)} + \frac{1}{2}f_{\sigma(l+1)}e_2 + \frac{1}{2}f_{\sigma(l+2)}e_3 - \frac{1}{2}f_{\sigma(l+3)}e_4$. Let $k = l - n$. Therefore, the expression

$$F'_{n,\sigma} F''_{l,\sigma} - \frac{\mathbf{t}(F'_{n,\sigma} F''_{l,\sigma})}{2}$$

depends only of k .

Proof. For $F'_{n,\sigma} = \frac{1}{2}f_{n+3} - \frac{1}{2}f_n e_2 - \frac{1}{2}f_{n+1} e_3 + \frac{1}{2}f_{n+2} e_4$ and $F''_{l,\sigma} = \frac{1}{2}f_{l+3} + \frac{1}{2}f_l e_2 + \frac{1}{2}f_{l+1} e_3 - \frac{1}{2}f_{l+2} e_4$ we apply Corollary 4.4 and Remark 4.5 and we have

$$\begin{aligned}
F'_{n,\sigma} F''_{l,\sigma} &= \frac{1}{4} (f_{n+3} f_{l+3} + f_n f_l + 2f_{n+1} f_{l+1} + 2f_{n+2} f_{l+2}) + \\
&+ \frac{1}{4} (f_{n+3} f_l - f_n f_{l+3} + 2f_{n+1} f_{l+2} - 2f_{n+2} f_{l+1}) e_2 + \\
&+ \frac{1}{4} (f_{n+3} f_{l+1} - f_{n+1} f_{l+3} - f_n f_{l+2} + f_{n+2} f_l) e_3 + \\
&+ \frac{1}{4} (-f_{n+3} f_{l+2} + f_{n+2} f_{l+3} - f_n f_{l+1} + f_{n+1} f_l) e_4.
\end{aligned}$$

From the properties of Fibonacci numbers and Remark 5.6., we get

$$F'_{n,\sigma} F''_{l,\sigma} = \frac{1}{4} (2f_{n+l+3} + 2f_{n+l+3}) + \frac{1}{4} [(-1)^{n-3} f_3 f_{l-n} + 2(-1)^n f_1 f_{l-n+2}] e_2 +$$

$$\begin{aligned}
& +\frac{1}{4} \left[(-1)^{n+1} f_2 f_{l-n} + (-1)^n f_1 f_{l-n} \right] e_3 + \frac{1}{4} \left[(-1)^{n+2} f_1 f_{l-n} + (-1)^n f_1 f_{l-n} \right] e_4 \\
& = f_{n+l+3} + \frac{1}{2} (-1)^n f_{l-n+1} e_2 + \frac{1}{2} (-1)^n f_{l-n} e_4.
\end{aligned}$$

It results that $\mathbf{t}(F'_{n,\sigma} F''_{l,\sigma}) = 2f_{n+l+3}$. Therefore, we obtain that

$$F'_{n,\sigma} F''_{l,\sigma} - \frac{\mathbf{t}(F'_{n,\sigma} F''_{l,\sigma})}{2} = \frac{1}{2} (-1)^n f_{k+1} e_2 + \frac{1}{2} (-1)^n f_k e_4.$$

Conclusions. In this paper we have found an interesting connection between quaternion algebras and some quaternary quadratic forms, arising from some results obtained by Ramanujan in [Ra; 1917], we have define a monoid structure over a finite set on which we will prove that the defined Fibonacci sequence is stationary, we have proved some results regarding the arithmetic of integer quaternions defined on some special division quaternion algebras and we have defined and gave properties of some special quaternions by using Fibonacci sequences.

References

- [Co, Sm; 03] J.H. Conway, D.A. Smith, *On Quaternions and Octonions*, A.K. Peters, Natick, Massachusetts, 2003.
- [Da, Sa, Va; 03] G. Davidoff, P. Sarnak, A. Valette, *Elementary Number Theory, Group Theory, and Ramanujan Graphs*, Cambridge University Press, 2003.
- [Fib.] <http://www.maths.surrey.ac.uk/hosted-sites/R.Knott/Fibonacci/fib.html>
- [Fl; 20] C. Flaut, Some applications of MV algebras, <https://arxiv.org/pdf/2001.03971.pdf>
- [Fl, Sa; 15] C. Flaut, D. Savin, *Quaternion Algebras and Generalized Fibonacci-Lucas Quaternions*, Adv. Appl. Clifford Algebras, 25(4)(2015), 853-862.
- [FSZ; 20] C. Flaut, D. Savin, G. Zaharia, *Properties and applications of some special integer number sequences*, accepted in Mathematical Methods in the Applied Sciences, DOI: 10.1002/mma.6257 (<https://onlinelibrary.wiley.com/doi/abs/10.1002/mma.6257>)
- [Fl, Sh; 13] C. Flaut, V. Shpakivskyi, *On Generalized Fibonacci Quaternions and Fibonacci-Narayana Quaternions*, Adv. Appl. Clifford Algebras, 23(3)(2013), 673-688.
- [GF; 10] F. Ghaboussi, J. Freudenberger, *Codes over Gaussian integer rings*, 18th Telecommunications forum TELFOR 2010, 662-665.
- [Gi, Sz; 06] P. Gille, T., Szamuely, *Central Simple Algebras and Galois Cohomology*, Cambridge University Press, 2006.

- [Gu; 13] M. Güzeltepe, *Codes over Hurwitz integers*, Discrete Math., **313(5)(2013)**, 704-714.
- [Ho; 63] A. F. Horadam, *Complex Fibonacci Numbers and Fibonacci Quaternions*, Amer. Math. Monthly, 70(1963), 289–291.
- [Hu; 94] K. Huber, *Codes over Gaussian integers*, IEEE Trans. Inform. Theory, 40(1994), 207–216.
- [Ko] D. R. Kohel, *Quaternion algebras*, available online at <http://www.i2m.univ-amu.fr/perso/david.kohel/alg/doc/AlgQuat.pdf>.
- [Ma; 07] K. Martin, Classification of quaternion algebras, <http://www2.math.ou.edu/~kmartin/quaint/ch5.pdf>, 2017.
- [Mag] <http://magma.maths.usyd.edu.au/magma/handbook/>
- [Ra; 1917] S. Ramanujan, *On the expression of a number in the form $ax^2 + by^2 + cz^2 + du^2$* , Proc. Cambridge Philos. Soc. 19(1917), 11-21. Collected Papers, AMS Chelsea Publ., Providence, RI, 2000, 169-178.
- [Ri; 95] J. Rifà, *Groups of Complex Integer Used as QAM Signals*, IEEE Transactions on Information Theory, **41(5)(1995)**, 1512-1517.
- [Sa; 14] D. Savin, *About some split central simple algebras*, An. St. Univ. Ovidius Constanta Mat. Ser. 22(1) (2014), 263–272.
- [Sa; 17] D. Savin, *About special elements in quaternion algebras over finite fields* Adv. Appl. Clifford Algebras 27(2) (2017), 1801–1813.
- [Sa; 19] D. Savin, *Special numbers, special quaternions and special symbol elements*, chapter in the book Models and Theories in Social Systems, vol. 179, Springer 2019, ISBN-978-3-030-00083-7, p. 417-430.
- [Si; 70] W. Sierpinski, *250 Problems in elementary number theory*, American Elsevier Pub. Co (1970).
- [Vi; 80] M. F. Vigneras, *Arithmetique des algebres de quaternions*, Lecture Notes in Math., vol. 800, Springer, Berlin, 1980.

Cristina FLAUT

Faculty of Mathematics and Computer Science,
 Ovidius University of Constanța, România,
 Bd. Mamaia 124, 900527,
<http://www.univ-ovidius.ro/math/>
 e-mail: cflaut@univ-ovidius.ro; cristina_flaut@yahoo.com

Diana SAVIN

Faculty of Mathematics and Computer Science,
 Ovidius University of Constanța, România,
 Bd. Mamaia 124, 900527,
<http://www.univ-ovidius.ro/math/>
 e-mail: savin.diana@univ-ovidius.ro, dianet72@yahoo.com