

A note on a Bonnet-Myers type diameter bound for graphs with positive entropic Ricci curvature

S. Kamtue¹

¹Department of Mathematical Sciences, Durham University

March 4, 2020

Abstract

An equivalent definition of entropic Ricci curvature on discrete spaces was given in terms of the global gradient estimate in [1, Theorem 3.1]. With a particular choice of the density function ρ , we obtain a localized gradient estimate, which in turns allow us to apply the same technique as in [4] to derive a Bonnet-Myers type diameter bound for graphs with positive entropic Ricci curvature. However, the case of the hypercubes indicates that the bound may be not optimal (where θ is chosen to be logarithmic mean by default). If θ is arithmetic mean, the Bakry-Émery criterion can be recovered and the diameter bound is optimal as it can be attained by the hypercubes.

1 Introduction

The notion of entropic Ricci curvature on discrete spaces (i.e., Markov chains, or graphs) was introduced by Erbar and Maas [2] inspired by work of Sturm [10] and Lott and Villani [6] which describes the lower bound of Ricci curvature via the displacement convexity of the entropy functional. Equivalent definitions of Entropic curvature are also given in terms of Bochner's inequality [2] and in terms of gradient estimates [1]. Detailed notations and definitions are given in Section 2 and proofs are presented in Section 3.

Definition 1.1 (Entropic curvature). An irreducible and reversible Markov kernel Q on a finite discrete space X (with a steady state π) **has entropic curvature at least** $\kappa \in \mathbb{R}$, written as $\text{Ric}(Q) \geq \kappa$, if and only if the entropy functional \mathcal{H} satisfies the following κ -convexity inequality:

$$\mathcal{H}(\rho_t) \leq (1-t)\mathcal{H}(\rho_0) + t\mathcal{H}(\rho_1) - \frac{\kappa}{2}t(1-t)\mathcal{W}(\rho_0, \rho_1)^2 \quad (1.1)$$

for every constant speed geodesic $(\rho_t)_{t \in [0,1]}$ in the (modified) L^2 -Wasserstein space of probability densities $(\mathcal{D}(X), \mathcal{W})$.

Theorem 1.1 (Equivalent definitions). *For a finite set X equipped with an irreducible and reversible Markov kernel Q (and the steady state π), the following properties are equivalent:*

- (1) $\text{Ric}(Q) \geq \kappa$;

(2) Bochner's inequality holds for all $\rho \in \mathcal{D}(X)$ and $f \in C(X)$:

$$\mathcal{B}(\rho, f) \geq \kappa \mathcal{A}(\rho, f), \quad (1.2)$$

where

$$\mathcal{A}(\rho, f) := \langle \hat{\rho} \cdot \nabla f, \nabla f \rangle_\pi,$$

and

$$\mathcal{B}(\rho, f) := \frac{1}{2} \langle \hat{\Delta} \rho \cdot \nabla f, \nabla f \rangle_\pi - \langle \hat{\rho} \cdot \nabla f, \nabla \Delta f \rangle_\pi,$$

where

$$\hat{\rho}(x, y) := \theta(\rho(x), \rho(y)),$$

$$\hat{\Delta} \rho := \partial_1 \theta(\rho(x), \rho(y)) \Delta \rho(x) + \partial_2 \theta(\rho(x), \rho(y)) \Delta \rho(y),$$

and θ is a logarithmic mean: $\theta(a, b) = \frac{a-b}{\log a - \log b}$;

(3) Global gradient estimate holds for all $\rho \in \mathcal{D}(X)$ and $f \in C(X)$ and $t \geq 0$,

$$|\nabla P_t f|_\rho^2 \leq e^{-2\kappa t} |\nabla f|_{P_t \rho}^2,$$

or more explicitly

$$\frac{1}{2} \sum_{u,v} (P_t f(v) - P_t f(u))^2 \hat{\rho}(u, v) Q(u, v) \pi(u) \leq e^{-2\kappa t} \frac{1}{2} \sum_{u,v} (f(v) - f(u))^2 \widehat{P_t \rho}(u, v) Q(u, v) \pi(u) \quad (1.3)$$

where $P_t = e^{t\Delta}$ denotes the heat semigroup.

The contribution of this paper is to follow ideas from [1] to derive a similar local gradient estimate, and to apply the same technique as in [4] to prove a Bonnet-Myers type diameter bound on the underlying graph X (when Q is a simple random walk) in the case of a positive lower bound of entropic curvature.

Theorem 1.2 (Local gradient estimate). *If $\text{Ric}(Q) \geq \kappa$, then for all $x \in X$ and $f \in C(X)$ and all $\varepsilon \geq 0$ we have*

$$\Gamma(P_t f)(x) \pi(x) \leq \frac{e^{-2\kappa t}}{2\theta(1, \varepsilon)} [P_t \Gamma(f)(x) \pi(x) + \varepsilon \sum_{y \in N(x)} P_t \Gamma(f)(y) \pi(y)]. \quad (1.4)$$

where $\Gamma(f)(x) := \sum_y (f(y) - f(x))^2 Q(x, y)$.

Theorem 1.3 (Diameter bound). *Let Q represent a simple random walk on X with strictly positive entropic Ricci curvature $\text{Ric}(Q) \geq \kappa > 0$. Then the diameter is bounded from above by $\text{diam}(X) \leq \frac{2}{\kappa} \sqrt{\frac{D \log D}{D-1}}$ where D is the maximal (vertex) degree.*

Remark 1.4. It is known that the entropic curvature of (a simple random walk on) the discrete hypercube \mathcal{Q}^n is $\frac{2}{n}$ (see [2, Example 5.7]). Therefore, in view of the hypercubes, the bound in Theorem 1.3 is not optimal:

$$n = \text{diam}(\mathcal{Q}^n) \leq n \sqrt{\frac{n}{n-1} \log n}.$$

However, if we replace the logarithmic mean by the arithmetic mean for θ in Bochner's formula (1.2) and inequality (1.3), the local gradient estimate in Theorem 1.2 would imply (by taking $\varepsilon = 0$) the Bakry-Émery curvature criterion $CD(\kappa, \infty)$: $\Gamma(P_t f)(x) \leq e^{-2\kappa t} P_t \Gamma(f)(x)$ in the sense of [3, Corollary 3.3]. This implication has already been mentioned in the survey [8]. Consequently, we obtain a diameter bound: $\text{diam}(X) \leq \frac{2}{\kappa}$, which is sharp and the equality is attained if and only if X is a hypercube \mathcal{Q}^n (for details see [4] and [5]).

2 Setup and notations

2.1 Notions associated to a discrete Markov chain

We start with a Markov chain (X, Q) , where X is a finite set and $Q : X \times X \rightarrow \mathbb{R}^+ \cup \{0\}$ is a Markov kernel Q , i.e. $\sum_{y \in X} Q(x, y) = 1$ for all $x \in X$. Furthermore, we assume that Q is irreducible and reversible, which implies that there exists a unique stationary probability measure π on X satisfying $\sum_{x \in X} \pi(x) = 1$ and the detail balanced equations:

$$Q(x, y)\pi(x) = Q(y, x)\pi(y) \quad \forall x, y \in X.$$

The set of probability densities (with respect to π) is defined as

$$\mathcal{D}(X) := \left\{ \rho : X \rightarrow \mathbb{R}^+ \cup \{0\} \mid \sum_{x \in X} \pi(x)\rho(x) = 1 \right\}.$$

The entropy functional, defined on $\mathcal{D}(X)$, is given by

$$\mathcal{H}(\rho) := \sum_{x \in X} \rho(x) \log \rho(x) \pi(x).$$

The *discrete gradient* $\nabla : C(X) \rightarrow C(X \times X)$, *discrete divergence* $\nabla \cdot : C(X \times X) \rightarrow C(X)$, and *laplacian* $\Delta := \nabla \cdot \nabla$ are defined as follows:

Definition 2.1. For all $f, g \in C(X)$ and $U, V \in C(X \times X)$,

$$\begin{aligned} \nabla f(x, y) &:= f(y) - f(x), \\ \nabla \cdot V(x) &:= \frac{1}{2} \sum_{y \in X} (V(x, y) - V(y, x)) Q(x, y), \\ \Delta f(x) &:= \sum_{y \in X} (f(y) - f(x)) Q(x, y). \end{aligned}$$

The inner products are defined as

$$\begin{aligned} \langle f, g \rangle_\pi &:= \sum_{x \in X} f(x)g(x)\pi(x), \\ \langle U, V \rangle_\pi &:= \frac{1}{2} \sum_{x, y \in X} U(x, y)V(x, y)Q(x, y)\pi(x), \end{aligned}$$

and for all $\rho \in \mathcal{D}(X)$,

$$\langle U, V \rangle_\rho := \langle \hat{\rho} \cdot U, V \rangle_\pi = \frac{1}{2} \sum_{x, y \in X} U(x, y) V(x, y) \hat{\rho}(x, y) Q(x, y) \pi(x),$$

where $\hat{\rho}(x, y) := \theta(\rho(x), \rho(y))$ and θ is a *suitable mean* satisfying Assumption 2.1 in [2]. By default, θ is chosen to be the logarithmic mean: $\theta(a, b) = \frac{a-b}{\log a - \log b}$.

Furthermore, we already introduced

$$\begin{aligned} \mathcal{A}(\rho, f) &:= |\nabla f|_\rho^2 = \langle \hat{\rho} \cdot \nabla f, \nabla f \rangle_\pi, \\ \mathcal{B}(\rho, f) &:= \frac{1}{2} \langle \hat{\Delta} \rho \cdot \nabla f, \nabla f \rangle_\pi - \langle \hat{\rho} \cdot \nabla f, \nabla \Delta f \rangle_\pi, \end{aligned}$$

where $\hat{\Delta} \rho := \partial_1 \theta(\rho(x), \rho(y)) \Delta \rho(x) + \partial_2 \theta(\rho(x), \rho(y)) \Delta \rho(y)$.

Definition 2.2 (Discrete transport metric). For $\bar{\rho}_0, \bar{\rho}_1 \in \mathcal{D}(X)$,

$$\mathcal{W}(\bar{\rho}_0, \bar{\rho}_1) := \inf \left\{ \int_0^1 \mathcal{A}(\rho_t, f_t) dt \mid (\rho_t, f_t) \in CE(\bar{\rho}_0, \bar{\rho}_1) \right\}^{\frac{1}{2}},$$

where the infimum is taken over the set $CE(\rho_0, \rho_1)$ which consists of all sufficiently regular curves $(\rho_t)_{t \in [0,1]}$ on $\mathcal{D}(X)$ and $(f_t)_{t \in [0,1]}$ on $C(X)$ which satisfy the *continuity equation* $\partial_t \rho_t + \nabla \cdot (\hat{\rho}_t \nabla f_t) = 0$ and $\rho_0 = \bar{\rho}_0, \rho_1 = \bar{\rho}_1$. It was shown that \mathcal{W} is a metric on $\mathcal{D}(X)$. We refer to [2, 7] for further details. Note that this notion of the \mathcal{W} -metric is relevant in Definition 1.1 of entropic curvature.

2.2 Graph theoretical notions

The kernel Q induces a graph structure on X by assigning an edge $x \sim y$ if and only if $Q(x, y) > 0$. Note that the graph is connected and undirected, due to irreducibility and reversibility of Q , respectively. For a vertex $x \in X$, denote $N(x) := \{y \in X \mid x \sim y\}$ the set of neighbors of x , and $d_x := |N(x)|$ the degree of x , and $D := \max_{x \in X} d_x$ the maximal degree. The graph is equipped with the usual *combinatorial distance function* d where $d(x, y)$ is the length of shortest path(s) between x and y , and the diameter of the graph is defined as $\text{diam}(X) = \max_{x, y \in X} d(x, y)$. In this note, we restrict Q to be a simple random walk, which is given by $Q(x, y) = \frac{1}{d_x}$ for all $y \in N(x)$, and $\pi(x) = d_x / (\sum_{v \in X} d_v)$.

3 Proofs

Proof of Theorem 1.1. Equivalence (1) \Leftrightarrow (2) was stated in [2, Theorem 4.5] under the assumption that θ is the logarithmic mean. Equivalence (2) \Leftrightarrow (3) was stated in [1, Theorem 3.1], regardless of the choice of a ‘suitable’ mean for θ . \square

Proof of Theorem 1.2.

The proof follows ideas of [1, Corollary 3.4]. Note that the inequality (1.3) is homogeneous in ρ . Therefore, we can drop the requirement that ρ is a probability density: $\sum_x \rho(x) \pi(x) =$

1. We localize inequality (1.3) by choosing $\rho = \mathbf{1}_x + \varepsilon \sum_{y \in N(x)} \mathbf{1}_y$ for a fixed $x \in V$ and a parameter $\varepsilon \in [0, \infty)$.

In particular we know that $\hat{\rho}(x, y) = \hat{\rho}(y, x) = \theta(1, \varepsilon)$ for all $y \in N(x)$. The left-hand-side of (1.3) has the following lower bound

$$L.H.S. \geq \sum_{y \in N(x)} (P_t f(y) - P_t f(x))^2 \theta(1, \varepsilon) Q(x, y) \pi(x) = \theta(1, \varepsilon) \cdot \Gamma(P_t f)(x) \pi(x) \quad (3.1)$$

On the other hand, we have the following bound on the right-hand-side of (1.3) by

$$\begin{aligned} R.H.S. &\leq e^{-2\kappa t} \frac{1}{2} \sum_{u, v} (f(v) - f(u))^2 \frac{P_t \rho(u) + P_t \rho(v)}{2} Q(u, v) \pi(u) \\ &= e^{-2\kappa t} \frac{1}{2} \sum_{u, v} (f(v) - f(u))^2 P_t \rho(u) Q(u, v) \pi(u) \end{aligned} \quad (3.2)$$

due to $\theta(s, t) \leq (s + t)/2$ and symmetry from interchanging u and v .

We now apply the heat kernel $p_t(\cdot, \cdot)$ given by $P_t g(u) = \sum_z p_t(u, z) g(z) \pi(z)$ for every function g . With our chosen ρ , we obtain $P_t \rho(u) = p_t(u, x) \rho(x) \pi(x) + \varepsilon \sum_{y \in N(x)} p_t(u, y) \rho(y) \pi(y)$, which we substitute into (3.2) and use the symmetry of heat kernel: $p_t(u, v) = p_t(v, u)$ to derive

$$\begin{aligned} R.H.S. &\leq \frac{e^{-2\kappa t}}{2} \left[\pi(x) \sum_u p_t(u, x) \pi(u) \sum_v (f(v) - f(u))^2 Q(u, v) + \right. \\ &\quad \left. \varepsilon \sum_{y \in N(x)} \pi(y) \sum_u p_t(u, y) \pi(u) \left(\sum_v (f(v) - f(u))^2 Q(u, v) \right) \right] \\ &= \frac{e^{-2\kappa t}}{2} \left[\pi(x) P_t \Gamma(f)(x) + \varepsilon \sum_{y \in N(x)} \pi(y) P_t \Gamma(f)(y) \right]. \end{aligned} \quad (3.3)$$

The desired inequality then follows from (3.1) and (3.3). \square

For the underlying graph X with Q representing a simple random walk, we have the following corollary as an immediate consequence of Theorem 1.2.

Corollary 3.1. *Let Q represent a simple random walk on X with entropic Ricci curvature $\text{Ric}(Q) \geq \kappa$. Then we have the following gradient estimate:*

$$\Gamma(P_t f)(x) \leq c \cdot e^{-2\kappa t} \|P_t \Gamma(f)\|_\infty \quad (3.4)$$

where $c := \frac{D \log D}{D-1}$ and D is the maximal degree.

Proof of Corollary 3.1. Theorem 1.2 implies that $\Gamma(P_t f)(x) \leq c_{\varepsilon, x} \cdot e^{-2\kappa t} \|P_t \Gamma(f)\|_\infty$.

where $c_{\varepsilon, x} := \frac{1 + \varepsilon \sum_{y \in N(x)} \frac{d_y}{d_x}}{2\theta(1, \varepsilon)} \leq \frac{1 + \varepsilon D}{2\theta(1, \varepsilon)}$. In particular when $\varepsilon = \frac{1}{D}$, we have $c_{\varepsilon, x} \leq c$. \square

Finally, we present the proof of the diameter bound in the case of strictly positive entropic curvature.

Proof of Theorem 1.3.

The proof follows ideas of [4, Theorem 2.1]. Consider a particular choice of function $f \in C(X)$ given by $f(x) = d(x, x_0)$ for an arbitrary reference point $x_0 \in X$. Since f is a Lipschitz function with constant 1, it follows that $\Gamma(f)(x) = \sum_{y \in N(x)} \frac{1}{d_x} (f(y) - f(x))^2 \leq 1$ for all $x \in X$, i.e., $\|\Gamma(f)\|_\infty \leq 1$, which then implies $\|P_t \Gamma(f)\|_\infty \leq \|\Gamma(f)\|_\infty \leq 1$.

Moreover, Cauchy-Schwartz and inequality (3.4) give

$$|\Delta P_t f(x)|^2 \leq \frac{1}{d_x} \sum_{y \in N(x)} (P_t f(y) - P_t f(x))^2 = \Gamma(P_t f)(x) \leq e^{-2\kappa t} c \|P_t \Gamma(f)\|_\infty \leq c e^{-2\kappa t}. \quad (3.5)$$

From the fundamental theorem of calculus and the definition of P_t , we then obtain

$$|f(x) - P_T f(x)| \leq \int_0^T \left| \frac{\partial}{\partial t} P_t f(x) \right| dt = \int_0^T |\Delta P_t f(x)| dt \leq \int_0^T \sqrt{c} e^{-\kappa t} dt \leq \frac{\sqrt{c}}{\kappa},$$

which holds true for all $T > 0$.

Moreover, the chain of inequalities in (3.5) implies that $|P_t f(y) - P_t f(x)|^2 \leq d_x \cdot c e^{-2\kappa t} \rightarrow 0$ as $t \rightarrow \infty$ for all neighbors $y \sim x$. Therefore, $|P_t f(y) - P_t f(x)| \rightarrow 0$ as $t \rightarrow \infty$ for an arbitrary pair of x, y (by considering a connected path from x to y).

Passing to the limit $T \rightarrow \infty$, we can conclude from triangle inequality that

$$d(x, x_0) = |f(x) - f(x_0)| \leq |f(x) - P_T f(x)| + |f(x_0) - P_T f(x_0)| + |P_T f(x) - P_T f(x_0)| \leq \frac{2\sqrt{c}}{\kappa}.$$

Since x, x_0 are arbitrary, we obtain the desired diameter bound. \square

References

- [1] M. Erbar and M. Fathi, *Poincaré, modified logarithmic Sobolev and isoperimetric inequalities for Markov chains with non-negative Ricci curvature*, J. Funct. Anal. **274**(11) (2018), 3056–3089.
- [2] M. Erbar and J. Maas, *Ricci Curvature of Finite Markov Chains via Convexity of the Entropy*, Arch. Rational Mech. Anal. **206** (2012) 997–1038
- [3] S. Liu and Y. Lin, *Equivalent Properties of CD Inequality on Graph*, arXiv:1512.02677
- [4] S. Liu, F. Münch and N. Peyerimhoff, *Bakry–Émery curvature and diameter bounds on graphs*, Calc. Var. Partial Differential Equations **57**(2) (2018), Art. 67, 9.
- [5] S. Liu, F. Münch and N. Peyerimhoff, *Rigidity properties of the hypercube via Bakry–Émery curvature*, arXiv:1705.06789.
- [6] L. Lott and C. Villani, *Ricci curvature for metric-measure spaces via optimal transport*, Ann. Math. (2) **169**(3) (2009), 903–991
- [7] J. Maas, *Gradient flows of the entropy for finite Markov chains*, J. Funct. Anal. **261**(8) (2011), 2250–2292.

- [8] J. Maas, *Entropic Ricci Curvature for Discrete Spaces*, Modern Approaches to Discrete Curvature, Lecture Notes in Mathematics, vol 2184 (2017), L. Najman and P. Romon (eds), Springer, Cham.
- [9] S. B. Myers, *Riemannian manifolds with positive mean curvature*, Duke Math. J. **8** (1941), 401–404.
- [10] K.-Th. Sturm, *On the geometry of metric measure spaces I and II*, Acta Math. **196**(1) (2006), 65–177.